

(19)



(11)

EP 2 143 436 A1

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art. 153(4) EPC

(43) Date of publication:

13.01.2010 Bulletin 2010/02

(51) Int Cl.:

A61K 38/17 (2006.01)

A61P 31/00 (2006.01)

(21) Application number: **08750412.2**

(86) International application number:

PCT/ES2008/000177

(22) Date of filing: **27.03.2008**

(87) International publication number:

WO 2008/119851 (09.10.2008 Gazette 2008/41)

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT
RO SE SI SK TR**

(72) Inventors:

- **SARRIAS FORNÉS, María Rosa**
E-08036 Barcelona (ES)
- **LOZANO SOTO, Francisco**
E-08029 Barcelona (ES)

(30) Priority: **28.03.2007 ES 200700893**

(71) Applicant: **Universidad de Barcelona**
08028 Barcelona (ES)

(74) Representative: **Jané Bonet, Montserrat et al**
Zea, Barlocchi & Markvardsen, S.L.
Plaza Cataluña no. 1
08002 Barcelona (ES)

(54) **PROTEIN PRODUCT FOR TREATMENT OF INFECTIOUS DISEASES AND RELATED INFLAMMATORY PROCESSES**

(57) The inventors have found that CD6, a member of the Scavenger Receptor Cysteine-Rich (SRSR) superfamily expressed on human lymphocytes binds to Gram-positive and Gram-negative bacteria, as well as to other microbial structures. Thus, a CD6 product is useful for the manufacture of a medicament for therapeutic and/or preventive treatment of an infectious disease or

of an inflammatory condition related to an infectious disease or to the presence of a product derived from an infectious agent in a mammal including a human. Examples of such inflammatory conditions are systemic inflammatory response syndrome (SIRS), sepsis, severe sepsis and septic shock.

EP 2 143 436 A1

Description

[0001] This invention relates to the field of medicine, and specifically to compounds of protein nature for the manufacture of medicaments for therapeutic and/or preventive treatment of infectious diseases and of inflammatory conditions related thereto.

BACKGROUND ART

[0002] Systemic microbial infection and release of microbial products induces the hyperactivation of host immune cells, subsequently generating an exacerbated inflammatory response. Sepsis may result in the inability of the immune system to control this inflammation, and can end in tissue injury, increased vascular permeability and it can cause multi-organ failure and shock (septic shock).

[0003] Sepsis can result from many causes but is typically triggered by pneumonia, trauma, surgery, and burns or by conditions such as cancer or AIDS. Sepsis usually begins with tremor, fever, falling blood pressure (septic shock), rapid breathing, rapid heart rate, and skin lesions. Within hours, sepsis may cause spontaneous clotting in blood vessels, severe hypotension, multiple organ failure, shock, gangrene and eventually death. Sepsis causes high morbidity and mortality in humans and other animals (mortality of up to 70% in septic patients). In the United States and Europe, 1.5 million people develop sepsis annually. 30% of these patients die after one month and a 20% after six months. In the United States, sepsis is the 10th cause of death, which represents a mortality higher than those caused by infarct, breast cancer or lung cancer.

[0004] The most important intervention in sepsis is quick diagnosis and treatment. Diagnosing sepsis can be difficult. Some of its symptoms, such as fever, rapid pulse, and respiratory difficulty occur frequently and can be confused as being due to other disorders. Patients diagnosed with severe sepsis are usually placed in the intensive care unit (ICU) of the hospital for special treatment. The first line of treatment is to identify and eliminate the underlying infection with anti-infection agents or surgery to drain the site of infection.

[0005] Current methods for treating sepsis include antibacterials, antibodies, peptides, and a recombinant human activated protein C named drotrecogin alpha, marketed by Eli Lilly as Xigris®. However, drotrecogin alpha only reduces the mortality associated with severe sepsis in a 5%, and not all the patients respond positively to this drug. Steroids have also been recently shown to be valuable in patients with septic shock. The doctor also administers intravenous fluids to prevent blood pressure from dropping too much. In some cases, vasopressor medications (which constrict blood vessels) are needed to achieve an adequate blood pressure. Finally, if organ failures occur, appropriate supportive care is provided (e.g., dialysis for kidney failure, mechanical ventilation for respiratory failure, etc.).

[0006] Due to the high level of redundancy regarding molecular mediators in the sepsis response, new approaches are likely to focus on intervening at multiple points in the sepsis cascade. Some drug candidates in early phase of development are a Triggering Receptor Expressed on Myeloid cells-1 (TREM-1) receptor antagonist (Merck & Co Inc and BioXell SpA); a super-antigen antagonist (Atox Bio Ltd.), a short peptide which blocks the action of a family of deadly bacterial toxins produced by Staphylococcus aureus and Streptococcus pyogenes, termed as super-antigens; Immune Regulating Hormone (IRH, Hollis-Eden Pharmaceuticals Inc.), an autoimmune and anti-inflammatory drug which controls immune system and metabolic functions; and an Adenosine A1 receptor antagonist as a treatment for Gram-negative septicemia (Endacea Inc.). Other molecules under development are Toll-like Receptor-4 antagonists (Takeda and Eisai); anti-TNF- α polyclonal antibody fragment (Protherics); bovine intestine-derived alkaline phosphatase (AM-Pharma); Norathiol (Medinor), which neutralizes nitric oxide; and transgenic antithrombin III ATryn® (GTC Biotherapeutics) which received marketing approval from European regulatory authorities in 2006 and it is in late-stage clinical trials in the United States.

[0007] Other approaches have been proposed for treating sepsis, such as: anti-IL-8 antibodies (US Patent application 2003002178A), anti-IL-18 antibodies (US 20030008822A), anti-C5a antibodies and C-terminal truncated C5a peptides (US 20020165138A), chemokines and chemokine fragments (US 20020155094A), a combination of protein C and BPI antibodies (US 20020044929A), COX-2 inhibitors (US 20020006915A), algae lipopolysaccharides (US 6534648) and using an antibody to TNF- α and an antibody to bacterial lipopolysaccharide (US 6315999). However, despite the major advances of the past several decades in the treatment of serious infections, the incidence of sepsis and mortality due to sepsis continue to increase. Therefore, it seems desirable to provide new methods and compositions for the prevention and treatment of infectious diseases and of inflammatory conditions related to these infectious diseases.

SUMMARY OF THE INVENTION

[0008] Inventors have found that the ectodomain of human CD6, a cell surface receptor mainly expressed by cells of the lymphoid lineage, is able to bind to conserved microbial structures such as lipopolysaccharide (LPS) and lipoteichoic acid (LTA) from Gram-negative and Gram-positive bacteria, respectively. Surprisingly, the LPS-CD6 interaction show

a relative high affinity similar in magnitude to that reported for the interaction of LPS with CD14, the most important LPS receptor in mammalian cells. CD6 binds to Re-LPS about 10-fold more tightly than Sp α , another receptor with Scavenger Receptor Cysteine-Rich (SRCR) domains. These data lead to the finding that i.p. administration of a recombinant form of CD6 abolishes the lethal effects caused by LPS-induced septic shock in mice. Accordingly, CD6 has therapeutic

potential for the intervention of septic shock syndrome and of other inflammatory diseases related to infectious diseases. **[0009]** Innate immune responses rely on the ability of multiple non-polymorphic germline-encoded receptors to recognize the so-called Pathogen-Associated Molecular Patterns (PAMPs), which are conserved products of microbial pathogens, not shared by the host, and essential for their survival. Pattern-Recognition Receptors (PRRs) are mainly expressed by phagocytic cells (granulocytes, macrophages, dendritic cells) and cells of epithelial barriers. Some of them have been shown to directly interact with a variety of microbial components. Herein the inventors have surprisingly found that CD6, a member of the SRCR superfamily expressed on human lymphocytes, binds to Gram-positive and Gram-negative bacteria, as well as to other microbial (viral, fungal) structures. The main reported role of CD6 is the modulation of T cell activation and differentiation signals. It is surprising that in addition, CD6 has the ability of sensing the presence of microbial components. This is a surprising activity of lymphocytes, since this activity has been related to macrophages, which express receptors such as CD14 and Sp α . Interestingly, competition binding experiments show that the interaction of CD6 with the bacterial surfaces is specific. In contrast to CD6, the recombinant form encompassing the ectodomain of CD5 do not bind to bacteria or to LPS. Data also show that CD6 binds to LTA and LPS through independent and nonoverlapping sites of the molecule.

[0010] Presence of CD6 induces bacterial aggregation. The bacterial aggregation data, together with the aggregation of LPS induced by CD6 in the presence of Ca²⁺, indicate that CD6 may contribute to increase the size of invading bacteria as well as of LPS particles. This would facilitate particle clearance from the circulation, and reduce subsequent inflammatory processes, which in cases such as sepsis may even result fatal. Accordingly, administration of a single dose (25 μ g) of CD6 into mice one hour prior to LPS challenge significantly enhanced their survival rate (up to 70 %) as compared to CD5 or saline treatment, and concomitantly induced a significant reduction on the serum levels of pro-inflammatory cytokines TNF- α , IL-1 β and IL-6 in these mice.

[0011] Low levels of soluble CD6 have been detected in normal human sera by ELISA assays, but its biochemical characterization has not been achieved until present. By using affinity chromatography techniques, a natural soluble CD6 (nsCD6) protein has been purified from pooled human sera with similar molecular weight (MW), antibody reactivity and cell binding characteristics than the recombinant form of CD6, and exclusively composed of the ectodomain of human CD6. These data, together with the shared ability to bind to LPS in ELISA assays, indicate that CD6 retains the biological activity of the circulating form of CD6 and validate the use of recombinant CD6 in the studies given the low availability of natural CD6 (it is present in serum at concentrations around 5.27 ng/ml). These results also validate the potential use of a recombinant form of CD6 to be administered to a patient.

[0012] Accordingly, the present invention relates to the use of a CD6 product for the manufacture of a medicament for therapeutic and/or preventive treatment, in a mammal including a human, of an infectious disease, or of an inflammatory condition related to an infectious disease, or of an inflammatory disease related to the presence of a product derived from an infectious agent.

[0013] The invention may alternatively be formulated as a method for therapeutic and/or preventive treatment in a mammal including a human, of an infectious disease, or of an inflammatory condition related to an infectious disease, or of an inflammatory disease related to the presence of a product derived from an infectious agent, comprising administering to said mammal in need thereof an effective amount of CD6 product.

[0014] In a particular embodiment of the invention, the infectious disease is a microbial infection. In more particular embodiments, the microbial infection is selected from the group consisting of a bacterial infection (either Gram-negative or Gram-positive bacteria), a parasitic infection, a viral infection, a fungal infection and combinations thereof (polymicrobial infection).

[0015] In another particular embodiment, the infectious disease is a septicemia. As used herein, the term "septicemia" refers to the presence of any microbe in blood stream. Particularly, the septicemia is selected from the group consisting of a bacteremia, a viremia, a fungemia, a parasitemia and combinations thereof.

[0016] The presence of viable microbes is found in most cases of inflammatory conditions related to an infectious disease, whereas 20% to 30% of patients do not have microbes identified from any source but products derived from them. Thus, in another embodiment, the inflammatory condition is related to a product derived from an infectious agent. Particularly, the infectious agent is selected from the group consisting of a bacterium, a parasite, a virus, a fungus, and combinations thereof.

[0017] Sepsis is defined as the presence or presumed presence of an infection accompanied by evidence of a systemic response called the systemic inflammatory response syndrome (SIRS). For sepsis definition, reference is made to the article "Severe sepsis and septic shock: review of the literature and emergency department management guidelines", H.B. Nguyen et al., Ann. Emergency Med. 2006, vol. 48, pp. 28-54. Sepsis is usually caused by bacterial infections (either Gram-negative or Gram-positive bacteria) but can also be caused by other pathogens. Most often however,

sepsis is caused by Gram-negative bacteria infections. In this case, however, the injury and symptoms attributable to sepsis are not only caused by the bacteria but are also caused by a component of the bacteria cell wall known as endotoxin or LPS. LPS molecules are glycolipids that are ubiquitous in the outer membrane of Gram-negative bacteria. LPS is released when the immune system destroys the invading bacteria. The released LPS binds to monocytes, macrophages, and endothelial cells and triggers the production of various mediators such as TNF- α and interleukins (IL-1, IL-6, and IL-8). Production of excessive TNF- α , IL-1, IL-6, and IL-8 is a major cause of severe forms of sepsis.

[0018] In a particular embodiment of the invention, the inflammatory condition is SIRS. In another particular embodiment, the inflammatory condition is sepsis. SIRS is defined as the presence of two or more of the following: (1) temperature greater than 38°C or less than 36 °C; (2) pulse rate greater than 90 beats/min; (3) respiratory rate greater than 20 breaths/min (or PCO₂ less than 32 torr); and (4) white blood cells count greater than 12000/mm³ or less than 4000/mm³, or greater than 10% immature band forms.

[0019] In another particular embodiment, the inflammatory condition is severe sepsis. Severe sepsis is defined as the sepsis which is accompanied by one or more organ dysfunctions. Organ dysfunction can be defined as acute lung injury; coagulation abnormalities; thrombocytopenia; altered mental status; renal, liver, or cardiac failure; or hypoperfusion with lactic acidosis.

[0020] Finally, in another particular embodiment, the inflammatory condition is septic shock. Septic shock is defined as the presence of sepsis and refractory hypotension, i.e., systolic blood pressure less than 90 mmHg, mean arterial pressure less than 65 mmHg, or a decrease of 40 mmHg in systolic blood pressure compared to baseline unresponsive to a crystalloid fluid challenge of 20 to 40 ml/kg. Thus, septic shock is effectively a form of severe sepsis.

[0021] The source of the infection can be any of a number of places throughout the body. Common sites of infection that can lead to sepsis comprise the following:

- inflammation of the appendix (appendicitis), diverticulitis, bowel problems, infection of the abdominal cavity (peritonitis), and gallbladder or liver infections;
- inflammation or infections of the brain or the spinal cord;
- lung infections such as pneumonia;
- skin infections through wounds or through openings made with intravenous catheters, cellulitis (inflammation of the skin's connective tissue);
- urinary tract infections, especially if the patient has a urinary catheter to drain urine;
- dental and gynecological examinations or treatments;
- blunt or penetrating trauma, surgery, and endocarditis.

Definition of CD6 product

[0022] The CD6 receptor is a lymphoid-specific surface glycoprotein expressed on the membrane of thymocytes, mature T cells, and the B1a B cell subset, although CD6 expression has also been reported on certain regions of the brain. The CD6 receptor belongs to the SRCR superfamily characterized by the presence of one or several repeats of a cysteine-rich extracellular domain named SRCR (cf. M.R. Sarrias et al., "The Scavenger Receptor Cysteine-Rich (SRCR) domain: an ancient and highly conserved protein module of the innate immune system", Crit. Rev. Immunol. 2004, vol. 24, pp. 1-37). Its extracellular region is exclusively composed of three consecutive SRCR domains. Functionally, it is physically associated to the antigen-specific receptor complex present on T (TCR/CD3) and B (BCR) cells, where CD6 contributes to either positive or negative modulation of the activation and differentiation signals delivered by that receptor complex. It is well accepted that CD6 binds to its natural ligand ALCAM ("Activated Leukocyte Cell Adhesion Molecule", also known as CD166) (cf. M.A. Bowen et al., "Analysis of domain-domain interactions between CD6 and activated leukocyte cell adhesion molecule (ALCAM)", Tissue Antigens 1996, vol. 48, pp. AS401.32), a broadly expressed adhesion molecule of the Ig superfamily.

[0023] As used herein, the term "CD6 product" means a product comprising the CD6 ectodomain or a fragment thereof. Ectodomain refers to the three SRCR domains with the intervening sequences and the stalk region which separates it from the membrane. Suitable CD6 products include natural, synthetic, or recombinant biologically active polypeptide of CD6 ectodomain or fragments thereof; biologically active polypeptide variants of CD6 ectodomain or fragments thereof, including hybrid fusion proteins or dimers; or to biologically active polypeptide analogs of CD6 ectodomain or fragments thereof. Analogs include products where one or more amino acid residues have been replaced by a different amino acid. Conservative amino acid substitutions are preferred.

[0024] CD6 product is from mammalian origin and more preferably from human origin.

[0025] Human CD6 is 100 to 130 kDa. The human full length CD6 protein described in GenBank Accession number NP_006716 has 668 amino acids. The ectodomain is composed of three SRCR domains, the intervening sequences and a stalk region.

[0026] In a particular embodiment, the CD6 product of the invention comprises the amino acid sequence SEQ ID NO:

1. This sequence includes the three SRCR domains, the intervening sequences and the stalk region. In another embodiment, the CD6 product is a murine CD6 product.

[0027] Due to the low plasma levels of CD6, it is not industrially viable to obtain CD6 from purification plasma or serum. Thus, for the purpose of the invention, it is preferred to produce the CD6 product by genetic engineering methods. Any method commonly used in the art can be employed to produce recombinant CD6, hereafter referred as rCD6. A preferred method to express and purify rCD6 is described below in the examples. This method allows to produce rCD6 for experimental purposes, so industrial scaling-up is necessary to produce large amounts of rCD6. The CD6 of the present invention may also be expressed as a fusion protein.

[0028] To test whether a CD6 product is suitable for the purpose of the invention, a microbial binding assay may be used. A suitable assay is described in Example 1.

[0029] According to the teaching of the present invention, the CD6 product can be administered to a mammalian, preferably a human. The purpose of the administration of the CD6 product may be preventive (to avoid the development of these diseases) and/or therapeutic (to treat these diseases once they have been developed/installed).

[0030] It is to be understood that the CD6 product is administered in a pharmaceutically acceptable form. Those skilled in the art may ascertain the proper dose using standard procedures. It is understood that the dose should be an effective amount of CD6 product in the sense that a reduced inflammatory response is seen in the treated subject.

[0031] The CD6 product of the invention can be administered alone or in a composition with pharmaceutically acceptable carriers or excipients. The skilled in the art will adapt the composition depending on the particular mode of administration. The compositions may comprise the CD6 product as a single agent against the infectious diseases or the inflammatory conditions related thereto, combinations of such agents, or combinations with other therapeutic agents depending on the condition.

[0032] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skilled in the art to which this invention belongs. Methods and materials similar or equivalent to those described herein can be used in the practice of the present invention. Throughout the description and claims the word "comprise" and its variations are not intended to exclude other technical features, additives, components, or steps. Additional objects, advantages and features of the invention will become apparent to those skilled in the art upon examination of the description or may be learned by practice of the invention. The following examples and drawings are provided by way of illustration, and they are not intended to be limiting of the present invention.

[0033] In the examples and drawings, rsCD6 refers to the recombinantly soluble obtained CD6 ectodomain to be distinguished from the CD6 receptor bound to membrane.

DESCRIPTION OF THE DRAWINGS

[0034]

FIG. 1 shows binding of rsCD6 to Gram-positive and Gram-negative bacteria. A) Western blot analysis of the affinity-purified biotin-labeled proteins with streptavidin- HRP. B) Binding of biotinylated proteins (BSA, rsCD5, rSpα and rsCD6) to *E. coli* and *S. aureus*. C) Calcium-dependence of the binding of biotin-labeled rsCD6 and rSpα to *E. coli* and *S. aureus*. TPA means "total protein added". D) Competition binding assays of rsCD6-biotin to *E. coli* and *S. aureus* in the presence of increasing concentrations of LPS or LTA. C means "competitor".

FIG. 2 shows characterization of affinity-purified circulating CD6 from human serum. A) Coomassie blue staining of affinity-purified nsCD6 from human serum. B) Western blot analysis of biotin-labeled purified nsCD6 and rsCD6 proteins and membrane CD6 immunoprecipitated from surface biotinylated HUT-78 T cells with streptavidin-HRP. C) Western blot analysis of the same biotin-labeled proteins as in B) with a rabbit polyclonal antiserum specific for the extracellular region of CD6. D) Flow cytometry analysis of the reactivity of biotin-labeled rsCD5, rsCD6, nsCD6 or BSA with the K652 and Raji cells. Bound protein was detected with streptavidin-Tricolor. F.I. means "fluorescence intensity" and C.N. means "cell number".

FIG. 3 refers to the binding of rsCD6 to LPS. A) ELISA showing direct binding of affinity-purified nsCD6 and rsCD6 to LPS. LPS purified from *E. coli* was coated into 96-well microtiter plates. Several concentrations of biotin-labeled rsCD6, nsCD6 or BSA, were then added to the wells, and bound protein was detected with streptavidin-HRP. Binding of rsCD6 or rsCD5 to Re-LPS was then monitored by changes in FITC-Re-LPS fluorescent properties. P.a. means "protein added". B) rsCD6, but not rsCD5, induces a significant increase in fluorescence anisotropy (referred as "A") upon binding to FITC-Re-LPS, which increases with increasing rsCD6 concentration. C) Net change in fluorescence emission intensity (referred as "F") of FITC-Re-LPS at 520 nm upon addition of increasing amounts of rsCD6 or rsCD5. The apparent K_d for FITC-Re-LPS/sCD6 complexes, calculated from the saturation curve fitted to a rectangular hyperbola, was $2.69 \pm 0.32 \times 10^{-8}$ M.

FIG. 4 shows the binding of LPS from *E. coli* to cell surface CD6. A) Flow cytometry analysis showing direct binding of increasing amounts of LPS-FITC (0, 10, 20 μg) to parental and CD6.wt-transfected 2G5 cells. C.N. means "cell

number" and F.I. "fluorescence intensity". B) To ease comparison, mean fluorescence intensities of A) were plotted against the amount of LPS-FITC added to each cell line. G.M. means "Geo mean". C) Competition studies of LPS-FITC binding to the 2G5-CD6.wt transfectants. Cells were incubated with 15 μ g of FITC-LPS in the presence of increasing amounts (0, 10, 20 μ g) of rsCD6 (left) or with 15 μ g of either rsCD6 or rsCD5 (right). Fluorescence intensity was analyzed by flow cytometry.

FIG. 5 shows that rsCD6 induces bacterial aggregation. A) FITC-labeled *E. coli* and *S. aureus* bacterial suspensions were incubated overnight at room temperature (RT) with rsCD6 or rsCD5 (2 μ M) in the presence of 5 mM Ca^{2+} . Equimolar concentrations of rSp α and HSA were used as positive and negative control, respectively. Aggregation was observed by direct examination on a fluorescence microscope. B) Kinetics of Ca^{2+} -dependent Re-LPS aggregation in the absence ($\bullet$) and presence of increasing concentration of rsCD6. The final concentrations of Re-LPS, calcium, and EDTA were 100 μ g/mL, 2.5 mM and 5 mM, respectively. The final concentrations of rsCD6 were 0.25 (O), 0.50 (Δ), 1.0 ($\bullet$), 2.0 ($\square$) μ g/ml. One representative experiment of two performed is shown. Time is indicated as T (min).

FIG. 6 shows the effect of rsCD6 and rsCD5 on survival rate and cytokine serum levels following LPS-induced septic shock. A) Survival (S) graph. C57BL/6J mice (8-weeks old) were injected i.p. with a lethal dose of LPS from *E. coli* 0111 :B4 (30 mg/kg) 1 h after i.p. administration of either sterile saline solution (referred as "S") (n=26) or rsCD5 (n=10) or rsCD6 (n=16) (25 μ g, each). The percentage of survival mice was analyzed using Gradpath prism 4.0 and the log-rank t-test P-values were calculated. B) Circulating levels of cytokines in LPS-challenged mice. Plasma levels of TNF- α , IL-1 β and IL-6 were quantified by ELISA at different times after LPS injection. Data are expressed as mean $\pm$ SEM. Statistical differences in the results were evaluated by the two-tailed Student's t test. *, Statistically significant difference (P<0.05). Time is indicated as T (min)

FIG. 7 shows the binding of rsCD6 to zymosan extracted from yeast (*Saccharomyces cerevisiae*). Results of ELISA assay where A means "Absorbance", Z "Zymosan", and P "peptidoglycan" purified from *S. aureus*. LPS purified from *E. coli* strain 0111:B4 and BSA are the positive and negative controls.

FIG. 8 shows the binding of rsCD6 binds to human cytomegalovirus particles. Results of ELISA assay.

FIG. 9 shows the binding of rsCD6 binds to HIV surface protein gp120. Results of ELISA assay.

DETAILED DESCRIPTION OF PARTICULAR EMBODIMENTS

Cells

[0035] The human lymphoblastoid B cell line Raji, the erythromyeloid cell line K562, as well as the leukemic T cell line HUT-78, were obtained from the American Type Culture Collection (ATCC, Manassas, VA). The CD5- and CD6-negative Jurkat 2G5 cells were obtained by Fluorescent Activated Cell Sorting (FACS) and further cloning of Jurkat cells, as reported (M. Simarro et al., "The cytoplasmic domain of CD5 mediates both TCR/CD3-dependent and - independent diacylglycerol production", J. Immunol. 1997, vol. 159, pp. 4307-4315). The 2G5 Jurkat cells were stably transfected with the pH β -CD6.wt construct. Briefly, the expression construct coding for wild-type CD6 (pH β -CD6.wt) was obtained by cloning *SalI/EcoRI*- and *EcoRI/BamHI*-restricted (Fermentas MBI) fragments corresponding to the extracellular and cytoplasmic regions of CD6, respectively, into *SalI/BamHI*-restricted pH β APr-1-neo mammalian expression vector. The extracellular portion of CD6 was obtained by PCR amplification using the 5'TCTCGTCGACATGTGGC-TCTTCTTCGGGAT3' (SEQ ID NO: 2) and 5'AACTTCTTTGGGGATGGTGATGGG3' (SEQ ID NO: 3) primers and the CD6-PB1 cDNA sequence cloned into pBNeo as a template. The intracellular region of CD6 was obtained by PCR amplification of HUT78 cDNA with the 5'GTCACTATAGAATCTTCTGTG3' (SEQ ID NO: 4) and 5'AAAGGATCCCT-AGGCTGCGCTGATGTCATC3' (SEQ ID NO: 5) primers (cf. I. Gimferrer et al., "The accessory molecules CD5 and CD6 associate on the membrane of lymphoid T cells", J. Biol. Chem. 2003, vol. 278, pp. 8564-71).

[0036] Unless indicated, all cells used in this study were grown in RPMI 1640 medium (Life Technologies, Gaithersburg, MD) supplemented with 100 U/ml penicillin, 100 μ g/ml streptomycin and 10% FCS (GIBCO Invitrogen, Paisley, UK). The human embryonic kidney epithelial cell line HEK 293-EBNA (Invitrogen Life Technologies, Paisley, U.K.) was grown in Dulbecco's modified Eagle's medium (DMEM/F12; Invitrogen Life Technologies), supplemented with 100 U/ml penicillin, 100 μ g/ml streptomycin, 250 μ g/ml geneticin (G418), and 10% FCS.

Antibodies and reagents

[0037] The mAbs Cris-1 (anti-CD5, IgG2a) and 161.8 (anti-CD6, IgG1) were produced by Dr. R. Vilella (Hospital Clinic, Barcelona, Spain). The mouse anti-human CD6 mAbs MAE1-C10 (IgG1) and SPV-L14.2 (IgG1) were provided by Drs. F. Sánchez-Madrid (Hospital de la Princesa, Madrid, Spain) (cf. L. Cardenas et al., "Phosphorylation-Dephosphorylation of the CD6 Glycoprotein Renders 2 Isoforms of 130 and 105 Kilodaltons - Effect of Serum and Protein-Kinase-C Activators" Journal of Immunology 1990, vol. 145, pp. 1450-55) and Jo Hilders (Bioprobe B.V., The Netherlands), respectively. The

rabbit polyclonal antiserum against the extracellular region of human CD6 was produced in the laboratory by immunization with rsCD6 (cf. I. Gimferrer et al., J. Biol. Chem. 2003, vol. 278, pp. 8564-71).

[0038] The following reagents were purchased: HRP-conjugated streptavidin (DakoCytomation, Glostrup, Denmark); FITC-conjugated rabbit anti-mouse Ig (Sigma-Aldrich, St. Louis, MO, USA); purified LPS from *E. coli* (055:B5, 026:B6, 0111:B4) and Lipoteichoic Acid (LTA) from *S. aureus*, as well as FITC-LPS from *E. coli* 0111:B4 (Sigma-Aldrich). PBS (Roche Diagnostics, Indianapolis, USA) is 137 mM NaCl, 3 mM KCl, 8 mM Na₂HPO₄, 1.5 mM KH₂PO₄, pH 7.4; TBS is 140 mM NaCl, 50 mM Tris-HCl, pH 7.4. HSA was from Grifols (Grifols, Barcelona, Spain). The Re-LPS, a Re595 mutant of LPS from *Salmonella* Minnesota, was from Sigma. Fluorescein, and fluorescein-5-isothiocyanate (FITC, isomer I) were from Molecular Probes (Eugene, OR). Methanol and chloroform used to dissolve Re-LPS were HPLC-grade (Scharlau, Barcelona).

Expression and affinity-purification of recombinant soluble proteins

[0039] The ectodomain of human CD6 (rsCD6 amino acids D25 to E384, immature protein numbering) was expressed using an episomal expression system in human embryonic kidney cells (HEK 293-EBNA). These cells constitutively express the Epstein Barr viral protein EBNA-1, allowing episomal replication of the pCEP-Pu vector, a kind gift from Drs T. Sasaki and R. Timpl (Max Planck Institute for Biochemistry, Martinsried, Germany) (cf. E. Kohfeldt et al., "Properties of the extracellular calcium binding module of the proteoglycan testican", 1997, FEBS Lett. Vol. 414, pp. 557-61). The extracellular region of CD6 was PCR amplified by using the 5'CTTCTAGATGACCAGCTCAACACCACCAGCA3' (SEQ ID NO: 6) and 5'GCGGATCCCTA TTCTATAGTGACTGTCTGAACA3' (SEQ ID NO: 7) and the CD6-PB1 cDNA as a template (cf. W.H. Robinson et al., "Human CD6 possesses a large, alternatively spliced cytoplasmic domain", Eur. J. Immunol. 1995, vol. 25, pp. 276). The PCR product was cloned into the pCEP-Pu vector. The resulting constructs were transfected into HEK 293-EBNA cells. Briefly, 10⁶ cells in 10-cm culture dishes were transfected using the calcium-phosphate method with 20 µg of the plasmids. Transfectants were selected with 1 µg/ml puromycin (Sigma, St. Louis, MO) in the culture medium. Cell transfectants expressing rsCD6 were grown to confluence in DMEM/FCS, washed twice with phosphate buffered saline (PBS, Roche Diagnostics, Indianapolis, USA) and exchanged to serum-free medium (DMEM/F12), containing 100 U/ml penicillin, 100 µg/ml streptomycin, 250 µg/ml G418, and 1 µg/ml puromycin. The medium was collected every 48-72 h over a period of 15 days. The recombinant protein was affinity-purified over a CNBr-activated Sepharose 4B column covalently coupled to mAb 168.1 (anti-CD6). Unbound protein was then washed off with PBS containing 0.5 M NaCl, and 1 % NP40. The protein was eluted with PBS containing 3.5 M MgCl₂, and dialyzed to PBS. The purity of the purified protein, was assessed by SDS-PAGE under reducing conditions, and staining with Coomassie blue.

Protein biotinylation

[0040] Protein biotinylation was performed with EZ-Link PEO-maleimide-activated biotin (Pierce, Perbio Science, Cheshire, UK) following the manufacturer's instructions and as before (cf. M.R. Sarrias et al., "A role for human SP alpha as a pattern recognition receptor", J. Biol. Chem. 2005, vol. 280, pp. 35391-8). Free biotin was removed by exchanging the samples to PBS over a HiTrap™ desalting column (Amersham Pharmacia Biotech). Monitoring of the biotinylation reactions was performed by Western blotting analysis.

Bacterial strains and bacterial binding studies

[0041] The *E. coli* and *Staphylococcus aureus* bacterial strains used in this study are clinical isolates characterized by the Department of Microbiology of the Hospital Clinic of Barcelona using standard biochemical procedures. Bacteria were grown overnight in Luria Bertoni broth (LB) at 37 °C with aeration and then harvested by centrifugation at 3500 x g for 10 min. Bacterial pellets were re-suspended in TBS (140 mM NaCl, 50 mM Tris-HCl, pH 7.4) to a final density of 10¹⁰ bacteria per ml. Quantification was done by plating bacteria dilutions on agar. Binding of rsCD6 to bacteria was studied following a method described previously (cf. M.R. Sarrias et al., J. Biol. Chem. 2005, vol. 280, pp. 35391-8).

Purification of soluble CD6 from human serum

[0042] One liter of human plasma pooled from healthy blood donors was obtained from the Blood Bank of the Hospital Clinic de Barcelona. The plasma was centrifuged at 10,000g for 15 min and filtered through a 0.22 µm filter (Millipore, Billerica, MA, USA). The plasma was then precipitated with 20% (w/v) (NH₄)₂SO₄ at 4 °C for 30 min in the presence of 2 mg/ml aprotinin and leupeptin, 1 mM PMSF, 0.02% sodium azide and 0.5% NP-40, and then centrifuged at 10,000g for 30 min at 4 °C. The resulting supernatant was brought to 70% (w/v) (NH₄)₂SO₄ at 4 °C for 30 min, and then centrifuged at 10,000g for 30 min at 4 °C. The pellet was resuspended in PBS and subjected to dialysis at 4 °C against PBS

supplemented with 1 mM PMSF, and 0.02% sodium azide. The dialyzed protein was affinity-purified over a CNBr-activated Sepharose-4B column (Amersham Pharmacia Biotech) adsorbed with the anti-CD6 mAb SPV-L14.2. Presence of nsCD6 in the eluted fractions was assessed by sandwich ELISA with specific mAbs (cf. M. Ramos-Casals et al., "High circulating levels of soluble scavenger receptors (sCD5 and sCD6) in patients with primary Sjogren's syndrome" *Rheumatology* 2001, vol. 40, pp. 1056-9) and Western-blot. Purity was assessed by SDS-PAGE and Coomassie blue staining.

LPS-binding ELISA assays

[0043] Twelve μg of LPS purified from *E. coli* 055:B5, 0111:B4, or O26:B6 (Sigma) was used to coat 96-well microtiter plates (Nunc, Roskilde, Denmark) in PBS, overnight at 4 °C. Non-specific binding to the wells was prevented by the addition of PBS containing 1% BSA for 1 h at room temperature. Several concentrations of biotin-labeled BSA, rsCD6 or nsCD6 were then added to the wells and incubated for 1 h at RT. Bound protein was detected by the addition of a 1:1000 dilution of HRP-labeled streptavidin (DAKO, Glostrup, Denmark), incubating for 30 min at RT. Between each incubation step, unbound protein or HRP-streptavidin was washed off three times with PBS plus 0.01 % Tween-20. Color was developed by adding 3,3',5,5'-tetramethylbenzidine liquid substrate (Sigma), and the absorbance was read at 405 nm. The assay was repeated three times with similar results.

Binding assays of soluble proteins to FITC-Re-LPS

[0044] A fluorescent Re-LPS derivative (FITC-Re-LPS) was prepared in which the phosphoethanolamine group of Re-LPS was bound to FITC and prepared by a previously described method (cf. R.R. Skelly et al., "Stimulation of T-independent antibody responses by hapten-lipopolysaccharides without repeating polymeric structure", *Infect. Immun.* 1979, vol. 23, pp. 287-93). Fluorescence measurements were carried out using an SLM-Aminco AB-2 spectrofluorimeter with a thermostated cuvette holder (± 0.1 °C), using 5x5 mm path-length quartz cuvettes. Fluorescence emission spectra of FITC-Re-LPS (0.5 $\mu\text{g}/\text{ml}$) were measured in the presence and absence of either rsCD6 or rsCD5 in 100 mM NaCl, 2 mM EDTA, 5 mM Tris-HCl buffer (pH 8) at 15 °C. The blanks (protein alone) and FITC-Re-LPS samples (with and without protein) were excited at 470 nm and emission spectra recorded from 500 to 650 nm. The apparent K_d for FITC-Re-LPS/protein complexes was obtained by analyzing the time dependence of the fluorescence change when 0.5 $\mu\text{g}/\text{ml}$ FITC-Re-LPS reacted with various concentrations of either rsCD6 or rsCD5 at 15 °C. Fluorescence emission was monitored at 520 nm for 30 min. These experiments were performed twice and in duplicates, as previously described for K_d determination of the Re-LPS interaction with LBP, CD14, surfactant protein A, and rSp α (cf. M.R. Sarrias et al., *J. Biol. Chem.* 2005, vol. 280, pp. 35391-8; P.S. Tobias et al., "Lipopolysaccharide binding protein-mediated complexation of lipopolysaccharide with soluble CD14", *J. Biol. Chem.* 1995, vol. 270, pp. 10482-8; and I. Garcia-Verdugo et al., "Interaction of SP-A (surfactant protein A) with bacterial rough lipopolysaccharide (Re-LPS), and effects of SP-A on the binding of Re-LPS to CD14 and LPS-binding protein", *Biochemical Journal* 2005, vol. 391, pp. 115-24).

[0045] Fluorescence emission anisotropy measurements were obtained with Glan Prism polarizers as previously described (cf. I. Garcia-Verdugo et al., *Biochemical Journal* 2005, vol. 391, pp. 115-24). Excitation and emission wavelengths were set at 470 and 520 nm, respectively.

Bacteria and LPS aggregation assays

[0046] The bacteria aggregation assays were performed as previously described (cf. M.R. Sarrias et al., *J. Biol. Chem.* 2005, vol. 280, pp. 35391-8). LPS aggregation induced by rsCD6 was studied at 15 °C by measuring the change in absorbance at 400 nm in a Beckman DU-640 spectrophotometer as described (cf. I. Garcia-Verdugo et al., *Biochemical Journal* 2005, vol. 391, pp. 115-24). Briefly, the sample and reference cuvettes were first filled with Re-LPS (100 $\mu\text{g}/\text{ml}$, final concentration) in 5 mM Tris-HCl buffer (pH 7.4), 150 mM NaCl, 0.2 mM EDTA. After a 10 min equilibration period at 15 °C, rsCD6 was added to the sample cuvette at the indicated concentration, and the change in absorbance at 400 nm was monitored. Next, Ca^{2+} (2.5 mM) was added to both the sample and reference cuvettes, and the change in absorbance was monitored again. Ca^{2+} -dependent LPS aggregation was reversed by adding EDTA (5 mM, final concentration).

Flow cytometry assays

[0047] Cell-binding properties of soluble proteins were assessed as previously described (cf. J. Calvo et al., "Identification of a natural soluble form of human CD5" *Tissue Antigens* 1999, vol. 54, pp. 128-37). Binding of LPS to cell surface CD6 was assessed by using the 2G5-CD6.wt cell line. Briefly, 2×10^5 cells were incubated with different amounts of LPS-FITC from *E. coli* 0111:B4 (Sigma) in the presence of blocking buffer (PBS plus 10% human AB serum, 2% FCS and 0.02% sodium azide). After 1 h of incubation at 4 °C, the cells were washed twice with washing buffer (PBS

plus 2% FCS and 0.02% sodium azide). For competition studies, 15 µg of FITC-LPS were incubated for 30 min on ice with 2x10⁵ 2G5-CD6.wt cells in the presence of different amounts of rsCD5 or rsCD6.

CD6 immunoprecipitation

[0048] For immunoprecipitation of cell surface bound CD6 (mCD6), 1×10⁶ HUT-78 T cells were surface labeled with EZ-Link PEO-maleimide-activated biotin (Pierce, Perbio Science, Cheshire, UK) following the manufacturer's instructions. Then, membranes were solubilized with Triton X-100 detergent and proteins immunoprecipitated for 2 h at 4 °C with 1 µg of anti-CD6 mAb (161.8) plus 20 µl of 50% Protein A Sepharose CL-4B beads (Amersham Biosciences). Immune complexes were analyzed by Western Blotting as described (cf. I. Gimferrer et al., J. Biol. Chem. 2003, vol. 278, pp. 8564-71).

LPS-induced endotoxic shock

[0049] C57BL/6J mice (8-weeks old) were injected i.p. with a lethal dose of LPS from *E. coli* 0111 :B4 (Sigma, St. Louis, MO) (30 mg/kg) in a volume of 250 µl sterile saline solution. Administration of 25 µg of either rsCD5 (10 mice) or rsCD6 (16 mice) was carried out 1 h prior to the LPS challenge. Control mice received the same volume of sterile saline solution (26 mice). The percentage of survival mice was analyzed using Gradpath prism 4.0 and the log-rank t-test P-values were calculated.

[0050] The experimental procedure was approved by the ethics committee of the University of Murcia and performed in accordance with institutional animal care guidelines that comply with regulations in Spain (RD 1201/2005), Europe (86/609) and the National Institutes of Health's Guide for the Care and Use of Laboratory Animals.

Determination of cytokine serum levels

[0051] The systemic release of TNF-α, IL-1β and IL-6 cytokines was determined in pooled serum samples from 6 mice of each group by ELISA according to manufacturer's instructions (R&D Systems, Minneapolis, MN). Data are expressed as mean ± SEM. Statistical differences in the results were evaluated by the two-tailed Student's t test.

EXAMPLE 1: rsCD6 binds to Gram-positive and Gram-negative bacteria

[0052] To determine whether the ectodomain of human CD6 could directly bind to the surface of whole bacteria, the approach described in D.W. Dunne et al., "The Type-I Macrophage Scavenger Receptor Binds to Gram-Positive Bacteria and Recognizes Lipoteichoic Acid", Proceedings of the National Academy of Sciences of the United States of America 1994, vol. 91, pp. 1863-7 was used. Thus, biotin-labeled recombinant soluble proteins encompassing the ectodomains of human CD5, CD6 and Spα (rsCD5, rsCD6, and rSpα) (FIG. 1A) were incubated with bacterial suspensions and their binding to bacterial pellets further assayed by SDS-PAGE and western blotting against streptavidin-HRP. The results show that rsCD6 bound to Gram-positive and - negative bacteria (FIG. 1 B), indicating that this protein possesses bacterial binding activity. In contrast, neither rsCD5 nor the negative control BSA bound to bacterial suspensions. As illustrated by FIG. 1C, the presence of biotin-labeled rsCD6 was greatly reduced in *E. coli* and *S. aureus* bacterial cell pellets in the presence of EDTA. This indicates that rsCD6 mediates Ca²⁺-dependent recognition of cell wall components from Gram-positive and - negative bacteria.

[0053] To determine whether the observed binding of rsCD6 to bacteria was specific and to identify which bacterial cell surface structures were being recognized, competition experiments were designed in which biotin-labeled rsCD6 was incubated with increasing concentrations of purified LPS, or LTA, before the addition of a suspension of either *E. coli* or *S. aureus* (5×10⁷ cells). LPS and LTA were assayed because they are ubiquitous cell surface components of these microorganisms. As illustrated by FIG. 1 D, binding of biotin-labeled rsCD6 to *E. coli* was competed in a dose-dependent manner by LPS (from *E. coli*), but not by LTA (from *S. aureus*). On the contrary, when the binding of rsCD6 to *S. aureus* was studied, LPS did not affect such interaction. Interestingly, this binding was competed in a dose-dependent manner by LTA from *S. aureus*.

EXAMPLE 2: Purification of nsCD6 from human serum

[0054] Natural sCD6 was affinity-purified from 1 l of pooled plasma. This yielded 6 µg of a single protein with a molecular weight (MW) of 80 kDa as deduced from SDS-PAGE analysis and Coomassie blue staining (FIG. 2A). The observed MW closely resembles that of recombinant soluble CD6 (rsCD6) produced in the laboratory, which is exclusively composed of the three extracellular SRCR domains of CD6, and is in contrast to that of the membrane form of CD6 (mCD6), which ranges from 105 to 130 kDa, depending on its degree of phosphorylation. The observed MW of the three different

CD6 forms, i.e. rsCD6, nsCD6 and mCD6 immunoprecipitated from human HUT-78 T cells is shown in FIG. 2B. The purified nsCD6 protein was identified as CD6 by Western blotting assays with a polyclonal antiserum raised against the extracellular region of human CD6 (FIG. 2C). In cell binding experiments, both biotin-labeled rsCD6 and nsCD6 bound to Raji B cells but not to K562 erythroleukemic cells, in accordance with the differential expression of the CD6 ligand (ALCAM/CD166) (FIG. 2D).

EXAMPLE 3: Binding of rsCD6 to LPS and kinetics of the rsCD6-LPS interaction

[0055] Further confirmation of the rsCD6-LPS interaction was obtained from direct binding ELISA assays in which plates were coated with LPS purified from three different *E. coli* strains (O55:B5, O111:B4, or O26:B6), and assayed for binding of biotin-labeled rsCD6, nsCD6, or BSA. The results presented in FIG. 3A show that, in accordance with the bacterial binding experiments in FIG. 1, both natural and recombinant soluble CD6 forms bound to LPS in a dose-dependent fashion. No BSA-LPS interaction could be observed.

[0056] The binding of rsCD6 and rsCD5 to a rough mutant (Re595) of LPS (Re-LPS) in solution was studied next by analyzing the changes in fluorescent properties of FITC-Re-LPS such as anisotropy and intensity. FIG. 3B shows the binding of rsCD6 and rsCD5 to FITC-Re-LPS by measuring fluorescence anisotropy of the labeled LPS molecule. Fluorescence anisotropy measurements depend on the rate and extent of the rotational motion of the fluorophore during the lifetime of the excited state. Addition of different amounts of rsCD6 to FITC-Re-LPS caused a protein concentration-dependent increase of the anisotropy values of FITC-Re-LPS, indicating that the binding of rsCD6 to Re-LPS caused mechanical restrictions of the rotational mobility of the dye. Control experiments were done with free fluorescein to demonstrate that all of these changes did not result from the interaction of rsCD6 with the dye (fluorescein), but with the LPS molecule (data not shown). The fluorescence emission anisotropy of free fluorescein was very low and was not affected by addition of 3-fold excess of rsCD6 (data not shown). On the other hand, rsCD5 did not cause any change in FITC-Re-LPS fluorescence anisotropy, indicating that this protein does not bind to Re-LPS.

[0057] Addition of rsCD6, but not rsCD5, to FITC-Re-LPS in solution also produced an increase of total fluorescence emission intensity of fluorescent LPS. The time dependence of the change in fluorescence emission intensity at 520 nm when FITC-Re-LPS reacted with increasing amounts of either rsCD6 or rsCD5 was determined. FIG. 3C shows that the magnitude of the fluorescence intensity change increased as a function of rsCD6 concentration, but not rsCD5 concentration, and was saturable. These results allowed to determine the affinity of rsCD6 binding to LPS. The apparent K_d for FITC-Re-LPS/rsCD6 complexes, calculated from the saturation curve fitted to a rectangular hyperbola, was 2.69 (± 0.32) 10^{-8} M.

EXAMPLE 4: Binding of LPS to cell-surface CD6

[0058] To determine whether the LPS-CD6 interaction occurs as well with the receptor expressed on the cell surface, studies were performed by staining with FITC-labeled LPS of 2G5 cells, a Jurkat cell derivative selected for deficient CD5 and CD6 expression. As shown by the flow cytometry studies displayed in FIG. 4A and 4B, fluorescence intensity was higher in 2G5 cells stably expressing wild-type CD6 (2G5-CD6.wt) compared to parental untransfected 2G5 cells. Further confirmation of these results was obtained from competition binding experiments. 2G5-CD6.wt cells were stained with a single amount of FITC-LPS in the presence of increasing amounts of rsCD6. In these experiments, binding of FITC-LPS to 2G5-CD6.wt cells was inhibited in a dose-dependent manner by rsCD6, but not with rsCD5, used as a negative control (FIG. 4C), indicating that the inhibition was specific. From these data it is concluded that LPS is able to interact with CD6 on the cell surface.

EXAMPLE 5: Binding of rsCD6 leads to both bacteria and LPS aggregation

[0059] FIG. 5A shows that presence of rsCD6 induced aggregation of Gram-negative (*E. coli*) as well as Gram-positive (*S. aureus*) bacteria. In accordance with its inability to bind bacteria, rsCD5 was also unable to induce their aggregation, as it did not the negative control HSA.

[0060] Since binding of rsCD6 to whole Gram-negative bacteria was enhanced by the presence of Ca²⁺ in the binding buffer (see FIG. 1) it was further explored the process of Re-LPS aggregation induced by rsCD6 in the presence of Ca²⁺. This was analyzed by measuring changes in light absorbance at 400 nm (FIG. 5B). These experiments were carried out under the same ionic conditions as binding studies with fluorescent LPS, except that Ca²⁺ as well as concentrations of Re-LPS, 200 times higher were needed to produce detectable light absorption at 400 nm. FIG. 5B shows that LPS molecules were able to aggregate in buffers containing Ca²⁺ and that low concentrations of rsCD6 induced a further aggregation of LPS. Together, these data suggest that, in the presence of Ca²⁺, rsCD6 may contribute to increase the size of bacterial as well as of LPS aggregates.

EXAMPLE 6: rsCD6 prevents LPS-induced septic shock in mice

[0061] The data on FIG. 5 lead to the notion that circulating CD6 (nsCD6) may contribute to the clearance of LPS particles, reducing their inflammatory effect. Increasing the amount of circulating rsCD6 might therefore have a protective effect in situations in which an excess of LPS could be lethal. To test this, it was assessed whether administration of the rsCD6 into mice would improve their survival in front of LPS-induced septic shock. As observed in FIG. 6A, administration of a single i.p. dose of 25 µg of rsCD6, but not rsCD5 in mice one hour prior to i.p. LPS challenge significantly enhanced their survival rate as compared with the saline control (up to 70 %). In accordance with this data, administration of rsCD6 induced a significant reduction on the levels of plasma pro-inflammatory cytokines TNF-α, IL-1β and IL-6 in these mice (FIG. 6B).

EXAMPLE 7: rsCD6 binds to zymosan extracted from yeast (*Saccharomyces cerevisiae*)

[0062] Twelve µg of LPS purified from *E. coli* 0111:B4 (Sigma), Zymosan from *Saccharomyces cerevisiae* (Sigma), peptidoglycan purified from *S. aureus* (Fluka) or Bovine Serum Albumin (BSA), were used to coat 96-well microtiter plates (Nunc, Roskilde, Denmark) in PBS, overnight at 4 °C. Non-specific binding to the wells was prevented by the addition of PBS containing 1% BSA for 1 h at room temperature. Several concentrations of rsCD6 were then added to the wells and incubated for 3 h at room temperature. Bound protein was detected by the addition of a 1:200 dilution of biotin-labeled mAb anti-CD6 (MAE) for 1 h at RT, followed by a 1:1000 dilution of HRP-labeled streptavidin (DAKO, Glostrup, Denmark), which was incubated for 30 min at RT. Between each incubation step, unbound protein or HRP-streptavidin was washed off three times with PBS plus 0.01 % Tween-20. Color was developed by adding 3,3',5,5'-tetramethylbenzidine liquid substrate (Sigma), and the absorbance was read at 405 nm. The assay was repeated three times with similar results.

[0063] The results presented in FIG. 7 show that rsCD6 bound to zymosan, peptidoglycan and LPS in a dose-dependent fashion. No BSA-rsCD6 interaction could be observed. This ELISA experiment indicates that peptidoglycan, a highly abundant structure from the surface of Gram- positive bacteria, is also recognized by rsCD6. Interestingly, rsCD6 is also able to bind to yeast zymosan.

EXAMPLE 8: rsCD6 binds to human cytomegalovirus particles

[0064] One microgram of a polyclonal anti-CMV antibody (BiosPacific, Emeryville, CA, USA) was used to coat 96-well microtiter plates (Nunc, Roskilde, Denmark) in PBS, overnight at 4 °C. Non-specific binding to the wells was prevented by the addition of PBS containing 3% BSA for 1 h at RT. One microgram of a suspension of CMV particles (ABI Advanced Biotechnologies, Maryland, USA) then added to the wells and incubated for 1 h at RT. Then, different concentrations of rsCD6 protein were added to the wells and incubated for 3 h at RT. Bound protein was detected by the addition of a 1:200 dilution of biotin-labeled mAb anti-CD6 (MAE) for 1 h at RT, followed by a 1:1000 dilution of HRP-labeled streptavidin (DAKO, Glostrup, Denmark), which was incubated for 30 min at RT. Between each incubation step, unbound protein or HRP-streptavidin was washed off three times with PBS plus 0.01 % Tween-20. Color was developed by adding 3,3',5,5'-tetramethylbenzidine liquid substrate (Sigma), and the absorbance was read at 405 nm. The assay was repeated three times with similar results.

[0065] The results presented in FIG. 8 show that rsCD6 bound to CMV particles in a dose-dependent fashion. Background non-specific binding of CD6 to antibody-coated wells was subtracted. These results suggest that rsCD6 is able to bind to the surface of human cytomegalovirus.

EXAMPLE 9: rsCD6 binds to HIV surface protein gp120

[0066] 100 µl of serum-free culture supernatant from cell transfectants expressing rsCD6 were used to coat 96-well microtiter plates (Nunc, Denmark), o/n at 4 °C. Non-specific binding to the wells was prevented by the addition of PBS containing 3% BSA for 1 h at RT. Then, different concentrations of recombinant gp120 protein from HIV (Immunodiagnos- tics, through the NIH AIDS Research and Reference Reagent Program, Germantown, Maryland, USA) were added to the wells and incubated for 1 h at RT. Bound protein was detected by the addition of 500 ng of an anti-gp120 mAb (hm Ab 2G12, Immunodiagnos- tics) in 2% heat- inactivated human serum for 1 h at RT, followed by a 1:200 dilution of HRP-labeled anti-human IgG antibody (DAKO, Glostrup, Denmark), which was incubated for 30 min at RT. Between each incubation step, unbound protein or HRP-streptavidin was washed off three times with PBS plus 0.01 % Tween- 20. Color was developed by adding 3,3',5,5'-tetramethylbenzidine liquid substrate (Sigma), and the absorbance was read at 405 nm. The assay was repeated three times with similar results.

[0067] The results presented in FIG. 9 show that rsgp120 bound to rsCD6 in a dose-dependent fashion. Background non-specific binding of rsgp120 to blocking buffer on the wells was subtracted. These results suggest that rsCD6 is able

EP 2 143 436 A1

to bind to rgp120, the major outer protein of Human Immunodeficiency Virus.

5

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

<110> Universidad de Barcelona
 5 <120> Protein product for the treatment of infectious diseases and of
 inflammatory conditions related thereto
 <130> WO AVCRI012
 <150> ES200700893
 10 <151> 2007-03-28
 <160> 7
 <170> PatentIn version 3.3
 <210> 1
 15 <211> 360
 <212> PRT
 <213> Homo sapiens (fragment)
 <400> 1
 20 Asp Gln Leu Asn Thr Ser Ser Ala Glu Ser Glu Leu Trp Glu Pro Gly
 1 5 10 15
 Glu Arg Leu Pro Val Arg Leu Thr Asn Gly Ser Ser Ser Cys Ser Gly
 20 25 30
 25 Thr Val Glu Val Arg Leu Glu Ala Ser Trp Glu Pro Ala Cys Gly Ala
 35 40 45
 30 Leu Trp Asp Ser Arg Ala Ala Glu Ala Val Cys Arg Ala Leu Gly Cys
 50 55 60
 Gly Gly Ala Glu Ala Ala Ser Gln Leu Ala Pro Pro Thr Pro Glu Leu
 65 70 75 80
 35 Pro Pro Pro Pro Ala Ala Gly Asn Thr Ser Val Ala Ala Asn Ala Thr
 85 90 95
 40 Leu Ala Gly Ala Pro Ala Leu Leu Cys Ser Gly Ala Glu Trp Arg Leu
 100 105 110
 Cys Glu Val Val Glu His Ala Cys Arg Ser Asp Gly Arg Arg Ala Arg
 115 120 125
 45 Val Thr Cys Ala Glu Asn Arg Ala Leu Arg Leu Val Asp Gly Gly Gly
 130 135 140
 Ala Cys Ala Gly Arg Val Glu Met Leu Glu His Gly Glu Trp Gly Ser
 145 150 155 160
 50 Val Cys Asp Asp Thr Trp Asp Leu Glu Asp Ala His Val Val Cys Arg
 165 170 175
 55 Gln Leu Gly Cys Gly Trp Ala Val Gln Ala Leu Pro Gly Leu His Phe
 180 185 190

EP 2 143 436 A1

	Thr	Pro	Gly	Arg	Gly	Pro	Ile	His	Arg	Asp	Gln	Val	Asn	Cys	Ser	Gly	
			195					200					205				
5	Ala	Glu	Ala	Tyr	Leu	Trp	Asp	Cys	Pro	Gly	Leu	Pro	Gly	Gln	His	Tyr	
		210					215					220					
10	Cys	Gly	His	Lys	Glu	Asp	Ala	Gly	Val	Val	Cys	Ser	Glu	His	Gln	Ser	
	225					230					235					240	
	Trp	Arg	Leu	Thr	Gly	Gly	Ala	Asp	Arg	Cys	Glu	Gly	Gln	Val	Glu	Val	
					245					250					255		
15	His	Phe	Arg	Gly	Val	Trp	Asn	Thr	Val	Cys	Asp	Ser	Glu	Trp	Tyr	Pro	
				260					265					270			
20	Ser	Glu	Ala	Lys	Val	Leu	Cys	Gln	Ser	Leu	Gly	Cys	Gly	Thr	Ala	Val	
			275					280					285				
	Glu	Arg	Pro	Lys	Gly	Leu	Pro	His	Ser	Leu	Ser	Gly	Arg	Met	Tyr	Tyr	
		290					295					300					
25	Ser	Cys	Asn	Gly	Glu	Glu	Leu	Thr	Leu	Ser	Asn	Cys	Ser	Trp	Arg	Phe	
	305					310					315					320	
30	Asn	Asn	Ser	Asn	Leu	Cys	Ser	Gln	Ser	Leu	Ala	Ala	Arg	Val	Leu	Cys	
					325					330					335		
	Ser	Ala	Ser	Arg	Ser	Leu	His	Asn	Leu	Ser	Thr	Pro	Glu	Val	Pro	Ala	
				340					345					350			
35	Ser	Val	Gln	Thr	Val	Thr	Ile	Glu									
			355					360									
40	<210>	2															
	<211>	30															
	<212>	DNA															
	<213>	Artificial															
	<220>																
	<223>	PCR primer															
45	<400>	2															
	tctcgtc	gac	atgtgg	ctct	tcttc	gggat											30
50	<210>	3															
	<211>	24															
	<212>	DNA															
	<213>	Artificial															
	<220>																
	<223>	PCR primer															
55	<400>	3															
	aacttct	tttg	gggat	ggtga	tggg												24

5
 <210> 4
 <211> 21
 <212> DNA
 <213> Artificial
 <220>
 <223> PCR primer
 <400> 4
 gtcactatag aatcttctgt g 21
 10
 <210> 5
 <211> 30
 <212> DNA
 <213> Artificial
 15
 <220>
 <223> PCR primer
 <400> 5
 aaagatccc taggctgcmc tgatgcatc 30
 20
 <210> 6
 <211> 31
 <212> DNA
 <213> Artificial
 25
 <220>
 <223> PCR primer
 <400> 6
 cttctagatg accagctcaa caccaccagc a 31
 30
 <210> 7
 <211> 33
 <212> DNA
 <213> Artificial
 35
 <220>
 <223> PCR primer
 <400> 7
 gcggatccct attctatagt gactgtctga aca 33
 40
 45

Claims

1. Use of a CD6 product for the manufacture of a medicament for therapeutic and/or preventive treatment, in a mammal including a human, of an infectious disease, or of an inflammatory condition related to an infectious disease, or of an inflammatory disease related to the presence of a product derived from an infectious agent.
2. The use according to claim 1, wherein the infectious disease is a microbial infection.
3. The use according to claim 2, wherein the microbial infection is selected from the group consisting of a bacterial infection, a parasitic infection, a viral infection, a fungal infection and combinations thereof.
4. The use according to claim 1, wherein the infectious disease is a septicemia.

EP 2 143 436 A1

5. The use according to claim 4, wherein the septicemia is selected from the group consisting of a bacteremia, a viremia, a fungemia, a parasitemia and combinations thereof.
6. The use according to claim 1, wherein the infectious agent is selected from the group consisting of a bacterium, a parasite, a virus, a fungus, and combinations thereof.
7. The use according to claim 1, wherein the inflammatory condition is systemic inflammatory response syndrome.
8. The use according to claim 1, wherein the inflammatory condition is sepsis.
9. The use according to claim 8, wherein the sepsis is severe sepsis.
10. The use according to claim 9, wherein the severe sepsis is septic shock.
11. The use according to any of the claims 1-10, wherein the CD6 product is a recombinant CD6.
12. The use according to claim 11, wherein the CD6 product is a human CD6 product.
13. The use according to claim 12, wherein the CD6 product comprises the amino acid sequence SEQ ID NO: 1.
14. The use according to claim 11, wherein the CD6 product is a murine CD6 product.




FIG. 1**A****B**

FIG. 1

C



D




FIG. 2





FIG. 3


FIG. 4



FIG. 5


FIG. 6




FIG. 7**FIG. 8**

FIG. 9

INTERNATIONAL SEARCH REPORT

International application No.
PCT/ ES 2008/000177

A. CLASSIFICATION OF SUBJECT MATTER

see extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K,A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CIBEPAT,EPODOC,WPI, TXTE, XPESP, BIOSIS, MEDLINE, NPL, EMBASE, EBI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	Sarrias M.R. et al. CD6 binds to pathogen-associated molecular patterns and protects from LPS-induced septic shock. Proceedings of the National Academy of Sciences. 10-07-2007, Vol. 104, N° 28, pages 11724-11729 (the whole document).	1-14
A	Stelter F. et al. Different efficacy of soluble CD14 treatment in high- and low-dose LPS model. European Journal of Clinical Investigation. 1998, Vol. 28, N° 3, pages 205-213 (the whole document).	1-14
A	WO 93/19772 A1 (NORTH SHORE UNIVERSITY HOSPITAL RESEARCH CORPORATION) 14.10.1993, (the whole document)	1-14
A	US 2003/0114377 A1 (KIRKLAND et al) 19.06.2003, (the whole document)	1-14

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance.		
"E" earlier document but published on or after the international filing date		
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"O" document referring to an oral disclosure use, exhibition, or other means	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other documents, such combination being obvious to a person skilled in the art
"P" document published prior to the international filing date but later than the priority date claimed		
	"&"	document member of the same patent family

Date of the actual completion of the international search

31.July.2008 (31.07.2008)

Date of mailing of the international search report

(18-08-2008)

Name and mailing address of the ISA/
O.E.P.M.Paseo de la Castellana, 75 28071 Madrid, España.
Facsimile No. 34 91 3495304

Authorized officer

M. Cumbreño Galindo

Telephone No. +34 91 349 96880

Form PCT/ISA/210 (second sheet) (April 2007)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/ES 2008/000177

C (continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of documents, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2006/056492 A1 (BIOXELL SPA AND UNIVERSITE HENRI POINCARÉ-NANCY 1) 01.06.2006, (the whole document)	1-14

Form PCT/ISA/210 (continuation of second sheet) (April 2007)

EP 2 143 436 A1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/ES 2008/000177

Patent document cited in the search report	Publication date	Patent family member(s)	Publication date
WO 9319772 A	14.10.1993	CA 2133758 A EP 0634935 A EP 19930911583 US 5804189 A	14.10.1993 25.01.1995 06.04.1993 08.09.1998
WO 2006056492 A	01.06.2006	CA 2526684 A JP 2006149365 A AU 2005234705 A SG 122903 A CN 1817901 A US 2006246082 A GB 2433936 A EP 1817337 A EP 20050752687 NZ 543660 A NZ 552065 A	29.05.2006 15.06.2006 15.06.2006 29.06.2006 16.08.2006 02.11.2006 11.07.2007 15.08.2007 20.05.2005 31.01.2008 30.04.2008
US 2003114377 A	19.06.2003	NONE	-----

Form PCT/ISA/210 (patent family annex) (April 2007)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/ES 2008/000177

CLASSIFICATION OF SUBJECT MATTER

A61K 38/17 (2006.01)

A61P 31/00 (2006.01)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 2003002178 A [0007]
- US 20030008822 A [0007]
- US 20020165138 A [0007]
- US 20020155094 A [0007]
- US 20020044929 A [0007]
- US 20020006915 A [0007]
- US 6534648 B [0007]
- US 6315999 B [0007]

Non-patent literature cited in the description

- **H.B. Nguyen et al.** Severe sepsis and septic shock: review of the literature and emergency department management guidelines. *Ann. Emergency Med.*, 2006, vol. 48, 28-54 [0017]
- **M.R. Sarrias et al.** The Scavenger Receptor Cysteine-Rich (SRCR) domain: an ancient and highly conserved protein module of the innate immune system. *Crit. Rev. Immunol.*, 2004, vol. 24, 1-37 [0022]
- **M.A. Bowen et al.** Analysis of domain-domain interactions between CD6 and activated leukocyte cell adhesion molecule (ALCAM). *Tissue Antigens*, 1996, vol. 48, AS401.32 [0022]
- **M. Simarro et al.** The cytoplasmic domain of CD5 mediates both TCR/CD3-dependent and - independent diacylglycerol production. *J. Immunol.*, 1997, vol. 159, 4307-4315 [0035]
- **I. Gimferrer et al.** The accessory molecules CD5 and CD6 associate on the membrane of lymphoid T cells. *J. Biol. Chem.*, 2003, vol. 278, 8564-71 [0035]
- **L. Cardenas et al.** Phosphorylation-Dephosphorylation of the CD6 Glycoprotein Renders 2 Isoforms of 130 and 105 Kilodaltons - Effect of Serum and Protein-Kinase-C Activators. *Journal of Immunology*, 1990, vol. 145, 1450-55 [0037]
- **I. Gimferrer et al.** *J. Biol. Chem.*, 2003, vol. 278, 8564-71 [0037] [0048]
- **E. Kohfeldt et al.** Properties of the extracellular calcium binding module of the proteoglycan testican. *FEBS Lett.*, 1997, vol. 414, 557-61 [0039]
- **W.H. Robinson et al.** Human CD6 possesses a large, alternatively spliced cytoplasmic domain. *Eur. J. Immunol.*, 1995, vol. 25, 276 [0039]
- **M.R. Sarrias et al.** A role for human SP alpha as a pattern recognition receptor. *J. Biol. Chem.*, 2005, vol. 280, 35391-8 [0040]
- **M.R. Sarrias et al.** *J. Biol. Chem.*, 2005, vol. 280, 35391-8 [0041] [0044] [0046]
- **M. Ramos-Casals et al.** High circulating levels of soluble scavenger receptors (sCD5 and sCD6) in patients with primary Sjogren's syndrome. *Rheumatology*, 2001, vol. 40, 1056-9 [0042]
- **R.R. Skelly et al.** Stimulation of T-independent antibody responses by hapten-lipopolysaccharides without repeating polymeric structure. *Infect. Immun.*, 1979, vol. 23, 287-93 [0044]
- **P.S. Tobias et al.** Lipopolysaccharide binding protein-mediated complexation of lipopolysaccharide with soluble CD14. *J. Biol. Chem.*, 1995, vol. 270, 10482-8 [0044]
- **I. Garcia-Verdugo et al.** Interaction of SP-A (surfactant protein A) with bacterial rough lipopolysaccharide (Re-LPS), and effects of SP-A on the binding of Re-LPS to CD14 and LPS-binding protein. *Biochemical Journal*, 2005, vol. 391, 115-24 [0044]
- **I. Garcia-Verdugo et al.** *Biochemical Journal*, 2005, vol. 391, 115-24 [0045] [0046]
- **J. Calvo et al.** Identification of a natural soluble form of human CD5. *Tissue Antigens*, 1999, vol. 54, 128-37 [0047]
- **D.W. Dunne et al.** The Type-I Macrophage Scavenger Receptor Binds to Gram-Positive Bacteria and Recognizes Lipoteichoic Acid. *Proceedings of the National Academy of Sciences of the United States of America*, 1994, vol. 91, 1863-7 [0052]