



(12) UK Patent (19) GB (11) 2 062 665 B

(54) Title of invention

Phthaloperinone dyestuffs

(51) INT CL³: C09B 57/02

(21) Application No
8033855

(22) Date of filing
21 Oct 1980.

(30) Priority data

(31) 9590/79

(32) 25 Oct 1979

(33) Switzerland (CH)

(43) Application published
28 May 1981

(45) Patent published
18 May 1983

(52) Domestic classification
C4P 104 D1M2 D1M
C3R 3D13 3D22 3D2E 3N1 SM

(56) Documents cited
GB 1445049
GB 1062368

(58) Field of search
C4P

(73) Proprietor
Sandoz Ltd.
(Switzerland),
Lichtstrasse 35, 4002 Basle,
Switzerland

(72) Inventor
Paul Buecheler

(74) Agent
B. A. Yorke and Co.,
98 The Centre, Feltham,
Middlesex TW13 4EP


SPECIFICATION

Phthaloperinone dyestuffs

The present invention relates to phthaloperinone compounds, their production and use as dyestuffs.

5 More particularly, the present invention provides dyestuffs of formula I

5



(I)

in which R₁ is a mono- or bicyclic aromatic radical, and the ring A may bear up to 4 substituents selected from chlorine, bromine and nitro (only 1 nitro group)

10 in which the dyestuff is free from sulphonic and carboxylic acid groups, and mixtures of such dyestuffs.

10

R₁ is preferably R₁', where R₁' is naphthyl or phenyl which phenyl is optionally substituted by up to two substituents selected from the group consisting of chlorine, bromine, methyl, methoxy, ethoxy, alkoxyC₁₋₂carbonyl (up to two of each of these), nitro, cyano and aminocarbonyl (only one of any of these).

15 these).

15

More preferably, R₁ is R₁'', where R₁'' is phenyl optionally substituted by up to two substituents selected from the group consisting of chlorine, methyl, methoxy, ethoxy (up to two of each of these) aminocarbonyl and nitro (only one of any of these). More preferably, R₁ is R₁'', where R₁''' is phenyl optionally substituted by up to two substituents selected from the group consisting of methyl, methoxy, ethoxy and aminocarbonyl (only one of the latter).

20 ethoxy and aminocarbonyl (only one of the latter).


20

The ring A is preferably A', where the ring A' is unsubstituted or is substituted by up to four substituents selected from chlorine, bromine and nitro (only 1 nitro group). More preferably the ring A is A'', where the ring A'' is unsubstituted or substituted by up to 4 chlorine atoms. Most preferably the ring A is A''', where the ring A''' is unsubstituted or is substituted by two or four chlorine atoms.

25 Thus preferred compounds and mixtures according to the invention are those of formula I in which

R₁ is R₁' and the ring A is A'. More preferred compounds are those where R₁ is R₁' and A is A'', especially those compounds and mixtures where the SO₂NHR₁ group is in the 4- and/or 6-position.


The present invention also provides a process for the production of compounds of formula I, comprising condensing a compound of formula II



(II)

30

or a mixture thereof with an amine of formula III



(III)

35 The condensation reaction is effected in accordance with known methods. The compounds of formula II and III are either known or are prepared in accordance with known methods from available starting materials. When the compounds of formula II are prepared by sulphonylation of the corresponding phthaloperinones a mixture of isomers is obtained, the predominant isomers being those where the sulphonylchloride group is in the 4- and 6-positions. If pure compounds of formula II are desired, the isomers can be separated, however, in general for commercial purposes no separation step would be carried out. Thus, the present invention primarily provides mixtures of compounds of formula I. 40 Further, when the ring A bears up to three substituents, that is when the compound of formula II is produced from the correspondingly substituted phthalic acid anhydride and 1,8-diaminonaphthalene, the compound of formula II and hence the compound of formula I will be obtained as a multiple isomeric mixture.

45 The dyestuffs of formula I and mixtures thereof are useful primarily for spin dyeing plastics in the mass, especially linear, aromatic polyesters. Such polyesters are polycondensates of terephthalic acid, and optionally isophthalic acid, with ethylene glycol and/or cyclohexane-diol. Dyeing in the mass is carried out in accordance with known methods.

The preferred method of producing the mass-dyed polyester is first to mix the dyestuff, suitably

after having been purified and/or ground with a saturated linear, aromatic polyester to form a concentrate or so-called "master-batch". Generally such master-batch contains from 10 to 50 percent by weight of dyestuff and 50 to 90 percent by weight polyester. This master-batch is then added in molten form, in the desired amount, depending on the depth of dyeing desired, to the molten high molecular weight polyester and distributing the same throughout the melt. The preparation of the master-batch is suitably effected by grinding together to form a fine powder and, optionally processing the powder through an extruder to form chippings or granulates. The polyester for the master-batch formation may be one which has a low melting point i.e. a melting point in the range of from 75°C to 230°C and a softening point in the range of from 60°C to 80°C.

After addition of the master batch to the high-molecular weight polyester, the resulting mix can be formed into shaped articles, such as films, foils, fibres or filaments in conventional manner, e.g. by extrusion or spinning and such articles further processed, e.g. into yarn, cord, rope, woven, non-woven and knitted goods, or the mix can be formed into granules or chippings for subsequent melting and formation into such shaped articles.

The dyestuffs as well as being employable for the mass dyeing of polyester by the method described above, may also be incorporated into the polyester by co-condensation with the polymer precursors.

The dyestuffs of formula I dissolve in the polymer melt and remain in the same dissolved form after cooling of the melt.

The dyeings obtained have good fastnesses, especially to light, migration, gas-fumes, ozone and sublimation as well as notable wet-fastnesses. Of particular interest is the stability of the compounds of formula I during the processing of the polyesters e.g. extrusion and exposure to heat.

The following Examples further serve to illustrate the invention. In the Examples all parts are by weight and all temperatures in degrees Centigrade.

25 EXAMPLE 1

54 Parts phthaloperinone are added at 22 to 26° with stirring to 385 parts chlorosulphonic acid. The reaction mixture is stirred for 1 hour at this temperature and then poured onto 700 parts ice. The temperature of the precipitate is kept at 0° with ice. The suspension is subsequently filtered, the residue washed with a mixture containing 1000 parts ice, 1000 parts water and 30 parts 30% aqueous hydrochloric acid and added, with rapid stirring, to a mixture of 400 parts ice, 100 parts water and 23 parts aniline. Cooling is effected to keep the temperature at 1 to 3°. The pH of the reaction mixture is 5.3 and is raised to 7.1 by the addition of 30% aqueous sodium hydroxide. Over a period of 2 hours the temperature is allowed to rise to 25° and stirring is effected at this temperature for 4 hours. The precipitate is suction filtered, washed with water and dried at 100°. 77 Parts of product of formula I in which R₁ is phenyl and the ring A is unsubstituted are obtained. The product is predominantly (96%) a mixture of the two isomers, one where SO₂NHR₁ is bound to the 6-position and one where such group is bound to the 4-position. The product dyes linear, aromatic polyesters in the mass in golden-yellow shades.

Instead of neutralizing with 30% aqueous sodium hydroxide, NaHCO₃ or Na₂CO₃ can advantageously be employed.

EXAMPLE 2

In analogous manner to the procedure described in Example 1, a dyestuff is prepared from the dichlorophthaloperinone mixture (which has been prepared from 1,8-diaminonaphthalene and 3,4-dichlorophthalic acid anhydride) by sulphochlorinating the phthaloperinone and condensing with aniline.

The product essentially comprises 4 isomers, one where the SO₂NHR₁ group is in the 4-position and the 2'- and 3'-positions of ring A bear chlorine atoms, one where the SO₂NHR₁ is in the same position and the 4'- and 5'-positions of the ring A bear chlorine atoms, one where the SO₂NHR₁ group is in the 6-position and the chlorine atoms are in the 2' - and 3'-position and the last where the SO₂NHR₁ group is in the same position and the chlorine atoms are in the 4'- and 5'-positions.

The dyes in the following Table 1 are prepared in analogous manner to the procedure described in Example 1. They correspond to formula I wherein the ring A is unsubstituted, R₁ is a phenyl radical and the substituents thereon are given in the Table. As in the case of the dyestuff of Example 1 a mixture of isomers is obtained, the mixture essentially consists of (96%) the compounds where the SO₂NHR₁ group is bound to the 4- and 6-positions.

TABLE 1

Example No.	Substituent on R ₁	Shade of the substance	shade in polyester
3	3,4-dichloro	yellow-orange	yellow
4	2-CH ₃	yellow-brown	brown
5	4-CH ₃	ochre-yellow	yellow
6	2,5-dimethyl	yellow	do.
7	2-CH ₃ -4-Cl	orange-yellow	do.
8	2-OCH ₃	yellow-orange	do.
9	2-OCH ₃ -4-CH ₃	orange-yellow	do.
10	2-OCH ₃ -4-NO ₂	ochre-yellow	brown
11	3-CONH ₂	orange-yellow	yellow

EXAMPLE 12

26.3 Parts 1,8-diaminonaphthalene and 38 parts of chlorinated phthalic acid anhydride (prepared as described below) are heated to boiling point in 110 parts N-methylpyrrolidone over the period of 1 hour and are kept at this temperature for 1 hour, the reaction-water being distilled off. The mixture is cooled to room temperature and then the mixture, with intensive stirring, is slowly poured onto 1000 parts water, the residue is filtered off and dried. In place of the N-methylpyrrolidone, nitrobenzene or orthodichlorobenzene may also be used.

The chlorination of the phthalic acid is effected in analogy with known methods in water at 45 to 50° by admitting chlorine gas with the pH value being kept at approximately 5 by the addition of aqueous sodium hydroxide. After approximately 10 hours reacting, a mixture containing approximately 63% 4,5-dichlorophthalic acid, 17% 3,4-dichlorophthalic acid, 9% 3,6-dichlorophthalic acid, 6% trichlorophthalic acid and 4% 4-chlorophthalic acid (chromatographic analysis) is obtained. The mixture is heated under low vacuum to approximately 200° to yield the corresponding chlorophthalic acid anhydride. The product is employed as such for the reaction with the 1,8-diaminonaphthalene.

34 Parts of the perinone mixture are added over 30 minutes to 178 parts chlorosulphonic acid and reacted further as described in Example 1. A dyestuff which dyes linear, aromatic polyesters in the mass in yellow-brown shades is obtained.

The dyestuffs in the following Table 2 are prepared in analogous manner. They correspond to formula I wherein R₁ is unsubstituted phenyl and also predominantly consist of the 4- and 6-position isomers.

TABLE 2

Example No.	Substituents on A	Shade on polyester
13	3',4'-dichloro	orange
14	mixture of 3'- and 4'-chloro	yellow-brown
15	2',3',4',5'-tetrachloro	brown-red
16	2',3',4',5'-tetrabromo	do.
17	mixture of 3'- and 4'-nitro	brown

EXAMPLE 18

In analogy with the method of the preceding Examples 13.5 parts phthaloperinone and 20.4 parts tetrachlorophthaloperinone were sulphochlorinated and condensed with 40 parts aniline. The dyestuff obtained dyes linear, aromatic polyesters in orange shades.

Application Example A

1000 Parts conventional polyester obtained from terephthalic acid and ethylene glycol are ground to a powder in a spin mill, the polyester particles then have a diameter of between 300 and 600 μ . This powder is then mixed well with 150 parts dyestuff of Example 1 at room temperature and then

5 processed in an extruder at 130° to form a cable which is then cut to granulate.

5

The concentrate is melted in the shunt current of a helical spinning machine and is added by a metering device to further such polyester in the primary current of the spinning machine heated to 270 to 275°. The metering device adds 4 parts master batch or concentrate to 48 parts polyethylene terephthalate. The mixture is then spun at 270 to 275° at a winding speed of 200 metres per minute,

10 the spun fibres are then stretched at 90° in a drawing machine in a ratio of 1:4 and are twisted in the usual manner in a ring twister. A yellow mass-dyed yarn with good fastnesses is obtained.

10

The dyestuffs of Examples 2 to 18 may also be employed in the same manner to obtain dyeings of the shades indicated in those Examples.

Application Example B

15 1360 Parts of ethylene glycol and 1700 parts dimethyl terephthalate are stirred with 0.55 parts manganese acetate for 3½ hours at 180° and the methanol produced is distilled off.

15

The mass is then transferred to a vacuum container suitable for polycondensation and a mixture of 80 parts ethylene glycol, 0.45 parts antimony trioxide, 20 parts trinonyl phenyl phosphite and 17 parts of the dyestuff of Example 1 (in powder form) are added thereto. With stirring and successively

20 decreasing vacuum to 1 Torr, the whole is heated to 275° until an intrinsic viscosity of $\eta = 0.70$ is reached with ethylene glycol being distilled off.

20


The so-dyed polyester is then extruded into water to cool and is granulated. The granules are vacuum dried at 140° for 16 hours and finally spun, stretched and twisted as described in Example A. A golden-yellow yarn is obtained.

25 The dyestuffs of Examples 2 to 18 may also be employed in the same manner.

25

CLAIMS

1. A dyestuff of formula I,



(I)

in which R₁ is a mono- or bicyclic aromatic radical, and

30 the ring A may bear up to 4 substituents selected from chlorine, bromine and nitro (only 1 nitro group),

30

in which the dyestuff is free from sulpho and carboxy groups, and mixtures of such dyestuffs.

2. A dyestuff according to Claim 1, in which R₁ is a phenyl or naphthyl group and such phenyl is optionally substituted by up to two substituents selected from the group consisting of chlorine, bromine, methyl, methoxy, ethoxy, alkoxyC₁₋₂carbonyl (up to two of each of these), nitro, cyano and aminocarbonyl (only one of any of these).

3. A dyestuff according to Claim 1 or Claim 2, in which the ring A is substituted by up to four substituents selected from chlorine, bromine and nitro (only 1 nitro group).

4. A dyestuff according to any one of Claims 1 to 3, in which R₁ is phenyl optionally substituted by up to two substituents selected from the group consisting of chlorine, methyl, methoxy and ethoxy (up to two of each of these) aminocarbonyl and nitro (only 1 of any of these).

40

5. A dyestuff according to any one of Claims 1 to 4, in which R₁ is phenyl optionally substituted by up to two substituents selected from the group consisting of methyl methoxy, ethoxy and aminocarbonyl (only one of the latter).


6. A dyestuff according to any one of Claims 1 to 5, in which the ring A is unsubstituted or substituted by up to 4 chlorine atoms.

45


7. A dyestuff according to any one of Claims 1 to 6, in which the ring A is unsubstituted or is substituted by two or four chlorine atoms.

50 8. A compound of formula


50




9. A compound of formula



10. A compound of formula



11. A compound according to any one of Claims 1 to 7, in which the aminosulphonyl group is in the 4-position. 5
12. A compound according to any one of Claims 1 to 7, in which the aminosulphonyl group is in the 6-position.
13. A mixture of compounds according to Claim 8, in which the $\text{SO}_2\text{NHC}_6\text{H}_5$ group is the 4- and 6- 10 position.
14. A mixture of the compounds of Claim 9 and 10, in which the $-\text{SO}_2\text{NHC}_6\text{H}_5$ group is the 4- and 6-position. 10
15. A mixture of compounds according to any one of Claims 1 to 7 in which the aminosulphonyl group is in the 4- and 6-position.
16. A dyestuff according to any one of Examples 3 to 18. 15
17. A process for the production of a dyestuff of formula I, as defined in Claim 1, or a mixture thereof, comprising condensing a compound of formula II



or a mixture thereof, with an amine of formula III,

- 20 R_1-NH_2 III 20

18. A process for the production of a dyestuff of formula I or a mixture thereof, substantially as hereinbefore described with reference to any one of Examples 1 to 18.
19. A dyestuff whenever obtained by a process according to Claim 17 or Claim 18.
20. A process for dyeing organic plastics in the mass comprising incorporating a dyestuff 25 according to any one of Claims 1 to 4 or 19 therein. 25
21. A process according to Claim 20, in which the substrate is a linear, aromatic polyester.
22. A process for dyeing linear, aromatic polyesters in the mass substantially as hereinbefore described with reference to Application Example A or B.
23. Mass dyed organic plastics whenever obtained by a process according to any one of Claims 20 30 to 22. 30

2062665 Dated: 21 October 1980 Application No.: 8033855 Published: 28 May 1981
Priority: 25 October 1979 Switzerland 9590/79

SANDOZ LTD., of Lichtstrasse 35, 4002 Basle, Switzerland, a Swiss Body Corporate,

PAUL BUECHELER, Ob. Rebbergweg 93, 4153 Reinach, Switzerland

Improvements in or relating to organic compounds:

AMENDED TITLE AT GRANT: Phthaloperinone dyestuffs: ^{LA} 22.1.86

Address for Service:

B A Yorke & Co., 98 The Centre, Feltham, Middlesex TW13 4EP

Request for examination: 21 OCT 1980		
Application refused		
or withdrawn:		
Patent granted:	WITH EFFECT FROM SECTION 14(1)	18 MAY 1983
Renewal Fee paid in respect of		
5th Year		
6th Year		
7th Year		
8th Year		
9th Year		
10th Year		
11th Year		
12th Year		
13th Year		
14th Year		
15th Year		
16th Year		
17th Year		
18th Year		
19th Year		
20th Year		
Patent ceased or expired:		