



(86) Date de dépôt PCT/PCT Filing Date: 2003/10/08
(87) Date publication PCT/PCT Publication Date: 2004/10/07
(85) Entrée phase nationale/National Entry: 2005/09/22
(86) N° demande PCT/PCT Application No.: EP 2003/011138
(87) N° publication PCT/PCT Publication No.: 2004/084906
(30) Priorité/Priority: 2003/03/25 (PCT/EP03/03096) EP

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 31/519, A61P 9/12, A61P 9/10,
A61P 43/00, A61P 3/10, A61P 39/06, A61P 3/06,
A61P 9/04


(71) Demandeur/Applicant:
VASOPHARM BIOTECH GMBH, DE

(72) Inventeurs/Inventors:
DOBLHOFFER, ROBERT, DE;
TEGTMEIER, FRANK, DE

(74) Agent: BERESKIN & PARR

(54) Titre : UTILISATION DE DERIVES DE PTERIDINE POUR LE TRAITEMENT DE PRESSION INTRACRANIEUNE
ACCRUE, D'ISCHEMIE SECONDAIRE, ET DE TROUBLES ASSOCIES A UN NIVEAU ACCRU D'ESPECES
REACTIVES CYTOTOXIQUES DE L'OXYGENE

(54) Title: USE OF PTERIDINE DERIVATIVES FOR THE TREATMENT OF INCREASED INTRACRANIAL PRESSURE,
SECONDARY ISCHEMIA, AND DISORDERS ASSOCIATED WITH AN INCREASED LEVEL OF CYTOTOXIC
REACTIVE OXYGEN SPECIES



(57) Abrégé/Abstract:

The present invention relates to the use of pteridine derivatives for the treatment of increased intracranial pressure, secondary ischemia, and disorders associated with an increased level of cytotoxic reactive oxygen species.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
7 October 2004 (07.10.2004)

PCT

(10) International Publication Number
WO 2004/084906 A1

(51) International Patent Classification⁷: **A61K 31/519**,
A61P 3/06, 3/10, 9/04, 9/10, 9/12, 39/06, 43/00

(21) International Application Number:
PCT/EP2003/011138

(22) International Filing Date: 8 October 2003 (08.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PCT/EP03/03096 25 March 2003 (25.03.2003) EP

(71) Applicant (for all designated States except US):
VASOPHARM BIOTECH GMBH [DE/DE];
Friedrich-Bergius-Ring 15, 97076 Würzburg (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DOBLHOFER, Robert** [AT/DE]; Tannenstrasse 48, 85640 Putzbrunn (DE). **TEGTMEIER, Frank** [DE/DE]; Am Sprenger 61, 41516 Grevenbroich (DE).

(74) Agent: **VIERING, Hans-Martin**; Viering, Jentschura & Partner, Steinsdorfstr. 6, 80358 München (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF PTERIDINE DERIVATIVES FOR THE TREATMENT OF INCREASED INTRACRANIAL PRESSURE, SECONDARY ISCHEMIA, AND DISORDERS ASSOCIATED WITH AN INCREASED LEVEL OF CYTOTOXIC REACTIVE OXYGEN SPECIES



(57) Abstract: The present invention relates to the use of pteridine derivatives for the treatment of increased intracranial pressure, secondary ischemia, and disorders associated with an increased level of cytotoxic reactive oxygen species.

WO 2004/084906 A1

Use of pteridine derivatives for the treatment of increased intracranial pressure, secondary ischemia, and disorders associated with an increased level of cytotoxic reactive oxygen species

5 The present invention refers to the use of pteridine derivatives for the treatment of increased intracranial pressure and/or secondary ischemia. In particular, these pteridine derivatives provide for an effective treatment of increased intracranial pressure caused by closed cranio-cerebral trauma (CCT). The invention also relates to the use of pteridine derivatives for the treatment of a disorder associated with an increased level of cytotoxic
10 reactive oxygen species.

Traumatic brain injury (TBI) remains a major public health problem globally. In the United States the incidence of closed head injuries admitted to hospitals is conservatively estimated to be 200 per 100,000 population, and the incidence of penetrating head injury is
15 estimated to be 12 per 100,000 population. This yields an approximate number of 500,000 new cases each year, a sizeable proportion of which suffers from significant long-term disabilities.

From the diagnostic point of view, it is distinguished between open and closed CCTs. An
20 open CCT is considered to be an injury in which the cerebral meninges (Dura mater) is mechanically destroyed and the brain is in contact with the environment through this opening. Often, an open CCT is associated with the exit of liquor and brain tissue debris. In a closed CCT the skull remains intact, and the primary damage of the brain (trauma) is characterized by local lesions such as contusions or hematomas and/or diffuse cerebral
25 tissue damage. This primary brain damage, together with possible cardiovascular and respiratory depression, is accompanied by secondary damages, in particular necrosis, apoptosis, edema (vasogenic and/or cellular), secondary bleeding, alterations of cerebral blood volume, disturbed autoregulation of cerebral blood flow and ischemia. Edema, bleeding and an increase of cerebral blood volume as space demanding processes lead to
30 an increase in the intracranial pressure because no volume compensation can occur due to the intact skull cap. This increase in intracranial pressure (ICP) can in turn result in possible acute respiratory depression and thus can be life-threatening itself. Furthermore, the increase in the intracranial pressure can also lead to further secondary brain damages such as the destruction of axons and dendrites of nerve cells due to shearing forces as a
35 result of increased ICP and thus to irreversible permanent disorders of cerebral function of a sensory, motor or intellectual type.

Up to present, no drug is known that can be used for lowering the intracranial pressure or for diminishing the detrimental effects associated with the increase in the intracranial pressure. The pharmaceutical approaches that have been described in conjunction with cranio-cerebral traumas exclusively refer to the prevention of the cell death, in particular to the death of neuronal cells, which occurs as secondary brain damage.

For example, US patent 5,409,935 describes the use of xanthine derivatives for the treatment of secondary nerve cell damage and functional disorders after cranio-cerebral disorders by inhibiting the formation of cell damaging free radicals. According to this US patent, the xanthine derivatives disclosed therein exhibit a potent inhibition of the formation of free radicals in peripheral macrophages and in cultures of activated microglia cells of the brain, i.e. exactly in those two cell types the activation of which has been observed in a multiplicity of neuropathological processes which accompany the death of cerebral tissue in the course of post traumatic brain damage.

The German Offenlegungsschriften 197 40 785 and 197 54 573 as well as the PCT application WO 99/29346 disclose the combined use of 5-HT_{1A} receptor antagonists and calcium channel antagonists for treating cerebral apoplexy and cranio-cerebral trauma. By this approach, the protective effect of 5-HT_{1A} receptor antagonists such as dihydropyridines and calcium channel antagonists such as 2-aminomethyl-chromanes which block the L-type calcium channels of neuronal cells is improved according to these publications. WO 02/069972 discloses triazole compounds which are also said to be specific antagonists of the 5-HT_{1A} receptor and thus to be useful in the prophylaxis and therapy of neurodegenerative diseases, cerebral trauma and cerebral ischemia.

In US Patents 6,469,054 and 6,462,074 aryl sulphonamides and substituted alpha, beta annellated butyrolactones, respectively are disclosed which are to target the CB1 receptor in order to be effective for the treatment of neuronal damage of various cause, for example, cerebral ischemia, cerebral vasospasms or atherosclerotic change.

In addition, the US Patent 6,448,270 discloses 4-substitued piperidine analogs, including hydroxypiperidine and tetrahydropyridine which are said to be selectively active as antagonists of N-methyl-D-aspartate (NDMA) receptor subtypes. According to US Patent 6,448,270 these compounds inhibit the excitotoxic action of excitatory amino acids glutamate and aspartate at the N-methyl-D-aspartate (NDMA) receptor, which excitotoxic action is considered responsible for the loss of neurons in cerebrovascular disorders which result from conditions such as cerebral vasospasms, hypoglycemia or cerebral trauma.

However, as mentioned above, no efforts have been taken yet to pharmacologically overcome the increase in the intracranial pressure. Up to now, this increase can only be prevented either by surgically opening the skull of the affected patient in order to provide a volume compensation by surgical drainage of the ventricles, or by the use of osmotherapeutics such as mannitol or sorbitol. These osmotherapeutics are infused into the blood stream where they provide for a difference in the osmotic pressure between the blood stream and the intracranial parenchymal region. This gradient leads to a decrease of the intracranial pressure (cf. McGraw CP, Alexander E Jr, Howard G Surg Neurol 1978 Aug, 10(2):127-30 or McGraw CP, Howard G., Neurosurgery 1983 Sep, 13(3):269-71 for example).

However, both methods suffer from serious disadvantages. Opening of the skull is a severe surgical intervention which can obviously jeopardize the life especially of patients with severe closed CCT and increases the risk for serious bacterial infections and requires specific arrangements to prevent sepsis. Moreover, this surgery requires post-operative treatment specially trained personal so that it is only carried out at selected clinics.

Administration of osmotherapeutics only provides a rather short term temporary effect, followed by a subsequent new increase in the intracranial pressure that can sometimes already be observed within about 10 to 15 minutes after administration. The osmotherapeutical approach is further complicated by the fact that the applied initial amount of osmolyte, e.g. mannitol, strongly influences the size of subsequent doses of the osmolyte. Initial administration of more osmolyte than absolutely needed will lead to larger doses being required to control the ICP. Therefore, the initial dose must be kept as low as possible which limits the osmotherapeutical approach. And, since the difference in the osmotic pressure of the blood and the brain gets smaller by each application, osmotherapeutics can usually only be administered two or three times. Finally, this non-surgical treatment bears the risk of brain ischemia and rebound effects on the intracranial pressure. Moreover this treatment is believed to harm the integrity of the blood brain barrier and induce secondary brain inflammation due to infiltration of the cerebral parenchyma with blood constituents.

Therefore, there is a need for a drug that reduces the intracranial pressure caused by traumatic brain injury and that alleviates the pathophysiological conditions associated with the increased intracranial pressure.

This problem is solved by the use of a pteridine derivative of one of the general formulae (Ia) to (Ie). Compounds of formulae (Ia) to (Ie) are known to be inhibitors of nitric oxide (NO) synthase (see for, example, US Patent 5,902,810; WO 95/31987; WO 95/32203; WO 01/21619; or US Patent 5,922,713).

5

On the molecular level, the present invention is based on the finding that the ability of the compounds according to formulae (Ia) to (Ie) to be inhibitors of nitric oxide (NO) synthases can be used to inhibit the NO production preferentially in areas of increased activity of NO synthase, in particular that of the endothelial NO synthase (e-NOS), which
10 is hyper activated in the region of trauma.

By doing so, the vasodilatory effect of NO on the blood vessels in the brain is prevented. In the event of a cranio-cerebral trauma this leads to the effect that the blood vessels in the brain do not swell but rather maintain a constant volume or even re-contract whereby an
15 already developed dilatation will be partly overcome. Accordingly, additional intracranial space is created which can be occupied by the damaged cerebral tissue which is accompanied with a reduction (in the increase) of the intracranial pressure. To the surprise of the inventors some of the pteridine compounds used in the present invention have only a very little or no tendency at all to cross the blood-brain-barrier. This property in connection
20 with the specificity for NO synthases allows administering the compounds of formulae (Ia) to (Ie) in high dosages rendering them ideal candidates for use in emergency situations and thus for decreasing the critically elevated intracranial pressure that usually occurs after closed cranio-cerebral trauma or non-traumatic brain injury. On the other hand, as a long term effect of closed cranio-cerebral trauma or non-traumatic brain injury, usually the
25 blood-brain-barrier becomes permissive. This in turn enables the pteridine derivatives used in the present invention to pass (penetrate through) the blood-brain-barrier and to target the NO synthases (neuronal NOS, inducible NOS, mitochondrial NOS) particularly expressed in the brain parenchyma. Thereby, as a secondary use of the pteridine derivatives described here, secondary brain damage such as inflammatory processes, acute cell death via
30 necrosis, cell death via necrosis and apoptosis and formation of edema can be antagonized (cf. Fig. 1.).

The compounds disclosed in the present invention are thus useful for the treatment of increased intracranial pressure, in particular an elevated intracranial pressure that is caused
35 by closed cranio-cerebral trauma. The compounds can especially be administered in case of critically increased intracranial pressure and also secondary ischemia. The use of the compounds of formulae (Ia) to (Ie) for the treatment of increased intracranial pressure that

is caused by non-traumatic brain injury, e.g. by stroke or cold lesion, is also contemplated in the present invention.

It is further known that tetrahydrobiopterin is unstable at physiological pH and prone to decompose in oxygenated solutions. The loss of enzyme-associated tetrahydrobiopterin cofactor leads to monomerisation and inactivation of NOS. Although, the underlying mechanism by which tetrahydrobiopterin stabilizes the NOS dimer complex is unclear and appears to be isoform-specific, tetrahydrobiopterin may act as a molecular clip to prevent NOS subunit dissociation (cf. Crane, et al., (1998) Science 279: 2121-2126). Apparently, tetrahydrobiopterin induces a large conformational change upon binding to the NOS oxygenase domain. Decoupling of NO-synthase can occur when its co-factor tetrahydrobiopterin is depleted or oxidized and it has been shown to happen in the brain after TBI and similar pathophysiologies such as cerebral ischemia. It is also an important mechanism in endothelial dysfunction and its related diseases.

The conversion of arginine to N-hydroxy-L-arginine (NHA) and of NHA to L-citrulline and NO both depend on the presence of tetrahydrobiopterin. In the absence of sufficient tetrahydrobiopterin, NOS will switch from NO synthesis to superoxide anion generation (Knowles, R.G. and Moncada, S. (1994) Biochem. J. 298: 249-258; Pou, S. et al., (1992) J. Biol. Chem. 267: 24173-24176), subsequently leading to NO depletion and oxidant accumulation (Beckman, J.S. et al., Methods of Enzymology, Vol.233, Part C: Oxygen Radicals in Biological Systems. L. Packer (Ed.), Academic Press, Inc., San Diego, CA 229-240) which causes further decoupling of NOS. This vicious cycle leads to a burst of cytotoxic reactive oxygen species (ROS) such as superoxide.

Superoxide, the main product of decoupled NOS, quickly reacts with NO to form the even more highly cytotoxic and stable peroxynitrite which is likely to be responsible for most of the cell damage that occurs in the injured brain. This reaction is even faster than the one of superoxide anion with superoxide dismutase to form hydrogen peroxide and O₂. Any peroxynitrite produced when protonated forms peroxynitrous acid which usually undergoes isomerisation to form hydrogen cation and nitrate anion. Furthermore, peroxynitrous acid may undergo homolytic cleavage to a hydroxyl free radical and nitrogen dioxide free radical, or heterolytic cleavage to a nitronium cation and hydroxide anion. Three of these cleavage products (hydroxyl free radical, nitronium cation and nitrogen oxide free radicals), are among the most reactive and damaging species in biological systems.

The NOS-catalyzed formation of superoxide anion and its subsequent transformation into peroxynitrous acid cleavage products, or its dismutation into hydrogen peroxide and hydroxyl free radical plays a central role in the endothelial dysfunction and oxidative vascular injury described in a number of vascular diseases.

5

In addition to their ability to inhibit the production of NO by NO-synthases, the pteridines of formulae (Ia) to (Ie) can inhibit the production of cytotoxic reactive oxygen species such as superoxide from decoupled NO-synthase. The pteridines of formulae (Ia) to (Ie) act as antagonists of endogenous tetrahydrobiopterin. Like tetrahydrobiopterin they have the ability to stabilize the NOS-homodimers, but in contrast to the endogenous co-factor, they inhibit the production of NO. On the other hand, by stabilizing the dimeric structure of the enzyme, the pteridines also inhibit decoupling of NO-synthase and thus the production of cytotoxic peroxynitrite. This dual mechanism of action leads to beneficial effects of the compounds both on the vascular and on the cellular level.

15

The compounds disclosed in the present invention are thus not only useful for the treatment of increased intracranial pressure, in particular an elevated intracranial pressure that is caused by closed cranio-cerebral trauma. Rather, they can also be used in the treatment of other disorders which are associated with an increased level of cytotoxic reactive oxygen species generated by a decoupled NO-synthase. Accordingly, the use of the compounds of formulae (Ia) to (Ie) for the treatment of increased intracranial pressure that is caused by non-traumatic brain injury, e.g. by stroke or cold lesion, and for alleviation of oxidative stress derived from decoupled NO-synthase is also contemplated in the present invention.

20

25 In the compounds of formulae (Ia) to (Ie) used in the present invention, the following definitions apply, if not stated otherwise.

If groups or substituents occur more than once in the compounds of the following formulae (Ia) to (Ie), they may all, independently of one another, have the stated meanings and may in each case be identical or different.

30

Alkyl radicals in the compounds used in the present invention may be straight-chain or branched. This also applies if they are present in other groups, for example in alkoxy groups, alkoxy carbonyl groups or in amino groups, or if they are substituted. Alkyl radicals normally contain one to twenty carbon atoms, preferably one to ten carbon atoms. Examples of alkyl groups are methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl,

35

nonyl, decyl, the n isomers of these radicals, isopropyl, isobutyl, isopentyl, sec-butyl, tert-butyl, neopentyl, 3,3-dimethylbutyl.

5 Examples of alkenyl radicals are straight-chain or branched hydrocarbon radicals which contain one or more double bonds. Alkenyl radicals normally contain two to twenty carbon atoms and one or two double bonds, preferably two to ten carbon atoms and one double bond.

10 Examples of alkynyl radicals are straight-chain or branched hydrocarbon radicals which contain one or more triple bonds. Alkynyl radicals normally contain two to twenty carbon atoms and one or two triple bonds, preferably two to ten carbon atoms and one triple bond.

15 Examples of alkenyl radicals are the vinyl radical, the 2-propenyl radical (allyl radical), the 2-butenyl radical and the 2-methyl-2-propenyl radical.

20 Examples of alkynyl radicals are the ethynyl radical, the 2-propynyl radical (propargyl radical) or the 3-butenyl radical.

25 Cycloalkyl radicals are saturated cyclic hydrocarbons which normally contain three to eight ring carbon atoms, preferably five or six ring carbon atoms. Cycloalkyl radicals may in turn be substituted.

30 Examples of cycloalkyl radicals are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, all of which may also be substituted for example by one or more identical or different (C₁-C₄)-alkyl radicals, in particular by methyl. Examples of such substituted cycloalkyl radicals are 4-methylcyclohexyl or 2,3-dimethylcyclopentyl.

35 Cycloalkenyl radicals are unsaturated cyclic hydrocarbons which normally contain three to eight ring carbon atoms, preferably five or six ring carbon atoms. Cycloalkenyl radicals preferably have a double bond in the ring system. Cycloalkenyl radicals may in turn be substituted.

40 Cycloalkylalkyl radicals are saturated hydrocarbons which are derived from a cycloalkyl-substituted alkyl group. The cycloalkyl group normally has five to six ring carbon atoms. Examples of cycloalkylalkyl radicals are cyclopentylmethyl, cyclopentylethyl, cyclohexylethyl and, in particular, cyclohexylmethyl. Cycloalkylalkyl radicals may in turn be substituted.

Aryl is a carbocyclic or heterocyclic aromatic radical, preferably phenyl, naphthyl or heteroaryl. Aryl radicals may be unsubstituted or substituted. Substituents are one or more identical or different monovalent organic radicals, for example or from the series halogen, alkyl, phenyl, -OH, -O-alkyl, alkylendioxy, -NR₈R₉, -NO₂, -CO-(C₁-C₅)-alkyl, -CF₃, -CN, -CONR₈R₉, -COOH, -CO-O-(C₁-C₅)-alkyl, -S(O)_n-(C₁-C₅)-alkyl, -SO₂-NR₈R₉.

Alkylaryl is an alkyl-substituted aryl radical, preferably (C₁-C₃)-alkylaryl, in particular methylphenyl.

Arylalkyl is an aryl-substituted alkyl radical, preferably phenylmethyl or 2-phenylethyl.

Heteroaryl or a heterocyclic aromatic radical is preferably a 5- to 7-membered unsaturated heterocycle which has one or more heteroatoms from the series O, N, S.

Examples of heteroaryls from which the radicals occurring in compounds of the formula I may be derived are pyrrole, furan, thiophene, imidazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, 1,3-oxazole, 1,2-oxazole, 1,3-thiazole, 1,2-thiazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, pyran, thiopyran, 1,4-dioxin, 1,2-oxazine, 1,3-oxazine, 1,4-oxazine, 1,2-thiazine, 1,3-thiazine, 1,4-thiazine, 1,2,3-triazine, 1,2,4-triazine, 1,3,5-triazine, 1,2,4,5-tetrazine, azepine, 1,2-diazepine, 1,3-diazepine, 1,4-diazepine, 1,3-oxazepine or 1,3-thiazepine.


The radicals derived from the heterocycles may be bonded via any suitable carbon atom.

Nitrogen heterocycles which have a hydrogen atom (or a substituent) on a ring nitrogen atom, for example pyrrole, imidazole, etc, may also be bonded via a ring nitrogen atom, especially if the relevant nitrogen heterocycle is bonded to a carbon atom. A thienyl radical may, for example, be in the form of a 2-thienyl radical or 3-thienyl radical, a furan radical in the form of a 2-furyl radical or 3-furyl radical, a pyridyl radical in the form of a 2-pyridyl radical, 3-pyridyl radical or 4-pyridyl radical.

Halogen is fluorine, chlorine, bromine or iodine, preferably fluorine or chlorine.

In preferred embodiments of the invention the use of pteridine derivatives having an amino group, either unsubstituted or substituted, in the 2- and/or the 4-position of the ring system is preferred.

In one of these preferred embodiments, the invention relates to the use of pteridines of formula (Ia)




- 5 for treating increased intracranial pressure and/or secondary ischemia, wherein in formula (Ia) R_1 , R_3 are independently selected from H or OH, R_4 is H, CH_3 , CH_2OH , CHO, R_2 is H, CH_3 , CH_2OH , CHO or a lower C1-C9 alkyl radical, which can be a straight or a branched chain, as well as $(CH(OH))_n-Y$ or $(CH(OH))_n-(CH_2)_m-W$, wherein Y is hydrogen or a lower alkyl radical, W is hydrogen or a hydroxyl group, an n and m are
 10 independently from each other 1-20.

The compounds of formula (Ia) can be prepared as described in US Patent 5,922,713 or Werner, Ernst R. et al, Biochem. J., Vol. 320, Nov. 1996, 193-196. Moreover, these compounds are also commercially available and for example, can be purchased from
 15 Schircks Laboratories, Jona, Switzerland.

In a particularly preferred embodiment, the compounds of formula (Ia) R_1 and R_4 (Ia) are hydrogen, R_2 is methyl and R_3 is hydroxyl. A compound that is preferably used in the present invention is the 6-(L-erythro) isomer, i.e. 2,4-diamino-5,6,7,8-tetrahydro-6-(L-
 20 erythro-1,2-dihydroxypropyl)-pteridine.

A further preferred embodiment of 2,4-diamino-substituted pteridines for treating increased intracranial pressure and/or secondary ischemia is directed to the use of compounds of the formula (Ib)

10



(Ib)

5


wherein in formula (Ib) R_1 , R_2 , R_3 and R_4 are independently from each other H or OH, R_5 is H, CH_3 , CH_2OH , CHO or a lower C_1 - C_9 alkyl radical, which can be a straight or a branched chain, as well as $(CH(OH))_n-Y$ or $(CH(OH))_n-(CH_2)_m-W$, wherein Y is hydrogen or a lower alkyl radical, W is hydrogen or a hydroxyl group, an n and m are

10

In a preferred compound of formula (Ib) R_5 is methyl, R_3 is hydrogen, R_2 and R_4 are both hydrogen and R_1 is either hydrogen or hydroxyl. The compounds of formula (Ib) can be prepared as described in EP 0 906 913.

15

In a further embodiment the present invention relates to the use of 4-aminopteridines of the general formula (Ic)




(Ic)

20

for treating increased intracranial pressure and/or secondary ischemia, wherein in formula (Ic)

25

A is a bridge of the form



R₁ is hydrogen, (C₁-C₂₀)-alkyl, (C₁-C₂₀)-alkenyl, (C₁-C₂₀)-alkynyl, preferably (C₁-C₁₀)-alkyl, cycloalkyl, cycloalkenyl, preferably (C₃-C₈)-cycloalkyl, cycloalkylalkyl, aryl, alkylaryl, preferably (C₁-C₃)-alkylaryl or arylalkyl, where the organic radicals, preferably the alkyl radicals, may be substituted by one or more substituents, preferably by substituents R₆,

R₂ is, independently of R₁, hydrogen, (C₁-C₂₀)-alkyl, (C₁-C₂₀)-alkenyl, (C₁-C₂₀)-alkynyl, preferably (C₁-C₁₀)-alkyl, cycloalkyl, cycloalkenyl, preferably (C₃-C₈)-cycloalkyl, cycloalkylalkyl, aryl, alkylaryl, preferably (C₁-C₃)-alkylaryl, or arylalkyl, where the organic radicals, preferably the alkyl radicals, may be substituted by one or more substituents, preferably by substituents R₆,

R₁ and R₂ may, together with the nitrogen atom bearing them, form a 3-8-membered ring which may optionally contain 0, 1 or 2 further heteroatoms from the series N, O, S and which is optionally substituted by one or more radicals, preferably R₆ radicals,

R₃ is hydrogen, -CO-alkyl, preferably -CO-(C₁-C₇)-alkyl, -CO-alkylaryl, preferably -CO-(C₁-C₃)-alkylaryl or -CO-aryl,

R₄ is alkyl, alkenyl, alkynyl, preferably (C₁-C₁₀)-alkyl, cycloalkyl, cycloalkenyl, preferably (C₃-C₈)-cycloalkyl, cycloalkylalkyl, aryl or alkylaryl, preferably (C₁-C₃)-alkylaryl, arylalkyl, -CO-O-alkyl, preferably -CO-O-(C₁-C₅)-alkyl, -CO-O-aryl, -CO-alkyl, preferably -CO-(C₁-C₅)-alkyl or -CO-aryl, where the organic radicals, preferably the alkyl radicals, may be substituted by one or more substituents, in particular by substituents R₇,

R₅ is, independently of R₃, hydrogen, -CO-alkyl, preferably -CO-(C₁-C₇)-alkyl, -CO-alkylaryl, preferably -CO-(C₁-C₃)-alkylaryl or -CO-aryl,

R₆ is -F, -OH, -O-(C₁-C₁₀)-alkyl, -O-phenyl, -O-CO-(C₁-C₁₀)-alkyl, -O-CO-aryl, -NR₈R₉, oxo, phenyl, -CO-(C₁-C₅)-alkyl, -CF₃, -CN, -CONR₈R₉, -COOH, -CO-O-(C₁-C₅)-alkyl, -CO-O-aryl, -S(O)_n-(C₁-C₅)-alkyl, -SO₂-NR₈R₉,

5

R₇ has, independently of R₆, one of the meanings of R₆,

R₈ is hydrogen or (C₁-C₂₀)-alkyl, preferably (C₁-C₅)-alkyl,

10 R₉ is hydrogen, (C₁-C₂₀)-alkyl, preferably (C₁-C₅)-alkyl or aryl, preferably phenyl,

R₁₀ is hydrogen, (C₁-C₂₀)-alkyl, preferably (C₁-C₅)-alkyl, alkoxy or aryl,

15 aryl is preferably phenyl, naphthyl or heteroaryl, each of which may be unsubstituted or substituted, for example may be substituted by one or more identical or different substituents from the series halogen, (C₁-C₂₀)-alkyl, preferably (C₁-C₅)-alkyl or phenyl, -OH, -O-(C₁-C₂₀)-alkyl, preferably -O-(C₁-C₅)-alkyl, (C₁-C₂₀)-alkylenedioxy, preferably (C₁-C₂)-alkylenedioxy, -N₈R₉, -NO₂, -CO-(C₁-C₅)-alkyl, -CF₃, -CN, -CONR₈R₉, -COOH, -CO-O-(C₁-C₅)-alkyl, -S(O)_n-(C₁-C₅)-alkyl, 20 -SO₂-NR₈R₉,

heteroaryl is a 5- to 7-membered unsaturated heterocycle which contains one or more heteroatoms from the series O, N, S,

25 n is 0, 1 or 2,

in all their stereoisomeric and tautomeric forms and mixtures thereof in all ratios, and their physiologically tolerated salts, hydrates and esters. The compounds of formula (Ic) can be synthesized as described in WO 01/21619 or WO 00/39129. 2,4-diaminopteridine derivatives of formula (Ic) can also be obtained by the process described in WO 97/21711 30 using for example 2,4,5,6-tetraaminopyrimidine-dihydrochloride as starting material which is reacted with a respective oximine-derivative according to formula (II) of WO 97/21711.

In preferred embodiments of formula (Ic)

35

R₁ is preferably hydrogen, (C₂-C₄)-alkyl which may be substituted by one or more substituents R₆, or (C₁-C₂)-alkylaryl, and R₁ is particularly preferably arylmethyl

R₂ is preferably (C₂-C₄)-alkyl which may be substituted by one or more substituents R₆, or (C₁-C₂)-alkylaryl, and R₂ is particularly preferably arylmethyl

5 in addition, R₁ and R₂ preferably form, together with the nitrogen atom bearing them, a 5-7-membered ring which preferably contains no or only one other heteroatom from the series N, O, S. Very particularly preferred rings of this type are pyrrolidine, piperidine, morpholine, dimethylmorpholine, thiomorpholine or N-(C₁-C₂)-alkylpiperazine, where these rings themselves may also be substituted, for
10 example by -OH, -O-(C₁-C₃)-alkyl, -NR₈R₉ or -COOH.

R₃ is preferably hydrogen, CO-(C₁-C₃)-alkyl or CO-aryl, and R₃ is very particularly preferably hydrogen.

15 R₄ is preferably aryl, (C₁-C₃)-alkyl which may be substituted by one or more substituents R₇, or -CO-O-aryl. Particularly preferred R₄ radicals are aryl and 1,2-dihydroxypropyl.

R₅ is preferably hydrogen.

20

R₆ is preferably -OH, -O-(C₁-C₃)-alkyl, -NR₈R₉ or -COOH.

R₇ is preferably -OH, -O-(C₁-C₁₀)-alkyl, phenoxy, oxo, particularly preferably -OH, decyloxy and phenoxy.

25


R₁₀ is preferably hydrogen or methyl.

aryl is preferably phenyl, thiophenyl, furyl and pyridyl, and phenyl is particularly preferred, all of which can be substituted as described. Preferred substituents are
30 (C₁-C₃)-alkyl, halogen and (C₁-C₃)-alkyloxy and (C₁-C₂)-alkylenedioxy. The preferred number of substituents on aryl radicals is 0, 1 or 2; phenyl substituents are preferably in the meta or para position, and in the case of two substituents in the 3 and 4 positions.

35 n is preferably 0 and 2

Particularly preferred is the use of 5,6,7,8-tetrahydropteridines of formula (Ic) in which R₁ and R₂ are both methyl, ethyl, propyl or R₁ and R₂ form together with the nitrogen atom bearing them, a 5 or 6-membered ring which preferably contains no or only one other heteroatom from the series N, O, S, and wherein R₄ is benzyl, or thiophenyl which is optionally substituted with (C₁-C₃)-alkyl, halogen and (C₁-C₃)-alkyloxy and (C₁-C₂)-alkylenedioxy.


In a further aspect, the invention relates to the use of compounds of the general formula (Id)



for treating increased intracranial pressure and/or secondary ischemia, wherein in formula (Id)

- | | | |
|----|----------------|--|
| 15 | X | is O or NH; |
| | R ₁ | is hydrogen, methyl, (C ₁ -C ₅)-alkanoyl, nicotinoyl or (1-methyl-3-pyridinio)carbonyl; |
| | R ₂ | is hydrogen or methyl; |
| 20 | R ₃ | is hydrogen, methyl, ethyl, benzyl, (C ₁ -C ₅)-alkanoyl, unsubstituted benzoyl, substituted benzoyl, pyridoyl, thienylcarbonyl, one of the radicals |

15



the radical $R_9R_{9a}N-CO-$, the radical $R_9R_{9a}N-CS-$, phenoxycarbonyl, or
 5 benzyloxycarbonyl;

R_4 is hydrogen, (C_2-C_5) -alkyl, unsubstituted phenyl, substituted phenyl or the
 radical $R_{4a}-CH_2-$;

R_{4a} is hydrogen, (C_1-C_4) -alkylmercapto, the radical $-S(O)_mR_{10}$, where m is the
 number 1 or 2, the radical $-NR_{11}R_{12}$ or the radical $-OR_{13}$, or

10 R_3 and R_{4a} together are the group $-CO-O-$, its carbonyl carbon atom being bonded
 to the 5-position of the pteridine molecule;

R_5 is hydrogen or phenyl;

R_6 is hydrogen;

R_7 is hydrogen or methyl;

15 R_8 is (C_1-C_{10}) -alkyl or benzyl;

R_9 is hydrogen, (C_1-C_6) -alkyl, cyclohexyl, phenyl or benzoyl;

R_{9a} is hydrogen, methyl or ethyl;

R_{10} is methyl;


R_{11} and R_{12} independently of one another are hydrogen or methyl;

20 R_{13} is hydrogen, (C_1-C_{10}) -alkyl, 2-methoxyethyl, phenyl, 3-phenylpropyl, 3-
 cyclohexylpropyl, (C_1-C_5) -alkanoyl, hydroxyacetyl, 2-amino- (C_2-C_6) -
 alkanoyl, which is unsubstituted or substituted in the alkyl moiety by a
 phenyl radical, or $((C_1-C_2)$ -alkoxy)carbonyl;

A is a pharmacologically tolerable anion; and their tautomeric forms and their pharmacologically tolerable salts.

Preferred compounds of formula (Id) are the compounds in which X = O, R₁, R₂, R₅ and R₆ represent H, R₃ represent H, methyl, or ethyl and R₄ is CH₂NH₂, CH₂NHMe, or CH₂NMe₂.

In yet another embodiment, the invention relates to the use of compounds of the general (Ie)



for treating increased intracranial pressure and/or secondary ischemia, wherein in formula (Ie)

X is O, NH or N-(C₁-C₅)-alkanoyl;

R is hydrogen and

R₁ is hydrogen or (C₁-C₅)-alkanoyl or R and R₁ together

with the nitrogen atom to which they are bonded from a dimethylaminomethyleneamino group;

R₂ is hydrogen, methyl, phenyl, hydroxyl, methoxy or amino;

R₃ is the radical -OR₄, -NR₅R₆ or -S(O)_mR₇, where m represent the numbers 0, 1 or 2;

R₄ is hydrogen, (C₁-C₁₀)-alkyl, cyclohexyl, benzyl, phenyl which is unsubstituted or substituted by chlorine or the radical -COR₈, aminocarbonylmethyl which is unsubstituted or substituted on the nitrogen by one or two identical or different (C₁-C₄)-alkyl radicals, 2-methoxyethyl, the (2,2-dimethyl-1,3-dioxolan-4-yl)methyl radical or the radical -COR₉;

R₅ is hydrogen, methyl, ethyl, 2-hydroxyethyl, 2-chloroethyl, benzyl, pyridylmethyl, phenylethyl, pyridylethyl or acetyl;


R₆ independently of the meaning of R₅ has the meanings indicated for R₅ or, if R₅ is hydrogen or methyl, is also cyclohexyl, 3-(2-ethoxyethoxy)-propyl, benzyl which carries

one or two chlorine atoms or the radical $-\text{COR}_{10}$ on the phenyl ring, (C_1-C_5) -alkanoyl, the radical $-\text{COR}_{10}$ or the radical $-(\text{CH}_2)_4-\text{COR}_{10}$;


R_7 is (C_1-C_4) -alkyl, benzyl, phenyl which is unsubstituted or substituted by chlorine, the radical $-\text{COR}_8$ or the radical $-\text{CO}-\text{O}-\text{CO}-(\text{C}_1-\text{C}_4)$ -alkyl or naphthyl;

5 R_8 is hydrogen, methoxy, amino or R_{10} ;

R_9 is (C_1-C_4) -alkyl, hydroxymethyl, trifluoromethyl, (C_1-C_2) -alkoxy or R_{11} ; R_{10} is the radical



R_{11} is the radical



10

R_{12} is hydroxyl or (C_1-C_2) -alkoxy;

R_{13} is (C_1-C_4) -alkyl or benzyl;

R_{14} is hydrogen or benzyloxycarbonyl; and their tautomeric forms and their pharmacologically tolerable salts.

15

In preferred embodiments, the use of compounds wherein R is hydrogen, and X is O or NH is preferred. Furthermore, the use of compounds of formula (1e) with R_1 and/or R_2 is hydrogen is also preferred.

20

As mentioned above, the present invention also encompasses the use of corresponding physiologically or toxicologically acceptable salts, in particular the pharmaceutically usable salts of the compounds according to formulae (Ia) to (Ie). Thus, the compounds of any of the formulae (Ia) to (Ie) which contain acidic groups may, for example, be in the form of alkali metal salts, alkaline earth metal salts or of ammonium salts and these groups can be used according to the invention. Examples of such salts are sodium salts, potassium salts, calcium salts, magnesium salts or salts with ammonia or organic amines such as, for example, ethylamine, ethanolamine, triethanolamine or amino acids. Compounds of the formulae (Ia) to (Ie) which contain one or more basic, that is protonatable, groups may be used in the form of their acid addition salts with physiologically tolerated inorganic or organic acids and used according to the invention, for example as salts with hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, nitric acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acids, oxalic acid, acetic acid, tartaric acid,

25
30

lactic acid, salicylic acid, benzoic acid, formic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, malic acid, sulfamic acid, phenylpropionic acid, gluconic acid, ascorbic acid, isonicotinic acid, citric acid, adipic acid etc.

5

If a compound according to any of the formulae (Ia) to (Ie) contain both acidic and basic groups in the molecule, the invention also includes inner salts or betaines (zwitterions) in addition to the salt forms described.

10

Salts can be obtained from compounds of the formula I by conventional processes known to the person skilled in the art, for example by combining with an organic or inorganic acid or base in a solvent or dispersant, or else by anion exchange or cation exchange from other salts. The present invention further encompasses the use of all solvates of compounds of formulae (Ia) to (Ie), for example hydrates or adducts with alcohols, and derivatives of the compounds of the formula I such as, for example, esters, and prodrugs and active metabolites.

15

In accordance with the above disclosure, the present invention is also directed to a method of treating a subject having an increased intracranial pressure and secondary ischemia comprising administering to the subject a therapeutically sufficient amount of a compound of any of the general formulae (Ia) to (Ie). The application of a compound described here is preferred, when the increased intracranial pressure is caused by a closed craniocerebral trauma.

20

25

In another aspect, the present invention is directed to a method of treating a subject having a disorder associated with an increased level of cytotoxic reactive oxygen species generated by a decoupled NO-synthase. For example, these cytotoxic reactive oxygen species encompass superoxide anion, peroxinitrite, hydroxyl free radical, nitrogen dioxide free radical and nitronium cation. Disorders which are associated with an increased level of cytotoxic reactive oxygen species generated by a decoupled NO-synthase include, but are not limited to, elevated intracranial pressure that is caused by closed craniocerebral trauma or non-traumatic brain injury. These disorders also include any disorder that is associated with endothelial dysfunction. Examples of such a disorder are hypercholesterolemia, endothelial damage due to mechanical und chemical noxia, peripheral arterial disease and cardiovascular diseases such as arteriosclerosis, heart disease, coronary heart disease, diabetic micro- and macroangiopathy, heart failure, stroke, angina pectoris, hypertension, ischemia/reperfusion injury, and insulin-resistant diabetes. The term "endothelial

30

35

dysfunction” is used herein in accordance with its regular meaning as given, for example, Widlansky M.E. et al., J. Am. Coll. Cardiol. 2003 Oct 1; 42(7):1149-60. Thus, endothelial dysfunction means that the endothelium of the vessels cannot function in a normal way anymore, e.g. due to oxidative stress (the presence of reactive oxygen species).

5

The compounds of the general formulae (Ia) to (Ie) exert this function by inhibiting decoupling of NO-synthase and thus the production of cytotoxic peroxynitrite, superoxide and related reactive oxygen species. In one of these embodiments, the endothelial NO-synthase is inhibited.

10

Subjects that are treated in the present invention are preferably mammals such as humans, monkeys, cats, dogs, mice or rat with humans being preferred.

15

For lowering the (enhanced) intracranial pressure, the compounds of formulae (Ia) to (Ie) can be administered orally, for example in the form of pills, tablets, film-coated tablets, sugar-coated tablets, granules, hard and soft gelatin capsules, aqueous, alcoholic or oily solutions, syrups, emulsions or suspensions, or rectally, for example in the form of suppositories. The administration can also take place parenterally, for example subcutaneously, intramuscularly or intravenously in the form of injection solutions or infusion solutions. Further suitable administration forms are, for example, percutaneous or topical administration, for example in the form of ointments, tinctures, sprays or transdermal therapeutic systems, or inhalational administration in the form of nasal sprays or aerosol mixtures, or, for example, microcapsules, implants or rods. One preferred mode of administration, especially in case of emergencies, is by injection or infusion in an aqueous pharmaceutically acceptable solution.

20

The corresponding pharmaceutical compositions that are used according to the invention can be produced by the standard processes known for producing pharmaceutical products.

30

For this purpose, one or more compounds of the formulae (Ia) to (Ie) and/or their physiologically tolerated salts, esters and hydrates are converted together with one or more solid or liquid pharmaceutical carriers and/or additives or excipients and, if desired, in combination with other active pharmaceutical ingredients with therapeutic or prophylactic action into a suitable administration form or dosage form, which can then be used as pharmaceutical in human medicine or veterinary medicine. The pharmaceutical products comprise a therapeutically or prophylactically effective dose of the compounds of the

35

formulae (Ia) to (Ie) and/or their physiologically tolerated salts, esters and hydrates, which normally amounts to from 0.5 to 90% by weight of the pharmaceutical product.

To produce, for example, pills, tablets, sugar-coated tablets and hard gelatin capsules it is possible to use lactose, starch, for example corn starch or starch derivatives, talc, stearic acid or salts thereof etc. Carriers for soft gelatin capsules and suppositories are for example fats, waxes, semisolid and liquid polyols, natural or hydrogenated oils etc. Examples of carriers suitable for producing solutions, for example injection solutions, or emulsions or syrups are water, physiological saline, alcohols such as ethanol, glycerol, polyols, sucrose, invert sugar, glucose, mannitol, cyclodextrine, vegetable oils etc. The compounds of the formulae (Ia) to (Ie) and their physiologically tolerated salts, esters and hydrates may also be lyophilized, and the resulting lyophilizates can be used together with a reconstitution solution, for example, for producing products for injection or products for infusion. Examples of carriers suitable for microcapsules, implants or rods are copolymers of glycolic acid and lactic acid.

The pharmaceutical products may besides the active ingredients and carriers also comprise conventional additives, for example fillers, disintegrants, binders, lubricants, wetting agents, stabilizers, emulsifiers, dispersants, preservatives, sweeteners, colorants, flavoring or aromatizing agents, thickeners, diluents, buffer substances, also solvents or solubilizers or means to achieve a depot effect, salts to alter the osmotic pressure, coating agents or antioxidants.

The dosage of the active ingredient of the formula I to be administered, and/or of a physiologically tolerated salt, ester or hydrate thereof depends on the individual case and should be adapted to the individual circumstances for an optimal effect in the conventional way. Thus, it depends on the nature and severity of the disease to be treated and on the sex, age, weight and individual response of the human or animal to be treated, on the potency and duration of action of the compounds employed, or on whether other active ingredients are administered in addition to compounds of the formula I. In general, a daily dose of about 0.01 to 100 mg/kg, preferably 0.1 to 10 mg/kg, in particular 0.3 to 5 mg/kg (in each case mg per kg of body weight) is appropriate on administration to an adult weighing about 75 kg to achieve the desired effect. The daily dose may be administered in a single dose or, especially on administration of larger amounts, be divided into a plurality of, for example two, three or four, single doses. It may, depending on the individual characteristics, be necessary where appropriate to deviate upward or downward from the stated daily dose. Pharmaceutical products normally contain 0.2 to 500 mg, preferably 1 to 200 mg, of an

active ingredient of any of the formulae (Ia) to (Ie) and/or its physiologically tolerated salts.

5 The present invention will be further illustrated by the attached Figures and the following non-limiting Examples.

Fig. 1 is a scheme that illustrates the course of the events in the treatment of increased intracranial pressure and secondary ischemia as disclosed here. First, usually within the first 6 hours after cerebral trauma such as CCT or non-traumatic brain injury the intracranial pressure increases to a critically elevated level, i.e. a level that becomes life-threatening for example because it leads to possible acute respiratory depression and unconsciousness. As the inventors have found out here, the administration of the pteridine derivatives described herein leads to an inhibition of the endothelial NO synthase (e-NOS) in particular in the region of the brain where the trauma occurred and the e-NOS activity is increased. This is in turn accompanied in a contraction of blood vessel in the brain and thus leads to a decrease in the ICP within these first hours after the trauma. As a secondary effect, once the blood-brain-barrier has become permissive as a further result of the trauma, the pteridines used in the present invention are able to cross the blood-brain-barrier and to inhibit the neuronal NO synthase (n-NOS), the inducible NO synthase (i-NOS) and the mitochondrial NO synthase (m-NOS), the activities of which are increased after trauma and in secondary ischemia. Thereby, acute cell death via necrosis, inflammatory processes, cell death via necrosis and apoptosis as well as the formation of edema can be prevented or treated. Moreover, the synthesis of NO in the parenchyma of the brain is blocked. This in turn antagonizes the dilatation of vessels steered by NO coming from the brain parenchyma.

Figure 2 shows the contractile effect of 2,4-diamino-5,6,7,8-tetrahydro-6-(*L-erythro*-1,2-dihydroxypropyl)-pteridine in isolated rat basilar arteries (BA, open circles) and middle cerebral arteries (MCA, closed circles) tested as described in Example 4. The contraction is measured in mN force and expressed in percent contraction relative to the contraction of the arteries caused by 124 mM K⁺-Krebs solution. The concentration of 2,4-diamino-5,6,7,8-tetrahydro-6-(*L-erythro*-1,2-dihydroxypropyl)-pteridine is given on a semilogarithmic scale.

35 Example 1

Preparation of 2,4-diamino-5,6,7,8-tetrahydro-6-(*L-erythro*-1,2-dihydroxypropyl)-pteridine (H₄-aminobiopterin)

2,4-Diamino-5,6,7,8-tetrahydro-6-(*L-erythro*-1,2-dihydroxypropyl)-pteridine was synthesized as follows:

5 Azodicarbonic acid ethylester was added dropwise to a mixture of N²,1',2'-triacetylbipterin (7,9 g), 2-(4-nitrophenyl)ethanol (5,1 g) and triphenylphosphane (8,4 g) in dioxane (150 ml) and was then stirred at room temperature for 20 hours. The solution was rotary evaporated, the remainder was then solved in CHCl₃ and put on a silica gel column (18 x 6 cm). To remove triphenylphosphane oxide, the solution was first
10 chromatographed using EtOAc/n-hexane (1:1) and EtOAc/n-hexane (4:1), and then EtOAc (1,5 l) was used for chromatography to eluate the reaction product. Concentration of the solution yields the pure product: 8,63 g (75%) N²,1',2'-triacetyl-O⁴-2-(4-nitrophenyl)ethylbipterin.

15 N²,1',2'-triacetyl-O⁴-2-(4-nitrophenyl)ethylbipterin (9,0 g) was suspended in a mixture of methanol (90 ml) and concentrated ammonia (90 ml) and the resulting solution was then stirred at room temperature for 36 hours. The solution was concentrated to half the volume, extracted using CH₂Cl₂, and the aqueous phase was then further concentrated. The remainder was treated with warm ethanol. After cooling down, collecting the precipitation
20 and washing with ether, the product was dried over P₄O₁₀ in a vacuum desiccator. Yield: 3,3 g (77%) 4-amino-4-desoxybipterin. Further purification can optionally be accomplished by re-crystallization using a large amount of ethanol.

PtO₂ (0,15 g) were pre-hydrogenated in a solution of trifluoroacetic acid (50 ml) placed in
25 a shaker under hydrogen atmosphere. 4-Amino-4-desoxybipterin (1,0 g) were then added and the reduction process was continued under shaking. After 3 hours the H₂-uptake (200 ml, read off from the burette) was completed. The catalyst was sucked off under nitrogen atmosphere, the solution was concentrated up to dryness and dioxane/HCl-gas (10 ml) and ether (100 ml) were added to the syrup-like remainder and the resulting solution was
30 stirred for several hours, whereby a colourless precipitation deposited. The precipitation was sucked off, washed with ether and dried over solid KOH and P₄O₁₀ in a vacuum desiccator. Yield: 1,23 g (90%) 4-amino-4-desoxy-5,6,7,8-tetrahydrobipterin.

C₉H₁₆N₆O₂ x 2HCl x 2H₂O (348,3); theoretical: C 31.03, H 6.37, N 24.13; determined: C
35 31.23, H 6.18, N 23.95.

Example 2

Determination of the binding specificity of H₄-aminobiopterin by radioactive binding tests

The binding behaviour of H₄-aminobiopterin to various receptors of physiological relevance was determined in competitive binding assays using a radioactively labelled ligand. The determination of IC₅₀ values for NO-synthase was performed in accordance with the protocol published in POLLOCK, J.S., FORSTERMANN, U., MITCHELL, J.A., WARNER, T.D., SCHMIDT, H.H.H., NAKANE, M. and MURAD, F. (1991) Proc. Natl. Acad. Sci. USA, 88 10480-10484 (cf. also test ref. number 766-c of the catalogue of Cerep, Paris, France, edition 2002).

The determination of the IC₅₀ value for each of the other receptors which were chosen for determining the binding specificity of H₄-aminobiopterin was carried out as also described in the catalogue of Cerep, Paris, France, edition 2002.

The results are listed in Table 1. As can be seen from Table 1, H₄-aminobiopterin does not have any measurable binding affinity towards any other investigated physiological important receptor, in particular towards receptors such as receptors of the 5-HT type, the N-methyl-D-aspartate (NDMA) receptor or the GABA receptor, who are involved in the death of neuronal cells after CCT. This clearly suggests that the pteridine compounds used in the present invention display their activity preferentially through binding to NOS.

Table 1

Effects of H₄-aminobiopterin on the specific radioligand binding to receptors studied and IC₅₀ values for reference compounds with known binding affinity to the given receptor

Cerep catalog ref.#/page	Receptor	H ₄ - aminobiopte rin	Reference compound		
		10 μ M		IC ₅₀ (nM)	(nH)
802-1a/35	α_1 (non-selective)	-	Prazosin	0.31	(1.2)
802-1b/36	α_2 (non-selective)	-	yohimbine	74	(0.8)
802-2ah/38	β_1 (h)	-	Atenolol	1,300	(0.9)

Cerep catalog ref.#/page	Receptor	H ₄ - aminobiopte rin	Reference compound		
		10 μ M		IC ₅₀ (nM)	(nH)
802-2bh/38	β_2 (h)	-	ICI 118551	2.3	(1.6)
821-1h/39	AT ₁ (h)	-	saralasin	1.9	(1.0)
821-2h/40	AT ₂ (h)	16	saralasin	0.69	(0.9)
803-1h/45	D1 (h)	-	SCH 23390	0.78	(1.1)
803-2h/45	D2 (h)	-	(+)butaclamol	5.5	(1.1)
825-1h/47	ET _A (h)	-	endothelin-1	0.13	(1.0)
825-2h/47	ET _B (h)	-	endothelin-3	0.12	(1.4)
804/48	GABA (non-selective)	-	GABA	27	(1.3)
804-1/48	GABA _A	-	Muscimol	7.2	(0.8)
804-2/48	GABA _B	-	Baclofen	108	(1.1)
895-1/49	AMPA	-	L-glutamate	850	(0.8)
895-2/50	Kainate	-	kainic acid	26	(1.0)
895-3/50	NMDA	-	CGS 19755	222	(0.9)
882-h/51	EGF (h)	10	EGF	3.1	(1.5)
881/51	PDGF	-	PDGF BB	0.10	(1.9)
805-1c/54	H ₁ (central)	-	pyrilamine	1.8	(1.3)
805-2/55	H ₂	-	cimetidine	781	(0.6)
861-5/56	IP ₃	-	D-(1,4,5)IP ₃	33	(1.0)
806-1h/60	M ₁ (h)	-	pirenzepine	11	(1.0)
806-2h/61	M ₂ (h)	-	methoctramine	24	(0.8)
806-3h/61	M ₃ (h)	10	4-DAMP	1.5	(1.4)

Cerep catalog ref.#/page	Receptor	H ₄ - aminobiopte rin	Reference compound		
		10 μ M		IC ₅₀ (nM)	(nH)
830/68	PAF	-	WEB 2086	8.9	(1.0)
808-1b/71	5-HT _{1B}	-	serotonin	14	(0.9)
808-1d/71	5-HT _{1D}	-	serotonin	2.8	(0.9)
808-2h/71	5-HT _{2A} (h)	-	ketanserin	2.9	(1.1)
808-2bh/72	5-HT _{2B} (h)	-	serotonin	42	(0.8)
813/78	Estrogen (ER)	-	17- β -estradiol	1.8	(1.3)
848-h/79	Vitamin D ₃ (h)	-	1 α ,25-(OH) ₂ D ₃	4.2	(1.8)
861-L1/82	Ca ²⁺ channel (L, DHP site)	-	nitrendipine	1.4	(1.4)
861-L2/82	Ca ²⁺ channel (L, diltiazem site)	-	diltiazem	27	(0.8)
861-L3/83	Ca ²⁺ channel (L, verapamil site)	-	D 600	12	(0.9)
863-1/83	K ⁺ _{ATP} channel	-	glibenclamide	1.3	(1.4)
863-2/84	K ⁺ _v channel	-	α -dendrotoxin	0.50	(1.0)
863-3/84	SK ⁺ _{Ca} channel	-	α -dendrotoxin	0.50	(1.0)

For H₄-aminobiopterin, the results are expressed as a percent inhibition of control specific binding (mean values ; n = 2). The symbol - indicates an inhibition of less than 10%.

Example 3

5 In vitro specificity of H₄-aminobiopterin

The physiological effect of H₄-aminobiopterin on NO synthases was further examined in cell culture tests with cells expressing relevant receptors by use of the protocol described in the catalogue of Cerep, Paris, France, edition 2002 (see Table 2). The determination of the IC₅₀ value for each of the other receptors which were chosen for determining the binding specificity of H₄-aminobiopterin was again also performed as described in the catalogue of

Cerep, Paris, France, edition 2002. Furthermore, the modulatory effect of H₄-aminobiopterin on the β -adrenoceptor-G protein coupling (Table 3) was studied by means of the assay described in ref. 758-2a (p. 134) and 758-2b (p. 134) of the 2002 edition of the Cerep catalogue.

5

The results of these experiments are summarized in Tables 2 and 3. These data also confirm that H₄-aminobiopterin has a preferred binding affinity to NOS.

Table 2

10

Effects of H₄-aminobiopterin in studied cell biology assays
and IC₅₀ values for reference compounds

Cerep catalo ref.# / page	Assays	H ₄ -amino- biopterin 10 μ M	Reference compounds		
				IC ₅₀ (μ M)	(<i>nH</i>)
766-i1/ 112	NOS inducible (whole cells/spectrophoto.)	-	L-NMMA	185	(1.0)
766-c/ 112	NOS constitutive (<i>h</i>) (endothelial)	76	L-NMMA	0.14	(2.7)
752-ch/ 114	Phosphodiesterase III (<i>h</i>)	-	milrinone	0.31	(1.0)
752-e/ 114	Phosphodiesterase V (<i>h</i>)	-	dipyridamole	0.35	(1.0)
735/122	Ca ²⁺ -ATPase (pump)	-	thapsigargin	0.087	(>3)
740-1/ 123	L-type Ca ²⁺ channel (stimulated)	-	nitrendipine	0.0038	(0.8)
739-1/ 123	Na ⁺ channel (<i>h</i>) (stimulated)	12	tetrodotoxin	0.0068	(0.9)
734/124	Na ⁺ /K ⁺ ATPase (pump)	-	ouabain	76	(0.8)
706-2/ 127	Protein kinase A (stimulated)	-	H89	0.10	(0.6)
705-dh/ 128	Protein kinase C- δ (<i>h</i>)		Staurosporine		

Cerep catalo ref.# / page	Assays	H ₄ -amino- biopterin	Reference compounds	
		10 μ M		IC ₅₀ (nH) (μ M)
711/141	NE uptake	-	Protriptyline	0.0019
712/141	DA uptake	-	GBR 12909	0.0018
710/141	5-HT uptake	-	imipramine	0.044
791-4/ 146	Cell proliferation (h) (EGF-stimulated A-431)	23	Staurosporine	0.0025
791-3/ 146	Cell proliferation (PDGF-stimulated Balb/c 3T3)	-	Cycloheximide	0.19
760/155	ATPase (Na ⁺ /K ⁺)	-	ouabain	0.28 (1.3)
761-h/ 156	Acetylcholinesterase (h)	-	neostigmine	0.056 (1.0)

For VAS 203, the results are expressed as a percent inhibition of control activity (mean values ; n = 2). The symbol - indicates an inhibition of less than 10%.

Table 3a

Effects of H₄-aminobiopterin in the studied cell biology assays
and IC₅₀ or EC₅₀ values for the reference compound

5

Cerep catalog ref.# / page	Assay	H ₄ -aminobiopterin	Reference compound	
		10 μ M		EC ₅₀ (μ M)
758-2a/ 134	β -adrenoceptor - G protein coupling (agonist effect)	11	isoproterenol	2.4

For H₄-aminobiopterin, the results are expressed as a stimulation of control activity (mean values ; n = 2).

Table 3b

Effects of H₄-aminobiopterin in the studied cell biology assays

10

and IC₅₀ or EC₅₀ values for the reference compound

Cerep catalog Assay ref.# / page		H ₄ -aminobiopterin 10 μ M	Reference compound IC ₅₀ (nH) (μ M)		
758-2b/ 134	β -adrenoceptor -G protein coupling (antagonist effect)	31	ICI 89406	21	(0.7)
For H ₄ -aminobiopterin, the results are expressed as a percent of inhibition of control activity (mean values ; n = 2).					

Example 4

- 5 Characterization of the contractile action of H₄-aminobiopterin in isolated rat cerebral arteries

The contractile effect of 2,4-diamino-5,6,7,8-tetrahydro-6-(L-erythro-1,2-dihydroxypropyl)-pteridine in isolated rat basilar arteries (BA) and middle cerebral arteries (MCA) was tested as described by Schilling et al., Peptides 21 (2000), 91-99.

For these experiments ring segments of cerebral blood vessels from adult male Sprague-Dawley rats (300-450 g) were prepared as described in Schilling et al., 2000, supra. After sacrificing the animals by bleeding from the carotid arteries the brain was quickly removed and immersed in precooled modified Krebs-Högestätt solution having the composition (mM): NaCl, 119; KCl, 3.0; Na₂HPO₄, 1.2; CaCl₂ x 2 H₂O, 1.5; MgCl₂ x 6H₂O, 1.2; NaHCO₃, 20, glucose, 10. By use of a binocular microscope the respective artery was meticulously dissected and cut into four ring segments that were mounted on two L-shaped stainless steel wires (70 μ m) in an organ bath of 5 ml volume. One wire was firmly attached to the wall of the bath whereas the other one was connected to a force transducer mounted on a microdrive (M33, Märzhäuser, Wetzlar, Germany). Changes in isometric forces were measured by a transducer coupled to an amplifier (Q11 amplifier and QT233 amplifier, each by Hottinger Baldwin, Darmstadt, Germany) and displayed on a pen chart recorder.

25

After mounting, the segments received a 90 min accommodation period with gradually increasing the bath temperature to 37°C. Bath solution was constantly bubbled with a humidified gas mixture of 93 % O₂/7 % CO₂ resulting in a pH around 7.35 at 37°C.

During accommodation the segments were repeatedly stretched and resting tension was adjusted to 2.5 to 3.5 milli-Newton force (mN). Thereafter, a reference contraction was obtained by exchanging the bath solution with 124 mM K⁺-Krebs solution (NaCl replaced by KCl). Segments that developed less than 2.5 mN were discarded. The functional state of the endothelium was tested with cumulative application of acetylcholine (10⁻⁸ to 10⁻⁴ M) after precontraction with serotonin (5-HT, 10⁻⁶ M) followed by exchanging the bath solution back to modified Krebs-Högestätt solution as described above. As described by Schilling et al, 2000, supra only segments in which relaxation exceeded 25 % of precontraction were considered to carry a functional endothelium and included in the analysis. In each segment H₄-aminobiopterin was then applied cumulatively (10⁻¹⁰ to 10⁻³ M in a first experiment in order to determine the effective concentration, later in a narrower concentration range) into the bath solution, wherein the segments were incubated under resting tension to characterize contraction. The contraction was measured in mN force and expressed in percent reference contraction (contraction caused by 124 mM K⁺-Krebs solution, see above).

The results are shown in Fig.2. As can be seen from Fig.2, H₄-aminobiopterin caused a clear concentration dependent contraction of both rat basilar arteries and middle cerebral arteries. Accordingly, these results clearly indicate the vasoconstrictorial effect of H₄-aminobiopterin on the brain vessels and thus its suitability to lower the increased intracranial pressure by recontracting dilated blood vessels in the brain.

03-05-2005

EP0311138


1

PCT Application PCT/EP03/11138
Vasopharm Biotech GmbH

Claims

5

1. Use of a compound of the general formula (Ia)



10

for the manufacture of a pharmaceutical composition for treating increased intracranial pressure, wherein in formula (Ia) R_1 , R_3 are independently selected from H or OH, R_4 is H, CH_3 , CH_2OH , CHO, R_2 is H, CH_3 , CH_2OH , CHO or a lower C_1 - C_9 alkyl radical, which can be a straight or a branched chain, as well as $(CH(OH))_n-Y$ or $(CH(OH))_n-(CH_2)_m-W$,
 15 wherein Y is hydrogen or a lower alkyl radical, W is hydrogen or a hydroxyl group, an n and m are independently from each other 1-20.

15

2. The use of claim 1, wherein R_1 and R_4 in formula (Ia) are hydrogen, R_2 is methyl and R_3 is hydroxyl.

20

3. The use of claim 2, wherein the compound is 2,4-diamino-5,6,7,8-tetrahydro-6-(L-erythro-1,2-dihydroxypropyl)-pteridine.


4. The use of a compound of the general formula (Ib)

25

03-05-2005

EP0311138

2




for the manufacture of a pharmaceutical composition for treating increased intracranial pressure, wherein in formula (Ib) R_1 , R_2 , R_3 and R_4 are independently from each other H or OH, R_5 is H, CH_3 , CH_2OH , CHO or a lower C_1 - C_9 alkyl radical, which can be a straight or a branched chain, as well as $(CH(OH))_n-Y$ or $(CH(OH))_n-(CH_2)_m-W$, wherein Y is hydrogen or a lower alkyl radical, W is hydrogen or a hydroxyl group, an n and m are independently from each other 1-20.

5 10 5. The use of claim 4, wherein R_5 is methyl, R_3 is hydrogen, R_2 and R_4 are both hydrogen and R_1 is either hydrogen or hydroxyl.

6. The use according to any of claims 1 to 5, wherein the increased intracranial pressure is caused by closed craniocerebral trauma or by non-traumatic brain injury.

15 7. Use of a compound of the general formula (Ia)



20 for the manufacture of a pharmaceutical composition for treating secondary brain damage, wherein the secondary brain damage is an inflammatory process, wherein in formula (Ia) R_1 , R_3 are independently selected from H or OH, R_4 is H, CH_3 , CH_2OH , CHO, R_2 is H, CH_3 , CH_2OH , CHO or a lower C_1 - C_9 alkyl radical, which can be a straight or a branched chain, as well as $(CH(OH))_n-Y$ or $(CH(OH))_n-(CH_2)_m-W$, wherein Y is

03-05-2005

EP0311138

3


hydrogen or a lower alkyl radical, W is hydrogen or a hydroxyl group, an n and m are independently from each other 1-20.

8. The use of claim 7, wherein R_1 and R_4 in formula (Ia) are hydrogen, R_2 is methyl and R_3 is hydroxyl.

9. The use of claim 8 wherein the compound is 2,4-diamino-5,6,7,8-tetrahydro-6-(L-erythro-1,2-dihydroxypropyl)-pteridine.

10

10. The use of a compound of the general formula (Ib)



15

for the manufacture of a pharmaceutical composition for treating secondary brain damage, wherein the secondary brain damage is an inflammatory process, wherein in formula (Ib) R_1 , R_2 , R_3 and R_4 are independently from each other H or OH, R_5 is H, CH_3 , CH_2OH , CHO or a lower C_1 - C_9 alkyl radical, which can be a straight or a branched chain, as well as $(CH(OH))_n$ -Y or $(CH(OH))_n$ -(CH_2) $_m$ -W, wherein Y is hydrogen or a lower alkyl radical, W is hydrogen or a hydroxyl group, an n and m are independently from each other 1-20.

11. The use of claim 10, wherein R_5 is methyl, R_3 is hydrogen, R_2 and R_4 are both hydrogen and R_1 is either hydrogen or hydroxyl.

12. The use according to any of claims 7 to 11, wherein the inflammatory process is caused by closed craniocerebral trauma or by non-traumatic brain injury.

13. The use of any of claims 1 to 12 for the treatment of a mammal, especially a human.




Fig. 1

2/2




Fig. 2

