

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number
WO 00/76532 A1

(51) International Patent Classification⁶: A61K 38/00,
C07K 14/435, 14/00, C12N 15/12

Lake City, UT 84101 (US). **JONES, Robert, M.**; 44 West Broadway #2103 South, Salt Lake City, UT 84101 (US).

(21) International Application Number: PCT/US00/15779

(74) Agents: **IHNEN, Jeffrey, L.** et al.; Rothwell, Figg, Ernst & Manbeck, Suite 701 East, 555 13th Street N.W., Columbia Square, Washington, DC 20004 (US).

(22) International Filing Date: 9 June 2000 (09.06.2000)

(25) Filing Language: English

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
60/138,507 10 June 1999 (10.06.1999) US

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


(71) Applicants: **UNIVERSITY OF UTAH RESEARCH FOUNDATION** [US/US]; Suite 110, 615 Arapeen Drive, Salt Lake City, UT 84108 (US). **COGNETIX, INC.** [US/US]; Suite 201, 421 Wakara Way, Salt Lake City, UT 84108 (US).

(72) Inventors: **OLIVERA, Baldomero, M.**; 1370 Bryan Avenue, Salt Lake City, UT 84108 (US). **MCINTOSH, J., Michael**; 1151 South 2000 East, Salt Lake City, UT 84108 (US). **MCCABE, R., Tyler**; 715 Freeze Creek Circle, Salt Lake City, UT 84108 (US). **GARRETT, James, E.**; 1584 East 3150 South, Salt Lake City, UT 84106 (US). **LAYER, Richard, T.**; 9024 Sunburst Court, Sandy, UT 84093 (US). **WAGSTAFF, John, D.**; 44 West Broadway, #2103S, Salt

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 00/76532 A1

(54) Title: **μO-CONOPEPTIDES AND THEIR USE AS LOCAL ANESTHETICS**

(57) Abstract: The present invention is directed to the new μ O-conopeptides, their coding sequences and their propeptides and to the use of μ O-conopeptides as a local anesthetic for treating pain. The μ O-conopeptides have long lasting anesthetic activity and are particularly useful for spinal anesthesia, either administered acutely for post-operatively pain or via an intrathecal pump for severe chronic pain situations or for treatment of pain in epithelial tissue.

5 TITLE OF THE INVENTION

MuO-CONOPEPTIDES AND THEIR USE AS LOCAL ANESTHETICS

This invention was made with Government support under Grant No. GM48677 awarded by the National Institute of General Medical Sciences, National Institutes of Health, Bethesda, Maryland. The United States Government has certain rights in the invention.

BACKGROUND OF THE INVENTION

The present invention is directed to the use of μ O-conopeptides as a local anesthetic for treating pain. The μ O-conopeptides have long-lasting anesthetic activity and are particularly useful for spinal anesthesia, administered either acutely for post-operative pain or via an intrathecal pump for severe chronic pain situations. The present invention is further directed to new μ O-conopeptides, their coding sequences and their propeptides.

The publications and other materials used herein to illuminate the background of the invention, and in particular, cases to provide additional details respecting the practice, are incorporated herein by reference, and for convenience, are referenced by author and date in the following text and respectively grouped in the appended List of References.

Conus is a genus of predatory marine gastropods (snails) which envenomate their prey. Venomous cone snails use a highly developed projectile apparatus to deliver their cocktail of toxic conotoxins into their prey. In fish-eating species such as *Conus magus* the cone detects the presence of the fish using chemosensors in its siphon and when close enough extends its proboscis and fires a hollow harpoon-like tooth containing venom into the fish. This immobilizes the fish and enables the cone snail to wind it into its mouth via an attached filament. For general information on *Conus* and their venom see the website address <http://grimwade.biochem.unimelb.edu.au/cone/referenc.html>. Prey capture is accomplished through a sophisticated arsenal of peptides which target specific ion channel and receptor subtypes. Each *Conus* species venom appears to contain a unique set of 50-200 peptides. The composition of the venom differs greatly between species and between individual snails within each species, each optimally evolved to paralyse it's prey. The active

components of the venom are small peptides toxins, typically 12-30 amino acid residues in length and are typically highly constrained peptides due to their high density of disulphide bonds.

The venoms consist of a large number of different peptide components that when separated exhibit a range of biological activities: when injected into mice they elicit a range of physiological responses from shaking to depression. The paralytic components of the venom that have been the focus of recent investigation are the α -, ω - and μ -conotoxins. All of these conotoxins act by preventing neuronal communication, but each targets a different aspect of the process to achieve this. The α -conotoxins target nicotinic ligand gated channels, the μ -conotoxins target the voltage-gated sodium channels and the ω -conotoxins target the voltage-gated calcium channels (Olivera et al., 1985). For example a linkage has been established between α -, α A- & ϕ -conotoxins and the nicotinic ligand-gated ion channel; ω -conotoxins and the voltage-gated calcium channel; μ -conotoxins and the voltage-gated sodium channel; δ -conotoxins and the voltage-gated sodium channel; κ -conotoxins and the voltage-gated potassium channel; conantokins and the ligand-gated glutamate (NMDA) channel. For a partial list of *Conus* peptides and their amino acid sequences see the website address <http://pir.georgetown.edu>.

However, the structure and function of only a small minority of these peptides have been determined to date. For peptides where function has been determined, three classes of targets have been elucidated: voltage-gated ion channels; ligand-gated ion channels, and G-protein-linked receptors.

Conus peptides which target voltage-gated ion channels include those that delay the inactivation of sodium channels, as well as blockers specific for sodium channels, calcium channels and potassium channels. Peptides that target ligand-gated ion channels include antagonists of NMDA and serotonin receptors, as well as competitive and noncompetitive nicotinic receptor antagonists. Peptides which act on G-protein receptors include neuropeptides and vasopressin receptor agonists. The unprecedented pharmaceutical selectivity of conotoxins is at least in part defined by a specific disulfide bond frameworks combined with hypervariable amino acids within disulfide loops (for a review see McIntosh et al., 1998).

The pain response is a protective reflex system warning an individual of hostile situations and tissue injury. The origins of clinically significant acute and chronic pain in a mammal are different, but the biochemical and neurological pathways are similar. In the following discussion on pain and its management, the focus is primarily on humans, however, it should be understood

that the concepts of pain are applicable to mammalian animals and the management of such pain is applicable to veterinary medicine.

Acute pain is often associated with surgery and with trauma. The intensity of acute postoperative pain varies considerably depending on the extent of the surgical procedure performed, 5 on the individual's pain sensitivity, and on the type of anesthetic management employed during surgery. In general, major operations on the thorax and the upper abdominal region induce the most intensive postoperative pain. Extensive orthopedic operations also produce strong postoperative pain.

Chronic pain is persistent pain which has long outlasted the onset of any known or suspected 10 physical cause. It can occur after a known injury or disease, or it can occur without any known physical cause whatsoever. Moreover, it can be accompanied by known tissue pathology, such as chronic inflammation that occurs in some types of arthritis, or it can occur long after the healing of the injured tissue which is suspected or known to be the cause of chronic pain. Chronic pain is a 15 very general concept and there are several varieties of chronic pain related to the musculoskeletal system, visceral organs, skin, and nervous system.

Neuropathic pain can occur as a form of chronic pain and can also occur under acute conditions such as those following surgery or accidental trauma. Neuropathic pain can be defined 20 as pain that results from an abnormal functioning of the peripheral and/or central nervous system. A critical component of this abnormal functioning is an exaggerated response of pain-related nerve cells either in the peripheral or in the central nervous system. This exaggerated responsiveness is manifested behaviorally as increased sensitivity to pain, i.e., as hyperalgesia or allodynia, both of 25 which can occur in chronic neuropathic and acute inflammatory pains. An example is the pain from causalgia wherein even a light touch to the skin is felt as an excruciating burning pain (allodynia) or a normally mild pain is experienced as an excruciating one (hyperalgesia). Neuropathic pain is thought to be a consequence of damage to peripheral nerves or to regions of the central nervous 30 system. However, abnormal functioning of pain-related regions of the nervous system can also occur with chronic inflammatory conditions such as certain types of arthritis and metabolic disorders such as diabetes as well as with acute inflammatory conditions. Thus, many types of chronic pains that are related to inflammation as well as acute pains that are related to inflammation can be considered to be at least partly neuropathic pains.

The modern concept of pain treatment emphasizes the significance of prophylactic prevention of pain, as pain is more easily prevented than relieved. Additionally, the hormonal stress

responses associated with pain are considered harmful to the patient, impair the healing process and overall recovery, and generally are to be avoided.

While compounds utilized as general anesthetics reduce pain by producing a loss of consciousness, local anesthetics act to induce a loss of sensation in the localized area of administration in the body. The mechanism by which local anesthetics induce their effect, while not having been determined definitively, is generally thought to be based upon the ability to interfere with the initiation and transmission of the nerve impulse conduction along an axon through a reversible blockade of sodium channels. Currently used local anesthetics have durations of action lasting only several hours. While this length of duration meets many needs, particularly the control of acute pain, local anesthetic agents with longer duration of action would have broad clinical application for the treatment of postoperative and chronic pain (Kuzma et al., 1997).

The duration of action of a local anesthetics is proportional to the time during which it is in actual contact with the nervous tissues. In an effort to increase the duration of action, procedures or formulations that maintain localization of the drug at the nerve greatly prolong anesthesia. All local anesthetics are potentially toxic, and therefore it is of great importance that the choice of drug, concentration, rate and site of administration, as well as other actors, be considered in their use. On the other hand, a local anesthetic must remain at the site long enough to allow sufficient time for the localized pain to subside. Different devices and formulations are known in the art for administration of local anesthetics. See U.S. Patent No. 5,747,060, which discloses such devices and formulations.

Side effects which have been associated with the use of different drugs for treating pain or as local anesthetics includes include respiratory depression, reduced cough reflex, bronchial spasms, nausea, vomiting, release of histamine, peripheral vasodilation, orthostatic hypotension, vagal impact on the heart, contraction of smooth muscles (sphincters), reduced peristaltic motility in the gastrointestinal tract, urinary retention, stimulated release of adrenalin, anti-diuretic hormone, changes in the regulation of body temperature and sleep pattern, tolerance, addiction, tachycardia, increase in blood pressure, and agitation. Not all of these side effects are seen with any given drug used to treat pain.

Thus, there is a need to develop additional drugs and methods which can be used for the treatment of pain, which can act as local anesthetics, which have a longer duration of action and which have reduced side effects. Accordingly, an object of the invention is to provide methods and compositions for the treatment of acute or chronic pain which provide effective control of pain with longer duration of action and reduced side effects associated with traditional analgesics.

SUMMARY OF THE INVENTION

The present invention is directed to the new μ O-conopeptides, their coding sequences and their propeptides and to the use of μ O-conopeptides as a local anesthetic for treating pain. The μ O-conopeptides have long lasting anesthetic activity and are particularly useful for spinal anesthesia, either administered acutely for post-operative pain or via an intrathecal pump for severe chronic pain situations or for treatment of pain in epithelial tissue.

More specifically, the present invention is directed to μ O-conopeptides having the general formula I:

Xaa₁-Xaa₂-Cys-Xaa₃-Xaa₄-Xaa₅-Xaa₆-Xaa₇-Xaa₈-Cys-Xaa₉-Xaa₁₀-Xaa₁₁-Xaa₁₂-Xaa₁₃-Xaa₁₄-
10 Xaa₁₅-Xaa₁₆-Xaa₁₇-Cys-Cys-Xaa₁₈-Xaa₁₉-Xaa₂₀-Xaa₂₁-Cys-Xaa₂₂-Xaa₂₃-Xaa₂₄-Xaa₂₅-Cys-Xaa₂₆-
Xaa₂₇-Xaa₂₈-Xaa₂₉-Xaa₃₀ (SEQ ID NO:1),

wherein Xaa₁ is des-Xaa₁, Pro, hydroxy-Pro (Hyp), Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₂ is des-Xaa₂, Ala, Gly, Asp, Glu, γ -carboxy-glutamate (Gla), any synthetic acidic amino acid, Thr, Ser, g-Thr (where g is glycosylation), g-Ser, Trp (D or L), neo-Trp or halo-Trp (D or L) or Xaa₂ may be pyroglutamate if Xaa₁ is des-Xaa₁; Xaa₃ is Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ser, Thr, g-Ser, g-Thr, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, His, Glu, Gln, Gla, Asp, Asn or any synthetic acidic amino acid; Xaa₄ is Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Ser, Thr, Pro, Hyp, g-Ser, g-Thr, g-Hyp or any synthetic hydroxylated amino acid; Xaa₅ is Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Pro or Hyp; Xaa₆ is Trp (D or L), neo-Trp, halo-Trp (D or L), Gly, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr,

di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid; Xaa₇ is Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Met, norleucine (Nle), Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₈ is Leu, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₉ is Pro, Hyp, Gly, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₁₀ is Thr, Ser, g-Thr, g-Ser, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₁₁ is Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Thr or any hydroxylated amino acid; Xaa₁₂ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₃ is Pro, Hyp, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₄ is Gly, His, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₅ is des-Xaa₁₅, Ser, Thr, g-Ser, g-Thr, Val, Asn, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₁₆ is Met, Nle, Leu, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₇ is Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Thr, any

hydroxylated amino acid, Ala, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, His or Gly; Xaa₁₈ is Gly, Asn or Gln; Xaa₁₉ is Leu, Trp (D or L), neo-Trp or halo-Trp (D or L); Xaa₂₀ is des-Xaa₂₀, Leu or Trp (D or L), neo-Trp or halo-Trp (D or L); Xaa₂₁ is des-Xaa₂₁ or an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₂₂ is des-Xaa₂₂, Gly, Met, Nle, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₃ is des-Xaa₂₃, Pro, Hyp, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, 10 Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₄ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), 15 neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₅ is Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa₂₆ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₂₇ is des-Xaa₂₇, Asp, Glu, Gla, Pro, Hyp, Ser, Thr, 20 g-Hyp, g-Ser, g-Ser or any synthetic hydroxylated amino acid; Xaa₂₈ is des-Xaa₂₈, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ile, Ser, Thr, g-Ser or g-Thr; Xaa₂₉ is des-Xaa₂₉, Pro, Hyp, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa₃₀ is des-Xaa₃₀ or Phe, with the proviso that the peptide is not MrVIA/B as defined below. The Cys residues may be in D or L configuration and may optionally be substituted with homocysteine (D or L). The Tyr residues may be substituted with the 3-hydroxyl or 2-hydroxyl isomers and corresponding O-sulpho- and O-phospho-derivatives. The acidic amino acid residues may be substituted with any synthetic acidic amino acid, e.g., tetrazolyl derivatives of Gly and Ala. The nonnatural derivatives of the aliphatic amino acids include those synthetic derivatives bearing non-natural aliphatic branched or linear side

chains C_nH_{2n+2} up to and including n=8. The halogen is iodo, chloro, fluoro or bromo; preferably iodo for halogen substituted-Tyr and bromo for halogen-substituted Trp.

MrVIA/B has the sequence: Ala-Cys-Xaa₃₁-Lys-Lys-Trp-Glu-Tyr-Cys-Ile-Val-Xaa₃₂-Ile-Xaa₃₃-Gly-Phe-Xaa₃₄-Tyr-Cys-Cys-Xaa₃₂-Gly-Leu-Ile-Cys-Gly-Xaa₃₂-Phe-Val-Cys-Val, wherein Xaa₃₁ is Arg or Ser, Xaa₃₂ is Pro or hydroxy-Pro, Xaa₃₃ is Ile or Leu and Xaa₃₄ is Ile or Val (SEQ ID NO:2).

The present invention is also directed to novel specific conotoxin peptides within general formula I having the formulas:

Ala-Cys-Arg-Gln-Xaa₁-Xaa₂-Xaa₃-Phe-Cys-Leu-Val-Xaa₄-Ile-Ile-Gly-Xaa₂-Ile-Xaa₂-Cys-Cys-Ala-Gly-Leu-Ile-Cys-Gly-Xaa₄-Phe-Val-Cys-Leu (SEQ ID NO:3);

Xaa₄-Thr-Cys-Leu-Xaa₁-Gln-Asp-Xaa₁-Phe-Cys-Ile-Ile-Xaa₄-Leu-Ile-Gly-Thr-Leu-Xaa₂-Cys-Cys-Ser-Gly-Leu-Ile-Cys-Gly-Phe-Phe-Val-Cys-Val-Xaa₄-Xaa₁-Xaa₄-Phe (SEQ ID NO:4);

Asp-Cys-Xaa₃-Ala-Asp-Gly-Ala-Phe-Cys-Gly-Ile-Val-Xaa₁-Asn-Xaa₅-Met-Cys-Cys-Ser-Asn-Leu-Cys-Ile-Phe-Ala-Cys-Val-Xaa₄-Xaa₃-Xaa₂ (SEQ ID NO:5);

Asp-Cys-His-Xaa₃-Arg-Xaa₅-Asp-Xaa₅-Cys-Xaa₄-Ala-Ser-Ile-Leu-Gly-Val-Ile-Xaa₂-Cys-Cys-Xaa₃-Gly-Leu-Ile-Cys-Phe-Ile-Ala-Phe-Cys-Ile (SEQ ID NO:6);

Asp-Cys-Gln-Xaa₃-Xaa₁-Xaa₅-Xaa₃-Phe-Cys-Ile-Val-Xaa₄-Ile-Leu-Gly-Phe-Val-Xaa₂-Cys-Cys-Xaa₄-Gly-Leu-Ile-Cys-Gly-Xaa₄-Phe-Val-Cys-Val-Asp-Ile (SEQ ID NO:7);

Xaa₄-Thr-Cys-Val-Ser-Xaa₂-Asn-Val-Phe-Cys-Gly-Val-Xaa₄-Leu-Val-Gly-Thr-Xaa₂-Leu-Cys-Cys-Ser-Gly-Leu-Val-Cys-Leu-Val-Val-Cys-Ile (SEQ ID NO:8);

Cys-Arg-Xaa₄-Arg-Gly-Met-Phe-Cys-Gly-Phe-Xaa₄-Xaa₁-Xaa₄-Gly-Xaa₄-Xaa₂-Cys-Cys-Asn-Gly-Xaa₅-Cys-Phe-Phe-Val-Cys-Ile (SEQ ID NO:9);

Arg-Xaa₅-Cys-Ala-Leu-Asp-Gly-Xaa₃-Leu-Cys-Ile-Ile-Xaa₄-Val-Ile-Gly-Ser-Ile-Phe-Cys-Cys-His-Gly-Ile-Cys-Met-Ile-Xaa₂-Cys-Val (SEQ ID NO:10);

Asp-Cys-Arg-Xaa₄-Val-Gly-Gln-Xaa₂-Cys-Gly-Ile-Xaa₄-Xaa₂-Xaa₁-His-Asn-Xaa₅-Arg-Cys-Cys-Ser-Gln-Leu-Cys-Ala-Ile-Ile-Cys-Val-Ser (SEQ ID NO:11); and

Gly-Cys-Leu-Asp-Xaa₄-Gly-Xaa₂-Phe-Cys-Gly-Thr-Xaa₄-Phe-Leu-Gly-Ala-Xaa₂-Cys-Cys-Gly-Gly-Ile-Cys-Leu-Ile-Val-Cys-Ile-Xaa₃-Thr (SEQ ID NO:12),

wherein Xaa₁ is Lys, N-methy-Lys, N,N-dimethyl-Lys or N,N,N-trimethyl-Lys; Xaa₂ is Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa₃ is Glu or gamma-carboxy-Glu (Gla); Xaa₄ is Pro or hydroxy-Pro; Xaa₅ is Trp or halo-Trp; and the C-terminus contains a carboxyl or amide group. The halo is preferably chlorine or iodine, more preferably iodine. In

addition, the Arg residues may be substituted by Lys, ornithine, homoarginine, nor-Lys, N-methyl-Lys, N,N-dimethyl-Lys, N,N,N-trimethyl-Lys or any synthetic basic amino acid; the Xaa₁ residues may be substituted by Arg, ornithine, homoarginine, nor-Lys, or any synthetic basic amino acid; the Tyr residues may be substituted with any synthetic hydroxy containing amino acid; the Ser residues 5 may be substituted with Thr or any synthetic hydroxylated amino acid; the Thr residues may be substituted with Ser or any synthetic hydroxylated amino acid; the Phe and Trp residues may be substituted with any synthetic aromatic amino acid; and the Asn, Ser, Thr or Hyp residues may be glycosylated. The Cys residues may be in D or L configuration and may optionally be substituted with homocysteine (D or L). The Tyr residues may also be substituted with the 3-hydroxyl or 2-hydroxyl isomers (meta-Tyr or ortho-Tyr, respectively) and corresponding O-sulpho- and O-phospho-derivatives. The acidic amino acid residues may be substituted with any synthetic acidic 10 amino acid, e.g., tetrazolyl derivatives of Gly and Ala. The aliphatic amino acids may be substituted by synthetic derivatives bearing non-natural aliphatic branched or linear side chains C_nH_{2n+2} up to and including n=8.

15 More specifically, the present invention is directed to the following μ O-conopeptides within general formula I:

MrVIA:	SEQ ID NO:2, whererin Xaa ₃₀ is Arg, Xaa ₃₁ is Ile and Xaa ₃₂ is Ile;
MrVIB:	SEQ ID NO:2, wherein Xaa ₃₀ is Ser, Xaa ₃₁ is Leu and Xaa ₃₂ is Val;
A657:	SEQ ID NO:3, wherein Xaa ₁ is Lys, Xaa ₂ is Tyr, Xaa ₃ is Glu and Xaa ₄ is Pro;
20 F079:	SEQ ID NO:4, wherein Xaa ₁ is Lys, Xaa ₂ is Tyr and Xaa ₄ is Pro;
Ca6.1:	SEQ ID NO:5, wherein Xaa ₁ is Lys, Xaa ₂ is Tyr, Xaa ₃ is Glu, Xaa ₄ is Pro and Xaa ₅ is Trp;
Tx6.12:	SEQ ID NO:6, wherein Xaa ₂ is Tyr, Xaa ₃ is Glu, Xaa ₄ is Pro and Xaa ₅ is Trp;
Tx6.13:	SEQ ID NO:7, wherein Xaa ₁ is Lys, Xaa ₂ is Tyr, Xaa ₃ is Glu, Xaa ₄ is Pro and Xaa ₅ is Trp;
25 G28:	SEQ ID NO:8, wherein Xaa ₂ is Tyr and Xaa ₄ is Pro;
F763:	SEQ ID NO:9, wherein Xaa ₁ is Lys, Xaa ₂ is Tyr, Xaa ₄ is Pro and Xaa ₅ is Trp;
F080:	SEQ ID NO:10, wherein Xaa ₂ is Tyr, Xaa ₃ is Glu, Xaa ₄ is Pro and Xaa ₅ is Trp;
30 F008:	SEQ ID NO:11, wherein Xaa ₁ is Lys, Xaa ₂ is Tyr, Xaa ₄ is Pro and Xaa ₅ is Trp; and
G18:	SEQ ID NO:12, wherein Xaa ₂ is Tyr, Xaa ₃ is Glu and Xaa ₄ is Pro.

Examples of synthetic aromatic amino acid include, but are not limited to, such as nitro-Phe, 4-substituted-Phe wherein the substituent is C₁-C₃ alkyl, carboxyl, hydroxymethyl, sulphomethyl, halo, phenyl, -CHO, -CN, -SO₃H and -NHAc. Examples of synthetic hydroxy containing amino acid, include, but are not limited to, such as 4-hydroxymethyl-Phe, 4-hydroxyphenyl-Gly, 2,6-dimethyl-Tyr and 5-amino-Tyr. Examples of synthetic basic amino acids include, but are not limited to, N-1-(2-pyrazolinyl)-Arg, 2-(4-piperinyl)-Gly, 2-(4-piperinyl)-Ala, 2-[3-(2S)pyrrolininyl]-Gly and 2-[3-(2S)pyrrolininyl]-Ala. These and other synthetic basic amino acids, synthetic hydroxy containing amino acids or synthetic aromatic amino acids are described in Building Block Index, Version 3.0 (1999 Catalog, pages 4-47 for hydroxy containing amino acids and aromatic amino acids and pages 66-87 for basic amino acids; see also <http://www.amino-acids.com>), incorporated herein by reference, by and available from RSP Amino Acid Analogues, Inc., Worcester, MA. Examples of synthetic acid amino acids include those derivatives bearing acidic functionality, including carboxyl, phosphate, sulfonate and synthetic tetrazolyl derivatives such as described by Ornstein et al. (1993) and in U.S. Patent No. 5,331,001, each incorporated herein by reference.

Optionally, in the peptides of general formula I and the specific peptides described above, the Asn residues may be modified to contain an N-glycan and the Ser, Thr and Hyp residues may be modified to contain an O-glycan (e.g., g-N, g-S, g-T and g-Hyp). In accordance with the present invention, a glycan shall mean any N-, S- or O-linked mono-, di-, tri-, poly- or oligosaccharide that can be attached to any hydroxy, amino or thiol group of natural or modified amino acids by synthetic or enzymatic methodologies known in the art. The monosaccharides making up the glycan can include D-allose, D-altrose, D-glucose, D-mannose, D-gulose, D-idose, D-galactose, D-talose, D-galactosamine, D-glucosamine, D-N-acetyl-glucosamine (GlcNAc), D-N-acetyl-galactosamine (GalNAc), D-fucose or D-arabinose. These saccharides may be structurally modified, e.g., with one or more O-sulfate, O-phosphate, O-acetyl or acidic groups, such as sialic acid, including combinations thereof. The glycan may also include similar polyhydroxy groups, such as D-penicillamine 2,5 and halogenated derivatives thereof or polypropylene glycol derivatives. The glycosidic linkage is beta and 1-4 or 1-3, preferably 1-3. The linkage between the glycan and the amino acid may be alpha or beta, preferably alpha and is 1-.

Core O-glycans have been described by Van de Steen et al. (1998), incorporated herein by reference. Mucin type O-linked oligosaccharides are attached to Ser or Thr (or other hydroxylated residues of the present peptides) by a GalNAc residue. The monosaccharide building blocks and the linkage attached to this first GalNAc residue define the "core glycans," of which eight have been

identified. The type of glycosidic linkage (orientation and connectivities) are defined for each core glycan. Suitable glycans and glycan analogs are described further in U.S. Serial No. 09/420,797 filed 19 October 1999 and in PCT Application No. PCT/US99/24380 filed 19 October 1999 (PCT Published Application No. WO 00/23092), each incorporated herein by reference. A preferred 5 glycan is Gal(β1→3)GalNAc(α1→).

Optionally, in the peptides of general formula I and the specific peptides described above, pairs of Cys residues may be replaced pairwise with isoteric lactam or ester-thioether replacements, such as Ser/(Glu or Asp), Lys/(Glu or Asp) or Cys/Ala combinations. Sequential coupling by known methods (Barnay et al., 2000; Hruby et al., 1994; Bitan et al., 1997) allows replacement of 10 native Cys bridges with lactam bridges. Thioether analogs may be readily synthesized using halo-Ala residues commercially available from RSP Amino Acid Analogues.

The present invention is also directed to the identification of the nucleic acid sequences encoding these peptides and their propeptides and the identification of nucleic acid sequences of additional related μO-conopeptides.

15 The present invention is further directed to a method of reducing/alleviating/decreasing the perception of pain by a subject or for inducing analgesia, particularly local analgesia, in a subject comprising administering to the subject an effective amount of the pharmaceutical composition comprising a therapeutically effective amount of a μO-conotoxin peptide described herein or a pharmaceutically acceptable salt or solvate thereof, including MrVIA and MrVIB. The present 20 invention is also directed to a pharmaceutical composition comprising a therapeutically effective amount of a μO-conotoxin peptide described herein or a pharmaceutically acceptable salt or solvate thereof and a pharmaceutically acceptable carrier.

BRIEF DESCRIPTION OF THE FIGURES

25 Figure 1 shows μO-conopeptide MrVIB inhibits skin flinch sensitivity in the Guinea pig intracutaneous wheal assay with greater potency than lidocaine or bupivacaine. Data represent the number of flinches observed after 36 pin pricks in a 30 minutes test period. Each point represents the mean of at least three observations.

30 Figure 2 shows μO-conopeptide MrVIB produces a long-lasting inhibition of skin flinch sensitivity relative to either lidocaine or bupivacaine in the Guinea pig intracutaneous wheal assay. Data represent the percentage of flinches observed out of six total at each time point. Each point represents the mean of at least three observations.

SUMMARY OF THE SEQUENCE LISTING

SEQ ID NO:1 is a generic formula for μ O-conopeptides. SEQ ID NO:2 is a generic formula for μ O-conopeptides MrVIA and MrVIB. SEQ ID NO:3 is a generic formula for μ O-conopeptide A657. SEQ ID NO:4 is a generic formula for μ O-conopeptide F079. SEQ ID NO:5 is a generic formula for μ O-conopeptide Ca6.1. SEQ ID NO:6 is a generic formula for μ O-conopeptide Tx6.12. SEQ ID NO:7 is a generic formula for μ O-conopeptide Tx6.13. SEQ ID NO:8 is a generic formula for the μ O-conopeptide G28. SEQ ID NO:9 is a generic formula for the μ O-conopeptide F763. SEQ ID NO:10 is a generic formula for the μ O-conopeptide F080. SEQ ID NO:11 is a generic formula for the μ O-conopeptide F008. SEQ ID NO:12 is a generic formula for the μ O-conopeptide G18. SEQ ID NO:13 is a primer for amplifying “O-Superfamily” conotoxins. SEQ ID NO:14 is a primer for amplifying “O-Superfamily” conotoxins. SEQ ID NO:15 is a nucleotide sequence for the gene coding for the A657 propeptide. SEQ ID NO:16 is an amino acid sequence of the A657 propeptide. SEQ ID NO:17 is a nucleotide sequence for the gene coding for the F079 propeptide. SEQ ID NO:18 is an amino acid sequence of the F079 propeptide. SEQ ID NO:19 is a nucleotide sequence for the gene coding for the Ca6.1 propeptide. SEQ ID NO:20 is an amino acid sequence of the Ca6.1 propeptide. SEQ ID NO:21 is a nucleotide sequence for a portion of the gene coding for the Tx6.12 propeptide. SEQ ID NO:22 is an amino acid sequence of a portion of the Tx6.12 propeptide. SEQ ID NO:23 is a nucleotide sequence for a portion of the gene coding for the Tx6.13 propeptide. SEQ ID NO:24 is an amino acid sequence of a portion of the Tx6.13 propeptide. SEQ ID NO:25 is a nucleotide sequence for the gene coding for the G28 propeptide. SEQ ID NO:26 is an amino acid sequence of the G28 propeptide. SEQ ID NO:27 is a nucleotide sequence for the gene coding for the F763 propeptide. SEQ ID NO:28 is an amino acid sequence of the F763 propeptide. SEQ ID NO:29 is a nucleotide sequence for the gene coding for the F080 propeptide. SEQ ID NO:30 is an amino acid sequence of the F080 propeptide. SEQ ID NO:31 is a nucleotide sequence for the gene coding for the F008 propeptide. SEQ ID NO:32 is an amino acid sequence of the F008 propeptide. SEQ ID NO:33 is a nucleotide sequence for the gene coding for the G18 propeptide. SEQ ID NO:34 is an amino acid sequence of the G18 propeptide.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to the new μ O-conopeptides, their coding sequences and their propeptides and to the use of μ O-conopeptides as a local anesthetic for treating pain. The μ O-conopeptides have long lasting anesthetic activity and are particularly useful for spinal anesthesia,

either administered acutely for post-operative pain or via an intrathecal pump for severe chronic pain situations or for treatment of pain in epithelial tissue.

The present invention, in another aspect, relates to a pharmaceutical composition comprising an effective amount of a conotoxin peptide described herein or a pharmaceutically acceptable salt or solvate thereof. Such a pharmaceutical composition has the capability of acting as analgesic agents.

The present invention also provides for a method provides local anesthesia to a patient having pain. In one embodiment, the pain results from surgical or medical procedures, and the compounds are administered to the central nervous system (CNS), e.g. to the spine for spinal analgesia. In a second embodiment, the pain is in an epithelial tissue region associated with damage or loss of epithelial tissue as a result of, for example, plastic surgery, canker sores, burns, sore throats, genital lesions, upper or lower gastrointestinal bronchoscopy or endoscopy, intubation, dermatologic abrasions or chemical skin peels, and the compounds are administered to alleviate the associated pain.

The conotoxin peptides described herein are sufficiently small to be chemically synthesized. General chemical syntheses for preparing the foregoing conotoxin peptides are described hereinafter. Various ones of the conotoxin peptides can also be obtained by isolation and purification from specific *Conus* species using the technique described in U.S. Patent No. 4,447,356 (Olivera et al., 1984), the disclosure of which is incorporated herein by reference.

Although the conotoxin peptides of the present invention can be obtained by purification from cone snails, because the amounts of conotoxin peptides obtainable from individual snails are very small, the desired substantially pure conotoxin peptides are best practically obtained in commercially valuable amounts by chemical synthesis using solid-phase strategy. For example, the yield from a single cone snail may be about 10 micrograms or less of conotoxin peptide. By "substantially pure" is meant that the peptide is present in the substantial absence of other biological molecules of the same type; it is preferably present in an amount of at least about 85% purity and preferably at least about 95% purity. Chemical synthesis of biologically active conotoxin peptides depends of course upon correct determination of the amino acid sequence.

The conotoxin peptides can also be produced by recombinant DNA techniques well known in the art. Such techniques are described by Sambrook et al. (1989). The peptides produced in this manner are isolated, reduced if necessary, and oxidized to form the correct disulfide bonds.

One method of forming disulfide bonds in the peptides of the present invention is the air oxidation of the linear peptides for prolonged periods under cold room temperatures or at room temperature. This procedure results in the creation of a substantial amount of the bioactive, disulfide-linked peptides. The oxidized peptides are fractionated using reverse-phase high performance liquid chromatography (HPLC) or the like, to separate peptides having different linked configurations. Thereafter, either by comparing these fractions with the elution of the native material or by using a simple assay, the particular fraction having the correct linkage for maximum biological potency is easily determined. However, because of the dilution resulting from the presence of other fractions of less biopotency, a somewhat higher dosage may be required.

The peptides are synthesized by a suitable method, such as by exclusively solid-phase techniques, by partial solid-phase techniques, by fragment condensation or by classical solution couplings.

In conventional solution phase peptide synthesis, the peptide chain can be prepared by a series of coupling reactions in which constituent amino acids are added to the growing peptide chain in the desired sequence. Use of various coupling reagents, e.g., dicyclohexylcarbodiimide or diisopropylcarbonyldimidazole, various active esters, e.g., esters of N-hydroxyphthalimide or N-hydroxy-succinimide, and the various cleavage reagents, to carry out reaction in solution, with subsequent isolation and purification of intermediates, is well known classical peptide methodology. Classical solution synthesis is described in detail in the treatise, "Methoden der Organischen Chemie (Houben-Weyl): Synthese von Peptiden," (1974). Techniques of exclusively solid-phase synthesis are set forth in the textbook, "Solid-Phase Peptide Synthesis," (Stewart and Young, 1969), and are exemplified by the disclosure of U.S. Patent 4,105,603 (Vale et al., 1978). The fragment condensation method of synthesis is exemplified in U.S. Patent 3,972,859 (1976). Other available syntheses are exemplified by U.S. Patents No. 3,842,067 (1974) and 3,862,925 (1975). The synthesis of peptides containing γ -carboxyglutamic acid residues is exemplified by Rivier et al. (1987), Nishiuchi et al. (1993) and Zhou et al. (1996).

Common to such chemical syntheses is the protection of the labile side chain groups of the various amino acid moieties with suitable protecting groups which will prevent a chemical reaction from occurring at that site until the group is ultimately removed. Usually also common is the protection of an α -amino group on an amino acid or a fragment while that entity reacts at the carboxyl group, followed by the selective removal of the α -amino protecting group to allow subsequent reaction to take place at that location. Accordingly, it is common that, as a step in such

a synthesis, an intermediate compound is produced which includes each of the amino acid residues located in its desired sequence in the peptide chain with appropriate side-chain protecting groups linked to various ones of the residues having labile side chains.

As far as the selection of a side chain amino protecting group is concerned, generally one is chosen which is not removed during deprotection of the α -amino groups during the synthesis. However, for some amino acids, e.g., His, protection is not generally necessary. In selecting a particular side chain protecting group to be used in the synthesis of the peptides, the following general rules are followed: (a) the protecting group preferably retains its protecting properties and is not split off under coupling conditions, (b) the protecting group should be stable under the reaction conditions selected for removing the α -amino protecting group at each step of the synthesis, and (c) the side chain protecting group must be removable, upon the completion of the synthesis containing the desired amino acid sequence, under reaction conditions that will not undesirably alter the peptide chain.

It should be possible to prepare many, or even all, of these peptides using recombinant DNA technology. However, when peptides are not so prepared, they are preferably prepared using the Merrifield solid-phase synthesis, although other equivalent chemical syntheses known in the art can also be used as previously mentioned. Solid-phase synthesis is commenced from the C-terminus of the peptide by coupling a protected α -amino acid to a suitable resin. Such a starting material can be prepared by attaching an α -amino-protected amino acid by an ester linkage to a chloromethylated resin or a hydroxymethyl resin, or by an amide bond to a benzhydrylamine (BHA) resin or para-methylbenzhydrylamine (MBHA) resin. Preparation of the hydroxymethyl resin is described by Bodansky et al. (1966). Chloromethylated resins are commercially available from Bio Rad Laboratories (Richmond, CA) and from Lab. Systems, Inc. The preparation of such a resin is described by Stewart and Young (1969). BHA and MBHA resin supports are commercially available, and are generally used when the desired polypeptide being synthesized has an unsubstituted amide at the C-terminus. Thus, solid resin supports may be any of those known in the art, such as one having the formulae -O-CH₂-resin support, -NH BHA resin support, or -NH-MBHA resin support. When the unsubstituted amide is desired, use of a BHA or MBHA resin is preferred, because cleavage directly gives the amide. In case the N-methyl amide is desired, it can be generated from an N-methyl BHA resin. Should other substituted amides be desired, the teaching of U.S. Patent No. 4,569,967 (Kornreich et al., 1986) can be used, or should still other groups than

the free acid be desired at the C-terminus, it may be preferable to synthesize the peptide using classical methods as set forth in the Houben-Weyl text (1974).

5 The C-terminal amino acid, protected by Boc or Fmoc and by a side-chain protecting group, if appropriate, can be first coupled to a chloromethylated resin according to the procedure set forth in Horiki et al. (1978), using KF in DMF at about 60°C for 24 hours with stirring, when a peptide having free acid at the C-terminus is to be synthesized. Following the coupling of the BOC-protected amino acid to the resin support, the α -amino protecting group is removed, as by using trifluoroacetic acid (TFA) in methylene chloride or TFA alone. The deprotection is carried out at a temperature between about 0°C and room temperature. Other standard cleaving reagents, such as 10 HCl in dioxane, and conditions for removal of specific α -amino protecting groups may be used as described in Schroder & Lubke (1965).

15 After removal of the α -amino-protecting group, the remaining α -amino- and side chain-protected amino acids are coupled step-wise in the desired order to obtain the intermediate compound defined hereinbefore, or as an alternative to adding each amino acid separately in the synthesis, some of them may be coupled to one another prior to addition to the solid phase reactor. Selection of an appropriate coupling reagent is within the skill of the art. Particularly suitable as a coupling reagent is N,N'-dicyclohexylcarbodiimide (DCC, DIC, HBTU, HATU, TBTU in the presence of HoBt or HoAt).

20 The activating reagents used in the solid phase synthesis of the peptides are well known in the peptide art. Examples of suitable activating reagents are carbodiimides, such as N,N'-diisopropylcarbodiimide and N-ethyl-N'-(3-dimethylaminopropyl)carbodiimide. Other activating reagents and their use in peptide coupling are described by Schroder & Lubke (1965) and Kapoor (1970).

25 Each protected amino acid or amino acid sequence is introduced into the solid-phase reactor in about a twofold or more excess, and the coupling may be carried out in a medium of dimethylformamide (DMF):CH₂Cl₂ (1:1) or in DMF or CH₂Cl₂ alone. In cases where intermediate coupling occurs, the coupling procedure is repeated before removal of the α -amino protecting group prior to the coupling of the next amino acid. The success of the coupling reaction at each stage of the synthesis, if performed manually, is preferably monitored by the ninhydrin reaction, as described 30 by Kaiser et al. (1970). Coupling reactions can be performed automatically, as on a Beckman 990 automatic synthesizer, using a program such as that reported in Rivier et al. (1978).

After the desired amino acid sequence has been completed, the intermediate peptide can be removed from the resin support by treatment with a reagent, such as liquid hydrogen fluoride or TFA (if using Fmoc chemistry), which not only cleaves the peptide from the resin but also cleaves all remaining side chain protecting groups and also the α -amino protecting group at the N-terminus if it was not previously removed to obtain the peptide in the form of the free acid. If Met is present in the sequence, the Boc protecting group is preferably first removed using trifluoroacetic acid (TFA)/ethanedithiol prior to cleaving the peptide from the resin with HF to eliminate potential S-alkylation. When using hydrogen fluoride or TFA for cleaving, one or more scavengers such as anisole, cresol, dimethyl sulfide and methylethyl sulfide are included in the reaction vessel.

Cyclization of the linear peptide is preferably affected, as opposed to cyclizing the peptide while a part of the peptido-resin, to create bonds between Cys residues. To effect such a disulfide cyclizing linkage, fully protected peptide can be cleaved from a hydroxymethylated resin or a chloromethylated resin support by ammonolysis, as is well known in the art, to yield the fully protected amide intermediate, which is thereafter suitably cyclized and deprotected. Alternatively, deprotection, as well as cleavage of the peptide from the above resins or a benzhydrylamine (BHA) resin or a methylbenzhydrylamine (MBHA), can take place at 0°C with hydrofluoric acid (HF) or TFA, followed by oxidation as described above.

The peptides are also synthesized using an automatic synthesizer. Amino acids are sequentially coupled to an MBHA Rink resin (typically 100 mg of resin) beginning at the C-terminus using an Advanced Chemtech 357 Automatic Peptide Synthesizer. Couplings are carried out using 1,3-diisopropylcarbodiimide in N-methylpyrrolidinone (NMP) or by 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) and diethylisopro- pylethylamine (DIEA). The Fmoc protecting group is removed by treatment with a 20% solution of piperidine in dimethylformamide(DMF). Resins are subsequently washed with DMF (twice), followed by methanol and NMP.

Additional conotoxin peptides are identified by cloning by reverse transcription-polymerase chain reaction (RT-PCR) from cone snail venom duct mRNA. The PCR primers are based on the DNA sequences coding for the precursor peptides of the “O-Superfamily” as described herein. RT-PCR of venom duct mRNA produces a product of about 250-300 nucleotides in *Conus* species that express conotoxin genes. The PCR product is then cloned into a plasmid vector and individual clones are sequenced to determine the sequence of various conotoxin genes. Alternatively, cDNA libraries are prepared from *Conus* venom duct using conventional techniques. DNA from single

clones is amplified by conventional techniques using primers which correspond approximately to the M13 universal priming site and the M13 reverse universal priming site. Clones having a size of approximately 250 nucleotides are sequenced and screened for similarity in sequence to the propeptide described herein. In this manner, conotoxins having the basic structure and activity described herein are cloned from many *Conus* species.

Muteins, analogs or active fragments (collectively referred to herein as derivatives) of μ O-conopeptides are also contemplated for use as local anesthetics. See, e.g., Hammerland et al. (1992). Derivative muteins, analogs or active fragments of μ O-conopeptides may be synthesized according to known techniques, including conservative amino acid substitutions, such as outlined in U.S. Patent Nos. 5,545,723; 5,534,615 and 5,364,769. The derivative muteins, analogs or active fragments may be conveniently assayed for activity by using a hindlimb paralysis test such as described in Example 2 or a local anesthetic test such as described in Example 3.

A variety of peptides from *Conus* target sodium channels. μ -Conopeptides (i.e., GVIA) block sodium channels expressed by muscle cells (Olivera et al., 1990). δ -Conopeptides (i.e., GmVIA) delay the inactivation of neuronal sodium channels (Olivera et al., 1990). Another class of conopeptide (i.e., μ -PnIVA and μ -PnIVB; unfortunately also called μ but having a distinct cysteine framework from that which is considered a μ -conopeptide) blocks sodium channels in molluscan neurons, but has no effect on sodium currents in bovine chromaffin cells or in rat brain synaptosomes (Fainzilber et al., 1995). Finally, the μ O-conopeptides (MrVIA and MrVIB) block mammalian sodium channels (McIntosh et al., 1995).

Since the μ O-conopeptides have been shown to have a slow and incomplete washout from *Xenopus* oocytes expressing cloned rat type II sodium channels (Terlau et al., 1996), the present invention examined whether the μ O-conopeptides might represent a candidate for a long-lasting local anesthetic.

Thus, the present invention is directed to a method for inducing local analgesia by administering the μ O-conopeptides described herein. In one embodiment,

In a second embodiment, μ O-conopeptides are used to provide local anesthesia for pain associated with any epithelial tissue region in a subject, for example, pain associated with epithelial ulcers, such as a canker sore or genital lesions. Canker sores can occur alone or in groups on the inside of the cheek or lip or underneath the tongue. Severely affected people have continuously recurring ulcers which last for one to two weeks (Clayman). Genital ulcers are usually caused by sexually transmitted diseases, including herpes and syphilis. The early stages of syphilis are

characterized by a hard chancre, a painful ulcer where bacteria has penetrated the skin. This may be followed by shallow, elongated ulcers once the chancre has healed. Such ulcers are painful. Genital ulceration may also be a side effect of drugs taken orally or caused by solutions applied to genital warts. Pain in epithelial tissue is also caused by burns. Burns affecting the epidermal layer are usually associated with pain, restlessness and fever. Treatment of such a burn in accordance with the method of the invention can provide relief from the attendant pain. Pain as a result of damage to or loss of epithelial tissue is also associated with other conditions and procedures, such as sore throats and plastic surgery, for example carbon dioxide laser surgery to remove for skin resurfacing and removal of wrinkles (Rosenberg et al., 1996), burns, genital lesions, upper or lower 5 gastrointestinal bronchoscopy or endoscopy, intubation, dermatologic abrasions or chemical skin peels. The μ O-conopeptides administered in accordance with the method of the invention is 10 beneficial in relieving pain associated with such damaged tissues.

Pharmaceutical compositions containing a μ O-conopeptide or pharmaceutically acceptable salts thereof as the active ingredient (agent) can be prepared according to conventional pharmaceutical compounding techniques. See, for example, *Remington's Pharmaceutical Sciences*, 15 18th Ed. (1990, Mack Publishing Co., Easton, Pa.). Typically, a therapeutically effective amount of the active ingredient will be admixed with a pharmaceutically acceptable carrier. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., intravenous, oral or parenteral. The compositions may further contain antioxidantizing agents, 20 stabilizing agents, preservatives and the like.

"Pharmaceutical composition" means physically discrete coherent portions suitable for medical administration. "Pharmaceutical composition in dosage unit form" means physically discrete coherent units suitable for medical administration, each containing a daily dose or a multiple (up to four times) or a sub-multiple (down to a fortieth) of a daily dose of the active 25 compound in association with a carrier and/or enclosed within an envelope. Whether the composition contains a daily dose, or for example, a half, a third or a quarter of a daily dose, will depend on whether the pharmaceutical composition is to be administered once or, for example, twice, three times or four times a day, respectively.

The term "salt", as used herein, denotes acidic and/or basic salts, formed with inorganic or 30 organic acids and/or bases, preferably basic salts. While pharmaceutically acceptable salts are preferred, particularly when employing the compounds of the invention as medicaments, other salts

find utility, for example, in processing these compounds, or where non-medicament-type uses are contemplated. Salts of these compounds may be prepared by art-recognized techniques.

Examples of such pharmaceutically acceptable salts include, but are not limited to, inorganic and organic addition salts, such as hydrochloride, sulphates, nitrates or phosphates and acetates, trifluoroacetates, propionates, succinates, benzoates, citrates, tartrates, fumarates, maleates, methane-sulfonates, isothionates, theophylline acetates, salicylates, respectively, or the like. Lower alkyl quaternary ammonium salts and the like are suitable, as well.

As used herein, the term "pharmaceutically acceptable" carrier means a non-toxic, inert solid, semi-solid liquid filler, diluent, encapsulating material, formulation auxiliary of any type, or simply a sterile aqueous medium, such as saline. Some examples of the materials that can serve as pharmaceutically acceptable carriers are sugars, such as lactose, glucose and sucrose, starches such as corn starch and potato starch, cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt, gelatin, talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol, polyols such as glycerin, sorbitol, mannitol and polyethylene glycol; esters such as ethyl oleate and ethyl laurate, agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline, Ringer's solution; ethyl alcohol and phosphate buffer solutions, as well as other non-toxic compatible substances used in pharmaceutical formulations.

Wetting agents, emulsifiers and lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator. Examples of pharmaceutically acceptable antioxidants include, but are not limited to, water soluble antioxidants such as ascorbic acid, cysteine hydrochloride, sodium bisulfite, sodium metabisulfite, sodium sulfite, and the like; oil soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, aloha-tocopherol and the like; and the metal chelating agents such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid and the like.

For oral administration, the compounds can be formulated into solid or liquid preparations such as capsules, pills, tablets, lozenges, melts, powders, suspensions or emulsions. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents,

suspending agents, and the like in the case of oral liquid preparations (such as, for example, suspensions, elixirs and solutions); or carriers such as starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations (such as, for example, powders, capsules and tablets). Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar-coated or enteric-coated by standard techniques. The active agent can be encapsulated to make it stable to passage through the gastrointestinal tract while at the same time allowing for passage across the blood brain barrier. See for example, WO 96/11698.

For parenteral administration, the compound may be dissolved in a pharmaceutical carrier and administered as either a solution or a suspension. Illustrative of suitable carriers are water, saline, dextrose solutions, fructose solutions, ethanol, or oils of animal, vegetative or synthetic origin. The carrier may also contain other ingredients, for example, preservatives, suspending agents, solubilizing agents, buffers and the like. When the compounds are being administered intrathecally, they may also be dissolved in cerebrospinal fluid.

For topical administration, the compound may be formulated as an ointment, cream, gel or paste comprising the compound to be administered in a pharmaceutical acceptable carrier. One means of topical administration is a transdermal patch containing the compound to be administered.

A variety of administration routes are available. The particular mode selected will depend of course, upon the particular drug selected, the severity of the disease state being treated and the dosage required for therapeutic efficacy. The methods of this invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of the active compounds without causing clinically unacceptable adverse effects. Such modes of administration include oral, rectal, sublingual, topical, nasal, transdermal or parenteral routes. The term "parenteral" includes subcutaneous, intravenous, epidural, irrigation, intramuscular, release pumps, or infusion.

For example, administration of the active agent according to this invention may be achieved using any suitable delivery means, including:

- (a) pump (see, e.g., Lauer & Hatton (1993), Zimm et al. (1984) and Ettinger et al. (1978));
- (b) microencapsulation (see, e.g., U.S. Patent Nos. 4,352,883; 4,353,888; and 5,084,350);
- (c) continuous release polymer implants (see, e.g., U.S. Patent No. 4,883,666);

(d) macroencapsulation (see, e.g., U.S. Patent Nos. 5,284,761, 5,158,881, 4,976,859 and 4,968,733 and published PCT patent applications WO92/19195, WO 95/05452);

(e) naked or unencapsulated cell grafts to the CNS (see, e.g., U.S. Patent Nos. 5,082,670 and 5,618,531);

5 (f) injection, either subcutaneously, intravenously, intra-arterially, intramuscularly, or to other suitable site;

(g) oral administration, in capsule, liquid, tablet, pill, or prolonged release formulation; or

(h) topical (see, e.g., U.S. Patent Nos. 6,046,187 and 6,030,974).

In one embodiment of this invention, an active agent is delivered directly into the CNS, 10 preferably to the brain ventricles, brain parenchyma, the intrathecal space or other suitable CNS location, most preferably intrathecally.

Alternatively, targeting therapies may be used to deliver the active agent more specifically to certain types of cells, by the use of targeting systems such as antibodies or cell-specific ligands. Targeting may be desirable for a variety of reasons, e.g. if the agent is unacceptably toxic, if it 15 would otherwise require too high a dosage, or if it would not otherwise be able to enter target cells.

The active agents, which are peptides, can also be administered in a cell based delivery system in which a DNA sequence encoding an active agent is introduced into cells designed for implantation in the body of the patient, especially in the spinal cord region. Suitable delivery systems are described in U.S. Patent No. 5,550,050 and published PCT Application Nos. WO 20 92/19195, WO 94/25503, WO 95/01203, WO 95/05452, WO 96/02286, WO 96/02646, WO 96/40871, WO 96/40959 and WO 97/12635. Suitable DNA sequences can be prepared synthetically for each active agent on the basis of the developed sequences and the known genetic code.

The active agent is preferably administered in an therapeutically effective amount. By a 25 "therapeutically effective amount" or simply "effective amount" of an active compound is meant a sufficient amount of the compound to treat or alleviate pain or to induce analgesia at a reasonable benefit/risk ratio applicable to any medical treatment. The actual amount administered, and the rate and time-course of administration, will depend on the nature and severity of the condition being treated. Prescription of treatment, e.g. decisions on dosage, timing, etc., is within the responsibility 30 of general practitioners or specialists, and typically takes account of the disorder to be treated, the condition of the individual patient, the site of delivery, the method of administration and other factors known to practitioners. Examples of techniques and protocols can be found in *Remington's Pharmaceutical Sciences*.

For the treatment of pain, if the route of administration is directly to the CNS, the dosage contemplated is from about 1 ng to about 100 mg per day, preferably from about 100 ng to about 10 mg per day, more preferably from about 1 μ g to about 100 μ g per day. If administered peripherally, the dosage contemplated is somewhat higher, from about 100 ng to about 1000 mg per day, preferably from about 10 μ g to about 100 mg per day, more preferably from about 100 μ g to about 10 mg per day.

If the μ O-conopeptide is delivered by continuous infusion (e.g., by pump delivery, biodegradable polymer delivery or cell-based delivery), then a lower dosage is contemplated than for bolus delivery.

However, it will be understood that the amount of the active compound actually administered will be determined by a physician, in the light of the relevant circumstances including the condition to be treated, the chosen route of administration, the age, weight, and response of the individual patient, and the severity of the patient's symptoms, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way. As used herein the terms "pharmaceutical compositions" and "pharmaceutically acceptable" include compositions and ingredients for both human and veterinary use.

The present data suggest that μ O-conopeptides are extremely potent and long-lasting local anesthetic agents, most likely due to their ability to block neuronal sodium channels. Moreover, since μ O-conopeptides probably act at a site on sodium channels distinct from other local anesthetics or guanidinium toxins like tetrodotoxin (since they are likely to act at an extracellular target, but do compete for [3 H]saxitoxin at site I) (Terlau et al., 1996), and probably do not affect sodium channels in the muscles or heart (since i.p. injection of 10 nmol is without effect in mice (McIntosh et al., 1995), these peptides lack the untoward side effects of clinically used local anesthetics.

Despite the high hydrophobicity of these peptides, there is a cluster of charged amino acid residues at the amino terminus. This cluster of charge, combined with the size of the peptides, probably results in poor permeation of the nerve sheath and thus accounts for the poor efficacy in the tail withdrawal assay. In contrast, when the nerve sheath is not a barrier, such as following intrathecal injection or intracutaneous injection, μ O-conopeptides are effective and long-lasting. These facts establish that μ O-conopeptides are novel candidates for spinal anesthesia, either

administered acutely for post-operative pain or via an intrathecal pump for severe chronic pain situations.

The practice of the present invention employs, unless otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA, genetics, immunology, cell biology, cell culture and transgenic biology, which are within the skill of the art. See, e.g., Maniatis *et al.*, 1982; Sambrook *et al.*, 1989; Ausubel *et al.*, 1992; Glover, 1985; Anand, 1992; Guthrie and Fink, 1991; Harlow and Lane, 1988; Jakoby and Pastan, 1979; *Nucleic Acid Hybridization* (B. D. Hames & S. J. Higgins eds. 1984); *Transcription And Translation* (B. D. Hames & S. J. Higgins eds. 1984); *Culture Of Animal Cells* (R. I. Freshney, Alan R. Liss, Inc., 1987); *Immobilized Cells And Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide To Molecular Cloning* (1984); the treatise, *Methods In Enzymology* (Academic Press, Inc., N.Y.); *Gene Transfer Vectors For Mammalian Cells* (J. H. Miller and M. P. Calos eds., 1987, Cold Spring Harbor Laboratory); *Methods In Enzymology*, Vols. 154 and 155 (Wu *et al.* eds.), *Immunochemical Methods In Cell And Molecular Biology* (Mayer and Walker, eds., Academic Press, London, 1987); *Handbook Of Experimental Immunology*, Volumes I-IV (D. M. Weir and C. C. Blackwell, eds., 1986); Riott, *Essential Immunology*, 6th Edition, Blackwell Scientific Publications, Oxford, 1988; Hogan *et al.*, Manipulating the Mouse Embryo, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986).

20

EXAMPLES

The present invention is further detailed in the following Examples, which are offered by way of illustration and are not intended to limit the invention in any manner. Standard techniques well known in the art or the techniques specifically described below are utilized.

25

EXAMPLE 1

Isolation of μ O-conopeptides A657 and F079

PCR primers designed to amplify “O Superfamily” conotoxin genes were used in RT-PCR amplification of venom duct cDNA from a variety of *Conus* species. The primers have the following sequences:

Forward Primer: OCon6 CAGGATCCATGAAACTGACGTGYRTGGTG (SEQ ID NO:13)

Reverse Primer: OCon7 ATCTCGAGCACAGGTATGGATGACTCAGG (SEQ ID NO:14).

Amplification products in the appropriate size range were cloned and sequenced. A range of “O-Superfamily” gene sequences were identified. The novel genes, A657 from *C. skinneri*, F079, F080 and G28 from *C. tessulatus*, F763 from *C. atlanticus*, F008 from *C. arenatus*, Tx6.12 and Tx6.13 from *C. textile* and G18 from *C. generalis*, were identified as μ O-conopeptides on the basis of their similarity to the μ -O conopeptides MrVIA and MrVIB. This similarity was much greater than the similarity with any of the ω -, κ - or δ -conopeptides that comprise the “O Superfamily” peptides. The cDNA and amino acid sequence for the A657, F079, Ca6.1, Tx6.12 (portion), Tx6.13 (portion), G28, F763, F080, F008 and G18 propeptides are set forth in Tables 1-10, respectively. The amino acid sequences of the mature μ O-conopeptides are as shown above.

10

TABLE 1

DNA Sequence (SEQ ID NO:15) and Protein Sequence (SEQ ID NO:16) of A657

atg aaa ctg acg tgt gtg atc gtt gct gtg ctg ttc ttg acc gcc
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Phe Leu Thr Ala
 15
 tgg aca ttc gtc atg gct gat gac ccc aga gat gga gcg gag att aga
 Trp Thr Phe Val Met Ala Asp Asp Pro Arg Asp Gly Ala Glu Ile Arg
 20
 agc atg gta agg ggg gaa cct ctg tcg aag gca cgt gac gaa atg aac
 Ser Met Val Arg Gly Glu Pro Leu Ser Lys Ala Arg Asp Glu Met Asn
 ccc gaa gcc tct aaa ttg gag aaa agg gcg tgc cgc caa aaa tac gaa
 Pro Glu Ala Ser Lys Leu Glu Lys Arg Ala Cys Arg Gln Lys Tyr Glu
 25
 ttt tgt cta gta ccg atc att gga tac ata tat tgc tgc gct ggc tta
 Phe Cys Leu Val Pro Ile Ile Gly Tyr Ile Tyr Cys Cys Ala Gly Leu
 atc tgt ggt cct ttc gtc tgc ctt tgatagtat gtcttctact gccatctgtg
 Ile Cys Gly Pro Phe Val Cys Leu
 30
 ctacccctgg cttgatcttt gataggcggtt gttgccttc actggtttat gaaccctctg
 atcatactct ctggaccctt ggggtccaa catccaaata aagcgacatc ccaaaaaaaaa
 35
 aaaaaaaaaaa

TABLE 2

DNA Sequence (SEQ ID NO:17) and Protein Sequence (SEQ ID NO:18) of F079

40
 gga tcc atg aaa ctg acg tgc atg gtg atc gtt gtt gtg ctg ttg ttg
 Gly Ser Met Lys Leu Thr Cys Met Val Ile Val Val Leu Leu Leu
 aac gcc tgg aca ttc gtc tcc ata aat gga aag gcg aat cgt ttt tgg
 Asn Ala Trp Thr Phe Val Ser Ile Asn Gly Lys Ala Asn Arg Phe Trp
 45
 aag gca cgt gac gaa atg aag gac tcc gaa gtt tct gaa ttg gag aaa
 Lys Ala Arg Asp Glu Met Lys Asp Ser Glu Val Ser Glu Leu Glu Lys

agg agg aaa ccg acc tgc ctg aag cag gac aag ttt tgc ata ata ccg
 Arg Arg Lys Pro Thr Cys Leu Lys Gln Asp Lys Cys Ile Ile Pro
 5
 ctc att gga acc ctt tat tgc tgc agt ggg tta atc tgt ggg ttt ttt
 Leu Ile Gly Thr Leu Tyr Cys Cys Ser Gly Leu Ile Cys Gly Phe Phe
 gtc tgc gtc cca aag ccg ttc tgatgtcttc tactgccatc tgtgctaccc
 Val Cys Val Pro Lys Pro Phe
 10
 ctggcttgat ctttgattgg cgtgtgcct tcactggta tgaacccttc tgatcctact
 gtctggacgc ctcggcgtc caacgtccaa ataaagcgac atcccaataa aaaaaaaaaa
 15
 aaaaaaaaa

TABLE 3

DNA Sequence (SEQ ID NO:19) and Protein Sequence (SEQ ID NO:20) of Ca6.1

20 atg aaa ctg acg tgc gtg atg atc gtt gct gtg ctg ttc ttg acc gcc
 Met Lys Leu Thr Cys Val Met Ile Val Ala Val Leu Phe Leu Thr Ala
 tgg aca ttc gtc acg gct gat gac tcc att aat gca ctg gag gat ctt
 Trp Thr Phe Val Thr Ala Asp Asp Ser Ile Asn Ala Leu Glu Asp Leu
 25 ttt tcg aag gca cgt gac gaa atg gaa aac ggc gaa gct tct aca ttg
 Phe Ser Lys Ala Arg Asp Glu Met Glu Asn Gly Glu Ala Ser Thr Leu
 aac gag aga gac tgc gaa gca gat ggt gca ttt tgt ggt atc cca att
 30 Asn Glu Arg Asp Cys Glu Ala Asp Gly Ala Phe Cys Gly Ile Pro Ile
 gtg aag aac tgg atg tgc tgc agt aac ttg tgt att ttt gcc tgc gta
 Val Lys Asn Trp Met Cys Ser Asn Leu Cys Ile Phe Ala Cys Val
 35 ccc gag tat taagactgcc gtgatgtctt ctccctccct c
 Pro Glu Tyr

TABLE 4

DNA Sequence (SEQ ID NO:21) and Protein Sequence (SEQ ID NO:22) of Tx6.12

40 a ttg gag aaa agg gat tgc cac gaa agg tgg gat tgg tgt cca gca tca
 Leu Glu Lys Arg Asp Cys His Glu Arg Trp Asp Trp Cys Pro Ala Ser
 atc ctt gga gtg ata tat tgc tgc gag gga tta att tgt ttt att gcc
 Ile Leu Gly Val Ile Tyr Cys Cys Glu Gly Leu Ile Cys Phe Ile Ala
 45 ttc tgc att tgatagtgat gtcttctcct cccctc
 Phe Cys Ile

TABLE 5

DNA Sequence (SEQ ID NO:23) and Protein Sequence (SEQ ID NO:24) of Tx6.13

a ttg gag aaa agg gat tgc caa gag aaa tgg gag ttt tgt ata gta ccg
 Leu Glu Lys Arg Asp Cys Gln Glu Lys Trp Glu Phe Cys Ile Val Pro
 55 atc ctt gga ttt gta tat tgc tgc cct ggc tta atc tgt ggc cct ttt
 Ile Leu Gly Phe Val Tyr Cys Cys Pro Gly Leu Ile Cys Gly Pro Phe

5 gtc tgc gtt gat atc tgatgtcttc tcctccatc
Val Cys Val Asp Ile

5 TABLE 6

DNA Sequence (SEQ ID NO:25) and Protein Sequence (SEQ ID NO:26) of G28

ggatcc atg aaa ctg acg tgt gtg atc gtt gtt gtg ctg ttg ttg
Met Lys Leu Thr Cys Val Val Ile Val Val Val Leu Leu Leu

10 aac gcc tgg aca ttc gtc tcc ata aat gga aag gcg aat cct ttt tgg
Asn Ala Trp Thr Phe Val Ser Ile Asn Gly Lys Ala Asn Pro Phe Trp

15 aag gca cgt gac gaa atg aag gac tcc gaa gtt tct gag ttg gag aaa
Lys Ala Arg Asp Glu Met Lys Asp Ser Glu Val Ser Glu Leu Glu Lys

15 agg agg aaa ccg acc tgc gtg tcg tat aac gtg ttt tgc gga gta ccg
Arg Arg Lys Pro Thr Cys Val Ser Tyr Asn Val Phe Cys Gly Val Pro

20 ctc gtt gga acc tac ctt tgc tgc agt ggc tta gtc tgt ctc gta gtc
Leu Val Gly Thr Tyr Leu Cys Cys Ser Gly Leu Val Cys Leu Val Val

tgc atc tagtactgat gtcttctact cccatctgtg ctaccctcg ag
Cys Ile

25

TABLE 7

DNA Sequence (SEQ ID NO:27) and Protein Sequence (SEQ ID NO:28) of F763

ggatcc atg aaa ctg acg tgc gtg gtg atc gtt gct gtg ctg ttc ttg
Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu

30 acc gcc tgg aca ttc gtc acg gct gat gac tcc ata aat ggg ttg gag
Thr Ala Trp Thr Phe Val Thr Ala Asp Asp Ser Ile Asn Gly Leu Glu

35 aat ctt ttt ccg aag gca cgt cac gaa atg agg aaa ccc gaa gcc tct
Asn Leu Phe Pro Lys Ala Arg His Glu Met Arg Lys Pro Glu Ala Ser

aga tcg aga ggg agg tgc cgt cct cgt ggt atg ttc tgt ggc ttt ccg
Arg Ser Arg Gly Arg Cys Arg Pro Arg Gly Met Phe Cys Gly Phe Pro

40 aaa cct gga cca tac tgc tgc aat ggc tgg tgc ttt ttc gtc tgc atc
Lys Pro Gly Pro Tyr Cys Cys Asn Gly Trp Cys Phe Phe Val Cys Ile

taaaaactgcc gtgatgtgtt ctactccat ctgtgctacc cctcgag

45

TABLE 8

DNA Sequence (SEQ ID NO:29) and Protein Sequence (SEQ ID NO:30) of F080

ggatcc atg aaa ctg acg tgc gtg gtg gtc gtt gct gtg ctg ttc ttg
Met Lys Leu Thr Cys Val Val Val Ala Val Leu Phe Leu

50 aac gcc tgg aca ttc gcc acg gct gtt gac tcc aaa cat gca ctg gcg
Asn Ala Trp Thr Phe Ala Thr Ala Val Asp Ser Lys His Ala Leu Ala

55 aaa ctt ttt atg aag gca cgt gac gaa atg tat aac ccc gat gcc act
Lys Leu Phe Met Lys Ala Arg Asp Glu Met Tyr Asn Pro Asp Ala Thr

aaa ttg gac gat aag aga tgg tgc gct tta gat ggt gaa ctt tgt atc
Lys Leu Asp Asp Lys Arg Trp Cys Ala Leu Asp Gly Glu Leu Cys Ile

ata ccg gtc att ggg tcc ata ttt tgc tgc cat ggc ata tgt atg atc
 Ile Pro Val Ile Gly Ser Ile Phe Cys Cys His Gly Ile Cys Met Ile
 5 tac tgc gtc tagttgaact gccgtgatgt cttctactcc cctctgtgct
 Tyr Cys Val
 acccctgggtt tgatcttga ttgcctgtg cccttcactg attatgaatc cctctgatcc
 10 tactctctga agacctcttg gggccaaca tccaaataaa gcgacatccc aaaaaaaaaa
 aaaaaaaaaa

TABLE 9

15 DNA Sequence (SEQ ID NO:31) and Protein Sequence (SEQ ID NO:32) of F008

ggatcc atg aaa ctg acg tgt gtg gtg atc gtt gct gtg ctg ttc ttg
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Phe Leu
 20 acc gcc tgg aca ttc gtc acg gct gac tcc ata cgt gca ctg gag gat
 Thr Ala Trp Thr Phe Val Thr Ala Asp Ser Ile Arg Ala Leu Glu Asp
 ttt ttt gcg aag gca cgt gac gaa atg gaa aac agc gga gct tct cca
 Phe Phe Ala Lys Ala Arg Asp Glu Met Glu Asn Ser Gly Ala Ser Pro
 25 ttg aac gag aga gac tgc cga cct gta ggt caa tat tgt ggc ata ccg
 Leu Asn Glu Arg Asp Cys Arg Pro Val Gly Gln Tyr Cys Gly Ile Pro
 tat aag cac aac tgg cga tgc agt cag ctt tgt gca att atc tgt
 Tyr Lys His Asn Trp Arg Cys Ser Gln Leu Cys Ala Ile Ile Cys
 30 gtt tcc taaccctct gatcctactc tctgaagacc tccggattc aacatccaa
 Val Ser
 taaagcgaca tcccgatnaa aaaaaangaa aaaaaaaaaa aaaa

TABLE 10

35 DNA Sequence (SEQ ID NO:33) and Protein Sequence (SEQ ID NO:34) of G18

ggatcc atg aaa ctg acg tgt gtg gtg atc gtt gct gtg cta ttc ttg
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Phe Leu
 acc gcc tgg aca ttc gtc acg gct gat gac acc aga tat aaa ctg gag
 Thr Ala Trp Thr Phe Val Thr Ala Asp Asp Thr Arg Tyr Lys Leu Glu
 40 aat cct ttt ctg aag gca cgc aac gaa ctg cag aaa cac gaa gcc tct
 Asn Pro Phe Leu Lys Ala Arg Asn Glu Leu Gln Lys His Glu Ala Ser
 caa ctg aac gag aga ggc tgc ctt gac cca ggt tac ttc tgt ggg acg
 Gln Leu Asn Glu Arg Gly Cys Leu Asp Pro Gly Tyr Phe Cys Gly Thr
 45 ccg ttt ctt gga gca tac tgc tgc ggt ggc att tgc ctt att gtc tgc
 Pro Phe Leu Gly Ala Tyr Cys Cys Gly Gly Ile Cys Leu Ile Val Cys
 ata gaa acg taaaggcttg atgtcttcta ctcccatctg tgctaccct cgag
 55 Ile Glu Thr

EXAMPLE 2

Effect of Intrathecal Administration of MrVIB

Male C57 black mice (20-25g) were obtained from Charles River Laboratories. These mice and the animals used in the other examples were housed in a temperature controlled (23° ± 3° C) room with a 12 hour light-dark cycle with free access to food and water. All animals were euthanized in accordance with Public Health Service policies on the humane care of laboratory animals.

Intrathecal (it) drug injections were performed as described (Hylden and Wilcox, 1980). MrVIB (10 nmol) or vehicle was administered in a volume of 5 µl. Duration of hind-limb paralysis was assessed. This experiment revealed that injection of 10 nmols of MrVIB into the intrathecal space of C57 black mice produced a long-lasting paralysis (>20 hrs) of the animal. The injection initially produced a paralysis of the hind-limbs, but over the following 30 minutes resolved into paralysis of the entire animal. Despite the long duration of anesthesia, the animals in this experiment recovered fully. Similar results were obtained with MrVIA. Similar results are also obtained with A657, F079, Ca6.1, Tx6.12, Tx6.13, G28, F763 and F080.

EXAMPLE 3

Effect of MrVIB as a Local Anesthetic

Male Hartley guinea pigs (retired breeders) were obtained from Charles River Laboratories. The local anesthetic test was performed essentially as described (Bulbring and Wajda, 1945). On the day prior to test day, a patch on the back of the guinea pig was denuded of hair, first by shaving with electric clippers and subsequently with depilatory cream (Nair®). Depilatory cream was applied for five minutes and removed with a warm washcloth. The guinea pigs were dried and returned to their cages. On the following day, intradermal injections (0.1 ml vols) of lidocaine, bupivacaine, MrVIB or vehicle (0.5% cyclodextran) were made into the denuded patch. The injection produced a raised wheal on the surface of the skin which was circled with a felt-tipped pen. Typically, four injections were made on the back of each guinea pig. In some cases, guinea pigs were reused following at least one week of recovery and injecting into an unused portion of the skin.

The stimulus consisted of mild pin pricks (not hard enough to break the skin) with a 26G needle. The response is a localized skin twitch caused by contraction of cutaneous muscles. A unit test consisted of six uniform pin pricks, 3-5 seconds apart, within the injected area. Unit scores ranged from 0 (complete anesthesia) to 6 (no anesthesia). For potency experiments, the unit test was

repeated at each site at five minute intervals for 30 minutes, and unit test scores summed (with 36 representing no anesthesia to 0 representing complete anesthesia. For duration experiments, unit tests were performed as described over the course of several hours to days.

MrVIB produced a potent (Figure 1) and long lasting (Figure 2) local anesthetic effect in the 5 intracutaneous wheal test in the guinea pig. The ED₅₀ for this response (\approx 100 pmol) was at least two orders of magnitude greater than the ED₅₀'s for lidocaine and bupivacaine. Moreover, the duration of roughly equieffective doses of MrVIB (roughly 24 and 48 hours for full recovery following 1 and 10 nmol, respectively) was much longer than that of lidocaine and bupivacaine (\approx 10 30 and 90 minutes for full recovery, respectively). As expected, bupivacaine had a slightly longer duration than lidocaine, consistent with clinical observations. It was seen during the experiment that the intracutaneous wheal consistently turned red several hours following injection of MrVIB, possibly suggesting an antigenic action. Similar results are obtained with MrVIA, A657, F079, Ca6.1, Tx6.12, Tx6.13, G28, F763 and F080.

15 While the invention has been disclosed in this patent application by reference to the details of preferred embodiments of the invention, it is to be understood that the disclosure is intended in an illustrative rather than in a limiting sense, as it is contemplated that modifications will readily occur to those skilled in the art, within the spirit of the invention and the scope of the appended 20 claims.

LIST OF REFERENCES

Barnay, G. et al. (2000). *J. Med. Chem.*

Bitan, G. et al. (1997). *J. Peptide Res.* **49**:421-426.

Bodansky et al. (1966). *Chem. Ind.* **38**:1597-98.

Bulbring, W. and Wajda, J. (1945). *J. Pharmacol. Exp. Ther.* **85**:78-84.

Ettinger, L.J. et al. (1978). *Cancer* **41**:1270-1273.

Fainzilber, M. et al. (1995). *Biochemistry* **34**:8649-8656.

Hammerland et al. (1992). *Eur. J. Pharmacol.* **226**:239-242.

Horiki, K. et al. (1978). *Chemistry Letters* 165-68.

Hubry, V. et al. (1994). *Reactive Polymers* **22**:231-241.

Hylden, J.L.K. and Wilcox, G. (1980). *Eur. J. Pharmacol.* **67**:313-316.

Kaiser et al. (1970). *Anal. Biochem.* **34**:595.

Kapoor (1970). *J. Pharm. Sci.* **59**:1-27.

Kornreich, W.D. et al. (1986). U.S. Patent No. 4,569,967.

Kuzma, P.K. et al. (1997). *Regional Anesthesia* **22**:543-551.

Luer, M.S. et al. (1993). *Annals of Pharmacotherapy*, **27**:912-921.

5 McIntosh, J.M. et al. (1995). *J. Biol. Chem.* **270**:16796-16802.

McIntosh, J. M. et al. (1998). *Methods Enzymol.* **294**:605-624.

Methoden der Organischen Chemie (Houben-Weyl): Synthese von Peptiden, E. Wunsch (Ed.), Georg Thieme Verlag, Stuttgart, Ger. (1974).

Nishiuchi, Y. et al. (1993). *Int. J. Pept. Protein Res.* **42**:533-538.

10 Nowak, L. et al. (1984). *Nature* **307**:462-465.

Olivera, B.M. et al. (1984). U.S. Patent 4,447,356.

Olivera, B.M. et al. (1985). *Science* **230**:1338-1343.

Olivera, B.M. et al. (1990). *Science* **249**:257-263.

Ornstein, et al. (1993). *Biorganic Medicinal Chemistry Letters* **3**:43-48.

15 *Remington's Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co., Easton, PA (1990).

Rivier, J.R. et al. (1978). *Biopolymers* **17**:1927-38.

Rivier, J.R. et al. (1987). *Biochem.* **26**:8508-8512.

Rosenberg, G.J. et al. (1996). *Clinics in Plastic Surgery* **23**:29.

20 Sambrook, J. et al. (1989). *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.

Schroder & Lubke (1965). *The Peptides* **1**:72-75, Academic Press, NY.

Stewart and Young, *Solid-Phase Peptide Synthesis*, Freeman & Co., San Francisco, CA (1969).

Terlau et al. (1996). *J. Neurophysiol.* **76**:1423-1429.

Vale et al. (1978). U.S. Patent 4,105,603.

25 Van de Steen, P. et al. (1998). *Critical Rev. in Biochem. and Mol. Biol.* **33**:151-208.

Zhou L.M., et al. (1996). *J. Neurochem.* **66**:620-628.

Zimm, S. et al. (1984). *Cancer Research*, **44**:1698-1701.

U.S. Patent No. 3,842,067.

U.S. Patent No. 3,862,925.

30 U.S. Patent No. 3,972,859.

U.S. Patent No. 4,352,883

U.S. Patent No. 4,353,888

U.S. Patent No. 4,883,666
U.S. Patent No. 4,968,733
U.S. Patent No. 4,976,859
U.S. Patent No. 5,082,670
5 U.S. Patent No. 5,084,350
U.S. Patent No. 5,158,884
U.S. Patent No. 5,284,761
U.S. Patent No. 5,364,769
U.S. Patent No. 5,514,774.
10 U.S. Patent No. 5,531,001.
U.S. Patent No. 5,534,615
U.S. Patent No. 5,545,723
U.S. Patent No. 5,550,050.
U.S. Patent No. 5,591,821.
15 U.S. Patent No. 5,618,531
U.S. Patent No. 5,719,264
U.S. Patent No. 5,844,077.
U.S. Patent No. 5,859,186.
PCT Published Application WO 92/19195.
20 PCT Published Application WO 94/25503.
PCT Published Application WO 95/01203.
PCT Published Application WO 95/05452.
PCT Published Application WO 96/02286.
PCT Published Application WO 96/02646.
25 PCT Published Application WO 96/11698.
PCT Published Application WO 96/40871.
PCT Published Application WO 96/40959.
PCT Published Application WO 97/12635.
PCT Published Application WO 98/03189.
30 PCT Published Application WO 00/23092.

WHAT IS CLAIMED IS:

1. A method of alleviating pain which comprises administering to a mammal that is either exhibiting pain or is about to be subjected to a pain-causing event a pain-alleviating amount of an active agent comprising a μ O-conopeptide, derivative or pharmaceutically acceptable salt or solvate thereof.
2. The method of claim 1, wherein said μ O-conopeptide has the general formula I:

10 Xaa₁-Xaa₂-Cys-Xaa₃-Xaa₄-Xaa₅-Xaa₆-Xaa₇-Xaa₈-Cys-Xaa₉-Xaa₁₀-Xaa₁₁-Xaa₁₂-Xaa₁₃-Xaa₁₄-Xaa₁₅-Xaa₁₆-Xaa₁₇-Cys-Cys-Xaa₁₈-Xaa₁₉-Xaa₂₀-Xaa₂₁-Cys-Xaa₂₂-Xaa₂₃-Xaa₂₄-Xaa₂₅-Cys-Xaa₂₆-Xaa₂₇-Xaa₂₈-Xaa₂₉-Xaa₃₀ (SEQ ID NO:1),

15 wherein Xaa₁ is des-Xaa₁, Pro, hydroxy-Pro (Hyp), Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₂ is des-Xaa₂, Ala, Gly, Asp, Glu, γ -carboxy-glutamate (Gla), any synthetic acidic amino acid, Thr, Ser, g-Thr (where g is glycosylation), g-Ser, Trp (D or L), neo-Trp or halo-Trp (D or L) or Xaa₂ may be pyroglutamate if Xaa₁ is des-Xaa₁; Xaa₃ is Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ser, Thr, g-Ser, g-Thr, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, His, Glu, Gln, Gla, Asp, Asn or any synthetic acidic amino acid; Xaa₄ is Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Ser, Thr, Pro, Hyp, g-Ser, g-Thr, g-Hyp or any synthetic hydroxylated amino acid; Xaa₅ is Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Glu, Gla, Gln, Asp, Asn, any synthetic

acidic amino acid, Pro or Hyp; Xaa₆ is Trp (D or L), neo-Trp, halo-Trp (D or L), Gly, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid; Xaa₇ is Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Met, norleucine (Nle), Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₈ is Leu, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₉ is Pro, Hyp, Gly, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₁₀ is Thr, Ser, g-Thr, g-Ser, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₁₁ is Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Thr or any hydroxylated amino acid; Xaa₁₂ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₃ is Pro, Hyp, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₄ is Gly, His, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₅ is des-Xaa₁₅, Ser, Thr, g-Ser, g-Thr, Val, Asn, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp,

halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₁₆ is Met, Nle, Leu, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₇ is Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Thr, any hydroxylated amino acid, Ala, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, His or Gly; Xaa₁₈ is Gly, Asn or Gln; Xaa₁₉ is Leu, Trp (D or L), neo-Trp or halo-Trp (D or L); Xaa₂₀ is des-Xaa₂₀, Leu or Trp (D or L), neo-Trp or halo-Trp (D or L); Xaa₂₁ is des-Xaa₂₁ or an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₂₂ is des-Xaa₂₂, Gly, Met, Nle, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₃ is des-Xaa₂₃, Pro, Hyp, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₄ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₅ is Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₆ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₂₇ is des-Xaa₂₇, Asp, Glu, Gla, Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Ser or any synthetic hydroxylated amino acid; Xaa₂₈ is des-Xaa₂₈, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ile, Ser, Thr, g-Ser or g-Thr; Xaa₂₉ is des-Xaa₂₉, Pro, Hyp, Tyr,

meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa₃₀ is des-Xaa₃₀ or Phe.

3. The method of claim 2, wherein said μ O-conopeptide is selected from the group consisting of:

5 Ala-Cys-Arg-Gln-Xaa₁-Xaa₂-Xaa₃-Phe-Cys-Leu-Val-Xaa₄-Ile-Ile-Gly-Xaa₂-Ile-Xaa₂-Cys-Cys-Ala-Gly-Leu-Ile-Cys-Gly-Xaa₄-Phe-Val-Cys-Leu (SEQ ID NO:3);

10 Xaa₄-Thr-Cys-Leu-Xaa₁-Gln-Asp-Xaa₁-Phe-Cys-Ile-Ile-Xaa₄-Leu-Ile-Gly-Thr-Leu-Xaa₂-Cys-Cys-Ser-Gly-Leu-Ile-Cys-Gly-Phe-Phe-Val-Cys-Val-Xaa₄-Xaa₁-Xaa₄-Phe (SEQ ID NO:4);

15 Asp-Cys-Xaa₃-Ala-Asp-Gly-Ala-Phe-Cys-Gly-Ile-Xaa₄-Ile-Vla-Xaa₁-Asn-Xaa₅-Met-Cys-Cys-Ser-Asn-Leu-Cys-Ile-Phe-Ala-Cys-Val-Xaa₄-Xaa₃-Xaa₂ (SEQ ID NO:5);

20 Asp-Cys-His-Xaa₃-Arg-Xaa₅-Asp-Xaa₅-Cys-Xaa₄-Ala-Ser-Ile-Leu-Gly-Val-Ile-Xaa₂-Cys-Cys-Xaa₃-Gly-Leu-Ile-Cys-Phe-Ile-Ala-Phe-Cys-Ile (SEQ ID NO:6);

25 Asp-Cys-Gln-Xaa₃-Xaa₁-Xaa₅-Xaa₃-Phe-Cys-Ile-Val-Xaa₄-Ile-Leu-Gly-Phe-Val-Xaa₂-Cys-Cys-Xaa₄-Gly-Leu-Ile-Cys-Gly-Xaa₄-Phe-Val-Cys-Val-Asp-Ile (SEQ ID NO:7);

30 Xaa₄-Thr-Cys-Val-Ser-Xaa₂-Asn-Val-Phe-Cys-Gly-Val-Xaa₄-Leu-Val-Gly-Thr-Xaa₂-Leu-Cys-Ser-Gly-Leu-Val-Cys-Leu-Val-Val-Cys-Ile (SEQ ID NO:8);

Arg-Xaa₅-Cys-Ala-Leu-Asp-Gly-Xaa₃-Leu-Cys-Ile-Ile-Xaa₄-Val-Ile-Gly-Ser-Ile-Phe-Cys-Cys-His-Gly-Ile-Cys-Met-Ile-Xaa₂-Cys-Val (SEQ ID NO:10);

Asp-Cys-Arg-Xaa₄-Val-Gly-Gln-Xaa₂-Cys-Gly-Ile-Xaa₄-Xaa₂-Xaa₁-His-Asn-Xaa₅-Arg-Cys-Cys-Ser-Gln-Leu-Cys-Ala-Ile-Ile-Cys-Val-Ser (SEQ ID NO:11); and

Gly-Cys-Leu-Asp-Xaa₄-Gly-Xaa₂-Phe-Cys-Gly-Thr-Xaa₄-Phe-Leu-Gly-Ala-Xaa₂-Cys-Cys-Gly-Gly-Ile-Cys-Leu-Ile-Val-Cys-Ile-Xaa₃-Thr (SEQ ID NO:12),

30 wherein Xaa₁ is Lys, N-methy-Lys, N,N-dimethyl-Lys or N,N,N-trimethyl-Lys; Xaa₂ is Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa₃ is Glu or gamma-carboxy-Glu (Gla); Xaa₄ is Pro or hydroxy-Pro; Xaa₅ is Trp or halo-Trp; and the C-terminus contains a carboxyl or amide group.

4. The method of claim 1, wherein the pain is chronic pain or acute inflammatory pain.

5. The method of claim 1, wherein the pain is neuropathic pain.
6. The method of claim 1, wherein the active agent is administered prior to surgery.
- 5 7. The method of claim 1, wherein the active agent is administered as a spinal anesthetic.
8. The method of claim 1, wherein the active agent is administered as a local anesthetic.
9. The method of claim 1, wherein said active agent is administered in an amount from about 10 1 ng to about 1000 mg per day.
10. The method of claim 1, wherein said active agent is administered in an amount from about 100 ng to about 100 mg per day.
- 15 11. The method of claim 1, wherein said active agent is administered in an amount from about 1 μ g to about 10 mg per day.
12. An isolated nucleic acid comprising a nucleic acid coding for a μ O-conopeptide precursor comprising an amino acid sequence selected from the group of amino acid sequences set 20 forth in Tables 1-10.
13. The nucleic acid of claim 12 wherein the nucleic acid comprises a nucleotide sequence selected from the group of nucleotide sequences set forth in Tables 1-10 or their complements.
- 25 14. A substantially pure μ O-conopeptide precursor comprising an amino acid sequence selected from the group of amino acid sequences set forth in Tables 1-10.
15. A substantially pure μ O-conopeptide having the generic formula I: Xaa₁-Xaa₂-Cys-Xaa₃-Xaa₄-Xaa₅-Xaa₆-Xaa₇-Xaa₈-Cys-Xaa₉-Xaa₁₀-Xaa₁₁-Xaa₁₂-Xaa₁₃-Xaa₁₄-Xaa₁₅-Xaa₁₆-Xaa₁₇-Cys-Cys-Xaa₁₈-Xaa₁₉-Xaa₂₀-Xaa₂₁-Cys-Xaa₂₂-Xaa₂₃-Xaa₂₄-Xaa₂₅-Cys-Xaa₂₆-Xaa₂₇-Xaa₂₈-Xaa₂₉-Xaa₃₀ (SEQ ID NO:1), wherein Xaa₁ is des-Xaa₁, Pro, hydroxy-Pro (Hyp), Arg, Lys, 30

ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₂ is des-Xaa₂, Ala, Gly, Asp, Glu, γ -carboxy-glutamate (Gla), any synthetic acidic amino acid, Thr, Ser, g-Thr (where g is glycosylation), g-Ser, Trp (D or L), neo-Trp or halo-Trp (D or L) or Xaa₂ may be pyroglutamate if Xaa₁ is des-Xaa₁; Xaa₃ is Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ser, Thr, g-Ser, g-Thr, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, His, Glu, Gln, Gla, Asp, Asn or any synthetic acidic amino acid; Xaa₄ is Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Ser, Thr, Pro, Hyp, g-Ser, g-Thr, g-Hyp or any synthetic hydroxylated amino acid; Xaa₅ is Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Pro or Hyp; Xaa₆ is Trp (D or L), neo-Trp, halo-Trp (D or L), Gly, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid; Xaa₇ is Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Met, norleucine (Nle), Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₈ is Leu, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₉ is Pro, Hyp, Gly, an aliphatic amino acids bearing

linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₁₀ is Thr, Ser, g-Thr, g-Ser, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₁₁ is Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Thr or any hydroxylated amino acid; Xaa₁₂ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₃ is Pro, Hyp, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₄ is Gly, His, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₅ is des-Xaa₁₅, Ser, Thr, g-Ser, g-Thr, Val, Asn, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₁₆ is Met, Nle, Leu, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys or any synthetic basic amino acid; Xaa₁₇ is Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Thr, any hydroxylated amino acid, Ala, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, His or Gly; Xaa₁₈ is Gly, Asn or Gln; Xaa₁₉ is Leu, Trp (D or L), neo-Trp or halo-Trp (D or L); Xaa₂₀ is des-Xaa₂₀, Leu or Trp (D or L), neo-Trp or halo-Trp (D or L); Xaa₂₁ is des-Xaa₂₁ or an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₂₂ is des-Xaa₂₂, Gly, Met, Nle, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid.

amino acid; Xaa₂₃ is des-Xaa₂₃, Pro, Hyp, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₄ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa₂₅ is Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa₂₆ is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid; Xaa₂₇ is des-Xaa₂₇, Asp, Glu, Gla, Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Ser or any synthetic hydroxylated amino acid; Xaa₂₈ is des-Xaa₂₈, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N"-trimethyl-Lys, any synthetic basic amino acid, Ile, Ser, Thr, g-Ser or g-Thr; Xaa₂₉ is des-Xaa₂₉, Pro, Hyp, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa₃₀ is des-Xaa₃₀ or Phe, with the proviso that said μ O-conopeptide is not MrVIA/B.

16. The substantially pure μ O-conopeptide of claim 15 selected from the group consisting of:

Ala-Cys-Arg-Gln-Xaa₁-Xaa₂-Xaa₃-Phe-Cys-Leu-Val-Xaa₄-Ile-Ile-Gly-Xaa₂-Ile-Xaa₂-Cys-Cys-Ala-Gly-Leu-Ile-Cys-Gly-Xaa₄-Phe-Val-Cys-Leu (SEQ ID NO:3);
Xaa₄-Thr-Cys-Leu-Xaa₁-Gln-Asp-Xaa₁-Phe-Cys-Ile-Ile-Xaa₄-Leu-Ile-Gly-Thr-Leu-Xaa₂-Cys-Cys-Ser-Gly-Leu-Ile-Cys-Gly-Phe-Phe-Val-Cys-Val-Xaa₄-Xaa₁-Xaa₄-Phe (SEQ ID NO:4);
Asp-Cys-Xaa₃-Ala-Asp-Gly-Ala-Phe-Cys-Gly-Ile-Xaa₄-Ile-Vla-Xaa₁-Asn-Xaa₅-Met-Cys-Cys-Ser-Asn-Leu-Cys-Ile-Phe-Ala-Cys-Val-Xaa₄-Xaa₃-Xaa₂ (SEQ ID NO:5);

Asp-Cys-His-Xaa₃-Arg-Xaa₅-Asp-Xaa₅-Cys-Xaa₄-Ala-Ser-Ile-Leu-Gly-Val-Ile-Xaa₂-Cys-Cys-Xaa₃-Gly-Leu-Ile-Cys-Phe-Ile-Ala-Phe-Cys-Ile (SEQ ID NO:6);

Asp-Cys-Gln-Xaa₃-Xaa₁-Xaa₅-Xaa₃-Phe-Cys-Ile-Val-Xaa₄-Ile-Leu-Gly-Phe-Val-Xaa₂-Cys-Cys-Xaa₄-Gly-Leu-Ile-Cys-Gly-Xaa₄-Phe-Val-Cys-Val-Asp-Ile (SEQ ID NO:7);

Xaa₄-Thr-Cys-Val-Ser-Xaa₂-Asn-Val-Phe-Cys-Gly-Val-Xaa₄-Leu-Val-Gly-Thr-Xaa₂-Leu-Cys-Cys-Ser-Gly-Leu-Val-Cys-Leu-Val-Val-Cys-Ile (SEQ ID NO:8);

Cys-Arg-Xaa₄-Arg-Gly-Met-Phe-Cys-Gly-Phe-Xaa₄-Xaa₁-Xaa₄-Gly-Xaa₄-Xaa₂-Cys-Cys-Asn-Gly-Xaa₅-Cys-Phe-Phe-Val-Cys-Ile (SEQ ID NO:9);

Arg-Xaa₅-Cys-Ala-Leu-Asp-Gly-Xaa₃-Leu-Cys-Ile-Ile-Xaa₄-Val-Ile-Gly-Ser-Ile-Phe-Cys-Cys-His-Gly-Ile-Cys-Met-Ile-Xaa₂-Cys-Val (SEQ ID NO:10);

Asp-Cys-Arg-Xaa₄-Val-Gly-Gln-Xaa₂-Cys-Gly-Ile-Xaa₄-Xaa₂-Xaa₁-His-Asn-Xaa₅-Arg-Cys-Cys-Ser-Gln-Leu-Cys-Ala-Ile-Ile-Cys-Val-Ser (SEQ ID NO:11); and

Gly-Cys-Leu-Asp-Xaa₄-Gly-Xaa₂-Phe-Cys-Gly-Thr-Xaa₄-Phe-Leu-Gly-Ala-Xaa₂-Cys-Cys-Gly-Gly-Ile-Cys-Leu-Ile-Val-Cys-Ile-Xaa₃-Thr (SEQ ID NO:12),

wherein Xaa₁ is Lys, N-methy-Lys, N,N-dimethyl-Lys or N,N,N-trimethyl-Lys; Xaa₂ is Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa₃ is Glu or gamma-carboxy-Glu (Gla); Xaa₄ is Pro or hydroxy-Pro; Xaa₅ is Trp or halo-Trp; and the C-terminus contains a carboxyl or amide group.

17. A pharmaceutical composition comprising the μ O-conotopeptide of claim 15 or a pharmaceutically acceptable salt or solvate thereof and a pharmaceutically acceptable carrier.

18. A pharmaceutical composition comprising the μ O-conotopeptide of claim 16 or a pharmaceutically acceptable salt or solvate thereof and a pharmaceutically acceptable carrier.

1/2




FIG. 1

2/2




FIG. 2

SEQUENCE LISTING

5 <110> Olivera, Baldomero M.
McIntosh, J. Michael
McCabe, R. Tyler
Garrett, James E.
Layer, Richard T.
Wagstaff, John D.
Jones, Robert M.
10 Cognetix, Inc.
University of Utah Research Foundation

15 <120> MuO-Conopeptides and Their Use as Local Anesthetics

20 <130> MuO-Conotoxins

<140>
<141>

25 <150> US 60/138,507
<151> 1999-06-10

<160> 34

30 <170> PatentIn Ver. 2.0

<210> 1
<211> 36
<212> PRT
35 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:generic
MuO-conopeptide sequence

40 <220>
<221> PEPTIDE
<222> (1)
<223> Xaa at residue 1 is des-Xaa, Pro, hydroxy-Pro
(Hyp), Arg, Lys, ornithine, homo-Lys,
homoarginine, nor-Lys, N-methyl-Lys,
N,N'-dimethyl-Lys, N,N',N''-trimethyl-Lys or any
synthetic basic amino acid

45 <220>
<221> PEPTIDE
<222> (2)
<223> Xaa at residue 2 is des-Xaa, Ala, Gly, Asp, Glu,
gamma-carboxy-glutamate (Gla), any synthetic
50 acidic amino acid, Thr, Ser, g-Thr (where g is
glycosylation), g-Ser, Trp (D or L), neo-Trp or

55 <220>
<221> PEPTIDE
<222> (2)..(4)
<223> or halo-Trp (D or L) or Xaa2 may be pyroglutamate
if Xaa at residue 1 is des-Xaa; Xaa at residue 4
is Arg, Lys, ornithine, homo-Lys, homoarginine,
nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys,

60 <220>
<221> PEPTIDE
<222> (4)
<223> N,N',N''-trimethyl-Lys, any synthetic basic amino

acid, Ser, Thr, α -Ser, α -Thr, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val

5

<220>
<221> PEPTIDE
<222> (4)..(5)
10 <223> or non-natural derivatives of the aliphatic amino acid, His, Glu, Gln, Gla, Asp, Asn or any synthetic acidic amino acid; Xaa at residue 5 is Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid,

15

<220>
<221> PEPTIDE
<222> (5)
20 <223> Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N''-trimethyl-Lys, any synthetic basic amino acid, Ala, an aliphatic amino acids bearing linear or branched

25

<220>
<221> PEPTIDE
<222> (5)
30 <223> saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Ser, Thr, Pro, Hyp, α -Ser, α -Thr, α -Hyp or any synthetic hydroxylated amino acid;

35

<220>
<221> PEPTIDE
<222> (6)
40 <223> Lys, Arg, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N''-trimethyl-Lys, any synthetic basic amino acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr,

50

<220>
<221> PEPTIDE
<222> (6)
45 <223> di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of

55

<220>
<221> PEPTIDE
<222> (6)..(7)
55 <223> of the aliphatic amino acid, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, Pro or Hyp; Xaa at residue 7 is Trp (D or L), neo-Trp, halo-Trp (D or L), Gly, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr,

60

<220>
<221> PEPTIDE
<222> (7)..(8)
60 <223> mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Glu, Gla, Gln, Asp, Asn,

any synthetic acidic amino acid; Xaa at residue 8
is Glu, Gla, Gln, Asp, Asn, any

5 <220>
<221> PEPTIDE
<222> (8)
<223> synthetic acidic amino acid, Met, norleucine
(Nle), Ala, an aliphatic amino acids bearing
10 linear or branched saturated hydrocarbon chains
such as Leu (D or L), Ile and Val or

15 <220>
<221> PEPTIDE
<222> (8)
<223> non-natural derivatives of the aliphatic amino
acid, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr,
mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr,
20 O-phospho-Tyr, nitro-Tyr, Lys, Arg, ornithine,
homo-Lys, homoarginine,

25 <220>
<221> PEPTIDE
<222> (8)..(9)
<223> nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys,
N,N',N'''-trimethyl-Lys or any synthetic basic
amino acid; Xaa at residue 9 is Leu, Phe, Tyr,
meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr,
di-halo-Tyr,

30 <220>
<221> PEPTIDE
<222> (9)..(11)
<223> O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or
35 L), neo-Trp, halo-Trp (D or L) or any synthetic
aromatic amino acid; Xaa at residue 11 is Pro,
Hyp, Gly, an aliphatic amino acids bearing linear
or

40 <220>
<221> PEPTIDE
<222> (11)..(12)
<223> branched saturated hydrocarbon chains such as Leu
(D or L), Ile and Val or non-natural derivatives
45 of the aliphatic amino acid; Xaa at residue 12 is
Thr, Ser, g-Thr, g-Ser, Ala, an aliphatic amino

50 <220>
<221> PEPTIDE
<222> (12)
<223> acids bearing linear or branched saturated
hydrocarbon chains such as Leu (D or L), Ile and
Val or non-natural derivatives of the aliphatic
amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr,
nor-Tyr,

55 <220>
<221> PEPTIDE
<222> (12)..(13)
<223> mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr,
O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp,
60 halo-Trp (D or L) or any synthetic aromatic amino
acid; Xaa at residue 13 is Pro, Hyp, Ser, Thr,
g-Hyp,

5 <220>
 <221> PEPTIDE
 <222> (13)..(14)
 <223> g-Ser, g-Thr or any hydroxylated amino acid; Xaa
 at residue 14 is an aliphatic amino acids bearing
 linear or branched saturated hydrocarbon chains
 such as Leu (D or L), Ile and Val or non-natural

10 <220>
 <221> PEPTIDE
 <222> (14)
 <223> derivatives of the aliphatic amino acid, Phe, Tyr,
 meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr,
 di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr,
 15 nitro-Tyr, Lys, Arg, ornithine, homo-Lys,
 homoarginine,

20 <220>
 <221> PEPTIDE
 <222> (14)..(15)
 <223> nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys,
 N,N',N'''-trimethyl-Lys or any synthetic basic
 amino acid; Xaa at residue 15 is Pro, Hyp, an
 25 aliphatic amino acids bearing linear or branched
 saturated

30 <220>
 <221> PEPTIDE
 <222> (15)
 <223> hydrocarbon chains such as Leu (D or L), Ile and
 Val or non-natural derivatives of the aliphatic
 amino acid, Lys, Arg, ornithine, homo-Lys,
 homoarginine, nor-Lys, N-methyl-Lys,
 35 N,N'-dimethyl-Lys,

40 <220>
 <221> PEPTIDE
 <222> (15)..(16)
 <223> N,N',N'''-trimethyl-Lys or any synthetic basic
 amino acid; Xaa at residue 16 is Gly, His, Lys,
 Arg, ornithine, homo-Lys, homoarginine, nor-Lys,
 N-methyl-Lys, N,N'-dimethyl-Lys,
 N,N',N'''-trimethyl-Lys

45 <220>
 <221> PEPTIDE
 <222> (16)..(17)
 <223> or any synthetic basic amino acid; Xaa at residue
 50 17 is des-Xaa, Ser, Thr, g-Ser, g-Thr, Val, Asn,
 Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr,
 mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr,
 O-phospho-Tyr,

55 <220>
 <221> PEPTIDE
 <222> (17)..(18)
 <223> nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or
 L) or any synthetic aromatic amino acid; Xaa at
 residue 18 is Met, Nle, Leu, Phe, Tyr, meta-Tyr,
 60 ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr,

<220>
 <221> PEPTIDE
 <222> (18)

5 <223> O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Arg, Lys, ornithine, homo-Lys, homoarginine, nor-Lys, N-methyl-Lys, N,N'-dimethyl-Lys, N,N',N''-trimethyl-Lys or any synthetic basic amino acid

10 <220>

<221> PEPTIDE

<222> (19)..(22)

10 <223> Pro, Hyp, Ser, Thr, g-Hyp, g-Ser, g-Thr, any hydroxylated amino acid, Ala, Glu, Gla, Gln, Asp, Asn, any synthetic acidic amino acid, His or Gly; Xaa at residue 22 is Gly, Asn or Gln

15 <220>

<221> PEPTIDE

<222> (23)..(24)

20 <223> Xaa at residue 23 is Leu, Trp (D or L), neo-Trp or halo-Trp (D or L); Xaa at residue 24 is des-Xaa, Leu or Trp (D or L), neo-Trp or halo-Trp (D or L)

25 <220>

<221> PEPTIDE

<222> (25)

30 <223> Xaa at residue 25 is des-Xaa or an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino

35 <220>

<221> PEPTIDE

<222> (25)..(27)

40 <223> acid; Xaa at residue 27 is des-Xaa, Gly, Met, Nle, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp,

45 <220>

<221> PEPTIDE

<222> (27)..(28)

50 <223> halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa at residue 28 is des-Xaa, Pro, Hyp, Ala, an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as

55 <220>

<221> PEPTIDE

<222> (28)

55 <223> Leu (D or L), Ile and Val or non-natural derivatives of the aliphatic amino acid, Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr,

60 <220>

<221> PEPTIDE

<222> (28)..(29)

60 <223> Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; Xaa at residue 29 is an aliphatic amino acids bearing linear or branched saturated hydrocarbon chains such as

Xaa Xaa Xaa Cys Cys Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Cys Xaa
20 25 30

5 Xaa Xaa Xaa Xaa
35

<210> 2

<211> 31

10 <212> PRT

<213> Conus magus

<220>

<221> PEPTIDE

15 <222> (3)..(27)

<223> Xaa at residue 3 may be Arg or Ser; Xaa at
residues 12, 21 and 27 5 may be Pro or
hydroxy-Pro; Xaa at residue 14 may be Ile or Leu ;
Xaa at residue 17 may be Ile or Val

20

<400> 2

Ala Cys Xaa Lys Lys Trp Glu Tyr Cys Ile Val Xaa Ile Xaa Gly Phe
1 5 10 15

25 Xaa Tyr Cys Cys Xaa Gly Leu Ile Cys Gly Xaa Phe Val Cys Val
20 25 30

<210> 3

<211> 31

<212> PRT

<213> Conus skinneri

<220>

<221> PEPTIDE

35 <222> (5)..(18)

<223> Xaa at residue 5 is Lys, N-methyl-Lys,
N,N-dimethyl-Lys or N,N,N-trimethyl-Lys; Xaa at
residue 6, 16 and 18 may be Tyr, mono-halo-Tyr,
40 di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or
nitro-Tyr.

<220>

<221> PEPTIDE

45 <222> (7)..(27)

<223> Xaa at residue 7 may be Glu or gamma-carboxy-Glu;
Xaa at residues 12 and 27 may be Pro or
hydroxy-Pro.

50

<400> 3

Ala Cys Arg Gln Xaa Xaa Xaa Phe Cys Leu Val Xaa Ile Ile Gly Xaa
1 5 10 15

55

Ile Xaa Cys Cys Ala Gly Leu Ile Cys Gly Xaa Phe Val Cys Leu
20 25 30

<210> 4

<211> 36

60 <212> PRT

<213> Conus tessulatus

<220>

<221> PEPTIDE

5 <222> (1)..(34)
<223> Xaa at residue 1 may be Glu or gamma-carboxy-Glu;
Xaa at residues 5, 8 and 34 may be Lys,
N-methyl-Lys, N,N-dimethyl-Lys or
N,N,N-trimethyl-Lys.

10 <220>
<221> PEPTIDE
<222> (13)..(35)
<223> Xaa at residues 1, 33 and 35 may be Pro or
hydroxy-Pro ; Xaa at residue 19 may be Tyr,
mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr,
O-phospho-Tyr or nitro-Tyr.

15 <400> 4
Xaa Thr Cys Leu Xaa Gln Asp Xaa Phe Cys Ile Ile Xaa Leu Ile Gly
1 5 10 15
20 Thr Leu Xaa Cys Cys Ser Gly Leu Ile Cys Gly Phe Phe Val Cys Val
20 25 30
Xaa Xaa Xaa Phe
35

25 <210> 5
<211> 32
<212> PRT
<213> Conus characteristicus
30
<220>
<221> PEPTIDE
<222> (3)..(31)
<223> Xaa at residues 3 and 31 may be Glu or
gamma-carboxy-Glu; Xaa at residues 12 and 30 may
be Pro or hydroxy-Pro; Xaa at residue 15 may be
Lys, N-methyl-Lys, N,N-dimethyl-Lys or
N,N,N-trimethyl-Lys.

40 <220>
<221> PEPTIDE
<222> (14)..(32)
<223> Xaa at residue 14 may be Trp or bromo-Trp; Xaa at
residue 32 may be Tyr, mono-halo-Tyr, di-halo-Tyr,
O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr.

45 <400> 5
Asp Cys Xaa Ala Asp Gly Ala Phe Cys Gly Ile Xaa Ile Val Xaa Asn
1 5 10 15
50 Xaa Met Cys Cys Ser Asn Leu Cys Ile Phe Ala Cys Val Xaa Xaa Xaa
20 25 30

55 <210> 6
<211> 31
<212> PRT
<213> Conus textile
60 <220>
<221> PEPTIDE
<222> (4)..(21)
<223> Xaa at residues 4 and 21 is Glu or
gamma-carboxy-Glu; Xaa at residues 6 and 8 is Trp

or halo-Trp; Xaa at residue 10 is Pro or hydroxy-Pro; Xaa at residue 18 is Tyr, mono-halo-Tyr, di-halo-Tyr,

5 <220>
<221> PEPTIDE
<222> (4)..(21)
<223> O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr

10 <400> 6
Asp Cys His Xaa Arg Xaa Asp Xaa Cys Xaa Ala Ser Ile Leu Gly Val
1 5 10 15

15 Ile Xaa Cys Cys Xaa Gly Leu Ile Cys Phe Ile Ala Phe Cys Ile
20 25 30

20 <210> 7
<211> 33
<212> PRT
<213> Conus textile

25 <220>
<221> PEPTIDE
<222> (4)..(27)
<223> Xaa at residues 4 and 7 is Glu or gamma-carboxy-Glu; Xaa at residue 5 is Lys, N-methy-Lys, N,N-dimethyl-Lys or N,N,N-trimethyl-Lys; Xaa at residue 6 is Trp or
30 halo-Trp; Xaa at residues 12, 21 and 27

35 <220>
<221> PEPTIDE
<222> (4)..(27)
<223> is Pro or hydroxy-Pro; Xaa at residue 18 is Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr

40 <400> 7
Asp Cys Gln Xaa Xaa Xaa Xaa Phe Cys Ile Val Xaa Ile Leu Gly Phe
1 5 10 15

45 Val Xaa Cys Cys Xaa Gly Leu Ile Cys Gly Xaa Phe Val Cys Val Asp
20 25 30

Ile

50 <210> 8
<211> 31
<212> PRT
<213> Conus tessulatus

55 <220>
<221> PEPTIDE
<222> (1)..(18)
<223> Xaa at residues 1 and 13 is Pro or hydroxy-Pro; Xaa at residues 6 and 18 is Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or
60 nitro-Tyr

<400> 8
Xaa Thr Cys Val Ser Xaa Asn Val Phe Cys Gly Val Xaa Leu Val Gly
1 5 10 15

10

Thr Xaa Leu Cys Cys Ser Gly Leu Val Cys Leu Val Val Cys Ile
20 25 30

5 <210> 9
<211> 27
<212> PRT
<213> Conus atlanticus

10 <220>
<221> PEPTIDE
<222> (3)..(21)
<223> Xaa at residues 3, 11, 13 and 15 is Pro or
hydroxy-Pro; Xaa at residue 12 is Lys,
15 N-methy-Lys, N,N-dimethyl-Lys or
N,N,N-trimethyl-Lys; Xaa at residue 16 is Tyr,
mono-halo-Tyr, di-halo-Tyr,

20 <220>
<221> PEPTIDE
<222> (3)..(21)
<223> O-sulpho-Tyr, O-phospho-Tyr or nitro-Tyr; Xaa at
residue 21 is Trp or halo-Trp

25 <400> 9
Cys Arg Xaa Arg Gly Met Phe Cys Gly Phe Xaa Xaa Xaa Gly Xaa Xaa
1 5 10 15

30 Cys Cys Asn Gly Xaa Cys Phe Phe Val Cys Ile
20 25

35 <210> 10
<211> 30
<212> PRT
<213> Conus tessulatus

40 <220>
<221> PEPTIDE
<222> (2)..(13)
<223> Xaa at residue 2 is Trp or halo-Trp; Xaa at residue
8 is Glu or gamma-carboxy-Glu; Xaa 13 is Pro or
hydroxy-Pro

45 <400> 10
Arg Xaa Cys Ala Leu Asp Gly Xaa Leu Cys Ile Ile Xaa Val Ile Gly
1 5 10 15

50 Ser Ile Phe Cys Cys His Gly Ile Cys Met Ile Xaa Cys Val
20 25 30

55 <210> 11
<211> 30
<212> PRT
<213> Conus arenatus

60 <220>
<221> PEPTIDE
<222> (4)..(17)
<223> Xaa at residues 4 and 12 is Pro or hydroxy-Pro;
Xaa at residues 8 and 13 is Tyr, mono-halo-Tyr,
di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or
nitro-Tyr; Xaa at residue 14 is Lys, N-methy-Lys,

5 <220>
 <221> PEPTIDE
 <222> (4)..(17)
 <223> N,N-dimethyl-Lys or N,N,N-trimethyl-Lys; Xaa at
 residue 17 is Trp or halo-Trp

10 <400> 11
 Asp Cys Arg Xaa Val Gly Gln Xaa Cys Gly Ile Xaa Xaa Xaa His Asn
 1 5 10 15

10 Xaa Arg Cys Cys Ser Gln Leu Cys Ala Ile Ile Cys Val Ser
 20 25 30

15 <210> 12
 <211> 30
 <212> PRT
 <213> Conus generalis

20 <220>
 <221> PEPTIDE
 <222> (5)..(29)
 <223> Xaa at residues 5 and 12 is Pro or hydroxy-Pro;
 Xaa at residues 7 and 17 is Tyr, mono-halo-Tyr,
25 di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr or
 nitro-Tyr; Xaa at residue 29 is Glu or
 gamma-carboxy-Glu

30 <400> 12
 Gly Cys Leu Asp Xaa Gly Xaa Phe Cys Gly Thr Xaa Phe Leu Gly Ala
 1 5 10 15

35 Xaa Cys Cys Gly Gly Ile Cys Leu Ile Val Cys Ile Xaa Thr
 20 25 30

35 <210> 13
 <211> 29
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> Description of Artificial Sequence:amplification
 primer

45 <400> 13
 caggatccat gaaactgacg tgyrtgg 29

50 <210> 14
 <211> 29
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> Description of Artificial Sequence:amplification
 primer

60 <400> 14
 atctcgagca caggtatgga tgactcagg 29

60 <210> 15
 <211> 424
 <212> DNA

<213> Conus skinneri

<220>

<221> CDS

5 <222> (1)..(264)

<400> 15

atg aaa ctg acg tgt gtg gtg atc gtt gct gtc ctg ttc ttg acc gcc 48
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu Thr Ala
 1 5 10 15

tgg aca ttc gtc atg gct gat gac ccc aga gat gga gcg gag att aga 96
 Trp Thr Phe Val Met Ala Asp Asp Pro Arg Asp Gly Ala Glu Ile Arg
 20 25 30

15 agc atg gta agg ggg gaa cct ctg tcg aag gca cgt gac gaa atg aac 144
 Ser Met Val Arg Gly Glu Pro Leu Ser Lys Ala Arg Asp Glu Met Asn
 35 40 45

20 ccc gaa gcc tct aaa ttg gag aaa agg gcg tgc cgc caa aaa tac gaa 192
 Pro Glu Ala Ser Lys Leu Glu Lys Arg Ala Cys Arg Gln Lys Tyr Glu
 50 55 60

25 ttt tgt cta gta ccg atc att gga tac ata tat tgc tgc gct ggc tta 240
 Phe Cys Leu Val Pro Ile Ile Gly Tyr Ile Tyr Cys Cys Ala Gly Leu
 65 70 75 80

30 atc tgt ggt cct ttc gtc tgc ctt tgatagtat gtcttctact gccatctgtg 294
 Ile Cys Gly Pro Phe Val Cys Leu
 85

ctaccctgg cttgatcttt gataggcggtt gttgccttc actggtttat gaaccctctg 354

35 atcatactct ctggaccctt gggggtccaa catccaaata aagcgacatc ccaaaaaaaaaa 414
 aaaaaaaaaa 424

<210> 16

40 <211> 88

<212> PRT

<213> Conus skinneri

<400> 16

45 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu Thr Ala
 1 5 10 15

Trp Thr Phe Val Met Ala Asp Asp Pro Arg Asp Gly Ala Glu Ile Arg
 20 25 30

50 Ser Met Val Arg Gly Glu Pro Leu Ser Lys Ala Arg Asp Glu Met Asn
 35 40 45

55 Pro Glu Ala Ser Lys Leu Glu Lys Arg Ala Cys Arg Gln Lys Tyr Glu
 50 55 60

Phe Cys Leu Val Pro Ile Ile Gly Tyr Ile Tyr Cys Cys Ala Gly Leu
 65 70 75 80

60 Ile Cys Gly Pro Phe Val Cys Leu
 85

<210> 17

<211> 418
 <212> DNA
 <213> Conus tessulatus

<220>
 <221> CDS
 <222> (1)..(261)

<400> 17

gga tcc atg aaa ctg acg tgc atg gtg atc gtt gtt gtg ctg ttg ttg	48
Gly Ser Met Lys Leu Thr Cys Met Val Ile Val Val Val Leu Leu Leu	
1 5 10 15	

aac gcc tgg aca ttc gtc tcc ata aat gga aag gcg aat cgt ttt tgg 96

Asn Ala Trp Thr Phe Val Ser Ile Asn Gly Lys Ala Asn Arg Phe Trp	
20 25 30	

aag gca cgt gac gaa atg aag gac tcc gaa gtt tct gaa ttg gag aaa 144

Lys Ala Arg Asp Glu Met Lys Asp Ser Glu Val Ser Glu Leu Glu Lys	
35 40 45	

agg agg aaa ccg acc tgc ctg aag cag gac aag ttt tgc ata ata ccg 192

Arg Arg Lys Pro Thr Cys Leu Lys Gln Asp Lys Phe Cys Ile Ile Pro	
50 55 60	

ctc att gga acc ctt tat tgc tgc agt ggg tta atc tgt ggg ttt ttt 240

Leu Ile Gly Thr Leu Tyr Cys Cys Ser Gly Leu Ile Cys Gly Phe Phe	
65 70 75 80	

gtc tgc gtc cca aag ccg ttc tgatgtcttc tactgccatc tgtgctaccc 291

Val Cys Val Pro Lys Pro Phe	
85	

ctggcttgat cttgattgg cgtgtgccct tcactggta tgaacccttc tgatcctact 351

gtctggacgc ctcggcgtc caacgtccaa ataaagcgac atcccaataaa aaaaaaaaaa 411

aaaaaaaaaa 418

<210> 18
 <211> 87
 <212> PRT
 <213> Conus tessulatus

<400> 18

Gly Ser Met Lys Leu Thr Cys Met Val Ile Val Val Val Leu Leu Leu	
1 5 10 15	

Asn Ala Trp Thr Phe Val Ser Ile Asn Gly Lys Ala Asn Arg Phe Trp 88

20 25 30	
----------	--

Lys Ala Arg Asp Glu Met Lys Asp Ser Glu Val Ser Glu Leu Glu Lys 88

35 40 45	
----------	--

Arg Arg Lys Pro Thr Cys Leu Lys Gln Asp Lys Phe Cys Ile Ile Pro 88

50 55 60	
----------	--

Leu Ile Gly Thr Leu Tyr Cys Cys Ser Gly Leu Ile Cys Gly Phe Phe 88

65 70 75 80	
-------------	--

Val Cys Val Pro Lys Pro Phe 88

85	
----	--

5 <210> 19
 <211> 280
 <212> DNA
 <213> *Conus characteristicus*
 10 <220>
 <221> CDS
 <222> (1)...(249)
 15 <400> 19
 atg aaa ctg acg tgc gtg atg atc gtt gct gtg ctg ttc ttg acc gcc 48
 Met Lys Leu Thr Cys Val Met Ile Val Ala Val Leu Phe Leu Thr Ala
 1 5 10 15
 20 tgg aca ttc gtc acg gct gat gac tcc att aat gca ctg gag gat ctt 96
 Trp Thr Phe Val Thr Ala Asp Asp Ser Ile Asn Ala Leu Glu Asp Leu
 25 20 25 30
 25 ttt tcg aag gca cgt gac gaa atg gaa aac ggc gaa gct tct aca ttg 144
 Phe Ser Lys Ala Arg Asp Glu Met Glu Asn Gly Glu Ala Ser Thr Leu
 35 35 40 45
 30 aac gag aga gac tgc gaa gca gat ggt gca ttt tgt ggt atc cca att 192
 Asn Glu Arg Asp Cys Glu Ala Asp Gly Ala Phe Cys Gly Ile Pro Ile
 50 55 60
 35 gtg aag aac tgg atg tgc tgc agt aac ttg tgt att ttt gcc tgc gta 240
 Val Lys Asn Trp Met Cys Cys Ser Asn Leu Cys Ile Phe Ala Cys Val
 65 70 75 80
 40 ccc gag tat taagactgcc gtgatgtctt ctcctccct c 280
 Pro Glu Tyr
 45 <210> 20
 <211> 83
 <212> PRT
 <213> *Conus characteristicus*
 50 <400> 20
 Met Lys Leu Thr Cys Val Met Ile Val Ala Val Leu Phe Leu Thr Ala 40
 1 5 10 15
 55 Trp Thr Phe Val Thr Ala Asp Asp Ser Ile Asn Ala Leu Glu Asp Leu
 20 25 30
 60 Phe Ser Lys Ala Arg Asp Glu Met Glu Asn Gly Glu Ala Ser Thr Leu
 35 40 45
 65 Asn Glu Arg Asp Cys Glu Ala Asp Gly Ala Phe Cys Gly Ile Pro Ile
 50 55 60
 70 Val Lys Asn Trp Met Cys Cys Ser Asn Leu Cys Ile Phe Ala Cys Val
 65 70 75 80
 75 Pro Glu Tyr
 80 <210> 21
 <211> 132
 <212> DNA
 <213> *Conus textile*

<220>
<221> CDS
<222> (2)..(106)

5 <400> 21
a ttg gag aaa agg gat tgc cac gaa agg tgg gat tgg tgt cca gca tca 49
Leu Glu Lys Arg Asp Cys His Glu Arg Trp Asp Trp Cys Pro Ala Ser
1 5 10 15

10 atc ctt gga gtg ata tat tgc tgc gag gga tta att tgt ttt att gcc 97
Ile Leu Gly Val Ile Tyr Cys Cys Glu Gly Leu Ile Cys Phe Ile Ala
20 25 30

15 ttc tgc att tgatagtgtat gtcttctcct cccctc 132
Phe Cys Ile
35

20 <210> 22
<211> 35
<212> PRT
<213> Conus textile

25 <400> 22
Leu Glu Lys Arg Asp Cys His Glu Arg Trp Asp Trp Cys Pro Ala Ser
1 5 10 15

Ile Leu Gly Val Ile Tyr Cys Cys Glu Gly Leu Ile Cys Phe Ile Ala
20 25 30

30 Phe Cys Ile
35

35 <210> 23
<211> 132
<212> DNA
<213> Conus textile

40 <220>
<221> CDS
<222> (2)..(112)

<400> 23
a ttg gag aaa agg gat tgc caa gag aaa tgg gag ttt tgt ata gta ccg 49
Leu Glu Lys Arg Asp Cys Gln Glu Lys Trp Glu Phe Cys Ile Val Pro
1 5 10 15

45 atc ctt gga ttt gta tat tgc tgc cct ggc tta atc tgt ggc cct ttt 97
Ile Leu Gly Phe Val Tyr Cys Cys Pro Gly Leu Ile Cys Gly Pro Phe
20 25 30

50 gtc tgc gtt gat atc tgatgtcttc tcctccatc 132
Val Cys Val Asp Ile
55 35

60 <210> 24
<211> 37
<212> PRT
<213> Conus textile

<400> 24

Leu Glu Lys Arg Asp Cys Gln Glu Lys Trp Glu Phe Cys Ile Val Pro
 1 5 10 15

5 Ile Leu Gly Phe Val Tyr Cys Cys Pro Gly Leu Ile Cys Gly Pro Phe
 20 25 30

Val Cys Val Asp Ile
 35

10 <210> 25
 <211> 288
 <212> DNA
 <213> Conus tessulatus

15 <220>
 <221> CDS
 <222> (7)..(246)

20 <400> 25
 ggatcc atg aaa ctg acg tgt gtg gtg atc gtt gtt gtg ctg ttg ttg 48
 Met Lys Leu Thr Cys Val Val Ile Val Val Leu Leu Leu
 1 5 10

25 aac gcc tgg aca ttc gtc tcc ata aat gga aag gcg aat cct ttt tgg 96
 Asn Ala Trp Thr Phe Val Ser Ile Asn Gly Lys Ala Asn Pro Phe Trp
 15 20 25 30

30 aag gca cgt gac gaa atg aag gac tcc gaa gtt tct gag ttg gag aaa 144
 Lys Ala Arg Asp Glu Met Lys Asp Ser Glu Val Ser Glu Leu Glu Lys
 35 40 45

35 agg agg aaa ccg acc tgc gtg tcg tat aac gtg ttt tgc gga gta ccg 192
 Arg Arg Lys Pro Thr Cys Val Ser Tyr Asn Val Phe Cys Gly Val Pro
 50 55 60

40 ctc gtt gga acc tac ctt tgc agt ggc tta gtc tgt ctc gta gtc 240
 Leu Val Gly Thr Tyr Leu Cys Cys Ser Gly Leu Val Cys Leu Val Val
 65 70 75

45 tgc atc tagtactgat gtcttctact cccatctgtg ctacccctcg ag 288
 Cys Ile
 80

50 <210> 26
 <211> 80
 <212> PRT
 <213> Conus tessulatus

55 <400> 26
 Met Lys Leu Thr Cys Val Val Ile Val Val Leu Leu Leu Asn Ala
 1 5 10 15

55 Trp Thr Phe Val Ser Ile Asn Gly Lys Ala Asn Pro Phe Trp Lys Ala
 20 25 30

60 Arg Asp Glu Met Lys Asp Ser Glu Val Ser Glu Leu Glu Lys Arg Arg
 35 40 45

Lys Pro Thr Cys Val Ser Tyr Asn Val Phe Cys Gly Val Pro Leu Val
 50 55 60

Gly Thr Tyr Leu Cys Cys Ser Gly Leu Val Cys Leu Val Val Cys Ile
 65 . 70 . 75 80

5 <210> 27
 <211> 287
 <212> DNA
 <213> Conus atlanticus

10 <220>
 <221> CDS
 <222> (7)..(240)

15 <400> 27
 ggatcc atg aaa ctg acg tgc gtg gtg atc gtt gct gtg ctg ttc ttg 48
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu
 1 5 10

20 acc gcc tgg aca ttc gtc acg gct gat gac tcc ata aat ggg ttg gag 96
 Thr Ala Trp Thr Phe Val Thr Ala Asp Asp Ser Ile Asn Gly Leu Glu
 15 20 25 30

25 aat ctt ttt ccg aag gca cgt cac gaa atg agg aaa ccc gaa gcc tct 144
 Asn Leu Phe Pro Lys Ala Arg His Glu Met Arg Lys Pro Glu Ala Ser
 35 40 45

30 aga tcg aga ggg agg tgc cgt cct cgt ggt atg ttc tgt ggc ttt ccg 192
 Arg Ser Arg Gly Arg Cys Arg Pro Arg Gly Met Phe Cys Gly Phe Pro
 50 55 60

35 aaa cct gga cca tac tgc tgc aat ggc tgg tgc ttt ttc gtc tgc atc 240
 Lys Pro Gly Pro Tyr Cys Cys Asn Gly Trp Cys Phe Val Cys Ile
 65 70 75

40 taaaactgcc gtgatgtgtt ctactcccat ctgtgctacc cctcgag 287

45 <210> 28
 <211> 78
 <212> PRT
 <213> Conus atlanticus

50 <400> 28
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu Thr Ala 15
 1 5 10 15

55 Trp Thr Phe Val Thr Ala Asp Asp Ser Ile Asn Gly Leu Glu Asn Leu
 20 25 30

60 Phe Pro Lys Ala Arg His Glu Met Arg Lys Pro Glu Ala Ser Arg Ser
 35 40 45

Arg Gly Arg Cys Arg Pro Arg Gly Met Phe Cys Gly Phe Pro Lys Pro
 50 55 60

65 Gly Pro Tyr Cys Cys Asn Gly Trp Cys Phe Phe Val Cys Ile
 65 70 75

70 <210> 29
 <211> 419
 <212> DNA
 <213> Conus tessulatus

<220>
 <221> CDS
 <222> (7)...(249)

5 <400> 29
 ggatcc atg aaa ctg acg tgc gtg gtg gtc gtt gct gtg ctg ttc ttg 48
 Met Lys Leu Thr Cys Val Val Val Ala Val Leu Phe Leu
 1 5 10

10 aac gcc tgg aca ttc gcc acg gct gtt gac tcc aaa cat gca ctg gcg 96
 Asn Ala Trp Thr Phe Ala Thr Ala Val Asp Ser Lys His Ala Leu Ala
 15 20 25 30

15 aaa ctt ttt atg aag gca cgt gac gaa atg tat aac ccc gat gcc act 144
 Lys Leu Phe Met Lys Ala Arg Asp Glu Met Tyr Asn Pro Asp Ala Thr
 35 40 45

20 aaa ttg gac gat aag aga tgg tgc gct tta gat ggt gaa ctt tgt atc 192
 Lys Leu Asp Asp Lys Arg Trp Cys Ala Leu Asp Gly Glu Leu Cys Ile
 50 55 60

25 ata ccg gtc att ggg tcc ata ttt tgc tgc cat ggc ata tgt atg atc 240
 Ile Pro Val Ile Gly Ser Ile Phe Cys Cys His Gly Ile Cys Met Ile
 65 70 75

30 tac tgc gtc tagttgaact gccgtgatgt cttctactcc cctctgtgct 289
 Tyr Cys Val
 80

35 accctgggtt tgatcttga ttgcctgtg cccttcactg attatgaatc cctctgatcc 349
 tactctctga agaccttttg gggccaaca tccaaataaa gcgacatccc aaaaaaaaaaa 409
 aaaaaaaaaaa 419

40 <210> 30
 <211> 81
 <212> PRT
 <213> *Conus tessulatus*

45 <400> 30
 Met Lys Leu Thr Cys Val Val Val Ala Val Leu Phe Leu Asn Ala 15
 1 5 10 15

50 Trp Thr Phe Ala Thr Ala Val Asp Ser Lys His Ala Leu Ala Lys Leu
 20 25 30

55 Phe Met Lys Ala Arg Asp Glu Met Tyr Asn Pro Asp Ala Thr Lys Leu
 35 40 45

60 Asp Asp Lys Arg Trp Cys Ala Leu Asp Gly Glu Leu Cys Ile Ile Pro
 50 55 60

65 Val Ile Gly Ser Ile Phe Cys Cys His Gly Ile Cys Met Ile Tyr Cys
 65 70 75 80

70 Val

75 <210> 31
 <211> 340
 <212> DNA

<213> Conus arenatus

<220>

<221> CDS

5 <222> (7)...(246)

<400> 31

ggatcc atg aaa ctg acg tgt gtg gtg atc gtt gct gtg ctg ttc ttg 48
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu
 10 1 5 10

acc gcc tgg aca ttc gtc acg gct gac tcc ata cgt gca ctg gag gat 96
 Thr Ala Trp Thr Phe Val Thr Ala Asp Ser Ile Arg Ala Leu Glu Asp
 15 20 25 30

15 ttt ttt gcg aag gca cgt gac gaa atg gaa aac agc gga gct tct cca 144
 Phe Phe Ala Lys Ala Arg Asp Glu Met Glu Asn Ser Gly Ala Ser Pro
 35 40 45

20 ttg aac gag aga gac tgc cga cct gta ggt caa tat tgt ggc ata ccg 192
 Leu Asn Glu Arg Asp Cys Arg Pro Val Gly Gln Tyr Cys Gly Ile Pro
 50 55 60

25 tat aag cac aac tgg cga tgc tgc agt cag ctt tgt gca att atc tgt 240
 Tyr Lys His Asn Trp Arg Cys Cys Ser Gln Leu Cys Ala Ile Ile Cys
 65 70 75

30 gtt tcc taaccctct gatcctactc tctgaagacc tccgggattc aacatccaaa 296
 Val Ser
 80

taaagcgaca tcccgatnaa aaaaaangaa aaaaaaaaaa aaaa 340

35 <210> 32
 <211> 80
 <212> PRT
 <213> Conus arenatus

40 <400> 32
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu Thr Ala
 1 5 10 15

45 Trp Thr Phe Val Thr Ala Asp Ser Ile Arg Ala Leu Glu Asp Phe Phe
 20 25 30

Ala Lys Ala Arg Asp Glu Met Glu Asn Ser Gly Ala Ser Pro Leu Asn
 35 40 45

50 Glu Arg Asp Cys Arg Pro Val Gly Gln Tyr Cys Gly Ile Pro Tyr Lys
 50 55 60

His Asn Trp Arg Cys Cys Ser Gln Leu Cys Ala Ile Ile Cys Val Ser
 65 70 75 80

55 <210> 33
 <211> 293
 <212> DNA
 60 <213> Conus generalis

<220>

<221> CDS

<222> (7)...(249)

<400> 33
 ggatcc atg aaa ctg acg tgt gtg atc gtt gct gtg cta ttc ttg 48
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu
 1 5 10

5
 acc gcc tgg aca ttc gtc acg gct gat gac acc aga tat aaa ctg gag 96
 Thr Ala Trp Thr Phe Val Thr Ala Asp Asp Thr Arg Tyr Lys Leu Glu
 15 20 25 30

10 aat cct ttt ctg aag gca cgc aac gaa ctg cag aaa cac gaa gcc tct 144
 Asn Pro Phe Leu Lys Ala Arg Asn Glu Leu Gln Lys His Glu Ala Ser
 35 40 45

15 caa ctg aac gag aga ggc tgc ctt gac cca ggt tac ttc tgt ggg acg 192
 Gln Leu Asn Glu Arg Gly Cys Leu Asp Pro Gly Tyr Phe Cys Gly Thr
 50 55 60

20 ccg ttt ctt gga gca tac tgc tgc ggt ggc att tgc ctt att gtc tgc 240
 Pro Phe Leu Gly Ala Tyr Cys Cys Gly Ile Cys Leu Ile Val Cys
 65 70 75

25 ata gaa acg taaaggcttg atgtcttcta ctcccatctg tgctaccctt cgag 293
 Ile Glu Thr
 80

30 <210> 34
 <211> 81
 <212> PRT
 <213> Conus generalis

<400> 34
 Met Lys Leu Thr Cys Val Val Ile Val Ala Val Leu Phe Leu Thr Ala 15
 1 5 10 15

35 Trp Thr Phe Val Thr Ala Asp Asp Thr Arg Tyr Lys Leu Glu Asn Pro 30
 20 25 30

40 Phe Leu Lys Ala Arg Asn Glu Leu Gln Lys His Glu Ala Ser Gln Leu
 35 40 45

Asn Glu Arg Gly Cys Leu Asp Pro Gly Tyr Phe Cys Gly Thr Pro Phe
 50 55 60

45 Leu Gly Ala Tyr Cys Cys Gly Gly Ile Cys Leu Ile Val Cys Ile Glu 80
 65 70 75 80

Thr

50

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/15779

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 38/00; C07K 14/435, 14/00; C12N 15/12
US CL : 514/2; 530/324, 325

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2; 530/324, 325

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BECKER, S. Synthesis and characterization of mu-conotoxins IIIa. European Journal of Biochem. June 1989, Vol. 185, pages 79-84, especially page 81-83.	1-18
Y	BRAGA, M.F.M. et al. Interactions between suxamethonium and non-depolarizing neuromuscular blocking drugs. British Journal of Anaesthesia. February 1994, Vol. 72, No. 2, pages 198-204, especially pages 201-203.	1-18
Y	SOSA, M.A. et al. Use of mu-conotoxin GIIIA for the study of synaptic transmission at the frog neuromuscular junction. Neuroscience Letters. July 1993, Vol. 157, No. 2, pages 235-238, entire article	1-18
Y	STEPHAN, M.M. et al. The muI Skeletal Muscle Sodium Channel: Mutation E403Q Eliminates Sensitivity to tetrodotoxin but not to mu-conotoxins GIIIA and GIIIB. Journal of Membrane Biology. January 1994, Vol. 137, No. 1, pages 1-8, entire document	1-18
X	MCINTOSH, J.M. et al. A new family of Conotoxins that block voltage-gated sodium channels. Journal of Biological Chemistry. 14 July 1995, Vol. 270, No. 28, pages 16796-16802, entire document	1-18

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 26 July 2000 (26.07.2000)	Date of mailing of the international search report 15 AUG 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703)305-3230	Authorized officer Patricia Robinson Christopher Low Telephone No. 703-308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/15779

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claim Nos.: 2,3,15 and 16 (IN-PART)
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The CRF disk that was submitted was defective.

3. Claim Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/15779

Continuation of B. FIELDS SEARCHED Item 3: STN: Biosis, Medline, Caplus
Conotoxins, mu, pain, neuro blocking,