



(86) Date de dépôt PCT/PCT Filing Date: 2012/05/24
(87) Date publication PCT/PCT Publication Date: 2012/11/29
(45) Date de délivrance/Issue Date: 2020/12/29
(85) Entrée phase nationale/National Entry: 2013/10/31
(86) N° demande PCT/PCT Application No.: EP 2012/059787
(87) N° publication PCT/PCT Publication No.: 2012/160179
(30) Priorités/Priorities: 2011/05/25 (EP11167552.6);
2011/05/26 (EP11167732.4)

(51) Cl.Int./Int.Cl. A61K 9/107 (2006.01),
A61K 31/436 (2006.01), A61K 31/573 (2006.01),
A61K 47/00 (2006.01), A61K 47/06 (2006.01),
A61K 47/24 (2006.01), A61K 9/00 (2006.01),
A61P 17/06 (2006.01), A61P 17/10 (2006.01),
A61P 17/12 (2006.01), A61P 17/14 (2006.01),
A61P 31/04 (2006.01), A61P 31/10 (2006.01)

(72) Inventeurs/Inventors:
GUNTHER, BERNHARD, DE;
THEISINGER, BASTIAN, DE;
THEISINGER, SONJA, DE;
SCHERER, DIETER, CH

(73) Propriétaire/Owner:

(54) Titre : COMPOSITION PHARMACEUTIQUE A BASE D'ALCANES SEMIFLUORE
(54) Title: TOPICAL PHARMACEUTICAL COMPOSITION BASED ON SEMIFLUORINATED ALKANES

(57) Abrégé/Abstract:

The invention provides liquid or semi-solid pharmaceutical compositions for topical administration comprising a semifluorinated alkane. The compositions are useful for the delivery of active ingredients into the deep layers of the skin or skin appendages. Various active ingredients may be incorporated, such as immunosuppressants, antiinfectives, antifungal agents, antiinflammatory agents, and retinoids.

(73) Propriétaires(suite)/Owners(continued):NOVALIQ GMBH, DE

(74) Agent: GOWLING WLG (CANADA) LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(43) International Publication Date
29 November 2012 (29.11.2012)

WIPO | PCT

(10) International Publication Number

WO 2012/160179 A3

(51) International Patent Classification:

<i>A61K 9/07</i> (2006.01)	<i>A61P 17/14</i> (2006.01)
<i>A61K 47/06</i> (2006.01)	<i>A61P 17/12</i> (2006.01)
<i>A61K 31/436</i> (2006.01)	<i>A61P 31/10</i> (2006.01)
<i>A61K 31/573</i> (2006.01)	<i>A61P 31/04</i> (2006.01)
<i>A61P 17/06</i> (2006.01)	<i>A61K 47/24</i> (2006.01)
<i>A61P 17/10</i> (2006.01)	<i>A61K 9/00</i> (2006.01)

(74) Agent: **CREMER, Karsten**; Pharma Concepts GmbH, 55, rue des Bruyères, L-1274 Howald (LU).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/EP2012/059787

(22) International Filing Date:

24 May 2012 (24.05.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

11167552.6	25 May 2011 (25.05.2011)	EP
11167732.4	26 May 2011 (26.05.2011)	EP

(71) **Applicant** (for all designated States except US): **NOVALIQ GMBH** [DE/DE]; Im Neuenheimer Feld 515, 69120 Heidelberg (DE).

(72) Inventors; and

(75) **Inventors/Applicants** (for US only): **GÜNTHER, Bernhard** [DE/DE]; Schriesheimerstrasse 19, 69221 Dossenheim (DE). **THEISINGER, Bastian** [DE/DE]; Zur Waldau 23, 68239 Mannheim (DE). **THEISINGER, Sonja** [DE/DE]; Zur Waldau 23, 68239 Mannheim (DE). **SCHERER, Dieter** [DE/CH]; Im Tschambol 18, CH-4242 Laufen (CH).

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

17 January 2013

WO 2012/160179 A3

(54) **Title:** TOPICAL PHARMACEUTICAL COMPOSITION BASED ON SEMIFLUORINATED ALKANES

(57) **Abstract:** The invention provides liquid or semi-solid pharmaceutical compositions for topical administration comprising a semifluorinated alkane. The compositions are useful for the delivery of active ingredients into the deep layers of the skin or skin appendages. Various active ingredients may be incorporated, such as immunosuppressants, antiinfectives, antifungal agents, antiinflammatory agents, and retinoids.

TOPICAL PHARMACEUTICAL COMPOSITION BASED ON SEMIFLUORINATED ALKANES

Description

BACKGROUND OF THE INVENTION

5 The present invention is in the field of pharmacotherapy. More specifically, it relates to the treatment of diseases and conditions affecting the skin and skin appendages.

The pharmacological treatment of cutaneous disorders may involve systemic or local administration of an effective medicament. Typically, whenever local (e.g. 10 epicutaneous) administration is possible and leads to the desired outcome, it is preferred over systemic therapy which is, quite plausibly, more often associated with adverse systemic effects. However, effective treatment via the topical route of administration requires that the drug substance is soluble in, and/or otherwise compatible with a suitable carrier or vehicle, and that it reaches its target tissue 15 layer(s) in an effective amount or concentration. In fact, in some cases, effective epicutaneous treatment is particularly difficult to achieve.

The skin consists of three main layers - the epidermis, the dermis and the hypodermis. The epidermis comprises the stratum corneum (the outermost superficial layer of skin, consisting of dead cells) and the vital (living) epidermis. 20 Depending on the cutaneous disease or condition, one or more layers of the skin may become affected. In some dermatological conditions, such as ichthyosis, callus, or plaque psoriasis, the stratum corneum is thicker than in the healthy state and can provide a significantly greater barrier to the penetration of a therapeutical compound, thus reducing its efficacy in the treatment of the deeper layers of the skin. 25 Moreover, it has been shown recently that with increasing age, the skin of a human also becomes more resistant to the penetration by drug substances.

The use of penetration enhancers has been proposed in order to achieve a sufficient and consistent availability of various therapeutic compounds at the site of action during the treatment of cutaneous diseases. For example, U.S. Patent No.

5,326,566 describes a composition of a pharmacological agent in combination with dibutyl adipate, or a mixture of dibutyl adipate and isopropyl myristate, which could enhance the penetration through the skin and even increase the amount absorbed into the systemic circulation, if that is desired.

5 Even though a variety of penetration enhancers have been used for enhancing the absorption of therapeutic agents into and through the skin, substantial problems may arise when the penetration enhancers are incompatible with a particular drug substance, leading to drug instability and degradation into potentially harmful degradants.

10 US 2005/0079210 A1 proposes the use of liposomes for epicutaneous administration of drugs and cosmetically useful agents. However, liposomes are difficult to manufacture cost-effectively and in a reproducible manner.

15 Other novel formulation strategies for topical medications require the occlusion of the skin after administration, which is however perceived as inconvenient by many patients

EP0670159A1 discloses stable oily emulsions comprising surfactant and semifuorinated alkane, which are proposed to be suitable for transporting oxygen to the skin or vascular system or for use as a standard for blood-gas analysis. The use of these compositions as carriers of active agent was also suggested, although no mention is made of whether these compositions are capable of effectively delivering active agents in particular to the deeper layers of the skin.

20 There clearly remains a need for pharmaceutical formulations and vehicles which allow the effective treatment of cutaneous disorders by topical administration. It is therefore an object of the present invention to provide such improved compositions which overcome one or more disadvantages of known compositions. In particular, it is an object of the invention to provide compositions of topically active drug substances which are effective and convenient. Further objects of the invention will become clear on the basis of the description of the invention below, including the examples.

SUMMARY OF THE INVENTION

The present invention provides a novel pharmaceutical composition for administration to the skin or a skin appendage. The composition comprises an effective amount of an active ingredient and a semifluorinated alkane (SFA) according to formula RFRH or RFRHRF, wherein RF is a perfluorinated hydrocarbon segment with 20 or less carbon atoms and RH is a non-fluorinated hydrocarbon segment with 3 to 20 carbon atoms. In a preferred embodiment, the SFA is selected from F4H5, F4H6, F4H8, F6H6 and F6H8.

It has been surprisingly found by the inventors that SFAs are capable of delivering active pharmaceutical ingredients effectively into the deeper layers of the skin, i.e. into the vital layers below the stratum corneum of the epidermis. Moreover, they are capable of delivering active ingredients deep into nails.

The composition may, for example, be in the form of a solution or an emulsion. They are useful for administering a wide range of topical agents indicated in a number of cutaneous diseases and conditions.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the amount of tacrolimus found in different layers of the skin (SC=stratum corneum; EP=vital epidermis; DR=dermis; AK=hypodermis) 30 min, 300 min, and 1,000 min after administration as an SFA-based formulation to human skin, relative to the administered dose. For further details, see Example 1.

Figure 2 shows the amount of tacrolimus found in different layers of the skin (SC=stratum corneum; EP=vital epidermis; DR=dermis; AK=hypodermis) 30 min, 300 min, and 1,000 min after administration of a commercial formulation (Protopic[®] 0.1%) to human skin, relative to the administered dose. For further details, see Example 1.

Figure 3 shows the penetration profiles of tacrolimus 30 min, 300 min, and 1,000 min after administration as an SFA-based formulation to human skin, respectively. For further details, see Example 1.

Figure 4 shows the penetration profiles of tacrolimus 30 min, 300 min, and 1,000 min after administration a commercial formulation (Protopic® 0.1%) to human skin, respectively. For further details, see Example 1.

Figure 5 shows the concentration of D,L- α -tocopherol found in different layers 5 of the skin (SC=stratum corneum; EP=vital epidermis; DR=dermis) 100 min after administration as SFA-based formulation (SFA), Unguentum emulsificans aquosum DAB (WHS), Basiscreme DAC (BC), Lanae alcoholum unguentum aquosum DAB (WWS). For further details, see Example 2.

Figure 6 shows the concentration of betamethasone-17,21-dipropionate in 10 different layers of the skin (SC=stratum corneum; EP=vital epidermis; DR=dermis) 100 min after administration as SFA-based formulation (SFA), Unguentum emulsificans aquosum DAB (WHS), Basiscreme DAC (BC), Lanae alcoholum unguentum aquosum DAB (WWS). For further details, see Example 3.

DETAILED DESCRIPTION OF THE INVENTION

15 The invention provides a novel pharmaceutical composition for use as a topical formulation which is to be administered to the skin or a skin appendage. The composition comprises an effective amount of an active ingredient and a semifluorinated alkane according to formula RFRH or RFRHRF, wherein RF is a perfluorinated hydrocarbon segment with 20 or less carbon atoms and RH is a non- 20 fluorinated hydrocarbon segment with 3 to 20 carbon atoms, and in particular, wherein the perfluorinated segment is linear and comprises from 4 to 12 carbon atoms, and/or the non-fluorinated segment is linear and comprises from 4 to 8 carbon atoms.

25 The invention is based on the surprising discovery that the semifluorinated alkanes (SFAs) as defined above are highly suitable carriers for topically administered drugs and capable of delivering such drugs very effectively to the layers of the skin below the stratum corneum, i.e. the viable epidermis, but also to the dermis and hypodermis. This was particularly unexpected in view of Hardung (Doctoral Thesis, Albert-Ludwigs-University of Freiburg, Germany, 2008), who

investigated the suitability of SFAs as carriers for topical medications using e.g. testosterone as a model drug, and who came to the conclusion that an SFA such as F6H8 (see below for an explanation of the terminology of SFAs) does not affect an improved penetration into the skin, interact with the stratum corneum or exhibit any

5 penetration-enhancing effect (p. 103, first full paragraph). However, in spite of such discouraging statements in the prior art, the inventors have further investigated the impact of SFAs and SFA-based carriers on the delivery of drugs into the deeper layers of the skin and nails using excised human skin (in contrast to Hardung who used pig skin), and have now surprisingly found that SFAs are indeed capable of enhancing the

10 penetration of drugs into the relevant skin layers, as will be shown in more detail below.

As used herein, a pharmaceutical composition is any composition comprising an active ingredient useful for the diagnosis, prevention, management or therapy of a disease, symptom or health condition and at least one carrier or excipient. A topical

15 formulation is a composition which is in a form suitable for topical administration. In the present invention, the composition is suitable for administration to the skin or a skin appendage of a subject, which subject may be a human or an animal. The skin may be intact skin or injured, bruised, damaged or otherwise affected skin. A skin appendage is a structure derived from the skin, and usually adjacent to it, such as a

20 hair, a sweat or sebaceous gland. Semifluorinated alkanes are linear or branched alkanes some of whose hydrogen atoms have been replaced by fluorine. In a preferred embodiment, the semifluorinated alkanes (SFA's) used in the present invention are composed of at least one non-fluorinated hydrocarbon segment and at least one perfluorinated hydrocarbon segment. Particularly useful are SFA's which

25 have one non-fluorinated hydrocarbon segment attached to one perfluorinated hydrocarbon segment, according to the general formula $F(CF_2)_n(CH_2)_mH$, or two perfluorinated hydrocarbon segments separated by one non-fluorinated hydrocarbon segment, according to the general formula $F(CF_2)_n(CH_2)_m(CF_2)_oF$.

Another nomenclature which is used herein refers to the above-mentioned

30 SFA's having two or three segments as RFRH and RFRHRF, respectively, wherein R_F designates a perfluorated hydrocarbon segment, R_H designates a non-fluorinated

segment. Alternatively, the compounds may be referred to as FnHm and FnHmFo, respectively, wherein F means a perfluorinated hydrocarbon segment, H means a non-fluorinated segment, and n, m and o is the number of carbon atoms of the respective segment. For example, F3H3 is used for perfluoropropylpropane. Moreover, this type

5 of nomenclature is usually used for compounds having linear segments. Therefore, unless otherwise indicated, it should be assumed that F3H3 means 1-perfluoropropylpropane, rather than 2-perfluoropropylpropane, 1-perfluoroisopropylpropane or 2-perfluoroisopropylpropane.

Preferably, the semifluorinated alkanes according to the general formulas

10 $F(CF_2)_n(CH_2)_mH$ and $F(CF_2)_n(CH_2)_m(CF_2)_oF$ have segment sizes ranging from 3 to 20 carbon atoms, i.e. n, m and o are independently selected in the range from 3 to 20. SFAs which are useful in the context of the present invention are also described in EP-A 965 334, EP-A 965329 and EP-A 2110126.

15 In a further embodiment, the semifluorinated alkane is a compound according to the formula RFRH, whose segments R_F and R_H are linear and each - but independently from one another - have from 3 to 20 carbon atoms. In particular, the perfluorinated segment is linear and comprises from 4 to 12 carbon atoms, and/or the non-fluorinated segment is linear and comprises from 4 to 8 carbon atoms.

20 Preferred SFA's include in particular the compounds F4H5, F4H6, F4H8, F6H4, F6H6, F6H8, and F6H10. Presently most preferred for carrying out the invention are F4H5, F4H6, F6H6 and F6H8.

25 Optionally, the composition may comprise more than one SFA. It may be useful to combine SFA's, for example, in order to achieve a particular target property such as a certain density, viscosity, or solubilising capacity for a particular active ingredient. If a mixture of SFA's is used, it is furthermore preferred that the mixture comprises at least one of F4H5, F4H6, F4H8, F6H4, F6H6, F6H8, and F6H10, and in particular one of F4H5, F4H6, F4H8, F6H6 and F6H8. In another embodiment, the mixture comprises at least two members selected from F4H5, F4H6, F4H8, F6H4, F6H6, F6H8,

30 and F6H10, and in particular at least two members selected from F4H5, F6H6, F4H8, and F6H8. Furthermore, mixtures of a semifluorinated alkane, e.g. F4H5, with a

perfluorinated compound, such as perfluoroctylbromide or perfluorodecalin, or with another oily material are also contemplated. Other oily materials would, for example, include silicone oils, triglycerides (e.g. native or synthetic medium-chain triglycerides), or compounds such as isopropyl myristate. In some preferred 5 embodiments, however, perfluorinated compounds are absent.

Liquid SFA's are chemically and physiologically inert, colourless and stable. Their typical densities range from 1.1 to 1.7 g/cm³, and their surface tension may be as low as 19 mN/m. SFA's of the RFRH type are insoluble in water but also somewhat 10 amphiphilic, with increasing lipophilicity correlating with an increasing size of the non-fluorinated segment. Again, for practising the current invention, an SFA having a density of at least 1.2 g/cm³ should be selected.

Liquid SFA's of the RFRH type are being used commercially for unfolding and reapplying a retina, for long-term tamponade as vitreous humor substitute (H. Meinert et al., European Journal of Ophthalmology, Vol. 10(3), pp. 189-197, 2000), 15 and as wash-out solutions for residual silicon oil after vitreo-retinal surgery. Experimentally, they have also been used as blood substitutes (H. Meinert et al., Biomaterials, Artificial Cells, and Immobilization Biotechnology, Vol. 21(5), pp. 583-95, 1993). These applications have established SFA's as physiologically well tolerated compounds. On the other hand, SFA's have not been used as excipients in approved 20 drug products as of today.

For allowing convenient administration to the skin or skin appendages, the composition is preferably liquid or semi-solid. As used herein, semi-solid means that a material behaves like a solid upon the exertion of low shear force, but above a certain force threshold, the so-called "yield point", it behaves like a viscous fluid. 25 Whether liquid or semi-solid, the composition may represent a single phase (i.e. a solution), or consist of two or more phases.

In a particular embodiment, for example, the composition is in the form of an emulsion. As used herein, an emulsion is a system comprising a dispersed (or inner, or emulsified, or discontinuous) liquid phase within a continuous (or outer, or 30 coherent) liquid or semisolid phase. The two phases are not miscible. In an O/W-

emulsion (also referred to as oil-in-water emulsion), a water-immiscible organic liquid phase, which does not have to be an "oil" by any specific definition, is dispersed in a water-miscible continuous phase which may or may not be substantially comprised of water itself. Within emulsions, O/W-emulsions are preferred for

5 carrying out the invention.

In another embodiment, the composition is in the form of a microemulsion. A microemulsion is a clear, thermodynamically stable, optically isotropic mixture of a lipophilic component, a hydrophilic component, and an amphiphilic component.

Typically, a microemulsion forms spontaneously when the components are combined

10 and mixed with each other, without requiring high energy input as is normally required for the formation of an "ordinary" emulsion. Microemulsions may have a colloidal lipophilic phase dispersed in a hydrophilic phase, or a hydrophilic phase colloidally dispersed in a lipophilic phase. The size of the dispersed phases is usually in the range from about 5 nm to about 400 nm, and most often below about 200 nm.

15 In one of the preferred embodiments of the invention, the particle size is from about 5 nm to about 100 nm. In terms of its rheological properties, the microemulsion may be in the form of a liquid or a gel, i.e. in liquid or semisolid form. In a preferred embodiment, the microemulsion is in liquid form.

For microemulsions according to the invention, it is preferred to use an oil

20 phase comprising from about 5 to about 95 wt.-% of SFA, and more preferably from about 10 to about 80 wt.-% of SFA, the remainder of the oil phase being another oil such as isopropylmyristate. With respect to the incorporation of aqueous components, surfactants and optionally cosurfactants, commonly known constituents of microemulsions may be used to formulate the composition.

25 The composition, whether formulated as a solution, microemulsion or conventional emulsion, may contain any further pharmaceutically acceptable ingredients as required or useful in view of the intended application. In a particular embodiment, an additional solvent or co-solvent may be incorporated, for example in order to achieve a higher solubility for a specific active ingredient, to modify the

30 viscosity or stability of the formulation, or to further enhance the cutaneous penetration of the active ingredient. Such co-solvent should of course be selected in

type, quality and amount such as to maintain the physiological tolerability of the formulation. Potentially suitable co-solvents include ethanol, acetone, ethyl acetate, isopropyl alcohol, glycerol, propylene glycol, pentylene glycol, polyethylene glycol, liquid paraffin, triglyceride oils, hydrofluorocarbons such as HFA 134a and/or HFA 227, and liquid mono- or diglycerides. Among these, ethanol and pentylene glycol are among the particularly preferred solvents. It has been found that relatively small amounts of ethanol may substantially modify the solubilising properties of SFAs and allow the incorporation of a number of active ingredients at higher concentrations. For example, the composition may comprise up to about 10 wt.-% of ethanol, or up to about 5 wt.-% of ethanol. Even at a low concentration of up to about 3 wt.-%, ethanol may be a very useful additive in the composition, depending on the specific drug substance to be delivered. Pentylene glycol is also very useful as co-solvent in general and, in particular, it has been found that its incorporation is advantageous in the formulation of microemulsions.

Moreover, the composition may comprise one or more stabilisers, surfactants (in particular if the composition is in the form of an emulsion or microemulsion), cosurfactants (in particular if it is in the form of a microemulsion), colouring agents, antioxidants (for example, α -tocopherol), thickeners (viscosity-increasing agents such as bentonite), and fragrances.

If a surfactant is present, it may, for example, be selected from physiologically acceptable phospholipids (such as phosphatidylcholines); nonionic surfactants such as fatty alcohols, pegylated glycerides, pegylated fatty acids, pegylated fatty alcohols, pegylated sorbitan fatty acid esters, and poloxamers; anionic surfactants such as sodium lauryl sulphate, docosate sodium, and sodium deoxycholate.

The composition may be prepared by commonly known techniques for the manufacture of pharmaceutical solutions, emulsions, and microemulsions. Depending on their consistency and specific use, they may be presented in bottles, spray bottles, or tubes.

The active ingredient incorporated within the composition may, in principle, be selected from drug substances which are useful for the prevention, management, or therapy of a cutaneous disease or condition.

In one of the specific embodiments, the active ingredient is poorly water-soluble. In particular, its water solubility is not more than about 1 mg/mL. In other preferred embodiments, the water solubility is not higher than about 0.1 mg/mL, or not more than about 10 µg/mL, respectively. The invention is particularly useful for delivering such active ingredients because it allows the administration of effective doses in relatively small volumes, which is at least partly due to the surprisingly high solubilisation capacity of semifluorinated alkanes for many poorly water-soluble drug substances.

For the avoidance of doubt, it is affirmed that the invention is not restricted to poorly water-soluble drug substances. It has been found that certain bioactive agents have substantial aqueous solubility, but can also be advantageously formulated in SFA-based carriers.

The active ingredient may be selected from drug substances useful for the prevention and in particular for the management and therapy of cutaneous diseases and conditions, such as psoriasis, dermatitis, erythema, acne, actinic keratosis, verrucae, and rosacea. The invention appears generally useful for delivering drug substances of various therapeutic categories to the deep layers of the skin, i.e. to the vital epidermis, the dermis, and/or to the hypodermis which may also be understood as being part of the integumentary system. Topical administration of the compositions to the skin may result in the delivery of the drug substance to the stratum corneum, as well as penetration through the stratum corneum and delivery of the active drug substance to these deeper skin layers. Cutaneous diseases or conditions may affect the stratum corneum, as well as one or more of the deeper skin layers. The composition may thus be used for the prevention or treatment of diseases or conditions that affect also the vital epidermis, the dermis and/or the hypodermis. In one of the specific embodiments, the composition is used in the therapy of psoriasis, and incorporates an active ingredient which may be used to control psoriasis or a symptom thereof. Such active ingredient may, for example, be an

immunosuppressant drug, such as a macrolide immunosuppressant. Examples of potentially suitable macrolide immunosuppressants include tacrolimus, sirolimus, everolimus, tacrolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A; or retinoids such as retinol, retinal, tretinoin, isotretinoin, alitretinoin, etretinate, acitretin, tazarotene, bexarotene and adapalene. Tacrolimus is one of the particularly preferred drugs in this group. It has been found by the inventors that tacrolimus can be very effectively (i.e. faster and deeper than using conventional formulations) delivered to the deep and viable layers of the skin using the composition of the invention (see Example 1).

10 As illustrated in Example 1, the composition of the invention may be used to effectively deliver an active ingredient such as a macrolide immunosuppressant, as for example tacrolimus, to the deeper layers of the skin. For example, at least about 5% of the topically administered amount of drug may be delivered to the vital epidermis within only about 30 minutes. The composition may also be used to deliver 15 at least 5% of the topically administered drug to the dermis within about 30 minutes, or even at least about 10% of the administered dose. Also in terms of delivery to the hypodermis, the composition may be used to deliver at least 5% of the topically administered amount of drug to this layer within the same period of time, or even 10% or more.

20 Taking into account the typical thickness of the respective skin layers, the composition may be used to deliver within about 30 minutes at least about 15% of the topically administered drug dose to a depth of at least about 10 μm . In other embodiments, at least about 20% of the dose is delivered to a depth of at least about 10 μm . The composition may also be used to deliver at least about 10% of the 25 topically administered drug to a depth of 100 μm or more within about 30 minutes; or at least about 15% or at least about 20% of the dose, respectively.

Another therapeutic indication in which it is of great importance that the therapeutic agent penetrates the deeper layers of the affected skin is cutaneous mycosis, such as athlete's foot, candidal vulvovaginitis, and other fungal infections of 30 the integument. in the context, the invention may be used by preparing and administering an SFA-based composition comprising a suitable topically active

antifungal agent, in particular an azole such as clotrimazole, ketoconazole, or miconazole; or an allylamine, such as terbinafine, naftifine, or butenafine.

In a further embodiment, the disease to be treated is acne, such as acne vulgaris or other forms of acneiform eruptions. In this case, the active ingredient may, for 5 example, be selected from retinoids, such as retinol, retinal, tretinoin, isotretinoin, alitretinoin, etretinate, acitretin, tazarotene, bexarotene and adapalene; antiinfective agents such as azithromycin, erythromycin, clindamycin, or minocycline; or spironolactone.

According to another embodiment, the invention is used in the treatment of 10 actinic keratosis, which is a premalignant condition of the skin characterised by the presence of thick, scaly, or crusty patches. In this case, the active ingredient may be selected from NSAIDs such as diclofenac, ketoprofene and flurbiprofene, but also 5-fluorouracil and imiquimod.

Moreover, the invention is suitable for the treatment of viral infections of the 15 skin, such as herpes simplex, herpes zoster, and varicella. For this indication, an antiviral compound is incorporated in the SFA-based composition, such as acyclovir, penciclovir, or docosanol.

In a further embodiment, the composition of the invention comprises a local 20 anaesthetic agent such as benzocaine, butamben, dibucaine, lidocaine, oxybuprocaine, pramoxine, proparacaine, proxymetacaine, and tetracaine. Such composition may be used to relieve pain or itching caused by conditions such as sunburn or other minor burns, insect bites or stings, or in preparation of minor surgical procedures.

A large number of cutaneous diseases and conditions involve symptoms related 25 to inflammatory responses, such as most forms of dermatitis, including atopic dermatitis, seborrhoeic dermatitis, allergic dermatitis, contact dermatitis and the like, but also certain forms of pruritus, psoriasis, allergies, nummular eczema, xerotic eczema, lichen sclerosis, lichen planus, discoid lupus, chapped feet, lichen simplex chronicus, poison ivy reactions, alopecia areata, nummular eczema etc. Often these conditions require therapy or symptomatic control with topical corticosteroids,

which can also be formulated and administered advantageously according to the invention. Suitable corticosteroid which may be incorporated in an SFA-based composition include amcinonide, betamethasone dipropionate, clobetasol propionate, desonide, desoximetasone, diflorasone diacetate, fluocinolone acetonide, 5 fluocinonide, flurandrenolide, fluticasone propionate, halcinonide, halobetasol propionate, hydrocortisone, hydrocortisone butyrate, hydrocortisone valerate, mometasone furoate, prednicarbate, and triamcinolone acetonide.

The invention is further illustrated by the following examples.

Examples

10 Example 1

Tacrolimus was dissolved in a liquid carrier comprising F6H8 and 3.8 wt.-% ethanol to obtain a clear solution having a tacrolimus concentration of 1 mg/mL.

Excised human skin was obtained from a breast reduction surgical centre. Skin samples were washed with isotonic sodium chloride solution. The subcutaneous 15 adipose layers were removed. Circular pieces of 20 mm in diameter (3.14 cm^2) were punched out, supported by filter gauze, and mounted in FRANZ diffusion cells in such a way that the dermal side was in contact with the acceptor fluid (PBS pH 7.4), which was continuously stirred and kept at 32 °C. Loss through evaporation was avoided by means of a glass lid.

20 On each skin sample (epidermal side), an amount of 20 µL of the tacrolimus solution was placed and spread out evenly. After a testing time of 30 min, 300 min, or 1,000 min, respectively, the remaining tacrolimus solution on the samples was collected and the skin samples were removed. From each specimen, three small punch biopsies of 6 mm diameter (0.2827 cm^2) were obtained and frozen. At -40 °C, 25 the punch biopsies were cryomicrotomed as follows:

- 1 slice of 10 µm thickness, corresponding to the stratum corneum
- 4 slices of 20 µm each, corresponding to the vital epidermis
- 5 x 5 slices of 40 µm each, corresponding to 5 layers of dermis

All slices and the remaining stumps were extracted, and the amount of tacrolimus was determined by HPLC-MS. Moreover, the acceptor fluid and the fluid removed from the stratum corneum after each test were also analysed for tacrolimus. Each test was performed in triplicate, using skin from three different donors.

5 For comparison, a similar series of experiments using a commercial tacrolimus ointment (Protopic® 0.1 %, Astellas Pharma, 20 mg per skin sample and FRANZ cell) was conducted.

In result, it was found that the penetration of tacrolimus into the relevant skin layers was clearly superior in the case of the solution according to the invention. For 10 example, already after 30 minutes approx. 9% of the administered dose was found in the vital epidermis, and approx. 12% was found in the layers of the dermis (see fig. 1). For comparison, the commercial formulation delivered only about 2.5% of the dose to the vital epidermis and only about 5% to the dermis during the same period of time (see fig. 2). After 1,000 minutes, the amount in the dermis was about 30% of the dose 15 in the case of the solution of the invention (see fig. 1), but only about 12.5% in the case of the commercial preparation (see fig. 2). At all times, the penetration depth was much higher for the composition of the invention (see fig. 3) than for the commercial product (see fig. 4).

Example 2

20 A solution of radiolabeled D,L- α -tocopherol in F6H8 having a concentration of 10 mg/mL was prepared and used for a series of skin penetration experiments in analogy to Example 1, using excised human skin, with a few differences to the procedure of Example 1 as described below.

The amount of the formulation applied to each skin sample was 6 mg. The test 25 period was 100 min. After the testing period, the stratum corneum of each sample was removed by the film stripping method (20 strips). Slices of the vital epidermis and the dermis were however obtained by microtomy at -40 °C.

For comparison, three pharmacopoeial ointment formulations incorporating the same concentration of D,L- α -tocopherol were prepared and tested: "Unguentum

"emulsificans aquosum" (DAB), "Basiscreme" (DAC), and "Lanae alcoholum unguentum aquosum" (DAB).

In result, it was surprising to find that the SFA-based formulation resulted in a much higher concentration of tocopherol in the vital epidermis than the three 5 conventional formulations (see fig. 5).

Example 3

The series of experiments of Example 2 were repeated in analogy, but using radiolabeled betamethasone-17,21-dipropionate as active ingredient at a concentration of 1 mg/mL instead of tocopherol. Another difference was that the SFA-10 based formulation of the drug contained 2.5 wt.-% ethanol.

Here it was found that the drug concentration in the skin layers was highest in the case of the SFA-based formulation of the invention, with a clear difference between the tested formulations with respect to the stratum corneum and the vital epidermis, but somewhat less differentiation in the dermis (see fig. 6).

15 Example 4

An SFA-based microemulsion with tacrolimus was prepared. Basically, 10 mg of tacrolimus were dissolved in 1.5 mg of Tween® 80 and 2 g of Brij® 30. Then 2 g of 1,2-pentandiole were added, followed by 0.3 g of F6H8 and 0.7 g of isopropyl myristate. After gentle stirring, 4 g of water were added and the mixture was homogenised until 20 it had a transparent appearance.

Claims

1. A use of a topical pharmaceutical composition comprising an active ingredient and a semifluorinated alkane according to formula

RFRH

wherein RF is a linear perfluorinated hydrocarbon segment with 4 to 12 carbon atoms, and RH is a linear alkyl group with 4 to 8 carbon atoms for the prevention or treatment of a disease or condition affecting the skin or a skin appendage:

i) wherein:

- a) the skin is affected by psoriasis, and wherein the active ingredient is an immunosuppressant selected from the group consisting of: tacrolimus, sirolimus, everolimus, tacrolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A; or
- b) the skin is affected by acne, and wherein the active ingredient is an anti-infective agent selected from the group consisting of: azithromycin, erythromycin, clindamycin, and minocycline,

or

ii) wherein:

- a) the skin is affected by pain or itching and the active ingredient is a local anaesthetic agent; or
- b) the skin is affected by a cutaneous disease or condition involving symptoms related to inflammatory responses and the active ingredient is a corticosteroid,

or

iii) wherein:

- a) the disease affecting the skin is acne and the active ingredient is an anti-infective selected from azithromycin, erythromycin, clindamycin, and minocycline; or
- b) the disease affecting the skin is psoriasis and the active ingredient is a macrolide immunosuppressant selected from the group consisting of: tacrolimus, sirolimus, everolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A,

or

iv) wherein:

- a) the disease affecting the skin is cutaneous mycosis, or candidal vulvovaginitis, and the active ingredient is an anti-fungal agent; or

- b) the disease affecting the skin is a viral infection, and the active ingredient is an antiviral agent; or
- c) the disease affecting the skin is actinic keratosis and the active ingredient is a NSAID; or 5-fluorouracil or imiquimod; or
- d) the disease affecting the skin is acne or psoriasis and the active ingredient is a retinoid.

2. Use of an active ingredient and a semifluorinated alkane according to formula

RFRH

wherein RF is a linear perfluorinated hydrocarbon segment with 4 to 12 carbon atoms, and RH is a linear alkyl group with 4 to 8 carbon atoms, in the manufacture of a topical pharmaceutical composition for the prevention or treatment of a disease or condition affecting the skin or a skin appendage:

- i) wherein:
 - a) the skin is affected by psoriasis, and wherein the active ingredient is an immunosuppressant selected from the group consisting of: tacrolimus, sirolimus, everolimus, tacrolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A; or
 - b) the skin is affected by acne, and wherein the active ingredient is an anti-infective agent selected from the group consisting of: azithromycin, erythromycin, clindamycin, and minocycline,

or

- ii) wherein:
 - a) the skin is affected by pain or itching and the active ingredient is a local anaesthetic agent; or
 - b) the skin is affected by a cutaneous disease or condition involving symptoms related to inflammatory responses and the active ingredient is a corticosteroid,

or

- iii) wherein:
 - a) the disease affecting the skin is acne and the active ingredient is an anti-infective selected from azithromycin, erythromycin, clindamycin, and minocycline; or
 - b) the disease affecting the skin is psoriasis and the active ingredient is a macrolide immunosuppressant selected from the group consisting of: tacrolimus, sirolimus, everolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A,

or

- iv) wherein:
 - a) the disease affecting the skin is cutaneous mycosis, or candidal vulvovaginitis, and the active ingredient is an anti-fungal agent; or
 - b) the disease affecting the skin is a viral infection, and the active ingredient is an antiviral agent; or
 - c) the disease affecting the skin is actinic keratosis and the active ingredient is a NSAID; or 5-fluorouracil or imiquimod; or
 - d) the disease affecting the skin is acne or psoriasis and the active ingredient is a retinoid.

3. A topical pharmaceutical composition for use in the prevention or treatment of a disease or condition affecting the skin or a skin appendage, comprising an active ingredient and a semifluorinated alkane according to formula

RFRH

wherein RF is a linear perfluorinated hydrocarbon segment with 4 to 12 carbon atoms, and RH is a linear alkyl group with 4 to 8 carbon atoms;

- i) wherein:
 - a) the skin is affected by psoriasis, and wherein the active ingredient is an immunosuppressant selected from the group consisting of: tacrolimus, sirolimus, everolimus, tacrolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A; or
 - b) the skin is affected by acne, and wherein the active ingredient is an anti-infective agent selected from the group consisting of: azithromycin, erythromycin, clindamycin, and minocycline,

or

- ii) wherein:
 - (a) the skin is affected by pain or itching and the active ingredient is a local anaesthetic agent; or
 - (b) the skin is affected by a cutaneous disease or condition involving symptoms related to inflammatory responses and the active ingredient is a corticosteroid,

or

- iii) wherein:
 - a) the disease affecting the skin is acne and the active ingredient is an anti-infective selected from azithromycin, erythromycin, clindamycin, and minocycline; or
 - b) the disease affecting the skin is psoriasis and the active ingredient is a macrolide immunosuppressant selected from the group consisting of: tacrolimus, sirolimus,

everolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A,

or

iv) wherein

- a) the disease affecting the skin is cutaneous mycosis, or candidal vulvovaginitis, and the active ingredient is an anti-fungal agent; or
- b) the disease affecting the skin is a viral infection, and the active ingredient is an antiviral agent; or
- c) the disease affecting the skin is actinic keratosis and the active ingredient is a NSAID; or 5-fluorouracil or imiquimod; or
- d) the disease affecting the skin is acne or psoriasis and the active ingredient is a retinoid.

4. Use of a topical pharmaceutical composition formulated as a solution in the prevention or treatment of a disease or condition affecting the layers of the skin below the stratum corneum, comprising an effective amount of an active ingredient and a semifluorinated alkane according to formula

RFRH

wherein RF is a linear perfluorinated hydrocarbon segment with 4 to 12 carbon atoms, and RH is a linear alkyl group with 4 to 8 carbon atoms: wherein upon use of the pharmaceutical composition the active ingredient penetrates through the stratum corneum and is delivered to the layers of the skin below the stratum corneum.

5. Use of an of active ingredient and a semifluorinated alkane according to formula

RFRH

wherein RF is a linear perfluorinated hydrocarbon segment with 4 to 12 carbon atoms, and RH is a linear alkyl group with 4 to 8 carbon atoms, in the manufacture of a topical pharmaceutical composition for the prevention or treatment of a disease or condition affecting the skin or a skin appendage, wherein upon use of the pharmaceutical composition the active ingredient penetrates through the stratum corneum and is delivered to the layers of the skin below the stratum corneum.

6. A topical pharmaceutical composition formulated as a solution for use in the prevention or treatment of a disease or condition affecting the layers of the skin below the stratum corneum, comprising an effective amount of an active ingredient and a semifluorinated alkane according to formula

RFRH

wherein RF is a linear perfluorinated hydrocarbon segment with 4 to 12 carbon atoms, and RH is a linear alkyl group with 4 to 8 carbon atoms; wherein upon use of the

pharmaceutical composition the active ingredient penetrates through the stratum corneum and is delivered to the layers of the skin below the stratum corneum.

7. The use of claim 4 or 5 or the topical pharmaceutical composition for use according to claim 6,

wherein the skin is affected by psoriasis, and wherein the active ingredient is an immunosuppressant selected from the group consisting of: tacrolimus, sirolimus, everolimus, tacrolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A; or

wherein the skin is affected by acne, and wherein the active ingredient is an anti-infective agent selected from the group consisting of: azithromycin, erythromycin, clindamycin, and minocycline; or

wherein the skin is affected by pain or itching and the active ingredient is a local anaesthetic agent; or

wherein the skin is affected by a cutaneous disease or condition involving symptoms related to inflammatory responses and the active ingredient is a corticosteroid; or

wherein the disease affecting the skin is acne and the active ingredient is an anti-infective selected from azithromycin, erythromycin, clindamycin, and minocycline; or

wherein the disease affecting the skin is psoriasis and the active ingredient is a macrolide immunosuppressant selected from the group consisting of: tacrolimus, sirolimus, everolimus, pimecrolimus, ridaforolimus, temsirolimus, zotarolimus, and ciclosporin A; or

wherein the disease affecting the skin is cutaneous mycosis, or candidal vulvovaginitis, and the active ingredient is an anti-fungal agent; or

wherein the disease affecting the skin is a viral infection, and the active ingredient is an antiviral agent; or

wherein the disease affecting the skin is actinic keratosis and the active ingredient is a NSAID; or 5-fluorouracil or imiquimod; or

wherein the disease affecting the skin is acne or psoriasis and the active ingredient is a retinoid.

8. The use according to any one of claims 1, 2, 4 or 5 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 wherein the semifluorinated alkane is selected from F4H5, F4H6, F4H8, F6H6 and F6H8, wherein F means a perfluorinated hydrocarbon segment, H means a non-fluorinated segment, and the number is the number of carbon atoms of the respective segment.

9. The use according to any one of claims 1, 2, 4, 5 or 7, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 7, wherein the composition is in the form of a liquid or semisolid solution, emulsion, or microemulsion or the topical composition for use according to claim 6, wherein wherein the composition is formulated in the form of a liquid solution.

10. The use according to any one of claims 1, 2, 4, 5, or 7 to 9, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 8, wherein the composition comprises a physiologically acceptable cosolvent.
11. The use or the topical pharmaceutical composition for use according to claim 10, wherein the physiologically acceptable cosolvent is selected from the group consisting of: ethanol, acetone, ethyl acetate, isopropyl alcohol, glycerol, propylene glycol, pentylene glycol, polyethylene glycol, liquid paraffin, triglyceride oils, hydrofluorocarbons, and liquid mono- or diglycerides.
12. The use or the topical pharmaceutical composition for use according to claim 11, wherein the hydrofluorocarbons are selected from one or more of HFA 134a and HFA 227.
13. The use according to any one of claims 1, 2, 4, 5 or 8 to 12, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 12, wherein an effective amount of the active ingredient is present in the composition.
14. The use according to any one of claims 1, 2, 4, 5 or 7 to 13, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 12, wherein the disease or condition affects one or more of: the vital epidermis, dermis, and hypodermis of the skin.
15. The use according to any one of claims 1, 2, 4, 5 or 7 to 14 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 14, wherein the active ingredient is delivered to one or more of: the vital epidermis, dermis, and hypodermis of the skin.
16. The use according to any one of claims 1, 2, 4, 5 or 7 to 15 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 15, wherein the skin is affected by pain or itching, and wherein the local anaesthetic agent is selected from the group consisting of: benzocaine, butaben, dibucaine, lidocaine, oxybuprocaine, pramoxine, proparacaine, proxymetacaine, and tetracaine.
17. The use or topical pharmaceutical composition for use according to claim 16, wherein the pain or itching is caused by sunburn, minor burns, insect bites or stings.
18. The use or topical pharmaceutical composition for use according to claim 16, wherein the topical pharmaceutical composition comprises the local anesthetic agent as an active ingredient and the topical pharmaceutical composition is used in preparation of minor surgical procedures to prevent or treat the pain or itching of the skin.
19. The use according to any one of claims 1, 2, 4, 5 or 7 to 15 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 15, wherein the cutaneous disease or condition involving symptoms relating to inflammatory responses affecting the skin is selected from the group consisting of: dermatitis, atopic dermatitis, seborrheic dermatitis, allergic dermatitis, contact dermatitis, pruritis, psoriasis, allergies, nummular eczema, xerotic eczema, lichen sclerosis, lichen planus, discoid lupus, chapped feet, lichen simplex chronicus, poison ivy reactions, and alopecia areata.
20. The use according to any one of claims 1, 2, 4, 5 or 7 to 15 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 15, wherein the active

ingredient is a corticosteroid selected from the group consisting of: amcinonide, betamethasone dipropionate, clobetasol propionate, desonide, desoximetasone, diflorasone diacetate, fluocinolone acetonide, fluocinonide, flurandrenolide, fluticasone propionate, halcinonide, halobetasol propionate, hydrocortisone, hydrocortisone butyrate, hydrocortisone valerate, mometasone furoate, prednicarbate, and triamcinolone acetonide.

21. The use according to any one of claims 1, 2, 4, 5 or 7 to 15 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 15, wherein the topical pharmaceutical composition is a topical corticosteroid composition.
22. The use according to any one of claims 1, 2, 4, 5 or 7 to 15 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 15, wherein the disease affecting the skin is cutaneous mycosis, or candidal vulvovaginitis, and the active ingredient is an anti-fungal agent selected from azoles; and an allylamine.
23. The use or the topical pharmaceutical composition for use according to claim 22, wherein the cutaneous mycosis is athlete's foot.
24. The use or the topical pharmaceutical composition for use according to claim 22 or 23, wherein: (i) the azole is selected from the group consisting of clotrimazole, ketoconazole and miconazole and (ii) the allylamine is selected from the group consisting of terbinafine, naftifine and butenafine.
25. The use according to any one of claims 1, 2, 4, 5 or 7 to 15 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 15, wherein the disease affecting the skin is a viral infection, and the active ingredient is an antiviral compound selected from acyclovir, penciclovir and docosanol.
26. The use or the topical pharmaceutical composition for use according to claim 25, wherein the viral infection is selected from the group consisting of herpes simplex, herpes zoster, and varicella.
27. The use according to any one of claims 1, 2, 4, 5 or 7 to 15 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 15, wherein the disease affecting the skin is actinic keratosis and the active ingredient is a NSAID selected from diclofenac, ketoprofene and flurbiprofene.
28. The use according to any one of claims 1, 2, or 7 to 15 or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 15, wherein the disease affecting the skin is acne or psoriasis and the active ingredient is a retinoid selected from the group consisting of retinal, tretinoin, isotretinoin, alitretinoin, etretinate, acitretin, tazarotene, bexarotene and adapalene.
29. The use according to any one of claims 1, 2, 4, 5 or 7 to 28, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 28, wherein the use is for delivering
 - i. at least 5% of a total amount of the active ingredient in a dose to the vital epidermis within about 30 minutes, or

- ii. at least 5% of a total amount of the active ingredient in a dose to the dermis within about 30 minutes; or
- iii. at least 10% of a total amount of the active ingredient in a dose to the dermis within about 30 minutes; or
- iv. at least 5% of a total amount of the active ingredient in a dose to the hypodermis within about 30 minutes; or
- v. at least 10% of a total amount of the active ingredient in a dose to the hypodermis within about 30 minutes.

30. The use according to any one of claims 1, 2, 4, 5 or 7 to 28, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 28, for delivering at least 15% of the total amount of the active ingredient in the dose to a depth of at least 10 µm within about 30 minutes.

31. The use according to any one of claims 1, 2, 4, 5 or 7 to 28, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 28, for delivering at least 20% of the total amount of the active ingredient in the dose to a depth of at least 10 µm within about 30 minutes.

32. The use according to any one of claims 1, 2, 4, 5 or 7 to 28, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 28, for delivering at least 10% or at least 15% or at least 20% of the total amount of the active ingredient in the dose to a depth of at least 100 µm within about 30 minutes.

33. The use according to claim 4 or 5 or the topical pharmaceutical composition for use according to claim 6, wherein the disease or condition is selected from the group consisting of: psoriasis, dermatitis, erythema, acne, actinic keratosis, verrucae, and rosacea.

34. The use according to any one of claims 4, 5 or 7 to 32, or the topical pharmaceutical composition for use according to any one of claims 6 to 32, wherein the active ingredient has a water solubility of not more than 1 mg/mL.

35. The use according to any one of claims 1, 2, 4, 5 or 7 to 34, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 34, comprising up to 10 wt.-% of ethanol.

36. The use according to any one of claims 1, 2, 4, 5 or 7 to 35, or the topical pharmaceutical composition for use according to any one of claims 3 or 6 to 35, wherein the use is in a human.

37. The use according to any one of claims 4 or 5, or the topical pharmaceutical composition for use according to claim 6, wherein:
 - a. the disease or condition is psoriasis and wherein the active ingredient is selected from immunosuppressants or retinoids; or
 - b. the disease or condition is acne and wherein the active ingredient is selected from retinoids or antiinfective agents or; or
 - c. the disease or condition is a fungal infection and wherein the active ingredient is an antifungal agent; or
 - d. the disease or condition is a viral infection and wherein the active ingredient is an antiviral compound; or
 - e. the disease or condition is involving symptoms related to inflammatory responses and wherein the active ingredient is a corticosteroid; or
 - f. the disease or condition is related to pain or itching caused by conditions; or
 - g. the disease or condition is itching or pain of the skin and the topical pharmaceutical composition is used in preparation of minor surgical procedures to prevent or treat the itching or pain of the skin and wherein the active ingredient is a local anaesthetic.
38. The use or topical pharmaceutical composition for use of claim 37, wherein the disease or condition is related to pain or itching caused by conditions selected from the group consisting of: sunburn, minor burns other than a sunburn, insect bites and insect stings.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6