



(22) Date de dépôt/Filing Date: 1991/07/16

(41) Mise à la disp. pub./Open to Public Insp.: 1992/01/19

(45) Date de délivrance/Issue Date: 2002/10/08

(30) Priorité/Priority: 1990/07/18 (9015825.4) GB

(51) Cl.Int.⁵/Int.Cl.⁵ C12N 15/62, C12P 21/00, C07K 19/00

(72) Inventeurs/Inventors:

Heim, Jutta, CH;
Seebboth, Peter, DE;
Takabayashi, Kenji, CH

(73) Propriétaire/Owner:

NOVARTIS AG, CH

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : TRAITEMENT IN VITRO DE PROTEINES DE FUSION

(54) Title: VITRO PROCESSING OF FUSION PROTEINS

(57) **Abrégé/Abstract:**

There is provided a novel method for the production of a biologically active protein comprising treating a fusion protein consisting of 1, one or multiple successive protein segment(s) each consisting of said biologically active protein the C-terminal amino acid of which is joined to a linker polypeptide sequence L and 2. a polypeptide tag joined to the C-terminal amino acid of said successive protein segment(s), or consisting of 1. a polypeptide tag joined to the N-terminal amino acid of 2. multiple successive protein segments each consisting of a linker polypeptide sequence L and said biologically active protein, with soluble yeast endoprotease yscF and with soluble yeast carboxypeptidase ysc α , and isolating said biologically active protein.



In vitro processing of fusion proteinsAbstract of the disclosure

There is provided a novel method for the production of a biologically active protein comprising treating a fusion protein consisting of 1. one or multiple successive protein segment(s) each consisting of said biologically active protein the C-terminal amino acid of which is joined to a linker polypeptide sequence L and 2. a polypeptide tag joined to the C-terminal amino acid of said successive protein segment(s), or consisting of 1. a polypeptide tag joined to the N-terminal amino acid of 2., multiple successive protein segments each consisting of a linker polypeptide sequence L and said biologically active protein, with soluble yeast endoprotease yscF and with soluble yeast carboxypeptidase ysc α , and isolating said biologically active protein.

4-18154/AIn vitro processing of fusion proteins

The present invention relates to a method for the production of mature proteins by in vitro processing of fusion proteins, and to means required for the preparation of such fusion proteins.

The production of pharmaceutically applicable or enzymatically active proteins is a key area in the rapidly developing biotechnology industry. Since the beginning of the era of recombinant DNA technology a great number of valuable heterologous proteins have been expressed in prokaryotic and eukaryotic host cells which had been transformed with suitable expression vectors containing DNA sequences coding for said proteins. In many cases, however, great difficulties are encountered in producing sufficient amounts of pure products. It is frequently observed, especially in the case of low molecular weight proteins, that the product is impure due to the presence of varying amounts of degraded, in particular C-terminal degraded, byproducts lowering the yield and rendering the purification a difficult task. In other cases the overall yield in the desired protein is unsatisfactory and cannot significantly be raised by modification in the process (e.g. choice of particular vectors and host strains, condition of cultivation). Furthermore, protein purification is not a trivial task. Frequently, the primary isolation mixture contains only an utmost minor amount of the desired protein while host-specific proteins are dominating. Separation of these host-specific proteins while maintaining the biological activity and structural entirety of the desired protein may cause severe problems.

One known approach to overcome the difficulties mentioned is to express the desired protein as a fusion protein, i.e. to the C-terminus or especially N-terminus of the desired protein is bound a stabilizing and/or protective polypeptide. This polypeptide is chosen depending on the problems to be solved (degradation, purification etc.) and the host used for the expression of the fusion protein. Such fusion proteins have proved effective in preventing desired proteins from degradation and in facilitating the purification procedure. There are described numerous polypeptides which have been used as fusion partners in fusion proteins, see for example H.M. Sassenfeld, Trends in Biotechnology 8, 88-93 (1990). Exemplary polypeptides include β -galactosidase, protein A and chloramphenicol

acetyltransferase. As in most cases and for obvious reasons (antigenicity etc.) the fusion protein has no practical utility as such it is required to remove the fusion partner from the desired protein after expression and purification (alternatively, the fusion protein can be another heterologous protein and separation leads to two desired proteins). This is generally done by means of a linker sequence linking the desired protein to the fusion partner and containing a cleavage site which can selectively be cleaved by chemical or enzymatic means. Such cleavage sites include, for example, a methionyl radical which is susceptible to the attack of cyanogen bromide, or a polypeptide chain including the tetrapeptidyl radical Asp-Asp-Asp-Lys which is cleaved by enterokinase after Lys. It is obvious that such amino acid subsequences must not occur at the surface of the desired proteins. As the convenient removal of the fusion partner still is the most significant problem to be solved there is a need for alternative means which allow the specific and gentle freeing of the desired protein from the fusion partner. It is an object of the invention to provide such means.

The protease yscF (or KEX2 encoded endoprotease) and the peptidase ysc α (or KEX1 encoded carboxypeptidase) have been identified to be responsible for the maturation of the mating α -factor precursor in yeast to yield the mature α -factor pheromone. Protease yscF is a membrane-bound endoprotease which is active in the neutral pH range and is completely dependent on Ca²⁺ ions. It has, near the carboxyterminus, a hydrophobic region which has been identified to be responsible for membrane binding. Removal of this membrane-binding domain yields a soluble yscF derivative which still retains its enzymatic activity and specificity, viz. cleavage at the C-terminal side of a pair of basic amino acids, such as Lys-Arg or Arg-Arg [cf. R.S. Fuller et al., Proc. Natl. Acad. Sci. USA 86, 1434-1438 (1989)]. On the other hand, protease ysc α is a membrane-bound carboxyexopeptidase which is highly active towards C-terminal basic amino acids (Arg, Lys) at pH between 6 and 7.5. Like yscF, ysc α contains a hydrophobic membrane-binding segment in the vicinity of the C-terminus. Removal of this membrane-binding domain leads to a soluble ysc α derivative with retained enzymatic activity and specificity [A. Cooper and H. Bussey, Mol. Cell. Biol. 9, 2706-2714 (1989)]. It has now been found that proteases yscF and ysc α can advantageously be used in combination for the preparation of mature proteins by in vitro processing of suitably tailored fusion proteins.

Accordingly, the present invention concerns a method for the production of a biologically active protein comprising treating a fusion protein consisting of

1. one or multiple successive protein segment(s) each consisting of said biologically active protein the C-terminal amino acid of which is joined to a linker polypeptide sequence L the N-terminal first and second and, in the case of multiple successive protein segments, also the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg, and
2. a polypeptide tag joined to the C-terminal amino acid of said successive protein segment(s),

or consisting of

1. a polypeptide tag joined to the N-terminal amino acid of
2. multiple successive protein segments each consisting of a linker polypeptide sequence L the N-terminal first and second as well as the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg and the ultimate basic amino acid of said linker polypeptide sequence L being joined to said biologically active protein,

with soluble yeast endoprotease yscF and with soluble yeast carboxypeptidase ysc α , and isolating said biologically active protein.

The fusion protein may be represented by the formula

$$\begin{array}{ll} (P-L)_m-T & \text{(I) or} \\ T-(L-P)_n & \text{(II),} \end{array}$$

in which P is the biologically active protein, L is a linker polypeptide sequence as defined above, T is a polypeptide tag, m is an integer from 1 to 10 and n is an integer from 2 to 10.

The biologically active protein may be any protein of biological interest and of prokaryotic or especially eukaryotic, in particular higher eukaryotic such as mammalian (including animal and human), origin and is, for example, an enzyme which can be used, for example, for the production of nutrients and for performing enzymatic reactions in chemistry or molecular biology, or a protein which is useful and valuable for the treatment of human and animal diseases or for the prevention thereof, for example a hormone,

polypeptide with immunomodulatory, anti-viral and anti-tumor properties, an antibody, viral antigen, blood clotting factor, a fibrinolytic agent, a growth regulation factor, furthermore a foodstuff and the like.

Example of such proteins are e.g. hormones such as secretin, thymosin, relaxin, calcitonin, luteinizing hormone, parathyroid hormone, adrenocorticotropin, melanocyte-stimulating hormone, β -lipotropin, urogastrone, insulin, growth factors, such as epidermal growth factor (EGF), insulin-like growth factor (IGF), e.g. IGF-I and IGF-II, mast cell growth factor, nerve growth factor, glia derived nerve cell growth factor, platelet derived growth factor (PDGF), or transforming growth factor (TGF), such as TGF β , growth hormones, such as human or bovine growth hormones, interleukin, such as interleukin-1 or -2, human macrophage migration inhibitory factor (MIF), interferons, such as human α -interferon, for example interferon- α A, α B, α D or α F, β -interferon, γ -interferon or a hybrid interferon, for example an α A- α D- or an α B- α D-hybrid interferon, especially the hybrid interferon BDBB, proteinase inhibitors such as α_1 -antitrypsin, SLPI and the like, hepatitis virus antigens, such as hepatitis B virus surface or core antigen or hepatitis A virus antigen, or hepatitis nonA-nonB antigen, plasminogen activators, such as tissue plasminogen activator or urokinase, hybrid plasminogen activators, such as K₂tuPA, tick anticoagulant peptide (TAP), tumour necrosis factor, somatostatin, renin, immunoglobulins, such as the light and/or heavy chains of immunoglobulin D, E or G, or human-mouse hybrid immunoglobulins, immunoglobulin binding factors, such as immunoglobulin E binding factor, human calcitonin-related peptide, blood clotting factors, such as factor IX or VIIIc, platelet factor 4, erythropoietin, eglin, such as eglin C, desulfatohirudin, such as desulfatohirudin variant HV1, HV2 or PA, corticostatin, echistatin, cystatins, human superoxide dismutase, viral thymidin kinase, β -lactamase or glucose isomerase. Preferred genes are those coding for a human α -interferon e.g. interferon α B, or hybrid interferon, particularly hybrid interferon BDBB (see EP 205,404), human tissue plasminogen activator (t-PA), human single chain urokinase-type plasminogen activator (scu-PA), hybrid plasminogen activator K₂tuPA (see EP 277,313), transforming growth factor β , human calcitonin, insulin-like growth factor I and II and desulfatohirudin, e.g. variant HV1. Proteins containing a pair of basic amino acids, such as Arg-Arg, Lys-Arg, Lys-Lys and Arg-Lys, exposed on the protein surface and therefore amenable to proteolytic cleavage, are not suited for the process according to the invention and will have to be mutated such that one of the consecutive basic amino acids is replaced by another non-basic amino acid without affecting the biological activity. Proteins having a C-terminal basic amino acid which is essential for biological activity cannot be produced

by the process according to the invention (because this basic amino acid is removed by soluble ysc α) unless an additional non-basic protective amino acid is added to the C-terminus of such proteins.

The linker polypeptide sequence L comprises 2 to about 20, especially 2 to 12, amino acid residues and contains one or multiple, especially 1 to 5, pairs of basic amino acids, such as Arg-Arg, Lys-Arg, Lys-Lys or Arg-Lys, provided that in compounds of the formula I ($m > 1$) and II the N-terminal first and second amino acids as well as the C-terminal penultimate and ultimate amino acids represent such a pair of basic amino acids while in compounds of the formula I ($m = 1$) it is sufficient that solely the N-terminal first and second amino acids represent such a pair of basic amino acids. The choice of the amino acid residues linking the individual pairs of basic amino acids and/or joining the pair(s) of basic amino acid residues to the polypeptide tag (cf. compounds of the formula I with $m = 1$) is not crucial. For example, any of the genetically encoded neutral amino acids may be chosen for that purpose. The most simple linker polypeptide sequence L is a dipeptidyl radical selected from Arg-Arg, Lys-Arg, Lys-Lys and Arg-Lys.

Since the extension of a small or medium-sized biologically active protein has proved to have a favourable influence on stability the polypeptide tag T may, in principle, be any synthetic polypeptide imaginable or any naturally occurring polypeptide or part thereof. Preferably, the polypeptide tag T consists of about 10 to about 1000, in particular 30 to 300, amino acid residues and represents a full-length polypeptide which is well-expressed in the host used for expression of the fusion protein (see below) or a part thereof. If it is the main object to facilitate the purification it is preferable to use a polypeptide tag which is susceptible for affinity chromatography (the polypeptide tag is, for example, recognized by an available monoclonal or polyclonal antibody or is bound by a specific material such as cellulose) or ion exchange chromatography (the polypeptide tag comprises a large number of acidic or basic amino acid residues). Examples of such polypeptide tags include, for example, yeast acid phosphatase PH05, yeast invertase or yeast carboxypeptidase Y especially when yeast is used as the host, polymerase of MS2 phage, β -galactosidase or E. coli acid phosphatase especially when E. coli is used as the host, neomycin, phosphotransferase, dehydrofolate reductase, an immunoglobulin or a lectin especially when mammalian host cells are used, furthermore desulfatohirudin, eglin C, chymosin, interleukin I, the Exg protein of Cellulomonas fimi, and the like, it also being possible to use fragments of these polypeptides.

Preferably, m is an integer from 1 to 5 and n is an integer from 2 to 5.

Soluble yeast endoprotease yscF and yeast carboxypeptidase ysc α are muteins of yscF and ysc α in which the hydrophobic membrane binding sites have been deleted. The amino acid sequence of the 814-residue protease yscF is known from K. Mizuno et al. [Biochem. Biophys. Res. Commun. 156, 246-254 (1988)]. The membrane binding site is located in the region Tyr⁶⁷⁹ to Met⁶⁹⁹. In the soluble yscF endoproteases according to the invention the membrane binding site has selectively been removed hence the C-terminus starting with, for example, amino acid 700 (Lys) is still present, or the whole C-terminus including the membrane binding site, i.e. 136 to approximately 200 amino acids from the C-terminus, has been removed. Such soluble yscF deletion muteins are described, for example, in EP 327,377 or in R.S. Fuller et al. (supra). The amino acid sequence of the 729-residue peptidase ysc α is likewise known [A. Dmochowska et al., Cell 50, 573-584 (1987)]. The membrane binding site is located in the region Ala⁶¹⁹ to Tyr⁶³⁷. As above, in the soluble ysc α carboxypeptidases according to the invention the membrane binding site has selectively been removed hence leaving the C-terminus starting with, for example, amino acid 638 (Asp) intact, or the whole C-terminus including the membrane binding site, i.e. 93 to approximately 110 amino acids from the C-terminus, has been removed. One such soluble ysc α carboxypeptidase mutein has been described by A. Cooper and H. Bussey (supra). The preferred soluble yscF carboxypeptidase according to the invention has the sequence depicted in the sequence listing under SEQ ID No. 1 while the preferred soluble ysc α carboxypeptidase has the sequence depicted in the sequence listing under SEQ ID No. 2.

The digestion of the fusion proteins with soluble yscF and soluble ysc α is performed using conditions under which yscF and ysc α are known to work best, i.e. in a buffered solution at pH from about 6.0 to about 7.5, preferably at about 7.0, in a temperature range of from about 25°C to about 37°C and for about 1 to 4 hours, preferably until the digestion is complete as judged by HPLC control. As yscF is strongly dependent on Ca²⁺ ions, a calcium salt, such as calcium chloride, is added to the digestion mixture. The molar ratios of fusion protein : soluble yscF : soluble ysc α is in the range 1:1:1 to 10⁴:1:1. The molar concentration of Ca²⁺ ions is in the range of from 0.1 mM to 10 mM. Optionally, a low concentration (< 1 %) of a non-ionic detergent, such as Triton X-100, may be added to the mixture as yscF is known to be activated thereby. The fusion protein may be treated with a digestive mixture containing both soluble yscF and soluble ysc α , or the fusion protein may first be treated with soluble yscF and, when the digestion has sufficiently

proceeded or is complete, thereupon with soluble ysc α , preferably in situ, i.e. without isolating the product of the first digestion step.

The biologically active protein can be isolated from the digestion mixture using conventional means. Suitable purification steps include, for example, desalination, chromatographic processes such as ion exchange chromatography, gel filtration chromatography, partition chromatography, HPLC, reversed phase HPLC, gel electrophoresis, carrier-free electrophoresis, affinity chromatography such as affinity chromatography with monoclonal antibodies coupled to an insoluble matrix, and the like, or any combination thereof.

In case the desired protein does not assume the correct three-dimensional structure due, for example, to an incorrect formation of disulfide bonds (if cysteine residues are present), it may be necessary to solubilize it under conditions which are suitable in maintaining it in its denatured form and subsequently to refold it with the concomitant coupling of disulfide bonds (if cysteine residues are present). Appropriate methods are known for a large number of proteins. Otherwise, it is required to adapt methods known in the art to the specific problems encountered.

As mentioned above some soluble yscF and ysc α deletion muteins are known from the literature. Further deletion muteins according to the invention can be prepared using methods known in the art, for example by preparing a corresponding DNA coding for said mutein, inserting it in a suitable vector DNA under the control of an expression control sequence, transforming suitable host microorganisms with the expression vector formed, culturing the transformed host microorganism in a suitable culture medium and isolating the produced mutein. The DNA coding for any of said muteins can be produced for example, by taking a plasmid containing the DNA coding for yscF (KEX2) or ysc α (KEX1) and 1. digesting it with a restriction enzyme which cleaves within or 3' of the DNA region coding for the membrane binding site (for example, EcoRI, BstXI or NarI in the case of KEX2 and HgiAI, TaqII or ClaI in the case of KEX1), digesting the cleaved DNA with a suitable endonuclease, for example Bal31, such that said DNA region is removed and recircularizing the linearized plasmid by blunt end ligation or the like, or 2. choosing or creating (for example by site-directed mutagenesis) one restriction site 5' to and one restriction site 3' to the DNA region coding for the membrane binding site (for example PvuII and NarI or EcoRI in the case of KEX2, XhoI, StuI or XcaI in the case of KEX1; the 3' restriction site may also be located within the plasmid DNA adjacent to the

translation stop signal of the KEX2 or KEX1 gene), digesting the plasmid with two restriction enzymes recognizing said restricting sites and recircularizing the linearized plasmid by blunt end ligation or the like, or 3. deleting the DNA region coding for the membrane binding site by using loop-out mutagenesis, or 4. totally deleting the C-terminus by digesting with PvuII in the case of KEX2 and with XhoI, StuI or SciI in the case of KEX1, respectively, and recircularizing the linearized plasmid by blunt end ligation or the like. As the DNA sequences of KEX2 and KEX1 are known (K. Mizumo et al., A. Dmochowska et al., supra) a suitable mutagenic oligonucleotide can easily be devised and used to delete said DNA region applying the M13 cloning system. Care must be taken that the mutated KEX2 or KEX1 genes include a translation stop signal (TAA, TAG or TGA). If necessary, such a stop signal can be introduced at the desired place via a synthetic linker DNA or it may be provided by the adjacent vector DNA. Accordingly, the mutated KEX2 and KEX1 genes may include at their 3' ends codons derived from the vector DNA which code for a few (such as 2 to 20) additional amino acids at the C-termini of the soluble yscF and ysc α muteins. All of these methods make use of conventional techniques.

Preparation of the fusion protein

The fusion proteins according to the invention can be prepared by recombinant DNA technique comprising culturing a transformed host under conditions which allow expression thereof and isolating the fusion protein. More specifically, the desired compounds are manufactured by

- a) providing an expression vector comprising an expression cassette containing a DNA sequence coding for said fusion protein,
- b) transferring the expression vector into a recipient host,
- c) culturing the transformed host under conditions which allow expression of the fusion protein, and
- d) isolating the fusion protein.

The steps involved in the preparation of the fusion proteins by recombinant DNA technique will be discussed in more detail hereinbelow.

Expression vectors

The invention relates to expression vectors comprising an expression cassette containing a DNA sequence coding for a fusion protein consisting of 1. one or multiple successive protein segment(s) each consisting of a biologically active protein the C-terminal amino acid of which is joined to a linker polypeptide sequence L the N-terminal first and second and, in the case of multiple successive protein segments, also the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg, and 2. a polypeptide tag joined to the C-terminal amino acid of said successive protein segment(s), or consisting of 1. a polypeptide tag joined to the N-terminal amino acid of 2. multiple successive protein segments each consisting of a linker polypeptide sequence L the N-terminal first and second as well as the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg and the ultimate basic amino acid of said linker polypeptide sequence L being joined to said biologically active protein, and a method for the preparation thereof.

The expression cassette may be represented by the formula

$$\text{Pr-S-(D}_P\text{-D}_L\text{-D}_T\text{)}_m\text{-T} \quad (\text{III})$$

wherein Pr is the expression control sequence, S is a bond or represents a DNA sequence encoding a signal peptide, D_P represents a DNA sequence coding for a biologically active protein, D_L is a DNA sequence coding for the linker polypeptide L wherein the first and second codons adjacent to the 3' end of D_P and, if m is different from 1, also the penultimate and ultimate codons adjacent to the 5' end of D_T code for basic amino acids selected from Lys and Arg, D_T represents a DNA sequence coding for a polypeptide tag and T represents a DNA sequence containing transcription termination signals, wherein S, D_P, D_L and D_T are in the same reading frame, and m is an integer from 1 to 10, or by the formula

$$\text{Pr-S-(D}_T\text{-D}_L\text{-D}_P\text{)}_n\text{-T} \quad (\text{IV})$$

wherein Pr, S, D_T, D_L, D_P and T have the meanings given above, and wherein S, D_T, D_L and D_P are in the same reading frame, and n is an integer from 2 to 10.

Constructs in which S represents a DNA sequence encoding a signal peptide are preferred.

The vector is selected depending on the host cells envisaged for transformation. In principle, all vectors which replicate and express the gene according to the invention in the chosen host are suitable. Examples of suitable hosts are prokaryotes and eukaryotes, which are devoid of or poor in restriction enzymes or modification enzymes and which are devoid of yscF or yscF-related activity, such as bacteria, for example Escherichia coli or Bacillus subtilis, yeasts, for example Saccharomyces cerevisiae, especially kex2 mutants thereof, and furthermore mammalian cells, in particular established human or animal cell lines, e.g. myeloma cells, human embryonic lung fibroblasts L-132, mice LTK cells, human malignant melanoma Bowes cells, HeLa cells, SV-40 virus transformed kidney cells of African green monkey COS-7 or chinese hamster ovary (CHO) cells and variants thereof. Chinese hamster ovary cells and strains of Escherichia coli and Saccharomyces cerevisiae are preferred as the host microorganism.

Vectors for use in yeast and E. coli.

Examples of vectors that are suitable for the expression of the fusion protein gene in an E. coli strain are bacteriophages, for example derivatives of the bacteriophage λ , or plasmids, such as the plasmid colE1 and its derivatives, for example pMB9, pSF2124, pBR317 or pBR322. Suitable vectors contain a complete replicon and a marker gene, which renders possible the selection and identification of the microorganisms transformed by the expression plasmids by means of a phenotype feature. Suitable marker genes impart to the microorganism, for example, resistance to heavy metals, antibiotics such as ampicillin or tetracyclin, and the like.

Several expression control sequences Pr can be used for regulating the expression cassette in E. coli. Especially promoters of strongly expressed genes are used. Suitable promoters are the lac, tac, trp and lpp promoters, furthermore the phage λ N or the phage λ pL promoter, and others. In the present invention, the preferred promoter for use in E. coli is the λ pL, trp and the lac promoter.

Vectors suitable for replication and expression in S. cerevisiae contain a yeast-replication origin and a selective genetic marker for yeast. Hybrid vectors that contain a yeast replication origin, for example the chromosomal autonomously replicating segment (ARS), are retained extrachromosomally within the yeast cell after transformation and are replicated autonomously during mitosis. Also, hybrid vectors that contain sequences

homologous to the yeast 2 μ plasmid DNA or that contain ARS and a sequence of a chromosomal centromere, for example CEN4, can be used. Suitable marker genes for yeast are especially those that impart antibiotic resistance to the host or, in the case of auxotrophic yeast mutants, genes that complement the host lesions. Corresponding genes impart, for example, resistance to the antibiotic cycloheximide or provide for prototrophy in an auxotrophic yeast mutant, for example the URA3, LEU2, HIS3 or the TRP1 gene.

Preferably, yeast hybrid vectors furthermore contain a replication origin and a marker gene for a bacterial host, especially E. coli, so that the construction and the cloning of the hybrid vectors and their precursors can be carried out in E. coli. Expression control sequences Pr suitable for expression in yeast are, for example, those of the CYC1, GAL1/10, or PH05 gene, and also promoters involved in glycolysis, for example the PGK or the GAP (including 5' truncated GAP) promoter, furthermore the α -factor promoter and hybrid promoters, such as hybrid PH05-GAP promoters.

The DNA sequence encoding a signal peptide S ("signal sequence") is derived from a gene of the microbial host coding for a polypeptide which is ordinarily secreted. When E. coli is used as the host microorganism the ompA, lpp, maltose binding protein, λ receptor, acid phosphatase or β -lactamase signal sequence may be chosen. Suitable signal sequences S for use in yeast are, for example, the signal and prepro sequences of the yeast invertase, pheromone peptidase (KEX1), "killer toxin" and repressible acid phosphatase (PH05) genes, the α -factor leader and the glucoamylase signal sequence from Aspergillus awamori. Alternatively, fused signal sequences may be constructed by ligating part of the signal sequence (if present) of the gene naturally linked to the promoter used (for example PH05), with part of the signal sequence of the biologically active protein (if present). Those combinations are favoured which allow a precise cleavage between the signal peptide and the fusion protein amino acid sequence.

The DNA sequence containing transcription termination signals T is preferably the 3' flanking sequence of a gene derived from the selected microbial host which contains proper signals for transcription termination. Suitable 3' flanking sequences are, for example, those of the gene naturally linked to the promoter used.

Vectors for use in mammalian cells

Vectors for replication and expression in mammalian cells are frequently provided with

DNA from viral origin, e.g. from simian virus 40 (SV 40), Rous sarcoma virus (RSV), adenovirus 2, bovine papilloma virus (BPV), papovavirus BK mutant (BKV), or mouse or human cytomegalovirus (MCMV and HCMV, respectively).

Expression control sequences P_r which are suitable for use in mammalian cells include, inter alia, the early and late promoters of SV40, the major late promoter of adenovirus, the promoter of the murine metallothionein gene and the enhancer-promoter region of the mouse or human cytomegalovirus major immediate-early gene, the human immunoglobulin enhancer-promoter region, the human α -globin promoter optionally combined with the SV40 enhancer and promoters derived from the heat shock genes. Signal sequences S for use in mammalian cells are, for example, the signal sequences derived from influenza haemagglutinin, human t-PA and prepro-insulin.

Suitable marker genes for mammalian cells are, for example, the neo and ble genes from transposon Tn5 which confer resistance to the antibiotic G418 and to bleomycin-type antibiotics, respectively, the E.coli gene for hygromycin-B resistance, the dihydrofolate reductase gene (dhfr) from mammalian cells or E.coli which changes the phenotype of DHFR⁻ cells into DHFR⁺ cells and/or confer resistance to methotrexate, and the thymidine kinase gene of herpes simplex virus which makes TK⁻ cells phenotypically TK⁺ cells.

Preferably, the hybrid vectors for mammalian cells contain the 3' untranslated region of a mammalian gene containing signals for proper transcription termination and polyadenylation (T), such as, for example, the 3' flanking region of the β -globin gene. Advantageously, the regions flanking the polypeptide coding region include one or more native introns having the appropriate splicing signals at their termini. Such additions are deemed necessary as cDNAs and prokaryotic DNAs such as the above selection genes, generally lack such transcription and processing signals.

Preferably, such vectors contain an origin of replication and an antibiotic resistance gene for propagation in E.coli. A mammalian origin of replication may be provided either by including in the construction of the vector a eukaryotic origin, such as derived from SV40 or from another viral source, or may be provided by the host cell chromosome upon integration of the vector into the host cell chromosome.

The DNA sequences D_p and D_T coding for the biologically active protein and the polypeptide tag are known or, if not, can be deduced from known amino acid sequences and

produced synthetically applying methods known per se. The DNA sequence D_L coding for the linker polypeptide L is preferably a synthetic DNA sequence containing pair(s) of basic amino acids as specified above and is prepared synthetically using conventional synthetic methods.

In the expression cassette according to the invention the coding sequences S, D_P , D_L and D_T are joined together in one uninterrupted reading frame starting with an ATG at the 5' end and terminating with a translation stop signal (TAA, TAG or TGA) at the 3' end. The coding sequences $S-(D_P-D_L-D_T)_m$ and $S-(D_T-D_L-D_P)_n$ are operably linked to the expression control sequence Pr.

The expression vectors according to the invention are prepared by methods known in the art applying conventional ligation techniques, for example by linking the expression cassette (prepared previously) as such or the components of the expression cassette successively in the predetermined order to the vector DNA. The components of the expression vectors are linked through common restriction sites and/or by means of synthetic linker molecules and/or by blunt end ligation.

Transformed hosts

Another aspect of the present invention involves host cells transformed with an expression vector comprising an expression cassette containing a DNA sequence coding for a fusion protein consisting of 1. one or multiple successive protein segment(s) each consisting of a biologically active protein the C-terminal amino acid of which is joined to a linker polypeptide sequence L the N-terminal first and second and, in the case of multiple successive protein segments, also the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg, and 2. a polypeptide tag joined to the C-terminal amino acid of said successive protein segment(s), or consisting of 1. a polypeptide tag joined to the N-terminal amino acid of 2. multiple successive protein segments each consisting of a linker polypeptide sequence L the N-terminal first and second as well as the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg and the ultimate basic amino acid of said linker polypeptide sequence L being joined to said biologically active protein, and a method for the preparation thereof.

Examples of suitable hosts including prokaryotic and eukaryotic hosts are those specified

above. The method for the preparation of transformed host cells comprises transforming host cells with the above expression vector.

The transformation of the eukaryotic host cells is accomplished by methods known in the art. For example, the transformation of yeast with the hybrid vectors may be accomplished according to the method described by Hinnen et al [Proc. Natl. Acad. Sci. USA 75, 1919(1978)]. This method can be divided into three steps:

- (1) Removal of the yeast cell wall or parts thereof.
- (2) Treatment of the "naked" yeast cells (spheroplasts) with the expression vector in the presence of PEG (polyethyleneglycol) and Ca^{2+} ions.
- (3) Regeneration of the cell wall and selection of the transformed cells in a solid layer of agar.

The introduction of expression vectors into mammalian cells is done by transfection in the presence of helper compounds, e.g. diethylaminoethyl-dextran, dimethyl sulfoxide, glycerol, polyethylene glycol or the like, or as co-precipitates of vector DNA and calcium phosphate. Further suitable methods include direct microinjection of vector DNA into the cell nucleus and electroporation, i.e. introduction of DNA by a short electric pulse increasing the permeability of cell membranes. The subsequent selection of transfected cells can be done using a selection marker which is either covalently integrated into the expression vector or added as a separate entity. The selection markers include genes which confer resistance to antibiotics or genes which complement a genetic lesion of the host cell (supra).

The transformation of the bacterial host strains with the expression vectors according to the invention is carried out, for example, in the manner described in the literature for B. subtilis [Anagnostopoulos et al., J. Bacteriol. 81, 741 (1961)] and E. coli [M. Mandel et al., J. Mol. Biol. 53, 159 (1970)]. The isolation of the transformed host cells is effected advantageously from a selective nutrient medium to which there has been added, for example, the biocide against which the marker gene contained in the expression plasmid imparts resistance. If, for example, the hybrid vectors contain the amp^R gene, ampicillin is accordingly added to the nutrient medium. Cells that do not contain the hybrid vector are destroyed in such a medium.

Cultivation of transformed host cells

The invention concerns furthermore a method for the production of a fusion protein consisting of 1. one or multiple successive protein segment(s) each consisting of a biologically active protein the C-terminal amino acid of which is joined to a linker polypeptide sequence L the N-terminal first and second and, in the case of multiple successive protein segments, also the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg, and 2. a polypeptide tag joined to the C-terminal amino acid of said successive protein segment(s) or consisting of 1. a polypeptide tag joined to the N-terminal amino acid of 2. multiple successive protein segments each consisting of a linker polypeptide sequence L the N-terminal first and second as well as the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg and the ultimate basic amino acid of said linker polypeptide sequence L being joined to said biologically active protein, comprising culturing under appropriate nutrient conditions a transformed host cells containing an expression vector comprising a DNA sequence coding for said fusion protein and isolating said fusion protein.

The transformed host cells are cultured by methods known in the art in a liquid medium containing assimilable sources of carbon, nitrogen and inorganic salts. Various sources of carbon can be used for culture of the transformed *E. coli* and yeast cells according to the invention. Examples of preferred sources of carbon are assimilable carbohydrates, such as glucose, maltose, mannitol or lactose, or an acetate, which can be used either by itself or in suitable mixtures. Examples of suitable sources of nitrogen are amino acids, such as casaminoacids, peptides and proteins and their degradation products, such as tryptone, peptone or meat extracts, yeast extracts, malt extract and also ammonium salts, for example ammonium chloride, sulfate or nitrate, which can be used either by themselves or in suitable mixtures. Inorganic salts which can also be used are, for example, sulfates, chlorides, phosphates and carbonates of sodium, potassium, magnesium and calcium. The medium furthermore contains, for example, growth-promoting substances, such as trace elements, for example iron, zinc, manganese and the like, and preferably substances which exert a selection pressure and prevent the growth of cells which have lost the expression plasmid. Thus, for example, if a yeast strain which is auxotrophic in, for example, an essential amino acid, is used as the host microorganism, the plasmid preferably contains a

gene coding for an enzyme which complements the host defect. Cultivation of the yeast strain is performed in a minimal medium deficient in said amino acid.

Culturing is effected by processes which are known in the art. The culture conditions, such as temperature, pH value of the medium and fermentation time, are chosen such that a maximum titre of the fusion proteins of the invention is obtained. Thus, the yeast strain is preferably cultured under aerobic conditions by submerged culture with shaking or stirring at a temperature of about 20 to 40°C, preferably about 30°C, and a pH value of 5 to 8, preferably at about pH 7, for about 4 to 30 hours, preferably until maximum yields of the proteins of the invention are reached. Mammalian cells are grown under tissue culture conditions using commercially available media optionally supplemented with growth-promoting substances and/or mammal sera. The cells are grown either attached to a solid support, e.g. a microcarrier or porous glass fibres, or free-floating in appropriate culture vessels. The culture medium is selected in such a way that selection pressure is exerted and only those cells survive which still contain the hybrid vector DNA including the genetic marker. Thus, for example, an antibiotic is added to the medium when the hybrid vector includes the corresponding antibiotic resistance gene.

When the cell density has reached a sufficient value culturing is interrupted and the protein isolated. If the expression cassette comprises a signal sequence the desired fusion protein may be secreted into the medium. The medium containing the product is separated from the cells which can be provided with fresh medium and used for continuous production. When yeast cells or *E. coli* cells are used the protein can also accumulate within the cells, especially in the periplasmic space. In the latter case the first step for the recovery of the fusion protein consists in liberating the protein from the cell interior. The cell wall is first removed by enzymatic digestion with glucosidases (supra) or, alternatively, the cell wall is removed by treatment with chemical agents, i.e. thiol reagents or EDTA, which give rise to cell wall damages permitting the produced protein to be released. The resulting mixture is enriched for fusion protein by conventional means, such as removal of most of the non-proteinaceous material by treatment with polyethyleneimine, precipitation of the proteins using ammonium sulphate, gel electrophoresis, dialysis, chromatography, for example, ion exchange chromatography (especially preferred when the polypeptide tag of the fusion protein includes a large number of acidic

or basic amino acids), size-exclusion chromatography, HPLC or reverse phase HPLC, molecular sizing on a suitable Sephadex[®] column, or the like. The final purification of the pre-purified product is achieved, for example, by means of affinity chromatography, for example antibody affinity chromatography, especially monoclonal antibody affinity chromatography using antibodies fixed on an insoluble matrix by methods known in the art, which antibodies recognize, for example, epitopes located within the polypeptide tag of the fusion protein.

The invention concerns furthermore the fusion proteins according to the invention per se which are valuable intermediates for the production of biologically active proteins.

The invention concerns especially the expression vectors, the transformed hosts, the fusion proteins and the methods for the preparation thereof and the method for the preparation of a biologically active protein as described in the examples.

The following examples serve to illustrate the invention but should not be construed as a limitation thereof.

Example 1: Construction of a shortened KEX2 gene encoding soluble yscF variant

The yeast KEX2 gene codes for an endoprotease called yscF, which is a membrane bound protein localized in the Golgi apparatus. The yscF protein consists of a N-terminal catalytic domain, a Ser/Thr rich domain, a membrane spanning domain and a C-terminal tail responsible for Golgi localization of the yscF protein. Mutant yscF enzyme lacking 200 C-terminal amino acids, including the Ser/Thr rich domain, the membrane spanning domain and the C-terminal tail still retains protease activity [Fuller et al., 1989, Proc. Natl. Acad. Sci. 86, 1434-1438; Fuller et al., 1989, Science 246, 482-485]. In order to get a soluble yscF protease activity, a mutant KEX2 gene lacking 600 bp, coding for the C terminal 200 amino acids, is constructed. The truncated gene is under the control of the KEX2 promoter reaching from -1 to -502. Translation is terminated at a stop codon (TAA) originating from the polylinker of pUC18.

In detail, plasmid pUC19 [Boehringer Mannheim GmbH, FRG] is digested to completion with HindIII and the 2686 bp fragment is isolated. The ends are filled in and the fragment is religated. An aliquot of the ligation mixture is added to calcium-treated, transformation competent E.coli JM101 [Invitrogen, San Diego, USA] cells. 12 trans-

formed ampicillin resistant E.coli transformants are grown in the presence of 100 µg/ml ampicillin. Plasmid DNA is prepared and analysed by digestion with HindIII as well as with BamHI. The plasmid lacking the HindIII site is designated pUC19woH.

A 3207 bp BalI-AhaIII KEX2 fragment (obtainable from total genomic yeast DNA) is provided at both ends with BamHI linkers followed by a complete digestion with BamHI. Plasmid pUC19woH is cut to completion with BamHI, the linear 2690 bp fragment is isolated and ligated to the BamHI KEX2 fragment described above. An aliquot of the ligation mixture is transformed into E.coli JM101 cells. 12 transformed, ampicillin resistant colonies are grown in ampicillin (100 µg/ml) containing LB medium, plasmid DNA is extracted and analyzed by BamHI digests. One clone with the expected restriction fragments is selected and called pKS301b.

The 2 µm yeast vector pAB24 which corresponds essentially to plasmid pDP34 is cut to completion with BamHI and the linear pAB24 fragment is isolated. Plasmid pKS301b is digested with BamHI and the fragment containing the complete KEX2 gene is isolated and ligated to the linearized yeast vector pAB24. An aliquot of the ligation mixture is transformed into E.coli JM101 and plasmid DNA of twelve positive clones is examined by BamHI digests. One clone with the expected restriction fragments is referred to as pAB226.

Plasmid pKS301b is digested to completion with SphI, PvuII and ScaI. The 2.37 kb SphI-PvuII fragment containing KEX2 sequences from -502 to +1843 and a part of the pUC19 polylinker is isolated. Plasmid pUC18 [Boehringer Mannheim, FRG] is cut to completion with SphI and SmaI. The 2660 bp SphI-SmaI pUC18 fragment is ligated to the 2.37 kb SphI-PvuII KEX2 fragment by SphI/SphI and PvuII/SmaI ligation. The PvuII/SmaI ligation results in the fusion of the KEX2 ORF coding for 614 amino acids to an ORF in the pUC18 sequences which codes for 7 additional C-terminal amino acids (-G-V-P-S-S-N-S) and is followed by a stop codon (TAA). An aliquot of the ligation mixture is transformed into E.coli JM101. Plasmid DNA is isolated from ampicillin resistant E.coli transformants and analyzed by digestion with SphI and EcoRI as well as with HindIII. One clone with the expected restriction pattern is referred to as p18kexp. In the sequence listing under SEQ ID No. 1 the ORF encoding soluble yscF with KEX2-derived DNA is shown.

Plasmid p18kexp is cut to completion with PvuII, SalI and ScaI. The 2552 bp SalI-PvuII

fragment containing the KEX2 sequences reaching from -502 to +1843 as well as 206 bp of pUC18 sequences is isolated. Plasmid pDP34 is digested with BamHI and the ends of the linearized plasmid are filled in. After inactivation of T4 polymerase the linearized filled-in plasmid is cut with SalI and the 11.78 kb fragment is isolated. The pDP34 BamHI*-SalI fragment is ligated to the 2552 bp SalI-PvuII fragment by SalI/SalI and BamHI*/PvuII ligation (BamHI*: filled-in BamHI). An aliquot of the ligation mixture is transformed into transformation competent E.coli JM101 cells. Plasmid DNA is extracted from ampicillin resistant cells and analyzed by restriction analysis with SalI, NcoI, SmaI, XbaI, EcoRI. One clone with the expected restriction fragments is referred to as pDPkexp.

Example 2: Assay of endoprotease activity

Cells are cultured as described in Example 3. The culture broth is centrifuged and separated in cells and culture medium. Endoprotease activity is determined in the culture medium, on the surface of intact cells and in "permeabilized" cells. The cells are washed once with one volume 0.2 M HEPES (4-(2-Hydroxyethyl)-piperazine-1-ethane sulfonic acid, Fluka, Switzerland) pH 7.0 and resuspended in one volume 0.2 M HEPES pH 7.0 for measurements on intact cells or resuspended in 0.2 M HEPES pH 7.0 containing 0.1% Triton X-100 in order to produce "permeabilized" cells. Endoprotease activity is measured as follows: 900 µl of assaymix (0.2 M HEPES pH 7.0, 1 mM CaCl₂, 0.5 mM PMSF [for measuring activity of permeabilized cells additional 0.1% Triton X-100]) is incubated with 50 µl sample at 37°C for 2 minutes and A_{405nm} increase during this incubation period is spectrophotometrically determined. No change in A_{405} can be observed during this preincubation without substrate. The reaction is started by adding the substrate (50 µl of 10 mM benzyloxycarbonyl-L-tyrosyl-L-lysyl-L-arginine-4-nitroanilide, Bachem, Bubendorf, Switzerland) and the increase in A_{405} is taken for calculation of endoprotease activity. 1U endoprotease activity is defined as 1 µmol 4-nitroanilide produced per minute under the described assay conditions. About 90% of yscF activity is found in the culture medium.

Example 3: Transformation of *S.cerevisiae* strain AB110

Saccharomyces cerevisiae strain AB110 (ATCC 20796) is transformed with plasmids pDP34, pAB226 and pDPkexp using the transformation protocol described by Dohmen et al. [1989, Curr. Genet. 15, 319-325]. Transformed yeast cells are selected on yeast

minimal plates supplemented with an amino acid/nucleotide mix deficient in uracil (amino acid/nucleotide mix supplements the culture medium with adenine 12 mg/l, arginine 40 mg/l, histidine 40 mg/l, isoleucine 60 mg/l, leucine 60 mg/l, lysine 60 mg/l, methionine 40 mg/l, phenylalanine 50 mg/l, threonine 60 mg/l, tryptophan 40 mg/l, tyrosine 15 mg/l, uracil 40 mg/l, valine 60 mg/l). Single transformed yeast clones are isolated and referred to as:

Saccharomyces cerevisiae AB110/pDP34

Saccharomyces cerevisiae AB110/pAB226

Saccharomyces cerevisiae AB110/pDPkexp.

Example 4: Fermentation of transformed yeast strains on a laboratory scale

Cells of *S. cerevisiae* AB110/pDP34, *S. cerevisiae* AB110/pAB226 and *S. cerevisiae* AB110/pDPkexp are grown in a 10 ml preculture composed of

Difco Yeast Nitrogen Base w/o Amino Acids	6.7 g/l
glucose	20.0 g/l
HEPES	24.0 g/l.

The culture medium is supplemented with an amino acid/nucleotide mix deficient in leucine (described in Example 3) and the pH is set to 7.0.

The precultures are grown for 48 h at 30°C and 180 r.p.m..

The main culture medium is composed of:

Difco Yeast Nitrogen Base w/o Amino Acids	6.7 g/l
glucose	40.0 g/l
arginine	8.0 g/l
Difco Casaminoacids	8.5 g/l
HEPES	24.0 g/l,
pH is set to 7.0.	

The main cultures are inoculated with about 1% v/v of the precultures and incubated at 30°C and 180 r.p.m.. At several time points during the fermentation aliquots of the cultures are taken and analyzed for cell density, pH of the culture broth and

endoprotease activity. Figure 1 shows the intracellular and extracellular yscF activities synthesized by the S.cerevisiae AB110/pDPkexp transformant proving that a soluble secreted yscF variant is expressed in comparison to the transformant AB110/pDP34 which harbors only the yeast vector and the transformant AB110/pAB226 which expresses the entire yscF protein.

Example 5: Immunological demonstration of a truncated soluble yscF protein in the culture medium of S.cerevisiae AB110/pDPkexp transformants

Strains S.cerevisiae AB110/pDPkexp and AB110/pDP34 are grown as described in example 4. The culture broths are separated in cells and culture supernatants by centrifugation. The culture supernatants are concentrated using Centricon 30000 filters (Amicon) to 1/5 of the original volume. Different amounts of the concentrated supernatants are reduced with 1,4-dithio-DL-threitol and separated on a 8% polyacrylamide gel under denaturing conditions. The proteins are either stained with Coomassie brilliant blue or blotted onto a nitrocellulose filter (Western blot). The methods for PAGE and Western blot are described in Sambrook et al., 1989, Molecular Cloning, 2nd edition, Cold Spring Harbor Laboratory Press, New York.

The nitrocellulose filter is incubated at 37°C for 2h in 10 mM NaH₂PO₄, 150 mM NaCl, 1% BSA, pH7.2 followed by a second incubation under the same conditions in the same solution which now contains additional polyclonal antibodies raised in rabbits against a lacZ-yscF fusion. (Production and purification of antibodies is described in Sambrook et al., 1989, Molecular Cloning, 2nd edition, Cold Spring Harbor Laboratory Press, New York.) The filter is washed three times for 30 minutes at 37°C in 10 mM NaH₂PO₄, 150 mM NaCl, 1% Triton X-100*, 0.1% BSA, pH 7.2. These washings are followed by a hybridization reaction with Goat anti-rabbit Ig's alkaline phosphatase (TAGO, Inc., Burlingame, CA), an alkaline phosphatase which is coupled to an antibody specific for the constant regions of antibodies raised in rabbits. This reaction is performed in 10 mM NaH₂PO₄, 150 mM NaCl, 0.01% Tween 20*, 0.1% BSA, pH 7.2, containing the anti-rabbit Ig's alkaline phosphatase. The blot is washed three times for 30 minutes in 10 mM NaH₂PO₄, 150 mM NaCl, 0.01% Tween 20*, 0.1% BSA, pH 7.2. Visualization of the specific binding occurs by incubation of the blot in a solution composed of: 0.1 M Tris-HCl pH8.8, 100 mM NaCl, 2mM MgCl₂, 10 mg/100 ml 5-Brom-4-Chlorindolyl-Phosphat (BCIP) and 100 mg/100ml Nitroblue-Tetrazolium-Chloride (NBT). An additional 68 kDa protein is found in the culture supernatants of S.cerevisiae AB110/pDPkexp compared to the culture supernatants of S.cerevisiae

*Trade-mark

AB110/pDP34. This 68 kDa protein reacts with the antibodies raised against a part of the yscF protein.

Example 6: In vitro processing of pre-pro-IGF1

IGF1 is expressed in *S.cerevisiae* as an intracellular fusion protein composed of mature IGF1 fused to the leader sequence of the *S.cerevisiae* α -factor precursor. The fusion protein has the following structure:

N-(α -factor leader)-IGF1-C

The amino acid sequence Tyr-Lys-Arg is the C-terminal end of the 89 amino acid α -factor leader sequence [Kurjan, J. and Herskowitz, I. 1982, Cell 30, 933-943] and a substrate for the yscF endoprotease. The fusion protein is cleaved by the truncated soluble yscF protease which leads to the digestion products N-pre-pro- α -factor-C and mature IGF1.

S.cerevisiae crude extract samples each containing 50 pmol of the α -factor leader IGF1 fusion protein are incubated with culture supernatant samples of *S.cerevisiae* AB110/pDPkexp each containing 5 mU soluble yscF endoprotease activity (calculated from the endoprotease activity assay described in example 2). The reactions are performed in 10 μ l 50 mM Na-phosphate buffer pH 7.5 for different time periods. Mature IGF1 derived from the in vitro processing procedure is detected by "Western blot" analysis. This method is described in example 5. Hybridization is performed using polyclonal antibodies raised in rabbits against mature IGF1. Detection of specific binding is described in example 5. Western blot analysis shows the processing of pre-pro-IGF1 with a culture supernatant containing the soluble yscF protease variant secreted from the *S.cerevisiae* AB110/pDPkexp transformant. The pre-pro-IGF1 protein is processed to the mature IGF1 protein within 30 to 120 min using the described assay conditions.

Example 7: Construction of a hybrid gene containing the GAPFL promoter and the KEX1 structural gene.

As disclosed in European Patent Application No. 341,215 the full-length KEX1 gene is isolated from total genomic *Saccharomyces cerevisiae* DNA on a 3.1 kb HindIII fragment. This fragment is cloned into the unique HindIII site of the yeast plasmid pJDB207 [Beggs,

- 23 -

J.D., 1981, in D. von Wettstein et al. (ed.), Molecular Genetics in Yeast, Alfred Benzon Symposium 16. Munsgaard, Copenhagen] . E. coli HB101 is transformed with the resulting plasmid pJDB207/KEX1. The transformed E. coli strain is designated E. coli HB101/KEX1.

In order to obtain high level expression of a secreted soluble form of ysc α (the protein encoded by KEX1), the very weak KEX1-promoter present on the HindIII fragment is exchanged for the strong constitutive promoter element (GAPFL) of the yeast glyceraldehyd-3-phosphate dehydrogenase promoter (European Patent Application No. 225,633). Since the GAPFL promoter is contained on a 478 bp SalI-EcoRI fragment, first an EcoRI-site in the KEX1 coding region is eliminated by in vitro mutagenesis, followed by introduction of a new EcoRI site directly 5' to the KEX1 ATG start-codon to allow proper fusion of the GAPFL promoter to the KEX1 coding region.

In detail, the 3.1 kb HindIII fragment containing KEX1 is subcloned into the HindIII site of the M13-derived vector pBluescript KS+ (Stratagene, La Jolla, Ca. USA) to give KS+/KEX1. To remove the internal EcoRI site located 420 bp upstream of the 5'-HindIII site, the following oligonucleotide is synthesized:

5'- CCTTTTAGGGTCAATTCAGACGGT -3'

Site-directed in vitro mutagenesis is carried out as described (Bio-Rad Muta-Gene* M13 kit, Bio-Rad, Richmond, Ca. USA). First, KS+/KEX1 is transfected into E. coli CJ 236 to incorporate uracil (Muta-Gene kit, supra). Single-stranded DNA from transfected E. coli CJ 236 is isolated using M13 helper phage (Stratagene, supra).

200 pmoles of the oligonucleotide primer are phosphorylated in a total volume of 30 μ l containing 3 μ l 1 M Tris-HCl pH 8.0, 0.3 μ l 1 M MgCl₂, 0.75 μ l 0.2 M DTT, 0.6 μ l 20 mM ATP. 4.5 units T4 polynucleotide kinase are added and the mixture incubated at 37°C for 45 min and at 65°C for 10 min.

The phosphorylated oligonucleotide is then annealed to the template DNA under the following conditions: 0.1 pmoles of uracil containing DNA derived from KS+/KEX1 are incubated with 2 pmoles of phosphorylated primer in a total volume of 10 μ l annealing buffer (20 mM Tris-HCl pH 7.4, 2 mM MgCl₂, 50 mM NaCl). The mixture is heated in a water bath to 80°C and then allowed to cool slowly until ambient temperature is reached.

*Trade-mark

The complementary strand is then formed under the following conditions: 10 µl of the annealing mixture are incubated with 4 µl 2 mM dNTP's, 0.75 µl 0.5 M Tris-HCl pH 7.4, 0.75 µl 0.1 M MgCl₂, 2.15 µl 0.2 M DTT 1 unit T4 DNA polymerase and 2 units T4 DNA ligase. The reaction mixture is first incubated on ice for 5 min, then at 25°C for 5 min and finally at 37°C for 90 min. The resulting double-stranded DNA's are transformed into E. coli JM 101, a strain which efficiently removes the uracil-containing template, leaving the mutagenized complementary strand to replicate (Bio-Rad, supra). Plasmids are prepared and analysed for the absence of the EcoRI site. Correct mutagenesis is further confirmed by sequence analysis. One plasmid with the correct disruption of the internal EcoRI site is designated as KS+/KEX1(-EcoRI).

A new EcoRI site is now introduced into KS+/KEX1(-EcoRI) using the following oligonucleotide as primer:

5'- AGATAAAGACCTGAATTCAGATGTTTTACAAT -3'

In vitro mutagenesis is carried out exactly as described supra. Plasmids are checked by restriction analysis for the presence of the new EcoRI site immediately adjacent to the start codon and by sequence analysis. One plasmid with the correct new EcoRI site is designated as KS+/KEX1(EcoRInew).

KS+/KEX1(EcoRInew) is digested with EcoRI and HindIII and the 3.0 kb KEX1 containing fragment separated on a 0.8 % preparative agarose gel and isolated using GeneClean*(Bio 101 Inc., La Jolla, CA, USA).

To obtain the GAPFL promoter fragment plasmid pJDB207/GAPFL-HIR (European Patent Application No 225,633) is digested with SalI and EcoRI and the 478 bp fragment isolated using the same procedure as described (GeneClean*, supra).

0.2 pmoles of the 478 bp SalI-EcoRI promoter fragment, 0.2 pmoles of the EcoRI-HindIII fragment containing the full-length KEX1 structural gene and 0.1 pmoles of SalI-HindIII cut yeast multicopy vector pDP34 (cf. European Patent Application No. 340,170) are ligated and plasmids prepared from transformed E. coli JM 101 cells. One correct plasmid is referred to as pDP34/GAPFL-KEX1.

*Trade-mark

Example 8: Construction of a shortened KEX1 gene encoding soluble ysc α activity

KEX1 encodes a membrane-bound carboxypeptidase - ysc α - which is localised within the secretory machinery, most probably in the Golgi-apparatus. The ysc α protein consists of an amino-terminal catalytic domain, followed by an Asp-Glu rich acidic region, a transmembrane domain and a C-terminal tail. To allow for secretion into the medium via the default pathway, truncation at the C-terminus is the method of choice. A convenient unique XhoI site is located shortly upstream of the transmembrane domain of the encoded protein at the end of the acidic region.

KS+/KEX1 (EcoRI new) is digested with EcoRI and XhoI and the 1.7 kb fragment containing the C-terminally truncated KEX1 gene isolated. This 1.7 kb KEX1 encoding fragment and the 478 bp SalI/EcoRI GAPFL promoter fragment (Example 7) are then ligated into the unique SalI site of pDP34. The resulting plasmid is referred to as pDP34/GAPFL-KEX1*. The ORF encoding soluble ysc α * with KEX1-derived sequences is depicted in the sequence listing under SEQ ID No. 2.

Example 9: Transformation of *S. cerevisiae* strain Tr 1176

a. Crossing of *S. cerevisiae* kex1 mutant strain 96 with *S. cerevisiae* strain BYS and analysis of the spores on α -factor secretory capacity and carboxypeptidase ysc α (KEX1 gene) activity

The *S. cerevisiae* kex1 mutant strain 96 (a, kex1, ade2, thr1), which is obtained from the yeast Genetic Stock Center, Berkeley, USA, is crossed into *S. cerevisiae* strain BYS232-31-42 (α , prb1-1, prel-1, cps1-3, lys2, leu2, his7) [Achstetter, T. and Wolf, D.H. (1985) EMBO J. 4, 173 - 177; Wolf, D.H. and Ehmann, C. (1981) J. Bacteriol. 147, 418 - 426] carrying the wild-type KEX1 allele. Diploid heterozygous cells of the genotype kex1/KEX1 are isolated from this cross. The tetrads which derive from the diploid cells are dissected according to standard genetic techniques [Hawthorne, D.C. and Mortimer, R.K. (1960) Genetics 45, 1085 - 1110; Methods in Yeast Genetics 1986 (Sherman, F. et al., eds.) Cold Spring Harbor Laboratory, N.Y.].

The four spores of each tetrad are tested for their ability to secrete α -factor. To distinguish between KEX1 wild-type and kex1 mutant colonies, the pheromone-supersensitive tester strain *S. cerevisiae* RC629 (a, sst-2, ade2-1, ura1, his6, met1, can1, cyh2, rme) is used

[Chan, R.K. and Otte, C.K. (1982) Mol. Cell. Biol. 2, 11 - 20; Chan, R.K. and Otte, C.K. (1982) Mol. Cell. Biol. 2, 21 - 29]. As expected from traits coded for by single nuclear genes, from all tetrads analysed, two spores of each tetrad secrete the a-factor, whereas the two other spores secrete α -factor. Wild-type KEX1 colonies of the α -mating type inhibit growth of the tester strain to a large extent and thus produce a large halo around themselves, since they are able to process the α -factor precursor completely and produce four active α -factor molecules from one precursor molecule. In contrast, *kex1* mutant colonies inhibit the growth of the tester strain to a less extent and thus produce a small halo around themselves, since they are only able to produce one mature α -factor molecule from one precursor molecule.

The spores of several complete tetrads which are identified as defective at the *kex1* gene by the above described pheromone assay, are finally tested for specific activity of carboxypeptidase *ysc α* . Cells are grown, membranes thereof are prepared and tested for carboxypeptidase *ysc α* activity using Chz-Tyr-Lys-Arg as substrate as described [Wagner, J.C. and Wolf, D.H. (1987) FEBS Lett. 221, 2, 423 - 426]. The fact that activity of carboxypeptidase *ysc α* is lacking in *kex1* mutant cells, indicates that KEX1 is the structural gene of this enzyme. This implies that carboxypeptidase *ysc α* is indeed involved in carboxy-terminal processing of α -factor.

b. Classification of confirmed *kex1* mutants on additional deficiency of proteases *yscB*, *yscY* and *yscS*

S. cerevisiae *kex1* mutants are classified with regard to the deficiency of other proteases (proteinase *yscB*, carboxypeptidase *yscY* and carboxy-peptidase *yscS*) and additional growth factor requirements.

Cell material of *kex1* mutants which are prepared from stationary phase in YPD (Difco) medium is suspended in the 200 μ l 20 mM Tris-HCl buffer, pH 7.2 in Eppendorf microfuge and glass beads (0.4 mm in diameter) are added up to two thirds of the volume. The suspension is heavily shaken three times for 1 min on a vortex mixer with intermittent cooling on ice.

Centrifugation for 5 min allows recovery of the supernatant crude extracts. These extracts are dialysed against 0.1 M imidazole-HCl buffer pH 5.2 with 0.1 mM ZnCl_2 in order to activate proteases and to remove free amino acids from the extracts.

Proteinase yscB activities are measured according to the Azocoll- test [R.E. Ulane et al. (1976) J. Biol. Chem. 251, 3367; E. Cabib et al. (1973) Biochem. Biophys. Res. Commun. 50, 186; T. Saheki et al. (1974) Eur. J. Biochem. 42, 621]. After the protein concentration measurements, an aliquot of each sample is filled with 0.1 M sodium phosphate (NaPi) buffer pH 7.0 up to 100 µl to adjust the required equal protein amounts. To the protein solution, a suspension of 500 µl Azocoll (240 mg in 10 ml 0.1 M NaPi buffer, pH 7.0) is added. These mixtures are incubated at 30°C for one hour with agitation. After the addition of 500 µl 10 % tri-chloroacetic acid which stops the reaction, the mixtures are centrifuged two times and the absorption spectra of the supernatants at 520 nm are measured.

The activities of carboxypeptidase yscY and yscS are measured using the chromogenic substrate Cbz-Gln-Leu [cf. D.H. Wolf et al. (1978) FEBS Lett. 91, 59; D.H. Wolf et al. (1977) Eur. J. Biochem. 73, 553]. The dialysed extracts are divided into three portions and to two of them phenyl-methylsulfonyl fluoride (PMSF) at a final concentration of 1 mM or EDTA at a final concentration of 5 mM is added to block the two protease activities selectively. Namely PMSF inhibits carboxypeptidase yscY activity and EDTA inhibits that of carboxypeptidase yscS. The mixtures with inhibitors are each incubated at 25°C for one hour to complete the inhibition. After the determination of the protein concentration, two aliquots with inhibitor and one aliquot without inhibitor as a control of each sample are filled with 0.1 M NaPi buffer pH 7.4 up to 50 µl in order to receive equal protein amounts. To these protein solutions the following test solutions are added.

500 µl test solution I:

L-amino acid oxidase	0.24 mg/ml
horseradish peroxidase	0.40 mg/ml
0.01 mM MnCl ₂	
in 0.1 M NaPi buffer, pH 7.4	

50 µl test solution II

o-dianisidin	2 mg/ml
in water	

500 µl test solution III

20 mM Cbz-Gly-Leu	
in 0.2 M potassium phosphate buffer, pH 7.4	

The mixtures are incubated at 28°C for one hour and after the addition of 100 µl 20 % Triton X-100 to stop the reaction, the absorbances at 405 nm are measured.

For the purpose of the subsequent transformation, an amino acid auxotrophic marker for leucine is scored with the replica-technique on minimal plates supplied with adenine, threonine, lysine and histidine, and with or without leucine.

By means of the above described assays, mutants are isolated designated *S. cerevisiae* BYSkex1, which exhibit a quadruple protease-deficiency (α , prb-1, pre-1, cps-3, kex1) and an additional requirement for leucine.

c. Production of ura3-deficient yeast strain TR1176

Tr 1176 is a ura3- derivative of strain BYSkex1 described above. Disruption of the URA3 gene in BYSkex1 is carried out as detailed in European Patent Application Nr. 340,170. Tr 1176 has the following genetic markers: MAT α , prb-1, pre-1, cps-1, kex-1, ade2, leu2, ura3.

S. cerevisiae Tr 1176 is transformed with plasmids pDP34/GAPFL-KEX1 and pDP34/GAPFL-KEX1* using the transformation protocol described by Dohmen (supra). Transformed yeast cells are selected on yeast minimal media plates supplemented with leucine and adenine and deficient in uracil. Single transformed yeast colonies are isolated and referred to as:

Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1

and

Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1*

Example 10: Biological activity of soluble ysc α

Cells of Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1 and Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1* are grown in minimal medium composed of :

Difco Yeast Nitrogen Base w/o amino acids	8.4 g/l glucose
	20.0 g/l L-asparagine
	10.0 g/l adenine
	0.1 g/l

for 48 h at 30°C and 180 r.p.m.

The cells are separated from the medium by centrifugation, cells are resuspended in 0.9 % NaCl and tested for ysc α activity by halo-formation on a lawn of α -factor supersensitive a-cells (Chan, R.K. and Ote, C.K. (1982) Mol.Cell.Biol.2, 11-20). Strain Tr 1176 without plasmid shows only a minute halo on the a-cell lawn due to the lack of ysc α activity.

Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1 and Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1* both show halo-formation, soluble ysc α has therefore retained its normal biological function, i.e. α -factor maturation.

Supernatants of Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1 and Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1* fermentations (supra) are then analysed for the presence of ysc α activity in the medium using the synthetic peptide substrate Cbz-Tyr-Lys-Arg and an assay as described (Wolf, D.H. and Weiser,U. (1977) Eur.J.Biochem. 73, 553-556). Only supernatants of Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1* show proteolytic degradation of this substrate, whereas supernatants of Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1 have no activity on Cbz-Tyr-Lys-Arg. It can therefore be concluded that KEX1* encodes biologically active ysc α which is released into the medium. The new protein is referred to as ysc α *.

Example 11: Purification of ysc α *

5 ml of the ion exchanger Fractogel TSK DEAE 650 (Merck AG, Darmstadt, FRG) are packed into a column and equilibrated with 50 mM sodium phosphate buffer, pH 7.0. 80 ml of culture broth of Saccharomyces cerevisiae Tr 1176/pDP34/GAPFL-KEX1* (supra) are adjusted to pH 7.0 and loaded onto the column overnight. ysc α * under those conditions binds to the column. Bound proteins are then eluted with a combined pH/salt gradient (40 ml 50 mM sodium phosphate buffer pH 7.0 and 40 ml 50 mM sodium-phosphate buffer pH 4.0 + 1 M NaCl). At about 0.2 M NaCl ysc α * elutes in essentially pure form with an apparent molecular weight of about 66 kDa (calculated molecular weight 65 kDa). When compared with porcine pancreas carboxypeptidase B (Boehringer Mannheim, Mannheim, FRG) ysc α * at the same protein concentration has the equivalent activity on the substrate Cbz-Tyr-Lys-Arg at pH 7.0 and slightly higher activity at pH 4.0.

Example 12: Construction of a recombinant fusion peptide expressing hirudin and human calcitonin simultaneously

A fusion peptide is constructed which allows for simultaneous expression of hirudin and of precursor human calcitonin. Between the two peptides the KEX2/KEX1 recognition site Lys-Arg is introduced to allow for in vitro processing with soluble yscF and soluble ysc α * (supra).

In detail, plasmid pJDB207/GAPFL-HIR (supra, Example 7) is digested with Sall and BamHI and the 0.7 kb fragment subcloned into Sall-BamHI of pBluescript KS⁺. The gene encoding the C-terminal glycine-precursor of human calcitonin is assembled from individual oligonucleotides as described for hirudin (cf. European Patent Application No. 168342) and subcloned into the PstI site of pUC19. The sequence of human precursor calcitonin with a 5'-extension to provide for in vitro processing is depicted in the sequence listing under SEQ ID No. 3 (as confirmed by sequence analysis).

The 132 bp PstI fragment is excised and cloned into the PstI site of the hirudin containing Bluescript vector.

Transformants are checked for proper orientation of the insert and one correct plasmid is referred to as KS+/GAPFL-HIR-CALC.

Plasmid pJDB207/GAPFL-HIR is digested with Sall and HindIII and the 7 kb large vector

fragment isolated. Plasmid pJDB207/GAPFL-HIR is digested with BamHI and HindIII to obtain the 350 bp PH05 terminator fragment.

Finally, pJDB207 vector fragment, terminator fragment and the SalI-BamHI fragment of KS+/GAPFL-HIR-CALC are ligated in a triple ligation to yield plasmid pJDB207/GAPFL-HIR-CALC.

Saccharomyces cerevisiae strain HT246 (DSM 4084) is transformed with plasmid pJDB207/GAPFL-HIR-CALC according to Example 3. Transformed yeast cells are selected on yeast minimal medium plates deficient in leucine. Single transformed yeast cells are isolated and referred to as S. cerevisiae HT246/pJDB207/GAPFL-HIR-CALC.

For the expression and secretion of the fusion protein S. cerevisiae HT246/pJDB207/GAPFL-HIR-CALC is cultivated as disclosed in European Patent Application No. 340170 (example 10 therein). After 48h and 72h of cultivation samples are withdrawn, the cells removed by centrifugation and the culture supernatant containing the fusion protein taken for further processing. First, the supernatant containing the hirudin-calcitonin fusion protein is endoproteolytically split by digestion with truncated soluble yscF protein according to Example 6. This cleavage results in full-length precursor human calcitonin and a hirudin variant which still contains a lysine-arginine extension at its C-terminus. This lysine-arginine extension is removed by digestion with ysc α * using purified enzyme (see Example 11). Correct removal of the lysine-arginine extension is demonstrated by reversed phase HPLC of the reaction mixture as disclosed in European Patent Application No. 340170.

Example 13: Construction of a recombinant fusion protein expressing eglin C

A fusion peptide is constructed which allows the expression of a fusion protein according to the formula P-L-T in which P is the eglin C polypeptide (Rink H. et al., 1984 Nucl. Acids Res. 12, 6369-6387), L the linker sequence

Lys-Arg-Glu-Ala-Glu-Ala-Trp-Val-Pro and T a cellulose binding domain of the Cellulomonas fimi Exg protein encoded in the Cellulomonas fimi cex gene (Neill G.O. et al., 1986 Gene 44, 325-330).

In detail, plasmid pEC-1 (Neill G.O. et al., 1986 Gene 44, 325-330) is cut to completion with SmaI and StuI. The 433 bp fragment encoding the cellulose binding domain of the

cellulomonas cex gene is isolated. Plasmid p18kexp (example 1) is cut with Asp718 and the sticky ends are filled in using Klenow polymerase resulting in blunt ends. The linearised plasmid is ligated with the 433 bp SmaI-StuI cex fragment and transformed into *E.coli* HB101. The clones are checked for the orientation of the 433 bp cex fragment. One clone with the orientation that couples the reading frame of the KEX2 gene to the reading frame of the cex gene is referred to as p18kexpcex.

Plasmid pUC19 is cut with Sall and BamHI and the 276 bp Sall-BamHI fragment of pBR322 is inserted. The resulting plasmid pUC19pBR is cut with BamHI and EcoRI and ligated in the presence of the BamHI-EcoRI GAPDH promoter fragment resulting in plasmid pTM1. pTM1 is cut with EcoRI and the following linker is inserted

5' -AATTCATGCA TGCATGCAGA TCT-3'

3' -GTACGT ACGTACGTCT AGATTAA-5'

in such a way that the EcoRI site is reconstituted adjacent to the promoter sequences. The resulting plasmid is referred to as pTM2.

Plasmid pML147 (Rink H. et al., 1984 Nucl. Acids Res. 12, 6369-6387) is cut to completion with EcoRI and BamHI. The EcoRI-BamHI eglin C fragment is cloned into EcoRI/BamHI cut pBR322. One plasmid with the desired sequence is called pML141.

Using the oligonucleotides

5'-CCGGTACCCA AGCTTCGGCT TCTCTCTTAC CAACATGCGG AACATG-3'

and

5'-CGGGA'TCCAA GAATTCATGA CTGAATT-3'

(the sequence coding for the linker sequence L referred to above is underlined) a PCR reaction is performed on pML141 using the PCR-Kit from Perkin- Elmer-Cetus and following strictly the instructions of the supplier. The PCR reaction results in the amplification of a 257 bp fragment containing the eglin C gene joined to the linker sequence L. The fragment is isolated from an agarose gel and cut to completion with BamHI and Asp718. Plasmid p18kexpcex (see above) is cut to completion with BamHI and Asp718 and the 3.1 kb fragment is isolated. The 3.1 kb BamHI-Asp718 p18kexpcex fragment containing the cex gene fragment and pUC19 sequences is ligated with the PCR generated BamHI-Asp718 eglin C fragment and the resulting plasmid is called p18eglinecex. The plasmid contains an open reading frame reaching from the first amino acid of eglin C to the last amino acid of the Exg protein. The two domains, i.e. eglin C and the 133 C-terminal amino acids of Exg (including an extra proline), are separated by the linker sequence given above. The nucleotide sequence encoding the fusion protein (see

SEQ ID No. 4) is confirmed by DNA sequencing.

Plasmid pTM2 (see above) is cut with EcoRI and the 5'-ends are dephosphorylated. Plasmid p18eglin-cex is cut with EcoRI and the 686 bp eglin-cex fragment is isolated. This 686 bp fragment is ligated into the EcoRI cut and dephosphorylated vector pTM2. The resulting clones are checked for the orientation of the eglin-cex fragment. One clone with the desired orientation resulting in the following order of the sequences GAPDH-eglin-cex is called p19/GAPDH-EGLIN-CEX.

Plasmid p19/GAPDH-EGLIN-CEX is digested with BamHI and BglII and the 1.1 kb fragment encoding the GAPDH-eglin-cex sequences is isolated. Plasmid pDP34 is cut with BamHI, the 5'-ends are dephosphorylated and the resulting fragment is ligated with the 1.1 kb BamHI-BglII fragment. The clones are checked for the orientation of the 1.1 kb BamHI-BglII fragment and referred to as pDP34GAPDH-eglin-cex-1 and pDP34GAPDH-eglin-cex-2 wherein index 1 indicates that the insert GAPDH-eglin-cex is clockwise oriented in plasmid pDP34 while index 2 indicates a counterclockwise orientation.

pDP34GAPDH-eglin-cex-1 is transformed into Saccharomyces cerevisiae strain AB110 (ATCC 20796) according to example 3. Cells of AB110/pDP34GAPDH-eglin-cex-1 are grown in 10 ml minimal medium (Difco Yeast Nitrogen Base w/o Amino Acids 6.7 g/l, glucose 40 g/l, amino acid/nucleotide mix according to example 3 w/o leucine) for 24 h at 30 °C with 180 rpm. The main cultures, 250 ml of minimal medium are inoculated with the preculture and grown at 30 °C for 24 h with 180 rpm. The production culture is set up as follows: The cells of the main culture are separated from the culture broth by centrifugation, resuspended in 250 ml of double minimal medium (Difco Yeast Nitrogen Base w/o Amino Acids 13.4 g/l, glucose 40 g/l, 2x amino acid/nucleotide mix according to example 3 w/o leucine) and fermented at 30 °C for 24 h with 180 rpm.

The cells of the production culture are separated from the culture medium by centrifugation, washed first in 200ml and then in 40 ml buffer A (buffer A: 50 mM Tris-HCl pH 7.0, 100 mM NaCl). Cells are broken with glass beads in repeated steps each of which in about 10 ml buffer A and the crude extract of a total volume of 40 ml is cleared by centrifugation (25000 x g, 15 min, 4 °C). 5 g (dry weight) cellulose (Avicel or fibrous cellulose) is added to the crude extract and incubated on a turning shaker for 1.5 h at 4 °C.

The cellulose bound fusion protein is eluted with water as a fusion protein and afterwards digested with yscF resulting in eglin C with the C-terminal extension Lys-Arg. These two C-terminal amino acids are removed by digestion with ysc α . In detail: The cellulose after incubation in the crude extract is filled into a chromatography column (1.6 x 20 cm) and washed on the column with three column bed volumes buffer A. Then a gradient of buffer A and water is applied. The gradient is linear and runs from 100% buffer A, 0% water to 0% buffer A, 100 % water within three column bed volumes. Fractions are collected and tested for the fusion protein by the immunological method as described in example 5. Fractions containing the fusion protein are subjected to a treatment with the soluble protease yscF as described in example 6 resulting in the cleavage product eglinC-Lys-Arg which is afterwards digested with soluble ysc α * to remove the Lys-Arg tail.

Alternatively, the cellulose bound fusion protein is digested with yscF while bound to cellulose and a protein consisting of eglin C with the C-terminal extension Lys-Arg is set free. This protein is afterwards digested with ysc α * to remove these two C-terminal amino acids. In detail: Different amounts (10 mg to 500 mg) of the cellulose after incubation with the crude extract are incubated with the soluble protease yscF as described in example 6 whereby eglinC-Lys-Arg is set free. Eglin C-Lys-Arg is further digested with soluble ysc α * in order to remove the Lys-Arg tail. The obtained eglin C is identical in structure to natural eglin C as evidenced by HPLC.

Deposition of microorganisms

The following microorganism strains were deposited at the Deutsche Sammlung von Mikroorganismen (DSMZ), Mascheroder Weg 1b, D-3300 Braunschweig (deposition dates and accession numbers given):

Escherichia coli JM109/pDP34: March 14, 1988, DSM 4473.

Saccharomyces cerevisiae BYS232-31-42: May 6, 1988, DSM 4583.

Escherichia coli JM101/pKS301b: June 25, 1990, DSM 6028.

Escherichia coli HB101/KEX1: June 25, 1990, DSM 6027.

Escherichia coli JM101/p19/GAPDH-EGLIN-CEX: June 3, 1991, DSM6546.

SEQ ID No. 1

Sequence type: Polynucleotide with corresponding polypeptide

Sequence length: 1866 base pairs

Strandedness: double

Topology: linear

Source: yeast genomic DNA

Immediate experimental source: E.coli JM101/pKS301b (DSM6028)

Features: from 1 to 1866 coding region for soluble yscF

ATG	AAA	GTG	AGG	AAA	TAT	ATT	ACT	TTA	TGC	TTT	TGG	TGG	39
Met	Lys	Val	Arg	Lys	Tyr	Ile	Thr	Leu	Cys	Phe	Trp	Trp	
1				5					10				
GCC	TTT	TCA	ACA	TCC	GCT	CTT	GTA	TCA	TCA	CAA	CAA	ATT	78
Ala	Phe	Ser	Thr	Ser	Ala	Leu	Val	Ser	Ser	Gln	Gln	Ile	
	15					20				25			
CCA	TTG	AAG	GAC	CAT	ACG	TCA	CGA	CAG	TAT	TTT	GCT	GTA	117
Pro	Leu	Lys	Asp	His	Thr	Ser	Arg	Gln	Tyr	Phe	Ala	Val	
			30					35					
GAA	AGC	AAT	GAA	ACA	TTA	TCC	CGC	TTG	GAG	GAA	ATG	CAT	156
Glu	Ser	Asn	Glu	Thr	Leu	Ser	Arg	Leu	Glu	Glu	Met	His	
40						45				50			
CCA	AAT	TGG	AAA	TAT	GAA	CAT	GAT	GTT	CGA	GGG	CTA	CCA	195
Pro	Asn	Trp	Lys	Tyr	Glu	His	Asp	Val	Arg	Gly	Leu	Pro	
		55					60				65		
AAC	CAT	TAT	GTT	TTT	TCA	AAA	GAG	TTG	CTA	AAA	TTG	GGC	234
Asn	His	Tyr	Val	Phe	Ser	Lys	Glu	Leu	Leu	Lys	Leu	Gly	
			70						75				
AAA	AGA	TCA	TCA	TTA	GAA	GAG	TTA	CAG	GGG	GAT	AAC	AAC	279
Lys	Arg	Ser	Ser	Leu	Glu	Glu	Leu	Gln	Gly	Asp	Asn	Asn	
	80						85				90		

GAC	CAC	ATA	TTA	TCT	GTC	CAT	GAT	TTA	TTC	CCG	CGT	AAC	312
Asp	His	Ile	Leu	Ser	Val	His	Asp	Leu	Phe	Pro	Arg	Asn	
			95									100	

GAC	CTA	TTT	AAG	AGA	CTA	CCG	GTG	CCT	GCT	CCA	CCA	ATG	351
Asp	Leu	Phe	Lys	Arg	Leu	Pro	Val	Pro	Ala	Pro	Pro	Met	
105						110					115		

GAC	TCA	AGC	TTG	TTA	CCG	GTA	AAA	GAA	GCT	GAG	GAT	AAA	390
Asp	Ser	Ser	Leu	Leu	Pro	Val	Lys	Glu	Ala	Glu	Asp	Lys	
		120					125					130	

CTC	AGC	ATA	AAT	GAT	CCG	CTT	TTT	GAG	AGG	CAG	TGG	CAC	429
Leu	Ser	Ile	Asn	Asp	Pro	Leu	Phe	Glu	Arg	Gln	Trp	His	
				135					140				

TTG	GTC	AAT	CCA	AGT	TTT	CCT	GGC	AGT	GAT	ATA	AAT	GTT	468
Leu	Val	Asn	Pro	Ser	Phe	Pro	Gly	Ser	Asp	Ile	Asn	Val	
	145						150					155	

CTT	GAT	CTG	TGG	TAC	AAT	AAT	ATT	ACA	GGC	GCA	GGG	GTC	507
Leu	Asp	Leu	Trp	Tyr	Asn	Asn	Ile	Thr	Gly	Ala	Gly	Val	
			160						165				

GTG	GCT	GCC	ATT	GTT	GAT	GAT	GGC	CTT	GAC	TAC	GAA	AAT	546
Val	Ala	Ala	Ile	Val	Asp	Asp	Gly	Leu	Asp	Tyr	Glu	Asn	
170						175					180		

GAA	GAC	TTG	AAG	GAT	AAT	TTT	TGC	GCT	GAA	GGT	TCT	TGG	585
Glu	Asp	Leu	Lys	Asp	Asn	Phe	Cys	Ala	Glu	Gly	Ser	Trp	
		185					190					195	

GAT	TTC	AAC	GAC	AAT	ACC	AAT	TTA	CCT	AAA	CCA	AGA	TTA	624
Asp	Phe	Asn	Asp	Asn	Thr	Asn	Leu	Pro	Lys	Pro	Arg	Leu	
					200							205	

TCT	GAT	GAC	TAC	CAT	GGT	ACG	AGA	TGT	GCA	GGT	GAA	ATA	663
Ser	Asp	Asp	Tyr	His	Gly	Thr	Arg	Cys	Ala	Gly	Glu	Ile	
	210					215					220		
GCT	GCC	AAA	AAA	GGT	AAC	AAT	TTT	TGC	GGT	GTC	GGG	GTA	702
Ala	Ala	Lys	Lys	Gly	Asn	Asn	Phe	Cys	Gly	Val	Gly	Val	
			225					230					
GGT	TAC	AAC	GCT	AAA	ATC	TCA	GGC	ATA	AGA	ATC	TTA	TCC	741
Gly	Tyr	Asn	Ala	Lys	Ile	Ser	Gly	Ile	Arg	Ile	Leu	Ser	
235						240					245		
GGT	GAT	ATC	ACT	ACG	GAA	GAT	GAA	GCT	GCG	TCC	TTG	ATT	780
Gly	Asp	Ile	Thr	Thr	Glu	Asp	Glu	Ala	Ala	Ser	Leu	Ile	
		250					255					260	
TAT	GGT	CTA	GAC	GTA	AAC	GAT	ATA	TAT	TCA	TGC	TCA	TGG	819
Tyr	Gly	Leu	Asp	Val	Asn	Asp	Ile	Tyr	Ser	Cys	Ser	Trp	
				265					270				
GGT	CCC	GCT	GAT	GAC	GGA	AGA	CAT	TTA	CAA	GGC	CCT	AGT	858
Gly	Pro	Ala	Asp	Asp	Gly	Arg	His	Leu	Gln	Gly	Pro	Ser	
	275						280					285	
GAC	CTG	GTG	AAA	AAG	GCT	TTA	GTA	AAA	GGT	GTT	ACT	GAG	897
Asp	Leu	Val	Lys	Lys	Ala	Leu	Val	Lys	Gly	Val	Thr	Glu	
			290						295				
GGA	AGA	GAT	TCC	AAA	GGA	GCG	ATT	TAC	GTT	TTT	GCC	AGT	936
Gly	Arg	Asp	Ser	Lys	Gly	Ala	Ile	Tyr	Val	Phe	Ala	Ser	
300						305					310		
GGA	AAT	GGT	GGA	ACT	CGT	GGT	GAT	AAT	TGC	AAT	TAC	GAC	975
Gly	Asn	Gly	Gly	Thr	Arg	Gly	Asp	Asn	Cys	Asn	Tyr	Asp	
		315					320					325	

GGC	TAT	ACT	AAT	TCC	ATA	TAT	TCT	ATT	ACT	ATT	GGG	GCT	1014
Gly	Tyr	Thr	Asn	Ser	Ile	Tyr	Ser	Ile	Thr	Ile	Gly	Ala	
			330						335				
ATT	GAT	CAC	AAA	GAT	CTA	CAT	CCT	CCT	TAT	TCC	GAA	GGT	1053
Ile	Asp	His	Lys	Asp	Leu	His	Pro	Pro	Tyr	Ser	Glu	Gly	
	340					345					350		
TGT	TCC	GCC	GTC	ATG	GCA	GTC	ACG	TAT	TCT	TCA	GGT	TCA	1092
Cys	Ser	Ala	Val	Met	Ala	Val	Thr	Tyr	Ser	Ser	Gly	Ser	
			355					360					
GGC	GAA	TAT	ATT	CAT	TCG	AGT	GAT	ATC	AAC	GGC	AGA	TGC	1131
Gly	Glu	Tyr	Ile	His	Ser	Ser	Asp	Ile	Asn	Gly	Arg	Cys	
365					370					375			
AGT	AAT	AGC	CAC	GGT	GGA	ACG	TCT	GCG	GCT	GCT	CCA	TTA	1170
Ser	Asn	Ser	His	Gly	Gly	Thr	Ser	Ala	Ala	Ala	Pro	Leu	
	380					385					390		
GCT	GCC	GGT	GTT	TAC	ACT	TTG	TTA	CTA	GAA	GCC	AAC	CCA	1209
Ala	Ala	Gly	Val	Tyr	Thr	Leu	Leu	Leu	Glu	Ala	Asn	Pro	
			395					400					
AAC	CTA	ACT	TGG	AGA	GAC	GTA	CAG	TAT	TTA	TCA	ATC	TTG	1248
Asn	Leu	Thr	Trp	Arg	Asp	Val	Gln	Tyr	Leu	Ser	Ile	Leu	
	405					410				415			
TCT	GCG	GTA	GGG	TTA	GAA	AAG	AAC	GCT	GAC	GGA	GAT	TGG	1287
Ser	Ala	Val	Gly	Leu	Glu	Lys	Asn	Ala	Asp	Gly	Asp	Trp	
			420					425					
AGA	GAT	AGC	GCC	ATG	GGG	AAG	AAA	TAC	TCT	CAT	CGC	TAT	1326
Arg	Asp	Ser	Ala	Met	Gly	Lys	Lys	Tyr	Ser	His	Arg	Tyr	
430					435					440			

GGC	TTT	GGT	AAA	ATC	GAT	GCC	CAT	AAG	TTA	ATT	GAA	ATG	1365
Gly	Phe	Gly	Lys	Ile	Asp	Ala	His	Lys	Leu	Ile	Glu	Met	
		445				450					455		
TCC	AAG	ACC	TGG	GAG	AAT	GTT	AAC	GCA	CAA	ACC	TGG	TTT	1404
Ser	Lys	Thr	Trp	Glu	Asn	Val	Asn	Ala	Gln	Thr	Trp	Phe	
			460					465					
TAC	CTG	CCA	ACA	TTG	TAT	GTT	TCC	CAG	TCC	ACA	AAC	TCC	1449
Tyr	Leu	Pro	Thr	Leu	Tyr	Val	Ser	Gln	Ser	Thr	Asn	Ser	
	470					475					480		
ACG	GAA	GAG	ACA	TTA	GAA	TCC	GTC	ATA	ACC	ATA	TCA	GAA	1482
Thr	Glu	Glu	Thr	Leu	Glu	Ser	Val	Ile	Thr	Ile	Ser	Glu	
			485					490					
AAA	AGT	CTT	CAA	GAT	GCT	AAC	TTC	AAG	AGA	ATT	GAG	CAC	1521
Lys	Ser	Leu	Gln	Asp	Ala	Asn	Phe	Lys	Arg	Ile	Glu	His	
495					500					505			
GTC	ACG	GTA	ACT	GTA	GAT	ATT	GAT	ACA	GAA	ATT	AGG	GGA	1560
Val	Thr	Val	Thr	Val	Asp	Ile	Asp	Thr	Glu	Ile	Arg	Gly	
		510					515					520	
ACT	ACG	ACT	GTC	GAT	TTA	ATA	TCA	CCA	CCG	GGG	ATA	ATT	1599
Thr	Thr	Thr	Val	Asp	Leu	Ile	Ser	Pro	Ala	Gly	Ile	Ile	
			525						530				
TCA	AAC	CTT	GGC	GTT	GTA	AGA	CCA	AGA	GAT	GTT	TCA	TCA	1638
Ser	Asn	Leu	Gly	Val	Val	Arg	Pro	Arg	Asp	Val	Ser	Ser	
	535					540					545		
GAG	GGA	TTC	AAA	GAC	TGG	ACA	TTC	ATG	TCT	GTA	GCA	CAT	1677
Glu	Gly	Phe	Lys	Asp	Trp	Thr	Phe	Met	Ser	Val	Ala	His	
			550					555					

TGG	GGT	GAG	AAC	GGC	GTA	GGT	GAT	TGG	AAA	ATC	AAG	GTT	1716
Trp	Gly	Glu	Asn	Gly	Val	Gly	Asp	Trp	Lys	Ile	Lys	Val	
560					565						570		
AAG	ACA	ACA	GAA	AAT	GGA	CAC	AGG	ATT	GAC	TTC	CAC	AGT	1755
Lys	Thr	Thr	Glu	Asn	Gly	His	Arg	Ile	Asp	Phe	His	Ser	
		575					580					585	
TGG	AGG	CTG	AAG	CTC	TTT	GGG	GAA	TCC	ATT	GAT	TCA	TCT	1794
Trp	Arg	Leu	Lys	Leu	Phe	Gly	Glu	Ser	Ile	Asp	Ser	Ser	
				590						595			
AAA	ACA	GAA	ACT	TTC	GTC	TTT	GGA	AAC	GAT	AAA	GAG	GAG	1833
Lys	Thr	Glu	Thr	Phe	Val	Phe	Gly	Asn	Asp	Lys	Glu	Glu	
	600					605					610		
GTT	GAA	CCA	GGG	GTA	CCG	AGC	TCG	AAT	TCG	TAA			1866
Val	Glu	Pro	Gly	Val	Pro	Ser	Ser	Asn	Ser				
			615					620					

SEQ ID No. 2

Sequence type: Polynucleotide with corresponding polypeptide

Sequence length: 1797 base pairs

Strandedness: double

Topology: linear

Source: yeast genomic DNA

Immediate experimental source: E.coli HB101/KEX1 (DSM6027) Features: From 1 to 1797 coding region for soluble ysc α *

ATG	TTT	TAC	AAT	AGG	TGG	CTC	GGA	ACG	TGG	CTA	GCC	ATG	39
Met	Phe	Tyr	Asn	Arg	Trp	Leu	Gly	Thr	Trp	Leu	Ala	Met	
1				5					10				
TCT	GCT	TTA	ATA	AGG	ATC	TCA	GTA	TCC	CTT	CCG	TCA	TCT	78
Ser	Ala	Leu	Ile	Arg	Ile	Ser	Val	Ser	Leu	Pro	Ser	Ser	
	15				20						25		
GAG	GAG	TAC	AAA	GTG	GCA	TAT	GAG	CTG	TTG	CCA	GGG	TTA	117
Glu	Glu	Tyr	Lys	Val	Ala	Tyr	Glu	Leu	Leu	Pro	Gly	Leu	
			30					35					
TCT	GAA	GTG	CCA	GAC	CCT	TCA	AAT	ATT	CCA	CAG	ATG	CAT	156
Ser	Glu	Val	Pro	Asp	Pro	Ser	Asn	Ile	Pro	Gln	Met	His	
	40				45					50			
GCT	GGC	CAT	ATT	CCT	TTA	CGT	TCC	GAA	GAT	GCG	GAT	GAA	195
Ala	Gly	His	Ile	Pro	Leu	Arg	Ser	Glu	Asp	Ala	Asp	Glu	
		55				60					65		
CAG	GAT	AGC	TCT	GAC	TTG	GAG	TAC	TTT	TTT	TGG	AAG	TTT	234
Gln	Asp	Ser	Ser	Asp	Leu	Glu	Tyr	Phe	Phe	Trp	Lys	Phe	
				70				75					
ACC	AAT	AAC	GAC	TCT	AAT	GGC	AAT	GTC	GAC	CGT	CCC	TTA	279
Thr	Asn	Asn	Asp	Ser	Asn	Gly	Asn	Val	Asp	Arg	Pro	Leu	
	80					85					90		

ATT	ATA	TGG	TTA	AA	GGT	GGA	CCC	GGT	TGC	TCT	TCC	ATG	312
Ile	Ile	Trp	Leu	Asn	Gly	Gly	Pro	Gly	Cys	Ser	Ser	Met	
		95						100					
GAT	GGT	GCC	TTG	GTC	GAA	TCC	GGC	CCT	TTT	AGG	GTG	AAT	351
Asp	Gly	Ala	Leu	Val	Glu	Ser	Gly	Pro	Phe	Arg	Val	Asn	
105					110					115			
TCA	GAC	GGT	AAA	CTT	TAT	CTA	AAT	GAA	GGG	TCC	TGG	ATA	390
Ser	Asp	Gly	Lys	Leu	Tyr	Leu	Asn	Glu	Gly	Ser	Trp	Ile	
		120					125					130	
TCC	AAA	GGC	GAT	CTT	TTA	TTT	ATC	GAC	CAA	CCT	ACT	GGT	429
Ser	Lys	Gly	Asp	Leu	Leu	Phe	Ile	Asp	Gln	Pro	Thr	Gly	
				135					140				
ACT	GGG	TTT	TCT	GTC	GAA	CAA	AAT	AAA	GAC	GAA	GGT	AAA	468
Thr	Gly	Phe	Ser	Val	Glu	Gln	Asn	Lys	Asp	Glu	Gly	Lys	
	145					150					155		
ATC	GAC	AAA	AAC	AAA	TTT	GAC	GAA	GAC	CTA	GAA	GAT	GTG	507
Ile	Asp	Lys	Asn	Lys	Phe	Asp	Glu	Asp	Leu	Glu	Asp	Val	
			160					165					
ACC	AAG	CAT	TTT	ATG	GAT	TTT	CTG	GAG	AAC	TAT	TTC	AAG	546
Thr	Lys	His	Phe	Met	Asp	Phe	Leu	Glu	Asn	Tyr	Phe	Lys	
170					175					180			
ATT	TTT	CCA	GAA	GAC	CTC	ACT	AGG	AAA	ATC	ATA	CTA	TCG	585
Ile	Phe	Pro	Glu	Asp	Leu	Thr	Arg	Lys	Ile	Ile	Leu	Ser	
		185					190					195	
GGT	GAA	AGT	TAC	GCT	GGC	CAA	TAC	ATA	CCA	TTC	TTT	GCC	624
Gly	Glu	Ser	Tyr	Ala	Gly	Gln	Tyr	Ile	Pro	Phe	Phe	Ala	
				200					205				

AAT	GCA	ATT	TTG	AAC	CAT	AAC	AAA	TTT	TCA	AAG	ATC	GAC	663
Asn	Ala	Ile	Leu	Asn	His	Asn	Lys	Phe	Ser	Lys	Ile	Asp	
	210					215					220		
GGG	GAT	ACA	TAC	GAC	TTG	AAG	GCG	CTA	TTG	ATT	GGT	AAC	702
Gly	Asp	Thr	Tyr	Asp	Leu	Lys	Ala	Leu	Leu	Ile	Gly	Asn	
		225					230						
GGT	TGG	ATT	GAC	CCC	AAT	ACA	CAG	TCC	CTA	TCG	TAC	CTT	741
Gly	Trp	Ile	Asp	Pro	Asn	Thr	Gln	Ser	Leu	Ser	Tyr	Leu	
235					240					245			
CCG	TTT	GCT	ATG	GAG	AAG	AAA	CTG	ATT	GAT	GAA	AGC	AAC	780
Pro	Phe	Ala	Met	Glu	Lys	Lys	Leu	Ile	Asp	Glu	Ser	Asn	
	250					255					260		
CCC	AAT	TTC	AAA	CAC	TTA	ACG	AAC	GCA	CAC	GAG	AAT	TGC	819
Pro	Asn	Phe	Lys	His	Leu	Thr	Asn	Ala	His	Glu	Asn	Cys	
			265				270						
CAG	AAT	CTG	ATA	AAC	TCT	GCC	AGT	ACA	GAT	GAG	GCA	GCC	858
Gln	Asn	Leu	Ile	Asn	Ser	Ala	Ser	Thr	Asp	Glu	Ala	Ala	
	275					280				285			
CAT	TTT	TCG	TAT	CAG	GAA	TGT	GAG	AAT	ATT	TTA	AAC	CTT	897
His	Phe	Ser	Tyr	Gln	Glu	Cys	Glu	Asn	Ile	Leu	Asn	Leu	
		290					295						
CTA	TTG	TCT	TAT	ACC	AGG	GAA	TCT	TCG	CAA	AAG	GGA	ACA	936
Leu	Leu	Ser	Tyr	Thr	Arg	Glu	Ser	Ser	Gln	Lys	Gly	Thr	
300					305					310			
GCG	GAT	TGC	TTG	AAC	ATG	TAT	AAC	TTC	AAT	TTA	AAA	GAT	975
Ala	Asp	Cys	Leu	Asn	Met	Tyr	Asn	Phe	Asn	Leu	Lys	Asp	
	315						320				325		

AGT TAT CCA TCT TGT GGT ATG AAT TGG CCG AAA GAT ATT 1014
 Ser Tyr Pro Ser Cys Gly Met Asn Trp Pro Lys Asp Ile
 330 335

TCC TTT GTC AGT AAA TTC TTC AGC ACA CCT GGT GTT ATT 1053
 Ser Phe Val Ser Lys Phe Phe Ser Thr Pro Gly Val Ile
 340 345 350

GAT TCG TTG CAT CTT GAT TCT GAT AAA ATT GAT CAT TGG 1092
 Asp Ser Leu His Leu Asp Ser Asp Lys Ile Asp His Trp
 355 360

AAG GAA TGC ACT AAT AGC GTT GGA ACT AAA TTG TCT AAT 1131
 Lys Glu Cys Thr Asn Ser Val Gly Thr Lys Leu Ser Asn
 365 370 375

CCT ATT TCA AAG CCA TCT ATC CAT TTA TTA CCT GGT CTA 1170
 Pro Ile Ser Lys Pro Ser Ile His Leu Leu Pro Gly Leu
 380 385 390

CTT GAA AGT GGA ATA GAG ATT GTC TTG TTC AAT GGT GAC 1209
 Leu Glu Ser Gly Ile Glu Ile Val Leu Phe Asn Gly Asp
 395 400

AAA GAC TTG ATT TGT AAT AAC AAG GGC GTA TTA GAT ACT 1248
 Lys Asp Leu Ile Cys Asn Asn Lys Gly Val Leu Asp Thr
 405 410 415

ATA GAC AAT CTA AAA TGG GGT GGA ATA AAG GGA TTT AGC 1287
 Ile Asp Asn Leu Lys Trp Gly Gly Ile Lys Gly Phe Ser
 420 425

GAC GAT GCT GTT TCG TTC GAT TGG ATC CAT AAA TCG AAG 1326
 Asp Asp Ala Val Ser Phe Asp Trp Ile His Lys Ser Lys
 430 435 440

AGT	ACA	GAC	GAC	AGC	GAA	GAA	TTT	AGC	GGA	TAC	GTG	AAG	1365
Ser	Thr	Asp	Asp	Ser	Glu	Glu	Phe	Ser	Gly	Tyr	Val	Lys	
		445					450					455	
TAT	GAT	AGA	AAT	TTG	ACG	TTT	GTT	AGC	GTT	TAT	AAT	GCT	1404
Tyr	Asp	Arg	Asn	Leu	Thr	Phe	Val	Ser	Val	Tyr	Asn	Ala	
			460						465				
TCT	CAC	ATG	GTA	CCC	TTC	GAT	AAA	AGT	TTA	GTG	AGT	AGA	1443
Ser	His	Met	Val	Pro	Phe	Asp	Lys	Ser	Leu	Val	Ser	Arg	
	470					475					480		
GGC	ATT	GTC	GAT	ATT	TAC	TCG	AAC	GAT	GTT	ATG	ATC	ATT	1482
Gly	Ile	Val	Asp	Ile	Tyr	Ser	Asn	Asp	Val	Met	Ile	Ile	
			485					490					
GAC	AAC	AAT	GGG	AAA	AAT	GTT	ATG	ATT	ACT	ACT	GAC	GAC	1521
Asp	Asn	Asn	Gly	Lys	Asn	Val	Met	Ile	Thr	Thr	Asp	Asp	
495					500					505			
GAT	AGT	GAT	CAA	GAT	GCT	ACT	ACT	GAA	AGC	GGT	GAT	AAG	1560
Asp	Ser	Asp	Gln	Asp	Ala	Thr	Thr	Glu	Ser	Gly	Asp	Lys	
		510				515					520		
CCA	AAA	GAA	AAC	CTC	GAA	GAG	GAA	GAA	CAG	GAA	GCG	CAG	1599
Pro	Lys	Glu	Asn	Leu	Glu	Glu	Glu	Glu	Gln	Glu	Ala	Gln	
			525						530				
AAT	GAG	GAA	GGA	AAG	GAA	AAA	GAA	GGC	AAT	AAA	GAT	AAA	1638
Asn	Glu	Glu	Gly	Lys	Glu	Lys	Glu	Gly	Asn	Lys	Asp	Lys	
	535					540					545		
GAT	GGC	GAT	GAT	GAT	AAC	GAC	AAT	GAT	GAC	GAC	GAT	GAA	1677
Asp	Gly	Asp	Asp	Asp	Asn	Asp	Asn	Asp	Asp	Asp	Asp	Glu	
			550					555					

GAC GAT CAC AAC TCC GAG GGC GAC GAC GAT GAT GAC GAT . 1716
Asp Asp His Asn Ser Glu Gly Asp Asp Asp Asp Asp Asp
560 565 570

GAC GAT GAT GAA GAC GAT AAT AAT GAA AAA CAA AGT AAT 1755
Asp Asp Asp Glu Asp Asp Asn Asn Glu Lys Gln Ser Asn
575 580 585

CAA GGC CTC GAC TAC GTC GTA AGG CCG TTT CTG ACA GAG 1794
Gln Gly Leu Asp Tyr Val Val Arg Pro Phe Leu Thr Glu
590 595

TAA 1797

SEQ ID No. 3

Sequence type: Polynucleotide with corresponding polypeptide

Sequence length: 132 base pairs

Strandedness: double

Topology: linear

Immediate experimental source: *S. cerevisiae* HT246/pJDB207/GAPFL-HIR-CALC

Features: From 1 to 7 linker sequence including PstI restriction site and coding region for linker cleavable by yscF and ysc α^* , from 8 to 127 coding region for C-terminal glycine precursor of human calcitonin, from 125 to 132 sequence including PstI restriction site

G AAG AGG GTA CAG CTG GAT AAA AGA TGT GGT AAC TTG TCT	40
Lys Arg Val Gln Leu Asp Lys Arg Cys Gly Asn Leu Ser	
510	
ACC TGT ATG TTG GGT ACC TAC ACC CAA GAC TTC AAC AAG	79
Thr Cys Met Leu Gly Thr Thr Thr Gln Asp Phe Asn Lys	
152025	
TTC CAC ACC TTC CCA CAA ACC GCT ATC GGT GTT GGT GCT	118
Phe His Thr Phe Pro Gln Thr Ala Ile Gly Val Gly Ala	
3035	
CCA GGT TGA CTGCA	132
Pro Gly	
40	

SEQ ID No. 4

Sequence type: Polypeptide

Sequence length: 212 amino acids

Topology: linear

Immediate experimental source: S.cerevisiae AB110/pDP34GAPDH-eglinex-1

Features: From 1 to 71 amino acid sequence of eglin C, from 72 to 79 linker sequence,
from 80 to 212 amino acid sequence of the cellulose binding domain of C.fimi Exg protein

Met Thr Glu Phe Gly Ser Glu Leu Lys Ser Phe Pro Glu
5 10

Val Val Gly Lys Thr Val Asp Gln Ala Arg Glu Tyr Phe
15 20 25

Thr Leu His Tyr Pro Gln Tyr Asp Val Tyr Phe Leu Pro
30 35

Glu Gly Ser Pro Val Thr Leu Asp Leu Arg Tyr Asn Arg
40 45 50

Val Arg Val Phe Tyr Asn Pro Gly Thr Asn Val Val Asn
55 60 65

His Val Pro His Val Gly Lys Arg Glu Ala Glu Ala Trp
70 75

Val Pro Phe Gly Ala Ser Pro Thr Pro Thr Pro Thr Thr
80 85 90

Pro Thr Pro Thr Pro Thr Thr Pro Thr Pro Thr Pro Thr
95 100

Ser Gly Pro Ala Gly Cys Gln Val Leu Trp Gly Val Asn
105 110 115

Gln Trp Asn Thr Gly Phe Thr Ala Asn Val Thr Val Lys
120 125 130

Asn Thr Ser Ser Ala Pro Val Asp Gly Trp Thr Leu Thr
135 140

Phe Ser Phe Pro Ser Gly Gln Gln Val Thr Gln Ala Trp
145 150 155

Ser Ser Thr Val Thr Gln Ser Gly Ser Ala, Val Thr Val
160 165

Arg Asn Ala Pro Trp Asn Gly Ser Ile Pro Ala Gly Gly
170 175 180

Thr Ala Gln Phe Gly Phe Asn Gly Ser His Thr Gly Thr
185 190 195

Asn Ala Ala Pro Thr Ala Phe Ser Leu Asn Gly Thr Pro
200 205

Cys Thr Val Gly
210

Claims

1. A method for the production of a biologically active protein comprising treating a fusion protein consisting of

1. one or multiple successive protein segment(s) each consisting of said biologically active protein the C-terminal amino acid of which is joined to a linker polypeptide sequence L the N-terminal first and second and, in the case of multiple successive protein segments, also the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg, and

2. a polypeptide tag joined to the C-terminal amino acid of said successive protein segment(s),

or consisting of

1. a polypeptide tag joined to the N-terminal amino acid of

2. multiple successive protein segments each consisting of a linker polypeptide sequence L the N-terminal first and second as well as the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg and the ultimate basic amino acid of said linker polypeptide sequence L being joined to said biologically active protein,

with soluble yeast endoprotease yscF and with soluble yeast carboxypeptidase ysc α , and isolating said biologically active protein.

2. A method according to claim 1 in which the fusion protein has the formula

$$\begin{array}{ll} (P-L)_m-T & \text{(I) or} \\ T-(L-P)_n & \text{(II),} \end{array}$$

in which P is the biologically active protein, L has the meaning given in claim 1, T is a polypeptide tag, m is an integer from 1 to 10 and n is an integer from 2 to 10.

3. A method according to claim 1 in which the fusion protein has the formula



in which P, L and T have the meanings given above and m is 1.

4. A method according to claim 1 in which the biologically active protein is of prokaryotic or eukaryotic origin.

5. A method according to claim 1 in which the biologically active protein is of mammalian origin.

6. A method according to claim 1 in which the biologically active protein is a hormone, a polypeptide with immunomodulatory, anti-viral and anti-tumor properties, an antibody, viral antigen, blood clotting factor, a fibrinolytic agent or a growth regulation factor.

7. A method according to claim 1 in which the biologically active protein is selected from the group consisting of a human α interferon, a hybrid interferon, human tissue plasminogen activator, human single chain urokinase type plasminogen activator, a hybrid plasminogen activator, transforming growth factor β , human calcitonin, insulin-like growth factor I and II and desulfatohirudin.

8. A method according to claim 1 in which the linker polypeptide sequence L comprises 2 to about 20 amino acid residues and contains one or multiple pairs of basic amino acids, provided that the N-terminal first and second amino acids as well as the C-terminal penultimate and ultimate amino acids represent such a pair of basic amino acids.

9. A method according to claim 1 in which the linker polypeptide sequence L is a dipeptidyl radical selected from Arg-Arg, Lys-Arg, Lys-Lys and Arg-Lys.

10. A method according to claim 1 in which the polypeptide tag T consists of about 10 to about 1000 amino acid residues.

11. A method according to claim 1 in which m is an integer from 1 to 5.

12. A method according to claim 1 in which n is an integer from 2 to 5.

21489-8284

52

13. A method according to claim 1 in which soluble yscF is a mutein of yeast endoprotease yscF in which the hydrophobic membrane binding site has been deleted.

14. A method according to claim 1 in which soluble ysc α is a mutein of yeast carboxypeptidase ysc α in which the hydrophobic membrane binding site has been deleted.

15. A method according to claim 1 which is performed in a buffered solution at pH from about 6.0 to about 7.5 in a temperature range of from about 25°C to about 37°C.

16. A method according to claim 1 in which the digestion with soluble yscF is performed in the presence of Ca²⁺ ions.

17. A method according to claim 1 in which the fusion protein is treated with a digestive mixture containing both soluble yscF and soluble ysc α .

18. A method according to claim 1 in which the fusion protein is first treated with soluble yscF and, when the digestion has sufficiently proceeded or is complete, thereupon with soluble ysc α .

19. A method for the production of a biologically active protein, comprising:

culturing under appropriate nutrient conditions host cells transformed with an expression vector comprising an expression cassette containing a DNA sequence coding for a fusion protein, which consists of

(1) one or multiple successive protein segment(s) each consisting of said biologically active protein the C-terminal amino acid of which is joined to a linker polypeptide sequence L the N-terminal first and second and,

21489-8284

53

in the case of multiple successive protein segments, the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg, and

5 (2) a polypeptide tag joined to the C-terminal amino acid of said successive protein segment(s), or consisting of

(1) a polypeptide tag joined to the N-terminal amino acid of

10 (2) multiple successive protein segments each consisting of a linker polypeptide sequence L the N-terminal first and second as well as the C-terminal penultimate and ultimate amino acid residues of said linker polypeptide sequence L being basic amino acids selected from Lys and Arg
15 and the ultimate basic amino acid of said linker polypeptide sequence L being joined to said biologically active protein,

isolating said fusion protein,

treating said fusion protein with soluble yeast endoprotease yscF and with soluble yeast carboxypeptidase
20 ysc α , and

isolating said biologically active protein.

20. The method of claim 19 wherein the fusion protein is represented by the formula $(P-L)_m-T(I)$ or $T-(L-P)_n(II)$, in which P is the biologically active protein, L has the
25 meaning given in claim 18, T is a polypeptide tag, m is an integer from 1 to 10 and n is an integer from 2 to 10.

FETHERSTONHAUGH & CO.

OTTAWA, CANADA

PATENT AGENTS