

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 December 2007 (21.12.2007)

PCT

(10) International Publication Number
WO 2007/144669 A1

(51) International Patent Classification:
C07D 487/04 (2006.01) **A61P 9/10** (2006.01)
A61K 31/519 (2006.01)

(74) Agent: **HUSSAIN, Deeba**; European Patent Department,
Hoddesdon, Hertfordshire EN11 9BU (GB).

(21) International Application Number:
PCT/GB2007/050332

(22) International Filing Date: 12 June 2007 (12.06.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0611836.8 15 June 2006 (15.06.2006) GB
0701780.9 31 January 2007 (31.01.2007) GB

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): **ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI SPA** [IT/IT]; Via Pontina KM 30.600, I-00040 Pomezia (Rome) (IT).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **JONES, Philip** [GB/IT]; Irbm, Via Pontina KM 30.600, I-00040 Pomezia (Rome) (IT). **MURAGLIA, Ester** [IT/IT]; Irbm, Via Pontina KM 30.600, I-00040 Pomezia (Rome) (IT). **ONTORIA ONTORIA, Jesus Maria** [ES/IT]; Irbm, Via Pontina KM 30.600, I-00040 Pomezia (Rome) (IT).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRAZOLO[1,5-A]QUINAZOLIN-5(4H)-ONES AS INHIBITORS OF POLY(ADP-RIBOSE)POLYMERASE (PARP)



(I)

(57) Abstract: The present invention relates to compounds of formula (I): and pharmaceutically acceptable salts or tautomers thereof which are inhibitors of poly(ADP- ribose)polymerase (PARP) and thus useful for the treatment of cancer, inflammatory diseases, reperfusion injuries, ischemic conditions, stroke, renal failure, cardiovascular diseases, vascular diseases other than cardiovascular diseases, diabetes mellitus, neurodegenerative diseases, retroviral infections, retinal damage, skin senescence and UV-induced skin damage, and as chemo- or radiosensitizers for cancer treatment.

**Pyrazolo[1,5-*a*]quinazolin-5(4*H*)-ones as
inhibitors of poly(ADP-ribose)polymerase (PARP)**

The present invention relates to pyrazolo[1,5-*a*]quinazolin-5(4*H*)-ones which are inhibitors of the enzyme poly(ADP-ribose)polymerase (PARP), previously known as poly(ADP-ribose)synthase and poly(ADP-ribosyl)transferase. The compounds of the present invention are useful as mono-therapies in tumors with specific defects in DNA-repair pathways and as enhancers of certain DNA-damaging agents such as anticancer agents and radiotherapy. Further, the compounds of the present invention are useful for reducing cell necrosis (in stroke and myocardial infarction), down regulating inflammation and tissue injury, treating retroviral infections and protecting against the toxicity of chemotherapy.

Poly(ADP-ribose) polymerase (PARP) constitute a super family of eighteen proteins containing PARP catalytic domains (*Bioessays* (2004) **26**:1148). These proteins include PARP-1, PARP-2, PARP-3, tankyrase-1, tankyrase-2, vaultPARP and TiPARP. PARP-1, the founding member, consists of three main domains: an amino (N)-terminal DNA-binding domain (DBD) containing two zinc fingers, the automodification domain, and a carboxy (C)-terminal catalytic domain.

PARP are nuclear and cytoplasmic enzymes that cleave NAD^+ to nicotinamide and ADP-ribose to form long and branched ADP-ribose polymers on target proteins, including topoisomerases, histones and PARP itself (*Biochem. Biophys. Res. Commun.* (1998) **245**:1-10).

Poly(ADP-ribosyl)ation has been implicated in several biological processes, including DNA repair, gene transcription, cell cycle progression, cell death, chromatin functions and genomic stability.

The catalytic activity of PARP-1 and PARP-2 has been shown to be promptly stimulated by DNA strand breakages (see *Pharmacological Research* (2005) **52**:25-33). In response to DNA damage, PARP-1 binds to single and double DNA nicks. Under normal physiological conditions there is minimal PARP activity, however, upon DNA damage an immediate activation of PARP activity of up to 500-fold occurs. Both PARP-1 and PARP-2 detect DNA strand interruptions acting as nick sensors, providing rapid signals to halt transcription and recruiting the enzymes required for DNA repair at the site of damage. Since radiotherapy and many chemotherapeutic approaches to cancer therapy act by inducing DNA damage, PARP inhibitors are useful as chemo- and radiosensitizers for cancer treatment. PARP inhibitors have been reported to be effective in radio sensitizing hypoxic tumor cells (US 5,032,617, US 5,215,738 and US 5,041,653).

Most of the biological effects of PARP relate to this poly (ADP-ribosyl)ation process which influences the properties and function of the target proteins; to the PAR oligomers that, when cleaved from poly(ADP-ribosyl)ated proteins, confer distinct cellular effects; the physical

association of PARP with nuclear proteins to form functional complexes; and the lowering of the cellular level of its substrate NAD^+ (*Nature Review* (2005) **4**:421-440).

Besides being involved in DNA repair, PARP may also act as a mediator of cell death. Its excessive activation in pathological conditions such as ischemia and reperfusion injury can result in substantial depletion of the intercellular NAD^+ , which can lead to the impairment of several NAD^+ dependent metabolic pathways and result in cell death (see *Pharmacological Research* (2005) **52**:44-59). As a result of PARP activation, NAD^+ levels significantly decline. Extensive PARP activation leads to severe depletion of NAD^+ in cells suffering from massive DNA damage. The short half-life of poly(ADP-ribose) results in a rapid turnover rate, as once poly(ADP-ribose) is formed, it is quickly degraded by the constitutively active poly(ADP-ribose) glycohydrolase (PARG). PARP and PARG form a cycle that converts a large amount of NAD^+ to ADP-ribose, causing a drop of NAD^+ and ATP to less than 20% of the normal level. Such a scenario is especially detrimental during ischemia when deprivation of oxygen has already drastically compromised cellular energy output. Subsequent free radical production during reperfusion is assumed to be a major cause of tissue damage. Part of the ATP drop, which is typical in many organs during ischemia and reperfusion, could be linked to NAD^+ depletion due to poly(ADP-ribose) turnover. Thus, PARP inhibition is expected to preserve the cellular energy level thereby potentiating the survival of ischemic tissues after insult. Compounds which are inhibitors of PARP are therefore useful for treating conditions which result from PARP mediated cell death, including neurological conditions such as stroke, trauma and Parkinson's disease.

PARP inhibitors have been demonstrated as being useful for the specific killing of BRCA-1 and BRCA-2 deficient tumors (*Nature* (2005) **434**:913-916 and 917-921; and *Cancer Biology & Therapy* (2005) **4**:934-936).

PARP inhibitors have been demonstrated to be selective in killing cells with defects in ATM, DNA_PK or KU80 (*Nucleic Acid Research* (2006) **34**:1685-1691).

PARP inhibitors have been shown to enhance the efficacy of anticancer drugs (*Pharmacological Research* (2005) **52**:25-33), including platinum compounds such as cisplatin and carboplatin (*Cancer Chemother Pharmacol* (1993) **33**:157-162 and *Mol Cancer Ther* (2003) **2**:371-382). PARP inhibitors have been shown to increase the antitumor activity of topoisomerase I inhibitors such as Irinotecan and Topotecan (*Mol Cancer Ther* (2003) **2**:371-382; and *Clin Cancer Res* (2000) **6**:2860-2867) and this has been demonstrated in *in vivo* models (*J Natl Cancer Inst* (2004) **96**:56-67).

PARP inhibitors have been shown to restore susceptibility to the cytotoxic and antiproliferative effects of temozolomide (TMZ) (see *Curr Med Chem* (2002) **9**:1285-1301 and *Med Chem Rev Online* (2004) **1**:144-150). This has been demonstrated in a number of *in vitro* models (*Br J Cancer* (1995) **72**:849-856; *Br J Cancer* (1996) **74**:1030-1036; *Mol Pharmacol* (1997) **52**:249-258; *Leukemia* (1999) **13**:901-909; *Glia* (2002) **40**:44-54; and *Clin Cancer Res* (2000) **6**:2860-2867 and (2004) **10**:881-889) and *in vivo* models (*Blood* (2002) **99**:2241-2244;

Clin Cancer Res (2003) **9**:5370-5379 and *J Natl Cancer Inst* (2004) **96**:56-67). PARP inhibitors have also been shown to prevent the appearance of necrosis induced by selective N3-adenine methylating agents such as MeOSO₂(CH₂)-lexitropsin (Me-Lex) (*Pharmacological Research* (2005) **52**:25-33).

5 PARP inhibitors have been shown to act as radiation sensitizers. PARP inhibitors have been reported to be effective in radiosensitizing (hypoxic) tumor cells and effective in preventing tumor cells from recovering from potentially lethal (*Br. J. Cancer* (1984) **49**(Suppl. VI):34-42; and *Int. J. Radiat. Biol.* (1999) **75**:91-100) and sub-lethal (*Clin. Oncol.* (2004) **16**(1):29-39) damage of DNA after radiation therapy, presumably by their ability to prevent DNA strand break
10 rejoining and by affecting several DNA damage signaling pathways.

PARP inhibitors have also been shown to be useful for treating acute and chronic myocardial diseases (see *Pharmacological Research* (2005) **52**:34-43). For instance, it has been demonstrated that single injections of PARP inhibitors have reduced the infarct size caused by ischemia and reperfusion of the heart or skeletal muscle in rabbits. In these studies, a single
15 injection of 3-amino-benzamide (10 mg/kg), either one minute before occlusion or one minute before reperfusion, caused similar reductions in infarct size in the heart (32-42%) while 1,5-dihydroxyisoquinoline (1 mg/kg), another PARP inhibitor, reduced infarct size by a comparable degree (38-48%). These results make it reasonable to assume that PARP inhibitors could salvage previously ischemic heart or reperfusion injury of skeletal muscle tissue (*PNAS* (1997)
20 **94**:679-683). Similar findings have also been reported in pigs (*Eur. J. Pharmacol.* (1998) **359**:143-150 and *Ann. Thorac. Surg.* (2002) **73**:575-581) and in dogs (*Shock*. (2004) **21**:426-32).

PARP inhibitors have been demonstrated as being useful for treating certain vascular diseases, septic shock, ischemic injury and neurotoxicity (*Biochim. Biophys. Acta* (1989) **1014**:1-7; *J. Clin. Invest.* (1997) **100**: 723-735). Oxygen radical DNA damage that leads to strand
25 breaks in DNA, which are subsequently recognized by PARP, is a major contributing factor to such disease states as shown by PARP inhibitor studies (*J. Neurosci. Res.* (1994) **39**:38-46 and *PNAS* (1996) **93**:4688-4692). PARP has also been demonstrated to play a role in the pathogenesis of hemorrhagic shock (*PNAS* (2000) **97**:10203-10208).

PARP inhibitors have been demonstrated as being useful for treatment of inflammation
30 diseases (see *Pharmacological Research* (2005) **52**:72-82 and 83-92).

It has also been demonstrated that efficient retroviral infection of mammalian cells is blocked by the inhibition of PARP activity. Such inhibition of recombinant retroviral vector infections has been shown to occur in various different cell types (*J. Virology*, (1996) **70**(6):3992-4000). Inhibitors of PARP have thus been developed for use in anti-viral therapies
35 and in cancer treatment (WO 91/18591).

In vitro and *in vivo* experiments have demonstrated that PARP inhibitors can be used for the treatment or prevention of autoimmune diseases such as Type I diabetes and diabetic complications (*Pharmacological Research* (2005) **52**:60-71).

PARP inhibition has been speculated as delaying the onset of aging characteristics in human fibroblasts (*Biochem. Biophys. Res. Comm.* (1994) **201**(2):665-672 and *Pharmacological Research* (2005) **52**:93-99). This may be related to the role that PARP plays in controlling telomere function (*Nature Gen.*, (1999) **23**(1):76-80).


The vast majority of PARP inhibitors to date interact with the nicotinamide binding domain of the enzyme and behave as competitive inhibitors with respect to NAD^+ (*Expert Opin. Ther. Patents* (2004) **14**:1531-1551). Structural analogues of nicotinamide, such as benzamide and derivatives were among the first compounds to be investigated as PARP inhibitors. However, these molecules have a weak inhibitory activity and possess other effects unrelated to PARP inhibition. Thus, there is a need to provide potent inhibitors of the PARP enzyme.

EP 0374781 discloses pyrazolo[1,5-*a*]quinazolin-5(4*H*)-ones as couplers for photography.

Molecular Diversity (2003) **7**:161 and *J. Het. Chem.* (1969) **6**:947 describe methods for the synthesis of pyrazoloquinazolinones.

It has now surprisingly been discovered that pyrazolo(1,5-*a*)quinazolin-5(4*H*)-one derivatives of the present invention exhibit high levels of inhibition of the activity of PARP.

The compounds of this invention are useful in the inhibition of poly(ADP-ribose)polymerase (PARP). The present invention provides the use of a compound of formula I:



wherein:

a is 0, 1, 2, 3 or 4;

each b is independently 0, 1, 2 or 3;

c is 0 or 1;

d is 0, 1, 2, 3 or 4;

A is C_{3-10} cycloalkyl, C_{6-10} aryl, a 4 membered saturated heterocycle containing one N atom, a 5, 6 or 7 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic

ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, or a 7-13 membered unsaturated or partially saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S;

each R^1 is independently hydroxy, halogen, cyano, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy, carboxy, C_{1-6} alkoxycarbonyl, nitro, amino, C_{1-6} alkylamino or di(C_{1-6} alkyl)amino;

each Y is independently a direct bond, O, $O(C=O)$, $(C=O)O$, $C=O$, $NR^7(C=O)$, $(C=O)NR^7$, $S(O)_r$, $NR^7S(O)_r$ or $S(O)_rNR^7$;

each r is independently 0, 1 or 2;

each R^2 is independently hydrogen, hydroxy, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, halo C_{1-6} alkoxy, C_{2-10} alkenyl, C_{2-10} alkynyl, nitro, $N(R^a)_2$, $N(R^a)CON(R^a)_2$, $N(R^a)COOR^a$, $N(R^a)S(O)_rR^b$, $N(R^a)S(O)_rN(R^a)_2$ or a ring which is: C_{3-10} cycloalkyl, C_{6-10} aryl, C_{6-10} aryloxy, C_{6-10} arylcarbonyl, a 4 membered saturated heterocycle containing one N atom, a 5 or 6 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, or a 7, 8, 9 or 10 membered unsaturated, partially saturated or saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from $(CH_2)_eR^4$;

R^3 is halogen, cyano, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkoxy or halo C_{1-6} alkoxy;

each R^a is independently hydrogen, C_{1-6} alkyl, halo C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{6-10} aryl or C_{6-10} aryl C_{1-6} alkyl;

each R^b is independently C_{1-6} alkyl or C_{1-6} alkylcarbonyl;

e is 0, 1, 2, 3 or 4; and

R^4 is hydroxy, oxo, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, nitro, $N(R^a)_2$, $S(O)_rN(R^a)_2$, $CON(R^a)_2$, $N(R^a)COR^a$, $N(R^a)CON(R^a)_2$, $N(R^a)COOR^a$, $S(O)_rR^b$ or a ring which is: C_{3-10} cycloalkyl, C_{6-10} aryl, a 4 membered saturated heterocycle containing one N atom, a 5 or 6 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms or a 7, 8, 9 or 10 membered unsaturated, partially saturated or saturated heterocycle ring containing 1, 2 or 3 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from hydroxy, oxo, cyano, nitro, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy and $N(R^a)_2$;

R^7 is hydrogen, C_{1-6} alkyl or halo C_{1-6} alkyl;

each of R^8 and R^9 is independently hydrogen, C_{1-6} alkyl or C_{1-6} alkoxy;

or a pharmaceutically acceptable salt or tautomer thereof, for the manufacture of a medicament for the treatment or prevention of conditions which can be ameliorated by the inhibition of poly(ADP-ribose)polymerase (PARP).

The present invention also provides a method for the treatment or prevention of conditions which can be ameliorated by the inhibition of poly(ADP-ribose)polymerase (PARP), which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

In an embodiment, the conditions which can be ameliorated by the inhibition of poly(ADP-ribose)polymerase (PARP) include inflammatory diseases; reperfusion injuries; ischemic conditions; stroke; chronic and acute renal failure; vascular diseases other than cardiovascular diseases; cardiovascular diseases; diabetes mellitus; cancer, particularly cancer which is deficient in Homologous Recombination (HR) dependent DNA DSB repair activity, for example BRCA-1 or BRCA-2 deficient tumors; neurodegenerative diseases; retroviral infections; retinal damage; skin senescence; UV-induced skin damage; and premature aging.

In another embodiment is provided compounds of formula 1, or a pharmaceutically acceptable salt or tautomer thereof for the manufacture of a medicament for the treatment or prevention of reperfusion injuries; ischemic conditions; stroke; chronic and acute renal failure; vascular diseases other than cardiovascular diseases; cardiovascular diseases; diabetes mellitus; cancer, particularly cancer which is deficient in Homologous Recombination (HR) dependent DNA DSB repair activity, for example BRCA-1 or BRCA-2 deficient tumors; neurodegenerative diseases; retroviral infections; retinal damage; skin senescence; UV-induced skin damage; and premature aging.

In an embodiment:

each Y is independently a direct bond, O, $O(C=O)$, $(C=O)O$, $C=O$, $NH(C=O)$, $(C=O)NH$, $S(O)_r$, $NHS(O)_r$ or $S(O)_rNH$;

each R^2 is independently hydrogen, hydroxy, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, halo C_{1-6} alkoxy, C_{2-10} alkenyl, C_{2-10} alkynyl, nitro, $N(R^a)_2$, $N(R^a)CON(R^a)_2$, $N(R^a)S(O)_rR^b$ or a ring which is: C_{3-10} cycloalkyl, C_{6-10} aryl, C_{6-10} aryloxy, C_{6-10} arylcarbonyl, a 4 membered saturated heterocycle containing one N atom, a 5, 6 or 7 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, or a 7-10 membered unsaturated or partially saturated heterocyclic ring containing 1, 2 or 3 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from $(CH_2)_eR^4$;

R^4 is hydroxy, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, nitro, $N(R^a)_2$, $S(O)_rN(R^a)_2$, $CON(R^a)_2$, $N(R^a)CON(R^a)_2$ or $S(O)_rR^b$;

each of R^8 and R^9 is hydrogen; and

5 all other variables are as defined above.

The present invention also provides novel compounds of formula I wherein:

a, b, c, d, A, R^1 , R^2 , R^3 , R^8 , R^9 and Y are as defined above;

provided that:

- (a) when A is phenyl and either a is 0 or a is 1 and R^1 is methyl, bromine, fluorine, nitro or
10 carboxy, then d is not 0 and $Y-(CR^8R^9)_bR^2$ is not hydrogen, trifluoromethyl, chlorine or methoxy; and
- (b) when A is 2-thienyl or 2-furyl and either a is 0 or a is 1 and R^1 is methoxy, bromine, nitro or fluorine, then d is not 0 and $Y-(CR^8R^9)_bR^2$ is not hydrogen;

or a pharmaceutically acceptable salt or tautomer thereof.

15 In an embodiment, R^8 and R^9 are both hydrogen.

In an embodiment, when A is phenyl and a is 0 or 1, then d is not 0 and $Y-(CR^8R^9)_bR^2$ is not hydrogen, trifluoromethyl, chlorine or methoxy; and when A is 2-thienyl or 2-furyl and a is 0 or 1, then d is not 0 and $Y-(R^8R^9)_bR^2$ is not hydrogen.

In an embodiment a is 0.

20 In an embodiment b is 0.

In an embodiment c is 0. In another embodiment c is 1.

In an embodiment d is 0, 1, 2 or 3.

In an embodiment A is C_{3-10} cycloalkyl, C_{6-10} aryl, a 5 membered heteroaromatic ring containing 1, 2, 3 and 4 heteroatoms independently selected from N, O and S, not more than one
25 heteroatom of which is O or S or a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms.

In another embodiment A is a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms.

Preferably, A is phenyl, furyl, thienyl, pyridinyl, naphthyl, pyrazolyl or cyclohexyl.

30 In an embodiment A is not thienyl. In another embodiment A is not furyl.

More specifically A, as attached to the fused pyrazole ring is phenyl, 2-furyl, 2-thienyl, 3-furyl, pyridin-3-yl, pyridin-4-yl, 1-naphthyl, pyrazol-4-yl or cyclohexyl. A further specific A group is 2-naphthyl.

In an embodiment A is not 2-thienyl. In another embodiment A is not 2-furyl.

35 In an embodiment Y is a direct bond, O, C=O or (C=O)O. Further Y groups are (C=O)NH and (C=O)NCH₃.

In an embodiment R^2 is hydroxy, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, halo C_{1-6} alkoxy or a ring which is: C_{6-10} aryl or a 5, 6 or 7 membered saturated or partially saturated heterocycle

containing one, two or three N atoms and zero or one O atom, any of which rings being optionally substituted by one or more groups independently selected from $(\text{CH}_2)_e\text{R}^4$.

In an embodiment R^2 is hydroxy, cyano, halogen, halo C_{1-6} alkyl, halo C_{1-6} alkoxy or a ring which is phenyl, morpholinyl, diazepanyl, piperazinyl, octahydropyrrolopyrazinyl, piperidinyl, imidazolyl, pyrrolidinyl or hexahydropyrrolopyrrolyl; any of which rings being optionally substituted by one, two or three groups independently selected from $(\text{CH}_2)_e\text{R}^4$.

In an embodiment e is 0, 1 or 2.

In an embodiment, when R^4 is a ring it is optionally substituted by one, two or three independently selected groups. More particularly, when R^4 is a ring it is unsubstituted or monosubstituted.

In an embodiment R^4 is oxo, halogen, C_{1-6} alkyl, $\text{N}(\text{R}^a_2)$, C_{3-7} cycloalkyl, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S or a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, any of which rings being optionally substituted by one, two or three groups independently selected from halogen, C_{1-6} alkyl and halo C_{1-6} alkyl.

Particular R^4 groups are methyl, methylpyridinyl, methylthiazolyl, cyclopropyl, fluorine, oxo, dimethylamino and chlorine.

In an embodiment, when R^2 is a ring it is optionally substituted by one, two or three independently selected $(\text{CH}_2)_e\text{R}^4$ groups. Preferably, when R^2 is a ring it is unsubstituted, monosubstituted or disubstituted. More particularly when R^2 is a ring it is unsubstituted.

Preferably, R^2 is methyl, chlorine, fluorine, bromine, ethyl, isopropyl, trifluoromethoxy, trifluoromethyl, phenyl, cyano, morpholinyl or hydroxy. Further particular R^2 groups are methyldiazepanyl, methylpiperazinyl, dimethylamino, octahydropyrrolopyrazinyl, [(methylpyridinyl)methyl]piperazinyl, [(methylthiazolyl)methyl]piperazinyl, (cyclopropylmethyl)piperazinyl, difluorodiazepanyl, [(dimethylamino)ethyl](oxo)diazepanyl, [(dimethylamino)methyl]piperidinyl, methylimidazolyl, methylpyrrolidinyl, methylhexahydropyrrolopyrrolyl, (dichlorophenyl)piperazinyl, diazepanyl and (chlorophenyl)piperazinyl.

More specifically, R^2 is methyl, chlorine, fluorine, bromine, ethyl, isopropyl, trifluoromethoxy, trifluoromethyl, phenyl, cyano, morpholin-4-yl or hydroxy. Further specific R^2 groups are 4-methyl-1,4-diazepan-1-yl, 4-methylpiperazin-1-yl, dimethylamino, (8*aS*)octahydropyrrolo[1,2-*a*]pyrazin-2-yl, 4-[(3-methylpyridin-2-yl)methyl]piperazin-1-yl, 4-[(2-methyl-1,3-thiazol-4-yl)methyl]piperazin-1-yl, 4-(cyclopropylmethyl)piperazin-1-yl, 6,6-difluoro-1,4-diazepan-1-yl, 4-[(dimethylamino)ethyl]-5-oxo-1,4-diazepan-1-yl, 3-[(dimethylamino)methyl]piperidin-1-yl, 1-methyl-1*H*-imidazol-2-yl, 1-methylpyrrolidin-3-yl, 1-methyl-1,2,4,5,6,6*a*-hexahydropyrrolo[3,4-*b*]pyrrol-5-yl, 4-(2,4-dichlorophenyl)piperazin-1-yl, 1,4-diazepan-1-yl and 4-(4-chlorophenyl)piperazin-1-yl.

In an embodiment R^3 is C_{1-6} alkyl.

A particular R^3 group is methyl.


Particular R^7 groups are hydrogen and methyl.

In an embodiment each of R^8 and R^9 is independently selected from hydrogen and C_{1-6} alkyl.

5 In an embodiment R^8 is hydrogen or methyl and R^9 is hydrogen.

In an embodiment d is 1, 2, 3 or 4 and $Y-(CR^8R^9)_bR^2$ is not hydrogen, trifluoromethyl, chlorine or methoxy.

The present invention also provides compounds of formula II:



(II)

wherein:

a, b, c, A, R^1 , R^3 , R^8 and R^9 are as defined above;

f is 0, 1 or 2;

15 g is 1 or 2;

R^5 is independently hydroxy, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, halo C_{1-6} alkoxy, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, carboxy, nitro, $N(R^a)_2$, $S(O)_rN(R^a)_2$, $CON(R^a)_2$, $N(R^a)CON(R^a)_2$, $S(O)_rR^b$ or $N(R^a)S(O)_rR^b$;

20 each R^a is independently hydrogen, C_{1-6} alkyl, halo C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{6-10} aryl or C_{6-10} aryl C_{1-6} alkyl;

each R^b is independently C_{1-6} alkyl or C_{1-6} alkylcarbonyl;

Y^1 is a direct bond, $O(C=O)$, $(C=O)O$, $C=O$, $NR^7(C=O)$, $(C=O)NR^7$, $S(O)_r$, $NHR^7(O)_r$ or $S(O)_rNR^7$;

r is 0, 1 or 2;

25 R^6 is hydroxy, cyano, fluorine, bromine, iodine, C_{1-6} alkyl, C_{1-6} alkylcarbonyl, halo C_{1-6} alkoxy, C_{2-10} alkenyl, C_{2-10} alkynyl, halo C_{1-6} alkoxy, nitro, $N(R^a)_2$, $N(R^a)CON(R^a)_2$, $N(R^a)S(O)_rR^b$ or a ring which is: C_{3-10} cycloalkyl, C_{6-10} aryl, C_{6-10} aryloxy, C_{6-10} arylcarbonyl, 4 membered


saturated heterocycle containing one N atom, or a 5 or 6 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one O atom, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, or a 7, 8, 9 or 10 membered unsaturated, partially saturated or saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from $(\text{CH}_2)_e\text{R}^4$;

e is 0, 1, 2, 3 or 4; and

R^4 is hydroxy, oxo, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, nitro, $\text{N}(\text{R}^a)_2$, $\text{S}(\text{O})_r\text{N}(\text{R}^a)_2$, $\text{CON}(\text{R}^a)_2$, $\text{N}(\text{R}^a)\text{CON}(\text{R}^a)_2$, $\text{S}(\text{O})_r\text{R}^b$ or a ring which is: C_{3-10} cycloalkyl, C_{6-10} aryl, a 4 membered saturated heterocycle containing one N atom, a 5 or 6 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms or a 7, 8, 9 or 10 membered unsaturated, partially saturated or saturated heterocycle ring containing 1, 2 or 3 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from hydroxy, oxo, cyano, nitro, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy and $\text{N}(\text{R}^a)_2$;

or a pharmaceutically acceptable salt or tautomer thereof.

The present invention also provides compounds of formula III:



(III)

wherein:

a, b, c, f, g, R^1 , R^3 , R^5 , R^6 , R^8 , R^9 and Y^1 are as defined above;
or a pharmaceutically acceptable salt or tautomer thereof.

In an embodiment of compounds of formula II or III:

Y^1 is a direct bond, $O(C=O)$, $(C=O)O$, $C=O$, $NH(C=O)$, $(C=O)NH$, $S(O)_r$, $NHS(O)_r$ or

5 $S(O)_rNH$;

R^6 is hydroxy, cyano, fluorine, bromine, iodine, C_{1-6} alkyl, C_{1-6} alkylcarbonyl, halo C_{1-6} alkoxy, C_{2-10} alkenyl, C_{2-10} alkynyl, halo C_{1-6} alkoxy, nitro, $N(R^a)_2$, $N(R^a)CON(R^a)_2$, $N(R^a)S(O)_rR^b$ or a ring which is: C_{3-10} cycloalkyl, C_{6-10} aryl, C_{6-10} aryloxy, C_{6-10} arylcarbonyl, 4 membered saturated heterocycle containing one N atom, or a 5, 6 or 7 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one O atom, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, or a 7-10 membered unsaturated or partially saturated heterocyclic ring containing 1, 2 or 3 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from $(CH_2)_eR^4$;

R^4 is hydroxy, oxo, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, nitro, $N(R^a)_2$, $S(O)_rN(R^a)_2$, $CON(R^a)_2$, $N(R^a)CON(R^a)_2$ or $S(O)_rR^b$;

each of R^8 and R^9 is hydrogen; and

20 all other variables are as defined above.

The preferred identities with reference to formulae II and III are as defined previously for formula I *mutatis mutandis*.

In an embodiment g is 1.

In an embodiment each R^5 is independently halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, or
25 C_{1-6} alkoxy.

Specific R^5 groups are fluorine, methyl, methoxy and trifluoromethyl.

In an embodiment Y^1 is a direct bond, $C=O$ or $(C=O)O$. Further Y^1 groups are $(C=O)NH$ and $(C=O)NCH_3$.

In an embodiment R^6 is hydroxy, cyano, fluorine, bromine, iodine, C_{1-6} alkyl, halo C_{1-6} alkoxy or a ring which is: C_{6-10} aryl or a 5, 6 or 7 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one O atom, any of which rings being optionally substituted by one or more groups independently selected from $(CH_2)_eR^4$.

In an embodiment, when R^6 is a ring it is optionally substituted by one, two or three independently selected $(CH_2)_eR^4$ groups. Preferably, when R^2 is a ring it is unsubstituted, monosubstituted or disubstituted. More particularly when R^6 is a ring it is unsubstituted.
35

Preferably, R^6 is methyl, fluorine, bromine, iodine, ethyl, isopropyl, trifluoromethoxy, phenyl, cyano, morpholinyl or hydroxy. Further preferred R^6 groups are methyldiazepanyl, methylpiperazinyl, dimethylamino, octahydropyrrolopyrazinyl,

[(methylpyridinyl)methyl]piperazinyl, [(methylthiazolyl)methyl]piperazinyl, (cyclopropylmethyl)piperazinyl, difluorodiazepanyl, [(dimethylamino)ethyl](oxo)diazepanyl, [(dimethylamino)methyl]piperidinyl, methylimidazolyl, methylpyrrolidinyl, methylhexahydropyrrolopyrrolyl, (dichlorophenyl)piperazinyl, diazepanyl and (chlorophenyl)piperazinyl.

Specific R⁶ groups are methyl, fluorine, bromine, iodine, ethyl, isopropyl, trifluoromethoxy, phenyl, cyano, morpholin-4-yl, hydroxy, 4-methyl-1,4-diazepan-1-yl, 4-methylpiperazin-1-yl, dimethylamino, (8*aS*)octahydropyrrolo[1,2-*a*]pyrazin-2-yl, 4-[(3methylpyridin-2-yl)methyl]piperazin-1-yl, 4-[(2-methyl-1,3-thiazol-4-yl)methyl]piperazin-1-yl, 4-(cyclopropylmethyl)piperazin-1-yl, 6,6-difluoro-1,4-diazepan-1-yl, 4-[(dimethylamino)ethyl]-5-oxo-1,4-diazepan-1-yl, 3-[(dimethylamino)methyl]piperidin-1-yl, 1-methyl-1*H*-imidazol-2-yl, 1-methylpyrrolidin-3-yl, 1-methyl-1,2,4,5,6,6*a*-hexahydropyrrolo[3,4-*b*]pyrrol-5-yl, 4-(2,4-dichlorophenyl)piperazin-1-yl, 1,4-diazepan-1-yl and 4-(4-chlorophenyl)piperazin-1-yl.

The present invention also includes within its scope N-oxides of the compounds of formula I above. In general, such N-oxides may be formed on any available nitrogen atom. The N-oxides may be formed by conventional means, such as reacting the compound of formula I with oxone in the presence of wet alumina.

The present invention includes within its scope prodrugs of the compounds of formula I above. In general, such prodrugs will be functional derivatives of the compounds of formula I which are readily convertible in vivo into the required compound of formula I. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.


A prodrug may be a pharmacologically inactive derivative of a biologically active substance (the "parent drug" or "parent molecule") that requires transformation within the body in order to release the active drug, and that has improved delivery properties over the parent drug molecule. The transformation in vivo may be, for example, as the result of some metabolic process, such as chemical or enzymatic hydrolysis of a carboxylic, phosphoric or sulphate ester, or reduction or oxidation of a susceptible functionality.

The present invention includes within its scope solvates of the compounds of formula I and salts thereof, for example, hydrates.

The compounds of the present invention may have asymmetric centers, chiral axes, and chiral planes (as described in: E.L. Eliel and S.H. Wilen, *Stereochemistry of Carbon Compounds*, John Wiley & Sons, New York, 1994, pages 1119-1190), and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers, all such stereoisomers being included in the present invention.

The compounds disclosed herein may exist as tautomers and both tautomeric forms are intended to be encompassed by the scope of the invention, even though only one tautomeric

structure is depicted. For example, compounds of formula I may tautomerise into compounds of the following structure:



5

The compounds may exist in different isomeric forms, all of which are encompassed by the present invention.

The compounds may exist in a number of different polymorphic forms.

When any variable (e.g. R^1 and R^2 , etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every other occurrence. Also, combinations of substituents and variables are permissible only if such combinations result in stable compounds. Lines drawn into the ring systems from substituents represent that the indicated bond may be attached to any of the substitutable ring atoms.

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials. If a substituent is itself substituted with more than one group, it is understood that these multiple groups may be on the same carbon or on different carbons, so long as a stable structure results. The phrase “optionally substituted” should be taken to be equivalent to the phrase “unsubstituted or substituted with one or more substituents” and in such cases the preferred embodiment will have from zero to three substituents. More particularly, there are zero to two substituents. A substituent on a saturated, partially saturated or unsaturated heterocycle can be attached at any substitutable position.

As used herein, “alkyl” is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, “ C_{1-6} alkyl” is defined to include groups having 1, 2, 3, 4, 5 or 6 carbons in a linear or branched arrangement. For example, “ C_{1-6} alkyl” specifically includes methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, i-butyl, pentyl, hexyl, and so on. Preferred alkyl groups are methyl and ethyl. The term “cycloalkyl” means a monocyclic, bicyclic or polycyclic saturated aliphatic hydrocarbon group having the specified number of carbon atoms. For example, “ C_{3-10} cycloalkyl” includes

30

cyclopropyl, methyl-cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, cyclohexyl, and so on. In an embodiment of the invention the term "cycloalkyl" includes the groups described immediately above and further includes monocyclic unsaturated aliphatic hydrocarbon groups. For example, "cycloalkyl" as defined in this embodiment includes cyclopropyl, methyl-

5 cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, cyclohexyl, cyclopentenyl, cyclobutenyl, 7,7-dimethylbicyclo[2.2.1]heptyl and so on. Preferred cycloalkyl groups are cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

As used herein, the term " C_{2-10} alkenyl" refers to a non-aromatic hydrocarbon radical, straight or branched, containing from 2 to 10 carbon atoms and at least one carbon to carbon

10 double bond. Preferably one carbon to carbon double bond is present, and up to four non-aromatic carbon-carbon double bonds may be present. Alkenyl groups include ethenyl, propenyl, butenyl and 2-methylbutenyl. Preferred alkenyl groups include ethenyl and propenyl.

As used herein, the term " C_{2-10} alkynyl" refers to a hydrocarbon radical straight or branched, containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond.

15 Up to three carbon-carbon triple bonds may be present. Alkynyl groups include ethynyl, propynyl, butynyl, 3-methylbutynyl and so on. Preferred alkynyl groups include ethynyl and propynyl.

"Alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge. "Alkoxy" therefore encompasses the definitions of alkyl above.

20 Examples of suitable alkoxy groups include methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy and t-butoxy. The preferred alkoxy groups are methoxy and ethoxy.

The terms "halo C_{1-6} alkyl" and "halo C_{1-6} alkoxy" mean a C_{1-6} alkyl or C_{1-6} alkoxy group in which one or more (in particular, 1 to 3) hydrogen atoms have been replaced by halogen atoms, especially fluorine or chlorine atoms. Preferred are fluoro C_{1-6} alkyl and fluoro C_{1-6} alkoxy groups,

25 in particular fluoro C_{1-3} alkyl and fluoro C_{1-3} alkoxy groups, for example, CF_3 , CHF_2 , CH_2F , CH_2CH_2F , CH_2CHF_2 , CH_2CF_3 , OCF_3 , $OCHF_2$, OCH_2F , OCH_2CH_2F , OCH_2CHF_2 or OCH_2CF_3 , and most especially CF_3 , OCF_3 and $OCHF_2$.

As used herein, the term "hydroxy C_{1-6} alkyl" means a C_{1-6} alkyl group in which one or more (in particular, 1 to 3) hydrogen atoms have been replaced by hydroxy groups. Preferred

30 are CH_2OH , CH_2CHOH and $CHOHCH_3$.

As used herein, the term " C_{1-6} alkylcarbonyl" or " C_{1-6} alkoxycarbonyl" denotes a C_{1-6} alkyl or C_{1-6} alkoxy radical, respectively, attached via a carbonyl ($C=O$) radical. Suitable examples of C_{1-6} alkylcarbonyl groups include methylcarbonyl, ethylcarbonyl, propylcarbonyl, isopropylcarbonyl and *tert*-butylcarbonyl. Examples of C_{1-6} alkoxycarbonyl include

35 methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl and *tert*-butoxycarbonyl.

The rings present in the compounds of this invention may be monocyclic or multicyclic, particularly bicyclic. The multicyclic rings may be fused or spiro linked.

As used herein, "C₆₋₁₀aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of 6 to 10 atoms, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydronaphthyl, indanyl and tetrahydrobenzo[7]annulene. The preferred aryl group is phenyl or naphthyl, especially phenyl.

5 7-13 membered heterocycles include 7, 8, 9, 10, 11, 12 and 13 membered heterocycles.

Examples of particular heterocycles of this invention are benzimidazolyl, benzofurandionyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothienyl, benzoxazolyl, benzoxazolonyl, benzothiazolyl, benzothiadiazolyl, benzodioxolyl, benzoxadiazolyl, benzoisoxazolyl, benzoisothiazolyl, chromenyl, chromanyl, isochromanyl, carbazolyl, carbolinyl, cinnoliny, epoxidyl, furyl, furazanyl, imidazolyl, indoliny, indolyl, indoliziny, indoliny, isoindoliny, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridiny, oxadiazolyl, oxazolyl, oxazoliny, isoxazoliny, oxetanyl, puriny, pyranyl, pyraziny, pyrazolyl, pyridaziny, pyridopyridiny, pyridaziny, pyridiny, pyrimidiny, triaziny, tetraziny, pyrroly, quinazoliny, quinoliny, quinoxaliny, quinoliziny, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydroisoquinoliny, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, azetidiny, 1,4-dioxanyl, hexahydroazepiny, piperaziny, piperidyl, pyridin-2-onyl, pyrrolidiny, imidazoliny, imidazolidiny, pyrazoliny, pyrroliny, morpholiny, thiomorpholiny, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyraziny, dihydropyrazolyl, dihydropyridiny, dihydropyrimidiny, dihydropyrroly, dihydroquinoliny, dihydroisoquinoliny, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl, dihydroazetidiny, dihydroisochromenyl, dihydroimidazolonyl, dihydrotriazolonyl, dihydrobenzodioxiny, dihydrothiazolopyrimidiny, dihydroimidazopyraziny, methylenedioxybenzoyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydroquinoliny, thiazolidinonyl, imidazolonyl, isoindolinonyl, octahydroquinoliziny, octahydroisoindolyl, imidazopyridiny, azabicycloheptanyl, chromenonyl, triazolopyrimidiny, dihydrobenzoxaziny, thiazolotriazolyl, azoniabicycloheptanyl, azoniabicyclooctanyl, phthalaziny, naphthyridiny, quinazoliny, pteridiny, dihydroquinazoliny, dihydropthalaziny, benzisoxazolyl, tetrahydronaphthyridiny, dibenzo[*b,d*]furanyl, dihydrobenzothiazolyl, imidazothiazolyl, tetrahydroindazolyl, tetrahydrobenzothienyl, hexahydronaphthyridiny, tetrahydroimidazopyridiny, tetrahydroimidazopyraziny, pyrrolopyridiny, diazepamyl and N-oxides thereof. Further rings include octahydropyrrolopyraziny, hexahydropyrrolopyrroly, diazaspirononany, azepamyl, oxazepamyl, octahydroindoliziny, diazoniaspiro[4.4]nonany, hexahydropyrrolo[3,4-*b*]pyrroly, oxaazoniabicyclo[2.2.1]heptanyl, diazoniaspiro[3.3]heptanyl, diazoniaspiro[3.5]nonany, diazoniaspiro[4.5]decanyl, octahydropyrrolo[3,4-*c*]pyrroly, octahydropyrrolo[3,4-*b*]pyrroly, octahydrocyclopenta[*c*]pyrroly, dihydroindolyl, azoniaspiro[4.5]decanyl, diazoniabicyclo[2.2.2]octanyl, diazoniabicyclo[2.2.1]heptanyl,

10
15
20
25
30
35

diazoniabicyclo[3.2.1]octanyl, diazoniabicyclo[2.2.1]heptanyl and oxaazoniaspiro[4.5]decanyl and N-oxides thereof. Attachment of a heterocyclyl substituent can occur via a carbon atom or via a heteroatom.

A preferred 4 membered saturated heterocycle is azetidiny.

5 Preferred 5 or 6 membered saturated or partially saturated heterocycles are pyrrolidiny, piperidiny, piperaziny, morpholiny, tetrahydrofuran and thiomorpholiny.

A preferred 7 membered saturated heterocycle is diazepany. Further 7 membered saturated heterocycles are azepany and oxazepany.

10 Preferred 5 membered heteroaromatic rings are thienyl, thiazolyl, pyrazolyl, isoxazolyl, imidazolyl, thiadiazolyl, oxazolyl, oxadiazolyl, triazolyl, furyl, pyrrolyl and tetrazolyl.

Preferred 6 membered heteraromatic ring are pyridiny, pyrimidiny, pyridaziny and pyraziny.

Preferred 7-10 membered partially saturated or unsaturated heterocyclic rings are tetrahydroquinoliny, quinoliny, indolyl, imidazopyridiny, benzothiazolyl, quinoxaliny, 15 benzothiadiazolyl, benzoxazolyl, dihydrobenzodioxiny, benzotriazolyl, benzodioxolyl, dihydroisoindolyl, dihydroindolyl, tetrahydroisoquinoliny, isoquinoliny, benzoisothiazolyl, dihydroimidazopyraziny, benzothienyl, benzoxadiazolyl, thiazolotriazolyl, dihydrothiazolopyrimidiny, dihydrobenzoxaziny, dihydrobenzofurany, benzimidazolyl, benzofurany, dihydrobenzoxazolyl, dihydroquinazoliny, dihydrophthalaziny, indazolyl, 20 benzisoxazolyl, tetrahydronaphthyridiny, triazolopyrimidiny, dibenzo[*b,d*]furany, naphthyridiny, dihydroquinoliny, dihydroisochromenyl, dihydrochromenyl, dihydrobenzothiazolyl, imidazothiazolyl, tetrahydroindazolyl, tetrahydrobenzothienyl, hexahydronaphthyridiny, tetrahydroimidazopyridiny, tetrahydroimidazopyraziny, pyrrolopyridiny, quinazoliny and indoliziny. A further ring is hexahydropyrrolopyraziny.

25 A preferred 9 membered saturated heterocycle is octahydropyrrolopyraziny.

As used herein, the term "halogen" refers to fluorine, chlorine, bromine and iodine, of which fluorine and chlorine are preferred.

Particular compounds within the scope of the present invention are:

2-(2-methylphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
 30 2-(3-chlorophenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
 2-(3-fluorophenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
 2-(3-bromophenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
 2-(3-methylphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one
 2-(4-fluorophenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
 35 2-(4-bromophenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
 2-(4-methylphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
 2-(4-ethylphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
 2-(4-isopropoxyphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;

- 2-[4-(trifluoromethoxy)phenyl]pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
methyl 4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoate;
2-(3,4-difluorophenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(3,4-dimethoxyphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
5 2-(3,4-dimethylphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(3,5-dimethoxyphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(5-fluoro-2-methylphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(2,5-difluorophenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(2,3-dimethoxyphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
10 2-(2,4-dimethoxyphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(3,4,5-trimethoxyphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(4-methoxy-3,5-dimethylphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-[5-methyl-2-(trifluoromethyl)-3-furyl]pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(2-methyl-3-furyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
15 3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)pyridinium trifluoroacetate;
4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)pyridinium trifluoroacetate;
3-methyl-2-phenylpyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-biphenyl-2-ylpyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(1-naphthyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
20 4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzonitrile;
2-[1-methyl-5-(trifluoromethyl)-1*H*-pyrazol-4-yl]pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(3-fluoro-4-morpholin-4-ylphenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoic acid;
2-cyclohexylpyrazolo[1,5-*a*]quinazolin-5(4*H*)-one; and
25 pharmaceutically acceptable salts or tautomers thereof.

Further particular compounds within the scope of the present invention are:

- 1-methyl-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium
trifluoroacetate;
4-(3-{[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}propyl)morpholin-4-
30 ium trifluoroacetate;
1-methyl-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]piperazin-1-ium
trifluoroacetate;
N,N-dimethyl-2-{methyl[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-
yl)benzoyl]amino}ethanaminium trifluoroacetate;
35 (8*aS*)-2-[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]octahydropyrrolo[1,2-
a]pyrazin-5-ium trifluoroacetate;
1-[(3-methylpyridinium-2-yl)methyl]-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-
yl)benzoyl]piperazin-1-ium bis(trifluoroacetate);

- 1-[(2-methyl-1,3-thiazol-3-ium-4-yl)methyl]-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]piperazin-1-ium bis(trifluoroacetate);
- 1-(cyclopropylmethyl)-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate;
- 5 6,6-difluoro-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium trifluoroacetate;
- N,N*-dimethyl-2-{7-oxo-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-yl}ethanaminium trifluoroacetate;
- N,N*-dimethyl{1-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]piperidin-3-yl}methanaminium trifluoroacetate;
- 10 1-methyl-2-({methyl[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}methyl)-1*H*-imidazol-3-ium trifluoroacetate;
- 1-methyl-3-({[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}methyl)pyrrolidinium trifluoroacetate;
- 15 *N,N*-dimethyl-2-{[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}propan-1-aminium trifluoroacetate;
- 1-(cyclopropylmethyl)-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate;
- (8*aS*)-2-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]octahydropyrrolo[1,2-*a*]pyrazin-5-ium trifluoroacetate;
- 20 1-methyl-5-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]-1,2,4,5,6,6*a*-hexahydropyrrolo[3,4-*b*]pyrrol-1-ium trifluoroacetate;
- 4-(2,4-dichlorophenyl)-1-(2-{[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}ethyl)piperazin-1-ium trifluoroacetate;
- 25 1-methyl-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium trifluoroacetate;
- 1-methyl-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate;
- N,N*-dimethyl-2-{methyl[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}ethanaminium trifluoroacetate;
- 30 4-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium trifluoroacetate;
- 4-(2-{[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}ethyl)morpholin-4-ium trifluoroacetate;
- 35 1-(4-chlorophenyl)-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate; and
- pharmaceutically acceptable salts or tautomers thereof.

In an embodiment is provided PARP inhibitors selected from:

2-phenylpyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(4-chlorophenyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(2-furyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one;
2-(2-thienyl)pyrazolo[1,5-*a*]quinazolin-5(4*H*)-one; and

5 pharmaceutically acceptable salts or tautomers thereof.

Included in the instant invention is the free base of compounds of Formula I, as well as the pharmaceutically acceptable salts and stereoisomers thereof. Some of the specific compounds exemplified herein are the protonated salts of amine compounds. Compounds of Formula I with a heterocycle ring containing 2 or more N atoms may be protonated on any one,
10 some or all of the N atoms. The term "free base" refers to the amine compounds in non-salt form. The encompassed pharmaceutically acceptable salts not only include the salts exemplified for the specific compounds described herein, but also all the typical pharmaceutically acceptable salts of the free form of compounds of Formula I. The free form of the specific salt compounds described may be isolated using techniques known in the art. For example, the free form may be
15 regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia and sodium bicarbonate. The free forms may differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise pharmaceutically equivalent to their respective free forms for purposes of the invention.

20 The pharmaceutically acceptable salts of the instant compounds can be synthesized from the compounds of this invention which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts of the basic compounds are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various
25 combinations of solvents. Similarly, the salts of the acidic compounds are formed by reactions with the appropriate inorganic or organic base.

Thus, pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed by reacting a basic instant compound with an inorganic, organic acid or polymeric acid. For example, conventional
30 non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, hydroiodic, sulfuric, sulfurous, sulfamic, phosphoric, phosphorous, nitric and the like, as well as salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic, fumaric, toluenesulfonic, methanesulfonic, ethane
35 disulfonic, oxalic, isethionic, palmitic, gluconic, ascorbic, phenylacetic, aspartic, cinnamic, pyruvic, ethanesulfonic, ethane, disulfonic, valeric, trifluoroacetic and the like. Examples of suitable polymeric salts include those derived from the polymeric acids such as tannic acid, carboxymethyl cellulose. Preferably, a pharmaceutically acceptable salt of this invention

contains 1 equivalent of a compound of formula (I) and 1, 2 or 3 equivalent of an inorganic or organic acid. More particularly, pharmaceutically acceptable salts of this invention are the trifluoroacetate or the chloride salts, especially the trifluoroacetate salts.

When the compound of the present invention is acidic, suitable "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as arginine, lysine, betaine caffeine, choline, N,N¹-dibenzylethylenediamine, ethylamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, diethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine tripropylamine, tromethamine, dicyclohexylamine, butylamine, benzylamine, phenylbenzylamine, tromethamine and the like.

The preparation of the pharmaceutically acceptable salts described above and other typical pharmaceutically acceptable salts is more fully described by Berg *et al* (1977) *J. Pharm. Sci.*, 'Pharmaceutical Salts', **66**:1-19.

It will also be noted that the compounds of the present invention are potentially internal salts or zwitterions, since under physiological conditions a deprotonated acidic moiety in the compound, such as a carboxyl group, may be anionic, and this electronic charge might then be balanced off internally against the cationic charge of a protonated or alkylated basic moiety, such as a quaternary nitrogen atom.

The compounds of the invention can be used in a method of treatment of the human or animal body by therapy.

The invention provides compounds for use in the treatment or prevention of conditions which can be ameliorated by the inhibition of poly(ADP-ribose)polymerase (PARP).

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of conditions which can be ameliorated by the inhibition of poly(ADP-ribose)polymerase (PARP).

The present invention also provides a method for the treatment or prevention of conditions which can be ameliorated by the inhibition of poly(ADP-ribose)polymerase (PARP), which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The PARP inhibitors of the present invention are useful for the treatment of the diseases specified in WO 2005/082368.

The compounds of the invention are useful for the treatment of inflammatory diseases, including conditions resulting from organ transplant rejection, such as; chronic inflammatory diseases of the joints, including arthritis, rheumatoid arthritis, osteoarthritis and bone diseases associated with increased bone resorption; inflammatory bowel diseases such as ileitis, ulcerative colitis, Barrett's syndrome, and Crohn's disease; inflammatory lung diseases such as asthma, adult respiratory distress syndrome, and chronic obstructive airway disease; inflammatory diseases of the eye including corneal dystrophy, trachoma, onchocerciasis, uveitis, sympatheticophthalmitis and endophthalmitis; chronic inflammatory diseases of the gum, including gingivitis and periodontitis; tuberculosis; leprosy; inflammatory diseases of the kidney including uremic complications, glomerulonephritis and nephrosis; inflammatory diseases of the skin including sclerodermatitis, psoriasis and eczema; inflammatory diseases of the central nervous system, including chronic demyelinating diseases of the nervous system, multiple sclerosis, AIDS-related neurodegeneration and Alzheimer's disease, infectious meningitis, encephalomyelitis, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis and viral or autoimmune encephalitis; diabetic complications, including, but not limited to, immune-complex vasculitis, systemic lupus erythematosus (SLE); inflammatory diseases of the heart such as cardiomyopathy, ischemic heart disease, hypercholesterolemia, and atherosclerosis; as well as various other diseases that can have significant inflammatory components, including preeclampsia, chronic liver failure, brain and spinal cord trauma and multiple organ dysfunction syndrome (MODS) (multiple organ failure (MOF)). The inflammatory disease can also be a systemic inflammation of the body, exemplified by gram-positive or gram negative shock, hemorrhagic or anaphylactic shock, or shock induced by cancer chemotherapy in response to pro-inflammatory cytokines, e. g., shock associated with pro-inflammatory cytokines. Such shock can be induced, e. g. by a chemotherapeutic agent that is administered as a treatment for cancer.

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for treating or preventing inflammatory diseases.

The present invention also provides a method for the treatment or prevention of inflammatory diseases, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of the instant invention may also be useful in the treatment or prevention of reperfusion injuries, resulting from naturally occurring episodes and during a surgical procedure, such as intestinal reperfusion injury; myocardial reperfusion injury; reperfusion injury resulting from cardiopulmonary bypass surgery, aortic aneurysm repair surgery, carotid

endarterectomy surgery, or hemorrhagic shock; and reoxygenation injury resulting from transplantation of organs such as heart, lung, liver, kidney, pancreas, intestine, and cornea.

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of reperfusion injuries.

5 The present invention also provides a method for the treatment or prevention of reperfusion injuries, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

10 The compounds of the instant invention may also be useful in the treatment or prevention of ischemic conditions, including those resulting from organ transplantation, such as stable angina, unstable angina, myocardial ischemia, hepatic ischemia, mesenteric artery ischemia, intestinal ischemia, critical limb ischemia, chronic critical limb ischemia, cerebral ischemia, acute cardiac ischemia, ischemia kidney disease, ischemic liver disease, ischemic retinal disorder, septic shock, and an ischemic disease of the central nervous system, such as stroke or
15 cerebral ischemia.

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of ischemic conditions.

The present invention also provides a method for the treatment or prevention of ischemic conditions, which method comprises administration to a patient in need thereof of an effective
20 amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of the instant invention may also be useful for the treatment or prevention of chronic or acute renal failure

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of renal failure.

25 The present invention also provides a method for the treatment or prevention of renal failure, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of the instant invention may also be useful for the treatment or prevention of vascular diseases other than cardiovascular diseases, such as peripheral arterial
30 occlusion, thromboangitis obliterans, Reynaud's disease and phenomenon, acrocyanosis, erythromelalgia, venous thrombosis, varicose veins, arteriovenous fistula, lymphedema and lipedema.

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of vascular diseases other than cardiovascular
35 diseases.

The present invention also provides a method for the treatment or prevention of vascular diseases other than cardiovascular diseases, which method comprises administration to a patient

in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of the instant invention may also be useful for the treatment or prevention of cardiovascular diseases such as chronic heart failure, atherosclerosis, congestive heart failure, circulatory shock, cardiomyopathy, cardiac transplant, myocardial infarction, and a cardiac arrhythmia, such as atrial fibrillation, supraventricular tachycardia, atrial flutter, and paroxysmal atrial tachycardia.

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of cardiovascular diseases.

The present invention also provides a method for the treatment or prevention of cardiovascular diseases, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of this invention may also be useful for the treatment and prevention of diabetes mellitus, including Type I diabetes (Insulin Dependent Diabetes Mellitus), Type II diabetes (Non-Insulin Dependent Diabetes Mellitus), gestational diabetes, autoimmune diabetes, insulinopathies, diabetes due to pancreatic disease, diabetes associated with other endocrine diseases (such as Cushing's Syndrome, acromegaly, pheochromocytoma, glucagonoma, primary aldosteronism or somatostatinoma), Type A insulin resistance syndrome, Type B insulin resistance syndrome, lipatrophic diabetes, and diabetes induced by (3-cell toxins. The compounds of this invention may also be useful for the treatment or prevention of diabetic complications, such as diabetic cataract, glaucoma, retinopathy, nephropathy, (such as microalbuminuria and progressive diabetic nephropathy), polyneuropathy, gangrene of the feet, atherosclerotic coronary arterial disease, peripheral arterial disease, nonketotic hyperglycemic-hyperosmolar coma, mononeuropathies, autonomic neuropathy, foot ulcers, joint problems, and a skin or mucous membrane complication (such as an infection, a shin spot, a candidal infection or necrobiosis lipoidica diabetorum obesity), hyperlipidemia, hypertension, syndrome of insulin resistance, coronary artery disease, retinopathy, diabetic neuropathy, polyneuropathy, mononeuropathies, autonomic neuropathy, a foot ulcer, a joint problem, a fungal infection, a bacterial infection, and cardiomyopathy.

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of diabetes.

The present invention also provides a method for the treatment or prevention of diabetes, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of this invention may also be useful for the treatment or prevention of cancer including solid tumors such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma,

lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon cancer, colorectal cancer, kidney cancer, pancreatic cancer, bone cancer, breast cancer, ovarian cancer, prostate cancer, esophageal cancer, stomach cancer, oral cancer, nasal cancer, throat cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms'tumor, cervical cancer, uterine cancer, testicular cancer, small cell lung carcinoma, bladder carcinoma, lung cancer, epithelial carcinoma, skin cancer, melanoma, neuroblastoma and retinoblastoma; blood-borne cancers such as acute lymphoblastic leukemia("ALL"), acute lymphoblastic B-cell leukemia, acute lymphoblastic T-cell leukemia, acute myeloblastic leukemia ("AML"), acute promyelocytic leukemia("APL"), acute monoblastic leukemia, acute erythroleukemic leukemia, acute megakaryoblastic leukemia, acute myelomonocytic leukemia, acute nonlymphocytic leukemia, acute undifferentiated leukemia, chronic myelocytic leukemia("CML"), chronic lymphocytic leukemia("CLL"), hairy cell leukemia and multiple myeloma; acute and chronic leukemias such as lymphoblastic, myelogenous, lymphocytic, myelocytic leukemias; Lymphomas such as Hodgkin's disease, non-Hodgkin's Lymphoma, Multiple myeloma, Waldenstrom's macroglobulinemia, Heavy chain disease and Polycythemia vera; CNS and brain cancers such as glioma, pilocytic astrocytoma, astrocytoma, anaplastic astrocytoma, glioblastoma multiforme, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, vestibular schwannoma, adenoma, metastatic brain tumor, meningioma, spinal tumor and medulloblastoma.

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of cancer.

The present invention also provides a method for the treatment or prevention of cancer, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of the present invention may also be used for the treatment of cancer which is deficient in Homologous Recombination (HR) dependent DNA DSB repair (See WO 2004/080976).

BRCA-1 and BRCA-2 are known tumor suppressors whose wild-type alleles are frequently lost in tumors of heterozygous carriers (Jasin M. *Oncogene*, (2002) **21(58)**:8981-93; Tutt *et al*, *Trends Mol Med.*, (2002) **8(12)**:571-6). The association of BRCA-1 and/or BRCA-2 mutations with breast cancer has been well-characterized (Radice PJ. *Exp Clin Cancer Res.*, (2002) **21 (3 Suppl)**:9-12). Amplification of the EMSY gene, which encodes a BRCA-2 binding factor, is also known to be associated with breast and ovarian cancer. Carriers of mutations in BRCA-1 and/or BRCA-2 are also at elevated risk of cancer of the ovary, prostate and pancreas.

The detection of variation in BRCA-1 and BRCA-2 is well-known in the art and is described, for example in EP 699 754, EP 705 903, Neuhausen SL. and Ostrander EA. *Genet. Test* (1992) **1**:75-83; Chappnis, PO. *et al.*, *Cancer Treat Res* (2002) **107**:29-59; Janatova M. *et al.*, *Neoplasia* (2003) **50(4)**:246-50; Jancarkova N. *et al.*, (2003) **6(1)**:11-16). Determination of amplification of the BRCA-2 binding factor EMSY is described in Hughes-Davies *et al.*, *Cell* **115**:523-535. PARP inhibitors have been demonstrated as being useful for the specific killing of BRCA-1 and BRCA-2 deficient tumors (*Nature* (2005) **434**:913-916 and 917-920).

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for the treatment or prevention of BRCA-1 or BRCA-2 deficient tumors.

The present invention also provides a method for the treatment or prevention of BRCA-1 or BRCA-2 deficient tumors, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of this invention may be useful for the treatment or prevention of neurodegenerative diseases, including, polyglutamine-expansion-related neurodegeneration, Huntington's disease, Kennedy's disease, spinocerebellar ataxia, dentatorubral-pallidoluysian atrophy (DRPLA), protein-aggregation-related neurodegeneration, Machado-Joseph's disease, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, spongiform encephalopathy, a prion-related disease and multiple sclerosis (MS).

Thus, the present invention provides a compound of formula I for use in the manufacture of a medicament for treating or preventing neurodegenerative diseases.

The present invention also provides a method for treating or preventing neurodegenerative diseases, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I.

The compounds of the present invention may also be useful for the treatment or prevention of retroviral infection (US 5652260), retinal damage (Oku H. *et al Curr. Eye Res.* (2004), **29**:403), skin senescence and UV-induced skin damage (US5589483 and Farkas B *et al Biochem. Pharmacol* (2002) **63**:921).

The compounds of the invention are useful for the treatment or prevention of premature aging and postponing the onset of age-related cellular dysfunction (*Pharmacological Research* (2005) **52**:93-99).

The compounds of this invention may be administered to mammals, preferably humans, either alone or in combination with pharmaceutically acceptable carriers, excipients, diluents, adjuvants, fillers, buffers, stabilisers, preservatives, lubricants, in a pharmaceutical composition, according to standard pharmaceutical practice.

The compounds of this invention may be administered to a subject by any convenient route of administration, whether systemically/peripherally or at the site of desired action,

including but not limited to, oral (e.g. by ingestion); topical (including e.g. transdermal, intranasal, ocular, buccal, and sublingual); pulmonary (e.g. by inhalation or insufflation therapy using, e.g. an aerosol, e.g. through mouth or nose); rectal; vaginal; parenteral, (e.g. by injection, including subcutaneous, intradermal, intramuscular, intravenous, intraarterial, intracardiac, intrathecal, intraspinal, intracapsular, subcapsular, intraorbital, intraperitoneal, intratracheal, subcuticular, intraarticular, subarachnoid, and intrasternal); and by implant of a depot (e.g. subcutaneously or intramuscularly).

The subject may be a eukaryote, an animal, a vertebrate animal, a mammal, a rodent (e.g. a guinea pig, a hamster, a rat, a mouse), murine (e.g. a mouse), canine (e.g. a dog), feline (e.g. a cat), equine (e.g. a horse), a primate, simian (e.g. a monkey or ape), a monkey (e.g. marmoset, baboon), an ape (e.g. gorilla, chimpanzee, orangutang, gibbon), or a human.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, microcrystalline cellulose, sodium crosscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to mask the unpleasant taste of the drug or delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water soluble taste masking material such as hydroxypropyl-methylcellulose or hydroxypropylcellulose, or a time delay material such as ethyl cellulose, cellulose acetate butyrate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium

alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more
10 flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents
15 may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated hydroxyanisol or alpha-tocopherol.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents
20 and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a
25 mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally occurring phosphatides, for example soy bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavoring
30 agents, preservatives and antioxidants.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, flavoring and coloring agents and antioxidant.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous
35 solutions. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

The sterile injectable preparation may also be a sterile injectable oil-in-water microemulsion where the active ingredient is dissolved in the oily phase. For example, the

active ingredient may be first dissolved in a mixture of soybean oil and lecithin. The oil solution then introduced into a water and glycerol mixture and processed to form a microemulsion.

The injectable solutions or microemulsions may be introduced into a patient's blood stream by local bolus injection. Alternatively, it may be advantageous to administer the solution or microemulsion in such a way as to maintain a constant circulating concentration of the instant compound. In order to maintain such a constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension for intramuscular and subcutaneous administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Compounds of Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compound of Formula I are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Compounds of the present invention may also be delivered as a suppository employing bases such as cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

When a compound according to this invention is administered into a subject, the selected dosage level will depend on a variety of factors including, but not limited to, the activity of the particular compound, the severity of the individuals symptoms, the route of administration, the time of administration, the rate of excretion of the compound, the duration of the treatment, other

drugs, compounds, and/or materials used in combination, and the age, sex, weight, condition, general health, and prior medical history of the patient. The amount of compound and route of administration will ultimately be at the discretion of the physician, although generally the dosage will be to achieve local concentrations at the site of action which achieve the desired effect without causing substantial harmful or deleterious side-effects.

Administration in vivo can be effected in one dose, continuously or intermittently (e.g. in divided doses at appropriate intervals) throughout the course of treatment. Methods of determining the most effective means and dosage of administration are well known to those of skill in the art and will vary with the formulation used for therapy, the purpose of the therapy, the target cell being treated, and the subject being treated. Single or multiple administrations can be carried out with the dose level and pattern being selected by the treating physician.

In general, a suitable dose of the active compound is in the range of about 100 µg to about 250 mg per kilogram body weight of the subject per day. Where the active compound is a salt, an ester, prodrug, or the like, the amount administered is calculated on the basis of the parent compound and so the actual weight to be used is increased proportionately.

The instant compounds are also useful in combination with anti-cancer agents or chemotherapeutic agents.

The compounds of this invention may be useful as chemo- and radiosensitizers for cancer treatment. They are useful for the treatment of mammals who have previously undergone or are presently undergoing treatment for cancer. Such previous treatments include prior chemotherapy, radiation therapy, surgery or immunotherapy, such as cancer vaccines.

Thus, the present invention provides a combination of a compound of formula I and an anti-cancer agent for simultaneous, separate or sequential administration.

The present invention also provides a compound of formula I for use in the manufacture of a medicament for use as an adjunct in cancer therapy or for potentiating tumor cells for treatment with ionizing radiation or chemotherapeutic agents.

The present invention also provides a method of chemotherapy or radiotherapy, which method comprises administration to a patient in need thereof of an effective amount of a compound of formula I or a composition comprising a compound of formula I in combination with ionizing radiation or chemotherapeutic agents.

Examples of cancer agents or chemotherapeutic agents for use in combination with the compounds of the present invention can be found in *Cancer Principles and Practice of Oncology* by V.T. Devita and S. Hellman (editors), 6th edition (February 15, 2001), Lippincott Williams & Wilkins Publishers. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the cancer involved. Such anti-cancer agents include, but are not limited to, the following: HDAC inhibitors, estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic/cytostatic agents, antiproliferative agents, prenyl-protein transferase

inhibitors, HMG-CoA reductase inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors and other angiogenesis inhibitors, inhibitors of cell proliferation and survival signaling, apoptosis inducing agents and agents that interfere with cell cycle checkpoints. The instant compounds are particularly useful when co-administered with radiation therapy.

5 Examples of “HDAC inhibitors” include suberoylanilide hydroxamic acid (SAHA), LAQ824, LBH589, PXD101, MS275, FK228, valproic acid, butyric acid and CI-994.

“Estrogen receptor modulators” refers to compounds that interfere with or inhibit the binding of estrogen to the receptor, regardless of mechanism. Examples of estrogen receptor modulators include, but are not limited to, tamoxifen, raloxifene, idoxifene, LY353381,
10 LY117081, toremifene, fulvestrant, 4-[7-(2,2-dimethyl-1-oxopropoxy-4-methyl-2-[4-(1-piperidinyl)ethoxy]phenyl)-2*H*-1-benzopyran-3-yl]-phenyl-2,2-dimethylpropanoate, 4,4'-dihydroxybenzophenone-2,4-dinitrophenyl-hydrazone, and SH646.

“Androgen receptor modulators” refers to compounds which interfere or inhibit the binding of androgens to the receptor, regardless of mechanism. Examples of androgen receptor
15 modulators include finasteride and other 5 α -reductase inhibitors, nilutamide, flutamide, bicalutamide, liarozole, and abiraterone acetate.

“Retinoid receptor modulators” refers to compounds which interfere or inhibit the binding of retinoids to the receptor, regardless of mechanism. Examples of such retinoid receptor modulators include bexarotene, tretinoin, 13-cis-retinoic acid, 9-cis-retinoic acid, α -
20 difluoromethylornithine, ILX23-7553, trans-*N*-(4'-hydroxyphenyl) retinamide, and *N*-4-carboxyphenyl retinamide.

“Cytotoxic/cytostatic agents” refer to compounds which cause cell death or inhibit cell proliferation primarily by interfering directly with the cell's functioning or inhibit or interfere with cell mytosis, including alkylating agents, tumor necrosis factors, intercalators, hypoxia
25 activatable compounds, microtubule inhibitors/microtubule-stabilizing agents, inhibitors of mitotic kinesins, inhibitors of kinases involved in mitotic progression, antimetabolites; biological response modifiers; hormonal/anti-hormonal therapeutic agents, haematopoietic growth factors, monoclonal antibody targeted therapeutic agents, topoisomerase inhibitors, proteasome inhibitors and ubiquitin ligase inhibitors.

30 Examples of cytotoxic agents include, but are not limited to, sertenef, cachectin, ifosfamide, tasonermin, lonidamine, carboplatin, altretamine, prednimustine, dibromodulcitol, ranimustine, fotemustine, nedaplatin, oxaliplatin, temozolomide, heptaplatin, estramustine, improsulfan tosilate, trofosfamide, nimustine, dibrospidium chloride, pumitepa, lobaplatin, satraplatin, profiromycin, cisplatin, irofulven, dexifosfamide, cis-aminedichloro(2-methyl-
35 pyridine)platinum, benzylguanine, glufosfamide, GPX100, (trans, trans, trans)-bis-mu-(hexane-1,6-diamine)-mu-[diamine-platinum(II)]bis[diamine(chloro)platinum (II)]tetrachloride, diarizidinylspermine, arsenic trioxide, 1-(11-dodecylamino-10-hydroxyundecyl)-3,7-dimethylxanthine, zorubicin, idarubicin, daunorubicin, bisantrene, mitoxantrone, pirarubicin,

pinafide, valrubicin, amrubicin, antineoplaston, 3'-deamino-3'-morpholino-13-deoxo-10-hydroxycarminomycin, annamycin, galarubicin, elinafide, MEN10755, and 4-demethoxy-3-deamino-3-aziridiny-4-methylsulphonyl-daunorubicin (see WO 00/50032).

An example of a hypoxia activatable compound is tirapazamine.

5 Examples of proteasome inhibitors include but are not limited to lactacystin, bortezomib, epoxomicin and peptide aldehydes such as MG 132, MG 115 and PSI.

In an embodiment, the compounds of the present invention may be used in combination with HDAC inhibitors such as SAHA and proteasome inhibitors.

10 Examples of microtubule inhibitors/microtubule-stabilising agents include paclitaxel, vindesine sulfate, 3',4'-didehydro-4'-deoxy-8'-norvincal leukoblastine, docetaxol, rhizoxin, dolastatin, mivobulin isethionate, auristatin, cemadotin, RPR109881, BMS184476, vinflunine, cryptophycin, 2,3,4,5,6-pentafluoro-*N*-(3-fluoro-4-methoxyphenyl) benzene sulfonamide, anhydrovinblastine, *N,N*-dimethyl-L-valyl-L-valyl-*N*-methyl-L-valyl-L-prolyl-L-proline-*t*-butylamide, TDX258, the epothilones (see for example U.S. Pat. Nos. 6,284,781 and 6,288,237) and BMS188797.

Some examples of topoisomerase inhibitors are topotecan, hycaptamine, irinotecan, rubitecan, 6-ethoxypropionyl-3',4'-*O*-exo-benzylidene-chartreusin, 9-methoxy-*N,N*-dimethyl-5-nitropyrazolo[3,4,5-*kl*]acridine-2-(6*H*) propanamine, 1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1*H*,12*H*-benzo[*de*]pyrano[3',4':*b*,7]-indolizino[1,2*b*]quinoline-10,13(9*H*,15*H*)dione, lurtotecan, 7-[2-(*N*-isopropylamino)ethyl]-(2*S*)camptothecin, BNP1350, BNPI1100, BN80915, BN80942, etoposide phosphate, teniposide, sobuzoxane, 2'-dimethylamino-2'-deoxy-etoposide, GL331, *N*-[2-(dimethylamino)ethyl]-9-hydroxy-5,6-dimethyl-6*H*-pyrido[4,3-*b*]carbazole-1-carboxamide, asulacrine, (5*a*, 5*aB*, 8*aa*,9*b*)-9-[2-[*N*-[2-(dimethylamino)ethyl]-*N*-methylamino]ethyl]-5-[4-hydroxy-3,5-dimethoxyphenyl]-5,5*a*,6,8,8*a*,9-hexahydrofuro(3',4':6,7)naphtho(2,3-*d*)-1,3-dioxol-6-one, 2,3-(methylenedioxy)-5-methyl-7-hydroxy-8-methoxybenzo[*c*]-phenanthridinium, 6,9-bis[(2-aminoethyl)amino]benzo[*g*]isoguinoline-5,10-dione, 5-(3-aminopropylamino)-7,10-dihydroxy-2-(2-hydroxyethylaminomethyl)-6*H*-pyrazolo[4,5,1-*de*]acridin-6-one, *N*-[1-[2(diethylamino)ethylamino]-7-methoxy-9-oxo-9*H*-thioxanthen-4-ylmethyl]formamide, *N*-(2-(dimethylamino)ethyl)acridine-4-carboxamide, 6-[[2-(dimethylamino)ethyl]amino]-3-hydroxy-7*H*-indeno[2,1-*c*]quinolin-7-one, and dimesna.

35 Examples of inhibitors of mitotic kinesins, and in particular the human mitotic kinesin KSP, are described in PCT Publications WO 01/30768, WO 01/98278, WO 02/056880, WO 03/050,064, WO 03/050,122, WO 03/049,527, WO 03/049,679, WO 03/049,678, WO 03/039460, WO 03/079973, WO 03/099211, WO 2004/039774, WO 03/105855, WO 03/106417, WO 2004/087050, WO 2004/058700, WO 2004/058148 and WO 2004/037171 and US applications US 2004/132830 and US 2004/132719. In an embodiment inhibitors of mitotic kinesins include, but are not limited to inhibitors of KSP, inhibitors of MKLP1, inhibitors of

CENP-E, inhibitors of MCAK, inhibitors of Kif14, inhibitors of Mphosph1 and inhibitors of Rab6-KIFL.

“Inhibitors of kinases involved in mitotic progression” include, but are not limited to, inhibitors of aurora kinase, inhibitors of Polo-like kinases (PLK) (in particular inhibitors of PLK-1), inhibitors of bub-1 and inhibitors of bub-R1.

“Antiproliferative agents” includes antisense RNA and DNA oligonucleotides such as G3139, ODN698, RVASKRAS, GEM231, and INX3001, and antimetabolites such as enocitabine, carmofur, tegafur, pentostatin, doxifluridine, trimetrexate, fludarabine, capecitabine, galocitabine, cytarabine ocfosfate, fosteabine sodium hydrate, raltitrexed, paltitrexid, emitefur, tiazofurin, decitabine, nolatrexed, pemetrexed, nelzarabine, 2'-deoxy-2'-methylidenecytidine, 2'-fluoromethylene-2'-deoxycytidine, *N*-[5-(2,3-dihydro-benzofuryl)sulfonyl]-*N'*-(3,4-dichlorophenyl)urea, *N*6-[4-deoxy-4-[*N*2-[2(E),4(E)-tetradecadienoyl]glycylamino]-L-glycero-B-L-manno-heptopyranosyl]adenine, aplidine, ecteinascidin, troxacitabine, 4-[2-amino-4-oxo-4,6,7,8-tetrahydro-3*H*-pyrimidino[5,4-*b*][1,4]thiazin-6-yl-(*S*)-ethyl]-2,5-thienoyl-L-glutamic acid, aminopterin, 5-fluorouracil, alanosine, 11-acetyl-8-(carbamoyloxymethyl)-4-formyl-6-methoxy-14-oxa-1,11-diazatetracyclo(7.4.1.0.0)-tetradeca-2,4,6-trien-9-yl acetic acid ester, swainsonine, lometrexol, dexrazoxane, methioninase, 2'-cyano-2'-deoxy-*N*4-palmitoyl-1-*B*-D-arabino furanosyl cytosine and 3-aminopyridine-2-carboxaldehyde thiosemicarbazone.

Examples of monoclonal antibody targeted therapeutic agents include those therapeutic agents which have cytotoxic agents or radioisotopes attached to a cancer cell specific or target cell specific monoclonal antibody. Examples include Bexxar.

“HMG-CoA reductase inhibitors” refers to inhibitors of 3-hydroxy-3-methylglutaryl-CoA reductase. Examples of HMG-CoA reductase inhibitors that may be used include but are not limited to lovastatin (MEVACOR®; see U.S. Pat. Nos. 4,231,938, 4,294,926 and 4,319,039), simvastatin (ZOCOR®; see U.S. Pat. Nos. 4,444,784, 4,820,850 and 4,916,239), pravastatin (PRAVACHOL®; see U.S. Pat. Nos. 4,346,227, 4,537,859, 4,410,629, 5,030,447 and 5,180,589), fluvastatin (LESCOL®; see U.S. Pat. Nos. 5,354,772, 4,911,165, 4,929,437, 5,189,164, 5,118,853, 5,290,946 and 5,356,896) and atorvastatin (LIPITOR®; see U.S. Pat. Nos. 5,273,995, 4,681,893, 5,489,691 and 5,342,952). The structural formulas of these and additional HMG-CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", *Chemistry & Industry*, pp. 85-89 (5 February 1996) and US Patent Nos. 4,782,084 and 4,885,314. The term HMG-CoA reductase inhibitor as used herein includes all pharmaceutically acceptable lactone and open-acid forms (i.e., where the lactone ring is opened to form the free acid) as well as salt and ester forms of compounds which have HMG-CoA reductase inhibitory activity, and therefore the use of such salts, esters, open-acid and lactone forms is included within the scope of this invention.

“Prenyl-protein transferase inhibitor” refers to a compound which inhibits any one or any combination of the prenyl-protein transferase enzymes, including farnesyl-protein transferase

(FPTase), geranylgeranyl-protein transferase type I (GGPTase-I), and geranylgeranyl-protein transferase type-II (GGPTase-II, also called Rab GGPTase).

Examples of prenyl-protein transferase inhibitors can be found in the following publications and patents: WO 96/30343, WO 97/18813, WO 97/21701, WO 97/23478, WO 97/38665, WO 98/28980, WO 98/29119, WO 95/32987, U.S. Pat. No. 5,420,245, U.S. Pat. No. 5,523,430, U.S. Pat. No. 5,532,359, U.S. Pat. No. 5,510,510, U.S. Pat. No. 5,589,485, U.S. Pat. No. 5,602,098, European Patent Publ. 0 618 221, European Patent Publ. 0 675 112, European Patent Publ. 0 604 181, European Patent Publ. 0 696 593, WO 94/19357, WO 95/08542, WO 95/11917, WO 95/12612, WO 95/12572, WO 95/10514, U.S. Pat. No. 5,661,152, WO 95/10515, WO 95/10516, WO 95/24612, WO 95/34535, WO 95/25086, WO 96/05529, WO 96/06138, WO 96/06193, WO 96/16443, WO 96/21701, WO 96/21456, WO 96/22278, WO 96/24611, WO 96/24612, WO 96/05168, WO 96/05169, WO 96/00736, U.S. Pat. No. 5,571,792, WO 96/17861, WO 96/33159, WO 96/34850, WO 96/34851, WO 96/30017, WO 96/30018, WO 96/30362, WO 96/30363, WO 96/31111, WO 96/31477, WO 96/31478, WO 96/31501, WO 97/00252, WO 97/03047, WO 97/03050, WO 97/04785, WO 97/02920, WO 97/17070, WO 97/23478, WO 97/26246, WO 97/30053, WO 97/44350, WO 98/02436, and U.S. Pat. No. 5,532,359.

For an example of the role of a prenyl-protein transferase inhibitor on angiogenesis see *European J. of Cancer* (1999), **35(9)**:1394-1401.

“Angiogenesis inhibitors” refers to compounds that inhibit the formation of new blood vessels, regardless of mechanism. Examples of angiogenesis inhibitors include, but are not limited to, tyrosine kinase inhibitors, such as inhibitors of the tyrosine kinase receptors Flt-1 (VEGFR1) and Flk-1/KDR (VEGFR2), inhibitors of epidermal-derived, fibroblast-derived, or platelet derived growth factors, MMP (matrix metalloprotease) inhibitors, integrin blockers, interferon- α , interleukin-12, pentosan polysulfate, cyclooxygenase inhibitors, including nonsteroidal anti-inflammatories (NSAIDs) like aspirin and ibuprofen as well as selective cyclooxygenase-2 inhibitors like celecoxib and rofecoxib (*PNAS* (1992) **89**:7384; *JNCI* (1982) **69**:475; *Arch. Ophthalmol.* (1990) **108**:573; *Anat. Rec.* (1994) **238**:68; *FEBS Letters* (1995) **372**:83; *Clin. Orthop.* (1995) **313**:76; *J. Mol. Endocrinol.* (1996) **16**:107; *Jpn. J. Pharmacol.* (1997) **75**:105; *Cancer Res.* (1997) **57**:1625 (1997); *Cell* (1998) **93**:705; *Intl. J. Mol. Med.* (1998) **2**:715; *J. Biol. Chem.* (1999) **274**:9116)), steroidal anti-inflammatories (such as corticosteroids, mineralocorticoids, dexamethasone, prednisone, prednisolone, methylpred, betamethasone), carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl-fumagillol, thalidomide, angiostatin, troponin-1, angiotensin II antagonists (see Fernandez *et al* (1985) *J. Lab. Clin. Med.* **105**:141-145), and antibodies to VEGF (see, *Nature Biotechnology* (1999) **17**:963-968; Kim *et al* (1993) *Nature* **362**:841-844; WO 00/44777; and WO 00/61186).

Other therapeutic agents that modulate or inhibit angiogenesis and may also be used in combination with the compounds of the instant invention include agents that modulate or inhibit the coagulation and fibrinolysis systems (see review in *Clin. Chem. La. Med.* (2000) **38**:679-

692). Examples of such agents that modulate or inhibit the coagulation and fibrinolysis pathways include, but are not limited to, heparin (see *Thromb. Haemost.* (1998) **80**:10-23), low molecular weight heparins and carboxypeptidase U inhibitors (also known as inhibitors of active thrombin activatable fibrinolysis inhibitor [TAFIa]) (see *Thrombosis Res.* (2001) **101**:329-354).

5 TAFIa inhibitors have been described in PCT Publication WO 03/013,526 and U.S. Ser. No. 60/349,925 (filed January 18, 2002).

“Agents that interfere with cell cycle checkpoints” refer to compounds that inhibit protein kinases that transduce cell cycle checkpoint signals, thereby sensitizing the cancer cell to DNA damaging agents. Such agents include inhibitors of ATR, ATM, the Chk1 and Chk2 kinases and
10 cdk and cdc kinase inhibitors and are specifically exemplified by 7-hydroxystaurosporin, flavopiridol, CYC202 (Cyclacel) and BMS-387032.

“Inhibitors of cell proliferation and survival signaling pathway” refer to pharmaceutical agents that inhibit cell surface receptors and signal transduction cascades downstream of those surface receptors. Such agents include inhibitors of inhibitors of EGFR (for example gefitinib
15 and erlotinib), inhibitors of ERB-2 (for example trastuzumab), inhibitors of IGFR (for example those disclosed in WO 03/059951), inhibitors of cytokine receptors, inhibitors of MET, inhibitors of PI3K (for example LY294002), serine/threonine kinases (including but not limited to inhibitors of Akt such as described in (WO 03/086404, WO 03/086403, WO 03/086394, WO
20 03/086279, WO 02/083675, WO 02/083139, WO 02/083140 and WO 02/083138), inhibitors of Raf kinase (for example BAY-43-9006), inhibitors of MEK (for example CI-1040 and PD-098059) and inhibitors of mTOR (for example Wyeth CCI-779 and Ariad AP23573). Such agents include small molecule inhibitor compounds and antibody antagonists.

“Apoptosis inducing agents” include activators of TNF receptor family members (including the TRAIL receptors).

25 The invention also encompasses combinations with NSAID's which are selective COX-2 inhibitors. For purposes of this specification NSAID's which are selective inhibitors of COX-2 are defined as those which possess a specificity for inhibiting COX-2 over COX-1 of at least 100 fold as measured by the ratio of IC₅₀ for COX-2 over IC₅₀ for COX-1 evaluated by cell or microsomal assays. Such compounds include, but are not limited to those disclosed in U.S. Pat.
30 5,474,995, U.S. Pat. 5,861,419, U.S. Pat. 6,001,843, U.S. Pat. 6,020,343, U.S. Pat. 5,409,944, U.S. Pat. 5,436,265, U.S. Pat. 5,536,752, U.S. Pat. 5,550,142, U.S. Pat. 5,604,260, U.S. 5,698,584, U.S. Pat. 5,710,140, WO 94/15932, U.S. Pat. 5,344,991, U.S. Pat. 5,134,142, U.S. Pat. 5,380,738, U.S. Pat. 5,393,790, U.S. Pat. 5,466,823, U.S. Pat. 5,633,272, and U.S. Pat. 5,932,598, all of which are hereby incorporated by reference.

35 Inhibitors of COX-2 that are particularly useful in the instant method of treatment are 5-chloro-3-(4-methylsulfonyl)phenyl-2-(2-methyl-5-pyridinyl)pyridine; or a pharmaceutically acceptable salt thereof.

Compounds that have been described as specific inhibitors of COX-2 and are therefore useful in the present invention include, but are not limited to: parecoxib, CELEBREX[®] and BEXTRA[®] or a pharmaceutically acceptable salt thereof.

Other examples of angiogenesis inhibitors include, but are not limited to, endostatin, ukrain, ranpirnase, IM862, 5-methoxy-4-[2-methyl-3-(3-methyl-2-butenyl)oxiranyl]-1-oxaspiro[2,5]oct-6-yl(chloroacetyl)carbamate, acetyldinanaline, 5-amino-1-[[3,5-dichloro-4-(4-chlorobenzoyl)phenyl]methyl]-1*H*-1,2,3-triazole-4-carboxamide, CM101, squalamine, combretastatin, RPI4610, NX31838, sulfated mannopentaose phosphate, 7,7-(carbonyl-bis[imino-*N*-methyl-4,2-pyrrolocarbonylimino[*N*-methyl-4,2-pyrrole]-carbonylimino]-bis-(1,3-naphthalene disulfonate), and 3-[(2,4-dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416).

As used above, "integrin blockers" refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha_v\beta_3$ integrin, to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha_v\beta_5$ integrin, to compounds which antagonize, inhibit or counteract binding of a physiological ligand to both the $\alpha_v\beta_3$ integrin and the $\alpha_v\beta_5$ integrin, and to compounds which antagonize, inhibit or counteract the activity of the particular integrin(s) expressed on capillary endothelial cells. The term also refers to antagonists of the $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins. The term also refers to antagonists of any combination of $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\beta_5\alpha_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins.

Some specific examples of tyrosine kinase inhibitors include *N*-(trifluoromethylphenyl)-5-methylisoxazol-4-carboxamide, 3-[(2,4-dimethylpyrrol-5-yl)methylidenyl]indolin-2-one, 17-(allylamino)-17-demethoxygeldanamycin, 4-(3-chloro-4-fluorophenylamino)-7-methoxy-6-[3-(4-morpholinyl)propoxyl]quinazoline, *N*-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, BIBX1382, 2,3,9,10,11,12-hexahydro-10-(hydroxymethyl)-10-hydroxy-9-methyl-9,12-epoxy-1*H*-diindolo[1,2,3-*fg*:3',2',1'-*kl*]pyrrolo[3,4-*i*][1,6]benzodiazocin-1-one, SH268, genistein, STI571, CEP2563, 4-(3-chlorophenylamino)-5,6-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidinemethane sulfonate, 4-(3-bromo-4-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, 4-(4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, SU6668, STI571A, *N*-4-chlorophenyl-4-(4-pyridylmethyl)-1-phthalazinamine, and EMD121974.

Combinations with compounds other than anti-cancer compounds are also encompassed in the instant methods. For example, combinations of the instantly claimed compounds with PPAR- γ (i.e., PPAR-gamma) agonists and PPAR- δ (i.e., PPAR-delta) agonists are useful in the treatment of certain malignancies. PPAR- γ and PPAR- δ are the nuclear peroxisome proliferator-activated receptors γ and δ . The expression of PPAR- γ on endothelial cells and its involvement in angiogenesis has been reported in the literature (see *J. Cardiovasc. Pharmacol.* (1998) **31**:909-913; *J. Biol. Chem.* (1999) **274**:9116-9121; *Invest. Ophthalmol Vis. Sci.* (2000) **41**:2309-2317). More recently, PPAR- γ agonists have been shown to inhibit the angiogenic response to VEGF in vitro; both troglitazone and rosiglitazone maleate inhibit the development

of retinal neovascularization in mice. (*Arch. Ophthalmol.* (2001) **119**:709-717). Examples of PPAR- γ agonists and PPAR- γ/α agonists include, but are not limited to, thiazolidinediones (such as DRF2725, CS-011, troglitazone, rosiglitazone, and pioglitazone), fenofibrate, gemfibrozil, clofibrate, GW2570, SB219994, AR-H039242, JTT-501, MCC-555, GW2331, GW409544, NN2344, KRP297, NP0110, DRF4158, NN622, GI262570, PNU182716, DRF552926, 2-[(5,7-dipropyl-3-trifluoromethyl-1,2-benzisoxazol-6-yl)oxy]-2-methylpropionic acid (disclosed in USSN 09/782,856), and 2(R)-7-(3-(2-chloro-4-(4-fluorophenoxy)phenoxy)propoxy)-2-ethylchromane-2-carboxylic acid (disclosed in USSN 60/235,708 and 60/244,697).

Another embodiment of the instant invention is the use of the presently disclosed compounds in combination with anti-viral agents (such as nucleoside analogs including ganciclovir for the treatment of cancer. See WO 98/04290.

Another embodiment of the instant invention is the use of the presently disclosed compounds in combination with gene therapy for the treatment of cancer. For an overview of genetic strategies to treating cancer see Hall et al (*Am J Hum Genet* (1997) **61**:785-789) and Kufe et al (*Cancer Medicine*, 5th Ed, pp 876-889, BC Decker, Hamilton 2000). Gene therapy can be used to deliver any tumor suppressing gene. Examples of such genes include, but are not limited to, p53, which can be delivered via recombinant virus-mediated gene transfer (see U.S. Pat. No. 6,069,134, for example), a uPA/uPAR antagonist ("Adenovirus-Mediated Delivery of a uPA/uPAR Antagonist Suppresses Angiogenesis-Dependent Tumor Growth and Dissemination in Mice," *Gene Therapy*, August (1998) **5(8)**:1105-13), and interferon gamma (*J Immunol* (2000) **164**:217-222).

The compounds of the instant invention may also be administered in combination with an inhibitor of inherent multidrug resistance (MDR), in particular MDR associated with high levels of expression of transporter proteins. Such MDR inhibitors include inhibitors of p-glycoprotein (P-gp), such as LY335979, XR9576, OC144-093, R101922, VX853 and PSC833 (valspodar).

A compound of the present invention may be employed in conjunction with anti-emetic agents to treat nausea or emesis, including acute, delayed, late-phase, and anticipatory emesis, which may result from the use of a compound of the present invention, alone or with radiation therapy. For the prevention or treatment of emesis, a compound of the present invention may be used in conjunction with other anti-emetic agents, especially neurokinin-1 receptor antagonists, 5HT₃ receptor antagonists, such as ondansetron, granisetron, tropisetron, and zatisetron, GABA_B receptor agonists, such as baclofen, a corticosteroid such as Decadron (dexamethasone), Kenalog, Aristocort, Nasalide, Preferid, Benecorten or others such as disclosed in U.S. Patent Nos. 2,789,118, 2,990,401, 3,048,581, 3,126,375, 3,929,768, 3,996,359, 3,928,326 and 3,749,712, an antidopaminergic, such as the phenothiazines (for example prochlorperazine, fluphenazine, thioridazine and mesoridazine), metoclopramide or dronabinol. In an embodiment, an anti-emesis agent selected from a neurokinin-1 receptor antagonist, a 5HT₃

receptor antagonist and a corticosteroid is administered as an adjuvant for the treatment or prevention of emesis that may result upon administration of the instant compounds.

Neurokinin-1 receptor antagonists of use in conjunction with the compounds of the present invention are fully described, for example, in U.S. Pat. Nos. 5,162,339, 5,232,929, 5,242,930, 5,373,003, 5,387,595, 5,459,270, 5,494,926, 5,496,833, 5,637,699, 5,719,147; European Patent Publication Nos. EP 0 360 390, 0 394 989, 0 428 434, 0 429 366, 0 430 771, 0 436 334, 0 443 132, 0 482 539, 0 498 069, 0 499 313, 0 512 901, 0 512 902, 0 514 273, 0 514 274, 0 514 275, 0 514 276, 0 515 681, 0 517 589, 0 520 555, 0 522 808, 0 528 495, 0 532 456, 0 533 280, 0 536 817, 0 545 478, 0 558 156, 0 577 394, 0 585 913, 0 590 152, 0 599 538, 0 610 793, 0 634 402, 0 686 629, 0 693 489, 0 694 535, 0 699 655, 0 699 674, 0 707 006, 0 708 101, 0 709 375, 0 709 376, 0 714 891, 0 723 959, 0 733 632 and 0 776 893; PCT International Patent Publication Nos. WO 90/05525, 90/05729, 91/09844, 91/18899, 92/01688, 92/06079, 92/12151, 92/15585, 92/17449, 92/20661, 92/20676, 92/21677, 92/22569, 93/00330, 93/00331, 93/01159, 93/01165, 93/01169, 93/01170, 93/06099, 93/09116, 93/10073, 93/14084, 93/14113, 93/18023, 93/19064, 93/21155, 93/21181, 93/23380, 93/24465, 94/00440, 94/01402, 94/02461, 94/02595, 94/03429, 94/03445, 94/04494, 94/04496, 94/05625, 94/07843, 94/08997, 94/10165, 94/10167, 94/10168, 94/10170, 94/11368, 94/13639, 94/13663, 94/14767, 94/15903, 94/19320, 94/19323, 94/20500, 94/26735, 94/26740, 94/29309, 95/02595, 95/04040, 95/04042, 95/06645, 95/07886, 95/07908, 95/08549, 95/11880, 95/14017, 95/15311, 95/16679, 95/17382, 95/18124, 95/18129, 95/19344, 95/20575, 95/21819, 95/22525, 95/23798, 95/26338, 95/28418, 95/30674, 95/30687, 95/33744, 96/05181, 96/05193, 96/05203, 96/06094, 96/07649, 96/10562, 96/16939, 96/18643, 96/20197, 96/21661, 96/29304, 96/29317, 96/29326, 96/29328, 96/31214, 96/32385, 96/37489, 97/01553, 97/01554, 97/03066, 97/08144, 97/14671, 97/17362, 97/18206, 97/19084, 97/19942 and 97/21702; and in British Patent Publication Nos. 2 266 529, 2 268 931, 2 269 170, 2 269 590, 2 271 774, 2 292 144, 2 293 168, 2 293 169, and 2 302 689. The preparation of such compounds is fully described in the aforementioned patents and publications, which are incorporated herein by reference.

In an embodiment, the neurokinin-1 receptor antagonist for use in conjunction with the compounds of the present invention is selected from: 2-(*R*)-(1-(*R*)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(*S*)-(4-fluorophenyl)-4-(3-(5-oxo-1*H*,4*H*-1,2,4-triazolo)methyl)morpholine, or a pharmaceutically acceptable salt thereof, which is described in U.S. Pat. No. 5,719,147.

A compound of the instant invention may also be administered with an agent useful in the treatment of anemia. Such an anemia treatment agent is, for example, a continuous erythropoiesis receptor activator (such as epoetin alfa).

A compound of the instant invention may also be administered with an agent useful in the treatment of neutropenia. Such a neutropenia treatment agent is, for example, a hematopoietic

growth factor which regulates the production and function of neutrophils such as a human granulocyte colony stimulating factor, (G-CSF). Examples of a G-CSF include filgrastim.

A compound of the instant invention may also be administered with an immunologic-enhancing drug, such as levamisole, isoprinosine and Zadaxin.

5 A compound of the instant invention may also be useful for treating or preventing cancer, including bone cancer, in combination with bisphosphonates (understood to include bisphosphonates, diphosphonates, bisphosphonic acids and diphosphonic acids). Examples of bisphosphonates include but are not limited to: etidronate (Didronel), pamidronate (Aredia), alendronate (Fosamax), risedronate (Actonel), zoledronate (Zometa), ibandronate (Boniva),
10 incadronate or cimadronate, clodronate, EB-1053, minodronate, neridronate, piridronate and tiludronate including any and all pharmaceutically acceptable salts, derivatives, hydrates and mixtures thereof.

Thus, the scope of the instant invention encompasses the use of the instantly claimed compounds in combination with a second compound selected from: HDAC inhibitors, an
15 estrogen receptor modulator, an androgen receptor modulator, retinoid receptor modulator, a cytotoxic/cytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, a PPAR- γ agonist, a PPAR- δ agonist, an anti-viral agent, an inhibitor of inherent multidrug resistance, an anti-emetic agent, an agent useful in the treatment of anemia,
20 an agent useful in the treatment of neutropenia, an immunologic-enhancing drug, an inhibitor of cell proliferation and survival signaling, an agent that interferes with a cell cycle checkpoint, an apoptosis inducing agent and a bisphosphonate.

The term "administration" and variants thereof (e.g., "administering" a compound) in reference to a compound of the invention means introducing the compound or a prodrug of the
25 compound into the system of the animal in need of treatment. When a compound of the invention or prodrug thereof is provided in combination with one or more other active agents (e.g., a cytotoxic agent, etc.), "administration" and its variants are each understood to include concurrent and sequential introduction of the compound or prodrug thereof and other agents.

As used herein, the term "composition" is intended to encompass a product comprising
30 the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The term "therapeutically effective amount" as used herein means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue,
system, animal or human that is being sought by a researcher, veterinarian, medical doctor or
35 other clinician.

The term "treatment" refers to the treatment of a mammal afflicted with a pathological condition and refers to an effect that alleviates the condition by killing the cancerous cells, but also to an effect that results in the inhibition of the progress of the condition, and includes a

reduction in the rate of progress, a halt in the rate of progress, amelioration of the condition, and cure of the condition. Treatment as a prophylactic measure (i.e. prophylaxis) is also included.

The term "pharmaceutically acceptable" as used herein pertains to compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of a subject (e.g. human) without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. Each carrier, excipient, etc. must also be "acceptable" in the sense of being compatible with the other ingredients of the formulation.

The term "adjunct" refers to the use of compounds in conjunction with known therapeutic means. Such means include cytotoxic regimes of drugs and/or ionising radiation as used in the treatment of different cancer types. In particular, the active compounds are known to potentiate the actions of a number of cancer chemotherapy treatments, which include the topoisomerase class of poisons (e. g. topotecan, irinotecan, rubitecan), most of the known alkylating agents (e. g. DTIC, temozolamide) and platinum based drugs (e. g. carboplatin, cisplatin) used in treating cancer.

In an embodiment, the angiogenesis inhibitor to be used as the second compound is selected from a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP (matrix metalloprotease) inhibitor, an integrin blocker, interferon- α , interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, or an antibody to VEGF. In an embodiment, the estrogen receptor modulator is tamoxifen or raloxifene.

Also included in the scope of the claims is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with radiation therapy and/or in combination with a compound selected from: HDAC inhibitors, an estrogen receptor modulator, an androgen receptor modulator, retinoid receptor modulator, a cytotoxic/cytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, a PPAR- γ agonist, a PPAR- δ agonist, an anti-viral agent, an inhibitor of inherent multidrug resistance, an anti-emetic agent, an agent useful in the treatment of anemia, an agent useful in the treatment of neutropenia, an immunologic-enhancing drug, an inhibitor of cell proliferation and survival signaling, an agent that interferes with a cell cycle checkpoint, an apoptosis inducing agent and a bisphosphonate.

And yet another embodiment of the invention is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with paclitaxel or trastuzumab.

The invention further encompasses a method of treating or preventing cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with a COX-2 inhibitor.


The instant invention also includes a pharmaceutical composition useful for treating or preventing cancer that comprises a therapeutically effective amount of a compound of Formula I and a compound selected from: HDAC inhibitors, an estrogen receptor modulator, an androgen receptor modulator, a retinoid receptor modulator, a cytotoxic/cytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, a PPAR- γ agonist, a PPAR- δ agonist, an anti-viral agent, an inhibitor of cell proliferation and survival signaling, an agent that interferes with a cell cycle checkpoint, an apoptosis inducing agent and a bisphosphonate.

These and other aspects of the invention will be apparent from the teachings contained herein.


Abbreviations used in the description of the chemistry and in the Examples that follow are:

DMSO: dimethylsulfoxide; DMF: dimethylformamide; MeOH: methanol; EtOH: ethanol; AcOH: acetic acid; THF: tetrahydrofuran; DCM: dichloromethane; MeCN: acetonitrile; TFA: trifluoroacetic acid; HOBT: 1-hydroxybenzotriazole; PS-DCC: *N,N'*-dicyclohexylcarbodiimide polymer supported; min: minutes; h: hour(s); eq.: equivalent(s); M: molar; RT: room temperature; and RP-HPLC: reversed phase high-pressure liquid chromatography.

Compounds of formula 1 can be prepared by reacting a compound of formula 1A with a compound of formula IB:



(IA)




(IB)

wherein a, b, c, d, A, R¹, R², R³, R⁸, R⁹ and Y are as defined above. The reaction is generally carried out in the presence of a solvent such as AcOH at about 150°C, generally under microwave irradiation.

Where the synthesis of intermediates and starting materials is not described, these compounds are commercially available or can be made from commercially available compounds using standard methods or by extension of the Examples herein.

Compounds of formula I may be converted to other compounds of Formula I by known methods or by methods described in the Examples. For example, compounds containing ester moieties can be converted to corresponding carboxylic acid by standard hydrolysis methods such as using a base such as NaOH, generally in a solvent such as methanol and heating at 90°C under microwave irradiation.

Compounds of formula I wherein $Y-(R^8R^9)_bR^2$ is CO_2H can be converted to other compounds of formula I by reacting a compound of formula IC with a compound of formula ID:




wherein a, b, c, A, R^1 , R^2 , R^3 , R^8 and R^9 are as defined above and Y^2 is a direct bond, O or NR^7 , wherein R^7 is as defined above. The reaction is generally carried out in the presence of coupling agents such as HOBt in a solvent such as DMF at about room temperature.

During any of the synthetic sequences described herein it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protecting Groups in Organic Synthesis*, 3rd Edition, Greene, T. W. and Wuts, P. G. M.; Wiley Interscience, 1999 and Kocienski, P. J. *Protecting Groups*, Thieme, 1994. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The compounds of this invention were prepared according to the following schemes. All variables within the formulae are as defined above.

Scheme 1

In Scheme 1, pyrazolo[1,5-*a*]quinazolin-5(4*H*)-ones were prepared by reaction of α -cyanoketones with 2-hydrazino benzoic acids under microwave irradiation as described in *Molecular Diversity* **2003**, 7, 161.



wherein all variables are as defined above.

The exemplified compounds described herein were tested by the assay described below and were found to have an IC₅₀ value of less than 10 μ M.

PARP-1 SPA assay**Working Reagents**

Assay buffer: 100 mM Tris pH 8, 4 mM MgCl₂, 4 mM Spermine, 200 mM KCl, 0.04% Nonidet P-40.

Enzyme Mix: Assay buffer (12.5 μ l), 100 mM DTT (0.5 μ l), PARP-1 (5 nM, Trevigen 4668-500-01), H₂O (to 35 μ l).

Nicotinamide-adenine dinucleotide (NAD)/ DNA Mix: [³H-NAD] (250 uCi/ml, 0.4 μ l, Perkin-Elmer NET-443H), NAD (1.5 mM, 0.05 μ l, SIGMA N-1511), Biotinylated-NAD (250 μ M, 0.03 μ l, Trevigen 4670-500-01), Activated calf thymus (1mg/ml, 0.05 μ l, Amersham Biosciences 27-4575), H₂O (to 10 μ l).

Developing Mix: Streptavidin SPA beads (5mg/ml, Amersham Biosciences RPNQ 0007) dissolved in 500 mM EDTA.

Experimental Design

The reaction is performed in 96-well microplate with a final volume of 50 μ L/well. Add 5 μ l 5%DMSO/compound solution, add enzyme mix (35 μ l), start the reaction by adding NAD/DNA mix (10 μ L) and incubate for 2 hrs at RT. Stop the reaction by adding developing mix (25 μ l) and incubate 15 min at RT. Measure using a Packard TOP COUNT instrument.

EXAMPLE 1:2-phenylpyrazolo[1,5-*a*]quinazolin-5(4*H*)-one

A solution (0.66 M) of 2-hydrazinobenzoic acid hydrochloride (from Aldrich) in AcOH was treated with benzonitrile (1 eq.). The reaction mixture was heated at 150°C under microwave irradiation for 5 min. The resulting precipitate was filtered, washed with Et₂O and dried to afford (69%) the title compound as a solid.

¹H NMR (400 MHz, DMSO-*d*₆, 300K) δ 6.38 (s, 1H), 7.40 (t, *J* 7.6 Hz, 1H), 7.48 (t, *J* 7.6 Hz, 2H), 7.51 (t, *J* 7.6 Hz, 1H), 7.91 (t, *J* 7.6 Hz, 1H), 7.97 (d, *J* 7.6 Hz, 2H), 8.17-8.14 (m, 2H), 12.31 (s, 1H). MS (ES⁺) C₁₆H₁₁N₃O requires: 261, found: 262 (M+H)⁺.

The following Examples were made according to the process given in Example 1.

Example	Name	Molecular Ion [M+H] ⁺	Procedure
2	2-(2-methylphenyl)pyrazolo[1,5- <i>a</i>] quinazolin-5(4 <i>H</i>)-one	276	Example 1
3	2-(3-chlorophenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	296; 298	Example 1
4	2-(3-fluorophenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	280	Example 1
5	2-(3-bromophenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	340; 342	Example 1
6	2-(3-methylphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	276	Example 1
7	2-(4-fluorophenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	280	Example 1
8	2-(4-bromophenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	340, 342	Example 1
9	2-(4-chlorophenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	296, 298	Example 1
10	2-(4-methylphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	276	Example 1
11	2-(4-ethylphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	290	Example 1
12	2-(4-isopropoxyphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	320	Example 1

13	2-[4-(trifluoromethoxy)phenyl]pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	346	Example 1
14	methyl 4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoate	320	Example 1
15	2-(3,4-difluorophenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	298	Example 1
16	2-(2-furyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	252	Example 1
17	2-(3,4-dimethoxyphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	322	Example 1
18	2-(3,4-dimethylphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	290	Example 1
19	2-(3,5-dimethoxyphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	322	Example 1
20	2-(5-fluoro-2-methylphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	294	Example 1
21	2-(2,5-difluorophenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	298	Example 1
22	2-(2,3-dimethoxyphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	322	Example 1
23	2-(2,4-dimethoxyphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	322	Example 1
24	2-(3,4,5-trimethoxyphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	352	Example 1
25	2-(4-methoxy-3,5-dimethylphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	320	Example 1
26	2-(2-thienyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	268	Example 1
27	2-[5-methyl-2-(trifluoromethyl)-3-furyl]pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	334	Example 1
28	2-(2-methyl-3-furyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	266	Example 1
29	3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)pyridinium trifluoroacetate	263	Example 1
30	4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)pyridinium trifluoroacetate	263	Example 1
31	3-methyl-2-phenylpyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	276	Example 1
32	2-biphenyl-2-ylpyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	338	Example 1
33	2-(1-naphthyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	312	Example 1

34	4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzonitrile	287	Example 1
35	2-[1-methyl-5-(trifluoromethyl)-1 <i>H</i> -pyrazol-4-yl]pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	334	Example 1
36	2-(3-fluoro-4-morpholin-4-ylphenyl)pyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	365	Example 1
37	4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoic acid	306	hydrolysis of Example 14
38	2-cyclohexylpyrazolo[1,5- <i>a</i>]quinazolin-5(4 <i>H</i>)-one	268	Example 1

EXAMPLE 39:

1-methyl-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium trifluoroacetate

5 Step 1: methyl 4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoate

Starting from methyl 4-(cyanoacetyl)benzoate (from Aldrich) and following the procedure described in Example 1, the title compound was obtained (67%) as a solid.

¹H NMR (300 MHz, DMSO-*d*₆, 300K) δ 3.86 (s, 3H), 6.45 (s, 1H), 7.50 (t, *J* 7.5 Hz, 1H), 7.90 (t, *J* 7.6 Hz, 1H), 8.05 (dd, *J* 23.9 Hz, 8.3 Hz, 4H), 8.19-8.12 (m, 2H), 12.34 (s, 1H). MS (ES⁺)

10 C₁₈H₁₃N₃O₃ requires: 319, found: 320 (M+H)⁺.

Step 2: 4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoic acid

A solution (0.4 M) of the product from step 1 in MeOH was treated with 2N NaOH (3.5 eq.).

The reaction mixture was heated at 90°C under microwave irradiation for 5 min. The reaction mixture was poured into water and acidified with 6N HCl to pH 2, then it was centrifuged and

15 the aqueous supernatant solution was eliminated to afford (78%) the title compound as a solid.

¹H NMR (300 MHz, DMSO-*d*₆, 300K) δ 6.47 (s, 1H), 7.45-7.60 (m, 1H), 7.85-8.30 (m, 7H), 12.39 (bs, 1H), 12.97 (bs, 1H). MS (ES⁺) C₁₇H₁₁N₃O₃ requires: 305, found: 306 (M+H)⁺.

Step 3: 1-methyl-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium trifluoroacetate

20 A solution (0.06 M) of the product from step 2 and HOBt (1.7 eq.) in DMF was added to PS-DCC. The resulting suspension was stirred at RT for 45 min, then, a solution of 1-methyl-1,4-diazepane (0.7 eq.) in DMF was added. The reaction mixture was stirred at RT for 48 h. Then it was filtered and concentrated giving a residue that was purified by RP-HPLC (Conditions: Waters X-Bridge C18, 5 micron, 19 x 100 mm; flow: 20 mL/min; Gradient: A: H₂O + 0.1%

TFA; B: MeCN + 0.1% TFA; 10% A isocratic for 1 min, linear to 60% A in 9 min) to afford (24%) the title compound as a solid.

¹H NMR (400 MHz, DMSO-d₆, 300K) δ 2.00-2.20 (m, 2H), 2.83-2.90 (m, 3H), 3.16-3.20 (m, 2H), 3.37-3.75 (m, 5H), 4.05-4.18 (m, 1H), 6.44 (s, 1H), 7.49-7.57 (m, 3H), 7.92 (t, *J* 8.3 Hz, 1H), 8.11 (dd, *J* 44.2, *J* 8.6 Hz, 4H), 9.74 (bs, 1H), 12.35 (s, 1H). MS (ES⁺) C₂₃H₂₃N₅O₂ requires: 401, found: 402 (M+H)⁺.

EXAMPLE 40:

4-(3-{[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}propyl)morpholin-4-ium trifluoroacetate

Step 1: methyl 3-(bromoacetyl)benzoate

A solution of 3-(bromoacetyl)benzoic acid (from Fluorochem) in toluene/MeOH (7:3) was treated with trimethylsilyl diazomethane (2 M solution in Et₂O, 1 eq.). After stirring at RT for 5 min, the reaction mixture was concentrated to afford (66%) the title compound as a solid.

¹H NMR (300 MHz, DMSO-d₆, 300K) δ 3.89 (s, 3H), 5.00 (s, 2H), 7.67-7.76 (m, 1H), 8.18-8.28 (m, 2H), 8.49 (s, 1H). MS (ES⁺) C₁₀H₉BrO₃ requires: 256, found: 257 (M+H)⁺.

Step 2: methyl 3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoate

A solution (0.6 M) of the product from step 1 in EtOH/THF (2:1) was treated with an aqueous solution of KCN (6 N, 5.5 eq.). The reaction mixture was stirred at RT for 2 h, then, was poured onto ice/water and acidified with AcOH. After dilution with DCM the organic phase was separated and washed with brine and dried. Evaporation of the solvent gave a yellow solid which was treated with 2-hydrazinobenzoic acid hydrochloride as described in Example 1 to afford (66%) the title compound as a solid.

¹H NMR (300 MHz, DMSO-d₆, 300K) δ 3.90 (s, 3H), 6.44 (s, 1H), 7.51 (t, *J* 7.6Hz, 1H), 7.62 (t, *J* 7.6Hz, 1H), 7.87-8.02 (m, 2H), 8.12-8.26 (m, 3H), 8.52 (s, 1H), 12.35 (s, 1H). MS (ES⁺) C₁₈H₁₃N₃O₃ requires: 319, found: 320 (M+H)⁺.

Step 3: 3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoic acid

Starting from the product from step 2 and following the procedure described in Example 39, step 2, the title compound was obtained (62%) as a solid.

¹H NMR (300 MHz, DMSO-d₆, 300K) δ 6.43 (s, 1H), 7.51 (t, *J* 7.7Hz, 1H), 7.59 (t, *J* 7.7Hz, 1H), 7.85-8.00 (m, 2H), 8.12-8.24 (m, 3H), 8.51 (s, 1H), 12.33 (s, 1H), 13.08 (bs, 1H). MS (ES⁺) C₁₇H₁₁N₃O₃ requires: 305, found: 306(M+H)⁺.

Step 4: 4-(3-{[3-(5-oxo-4,5-dihydropyrazolo[1,5-*a*]quinazolin-2-yl)benzoyl]amino}propyl)morpholin-4-ium trifluoroacetate

Starting from the product from step 3 and following the procedure described in Example 39, step 3, it was obtained a residue that was purified by RP-HPLC (Conditions: Waters X-Bridge C18, 5 micron, 19 x 100 mm; flow: 20 mL/min; Gradient: A: H₂O + 0.1% TFA; B: MeCN + 0.1% TFA; 5% A isocratic for 1.5 min, linear to 50% A in 8.5 min) to afford (39%) the title compound as a solid.

MS (ES⁺) C₂₄H₂₅N₅O₃ requires: 431, found: 432 (M+H)⁺.


Example	Name	Molecular Ion [M+H] ⁺	Procedure
41	1-methyl-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate	388	Example 39
42	<i>N,N</i> -dimethyl-2-{methyl[4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]amino}ethanaminium trifluoroacetate	390	Example 39
43	(8 <i>aS</i>)-2-[4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]octahydropyrrolo[1,2- <i>a</i>]pyrazin-5-ium trifluoroacetate	414	Example 39
44	1-[(3-methylpyridinium-2-yl)methyl]-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]piperazin-1-ium bis(trifluoroacetate)	479	Example 39
45	1-[(2-methyl-1,3-thiazol-3-ium-4-yl)methyl]-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]piperazin-1-ium bis(trifluoroacetate)	485	Example 39
46	1-(cyclopropylmethyl)-4-[4-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate	428	Example 39
47	6,6-difluoro-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium trifluoroacetate	424	Example 40
48	<i>N,N</i> -dimethyl-2-{7-oxo-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-yl}ethanaminium trifluoroacetate	473	Example 40

49	<i>N,N</i> -dimethyl{1-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]piperidin-3-yl}methanaminium trifluoroacetate	430	Example 40
50	1-methyl-2-({methyl[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]amino } methyl)-1 <i>H</i> -imidazol-3-ium trifluoroacetate	413	Example 40
51	1-methyl-3-({ [3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]amino } methyl)pyrrolidinium trifluoroacetate	402	Example 40
52	<i>N,N</i> -dimethyl-2- { [3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]amino } propan-1-aminium trifluoroacetate	390	Example 40
53	1-(cyclopropylmethyl)-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate	428	Example 40
54	(8 <i>aS</i>)-2-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]octahydropyrrolo[1,2- <i>a</i>]pyrazin-5-ium trifluoroacetate	414	Example 40
55	1-methyl-5-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]-1,2,4,5,6,6 <i>a</i> -hexahydropyrrolo[3,4- <i>b</i>]pyrrol-1-ium trifluoroacetate	412	Example 40
56	4-(2,4-dichlorophenyl)-1-(2- { [3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]amino } ethyl)piperazin-1-ium trifluoroacetate	561, 563	Example 40
57	1-methyl-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium trifluoroacetate	402	Example 40
58	1-methyl-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate	388	Example 40

59	<i>N,N</i> -dimethyl-2-{methyl[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]amino}ethanaminium trifluoroacetate	390	Example 40
60	4-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]-1,4-diazepan-1-ium trifluoroacetate	388	Example 40
61	4-(2-{[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]amino}ethyl)morpholin-4-ium trifluoroacetate	418	Example 40
62	1-(4-chlorophenyl)-4-[3-(5-oxo-4,5-dihydropyrazolo[1,5- <i>a</i>]quinazolin-2-yl)benzoyl]piperazin-1-ium trifluoroacetate	484, 486	Example 40

CLAIMS

1. The use of a compound of formula I:



wherein:

a is 0, 1, 2, 3 or 4;

each b is independently 0, 1, 2 or 3;

c is 0 or 1;

d is 0, 1, 2, 3 or 4;

A is C₃₋₁₀cycloalkyl, C₆₋₁₀aryl, a 4 membered saturated heterocycle containing one N atom, a 5, 6 or 7 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, or a 7-13 membered unsaturated or partially saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S;

each R¹ is independently hydroxy, halogen, cyano, C₁₋₆alkyl, haloC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkoxy, haloC₁₋₆alkoxy, carboxy, C₁₋₆alkoxycarbonyl, nitro, amino, C₁₋₆alkylamino or di(C₁₋₆alkyl)amino;

each Y is independently a direct bond, O, O(C=O), (C=O)O, C=O, NR⁷(C=O), (C=O)NR⁷, S(O)_r, NR⁷S(O)_r or S(O)_rNR⁷;

each r is independently 0, 1 or 2;

each R² is independently hydrogen, hydroxy, cyano, halogen, C₁₋₆alkyl, haloC₁₋₆alkyl, C₁₋₆alkylcarbonyl, haloC₁₋₆alkoxy, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, nitro, N(R^a)₂, N(R^a)CON(R^a)₂, N(R^a)COOR^a, N(R^a)S(O)_rR^b, N(R^a)S(O)_rN(R^a)₂ or a ring which is: C₃₋₁₀cycloalkyl, C₆₋₁₀aryl, C₆₋₁₀aryloxy, C₆₋₁₀arylcarbonyl, a 4 membered saturated heterocycle containing one N atom, a 5 or 6 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic ring containing

1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, or a 7, 8, 9 or 10 membered unsaturated, partially saturated or saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from $(\text{CH}_2)_e\text{R}^4$;

R^3 is halogen, cyano, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkoxy or halo C_{1-6} alkoxy;

each R^a is independently hydrogen, C_{1-6} alkyl, halo C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{6-10} aryl or C_{6-10} aryl C_{1-6} alkyl;

each R^b is independently C_{1-6} alkyl or C_{1-6} alkylcarbonyl;

e is 0, 1, 2, 3 or 4; and

R^4 is hydroxy, oxo, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, nitro, $\text{N}(\text{R}^a)_2$, $\text{S}(\text{O})_r\text{N}(\text{R}^a)_2$, $\text{CON}(\text{R}^a)_2$, $\text{N}(\text{R}^a)\text{COR}^a$, $\text{N}(\text{R}^a)\text{CON}(\text{R}^a)_2$, $\text{N}(\text{R}^a)\text{COOR}^a$, $\text{S}(\text{O})_r\text{R}^b$ or a ring which is: C_{3-10} cycloalkyl, C_{6-10} aryl, a 4 membered saturated heterocycle containing one N atom, a 5 or 6 membered


saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms or a 7, 8, 9 or 10 membered unsaturated, partially saturated or saturated heterocycle ring containing 1, 2 or 3 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from hydroxy, oxo, cyano, nitro, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy and $\text{N}(\text{R}^a)_2$;

R^7 is hydrogen, C_{1-6} alkyl or halo C_{1-6} alkyl;

each of R^8 and R^9 is independently hydrogen, C_{1-6} alkyl or C_{1-6} alkoxy;

or a pharmaceutically acceptable salt or tautomer thereof, for the manufacture of a medicament for the treatment or prevention of conditions which can be ameliorated by the inhibition of poly(ADP-ribose)polymerase (PARP).

2. A compound of formula I:



wherein a, b, c, d, A, R¹, R², R³, R⁸, R⁹ and Y are as defined in claim 1;


5 provided that:

- (a) when A is phenyl and either a is 0 or a is 1 and R¹ is methyl, bromine, fluorine, nitro or carboxy, then d is not 0 and Y-(CR⁸R⁹)_bR² is not hydrogen, trifluoromethyl, chlorine or methoxy; and
- (b) when A is 2-thienyl or 2-furyl and either a is 0 or a is 1 and R¹ is methoxy, bromine, nitro or fluorine, then d is not 0 and Y-(CR⁸R⁹)_bR² is not hydrogen; or a pharmaceutically acceptable salt or tautomer thereof.
- 10

3. A compound of claim 2 wherein R² is hydroxy, cyano, halogen, haloC₁₋₆alkyl, haloC₁₋₆alkoxy or a ring which is phenyl, morpholinyl, diazepanyl, piperazinyl, octahydropyrrolopyrazinyl, piperidinyl, imidazolyl, pyrrolidinyl or hexahydropyrrolopyrrolyl; any of which rings being optionally substituted by one, two or three groups independently selected from (CH₂)_eR⁴.

15

4. A compound of claim 2 of formula II:



(II)

5 wherein:

a, b, c, A, R¹, R³, R⁸ and R⁹ are as defined in claim 2;

f is 0, 1 or 2;

g is 1 or 2;

10 R⁵ is independently hydroxy, cyano, halogen, C₁-₆alkyl, haloC₁-₆alkyl, C₁-₆alkylcarbonyl, haloC₁-₆alkoxy, C₂-₁₀alkenyl, C₂-₁₀alkynyl, C₁-₆alkoxy, C₁-₆alkoxycarbonyl, carboxy, nitro, N(Rᵃ)₂, S(O)ᵣN(Rᵃ)₂, CON(Rᵃ)₂, N(Rᵃ)CON(Rᵃ)₂, S(O)ᵣRᵇ or N(Rᵃ)S(O)ᵣRᵇ;

each Rᵃ is independently hydrogen, C₁-₆alkyl, haloC₁-₆alkyl, hydroxyC₁-₆alkyl, C₁-₆alkylcarbonyl, C₂-₁₀alkenyl, C₂-₁₀alkynyl, C₆-₁₀aryl or C₆-₁₀arylC₁-₆alkyl;

each Rᵇ is independently C₁-₆alkyl or C₁-₆alkylcarbonyl;

15 Y¹ is a direct bond, O(C=O), (C=O)O, C=O, NR⁷(C=O), (C=O)NR⁷, S(O)ᵣ, NHR⁷(O)ᵣ or S(O)ᵣNR⁷;

r is 0, 1 or 2;

20 R⁶ is hydroxy, cyano, fluorine, bromine, iodine, C₁-₆alkyl, C₁-₆alkylcarbonyl, haloC₁-₆alkoxy, C₂-₁₀alkenyl, C₂-₁₀alkynyl, haloC₁-₆alkoxy, nitro, N(Rᵃ)₂, N(Rᵃ)CON(Rᵃ)₂, N(Rᵃ)S(O)ᵣRᵇ or a ring which is: C₃-₁₀cycloalkyl, C₆-₁₀aryl, C₆-₁₀aryloxy, C₆-₁₀arylcarbonyl, 4 membered saturated heterocycle containing one N atom, or a 5 or 6 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one O atom, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms, or a 7, 8, 9 or 10 membered unsaturated, partially saturated or saturated heterocyclic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O

and S; any of which rings being optionally substituted by one or more groups independently selected from $(\text{CH}_2)_e\text{R}^4$;


e is 0, 1, 2, 3 or 4; and

R^4 is hydroxy, oxo, cyano, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy, C_{1-6} alkoxycarbonyl, nitro, $\text{N}(\text{R}^a)_2$, $\text{S}(\text{O})_r\text{N}(\text{R}^a)_2$, $\text{CON}(\text{R}^a)_2$, $\text{N}(\text{R}^a)\text{CON}(\text{R}^a)_2$, $\text{S}(\text{O})_r\text{R}^b$ or a ring which is: C_{3-10} cycloalkyl, C_{6-10} aryl, a 4 membered saturated heterocycle containing one N atom, a 5 or 6 membered saturated or partially saturated heterocycle containing one, two or three N atoms and zero or one heteroatom selected from O and S, a 5 membered heteroaromatic ring containing 1, 2, 3 or 4 heteroatoms independently selected from N, O and S, not more than one heteroatom of which is O or S, a 6 membered heteroaromatic ring containing 1, 2 or 3 N atoms or a 7, 8, 9 or 10 membered unsaturated, partially saturated or saturated heterocycle ring containing 1, 2 or 3 heteroatoms independently selected from N, O and S; any of which rings being optionally substituted by one or more groups independently selected from hydroxy, oxo, cyano, nitro, halogen, C_{1-6} alkyl, halo C_{1-6} alkyl, C_{1-6} alkoxy, halo C_{1-6} alkoxy and $\text{N}(\text{R}^a)_2$;

or a pharmaceutically acceptable salt or tautomer thereof.

5. A compound of any one of claims 2 to 4 wherein A is phenyl, furyl, thienyl, pyridinyl, naphthyl, pyrazolyl or cyclohexyl.

6. A compound of claim 4 of formula III:



(III)

wherein:

a, b, c, f, g, R^1 , R^3 , R^5 , R^6 , R^8 , R^9 and Y^1 are as defined in claim 4; or a pharmaceutically acceptable salt or tautomer thereof.

7. A compound of any one of claims 4 to 6 wherein R⁶ is methyl, fluorine, bromine, iodine, ethyl, isopropyl, trifluoromethoxy, phenyl, cyano, morpholinyl, hydroxy, methyldiazepanyl, methylpiperazinyl, dimethylamino, octahydropyrrolopyrazinyl, [(methylpyridinyl)methyl]piperazinyl, [(methylthiazolyl)methyl]piperazinyl, (cyclopropylmethyl)piperazinyl, difluorodiazepanyl, [(dimethylamino)ethyl](oxo)diazepanyl, [(dimethylamino)methyl]piperidinyl, methylimidazolyl, methylpyrrolidinyl, methylhexahydropyrrolopyrrolyl, (dichlorophenyl)piperazinyl, diazepanyl or (chlorophenyl)piperazinyl.

8. A pharmaceutical composition comprising a compound of any one of claims 2 to 7, or a pharmaceutically acceptable salt or tautomer thereof in association with a pharmaceutically acceptable carrier.

9. A compound as defined in any one of claims 1 to 7, or a pharmaceutically acceptable salt or tautomer thereof and an anti-cancer agent for simultaneous, separate or sequential administration.

10. A compound of any one of claims 2 to 7, or a pharmaceutically acceptable salt or tautomer thereof for use in therapy.

11. The use of a compound of any one of claims 2 to 7, or a pharmaceutically acceptable salt or tautomer thereof for the manufacture of a medicament for the treatment or prevention of conditions which can be ameliorated by the inhibition of poly(ADP-ribose)polymerase (PARP).

12. The use of a compound as defined in any one of claims 1 to 7, or a pharmaceutically acceptable salt or tautomer thereof for the manufacture of a medicament for the treatment or prevention of cancer, inflammatory diseases, reperfusion injuries, ischemic conditions, stroke, renal failure, cardiovascular diseases, vascular diseases other than cardiovascular diseases, diabetes mellitus, neurodegenerative diseases, retroviral infections, retinal damage, skin senescence or UV-induced skin damage.

13. The use of a compound as defined in any one of claims 1 to 7, or a pharmaceutically acceptable salt or tautomer thereof as a chemo-or radiosensitizer for cancer treatment.

14. A method of treating or preventing cancer, inflammatory diseases, reperfusion injuries, ischemic conditions, stroke, renal failure, cardiovascular diseases, vascular diseases

other than cardiovascular diseases, diabetes mellitus, neurodegenerative diseases, retroviral infections, retinal damage, skin senescence or UV-induced skin damage, which method comprises administration to a patient in need thereof of an effective amount of a compound of claim 1 or a composition comprising a compound of claim 1.

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2007/050332

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D487/04 A61K31/519 A61P9/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	VASQUEZ JR., T. E. ET AL.: MOLECULAR DIVERSITY, vol. 7, 2003, pages 161-164, XP002448151 cited in the application the whole document -----	1-14

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

29 August 2007

Date of mailing of the international search report

18/09/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Tabanella, Stefania

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2007/050332

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 13 and 14 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.