



(86) Date de dépôt PCT/PCT Filing Date: 2013/10/15
(87) Date publication PCT/PCT Publication Date: 2015/04/23
(85) Entrée phase nationale/National Entry: 2016/03/11
(86) N° demande PCT/PCT Application No.: US 2013/065020
(87) N° publication PCT/PCT Publication No.: 2015/057198

(51) CI.Int./Int.Cl. A61K 9/16 (2006.01),
A61K 31/20 (2006.01), A61K 31/70 (2006.01),
A61P 3/02 (2006.01)

(71) Demandeur/Applicant:
BENEMILK OY, FI

(72) Inventeurs/Inventors:
ARONEN, ILMO PELLERVO, FI;
HOLMA, MERJA BIRGITTA, FI;
WAN, FENG, US;
BUNTEL, CHRISTOPHER JOHN, SG

(74) Agent: REGEHR, HERBERT B.

(54) Titre : COMPOSITIONS ALIMENTAIRES CONTENANT UN HYDRATE DE CARBONE AINSI QUE LEURS METHODES DE
PREPARATION ET D'UTILISATION

(54) Title: CARBOHYDRATE-CONTAINING DIETARY COMPOSITIONS AND METHODS FOR THEIR PREPARATION AND USE





FIG. 1

(57) Abrégé/Abstract:

Dietary compositions containing at least one carbohydrate are described herein, as well as methods for their preparation and use. A premix dietary composition may include a fatty acid component and a carbohydrate component. The fatty acid component may include at least about 90% saturated fatty acid by weight. The fatty acid component may be present in the dietary composition in an amount of at least about 30% by weight of the dietary composition.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number
WO 2015/057198 A1

(43) International Publication Date
23 April 2015 (23.04.2015)

(51) International Patent Classification:
A61K 9/16 (2006.01) **A61K 9/50 (2006.01)**

(21) International Application Number:
PCT/US2013/065020

(22) International Filing Date:
15 October 2013 (15.10.2013)

(25) Filing Language:
English

(26) Publication Language:
English

(71) Applicant: **BENEMILK OY [FI/FI]**; Raisonkaari 55, Raisio 21200 (FI).

(72) Inventors: **ARONEN, Ilmo Pellervo**; Sonkintie 165, FI-27600 Hinnerjoki (FI). **HOLMA, Merja Birgitta**; Valo-


polku 1, FI-21200 Raisio (FI). **WAN, Feng**; 4209 257th Place SE, Issaquah, Washington 98029 (US). **BUNTEL, Christopher John**; 2 Ridgewood Close, #19-02, Singapore 276693 (SG).

(74) Agent: **PISANO, Anthony L.**; Pepper Hamilton LLP, Suite 5000, 500 Grant Street, Pittsburgh, Pennsylvania 15219-2507 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,

[Continued on next page]

(54) Title: CARBOHYDRATE-CONTAINING DIETARY COMPOSITIONS AND METHODS FOR THEIR PREPARATION AND USE



(57) Abstract: Dietary compositions containing at least one carbohydrate are described herein, as well as methods for their preparation and use. A premix dietary composition may include a fatty acid component and a carbohydrate component. The fatty acid component may include at least about 90% saturated fatty acid by weight. The fatty acid component may be present in the dietary composition in an amount of at least about 30% by weight of the dietary composition.

WO 2015/057198 A1

FIG. 1

WO 2015/057198 A1



MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,

Published:

— *with international search report (Art. 21(3))*

CARBOHYDRATE-CONTAINING DIETARY COMPOSITIONS AND METHODS FOR THEIR PREPARATION AND USE

BACKGROUND

[0001] Increasing production and fat content of milk obtained from lactating ruminants has been a major goal for dairy farmers. Additional milk production per ruminant is beneficial because it results in a higher yield, thereby increasing profits. Increased milk fat is desirable because it has a higher economic value and can be used in highly desirable food products, such as cheese, yogurt, and the like.

[0002] A common approach to increasing either or both production and milk fat content includes adjusting feed, nutrients, elements, vitamins, supplements, and/or the like provided to the ruminant. One such specific method includes feeding the ruminant a total mixed ration (TMR), which is a mix of grain and silage with some protein meals, such as, for example, soya bean meal and canola meal. Additional materials and trace elements, vitamins, extra nutrients, and the like may also be added to the TMR.

[0003] However, the current methods and feeds used to increase milk fat content tend to lower milk production, lower protein content, and/or have other detrimental effects on the ruminant. Furthermore, the methods and feeds often result in other undesired effects, such as increased trans fatty acid levels on the fatty acid profile of the milk fat.

SUMMARY

[0004] In an embodiment, a premix dietary composition may include a fatty acid component and a carbohydrate component. The fatty acid component may include at least about 90% saturated fatty acid by weight. The fatty acid component may be present in the dietary composition in an amount of at least about 30% by weight of the dietary composition. In an embodiment, a method of preparing a premix dietary composition for ruminants may include combining a fatty acid component and a carbohydrate component to form a mixture

and processing the mixture into a tablet, a capsule, a pellet, or a granular material. The fatty acid component may be present in the dietary composition in an amount of at least about 30% by weight of the dietary composition.

[0005] In an embodiment, a method of increasing milk fat content in ruminants may include combining a premix dietary composition with a feed to obtain a mixture and providing the mixture to a ruminant for ingestion. The premix dietary composition may include a fatty acid component and a carbohydrate component. The fatty acid component may be present in the premix dietary composition in an amount of at least about 30% by weight of the premix dietary composition.

[0006] In an embodiment, a premix composition for ruminants may include a fatty acid component and a carbohydrate component. The fatty acid component may include a palmitic acid compound in an amount of at least about 90% by weight of the fatty acid component. The fatty acid component may be present in the premix composition in an amount of at least about 30% by weight of the premix composition. The premix composition may be a solid having an average size of about 0.1 mm to about 3 mm. The premix composition may be configured to be mixed with a ruminant feed and provided to ruminants.

BRIEF DESCRIPTION OF THE DRAWINGS

[0007] FIG. 1 depicts a flow diagram of a method of preparing a dietary composition for ruminants according to an embodiment.

DETAILED DESCRIPTION

[0008] This disclosure is not limited to the particular systems, devices and methods described, as these may vary. The terminology used in the description is for the purpose of describing the particular versions or embodiments only, and is not intended to limit the scope.

[0009] As used in this document, the singular forms “a,” “an,” and “the” include plural references unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art. Nothing in this disclosure is to be construed as an admission that the embodiments described in this disclosure are not entitled to antedate such disclosure by virtue of prior invention. As used in this document, the term “comprising” means “including, but not limited to.”

[0010] The following terms shall have, for the purposes of this application, the respective meanings set forth below.

[0011] A ruminant is a class of mammal with a multiple chamber stomach that gives the animal an ability to digest cellulose-based food. The stomach of a ruminant has four morphologically distinct compartments: the rumen, the reticulum, the omasum, and the abomasum. Bacteria in the rumen enable the ruminant to digest cellulose-based food by softening it and regurgitating the semi-digested mass. The regurgitate, known as cud, is then chewed again by the ruminant. Specific examples of ruminants include, but are not limited to, cattle, bison, buffaloes, yaks, camels, llamas, giraffes, deer, pronghorns, antelopes, sheep, and goats. The milk produced by ruminants is widely used in a variety of dairy-based products. Dairy cows are of considerable commercial significance for the production of milk and processed dairy products such as, for example, yogurt, cheese, whey, and ice cream.

[0012] Silage refers to a feed that includes chopped green forage, such as, for example, grass, legumes, and field corn. The silage is placed in a structure or a container that is designed to exclude air. The silage is then fermented in the structure or container, thereby retarding spoilage. Silage can have a water content of about 60% to about 80% by weight.

[0013] A premix composition is a composition that contains a mixture of ingredients, such as, for example, a fatty acid component and a carbohydrate component. The premix

composition can be sold, marketed, packaged, transported, and/or the like without further modification. In some embodiments, the premix composition may be combined with one or more other ingredients. Combination with one or more other ingredients may occur before or after the premix composition is sold, marketed, packaged, transported, and/or the like. For example, the premix composition may be made, packaged, and shipped to an end user, and the end user may mix the premix composition with feed material.

[0014] The present disclosure relates generally to dietary compositions such as premix compositions that contain feed additives and the like. As used herein, “dietary composition,” “premix composition,” and “dietary premix composition” may be used interchangeably, as those skilled in the art will recognize that the various terms relate to the same thing. The dietary compositions described herein can be added to feed and fed to ruminants for purposes of affecting milk production in the ruminant. Particularly, the dietary compositions described herein may be fed to a ruminant to increase the amount of milk produced by the ruminant and/or to increase the fat content of the milk produced by the ruminant, as described in greater detail herein.

[0015] When a ruminant consumes feed, the fat in the feed is modified by the rumen to provide a milk fat profile that is different from the profile of fat in the feed. All fats which are not completely inert in the rumen may decrease rumen digestibility of the feed material. Milk composition and fat quality can be influenced by the ruminant’s diet. For example, oil feeding can have negative effects on both rumen function and milk formation. As a result of the oil feeding, the milk protein concentration is lowered, the fat concentration is decreased, and the proportion of trans fatty acids is increased. These have been connected especially to an increase in the harmful low-density lipoprotein (LDL) cholesterol and to a decrease in the beneficial high-density lipoprotein (HDL) cholesterol in human blood when the milk is consumed. In addition, the properties of the milk fat during industrial milk processing are

weakened. A high level of polyunsaturated fatty acids in milk can also cause taste defects and preservation problems. A typical fatty acid composition of milk fat may contain more than 70% saturated fatty acids, and the total amount of trans fatty acids may vary in the range of 3%-10%. When vegetable oil is added into the feed, the proportion of trans fatty acids may rise to more than 10%.

[0016] One solution to diminishing the detrimental effect of oil and fat is to prevent triglyceride fat hydrolysis. Fat hydrolysis can be decreased, for example, by protecting fats with formaldehyde treated casein. Another alternative is to make insoluble fatty acid calcium salts whereby hydrogenation in rumen can be avoided. However, fatty acid salts have a pungent taste, which can limit their usability in feeds and can result in decreased feed intake. The salts may also impact the pelletizing process of the feed.

[0017] Accordingly, the dietary composition described herein allows for the transfer of palmitic acid from the feed via the digestive tract into the blood circulation of a ruminant. This improves the energy efficiency of milk production of the ruminant. When the utilization of energy becomes more efficient, the milk production increases and the concentrations of protein and fat in the milk rise. Especially, the dietary composition enhances fat synthesis in the mammary gland by bringing milk fat components to the cell and therefore the energy consuming synthesis in the mammary gland may not be necessary. Thus, glucose may be more efficiently used for lactose production whereupon milk production increases. The milk protein content rises since there may be no need to produce glucose from amino acids. Thus, the ruminant may not lose as much weight at the beginning of the lactation period.

[0018] In the various embodiments described herein, the dietary composition may include at least one fatty acid component and at least one carbohydrate component. The fatty acid component may be primarily saturated fatty acid (such as palmitic acid) and may contain little or no unsaturated trans fatty acid, as described in greater detail herein. The fatty acid

component may be present in generally any concentration, such as, for example, about 30% to about 80% by weight of the dietary composition. In further examples, the fatty acid component may be present in the dietary composition in an amount of about 30% to about 50%, about 40% to about 60%, or about 60% to about 90% by weight of the dietary composition. The carbohydrate component may be present in generally any concentration, such as, for example, about 20% to about 70% by weight of the dietary composition. In further examples, the carbohydrate component may be present in the dietary composition in an amount of about 10% to about 40%, or about 50% to about 70% by weight of the composition. In some embodiments, the compositions described herein may be used as a booster or a supplement to other feed, such as a premix composition or the like.

[0019] FIG. 1 depicts a flow diagram of a representative method of preparing a dietary composition for a ruminant. In various embodiments, the dietary composition may be formulated in a manner so that when it is mixed with a feed and consumed by the ruminant, the dietary composition maximizes particular qualities in the milk produced by the ruminant, as well as an amount of milk produced by the ruminant, as described in greater detail herein. In particular embodiments, the dietary composition may be substantially a solid dietary composition, including, but not limited to, a capsule, a tablet, a pellet, or a granular material.

[0020] In various embodiments, the components described herein with respect to FIG. 1 may generally be combined in any order and/or any combination, and are not limited by the order described herein. In some embodiments, a dietary composition may be prepared by providing **105** a fatty acid component and adding **110** a carbohydrate component to the fatty acid component. Thus, processes **105** and **110** result in combining the fatty acid component to the carbohydrate component to obtain a mixture. In some embodiments, the dietary composition may consist essentially of the fatty acid component and the carbohydrate component. In other embodiments, the dietary composition may consist of the fatty acid

component and the carbohydrate component. In other embodiments, the dietary composition may include other components in addition to the fatty acid component and the carbohydrate component, as described in greater detail herein.

[0021] In various embodiments, the fatty acid component may generally include one or more free fatty acids and/or glycolipids. Free fatty acids may generally be unconjugated fatty acids, whereas glycolipids may be fatty acids conjugated with a carbohydrate. In some embodiments, the fatty acid component may be present in the dietary composition in an amount of at least about 30% by weight of the dietary composition, and can, for example, be about 30% by weight to about 80% by weight of the dietary composition. In some embodiments, the fatty acid component may be present in the dietary composition in an amount of at least about 50% by weight of the dietary composition. In particular embodiments, the fatty acid component may be present in the dietary composition in an amount of about 30% by weight, about 35% by weight, about 40% by weight, about 45% by weight, about 50% by weight, about 55% by weight, about 60% by weight, about 65% by weight, about 70% by weight, about 75% by weight, about 80% by weight, or any value or range between any two of these values. In some embodiments, the fatty acid component may represent about 30% to about 50%, about 30% to about 90%, or about 40% to about 60% by weight of the dietary composition.

[0022] In some embodiments, the fatty acid component may have a melting point equal to or greater than about 40°C. In some embodiments, the fatty acid component may have a melting point equal to or less than about 80°C. In some embodiments, the fatty acid component may have a melting point of about 40°C to about 80°C. In some embodiments, the fatty acid component may have a melting point of about 60°C to about 80°C. In some embodiments, the fatty acid component may have a melting point of about 63°C to about 65°C. In particular embodiments, the fatty acid component may have a melting point of

about 40°C, about 45°C, about 50°C, about 55°C, about 60°C, about 65°C, about 70°C, about 75°C, about 80°C, or any value or range between any two of these values. The melting point may generally be selected so that it is a temperature that ensures that the fatty acid is inert in the rumen environment.

[0023] In various embodiments, the fatty acid component may include at least one saturated fatty acid. For example, the fatty acid component may include 1, 2, 3, 4, 5, 6, or more different saturated fatty acids. In some embodiments, the saturated fatty acid may be present in the fatty acid component in an amount that results in a ruminant consuming the dietary composition to produce a desired quality and quantity of milk, as described in greater detail herein. Thus, in some embodiments, the saturated fatty acid may generally be present in any amount, such as an amount of at least about 90% by weight of the fatty acid component. In some embodiments, the saturated fatty acid may be present in an amount of about 90% by weight of the fatty acid component to about 100% by weight of the fatty acid component, including about 90% by weight, about 91% by weight, about 92% by weight, about 93% by weight, about 94% by weight, about 95% by weight, about 96% by weight, about 97% by weight, about 98% by weight, about 99% by weight, about 100% by weight, or any value or range between any two of these values. The saturated fatty acid is not limited by this disclosure, and may include any number of saturated fatty acids now known or later discovered, including all derivatives thereof. For example, derivatives of a saturated fatty acid may include salts, esters, amides, carbonates, carbamates, imides, anhydrides, alcohols, and/or the like.

[0024] As used herein, a salt of the fatty acid may be any acid addition salt, including, but not limited to, halogenic acid salts such as, for example, hydrobromic, hydrochloric, hydrofluoric, and hydroiodic acid salts; inorganic acid salts such as, for example, nitric, perchloric, sulfuric, and phosphoric acid salts; organic acid salts such as, for example,

sulfonic acid salts (methanesulfonic, trifluoromethane sulfonic, ethanesulfonic, benzenesulfonic, or p-toluenesulfonic), acetic, malic, fumaric, succinic, citric, benzoic, gluconic, lactic, mandelic, mucic, pamoic, pantothenic, oxalic, and maleic acid salts; and amino acid salts such as aspartic or glutamic acid salts. The acid addition salt may be a mono- or di-acid addition salt, such as a di-hydrohalogenic, di-sulfuric, di-phosphoric, or di-organic acid salt. In all cases, the acid addition salt is used as an achiral reagent which is not selected on the basis of any expected or known preference for interaction with or precipitation of a specific optical isomer of the products of this disclosure.

[0025] A fatty acid ester, as used herein, means an ester of a fatty acid. For example, the fatty acid ester may be in a form of RCOOR' . R may be any saturated or unsaturated alkyl group including, without limitation, C10, C12, C14, C16, C18, C20, and C24. R' may be any groups having from about 1 to about 1000 carbon atoms and with or without hetero atoms. In some embodiments, R' may have from about 1 to about 20, from about 3 to about 10, or from about 5 to about 15 carbon atoms. The hetero atoms may include, without limitation, N, O, S, P, Se, halogen, Si, and B. For example, R' may be a C1-6alkyl, such as methyl, ethyl or t-butyl; a C1-6alkoxyC1-6alkyl; a heterocyclyl, such as tetrahydrofuranyl; a C6-10aryloxyC1-6alkyl, such as benzyloxymethyl (BOM); a silyl, such as trimethylsilyl, t-butyldimethylsilyl and t-butyldiphenylsilyl; a cinnamyl; an allyl; a C1-6alkyl which is mono-, di- or trisubstituted by halogen, silyl, cyano or C1-6aryl, wherein the aryl ring is unsubstituted or substituted by one, two, or three residues selected from the group consisting of C1-7alkyl, C1-7alkoxy, halogen, nitro, cyano and CF₃; or a C1-2alkyl substituted by 9-fluorenyl.

[0026] As used herein, a fatty acid amide may generally include amides of fatty acids where the fatty acid is bonded to an amide group. For example, the fatty acid amide may have a formula of $\text{RCONR}'\text{R}''$. R may be any saturated or unsaturated alkyl group including,

without limitation, C10, C12, C14, C16, C18, C20, and C24. R' and R'' may each be any group having from about 1 to about 1000 carbon atoms and with or without hetero atoms. In some embodiments, R' may have from about 1 to about 20, from about 3 to about 10, or from about 5 to about 15 carbon atoms. The hetero atoms may include, without limitation, N, O, S, P, Se, halogen, Si, and B. For example, R' and R'' each may be an alkyl, an alkenyl, an alkynyl, an aryl, an aralkyl, a cycloalkyl, a halogenated alkyl, or a heterocycloalkyl group.

[0027] A fatty acid anhydride may generally refer to a compound which results from the condensation of a fatty acid with a carboxylic acid. Illustrative examples of carboxylic acids that may be used to form a fatty acid anhydride include acetic acid, propionic acid, benzoic acid, and the like.

[0028] An alcohol of a fatty acid refers to a fatty acid having straight or branched, saturated, radical groups with 3-30 carbon atoms and one or more hydroxy groups. The alkyl portion of the alcohol component can be propyl, butyl, pentyl, hexyl, iso-propyl, iso-butyl, sec-butyl, tert-butyl, or the like. One of skill in the art may appreciate that other alcohol groups may also be useful in the present disclosure.

[0029] In some embodiments, the saturated fatty acid may include a palmitic acid compound. The palmitic acid compound is not limited by this disclosure, and may include one or more of a conjugated palmitic acid, unconjugated palmitic acid, free palmitic acid, palmitic acid derivatives, and/or the like. Palmitic acid, also known as hexadecanoic acid, has a molecular formula of $\text{CH}_3(\text{CH}_2)_{14}\text{CO}_2\text{H}$. Specific examples of palmitic acid derivatives may include palmitic acid esters, palmitic acid amides, palmitic acid salts, palmitic acid carbonates, palmitic acid carbamates, palmitic acid imides, palmitic acid anhydrides, and/or the like. The palmitic acid compound may be present in the fatty acid component in generally any amount, such as an amount of at least about 60% by weight of the fatty acid component, including, for example, about 60% by weight of the fatty acid to about 100% by

weight of the fatty acid, including about 60% by weight, about 65% by weight, about 70% by weight, about 75% by weight, about 80% by weight, about 85% by weight, about 90% by weight, about 95% by weight, about 98% by weight, about 99% by weight, about 100% by weight, or any value or range between any two of these values. In some embodiments, the fatty acid component may consist essentially of the palmitic acid compound. In other embodiments, the fatty acid component may consist of or be entirely composed of the palmitic acid compound.

[0030] In some embodiments, the saturated fatty acid may include a stearic acid compound. The stearic acid compound is not limited by this disclosure, and may include conjugated stearic acid, unconjugated stearic acid, free stearic acid, stearic acid derivatives, and/or the like. Stearic acid, also known as octadecanoic acid, has a chemical formula of $\text{CH}_3(\text{CH}_2)_{16}\text{CO}_2\text{H}$. Specific examples of stearic acid derivatives may include stearic acid esters, stearic acid amides, stearic acid salts, stearic acid carbonates, stearic acid carbamates, stearic acid imides, stearic acid anhydrides, and/or the like. Because stearic acid in large amounts may hinder milk production capacity of the mammary gland, the amount of stearic acid may be present in the fatty acid component in an amount of about 30% or less by weight of the fatty acid component. In particular embodiments, the stearic acid compound may include about 30% by weight of the fatty acid component, about 25% by weight of the fatty acid component, about 20% by weight of the fatty acid component, about 15% by weight of the fatty acid component, about 10% by weight of the fatty acid component, about 5% by weight of the fatty acid component, or any value or range between any two of these values.

[0031] In some embodiments, the fatty acid component may include an unsaturated fatty acid. Unsaturated fatty acid, as used herein, refers to any mono- or polyunsaturated fat, and includes unsaturated trans fatty acids. The unsaturated fatty acids must contain at least one alkene bond and may contain two or more alkene groups in any position in the

hydrocarbon chain, and the unsaturation may or may not be present as a conjugated system of double bonds. The unsaturated fatty acid is not limited by this disclosure, and may include any number of unsaturated fatty acids now known or later discovered, including all derivatives thereof. For example, derivatives of an unsaturated fatty acid may include salts, esters, amides, anhydrides, alcohols, and/or the like, as previously described herein. In various embodiments, an amount of unsaturated fatty acid may be used in the fatty acid component to affect a desired quality of milk produced by the ruminant consuming the dietary composition, as described in greater detail herein. Thus, in some embodiments, the fatty acid component may be substantially free of unsaturated fatty acids. As used herein with respect to unsaturated fatty acids, the term “substantially free” is understood to mean substantially no amount of unsaturated fatty acids or about 10% or less by weight of unsaturated fatty acids, including trace amounts of unsaturated fatty acids. Accordingly, the unsaturated fatty acid may be present in the fatty acid component in an amount of about 10% or less by weight of the fatty acid component, including about 10% or less by weight, about 5% or less by weight, about 4% or less by weight, about 3% or less by weight, about 2% or less by weight, about 1% or less by weight, about 0.5% or less by weight, about 0% by weight, or any value or range between any two of these values.

[0032] In various embodiments, at least a portion of the fatty acid component may be contained. In some embodiments, the fatty acid component may be pre-contained prior to providing **105** the fatty acid component. In other embodiments, the fatty acid component may be contained as a result of the various processes **105, 110, 115, 120, 125** described herein. In some embodiments, the fatty acid may generally be contained by at least one supermolecular structure. Supermolecular structures may include vesicular structures such as microemulsions, liposomes (vesicles), micelles, and reverse micelles. The liposomes (vesicles) may contain an aqueous volume that is entirely enclosed by a membrane composed

of lipid molecules, such as phospholipids. In some embodiments, the liposomes may have a bilayer membrane. In some embodiments, the liposomes may include at least one surfactant. Examples of surfactants may include polyoxyethylene ethers and esters of fatty acids. The surfactant may have an hydrophilic-lipophilic balance (HLB) value of about 2 to about 12, including about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, or any range or value between any two of these values. Micelles and reverse micelles are microscopic vesicles that contain amphipathic constituents but do not contain an aqueous volume that is entirely enclosed by a membrane. In micelles, the hydrophilic part of the amphipathic compound is on the outside (on the surface of the vesicle). In reverse micelles, the hydrophobic part of the amphipathic compound is on the outside. The reverse micelles may thus contain a polar core that can solubilize both water and macromolecules within the inverse micelle. As the volume of the core aqueous pool increases, the aqueous environment may begin to match the physical and chemical characteristics of bulk water. The resulting inverse micelle may be referred to as a microemulsion of water in oil.

[0033] In some embodiments, at least a portion of the fatty acid component may be contained in a core of a micelle or a vesicle. The core may include any number of particles therein in addition to the fatty acid. The core composition may be made of a core material that includes at least one of a protein material, a cellulosic material, an amino acid, and an amino acid derivative.

[0034] In various embodiments, at least a portion of the fatty acid component may be encapsulated. In some embodiments, the fatty acid component may be pre-encapsulated prior to providing **105** the fatty acid component. In other embodiments, the fatty acid component may be encapsulated as a result of the various processes **105, 110, 115, 120, 125** described herein. In some embodiments, the fatty acid component may generally be encapsulated by a

capsule. The capsule may include a capsule shell, which is made up of at least one polysaccharide or protein. Illustrative examples of capsule shells as described herein may include capsule shells including agar, gelatin, starch casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phthalated gelatin, succinated gelatin, cellulosephthalate-acetate, polyvinylacetate, hydroxypropyl methylcellulose, polyvinylacetate-phthalate, polymerisates of acrylic esters, polymerisates of methacrylic esters, and/or any mixture thereof.

[0035] The carbohydrate component is not limited by this disclosure and may include any carbohydrates or combination of carbohydrates, particularly those used in animal feed and dietary compositions. In some embodiments, the carbohydrate component may generally provide a source of energy for the dietary composition. Illustrative examples of carbohydrate components may include molasses, sugar beet pulp, sugar cane, wheat bran, wheat middlings, wheat mill run, oat hulls, grain hulls, soya hulls, soybean hulls, peanut hulls, wood, brewery byproducts, beverage industry byproducts, forages, roughages, grass meal, hay meal, hay, alfalfa meal, alfalfa, straw, silages, sugars, starches, cellulose, hemicellulose, wheat, corn, oats, sorghum, millet, barley, barley fiber, barley hulls, barley middlings, barley bran, malting barley screenings, malting barley and fines, malt rootlets, maize bran, maize middlings, maize cobs, maize screenings, maize fiber, millet, rice, rice bran, rice middlings, rye, triticale, brewers grain, coffee grinds, tea leaf fines, citrus fruit pulp, rind residues, algae, algae meal, microalgae, and/or the like.

[0036] The carbohydrate component may be obtained from any carbohydrate source, and thus the source is not limited by this disclosure. In some embodiments, the carbohydrate may be obtained by breaking down a complex sugar source. Illustrative carbohydrate sources may include sugar, starch, cellulose, hemicellulose, and/or the like. In some embodiments,

the carbohydrate may be obtained from various crops that contain carbohydrates. Illustrative crops may include wheat, corn, oats, sorghum, millet, barley, and/or the like.

[0037] In various embodiments, one or more other ingredients may be added **115** to the mixture. The other ingredients may be added **115** at substantially the same time as processes **105** and **110**, may be added subsequent to processes **105** and **110**, may be added prior to processes **105** and **110**, or may be added during process **120** and **125**, as described in greater detail herein. Illustrative examples of other ingredients that may be added **115** include an emulsifier, a glucogenic precursor, an antioxidant, a vitamin, carnitine, an amino acid, a mineral, a nitrogen source material, a binding agent, a bulking agent, a filler, water, and the like, or a combination thereof. Any number and combination of ingredients may be added **115** to the mixture. The other ingredients may generally be added **115** in various amounts necessary to provide beneficial nutritional and dietary needs of the ruminant that is to consume the dietary composition. For example, other ingredients may include an amino acid and a mineral, each in an amount sufficient to provide beneficial nutritional and dietary needs of the ruminant.

[0038] The glucogenic precursor may include at least one of glycerol, propylene glycol, molasses, propionate, glycerine, propane diol, calcium propionate, propionic acid, octanoic acid, steam-exploded sawdust, steam-exploded wood chips, steam-exploded wheat straw, algae, algae meal, microalgae, and/or the like. The glucogenic precursor may generally be included in the dietary composition to provide an energy source to the ruminant so as to prevent gluconeogenesis from occurring within the ruminant's body.

[0039] The antioxidant is not limited by this disclosure and may include any antioxidants or combination of antioxidants, particularly those used in animal feed and dietary compositions. Illustrative examples of antioxidants may include alpha-carotene, beta-carotene, ethoxyquin, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT),

cryptoxanthin, lutein, lycopene, zeaxanthin, vitamin A, vitamin C, vitamin E, selenium, alpha-lipoic acid, and/or the like.

[0040] The vitamin may include any one or a combination of vitamins including, without limitation, vitamin A, vitamin B, vitamin C, vitamin D, vitamin E, vitamin K, and/or the like. The vitamin may include any vitamins from each particular vitamer group, including A vitamins, B vitamins, C vitamins, D vitamins, E vitamins, K vitamins, and/or the like. Specific examples of B vitamins include thiamine (vitamin B1), riboflavin (vitamin B2), niacin (vitamin B3), pantothenic acid (vitamin B5), pyridoxine (vitamin B6), biotin (vitamin B7), folic acid (vitamin B9), cobalamin (vitamin B12), and choline (vitamin Bp).

[0041] One additional ingredient that may be added **115** is carnitine. Carnitine may be included in the dietary composition to aid in the breakdown of fatty acids to generate metabolic energy in the ruminant. In some embodiments, carnitine may be provided as a portion of a carnitine premix composition.

[0042] In some embodiments, the amino acid may be an essential amino acid, including any one or a combination of leucine, lysine, histidine, valine, arginine, threonine, isoleucine, phenylalanine, methionine, tryptophan, and/or any derivative thereof. In some embodiments, the amino acid may be a non-essential amino acid, including any one or a combination of alanine, asparagine, aspartate, cysteine, glutamate, glutamine, glycine, proline, serine, tyrosine, and/or any derivative thereof. The amino acid and/or any derivative thereof may also include amino acids and derivatives of both non-essential and essential amino acids. The amino acid may generally be included in the dietary composition to provide a nutritional aid in various physiological processes in the ruminant, such as, for example, increasing muscle mass, providing energy, aiding in recovery, and/or the like. In some embodiments, the amino acid may be obtained from an amino acid premix composition.

[0043] The mineral may be any mineral that is a generally recognized as safe (GRAS) mineral or a combination of such minerals. The mineral may further be obtained from any mineral source that provides a bioavailable mineral. In some embodiments, the mineral may be one or more of calcium, sodium, magnesium, potassium, phosphorous, zinc, selenium, manganese, iron, cobalt, copper, iodine, molybdenum, and/or the like. In some embodiments, the mineral may be selected from one or more of a sodium salt, a calcium salt, a magnesium salt, a cobalt salt, a manganese salt, a potassium salt, an iron salt, a zinc salt, copper sulfate, copper oxide, selenium yeast, a chelated mineral, and/or the like. Illustrative examples of sodium salts include monosodium phosphate, sodium acetate, sodium chloride, sodium bicarbonate, disodium phosphate, sodium iodate, sodium iodide, sodium tripolyphosphate, sodium sulfate, sodium selenite, and/or the like. Illustrative examples of calcium salts include calcium acetate, calcium carbonate, calcium chloride, calcium gluconate, calcium hydroxide, calcium iodate, calcium iodobehenate, calcium oxide, anhydrous calcium sulfate, calcium sulfate dehydrate, dicalcium phosphate, monocalcium phosphate, tricalcium phosphate, and/or the like. Illustrative magnesium salts include magnesium acetate, magnesium carbonate, magnesium oxide, magnesium sulfate, and/or the like. Illustrative cobalt salts include cobalt acetate, cobalt carbonate, cobalt chloride, cobalt oxide, cobalt sulfate, and/or the like. Illustrative examples of manganese salts include manganese carbonate, manganese chloride, manganese citrate, manganese gluconate, manganese orthophosphate, manganese oxide, manganese phosphate, manganese sulfate, and/or the like. Illustrative examples of potassium salts include potassium acetate, potassium bicarbonate, potassium carbonate, potassium chloride, potassium iodate, potassium iodide, potassium sulfate, and/or the like. Illustrative examples of iron salts include iron ammonium citrate, iron carbonate, iron chloride, iron gluconate, iron oxide, iron phosphate, iron pyrophosphate,

iron sulfate, reduced iron, and/or the like. Illustrative examples of zinc salts include zinc acetate, zinc carbonate, zinc chloride, zinc oxide, zinc sulfate, and/or the like.

[0044] The nitrogen source material may generally include, for example, an oilseed meal. Oilseed meal is generally derived from residue that remains after reserved oil is removed from oilseeds. The oilseed meal may be rich in protein and variable in residual fats and oils. Illustrative examples of oilseed meal include soy meal, bean meal, rapeseed meal, soybean meal, sunflower meal, coconut meal, olive meal, linseed meal, grapeseed meal, cottonseed meal, camelina meal, mustard seed meal, crambe seed meal, safflower meal, rice meal, peanut meal, corn gluten meal, corn gluten feed, distillers dried grains, distillers dried grains with solubles, wheat gluten, and/or the like.

[0045] The binding agent may provide adhesive properties to the dietary composition, particularly so that the dietary composition does not fall apart in various forms such as pellet and tablet forms. Examples of binding agents include polysaccharides, proteins, and the like, or a combination thereof. The bulking agent may generally increase the bulk of the dietary composition without affecting the taste of the dietary composition. Examples of bulking agents may include silicate, kaolin, clay, and/or the like. The filler may generally be used to increase bulk, weight, viscosity, opacity, strength, and/or the like. Examples of filler may include gluten feed, sunflower hulls, distillers grains, guar hulls, wheat middlings, rice hulls, rice bran, oilseed meals, dried blood meal, animal byproduct meal, fish byproduct meal, dried fish solubles, feather meal, poultry byproducts, meat meal, bone meal, dried whey, soy protein concentrate, soy flour, yeast, wheat, oats, grain sorghum, corn feed meal, algae meal, rye, corn, barley, aspirated grain fractions, brewers dried grains, corn flower, corn gluten meal, feeding oat meal, sorghum grain flour, wheat mill run, wheat red dog, hominy feed, wheat flower, wheat bran, wheat germ meal, oat groats, rye middlings, cotyledon fiber, and/or ground grains.

[0046] In various embodiments, water may be present in the dietary composition. In some embodiments, the water may be inherently present in any of the ingredients in the dietary composition. In some embodiments, as previously described herein, an amount of water may be added **115**. The water may be included in an amount that is separate from any amounts of water that may be inherently present in any of the other ingredients described herein. The water may be present in the dietary composition in an amount that is about 3% or less by weight, including about 0.5% by weight, about 1% by weight, about 2% by weight, about 3% by weight, or any value or range between any two of these values.

[0047] In some embodiments, at least one cellulosic material may also be added **115**. The cellulosic material may generally provide a source of fiber for the ruminant to lower cholesterol levels and promote proper digestive function. Illustrative examples of cellulosic materials include wheat bran, wheat middlings, wheat mill run, oat hulls, oat bran, soya hulls, grass meal, hay meal, alfalfa meal, alfalfa, straw, hay, algae, algae meal, microalgae, and/or the like.

[0048] In some embodiments, a micronutrient mixture may be added **115**. Micronutrient mixtures are not limited by this disclosure and may generally contain any micronutrient mixture now known or later developed. The micronutrient mixture may include various components, such as at least one vitamin and at least one mineral, as described in greater detail herein. In some embodiments, the micronutrient mixture may be present in a micronutrient premix composition.

[0049] In various embodiments, the dietary composition may be processed **120**. In some embodiments, processing **120** may be completed prior to processes **105, 110, 115** to prepare various portions of the dietary composition for mixing and/or the like. In other embodiments, processing **120** may be completed after processes **105, 110, 115** to prepare a final product. In some embodiments, processing **120** may include forming the dietary

composition into a capsule, a shell, a pellet, a tablet, a granular material, and/or the like. Processing **120** may include pressing, molding, extruding, grinding, pelleting, encapsulating, granulating and/or the like. Pressing may include, for example, applying a pressure to an amount of the dietary composition. Molding may include, for example, open molding, compression molding, injection molding, centrifugal molding, or the like. Extruding may include, for example, forming an amount of the dietary composition by forcing the dietary composition through a die having a desired shape and size.

[0050] Grinding may be performed by various grinding devices known to those having ordinary skill in the art, such as a hammer mill, a roller mill, a disk mill, or the like. The dietary composition and/or portions thereof such as the carbohydrate component may be ground to various sizes, such as particle size (for instance, measured in millimeters), mesh sizes, surface areas, or the like. According to some embodiments, the dietary composition and/or portions thereof may be ground to an average particle size of about 0.05 mm to about 3 mm. In some embodiments, the average particle size may be about 0.1 mm to about 3 mm. More particularly, the dietary composition may be ground to produce a granular material having an average particle size of about 0.05 mm, about 0.1 mm, about 0.2 mm, about 0.5 mm, about 1 mm, about 2 mm, about 3 mm, or any value or range between any two of these values. In some embodiments, the dietary composition may be ground so that about 20% to 50% of the ground dietary composition is retained by a mesh having openings with a size of about 3 mm and so that about 70% to about 90% of the ground dietary composition is retained by a mesh having openings with a size of about 1 mm. In some embodiments, the dietary compositions and/or various portions thereof may have a varying distribution of particle sizes based upon the ingredients. For example, in embodiments containing one or more wheat ingredients, the particle size may be distributed so that about 95% of the ground wheat ingredients are retained by a mesh having openings with a size of about 0.0625 mm

and so that about 65% of the ground wheat ingredients are retained by a mesh having openings with a size of about 1.0 mm. In another example, such as embodiments containing one or more barley ingredients, the particle size may be distributed so that about 95% of the ground barley ingredients are retained by a mesh having openings with a size of about 0.0625 mm and so that about 60% of the ground barley ingredients are retained by a mesh having openings with a size of about 1.0 mm. The varying mesh sizes of each ingredient may be independent of mesh sizes for other ingredients.

[0051] Grinding may provide various benefits, such as improving certain characteristics of the carbohydrate component and/or the dietary composition formed therefrom. For instance, even and fine particle size may improve the mixing of different ingredients. According to certain embodiments, grinding may be configured to decrease a particle size of certain components of the dietary composition, for example, to increase the surface area open for enzymes in the gastrointestinal tract, which may improve the digestibility of nutrients, and/or to increase the palatability of the feed.

[0052] In some embodiments, the granular material or powder may be used in subsequent processes such as molding, extrusion, and/or tableting. In some embodiments, processing **120** may include drying the dietary composition and/or portions thereof. Drying may generally be completed to remove any excess water or other undesired materials, as well as to provide a material that is suitable for encapsulation, pelleting, extrusion, grinding, pressing, and/or the like.

[0053] Granular material, as used herein, refers to a conglomeration of discrete solid, macroscopic particles and is meant to encompass a wide variety of material types, shapes, and sizes. Granular material includes powders as a subset, but also includes groups of larger particles. Granular material may be particularly well-suited for tableting and encapsulation, as well as molding.

[0054] In various embodiments, additional steps may be completed **125**. The additional steps are not limited by this disclosure, and may include any additional steps necessary to prepare a dietary composition. An illustrative additional step may be to disperse the fatty acid composition in water. In some embodiments, dispersing the fatty acid composition in water may be completed so that the fatty acid composition and the carbohydrate composition can be adequately combined. Dispersing the fatty acid composition may include any amount of fatty acid and any amount of water sufficient to obtain an emulsion or a liquid suspension. For example, the fatty acid composition may include the fatty acid component and water in a volume/volume ratio from about 1:20 to about 1:1, from about 1:15 to about 2:1, from about 1:10 to about 3:1, or any value or range between any two of these values (including endpoints).

[0055] Another illustrative additional step that may be completed **125** may be to heat the fatty acid compound. In some embodiments, heating the fatty acid composition may be completed so that the fatty acid composition and the carbohydrate composition can be adequately combined. The fatty acid composition may generally be heated to a temperature at which the fatty acid composition melts to a semisolid or a liquid form. One illustrative temperature may be equal to or greater than about 40°C. Another illustrative temperature may be equal to or less than about 80°C. Another illustrative temperature may be about 40°C to about 80°C. Other illustrative temperatures may include about 40°C, about 45°C, about 50°C, about 55°C, about 60°C, about 65°C, about 70°C, about 75°C, about 80°C, or any value or range between any two of these values.

[0056] In various embodiments, a method of increasing milk fat content in ruminants may include providing at least the dietary composition as described herein to the ruminant for ingestion. In some embodiments, the dietary composition may be mixed with feed and then provided to the ruminant. In particular embodiments, the dietary composition may be mixed

with feed by an end user, such as a dairy farmer and/or the like. Thus, the end user may receive the dietary composition from a manufacturer, a distributor, and/or the like, may mix the dietary composition with the feed, and may provide the mixture to the ruminant. In other embodiments, the dietary composition may be directly fed to a ruminant without mixing with a feed.

[0057] In various embodiments, the dietary composition may be mixed with a feed in an amount such that the dietary composition is present in a premix/feed mixture in any ratio, including, for example, a ratio of about 1:25 dietary composition to feed to about 1:75 dietary composition to feed. Other illustrative ratios of dietary composition to feed may include 1:25, 1:30, 1:35, 1:40, 1:45, 1:50, 1:55, 1:60, 1:65, 1:70, 1:75, or any value or range between any two of these values (including endpoints). In some embodiments, the ratio may be dependent upon a concentration of the fatty acid component and/or the carbohydrate component in the dietary composition. Thus, for example, if the fatty acid component or the carbohydrate component is a higher concentration in the dietary composition, a higher ratio of dietary composition to feed may be used such that less dietary composition and more feed are used.

[0058] In various embodiments, the dietary composition may be provided to the ruminant in an amount such that the ruminant receives at least about 10 grams of fatty acid per kilogram of milk produced by the ruminant each day. The amount may be based on the previous day's milk production by the ruminant, an average day based on the previous week's milk production by the ruminant, an average day based on the previous month's milk production by the ruminant, an average production of milk by the ruminant when not provided with the dietary composition, and/or the like. In some embodiments, the ruminant may be provided with about 0.5 kg to about 1.5 kg of the dietary composition each day, including about 0.5 kg, about 0.75 kg, about 1.0 kg, about 1.25 kg, about 1.5 kg, or any value

or range between any two of these values (including endpoints). In some embodiments, the ruminant may be provided with additional amounts of the dietary composition to make up for portions of the dietary composition that are not consumed by the ruminant, such as amounts that are spilled by the ruminant when consuming the dietary composition, amounts that are consumed by other animals, and/or the like.

[0059] In some embodiments, providing the dietary composition to the ruminant for the ruminant to consume may result in an increase in production of milk and/or an increase in fat content of the milk produced. These increases may generally be relative to a similar ruminant that does not receive the dietary composition, an average of similar ruminants not receiving the dietary composition, an average of the milk production quantity and fat content of the same ruminant when not provided the dietary composition, and/or the like. In particular embodiments, the milk production may increase by an amount of at least about 1%. In some embodiments, the milk production may increase by an amount of about 1% to about 10%, including about 1%, about 2%, about 3%, about 4%, about 5%, about 6%, about 7%, about 8%, about 9%, about 10%, or any value or range between any two of these values. In particular embodiments, the milk fat content may increase by an amount of at least about 10%. In some embodiments, the milk fat content may increase by an amount of about 10% to about 15%, including about 10%, about 11%, about 12%, about 13%, about 14%, about 15%, or any value or range between any two of these values.

EXAMPLES

Example 1: Making a “Premix” Composition

[0060] A dietary composition to be used as a premix composition added to ruminant feed is made using a process of combining a fatty acid component and a carbohydrate component and grinding it into a granular material that can be mixed in the ruminant feed. The fatty acid component is heated to a temperature of about 50°C so that it melts and can be

easily mixed with the carbohydrate component. In addition, the carbohydrate component is ground using a standard commercial grinder so that it has an average particle size of about 2 mm.

[0061] The fatty acid component is in an amount that is about 50% by weight of the dietary composition. The fatty acid component includes about 90% by weight of a palmitic acid composition, about 10% by weight of a stearic acid composition, and no unsaturated trans fatty acids. The dietary composition also includes 50% by weight of a carbohydrate component to include additional nutrients not currently present and/or lacking in the ruminant's current feed. The carbohydrate component includes molasses, sugar beet pulp, algae and grass meal.

[0062] The resulting premix composition is packaged into bulk shipping containers that can be stored and/or shipped to distributors. The distributors will divide the bulk amounts into suitable amounts that are sold to end users such as dairy farmers and/or the like. Once the end user receives the premix composition, he/she may mix the premix composition with the ruminant feed prior to feeding the ruminant.

Example 2: Making a "Premix" Composition

[0063] A dietary composition to be used as a premix composition added to ruminant feed is made using a process of combining a fatty acid component and a carbohydrate component into a mixture. The mixture is ground into a granular material that is sufficiently sized and shaped such that can be sold and shipped to an end user such as a dairy farmer. The carbohydrate component portion of the mixture is ground using a standard commercial grinder so that it has an average particle size of about 1 mm.

[0064] The premix composition includes the fatty acid component in an amount that is about 60% by weight of the premix composition. The fatty acid component includes about 95% by weight of a palmitic acid composition, about 5% by weight of a stearic acid

composition, and no unsaturated trans fatty acids. The premix composition also includes 40% by weight of the carbohydrate component to include additional nutrients not currently present and/or lacking in the ruminant's current feed. The carbohydrate component includes molasses, sugar beet pulp, wheat bran, and oat hulls.

Example 3: Feeding a Dairy Cow

[0065] A premix-feed mixture is made by mixing 1 kg of a premix dietary composition as described above with respect to Example 1 with 45 kg of TMR for a cow to consume each day. Thus, the premix-feed mixture is a ratio of premix dietary composition to feed in an amount of about 1:45.

[0066] A dairy cow that has a normal (untreated) average daily production of 30 kg milk is provided with the premix-feed mixture each day for a month to increase the milk fat and the quantity of the milk produced. At the end of the month, it is observed that she produces 10% more milk than she did previously, and the milk that she produces contains 10% more milk fat content than the milk she produced previously.

Example 4: Providing to a Large Group of Cows

[0067] The dietary composition as described above with respect to Example 1 is mixed with a daily TMR feed and provided to a large group of cows on a commercial dairy farm to confirm its effectiveness. A group of 250 dairy cows from the commercial dairy farm are selected at random to provide a wide variety of variation in various characteristics, such as breed, weight, age of the cow, and the like. The 250 cows are divided into two groups: a sample cow group of 125 cows and a control cow group of 125 cows. Each day, the sample cow group is fed, ad libitum, a standard TMR feed with the "premix" dietary composition mixed therein. The control cow group is fed the standard TMR feed given to the sample group of cows ad libitum, but without the "premix" dietary composition. The 250 cows are

monitored for the amount of feed and/or dietary composition consumed, changes in weight, an amount of milk the cow produces each day, and the composition of the milk produced by the cow each day. Monitoring continues for a period of 60 days. A comparison of the two groups of cows over this period of time shows a statistically significant improvement from the group that consumed the "premix" composition over the control group that did not receive the "premix" composition.

Example 5: Two-month study of efficacy of "premix" dietary composition

[0068] An experiment is performed where conventional complete feed is mixed with a "premix" dietary composition according to the present disclosure. The experiment is continued for two months. The "premix" dietary composition includes the following ingredients and amounts (in percent by total weight of the solid dietary composition).

Palmitic Acid	40
Sugar beet pulp	15
Barley	15
Wheat bran	10
Oat bran	8
Molasses	2
Propylene glycol	7
Sodium bicarbonate	1
Biotin	1
Carnitine	0.5
Methionine	0.5

[0069] As shown in the table, the ingredients of the "premix" dietary composition include a fatty acid component consisting of palmitic acid and a carbohydrate component that includes sugar beet pulp, barley, wheat bran, oat bran, and molasses. Other ingredients

include a glucogenic precursor (propylene glycol), a mineral (sodium bicarbonate), a vitamin (biotin), carnitine, and an amino acid (methionine). The ingredients described above are mixed together and upon reaching an end user, are mixed with the cow's daily feed. Upon feeding the "premix"/daily feed combination to a cow, the following results are obtained from the milk produced by the cow. "Reference" refers to milk obtained from the same cow that is only fed the daily feed without the "premix" combination.

	Reference	Test Feed
Milk (kg/d)	29.5	32.5
Fat % by weight	3.98	4.43

[0070] As shown in the expected results above, milk fat concentrations and the amount of milk produced increase significantly when the cow consumes the test feed according to the present disclosure.

Example 6: Fatty Acid Composition

[0071] A fatty acid composition that is particularly suited to increase milk production and milk fat concentration in a ruminant primarily includes palmitic acid and includes, to a lesser extent, stearic acid. The fatty acid composition is mixed with a carbohydrate to obtain a premix dietary composition. The premix dietary composition is mixed with ruminant feed, as described in greater detail herein.

[0072] The following table describes such a fatty acid composition that is used to increase the volume of milk produced by a ruminant and the milk fat content of the milk produced by the ruminant.

<u>Fatty Acid</u>	<u>% of Fatty Acid Component (by weight)</u>
Palmitic Acid	≥90
Stearic Acid	≤10

Unsaturated Trans-fatty Acid	0
------------------------------	---

[0073] In the above detailed description, reference is made to the accompanying drawings, which form a part hereof. In the drawings, similar symbols typically identify similar components, unless context dictates otherwise. The illustrative embodiments described in the detailed description, drawings, and claims are not meant to be limiting. Other embodiments may be used, and other changes may be made, without departing from the spirit or scope of the subject matter presented herein. It will be readily understood that the aspects of the present disclosure, as generally described herein, and illustrated in the Figures, can be arranged, substituted, combined, separated, and designed in a wide variety of different configurations, all of which are explicitly contemplated herein.

[0074] The present disclosure is not to be limited in terms of the particular embodiments described in this application, which are intended as illustrations of various aspects. Many modifications and variations can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. Functionally equivalent methods and apparatuses within the scope of the disclosure, in addition to those enumerated herein, will be apparent to those skilled in the art from the foregoing descriptions. Such modifications and variations are intended to fall within the scope of the appended claims. The present disclosure is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled. It is to be understood that this disclosure is not limited to particular methods, reagents, compounds, compositions or biological systems, which can, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

[0075] With respect to the use of substantially any plural and/or singular terms herein, those having skill in the art can translate from the plural to the singular and/or from the

singular to the plural as is appropriate to the context and/or application. The various singular/plural permutations may be expressly set forth herein for sake of clarity.

[0076] It will be understood by those within the art that, in general, terms used herein, and especially in the appended claims (for example, bodies of the appended claims) are generally intended as “open” terms (for example, the term “including” should be interpreted as “including but not limited to,” the term “having” should be interpreted as “having at least,” the term “includes” should be interpreted as “includes but is not limited to,” et cetera). While various compositions, methods, and devices are described in terms of “comprising” various components or steps (interpreted as meaning “including, but not limited to”), the compositions, methods, and devices can also “consist essentially of” or “consist of” the various components and steps, and such terminology should be interpreted as defining essentially closed-member groups. It will be further understood by those within the art that if a specific number of an introduced claim recitation is intended, such an intent will be explicitly recited in the claim, and in the absence of such recitation no such intent is present. For example, as an aid to understanding, the following appended claims may contain usage of the introductory phrases “at least one” and “one or more” to introduce claim recitations. However, the use of such phrases should not be construed to imply that the introduction of a claim recitation by the indefinite articles “a” or “an” limits any particular claim containing such introduced claim recitation to embodiments containing only one such recitation, even when the same claim includes the introductory phrases “one or more” or “at least one” and indefinite articles such as “a” or “an” (for example, “a” and/or “an” should be interpreted to mean “at least one” or “one or more”); the same holds true for the use of definite articles used to introduce claim recitations. In addition, even if a specific number of an introduced claim recitation is explicitly recited, those skilled in the art will recognize that such recitation should be interpreted to mean at least the recited number (for example, the bare recitation of

"two recitations," without other modifiers, means at least two recitations, or two or more recitations). Furthermore, in those instances where a convention analogous to "at least one of A, B, and C, et cetera" is used, in general such a construction is intended in the sense one having skill in the art would understand the convention (for example, "a system having at least one of A, B, and C" would include but not be limited to systems that have A alone, B alone, C alone, A and B together, A and C together, B and C together, and/or A, B, and C together, et cetera). In those instances where a convention analogous to "at least one of A, B, or C, et cetera" is used, in general such a construction is intended in the sense one having skill in the art would understand the convention (for example, "a system having at least one of A, B, or C" would include but not be limited to systems that have A alone, B alone, C alone, A and B together, A and C together, B and C together, and/or A, B, and C together, et cetera). It will be further understood by those within the art that virtually any disjunctive word and/or phrase presenting two or more alternative terms, whether in the description, claims, or drawings, should be understood to contemplate the possibilities of including one of the terms, either of the terms, or both terms. For example, the phrase "A or B" will be understood to include the possibilities of "A" or "B" or "A and B."

[0077] In addition, where features or aspects of the disclosure are described in terms of Markush groups, those skilled in the art will recognize that the disclosure is also thereby described in terms of any individual member or subgroup of members of the Markush group.

[0078] As will be understood by one skilled in the art, for any and all purposes, such as in terms of providing a written description, all ranges disclosed herein also encompass any and all possible subranges and combinations of subranges thereof. Any listed range can be easily recognized as sufficiently describing and enabling the same range being broken down into at least equal halves, thirds, quarters, fifths, tenths, et cetera. As a non-limiting example, each range discussed herein can be readily broken down into a lower third, middle third and

upper third, et cetera. As will also be understood by one skilled in the art all language such as “up to,” “at least,” and the like include the number recited and refer to ranges which can be subsequently broken down into subranges as discussed above. Finally, as will be understood by one skilled in the art, a range includes each individual member. Thus, for example, a group having 1-3 cells refers to groups having 1, 2, or 3 cells. Similarly, a group having 1-5 cells refers to groups having 1, 2, 3, 4, or 5 cells, and so forth.

[0079] Various of the above-disclosed and other features and functions, or alternatives thereof, may be combined into many other different systems or applications. Various presently unforeseen or unanticipated alternatives, modifications, variations or improvements therein may be subsequently made by those skilled in the art, each of which is also intended to be encompassed by the disclosed embodiments.

CLAIMS

What Is Claimed Is:

1. A premix dietary composition comprising:
 - a fatty acid component, wherein the fatty acid component comprises at least about 90% saturated fatty acid by weight; and
 - a carbohydrate component,
wherein the fatty acid component is present in the dietary composition in an amount of at least about 30% by weight of the dietary composition.
2. The premix dietary composition of claim 1, wherein the premix dietary composition consists essentially of the fatty acid component and the carbohydrate component.
3. The premix dietary composition of claim 1, wherein the premix dietary composition consists of the fatty acid component and the carbohydrate component.
4. The premix dietary composition of claim 1, wherein the premix dietary composition is a solid in the form of a capsule, a tablet, a pellet, or a granular material having an average size of about 0.1 mm to about 3 mm.
5. The premix dietary composition of claim 1, wherein the carbohydrate component comprises at least one of sugar beet pulp, sugar cane, wheat bran, wheat middlings, wheat mill run, oat hulls, grain hulls, soya hulls, peanut hulls, wood, brewery byproduct, forages, roughages, grass meal, hay meal, alfalfa meal, alfalfa, straw, algae, or hay.
6. The premix dietary composition of claim 1, wherein the carbohydrate component comprises a carbohydrate obtained from at least one of a sugar, a starch, cellulose, or hemicellulose.
7. The premix dietary composition of claim 1, wherein the carbohydrate component comprises a carbohydrate obtained from at least one of wheat, corn, oats, sorghum, millet, or barley.

8. The premix dietary composition of claim 1, wherein the fatty acid component comprises a palmitic acid compound.

9. The premix dietary composition of claim 8, wherein the palmitic acid compound comprises free palmitic acid.

10. The premix dietary composition of claim 8, wherein the palmitic acid compound comprises a palmitic acid derivative selected from a palmitic acid ester, a palmitic acid amide, a palmitic acid salt, a palmitic acid carbonate, a palmitic acid carbamates, a palmitic acid imide, a palmitic acid anhydride, and a combination thereof.

11. The premix dietary composition of claim 1, wherein the fatty acid component comprises a palmitic acid compound in an amount of at least about 60% by weight of the fatty acid component.

12. The premix dietary composition of claim 1, wherein the fatty acid component comprises a palmitic acid compound in an amount of at least about 80% by weight of the fatty acid component.

13. The premix dietary composition of claim 1, wherein the fatty acid component comprises a palmitic acid compound in an amount of at least about 90% by weight of the fatty acid component.

14. The premix dietary composition of claim 1, wherein the fatty acid component comprises an unsaturated trans fatty acid in an amount of about 5% or less by weight of the fatty acid component.

15. The premix dietary composition of claim 1, wherein the fatty acid component comprises an unsaturated trans fatty acid in an amount of about 4% or less by weight of the fatty acid component.

16. The premix dietary composition of claim 1, wherein the fatty acid component comprises an unsaturated trans fatty acid in an amount of about 3% or less by weight of the fatty acid component.

17. The premix dietary composition of claim 1, wherein the fatty acid component comprises an unsaturated trans fatty acid in an amount of about 2% or less by weight of the fatty acid component.

18. The premix dietary composition of claim 1, wherein the fatty acid component comprises an unsaturated trans fatty acid in an amount of about 1% or less by weight of the fatty acid component.

19. The premix dietary composition of claim 1, wherein the fatty acid component comprises an unsaturated trans fatty acid in an amount of about 0.5% or less by weight of the fatty acid component.

20. The premix dietary composition of claim 1, wherein the fatty acid component is substantially free of unsaturated trans fatty acids.

21. The premix dietary composition of claim 1, wherein the saturated fatty acid is present in the fatty acid component in an amount of at least about 95% by weight of the fatty acid component.

22. The premix dietary composition of claim 1, wherein the saturated fatty acid is present in the fatty acid component in an amount of at least about 98% by weight of the fatty acid component.

23. The premix dietary composition of claim 1, wherein the fatty acid component has a melting point of about 60°C to about 80°C.

24. The premix dietary composition of claim 1, wherein the fatty acid component has a melting point of about 63 °C to about 65 °C.

25. The premix dietary composition of claim 1, wherein the fatty acid component has a melting point equal to or greater than about 40°C.

26. The premix dietary composition of claim 1, wherein the fatty acid component comprises about 30% or less of a stearic acid compound by weight of the fatty acid component.

27. The premix dietary composition of claim 1, wherein the fatty acid component is present in the dietary composition in an amount of at least about 50% by weight of the dietary composition.

28. The premix dietary composition of claim 1, further comprising at least one glucogenic precursor comprising glycerol, propylene glycol, molasses, propionate, glycerine, propane diol, calcium propionate, or a combination thereof.

29. The premix dietary composition of claim 1, further comprising at least one vitamin comprising vitamin A, vitamin B, vitamin C, vitamin D, vitamin E, vitamin K, or a combination thereof.

30. The premix dietary composition of claim 1, further comprising at least one vitamin comprising at least one B vitamin selected from thiamine, riboflavin, niacin, pantothenic acid, pyridoxine, biotin, folic acid, cobalamin, choline, and a combination thereof.

31. The premix dietary composition of claim 1, further comprising carnitine.

32. The premix dietary composition of claim 1, further comprising at least one amino acid comprising leucine, lysine, histidine, valine, arginine, threonine, isoleucine, phenylalanine, methionine, tryptophan, any derivative thereof, or a combination thereof.

33. The premix dietary composition of claim 1, further comprising at least one mineral comprising an ion of calcium, sodium, magnesium, potassium, phosphorus, zinc, selenium, manganese, iron, cobalt, copper, iodine, molybdenum, or a combination thereof.

34. The premix dietary composition of claim 1, further comprising at least one mineral comprising at least one sodium salt selected from monosodium phosphate, sodium acetate, sodium chloride, sodium bicarbonate, disodium phosphate, sodium iodate, sodium iodide, sodium tripolyphosphate, sodium sulfate, sodium selenite, and a combination thereof.

35. The premix dietary composition of claim 1, further comprising at least one mineral comprising at least one calcium salt selected from calcium acetate, calcium carbonate, calcium chloride, calcium gluconate, calcium hydroxide, calcium iodate, calcium iodobehenate, calcium oxide, anhydrous calcium sulfate, calcium sulfate dehydrate,

dicalcium phosphate, monocalcium phosphate, tricalcium phosphate, and a combination thereof.

36. The premix dietary composition of claim 1, further comprising at least one mineral comprising at least one magnesium salt selected from magnesium acetate, magnesium carbonate, magnesium oxide, magnesium sulfate, and a combination thereof.

37. The premix dietary composition of claim 1, further comprising at least one mineral comprising at least one cobalt salt selected from cobalt acetate, cobalt carbonate, cobalt chloride, cobalt oxide, cobalt sulfate, or and a combination thereof.

38. The premix dietary composition of claim 1, further comprising at least one mineral comprising at least one manganese salt selected from manganese carbonate, manganese chloride, manganese citrate, manganese gluconate, manganese orthophosphate, manganese oxide, manganese phosphate, manganese sulfate, and a combination thereof.

39. The premix dietary composition of claim 1, further comprising at least one mineral comprising at least one potassium salt selected from potassium acetate, potassium bicarbonate, potassium carbonate, potassium chloride, potassium iodate, potassium iodide, potassium sulfate, and a combination thereof.

40. The premix dietary composition of claim 1, further comprising at least one mineral comprising at least one iron salt selected from iron ammonium citrate, iron carbonate, iron chloride, iron gluconate, iron oxide, iron phosphate, iron pyrophosphate, iron sulfate, reduced iron, and a combination thereof.

41. The premix dietary composition of claim 1, further comprising at least one mineral comprising at least one zinc salt selected from zinc acetate, zinc carbonate, zinc chloride, zinc oxide, zinc sulfate, and a combination thereof.

42. The premix dietary composition of claim 1, further comprising at least one mineral comprising copper sulfate, copper oxide, selenium yeast, a chelated mineral, or a combination thereof.

43. The premix dietary composition of claim 1, further comprising at least one nitrogen source material comprising an oilseed meal.

44. The premix dietary composition of claim 43, wherein the oilseed meal is selected from at least one of soy meal, bean meal, rapeseed meal, sunflower meal, coconut meal, olive meal, linseed meal, grapeseed meal, cottonseed meal, camelina meal, mustard seed meal, crambe seed meal, safflower meal, rice meal, peanut meal, corn gluten meal, corn gluten feed, distillers dried grains, distillers dried grains with solubles, or wheat gluten.

45. A method of preparing a premix dietary composition for ruminants, the method comprising:

combining a fatty acid component and a carbohydrate component to form a mixture; and

processing the mixture into a tablet, a capsule, a pellet, or a granular material, wherein the fatty acid component is present in the dietary composition in an amount of at least about 30% by weight of the dietary composition.

46. The method of claim 45, wherein combining the fatty acid component and the carbohydrate component consists of combining substantially only the fatty acid component and the carbohydrate component.

47. The method of claim 45, wherein combining the fatty acid component and the carbohydrate component consists of combining only the fatty acid component and the carbohydrate component.

48. The method of claim 45, wherein processing the mixture comprises processing the mixture into a tablet, a capsule, a pellet, or a granular material having an average size of about 0.1 mm to about 3 mm.

49. The method of claim 45, further comprising grinding the carbohydrate component prior to combining with the fatty acid component.

50. The method of claim 45, further comprising dispersing the fatty acid component in water to obtain a liquid suspension prior to combining with the carbohydrate component.

51. The method of claim 45, further comprising dispersing the fatty acid component in water to obtain an emulsion prior to combining the carbohydrate component.

52. The method of claim 45, further comprising heating the fatty acid component to obtain a melted fatty acid component prior to combining with the carbohydrate component.

53. The method of claim 45, wherein processing comprises at least one of pressing, extruding, grinding, or pelleting the mixture into the tablet, the capsule, the pellet, or the granular material.

54. The method of claim 45, further comprising drying the mixture.

55. The method of claim 45, wherein the fatty acid component comprises saturated fatty acid in an amount of at least about 90% by weight of the fatty acid component.

56. The method of claim 45, wherein the fatty acid component comprises saturated fatty acid in an amount of at least about 95% by weight of the fatty acid component.

57. The method of claim 45, wherein the fatty acid component comprises saturated fatty acid in an amount of at least about 98% by weight of the fatty acid component.

58. The method of claim 45, wherein the fatty acid component is substantially free of unsaturated trans fatty acids.

59. The method of claim 45, wherein the fatty acid component comprises a glucogenic precursor comprising at least one of glycerol, propylene glycol, molasses, propionate, glycerine, propane diol, or calcium propionate.

60. The method of claim 45, further comprising combining at least one B vitamin with the mixture, wherein the B vitamin comprises thiamine, riboflavin, niacin, pantothenic acid, pyridoxine, biotin, folic acid, cobalamin, choline, or a combination thereof.

61. The method of claim 45, further comprising combining at least one vitamin with the mixture, wherein the vitamin comprises vitamin A, vitamin B, vitamin C, vitamin D, vitamin E, vitamin K, or a combination thereof.

62. The method of claim 45, further comprising combining at least one amino acid with the mixture, wherein the amino acid comprises leucine, lysine, histidine, valine, arginine, threonine, isoleucine, phenylalanine, methionine, tryptophan, any derivative thereof, or a combination thereof.

63. The method of claim 45, further comprising combining at least one mineral with the mixture, wherein the at least one mineral comprises calcium, sodium, magnesium, potassium, phosphorus, zinc, selenium, manganese, iron, cobalt, copper, iodine, molybdenum, or a combination thereof.

64. The method of claim 45, further comprising combining at least one nitrogen source material with the mixture, wherein the source material comprises an oilseed meal selected from at least one of soy meal, bean meal, rapeseed meal, sunflower meal, coconut meal, olive meal, linseed meal, grapeseed meal, cottonseed meal, camelina meal, mustard seed meal, crambe seed meal, safflower meal, rice meal, peanut meal, corn gluten meal, corn gluten feed, distillers dried grains, distillers dried grains with solubles, wheat gluten, or a combination thereof.

65. The method of claim 45, further comprising combining a cellulosic material with the mixture.

66. A method of increasing milk fat content in ruminants, the method comprising: combining a premix dietary composition with a feed to obtain a mixture, wherein the premix dietary composition comprises:

a fatty acid component, wherein the fatty acid component is present in the premix dietary composition in an amount of at least about 30% by weight of the premix dietary composition, and

a carbohydrate component; and
providing the mixture to a ruminant for ingestion.

67. The method of claim 66, wherein providing the mixture to the ruminant comprises providing an amount of the mixture such that the ruminant consumes about 0.5 kg to about 1.5 kg of the premix dietary composition daily.

68. The method of claim 66, wherein providing the mixture to the ruminant comprises providing the mixture to the ruminant at an amount such that the ruminant receives at least about 10 grams of fatty acid per kilogram of milk produced by the ruminant per day.

69. The method of claim 66, wherein providing the mixture to the ruminant results in at least one of an increase in production of milk by the ruminant or an increase in a fat

content in the milk produced by the ruminant, relative to a similar ruminant not provided with the mixture.

70. The method of claim 66, wherein providing the mixture to the ruminant results in at least one of: an at least about 1% increase in production of milk by the ruminant or an at least about 10% increase in a fat content in the milk produced by the ruminant, relative to a similar ruminant not provided with the mixture.

71. A premix composition for ruminants, the premix composition comprising:
a fatty acid component comprising a palmitic acid compound in an amount of at least about 90% by weight of the fatty acid component; and
a carbohydrate component,
wherein the fatty acid component is present in the premix composition in an amount of at least about 30% by weight of the premix composition,
wherein the premix composition is a solid having an average size of about 0.1 mm to about 3 mm, and
wherein the premix composition is configured to be mixed with a ruminant feed and provided to ruminants.

72. The dietary composition of claim 71, wherein the fatty acid component consists essentially of the palmitic acid compound.

73. The dietary composition of claim 71, wherein the fatty acid component consists of the palmitic acid compound.

1/1




FIG. 1