

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
23 July 2009 (23.07.2009)

PCT

(10) International Publication Number
WO 2009/090227 A2

(51) International Patent Classification:

A61K 38/00 (2006.01) *C07K 14/705* (2006.01)
A61K 38/08 (2006.01) *C07K 1/22* (2006.01)
A61P 25/28 (2006.01)

(74) Agent: MEYERS, Hans-Wilhelm; von Kreisler Selting Werner, P.O. Box 10 22 41, 50462 Cologne (DE).

(21) International Application Number:

PCT/EP2009/050446

(22) International Filing Date: 15 January 2009 (15.01.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

08100510.0 15 January 2008 (15.01.2008) EP

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (for all designated States except US): MDC MAX-DELBRÜCK-CENTRUM FÜR MOLEKULARE MEDIZIN BERLIN-BUCH [DE/DE]; Robert-Rössle-Strasse 10, 13125 Berlin-Buch (DE).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant and

(72) Inventor: KUNZE, Rudolf [DE/DE]; Bleicheroder Strasse 31D, 13178 Berlin (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): WALLUKAT, Gerd [DE/DE]; Wolkensteinstrasse 4, 13129 Berlin (DE). ROSENTHAL, Peter [DE/DE]; Höher Weg 27, 58511 Lüdenscheid (DE). STRAUBE, Richard [DE/DE]; Westerfelder Strasse 65, 58515 Lüdenscheid (DE).

Published:

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

WO 2009/090227 A2

(54) Title: PEPTIDES HAVING BINDING AFFINITY TO AN ANTIBODY WHICH RECOGNIZES AN EPITOPE ON AN $\alpha 1$ LOOP 2 OR $\beta 2$ LOOP 1 OF AN ADRENORECEPTOR

(57) Abstract: A peptide having an ED₅₀ of less than 500 nm, in particular 10 nM to an antibody which recognizes an epitope on an extracellular $\alpha 1$ loop 2 or $\beta 2$ loop 1 of a human adrenoceptor wherein the antibody's binding to the epitope results in an increase of γ -secretase activity, an increased release of β -amyloid molecules and/or cellular dysfunction of cerebral blood vessel cells, glia cells or neurons, wherein the ED₅₀ value is measured by bioassay.

Peptides having binding affinity to an antibody which recognizes an epitope on an $\alpha 1$ loop 2 or $\beta 2$ loop 1 of an adrenoreceptor

The invention pertains to a peptide having a binding affinity an antibody which

5 recognizes an epitope on an $\alpha 1$ loop 2 or $\beta 2$ loop1 of an adrenoreceptor, an isolated nucleic acid molecule, solid phases for affinity chromatography or solid-phase extraction consisting of organic, inorganic, synthetic polymers or of mixed polymers, a vector comprising a nucleic acid molecule, a host cell comprising the vector, a device for removing immunoglobulins from

10 immunoglobulin-containing fluids on solid phases, a pharmaceutical composition comprising a nucleic acid molecule, a peptide and/or a solid phase of the invention together with a pharmaceutically tolerable carrier, a kit comprising a nucleic acid molecule of the invention, a vector of the invention, a host cell of the invention, as well as an apparatus for chromatography,

15 comprising peptides of the invention.

Alzheimer's disease (AD), which is named after its discoverer Alois Alzheimer (1864-1915), is a progressive degenerative disease. It is characterized by changes in the cerebral cellular structure and function caused by deposits of

20 aggregated proteins with extensive consequences for essential brain functions. The following clinical and cognitive characteristics are essential:

- disorders of the short-term memory at first and of the long-term memory later on
- disorders of the perception of simple recognition patterns
- 25 • progressive loss of speech and articulateness
- progressive loss of sense of smell, of spatial orientation and decline of ability to perform routine tasks.

The diagnosis "AD" is obtained by different methods, for example, with imaging methods, such as computer tomography (CT) or magnetic resonance tomography (MRT) and by the chemical determination in the laboratory of the concentrations of the involved pathological molecules in the cerebrospinal fluid.

5 Electroencephalography (EEG) may also be resorted to. Cognitive tests, such as the mini mental status test (MMST) are increasingly gaining importance.

If untreated, the disease is a "one-way street" which unavoidably results in a deterioration of the clinical image over several years to end with death. The more advanced the stage of the disease, the more care and cost intensive the

10 patient becomes. This also holds for other forms of dementia (AD accounting for about two thirds of all dementia cases), which will not be dealt with here, however.

The prevalence of the disease is highly age-dependent. Among those of 60 to 70 years of age, it is about 1%, and among those of 70 to 80 years of age, it is as

15 high as about 10%. Since life expectancy of the humans in Germany is continuously increasing, another rise of the number of cases is to be expected.

Currently, the annual incidence for AD in Germany is 130 000 to 140 000 incidences (Hallauer, 2003), a number that needs interpretation, however. Since the mild form is little conspicuous, the majority of the patients will probably have

20 a medium severity of the disease.

Some years ago, the disease was still considered a stroke of fate that could not be cured. Severe cases have been treated with sedatives and neuroleptics for sedation. In the meantime, drugs of the first generation are available; they are mainly employed in the medium, moderate stage. Examples include approved

25 pharmaceuticals, such as Axura (active ingredient: memantine) or Exelon (active ingredient: rivastigmine). According to the published state of the study, these agents have a good effectiveness for mild and medium severities of AD.

However, the clinical effectiveness of Exelon and other choline-esterase inhibitors is denied according to metaanalyses. Although approved medicaments are available on the market and the preparations are prescribable, by far not all patients are treated.

- 5 Some prophylactic effect on AD seems to be exerted by anti-inflammatory drugs, such as acetylsalicylic acid (aspirin), ibuprofen, indomethazine or COX2 inhibitors. Evidently, local and systemic inflammations or inflammation mediators play a disease-promoting role in the organism/brain, probably due to an acceleration of neurofibril formation.
- 10 Cholesterol-reducing agents, such as statins, also seem to have a favorable influence on the risk of developing AD. At least, patients obtaining cholesterol-reducing agents exhibit a lesser incidence of AD. However, no hard scientific data are available to support this.

15 The situation is similar with hypotensive agents, such as inhibitors of angiotensin converting enzyme (ACE). In addition to a direct effect of these active substances, the reduction of the systolic blood pressure is also generally associated with the preventive effects observed.

20 Immunizations of patients with β -amyloid are in the experimental stage of a therapy. In response, the immune system produces antibodies which then destroy amyloid plaques by a local inflammatory reaction. This approach seems to be promising, but is not undisputed, because it involves high risks. Probably, the individual inflammatory immune responses that are used therapeutically cannot be dosed as precisely as would be required. Other experimental-therapeutic approaches are based on the compensation of a cerebral deficiency 25 of copper ions, which evidently plays a role in plaque formation.

The question of whether medicaments are basically able to achieve a standstill only, or also a clinical improvement of the patients' cognitive state is currently

still unanswered. The currently available means stabilize the clinical picture rather than improve it.

The pathogenesis of AD itself has not yet been finally clarified. A number of risk factors have been recognized. These include age, hypertension, high cholesterol 5 levels, overweight, apolipoprotein E4, copper deficiency, certain genetic predispositions, and possibly oxidative stress. The development of the disease is very probably multifactorial.

There is a plethora of references about molecules that play or could play a role in the development and progress of AD. As to a definition of term, it may be 10 mentioned here that most of the so-called pathological molecules are components of physiological structures and processes in the healthy organism. They are considered pathological if they exceed concentration limits, in the plasma or locally in the tissue, intercellular or intracellular. This holds for both dissolved proteins and aggregated molecules (e.g., β -amyloid in the form of 15 plaque).

The aggregation or deposit of two different, per se physiological protein molecules in the brain in intercellular or intracellular spaces causes fundamental neuronal functional disorders in signal transmission and the interaction of neurons. These proteins are essentially the neurotoxic β -amyloid peptide 20 (aggregated as a polymer in the form of amyloid plaque), and the Tau protein, a polymer aggregated as neurofibrillary tangles.

So-called amyloids are evidently involved causally in the development of AD. Thereafter, an increased formation of amyloid- β 40/42 occurs by enzymatic 25 cleavage of the molecule from the membrane of the nerve cells. This results in a locally increased concentration of single β -amyloid molecules, β -amyloid oligomers and its aggregation into amyloid plaques. These plaques disrupt the normal transmission of stimuli by the nerves and communication between the cells. The cleavage of aggregatable β -amyloids from the precursor molecules is

effected by the so-called γ -secretase. The activity of this enzyme is controlled by the $\beta 2$ -adrenoceptor.

The plaques in turn may also result in the generation of neurofibrillary tangles of the Tau protein, which normally is physiologically localized in the axonal tracts of 5 the nerve cells. The deposit of these aggregates takes place in a wide variety of cerebral areas. The process of plaque formation proceeds over many years from the beginning to the onset of the disease.

Both plaques and neurofibrillary tangles can be visualized well by microscopy in brain sections of dead AD patients and are an important part of diagnostics.

10 A pathomechanistic role of other molecules is also discussed in parallel, including the role of apolipoprotein E4. Further, rheologically relevant plasma proteins seem to be involved in the development of the clinical picture at least indirectly. These include increased or moderately increased systemic levels and local concentrations of alpha-2 macroglobulin, fibrinogen, but also oxidized plasma 15 proteins from oxidative stress reactions, such as oxidized low density lipoprotein (oxLDL).

20 The sum of effects of these molecules may contribute to a reduced blood circulation through the brain areas and thus induce the beginnings of the disease and promote its development and maintenance. Table 1 provides a survey of important pathological molecules.

Table 1: Survey of pathological molecules of Alzheimer's disease in the blood plasma

Pathological molecules in AD	MW	Comment
fragments of β -amyloid oligomers of β -amyloid	< 4,000 > 4,000	fragments of β -amyloid having neurotoxic properties
β -amyloid or fragments thereof bound to albumin	> 70,000	albumin is the most important plasma protein for the physiological clearance of noxious substances
apolipoprotein E, especially E4	ca. 34,000 (monomer)	predisposed for AD, may be contained in plaques
proinflammatory cytokines IL1 β , IL6, TNF α	16,000-50,000 (monomer and trimer, respectively)	increased levels in brain, neurotoxic
C-reactive protein (CRP)	120,000	involved in plaque formation, enhances cell degradation in necroses
fibrinogen	340,000	rheologically effective due to its size, reduces circulation through capillaries
L-kynurenine	208	neurotoxic precursor of 3OH-K, interacts with glutamate receptors
homocysteine, unbound	268	inhibits the degradation of ADMA and thus reduces vasodilatation
asymmetric dimethylarginine (ADMA)	202	inhibits NO synthetase and thus results in hypoperfusion
oxidative stress (reaction products)	variable, also microparticles	can be induced directly by β -amyloid, but are also formed by hypoperfusion

The most extensively studied group of diseases associated with agAAB against

5 GPCR are cardiac diseases. In myocarditis, dilated cardiomyopathy, peripartum cardiomyopathy, Chagas' disease, and ventricular arrhythmias, agAAB against the β_1 -AR can be detected in the sera of patients with clinically diagnosed diseases.

DCM and myocarditis are severe heart disorders, and it has been hypothesized that a chronic myocarditis may develop to a DCM. In both diseases an infection with an enterovirus or others may be the primary etiological agent, but the inflammatory response may have developed to an autoimmune 5 disease. In a later stage, the infectious agent may not be detectable, and even the inflammatory response may have declined, however, a chronic progressing heart failure is diagnosed. The DCM is characterized by a severe heart failure. In a subset of DCM, the idiopathic DCM, in which other etiological agents can be excluded, the prevalence of the β_1 -adrenergic AAB of up to 80 % is 10 remarkable.

Myocarditis and DCM are preferentially treated by antagonists of the β_1 -adrenergic and renin-angiotensin system and physical rest.

The therapeutic goal is to reduce cardiac overstimulation and prevent arrhythmias 15. The therapeutic use of betablockers in reducing adrenergic overstimulation is accompanied by a significant decrease in cardiac death and hospitalization in the patients subject to the study. Ventricular arrhythmias are a major cause of sudden death in dilated DCM and ventricular tachycardias have been shown to be strongly related with the detection of agAAB to the β_1 - 20 AR.

The chronic progression of heart failure in DCM leads to a stage of the disease where successful medication is not possible anymore. In this stage, only heart transplantation can be considered as a therapy sustainable for some year. DCM is the cause of about half of the heart transplantations. An extension of 25 therapeutic options might very likely prevent a substantial number of premature deaths in the case of severe heart failure.

Similar to DCM, peripartum cardiomyopathy is characterised by a dilation of the left ventricle and may develop into a life threatening disorder. The incidence is about 1:1000 in South African women, while the disease is rare 30 among white populations in western countries. The etiology of peripartum

cardiomyopathy is largely unknown. By definition it is diagnosed between the last month before and the 4th month after delivery. Recently, the presence of agAAB against the β_1 -AR in all sera of a sample of 10 patients suffering from peripartum cardiomyopathy was shown.

5 Chagas' disease is caused by the parasite *Trypanosoma cruzi* and very common in South America. Among other symptoms, the patients often develop a myocarditis as part of the dysautonomic syndrome that affects the whole cholinergic system. There is strong evidence that the dysautonomic syndrome is caused also by functional AAB against the muscarinic M₂-
10 acetylcholine receptor. The prevalence of agAAB against the M₂-receptor ranges from 50-94 %. In Chagas' disease, AAB against the β_1 -AR can also be found in about 53 % of the cases. This situation may represent the antigenic mimicry of originally the same antigen from *T. cruzi*, with a functional antagonism as an epiphomenon .

15 With respect to the pathogenetic nature of the described agAAB, only the signalling events with evidence of clinical relevance will be considered in the following paragraph. Activation of the β -AR leads to an elevation of cAMP by the activation of adenylate cyclase by stimulating G_s -proteins in the G-protein trimer. On the other hand, agonistic ligand binding to muscarinic and nicotinic
20 acetylcholine receptors inhibits adenylate cyclase by inhibiting G_i -proteins. However, this simple pattern has to be extended for the individual receptors.

Autoantibodies from sera of myocarditis or DCM patients show a positive chronotropic effect to the same extent as the agonist isoprenaline. Isoprenaline clearly increases the cAMP concentration in the cell and activates
25 the L-type Ca²⁺-channels. It has been shown that AAB from DCM patients activate the L-type Ca²⁺-channels. On the other hand, the effect of AAB on the accumulation of cAMP was rather marginal . It has been shown, however, that the cytosolic as well as the membrane bound protein kinase A in cardiomyocytes is activated upon AAB-stimulation. A monoclonal anti- β_1 -

receptor AB that induces apoptosis in rat cardiomyocytes also acts via cAMP and protein kinase A, MAP-kinases, esp. p38-kinase are activated downstream of several GPCRs but also in the signal transduction cascade of inflammatory cytokines. Thus, it may be suggested that stimulation of the β_1 -

5 AR by agAAB may also have a proinflammatory effect .

There are also important cAMP-independent signalling pathways. L-type Ca^{2+} -channels are not only activated through phosphorylation by protein kinase A, but also directly through subunits of G-proteins. Thus, the stimulation of Ca^{2+} -influx may be a significant pathway for the pathogenetic role of AAB in
10 autoimmune myocarditis and DCM. The action potential on rat and human cardiomyocytes is prolonged by the administration of agAAB against the β_1 -AR and activates in turn the electrogenic $\text{Na}^+/\text{Ca}^{2+}$ -exchanger, resulting in electrical instability of the heart. Intracellular calcium overload is also known
15 as a key event of remodelling, as is the β_1 -adrenergic stimulation. In a transgenic rat model, the β_1 -AR was crucial for the reorganization of actin mediated by enhanced transcription of the atrial natriuretic factor.

Taken altogether, β -adrenergic overstimulation, intracellular calcium overload, and the partial lack of down regulation of the β_1 -AR may be involved in the pathogenesis of myocarditis and DCM.

20 Results from animal experiments support the in vitro data about the pathogenic relevance of agAAB obtained in tissue culture experiments. Jahns et a. could induce agonistic antibodies in rats by immunisation with the second extracellular loop of the β_1 -AR. The animals developed a dilated cardiomyopathy similar to that of human dilated cardiomyopathy. The transfer
25 of the induced antibody/ immunoglobulins into healthy animals leads to a DCM also.

The induction of agonistic antibody against the β_1 -AR was also performed in rabbits and results in the occurrence of agAAB as well as DCM. A removal of agAAB from the animal's blood plasma by specific immunoabsorption using a

- 10 -

peptid column leads to an improvement of the heart muscle function, measured as left ventricular ejection fraction.

More as a dozen publications report about the theapeutical benefit of the theapeutical apheresis (immunoabsorption) in patients suffering from DCM. In 5 a clinical stage of the cardiac disease where no successful cardiac drug medication is possible, the theapeutical apheresis leads to a long lasting decline of the agAAB, a disappearance of activated inflammatory leukocytes from the heart muscle tissue and an increase of essential cardiac functions.

Other examples of diseases which are associated or caused by agAAB 10 preeclampsia and vascular necrotic kidney rejection (agAAB against the Angiotensin-1 receptor or the open angle glaucoma (agAAB against the β 2-adrenoceptor.

WO-A-02/093174 discloses a method and device for treating AD. The method involves the removal of circulating autoantibodies of a biochemical marker of 15 markers, specifically human glial fibrillary acidic protein (GFAP) and glyceraldehyde-3-phosphate dehydrogenase (GAPDH), in body fluid, preferably blood or a blood product. The invention further includes a device or process of immune system modulation effective for autoantibody removal.

D.M. Walsh et al. discloses in Biochemical Society Transactions (2005) Vol. 33, 20 part 5, PP 1087-1090 that soluble oligomers of A β are among the earliest effectors of Alzheimer's disease. They report about testing of compounds directed at three prominent amyloid-based therapeutic targets, inhibition of the secretases responsible for A β production, inhibition of A β aggregation and immunization against A β . In each case, compounds capable of reducing 25 oligomer production or antibodies that avidly bind A β oligomers also ameliorate the synaptotoxic effects of these natural, cell-derived oligomers.

Norman R. Relkin et al. discloses in *Neurobiology of Aging* (2008), Febr.20, E-publ. that natural occurring antibodies against β A are able to reduce the β A level in the body, and their intravenous application may improve the brain function in patients with mild AD.

5 Dennis J. Selkoe et al. Report in *Annu. Rev. Pharmacol. Toxicol.* 2003, 43:545-84 discusses molecular understanding predicting amyloid-based therapeutics. Degenerative diseases of the brain were long considered among the most obscure and intractable of human maladies. However, recent advances in understanding their mechanisms have brought us to the verge of potential

10 10 disease-modifying agents. This progress is perhaps best exemplified by the case of Alzheimer's disease. The application of molecular pathology and genetics has led to the recognition that the four genes implicated to date in familial Alzheimer's disease all chronically elevate cerebral levels of the Amyloid β -protein (A β). Accordingly, small molecule inhibitors of the β - and γ -secretases, the proteases that generate A β from its precursor, are under active development, and some have shown in vivo efficacy in mouse models. An alternative approach, active or passive immunization against A β , has received extensive pre-clinical validation in mice, but an effective preparation free of significant side effects in humans is still awaited. Several other

15 20 potential therapies are also reviewed here. If one or more of these varied approaches is ultimately proven to slow or prevent dementia, Alzheimer's disease will become a salient example of the successful application of reductionist biology to the most complex of organs, the human cerebral cortex.

25 EP-A-1832600 discloses an invention which relates to nucleic acid molecules encoding peptides which interact with autoantibodies associated with glaucoma, to the peptides themselves, to a pharmaceutical composition comprising said nucleic acid molecules and peptides, and to the use of said peptides - especially in apheresis - for the treatment of glaucoma. The

peptides represent an epitope which is a part of the extracellular loop2 of the β 2 adrenergic receptor.

WO-A-02/38592 discloses an invention which relates to peptides having a high affinity to immunoglobulins.

5 Y. Magnusson et al. report in Clin. Exp. Immunol. (1989) 78, 42-48 about an antigenic analysis of the second extra-cellular loop of the human beta-1 adrenergic receptor which is different to the beta-2 adrenergic receptor.

Yanxiang Ni et al. report about activation of β_2 -adrenergic receptor stimulating γ -secretase activity and accelerating amyloid plaque formation. They report
10 that β_2 -AR activation can stimulate γ -secretase activity and amyloid plaque formation and speculates that abnormal activation of β_2 -AR might contribute to A β accumulation in Alzheimer's disease pathogenesis.

An object of the invention was to provide a compound which is able to be used in the treatment of AD. Another object was to provide a compound which is
15 able to reduce the β 2-adrenoceptor mediated γ -secretase activity in individuals suffering from higher levels of γ -secretase due to auto-antibodies against adrenoreceptor.

The invention is based on the finding of agonistic autoantibodies (agAAB) directed against the G protein coupled receptors (GPCR), β 2 and α 1 adrenoceptor (AR), in the blood plasma of patients suffering from diseases
20 caused by an increase of γ -secretase activity, an increased release of β -amyloid molecules and/or cellular dysfunction of cerebral blood vessel cells, glia cells or neurons. In particular, the disease is AD.

In one embodiment of the present invention the object is solved by removing
25 immunoglobulins from patients being suspect of suffering from or suffering from a disease caused by an increase of γ -secretase activity, an increased

release of β -amyloid molecules and/or cellular dysfunction of cerebral blood vessel cells, glia cells or neurons. The removal of immunoglobulins may be performed by adsorption of the respective immunoglobulins on a solid support. These methods as such are well known in the art.

- 5 In a further embodiment the invention is concerned with the use of an immunoabsorption device for manufacturing a device for the treatment of a disease caused by an increase of γ -secretase activity, an increased release of β -amyloid molecules and/or cellular dysfunction of cerebral blood vessel cells, glia cells or neurons by removal of immunoglobulins from patients.
- 10 In another embodiment of the invention the object is solved by a peptide having an ED_{50} of less than 500 nM, in particular 10 nM to an antibody which recognizes an epitope on a human $\alpha 1$ loop 2 and $\beta 2$ loop 1 of an adrenoreceptor wherein the antibody's binding to the epitope results in increase of γ -secretase activity and/or increased pathological β -amyloid
- 15 release and/or cellular dysfunction of cerebral blood vessel cells, glia cells or neurons, wherein the ED_{50} value is measured by the bioassay which was used to identify the agAAB.

In particular, the peptide of the invention recognizes the human $\alpha 1$ loop epitopes of α -adrenoreceptor subtypes A, B, or C or combinations thereof.

In one embodiment of the invention the peptide of the invention consists of an amino acid sequence

X1-X2-X3-X4-X5-X6-X7-X8-X9

25 wherein

a)

X1 = amino group, amide, acetyl group, biotin group, marker, spacer, linker, C, GKK, SGKK or deletion,

X2 = A, G, α -Abu, F, Hph, Hyp, Nal, P, Pip, S, V, W, Y, or deletion,

X3 = Hyp, P, Pip, S, T, Y, or deletion,

X4 = Aad, Asu, D, E, Har, K, N, Orn, Q, R,

X5 = Aad, Asu, D, E, N, Q,

X6 = Aad, Asu, D, E, Har, K, N, Orn, Q, R,

5 X7 = F, Hph, Hyp, Nal, P, Pip, W, Y, or deletion,

X8 = A, α -Abu, β -A, G, S, T, V, Y, or deletion,

X9 = amide, hydrazide, azide, carbamate, marker, spacer, linker, C, GKK, SGKK or deletion, or

b)

10 X1 = amino group, amide, acetyl group, biotin group, marker, spacer, linker, C, GKK, SGKK or deletion,

X2 = F, Hph, Hyp, Nal, P, Pip, W, Y,

X3 = A, α -Abu, Dab, G, Hyl, K, Orn, R, S, V,

X4 = Aad, Asu, D, E, Hyp, N, P, Pip, Q,

15 X5 = Aad, Asu, D, E, Hyp, N, P, Pip, Q,

X6 = A, α -Abu, G, Hyl, K, Orn, R, S, V

X7 = F, Hph, Hyp, Nal, P, Pip, W, Y,

X8 = γ -Abu, Ahx, β -A, G, or deletion,

20 X9 = amide, hydrazide, azide, carbamate, marker, spacer, linker, C, GKK, SGKK or deletion, or

c)

X1 = amino group, amide, acetyl group, biotin group, marker, spacer, linker, C, GKK, SGKK or deletion,

25 X2 = A, α -Abu, Dab, G, Hyl, K, Orn, R, or deletion,

- 15 -

X3 = A, C, F, Hph, M, Nal, S, W, Y,

X4 = F, S, T, W, Y,

X5 = F, S, T, W, Y,

X6 = A, C, F, Hph, M, Nal, S, W, Y,

5 X7 = A, α -Abu, Dab, G, Hyl, K, Orn, R,

X8 = Abu, Ahx, β -A, G, or deletion,

X9 = amide, hydrazide, azide, carbamate, marker, spacer, linker, C, GKK, SGKK or deletion, or

10 d)

X1 = amino group, amide, acetyl group, biotin group, marker, spacer, linker, C, GKK, SGKK or deletion,

X2 = Aad, Asu, D, E, F, Hph, N, Nal, Q, W, Y,

X3 = A, C, G, Hcy, S, Sec,

15 X4 = D, E, F, N, Nal, Q, W, Y,

X5 = F, Hph, Nal, W, Y,

X6 = D, E, F, N, Nal, Q, W, Y,

X7 = A, C, G, Hcy, S, Sec,

X8 = Aad, Asu, D, E, F, Hph, N, Nal, Q, W, Y,

20 X9 = amide, hydrazide, azide, carbamate, marker, spacer, linker, C, GKK, SGKK or deletion

wherein Aad is α -amino adipic acid; α -Abu is α -aminobutyric acid; Ahx is ϵ -aminohexanoic acid; Asu is α -aminosuberic acid; β -A is β -Alanine; Dab is α,γ -diaminobutyric acid; γ -Abu is γ -aminobutyric acid; Har is homoarginine; Hcy is homocysteine; Hph is homophenylalanine; Hyl is δ -hydroxylysine; Hyp is hydroxyproline; Nal is β -(1- or 2-naphthyl)-alanine; Orn is ornithine; Pip is

pipecolic acid; Sec is selenocysteine and the single letter code of amino acids represents the common L-amino acids according IUPAC nomenclature as well as the corresponding D-amino acids.

- 5 In a further embodiment of the invention the peptide according the invention is characterized in that the linker and/or a spacer are selected from the group consisting of α -aminocarboxylic acids as well as homo- and heterooligomers thereof, α,ω -aminocarboxylic acids and branched - or heterooligomers thereof, other aliphatic and/or aromatic amino acids as well as linear and branched
- 10 homo- or heterooligomers; amino-oligoalkoxyalkylamines; maleimidocarboxylic acid derivatives; oligomers of alkylamines; 4-alkylphenyl derivatives; 4-oligoalkoxyphenyl or 4-oligoalkoxyphenoxy derivatives; 4-oligoalkyl-mercaptophenyl or 4-oligoalkylmercaptopheoxy derivatives; 4-oligoalkylaminophenyl or 4-oligoalkylaminophenoxy derivatives; (oligoalkylbenzyl)-
- 15 phenyl or 4-(oligoalkylbenzyl)phenoxy derivatives, as well as 4-(oligoalkoxybenzyl)phenyl or 4-(oligoalkoxy-benzyl)phenoxy derivatives; trityl derivatives; benzyloxyaryl or benzyloxyalkyl derivatives; xanthen-3-yloxyalkyl derivatives; (4-alkylphenyl)- or ω -(4-alkylphenoxy)alkanoic acid derivatives; oligoalkylphenoxyalkyl or oligoalkoxyphenoxyalkyl derivatives; carbamate
- 20 derivatives; amines; trialkylsilyl or dialkyialkoxysilyl derivatives; alkyl or aryl derivatives and/or combinations thereof.

In yet another embodiment the peptide according to the invention is characterized in that it is selected from the group consisting of:

- 25 a) a peptide consisting of the amino acid sequence

X1-X2-X3-X4-X5-X6-X7-X8-X9

wherein X1 to X9 have the same meaning as mentioned hereinabove;

b) a peptide consisting of an amino acid sequence having sufficient homology to be functionally analogous to an amino acid sequence in accordance with a);

5 c) a peptide according to an amino acid sequence a) or b) which is modified by deletions, additions, substitutions, translocations, inversions and/or insertions and functionally analogous to an amino acid sequence in accordance with a) or b);

10 d) a peptide according to amino acid sequence a), b) or c) which is modified by branch or extension with the same or another peptide according to amino acid sequence a), b) or c) to form a homooligomeric or heterooligomeric peptide.

e)

In a still further embodiment the peptide of the invention is characterized in that the amino acid sequence specified under b) has at least 40% homology to 15 any of the amino acid sequences specified under a). Moreover, the peptide of the invention is characterized in that the amino acid sequence specified under b) has at least 60%, preferably 70%, more preferably 80%, especially preferably 90% homology to any of the amino acid sequences specified under a).

20

In yet another embodiment of the invention the peptide of the invention is characterized in that it consists of the amino acid sequence APEDET; WKEPAP; PPDERF; KMWTFG or FGNFWCE.

25 According to the invention the peptide of the invention can be used as a medical active substance. The peptide according to the invention is able to bind to antibodies of patients suffering from Morbus Alzheimer.

The peptide of the invention is furthermore characterized in that it may be immobilized and/or fixed to magnetic, paramagnetic and/or non magnetic nanoparticles.

5 In still another embodiment of the invention, the peptide according to the invention is characterized in that it is bound to a solid phase.

The peptide of the invention may be present in a linear and branched as well as cyclic form. The peptide ring closure is e. g. effected through disulfide bridging when two cysteines are present, or through amide cyclization, which

10 is optionally effected through side chains, through the C to the N termini or through a combination of the latter.

The peptide according to the invention may be further characterized in that the immunoglobulins to which it binds are agAAB which interact with the
15 human β 2-AR and α -AR .

The peptide according to anyone of the foregoing claims, characterized in that it additionally comprises amino groups, amides, hydrazides, azides, carbamates, acetyl groups, biotin groups, markers, spacers and/or linkers.

20

Subject matter of the invention is also an isolated nucleic acid molecule comprising:

25

- a) a nucleic acid molecule comprising a nucleotide sequence which encodes at least one peptide selected from the group consisting of peptides according to claim 1;
- b) a nucleic acid molecule which is complementary to a nucleotide sequence in accordance with a);

- c) a nucleic acid molecule which undergoes hybridization with a nucleotide sequence according to a) or b) under stringent conditions;
- 5 d) a nucleic acid molecule comprising a nucleotide sequence having sufficient homology to be functionally analogous to a nucleotide sequence according to a), b) or c);
- e) a nucleic acid molecule which, as a consequence of the genetic code, is degenerated into a nucleotide sequence according to a) through d); and
- 10 f) a nucleic acid molecule according to a nucleotide sequence of a) through e) which is modified by deletions, additions, substitutions, translocations, inversions and/or insertions and functionally analogous to a nucleotide sequence according to a) through e).

The nucleic acid molecule according to the invention having the nucleotide sequence specified under d) comprises at least 40%, in particular at least 15 60%, preferably 70%, more preferably 80%, especially preferably 90% homology to any of the nucleotide sequences specified under a) through c).

The nucleic acid molecule of the invention is a genomic DNA, a cDNA and/or an RNA.

20 Subject matter of the present invention is also a vector comprising a nucleic acid molecule according to the invention.

Also a host cell is subject matter of the invention comprising the vector according to the invention.

25 Solid phases for affinity chromatography or solid-phase extraction are subject matter of the invention too, comprising organic, inorganic, synthetic polymers

- 20 -

or mixed polymers, preferably cross-linked agarose, cellulose, silica gel, polyamide and polyvinyl alcohols, which are optionally chemically activated, with peptides according to the invention, immobilized on the surface of the solid phase e. g. by covalent bonding or by adsorption. In particular, the 5 peptides can be covalently bound to the solid phase on position X1, X2, X3, X4, X5, X6, X7, X8 and/or X9. It may be advantageous that the peptides are distanced from the support surface by linkers or spacers.

Subject matter of the invention is also a device for removing immunoglobulins 10 from immunoglobulin-containing samples on solid phases, wherein the device contains the solid phase of the invention. Means for the entry of immunoglobulin-containing samples can also be provided.

Another subject matter of the invention is a pharmaceutical composition comprising the nucleic acid molecule according to the invention, the vector according to the invention, the host cell according to the invention, the peptide 15 according to the invention, and/or the solid phase according to the invention, optionally together with a pharmaceutically tolerable carrier substance.

Moreover, a kit comprising a nucleic acid molecule according to the invention, a vector according to the invention, a host cell according to the invention, a peptide according to the invention, a solid phase according to the invention 20 and/or a pharmaceutical composition according to claim 28, optionally together with instructions for combining the contents of the kit and/or providing a formulation.

Subject matter of the invention is also an apparatus for chromatography, comprising peptides according to the invention. In particular, the apparatus is 25 characterized in that the peptides are bound to the solid phase of the invention.

The invention is also concerned with the use of the nucleic acid molecule of the invention, the vector according to the invention, the host cell according to

the invention, the peptide according to the invention, the solid phase according to the invention, the pharmaceutical composition according to the invention, the kit according to the invention, the apparatus according to the invention in the prophylaxis, diagnosis, therapy of AD as well as in the 5 production or screening of a drug for the treatment of neurodegenerative diseases, such as AD .

The invention discloses also a method of treatment of Morbus Alzheimer by binding and/or removal of autoantibodies by means of peptides according to the invention bound to a solid phase.

10

Examples, Materials and Methods

Isolation and cultivation of neonatal heart cells were performed as described previously (Wallukat, G. and Wollenberger, A. 1987. Effects of the serum 15 gammaglobulin fraction of patients with allergic asthma and dilated cardiomyopathy on chronotropic β -adrenoceptor function in cultured rat heart myocytes. Biomed. Biochim. Acta. 78:634-639)

Cell culture. Briefly, single cells were dissociated from the minced ventricles of 20 Wistar rats (1-2 days old) with a 0.25% solution of crude trypsin and were cultured as monolayers with a density of 800 cells/mm² in Halle SM 20-I medium equilibrated with humidified air. The medium contained 10% heat-inactivated FCS and 2 μ mol/1 fluorodeoxyuridine (Serva, Heidelberg, Germany) the latter to prevent proliferation of nonmuscle cells.

25

On the third or fourth days, the cells were incubated for 2 h in 2 ml fresh serum-containing medium. Seven to 10 selected cells or synchronously contracting cell clusters per flask were counted for 15 s. The change of the pulsation rate were expressed as beats per minute. This procedure was 30 repeated twice in different cultures to yield results representing a total of up to 30 cells for each sample.

Immunoglobulin fractions, agonist and antagonist drugs, peptides etc. were added singly or cumulatively as indicated. The basal contraction rate of the spontaneously beating cardiomyocytes was 162 ± 4 min.

5 Preparation of the immunoglobulin fraction. The immunoglobulin fraction was isolated from 1- to 2-ml serum samples by ammonium sulfate precipitation at a saturation of 40%. The precipitates were washed and dissolved in dialysis buffer (154 mmol/l NaCl, 10 mmol/l sodium phosphate; pH 7.2). The procedure of precipitating, washing, and dissolving was repeated twice.

10

Finally, the immunoglobulins were taken up in 1 ml PBS (pH 7.2) and dialyzed at 4°C for 30 hours against 1 l of dialysis buffer. The buffer was changed five times during dialysis. For detection of autoantibodies, the immunoglobulin fractions were added at a dilution of 1:20 or 1:40 to the flasks.

15

After the dialysis, the IgG fractions feature a concentration of 10-15 mg/ml, a maximum of 10 µg/ml of which are specific receptor antibodies. 50 µl of this IgG solution are pre-incubated with the peptides with the following concentration of peptides: 5 µl of a buffered solution of 100 µg/ml per 50 µl

20

IgG solution. This mixture is pre-incubated at room temperature for 30 minutes and subsequently filled into the cell culture vials, after the former nutrient solution has been removed by pipet. The amount of the medium is 2 ml. The final concentration therefore is of 0.5-0.75 mg/ml for the IgG and 0.5 µg for the peptides. If the molecular weight of the peptide (10 mer) is set with

25

1 KD and that of the IgG is set with 150 KD, there is an excess of agAAB .

30

For the neutralization experiments, synthetic peptides corresponding to the sequence of the three extracellular loops of the human β 2 AR or α 1A-AR was added in excess (0.5 µg in 50 µl) to 50 µl of the immunoglobulin fraction. The mixtures were shaken and incubated at room temperature for 1 h. The 100-µl samples were then added to neonatal rat heart muscle cells cultured in 2 ml

medium to a final dilution of 1:40. The beating rate was counted for 15 s, about 60 min after the addition of the peptide/immunoglobulin mixture.

For identifying the autoantibody neutralizing epitopes synthetic overlapping

5 peptides corresponding to the sequence of the first extracellular loop of the human β 2-AR or second extracellular loop of α 1A-AR was added to the immunoglobulins. After 1h incubation at room temperature the samples were used as described above.

10 ED50 calculation. For the estimation of the 50% effective dosage of agAAB to stimulate neonatale rat cardiomyocytes the affinity chromatographic purified antibodies which bind to the epitopes in loop1 of the human β 2 adrenoceptor or loop2 of the human α 1 adrenocepotor (sub type A) the antibodies were given into the tissue culture containing neonatal rat cardiomyocytes
15 sequencially with increasing of antibody concentration until the maximum of the increase of the beating rate was observed. The ED50 was calculated by a computer program.

Patients. Sera of 14 patients with dementia of the Alzheimer type were
20 recruited from the Clinic of Psychiatry of the Hospital Luedenscheid. The controls are immunoglobulin fraction from the blood serum of healthy donors from the Max Delbrück Centre of Moleculare Medicine, Berlin.

For immunoglobulin preparation, the samples were centrifuged at 4,000 g for
25 30 min and stored at -20°C. Reagents was purchased from Sigma-Aldrich Chemie (Deisenhofen, Germany). All other chemicals were of analytical grade.

Results

The figures show the following:

30 Fig.1: Antibodies of patients (n = 11- 14) with Morbus Alzheimer stimulate rat cardiomyocytes via β 2 adrenergic receptor and α 1 adrenergic receptor.

The β 2 adrenergic antagonist ICI 118,551 blocks specifically but partially the stimulating effects of the agonistic antibodies. A more effective inhibition of cellular activity of contraction is observed if additional the α adrenergic antagonist prazosin was given into the cell culture. 11 from 14 serum samples 5 or immunoglobulins activate the cardiac myocytes.

Fig. 2: Antibodies from patients (n=6) with Morbus Alzheimer bind to the first extracellular loop of the β 2 Adrenoceptor. After coincubation of the extracellular loop peptides 1, 2 or 3 with the immunoglobulins of the patients 10 only the loop 1 neutralise the antibody mediated increase of the cellular beating rate.

Fig.3: Antibodies from patients (n = 6) with Morbus Alzheimer bind to a peptid epitope embedded in the first extracellular loop of the β 2-AR. After co- 15 incubation of the immunoglobulins with overlapping epitopes (2 amino acids) of the loop 1 peptides a neutralisation of antibody mediated increase of the beating rate was observed with the epitope FGNFWCE.

Fig.4: Antibodies of patients (n = 6) with Morbus Alzheimer and directed to 20 the β 2 Adrenoceptor belong to the immunoglobulin G subclass 1. After co-incubation of the immunoglobulins with monoclonal antibodies specific for the subclasses 1 - 4 only with the precipitate performed with the monoclonal antibody against IgG1 the antibody mediated activity disappeared.

25 Fig.5: Antibodies from patients (n=6) with Morbus Alzheimer bind to the second extracellular loop of the α 1 Adrenoceptor. After co-incubation of the loops 1, 2 or 3 with the immunoglobulins of the patients only the loop 2 neutralise the antibody mediated increase of the cellular beating rate.

30 Fig.6: Antibodies from patients (n = 5) with Morbus Alzheimer are neutralised by a peptide embedded in the second extracellular loop of the α 1A Adrenoceptor. Overlapping peptides (one or two aminoacids) were co-incubated

- 25 -

with the immunoglobulins. The peptide with the amino acid sequence APEDET neutralises the cellular activity of the immunoglobulins.

Fig.7: Antibodies of patients (n = 5) with Morbus Alzheimer and directed to the $\alpha 1A$ Adrenoceptor belong to the immunoglobulin G subclass 1. After co-incubation of the immunoglobulins with monoclonal antibodies specific for the subclasses 1 - 4 only with the precipitate performed with the monoclonal antibody against IgG1 the antibody mediated activity disappeared.

Claims

1. A method of treating a disease caused by an increase of γ -secretase activity, an increased release of β -amyloid molecules and/or cellular dysfunction of cerebral blood vessel cells, glia cells or neurons, comprising the removal of immunoglobulins from patients suffering from the disease or being suspect to suffer from the disease.
5
2. The method of claim 1 wherein the disease is AD.
3. The method of any of the foregoing claims wherein the removal is performed by immunoabsorption of the immunoglobulin.
- 10 4. A peptide having an ED_{50} of less than 500 nm, in particular 10 nM to an antibody which recognizes an epitope on an extracellular $\alpha 1$ loop 2 or $\beta 2$ loop 1 of a human adrenoceptor (AR) wherein the antibody's binding to the epitope results in an increase of γ -secretase activity, an increased release of β -amyloid molecules and/or cellular dysfunction of cerebral blood vessel cells, glia cells or neurons, wherein the ED_{50} value is measured by a bioassay.
15
5. The peptide of claim 4 wherein the peptide recognizes the $\alpha 1$ extracellular loop 1 epitopes of human α -AR subtypes A, B, or C or combinations thereof, and the extracellular loop 1 of a $\beta 2$ AR, wherein the combinations are A and B; A and C; B and C; A and B and C.
20
6. The peptide of claim 4 or 5 consisting of an amino acid sequence

X1-X2-X3-X4-X5-X6-X7-X8-X9

wherein

a)

25 X1 = amino group, amide, acetyl group, biotin group, marker, spacer, linker, C, GKK, SGKK or deletion,
X2 = A, G, α -Abu, F, Hph, Hyp, Nal, P, Pip, S, V, W, Y, or deletion,
X3 = Hyp, P, Pip, S, T, Y, or deletion,

X4 = Aad, Asu, D, E, Har, K, N, Orn, Q, R,

X5 = Aad, Asu, D, E, N, Q,

X6 = Aad, Asu, D, E, Har, K, N, Orn, Q, R,

X7 = F, Hph, Hyp, Nal, P, Pip, W, Y, or deletion,

5 X8 = A, α -Abu, β -A, G, S, T, V, Y, or deletion,

X9 = amide, hydrazide, azide, carbamate, marker, spacer, linker, C,
GKK, SGKK or deletion, or

b)

10 X1 = amino group, amide, acetyl group, biotin group, marker, spacer,
linker, C, GKK, SGKK or deletion,

X2 = F, Hph, Hyp, Nal, P, Pip, W, Y,

X3 = A, α -Abu, Dab, G, Hyl, K, Orn, R, S, V,

X4 = Aad, Asu, D, E, Hyp, N, P, Pip, Q,

X5 = Aad, Asu, D, E, Hyp, N, P, Pip, Q,

15 X6 = A, α -Abu, G, Hyl, K, Orn, R, S, V

X7 = F, Hph, Hyp, Nal, P, Pip, W, Y,

X8 = γ -Abu, Ahx, β -A, G, or deletion,

X9 = amide, hydrazide, azide, carbamate, marker, spacer, linker, C,
GKK, SGKK or deletion, or

20

c)

X1 = amino group, amide, acetyl group, biotin group, marker, spacer,
linker, C, GKK, SGKK or deletion,

X2 = A, α -Abu, Dab, G, Hyl, K, Orn, R, or deletion,

25 X3 = A, C, F, Hph, M, Nal, S, W, Y,

X4 = F, S, T, W, Y,

X5 = F, S, T, W, Y,

X6 = A, C, F, Hph, M, Nal, S, W, Y,

X7 = A, α -Abu, Dab, G, Hyl, K, Orn, R,

5 X8 = Abu, Ahx, β -A, G, or deletion,

X9 = amide, hydrazide, azide, carbamate, marker, spacer, linker, C, GKK, SGKK or deletion, or

d)

10 X1 = amino group, amide, acetyl group, biotin group, marker, spacer, linker, C, GKK, SGKK or deletion,

X2 = Aad, Asu, D, E, F, Hph, N, Nal, Q, W, Y,

X3 = A, C, G, Hcy, S, Sec,

X4 = D, E, F, N, Nal, Q, W, Y,

15 X5 = F, Hph, Nal, W, Y,

X6 = D, E, F, N, Nal, Q, W, Y,

X7 = A, C, G, Hcy, S, Sec,

X8 = Aad, Asu, D, E, F, Hph, N, Nal, Q, W, Y,

20 X9 = amide, hydrazide, azide, carbamate, marker, spacer, linker, C, GKK, SGKK or deletion

25 wherein Aad is α -amino adipic acid; α -Abu is α -aminobutyric acid; Ahx is ϵ -aminohexanoic acid; Asu is α -aminosuberic acid; β -A is β -Alanine; Dab is α,γ -diaminobutyric acid; γ -Abu is γ -aminobutyric acid; Har is homoarginine; Hcy is homocysteine; Hph is homophenylalanine; Hyl is δ -hydroxylysine; Hyp is hydroxyproline; Nal is β -(1- or 2-naphthyl)-alanine; Orn is ornithine; Pip is pipecolic acid; Sec is selenocysteine and the single letter code of amino acids represents the common L-amino

acids according IUPAC nomenclature as well as the corresponding D-amino acids.

7. The peptide according to anyone of the foregoing claims, characterized in that the linker and/or a spacer are selected from the group

5 comprising: α -aminocarboxylic acids as well as homo- and heterooligomers thereof, α,ω -aminocarboxylic acids and branched - or heterooligomers thereof, other aliphatic and/or aromatic amino acids as well as linear and branched homo- or heterooligomers; amino-

10 oligoalkoxyalkylamines; maleimidocarboxylic acid derivatives; oligomers of alkylamines; 4-alkylphenyl derivatives; 4-

oligoalkoxyphenyl or 4-oligoalkoxyphenoxy derivatives; 4-

15 oligoalkylmercaptophenyl or 4-oligoalkylmercaptophenoxy derivatives; 4-oligo-alkylaminophenyl or 4-oligoalkylaminophenoxy derivatives;

(oligoalkylbenzyl)phenyl or 4-(oligoalkylbenzyl)phenoxy derivatives, as

15 well as 4-(oligo-alkoxybenzyl)phenyl or 4-(oligoalkoxy-benzyl)phenoxy derivatives; trityl derivatives; benzyloxyaryl or benzyloxyalkyl derivatives; xanthen-3-yloxyalkyl derivatives; (4-alkylphenyl)- or ω -(4-

20 alkylphenoxy)alkanoic acid derivatives; oligoalkylphenoxyalkyl or oligoalkoxyphenoxyalkyl derivatives; carbamate derivatives; amines; trialkylsilyl or dialkyalkoxysilyl derivatives; alkyl or aryl derivatives and/or combinations thereof.

8. The peptide according to anyone of the foregoing claims, characterized in that it is selected from the group consisting of:

a) a peptide consisting of the amino acid sequence

25 X1-X2-X3-X4-X5-X6-X7-X8-X9

according to claim 7;

b) a peptide consisting of an amino acid sequence having sufficient homology to be functionally analogous to an amino acid sequence in accordance with a);

5 c) a peptide according to an amino acid sequence a) or b) which is modified by deletions, additions, substitutions, translocations, inversions and/or insertions and functionally analogous to an amino acid sequence in accordance with a) or b);

d) a peptide according to amino acid sequence a), b) or c) which is modified by branch or extension with the same or another peptide according to amino acid sequence a), b) or c) to form a homooligomeric or heterooligomeric peptide.

10 9. The peptide according to anyone of the foregoing claims, characterized in that the amino acid sequence specified under b) has at least 40% homology to any of the amino acid sequences specified under a).

15 10. The peptide according to anyone of the foregoing claims, characterized in that the amino acid sequence specified under b) has at least 60%, preferably 70%, more preferably 80%, especially preferably 90% homology to any of the amino acid sequences specified under a).

11. The peptide according to anyone of the foregoing claims, characterized in that it consists of the amino acid sequence APEDET; WKEPAP; PPDERF; KMWTGF or FGNFWCE.

20 12. The peptide according to anyone of the foregoing claims for use as a medical active substance.

13. The peptide according to anyone of the foregoing claims, characterized in that the peptide binds to antibodies of patients suffering from AD.

25 14. The peptide according to anyone of the foregoing claims, characterized in that the peptide is immobilized and/or fixed to magnetic, paramagnetic and/or non magnetic nanoparticles.

15. The peptide according to anyone of the foregoing claims, characterized in that the peptide is bound to a solid phase.

16. The peptide according to anyone of the foregoing claims in a linear and branched as well as cyclic form, wherein the peptide ring closure is effected through disulfide bridging when two cysteines are present, or through amide cyclization, which is optionally effected through side chains, through the C to the N termini or through a combination of these last two possibilities.

5

17. The peptide according to anyone of the foregoing claims, characterized in that the immunoglobulins are agonistic autoantibodies which interact with the α -adrenoreceptor receptor.

10 18. The peptide according to anyone of the foregoing claims, characterized in that it additionally comprises amino groups, amides, hydrazides, azides, carbamates, acetyl groups, biotin groups, markers, spacers and/or linkers.

19. An isolated nucleic acid molecule selected from the group comprising:

15 a) a nucleic acid molecule comprising a nucleotide sequence which encodes at least one peptide selected from the group consisting of peptides according to claim 4;

b) a nucleic acid molecule which is complementary to a nucleotide sequence in accordance with a);

20 c) a nucleic acid molecule which undergoes hybridization with a nucleotide sequence according to a) or b) under stringent conditions;

d) a nucleic acid molecule comprising a nucleotide sequence having sufficient homology to be functionally analogous to a nucleotide sequence according to a), b) or c);

25 e) a nucleic acid molecule which, as a consequence of the genetic code, is degenerated into a nucleotide sequence according to a) through d); and

f) a nucleic acid molecule according to a nucleotide sequence of a) through e) which is modified by deletions, additions, substitutions, translocations, inversions and/or insertions and functionally analogous to a nucleotide sequence according to a) through e).

5 20. The nucleic acid molecule according to claim 19, characterized in that the nucleotide sequence specified under d) has at least 40% homology to any of the nucleotide sequences specified under a) through c).

10 21. The nucleic acid molecule according to claim 19, characterized in that the nucleotide sequence specified under d) has at least 60%, preferably 70%, more preferably 80%, especially preferably 90% homology to any of the nucleotide sequences specified under a) through c).

22. The nucleic acid molecule according to any of claims 19 to 21, characterized in that it is a genomic DNA, a cDNA and/or an RNA.

15 23. A vector comprising a nucleic acid molecule according to any of claims 19 to 22.

24. A host cell comprising the vector according to claim 23.

25. A peptide, said peptide being encoded by a nucleic acid molecule according to any of claims 19 to 22.

20 26. Solid phases for affinity chromatography or solid-phase extraction consisting of organic, inorganic, synthetic polymers or of mixed polymers, preferably cross-linked agarose, cellulose, silica gel, polyamide and polyvinyl alcohols, which are optionally chemically activated, with peptides according to at least one of claims 4 to 18, 25, immobilized on the surface of the solid phase.

27. The solid phases according to claim 26, wherein the peptides are bound to the solid support phase covalently or by adsorption.
28. The solid phases according to claim 26 and/or 27, wherein the peptides are covalently bound to the solid phase on position X1, X2, X3, X4, X5, 5 X6, X7, X8 and/or X9.
29. The solid phases according to any of claims 26 to 28, wherein the peptides are distanced from the support surface by linkers/spacers.
30. A device for removing immunoglobulins from immunoglobulin-containing samples on solid phases, wherein the device contains a solid phase 10 according to any of claims 26 to 29, and means for the entry of immunoglobulin-containing samples are provided.
31. A pharmaceutical composition comprising a nucleic acid molecule according to any of claims 19 to 22, a vector according to claim 23, a host cell according to claim 24, a peptide according to any of claims 4 to 15 17, 25, and/or a solid phase according to claims 27 to 29, optionally together with a pharmaceutically tolerable carrier.
32. A kit comprising a nucleic acid molecule according to any of claims 19 to 22, a vector according to claim 23, a host cell according to claim 24, a peptide according to any of claims 4 to 18, 25, a solid phase according to 20 claims 26 to 29 and/or a pharmaceutical composition according to claim 31, optionally together with instructions for combining the contents of the kit and/or providing a formulation.
33. An apparatus for chromatography, comprising peptides according to any of claims 4 to 18, 25.
- 25 34. The apparatus according to claim 33, characterized in that the peptides are bound to a solid phase according to claims 25 to 29.

35. Use of a nucleic acid molecule according to any of claims 19 to 22, a vector according to claim 23, a host cell according to claim 24, a peptide according to any of claims 4 to 18, 25, a solid phase according to claims 26 to 29, a pharmaceutical composition according to claim 31, a kit according to claim 32, an apparatus according to claim 33 or 34 in the prophylaxis, diagnosis, therapy, follow-up and/or aftercare of Morbus Alzheimer.
5
36. Use of a nucleic acid molecule according to any of claims 19 to 22, a vector according to claim 23, a host cell according to claim 24, a peptide according to any of claims 4 to 18, 25, a solid phase according to claims 26 to 29, a pharmaceutical composition according to claim 31, a kit according to claim 32, an apparatus according to claim 33 or 34 in the production or screening of a drug for the treatment of neurodegenerative diseases, such as AD.
10
37. Use of an immunoadsorption device for manufacturing a device for the treatment of a disease caused by an increase of γ -secretase activity, an increased release of β -amyloid molecules and/or cellular dysfunction of cerebral blood vessel cells, glia cells or neurons by removal of immunoglobulins from patients.
15
38. A method for the treatment of Morbus Alzheimer by binding and/or removal of autoantibodies by means of peptides according to any of claims 4 to 18, 25 bound to a solid phase.
20

- 1/1 -




Fig. 1




Fig. 2




Fig. 3




Fig. 4

- 2/2 -




Fig. 5




Fig. 6




Fig. 7