

REPUBLIC OF SOUTH AFRICA  
PATENTS ACT, 1978

PUBLICATION PARTICULARS AND ABSTRACT  
(Section 32(3)(a) – Regulations 22(i)(g) and 31)

OFFICIAL APPLICATION NO.		LODGING DATE	ACCEPTANCE DATE
21	01 <b>2006/03101</b>	23 19 April 2006	43 09-07-2007
INTERNATIONAL CLASSIFICATION		NOT FOR PUBLICATION	
51	C07D	CLASSIFIED BY :	
FULL NAME(S) OF APPLICANT(S)			
71	BAYER CROPSCIENCE AG		

FULL NAME(S) OF INVENTOR(S)	
72	DUNKEL, Ralf; ELBE, Hans-Ludwig; GREUL, Jörg, Nico; HARTMANN, Benoit; WACHENDORFF-NEUMANN, Ulrike; DAHMEN, Peter; KUCK, Karl-Heinz

EARLIEST PRIORITY CLAIMED	COUNTRY	NUMBER	DATE
NOTE: the country must be indicated by its International Abbreviation – see Schedule 4 of the Regulations	33 DE	31 10349497.9	32 23 October 2003

TITLE OF INVENTION	
54	N-SUBSTITUTED PYRAZOLYL CARBOXANILIDES

57	ABSTRACT (NOT MORE THAN 150 WORDS)	NUMBER OF PAGES	60
----	------------------------------------	-----------------	----

FOR ABSTRACT SEE THE NEXT SHEET

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum  
Internationales Büro



(43) Internationales Veröffentlichungsdatum  
12. Mai 2005 (12.05.2005)

PCT

(10) Internationale Veröffentlichungsnummer  
WO 2005/042480 A2

(51) Internationale Patentklassifikation<sup>7</sup>: C07D  
(21) Internationales Aktenzeichen: PCT/EP2004/011396  
(22) Internationales Anmeldedatum:  
12. Oktober 2004 (12.10.2004)  
(25) Einreichungssprache: Deutsch  
(26) Veröffentlichungssprache: Deutsch  
(30) Angaben zur Priorität:  
10349497.9 23. Oktober 2003 (23.10.2003) DE

(74) Gemeinsamer Vertreter: BAYER CROPSCIENCE  
AKTIENGESELLSCHAFT, Law and Patents, Patents  
and Licensing, 51368 Leverkusen (DE).

(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme  
von US): BAYER CROPSCIENCE AKTIENGE-  
SELLSCHAFT [DE/DE]; Alfred-Nobel-Str. 50, 40789  
Monheim (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): DUNKEL, Ralf  
[DE/DE]; Krischer Str. 22, 40789 Monheim (DE).  
ELBE, Hans-Ludwig [DE/DE]; Dasnöckel 59, 42329  
Wuppertal (DE). GREUL, Jörg, Nico [DE/DE]; Am  
Sandberg 30 a, 42799 Lanchlingen (DE). HARTMANN,  
Benoit [FR/DE]; Fährstr. 1, 40764 Langenfeld (DE).  
WACHENDORFF NEUMANN, Ulrike [DE/DE];  
Oberer Markengr. 85, 56566 Neuwied (DE). DAHMEN,  
Peter [DE/DE]; Altebrückerstr. 61, 41470 Neuss (DE).  
KUCK, Karl-Heinz [DE/DE]; Pastor-Löh-Str. 30a, 40764  
Langenfeld (DE).

(81) Bestimmungsstaaten (soweit nicht anders angegeben, für  
jede verfügbare nationale Schutzrechtsart): AE, AG, AL,  
AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,  
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES,  
FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.

(84) Bestimmungsstaaten (soweit nicht anders angegeben, für  
jede verfügbare regionale Schutzrechtsart): ARIPO (BW,  
GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,  
ZM, ZW), eurasisches (AM, AZ, BY, KG, KZ, MD, RU,  
TJ, TM), europäisches (AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT,  
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Veröffentlicht:

— ohne internationalen Recherchenbericht und erneut zu ver-  
öffentlichen nach Erhalt des Berichts

Zur Erklärung der Zweibuchstaben-Codes und der anderen Ab-  
kürzungen wird auf die Erklärungen ("Guidance Notes on Co-  
des and Abbreviations") am Anfang jeder regulären Ausgabe der  
PCT-Gazette verwiesen.

(54) Title: N-SUBSTITUTED PYRAZOLYL CARBOXANILIDES

(54) Bezeichnung: N-SUBSTITUIERTE PYRAZOLYL CARBOXANILIDE



(I)

(57) Abstract: The invention relates to novel N-substituted pyra-  
zoly carboxanilides of formula (I), in which R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup>  
are defined as cited in the description, to several methods for pro-  
ducing said substances, to their use for controlling undesirable mi-  
cro-organisms and to novel intermediate products and the produc-  
tion thereof.

(57) Zusammenfassung: Neue N-substituierte Pyrazolylcar-  
boxanilide der Formel (I), in welcher R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> und R<sup>4</sup> die in  
der Beschreibung angegebenen Bedeutungen haben, mehrere  
Verfahren zum Herstellen dieser Stoffe und deren Verwendung  
zum Bekämpfen von unerwünschten Mikroorganismen, sowie

WO 2005/042480 A2

neue Zwischenprodukte und deren Herstellung

RCS 033022

**N-substituted pyrazolyl carboxanilides**


The present invention relates to novel (N-substituted) pyrazolylcarboxanilides, to a plurality of processes for their preparation and to their use for controlling unwanted microorganisms.

5

It is already known that numerous carboxanilides have fungicidal properties (cf. WO 93/11117, EP-A 0 545 099, EP-A 0 824 099, JP 63-48269, WO 02/059086 and JP 8-176112). N-[2-(1,3-Dimethylbutyl)phenyl]-5-fluoro-1-methyl-N-prop-2-yn-1-yl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide and N-acetyl-N-[2-(1,3-dimethylbutyl)phenyl]-5-fluoro-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide (WO 02/059086) may be mentioned by way of example. The activity of these compounds is good; however, at low application rates it is sometimes unsatisfactory.

10

This invention now provides novel N-substituted pyrazolylcarboxanilides of the formula (I)



15 in which

R<sup>1</sup> represents methyl, trifluoromethyl or difluoromethyl,

R<sup>2</sup> represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl,

a) R<sup>3</sup> represents hydrogen,

20 R<sup>4</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms;

25

(C<sub>3</sub>-C<sub>8</sub>-cycloalkyl)carbonyl; (C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl)carbonyl having 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

b) R<sup>3</sup> represents halogen, C<sub>1</sub>-C<sub>8</sub>-alkyl or C<sub>1</sub>-C<sub>8</sub>-haloalkyl,

30 R<sup>4</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-

C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms;

(C<sub>1</sub>-C<sub>8</sub>-alkyl)carbonyl, (C<sub>1</sub>-C<sub>8</sub>-alkoxy)carbonyl, (C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>8</sub>-cycloalkyl)carbonyl; (C<sub>1</sub>-C<sub>6</sub>-haloalkyl)carbonyl, (C<sub>1</sub>-C<sub>6</sub>-haloalkoxy)carbonyl, (halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl)carbonyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

R<sup>5</sup> represents hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,

R<sup>6</sup> and R<sup>7</sup> independently of one another, each represent hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>8</sub>-haloalkyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,

R<sup>6</sup> and R<sup>7</sup> furthermore together with the nitrogen atom to which they are attached form a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen and C<sub>1</sub>-C<sub>4</sub>-alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and NR<sup>10</sup>,

R<sup>8</sup> and R<sup>9</sup> independently of one another, represent hydrogen, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>8</sub>-haloalkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,


R<sup>8</sup> and R<sup>9</sup> furthermore together with the nitrogen atom to which they are attached form a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen and C<sub>1</sub>-C<sub>4</sub>-alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and NR<sup>10</sup>,

R<sup>10</sup> represents hydrogen or C<sub>1</sub>-C<sub>6</sub>-alkyl.

If appropriate, the compounds according to the invention can be present as mixtures of various possible isomeric forms, in particular of stereoisomers, such as, for example, E and Z, threo and erythro, and also optical isomers, and, if appropriate, also of tautomers. What is claimed are both the E and Z isomers, and the threo and erythro and also the optical isomers, any mixtures of these isomers and the possible tautomeric forms.

Furthermore, it has been found that N-substituted pyrazolylcarboxanilides of the formula (I) are obtained when

a) carboxylic acid derivatives of the formula (II)




5 in which

R<sup>1</sup> is as defined above and

X<sup>1</sup> represents halogen or hydroxyl

are reacted with an aniline derivative of the formula (III)




10 in which R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as defined above,

if appropriate in the presence of a catalyst, if appropriate in the presence of a condensing agent, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

or

15 b) pyrazolylcarboxanilides of the formula (IV)




in which R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as defined above

are hydrogenated, if appropriate in the presence of a diluent and if appropriate in the presence of a catalyst,

20 or

c) pyrazolylcarboxanilides of the formula (Ia)



in which  $R^1$ ,  $R^2$  and  $R^3$  are as defined above  
are reacted with halides of the formula (V)



in which

$R^4$  is as defined above and

$X^2$  represents chlorine, bromine or iodine

in the presence of a base and in the presence of a diluent.

10

Finally, it has been found that the novel N-substituted pyrazolylcarboxanilides of the formula (I) have very good microbicidal properties and can be used for controlling unwanted microorganisms both in crop protection and in the protection of materials.

15 The formula (I) provides a general definition of the N-substituted pyrazolylcarboxanilides according to the invention. Preferred radical definitions of the formulae mentioned above and below are given below. These definitions apply to the end products of the formula (I) and likewise to all intermediates.

20  $R^1$  preferably represents methyl.

$R^1$  furthermore preferably represents trifluoromethyl.

$R^1$  furthermore preferably represents difluoromethyl.

$R^2$  preferably represents hydrogen.

25  $R^2$  furthermore preferably represents fluorine, where fluorine is particularly preferably located in the 4-, 5- or 6-position, very particularly preferably in the 4- or 6-position, especially in the 4-position, of the anilide radical [cf. formula (I) above].

$R^2$  furthermore preferably represents chlorine, where chlorine is particularly preferably located in the 5-position of the anilide radical [cf. formula (I) above].

30  $R^2$  furthermore preferably represents methyl, where methyl is particularly preferably located in the 3-position of the anilide radical (cf. formula (I) above).

$R^2$  furthermore preferably represents trifluoromethyl, where trifluoromethyl is particularly preferably located in the 4- or 5-position of the anilide radical [cf. formula (I) above].

If R<sup>3</sup> represents hydrogen,

R<sup>4</sup> preferably represents C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms;

10 (C<sub>3</sub>-C<sub>6</sub>-cycloalkyl)carbonyl; (C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl)carbonyl having 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>.

If R<sup>3</sup> represents hydrogen,

R<sup>4</sup> particularly preferably represents methyl, ethyl, n- or isopropyl, n-, iso-, sec- or tert-butyl, pentyl or hexyl, methylsulphinyl, ethylsulphinyl, n- or isopropylsulphinyl, n-, iso-, sec- or tert-butylsulphinyl, methylsulphonyl, ethylsulphonyl, n- or isopropylsulphonyl, n-, iso-, sec- or tert-butylsulphonyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl, trifluoromethyl, trichloromethyl, trifluoroethyl, difluoromethylthio, difluorochloromethylthio, trifluoromethylthio, trifluoromethylsulphinyl, trifluoromethylsulphonyl, trifluoromethoxymethyl; formyl, -CH<sub>2</sub>-CHO, -(CH<sub>2</sub>)<sub>2</sub>-CHO, -CH<sub>2</sub>-CO-CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-CO-CF<sub>3</sub>, -CH<sub>2</sub>-CO-CCl<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CCl<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH<sub>2</sub>CF<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CF<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CCl<sub>2</sub>CCl<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CF<sub>2</sub>CF<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CCl<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CCl<sub>2</sub>CCl<sub>3</sub>;

20 cyclopropylcarbonyl, cyclopentylcarbonyl, cyclohexylcarbonyl or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>.

If R<sup>3</sup> represents hydrogen,

30 R<sup>4</sup> very particularly preferably represents methyl, methoxymethyl, formyl, -CH<sub>2</sub>-CHO, -(CH<sub>2</sub>)<sub>2</sub>-CHO, -CH<sub>2</sub>-CO-CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub>, -C(=O)CHO, -C(=O)C(=O)CH<sub>3</sub>, -C(=O)C(=O)CH<sub>2</sub>OCH<sub>3</sub>, -C(=O)CO<sub>2</sub>CH<sub>3</sub>, -C(=O)CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>.

If

35 R<sup>3</sup> preferably represents fluorine, chlorine, bromine, iodine, C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>1</sub>-C<sub>6</sub>-haloalkyl having 1 to 13 fluorine, chlorine and/or bromine atoms,

particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, n-, isopropyl, n-, iso-, sec-, tert-butyl or C<sub>1</sub>-C<sub>4</sub>-haloalkyl having 1 to 9 fluorine, chlorine and/or bromine atoms, very particularly preferably represents methyl, ethyl, fluorine, chlorine, trifluoromethyl,

then

- 5 R<sup>4</sup> preferably represents C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halo-cycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; 10 halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms; (C<sub>1</sub>-C<sub>6</sub>-alkyl)carbonyl, (C<sub>1</sub>-C<sub>6</sub>-alkoxy)carbonyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>6</sub>-cycloalkyl)carbonyl; (C<sub>1</sub>-C<sub>4</sub>-haloalkyl)carbonyl, (C<sub>1</sub>-C<sub>4</sub>-haloalkoxy)carbonyl, (halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl)carbonyl having in each case 1 to 9 15 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, particularly preferably represents methyl, ethyl, n- or isopropyl, n-, iso, sec- or tert-butyl, pentyl or hexyl, methylsulphinyl, ethylsulphinyl, n- or isopropylsulphinyl, n-, iso-, sec- or tert-butylsulphinyl, methylsulphonyl, ethylsulphonyl, n- or isopropylsulphonyl, n-, iso-, sec- or tert-butylsulphonyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, 20 cyclopropyl, cyclopentyl, cyclohexyl, trifluoromethyl, trichloromethyl, trifluoroethyl, difluoromethylthio, difluorochloromethylthio, trifluoromethylthio, trifluoromethylsulphinyl, trifluoromethylsulphonyl, trifluoromethoxymethyl; formyl, -CH<sub>2</sub>-CHO, -CH<sub>2</sub>CH<sub>2</sub>-CHO, -CH<sub>2</sub>-CO-CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, 25 -CH<sub>2</sub>-CO<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>-CO-CF<sub>3</sub>, -CH<sub>2</sub>-CO-CCl<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CCl<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH<sub>2</sub>CF<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO-CH<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CF<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CCl<sub>3</sub>, -CH<sub>2</sub>-CO<sub>2</sub>CCl<sub>2</sub>CCl<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CF<sub>2</sub>CF<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CH<sub>2</sub>CCl<sub>3</sub>, -(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>CCl<sub>2</sub>CCl<sub>3</sub>; 30 methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, isopropylcarbonyl, tert-butylcarbonyl, methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl, cyclopropylcarbonyl; trifluoromethylcarbonyl, trifluoromethoxycarbonyl, or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, very particularly preferably represents methyl, methoxymethyl, formyl, -CH<sub>2</sub>-CHO, -(CH<sub>2</sub>)<sub>2</sub>-CHO, -CH<sub>2</sub>-CO-CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>-CO-CH(CH<sub>3</sub>)<sub>2</sub>, -C(=O)CHO, 35 -C(=O)C(=O)CH<sub>3</sub>, -C(=O)C(=O)CH<sub>2</sub>OCH<sub>3</sub>, -C(=O)CO<sub>2</sub>CH<sub>3</sub>, -C(=O)CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>.

- R<sup>5</sup> preferably represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms.
- R<sup>5</sup> particularly preferably represents hydrogen, methyl, ethyl, n- or isopropyl, tert-butyl, methoxy, ethoxy, n- or isopropoxy, tert-butoxy, cyclopropyl; trifluoromethyl, trifluoromethoxy.
- R<sup>6</sup> and R<sup>7</sup> independently of one another, preferably represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms.
- R<sup>6</sup> and R<sup>7</sup> furthermore together with the nitrogen atom to which they are attached preferably form a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of halogen and C<sub>1</sub>-C<sub>4</sub>-alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and NR<sup>10</sup>.
- R<sup>6</sup> and R<sup>7</sup> independently of one another, particularly preferably represent hydrogen, methyl, ethyl, n- or isopropyl, n-, iso-, sec- or tert-butyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl; trifluoromethyl, trichloromethyl, trifluoroethyl, trifluoromethoxymethyl.
- R<sup>6</sup> and R<sup>7</sup> furthermore together with the nitrogen atom to which they are attached particularly preferably form a saturated heterocycle from the group consisting of morpholine, thiomorpholine and piperazine which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of fluorine, chlorine, bromine and methyl, where the piperazine may be substituted by R<sup>10</sup> on the second nitrogen atom.
- R<sup>8</sup> and R<sup>9</sup> independently of one another, preferably represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms.
- R<sup>8</sup> and R<sup>9</sup> furthermore together with the nitrogen atom to which they are attached preferably form a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of halogen and C<sub>1</sub>-C<sub>4</sub>-alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and NR<sup>10</sup>.
- R<sup>8</sup> and R<sup>9</sup> independently of one another, particularly preferably represent hydrogen, methyl, ethyl, n- or isopropyl, n-, iso-, sec- or tert-butyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, cyclopropyl, cyclopentyl, cyclohexyl; trifluoromethyl, trichloromethyl, trifluoroethyl, trifluoromethoxymethyl.

R<sup>8</sup> and R<sup>9</sup> furthermore together with the nitrogen atom to which they are attached particularly preferably form a saturated heterocycle from the group consisting of morpholine, thiomorpholine and piperazine which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of fluorine, chlorine, bromine and methyl, where the piperazine may be substituted by R<sup>10</sup> on the second nitrogen atom.

R<sup>10</sup> preferably represents hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl.


R<sup>10</sup> particularly preferably represents hydrogen, methyl, ethyl, n- or isopropyl, n-, iso-, sec- or tert-butyl.

10

Emphasis is given to compounds of the formula (I) in which R<sup>4</sup> represents formyl.

Emphasis is furthermore given to compounds of the formula (I) in which R<sup>4</sup> represents -C(=O)C(=O)R<sup>5</sup>, where R<sup>5</sup> is as defined above.

15 Emphasis is furthermore given to N-substituted pyrazolylcarboxanilides of the formula (Ib)




in which

R<sup>4A</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms; (C<sub>3</sub>-C<sub>8</sub>-cycloalkyl)carbonyl; (C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl)carbonyl having 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, and R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined above.

25

Emphasis is furthermore given to N-substituted pyrazolylcarboxanilides of the formula (Ic)



in which

$R^{3B}$  represents halogen,  $C_1$ - $C_8$ -alkyl or  $C_1$ - $C_8$ -haloalkyl,

- $R^{4B}$  represents  $C_1$ - $C_8$ -alkyl,  $C_1$ - $C_6$ -alkylsulphinyl,  $C_1$ - $C_6$ -alkylsulphonyl,  $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -cycloalkyl;  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_4$ -haloalkylthio,  $C_1$ - $C_4$ -haloalkylsulphinyl,  $C_1$ - $C_4$ -haloalkylsulphonyl, halo- $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_8$ -halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -alkyl)carbonyl- $C_1$ - $C_3$ -alkyl, ( $C_1$ - $C_3$ -alkoxy)carbonyl- $C_1$ - $C_3$ -alkyl; halo-( $C_1$ - $C_3$ -alkyl)carbonyl- $C_1$ - $C_3$ -alkyl, halo-( $C_1$ - $C_3$ -alkoxy)carbonyl- $C_1$ - $C_3$ -alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms; ( $C_1$ - $C_8$ -alkyl)carbonyl, ( $C_1$ - $C_8$ -alkoxy)carbonyl, ( $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl)carbonyl, ( $C_3$ - $C_8$ -cycloalkyl)carbonyl; ( $C_1$ - $C_6$ -haloalkyl)carbonyl, ( $C_1$ - $C_6$ -haloalkoxy)carbonyl, (halo- $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkyl)carbonyl, ( $C_3$ - $C_8$ -halocycloalkyl)carbonyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; or  $-C(=O)C(=O)R^5$ ,  $-CONR^6R^7$  or  $-CH_2NR^8R^9$ , and  $R^1$ ,  $R^2$ ,  $R^5$ ,  $R^6$ ,  $R^7$ ,  $R^8$  and  $R^9$  are as defined above.

Saturated or unsaturated hydrocarbon radicals, such as alkyl or alkenyl, can in each case be straight-chain or branched as far as this is possible, including in combination with heteroatoms, such as, for example, in alkoxy. Optionally substituted radicals can be mono- or polysubstituted, where in the case of polysubstitution the substituents can be identical or different.


Halogen-substituted radicals, such as, for example, haloalkyl, are mono- or polyhalogenated. In the case of polyhalogenation, the halogen atoms can be identical or different. Here, halogen denotes fluorine, chlorine, bromine and iodine, in particular fluorine, chlorine and bromine.

However, the general or preferred radical definitions or illustrations given above can also be combined with one another as desired, i.e. including combinations between the respective ranges and preferred ranges. The definitions apply both to the end products and, correspondingly, to the precursors and intermediates.

The definitions mentioned can be combined with one another as desired. Moreover, individual definitions may not apply.

Preference, particular preference or very particular preference is given to compounds of the formula (I) which carry the substituents mentioned as preferred, particularly preferred and very particularly preferred, respectively.

- 5 Using 5-fluoro-1,3-dimethyl-1H-pyrazole-4-carbonyl chloride and N-(methoxymethyl)-2-(1,3,3-trimethylpentyl)aniline as starting materials, the process (a) according to the invention can be illustrated by the following formula scheme:




- 10 The formula (II) provides a general definition of the carboxylic acid derivatives required as starting materials for carrying out the process (a) of the invention. In this formula (II), R<sup>1</sup> preferably has those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred for R<sup>1</sup>. X<sup>1</sup> preferably represents chlorine, bromine or hydroxyl.

- 15 The carboxylic acid derivatives of the formula (II) are known and/or can be prepared by known processes (cf. WO 93/11117, EP-A 0 545 099, EP-A 0 589 301 and EP-A 0 589 313).

- 20 The formula (III) provides a general definition of the anilines furthermore required as starting materials for carrying out the process (a) according to the invention. In this formula (III), R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> preferably, particularly preferably and very particularly preferably have those meanings which have already been mentioned in connection with the description of the compounds of formula (I) according to the invention as being preferred, particularly preferred and very particularly preferred, respectively, for these radicals.

- 25 The aniline derivatives of the formula (III) are novel. They can be prepared by  
d) reacting alkylanilines of the formula (VI)



in which R<sup>2</sup> and R<sup>3</sup> are as defined above  
with halides of the formula (V)



in which

$R^4$  is as defined above and

$X^2$  represents chlorine, bromine or iodine,

5 if appropriate in the presence of a base and if appropriate in the presence of a diluent.

The formula (VI) provides a general definition of the alkyanilines required as starting materials for carrying out the process (d) according to the invention. In this formula (VI),  $R^2$  and  $R^3$  preferably, particularly preferably and very particularly preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred and very particularly preferred, respectively, for these radicals.


The alkyanilines of the formula (VI) are known and/or can be obtained by known methods (cf. WO 15 03/010149).

The formula (V) provides a general definition of the halides furthermore required as starting materials for carrying out the process (d) according to the invention. In this formula (V),  $R^4$  preferably, particularly preferably and very particularly preferably has those meanings which have already been given in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred and very particularly preferred, respectively, for  $R^4$ .

Halides of the formula (V) are known.

25

Using N-acetyl-5-fluoro-1,3-dimethyl-N-{2-[(1Z)-1,3,3-trimethylbut-1-en-1-yl]phenyl}-1H-pyrazole-4-carboxamide and hydrogen as starting materials and a catalyst, the course of the process (b) according to the invention can be illustrated by the following formula scheme:




30

The formula (IV) provides a general definition of the pyrazolylcarboxanilides required as starting materials for carrying out the process (b) according to the invention. In this formula (IV),  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  preferably, particularly preferably and very particularly preferably have those meanings which have already been given in connection with the description of the compounds of the formula (I)

according to the invention as being preferred, particularly preferred and very particularly preferred, respectively, for these radicals.

The compounds of the formula (IV) are novel. They can be prepared by

- 5 e) reacting pyrazolylcarboxanilides of the formula (VII)



in which  $R^1$ ,  $R^2$  and  $R^3$  are as defined above  
with halides of the formula (V)



- 10 in which

$R^4$  is as defined above and

$X^2$  represents chlorine, bromine or iodine,


if appropriate in the presence of a base and if appropriate in the presence of a diluent.

- 15 The formula (VII) provides a general definition of the pyrazolylcarboxanilides required as starting materials for carrying out the process (e) according to the invention. In this formula (VII),  $R^1$ ,  $R^2$  and  $R^3$  preferably, particularly preferably and very particularly preferably have those meanings which have already been given in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred and very particularly preferred,  
20 respectively, for these radicals.

The pyrazolylcarboxanilides of the formula (VII) are known (cf. WO 03/010149).

- The halides of the formula (V) furthermore required as starting materials for carrying out the process  
25 (e) according to the invention have already been described in connection with the process (d) according to the invention.

- Using 5-fluoro-1,3-dimethyl-N-[2-(1,3,3-trimethylbutyl)phenyl]-1H-pyrazole-4-carboxamide and ethyl chloro(oxo)acetate as starting materials, the course of the process (c) according to the invention  
30 can be illustrated by the following formula scheme:



Formula (Ia) provides a general definition of the pyrazolylcarboxanilides required as starting materials for carrying out the process (c) according to the invention. In this formula (Ia), R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> preferably, particularly preferably and very particularly preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred and very particularly preferred, respectively, for these radicals.

The pyrazolylcarboxanilides of the formula (Ia) are known (cf. WO 03/010149).

10

The halides of the formula (V) furthermore required as starting materials for carrying out the process (c) according to the invention have already been described in connection with the process (d) according to the invention.

15 Suitable diluents for carrying out the process (a) according to the invention are all inert organic solvents. These preferably include aliphatic, alicyclic or aromatic hydrocarbons, such as, for example, petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin; halogenated hydrocarbons, such as, for example, chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane; ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl tert-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole, or amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide.

25 The process (a) according to the invention is, if appropriate, carried out in the presence of a suitable acid acceptor. Suitable acid acceptors are all customary inorganic or organic bases. These preferably include alkaline earth metal or alkali metal hydrides, hydroxides, amides, alkoxides, acetates, carbonates or bicarbonates, such as, for example, sodium hydride, sodium amide, sodium methoxide, sodium ethoxide, potassium tert-butoxide, sodium hydroxide, potassium hydroxide, ammonium hydroxide, sodium acetate, potassium acetate, calcium acetate, ammonium acetate, sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate or ammonium carbonate, and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dimethylbenzylamine, pyridine, N-methylpiperidine, N-methylmorpholine, N,N-

30

dimethylaminopyridine, diazabicyclooctane (DABCO), diazabicyclononene (DBN) or diazabicycloundecene (DBU).

5 When carrying out the process (a) according to the invention, the reaction temperatures can be varied within a relatively wide range. In general, the process is carried out at temperatures of from 0°C to 150°C, preferably at temperatures of from 0°C to 80°C.

10 For carrying out the process (a) according to the invention for preparing the compounds of the formula (I), in general from 0.2 to 5 mol, preferably from 0.5 to 2 mol, of aniline derivatives of the formula (III) are employed per mole of the carboxylic acid derivative of the formula (II).

15 Suitable diluents for carrying out the process (b) according to the invention are all inert organic solvents. These preferably include aliphatic, alicyclic or aromatic hydrocarbons, such as, for example, petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin; ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl t-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole, or alcohols, such as methanol, ethanol, n- or i-propanol, n-, i-, sec- or tert-butanol, ethanediol, propane-1,2-diol, ethoxyethanol, methoxyethanol, diethylene glycol monomethyl ether or diethylene glycol monoethyl ether.

20

The process (b) according to the invention is, if appropriate, carried out in the presence of a catalyst. Suitable catalysts are all catalysts which are customarily used for hydrogenations. Examples which may be mentioned are: Raney nickel, palladium or platinum, if appropriate on a support, such as, for example, activated carbon.

25

Instead of in the presence of hydrogen in combination with a catalyst, the hydrogenation in the process (b) according to the invention can also be carried out in the presence of triethylsilane.

30 When carrying out the process (b) according to the invention, the reaction temperatures can be varied within a relatively wide range. In general, the process is carried out at temperatures of from 0°C to 150°C, preferably at temperatures of from 0°C to 80°C.

35 Suitable diluents for carrying out the process (c) according to the invention are all inert organic solvents. These preferably include aliphatic, alicyclic or aromatic hydrocarbons, such as, for example, petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin; halogenated hydrocarbons, such as, for example, chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane; ethers, such as

diethyl ether, diisopropyl ether, methyl tert-butyl ether, methyl tert-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole, or amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide.

5

The process (c) according to the invention is carried out in the presence of a base. Suitable bases are all customary inorganic or organic bases. These preferably include alkaline earth metal or alkali metal hydrides, hydroxides, amides, alkoxides, acetates, carbonates or bicarbonates, such as, for example, sodium hydride, sodium amide, sodium methoxide, sodium ethoxide, potassium tert-butoxide, sodium hydroxide, potassium hydroxide, ammonium hydroxide, sodium acetate, potassium acetate, calcium acetate, ammonium acetate, sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate or cesium carbonate, and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dimethylbenzylamine, pyridine, N-methylpiperidine, N-methylmorpholine, N,N-dimethylaminopyridine, diazabicyclooctane (DABCO), diazabicyclononene (DBN) or diazabicycloundecene (DBU).

When carrying out the process (c) according to the invention, the reaction temperatures can be varied within a relatively wide range. In general, the process is carried out at temperatures of from 0°C to 150°C, preferably at temperatures of from 20°C to 110°C.

20

For carrying out the process (c) according to the invention for preparing the compounds of the formula (I), in general from 0.2 to 5 mol, preferably from 0.5 to 2 mol, of halide of the formula (V) are employed per mole of the pyrazolylcarboxanilide of the formula (Ia).

25 Unless indicated otherwise, all processes according to the invention are generally carried out under atmospheric pressure. However, it is also possible to operate under elevated or reduced pressure – in general between 0.1 bar and 10 bar.

30 The substances according to the invention have potent microbicidal activity and can be employed for controlling unwanted microorganisms, such as fungi and bacteria, in crop protection and in the protection of materials.

Fungicides can be employed in crop protection for controlling Plasmodiophoromycetes, Oomycetes, Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes.

35

Bactericides can be employed in crop protection for controlling Pseudomonadaceae, Rhizobiaceae, Enterobacteriaceae, Corynebacteriaceae and Streptomyetaceae.

Some pathogens causing fungal and bacterial diseases which come under the generic names listed above may be mentioned as examples, but not by way of limitation:

- Xanthomonas species, such as, for example, *Xanthomonas campestris* pv. *oryzae*;
- 5 *Pseudomonas* species, such as, for example, *Pseudomonas syringae* pv. *lachrymans*;
- Erwinia* species, such as, for example, *Erwinia amylovora*;
- Pythium* species, such as, for example, *Pythium ultimum*;
- Phytophthora* species, such as, for example, *Phytophthora infestans*;
- Pseudoperonospora* species, such as, for example, *Pseudoperonospora humuli* or
- 10 *Pseudoperonospora cubensis*;
- Plasmopara* species, such as, for example, *Plasmopara viticola*;
- Bremia* species, such as, for example, *Bremia lactucae*;
- Peronospora* species, such as, for example, *Peronospora pisi* or *P. brassicae*;
- Erysiphe* species, such as, for example, *Erysiphe graminis*;
- 15 *Sphaerotheca* species, such as, for example, *Sphaerotheca fuliginea*;
- Podosphaera* species, such as, for example, *Podosphaera leucotricha*;
- Venturia* species, such as, for example, *Venturia inaequalis*;
- Pyrenophora* species, such as, for example, *Pyrenophora teres* or *P. graminea*  
(conidia form: *Drechslera*, syn: *Helminthosporium*);
- 20 *Cochliobolus* species, such as, for example, *Cochliobolus sativus*  
(conidia form: *Drechslera*, syn: *Helminthosporium*);
- Uromyces* species, such as, for example, *Uromyces appendiculatus*;
- Puccinia* species, such as, for example, *Puccinia recondita*;
- Sclerotinia* species, such as, for example, *Sclerotinia sclerotiorum*;
- 25 *Tilletia* species, such as, for example, *Tilletia caries*;
- Ustilago* species, such as, for example, *Ustilago nuda* or *Ustilago avenae*;
- Pellicularia* species, such as, for example, *Pellicularia sasakii*;
- Pyricularia* species, such as, for example, *Pyricularia oryzae*;
- Fusarium* species, such as, for example, *Fusarium culmorum*;
- 30 *Botrytis* species, such as, for example, *Botrytis cinerea*;
- Septoria* species, such as, for example, *Septoria nodorum*;
- Leptosphaeria* species, such as, for example, *Leptosphaeria nodorum*;
- Cercospora* species, such as, for example, *Cercospora canescens*;
- Alternaria* species, such as, for example, *Alternaria brassicae*; and
- 35 *Pseudocercospora* species, such as, for example, *Pseudocercospora herpotrichoides*,
- Rhizoctonia* species, such as, for example, *Rhizoctonia solani*.

The active compounds according to the invention also show a strong invigorating action in plants. Accordingly, they are suitable for mobilizing the internal defences of the plant against attack by unwanted microorganisms.

5 In the present context, plant-invigorating (resistance-inducing) compounds are to be understood as meaning substances which are capable of stimulating the defence system of plants such that, when the treated plants are subsequently inoculated with unwanted microorganisms, they display substantial resistance to these microorganisms.

10 In the present case, unwanted microorganisms are to be understood as meaning phytopathogenic fungi, bacteria and viruses. The compounds according to the invention can thus be used to protect plants within a certain period of time after treatment against attack by the pathogens mentioned. The period of time for which this protection is achieved generally extends for 1 to 10 days, preferably 1 to 7 days, from the treatment of the plants with the active compounds.

15

The fact that the active compounds are well tolerated by plants at the concentrations required for controlling plant diseases permits the treatment of above-ground parts of plants, of propagation stock and seeds, and of the soil.

20 The active compounds according to the invention can be used with particularly good results for controlling cereal diseases, such as, for example, against Puccinia species, and diseases in viticulture and the cultivation of fruits and vegetables, such as, for example, against Botrytis, Venturia or Alternaria species.

25 The active compounds according to the invention are also suitable for increasing the yield of crops. In addition, they show reduced toxicity and are well tolerated by plants.

If appropriate, the active compounds according to the invention can, at certain concentrations and application rates, also be employed as herbicides, for regulating plant growth and for controlling  
30 animal pests. If appropriate, they can also be used as intermediates or precursors in the synthesis of other active compounds.

According to the invention, it is possible to treat all plants and parts of plants. Plants are to be understood here as meaning all plants and plant populations, such as desired and undesired wild  
35 plants or crop plants (including naturally occurring crop plants). Crop plants can be plants which can be obtained by conventional breeding and optimization methods or by biotechnological and genetic engineering methods or combinations of these methods, including the transgenic plants and

including plant cultivars which can or cannot be protected by plant breeders' certificates. Parts of plants are to be understood as meaning all above-ground and below-ground parts and organs of plants, such as shoot, leaf, flower and root, examples which may be mentioned being leaves, needles, stems, trunks, flowers, fruit-bodies, fruits and seeds and also roots, tubers and rhizomes.

5 Parts of plants also include harvested material and vegetative and generative propagation material, for example seedlings, tubers, rhizomes, cuttings and seeds.

The treatment of the plants and parts of plants according to the invention with the active compounds is carried out directly or by action on their environment, habitat or storage area  
10 according to customary treatment methods, for example by dipping, spraying, evaporating, atomizing, broadcasting, brushing-on and, in the case of propagation material, in particular in the case of seeds, furthermore by one- or multilayer coating.

In the protection of materials, the compounds according to the invention can be employed for  
15 protecting industrial materials against infection with, and destruction by, unwanted microorganisms.

Industrial materials in the present context are understood as meaning non-living materials which have been prepared for use in industry. For example, industrial materials which are intended to be  
20 protected by active compounds according to the invention from microbial change or destruction can be tackifiers, sizes, paper and board, textiles, leather, wood, paints and plastic articles, cooling lubricants and other materials which can be infected with, or destroyed by, microorganisms. Parts of production plants, for example cooling-water circuits, which may be impaired by the proliferation of microorganisms may also be mentioned within the scope of the materials to be  
25 protected. Industrial materials which may be mentioned within the scope of the present invention are preferably tackifiers, sizes, paper and board, leather, wood, paints, cooling lubricants and heat-transfer liquids, particularly preferably wood.

Microorganisms capable of degrading or changing the industrial materials which may be  
30 mentioned are, for example, bacteria, fungi, yeasts, algae and slime organisms. The active compounds according to the invention preferably act against fungi, in particular moulds, wood-discolouring and wood-destroying fungi (Basidiomycetes) and against slime organisms and algae.

Microorganisms of the following genera may be mentioned as examples:

35 *Alternaria*, such as *Alternaria tenuis*,  
*Aspergillus*, such as *Aspergillus niger*,  
*Chaetomium*, such as *Chaetomium globosum*,

= 2006/031011

- Coniophora, such as Coniophora puetana,  
Lentinus, such as Lentinus tigrinus,  
Penicillium, such as Penicillium glaucum,  
Polyporus, such as Polyporus versicolor,  
5 Aureobasidium, such as Aureobasidium pullulans,  
Sclerophoma, such as Sclerophoma pityophila,  
Trichoderma, such as Trichoderma viride,  
Escherichia, such as Escherichia coli,  
Pseudomonas, such as Pseudomonas aeruginosa, and  
10 Staphylococcus, such as Staphylococcus aureus.

Depending on their particular physical and/or chemical properties, the active compounds can be converted into the customary formulations, such as solutions, emulsions, suspensions, powders, foams, pastes, granules, aerosols and microencapsulations in polymeric substances and in coating  
15 compositions for seeds, and ULV cool and warm fogging formulations.

These formulations are produced in a known manner, for example by mixing the active compounds with extenders, that is liquid solvents, liquefied gases under pressure, and/or solid carriers, optionally with the use of surfactants, that is emulsifiers and/or dispersants, and/or foam formers.  
20 If the extender used is water, it is also possible to employ, for example, organic solvents as auxiliary solvents. Essentially, suitable liquid solvents are: aromatics such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example petroleum fractions, alcohols such as butanol or glycol and  
25 their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethylformamide or dimethyl sulphoxide, or else water. Liquefied gaseous extenders or carriers are to be understood as meaning liquids which are gaseous at standard temperature and under atmospheric pressure, for example aerosol propellants such as halogenated hydrocarbons, or else butane, propane, nitrogen and carbon dioxide. Suitable  
30 solid carriers are: for example ground natural minerals such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground synthetic minerals such as finely divided silica, alumina and silicates. Suitable solid carriers for granules are: for example crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, or else synthetic granules of inorganic and organic meals, and granules of organic material such as  
35 sawdust, coconut shells, maize cobs and tobacco stalks. Suitable emulsifiers and/or foam formers are: for example nonionic and anionic emulsifiers, such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates,

alkyl sulphates, arylsulphonates, or else protein hydrolysates. Suitable dispersants are: for example lignosulphite waste liquors and methylcellulose.

5 Tackifiers such as carboxymethylcellulose, natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, or else natural phospholipids such as cephalins and lecithins and synthetic phospholipids can be used in the formulations. Other possible additives are mineral and vegetable oils.

10 It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyestuffs such as alizarin dyestuffs, azo dyestuffs and metal phthalocyanine dyestuffs, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

15 The formulations generally comprise between 0.1 and 95 per cent by weight of active compound, preferably between 0.5 and 90%.

20 The active compounds according to the invention can, as such or in their formulations, also be used in a mixture with known fungicides, bactericides, acaricides, nematocides or insecticides, to broaden, for example, the activity spectrum or to prevent development of resistance. In many cases, synergistic effects are obtained, i.e. the activity of the mixture is greater than the activity of the individual components.

Suitable mixing components are, for example, the following compounds:

**Fungicides:**

25 2-phenylphenol; 8-hydroxyquinoline sulphate; acibenzolar-S-methyl; aldimorph; amidoflumet; ampropylfos; ampropylfos-potassium; andoprim; anilazine; azaconazole; azoxystrobin; benalaxyl; benodanil; benomyl; bentiavalicarb-isopropyl; benzamacril; benzamacril-isobutyl; bilanafos; binapacryl; biphenyl; bitertanol; blasticidin-S; bromuconazole; bupirimate; buthiobate; butylamine; calcium polysulphide; capsimycin; captafol; captan; carbendazim; carboxin; 30 carpropamid; carvone; chinomethionat; chlombenthiazole; chlorfenazole; chloroneb; chlorothalonil; chlozolate; clozylacon; cyazofamid; cyflufenamid; cymoxanil; cyproconazole; cyprodinil; cyprofuram; Dagger G; debacarb; dichlofluanid; dichlone; dichlorophen; diclocymet; diclomezine; dicloran; diethofencarb; difenoconazole; diflumetorim; dimethirimol; dimethomorph; dimoxystrobin; diniconazole; diniconazole-M; dinocap; diphenylamine; dipyrithione; ditalimfos; dithianon; dodine; drazoxolon; edifenphos; epoxiconazole; ethaboxam; ethirimol; etridiazole; famoxadone; fenamidone; fenapanil; fenarimol; fenbuconazole; fenfuram; fenhexamid; fenitropan; 35 fenoxanil; fenpiclonil; fenpropidin; fenpropimorph; ferbam; fluazinam; flubenzimine; fludioxonil;

flumetover; flumorph; fluoromide; fluoxastrobin; fluquinconazole; flurprimidol; flusilazole; flusulphamide; flutolanil; flutriafol; folpet; fosetyl-Al; fosetyl-sodium; fuberidazole; furalaxyl; furametpyr; furcarbanil; furmecyclox; guazatine; hexachlorobenzene; hexaconazole; hymexazole; imazalil; imibenconazole; iminoctadine triacetate; iminoctadine tris(albesil); iodocarb; ipconazole; iprobenfos; iprodione; iprovalicarb; irumamycin; isoprothiolane; isovalledione; kasugamycin; kresoxim-methyl; mancozeb; maneb; meferimzone; mepanipyrim; mepronil; metalaxyl; metalaxyl-M; metconazole; methasulphocarb; methfuroxam; metiram; metominostrobin; metsulphovax; mildiomyacin; myclobutanil; myclozolin; natamycin; nicobifen; nitrothal-isopropyl; noviflumuron; nuarimol; ofurace; oryastrobin; oxadixyl; oxolinic acid; oxpoconazole; oxycarboxin; oxyfenthiin; paclobutrazole; pefurazoate; penconazole; pencycuron; phosdiphen; phthalide; picoxystrobin; piperalin; polyoxins; polyoxorim; probenazole; prochloraz; procymidone; propamocarb; propanosine-sodium; propiconazole; propineb; proquinazid; prothioconazole; pyraclostrobin; pyrazophos; pyrifenox; pyrimethanil; pyroquilon; pyroxyfur; pyrrolenitrine; quinconazole; quinoxifen; quintozene; simeconazole; spiroxamine; sulphur; tebuconazole; tecloftalam; tecnazene; tetcyclacis; tetraconazole; thiabendazole; thicyofen; thifluzamide; thiophanate-methyl; thiram; tioxymid; tolclofos-methyl; tolyfluanid; triadimefon; triadimenol; triazbutil; triazoxide; tricyclamide; tricyclazole; tridemorph; trifloxystrobin; triflumizole; triforine; triticonazole; uniconazole; validamycin A; vinclozolin; zineb; ziram; zoxamide; (2S)-N-[2-[4-[[3-(4-chlorophenyl)-2-propynyl]oxy]-3-methoxyphenyl]ethyl]-3-methyl-2-[[[(methylsulphonyl)amino]butanamide; 1-(1-naphthalenyl)-1H-pyrrole-2,5-dione; 2,3,5,6-tetrachloro-4-(methylsulphonyl)pyridine; 2-amino-4-methyl-N-phenyl-5-thiazolecarboxamide; 2-chloro-N-(2,3-dihydro-1,1,3-trimethyl-1H-inden-4-yl)-3-pyridinecarboxamide; 3,4,5-trichloro-2,6-pyridinedicarbonitrile; actinovate; cis-1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol; methyl 1-(2,3-dihydro-2,2-dimethyl-1H-inden-1-yl)-1H-imidazole-5-carboxylate; monopotassium carbonate; N-(6-methoxy-3-pyridinyl)-cyclopropanecarboxamide; N-butyl-8-(1,1-dimethylethyl)-1-oxaspiro[4.5]decane-3-amine; sodium tetrathiocarbonate; and copper salts and preparations, such as Bordeaux mixture; copper hydroxide; copper naphthenate; copper oxychloride; copper sulphate; cufraneb; copper oxide; mancopper; oxine-copper.

30 **Bactericides:**

bronopol, dichlorophen, nitrapyrin, nickel dimethyldithiocarbamate, kasugamycin, octhilinone, furancarboxylic acid, oxytetracyclin, probenazole, streptomycin, tecloftalam, copper sulphate and other copper preparations.

35 **Insecticides / acaricides / nematocides:**

abamectin, ABG-9008, acephate, acequinocyl, acetamiprid, acetoprole, acrinathrin, AKD-1022, AKD-3059, AKD-3088, alanycarb, aldicarb, aldoxycarb, allethrin, allethrin 1R-isomers, alpha-

cypermethrin (alphamethrin), amidoflumet, aminocarb, amitraz, avermectin, AZ-60541, azadirachtin, azamethiphos, azinphos-methyl, azinphos-ethyl, azocyclotin, *Bacillus popilliae*, *Bacillus sphaericus*, *Bacillus subtilis*, *Bacillus thuringiensis*, *Bacillus thuringiensis* strain EG-2348, *Bacillus thuringiensis* strain GC-91, *Bacillus thuringiensis* strain NCTC-11821, baculoviruses, 5 *Beauveria bassiana*, *Beauveria tenella*, benclothiaz, bendiocarb, benfuracarb, bensultap, benzoximate, beta-cyfluthrin, beta-cypermethrin, bifenazate, bifenthrin, binapacryl, bioallethrin, bioallethrin-S-cyclopentyl-isomer, bioethanomethrin, biopermethrin, bioresmethrin, bistrifluron, BPMC, brofenprox, bromophos-ethyl, bromopropylate, bromfeninfos (-methyl), BTG-504, BTG-505, bufencarb, buprofezin, butathiofos, butocarboxim, butoxycarboxim, butylpyridaben, 10 cadusafos, camphechlor, carbaryl, carbofuran, carbophenothion, carbosulphan, cartap, CGA-50439, chinomethionat, chlordane, chlordimeform, chloethocarb, chlorethoxyfos, chlorfenapyr, chlorfenvinphos, chlorfluazuron, chlormephos, chlorobenzilate, chloropicrin, chlorproxyfen, chlorpyrifos-methyl, chlorpyrifos (-ethyl), chlovaporthrin, chromafenozide, cis-cypermethrin, cis-resmethrin, cis-permethrin, clocythrin, cloethocarb, clofentezine, clothianidin, clothiazoben, codlemone, coumaphos, cyanofenphos, cyanophos, cycloprene, cycloprothrin, *Cydia pomonella*, cyfluthrin, cyhalothrin, cyhexatin, cypermethrin, cyphenothrin (1R-trans-isomer), cyromazine, DDT, deltamethrin, demeton-S-methyl, demeton-S-methylsulphone, diafenthiuron, dialifos, diazinon, dichlofenthion, dichlorvos, dicofol, dicrotophos, dicyclanil, diflubenzuron, dimefluthrin, dimethoate, dimethylvinphos, dinobuton, dinocap, dinotefuran, diofenolan, disulphoton, docusat-sodium, dofenapyn, DOWCO-439, eflusilanate, emamectin, emamectin-benzoate, empenthrin (1R-isomer), 20 endosulphan, *Entomophthora* spp., EPN, esfvalerate, ethiofencarb, ethiprole, ethion, ethoprophos, etofenprox, etoxazole, etrimfos, famphur, fenamiphos, fenazaquin, fenbutatin oxide, fenfluthrin, fenitrothion, fenobucarb, fenothiocarb, fenoxacrim, fenoxycarb, fenpropathrin, fenpyrad, fenpyrithrin, fenpyroximate, fensulphothion, fenthion, fentrifanil, fenvalerate, fipronil, flonicamid, fluacrypyrim, fluazuron, flubenzimine, flubrocycythrinate, flucyclohexuron, flucythrinate, flufenerim, flufenoxuron, flufenprox, flumethrin, flupyrazofos, flutenzin (flufenzine), fluvalinate, fonofos, formetanate, formothion, fosmethilan, fosthiazate, fubfenprox (fluproxyfen), furathiocarb, gamma-cyhalothrin, gamma-HCH, gossypure, grandlure, granulosis viruses, halfenprox, halofenozide, HCH, HCN-801, heptenophos, hexaflumuron, hexythiazox, hydramethylnone, hydroprene, IKA- 25 2002, imidacloprid, imiprothrin, indoxacarb, iodofenphos, iprobenfos, isazofos, isofenphos, isoprocarb, isoxathion, ivermectin, japonilure, kadethrin, nuclear polyhedrosis viruses, kinoprene, lambda-cyhalothrin, lindane, lufenuron, malathion, mecarbam, mesulphenfos, metaldehyde, metam-sodium, methacrifos, methamidophos, *Metharhizium anisopliae*, *Metharhizium flavoviride*, methidathion, methiocarb, methomyl, methoprene, methoxychlor, methoxyfenozide, metofluthrin, metolcarb, metoxadiazone, mevinphos, milbemectin, milbemycin, MKI-245, MON-45700, monocrotophos, moxidectin, MTI-800, naled, NC-104, NC-170, NC-184, NC-194, NC-196, niclosamide, nicotine, nitenpyram, nithiazine, NNI-0001, NNI-0101, NNI-0250, NNI-9768, 35

novaluron, noviflumuron, OK-5101, OK-5201, OK-9601, OK-9602, OK-9701, OK-9802, omethoate, oxamyl, oxydemeton-methyl, Paecilomyces fumosoroseus, parathion-methyl, parathion (-ethyl), permethrin (cis-, trans-), petroleum, PH-6045, phenothrin (1R-trans isomer), phenthoate, phorate, phosalone, phosmet, phosphamidon, phosphocarb, phoxim, piperonyl butoxide, pirimicarb, pirimiphos-methyl, pirimiphos-ethyl, potassium oleate, prallethrin, profenofos, profluthrin, promecarb, propaphos, propargite, propetamphos, propoxur, prothiofos, prothoate, protrifenbute, pymetrozine, pyraclofos, pyresmethrin, pyrethrum, pyridaben, pyridalyl, pyridaphenthion, pyridathion, pyrimidifen, pyriproxyfen, quinalphos, resmethrin, RH-5849, ribavirin, RU-12457, RU-15525, S-421, S-1833, salithion, sebufos, SI-0009, silafluofen, spinosad, spirodiclofen, spiromesifen, sulphuramid, sulphotep, sulprofos, SZI-121, tau-fluvalinate, tebufenozide, tebufenpyrad, tebupirimfos, teflubenzuron, tefluthrin, temephos, temvinphos, terbam, terbufos, tetrachlorvinphos, tetradifon, tetramethrin, tetramethrin (1R-isomer), tetrasul, theta-cypermethrin, thiacloprid, thiamethoxam, thiapronil, thiatriphos, thiocyclam hydrogenoxalate, thiodicarb, thiofanox, thiometon, thiosultap-sodium, thuringiensin, tolfenpyrad, tralocyrthrin, tralomethrin, transfluthrin, triarathene, triazamate, triazophos, triazuron, trichlophenidine, trichlorfon, Trichoderma atroviride, triflumuron, trimethacarb, vamidothion, vaniliprole, verbutin, Verticillium lecanii, WL-108477, WL-40027, YI-5201, YI-5301, YI-5302, XMC, xylylcarb, ZA-3274, zeta-cypermethrin, zolaprofos, ZXI-8901, the compound 3-methylphenyl propylcarbamate (tsumacide Z), the compound 3-(5-chloro-3-pyridinyl)-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane-3-carbonitrile (CAS-Reg. No. 185982-80-3) and the corresponding 3-endo-isomer (CAS-Reg. No. 185984-60-5) (cf. WO-96/37494, WO-98/25923), and preparations which comprise insecticidally active plant extracts, nematodes, fungi or viruses.

A mixture with other known active compounds, such as herbicides, or with fertilizers and growth regulators, safeners and/or semiochemicals is also possible.

In addition, the compounds of the formula (I) according to the invention also have very good antimycotic activity. They have a very broad antimycotic activity spectrum in particular against dermatophytes and yeasts, moulds and diphasic fungi (for example against *Candida* species such as *Candida albicans*, *Candida glabrata*) and *Epidermophyton floccosum*, *Aspergillus* species such as *Aspergillus niger* and *Aspergillus fumigatus*, *Trichophyton* species such as *Trichophyton mentagrophytes*, *Microsporon* species such as *Microsporon canis* and *audouinii*. The list of these fungi does by no means limit the mycotic spectrum which can be covered, but is only for illustration.

The active compounds can be used as such, in the form of their formulations or the use forms prepared therefrom, such as ready-to-use solutions, suspensions, wettable powders, pastes, soluble

powders, dusts and granules. Application is carried out in a customary manner, for example by watering, spraying, atomizing, broadcasting, dusting, foaming, spreading, etc. It is furthermore possible to apply the active compounds by the ultra-low volume method, or to inject the active compound preparation or the active compound itself into the soil. It is also possible to treat the  
5 seeds of the plants.

When using the active compounds according to the invention as fungicides, the application rates can be varied within a relatively wide range, depending on the kind of application. For the treatment of parts of plants, the active compound application rates are generally between 0.1 and  
10 10,000 g/ha, preferably between 10 and 1000 g/ha. For seed dressing, the active compound application rates are generally between 0.001 and 50 g per kilogram of seed, preferably between 0.01 and 10 g per kilogram of seed. For the treatment of the soil, the active compound application rates are generally between 0.1 and 10,000 g/ha, preferably between 1 and 5000 g/ha.

15 As already mentioned above, it is possible to treat all plants and their parts according to the invention. In a preferred embodiment, wild plant species and plant cultivars, or those obtained by conventional biological breeding, such as crossing or protoplast fusion, and parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant cultivars obtained by genetic engineering, if appropriate in combination with conventional methods (Genetically  
20 Modified Organisms), and parts thereof, are treated. The term "parts" or "parts of plants" or "plant parts" has been explained above.

Particularly preferably, plants of the plant cultivars which are in each case commercially available or in use are treated according to the invention. Plant cultivars are to be understood as meaning  
25 plants having new properties ("traits") and which have been obtained by conventional breeding, by mutagenesis or by recombinant DNA techniques. They can be cultivars, varieties, bio- or genotypes.

Depending on the plant species or plant cultivars, their location and growth conditions (soils,  
30 climate, vegetation period, diet), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the substances and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt  
35 content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage

stability and/or processability of the harvested products are possible which exceed the effects which were actually to be expected.

The transgenic plants or plant cultivars (i.e. those obtained by genetic engineering) which are preferably to be treated according to the invention include all plants which, in the genetic modification, received genetic material which imparted particularly advantageous useful properties ("traits") to these plants. Examples of such properties are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products. Further and particularly emphasized examples of such properties are a better defence of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or viruses, and also increased tolerance of the plants to certain herbicidally active compounds. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), maize, soya beans, potatoes, cotton, tobacco, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and grapes), and particular emphasis is given to maize, soya beans, potatoes, cotton, tobacco and oilseed rape. Traits that are emphasized are in particular increased defence of the plants against insects, arachnids, nematodes and slugs and snails by toxins formed in the plants, in particular those formed in the plants by the genetic material from *Bacillus thuringiensis* (for example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF and also combinations thereof) (hereinbelow referred to as "Bt plants"). Traits that are also particularly emphasized are the increased defence of the plants against fungi, bacteria and viruses by systemic acquired resistance (SAR), systemin, phytoalexins, elicitors and resistance genes and correspondingly expressed proteins and toxins. Traits that are furthermore particularly emphasized are the increased tolerance of the plants to certain herbicidally active compounds, for example imidazolinones, sulphonylureas, glyphosate or phosphinotricin (for example the "PAT" gene). The genes which impart the desired traits in question can also be present in combination with one another in the transgenic plants. Examples of "Bt plants" which may be mentioned are maize varieties, cotton varieties, soya bean varieties and potato varieties which are sold under the trade names YIELD GARD® (for example maize, cotton, soya beans), KnockOut® (for example maize), StarLink® (for example maize), Bollgard® (cotton), Nucoton® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are maize varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosate, for example maize, cotton, soya bean), Liberty Link® (tolerance to phosphinotricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulphonylureas, for example maize). Herbicide-resistant plants (plants bred in a

conventional manner for herbicide tolerance) which may be mentioned also include the varieties sold under the name Clearfield® (for example maize). Of course, these statements also apply to plant cultivars which have these genetic traits or genetic traits still to be developed, and which will be developed and/or marketed in the future.

5


The plants listed can be treated according to the invention in a particularly advantageous manner with the compounds of the general formula (I) or the active compound mixtures according to the invention. The preferred ranges stated above for the active compounds or mixtures also apply to the treatment of these plants. Particular emphasis is given to the treatment of plants with the compounds or mixtures specifically mentioned in the present text.

10

The preparation and the use of the active compounds according to the invention is illustrated by the examples below.

## Preparation examples

### Example 1




5

At room temperature, 56.0 mg (2.3 mmol) of sodium hydride are added to a solution of 300.0 mg (0.78 mmol) of 5-fluoro-1-methyl-3-(trifluoromethyl)-N-[2-(1,3,3-trimethylbutyl)phenyl]-1H-pyrazole-4-carboxamide in 10.0 ml of tetrahydrofuran. The reaction mixture is stirred at room temperature for 15 min, and 366.6 mg (4.7 mmol) of acetyl chloride and 2.2 g (21.2 mmol) of acetic anhydride are added. After 16 h of heating under reflux, the reaction mixture is added to 100 ml of methanol and 3 ml of triethylamine, and the mixture is filtered and concentrated under reduced pressure. Recrystallization from cyclohexane gives 182 mg (54.7% of theory) of N-acetyl-5-fluoro-1-methyl-3-(trifluoromethyl)-N-[2-(1,3,3-trimethylbutyl)phenyl]-1H-pyrazole-4-carboxamide of logP (pH 2.3) = 4.63.

15

### Example 2




20 1.66 g (5.0 mmol) of 5-fluoro-1,3-dimethyl-N-[2-(1,3,3-trimethylbutyl)phenyl]-1H-pyrazole-4-carboxamide and 1.02 g (7.4 mmol) of ethyl chloro(oxo)acetate were dissolved in 20 ml of carbon tetrachloride and stirred under reflux for 4 h. The reaction mixture was cooled to room temperature and concentrated. This gave 2 g (93% of theory) of ethyl [(5-fluoro-1,3-dimethyl-1H-pyrazol-4-yl)carbonyl][2-(1,3,3-trimethylbutyl)phenyl]amino(oxo)acetate of logP (pH 2.3) = 4.40.

25

The compounds of the formula (I) listed in table 1 below are obtained analogously to examples 1 and 2 and in accordance with the instructions in the general descriptions of the processes.


Table 1



Ex.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	logP
3	-CH <sub>3</sub>	-H	-CH <sub>3</sub>	-CH <sub>3</sub>	3.64
4	-CH <sub>3</sub>	-H	-H	-CH <sub>3</sub>	3.34
5	-CF <sub>3</sub>	-H	-CH <sub>3</sub>	-CH <sub>3</sub>	4.45
6	-CF <sub>3</sub>	-H	-CH <sub>3</sub>	-CH <sub>2</sub> -O-CH <sub>3</sub>	4.60

5 **Preparation of starting materials of the formula (III)**

**Example (III-1)**




10 243 mg (3.1 mmol) of acetyl chloride were added slowly to a solution of 500 mg (2.8 mmol) of 2-(1,3-dimethylbutyl)phenylamine in 35 ml of acetonitrile. The reaction mixture was stirred at room temperature for 1 h and then poured onto water. After extraction with ethyl acetate, the organic phases were dried over sodium sulphate and concentrated under reduced pressure. This gave 600 mg (98% of theory) of N-[2-(1,3-dimethylbutyl)phenyl]acetamide of logP (pH 2.3) = 2.65.

15

The compounds of the formula (III) listed in table 2 below are obtained analogously to example (III-1) and in accordance with the instructions in the general descriptions of the processes.

Table 2



Ex.	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	logP
III-2	-H	-H	-C(=O)H	2.78
III-3	-H	-H	-C(=O)CH <sub>2</sub> OCH <sub>3</sub>	3.24

5 The logP values given in the preparation examples and tables above are determined in accordance with EEC Directive 79/831 Annexe V.A8 by HPLC (High Performance Liquid Chromatography) on a reversed-phase column (C 18). Temperature: 43°C.

10 The determination in the acidic range is carried out at pH 2.3 using the mobile phases 0.1% aqueous phosphoric acid and acetonitrile; linear gradient from 10% acetonitrile to 90% acetonitrile.

Calibration was carried out using unbranched alkan-2-ones (with 3 to 16 carbon atoms) with known logP values (determination of the logP values by the retention times using linear interpolation between two successive alkanones).

15

The lambda max values were determined in the maxima of the chromatographic signals using the UV spectra from 200 nm to 400 nm.

**Use examples:**

**Example A**

**5 Podosphaera test (apple) / protective**

Solvents:                24.5 parts by weight of acetone  
                              24.5 parts by weight of dimethylacetamide  
Emulsifier :             1 part by weight of alkylaryl polyglycol ether

10

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvents and emulsifier, and the concentrate is diluted with water to the desired concentration.


15 To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, the plants are inoculated with an aqueous spore suspension of the apple mildew pathogen *Podosphaera leucotricha*. The plants are then placed in a greenhouse at about 23°C and a relative atmospheric humidity of about 70%.

20 Evaluation is carried out 10 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table A


**Podosphaera test (apple) / protective**

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
--	---	---------------



100

98



100

97

Example B

**Venturia test (apple) / protective**

- 5 Solvents:            24.5 parts by weight of acetone  
                         24.5 parts by weight of dimethylacetamide  
Emulsifier :            1 part by weight of alkylaryl polyglycol ether

10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvents and emulsifier, and the concentrate is diluted with water to the desired concentration.



15 To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, the plants are inoculated with an aqueous conidia suspension of the apple scab pathogen *Venturia inaequalis* and then remain in an incubation cabinet at about 20°C and 100% relative atmospheric humidity for one day.

20 The plants are then placed in a greenhouse at about 21°C and a relative atmospheric humidity of about 90%.

Evaluation is carried out 10 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table B

**Venturia test (apple) / protective**

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
	100	100
	100	100

Example C

**Botrytis test (bean) / protective**

- 5 Solvents:                    24.5 parts by weight of acetone  
                                  24.5 parts by weight of dimethylacetamide  
Emulsifier :                    1 part by weight of alkylaryl polyglycol ether

10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.


15 To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, 2 small pieces of agar colonized by *Botrytis cinerea* are placed onto each leaf. The inoculated plants are placed in a dark chamber at about 20°C and 100% relative atmospheric humidity.

20 2 days after the inoculation, the size of the infected areas on the leaves is evaluated. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table C


Botrytis test (bean) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
--	---	---------------



500

100



500

97

Example D

**Puccinia test (wheat) / protective**

- 5 Solvent: 25 parts by weight of N,N-dimethylacetamide  
Emulsifier: 0.6 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the  
10 desired concentration.




To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, the plants are sprayed with a conidia suspension of *Puccinia recondita*. The plants remain in an incubation cabinet at 20°C and 100%  
15 relative atmospheric humidity for 48 hours.

The plants are then placed in a greenhouse at a temperature of about 20°C and a relative atmospheric humidity of 80% to promote the development of rust pustules.

20 Evaluation is carried out 10 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table D

Puccinia test (wheat) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
	500	100
	500	100
	500	100

Example E

**Sphaerotheca test (cucumber) / protective**

- 5 Solvent: 49 parts by weight of N,N-dimethylformamide  
Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the  
10 desired concentration.


To test for protective activity, young cucumber plants are sprayed with the preparation of active compound at the stated application rate. 1 day after the treatment, the plants are inoculated with a spore suspension of *Sphaerotheca fuliginea*. The plants are then placed in a greenhouse at 70%  
15 relative atmospheric humidity and a temperature of 23°C.

Evaluation is carried out 7 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table E


Sphaerotheca test (cucumber) / protective

Active compound according to the invention	Application rate of active compound in g/ha	Efficacy in %
--	---	---------------



750

100




750

100

**Patent claims**

1. N-substituted pyrazolylcarboxanilides of the formula (I)



(I)

5 in which

R<sup>1</sup> represents methyl, trifluoromethyl or difluoromethyl,R<sup>2</sup> represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl,a) R<sup>3</sup> represents hydrogen,

R<sup>4</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)-carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms; (C<sub>3</sub>-C<sub>8</sub>-cycloalkyl)carbonyl; (C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl)carbonyl having 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

20 b) R<sup>3</sup> represents halogen, C<sub>1</sub>-C<sub>8</sub>-alkyl or C<sub>1</sub>-C<sub>8</sub>-haloalkyl,

R<sup>4</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)-carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms; (C<sub>1</sub>-C<sub>8</sub>-alkyl)carbonyl, (C<sub>1</sub>-C<sub>8</sub>-alkoxy)carbonyl, (C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl)-carbonyl, (C<sub>3</sub>-C<sub>8</sub>-cycloalkyl)carbonyl; (C<sub>1</sub>-C<sub>6</sub>-haloalkyl)carbonyl, (C<sub>1</sub>-C<sub>6</sub>-haloalkoxy)carbonyl, (halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>8</sub>-

30

halocycloalkyl)carbonyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; or  $-C(=O)C(=O)R^5$ ,  $-CONR^6R^7$  or  $-CH_2NR^8R^9$ ,

5  $R^5$  represents hydrogen,  $C_1-C_8$ -alkyl,  $C_1-C_8$ -alkoxy,  $C_1-C_4$ -alkoxy- $C_1-C_4$ -alkyl,  $C_3-C_8$ -cycloalkyl;  $C_1-C_6$ -haloalkyl,  $C_1-C_6$ -haloalkoxy, halo- $C_1-C_4$ -alkoxy- $C_1-C_4$ -alkyl,  $C_3-C_8$ -halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,

$R^6$  and  $R^7$  independently of one another, each represent hydrogen,  $C_1-C_8$ -alkyl,  $C_1-C_4$ -alkoxy- $C_1-C_4$ -alkyl,  $C_3-C_8$ -cycloalkyl;  $C_1-C_8$ -haloalkyl, halo- $C_1-C_4$ -alkoxy- $C_1-C_4$ -alkyl,  $C_3-C_8$ -halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,

10  $R^6$  and  $R^7$  furthermore together with the nitrogen atom to which they are attached form a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen and  $C_1-C_4$ -alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and  $NR^{10}$ ,

15  $R^8$  and  $R^9$  independently of one another, represent hydrogen,  $C_1-C_8$ -alkyl,  $C_3-C_8$ -cycloalkyl;  $C_1-C_8$ -haloalkyl,  $C_3-C_8$ -halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,

20  $R^8$  and  $R^9$  furthermore together with the nitrogen atom to which they are attached form a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen and  $C_1-C_4$ -alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and  $NR^{10}$ ,

$R^{10}$  represents hydrogen or  $C_1-C_6$ -alkyl.

2. N-substituted pyrazolylcarboxanilides of the formula (I) according to Claim 1 in which

25  $R^1$  represents methyl, trifluoromethyl or difluoromethyl,

$R^2$  represents hydrogen, fluorine, chlorine, methyl or trifluoromethyl,

a)  $R^3$  represents hydrogen,

30  $R^4$  represents  $C_1-C_6$ -alkyl,  $C_1-C_4$ -alkylsulphinyl,  $C_1-C_4$ -alkylsulphonyl,  $C_1-C_3$ -alkoxy- $C_1-C_3$ -alkyl,  $C_3-C_6$ -cycloalkyl;  $C_1-C_4$ -haloalkyl,  $C_1-C_4$ -haloalkylthio,  $C_1-C_4$ -haloalkylsulphinyl,  $C_1-C_4$ -haloalkylsulphonyl, halo- $C_1-C_3$ -alkoxy- $C_1-C_3$ -alkyl,  $C_3-C_6$ -halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl- $C_1-C_3$ -alkyl, ( $C_1-C_3$ -alkyl)carbonyl- $C_1-C_3$ -alkyl, ( $C_1-C_3$ -alkoxy)carbonyl- $C_1-C_3$ -alkyl; halo-( $C_1-C_3$ -alkyl)carbonyl- $C_1-C_3$ -alkyl, halo-( $C_1-C_3$ -alkoxy)carbonyl- $C_1-C_3$ -alkyl having  
35 in each case 1 to 13 fluorine, chlorine and/or bromine atoms;

(C<sub>3</sub>-C<sub>6</sub>-cycloalkyl)carbonyl; (C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl)carbonyl having 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

5 b) R<sup>3</sup> represents fluorine, chlorine, bromine, iodine, C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>1</sub>-C<sub>6</sub>-haloalkyl having 1 to 13 fluorine, chlorine and/or bromine atoms,

R<sup>4</sup> represents C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms;

10 (C<sub>1</sub>-C<sub>6</sub>-alkyl)carbonyl, (C<sub>1</sub>-C<sub>6</sub>-alkoxy)carbonyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>6</sub>-cycloalkyl)carbonyl; (C<sub>1</sub>-C<sub>4</sub>-haloalkyl)carbonyl, (C<sub>1</sub>-C<sub>4</sub>-haloalkoxy)carbonyl, (halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl)carbonyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

15 R<sup>5</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy, halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,

20 R<sup>6</sup> and R<sup>7</sup> independently of one another, each represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, halo-C<sub>1</sub>-C<sub>3</sub>-alkoxy-C<sub>1</sub>-C<sub>3</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,


25 R<sup>6</sup> and R<sup>7</sup> furthermore together with the nitrogen atom to which they are attached form a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of halogen and C<sub>1</sub>-C<sub>4</sub>-alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and NR<sup>10</sup>,

30 R<sup>8</sup> and R<sup>9</sup> independently of one another, represent hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl; C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms,

35 R<sup>8</sup> and R<sup>9</sup> furthermore together with the nitrogen atom to which they are attached form a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of halogen and C<sub>1</sub>-C<sub>4</sub>-alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and NR<sup>10</sup>,

R<sup>10</sup> represents hydrogen or C<sub>1</sub>-C<sub>4</sub>-alkyl.

3. N-substituted pyrazolylcarboxanilides of the formula (Ib)




5 in which

R<sup>4A</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms;

(C<sub>3</sub>-C<sub>8</sub>-cycloalkyl)carbonyl; (C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl)carbonyl having 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

15 and R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined in Claim 1.

4. N-substituted pyrazolylcarboxanilides of the formula (Ic)



20 in which

R<sup>3B</sup> represents halogen, C<sub>1</sub>-C<sub>8</sub>-alkyl or C<sub>1</sub>-C<sub>8</sub>-haloalkyl,

R<sup>4B</sup> represents C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphinyl, C<sub>1</sub>-C<sub>6</sub>-alkylsulphonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl; C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylthio, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphinyl, C<sub>1</sub>-C<sub>4</sub>-haloalkylsulphonyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; formyl, formyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, (C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl; halo-(C<sub>1</sub>-C<sub>3</sub>-alkyl)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl, halo-(C<sub>1</sub>-C<sub>3</sub>-alkoxy)carbonyl-C<sub>1</sub>-C<sub>3</sub>-alkyl having in each case 1 to 13 fluorine, chlorine and/or bromine atoms;

25

(C<sub>1</sub>-C<sub>8</sub>-alkyl)carbonyl, (C<sub>1</sub>-C<sub>8</sub>-alkoxy)carbonyl, (C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>8</sub>-cycloalkyl)carbonyl; (C<sub>1</sub>-C<sub>6</sub>-haloalkyl)carbonyl, (C<sub>1</sub>-C<sub>6</sub>-haloalkoxy)carbonyl, (halo-C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl)carbonyl, (C<sub>3</sub>-C<sub>8</sub>-halocycloalkyl)carbonyl having in each case 1 to 9 fluorine, chlorine and/or bromine atoms; or -C(=O)C(=O)R<sup>5</sup>, -CONR<sup>6</sup>R<sup>7</sup> or -CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,

and R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are as defined in Claim 1.

5. N-substituted pyrazolylcarboxanilides of the formula (I) according to Claim 1 in which R<sup>4</sup> represents formyl.


10

6. N-substituted pyrazolylcarboxanilides of the formula (I) according to Claim 1 in which R<sup>4</sup> represents -C(=O)C(=O)R<sup>5</sup> and R<sup>5</sup> is as defined in Claim 1.

7. Process for preparing compounds of the formula (I) according to Claim 1, characterized in that

15

a) carboxylic acid derivatives of the formula (II)




in which

R<sup>1</sup> is as defined above and

20

X<sup>1</sup> represents halogen or hydroxyl

are reacted with an aniline derivative of the formula (III)




in which R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as defined above,

if appropriate in the presence of a catalyst, if appropriate in the presence of a condensing agent, if appropriate in the presence of an acid binder and if appropriate in the presence of a diluent,

25

or


b) pyrazolylcarboxanilides of the formula (IV)



in which  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are as defined above  
are hydrogenated, if appropriate in the presence of a diluent and if appropriate in the presence of a catalyst,

5 or

c) pyrazolylcarboxanilides of the formula (Ia)



in which  $R^1$ ,  $R^2$  and  $R^3$  are as defined above  
are reacted with halides of the formula (V)



in which

$R^4$  is as defined above and

$X^2$  represents chlorine, bromine or iodine

in the presence of a base and in the presence of a diluent.

15


8. Compositions for controlling unwanted microorganisms, characterized in that they comprise at least one N-substituted pyrazolylcarboxanilide of the formula (I) according to Claim 1, in addition to extenders and/or surfactants.

20 9. Use of N-substituted pyrazolylcarboxanilides of the formula (I) according to Claim 1 for controlling unwanted microorganisms.

10. Method for controlling unwanted microorganisms, characterized in that N-substituted pyrazolylcarboxanilides of the formula (I) according to Claim 1 are applied to the  
25 microorganisms and/or their habitat.


11. Process for preparing compositions for controlling unwanted microorganisms, characterized in that N-substituted pyrazolylcarboxanilides of the formula (I) according to Claim 1 are mixed with extenders and/or surfactants.

- 5 12. Aniline derivatives of the formula (III)



in which R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as defined in Claim 1.

13. Pyrazolylcarboxanilides of the formula (IV)



10

in which R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are as defined in Claim 1.

14. N-substituted pyrazolylcarboxanilides of the formula (I) substantially as herein described with reference to and as exemplified in the process of Examples 1 to 6.
15. Process for preparing N-substituted pyrazolylcarboxanilides of the formula (I) substantially as herein described with reference to and as exemplified in any one of Examples 1 to 6.
16. Compositions for controlling unwanted microorganisms substantially as herein described with reference to and as exemplified in any one of Examples A to E.
17. Use of compositions as claimed in claim 14 for controlling unwanted micro organisms.
18. Method for controlling unwanted microorganisms substantially as herein described with reference to and as exemplified in any one of Examples A to E.

- 46a -

19. Process for preparing compositions for controlling unwanted microorganisms substantially as herein described with reference to and as exemplified in any one of Examples A to E.
  
20. Aniline derivatives of formula (III) substantially as herein described with reference to and as exemplified in any one of Examples III-1 to III-3.