

(19) **DANMARK**

(10) **DK/EP 3701944 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 9/20 (2006.01)** **A 61 K 31/565 (2006.01)** **A 61 P 5/00 (2006.01)**
A 61 P 15/18 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2022-03-14**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2021-12-08**
- (86) Europæisk ansøgning nr.: **20171407.8**
- (86) Europæisk indleveringsdag: **2016-06-17**
- (87) Den europæiske ansøgnings publiceringsdag: **2020-09-02**
- (30) Prioritet: **2015-06-18 EP 15172767**
- (62) Stamansøgningsnr: **16733341.8**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Estetra SRL, Rue Saint-Georges 5-7, 4000 Liège, Belgien**
- (72) Opfinder: **JASPART, Séverine Francine Isabelle, Rue des condruzes 19, 4560 Bois-et-Borsu, Belgien**
Platteeuw, Johannes Jan, Newtonplein 41, 5283 JH Boxtel, Holland
van den Heuvel, Denny Johan Marijn, Ganzeneiland 11, 6642 DJ Beuningen, Holland
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **ORO-DISPERGERBAR DOSERINGSENHED INDEHOLDENDE EN ESTETROLKOMPONENT**
- (56) Fremdragne publikationer:
WO-A1-00/42955
WO-A1-02/094276
WO-A1-2015/086643
US-A1- 2005 070 488
US-A1- 2007 286 819

DESCRIPTION

TECHNICAL FIELD OF THE INVENTION

[0001] The present invention provides an orodispersible solid pharmaceutical dosage unit having a weight of 30-1,000 mg and containing at least 0.1 mg of an estetrol component selected from estetrol, estetrol esters and combinations thereof. This solid dosage unit consists of:

- 0.1-25 wt.% of estetrol particles containing at least 80 wt.% of the estetrol component; and
- 75-99.9 wt.% of one or more pharmaceutically acceptable ingredients.

[0002] The invention also provides a process of preparing the aforementioned solid dosage unit.

[0003] Furthermore, the invention relates to the use of the solid dosage unit in medical treatment, female hormone replacement therapy and female contraception, said use comprising sublingual, buccal or sublabial administration of the solid dosage unit.

BACKGROUND OF THE INVENTION

[0004] Estetrol is a human steroid, produced by the fetal liver during pregnancy only. This natural hormone was discovered in urine of pregnant women by Diczfalussy and coworkers in 1965. Estetrol has the structure of an estrogenic steroid with four hydroxyl groups. Estetrol is synthesized in the fetal liver from estradiol and estriol by the two enzymes 15 α - and 16 α -hydroxylase. After birth the neonatal liver rapidly loses its capacity to synthesize estetrol because these two enzymes are no longer expressed.

[0005] Estetrol reaches the maternal circulation through the placenta and was already detected at nine weeks of pregnancy in maternal urine. During the second trimester of pregnancy high levels were found in maternal plasma, with steadily rising concentrations of unconjugated estetrol to about 1 ng/mL (> 3 nmol/L) towards the end of pregnancy. So far the physiological function of estetrol is unknown. The possible use of estetrol as a marker for fetal well-being has been studied quite extensively. However, due to the large intra- and inter-individual variation of maternal estetrol plasma levels during pregnancy this appeared not to be feasible.

[0006] Since 2001 estetrol has been studied extensively. In humans estetrol was shown to have a high and dose-proportional oral bioavailability and a long terminal elimination half-life of

about 28 hours. Results from *in vitro* studies showed that estetrol binds highly selective to the estrogen receptors with preference for the ER α form of the receptor, unlike the estrogens ethinyl estradiol and 17 β -estradiol. Also in contrast with ethinyl estradiol and especially with 17 β -estradiol, estetrol does not bind to sex hormone binding globulin (SHBG) and does not stimulate the production of SHBG *in vitro*.

[0007] The properties of estetrol have also been investigated in a series of predictive, well validated pharmacological *in vivo* rat models. In these models, estetrol exhibited estrogenic effects on the vagina, the uterus (both myometrium and endometrium), body weight, bone mass, bone strength, hot flushes and on ovulation (inhibition). All these effects of estetrol were dose-dependent with maximum effects at comparable dose levels. Surprisingly, estetrol prevented tumour development in a DMBA mammary tumour model to an extent and at a dose level similar to the anti-estrogen tamoxifen and to ovariectomy. This anti-estrogenic effect of estetrol in the presence of 17 β -estradiol has also been observed in *in vitro* studies using human breast cancer cells.

[0008] Buccal, sublingual or sublabial administration of estetrol is mentioned in a number of patent applications, including WO 2002/094275, WO 2002/094276, WO 2002/094278 and WO 2003/018026. Estetrol containing dosage units for buccal, sublingual or sublabial administration are not described in these publications.

[0009] WO 2010/033832 describes an oral dosage form comprising an estriol compound and a pharmaceutically acceptable matrix material, wherein the oral dosage form releases at least about 90% of the estriol compound in a time of less than about 300 seconds when contacted with saliva of the buccal and/or sublingual cavity.

[0010] US 2007/286829 describes an orally administered solid dosage form capable of delivering ethinyl estradiol with improved bioavailability, said solid dosage form comprising (i) about 0.5 μ g to about 50 μ g of ethinyl estradiol and (ii) an oral dissolution enhancing carrier that provides for at least 15% absorption of the ethinyl estradiol through the oral mucosa when said solid dosage form is orally administered to the patient with 2 ounces of water or less.

[0011] US 6,117,446 describes a buccal dosage unit for administering a combination of steroidal active agents, comprising a compressed tablet of a bioerodible polymeric carrier and therapeutically effective amounts of an androgenic agent selected from testosterone and pharmacologically acceptable esters thereof, a progestin and an estrogen. The examples describe buccal dosage units that were prepared by thoroughly mixing the following components: estrogen, progestogen, androgen, polyethylene oxide, carbomer and magnesium stearate. Next, the mixture was granulated by means of fluid bed granulation and the granulate so obtained was pressed into tablets.

[0012] Oral dosage units containing estetrol have been described in several patent publications.

[0013] WO 2002/094276 describes a pharmaceutical composition for use in a method of hormone replacement therapy, which method comprises administering to a person in need of such a therapy an effective amount of estetrol, said composition containing virtually no progestogen or anti-progestin. WO 2002/094276 describes the preparation of estetrol tablets having a weight of 185 mg, containing 1.5 mg estetrol, on the basis of the following formulation:

	mg
Estetrol	1.5
Polyvinylpyrrolidone (Kollidon 25 [®] ex BASF)	12.5
Lactose	135.795
Microcrystalline cellulose (Avicel PH 101 [®])	26.25
Glyceryl palmitostearate (Precirol [®])	2.775
Anhydrous colloidal silica (Aerosil 200 [®])	1.0
Crospovidone (Polyplasdone XL [®])	4.0
Coloring agent	0.18

[0014] WO 2002/094275 describes the use of an estetrol in a method of increasing libido in a woman, said method comprising administering to said woman an effective amount of estetrol. Oral administration is mentioned as a suitable mode of administration. This patent application describes the same estetrol tablet as WO 2002/094276.

[0015] WO 2002/094279 describes the use of estetrol in a method of contraception in mammalian females, which method comprises the oral administration of said estrogenic component and a progestogenic component to a female of childbearing capability in an effective amount to inhibit ovulation. The following formulation for a 185 mg estetrol tablet is described in this international patent application.

	mg
Estetrol	1.5
Levonorgestrel	0.15
Polyvinylpyrrolidone (Kollidon 25 [®] ex BASF)	13.5
Lactose	135.645
Microcrystalline cellulose (Avicel PH 101 [®])	26.25
Glyceryl palmitostearate (Precirol [®])	2.775
Anhydrous colloidal silica (Aerosil 200 [®])	1.0
Crospovidone (Polyplasdone XL [®])	4.0
Coloring agent	0.18

[0016] WO 2003/041718 describes the use of estetrol in a method of hormone replacement in mammals, which method comprises the oral administration of estetrol and a progestogenic component to a mammal in an effective amount to prevent or treat symptoms of hypoestrogenism. This patent application describes the same estetrol tablet as WO 2002/094279.

[0017] WO 2007/081206 describes the use of estetrol in a method of treating an acute vascular disorder in a mammal, said method comprising orally administering to said mammal, upon demand, an effective amount of the estetrol to the mammal. This patent application describes the preparation of hard gelatine capsules, containing 100 mg estetrol and 25 mg sildenafil citrate per capsule.

[0018] WO 2008/156365 describes the use of estetrol in the treatment of Meconium Aspiration Syndrome (MAS) in a newborn infant, said treatment comprising administering an effective amount of estrogen to said newborn infant within 7 days after birth. The international patent application describes a suppository for use in newborn infants comprising at least 1 µg of estrogen, said suppository further being characterized by a maximum diameter of less than 10 mm and a weight of less than 0.5 g. The excipient contained in the suppository may be based on lipid material that melts at body temperature or it may be based on a hydrophilic component that dissolves or disintegrates when it comes into contact with water.

SUMMARY OF THE INVENTION

[0019] The present invention provides an orodispersible solid pharmaceutical dosage unit containing an estetrol component. The dosage unit rapidly releases the estetrol in aqueous environment. The solid dosage unit is easy to manufacture and perfectly suited for sublingual, buccal or sublabial administration. Sublingual, buccal and sublabial administration each offer the advantages that the estetrol component does not have to pass through the digestive system and avoids first-pass liver exposure. Furthermore, these modes of administration provide a rapid onset of action.

[0020] The solid dosage unit according to the present invention has a weight between 30 and 1,000 mg; contains at least 100 µg of an estetrol component selected from estetrol, estetrol esters and combinations thereof; and consists of:

- 0.1-25 wt.% of estetrol particles containing at least 80 wt.% of the estetrol component; and
- 75-99.9 wt.% of one or more pharmaceutically acceptable ingredients.

[0021] This solid dosage is obtainable by a process comprising:

- providing estetrol particles containing at least 80 wt.% of the estetrol component and having a volume median diameter of 2 μm to 50 μm ;
- mixing the estetrol particles with one or more granulation excipients to produce a granulation mixture;
- mixing the granulation mixture with a granulation liquid to produce estetrol-containing granules, said granulation liquid containing at least 60 wt.% of liquid solvent;
- removing liquid solvent from the estetrol-containing granules to produce dry estetrol-containing granules;
- optionally mixing the dry granules with one or more tableting excipients; and
- forming the dry granules or the mixture of the dry granules and the one or more tableting excipients into a solid dosage unit.

[0022] Rapid and complete dissolution of the estetrol component into saliva is essential for efficient delivery of the component via sublingual, buccal or sublabial administration of the solid dosage unit. The inventors have unexpectedly found that the estetrol component is rapidly released and dispersed into saliva and absorbed through the mucosal lining of the oral cavity if it is present in the solid dosage unit in the form of very small particles.

[0023] The invention also provides a process of preparing the aforementioned solid dosage unit, said process comprising the steps of:

- providing estetrol particles containing at least 80 wt.% of an estetrol component selected from estetrol, estetrol esters and combinations thereof and having a volume median diameter of 2 μm to 50 μm ;
- mixing the estetrol particles with one or more granulation excipients to produce a granulation mixture;
- mixing the granulation mixture with a granulation liquid to produce estetrol-containing granules, said granulation liquid containing at least 60 wt.% of liquid solvent;
- removing liquid solvent from the estetrol-containing granules to produce dry estetrol-containing granules;
- optionally mixing the dry granules with one or more pharmaceutically acceptable excipients; and
- forming the dry granules or the mixture of the dry granules and the one or more tableting excipients into a solid dosage unit

[0024] The production of the estetrol-containing granules in this process may suitably be carried out, for instance, in a high shear granulator, a low shear granulator or a fluidized bed granulator.

BRIEF DESCRIPTION OF THE FIGURE

[0025] Figure 1 illustrates the manufacturing process flow chart used in Example 4.

DETAILED DESCRIPTION OF THE INVENTION

[0026] A first aspect of the invention relates to an orodispersible solid pharmaceutical dosage unit having a weight between 30 and 1,000 mg, said dosage unit consisting of:

- 0.1-25 wt.% of estetrol particles containing at least 80 wt.% of an estetrol component selected from estetrol, estetrol esters and combinations thereof; and
- 75-99.9 wt.% of one or more pharmaceutically acceptable ingredients;

the solid dosage unit comprising at least 100 µg of the estetrol component; wherein the solid dosage unit can be obtained by a process comprising:

- providing estetrol particles containing at least 80 wt.% of the estetrol component and having a volume median diameter of 2 µm to 50 µm;
- mixing the estetrol particles with one or more granulation excipients to produce a granulation mixture;
- mixing the granulation mixture with a granulation liquid to produce estetrol-containing granules, said granulation liquid containing at least 60 wt.% of liquid solvent;
- removing liquid solvent from the estetrol-containing granules to produce dry estetrol-containing granules;
- optionally mixing the dry granules with one or more tableting excipients; and
- forming the dry granules or the mixture of the dry granules and the one or more tableting excipients into a solid dosage unit.

[0027] The term 'estetrol' as used herein refers to 1,3,5 (10)-estratrien-3,15α,16α,17β-tetrol or 15α-hydroxyestriol as well as hydrates of estetrol, e.g. estetrol monohydrate.

[0028] The term 'orodispersible dosage unit' as used herein refers to a dosage unit that is designed to rapidly disintegrate in the oral cavity when it comes into contact with saliva and to disperse the estetrol component into the saliva so it may be absorbed through the mucosal lining of the oral cavity.

[0029] The terms 'pharmaceutically acceptable ingredients' as used herein include both pharmaceutically acceptable excipients and pharmaceutically active ingredients other than the estetrol component, as further defined below.

[0030] The term 'sublingual' as used herein refers to the pharmacological route of administration by which the estetrol component diffuses into the blood through tissues under the tongue.

[0031] The term 'buccal' as used herein refers to the pharmacological route of administration by which the estetrol component diffuses into the blood through tissues of the buccal vestibule, the area inside the mouth between the lining of cheek (the buccal mucosa) and the teeth / gums.

[0032] The term 'sublabial' as used herein refers to the pharmacological route of administration by which the estetrol component is placed between the lip and the gingiva.

[0033] The term "granulation" as used herein, unless indicated otherwise, relates to a process in which primary powder particles are made to adhere to form larger, multiparticle entities called "granules".

[0034] The term 'tableting excipient' as used herein refers to a pharmaceutically acceptable excipient that can be used in the manufacture of solid dosage units such as tablets.

[0035] Unless indicated otherwise, all percentages mentioned herein are percentages by weight.

[0036] Examples of solid dosage units encompassed by the present invention include tablets, dragees, lozenges and films. In accordance with a preferred embodiment, the dosage unit is a tablet, most preferably a compressed tablet.

[0037] The solid dosage unit typically has a weight between 40 and 500 mg, more preferably between 50 and 300 mg, and most preferably between 70 and 150 mg.

[0038] The solid dosage unit preferably comprises at least 1 wt.%, more preferably 2-25 wt.% and most preferably 2.2-15 wt.% of the estetrol component.

[0039] The amount of the estetrol component contained in the solid dosage unit preferably lies within the range of 0.3-100 mg, more preferably of 0.5-40 mg and most preferably of 1-20 mg.

[0040] The estetrol component of the present invention preferably is selected from the group consisting of estetrol, esters of estetrol wherein the hydrogen atom of at least one of the hydroxyl groups has been substituted by an acyl radical of a hydrocarbon carboxylic, sulfonic acid or sulfamic acid of 1-25 carbon atoms; and combinations thereof. Even more preferably, the estetrol component is estetrol (including estetrol hydrates). Most preferably, the estetrol component contained in the dosage unit is estetrol monohydrate.

[0041] The particle size of the estetrol particles in the solid dosage unit should be adequate for achieving sufficient absorption of the estetrol component after sublingual, buccal or sublabial administration. The estetrol particles within the solid dosage unit and (independently) the estetrol particles used in the preparation of the solid dosage unit preferably have a volume median diameter in the range of 3 μm to 35 μm , more preferably in the range of 4 μm to 25 μm .

and most preferably in the range of 5 μm to 15 μm .

[0042] The estetrol particles within the solid dosage unit and (independently) the estetrol particles used in the preparation of the solid dosage unit preferably contain not more than a limited amount of particles with a particle size in excess of 60 μm . Preferably, not more than 10 vol.% of more than 60 μm (D_{90}), more preferably not more than 5 vol.% of the estetrol particles have a particle size of more than 60 μm (D_{95}). Even more preferably, not more than 10 vol.% of more than 40 μm (D_{90}), more preferably not more than 5 vol.% of the estetrol particles have a particle size of more than 40 μm (D_{95}).

[0043] The particles size distribution of the estetrol particles, and of other particulate materials used in the present process, may suitably be determined by means of laser diffraction. The particle size distribution of the estetrol particles within the solid dosage unit can suitably be determined using spectroscopic techniques, e.g. Raman mapping.

[0044] The solid dosage unit of the present invention offers the advantage that the estetrol component is rapidly released when the dosage unit is introduced into the oral cavity and comes into contact with saliva. The rate of release of the estetrol component from the dosage unit can suitably be determined using the dissolution test described in the Examples, or a disintegration test according to Ph. Eur. 2.9.1 ("Disintegration of tablets and capsules") and USP <701> ("Disintegration"), also described in the Examples. The solid dosage unit of the present invention, when subjected to the aforementioned dissolution test, typically releases at least 50%, more preferably at least 70% and most preferably at least 80% of the estetrol component after 5 minutes. The solid dosage unit of the present invention, when subjected to the aforementioned disintegration test, typically disintegrates within less than 5 minutes, more preferably within less than 2 minutes, still more preferably within less than 1,5 minutes, still more preferably within less than 1 minute, still more preferably within less than 45 seconds, and most preferably within less than 30 seconds.

[0045] The estetrol particles employed in the solid dosage unit and in the present process preferably contain at least 90 wt.% of the estetrol component, more preferably at least 95 wt.% of the estetrol component and most preferably at least 99 wt.% of the estetrol component. Besides the estetrol component, the estetrol particles can suitably contain pharmaceutically acceptable excipients that aid dispersion of the dosage unit and dissolution and absorption of the estetrol component. Examples of such excipients include tensioactive agents, cosolvents, absorption enhancer, superdisintegrants and buffering agents.

[0046] The estetrol particles typically represent between 0.5-20 wt.% of the dosage unit. More preferably, the estetrol particles represent 1-18 wt.%, most preferably 5-15 wt.% of the dosage unit.

[0047] The solid dosage unit of the present invention preferably contains 50-99.8 wt.% of water-soluble carbohydrate selected from maltose, fructose, sucrose, lactose, glucose,

galactose, trehalose, xylitol, sorbitol, erythritol, maltitol, mannitol, isomalt and combinations thereof. More preferably, the water-soluble carbohydrate is selected from lactose, mannitol, erythritol and combinations thereof. Even more preferably, the water-soluble carbohydrate is selected from mannitol, erythritol and combinations thereof. Most preferably, the water-soluble carbohydrate is mannitol.

[0048] The water-soluble carbohydrate is preferably contained in the dosage unit in crystalline form.

[0049] According to another preferred embodiment, the dosage unit contains 0.1-15 wt.%, more preferably 0.2-10 wt.% and most preferably 1-5 wt.% of a disintegrating agent selected from modified starches (e.g. sodium salt of carboxymethyl starch), crosslinked polyvinylpyrrolidone, crosslinked carmellose and combinations thereof.

[0050] The estetrol particles are present in the solid dosage unit of the present invention as a component of granules that additionally contain one or more granulation excipients.

[0051] The dry estetrol-containing granules typically constitute 20-99.9 wt.% of the oral dosage unit. More preferably, these granules represent 50-99.8 wt.%, most preferably 60-99.7 wt.% of the oral dosage unit.

[0052] Typically, the dry estetrol-containing granules that contain the estetrol particles have a volume median diameter of 100-4,000 μm , more preferably of 150-1,000 μm and most preferably of 200-600 μm .

[0053] The dry estetrol-containing granules typically comprise 70-95 wt.% of one or more granulation excipients and 5-30 wt.% of the estetrol component. Even more preferably, these granules comprise 75-90 wt.% of the one or more granulation excipients and 10-25 wt.% of the estetrol component.

[0054] Other granulation excipients that may suitably be incorporated into the estetrol-containing granules include water-soluble carbohydrate, diluents/fillers (e.g. calcium salts, microcrystalline cellulose), binders, disintegrating agents, mucoadhesive agents, flavouring, colouring, glidants, lubricants and combinations thereof.

[0055] The dry estetrol-containing granules preferably contain at least 20 wt.%, more preferably at least 35 wt.% and most preferably at least 45 wt.% of the water-soluble carbohydrate as defined herein before.

[0056] The dry estetrol-containing granules preferably contain at least 30%, more preferably at least 40% and most preferably at least 50% of the water-soluble carbohydrate by weight of the one or more granulation excipients.

[0057] The dry estetrol-containing granules typically contain 0-20 wt.% of the binder as defined

herein before. Even more preferably these granules contain 0.1-15 wt.%, more preferably 0.2-10 wt.% of the binder.

[0058] In accordance with another preferred embodiment, the dry estetrol-containing granules comprise 0.1-20 wt.%, more preferably 0.2-10 wt.% of a disintegrating agent as defined herein before.

[0059] Besides dry estetrol-containing granules, the solid dosage unit of the present invention may suitably contain 0.1-80 wt.%, more preferably 0.2-50 wt.% and most preferably 0.3-40 wt.% of tableting excipients selected from lactose, mannitol, xylitol, microcrystalline cellulose, starch, croscarmellose sodium, polyvinyl pyrrolidone and combinations thereof. According to one particularly preferred embodiment, the tableting excipients comprises at least 50 wt.% mannitol. In accordance with another preferred embodiment, the tableting excipients comprise at least 50 wt.% lactose.

[0060] The solid dosage unit may contain one or more other pharmaceutically active ingredients besides the estetrol component. Examples of such other pharmaceutically active ingredients include steroid hormones. The solid dosage unit of the present invention preferably contains 0.05-10 mg, more preferably 0.1-5 mg of one or more progestogens, preferably one or more progestogens selected from progesterone, levonorgestrel, norgestimate, norethisterone, norethisteron-acetate (NETA), dydrogesterone, drospirenone, 3-beta-hydroxydesogestrel, 3-keto desogestrel (=etonogestrel), 17-deacetyl norgestimate, 19-norprogesterone, acetoxypregnenolone, allylestrenol, anagestone, chlormadinone, cyproterone, demegestone, desogestrel, dienogest, dihydrogesterone, dimethisterone, ethisterone, ethynodiol diacetate, flurogesterone acetate, gastrinon, gestodene, gestrinone, hydroxymethylprogesterone, hydroxyprogesterone, lynestrenol (=lynoestrenol), medrogestone, medroxyprogesterone, megestrol, melengestrol, nestorone, nomegestrol, nomegestrol-acetate (NOMAC), norethindrone (=norethisterone), norethynodrel, norgestrel (includes d-norgestrel and dl-norgestrel), norgestrienone, normethisterone, progesterone, quingestanol, (17alpha)-17-hydroxy-11-methylene-19-norpregna-4,15-diene-20-yn-3-one, tibolone, trimegestone, algestone acetophenide, nestorone, promegestone, 17-hydroxyprogesterone esters, 19-nor-17hydroxyprogesterone, 17alpha-ethinyl-testosterone, 17alpha-ethinyl-19-nor-testosterone, d-17beta-acetoxy-13beta-ethyl-17alpha-ethinyl-gon-4-en-3-one oxime and prodrugs of these compounds. Preferably the one or more progestogens used in accordance with the present invention is selected from the group consisting of progesterone, desogestrel, etonogestrel, gestodene, dienogest, levonorgestrel, norgestimate, norethisterone, norethisteron-acetate (NETA), nomegestrol, nomegestrol-acetate (NOMAC), drospirenone, trimegestone, nestorone and dydrogesterone.

[0061] The solid dosage unit according to the present invention preferably contains 0.05-100 mg, more preferably 0.1-50 mg of one or more androgens, preferably one or more androgens selected from testosterone, dehydroepiandrosterone (DHEA); DHEA-sulphate (DHEAS); testosterone esters (e.g. testosterone undecanoate, testosterone propionate, testosterone phenylpropionate, testosterone isohexanoate, testosterone enantate, testosterone bucanate,

testosterone decanoate, testosterone buciclate); methyltestosterone; mesterolone; stanozolol; androstenedione; dihydrotestosterone; androstenediol; metenolon; fluoxymesterone; oxymesterone; methandrostenolol; MENT and prodrugs of these compounds. Most preferably the one or more androgens are selected from the group consisting of testosterone, DHEA and MENT.

[0062] Another aspect of the present invention relates to the use of the aforementioned solid dosage unit in medical treatment, in female hormone replacement therapy or in female contraception, said use comprising sublingual, buccal or sublabial administration of the dosage unit. Examples of medical treatment in which the solid dosage unit of the present invention may suitably be used include treatment of osteoporosis and estrogen add-back treatment in endometriosis, breast cancer or prostate cancer. In accordance with a preferred embodiment, the solid dosage unit is used in female hormone replacement therapy or female contraception. Most preferably, the solid dosage is used in female hormone replacement therapy, especially to treat vulvovaginal atrophy and/or vasomotor symptoms.

[0063] The use of the solid dosage unit in medical treatment, in female hormone replacement therapy or in female contraception, typically comprises sublingual, buccal or sublabial administration of the dosage unit to provide at least 0.1 mg, more preferably 0.5-100 mg and most preferably 1-40 mg of the estetrol component.

[0064] To treat vulvovaginal atrophy the dosage unit is preferably administered in an amount sufficient to provide at least 0.1 mg of the estetrol component. More preferably, the administered dosage unit provides at least 0.5 mg, most preferably at least 1 mg of the estetrol component. In the treatment of vulvovaginal atrophy the dosage unit is preferably administered in an amount that provides no more than 50 mg, more preferably not more than 20 mg and most preferably not more than 10 mg of the estetrol component.

[0065] To treat vasomotor symptoms the dosage unit is preferably administered in an amount sufficient to provide at least 0.2 mg of the estetrol component. More preferably, the administered dosage unit provides at least 1 mg, most preferably of at least 2 mg of the estetrol component. In the treatment of vasomotor symptoms the dosage unit is preferably administered in an amount that provides no more than 100 mg, more preferably not more than 40 mg and most preferably not more than 20 mg of the estetrol component.

[0066] Typically, these uses of the solid dosage unit comprise once daily administration of the dosage unit during a period of at least 1 week, more preferably of at least 2 weeks. During these periods the solid dosage unit is preferably administered to provide a daily dose of at least 0.05 mg, more preferably of 0.1-40 mg and most preferably of 0.2-20 mg of the estetrol component.

[0067] To treat vulvovaginal atrophy the dosage unit is preferably administered to provide a daily dose of at least 0.1 mg of the estetrol component. More preferably, the dosage unit is administered to provide a daily dose of 0.5-20 mg, most preferably of 1-10 mg of the estetrol

component.

[0068] To treat vasomotor symptoms the dosage unit is preferably administered to provide a daily dose of at least 0.2 mg of the estetrol component. More preferably, the dosage unit is administered to provide a daily dose of 1-40 mg, most preferably 2-20 mg of the estetrol component.

[0069] Yet another aspect of the invention relates to a process of preparing a solid dosage unit as described herein before, said process comprising the steps of:

- providing estetrol particles containing at least 80 wt.% of an estetrol component selected from estetrol, estetrol esters and combinations thereof and having a volume median diameter of 2 μm to 50 μm ;
- mixing the estetrol particles with one or more granulation excipients to produce a granulation mixture;
- mixing the granulation mixture with a granulation liquid to produce estetrol-containing granules, said granulation liquid containing at least 60 wt.% of liquid solvent;
- removing liquid solvent from the estetrol-containing granules to produce dry estetrol-containing granules;
- optionally mixing the dry granules with one or more tableting excipients; and
- forming the dry granules or the mixture of the dry granules and the one or more tableting excipients into a solid dosage unit.

[0070] In the present process the granulation mixture is preferably produced by combining the estetrol particles with the one or more granulation excipients in a weight ratio that is in the range of 1:2 to 1:1000, more preferably in the range of 1:3 to 1:100 and most preferably in the range of 1:4 to 1:10.

[0071] The one or more granulation excipients used in the preparation of the granulation mixture preferably include water-soluble carbohydrate as defined herein before. Preferably, said water-soluble carbohydrate constitutes at least 50 wt.%, more preferably at least 60 wt.% and most preferably at least 80 wt.% of the one or more granulation excipients.

[0072] The one or more granulation excipients of the granulation mixture preferably include 0.1-20 wt.%, more preferably 0.15-10 wt.% and most preferably 0.2-5.0 wt.% of binder as defined herein before; all percentage calculated by weight of the granulation mixture.

[0073] The one or more granulation excipients of the granulation mixture may suitably include a disintegrating agent as defined herein before. Preferably, the disintegrating agent constitutes 0.1-20 wt.%, more preferably 0.2-10 wt.% of the granulation mixture.

[0074] The estetrol-containing granules are preferably produced by mixing the granulation mixture with the granulation liquid in a weight ratio that is in the range of 0.5:1 to 20:1. Even

more preferably, the granulation mixture and the granulation liquid are mixed in a weight ratio that is in the range of 0.8:1 to 12:1, even more preferably in the range of 1:1 to 10:1 and most preferably of 1.5:1 to 5:1. The aforementioned weight ratio is calculated on the basis of the total amount of granulation mixture and granulation liquid that is used in the preparation of the dry estetrol-containing granules.

[0075] The estetrol-containing granules are preferably produced by mixing the granulation mixture with the granulation liquid in a high shear granulator, a low shear granulator or a fluidized bed granulator. Most preferably, the granules are prepared in a low shear granulator.

[0076] The granulation liquid employed in the present process preferably contains at least 60 wt.%, more preferably at least 80 wt.% and most preferably at least 90 wt.% of polar solvent selected from water, methanol, ethanol, iso-propanol, acetone and combinations thereof.

[0077] According to a particularly preferred embodiment, the granulation liquid contains at least 60 wt.%, more preferably at least 80 wt.% and most preferably at least 90 wt.% of polar solvents selected from water, ethanol and combinations thereof. The polar solvent employed in the granulation liquid preferably contains at least 80 wt.% water, more preferably at least 90 wt.% water.

[0078] The granulation liquid employed in the present process may suitably contain further excipients besides liquid solvent. Examples of such further excipients include binders, disintegrating agents, mucoadhesive agents, colouring, flavouring and combinations thereof.

[0079] Preferably, the granulation liquid contains 0.5-40 wt.%, more preferably 1-25 wt.% and most preferably 2-20 wt.% of binder, said binder being selected from cellulose derivatives, starch and starch derivatives (e.g. pregelatinized starch), polyvinyl alcohol (PVA), polyvinylpyrrolidone (PVP), agar, gelatin, guar gum, gum Arabic, alginate, polyethylene glycol (PEG), glucose, sucrose, sorbitol and combinations thereof. Preferably, the binder employed in the present process is selected from cellulose derivatives, pregelatinized starch, polyvinylpyrrolidone and combinations thereof.

[0080] Examples of cellulose derivatives that may be employed in the dosage unit as binder include hydroxypropyl cellulose, hydroxyethyl cellulose, hydroxymethyl cellulose, hydroxypropylmethylcellulose, methylcellulose, ethylcellulose, carboxymethyl cellulose and combinations thereof.

[0081] In the present process liquid solvent is preferably removed after the mixing of the granulation mixture with the granulation liquid has been completed. In a preferred embodiment reduced pressure is applied (e.g. less than 200 mbar) to facilitate the removal of liquid solvent during production of the estetrol-containing granules. Typically, the liquid solvent is removed at elevated temperature (e.g. >50 °C).

[0082] The dry estetrol-containing granules produced in the present process typically have a

volume median diameter in the range of 100-4,000 μm . More preferably, the volume median diameter of these granules is in the range of 200-1,000 μm , most preferably in the range of 200-600 μm .

[0083] Examples of the one or more tableting excipients that are optionally combined with the dry estetrol-containing granules before the forming of the solid dosage unit include lactose, mannitol, xylitol, microcrystalline cellulose, starch, croscarmellose sodium, polyvinyl pyrrolidone and combinations thereof.

[0084] Preferably, in the present process the dry estetrol-containing granules are mixed with the one or more tableting excipients in a weight ratio that is the range of 1:4 to 9:1, more preferably in the range of 1:2.5 to 3:1 and most preferably of 1:1.5 to 1.5 to 1.

[0085] In the present process the solid dosage unit may suitably be formed by direct compression or compression moulding. Most preferably, the solid dosage unit is formed by direct compression.

[0086] The solid dosage units obtained by the present method can be packaged in different ways. Preferably, the dosage units are packaged in a blister pack containing at least 14 dosage units.

[0087] The invention is further illustrated by means of the following non-limiting examples.

EXAMPLES

Dissolution test

[0088] The dissolution test described below can be used to study the dissolution behaviour of orodispersible dosage units.

Dissolution apparatus

[0089]

- Paddle and basket dissolution tester VanKel VK 7010 or VK 7025, autosampler VK 8000, 1000 mL dissolution vessels and porous micron filters (35 pin)

Dissolution Medium

[0090]

- Transfer 9,000 ml of demineralised water into a volumetric flask of 10,000 ml.
- Add 68.05 g of KH_2PO_4 and 8.96 g NaOH and stir the solution until everything is dissolved.
- Mix the solution and adjust the pH to 6.8 with NaOH or phosphoric acid, if necessary and make up to volume with demineralised water.

Dissolution Procedure**[0091]**

- Transfer 900 ml of Dissolution Medium into each vessel of the paddle apparatus.
- Assemble the apparatus, warm the medium to 37 ± 0.5 °C, and remove the thermometer.
- Place in each of the six vessels one tablet at the bottom before starting the rotation of the paddles.
- Start the rotation of the paddles immediately.
- Use a stirring speed of 50 rpm.
- Take samples of 5 ml from the dissolution vessels after 5, 10, 20, 30, 45, 60, 75 and 90 minutes for a complete dissolution profile. Take the sample from a position midway between the surface of the dissolution medium and the top of the paddle blade and not less than 10 mm from the vessel wall. The removed dissolution volume is not replaced by fresh dissolution medium.

[0092] Estetrol concentrations in the samples were determined by means of HPLC using estetrol stock solutions as a reference.

Preparation of mobile phase (MP) phosphate buffer**[0093]**

- Transfer 1.15 g of $\text{NH}_4\text{H}_2\text{PO}_4$ (10 mM) into a 1,000 ml of demineralised water, dissolve it and adjust the pH to 3.0 with phosphoric acid.

HPLC apparatus

[0094]

- Alliance 2695 Separations module consisting of a quaternary solvent delivery system, a variable volume injector, a temperature controlled autosampler, column thermostat and Photodiode array detector 2996 (all by Waters)
- Analytical column: Symmetry C18, 3.9 × 150 mm, dp = 5 µm (ex Waters)
- Guard column: Security guard columg C18, 4×3 mm (Phenomenex)
- Flow: 1.0 mL/min
- Detection: UV @ 280 nm
- Column temperature: 30°C
- Autosampler temperature: 10°C
- Injection volume: 100 µL
- Run time: 12 min

Elution gradient**[0095]**

Time (min)	Acetonitrile (%)	Phosphate buffer (%)
0	20	80
9	75	25
10	20	80
12	20	80

[0096] The dissolution tests are conducted in triplicate.

Particle size measurements

[0097] Particle size distribution of estetrol monohydrate is performed using a MALVERN MASTERSIZER MICROPLUS laser particle size analyzer.

Preparation of dispersion medium:**[0098]**

- Weigh 1g of estetrol monohydrate and 1g of sorbitan trioleate into a flask.
- Add 1 litre of n-hexane and mix for at least 1 hour at room temperature
- Filter through a 0.45 µm filter.

Sample preparation:

[0099]

- Put 100 mg of sample in a 25 mL beaker.
- Add some drops of dispersion medium.
- Mix carefully with a glass rod to suspend well the powder.
- Add 10 mL of dispersion medium.
- Perform the analysis with the sample dispersion unit's speed at 3000-3500 rpm.

Analysis:

[0100] Particle size measurements are performed three times using the same dispersion. The final result is obtained by averaging the results of the three determinations.

Example 1

[0101] A sublingual tablet is prepared by means of the procedure described below.

[0102] A granulation mixture having the composition shown in Table 1 is prepared by dry blending of the ingredients.

Table 1

Ingredients	Wt. %
Milled estetrol ¹	25
Mannitol	40
Lactose	35
¹ D(v;0.5) = appr. 15 µm	

[0103] A granulation liquid having the composition shown in Table 2 is prepared by dispersing the pregelatinized starch into the water

Table 2

Ingredients	Wt. %
Pregelatinized starch	2
Purified water	98

[0104] The granulation mixture is granulated in a high shear granulator by gradually adding the aqueous granulation liquid. The total amount of granulation liquid added amounts to 25% by weight of the granulation mixture.

[0105] The granulate so obtained is dried in a vacuum drying oven at 40°C and subsequently screened over a 500 µm sieve.

[0106] A tableting mixture having the composition shown in Table 3 is prepared by mixing the dry granulate with the excipients for 15 minutes (except magnesium stearate). Magnesium stearate is added and mixing is continued for an additional 5 minutes.

Table 3

Ingredients	Wt. %
Granulated estetrol	50
Lactose	10
Ludiflash [®] 2	38.5
Magnesium stearate	1.5
2 A mixture of mannitol (90 wt. %), crospovidone (5 wt. %) and polyvinyl acetate (5 wt. %)	

[0107] The tableting mixture is compressed into 80 mg round tablets with a diameter of 6.5 mm. The estetrol content of these tablets is 10 mg.

Example 2

[0108] A sublingual tablet is prepared by means of the procedure described below.

[0109] A granulation mixture having the composition shown in Table 4 is prepared by dry blending of the ingredients.

Table 4

Ingredients	Wt. %
Milled estetrol ¹	25
Lactose	75

¹ $D_{(v;0,5)} = \text{appr. } 15 \mu\text{m}$

[0110] A granulation liquid having the composition shown in Table 5 is prepared by dispersing the polyvinylpyrrolidone (PVP) into the water.

Table 5

Ingredients	Wt.%
PVP (polyvinylpyrrolidone)	3
Purified water	97

[0111] The granulation mixture is granulated in a low shear granulator by gradually adding the granulation liquid. The total amount of granulation liquid added amounts to 25% by weight of the granulation mixture.

[0112] The granulate so obtained is dried in the low shear granulator at 40°C and is subsequently screened over a 500 μm sieve.

[0113] A tableting mixture having the composition shown in Table 6 is prepared by mixing the dry granulate with the excipients for 15 minutes (except magnesium stearate). Magnesium stearate is added and mixing is continued for an additional 5 minutes.

Table 6

Ingredients	Wt.%
Granulated estetrol	50
Mannitol	43.5
Crospovidone	5
Magnesium stearate	1.5

[0114] The blend is compressed into 80 mg round tablets with diameter 6.5 mm. The estetrol content of these tablets is approximately 10 mg.

Example 3

[0115] A sublingual tablet is prepared using the procedure described below.

[0116] A granulation mixture having the composition shown in Table 7 is prepared by dry blending of the ingredients.

Table 7

Ingredients	Wt. %
Milled estetrol ¹	25
Lactose	75
¹ D _(v;0,5) = appr. 15 µm	

[0117] A granulation liquid having the composition shown in Table 8 is prepared by dispersing the hydroxypropyl cellulose (HPC) into the water.

Table 8

Ingredients	Wt. %
HPC (hydroxypropyl cellulose)	2
Purified water	97

[0118] The granulation mixture is granulated in a fluid bed granulator by gradually adding the granulation liquid. The total amount of granulation liquid added amounts to 35% by weight of the granulation mixture.

[0119] The granulate so obtained is dried in the fluid bed granulator at 50°C and subsequently screened over a 500 µm sieve.

[0120] A tableting mixture having the composition shown in Table 9 is prepared by mixing the dry granulate with the excipients in Table 9 for 15 minutes (except magnesium stearate). Magnesium stearate is added and mixing is continued for an additional 5 minutes.

Table 9

Ingredients	Wt. %
Granulated estetrol	50
Mannitol	43.5
Sodium starch glycolate	5
Magnesium stearate	1.5

[0121] The blend is compressed into 80 mg round tablets with diameter 6.5 mm. The estetrol content of these tablets is approximately 10 mg.

Example 4

[0122] Seven different sets of sublingual tablets (formulations A to G) were prepared by means of the procedure described below and illustrated in Figure 1.

[0123] The target amounts of estetrol per tablet were as follows: 100 µg for formulation A, 250 mg for formulation B, 7,5 mg for formulation C, 1 mg for formulation D, and 10 mg for formulations E, F and G.

[0124] The target weights for the tablets were as follows: 30 mg for formulations A and C, 1000 mg for formulations B and D, and 80 mg for formulations E, F and G.

[0125] The estetrol was mixed with a part of the main diluent and screened over a 800 µm screen. All other excipients were also screened over a 800 µm screen.

[0126] A binder solution of 10% PVP (polyvinylpyrrolidone) in demineralized water was prepared (the PVP concentration in the binder solution used for formulations B, C and F was 15%).

[0127] Granulation was started by adding the binder solution to the dry mixture consisting of the intragranular excipients (with a batchsize of ± 100 gram). A granulate was formed after 30 seconds of mixing, and subsequently pure demineralized water was added until a suitable granulate was formed (as detected visually, 1 minute of granulation in total).

[0128] The granulate was dried overnight in a vacuum oven at 40°C and subsequently milled over a 800 µm screen. The mixture thus obtained was mixed for 15 minutes with the extragranular excipients (except for magnesium stearate). Finally, magnesium stearate was added and mixed for 3 minutes.

[0129] Compression was executed using a single punch machine equipped with proper punches (5 mm punch for 30 mg tablets (A and C), 6 mm for 80 mg tablets (E, F and G) and 15 mm for 1000 mg tablets (B and D)).

[0130] Disintegration time was quantified according to the known protocol described in Ph. Eur. 2.9.1 ("Disintegration of tablets and capsules"), and in USP <701> ("Disintegration") using water as the specified liquid.

[0131] Hardness was measured using the known protocol described in Ph. Eur. 2.9.8 ("Resistance to crushing of tablets").

[0132] The final formulations and corresponding tablet results can be found in Tables 10 and 11 below.

[0133] No problems were detected during the experiments, flowability of the blend was good and no sticking was observed.

Table 10 — details of the formulations in Wt. %

Formulation #	A	B	C	D	E	F	G
<i>Intragranular</i>							
Milled Estetrol ¹	0.33	24.92	25.00	0.10	12.49	12.48	12.49
Mannitol	91.17	65.89	65.49	91.90	79.25	18.99	
PVP (polyvinylpyrrolidone)	2.00	2.99	3.00	2.00	2.01		1.46
Pregelatinized starch						1.01	
Crospovidone	4.01	4.00	4.00	4.00	4.00		
Lactose						17.51	36.04
Demineralized water (per 100 gram granulation mixture)	22	26	26	23	19	16	24
<i>Extragranular</i>							
Crospovidone	0.99	1.10	1.00	1.01	0.99		5.00
Lactose						10.01	
Ludiflash ^{®2}						38.49	
Mannitol							43.49
Magnesium stearate	1.49	1.10	1.50	0.99	1.25	1.50	1.52
¹ D _(v;0.5) = 15µm ² Containing mannitol (90 wt.%), crospovidone (5 wt.%) and polyvinyl acetate (5 wt.%)							

Table 11 — experimentally determined characteristics of the Tablets

Test (average result of 6 samples)	Disintegration time	Hardness	Weight
Formulation #	(min:sec)	(N)	(mg)
A	0:39	14.17	28.2
B	1:44	85.85	1058.6
C	1:05	28.46	30.1
D	0:40	66.74	1048.9
E	0:32	29.70	78.9
F	0:41	30.97	82.2
G	0:33	22.69	78.1

[0134] It can be seen that all tablets were obtained with a final weight close to their target weight and that the disintegration times, even for the largest 1g tablets, were very short, in accordance with the intended sublingual, buccal or sublabial administration route for these tablets.

[0135] Finally, the hardness of all tablets was within a very acceptable range.

Example 5

[0136] A randomized, open-label, two-period, cross-over, pharmacokinetic study is conducted to compare sublingual bioavailability of 10 mg estetrol administered in one 80 mg tablet (having the same composition as the tablets described in Example 4, Formulation E) with oral availability of estetrol contained in a 83 mg tablet containing 10 mg estetrol. These tablets are administered sublingually and orally to healthy female volunteers under fasting conditions.

[0137] Ten healthy female subjects are selected on the basis of the following criteria: age of 45-65 years (inclusive), nonsmokers or past smokers (at least 6 months before dosing), body-mass index (BMI) = 18.5 to 30 kg/m² (inclusive at the time of the screening).

[0138] At the start of the first and the second period of the study, between 07:00 am and 07:28 am, 5 subjects receive a single dose of the sublingual formulation of estetrol by administering one estetrol tablet (tablet weight 80 mg; 10 mg estetrol) and 5 subjects receive a single oral dose of the oral estetrol formulation by administering one estetrol tablet (tablet weight 83 mg; 10 mg estetrol), ingested together with 200 ml water.

[0139] Subjects are required to fast for at least 10 hours prior to tablet administration and for at least 4 hours after administration. Drinking of water or beverages is not permitted within 1 hour before the drug administration. Subjects receive 200 ml of water 1 hour prior to and 2 hours after tablet administration. Subjects are free to drink water and fruit tea from 4 hours following the tablet administration. Standardized meals are provided 10.5 hours before and 4, 6, 9, and 13 hours after tablet administration.

[0140] The sequence of events that occurs during the first and second period is shown in Table 12:

Table 12

	Event
First period	
• Day 1	Confinement from 19:00
• Day 2	Dosing, blood and urine sampling, confinement
• Day 3	Exit procedure, confinement till 8 am
• Days 4-8	Return visits
• Days 9-13	Wash out

Second period	
• Day 14	Confinement from 19:00
• Day 15	Dosing, blood and urine sampling, confinement
• Day 16	Exit procedure, confinement till 8 am
• Days 17-21	Return visits
• Days 22-26	Wash out
• Day 27	Administration of a progestin
• Day 28	Phone call, progestin withdrawal test check

[0141] The blood and urine sampling schedule used in this study is shown in Table 13.

Table 13

Blood sampling	Blood collection (4 ml) is performed prior to administration of the tablet (0), and subsequently 0:10, 0:15, 0:20, 0:25, 0:30, 0:35, 0:40, 0:45, 0:50, 0:55, 1:00, 1:10, 1:20, 1:30, 2, 3, 4, 6, 10, 16, 24, 48, 72, 96, 120, 144 hours after administration.
	Total number of blood collections in each period is 27.
Urine sampling	Urine collection is performed prior to administration of the tablet and 2, 4, 8, 12, 24, 48, 72, 96, 120 and 144 hours after administration.
	Total number of urine collections in each period is 11.

[0142] The estetrol concentration in the collected blood samples is determined by means of HPLC/MS/MS. The concentrations of glucuronided estetrol (D-ring) in the urine samples is also determined with the help of HPLC/MS/MS.

[0143] Results of these analyses show that the bioavailability of sublingually administered estetrol is comparable or even superior to orally administered estetrol. Furthermore, the data suggest that sublingually administered estetrol has an earlier bioavailability compared to orally administered estetrol. Sublingual estetrol has less impact on a liver function parameter.

PREFERRED ASPECTS AND EMBODIMENTS

Aspect 1

[0144] Embodiment 1 of aspect 1 relates to an orodispersible solid pharmaceutical dosage unit having a weight between 30 and 1,000 mg, said dosage unit consisting of:

- 0.1-25 wt.% of estetrol particles containing at least 80 wt.% of an estetrol component selected from estetrol, estetrol esters and combinations thereof; and
- 75-99.9 wt.% of one or more pharmaceutically acceptable ingredients;

the solid dosage unit comprising at least 100 µg of the estetrol component; wherein the solid dosage unit can be obtained by a process comprising:

- providing estetrol particles containing at least 80 wt.% of the estetrol component and having a volume median diameter of 2 µm to 50 µm;
- mixing the estetrol particles with one or more granulation excipients to produce a granulation mixture;
- mixing the granulation mixture with a granulation liquid to produce estetrol-containing granules, said granulation liquid containing at least 60 wt.% of liquid solvent;
- removing liquid solvent from the estetrol-containing granules to produce dry estetrol-containing granules;
- optionally mixing the dry granules with one or more tableting excipients; and
- forming the dry granules or the mixture of the dry granules and the one or more tableting excipients into a solid dosage unit.

[0145] Embodiment 2 of aspect 1 is according to embodiment 1 of aspect 1 and is such that the dosage unit has a weight between 40 and 500 mg.

[0146] Embodiment 3 of aspect 1 is according to embodiment 1 or 2 of aspect 1 and is such that the dosage unit contains 0.5-25 wt.% of the estetrol component.

[0147] Embodiment 4 of aspect 1 is according to embodiment 1 or 2 or 3 of aspect 1 and is such that the dosage unit contains 0.3-100 mg of the estetrol component.

[0148] Embodiment 5 of aspect 1 is according to embodiment 1 or 2 or 3 or 4 of aspect 1 and is such that the estetrol component is estetrol.

[0149] Embodiment 6 of aspect 1 is according to embodiment 1 or 2 or 3 or 4 or 5 of aspect 1 and is such that the estetrol particles have a volume median diameter of 3-35 µm.

[0150] Embodiment 7 of aspect 1 is according to embodiment 1 or 2 or 3 or 4 or 5 or 6 of aspect 1 and is such that the dosage unit contains 50-99.9 wt.% of water-soluble carbohydrate selected from maltose, fructose, sucrose, lactose, glucose, galactose, trehalose, xylitol, sorbitol, erythritol, maltitol, mannitol, isomalt and combinations thereof.

[0151] Embodiment 8 of aspect 1 is according to embodiment 7 of aspect 1 and is such that the dosage unit contains at least 20 wt.% mannitol.

[0152] Embodiment 9 of aspect 1 is according to embodiment 1 or 2 or 3 or 4 or 5 or 6 or 7 or

8 of aspect 1 and is such that the dosage unit contains 0.1-20 wt.% of a disintegrating agent selected from modified starches, crosslinked polyvinylpyrrolidone, crosslinked carmellose and combinations thereof.

Aspect 2

[0153] Embodiment 1 of aspect 2 relates to a solid dosage unit according to any embodiment of aspect 1 for use in female hormone replacement therapy, said use comprising sublingual, buccal or sublabial administration of the dosage unit.

[0154] Embodiment 2 of aspect 2 is according to embodiment 1 of aspect 2 and is such that said use comprises once daily administration during a period of at least 1 week.

Aspect 3

[0155] Embodiment 1 of aspect 3 relates to a method of female contraception, said method comprising sublingual, buccal or sublabial administration of a dosage unit according to any embodiment of aspect 1.

[0156] Embodiment 2 of aspect 3 is according to embodiment 1 of aspect 3 and is such that said method comprises once daily administration during a period of at least 1 week.

Aspect 4

[0157] Embodiment 1 of aspect 4 relates to a process of preparing a solid dosage unit according to any embodiment of aspect 1 or aspect 2, said process comprising the steps of:

- providing estetrol particles containing at least 80 wt.% of an estetrol component selected from estetrol, estetrol esters and combinations thereof and having a volume median diameter of 2 μm to 50 μm ;
- mixing the estetrol particles with one or more granulation excipients to produce a granulation mixture;
- mixing the granulation mixture with a granulation liquid to produce estetrol-containing granules, said granulation liquid containing at least 60 wt.% of liquid solvent;
- removing liquid solvent from the estetrol-containing granules to produce dry estetrol-containing granules;
- optionally mixing the dry granules with one or more tableting excipients; and
- forming the dry granules or the mixture of the dry granules and the one or more tableting excipients into a solid dosage unit.

[0158] Embodiment 2 of aspect 4 is according to embodiment 1 of aspect 4 and is such that the estetrol particles have a volume median diameter of 3-35 μm .

[0159] Embodiment 3 of aspect 4 is according to embodiment 1 or 2 of aspect 4 and is such that the granulation mixture contains 0.1-20 wt.% of a disintegrating agent selected from modified starches, crosslinked PVP, crosslinked carmellose and combinations thereof.

[0160] Embodiment 4 of aspect 4 is according to embodiment 1 or 2 or 3 of aspect 4 and is such that the granulation mixture is produced by combining the estetrol particles with the one or more granulation excipients in a weight ratio that is in the range of 1:4 to 1:1,000.

[0161] Embodiment 5 of aspect 4 is according to embodiment 1 or 2 or 3 or 4 of aspect 4 and is such that the estetrol-containing granules are produced by mixing the granulation mixture with the granulation liquid in a weight ratio that is in the range of 0.5:1 to 20:1.

[0162] Embodiment 6 of aspect 4 is according to embodiment 1 or 2 or 3 or 4 or 5 of aspect 4 and is such that the estetrol-containing granules are produced by mixing the granulation mixture with the granulation liquid in a high shear granulator, a low shear granulator or a fluidized bed granulator.

[0163] Embodiment 7 of aspect 4 is according to embodiment 1 or 2 or 3 or 4 or 5 or 6 of aspect 4 and is such that the granulation liquid contains at least 60 wt.% of polar solvent selected from water, methanol, ethanol, iso-propanol, acetone and combinations thereof.

[0164] Embodiment 8 of aspect 4 is according to embodiment 7 of aspect 4 and is such that the granulation liquid contains at least 60 wt.% of polar solvents selected from water, ethanol and combinations thereof.

[0165] Embodiment 9 of aspect 4 is according to embodiment 7 or 8 of aspect 4 and is such that the polar solvent contains at least 80 wt.% water.

[0166] Embodiment 10 of aspect 4 is according to embodiment 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 of aspect 4 and is such that the granulation liquid contains 0.2-50 wt.% of binder, said binder being selected from cellulose derivatives, starch and starch derivatives, polyvinyl alcohol, polyvinylpyrrolidone, agar, gelatin, guar gum, gum Arabic, alginate, polyethylene glycol, glucose, sucrose, sorbitol and combinations thereof.

[0167] Embodiment 11 of aspect 4 is according to embodiment 10 of aspect 4 and is such that the binder is selected from cellulose derivatives, pregelatinized starch, polyvinylpyrrolidone and combinations thereof.

[0168] Embodiment 12 of aspect 4 is according to embodiment 10 or 11 of aspect 4 and is such that the cellulose derivatives are selected from hydroxypropyl cellulose, hydroxyethyl

cellulose, hydroxymethyl cellulose, hydroxypropylmethylcellulose, methylcellulose, ethylcellulose, carboxymethyl cellulose and combinations thereof.

[0169] Embodiment 13 of aspect 4 is according to embodiment 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 of aspect 4 and is such that the dry estetrol-containing granules have a volume median diameter in the range of 100-4,000 μm .

[0170] Embodiment 14 of aspect 4 is according to embodiment 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 of aspect 4 and is such that the one or more tableting excipients include lactose, mannitol, xylitol, microcrystalline cellulose, starch, croscarmellose sodium, polyvinyl pyrrolidone and combinations thereof.

[0171] Embodiment 15 of aspect 4 is according to embodiment 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 14 of aspect 4 and is such that the solid dosage unit is formed by direct compression or compression moulding.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2002094275A](#) [\[0008\]](#) [\[0014\]](#)
- [WO2002094276A](#) [\[0008\]](#) [\[0013\]](#) [\[0013\]](#) [\[0014\]](#)
- [WO2002094278A](#) [\[0008\]](#)
- [WO2003018026A](#) [\[0008\]](#)
- [WO2010033832A](#) [\[0009\]](#)
- [US2007286829A](#) [\[0010\]](#)
- [US6117446A](#) [\[0011\]](#)
- [WO2002094279A](#) [\[0015\]](#) [\[0016\]](#)
- [WO2003041718A](#) [\[0016\]](#)
- [WO2007081206A](#) [\[0017\]](#)
- [WO2008156365A](#) [\[0018\]](#)

Patentkrav

1. Oro-dispergerbar fast farmaceutisk doseringsenhed med en vægt mellem 30 og 1000 mg, fortrinsvis mellem 40 og 500 mg, hvor nævnte doseringsenhed består af:

- 5 • 0,1-25 vægtprocent, fortrinsvis 0,5-25 vægtprocent af estetrolpartikler indeholdende mindst 90 vægtprocent af en estetrolkomponent valgt fra estetrol, estetrolestere og kombinationer deraf, fortrinsvis er estetrolkomponenten estetrol; og
- 10 • 75-99,9 vægtprocent af en eller flere farmaceutisk acceptable ingredienser;
- den faste doseringsenhed omfattende mindst 100 µg af estetrolkomponenten; hvor den faste doseringsenhed kan opnås ved en proces omfattende:
- 15 • at tilvejebringe estetrolpartikler indeholdende mindst 90 vægtprocent af estetrolkomponenten og med en volumen-mediandiameter på 2 µm til 50 µm, fortrinsvis på 3 µm til 35 µm;
- 20 • at blande estetrolpartiklerne med et eller flere granulationshjælpestoffer for at fremstille en granulationsblanding;
- at blande granulationsblandingen med en granuleringsvæske for at fremstille estetrol-indeholdende granulater, nævnte granuleringsvæske indeholdende mindst 60 vægtprocent flydende solvent;
- 25 • at fjerne flydende solvent fra de estetrol-indeholdende granulater for at fremstille tørre estetrol-indeholdende granulater;
- eventuelt at blande de tørre granulater med et eller flere tableteringshjælpestoffer; og
- at forme de tørre granulater eller blandingen af de tørre granulater og det ene eller flere tableteringshjælpestoffer til en fast doseringsenhed, hvor doseringsenheden indeholder 0,3-100 mg af estetrol-komponenten.

- 2.** Fast doseringsenhed ifølge krav 1, hvor doseringsenheden indeholder 50-99,9 vægtprocent af vandopløseligt kulhydrat valgt fra maltose, fruktose, sukrose, 30 laktose, glukose, galaktose, trehalose, xylitol, sorbitol, erythritol, maltitol, mannitol, isomalt og kombinationer deraf.

- 3.** Fast doseringsenhed ifølge krav 2, hvor doseringsenheden indeholder mindst 20 vægtprocent mannitol.
- 4.** Fast doseringsenhed ifølge et hvilket som helst af de foregående krav, hvor doseringsenheden indeholder 0,1-20 vægtprocent af et desintegrerende middel valgt fra modificerede stivelser, tværbundet polyvinylpyrrolidon, tværbundet carmellose og kombinationer deraf.
- 5.** Fast doseringsenhed ifølge et hvilket som helst af de foregående krav, hvor nævnte farmaceutisk acceptable ingrediens omfatter en progestogen valgt fra gruppen bestående af: progesteron, desogestrel, etonogestrel, gestoden, dienogest, levonorgestrel, norgestimat, norethisteron, norethisteron-acetat (NETA), nomegestrol, nomegestrol-acetat (NOMAC), drospirenon, trimegeston, nestoron og dydrogesteron.
- 6.** Fast doseringsenhed ifølge krav 5, hvor nævnte progestogen er 0,05-10 mg af drospirenon.
- 7.** Fast doseringsenhed ifølge et hvilket som helst af de foregående krav til anvendelse i medicinsk behandling eller til anvendelse i hormonsubstitutionsterapi hos kvinder, hvor nævnte anvendelse omfatter sublingual, bukkal eller sublabial administration af doseringsenheden, fortrinsvis hvor nævnte anvendelse omfatter en daglig sublingual, bukkal eller sublabial administration af doseringsenheden under en periode på mindst 1 uge.
- 8.** Fremgangsmåde til antikonception hos kvinder, hvor nævnte fremgangsmåde omfatter sublingual, bukkal eller sublabial administration af en doseringsenhed ifølge et hvilket som helst af kravene 1-6, fortrinsvis hvor nævnte fremgangsmåde omfatter en daglig sublingual, bukkal eller sublabial administration af doseringsenheden under en periode på mindst 1 uge.
- 9.** Proces til fremstilling af en fast doseringsenhed ifølge kravene 1 til 6, hvor nævnte proces omfatter trinnene:
- at tilvejebringe estetrolpartikler indeholdende mindst 90 vægtprocent af en estetrolkomponent valgt fra estetrol, estetrolestere og kombinationer

deraf og med en volumen-mediandiameter på 2 µm til 50 µm, fortrinsvis på 3 µm til 35 µm;

- at blande estetrolpartiklerne med et eller flere granulationshjælpstoffer for at fremstille en granulationsblanding;
- 5 • at blande granulationsblandingen med en granuleringsvæske for at fremstille estetrol-indeholdende granulater, fortrinsvis i et vægtforhold som er i området fra 0,5:1 til 20:1, hvor nævnte granuleringsvæske indeholder mindst 60 vægtprocent af flydende solvent;
- at fjerne flydende solvent fra de estetrol-indeholdende granulater for at fremstille tørre estetrol-indeholdende granulater;
- 10 • eventuelt at blande de tørre granulater med et eller flere tableteringshjælpstoffer, fortrinsvis er nævnte hjælpestoffer valgt fra laktose, mannitol, xylitol, mikrokrySTALLinsk cellulose, stivelse, tvær-carmellosenatrium, polyvinylpyrrolidon og kombinationer deraf; og
- 15 • at danne de tørre granulater eller blandingen af de tørre granulater og det ene eller flere tableteringshjælpstoffer til en fast doseringsenhed, fortrinsvis ved direkte kompression eller formpresning.

10. Proces ifølge krav 9, hvor granulationsblandingen indeholder 0,1-20 vægtprocent af et desintegrerende middel valgt fra modificerede stivelser, tværbundet PVP, tværbundet carmellose og kombinationer deraf.

11. Proces ifølge et hvilket som helst af kravene 9 eller 10, hvor granulationsblandingen fremstilles ved at kombinere estetrolpartiklerne med det ene eller flere granulationshjælpstoffer i et vægtforhold, som er i området fra 1:4 til 1:1000.

12. Proces ifølge et hvilket som helst af kravene 9-11, hvor de estetrol-indeholdende granulater fremstilles ved at blande granulationsblandingen med granuleringsvæsken i en *high-shear*-granulator, en *low-shear*-granulator eller en fluidiseret leje-granulator.

13. Proces ifølge et hvilket som helst af kravene 9-12, hvor granuleringsvæsken indeholder mindst 60 vægtprocent af polært solvent valgt fra vand, methanol, ethanol, iso-propanol, acetone og kombinationer deraf, fortrinsvis valgt fra vand,

ethanol og kombinationer deraf.

14. Proces ifølge krav 13, hvor det polære solvent indeholder mindst 80 vægtprocent vand.

5

15. Proces ifølge et hvilket som helst af kravene 9-14, hvor granuleringsvæsken indeholder 0,2-50 vægtprocent af bindemiddel, hvor nævnte bindemiddel vælges fra cellulosederivater, stivelse og stivelsesderivater, polyvinylalkohol, polyvinylpyrrolidon, agar, gelatine, guar gummi, gummi arabikum, alginat, 10 polyethylenglykol, glukose, sukrose, sorbitol og kombinationer deraf, fortrinsvis valgt fra cellulosederivater, prægelatineret stivelse, polyvinylpyrrolidon og kombinationer deraf.

16. Proces ifølge krav 15, hvor cellulosederivaterne er valgt fra hydroxypropyl- 15 cellulose, hydroxyethylcellulose, hydroxymethylcellulose, hydroxypropylmethylcellulose, methylcellulose, ethylcellulose, carboxymethylcellulose og kombinationer deraf.

17. Proces ifølge et hvilket som helst af kravene 9-16, hvor de tørre estetrol- 20 indeholdende granulater har en volumen-mediandiameter i området fra 100-4000 µm.

DRAWINGS

