

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 October 2010 (07.10.2010)

PCT

(10) International Publication Number
WO 2010/114780 A1

(51) International Patent Classification:
A61K 31/44 (2006.01)

[CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN).

(21) International Application Number:
PCT/US2010/028989

(74) Common Representative: **MERCK SHARP & DOHME CORP.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(22) International Filing Date:
29 March 2010 (29.03.2010)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/165,643 1 April 2009 (01.04.2009) US

(71) Applicant (for all designated States except US): **MERCK SHARP & DOHME CORP.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHEN, Qi** [CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **CUI, Mingzhang** [CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **HU, Bin** [CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **HU, Shuangxi** [CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **PU, Zhifa** [CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **SANDERSON, Philip, E.** [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **SOLL, Richard** [US/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **TANG, Xue** [CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **WANG, Jiabing** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **YAN, Yongjin** [CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **YAO, Jinfeng** [CN/CN]; 288 Fute Zhong Road, Waigaoqiao Free Trade Zone, Shanghai 200131 (CN). **ZHANG, Hongjie**

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: INHIBITORS OF AKT ACTIVITY

(57) Abstract: The instant invention provides for substituted fused pyrimidine compounds that inhibit Akt activity. In particular, the compounds disclosed selectively inhibit one or two of the Akt isoforms. The invention also provides for compositions comprising such inhibitory compounds and methods of inhibiting Akt activity by administering the compound to a patient in need of treatment of cancer.



WO 2010/114780 A1

TITLE OF THE INVENTION

INHIBITORS OF AKT ACTIVITY

BACKGROUND OF THE INVENTION

5 The present invention relates to substituted fused pyrimidine compounds which are inhibitors of the activity of one or more of the isoforms of the serine/threonine kinase, Akt (also known as PKB; hereinafter referred to as "Akt"). The present invention also relates to pharmaceutical compositions comprising such compounds and methods of using the instant compounds in the treatment of cancer.

10 Apoptosis (programmed cell death) plays essential roles in embryonic development and pathogenesis of various diseases, such as degenerative neuronal diseases, cardiovascular diseases and cancer. Recent work has led to the identification of various pro- and anti-apoptotic gene products that are involved in the regulation or execution of programmed cell death. Expression of anti-apoptotic genes, such as Bcl2 or Bcl-x_L, inhibits apoptotic cell death
15 induced by various stimuli. On the other hand, expression of pro-apoptotic genes, such as Bax or Bad, leads to programmed cell death (Adams et al. *Science*, 281:1322-1326 (1998)). The execution of programmed cell death is mediated by caspase-1 related proteinases, including caspase-3, caspase-7, caspase-8 and caspase-9 etc (Thornberry et al. *Science*, 281:1312-1316 (1998)).

20 The phosphatidylinositol 3'-OH kinase (PI3K)/Akt pathway appears important for regulating cell survival/cell death (Kulik et al. *Mol. Cell. Biol.* 17:1595-1606 (1997); Franke et al, *Cell*, 88:435-437 (1997); Kauffmann-Zeh et al. *Nature* 385:544-548 (1997) Hemmings
25 *Science*, 275:628-630 (1997); Dudek et al., *Science*, 275:661-665 (1997)). Survival factors, such as platelet derived growth factor (PDGF), nerve growth factor (NGF) and insulin-like growth factor-1 (IGF-1), promote cell survival under various conditions by inducing the activity of PI3K (Kulik et al. 1997, Hemmings 1997). Activated PI3K leads to the production of
30 phosphatidylinositol (3,4,5)-triphosphate (PtdIns(3,4,5)-P₃), which in turn binds to, and promotes the activation of, the serine/threonine kinase Akt, which contains a pleckstrin homology (PH)-domain (Franke et al *Cell*, 81:727-736 (1995); Hemmings *Science*, 277:534 (1997); Downward, *Curr. Opin. Cell Biol.* 10:262-267 (1998), Alessi et al., *EMBO J.* 15: 6541-6551 (1996)). Specific inhibitors of PI3K or dominant negative Akt mutants abolish
35 survival-promoting activities of these growth factors or cytokines. It has been previously disclosed that inhibitors of PI3K (LY294002 or wortmannin) blocked the activation of Akt by upstream kinases. In addition, introduction of constitutively active PI3K or Akt mutants promotes cell survival under conditions in which cells normally undergo apoptotic cell death (Kulik et al. 1997, Dudek et al. 1997).

 Three members of the Akt subfamily of second-messenger regulated serine/threonine protein kinases have been identified and termed Akt1/ PKB α , Akt2/PKB β , and

Akt3/PKBy (hereinafter referred to as "Akt1", "Akt2" and "Akt3"), respectively. The isoforms are homologous, particularly in regions encoding the catalytic domains. Akts are activated by phosphorylation events occurring in response to PI3K signaling. PI3K phosphorylates membrane inositol phospholipids, generating the second messengers phosphatidyl-inositol 3,4,5-trisphosphate and phosphatidylinositol 3,4-bisphosphate, which have been shown to bind to the PH domain of Akt. The current model of Akt activation proposes recruitment of the enzyme to the membrane by 3'-phosphorylated phosphoinositides, where phosphorylation of the regulatory sites of Akt by the upstream kinases occurs (B.A. Hemmings, *Science* 275:628-630 (1997); B.A. Hemmings, *Science* 276:534 (1997); J. Downward, *Science* 279:673-674 (1998)).

Phosphorylation of Akt1 occurs on two regulatory sites, Thr³⁰⁸ in the catalytic domain activation loop and on Ser⁴⁷³ near the carboxy terminus (D. R. Alessi *et al.* *EMBO J.* 15:6541-6551 (1996) and R. Meier *et al.* *J. Biol. Chem.* 272:30491-30497 (1997)). Equivalent regulatory phosphorylation sites occur in Akt2 and Akt3. The upstream kinase, which phosphorylates Akt at the activation loop site has been cloned and termed 3'-phosphoinositide - dependent protein kinase 1 (PDK1). PDK1 phosphorylates not only Akt, but also p70 ribosomal S6 kinase, p90RSK, serum and glucocorticoid-regulated kinase (SGK), and protein kinase C. The upstream kinase phosphorylating the regulatory site of Akt near the carboxy terminus has not been identified yet, but recent reports imply a role for the integrin-linked kinase (ILK-1), a serine/threonine protein kinase, or autophosphorylation.

Analysis of Akt levels in human tumors showed that Akt2 is overexpressed in a significant number of ovarian (J. Q. Cheng *et al.* *Proc. Natl. Acad. Sci. U.S.A.* 89:9267-9271(1992)) and pancreatic cancers (J. Q. Cheng *et al.* *Proc. Natl. Acad. Sci. U.S.A.* 93:3636-3641 (1996)). Similarly, Akt3 was found to be overexpressed in breast and prostate cancer cell lines (Nakatani *et al.* *J. Biol. Chem.* 274:21528-21532 (1999)).

The tumor suppressor PTEN, a protein and lipid phosphatase that specifically removes the 3' phosphate of PtdIns(3,4,5)-P3, is a negative regulator of the PI3K/Akt pathway (Li *et al.* *Science* 275:1943-1947 (1997), Stambolic *et al.* *Cell* 95:29-39 (1998), Sun *et al.* *Proc. Natl. Acad. Sci. U.S.A.* 96:6199-6204 (1999)). Germline mutations of PTEN are responsible for human cancer syndromes such as Cowden disease (Liaw *et al.* *Nature Genetics* 16:64-67 (1997)). PTEN is deleted in a large percentage of human tumors and tumor cell lines without functional PTEN show elevated levels of activated Akt (Li *et al.* *supra*, Guldberg *et al.* *Cancer Research* 57:3660-3663 (1997), Risinger *et al.* *Cancer Research* 57:4736-4738 (1997)).

These observations demonstrate that the PI3K/Akt pathway plays important roles for regulating cell survival or apoptosis in tumorigenesis.

Inhibition of Akt activation and activity can be achieved by inhibiting PI3K with inhibitors such as LY294002 and wortmannin. However, PI3K inhibition has the potential to indiscriminately affect not just all three Akt isozymes but also other PH domain-containing signaling molecules that are dependent on PtdIns(3,4,5)-P3, such as the Tec family of tyrosine

kinases. Furthermore, it has been disclosed that Akt can be activated by growth signals that are independent of PI3K.

Alternatively, Akt activity can be inhibited by blocking the activity of the upstream kinase PDK1. No specific PDK1 inhibitors have been disclosed. Again, inhibition of PDK1 would result in inhibition of multiple protein kinases whose activities depend on PDK1, such as atypical PKC isoforms, SGK, and S6 kinases (Williams et al. *Curr. Biol.* 10:439-448 (2000)).

Inhibitors of Akt are known. WO2005/100344; WO2005/100356; WO2004/096135; WO2004/096129; WO2004/096130; WO2004/096131; WO2006/091395; WO2008/070134; WO2009/148916; WO2008/070016; WO2008/070041; WO2004/041162; WO2009/148887; WO2006/068796; WO2006/065601; WO2006/110638; WO2003/086394; WO2003/086403; WO2003/086404; WO2003/086279; WO2002/083139; WO2002/083675; WO2006/036395; WO2002/083138; WO2006/135627; and WO2002/083140. The compounds disclosed in these patent applications contain mono-, bi- and tri-cyclic core moieties.

Specific Akt inhibitors substituted with a methyl amine moiety are known. WO2006/135627; WO2008/070041; WO2008/070016; WO2008/070134; WO2009/148887; and WO2009/148916.

The compounds of the instant invention contain a bi-cyclic core moiety with internal nitrogen atoms where the rings fuse which have not been previously disclosed.

It is an object of the instant invention to provide novel compounds that are inhibitors of Akt.

It is also an object of the present invention to provide pharmaceutical compositions that comprise the novel compounds that are inhibitors of Akt.


It is also an object of the present invention to provide a method for treating cancer that comprises administering such inhibitors of Akt activity.

SUMMARY OF THE INVENTION

The instant invention provides for substituted fused pyrimidine compounds that inhibit Akt activity. In particular, the compounds disclosed selectively inhibit one or two of the Akt isoforms. The invention also provides for compositions comprising such inhibitory compounds and methods of inhibiting Akt activity by administering the compound to a patient in need of treatment of cancer.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of the instant invention are useful in the inhibition of the activity of the serine/threonine kinase Akt. In a first embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula A:



wherein:

A and B are independently selected from C or N, wherein at least one of A and B is N;

J and K are independently selected from C or N, wherein J and K are optionally substituted with R¹;

X, Y and Z are independently selected from C and N, wherein X, Y and Z are optionally substituted with R¹;

a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; p is independently 0, 1, 2, 3, 4 or 5;

Ring Q is selected from: (C₃-C₈)cycloalkyl, aryl, heteroaryl and heterocyclyl;

R¹ is independently selected from: H, oxo, (C=O)_aO_b(C₁-C₁₀)alkyl, (C=O)_aO_b-aryl, (C=O)_aO_b(C₂-C₁₀)alkenyl, (C=O)_aO_b(C₂-C₁₀)alkynyl, CO₂H, halo, OH, O_b(C₁-C₆)perfluoroalkyl, (C=O)_aNR⁷R⁸, CN, (C=O)_aO_b(C₃-C₈)cycloalkyl, S(O)_mNR⁷R⁸, SH, S(O)_m-(C₁-C₁₀)alkyl and (C=O)_aO_b-heterocyclyl, said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, and heterocyclyl is optionally substituted with one or more substituents selected from R⁶;

R² is independently selected from: OH, halo and (C₁-C₄)alkyl, said alkyl is optionally substituted with one or more substituents selected from OH and halo;

R³ and R^{3'} are independently selected from: H and (C₁-C₄)alkyl, said alkyl is optionally substituted with one or more substituents selected from OH, or R³ and R^{3'} can be taken together to form a (C₃-C₆)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkoxy, CO₂H, halo, CN, OH and NH₂;

R⁶ is: (C=O)_aO_bC₁-C₁₀ alkyl, (C=O)_aO_baryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aO_b heterocyclyl, CO₂H, halo, CN, OH, O_bC₁-C₆ perfluoroalkyl, O_a(C=O)_bNR⁷R⁸, oxo, CHO, (N=O)R⁷R⁸, S(O)_mNR⁷R⁸, SH, S(O)_m-(C₁-C₁₀)alkyl or (C=O)_aO_bC₃-C₈ cycloalkyl, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R^{6a};

R^{6a} is selected from: (C=O)_aO_b(C₁-C₁₀)alkyl, O_a(C₁-C₃)perfluoroalkyl, (C₀-C₆)alkylene-S(O)_mR^a, SH, oxo, OH, halo, CN, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, (C₃-C₆)cycloalkyl, (C₀-C₆)alkylene-aryl, (C₀-C₆)alkylene-heterocyclyl, (C₀-C₆)alkylene-N(R^b)₂, C(O)R^a, (C₀-C₆)alkylene-CO₂R^a, C(O)H, and (C₀-C₆)alkylene-CO₂H, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally substituted with up to three substituents

selected from R^b , OH, (C_1-C_6) alkoxy, halogen, CO_2H , CN, $O_a(C=O)_b(C_1-C_6)$ alkyl, oxo, and $N(R^b)_2$;


R^7 and R^8 are independently selected from: H, $(C=O)_aO_b(C_1-C_{10})$ alkyl, $(C=O)_aO_b(C_3-C_8)$ cycloalkyl, $(C=O)_aO_b$ -aryl, $(C=O)_aO_b$ -heterocyclyl, (C_2-C_{10}) alkenyl, (C_2-C_{10}) alkynyl, SH, SO_2R^a , and $(C=O)_aNR^b_2$, said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R^{6a} , or R^7 and R^8 can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R^{6a} ;

R^a is (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl, aryl, or heterocyclyl; and


R^b is independently: H, (C_1-C_6) alkyl, aryl, heterocyclyl, (C_3-C_6) cycloalkyl, $(C=O)_aO_b(C_1-C_6)$ alkyl, or $S(O)_mR^a$;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a second embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula A, wherein:



is selected from:



a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; p is independently 0, 1, 2, 3, 4 or 5;

Ring Q is phenyl;

R^1 is independently selected from: H, oxo, $(C=O)_aO_b(C_1-C_{10})$ alkyl, $(C=O)_aO_b$ -aryl, $(C=O)_aO_b(C_2-C_{10})$ alkenyl, $(C=O)_aO_b(C_2-C_{10})$ alkynyl, CO_2H , halo, OH, $O_b(C_1-C_6)$ perfluoroalkyl, $(C=O)_aNR^7R^8$, CN, $(C=O)_aO_b(C_3-C_8)$ cycloalkyl, $S(O)_mNR^7R^8$, SH, $S(O)_m(C_1-C_{10})$ alkyl and $(C=O)_aO_b$ -heterocyclyl, said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, and heterocyclyl is optionally substituted with one or more substituents selected from R^6 ;

R² is independently selected from: OH, halo and (C₁-C₄)alkyl, said alkyl is optionally substituted with one or more substituents selected from OH and halo;

R³ and R^{3'} are independently selected from: H and (C₁-C₄)alkyl, said alkyl is optionally substituted with one or more substituents selected from OH, or R³ and R^{3'} can be taken together to form a (C₃-C₆)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from OH and CH₃;

R⁶ is: (C=O)_aO_bC₁-C₁₀ alkyl, (C=O)_aO_baryl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, (C=O)_aO_b heterocyclyl, CO₂H, halo, CN, OH, O_bC₁-C₆ perfluoroalkyl, O_a(C=O)_bNR⁷R⁸, oxo, CHO, (N=O)R⁷R⁸, S(O)_mNR⁷R⁸, SH, S(O)_m-(C₁-C₁₀)alkyl or (C=O)_aO_bC₃-C₈ cycloalkyl, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R^{6a};

R^{6a} is selected from: (C=O)_aO_b(C₁-C₁₀)alkyl, O_a(C₁-C₃)perfluoroalkyl, (C₀-C₆)alkylene-S(O)_mR^a, SH, oxo, OH, halo, CN, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, (C₃-C₆)cycloalkyl, (C₀-C₆)alkylene-aryl, (C₀-C₆)alkylene-heterocyclyl, (C₀-C₆)alkylene-N(R^b)₂, C(O)R^a, (C₀-C₆)alkylene-CO₂R^a, C(O)H, and (C₀-C₆)alkylene-CO₂H, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally substituted with up to three substituents selected from R^b, OH, (C₁-C₆)alkoxy, halogen, CO₂H, CN, O_a(C=O)_b(C₁-C₆)alkyl, oxo, and N(R^b)₂;


R⁷ and R⁸ are independently selected from: H, (C=O)_aO_b(C₁-C₁₀)alkyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heterocyclyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, SH, SO₂R^a, and (C=O)_aNR^b₂, said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R^{6a}, or R⁷ and R⁸ can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R^{6a};

R^a is (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, aryl, or heterocyclyl; and


R^b is independently: H, (C₁-C₆)alkyl, aryl, heterocyclyl, (C₃-C₆)cycloalkyl, (C=O)_aO_b(C₁-C₆)alkyl, or S(O)_mR^a;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a third embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula A, wherein:



is selected from



a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; p is 0;

Ring Q is phenyl;


5 R¹ is independently selected from: H, oxo, (C=O)_aO_b(C₁-C₁₀)alkyl, (C=O)_aO_b(C₂-C₁₀)alkenyl, (C=O)_aO_b(C₂-C₁₀)alkynyl, CO₂H, halo, OH, O_b(C₁-C₆)perfluoroalkyl, (C=O)_aNR⁷R⁸, CN, (C=O)_aO_b(C₃-C₈)cycloalkyl, S(O)_mNR⁷R⁸, SH, S(O)_m-(C₁-C₁₀)alkyl, said alkyl, alkenyl, alkynyl and cycloalkyl is optionally substituted with
 10 NH₂;

R³ and R^{3'} are taken together to form cyclobutyl, said cyclobutyl is optionally substituted with one or more substituents selected from OH and CH₃; and

R⁷ and R⁸ are independently selected from: H and (C₁-C₆)alkyl;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

15 In a fourth embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula B:



wherein,

20 A and B are independently selected from C or N, wherein at least one of A and B is N;

J and K are independently selected from C or N, wherein J and K are optionally substituted with R¹;

X, Y and Z are independently selected from C and N, wherein X, Y and Z are optionally substituted with R¹;

a is 0 or 1; b is 0 or 1; m is 0, 1 or 2;

R¹ is independently selected from (C=O)_aO_b(C₁-C₁₀)alkyl, (C=O)_aO_b(C₁-C₁₀)alkenyl, (C=O)_aO_b(C₁-C₁₀)alkynyl, (C=O)_aO_b(C₃-C₁₀)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_mNR⁷R⁸, S(O)_m(C₁-C₁₀)alkyl, S(O)_m(C₁-C₁₀)alkenyl, S(O)_m(C₁-C₁₀)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R⁶;

R³ and R^{3'} are independently selected from H, (C₁-C₁₀)alkyl, (C₁-C₁₀)alkenyl, (C₁-C₁₀)alkynyl, (C₃-C₁₀)cycloalkyl, aryl, heteroaryl, heterocyclyl, (C=O)NR⁷R⁸, halo, OH, CF₃, CO₂H, CN, and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₁₀)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NR⁷R⁸, or R³ and R^{3'} can be taken together to form a (C₃-C₁₀)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from H, (C₁-C₁₀)alkyl, (C₁-C₁₀)alkenyl, (C₁-C₁₀)alkynyl, (C₃-C₁₀)cycloalkyl, aryl, heteroaryl, heterocyclyl, (C=O)NR⁷R⁸, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are further optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NR⁷R⁸;

R⁶ is selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_mNR⁷R⁸, S(O)_m(C₁-C₄)alkyl, S(O)_m(C₁-C₄)alkenyl, S(O)_m(C₁-C₄)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a};

R^{6a} is selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;

R⁷ and R⁸ are independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, N(R^b)₂,


(C=O)N(R^b)₂, (C=S)N(R^b)₂, S(O)_mN(R^b)₂, S(O)_m(C₁-C₆)alkyl, S(O)_m(C₁-C₆)alkenyl, S(O)_m(C₁-C₆)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a}, or R⁷ and R⁸ can be taken together to form a (C₃-C₁₀)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from H, (C₁-C₁₀)alkyl, (C₁-C₁₀)alkenyl, (C₁-C₁₀)alkynyl, (C₃-C₁₀)cycloalkyl, aryl, heteroaryl, heterocyclyl, (C=O)NR⁷R⁸, halo, OH, CF₃, CO₂H, CN, and NR⁷R⁸; and

R^b is independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl and (C=O)_aO_b-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;


or a tautomer thereof;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a fifth embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula B, wherein,



is selected from:



a is 0 or 1; b is 0 or 1; m is 0, 1 or 2;

R¹ is independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_mNR⁷R⁸, S(O)_m(C₁-C₄)alkyl, S(O)_m(C₁-C₄)alkenyl, S(O)_m(C₁-C₄)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl,

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R⁶;

R³ and R^{3'} are independently selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl and phenyl are optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂, or R³ and R^{3'} can be taken together to form a (C₃-C₅)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;

R⁶ is selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_mNR⁷R⁸, S(O)_m(C₁-C₄)alkyl, S(O)_m(C₁-C₄)alkenyl, S(O)_m(C₁-C₄)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a};

R^{6a} is selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;



R⁷ and R⁸ are independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, N(R^b)₂, (C=O)N(R^b)₂, (C=S)N(R^b)₂, S(O)_mN(R^b)₂, S(O)_m(C₁-C₆)alkyl, S(O)_m(C₁-C₆)alkenyl, S(O)_m(C₁-C₆)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a}; and

R^b is independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl and (C=O)_aO_b-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;

or a tautomer thereof;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a sixth embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula B, wherein:



a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; n is 0, 1, 2 or 3;

R^1 is independently selected from $(C=O)_aO_b(C_1-C_6)alkyl$, $(C=O)_aO_b(C_1-C_6)alkenyl$, $(C=O)_aO_b(C_1-C_6)alkynyl$, $(C=O)_aO_b(C_3-C_8)cycloalkyl$, $(C=O)_aO_b-aryl$, $(C=O)_aO_b-heteroaryl$, $(C=O)_aO_b-heterocyclyl$, halo, OH, CF_3 , CO_2H , CN, NR^7R^8 , $(C=O)NR^7R^8$, $S(O)_mNR^7R^8$, $S(O)_m(C_1-C_4)alkyl$, $S(O)_m(C_1-C_4)alkenyl$, $S(O)_m(C_1-C_4)alkynyl$, $S(O)_m(C_3-C_8)cycloalkyl$, $S(O)_m-aryl$, $S(O)_m-heteroaryl$, and $S(O)_m-heterocyclyl$, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^6 ;

R^3 and $R^{3'}$ are independently selected from H, $(C_1-C_4)alkyl$, $(C_1-C_4)alkenyl$, $(C_1-C_4)alkynyl$, $(C_3-C_5)cycloalkyl$, halo, OH, CF_3 , CO_2H , CN, phenyl and NH_2 , said alkyl, alkenyl, alkynyl, cycloalkyl and phenyl are optionally substituted with one or more substituents selected from H, $(C_1-C_4)alkyl$, $(C_1-C_4)alkenyl$, $(C_1-C_4)alkynyl$, $(C_3-C_5)cycloalkyl$, halo, OH, CF_3 , CO_2H , CN, phenyl and NH_2 , or R^3 and $R^{3'}$ can be taken together to form a $(C_3-C_5)cycloalkyl$, said cycloalkyl is optionally substituted with one or more substituents selected from H, $(C_1-C_4)alkyl$, $(C_1-C_4)alkenyl$, $(C_1-C_4)alkynyl$, $(C_3-C_5)cycloalkyl$, halo, OH, CF_3 , CO_2H , CN, phenyl and NH_2 ;

R^6 is selected from $(C=O)_aO_b(C_1-C_6)alkyl$, $(C=O)_aO_b(C_1-C_6)alkenyl$, $(C=O)_aO_b(C_1-C_6)alkynyl$, $(C=O)_aO_b(C_3-C_8)cycloalkyl$, $(C=O)_aO_b-aryl$, $(C=O)_aO_b-heteroaryl$, $(C=O)_aO_b-heterocyclyl$, halo, OH, CF_3 , CO_2H , CN, NR^7R^8 , $(C=O)NR^7R^8$, $S(O)_mNR^7R^8$, $S(O)_m(C_1-C_4)alkyl$, $S(O)_m(C_1-C_4)alkenyl$, $S(O)_m(C_1-C_4)alkynyl$, $S(O)_m(C_3-C_8)cycloalkyl$, $S(O)_m-aryl$, $S(O)_m-heteroaryl$, and $S(O)_m-heterocyclyl$ said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a} ;

R^{6a} is selected from $(C=O)_aO_b(C_1-C_6)alkyl$, $(C=O)_aO_b(C_1-C_6)alkenyl$, $(C=O)_aO_b(C_1-C_6)alkynyl$, $(C=O)_aO_b(C_3-C_8)cycloalkyl$, $(C=O)_aO_b-aryl$, $(C=O)_aO_b-heteroaryl$, $(C=O)_aO_b-heterocyclyl$, halo, OH, CF_3 , CO_2H , CN and NH_2 , said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more

substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;


R⁷ and R⁸ are independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, N(R^b)₂, (C=O)N(R^b)₂, (C=S)N(R^b)₂, S(O)_mN(R^b)₂, S(O)_m(C₁-C₆)alkyl, S(O)_m(C₁-C₆)alkenyl, S(O)_m(C₁-C₆)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a}; and

R^b is independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl and (C=O)_aO_b-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;


or a tautomer thereof;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

In a seventh embodiment of this invention, the inhibitors of Akt activity are illustrated by the Formula B, wherein:



is selected from



a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; n is 0, 1 or 2;

R¹ is independently selected from (C=O)_aO_b(C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₃-C₈)cycloalkyl, O_b-aryl, O_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_m(C₁-C₄)alkyl, said alkyl, alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R⁶;

R³ and R^{3'} are H, or R³ and R^{3'} can be taken together to form cyclobutyl, said cyclobutyl is optionally substituted with one or more substituents selected from H, OH, methyl and cyclopropyl;

R⁶ is selected from O_b(C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, O_b-aryl, halo, OH, CF₃, NR⁷R⁸, said alkyl and cycloalkyl are optionally substituted with one or more substituents selected from R^{6a};

R^{6a} is selected from O_b(C₁-C₆)alkyl, aryl, OH and NH₂, said alkyl and aryl are optionally substituted with one or more substituents selected from halo;

R⁷ and R⁸ are independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_a(C₃-C₈)cycloalkyl, (C=O)_a-aryl, heterocyclyl, (C=O)N(R^b)₂, (C=S)N(R^b)₂, S(O)_m(C₁-C₆)alkyl, and S(O)_m-aryl, said alkyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a}; and

R^b is independently selected from H, (C₁-C₆)alkyl and aryl;

or a tautomer thereof;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

Specific compounds of the instant invention include:

1-[4-(6-phenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine (**A-1**);

1-[4-(2-isopropyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine (**B-1**);

1-[4-(2-methyl-6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine (**C-1**);

1-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine (**D**);

3-amino-3-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]-1-methylcyclobutanol (**E**);

3-amino-1-methyl-3-[4-(2-methyl-3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanol (**F**);

1-[4-(2-methyl-3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine (**G**);

3-amino-1-methyl-3-[4-(2-methyl-6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol (**H**);

3-amino-1-methyl-3-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol (**I**);

1-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine (**J**);

1-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine (**K**);

3-amino-1-methyl-3-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol (**L**);

1-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]methanamine (**M**);

1-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]methanamine (**N**); and

1-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]methanamine (**O**);

or a pharmaceutically acceptable salt or stereoisomer thereof.

Specific compounds of the instant invention include:

1-[4-(6-phenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine (**A-1**);

1-[4-(2-isopropyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine (**B-1**);

1-[4-(2-methyl-6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine (**C-1**);

1-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine (**D**);

3-amino-3-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]-1-methylcyclobutanol (**E**);

3-amino-1-methyl-3-[4-(2-methyl-3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanol (**F**);

1-[4-(2-methyl-3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine (**G**);

- 3-amino-1-methyl-3-[4-(2-methyl-6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol (**H**);
- 3-amino-1-methyl-3-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol (**I**);
- 1-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine (**J**);
- 5 1-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine (**K**);
- 3-amino-1-methyl-3-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol (**L**);
- 1-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]methanamine (**M**);
- 1-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]methanamine (**N**);
- 10 1-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]methanamine (**O**);
- 4-(6-Phenyl-[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)-benzylamine (**3-6**);
- 3-Amino-1-cyclopropyl-3-[4-(6-phenyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutanol (**3-7**);
- 4-(6-Phenyl-pyrazolo[1,5-*a*]pyrimidin-5-yl)-benzylamine (**3-8**);
- 15 1-[4-(6-Phenyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**4-2**);
- 1-[4-(3-Bromo-6-phenyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**4-4**);
- 1-[4-(6-Phenyl-3-vinyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**4-6**);
- 1-{4-[3-(2-Methoxy-phenyl)-6-phenyl-imidazo[1,2-*a*]pyrimidin-7-yl]-phenyl}-cyclobutylamine (**4-8**);
- 20 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-*a*]pyrimidine-3-carbonitrile (**4-10**);
- 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-*a*]pyrimidine-3-carboxylic acid amide (**4-11**);
- 1-[4-(3-Chloro-6-phenyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**4-14**);
- 25 1-[4-(3-Methyl-6-phenyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**4-15**);
- 1-{4-[3-(2-Cyclopropyl-vinyl)-6-phenyl-imidazo[1,2-*a*]pyrimidin-7-yl]-phenyl}-cyclobutylamine (**4-16**);
- 1-[4-(2,6-Diphenyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**5-3**);
- 1-[4-(3-Bromo-2,6-diphenyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**5-5**);
- 30 1-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-2,6-diphenyl-imidazo[1,2-*a*]pyrimidin-7-yl]-phenyl}-cyclobutylamine (**5-7**);
- 1-[4-(2,6-Diphenyl-3-vinyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**5-9**);
- 1-[4-(2-Ethyl-6-phenyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**5-11**);
- 35 1-[4-(6-Phenyl-2-trifluoromethyl-imidazo[1,2-*a*]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**5-12**);
- 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-*a*]pyrimidine-2-carboxylic acid ethyl ester (**5-13**);
- 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-*a*]pyrimidine-2-carboxylic acid (**5-14**);

- 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-2-carboxylic acid methyl ester (5-15);
- 1-{4-[2-(4-Fluoro-phenyl)-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutylamine (5-16);
- 5 1-[4-(6-Phenyl-2-p-tolyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (5-17);
- 1-[4-(3-Bromo-6-phenyl-2-trifluoromethyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (5-18);
- 1-[4-(3-Methyl-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (5-19);
- 1-[4-(2,3,6-Triphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (5-20);
- 10 1-[4-(3-Cyclopropyl-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (5-21);
- 1-{4-[2,6-Diphenyl-3-(2,3,4-trifluoro-phenyl)-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutylamine (5-22);
- 1-[4-(2,6-Diphenyl-3-propenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (5-23);
- 15 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-ylamine (6-2);
- N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-acetamide (6-4);
- 1-[4-(2,6-Diphenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-7);
- 20 1-[4-(2-Phenoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-9);
- 1-[4-(2-Morpholin-4-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-10);
- 1-[4-(6-Phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-11);
- 25 1-[4-(2-Methyl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-12);
- Ethanesulfonic acid {5-[4-(1-amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-amide (6-13);
- N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-benzenesulfonamide (6-14);
- 30 N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-benzamide (6-15);
- Cyclopropanecarboxylic acid {5-[4-(1-amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-amide (6-16);
- 1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-3-phenyl-urea (6-17);
- 35 1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-3-phenyl-thiourea (6-18);

- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-carbamic acid ethyl ester (6-19);
- 1-{4-[2-(6-Methoxy-pyridin-2-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-20);
- 5 1-[4-(6-Phenyl-2-pyrazin-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-21);
- 1-{4-[2-(2,3-Dimethyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-22);
- 1-{4-[2-(3,4-Difluoro-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-23);
- 10 1-{4-[2-(5-Fluoro-2-methyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-24);
- 1-[4-(6-Phenyl-2-pyridin-4-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-25);
- 15 4-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-phenylamine (6-26);
- 1-[4-(6-Phenyl-2-p-tolyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-27);
- 1-{4-[2-(4-Methoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-28);
- 20 1-[4-(6-Phenyl-2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-29);
- 1-{4-[2-(2-Methoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-30);
- 1-{4-[2-(3-Chloro-4-fluoro-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-31);
- 25 1-{4-[2-(2-Fluoro-4-trifluoromethyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-32);
- 1-{4-[6-Phenyl-2-(2,3,4-trifluoro-phenyl)-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-33);
- 30 1-{4-[2-(2,4-Dimethoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-34);
- 1-[4-(2-Furan-3-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-35);
- 1-[4-(6-Phenyl-2-thiophen-3-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-36);
- 35 1-[4-(6-Phenyl-2-thiophen-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-37);

- 1-{4-[2-(6-Methoxy-naphthalen-2-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-38);
- 1-[4-(2-Dibenzofuran-4-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-39);
- 5 4-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-phenol (6-40);
- 1-[4-(6-Phenyl-2-m-tolyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-41);
- 1-{4-[2-(4-Phenoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-42);
- 10 1-[4-(2-Biphenyl-2-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-43);
- 1-[4-(2-Benzothiazol-2-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-44);
- 1-[4-(6-Phenyl-2-pyridin-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-45);
- 15 1-[4-(6-Phenyl-2-thiazol-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-46);
- 1-{4-[6-Phenyl-2-(pyridin-4-yloxy)-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-47);
- 20 1-[4-(2-Methoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-48);
- 1-[4-(2-Phenoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-49);
- 1-[4-(2-Methylsulfanyl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-50);
- 1-{4-[2-(Naphthalen-2-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-51);
- 25 1-{4-[2-(Biphenyl-2-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-52);
- 1-{4-[2-(Isoquinolin-7-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-53);
- 30 1-{4-[2-(4-Phenoxy-phenoxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-54);
- 1-{4-[2-(3,4-Dihydro-2H-quinolin-1-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-55);
- 1-{4-[2-(2,3-Dihydro-indol-1-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (6-56);
- 35 {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-methylamine (6-57);

- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-dimethyl-amine (6-58);
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-pyridin-3-yl-amine (6-59);
- 5 {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-diethyl-amine (6-60);
- N1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-ethane-1,2-diamine (6-61);
- 2-({5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-methyl-10 amino)-ethanol (6-62);
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-benzyl-methyl-amine (6-63);
- 1-[4-(2-{4-[Bis-(4-chloro-phenyl)-methyl]-piperazin-1-yl}-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-64);
- 15 {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-methyl-phenyl-amine (6-65);
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-(2-methoxyethyl)-amine (6-66);
- 3-Amino-3-[4-(2,6-diphenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol 20 (7-2);
- 3-Amino-1-methyl-3-[4-(6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutanol (7-3);
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methyl ester (8-2);
- 25 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid (8-4);
- 2-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-propan-2-ol (8-6);
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid 30 methylamide (8-8);
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-diphenyl-methanol (8-9);
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid amide (8-10);
- 35 {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-morpholin-4-yl-methanone (8-11);
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid phenethyl-amide (8-12);


- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid diethylamide (8-13);
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid phenylamide (8-14);
- 5 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid benzyl-methyl-amide (8-15);
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methoxy-methyl-amide (8-16);
- 10 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methyl-phenyl-amide (8-17);
- 1-[4-(2,6-Diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-2);
- 1-[4-(2,3,6-Triphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-4);
- 1-[4-(2,3,6-Triphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-6);
- 1-[4-(6-Phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-8);
- 15 1-[4-(2-Methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-9);
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-pyrazolo[1,5-a]pyrimidin-2-ol (9-10);
- 1-[4-(6-Phenyl-2-p-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-11);
- 1-{4-[2-(4-Methoxy-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (9-12);
- 20 1-[4-(6-Phenyl-2-o-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-13);
- 1-{4-[2-(4-Chloro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (9-14);
- 1-{4-[2-(3-Fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (9-15);
- 25 1-{4-[6-Phenyl-2-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (9-16);
- 1-{4-[2-(4-Fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (9-17);
- 1-[4-(3-Chloro-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-18);
- 30 1-[4-(3-Bromo-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-19);
- 1-[4-(2,6-Diphenyl-3-pyridin-4-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-20);
- 1-[4-(3-Methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-21);
- 1-{4-[3-(2,4-Difluoro-phenyl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (9-22);
- 35 1-[4-(3-Methyl-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-23);
- 1-[4-(3,6-Diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-24);
- 1-[4-(6-Phenyl-3-pyrimidin-5-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-25);

- 1-{4-[3-(4-Methoxy-phenyl)-2-methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (**9-26**);
- 1-{4-[3-(2-Methoxy-pyrimidin-5-yl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (**9-27**);
- 5 1-{4-[6-Phenyl-3-(4-trifluoromethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (**9-28**);
- 1-[4-(3-Naphthalen-1-yl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (**9-29**);
- 1-{4-[3-(2-Fluoro-5-trifluoromethyl-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (**9-30**);
- 10 1-{4-[3-(3-Chloro-4-fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (**9-31**);
- 1-{4-[3-(3-Fluoro-5-isopropoxy-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (**9-32**);
- 15 1-{4-[6-Phenyl-3-(2,3,4-trifluoro-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine (**9-33**);
- 3-Amino-3-[4-(2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol (**10-2**);
- 3-Amino-3-[4-(3-bromo-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol (**10-4**);
- 20 3-Amino-1-methyl-3-[4-(3-methyl-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutanol (**10-6**);
- 3-Amino-1-methyl-3-[4-(6-phenyl-2-o-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutanol (**10-7**);
- 25 3-Amino-3-{4-[2-(4-fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-1-methyl-cyclobutanol (**10-8**);
- 3-Amino-3-[4-(2,6-diphenyl-3-pyridin-3-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol (**10-9**);
- 3-Amino-3-{4-[3-(3-amino-phenyl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-1-methyl-cyclobutanol (**10-10**);
- 30 3-Amino-3-[4-(2,6-diphenyl-3-pyrimidin-5-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol (**10-11**); and
- 3-Amino-3-{4-[3-(2-amino-pyrimidin-5-yl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-1-methyl-cyclobutanol (**10-12**);
- 35 or a tautomer thereof;
- or a pharmaceutically acceptable salt or stereoisomer thereof.

The compounds of the present invention may have asymmetric centers, chiral axes, and chiral planes (as described in: E.L. Eliel and S.H. Wilen, *Stereochemistry of Carbon*

Compounds, John Wiley & Sons, New York, 1994, pages 1119-1190), and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers, all such stereoisomers being included in the present invention.

In addition, the compounds disclosed herein may exist as tautomers and both tautomeric forms are intended to be encompassed by the scope of the invention, even though only one tautomeric structure is depicted. For example the following is within the scope of the instant invention:



In particular, compounds disclosed in Formulas A, B and C may exist in the keto form (i.e. oxo substituent). An oxo group is included in the instant invention and where appropriate is a member of the substituent list (R¹, R², R³, R^{3'}, R^{3''}, R⁶, R⁷, R⁸, R^a, R^b). Tetrazoles exist as a mixture of 1H/2H tautomers. The tautomeric forms of the tetrazol moiety are also within the scope of the instant invention.

This invention is also intended to encompass pro-drugs of the compounds disclosed herein. A prodrug of any of the compounds can be made using well known pharmacological techniques.

When any variable (e.g. R¹, etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every other occurrence. Also, combinations of substituents and variables are permissible only if such combinations result in stable compounds.

Lines drawn into the ring systems from substituents represent that the indicated bond may be attached to any of the substitutable ring atoms. If the ring system is bicyclic, it is intended that the bond be attached to any of the suitable atoms on either ring of the bicyclic moiety.

It is understood that one or more silicon (Si) atoms can be incorporated into the compounds of the instant invention in place of one or more carbon atoms by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art from readily available starting materials. Carbon and silicon differ in their covalent radius leading to differences in bond distance and the steric arrangement when comparing analogous C-element and Si-element bonds. These differences lead to subtle changes in the size and shape of silicon-containing compounds when compared to carbon. One of ordinary skill in the art would understand that size and shape differences can lead to subtle or dramatic changes in potency, solubility, lack of off target activity, packaging properties, and so on. (Dias, J. O. *et al.* *Organometallics* (2006) 5:1188-1198; Showell, G.A. *et al.* *Bioorganic & Medicinal Chemistry Letters* (2006) 16:2555-2558).

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that

are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials. If a substituent is itself substituted with more than one group, it is understood that these multiple groups may be on the same carbon or on different carbons, so long as a stable structure results. The phrase

5 “optionally substituted with one or more substituents” should be taken to be equivalent to the phrase “optionally substituted with at least one substituent” and in such cases the preferred embodiment will have from zero to four substituents, and the more preferred embodiment will have from zero to three substituents.

As used herein, “alkyl” is intended to include both branched and straight-chain

10 saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, C₁-C₁₀, as in “(C₁-C₁₀)alkyl” is defined to include groups having 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbons in a linear or branched arrangement. For example, “(C₁-C₁₀)alkyl” specifically includes methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *t*-butyl, *i*-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, and so on.

15 The term “cycloalkyl” means a monocyclic saturated aliphatic hydrocarbon group having the specified number of carbon atoms. For example, “cycloalkyl” includes cyclopropyl, methyl-cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, cyclohexyl, and so on.

“Alkoxy” represents either a cyclic or non-cyclic alkyl group of indicated number of carbon atoms attached through an oxygen bridge. “Alkoxy” therefore encompasses the

20 definitions of alkyl and cycloalkyl above.

If no number of carbon atoms is specified, the term “alkenyl” refers to a non-aromatic hydrocarbon radical, straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to four non-aromatic carbon-carbon double bonds may be present. Thus, “(C₂-

25 C₁₀)alkenyl” means an alkenyl radical having from 2 to 10 carbon atoms. Alkenyl groups include ethenyl, propenyl, butenyl, 2-methylbutenyl and cyclohexenyl. The straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated.

The term “alkynyl” refers to a hydrocarbon radical straight, branched or cyclic,

30 containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond. Up to three carbon-carbon triple bonds may be present. Thus, “(C₂-C₁₀)alkynyl” means an alkynyl radical having from 2 to 10 carbon atoms. Alkynyl groups include ethynyl, propynyl, butynyl, 3-methylbutynyl and so on. The straight, branched or cyclic portion of the alkynyl group may contain triple bonds and may be substituted if a substituted alkynyl group is indicated.

35 In certain instances, substituents may be defined with a range of carbons that includes zero, such as (C₀-C₆)alkylene-aryl. If aryl is taken to be phenyl, this definition would include phenyl itself as well as -CH₂Ph, -CH₂CH₂Ph, CH(CH₃)CH₂CH(CH₃)Ph, and so on.

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydro-naphthyl, indanyl and biphenyl. In cases where the aryl substituent is bicyclic and one ring is non-aromatic, it is understood that

attachment is via the aromatic ring.


The term heteroaryl, as used herein, represents a stable monocyclic or bicyclic ring of up to 7 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Heteroaryl groups within the scope of this definition include but are not limited to: acridinyl, carbazoyl, cinnolinyl, quinoxalinyl, pyrazoyl, indolyl, benzotriazolyl, furanyl, thienyl, benzothienyl, benzofuranyl, quinolinyl, isoquinolinyl, oxazolyl, isoxazolyl, indolyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, tetrahydroquinoline. As with the definition of heterocycle below, "heteroaryl" is also understood to include the N-oxide derivative of any nitrogen-containing heteroaryl. In cases where the heteroaryl substituent is bicyclic and one ring is non-aromatic or contains no heteroatoms, it is understood that attachment is via the aromatic ring or via the heteroatom containing ring, respectively. Such heteroaryl moieties for substituent Q include but are not limited to: 2-benzimidazolyl, 2-quinolinyl, 3-quinolinyl, 4-quinolinyl, 1-isoquinolinyl, 3-isoquinolinyl and 4-isoquinolinyl.

The term "heterocycle" or "heterocyclyl" as used herein is intended to mean a 3- to 10-membered aromatic or nonaromatic heterocycle containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S, and includes bicyclic groups. "Heterocyclyl" therefore includes the above mentioned heteroaryls, as well as dihydro and tetrahydro analogs thereof. Further examples of "heterocyclyl" include, but are not limited to the following: benzoimidazolyl, benzoimidazolonyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazoyl, carbolinyl, cinnolinyl, furanyl, imidazolyl, indolinyl, indolyl, indolaziny, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinazolinyl, quinolyl, quinoxalinyl, tetrahydropyranyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, azetidiny, 1,4-dioxanyl, hexahydroazepinyl, piperazinyl, piperidinyl, pyridin-2-onyl, pyrrolidinyl, morpholinyl, thiomorpholinyl, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyrazinyl, dihydropyrazolyl, dihydropyridinyl, dihydropyrimidinyl, dihydropyrrolyl, dihydroquinolinyl, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl, dihydroazetidiny, methylenedioxybenzoyl, tetrahydrofuranyl, and tetrahydrothienyl, and N-oxides thereof. Attachment of a heterocyclyl substituent can occur via a carbon atom or via a heteroatom.


As appreciated by those of skill in the art, "halo" or "halogen" as used herein is intended to include chloro (Cl), fluoro (F), bromo (Br) and iodo (I).

In an embodiment of Formula A, X, Y and Z are independently selected from C and N, wherein X, Y and Z are optionally substituted with R¹.

5 In an embodiment of Formula A, the moiety illustrated by the formula:



includes the following structures:



In an embodiment, Ring Q is selected from: phenyl and heterocyclyl.

10 In yet another embodiment, Ring Q is phenyl.

In an embodiment, p is independently 0, 1, 2 or 3.

In a further embodiment, p is independently 0, 1 or 2.

In another embodiment, p is independently 1.

In another embodiment, p is 0.

15 In an embodiment, R¹ is selected from: oxo, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b-aryl, (C=O)_aO_b(C₂-C₆)alkenyl, (C=O)_aO_b(C₂-C₆)alkynyl, CO₂H, halo, OH, O_b(C₁-C₆)perfluoroalkyl, (C=O)_aNH₂, CN, (C=O)_aO_b(C₃-C₈)cycloalkyl, S(O)₂NH₂, SH, S(O)₂-(C₁-C₆)alkyl and (C=O)_aO_b-heterocyclyl, said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, and heterocyclyl is optionally substituted with halo, (C₁-C₆)alkyl, (C₁-C₆)alkyl-NH₂, OH, O(C₁-C₆)alkyl.

20 In another embodiment, R¹ is selected from: oxo, (C=O)_aO_b(C₁-C₁₀)alkyl, CO₂H, halo, OH, CN, (C₁-C₆)alkoxy, S(O)₂NH₂, SH, S(O)₂-(C₁-C₆)alkyl, O(C=O)(C₁-C₆)alkyl and NH₂, said alkyl is optionally substituted with halo, (C₁-C₆)alkyl, (C₁-C₆)alkyl-NH₂, OH, O(C₁-C₆)alkyl.

25 In another embodiment, R¹ is selected from: oxo, NH₂, OH, SH, O_a(C₁-C₆)alkyl, said alkyl is optionally substituted with halo, NH₂, OH, O(C₁-C₆)alkyl.

In another embodiment, when R¹ is aryl, said aryl is phenyl.

30 In an embodiment, R³ and R^{3'} are independently selected from: H and (C₁-C₄)alkyl, said alkyl is optionally substituted with one or more substituents selected from OH, or R³ and R^{3'} can be taken together to form cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

said cyclo-moiety is optionally substituted with one or more substituents selected from OH and CH₃.

In another embodiment, R³ and R^{3'} are independently selected from: H and CH₃, or R³ and R^{3'} can be taken together to form cyclobutyl, said cyclobutyl is optionally substituted with one or more substituents selected from OH and CH₃.

In another embodiment, R³ and R^{3'} are taken together to form cyclobutyl, said cyclobutyl is optionally substituted with one or more substituents selected from OH and CH₃.

In an embodiment of Formula C, R^{3'} and R^{3''} are independently H, OH, cyclopropyl and CH₃.

In an embodiment R² is independently selected from H, OH, halo, O(C₁-C₃)alkyl and (C₁-C₃)alkyl.

In an embodiment R² is H.

In an embodiment, R^a is (C₁-C₆)alkyl.

In an embodiment R^b is independently selected from H and (C₁-C₆)alkyl.

In another embodiment of Formula A, Ring Q is phenyl; R³ and R^{3'} are taken together to form cyclobutyl, which is optionally substituted with OH and CH₃; R¹ is selected from: H, heterocyclyl, (C₃-C₆)cycloalkyl, OH, (C₁-C₆)alkyl, NH(C₁-C₆)alkyl, NH-heterocyclyl, NH-cycloalkyl, said heterocyclyl, cycloalkyl and alkyl is optionally substituted with: halo, (C₁-C₆)alkyl, (C₁-C₆)alkyl-NH₂, OH, O(C₁-C₆)alkyl; and R² is H or F.

In another embodiment of Formula A, Ring Q is phenyl; p is 0; R³ and R^{3'} are taken together to form cyclobutyl, which is optionally substituted with OH and CH₃; and R¹ is selected from: H, heterocyclyl, (C₃-C₆)cycloalkyl, OH, (C₁-C₆)alkyl, NH(C₁-C₆)alkyl, NH-heterocyclyl, NH-cycloalkyl, said heterocyclyl, cycloalkyl and alkyl is optionally substituted with: halo, (C₁-C₆)alkyl, (C₁-C₆)alkyl-NH₂, OH, O(C₁-C₆)alkyl.

In an aspect of all embodiments of Formulas A and B, R³ and R^{3'} are H, or R³ and R^{3'} are taken together to form cyclobutyl, said cyclobutyl is optionally substituted with one or more substituents selected from OH, CH₃ and cyclopropyl.

In an aspect of all embodiments of Formulas A and B, R³ and R^{3'} are independently selected from H, (C₁-C₁₀)alkyl, (C₁-C₁₀)alkenyl, (C₁-C₁₀)alkynyl, (C₃-C₁₀)cycloalkyl, aryl, heteroaryl, heterocyclyl, (C=O)NR⁷R⁸, halo, OH, CF₃, CO₂H, CN, and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₁₀)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NR⁷R⁸, or R³ and R^{3'} can be taken together to form a (C₃-C₁₀)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from H, (C₁-C₁₀)alkyl, (C₁-C₁₀)alkenyl, (C₁-C₁₀)alkynyl, (C₃-C₁₀)cycloalkyl, aryl, heteroaryl, heterocyclyl, (C=O)NR⁷R⁸, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are

further optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NR⁷R⁸.

In an aspect of all embodiments of Formulas A and B, R⁶ is selected from O_b(C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, O_b-aryl, halo, OH, CF₃, NR⁷R⁸, said alkyl and cycloalkyl are optionally substituted with one or more substituents selected from R^{6a}.

In an aspect of all embodiments of Formulas A and B, R^{6a} is selected from O_b(C₁-C₆)alkyl, aryl, OH and NH₂, said alkyl and aryl are optionally substituted with one or more substituents selected from halo.

In an aspect of all embodiments of Formulas A and B, R⁷ and R⁸ are independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_a(C₃-C₈)cycloalkyl, (C=O)_a-aryl, heterocyclyl, (C=O)N(R^b)₂, (C=S)N(R^b)₂, S(O)_m(C₁-C₆)alkyl, and S(O)_m-aryl, said alkyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a}.

In an embodiment of Formula B, R⁷ and R⁸ are independently selected from H, (C₁-C₄)alkyl and phenyl.

In an embodiment of Formula B, R^b is independently selected from H, (C₁-C₄)alkyl, and phenyl.

Included in the instant invention is the free form of compounds of Formula A, as well as the pharmaceutically acceptable salts and stereoisomers thereof. Some of the isolated specific compounds exemplified herein are the protonated salts of amine compounds. The term "free form" refers to the amine compounds in non-salt form. The encompassed pharmaceutically acceptable salts not only include the isolated salts exemplified for the specific compounds described herein, but also all the typical pharmaceutically acceptable salts of the free form of compounds of Formula A. The free form of the specific salt compounds described may be isolated using techniques known in the art. For example, the free form may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia and sodium bicarbonate. The free forms may differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise pharmaceutically equivalent to their respective free forms for purposes of the invention.

The pharmaceutically acceptable salts of the instant compounds can be synthesized from the compounds of this invention which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts of the basic compounds are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents. Similarly, the salts of the acidic compounds are formed by reactions with the appropriate inorganic or organic base.

Thus, pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed by reacting a basic instant compound with an inorganic or organic acid. For example, conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like, as well as salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic (TFA) and the like.

When the compound of the present invention is acidic, suitable "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N¹-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

The preparation of the pharmaceutically acceptable salts described above and other typical pharmaceutically acceptable salts is more fully described by Berg *et al.*, "Pharmaceutical Salts," *J. Pharm. Sci.*, 1977:66:1-19.

It will also be noted that the compounds of the present invention are potentially internal salts or zwitterions, since under physiological conditions a deprotonated acidic moiety in the compound, such as a carboxyl group, may be anionic, and this electronic charge might then be balanced off internally against the cationic charge of a protonated or alkylated basic moiety, such as a quaternary nitrogen atom.

UTILITY

The compounds of the instant invention are inhibitors of the activity of Akt and are thus useful in the treatment of cancer, in particular cancers associated with irregularities in the activity of Akt and downstream cellular targets of Akt. Such cancers include, but are not limited to, ovarian, pancreatic, breast and prostate cancer, as well as cancers (including glioblastoma) where the tumor suppressor PTEN is mutated (Cheng *et al.*, *Proc. Natl. Acad. Sci.*

(1992) 89:9267-9271; Cheng et al., *Proc. Natl. Acad. Sci.* (1996) 93:3636-3641; Bellacosa et al., *Int. J. Cancer* (1995) 64:280-285; Nakatani et al., *J. Biol. Chem.* (1999) 274:21528-21532; Graff, *Expert. Opin. Ther. Targets* (2002) 6(1):103-113; and Yamada and Araki, *J. Cell Science.* (2001) 114:2375-2382; Mischel and Cloughesy, *Brain Pathol.* (2003) 13(1):52-61).

5 The compounds, compositions and methods provided herein are particularly deemed useful for the treatment of cancer. Cancers that may be treated by the compounds, compositions and methods of the invention include, but are not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: non-small cell lung, bronchogenic carcinoma (squamous

10 cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma),

15 small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma), colon, colorectal, rectal; Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate

20 (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's

25 sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma,

30 ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell

35 tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia,

chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles, dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above-identified conditions.

Cancers that may be treated by the compounds, compositions and methods of the invention include, but are not limited to: breast, prostate, colon, colorectal, lung, non-small cell lung, brain, testicular, stomach, pancreas, skin, small intestine, large intestine, throat, head and neck, oral, bone, liver, bladder, kidney, thyroid and blood.

Cancers that may be treated by the compounds, compositions and methods of the invention include: breast, prostate, colon, ovarian, colorectal, lung and non-small cell lung.

Cancers that may be treated by the compounds, compositions and methods of the invention include: breast, colon, (colorectal) and lung (non-small cell lung).

Cancers that may be treated by the compounds, compositions and methods of the invention include: lymphoma and leukemia.

The compounds of the instant invention are useful for the treatment of breast cancer.

The compounds of the instant invention are useful for the treatment of prostate cancer.

Akt signaling regulates multiple critical steps in angiogenesis. Shiojima and Walsh, *Circ. Res.* (2002) 90:1243-1250. The utility of angiogenesis inhibitors in the treatment of cancer is known in the literature, see J. Rak et al. *Cancer Research*, 55:4575-4580, 1995 and Dredge et al., *Expert Opin. Biol. Ther.* (2002) 2(8):953-966, for example. The role of angiogenesis in cancer has been shown in numerous types of cancer and tissues: breast carcinoma (G. Gasparini and A.L. Harris, *J. Clin. Oncol.*, 1995, 13:765-782; M. Toi et al., *Japan. J. Cancer Res.*, 1994, 85:1045-1049); bladder carcinomas (A.J. Dickinson et al., *Br. J. Urol.*, 1994, 74:762-766); colon carcinomas (L.M. Ellis et al., *Surgery*, 1996, 120(5):871-878); and oral cavity tumors (J.K. Williams et al., *Am. J. Surg.*, 1994, 168:373-380). Other cancers include, advanced tumors, hairy cell leukemia, melanoma, advanced head and neck, metastatic renal cell, non-Hodgkin's lymphoma, metastatic breast, breast adenocarcinoma, advanced melanoma, pancreatic, gastric, glioblastoma, lung, ovarian, non-small cell lung, prostate, small cell lung, renal cell carcinoma, various solid tumors, multiple myeloma, metastatic prostate, malignant glioma, renal cancer, lymphoma, refractory metastatic disease, refractory multiple myeloma, cervical cancer, Kaposi's sarcoma, recurrent anaplastic glioma, and metastatic colon cancer (Dredge et al., *Expert Opin. Biol. Ther.* (2002) 2(8):953-966). Thus, the Akt inhibitors disclosed in the instant application are also useful in the treatment of these angiogenesis related cancers.

Tumors which have undergone neovascularization show an increased potential for metastasis. In fact, angiogenesis is essential for tumor growth and metastasis. (S.P. Cunningham, et al., *Can. Research*, 61: 3206-3211 (2001)). The Akt inhibitors disclosed in the present application are therefore also useful to prevent or decrease tumor cell metastasis.

Further included within the scope of the invention is a method of treating or preventing a disease in which angiogenesis is implicated, which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of the present invention. Ocular neovascular diseases are an example of conditions where much of the resulting tissue damage can be attributed to aberrant infiltration of blood vessels in the eye (see WO 00/30651, published 2 June 2000). The undesirable infiltration can be triggered by ischemic retinopathy, such as that resulting from diabetic retinopathy, retinopathy of prematurity, retinal vein occlusions, etc., or by degenerative diseases, such as the choroidal neovascularization observed in age-related macular degeneration. Inhibiting the growth of blood vessels by administration of the present compounds would therefore prevent the infiltration of blood vessels and prevent or treat diseases where angiogenesis is implicated, such as ocular diseases like retinal vascularization, diabetic retinopathy, age-related macular degeneration, and the like.

Further included within the scope of the invention is a method of treating or preventing a non-malignant disease in which angiogenesis is implicated, including but not limited to: ocular diseases (such as, retinal vascularization, diabetic retinopathy and age-related macular degeneration), atherosclerosis, arthritis, psoriasis, obesity and Alzheimer's disease (Dredge et al., *Expert Opin. Biol. Ther.* (2002) 2(8):953-966). In another embodiment, a method of treating or preventing a disease in which angiogenesis is implicated includes: ocular diseases (such as, retinal vascularization, diabetic retinopathy and age-related macular degeneration), atherosclerosis, arthritis and psoriasis.

Further included within the scope of the invention is a method of treating hyperproliferative disorders such as restenosis, inflammation, autoimmune diseases and allergy/asthma.

Further included within the scope of the instant invention is the use of the instant compounds to coat stents and therefore the use of the instant compounds on coated stents for the treatment and/or prevention of restenosis (WO03/032809).

Further included within the scope of the instant invention is the use of the instant compounds for the treatment and/or prevention of osteoarthritis (WO03/035048).

Further included within the scope of the invention is a method of treating hyperinsulinism.

The compounds of the invention are also useful in preparing a medicament that is useful in treating the diseases described above, in particular cancer.

In an embodiment of the invention, the instant compound is a selective inhibitor whose inhibitory efficacy is dependent on the PH domain. In this embodiment, the compound

exhibits a decrease in *in vitro* inhibitory activity or no *in vitro* inhibitory activity against truncated Akt proteins lacking the PH domain.

In a further embodiment, the instant compound is selected from the group of a selective inhibitor of Akt1, a selective inhibitor of Akt2 and a selective inhibitor of both Akt1 and Akt2.

In another embodiment, the instant compound is selected from the group of a selective inhibitor of Akt1, a selective inhibitor of Akt2, a selective inhibitor of Akt3 and a selective inhibitor of two of the three Akt isoforms.

In another embodiment, the instant compound is a selective inhibitor of all three Akt isoforms, but is not an inhibitor of one, two or all of such Akt isoforms that have been modified to delete the PH domain, the hinge region or both the PH domain and the hinge region.

The present invention is further directed to a method of inhibiting Akt activity which comprises administering to a mammal in need thereof a pharmaceutically effective amount of the instant compound.

The compounds of this invention may be administered to mammals, including humans, either alone or, in combination with pharmaceutically acceptable carriers, excipients or diluents, in a pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, microcrystalline cellulose, sodium crosscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to mask the unpleasant taste of the drug or delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water soluble taste masking material such as hydroxypropylmethyl-cellulose or hydroxypropylcellulose, or a time delay material such as ethyl cellulose, cellulose acetate butyrate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethylene-oxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated hydroxyanisole or alpha-tocopherol.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsion. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphatides, for example soy bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example

polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring agents, preservatives and antioxidants.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent,
5 a preservative, flavoring and coloring agents and antioxidant.

The pharmaceutical compositions may be in the form of sterile injectable aqueous solutions. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

The sterile injectable preparation may also be a sterile injectable oil-in-water
10 microemulsion where the active ingredient is dissolved in the oily phase. For example, the active ingredient may be first dissolved in a mixture of soybean oil and lecithin. The oil solution then introduced into a water and glycerol mixture and processed to form a microemulsion.

The injectable solutions or microemulsions may be introduced into a patient's blood-stream by local bolus injection. Alternatively, it may be advantageous to administer the
15 solution or microemulsion in such a way as to maintain a constant circulating concentration of the instant compound. In order to maintain such a constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

The pharmaceutical compositions may be in the form of a sterile injectable
20 aqueous or oleagenous suspension for intramuscular and subcutaneous administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. In addition, sterile,
25 fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Compounds of Formula A may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug
30 with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

For topical use, creams, ointments, jellies, solutions or suspensions, etc.,
35 containing the compound of Formula A are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or via transdermal routes,

using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Compounds of the present invention may also be delivered as a suppository employing bases such as cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

When a composition according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

The dosage regimen utilizing the compounds of the instant invention can be selected in accordance with a variety of factors including type, species, age, weight, sex and the type of cancer being treated; the severity (i.e., stage) of the cancer to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to treat, for example, to prevent, inhibit (fully or partially) or arrest the progress of the disease. For example, compounds of the instant invention can be administered in a total daily dose of up to 10,000 mg. Compounds of the instant invention can be administered once daily (QD), or divided into multiple daily doses such as twice daily (BID), and three times daily (TID). Compounds of the instant invention can be administered at a total daily dosage of up to 10,000 mg, e.g., 2,000 mg, 3,000 mg, 4,000 mg, 6,000 mg, 8,000 mg or 10,000 mg, which can be administered in one daily dose or can be divided into multiple daily doses as described above.

For example, compounds of the instant invention can be administered in a total daily dose of up to 1,000 mg. Compounds of the instant invention can be administered once daily (QD), or divided into multiple daily doses such as twice daily (BID), and three times daily (TID). Compounds of the instant invention can be administered at a total daily dosage of up to 1,000 mg, e.g., 200 mg, 300 mg, 400 mg, 600 mg, 800 mg or 1,000 mg, which can be administered in one daily dose or can be divided into multiple daily doses as described above.

In addition, the administration can be continuous, i.e., every day, or intermittently. The terms "intermittent" or "intermittently" as used herein means stopping and starting at either regular or irregular intervals. For example, intermittent administration of a compound of the instant invention may be administration one to six days per week or it may mean administration in cycles (e.g. daily administration for two to eight consecutive weeks, then a rest period with no administration for up to one week) or it may mean administration on alternate days.

In addition, the compounds of the instant invention may be administered according to any of the schedules described above, consecutively for a few weeks, followed by a rest period. For example, the compounds of the instant invention may be administered according

to any one of the schedules described above from two to eight weeks, followed by a rest period of one week, or twice daily at a dose of 100 - 500 mg for three to five days a week. In another particular embodiment, the compounds of the instant invention may be administered three times daily for two consecutive weeks, followed by one week of rest.

5 Any one or more of the specific dosages and dosage schedules of the compounds of the instant invention, may also be applicable to any one or more of the therapeutic agents to be used in the combination treatment (hereinafter referred to as the "second therapeutic agent").

Moreover, the specific dosage and dosage schedule of this second therapeutic agent can further vary, and the optimal dose, dosing schedule and route of administration will be
10 determined based upon the specific second therapeutic agent that is being used.

Of course, the route of administration of the compounds of the instant invention is independent of the route of administration of the second therapeutic agent. In an embodiment, the administration for a compound of the instant invention is oral administration. In another embodiment, the administration for a compound of the instant invention is intravenous
15 administration. Thus, in accordance with these embodiments, a compound of the instant invention is administered orally or intravenously, and the second therapeutic agent can be administered orally, parenterally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery by catheter or stent, subcutaneously, intraadiposally,
20 intraarticularly, intrathecally, or in a slow release dosage form.

In addition, a compound of the instant invention and second therapeutic agent may be administered by the same mode of administration, i.e. both agents administered e.g. orally, by IV. However, it is also within the scope of the present invention to administer a compound of the instant invention by one mode of administration, e.g. oral, and to administer the second
25 therapeutic agent by another mode of administration, e.g. IV or any other ones of the administration modes described hereinabove.

The first treatment procedure, administration of a compound of the instant invention, can take place prior to the second treatment procedure, i.e., the second therapeutic agent, after the treatment with the second therapeutic agent, at the same time as the treatment
30 with the second therapeutic agent, or a combination thereof. For example, a total treatment period can be decided for a compound of the instant invention. The second therapeutic agent can be administered prior to onset of treatment with a compound of the instant invention or following treatment with a compound of the instant invention. In addition, anti-cancer treatment can be administered during the period of administration of a compound of the instant invention but does
35 not need to occur over the entire treatment period of a compound of the instant invention.

The instant compounds are also useful in combination with therapeutic, chemotherapeutic and anti-cancer agents. Combinations of the presently disclosed compounds with therapeutic, chemotherapeutic and anti-cancer agents are within the scope of the invention.

Examples of such agents can be found in *Cancer Principles and Practice of Oncology* by V.T. Devita and S. Hellman (editors), 6th edition (February 15, 2001), Lippincott Williams & Wilkins Publishers. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the cancer

involved. Such agents include the following: estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic/cytostatic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors and other angiogenesis inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors, inhibitors of cell proliferation and survival signaling, bisphosphonates, aromatase inhibitors, siRNA therapeutics, γ -secretase inhibitors, agents that interfere with receptor tyrosine kinases (RTKs) and agents that interfere with cell cycle checkpoints. The instant compounds are particularly useful when co-administered with radiation therapy.

“Estrogen receptor modulators” refers to compounds that interfere with or inhibit the binding of estrogen to the receptor, regardless of mechanism. Examples of estrogen receptor modulators include, but are not limited to, tamoxifen, raloxifene, idoxifene, LY353381, LY117081, toremifene, fulvestrant, 4-[7-(2,2-dimethyl-1-oxopropoxy-4-methyl-2-[4-[2-(1-piperidinyl)ethoxy]phenyl]-2H-1-benzopyran-3-yl)]-phenyl-2,2-dimethylpropanoate, 4,4'-dihydroxybenzophenone-2,4-dinitrophenyl-hydrazone, and SH646.

“Androgen receptor modulators” refers to compounds which interfere or inhibit the binding of androgens to the receptor, regardless of mechanism. Examples of androgen receptor modulators include finasteride and other 5 α -reductase inhibitors, nilutamide, flutamide, bicalutamide, liarozole, and abiraterone acetate.

“Retinoid receptor modulators” refers to compounds which interfere or inhibit the binding of retinoids to the receptor, regardless of mechanism. Examples of such retinoid receptor modulators include bexarotene, tretinoin, 13-cis-retinoic acid, 9-cis-retinoic acid, α -difluoromethylornithine, ILX23-7553, trans-N-(4'-hydroxyphenyl) retinamide, and N-4-carboxyphenyl retinamide.

“Cytotoxic/cytostatic agents” refer to compounds which cause cell death or inhibit cell proliferation primarily by interfering directly with the cell's functioning or inhibit or interfere with cell myosis, including alkylating agents, tumor necrosis factors, intercalators, hypoxia activatable compounds, microtubule inhibitors/microtubule-stabilizing agents, inhibitors of mitotic kinesins, histone deacetylase inhibitors, inhibitors of kinases involved in mitotic progression, inhibitors of kinases involved in growth factor and cytokine signal transduction pathways, antimetabolites, biological response modifiers, hormonal/anti-hormonal therapeutic agents, haematopoietic growth factors, monoclonal antibody targeted therapeutic agents, topoisomerase inhibitors, proteasome inhibitors, ubiquitin ligase inhibitors, and aurora kinase inhibitors.

Examples of cytotoxic/cytostatic agents include, but are not limited to, sertenef, cachectin, ifosfamide, tasonermin, lonidamine, carboplatin, altretamine, prednimustine, dibromodulcitol, ranimustine, fotemustine, nedaplatin, oxaliplatin, temozolomide, heptaplatin, estramustine, improsulfan tosilate, trofosfamide, nimustine, dibrospidium chloride, pumitepa, lobaplatin, satraplatin, proflomycin, cisplatin, irofulven, dexifosfamide, cis-aminedichloro(2-methyl-pyridine)platinum, benzylguanine, glufosfamide, GPX100, (trans, trans, trans)-bis-mu-(hexane-1,6-diamine)-mu-[diamine-platinum(II)]bis[diamine(chloro)platinum (II)]tetrachloride, diarizidinylspermine, arsenic trioxide, 1-(11-dodecylamino-10-hydroxyundecyl)-3,7-dimethylxanthine, zorubicin, idarubicin, daunorubicin, bisantrene, mitoxantrone, pirarubicin, pinafide, valrubicin, amrubicin, antineoplaston, 3'-deamino-3'-morpholino-13-deoxy-10-hydroxycarminomycin, annamycin, galarubicin, elinafide, MEN10755, 4-demethoxy-3-deamino-3-aziridinyl-4-methylsulphonyl-daunorubicin (see WO 00/50032), Raf kinase inhibitors (such as Bay43-9006) and mTOR inhibitors (such as Wyeth's CCI-779).

An example of a hypoxia activatable compound is tirapazamine.

Examples of proteasome inhibitors include but are not limited to lactacystin and MLN-341 (Velcade).

Examples of microtubule inhibitors/microtubule-stabilising agents include paclitaxel, vindesine sulfate, 3',4'-didehydro-4'-deoxy-8'-norvincaloblastine, docetaxol, rhizoxin, dolastatin, mivobulin isethionate, auristatin, cemadotin, RPR109881, BMS184476, vinflunine, cryptophycin, 2,3,4,5,6-pentafluoro-N-(3-fluoro-4-methoxyphenyl) benzene sulfonamide, anhydrovinblastine, N,N-dimethyl-L-valyl-L-valyl-N-methyl-L-valyl-L-prolyl-L-proline-t-butylamide, TDX258, the epothilones (see for example U.S. Pat. Nos. 6,284,781 and 6,288,237) and BMS188797. In an embodiment the epothilones are not included in the microtubule inhibitors/microtubule-stabilising agents.

Some examples of topoisomerase inhibitors are topotecan, hycaptamine, irinotecan, rubitecan, 6-ethoxypropionyl-3',4'-O-exo-benzylidene-chartreusin, 9-methoxy-N,N-dimethyl-5-nitropyrzolo[3,4,5-kl]acridine-2-(6H) propanamine, 1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4':b,7]-indolizino[1,2b]quinoline-10,13(9H,15H)dione, lurtotecan, 7-[2-(N-isopropylamino)ethyl]-(20S)camptothecin, BNP1350, BNPII 100, BN80915, BN80942, etoposide phosphate, teniposide, sobuzoxane, 2'-dimethylamino-2'-deoxy-etoposide, GL331, N-[2-(dimethylamino)ethyl]-9-hydroxy-5,6-dimethyl-6H-pyrido[4,3-b]carbazole-1-carboxamide, asulacrane, (5a, 5aB, 8aa,9b)-9-[2-[N-[2-(dimethylamino)ethyl]-N-methylamino]ethyl]-5-[4-hydroxy-3,5-dimethoxyphenyl]-5,5a,6,8,8a,9-hexahydrofuro(3',4':6,7)naphtho(2,3-d)-1,3-dioxol-6-one, 2,3-(methylenedioxy)-5-methyl-7-hydroxy-8-methoxybenzo[c]-phenanthridinium, 6,9-bis[(2-aminoethyl)amino]benzo[g]isoguinoline-5,10-dione, 5-(3-aminopropylamino)-7,10-dihydroxy-2-(2-hydroxyethylaminomethyl)-6H-pyrazolo[4,5,1-de]acridin-6-one, N-[1-[2(diethylamino)ethylamino]-7-methoxy-9-oxo-9H-thioxanthen-4-ylmethyl]formamide, N-(2-

(dimethylamino)ethyl)acridine-4-carboxamide, 6-[[2-(dimethylamino)ethyl]amino]-3-hydroxy-7H-indeno[2,1-c] quinolin-7-one, and dimesna.

Examples of inhibitors of mitotic kinesins, and in particular the human mitotic kinesin KSP, are described in Publications WO03/039460, WO03/050064, WO03/050122, WO03/049527, WO03/049679, WO03/049678, WO04/039774, WO03/079973, WO03/099211, WO03/105855, WO03/106417, WO04/037171, WO04/058148, WO04/058700, WO04/126699, WO05/018638, WO05/019206, WO05/019205, WO05/018547, WO05/017190, US2005/0176776. In an embodiment inhibitors of mitotic kinesins include, but are not limited to inhibitors of KSP, inhibitors of MKLP1, inhibitors of CENP-E, inhibitors of MCAK and inhibitors of Rab6-KIFL.

Examples of "histone deacetylase inhibitors" include, but are not limited to, SAHA, TSA, oxamflatin, PXD101, MG98 and scriptaid. Further reference to other histone deacetylase inhibitors may be found in the following manuscript; Miller, T.A. et al. *J. Med. Chem.* 46(24):5097-5116 (2003).

"Inhibitors of kinases involved in mitotic progression" include, but are not limited to, inhibitors of aurora kinase, inhibitors of Polo-like kinases (PLK; in particular inhibitors of PLK-1), inhibitors of bub-1 and inhibitors of bub-R1. An example of an "aurora kinase inhibitor" is VX-680.

"Antiproliferative agents" includes antisense RNA and DNA oligonucleotides such as G3139, ODN698, RVASKRAS, GEM231, and INX3001, and antimetabolites such as enocitabine, carmofur, tegafur, pentostatin, doxifluridine, trimetrexate, fludarabine, capecitabine, galocitabine, cytarabine ocfosfate, fosteabine sodium hydrate, raltitrexed, paltitrexid, emitofur, tiazofurin, decitabine, nolatrexed, pemetrexed, nelzarabine, 2'-deoxy-2'-methylidenecytidine, 2'-fluoromethylene-2'-deoxycytidine, N-[5-(2,3-dihydro-benzofuryl)sulfonyl]-N'-(3,4-dichlorophenyl)urea, N6-[4-deoxy-4-[N2-[2(E),4(E)-tetradecadienoyl]glycylamino]-L-glycero-B-L-manno-heptopyranosyl]adenine, aplidine, ecteinascidin, troxacitabine, 4-[2-amino-4-oxo-4,6,7,8-tetrahydro-3H-pyrimidino[5,4-b][1,4]thiazin-6-yl-(S)-ethyl]-2,5-thienoyl-L-glutamic acid, aminopterin, 5-fluorouracil, alanosine, 11-acetyl-8-(carbamoyloxymethyl)-4-formyl-6-methoxy-14-oxa-1,11-diazatetracyclo(7.4.1.0.0)-tetradeca-2,4,6-trien-9-yl acetic acid ester, swainsonine, lometrexol, dexrazoxane, methioninase, 2'-cyano-2'-deoxy-N4-palmitoyl-1-B-D-arabino furanosyl cytosine, 3-aminopyridine-2-carboxaldehyde thiosemicarbazone and trastuzumab.

Examples of monoclonal antibody targeted therapeutic agents include those therapeutic agents which have cytotoxic agents or radioisotopes attached to a cancer cell specific or target cell specific monoclonal antibody. Examples include Bexxar.

"HMG-CoA reductase inhibitors" refers to inhibitors of 3-hydroxy-3-methylglutaryl-CoA reductase. Examples of HMG-CoA reductase inhibitors that may be used include but are not limited to lovastatin (MEVACOR®; see U.S. Patent Nos. 4,231,938,

4,294,926 and 4,319,039), simvastatin (ZOCOR®; see U.S. Patent Nos. 4,444,784, 4,820,850 and 4,916,239), pravastatin (PRAVACHOL®; see U.S. Patent Nos. 4,346,227, 4,537,859, 4,410,629, 5,030,447 and 5,180,589), fluvastatin (LESCOL®; see U.S. Patent Nos. 5,354,772, 4,911,165, 4,929,437, 5,189,164, 5,118,853, 5,290,946 and 5,356,896), atorvastatin (LIPITOR®; see U.S. Patent Nos. 5,273,995, 4,681,893, 5,489,691 and 5,342,952) and cerivastatin (also known as rivastatin and BAYCHOL®; see US Patent No. 5,177,080). The structural formulas of these and additional HMG-CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", *Chemistry & Industry*, pp. 85-89 (5 February 1996) and US Patent Nos. 4,782,084 and 4,885,314. The term HMG-CoA reductase inhibitor as used herein includes all pharmaceutically acceptable lactone and open-acid forms (i.e., where the lactone ring is opened to form the free acid) as well as salt and ester forms of compounds which have HMG-CoA reductase inhibitory activity, and therefor the use of such salts, esters, open-acid and lactone forms is included within the scope of this invention.

"Prenyl-protein transferase inhibitor" refers to a compound which inhibits any one or any combination of the prenyl-protein transferase enzymes, including farnesyl-protein transferase (FPTase), geranylgeranyl-protein transferase type I (GGPTase-I), and geranylgeranyl-protein transferase type-II (GGPTase-II, also called Rab GGPTase).

Examples of prenyl-protein transferase inhibitors can be found in the following publications and patents: WO 96/30343, WO 97/18813, WO 97/21701, WO 97/23478, WO 97/38665, WO 98/28980, WO 98/29119, WO 95/32987, U.S. Patent No. 5,420,245, U.S. Patent No. 5,523,430, U.S. Patent No. 5,532,359, U.S. Patent No. 5,510,510, U.S. Patent No. 5,589,485, U.S. Patent No. 5,602,098, European Patent Publ. 0 618 221, European Patent Publ. 0 675 112, European Patent Publ. 0 604 181, European Patent Publ. 0 696 593, WO 94/19357, WO 95/08542, WO 95/11917, WO 95/12612, WO 95/12572, WO 95/10514, U.S. Patent No. 5,661,152, WO 95/10515, WO 95/10516, WO 95/24612, WO 95/34535, WO 95/25086, WO 96/05529, WO 96/06138, WO 96/06193, WO 96/16443, WO 96/21701, WO 96/21456, WO 96/22278, WO 96/24611, WO 96/24612, WO 96/05168, WO 96/05169, WO 96/00736, U.S. Patent No. 5,571,792, WO 96/17861, WO 96/33159, WO 96/34850, WO 96/34851, WO 96/30017, WO 96/30018, WO 96/30362, WO 96/30363, WO 96/31111, WO 96/31477, WO 96/31478, WO 96/31501, WO 97/00252, WO 97/03047, WO 97/03050, WO 97/04785, WO 97/02920, WO 97/17070, WO 97/23478, WO 97/26246, WO 97/30053, WO 97/44350, WO 98/02436, and U.S. Patent No. 5,532,359. For an example of the role of a prenyl-protein transferase inhibitor on angiogenesis see *European J. of Cancer*, Vol. 35, No. 9, pp.1394-1401 (1999).

"Angiogenesis inhibitors" refers to compounds that inhibit the formation of new blood vessels, regardless of mechanism. Examples of angiogenesis inhibitors include, but are not limited to, tyrosine kinase inhibitors, such as inhibitors of the tyrosine kinase receptors Flt-1 (VEGFR1) and Flk-1/KDR (VEGFR2), inhibitors of epidermal-derived, fibroblast-derived, or

platelet derived growth factors, MMP (matrix metalloprotease) inhibitors, integrin blockers, interferon- α , interleukin-12, pentosan polysulfate, cyclooxygenase inhibitors, including nonsteroidal anti-inflammatories (NSAIDs) like aspirin and ibuprofen as well as selective cyclooxygenase-2 inhibitors like celecoxib and rofecoxib (*PNAS*, Vol. 89, p. 7384 (1992); *JNCI*, Vol. 69, p. 475 (1982); *Arch. Ophthalmol.*, Vol. 108, p.573 (1990); *Anat. Rec.*, Vol. 238, p. 68 (1994); *FEBS Letters*, Vol. 372, p. 83 (1995); *Clin. Orthop.* Vol. 313, p. 76 (1995); *J. Mol. Endocrinol.*, Vol. 16, p.107 (1996); *Jpn. J. Pharmacol.*, Vol. 75, p. 105 (1997); *Cancer Res.*, Vol. 57, p. 1625 (1997); *Cell*, Vol. 93, p. 705 (1998); *Intl. J. Mol. Med.*, Vol. 2, p. 715 (1998); *J. Biol. Chem.*, Vol. 274, p. 9116 (1999)), steroidal anti-inflammatories (such as corticosteroids, mineralocorticoids, dexamethasone, prednisone, prednisolone, methylpred, betamethasone), carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl-fumagillol, thalidomide, angiostatin, troponin-1, angiotensin II antagonists (see Fernandez et al., *J. Lab. Clin. Med.* 105:141-145 (1985)), and antibodies to VEGF (see, *Nature Biotechnology*, Vol. 17, pp.963-968 (October 1999); Kim et al., *Nature*, 362, 841-844 (1993); WO 00/44777; and WO 00/61186).

Other therapeutic agents that modulate or inhibit angiogenesis and may also be used in combination with the compounds of the instant invention include agents that modulate or inhibit the coagulation and fibrinolysis systems (see review in *Clin. Chem. La. Med.* 38:679-692 (2000)). Examples of such agents that modulate or inhibit the coagulation and fibrinolysis pathways include, but are not limited to, heparin (see *Thromb. Haemost.* 80:10-23 (1998)), low molecular weight heparins and carboxypeptidase U inhibitors (also known as inhibitors of active thrombin activatable fibrinolysis inhibitor [TAFIa]) (see *Thrombosis Res.* 101:329-354 (2001)). TAFIa inhibitors have been described in U.S. Ser. Nos. 60/310,927 (filed August 8, 2001) and 60/349,925 (filed January 18, 2002).

“Agents that interfere with cell cycle checkpoints” refer to compounds that inhibit protein kinases that transduce cell cycle checkpoint signals, thereby sensitizing the cancer cell to DNA damaging agents. Such agents include inhibitors of ATR, ATM, the CHK1 and CHK2 kinases and cdk and cdc kinase inhibitors and are specifically exemplified by 7-hydroxystaurosporin, flavopiridol, CYC202 (Cyclacel) and BMS-387032.

“Agents that interfere with receptor tyrosine kinases (RTKs)” refer to compounds that inhibit RTKs and therefore mechanisms involved in oncogenesis and tumor progression. Such agents include inhibitors of c-Kit, Eph, PDGF, Flt3 and c-Met. Further agents include inhibitors of RTKs as described by Bume-Jensen and Hunter, *Nature*, 411:355-365, 2001.

“Inhibitors of cell proliferation and survival signalling pathway” refer to compounds that inhibit signal transduction cascades downstream of cell surface receptors. Such agents include inhibitors of serine/threonine kinases (including but not limited to inhibitors of Akt such as described in WO 02/083064, WO 02/083139, WO 02/083140, US 2004-0116432, WO 02/083138, US 2004-0102360, WO 03/086404, WO 03/086279, WO 03/086394, WO

03/084473, WO 03/086403, WO 2004/041162, WO 2004/096131, WO 2004/096129, WO 2004/096135, WO 2004/096130, WO 2005/100356, WO 2005/100344, US 2005/029941, US 2005/44294, US 2005/43361, 60/734188, 60/652737, 60/670469), inhibitors of Raf kinase (for example BAY-43-9006), inhibitors of MEK (for example CI-1040 and PD-098059), inhibitors of mTOR (for example Wyeth CCI-779), and inhibitors of PI3K (for example LY294002).

As described above, the combinations with NSAID's are directed to the use of NSAID's which are potent COX-2 inhibiting agents. For purposes of this specification an NSAID is potent if it possesses an IC₅₀ for the inhibition of COX-2 of 1 μM or less as measured by cell or microsomal assays.

The invention also encompasses combinations with NSAID's which are selective COX-2 inhibitors. For purposes of this specification NSAID's which are selective inhibitors of COX-2 are defined as those which possess a specificity for inhibiting COX-2 over COX-1 of at least 100 fold as measured by the ratio of IC₅₀ for COX-2 over IC₅₀ for COX-1 evaluated by cell or microsomal assays. Such compounds include, but are not limited to those disclosed in U.S. Patent 5,474,995, U.S. Patent 5,861,419, U.S. Patent 6,001,843, U.S. Patent 6,020,343, U.S. Patent 5,409,944, U.S. Patent 5,436,265, U.S. Patent 5,536,752, U.S. Patent 5,550,142, U.S. Patent 5,604,260, U.S. 5,698,584, U.S. Patent 5,710,140, WO 94/15932, U.S. Patent 5,344,991, U.S. Patent 5,134,142, U.S. Patent 5,380,738, U.S. Patent 5,393,790, U.S. Patent 5,466,823, U.S. Patent 5,633,272 and U.S. Patent 5,932,598, all of which are hereby incorporated by reference.

Inhibitors of COX-2 that are particularly useful in the instant method of treatment are: 3-phenyl-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone; and 5-chloro-3-(4-methylsulfonyl)phenyl-2-(2-methyl-5-pyridinyl)pyridine; or a pharmaceutically acceptable salt thereof.

Compounds that have been described as specific inhibitors of COX-2 and are therefore useful in the present invention include, but are not limited to, the following: parecoxib, BEXTRA® and CELEBREX® or a pharmaceutically acceptable salt thereof.

Other examples of angiogenesis inhibitors include, but are not limited to, endostatin, ukrain, ranpirnase, IM862, 5-methoxy-4-[2-methyl-3-(3-methyl-2-butenyl)oxiranyl]-1-oxaspiro[2,5]oct-6-yl(chloroacetyl)carbamate, acetyldinanaline, 5-amino-1-[[3,5-dichloro-4-(4-chlorobenzoyl)phenyl]methyl]-1*H*-1,2,3-triazole-4-carboxamide, CM101, squalamine, combretastatin, RPI4610, NX31838, sulfated mannopentase phosphate, 7,7-(carbonyl-bis[imino-N-methyl-4,2-pyrrolocarbonylimino[N-methyl-4,2-pyrrole]-carbonylimino]-bis-(1,3-naphthalene disulfonate), and 3-[(2,4-dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416).

As used above, "integrin blockers" refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the α_vβ₃ integrin, to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the α_vβ₅ integrin, to compounds which antagonize, inhibit or counteract binding of a physiological ligand to both the α_vβ₃ integrin and the α_vβ₅ integrin, and to compounds which

antagonize, inhibit or counteract the activity of the particular integrin(s) expressed on capillary endothelial cells. The term also refers to antagonists of the $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins. The term also refers to antagonists of any combination of $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins.

5 Some specific examples of tyrosine kinase inhibitors include N-(trifluoromethylphenyl)-5-methylisoxazol-4-carboxamide, 3-[(2,4-dimethylpyrrol-5-yl)methylidenyl]indolin-2-one, 17-(allylamino)-17-demethoxygeldanamycin, 4-(3-chloro-4-fluorophenylamino)-7-methoxy-6-[3-(4-morpholinyl)propoxyl]quinazoline, N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, BIBX1382, 2,3,9,10,11,12-hexahydro-10-(hydroxymethyl)-10-hydroxy-9-methyl-9,12-epoxy-1H-diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-i][1,6]benzodiazocin-1-one, SH268, genistein, STI571, CEP2563, 4-(3-chlorophenylamino)-5,6-dimethyl-7H-pyrrolo[2,3-d]pyrimidinemethane sulfonate, 4-(3-bromo-4-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, 4-(4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, SU6668, STI571A, N-4-chlorophenyl-4-(4-pyridylmethyl)-1-phthalazinamine, and EMD121974.

15 Combinations with compounds other than anti-cancer compounds are also encompassed in the instant methods. For example, combinations of the instantly claimed compounds with PPAR- γ (i.e., PPAR-gamma) agonists and PPAR- δ (i.e., PPAR-delta) agonists are useful in the treatment of certain malignancies. PPAR- γ and PPAR- δ are the nuclear peroxisome proliferator-activated receptors γ and δ . The expression of PPAR- γ on endothelial
20 cells and its involvement in angiogenesis has been reported in the literature (see *J. Cardiovasc. Pharmacol.* 1998; 31:909-913; *J. Biol. Chem.* 1999;274:9116-9121; *Invest. Ophthalmol Vis. Sci.* 2000; 41:2309-2317). More recently, PPAR- γ agonists have been shown to inhibit the angiogenic response to VEGF in vitro; both troglitazone and rosiglitazone maleate inhibit the development of retinal neovascularization in mice. (*Arch. Ophthalmol.* 2001; 119:709-717).
25 Examples of PPAR- γ agonists and PPAR- γ/α agonists include, but are not limited to, thiazolidinediones (such as DRF2725, CS-011, troglitazone, rosiglitazone, and pioglitazone), fenofibrate, gemfibrozil, clofibrate, GW2570, SB219994, AR-H039242, JTT-501, MCC-555, GW2331, GW409544, NN2344, KRP297, NP0110, DRF4158, NN622, GI262570, PNU182716, DRF552926, 2-[(5,7-dipropyl-3-trifluoromethyl-1,2-benzisoxazol-6-yl)oxy]-2-methylpropionic
30 acid (disclosed in USSN 09/782,856), and 2(R)-7-(3-(2-chloro-4-(4-fluorophenoxy)phenoxy)propoxy)-2-ethylchromane-2-carboxylic acid (disclosed in USSN 60/235,708 and 60/244,697).

Another embodiment of the instant invention is the use of the presently disclosed compounds in combination with gene therapy for the treatment of cancer. For an overview of
35 genetic strategies to treating cancer see Hall et al (*Am. J. Hum. Genet.* 61:785-789, 1997) and Kufe et al (Cancer Medicine, 5th Ed, pp 876-889, BC Decker, Hamilton 2000). Gene therapy can be used to deliver any tumor suppressing gene. Examples of such genes include, but are not limited to, p53, which can be delivered via recombinant virus-mediated gene transfer (see U.S.

Patent No. 6,069,134, for example), a uPA/uPAR antagonist ("Adenovirus-Mediated Delivery of a uPA/uPAR Antagonist Suppresses Angiogenesis-Dependent Tumor Growth and Dissemination in Mice," *Gene Therapy*, August 1998;5(8):1105-13), and interferon gamma (*J. Immunol.* 2000;164:217-222).

5 The compounds of the instant invention may also be administered in combination with an inhibitor of inherent multidrug resistance (MDR), in particular MDR associated with high levels of expression of transporter proteins. Such MDR inhibitors include inhibitors of p-glycoprotein (P-gp), such as LY335979, XR9576, OC144-093, R101922, VX853 and PSC833 (valspodar).

10 A compound of the present invention may be employed in conjunction with anti-emetic agents to treat nausea or emesis, including acute, delayed, late-phase, and anticipatory emesis, which may result from the use of a compound of the present invention, alone or with radiation therapy. For the prevention or treatment of emesis, a compound of the present invention may be used in conjunction with other anti-emetic agents, especially neurokinin-1
15 receptor antagonists, 5HT3 receptor antagonists, such as ondansetron, granisetron, tropisetron, and zatisetron, GABAB receptor agonists, such as baclofen, a corticosteroid such as Decadron (dexamethasone), Kenalog, Aristocort, Nasalide, Preferid, Benecorten or others such as disclosed in U.S. Patent Nos. 2,789,118, 2,990,401, 3,048,581, 3,126,375, 3,929,768, 3,996,359, 3,928,326 and 3,749,712, an antidopaminergic, such as the phenothiazines (for example prochlorperazine,
20 fluphenazine, thioridazine and mesoridazine), metoclopramide or dronabinol. In another embodiment, conjunctive therapy with an anti-emesis agent selected from a neurokinin-1 receptor antagonist, a 5HT3 receptor antagonist and a corticosteroid is disclosed for the treatment or prevention of emesis that may result upon administration of the instant compounds.

Neurokinin-1 receptor antagonists of use in conjunction with the compounds of
25 the present invention are fully described, for example, in U.S. Patent Nos. 5,162,339, 5,232,929, 5,242,930, 5,373,003, 5,387,595, 5,459,270, 5,494,926, 5,496,833, 5,637,699, 5,719,147; European Patent Publication Nos. EP 0 360 390, 0 394 989, 0 428 434, 0 429 366, 0 430 771, 0 436 334, 0 443 132, 0 482 539, 0 498 069, 0 499 313, 0 512 901, 0 512 902, 0 514 273, 0 514 274, 0 514 275, 0 514 276, 0 515 681, 0 517 589, 0 520 555, 0 522 808, 0 528 495, 0 532 456, 0
30 533 280, 0 536 817, 0 545 478, 0 558 156, 0 577 394, 0 585 913, 0 590 152, 0 599 538, 0 610 793, 0 634 402, 0 686 629, 0 693 489, 0 694 535, 0 699 655, 0 699 674, 0 707 006, 0 708 101, 0 709 375, 0 709 376, 0 714 891, 0 723 959, 0 733 632 and 0 776 893; PCT International Patent Publication Nos. WO 90/05525, 90/05729, 91/09844, 91/18899, 92/01688, 92/06079, 92/12151, 92/15585, 92/17449, 92/20661, 92/20676, 92/21677, 92/22569, 93/00330, 93/00331, 93/01159,
35 93/01165, 93/01169, 93/01170, 93/06099, 93/09116, 93/10073, 93/14084, 93/14113, 93/18023, 93/19064, 93/21155, 93/21181, 93/23380, 93/24465, 94/00440, 94/01402, 94/02461, 94/02595, 94/03429, 94/03445, 94/04494, 94/04496, 94/05625, 94/07843, 94/08997, 94/10165, 94/10167, 94/10168, 94/10170, 94/11368, 94/13639, 94/13663, 94/14767, 94/15903, 94/19320, 94/19323,

94/20500, 94/26735, 94/26740, 94/29309, 95/02595, 95/04040, 95/04042, 95/06645, 95/07886, 95/07908, 95/08549, 95/11880, 95/14017, 95/15311, 95/16679, 95/17382, 95/18124, 95/18129, 95/19344, 95/20575, 95/21819, 95/22525, 95/23798, 95/26338, 95/28418, 95/30674, 95/30687, 95/33744, 96/05181, 96/05193, 96/05203, 96/06094, 96/07649, 96/10562, 96/16939, 96/18643, 96/20197, 96/21661, 96/29304, 96/29317, 96/29326, 96/29328, 96/31214, 96/32385, 96/37489, 97/01553, 97/01554, 97/03066, 97/08144, 97/14671, 97/17362, 97/18206, 97/19084, 97/19942 and 97/21702; and in British Patent Publication Nos. 2 266 529, 2 268 931, 2 269 170, 2 269 590, 2 271 774, 2 292 144, 2 293 168, 2 293 169, and 2 302 689. The preparation of such compounds is fully described in the aforementioned patents and publications, which are incorporated herein by reference.

In an embodiment, the neurokinin-1 receptor antagonist for use in conjunction with the compounds of the present invention is selected from: 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo-1H,4H-1,2,4-triazolo)methyl)morpholine, or a pharmaceutically acceptable salt thereof, which is described in U.S. Patent No. 5,719,147.

A compound of the instant invention may also be administered with an agent useful in the treatment of anemia. Such an anemia treatment agent is, for example, a continuous erythropoiesis receptor activator (such as epoetin alfa).

A compound of the instant invention may also be administered with an agent useful in the treatment of neutropenia. Such a neutropenia treatment agent is, for example, a hematopoietic growth factor which regulates the production and function of neutrophils such as a human granulocyte colony stimulating factor, (G-CSF). Examples of a G-CSF include filgrastim.

A compound of the instant invention may also be administered with an immunologic-enhancing drug, such as levamisole, isoprinosine and Zadaxin.

A compound of the instant invention may also be useful for treating or preventing cancer in combination with P450 inhibitors including: xenobiotics, quinidine, tyramine, ketoconazole, testosterone, quinine, methyrapone, caffeine, phenelzine, doxorubicin, troleandomycin, cyclobenzaprine, erythromycin, cocaine, furafyline, cimetidine, dextromethorphan, ritonavir, indinavir, amprenavir, diltiazem, terfenadine, verapamil, cortisol, itraconazole, mibefradil, nefazodone and nelfinavir.

A compound of the instant invention may also be useful for treating or preventing cancer in combination with Pgp and/or BCRP inhibitors including: cyclosporin A, PSC833, GF120918, cremophorEL, fumitremorgin C, Ko132, Ko134, Iressa, Imatinib mesylate, EKI-785, CI1033, novobiocin, diethylstilbestrol, tamoxifen, reserpine, VX-710, tryprostatin A, flavonoids, ritonavir, saquinavir, nelfinavir, omeprazole, quinidine, verapamil, terfenadine, ketoconazole, nifedipine, FK506, amiodarone, XR9576, indinavir, amprenavir, cortisol, testosterone, LY335979, OC144-093, erythromycin, vincristine, digoxin and talinolol.

A compound of the instant invention may also be useful for treating or preventing cancer, including bone cancer, in combination with bisphosphonates (understood to include bisphosphonates, diphosphonates, bisphosphonic acids and diphosphonic acids). Examples of bisphosphonates include but are not limited to: etidronate (Didronel), pamidronate (Aredia),
5 alendronate (Fosamax), risedronate (Actonel), zoledronate (Zometa), ibandronate (Boniva), incadronate or cimadronate, clodronate, EB-1053, minodronate, neridronate, piridronate and tiludronate including any and all pharmaceutically acceptable salts, derivatives, hydrates and mixtures thereof.

A compound of the instant invention may also be useful for treating or preventing
10 breast cancer in combination with aromatase inhibitors. Examples of aromatase inhibitors include but are not limited to: anastrozole, letrozole and exemestane.

A compound of the instant invention may also be useful for treating or preventing cancer in combination with siRNA therapeutics.

The compounds of the instant invention may also be administered in combination
15 with γ -secretase inhibitors and/or inhibitors of NOTCH signaling. Such inhibitors include compounds described in WO 01/90084, WO 02/30912, WO 01/70677, WO 03/013506, WO 02/36555, WO 03/093252, WO 03/093264, WO 03/093251, WO 03/093253, WO 2004/039800, WO 2004/039370, WO 2005/030731, WO 2005/014553, USSN 10/957,251, WO 2004/089911, WO 02/081435, WO 02/081433, WO 03/018543, WO 2004/031137, WO 2004/031139, WO
20 2004/031138, WO 2004/101538, WO 2004/101539 and WO 02/47671 (including LY-450139).

Inhibitors of Akt, as disclosed in the following publications; WO 02/083064, WO 02/083139, WO 02/083140, US 2004-0116432, WO 02/083138, US 2004-0102360, WO 03/086404, WO 03/086279, WO 03/086394, WO 03/084473, WO 03/086403, WO 2004/041162, WO 2004/096131, WO 2004/096129, WO 2004/096135, WO 2004/096130, WO
25 2005/100356, WO 2005/100344, US 2005/029941, US 2005/44294, US 2005/43361, 60/734188, 60/652737, 60/670469, and including compounds of the instant invention, are also useful in combination with potassium salts, magnesium salts, beta-blockers (such as atenolol) and endothelin-a (ETa) antagonists with the goal of maintaining cardiovascular homeostasis.

Inhibitors of Akt, as disclosed in the following publications; WO 02/083064, WO
30 02/083139, WO 02/083140, US 2004-0116432, WO 02/083138, US 2004-0102360, WO 03/086404, WO 03/086279, WO 03/086394, WO 03/084473, WO 03/086403, WO 2004/041162, WO 2004/096131, WO 2004/096129, WO 2004/096135, WO 2004/096130, WO 2005/100356, WO 2005/100344, US 2005/029941, US 2005/44294, US 2005/43361, 60/734188, 60/652737, 60/670469, and including compounds of the instant invention, are also useful in
35 combination with insulin, insulin secretagogues, PPAR-gamma agonists, metformin, somatostatin receptor agonists such as octreotide, DPP4 inhibitors, sulfonylureas and alpha-glucosidase inhibitors with the goal of maintaining glucose homeostasis.

A compound of the instant invention may also be useful for treating or preventing cancer in combination with PARP inhibitors.

A compound of the instant invention may also be useful for treating cancer in combination with the following therapeutic agents: abarelix (Plenaxis depot®); aldesleukin (Prokine®); Aldesleukin (Proleukin®); Alemtuzumab (Campath®); alitretinoin (Panretin®);
 5 allopurinol (Zyloprim®); altretamine (Hexalen®); amifostine (Ethyol®); anastrozole (Arimidex®); arsenic trioxide (Trisenox®); asparaginase (Elspar®); azacitidine (Vidaza®); bevacuzimab (Avastin®); bexarotene capsules (Targretin®); bexarotene gel (Targretin®); bleomycin (Blenoxane®); bortezomib (Velcade®); busulfan intravenous (Busulfex®); busulfan
 10 oral (Myleran®); calusterone (Methosarb®); capecitabine (Xeloda®); carboplatin (Paraplatin®); carmustine (BCNU®, BiCNU®); carmustine (Gliadel®); carmustine with Polifeprosan 20 Implant (Gliadel Wafer®); celecoxib (Celebrex®); cetuximab (Erbix®); chlorambucil (Leukeran®); cisplatin (Platinol®); cladribine (Leustatin®, 2-CdA®); clofarabine (Clolar®); cyclophosphamide (Cytosan®, Neosar®); cyclophosphamide (Cytosan Injection®);
 15 cyclophosphamide (Cytosan Tablet®); cytarabine (Cytosar-U®); cytarabine liposomal (DepoCyt®); dacarbazine (DTIC-Dome®); dactinomycin, actinomycin D (Cosmegen®); Darbepoetin alfa (Aranesp®); daunorubicin liposomal (DanuoXome®); daunorubicin, daunomycin (Daunorubicin®); daunorubicin, daunomycin (Cerubidine®); Denileukin diftitox (Ontak®); dexrazoxane (Zinecard®); docetaxel (Taxotere®); doxorubicin (Adriamycin PFS®);
 20 doxorubicin (Adriamycin®, Rubex®); doxorubicin (Adriamycin PFS Injection®); doxorubicin liposomal (Doxil®); dromostanolone propionate (dromostanolone®); dromostanolone propionate (masterone injection®); Elliott's B Solution (Elliott's B Solution®); epirubicin (Ellence®); Epoetin alfa (epogen®); erlotinib (Tarceva®); estramustine (Emcyt®); etoposide phosphate (Etopophos®); etoposide, VP-16 (Vepesid®); exemestane (Aromasin®); Filgrastim (Neupogen®); floxuridine (intraarterial) (FUDR®); fludarabine (Fludara®); fluorouracil, 5-FU (Adrucil®); fulvestrant (Faslodex®); gefitinib (Iressa®); gemcitabine (Gemzar®); gemtuzumab ozogamicin (Mylotarg®); goserelin acetate (Zoladex Implant®); goserelin acetate (Zoladex®); histrelin acetate (Histrelin implant®); hydroxyurea (Hydrea®); Ibritumomab Tiuxetan (Zevalin®); idarubicin (Idamycin®); ifosfamide (IFEX®); imatinib mesylate (Gleevec®);
 30 interferon alfa 2a (Roferon A®); Interferon alfa-2b (Intron A®); irinotecan (Camptosar®); lenalidomide (Revlimid®); letrozole (Femara®); leucovorin (Wellcovorin®, Leucovorin®); Leuprolide Acetate (Eligard®); levamisole (Ergamisol®); lomustine, CCNU (CeeBU®); meclorothamine, nitrogen mustard (Mustargen®); megestrol acetate (Megace®); melphalan, L-PAM (Alkeran®); mercaptopurine, 6-MP (Purinethol®); mesna (Mesnex®); mesna (Mesnex tabs®); methotrexate (Methotrexate®); methoxsalen (Uvadex®); mitomycin C (Mutamycin®); mitotane (Lysodren®); mitoxantrone (Novantrone®); nandrolone phenpropionate (Durabolin-50®); nelarabine (Arranon®); Nofetumomab (Verluma®); Oprelvekin (Neumega®); oxaliplatin (Eloxatin®); paclitaxel (Paxene®); paclitaxel (Taxol®); paclitaxel protein-bound particles

(Abraxane®); palifermin (Kepivance®); pamidronate (Aredia®); pegademase (Adagen (Pegademase Bovine®); pegaspargase (Oncaspar®); Pegfilgrastim (Neulasta®); pemetrexed disodium (Alimta®); pentostatin (Nipent®); pipobroman (Vercyte®); plicamycin, mithramycin (Mithracin®); porfimer sodium (Photofrin®); procarbazine (Matulane®); quinacrine (Atabrine®); Rasburicase (Elitek®); Rituximab (Rituxan®); sargramostim (Leukine®); Sargramostim (Prokine®); sorafenib (Nexavar®); streptozocin (Zanosar®); sunitinib maleate (Sutent®); talc (Sclerosol®); tamoxifen (Nolvadex®); temozolomide (Temodar®); teniposide, VM-26 (Vumon®); testolactone (Teslac®); thioguanine, 6-TG (Thioguanine®); thiotepa (Thioplex®); topotecan (Hycamtin®); toremifene (Fareston®); Tositumomab (Bexxar®); Tositumomab/I-131 tositumomab (Bexxar®); Trastuzumab (Herceptin®); tretinoin, ATRA (Vesanoid®); Uracil Mustard (Uracil Mustard Capsules®); valrubicin (Valstar®); vinblastine (Velban®); vincristine (Oncovin®); vinorelbine (Navelbine®); zoledronate (Zometa®) and vorinostat (Zolinza®).

The compounds of the instant invention are useful for treating cancer in combination with taxanes.

The compounds of the instant invention are useful for treating cancer in combination with docetaxel (Taxotere®).

The compounds of the instant invention are useful for treating cancer in combination with vorinostat (Zolinza®).

The compounds of the instant invention are useful for treating cancer in combination with the aurora kinase inhibitor, MK-0457.

The compounds of the instant invention are useful for treating cancer in combination with the mTor inhibitor, AP 23573.

The compounds of the instant invention are useful for treating cancer in combination with the IGF1R inhibitor, MK-0646.

The compounds of the instant invention are useful for treating cancer in combination with satraplatin.

The compounds of the instant invention are useful for treating cancer in combination with lapatinib (Tykerb®).

Thus, the scope of the instant invention encompasses the use of the instantly claimed compounds in combination with a second compound selected from: an estrogen receptor modulator, an androgen receptor modulator, a retinoid receptor modulator, a cytotoxic/cytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, PPAR- γ agonists, PPAR- δ agonists, an inhibitor of inherent multidrug resistance, an anti-emetic agent, an agent useful in the treatment of anemia, an agent useful in the treatment of neutropenia, an immunologic-enhancing drug, an inhibitor of cell proliferation and survival signaling, a bisphosphonate, an aromatase inhibitor, an siRNA therapeutic, γ -secretase inhibitors, agents that

interfere with receptor tyrosine kinases (RTKs), an agent that interferes with a cell cycle checkpoint and any of the therapeutic agents listed above.

The term "administration" and variants thereof (e.g., "administering" a compound) in reference to a compound of the invention means introducing the compound or a prodrug of the compound into the system of the animal in need of treatment. When a compound of the invention or prodrug thereof is provided in combination with one or more other active agents (e.g., a cytotoxic agent, etc.), "administration" and its variants are each understood to include concurrent and sequential introduction of the compound or prodrug thereof and other agents.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The term "therapeutically effective amount" as used herein means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

The term "treating cancer" or "treatment of cancer" refers to administration to a mammal afflicted with a cancerous condition and refers to an effect that alleviates the cancerous condition by killing the cancerous cells, but also to an effect that results in the inhibition of growth and/or metastasis of the cancer.

In an embodiment, the angiogenesis inhibitor to be used as the second compound is selected from a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP (matrix metalloprotease) inhibitor, an integrin blocker, interferon- α , interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl-fumagillol, thalidomide, angiostatin, troponin-1, or an antibody to VEGF. In an embodiment, the estrogen receptor modulator is tamoxifen or raloxifene.

Also included in the scope of the claims is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of the instant invention in combination with radiation therapy and/or in combination with a second compound selected from: an estrogen receptor modulator, an androgen receptor modulator, a retinoid receptor modulator, a cytotoxiccytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, PPAR- γ agonists, PPAR- δ agonists, an inhibitor of inherent multidrug resistance, an anti-emetic agent, an agent useful in the treatment of anemia, an agent useful in the treatment of neutropenia, an immunologic-enhancing drug, an inhibitor of cell proliferation and survival signaling, a bisphosphonate, an aromatase inhibitor, an

siRNA therapeutic, γ -secretase inhibitors, agents that interfere with receptor tyrosine kinases (RTKs), an agent that interferes with a cell cycle checkpoint and any of the therapeutic agents listed above.

And yet another embodiment of the invention is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of the instant invention in combination with paclitaxel or trastuzumab.

The invention further encompasses a method of treating or preventing cancer that comprises administering a therapeutically effective amount of a compound of the instant invention in combination with a COX-2 inhibitor.


The instant invention also includes a pharmaceutical composition useful for treating or preventing cancer that comprises a therapeutically effective amount of a compound of the instant invention and a second compound selected from: an estrogen receptor modulator, an androgen receptor modulator, a retinoid receptor modulator, a cytotoxic/cytostatic agent, an antiproliferative agent, a prenyl-protein transferase inhibitor, an HMG-CoA reductase inhibitor, an HIV protease inhibitor, a reverse transcriptase inhibitor, an angiogenesis inhibitor, a PPAR- γ agonist, a PPAR- δ agonist, an inhibitor of cell proliferation and survival signaling, a bisphosphonate, an aromatase inhibitor, an siRNA therapeutic, γ -secretase inhibitors, agents that interfere with receptor tyrosine kinases (RTKs), an agent that interferes with a cell cycle checkpoint and any of the therapeutic agents listed above.

All patents, publications and pending patent applications identified are hereby incorporated by reference.


Abbreviations used in the description of the chemistry and in the Examples that follow are: HPLC (high performance liquid chromatography) and TFA (trifluoroacetic acid).

The compounds of this invention may be prepared by employing reactions as shown in the following Reaction Schemes, in addition to other standard manipulations that are known in the literature or exemplified in the experimental procedures. The illustrative Reaction Schemes below, therefore, are not limited by the compounds listed or by any particular substituents employed for illustrative purposes. Substituent numbering as shown in the Reaction Schemes does not necessarily correlate to that used in the claims and often, for clarity, a single substituent is shown attached to the compound where multiple substituents are allowed under the definitions of Formula A hereinabove.

Compound (I) in Reaction Schemes I can be prepared by the Suzuki coupling reaction between the heterocyclic chloride and the corresponding benzylamine boronic ester (IV).

Reaction Scheme IReaction Scheme II


Compound (XV) in Reaction Schemes II can be prepared via the general synthetic methods and procedures found in WO2009/021992. The requisite starting material (XVI) can be prepared according to Scheme I. Key intermediates are in some cases commercially available or can be prepared according to literature procedures.



Reaction Scheme III


Compound (XVIII) in Reaction Schemes III can be prepared via the general synthetic methods and procedures found in WO2009/021992. The requisite intermediates are in some cases commercially available or can be prepared according to literature procedures.

5


EXAMPLES

Examples and schemes provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be further illustrative of the invention and do not limit the reasonable scope thereof.

10


EXAMPLE A-1

1-[4-(6-phenylimidazo[1,2-a]pyrimidin-7-yl)phenyl]cyclobutanamine (A-1)




To a mixture of **1-B** (6g, for preparation see WO2009/021992) and *tert*-butyl {1-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]cyclobutyl}carbamate **1-A** (6g, for preparation see WO2008/070016) in 210mL of 1,2-dimethoxyethane; dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (0.96g) and 42mL of a 10% w/w sodium carbonate solution is added. The resulting mixture is heated to 80°C under an inert gas atmosphere for 15h. The work up is performed by diluting the reaction mixture with water and dichloromethane, separating the phases and extraction of the aqueous phase with dichloromethane. The combined organic layers are dried over sodium sulphate and evaporated. The residue is suspended in ethyl acetate and the resulting mixture stirred for 2h at room temperature. The product is collected by filtration and treated with 30mL TFA in 30mL dichloromethane for 1 hr. The resulting mixture is concentrated and purified via reverse phase HPLC to afford the desired product.

EXAMPLE B-1




15 1-[4-(2-isopropyl-6-phenyl[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)phenyl]cyclobutanamine (**B-1**)




To a mixture of **2-B** (6g, for preparation see WO2009/021992) and *tert*-butyl {1-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]cyclobutyl}carbamate **1-A** (6g, for preparation see WO2008/070016) in 210mL of 1,2-dimethoxyethane; dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (0.96g) and 42mL of a 10% w/w sodium carbonate solution is added. The resulting mixture is heated to 80°C under an inert gas atmosphere for 15h. The work up is performed by diluting the reaction mixture with water and dichloromethane, separating the phases and extraction of the aqueous phase with dichloromethane. The combined organic layers are dried over sodium sulphate and evaporated. The residue is suspended in ethyl acetate and the resulting mixture stirred for 2h at room temperature. The product is collected by filtration and treated with 30mL TFA in 30mL

dichloromethane for 1 hr. The resulting mixture is concentrated and purified via reverse phase HPLC to afford the desired product.

EXAMPLE C-1



5 1-[4-(2-methyl-6-phenylpyrazolo[1,5-a]pyrimidin-5-yl)phenyl]cyclobutanamine (C-1)



To a mixture of **3-B** (6g, for preparation see WO2009/021992) and *tert*-butyl {1-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]cyclobutyl} carbamate **1-A** (6g, for preparation see WO2008/070016) in 210mL of 1,2-dimethoxyethane; dichloro[1,1'-






10 bis(diphenylphosphino)ferrocene]palladium (II) dichloromethane adduct (0.96g) and 42mL of a 10% w/w sodium carbonate solution is added. The resulting mixture is heated to 80°C under an inert gas atmosphere for 15h. The work up is performed by diluting the reaction mixture with water and dichloromethane, separating the phases and extraction of the aqueous phase with dichloromethane. The combined organic layers are dried over sodium sulphate and evaporated.








15 The residue is suspended in ethyl acetate and the resulting mixture stirred for 2h at room temperature. The product is collected by filtration and treated with 30mL TFA in 30mL dichloromethane for 1 hr. The resulting mixture is concentrated and purified via reverse phase HPLC to afford the desired product.

Compounds disclosed in Table 1 are prepared according to procedures described

20 herein and in WO2009/021992 using appropriate starting materials.

Table 1

Example	Structure	Name	Molecular Weight
D		1-[4-(3,6-diphenylimidazo[1,2-a]pyrimidin-7-yl)phenyl]cyclobutanamine	416.20
E		3-amino-3-[4-(3,6-diphenylimidazo[1,2-a]pyrimidin-7-yl)phenyl]-1-methylcyclobutanol	446.21
F		3-amino-1-methyl-3-[4-(2-methyl-3,6-diphenylimidazo[1,2-a]pyrimidin-7-yl)phenyl]cyclobutanol	460.23
G		1-[4-(2-methyl-3,6-diphenylimidazo[1,2-a]pyrimidin-7-yl)phenyl]cyclobutanamine	430.22
H		3-amino-1-methyl-3-[4-(2-methyl-6-phenylpyrazolo[1,5-a]pyrimidin-5-yl)phenyl]cyclobutanol	384.20


I		3-amino-1-methyl-3-[4-(6-phenylpyrazolo[1,5-a]pyrimidin-5-yl)phenyl]cyclobutanol	370.18
J		1-[4-(6-phenylpyrazolo[1,5-a]pyrimidin-5-yl)phenyl]cyclobutanamine	340.17
K		1-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)phenyl]cyclobutanamine	355.18
L		3-amino-1-methyl-3-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)phenyl]cyclobutanol	385.19
M		1-[4-(3,6-diphenylimidazo[1,2-a]pyrimidin-7-yl)phenyl]methanamine	376.17
N		1-[4-(6-phenylpyrazolo[1,5-a]pyrimidin-5-yl)phenyl]methanamine	300.14
O		1-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)phenyl]methanamine	315.15

OTHER EXAMPLES


Examples and schemes provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be further illustrative of the invention and do not limit the reasonable scope thereof.

5

{1-[4-(3-Dimethylamino-2-phenyl-acryloyl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester
(1)



10

Reaction Scheme 1StepA: [1-(4-Cyano-phenyl)-cyclobutyl]-carbamic acid tert-butyl ester (1-2)

A mixture of [1-(4-Chloro-phenyl)-cyclobutyl]-carbamic acid tert-butyl ester **1-1** (prepared according to **WO2008/070016**; 1.57 g, 5.6 mmol), Zn(CN)_2 (395 mg, 3.37 mmol), $\text{Pd}_2(\text{dba})_3$ (257 mg, 0.28 mmol), dppf (311 mg, 0.56 mmol), Zinc powder (219 mg, 3.37 mmol) and 20 mL of DMF was heated to 100.deg.C for 2 hrs under nitrogen atmosphere.. After cooling to room temperature, the mixture was poured into water and extracted with EA for three times. The combined EA layer was washed with water, brine and concentrated to give a residue, which was purified by silica gel chromatography eluting with EA/PE (3/1) to give 1.23 g of compound

1-2. MS (ESI) m / e (M+H⁺): observed = 273.2, calculated = 273.2.

StepB: [1-(4-Phenylacetyl-phenyl)-cyclobutyl]-carbamic acid tert-butyl ester (1-3)


A solution of i-PrMgCl (9.65 mL, 19.3 mmol) in THF was added dropwise to a solution of compound **1-2** (5.24 g, 19.3 mmol) in dry THF at -5.deg.C. After stirring for 20 min, a solution of BnMgCl (29 mL, 2M, 58 mmol) in THF was added dropwise to the reaction mixture. After completion of addition, the reaction mixture was stirred for 1h at 0.deg.C, then warmed slowly to room temperature, and stirred for another 1h. The reaction was quenched with saturated NH₄Cl solution, extracted with ethyl acetate. Separated and collected the organic layer, and the aqueous solution was washed EA for two times. The combined organic layer was washed with brine, dried and concentrated in vacuo to get the crude product, which was purified by silica chromatography to give 5.5 g of ketone **1-3**.

MS (ESI) m / e (M+H⁺): observed = 388.2(+23)/332.2(-56) calculated = 388.2(+23)/332.2(-56)


StepC: {1-[4-(3-Dimethylamino-2-phenyl-acryloyl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (1)

To a solution of compound **1-3** (5.0 g, 13.7 mmol) in toluene (30 mL) was added DMFDMA (18.15 g, 68.4 mmol) and the mixture was heated at reflux for 4 hours. The TLC showed the starting material was disappeared. The mixture was cooled and the formed white solid was collected. After washed by PE, the solid was dried in vacuum to get the product **1**.

MS (ESI) m / e (M+H⁺): observed = 421.3; calculated = 421.2

{1-[4-(3-Dimethylamino-2-phenyl-acryloyl)-phenyl]-3-hydroxy-3-methyl-cyclobutyl}-carbamic acid tert-butyl ester (2)

Reaction Scheme 2



StepA: [1-(4-Cyano-phenyl)-3-hydroxy-3-methyl-cyclobutyl]-carbamic acid tert-butyl ester (2-2)

A mixture of [1-(4-Bromo-phenyl)-3-hydroxy-3-methyl-cyclobutyl]-carbamic acid tert-butyl ester 2-1 (prepared according to WO2008/070041; 3.56 g, 10 mmol), Zn (1 g, 15 mmol), Zn(CN)_2 (1.76 g, 15 mmol), $\text{Pd}_2(\text{dba})_3$ (0.5 g, 0.5 mmol) and DPPF (300 mg, 0.5 mmol) in 30 mL of DMF was stirred at 100°C for 2hrs under N_2 . After the mixture was cooled to rt, 100mL of water was added and product was extracted with ethyl acetate 100mL \times 3. The organic layer was combined, washed with brine, dried over sodium sulfate and concentrated under vacuum. The residue was purified by silica gel chromatography using EA/PE (1:2) as eluant to give compound 2-2.

MS (ESI) m / e ($\text{M}+\text{H}^+$): observed =303.3, calculated = 303.2.

StepB: [3-Hydroxy-3-methyl-1-(4-phenylacetyl-phenyl)-cyclobutyl]-carbamic acid tert-butyl ester (2-3)

A solution of $i\text{-PrMgCl}$ (9.65 mL, 19.3 mmol) in THF was added dropwise to a solution of compound 2-2 (5.84 g, 19.3 mmol) in dry THF at -5.deg.C. After stirring for 20 min, a solution of BnMgCl (58 mL, 2M, 116 mmol) in THF was added dropwise to the reaction mixture. After completion of addition, the reaction mixture was stirred for 1h at 0.deg.C, then warmed slowly to room temperature, and stirred for another 1h. The reaction was quenched with saturated NH_4Cl solution, extracted with ethyl acetate. Separated and collected the organic layer, and the aqueous solution was washed EA for two times. The combined organic layer was washed with brine, dried and concentrated in vacuo to get the crude product, which was purified by silica chromatography to afford 6.2 g of ketone 2-3.


MS (ESI) m / e ($\text{M}+\text{H}^+$): observed = 418.2(+23)/340.2(-56) calculated = 418.2(+23)/340.2(-56)

StepC: {1-[4-(3-Dimethylamino-2-phenyl-acryloyl)-phenyl]-3-hydroxy-3-methyl-cyclobutyl}-carbamic acid tert-butyl ester (**2**)


To a solution of compound **2-3** (3.96 g, 10 mmol) in toluene (20 mL) was added DMFDMA (9.2 g, 35 mmol) and the mixture was heated at reflux for 4 hours. The mixture was cooled and the formed white solid was collected. After washed by PE, the solid was dried in vacuum to get the product **2**.

MS (ESI) m/e ($M+H^+$): observed = 451.3; calculated = 451.2

Reaction Scheme 3



Example (3-6)



StepA: 6-Phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-5,7-diol (**3-2**)

A mixture of 1H-[1,2,4]Triazol-3-ylamine **SM** (500 mg, 5.95 mmol) and 2-Phenyl-malonic acid diethyl ester **3-1** (1.41 g, 5.95 mmol) in N,N-dibutylbutan-1-amine (5 mL) was stirred at 185°C overnight. The mixture was partitioned with two layers after cooling to room temperature. The top layer was removed and the lower layer was washed with cold PE. The resulting solution was concentrated and extracted with a mixture of diethyl ether and 10% w/w NaOH solution. The organic layer was discarded and aqueous layer acidified with concentrated HCl. The precipitated product was collected by filtration to afford compound **3-2**.

MS (ESI) m/e ($M+H^+$) 229.1

StepB: 5,7-Dichloro-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine (**3-3**)

Compound **3-2** (600 mg, 2.6 mmol) was suspended in 3 mL of $POCl_3$ and the mixture was stirred at 100°C for 6 h. The solvent was removed and the residue was dissolved in

a mixture of dichloromethane and ice water. The organic phase was separated and aqueous phase was extracted with dichloromethane twice. The combined organic phase was dried over Na_2SO_4 and evaporated to yield the crude product **3-3**, which was used without further purification.

5 **MS (ESI) m / e ($\text{M}+\text{H}^+$)** 266.1

StepC: 5-Chloro-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine (3-4)

To the solution of **3-3** (100 mg, 0.37 mmol) in a co-solvent of AcOH (0.2 mL), MeOH (0.2 mL) and THF (1.2 mL) was added Zn/Cu pair (73 mg, 0.6 mmol) and the mixture was stirred at r.t. for 3h. The mixture was filtered to remove the insoluble solid and the filtrate
10 was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq. and brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.TLC to give 43 mg of **3-4**.

MS (ESI) m / e ($\text{M}+\text{H}^+$) 231.4

StepD: [4-(6-Phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-benzyl]-carbamic acid tert-butyl ester (3-5)

A mixture of compound **3-4** (60 mg 0.24 mmol), [4-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzyl]-carbamic acid tert-butyl ester (100 mg, 0.3 mmol), aq. Na_2CO_3 (2M, 0.70 mmol, 0.35 mL) and Pd(DPPF) Cl_2 (10 mg) in dioxane (3 mL) was heated at 100°C in microwave system under N_2 atmosphere for 30 min. After cooling, the mixture was diluted with
20 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.TLC to give 46 mg of **3-5**. **MS (ESI) m / e ($\text{M}+\text{H}^+$)** 402.1

StepE: 4-(6-Phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-benzylamine (3-6)

To the solution of **3-5** (46 mg, 0.11 mmol) in 1 mL of DCM was added TFA (1
25 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **3-6**.

MS (ESI) m / e ($\text{M}+\text{H}^+$) 302.1


^1H NMR (400 MHz, MeOH): δ 8.50 (s, 1H), 7.61-7.50 (m, 3H), 7.42-7.33 (m, 5H), 7.28-7.22 (m, 2H), 4.13 (s, 2H)



30 The following compounds **3-7** to **3-8** were prepared in a similar manner to Example **3-6** using the appropriate materials commercially available.

Ex.	Structure	MW	Compound Name	LC/MS (ESI) observed $[\text{M}+1]^+$
-----	-----------	----	---------------	---------------------------------------

3-7		396.5	397.4
3-8		300.4	301.1

Reaction Scheme 4



**Example (4-2)**

5 **{1-[4-(6-Phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (4-1)**

Compound 1 (1.2 g, 2.9 mmol) and 2-aminoimidazole (0.77 g 4.3 mmol) were added into a solution of AcOH (2 mL) and EtOH (10 mL). The mixture was heated to reflux and stirred overnight. The solution was cooled to r.t. and diluted with 30 mL of DCM. The organic phase was washed with 0.1N HCl(aq) and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography to give 1.2 g of 4-1 as a yellow solid. **MS (ESI) m / e (M+H⁺): 441.3**


10 **1-[4-(6-Phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (4-2)**

To a solution of 4-1 (45 mg, 0.1 mmol) in 1 mL of MeOH was added 4M HCl/MeOH (2 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product 4-2. **MS (ESI) m / e (M+H⁺): 341.3**

15

¹H NMR (400 MHz, MeOH): δ 8.59 (s, 1H), 7.65 (d, J=1.2Hz, 1H), 7.64 (d, J=8.0Hz, 2H), 7.45 (d, J=8.4Hz, 2H), 7.40 (d, J=1.2Hz, 1H), 2.65-2.52 (m, 2H), 2.48-2.39 (m, 2H), 2.18-2.08 (m, 2H), 1.86-1.73 (m, 2H)

Example (4-4)



4-4

{1-[4-(3-Bromo-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (**4-3**)

To a solution of compound **4-1** (100 mg, 0.227 mmol) in CHCl₃ (5 mL) was added NBS (40 mg 0.227 mmol) and the mixture was heated to reflux for 2 hrs. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl(aq) and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 150 mg of **4-3**.

MS (ESI) m / e (M+H⁺) 520.3


1-[4-(3-Bromo-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**4-4**)

To a solution of **4-3** (52 mg, 0.1 mmol) in 1 mL of MeOH was added 4M HCl/MeOH (2 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **4-4**.

MS (ESI) m / e (M+H⁺): 420.3

¹H NMR (400 MHz, MeOH): δ 8.98 (s, 1H), 7.60 (brs, 4H), 7.28 (brs, 6H), 5.65 (d, J=17.2Hz, 1H), 5.49-7.42 (m, 1H), 5.07 (d, J=10.8Hz, 1H), 2.81-2.72 (m, 2H), 2.69-2.56 (m, 2H), 2.30-2.22 (m, 2H), 1.96-1.89 (m, 2H).

Example (4-6)



4-6

{1-[4-(6-Phenyl-3-vinyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (**4-5**)

A mixture of compound **4-3** (100 mg 0.193 mmol), Tributyl-vinyl-stannane (92 mg, 0.58 mmol), K_2CO_3 (27 mg, 0.193 mmol), Et_4NCl (31.8 mg 0.193 mmol) and $Pd(PPh_3)_2Cl_2$ (10 mg) in THF (5 mL) was heated at 100°C in microwave system under N_2 atmosphere for 45 min. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.TLC to give 78 mg of **4-5**.

MS (ESI) m / e ($M+H^+$): 467.3


1-[4-(6-Phenyl-3-vinyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**4-6**)

To the solution of **4-5** (47 mg, 0.1 mmol) in 1 mL of MeOH was added 4M HCl/MeOH (2 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **4-6**.

MS (ESI) m / e ($M+H^+$): 367.3

1H NMR (400 MHz, MeOD): δ 9.02 (s, 1H), 8.08 (s, 1H), 7.34 (d, $J=1.6$ Hz, 1H), 7.32-7.25 (m, 4H), 7.24-7.20 (m, 5H), 6.95 (dd, $J=16.4$ Hz, 8.4Hz, 2H), 6.51 (d, $J=8.4$ Hz, 1H), 3.50 (s, 3H), 2.62-2.55 (m, 2H), 2.53-2.49 (m, 2H), 2.25-2.18 (m, 1H), 1.96-1.89 (m, 1H).

Example (4-8)



(1-{4-[3-(2-Methoxy-phenyl)-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutyl)-carbamic acid tert-butyl ester (**4-7**)

A mixture of compound **4-3** (52 mg 0.1 mmol), 2-methoxyl phenyl boronic acid (30 mg, 0.15 mmol), aq. Na_2CO_3 (2M, 0.25 mmol, 0.12 mL) and $Pd(DPPF)Cl_2$ (10 mg) in dioxane (3 mL) was heated at 100°C in microwave system under N_2 atmosphere for 30 min. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.TLC to give 36 mg of **4-7**.


MS (ESI) m / e ($M+H^+$): 547.5

1-{4-[3-(2-Methoxy-phenyl)-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutylamine (**4-8**)

To a solution of **4-7** (55 mg, 0.1 mmol) in 1 mL of MeOH was added 4M HCl/MeOH (2 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **4-8**.
MS (ESI) m / e (M+H⁺): 447.5

- 5 **¹H NMR (400 MHz, MeOD):** δ 9.02 (s, 1H), 8.08 (s, 1H), 7.34 (d, J=1.6Hz, 1H), 7.32-7.25 (m, 4H), 7.24-7.20 (m, 5H), 6.95 (dd, J=16.4Hz, 8.4Hz, 2H), 6.51 (d, J=8.4Hz, 1H), 3.50 (s, 3H), 2.62-2.55 (m, 2H), 2.53-2.49 (m, 2H), 2.25-2.18 (m, 1H), 1.96-1.89 (m, 1H).

Example (4-10)



4-10

- 10 {1-[4-(3-Cyano-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (**4-9**)

A mixture of compound **4-3** (80 mg 0.154 mmol), Zn(CN)₂ (11 mg, 0.0906 mmol), Zn (1 mg 0.015 mmol) and Pd(dppf)Cl₂ (10 mg) in DMA (2 mL) was heated at 160°C in microwave system for 45 min. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl(aq) and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 60 mg of **4-9**.

MS (ESI) m / e (M+H⁺): 466.1


7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-3-carbonitrile (**4-10**)

- 20 To a solution of **4-9** (46 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **4-10**.

MS (ESI) m / e (M+H⁺): 366.1

- 25 **¹H NMR (400 MHz, MeOD):** δ 8.86 (s, 1H), 8.46 (s, 1H), 7.62 (dd, J=12.0Hz, 8.4Hz, 4H), 7.29-7.26 (m, 5H), 2.80-2.70 (m, 2H), 2.67-2.56 (m, 2H), 2.29-2.13 (m, 1H), 2.06-1.90 (m, 1H)

Example (4-11)



7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-3-carboxylic acid amide
(4-11)

To a suspension of compound **4-10** (40 mg 0.11 mmol) and K_2CO_3 (30 mg, 0.22 mmol) in DMSO (1.0 mL) was added 0.1 mL of H_2O_2 (30% aq.) and the mixture was heated at 60°C for 3h. After cooling, the mixture was diluted with 20 mL of EA, the combined organic phase was washed with sat. $NaHSO_3$ (aq) and brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.HPLC to give 19 mg of **4-11**.

MS (ESI) m / e (M+H⁺): 384.3


¹H NMR (400 MHz, MeOD): δ 8.87 (s, 1H), 8.12 (s, 1H), 7.68 (dd, J=7.6 Hz, 2H), 7.63 (d, J=8.0 Hz, 2H), 7.33-7.25 (m, 5H), 2.81-2.74 (m, 2H), 2.66-2.59 (m, 2H), 2.29-2.24 (m, 1H), 1.98-1.92 (m, 1H).


The following compound **4-14** was prepared similarly to Example 4-4; **4-15** was prepared similarly to Example 4-8; **4-16** was prepared similarly to Example 4-6.

Ex.	Structure	MW	Compound Name	LC/MS (ESI) found [M+1] ⁺
4-14		374.9	1-[4-(3-Chloro-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	375.7
4-15		354.5	1-[4-(3-Methyl-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	355.2

4-16		406.5	1-{4-[3-(2-Cyclopropyl-vinyl)-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutylamine	407.1
------	--	-------	--	-------

Reaction Scheme 5

Example (5-3)



5-3

{1-[4-(2-Amino-5-phenyl-pyrimidin-4-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (5-1)

To a solution of compound **1** (3 g, 7.2 mmol) and guanidine (2 g, 35 mmol) in DMF (15 mL) was added MeONa (1.8 g, 35 mmol) and the mixture was heated to 100°C and stirred for 4 h. The solution was cooled to r.t. and diluted with 80 mL of EA. The organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel column chromatography to give 2.1 g of **5-1** as a yellow solid.

MS (ESI) m / e (M+H⁺): 417.3

{1-[4-(2,6-Diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (5-2)

A mixture of **5-1** (150 mg, 0.36 mmol) and 2-Bromo-1-phenyl-ethanone (110 mg, 0.54 mmol) in ethanol (2 mL) was heated to 100°C and stirred for 45 mins under microwave radiation. The mixture was cooled to r.t. and diluted with 20 mL of DCM. The organic phase was washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 90 mg of **5-2**.

MS (ESI) m / e (M+H⁺): 517.4


1-[4-(2,6-Diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (5-3)

To the solution of **5-2** (52 mg, 0.1 mmol) in 1 mL of MeOH was added 4M HCl/MeOH (2 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **5-3**.

MS (ESI) m / e (M+H⁺): 417.4

¹H NMR (400 MHz, MeOD): δ 9.11 (s, 1H), 8.45 (s, 1H), 8.00 (d, J=7.6Hz, 2H), 7.68 (d, J=8.4Hz, 2H), 7.61-7.50 (m, 5H), 7.47-7.41 (m, 3H), 7.41-7.35 (m, 2H), 3.33-3.32 (m, 2H), 3.32-2.74 (m, 2H), 2.28-2.25 (m, 1H), 1.99-1.96 (m, 1H).

Example (5-5)

**5-5**

{1-[4-(3-Bromo-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (**5-4**)

To a solution of compound **5-2** (110 mg, 0.2 mmol) in CHCl_3 (5 mL) was added NBS (40 mg 0.22 mmol) and the mixture was heated to reflux for 1 hrs. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.TLC to give 71 mg of **5-4**.

MS (ESI) m/e ($M+H^+$) 596.3


1-[4-(3-Bromo-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**5-5**)

To the solution of **5-4** (59 mg, 0.1 mmol) in 1 mL of MeOH was added 4M HCl/MeOH (2 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **5-5**.

MS (ESI) m/e ($M+H^+$): 496.3

^1H NMR (400 MHz, MeOD): δ 8.26 (s, 1H), 7.80 (d, $J=7.6\text{Hz}$, 2H), 7.68 (d, $J=8.4\text{Hz}$, 2H), 7.61-7.50 (m, 5H), 7.47-7.41 (m, 3H), 7.41-7.35 (m, 2H), 3.33-3.32 (m, 2H), 3.32-2.74 (m, 2H), 2.28-2.25 (m, 1H), 1.99-1.96 (m, 1H).

Example (**5-7**)



(1-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutyl)-carbamic acid tert-butyl ester (**5-6**)

A mixture of compound **5-4** (60 mg 0.1 mmol), 3,5-ditrifluoromethyl phenyl boronic acid (38 mg, 0.15 mmol), aq. Na₂CO₃ (2M, 0.25 mmol, 0.12 mL) and Pd(DPPF)Cl₂ (10 mg) in dioxane (3 mL) was heated at 100°C in microwave system under N₂ atmosphere for 30 min. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 38 mg of **5-6**.

MS (ESI) m / e (M+H⁺): 728.6


1-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutylamine (**5-7**)

To the solution of **5-6** (38 mg, 0.05 mmol) in 1 mL of MeOH was added 4M HCl/MeOH (2 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **5-7**.

MS (ESI) m / e (M+H⁺): 628.3

¹H NMR (400 MHz, MeOD): δ 8.57 (s, 1H), 8.24 (s, 2H), 8.16 (s, 1H), 7.67 (d, J=8.4Hz, 2H), 7.60 (m, 2H), 7.49 (d, J=8.4Hz, 2H), 7.44-7.7.30 (m, 8H), 2.79 (m, 2H), 2.62(m, 2H), 2.26 (m, 1H), 2.00 (m, 1H)

Example (**5-9**)



5-9

{1-[4-(2,6-Diphenyl-3-vinyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (**5-8**)

A mixture of compound **5-4** (120 mg 0.2 mmol), Tributyl-vinyl-stannane (95 mg, 0.6 mmol), K₂CO₃ (30 mg, 0.2 mmol), Et₄NCl (35 mg 0.2 mmol) and Pd(PPh₃)₂Cl₂ (10 mg) in THF (3 mL) was heated at 100°C in microwave system under N₂ atmosphere for 45 min. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 57 mg of **5-8**.

MS (ESI) m / e (M+H⁺): 543.5

1-[4-(2,6-Diphenyl-3-vinyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine (**5-9**)

To the solution of **5-8** (54 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **5-9**.







MS (ESI) m/e ($M+H^+$): 443.5




1H NMR (400 MHz, MeOD): δ 8.94 (d, $J=2.0$ Hz, 1H), 7.74 (d, $J=7.2$ Hz, 2H), 7.55 (d, $J=8.0$ Hz, 2H), 7.47 (d, $J=7.6$ Hz, 2H), 7.38 (d, $J=8.4$ Hz, 2H), 7.30 (m, 5H), 6.95 (dd, $J=18.0$ Hz, 12.0 Hz, 1H), 5.84 (d, $J=18.0$ Hz, 1H), 5.68 (d, $J=12.0$ Hz, 1H), 2.65 (m, 2H), 2.50 (m, 2H), 2.14 (m,



5 1H), 1.86 (m, 1H)

The following compounds **5-11** to **5-17** were prepared in a similar manner to Example **5-3** using the appropriate commercially available materials; **5-18** was prepared similarly to Example **5-5**; compounds **5-19** to **5-22** were prepared similarly to Example **5-7**; **5-23** was prepared similarly to Example **5-9**.

Ex.	Structure	MW	Compound Name	LC/MS (ESI) found $[M+1]^+$
5-11		368.5	1-[4-(2-Ethyl-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	369.3
5-12		408.4	1-[4-(6-Phenyl-2-trifluoromethyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	409.2
5-13		412.5	7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-2-carboxylic acid ethyl ester	413.3
5-14		384.4	7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-2-carboxylic acid	385.2

5-15		398.5	7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-2-carboxylic acid methyl ester	399.4
5-16		434.5	1-[4-[2-(4-Fluorophenyl)-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl]-cyclobutylamine	435.2
5-17		430.6	1-[4-(6-Phenyl-2-p-tolyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	431.3
5-18		487.3	1-[4-(3-Bromo-6-phenyl-2-trifluoromethyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	397.4
5-19		430.6	1-[4-(3-Methyl-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	431.2
5-20		492.6	1-[4-(2,3,6-Triphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	493.3

5-21		456.6	1-[4-(3-Cyclopropyl-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	457.3
5-22		546.6	1-[4-{2,6-Diphenyl-3-(2,3,4-trifluoro-phenyl)-imidazo[1,2-a]pyrimidin-7-yl}-phenyl]-cyclobutylamine	547.3
5-23		456.6	1-[4-(2,6-Diphenyl-3-propenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine	457.3

Reaction Scheme 6Example (6-2)

5 {1-[4-(2-Amino-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (**6-1**)

A mixture of **1** (250 mg, 0.59 mmol) and **4H-[1,2,4]Triazole-3,5-diamine** (58.91 mg, 0.59 mmol) in acetic acid (0.4 mL) and ethanol (1.6 mL) was heated under reflux for 6 hours. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with NaHCO₃ aq. and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give **6-1**. MS (ESI) *m / e* (M+H⁺) 457.3


5 **5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-ylamine**
(**6-2**)

To the solution of **6-1** (46 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **6-2**.

MS (ESI) *m / e* (M+H⁺) 357.3

¹H NMR (400 MHz, MeOD): δ 8.59 (s, 1H), 8.51 (dd, J=17.2Hz,8.4Hz, 4H), 7.26-7.7.20 (m, 5H), 2.64 (m, 2H), 2.41 (m, 2H), 2.52 (m, 1H),1.85 (m, 1H)

Example (6-4)



6-4

{1-[4-(2-Acetylamino-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (6-3)

To the solution of **6-1** (100 mg, 0.22 mmol) and Et₃N (0.2 mL) DMAP (2 mg) in dioxane (2 mL) was added acetyl chloride (21mg, 0.25 mmol) and the mixture was heated at 100°C for 10 h. The mixture was diluted with 20 mL of DCM, the combined organic phase was washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 64 mg of **6-3**.

MS (ESI) *m / e* (M+H⁺) 499.4


25 **N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl]}-acetamide**
(**6-4**)

To the solution of **6-3** (50 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **6-4**.

MS (ESI) *m / e* (M+H⁺) 399.3

¹H NMR (400 MHz, DMSO): δ 8.533 (s, 1H), 8.491 (s, 1H), 7.337 (s, 4H), 7.254-7.245 (d, $J=3.6$ Hz, 3H), 7.189-7.172 (t, $J=6.8$ Hz, 2H), 2.385-2.325 (q, $J=24.0$ Hz, 4H), 1.981-1.939 (t, $J=16.8$ Hz, 1H), 1.792 (s, 4H)

Example (6-7)



6-7

{1-[4-(2-Bromo-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (6-5)

To a suspension of CuBr₂ (294 mg, 0.64 mmol) and t-BuONO (0.2 mL, 1.7 mmol) in CH₃CN (15 mL) at 50.°C was added a solution of compound **6-1** (500 mg, 1.1 mmol) in CH₃CN (8 mL) drop-wise under N₂ atmosphere and the resulting mixture was stirred for another 30 min. The mixture was cooled to room temperature and diluted with 60 mL of EA. The insoluble solid was filtered off and the mixture was washed with brine dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by silica gel chromatography (PE: EA=5:1) to afford 275 mg of **6-5**.

MS (ESI) m / e (M+H⁺) 521.4

{1-[4-(2,6-Diphenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (6-6)

A mixture of compound **6-5** (52 mg 0.1 mmol), phenyl boronic acid (19 mg, 0.15 mmol), aq.Na₂CO₃ (2M, 0.25 mmol, 0.12 mL) and Pd(dppf)Cl₂ (10 mg) in dioxane (2 mL) was heated at 100°C in microwave system under N₂ atmosphere for 30 min. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 26 mg of **6-6**.

MS (ESI) m / e (M+H⁺) 518.1

1-[4-(2,6-Diphenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-7)


To the solution of **6-6** (52 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **6-7**.

MS (ESI) m / e (M+H⁺) 418.1

¹H NMR (400 MHz, MeOD): δ 8.912 (s, 1H), 8.157-8.133 (q, $J=9.6$ Hz, 2H), 7.722-7.702 (d, $J=8.0$ Hz, 2H), 7.601-7.580 (d, $J=8.4$ Hz, 2H), 7.474-7.457 (d, $J=6.8$ Hz, 3H), 7.313-7.308 (d,

$J=2\text{Hz}$, 5H), 2.849-2.776 (m, 2H), 2.638-2.565 (m, 2H), 2.269-2.240 (m, 1H), 2.030-1.984 (m, 1H).

Example (6-9)



6-9

- 5 {1-[4-(2-Phenoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (6-8)

A mixture of 6-5 (52 mg, 0.1 mmol), Phenol (19 mg, 0.2 mmol) and K_2CO_3 (45 mg, 0.3 mmol) in DMF (1 mL) was heated at 110°C for 10 h. The mixture was diluted with 20 mL of DCM, the combined organic phase was washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.TLC to give 24 mg of 6-8.

MS (ESI) m/e ($M+H^+$) 534.4


1-[4-(2-Phenoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (6-9)

To the solution of 6-8 (24 mg, 0.04 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product 6-9.

MS (ESI) m/e ($M+H^+$) 434.4

$^1\text{H NMR}$ (400 MHz, MeOD): δ 8.720 (s, 1H), 7.724-7.703 (d, $J=8.4\text{Hz}$, 2H), 7.622-7.600 (d, $J=8.8\text{Hz}$, 2H), 7.384-7.368 (t, $J=6.4\text{Hz}$, 3H), 7.309-7.285 (m, 2H), 3.862-3.838 (t, $J=9.6\text{Hz}$, 4H), 3.646-3.622 (t, $J=8.8\text{Hz}$, 4H), 2.902-2.829 (m, 2H), 2.720-2.648 (m, 2H), 2.382-2.305 (m, 1H), 2.110-2.050 (m, 1H).

Example (6-10)



6-10

1-[4-(2-Morpholin-4-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine
(6-10)







A mixture of 6-5 (52 mg, 0.1 mmol), morph line (17 mg, 0.2 mmol) and Et₃N (52 mg, 0.5 mmol) in DMF (1 mL) was heated at 110°C for 10 h. The mixture was diluted with 20 mL of DCM, the combined organic phase was washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was used without purification in the following de-Boc reaction following the similar procedure from 6-8 to 6-9 to give 19 mg of 6-10 after prep.HPLC separation.







MS (ESI) m / e (M+H⁺) 427.3







¹H NMR (400 MHz, MeOD): δ 8.833 (s, 1H), 7.625-7.604 (d, J=8.4Hz, 2H), 7.524-7.503 (d, J=8.4Hz, 2H), 7.415-7.375 (t, J=16Hz, 2H), 7.325-7.304 (m, 3H), 7.294-7.203 (m, 5H), 2.781-2.709 (m, 2H), 2.598-2.524 (m, 2H), 2.274-2.183 (m, 1H), 2.000-1.931 (m, 1H).







The following examples 6-11 to 6-12 were prepared in a similar manner to Example 6-2 using the appropriate materials commercially available; examples 6-13 to 6-19 were prepared similarly to Example 6-4; examples 6-20 to 6-46 were prepared similarly to Example 6-7; examples 6-47 to 6-54 were prepared similarly to Example 6-9; examples 6-55 to 6-66 were prepared similarly to Example 6-10.

Ex.	Structure	MW	Compound Name	LC/MS (ESI) found [M+1] ⁺
6-11		341.4	1-[4-(6-Phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	342.2
6-12		355.4	1-[4-(2-Methyl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	356.2
6-13		448.6	Ethanesulfonic acid {5-[4-(1-amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-amide	449.4

6-14		496.6	N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-benzenesulfonamide	497.3
6-15		460.5	N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-benzamide	461.2
6-16		424.5	Cyclopropanecarboxylic acid {5-[4-(1-amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-amide	425.3
6-17		475.6	1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-3-phenyl-urea	476.4
6-18		491.6	1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-3-phenyl-thiourea	492.4
6-19		428.5	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-carbamic acid ethyl ester	429.3







6-20		448.5	1-{4-[2-(6-Methoxy-pyridin-2-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	449.3
6-21		419.5	1-[4-(6-Phenyl-2-pyrazin-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	420.3
6-22		445.6	1-{4-[2-(2,3-Dimethyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	446.3
6-23		453.5	1-{4-[2-(3,4-Difluoro-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	454.3
6-24		449.5	1-{4-[2-(5-Fluoro-2-methyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	450.3
6-25		418.5	1-[4-(6-Phenyl-2-pyridin-4-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	419.2







6-26		432.5	4-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-phenylamine	433.2
6-27		431.5	1-[4-(6-Phenyl-2-p-tolyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	432.2
6-28		447.5	1-[4-[2-(4-Methoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	448.2
6-29		418.5	1-[4-(6-Phenyl-2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	419.3
6-30		447.5	1-[4-[2-(2-Methoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	448.3
6-31		470.0	1-[4-[2-(3-Chloro-4-fluoro-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	471

6-32		503.5	1-{4-[2-(2-Fluoro-4-trifluoromethyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	504.3
6-33		471.5	1-{4-[6-Phenyl-2-(2,3,4-trifluoro-phenyl)-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	472.3
6-34		477.6	1-{4-[2-(2,4-Dimethoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	478.4
6-35		407.5	1-[4-(2-Furan-3-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	408.2
6-36		423.5	1-[4-(6-Phenyl-2-thiophen-3-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	424.3
6-37		423.5	1-[4-(6-Phenyl-2-thiophen-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	424.3

6-38		497.6	1-[4-[2-(6-Methoxynaphthalen-2-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	498.3
6-39		507.6	1-[4-(2-Dibenzofuran-4-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	508.2
6-40		433.5	4-[5-[4-(1-Aminocyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl]-phenol	434.3
6-41		431.5	1-[4-(6-Phenyl-2-m-tolyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	432.2
6-42		509.6	1-[4-[2-(4-Phenoxyphenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	510.3
6-43		493.6	1-[4-(2-Biphenyl-2-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	494.3


6-44		474.6	1-[4-(2-Benzothiazol-2-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	425.3
6-45		418.5	1-[4-(6-Phenyl-2-pyridin-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	419.1
6-46		424.5	1-[4-(6-Phenyl-2-thiazol-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	425.3
6-47		434.5	1-[4-[6-Phenyl-2-(pyridin-4-yloxy)-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	435.2
6-48		371.4	1-[4-(2-Methoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	372.2
6-49		433.5	1-[4-(2-Phenoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	434.3


6-50		387.5	1-[4-(2-Methylsulfanyl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	388.1
6-51		483.6	1-[4-[2-(Naphthalen-2-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	484.4
6-52		509.6	1-[4-[2-(Biphenyl-2-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	510.4
6-53		484.6	1-[4-[2-(Isoquinolin-7-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	485.3
6-54		525.6	1-[4-[2-(4-Phenoxyphenoxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	526.3
6-55		472.6	1-[4-[2-(3,4-Dihydro-2H-quinolin-1-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	472.4

6-56		458.6	1-{4-[2-(2,3-Dihydro-indol-1-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	459.4
6-57		370.5	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-methylamine	371.3
6-58		384.5	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-dimethylamine	385.3
6-59		433.5	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-pyridin-3-ylamine	434.2
6-60		412.5	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-diethylamine	413.2
6-61		399.5	N1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-ethane-1,2-diamine	400.2

6-62		414.5	2-({5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-methyl-amino)-ethanol	415.4
6-63		460.6	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-benzyl-methyl-amine	461.3
6-64		660.7	1-[4-(2-{4-[Bis-(4-chloro-phenyl)-methyl]-piperazin-1-yl}-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	661.5
6-65		446.6	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-methyl-phenyl-amine	447.2
6-66		414.5	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-(2-methoxy-ethyl)-amine	415.3

Reaction Scheme 7



Example (7-2)**7-2**

{1-[4-(2,6-Diphenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-3-hydroxy-3-methyl-cyclobutyl}-carbamic acid tert-butyl ester (7-1)

A mixture of **2** (270 mg, 0.6 mmol) and 5-Phenyl-4H-[1,2,4]triazol-3-ylamine (96 mg, 0.6 mmol) in acetic acid (1 mL) and ethanol (4 mL) was heated under reflux for 6 hours. After cooling, the mixture was diluted with 30 mL of DCM, the combined organic phase was washed with NaHCO₃ aq. and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give **7-1**.

MS (ESI) m / e (M+H⁺) 547.6

3-Amino-3-[4-(2,6-diphenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol (7-2)



To the solution of **7-1** (54 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **7-2**.

MS (ESI) m / e (M+H⁺) 447.6

¹H NMR (400 MHz, MeOD): δ 8.92 (s, 1H), 8.15-8.13(m, 2H), 7.72 (dd, J=8.4Hz,8.8Hz, 4H), 7.46 (m, 3H), 7.31 (s, 5H), 2.93-2.89 (d, J=14.8Hz, 2H), 2.74-2.70 (d, J=14.8Hz, 2H), 1.49 (s, 3H)

The following compound **7-3** was prepared in a similar manner to Example 7-2 using the appropriate materials commercially available.

Ex.	Structure	MW	Compound Name	LC/MS (ESI) found [M+1] ⁺
7-3		371.4	3-Amino-1-methyl-3-[4-(6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutanol	372.1

Reaction Scheme 8Example (8-2)

- 5 5-[4-(1-tert-Butoxycarbonylamino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methyl ester (**8-1**)

A mixture of **1** (420 mg, 1 mmol) and **5-Amino-4H-[1,2,4]triazole-3-carboxylic acid methyl ester** (215 mg, 1.5 mmol) in acetic acid (2.5 mL) and methanol (10 mL) was heated under reflux for 8 hours. After cooling, the mixture was diluted with 40 mL of DCM, the combined organic phase was washed with NaHCO₃ aq. and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by flash chromatography to give **8-1**.

MS (ESI) m / e (M+H⁺) 500.4


5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methyl ester (**8-2**)

To the solution of **8-1** (50 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **8-2** (35 mg, yield 87.8%).

MS (ESI) m / e (M+H⁺) 400.4

¹H NMR (400 MHz, MeOD): δ 9.04 (s, 1H), 7.63 (d, J=8.0Hz, 2H), 7.56 (d, J=8.0Hz, 2H), 7.31 (s, 5H), 3.95 (s, 3H), 2.74-2.81 (m, 2H), 2.56-2.63 (m, 2H), 2.19 (m, 1H), 1.97 (m, 1H)

Example (**8-4**)



8-4

5-[4-(1-tert-Butoxycarbonylamino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid (**8-3**)

A mixture of **8-1** (250 mg, 0.5 mmol) and LiOH (25 mg, 1.0 mmol) in THF (5 mL) and H₂O (2.5 mL) was stirred at room temperature overnight. The mixture was adjusted with 2N HCl aq to pH = 4 and then was extracted with EA twice. The combined organic phase was washed with H₂O and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.HPLC to give **8-3**.

MS (ESI) m / e (M+H⁺) 486.3


5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid (**8-4**)

To the solution of **8-3** (49 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **8-4**.

MS (ESI) m / e (M+H⁺) 386.3

¹H NMR (400 MHz, MeOD): δ 9.05 (s, 1H), 7.67 (d, J=8.0Hz, 2H), 7.58 (d, J=8.0Hz, 2H), 7.34 (s, 5H), 2.66-2.83 (m, 2H), 2.59-2.66 (m, 2H), 2.22-2.39 (m, 1H), 1.95-2.06 (m, 1H)

Example (8-6)



8-6

(1-{4-[2-(1-Hydroxy-1-methyl-ethyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutyl)-carbamic acid tert-butyl ester (**8-5**)

To a solution of **8-1** (100 mg, 0.2 mmol) in 4 mL of dry THF was added Methylmagnesium bromide (3 M THF solution, 0.3 mL, 0.9 mmol) dropwise at -20°C, then it was naturally warmed to 0°C and stirred for 3 h. The resulting solution was quenched by NH₄Cl aq, the mixture was extracted with EA twice. The combined organic phase was washed with NaHCO₃ aq. and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.HPLC to give **8-5**.

MS (ESI) m / e (M+H⁺) 500.4


2-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-propan-2-ol (**8-6**)

To the solution of **8-5** (50 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **8-6** (37 mg, yield 92.5%).

MS (ESI) m / e (M+H⁺) 400.4

¹H NMR (400 MHz, MeOD): δ 8.92 (s, 1H), 7.60 (d, J=8.0Hz, 2H), 7.52 (d, J=8.0Hz, 2H), 7.30-7.34 (m, 5H), 2.64-2.72 (m, 2H), 2.41-2.49 (m, 2H), 2.15-2.24 (m, 1H), 1.87-1.95 (m, 1H), 1.66 (s, 6H)

Example (8-8)

**8-8**

{1-[4-(2-Methylcarbamoyl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (**8-7**)

To a solution of **8-3** (243 mg, 0.5 mmol) in DMF (5 mL) was added Methanamine HCl salt (50 mg, 0.75mmol), HOBt (73 mg, 0.5 mmol), EDCI(192 mg, 1 mmol) and TEA (125 mg, 1.25 mmol) at room temperature and the mixture was stirred for overnight. The mixture was diluted with 15 mL of H₂O, extracted with EA, the combined organic phase was washed with NaHCO₃ aq. and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.HPLC to give **8-7**.

MS (ESI) m / e (M+H⁺) 499.4

5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methylamide (**8-8**)

To the solution of **8-7** (50 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **8-8**.



MS (ESI) m / e (M+H⁺) 399.4

¹H NMR (400 MHz, MeOD): δ 9.04 (s, 1H), 7.67 (d, J=8.0Hz, 2H), 7.59 (d, J=8.0Hz, 2H), 7.34-7.37 (m, 5H), 3.01 (s, 3H), 2.76-2.85 (m, 2H), 2.61-2.69 (m, 2H), 2.35-2.42 (m, 1H), 1.97-2.05 (m, 1H)


The following example **8-9** was prepared in a similar manner to Example **8-6** using the appropriate materials commercially available; examples **8-10** to **8-17** were prepared similarly to Example **8-8**.

Ex.	Structure	MW	Compound Name	LC/MS (ESI) found [M+1] ⁺
8-9		523.6	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-diphenyl-methanol	524.4


8-10		384.4	5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid amide	385.1
8-11		454.5	{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-morpholin-4-yl-methanone	455.3
8-12		488.6	5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid phenethyl-amide	489.4
8-13		440.6	5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid diethylamide	441.4
8-14		460.5	5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid phenylamide	461.3
8-15		488.6	5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid benzyl-methyl-amide	489.4

8-16		428.5	5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methoxy-methyl-amide	429.3
8-17		474.6	5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methyl-phenyl-amide	475.4

Reaction Scheme 9



Example (9-2)



9-2

{1-[4-(2,6-Diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (9-1)

A mixture of **1** (250 mg, 0.59 mmol) and 5-Phenyl-2H-pyrazol-3-ylamine (96 mg, 0.59 mmol) in acetic acid (0.4 mL) and ethanol (1.6 mL) was heated under reflux for 6 hours. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with NaHCO₃ aq. and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give **9-1**.

MS (ESI) m / e (M+H⁺) 517.4


1- [4-(2,6-Diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-2)

To the solution of **9-1** (52 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **9-2**.

MS (ESI) m / e (M+H⁺) 417.4

¹H NMR (400 MHz, MeOD): δ 8.57 (s, 1H), 7.89 (d, J=7.2Hz, 2H), 7.67 (d, J=8.0Hz, 2H), 7.56 (d, J=8.0Hz, 2H), 7.41-7.34 (m, 3H), 7.28-7.23 (m, 5H), 2.85-2.77 (m, 2H), 2.67-2.55 (m, 2H), 2.30-2.19 (m, 1H), 2.04-1.94 (m, 1H)

Example (9-4)



9-4

{1-[4-(3-Bromo-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (9-3)

To a solution of compound **9-1** (104 mg, 0.2 mmol) in CHCl₃ (5 mL) was added NBS (40 mg 0.22 mmol) and the mixture was stirred at room temperature overnight. The mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N

HCl aq and brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.TLC to give 83 mg of **9-3**.

MS (ESI) m / e ($\text{M}+\text{H}^+$) 595.9


1-[4-(2,3,6-Triphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-4)

To the solution of **9-3** (60 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **9-4**.

MS (ESI) m / e ($\text{M}+\text{H}^+$) 495.9

^1H NMR (400 MHz, MeOD): δ 8.64 (s, 1H), 7.97 (d, $J=7.2\text{Hz}$, 2H), 7.52 (d, $J=8.0\text{Hz}$, 2H), 7.47 (d, $J=8.0\text{Hz}$, 2H), 7.44-7.37 (m, 3H), 7.28-7.23 (m, 5H), 2.71-2.59 (m, 2H), 2.40-2.25 (m, 2H), 2.19-2.06 (m, 1H), 1.90-1.75 (m, 1H)

Example (9-6)



9-6

{1-[4-(2,3,6-Triphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (9-5)

A mixture of compound **9-3** (60 mg 0.1 mmol), phenyl boronic acid (19 mg, 0.15 mmol), aq. Na_2CO_3 (2M, 0.25 mmol, 0.12 mL) and $\text{Pd}(\text{DPPF})\text{Cl}_2$ (10 mg) in dioxane (2 mL) was heated at 100°C in microwave system under N_2 atmosphere for 30 min. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na_2SO_4 and concentrated. The residue was purified by prep.TLC to give 43 mg of **9-5**.

MS (ESI) m / e ($\text{M}+\text{H}^+$) 593.5






1-[4-(2,3,6-Triphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine (9-6)







To the solution of **9-5** (60 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **9-6**.







MS (ESI) m / e ($\text{M}+\text{H}^+$) 493.5







^1H NMR (400 MHz, MeOD): δ 8.50 (s, 1H), 7.59 (d, $J=8.0\text{Hz}$, 2H), 7.47 (d, $J=8.0\text{Hz}$, 2H), 7.42-7.38 (m, 4H), 7.31-7.21 (m, 11H), 2.77-2.65 (m, 2H), 2.51-2.47 (m, 2H), 2.22-2.11 (m, 1H), 1.96-1.83 (m, 1H).




The following examples 9-8 to 9-17 were prepared in a similar manner to Example 9-2 using the appropriate materials commercially available; examples 9-18 to 9-19 were prepared similarly to Example 9-4; examples 9-20 to 9-33 were prepared similarly to Example 9-6.

Ex.	Structure	MW	Compound Name	LC/MS (ESI) found [M+1] ⁺
9-8		340.4	1-[4-(6-Phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	341.2
9-9		354.5	1-[4-(2-Methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	435.3
9-10		356.4	5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-pyrazolo[1,5-a]pyrimidin-2-ol	357.2
9-11		430.6	1-[4-(6-Phenyl-2-p-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	431.4
9-12		446.6	1-[4-[2-(4-Methoxy-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	447.4


9-13		430.6	1-[4-(6-Phenyl-2-o-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	431.4
9-14		451.0	1-{4-[2-(4-Chlorophenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	452
9-15		434.5	1-{4-[2-(3-Fluorophenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	435.3
9-16		458.6	1-{4-[6-Phenyl-2-(2,4,6-trimethylphenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	459.4
9-17		434.5	1-{4-[2-(4-Fluorophenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	435.3
9-18		374.9	1-[4-(3-Chloro-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	375.7

9-19		419.3	1-[4-(3-Bromo-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	420.1
9-20		493.6	1-[4-(2,6-Diphenyl-3-pyridin-4-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	494.4
9-21		354.5	1-[4-(3-Methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	355.3
9-22		528.6	1-[4-(3-(2,4-Difluorophenyl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	529.4
9-23		430.6	1-[4-(3-Methyl-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	431.4
9-24		416.5	1-[4-(3,6-Diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	417.3


9-25		418.5	1-[4-(6-Phenyl-3-pyrimidin-5-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	419.3
9-26		460.6	1-[4-[3-(4-Methoxyphenyl)-2-methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	461.4
9-27		448.5	1-[4-[3-(2-Methoxypyrimidin-5-yl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	449.3
9-28		484.5	1-[4-[6-Phenyl-3-(4-trifluoromethylphenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	485.3
9-29		466.6	1-[4-(3-Naphthalen-1-yl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine	467.4
9-30		502.5	1-[4-[3-(2-Fluoro-5-trifluoromethylphenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl]-cyclobutylamine	503.3

9-31		469.0	1-{4-[3-(3-Chloro-4-fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	470
9-32		492.6	1-{4-[3-(3-Fluoro-5-isopropoxy-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	493.4
9-33		470.5	1-{4-[6-Phenyl-3-(2,3,4-trifluoro-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine	471.3

Reaction Scheme 10



Example (10-2)



5

{1-[4-(2,6-Diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-3-hydroxy-3-methyl-cyclobutyl}-carbamic acid tert-butyl ester (**10-1**)

A mixture of **2** (450 mg, 1 mmol) and 5-Phenyl-2H-pyrazol-3-ylamine (160 mg, 1 mmol) in acetic acid (2 mL) and ethanol (4 mL) was heated under reflux for 4 hours. After cooling, the mixture was diluted with 30 mL of DCM, the combined organic phase was washed with NaHCO₃ aq. and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by flash chromatography to give **10-1**.

MS (ESI) m / e (M+H⁺) 547.5


3-Amino-3-[4-(2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol
(10-2)

To the solution of **10-1** (54 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **10-2**.

MS (ESI) m / e (M+H⁺) 447.5

¹H NMR (400 MHz, MeOD): δ 8.57 (s, 1H), 7.90-7.88 (m, 2H), 7.65 (dd, J=8.4Hz,8.8Hz, 4H), 7.41-7.39 (m, 3H), 7.26 (s, 5H), 7.15 (s, 5H), 2.93-2.89 (d, J=14.8Hz, 2H), 2.73-2.69 (d, J=14.8Hz, 2H), 1.49 (s, 3H)

Example (10-4)



10-4

{1-[4-(3-Bromo-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-3-hydroxy-3-methyl-cyclobutyl}-carbamic acid tert-butyl ester (**10-3**)

To a solution of compound **10-1** (110 mg, 0.2 mmol) in CHCl₃ (5 mL) was added NBS (40 mg 0.22 mmol) and the mixture was stirred at room temperature overnight. The mixture was diluted with 15 mL of DCM, the combined organic phase was washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 81 mg of **10-3**.

MS (ESI) m / e (M+H⁺) 626.1


3-Amino-3-[4-(3-bromo-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol (**10-4**)

To the solution of **10-3** (62 mg, 0.1 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **9-4**.

MS (ESI) m / e (M+H⁺) 526.1

¹H NMR (400 MHz, MeOD): δ 8.70 (s, 1H), 7.98-7.96 (m, 2H), 7.68 (dd, J=8.4Hz,8.8Hz, 4H), 7.49-7.45 (m, 3H), 7.31 (s, 5H), 2.94-2.90 (d, J=14.8Hz, 2H), 2.74-2.70 (d, J=14.8Hz, 2H), 1.51 (s, 3H)

Example (10-6)

**10-6**

{3-Hydroxy-3-methyl-1-[4-(3-methyl-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutyl}-carbamic acid tert-butyl ester (**10-5**)

A mixture of compound **10-3** (62 mg, 0.1 mmol), Methyl boronic acid (60 mg, 1 mmol), aq. Na₂CO₃ (2M, 0.25 mmol, 0.12 mL) and Pd(DPPF)Cl₂ (10 mg) in dioxane (2 mL) was heated at 100°C in microwave system under N₂ atmosphere for 30 min. After cooling, the mixture was diluted with 15 mL of DCM, the combined organic phase was washed with 0.1N HCl aq and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by prep.TLC to give 23 mg of **10-5**.

MS (ESI) m / e (M+H⁺) 561.5

3-Amino-1-methyl-3-[4-(3-methyl-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutanol (**10-6**)






To the solution of **10-5** (23 mg, 0.04 mmol) in 1 mL of DCM was added TFA (1 mL) and the mixture was stirred at room temperature for 3 h. The mixture was concentrated by evaporation and the residue was purified by prep.HPLC to give the product **10-6**.

MS (ESI) m / e (M+H⁺) 461.5

¹H NMR (400 MHz, MeOD): δ 8.47 (s, 1H), 7.66-7.88 (m, 2H), 7.55 (dd, J=8.4Hz, 8.8Hz, 4H), 7.36-7.28 (m, 3H), 7.18 (s, 5H), 2.83-2.79 (d, J=14.8Hz, 2H), 2.60-2.59 (d, J=14.8Hz, 2H), 2.47 (s, 3H), 1.40 (s, 3H)

The following examples **10-7** to **10-8** were prepared in a similar manner to Example **10-2** using the appropriate materials commercially available; examples **10-9** to **10-12** were prepared similarly to Example **10-6**.

Ex.	Structure	MW	Compound Name	LC/MS (ESI) found [M+1] ⁺
10-7		460.6	3-Amino-1-methyl-3-[4-(6-phenyl-2-o-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutanol	461.4

10-8		464.5	3-Amino-3-[4-[2-(4-fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl]-1-methyl-cyclobutanol	465.3
10-9		523.6	3-Amino-3-[4-(2,6-diphenyl-3-pyridin-3-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol	524.4
10-10		537.7	3-Amino-3-[4-[3-(3-amino-phenyl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl]-1-methyl-cyclobutanol	538.5
10-11		524.6	3-Amino-3-[4-(2,6-diphenyl-3-pyrimidin-5-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol	525.4
10-12		539.7	3-Amino-3-[4-[3-(2-amino-pyrimidin-5-yl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl]-1-methyl-cyclobutanol	540.5

BIOLOGICAL EXAMPLE 1Cloning of the human Akt isoforms and ΔPH-Akt1

The pS2neo vector (deposited in the ATCC on April 3, 2001 as ATCC PTA-3253) was prepared as follows: The pRmHA3 vector (prepared as described in *Nucl. Acid Res.* 16:1043-1061 (1988)) was cut with BglII and a 2734 bp fragment was isolated. The pUCHsneo vector (prepared as described in *EMBO J.* 4:167-171 (1985)) was also cut with BglII and a 4029 bp band was isolated. These two isolated fragments were ligated together to generate a vector termed pS2neo-1. This plasmid contains a polylinker between a metallothionine promoter and an alcohol dehydrogenase poly A addition site. It also has a neo resistance gene driven by a heat shock promoter. The pS2neo-1 vector was cut with Psp5II and BsiWI. Two complementary oligonucleotides were synthesized and then annealed (CTGCGGCCGC (SEQ.ID.NO.: 1) and GTACGCGGCCGCAG (SEQ.ID.NO.: 2)). The cut pS2neo-1 and the annealed oligonucleotides were ligated together to generate a second vector, pS2neo. Added in this conversion was a NotI site to aid in the linearization prior to transfection into S2 cells.

Human Akt1 gene was amplified by PCR (Clontech) out of a human spleen cDNA (Clontech) using the 5' primer: 5'CGCGAATTCAGATCTACCATGAGCGACGTGGCTATTGTG 3' (SEQ.ID.NO.: 3), and the 3' primer: 5'CGCTCTAGAGGATCCTCAGGCCGTGCTGCTGGC3' (SEQ.ID.NO.: 4). The 5' primer included an EcoRI and BglII site. The 3' primer included an XbaI and BamHI site for cloning purposes. The resultant PCR product was subcloned into pGEM3Z (Promega) as an EcoRI/Xba I fragment. For expression/purification purposes, a middle T tag was added to the 5' end of the full length Akt1 gene using the PCR primer: 5'GTACGATGCTGAACGATATCTTCG 3' (SEQ.ID.NO.: 5). The resulting PCR product encompassed a 5' KpnI site and a 3' BamHI site which were used to subclone the fragment in frame with a biotin tag containing insect cell expression vector, pS2neo.

For the expression of a pleckstrin homology domain (PH) deleted (Δaa 4-129, which includes deletion of a portion of the Akt1 hinge region) version of Akt1, PCR deletion mutagenesis was done using the full length Akt1 gene in the pS2neo vector as template. The PCR was carried out in 2 steps using overlapping internal primers (5'GAATACATGCCGATGGAAAGCGACGGGGCTGAAGAGATGGAGGTG 3' (SEQ.ID.NO.: 6), and 5'CCCCTCCATCTCTTCAGCCCCGTCGCTTTCCATCGGCATG TATTC 3' (SEQ.ID.NO.: 7)) which encompassed the deletion and 5' and 3' flanking primers which encompassed the KpnI site and middle T tag on the 5' end. The final PCR product was digested with KpnI and SmaI and ligated into the pS2neo full length Akt1 KpnI/SmaI cut vector, effectively replacing the 5' end of the clone with the deleted version.

Human Akt3 gene was amplified by PCR of adult brain cDNA (Clontech) using the amino terminal oligo primer:

5' GAATTCAGATCTACCATGAGCGATGTTACCATTGTG 3' (SEQ.ID.NO.: 8); and the carboxy terminal oligo primer :

5' TCTAGATCTTATTCTCGTCCACTTGCCAGAG 3' (SEQ.ID.NO.: 9).

These primers included a 5' EcoRI/BglII site and a 3' XbaI/BglII site for cloning purposes. The resultant PCR product was cloned into the EcoRI and XbaI sites of pGEM4Z (Promega). For expression/purification purposes, a middle T tag was added to the 5' end of the full length Akt3 clone using the PCR primer:

5'GGTACCATGGAATACATGCCGATGGAAAGCGATGTTACCATTGTGAAG

3' (SEQ.ID.NO.: 10). The resultant PCR product encompassed a 5' KpnI site which allowed in frame cloning with the biotin tag containing insect cell expression vector, pS2neo.

Human Akt2 gene was amplified by PCR from human thymus cDNA (Clontech) using the amino terminal oligo primer:

5' AAGCTTAGATCTACCATGAATGAGGTGTCTGTC 3' (SEQ.ID.NO.: 11); and the carboxy terminal oligo primer: 5'GAATTCGGATCCTCACTCGCGGATGCTGGC 3' (SEQ.ID.NO.:

12). These primers included a 5' HindIII/BglII site and a 3' EcoRI/BamHI site for cloning purposes. The resultant PCR product was subcloned into the HindIII/EcoRI sites of pGem3Z (Promega). For expression/purification purposes, a middle T tag was added to the 5' end of the full length Akt2 using the PCR primer:

5'GGTACCATGGAATACATGCCGATGGAAAATGAGGTGTCTGTGCATCAAAG 3'

(SEQ.ID.NO.: 13). The resultant PCR product was subcloned into the pS2neo vector as described above.

EXAMPLE 2

Expression of human Akt isoforms and ΔPH-Akt1

The DNA containing the cloned Akt1, Akt2, Akt3 and ΔPH-Akt1 genes in the pS2neo expression vector was purified and used to transfect *Drosophila* S2 cells (ATCC) by the calcium phosphate method. Pools of antibiotic (G418, 500 μg/ml) resistant cells were selected. Cell were expanded to a 1.0 L volume (~7.0 x 10⁶ / ml), biotin and CuSO₄ were added to a final concentration of 50 μM and 50 mM respectively. Cells were grown for 72 h at 27°C and harvested by centrifugation. The cell paste was frozen at -70°C until needed.

EXAMPLE 3

Purification of human Akt isoforms and ΔPH-Akt1

Cell paste from one liter of S2 cells, described in Example 2, was lysed by sonication with 50 mls 1% CHAPS in buffer A: (50mM Tris pH 7.4, 1mM EDTA, 1mM EGTA, 0.2mM AEBSF, 10μg/ml benzamidine, 5μg/ml of leupeptin, aprotinin and pepstatin each, 10% glycerol and 1mM DTT). The soluble fraction was purified on a Protein G Sepharose fast flow (Pharmacia) column loaded with 9mg/ml anti-middle T monoclonal antibody and eluted with 75 μM EYMPME (SEQ.ID.NO.: 14) peptide in buffer A containing 25% glycerol. Akt/PKB containing fractions were pooled and the protein purity evaluated by SDS-PAGE. The purified

protein was quantitated using a standard Bradford protocol. Purified protein was flash frozen on liquid nitrogen and stored at -70°C.

Akt and Akt pleckstrin homology domain deletions purified from S2 cells required activation. Akt and Akt pleckstrin homology domain deletions were activated (Alessi et al. *Current Biology* 7:261-269) in a reaction containing 10 nM PDK1 (Upstate Biotechnology, Inc.), lipid vesicles (10 µM phosphatidylinositol-3,4,5-trisphosphate – Metreya, Inc, 100 µM phosphatidylcholine and 100 µM phosphatidylserine – Avanti Polar lipids, Inc.) and activation buffer (50 mM Tris pH7.4, 1.0 mM DTT, 0.1 mM EGTA, 1.0 µM Microcystin-LR, 0.1 mM ATP, 10 mM MgCl₂, 333 µg/ml BSA and 0.1mM EDTA). The reaction was incubated at 22°C for 4 hours. Aliquots were flash frozen in liquid nitrogen.

EXAMPLE 4

Akt Kinase Assays

Activated Akt isoforms and pleckstrin homology domain deletion constructs were assayed utilizing a GSK-derived biotinylated peptide substrate. The extent of peptide phosphorylation was determined by Homogeneous Time Resolved Fluorescence (HTRF) using a lanthanide chelate(Lance)-coupled monoclonal antibody specific for the phosphopeptide in combination with a streptavidin-linked allophycocyanin (SA-APC) fluorophore which will bind to the biotin moiety on the peptide. When the Lance and APC are in proximity (i.e. bound to the same phosphopeptide molecule), a non-radiative energy transfer takes place from the Lance to the APC, followed by emission of light from APC at 665 nm.

Materials required for the assay:

- A. Activated Akt isozyme or pleckstrin homology domain deleted construct
- B. Akt peptide substrate: GSK3α (S21) Peptide #3928 biotin-GGRARTSSFAEPG (SEQ.ID.NO.:15), Macromolecular Resources.
- 25 C. Lance labeled anti-phospho GSK3α monoclonal antibody (Cell Signaling Technology, clone # 27).
- D. SA-APC (Prozyme catalog no. PJ25S lot # 896067).
- E. Microfluor[®]B U Bottom Microtiter Plates (Dynex Technologies, Catalog no. 7205).
- 30 F. Discovery[®] HTRF Microplate Analyzer, Packard Instrument Company.
- G. 100 X Protease Inhibitor Cocktail (PIC): 1 mg/ml benzamidine, 0.5 mg/ml pepstatin, 0.5 mg/ml leupeptin, 0.5 mg/ml aprotinin.
- H. 10X Assay Buffer: 500 mM HEPES, pH 7.5, 1% PEG, mM EDTA, 1 mM EGTA, 1% BSA, 20 mM 9-Glycerol phosphate.
- 35 I. Quench Buffer: 50 mM HEPES pH 7.3, 16.6 mM EDTA, 0.1% BSA, 0.1% Triton X-100, 0.17 nM Lance labeled monoclonal antibody clone # 27, 0.0067 mg/ml SA-APC
- J. ATP/MgCl₂ working solution: 1X Assay buffer, 1 mM DTT, 1X PIC, 125 mM KCl, 5% Glycerol, 25 mM MgCl₂, 375 TM ATP

K. Enzyme working solution: 1X Assay buffer, 1 mM DTT, 1X PIC, 5% Glycerol, active Akt. The final enzyme concentrations were selected so that the assay was in a linear response range.

L. Peptide working solution: 1X Assay buffer, 1 mM DTT, 1X PIC, 5% Glycerol, 2

5 TM GSK3 biotinylated peptide # 3928

The reaction is assembled by adding 16 TL of the ATP/MgCl₂ working solution to the appropriate wells of a 96-well microtiter plate. Inhibitor or vehicle (1.0 TI) is added followed by 10 TI of peptide working solution. The reaction is started by adding 13 TI of the enzyme working solution and mixing. The reaction is allowed to proceed for 50 min and then stopped by the addition of 60 TI HTRF quench buffer. The stopped reactions were incubated at room temperature for at least 30 min and then read on the Discovery instrument.

Procedure for Streptavidin Flash Plate Assay:

Step 1:

A 1 µl solution of the test compound in 100% DMSO was added to 20 µl of 2X substrate solution (20 µM GSK3 Peptide, 300 µM ATP, 20 mM MgCl₂, 20 µCi / ml [³³P] ATP, 1X Assay Buffer, 5% glycerol, 1 mM DTT, 1X PIC, 0.1% BSA and 100 mM KCl). Phosphorylation reactions were initiated by adding 19 µl of 2X Enzyme solution (6.4 nM active Akt/PKB, 1X Assay Buffer, 5% glycerol, 1 mM DTT, 1X PIC and 0.1% BSA). The reactions were then incubated at room temperature for 45 minutes.

20 Step 2:

The reaction was stopped by adding 170 µl of 125 mM EDTA. 200 µl of stopped reaction was transferred to a Streptavidin Flashplate[®] PLUS (NEN Life Sciences, catalog no. SMP103). The plate was incubated for ≥10 minutes at room temperature on a plate shaker. The contents of each well was aspirated, and the wells rinsed 2 times with 200 µl TBS per well. The wells were then washed 3 times for 5 minutes with 200 µl TBS per well with the plates incubated at room temperature on a platform shaker during wash steps.

The plates were covered with sealing tape and counted using the Packard TopCount with the appropriate settings for counting [³³P] in Flashplates.

Procedure for Streptavidin Filter Plate Assay:

30 Step 1:

The enzymatic reactions as described in Step 1 of the Streptavidin Flash Plate Assay above were performed.

Step 2:

The reaction was stopped by adding 20 µl of 7.5M Guanidine Hydrochloride. 50 µl of the stopped reaction was transferred to the Streptavidin filter plate (SAM^{2™} Biotin Capture Plate, Promega, catalog no. V7542) and the reaction was incubated on the filter for 1-2 minutes before applying vacuum.

The plate was then washed using a vacuum manifold as follows: 1) 4 x 200 μ l/well of 2M NaCl; 2) 6 x 200 μ l/well of 2M NaCl with 1% H₃PO₄; 3) 2 x 200 μ l/well of diH₂O; and 4) 2 x 100 μ l/well of 95% Ethanol. The membranes were then allowed to air dry completely before adding scintillant.

5 The bottom of the plate was sealed with white backing tape, 30 μ l/well of Microscint 20 (Packard Instruments, catalog no. 6013621) was added. The top of the plate was sealed with clear sealing tape, and the plate then counted using the Packard TopCount with the appropriate settings for [³³P] with liquid scintillant.

Procedure for Phosphocellulose Filter Plate Assay:

10 Step 1:

The enzymatic reactions were performed as described in Step 1 of the Streptavidin Flash Plate Assay (above) utilizing KKGGRARTSSFAEPG (SEQ.ID.NO.: 16) as the substrate in place of biotin-GGRARTSSFAEPG.

Step 2:

15 The reaction was stopped by adding 20 μ l of 0.75% H₃PO₄. 50 μ l of stopped reaction was transferred to the filter plate (UNIFILTER™, Whatman P81 Strong Cation Exchanger, White Polystyrene 96 Well Plates, Polyfiltronics, catalog no. 7700-3312) and the reaction incubated on the filter for 1-2 minutes before applying vacuum.

20 The plate was then washed using a vacuum manifold as follows: 1) 9 x 200 μ l/well of 0.75% H₃PO₄; and 2) 2 x 200 μ l/well of diH₂O. The bottom of the plate was sealed with white backing tape, then 30 μ l/well of Microscint 20 was added. The top of the plate was sealed with clear sealing tape, and the plate counted using the Packard TopCount with the appropriate settings for [³³P] and liquid scintillant.

PKA assay:

25 Each individual PKA assay consists of the following components:

- A. 5X PKA assay buffer (200 mM Tris pH7.5, 100 mM MgCl₂, 5mM β -mercaptoethanol, 0.5 mM EDTA)
- B. 50 μ M stock of Kemptide (Sigma) diluted in water
- C. ³³P-ATP prepared by diluting 1.0 μ l ³³P-ATP [10 mCi/ml] into 200 Tl of a 50 μ M stock of unlabeled ATP
- 30 D. 10 μ l of a 70 nM stock of PKA catalytic subunit (UBI catalog # 14-114) diluted in 0.5 mg/ml BSA
- E. PKA/Kemptide working solution: equal volumes of 5X PKA assay buffer, Kemptide solution and PKA catalytic subunit.

35 The reaction is assembled in a 96 deep-well assay plate. The inhibitor or vehicle (10 Tl) is added to 10 Tl of the ³³P-ATP solution. The reaction is initiated by adding 30 Tl of the PKA/Kemptide working solution to each well. The reactions were mixed and incubated at room

temperature for 20 min. The reactions were stopped by adding 50 μ l of 100 mM EDTA and 100 mM sodium pyrophosphate and mixing.

The enzyme reaction product (phosphorylated Kemptide) was collected on p81 phosphocellulose 96 well filter plates (Millipore). To prepare the plate, each well of a p81 filter plate was filled with 75 mM phosphoric acid. The wells were emptied through the filter by applying a vacuum to the bottom of the plate. Phosphoric acid (75 mM, 170 μ l) was added to each well. A 30 μ l aliquot from each stopped PKA reaction was added to corresponding wells on the filter plate containing the phosphoric acid. The peptide was trapped on the filter following the application of a vacuum and the filters were washed 5 times with 75 mM phosphoric acid. After the final wash, the filters were allowed to air dry. Scintillation fluid (30 μ l) was added to each well and the filters counted on a TopCount (Packard).

PKC assay:

Each PKC assay consists of the following components:

- A. 10X PKC co-activation buffer: 2.5 mM EGTA, 4mM CaCl_2
- 15 B. 5X PKC activation buffer: 1.6 mg/ml phosphatidylserine, 0.16 mg/ml diacylglycerol, 100 mM Tris pH 7.5, 50 mM MgCl_2 , 5 mM 9-mercaptoethanol
- C. ^{33}P -ATP prepared by diluting 1.0 μ l ^{33}P -ATP [10 mCi/ml] into 100 μ l of a 100 μ M stock of unlabeled ATP
- D. Myelin basic protein (350 μ g/ml, UBI) diluted in water
- 20 E. PKC (50ng/ml, UBI catalog # 14-115) diluted into 0.5 mg/ml BSA
- F. PKC/Myelin Basic Protein working solution: Prepared by mixing 5 volumes each of PKC co-activation buffer and Myelin Basic protein with 10 volumes each of PKC activation buffer and PKC.

The assays were assembled in 96 deep-well assay plates. Inhibitor or vehicle (10 μ l) was added to 5.0 μ l of ^{33}P -ATP. Reactions were initiated with the addition of the PKC/Myelin Basic Protein working solution and mixing. Reactions were incubated at 30°C for 20 min. The reactions were stopped by adding 50 μ l of 100 mM EDTA and 100 mM sodium pyrophosphate and mixing. Phosphorylated Myelin Basic Protein was collected on PVDF membranes in 96 well filter plates and quantitated by scintillation counting.

Compounds of the instant invention described in the Schemes and Tables were tested in the assay described above and were found to have IC_{50} of ≤ 50 μ M against one or more of Akt1, Akt2 and Akt3.

EXAMPLE 5

Cell based Assays to Determine Inhibition of Akt/PKB

Cells (for example LnCaP or a PTEN^(-/-) tumor cell line with activated Akt/PKB) were plated in 100 mM dishes. When the cells were approximately 70 to 80% confluent, the cells were refed with 5 mls of fresh media and the test compound added in solution. Controls included untreated cells, vehicle treated cells and cells treated with either LY294002 (Sigma) or

wortmanin (Sigma) at 20 μ M or 200 nM, respectively. The cells were incubated for 2, 4 or 6 hrs, and the media removed, the cells were washed with PBS, scraped and transferred to a centrifuge tube. They were pelleted and washed again with PBS. Finally, the cell pellet was resuspended in lysis buffer (20 mM Tris pH8, 140 mM NaCl, 2 mM EDTA, 1% Triton, 1 mM Na

Pyrophosphate, 10 mM β -Glycerol Phosphate, 10 mM NaF, 0.5 mM NaVO₄, 1 μ M Microsystine, and 1x Protease Inhibitor Cocktail), placed on ice for 15 minutes and gently vortexed to lyse the cells. The lysate was spun in a Beckman tabletop ultra centrifuge at 100,000 x g at 4°C for 20min. The supernatant protein was quantitated by a standard Bradford protocol (BioRad) and stored at -70°C until needed.

Proteins were immunoprecipitated (IP) from cleared lysates as follows: For Akt1/PKB1, lysates are mixed with Santa Cruz sc-7126 (D-17) in NETN (100mM NaCl, 20mM Tris pH 8.0, 1mM EDTA, 0.5% NP-40) and Protein A/G Agarose (Santa Cruz sc-2003) was added. For Akt2/PKB β , lysates were mixed in NETN with anti-Akt2 agarose (Upstate Biotechnology #16-174) and for Akt3/PKBK, lysates were mixed in NETN with anti-Akt3 agarose (Upstate Biotechnology #16-175). The IPs were incubated overnight at 4°C, washed and separated by SDS-PAGE.

Western blots were used to analyze total Akt, pThr308 Akt1, pSer473 Akt1, and corresponding phosphorylation sites on Akt2 and Akt3, and downstream targets of Akt using specific antibodies (Cell Signaling Technology): Anti-Total Akt (cat. no. 9272), Anti-Phospho Akt Serine 473 (cat. no. 9271), and Anti-Phospho Akt Threonine 308 (cat. no. 9275). After incubating with the appropriate primary antibody diluted in PBS + 0.5% non-fat dry milk (NFDM) at 4 °C overnight, blots were washed, incubated with Horseradish peroxidase (HRP)-tagged secondary antibody in PBS + 0.5% NFDM for 1 hour at room temperature. Proteins were detected with ECL Reagents (Amersham/Pharmacia Biotech RPN2134).

EXAMPLE 6

Heregulin Stimulated Akt Activation

MCF7 cells (a human breast cancer line that is PTEN^{+/+}) were plated at 1x10⁶ cells per 100mm plate. When the cells were 70 – 80% confluent, they were refed with 5 ml of serum free media and incubated overnight. The following morning, compound was added and the cells were incubated for 1 – 2 hrs, after which time heregulin was added (to induce the activation of Akt) for 30 minutes and the cells were analyzed as described above.

EXAMPLE 7

Inhibition Of Tumor Growth

In vivo efficacy of an inhibitor of the growth of cancer cells may be confirmed by several protocols well known in the art.

Human tumor cell lines which exhibit a deregulation of the PI3K pathway (such as LnCaP, PC3, C33a, OVCAR-3, MDA-MB-468, A2780 or the like) are injected subcutaneously into the left flank of 6-10 week old female nude (also male mice [age 10-14

weeks] are used for prostate tumor xenografts [LnCaP and PC3]) mice (Harlan) on day 0. The mice are randomly assigned to a vehicle, compound or combination treatment group. Daily subcutaneous administration begins on day 1 and continues for the duration of the experiment. Alternatively, the inhibitor test compound may be administered by a continuous infusion pump.

5 Compound, compound combination or vehicle is delivered in a total volume of 0.2 ml. Tumors are excised and weighed when all of the vehicle-treated animals exhibited lesions of 0.5 - 1.0 cm in diameter, typically 4 to 5.5 weeks after the cells were injected. The average weight of the tumors in each treatment group for each cell line is calculated.

EXAMPLE 8

10 Spot Multiplex Assay

This procedure describes a sandwich immunoassay used to detect multiple phosphorylated proteins in the same well of a 96 well format plate. Cell lysates are incubated in 96-well plates on which different capture antibodies are placed on spatially distinct spots in the same well. Phosphorylation-specific rabbit polyclonal antibodies are added and the complex is

15 detected by an anti-rabbit antibody labeled with an electrochemiluminescent tag.

96-Well LNCaP plates +/- Compounds:

Spin in Beckman J6 1200 rpm 10 min, aspirate media. Add 50µl/well: TBS (Pierce #28376-20mM Tris pH 7.5, 150mM NaCl) + 1% Triton X-100 + Protease and Phosphatase Inhibitors. Wrap in plastic wrap, place in -70°C freezer until completely frozen.

20 Block Multiplex Plates (Meso Scale Discovery, Gaithersburg, MD) with 3% Blocker A in 1X Tris Wash Buffer, 150µl/well. Cover with plate sealer, incubate on Micromix shaker RT 2h (minimum). Wash with 1X RCM 51 (TTBS). Thaw cell lysate plates on ice, add 40µl lysate/well into blocked plates. Cover with plate sealer, incubate on Micromix shaker 4°C O/N, Wash with 1X RCM 51. Dilute Secondary Antibodies in 1% Blocker A in 1X Tris Wash Buffer:

25 Anti phospho AKT (T308), Anti phospho Tuberin (T1462), alone or in combination. Add 25µl/well, cover with plate sealer, incubate on Micromix shaker RT 3h. Wash with 1X RCM 51. Dilute Ru-GAR in 1% Blocker A in 1X Tris Wash Buffer. Add 25µl/well, cover with plate sealer, incubate on Micromix shaker RT 1h. Wash with 1X RCM 51. Dilute 4X Read Buffer T to 1X with Water, add 200µl diluted Read Buffer/well

30 Read on Sector 6000 Imager.

Protease and Phosphatase Inhibitors:

Microcystin-LR, Calbiochem # 475815 to 1 µM final concentration (stock=500µM)

Calbiochem # 524624, 100X Set I

Calbiochem # 524625, 100X Set II

35 Calbiochem # 539134, 100X Set III

Anti Phospho AKT (T308):

Cell Signaling Technologies # 9275

Anti Phospho Tuberin (T1462):

Covance Affinity Purified (Rabbits MS 2731/2732)

Ru-GAR = Ruthenylated Goat anti Rabbit

10X Tris Wash Buffer, Blocker A and 4X Read Buffer T

10X RCM 51 (10X TTBS, RCM 51)

5 1X = 20mM Tris pH 7.5, 140mM NaCl, 0.1% Tween-20

EXAMPLE 9

Cell-Based (In-vivo) Assay

This procedure describes a cell-based (in vivo) activity assay for the Akt serine/threonine kinase. Activated endogenous Akt is capable of phosphorylating a specific Akt substrate (GSK3 β) peptide which is biotinylated. Detection is performed by Homogeneous Time Resolved Fluorescence (HTRF) using a Europium Kryptate [Eu(K)] coupled antibody specific for the phosphopeptide and streptavidin linked XL665 fluorophore which will bind to the biotin moiety on the peptide. When the [Eu(K)] and XL665 are in proximity (i.e. bound to the same phosphopeptide molecule) a non-radiative energy transfer takes place from the Eu(K) to the XL665, followed by emission of light from XL665 at 665 nm.

The assay can be used to detect inhibitors of all three Akt isozymes (Akt1, Akt2, and Akt3) from multiple different species if specific antibodies to each exist.

MATERIALS AND REAGENTS

- A. Cell Culture Microtiter Flat Bottom 96 well plates, Corning Costar, Catalog no. 3598
- 20 B. Reacti-Bind Protein A Coated 96-well plates, Pierce, Catalog no 15130.
- C. Reacti-Bind Protein G Coated 96-well plates, Pierce, Catalog no 15131.
- D. Micromix 5 Shaker.
- E. Microfluor[®]B U Bottom Microtiter Plates, Dynex Technologies, Catalog no. 7205.
- F. 96 Well Plate Washer, Bio-Tek Instruments, Catalog no. EL 404.
- 25 G. Discovery[®] HTRF Microplate Analyzer, Packard Instrument Company.

BUFFER SOLUTIONS

- A. IP Kinase Cell Lysis Buffer: 1X TBS; 0.2% Tween 20; 1X Protease Inhibitor Cocktail III (Stock is 100X, Calbiochem, 539134); 1X Phosphatase Inhibitor Cocktail I (Stock is 100X, Calbiochem, 524624); and 1X Phosphatase Inhibitor Cocktail II (Stock is 100X, Calbiochem, 524625).
- 30 B. 10X Assay Buffer: 500 mM Hepes pH 7.5; 1% PEG; 1 mM EDTA; 1 mM EGTA; and 20 mM β -glycerophosphate.
- C. IP Kinase Assay Buffer: 1X Assay Buffer; 50 mM KCl; 150 μ M ATP; 10 mM MgCl₂; 5% Glycerol; 1 mM DTT; 1 Tablet Protease Inhibitor Cocktail per 50 ml Assay Buffer; and 0.1% BSA
- 35 D. GSK3 β Substrate Solution: IP Kinase Assay Buffer; and 500 nM Biotinylated GSK3 β peptide.
- E. Lance Buffer: 50 mM Hepes pH 7.5; 0.1% BSA; and 0.1% Triton X-100.

F. Lance Stop Buffer: Lance Buffer; and 33.3 mM EDTA.

G. Lance Detection Buffer: Lance Buffer; 13.3 µg/ml SA-APC; and 0.665 nM EuK Ab a-phospho (Ser-21) GSK3β

Multi-Step Immunoprecipitation Akt Kinase Assay

5 Day 1

A. Seed C33a cells Step: Plate 60,000 C33a cells/well in 96 well microtiter plate.

B. Incubate cells overnight at 37°C.

Day 2

10 D. Compound Addition Step: Add compounds in fresh media (alpha-MEM/10% FBS, room temp) to 96 well plate from above and incubate for 5 hrs in tissue culture incubator.

E. Cell Lysis Step: Aspirate media and add 100 µl of IP Kinase Cell Lysis Buffer.

F. Freeze 96 well microtiter plate at -70°C (NOTE: This step can be done for a minimum of 1 hour or overnight.)

Day 3

15 G. Coat Protein A/G 96 well plate Step: Add appropriate concentration of α-Akt antibody (Akt1, Akt2, or Akt3) in a 100 µl of PBS to the following wells:

α-Akt 1 (20 ng/well/100ul) B2 >>>>> B10 / rows B – G / Akt1 plate

α-Akt 2 (50 ng/well/100ul) B2 >>>>> B10 / rows B – G / Akt2 plate

Rabbit-IgG (150 ng/well/100 ul): B11 – G11 on every plate (Akt1 and Akt2)

20 H. Incubate in the cold room (+4°C) for 4 hours on the Micromix 5 (Form 20; Attitude 2) (NOTE: Attitude depends on which Micromix 5 machine).

I. Aspirate off α-Akt antibody solution and add 100 µl of PBS to each well.

J. Akt Immunoprecipitation Step: To the 100 µl of PBS from Step(I) add 5 µl of thawed cell lysate for Akt1 plates and 10 µl of thawed cell lysate for Akt2 plates. NOTE: Thaw cell lysate
25 on ice. Mix thawed lysate by pipetting up & down 10X before transferring to antibody plates. Keep the cell lysate plates on ice. After transfer of cell lysate to the antibody plates refreeze the cell lysate plates at -70°C.

K. Incubate in the cold room (+4°C) overnight on Micromix 5 shaker (form 20, attitude 3).

Day 4

30 L. Immunoprecipitation Plate Wash Step: Wash 96 well plates 1X with TTBS (RCM 51, 1X = 2 cycles) using the 96-Well Plate Washer. Fill wells with TTBS and incubate for 10 minutes. Wash 96 well plates 2X with TTBS. (NOTE: Prime plate washer before use: 1. Check buffer reservoirs, making sure they are full and 2. empty waste containers.

M. Manual Plate Wash Step: Add 180 µl of IP Kinase Assay buffer.

35 N. Start Akt Enzyme Reaction: Aspirate supernatant. Add 60 µl of GSK3β Substrate Solution.

O. Incubate for 2.5 hours on Micromix 5 shaker @ RT. NOTE: Time of incubation should be adjusted so that the ratio of Column 10 /Column 11 is not >10.

P. Combine 30 μ l of Lance Detection Buffer with 30 μ l of Lance Stop Buffer (60 μ l total/well) and add to Microfluor U bottom 96 well black plates.

Q. Stop Akt Enzyme Reaction: Transfer 40 μ l of Akt Enzyme Reaction Mix from Protein A/G 96 well plate from Step (O) to Microfluor U bottom 96 well black plates from Step (P).

- 5 U. Incubate at room temperature for 1-2 hrs on Micromix 5 shaker (form 20, attitude 3), then read with the Discovery HTRF Microplate Analyzer using Akt program.

IP Kinase Cell Lysis Buffer

100 μ l per well

	8 ml (1 Plate)		45 ml (6 Plates)
1X TBS	7744 μ l		NA
Tween 20	20 μ l		NA
1X Protease Inhibitor Cocktail III	80 μ l		NA
1X Phosphatase Inhibitor Cocktail I	80 μ l		450 μ l
1X Phosphatase Inhibitor Cocktail II	80 μ l		450 μ l
Microcystin LR (500X)			90 μ l

IP Kinase Assay Buffer

- 10 100 μ l per well

	8 ml (1 Plate)		50 ml (3 Plates)
10X Assay Buffer	800 μ l		5 ml
1 M KCl	400 μ l		2.5 ml
250 mM ATP	4.8 μ l		30 μ l
1M MgCl ₂	80 μ l		500 μ l
Glycerol	400 μ l		2.5 ml
1M DTT	8 μ l		50 μ l
Protease Inhibitor Cocktail	1 tablet/50 ml		1
10% BSA	80 μ l		500 μ l
di dH ₂ O	6227.2 μ l		38.9 ml

GSK3 β Substrate Solution

60 μ l per well

	5 ml (1 Plate)		7 ml
IP Kinase Assay Buffer	5 ml		-
1 mM GSK3 β peptide	2.5 μ l		3.5 μ l

Lance Stop Buffer

30 µl per well

	3 ml (1 Plate)	5 ml	5 ml
1X Lance Buffer EDTA 0.5 M	2800.2 µl 199.8 µl		


Lance Detection Buffer

30 µl per well

	3 ml (1 Plate)		5 ml
SA-APC (1 mg/ml in ddH ₂ O, dilute 1/75.2 in Lance Buffer)	40 µl		66.7 µl
EuK Ab α-phospho (Ser 21)GSK3β (680 nM, dilute 1/1133 in Lance Buffer)	2.7 µl		4.5 µl

WHAT IS CLAIMED IS:

1. A compound according to Formula B:



5 wherein,

A and B are independently selected from C or N, wherein at least one of A and B is N;

10 J and K are independently selected from C or N, wherein J and K are optionally substituted with R¹;

X, Y and Z are independently selected from C and N, wherein X, Y and Z are optionally substituted with R¹;

15 a is 0 or 1; b is 0 or 1; m is 0, 1 or 2;

R¹ is independently selected from (C=O)_aO_b(C₁-C₁₀)alkyl, (C=O)_aO_b(C₁-C₁₀)alkenyl, (C=O)_aO_b(C₁-C₁₀)alkynyl, (C=O)_aO_b(C₃-C₁₀)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_mNR⁷R⁸, S(O)_m(C₁-C₁₀)alkyl, S(O)_m(C₁-C₁₀)alkenyl, S(O)_m(C₁-C₁₀)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R⁶;

25 R³ and R^{3'} are independently selected from H, (C₁-C₁₀)alkyl, (C₁-C₁₀)alkenyl, (C₁-C₁₀)alkynyl, (C₃-C₁₀)cycloalkyl, aryl, heteroaryl, heterocyclyl, (C=O)NR⁷R⁸, halo, OH, CF₃, CO₂H, CN, and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₁₀)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NR⁷R⁸, or R³ and R^{3'} can be taken together to form a (C₃-C₁₀)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from H, (C₁-C₁₀)alkyl, (C₁-C₁₀)alkenyl, (C₁-C₁₀)alkynyl, (C₃-C₁₀)cycloalkyl, aryl, heteroaryl, heterocyclyl,

(C=O)NR⁷R⁸, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are further optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NR⁷R⁸;

5

R⁶ is selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_mNR⁷R⁸, S(O)_m(C₁-C₄)alkyl, S(O)_m(C₁-C₄)alkenyl, S(O)_m(C₁-C₄)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a};

10

R^{6a} is selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;

15

R⁷ and R⁸ are independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, N(R^b)₂, (C=O)N(R^b)₂, (C=S)N(R^b)₂, S(O)_mN(R^b)₂, S(O)_m(C₁-C₆)alkyl, S(O)_m(C₁-C₆)alkenyl, S(O)_m(C₁-C₆)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a}, or R⁷ and R⁸ can be taken together to form a (C₃-C₁₀)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from H, (C₁-C₁₀)alkyl, (C₁-C₁₀)alkenyl, (C₁-C₁₀)alkynyl, (C₃-C₁₀)cycloalkyl, aryl, heteroaryl, heterocyclyl, (C=O)NR⁷R⁸, halo, OH, CF₃, CO₂H, CN, and NR⁷R⁸; and

25

30


R^b is independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl and (C=O)_aO_b-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;

35


or a tautomer thereof;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

2. A compound according to Formula B of Claim 1, wherein,



is selected from:




10 a is 0 or 1; b is 0 or 1; m is 0, 1 or 2;

R^1 is independently selected from H, $(C=O)_aO_b(C_1-C_6)alkyl$, $(C=O)_aO_b(C_1-C_6)alkenyl$, $(C=O)_aO_b(C_1-C_6)alkynyl$, $(C=O)_aO_b(C_3-C_8)cycloalkyl$, $(C=O)_aO_b-aryl$, $(C=O)_aO_b-heteroaryl$, $(C=O)_aO_b-heterocyclyl$, halo, OH, CF_3 , CO_2H , CN, NR^7R^8 , $(C=O)NR^7R^8$, $S(O)_mNR^7R^8$, $S(O)_m(C_1-C_4)alkyl$, $S(O)_m(C_1-C_4)alkenyl$, $S(O)_m(C_1-C_4)alkynyl$, $S(O)_m(C_3-C_8)cycloalkyl$, $S(O)_m-aryl$, $S(O)_m-heteroaryl$, and $S(O)_m-heterocyclyl$, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^6 ;


20 R^3 and $R^{3'}$ are independently selected from H, $(C_1-C_4)alkyl$, $(C_1-C_4)alkenyl$, $(C_1-C_4)alkynyl$, $(C_3-C_5)cycloalkyl$, halo, OH, CF_3 , CO_2H , CN, phenyl and NH_2 , said alkyl, alkenyl, alkynyl, cycloalkyl and phenyl are optionally substituted with one or more substituents selected from H, $(C_1-C_4)alkyl$, $(C_1-C_4)alkenyl$, $(C_1-C_4)alkynyl$, $(C_3-C_5)cycloalkyl$, halo, OH, CF_3 , CO_2H , CN, phenyl and NH_2 , or R^3 and $R^{3'}$ can be taken together to form a $(C_3-C_5)cycloalkyl$, said
25 cycloalkyl is optionally substituted with one or more substituents selected from H, $(C_1-C_4)alkyl$, $(C_1-C_4)alkenyl$, $(C_1-C_4)alkynyl$, $(C_3-C_5)cycloalkyl$, halo, OH, CF_3 , CO_2H , CN, phenyl and NH_2 ;

- 5 R^6 is selected from $(C=O)_aO_b(C1-C6)alkyl$, $(C=O)_aO_b(C1-C6)alkenyl$, $(C=O)_aO_b(C1-C6)alkynyl$, $(C=O)_aO_b(C3-C8)cycloalkyl$, $(C=O)_aO_b-aryl$, $(C=O)_aO_b-heteroaryl$, $(C=O)_aO_b-heterocyclyl$, halo, OH, CF_3 , CO_2H , CN, NR^7R^8 , $(C=O)NR^7R^8$, $S(O)_mNR^7R^8$, $S(O)_m(C1-C4)alkyl$, $S(O)_m(C1-C4)alkenyl$, $S(O)_m(C1-C4)alkynyl$, $S(O)_m(C3-C8)cycloalkyl$, $S(O)_m-aryl$, $S(O)_m-heteroaryl$, and $S(O)_m-heterocyclyl$ said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a} ;
- 10 R^{6a} is selected from $(C=O)_aO_b(C1-C6)alkyl$, $(C=O)_aO_b(C1-C6)alkenyl$, $(C=O)_aO_b(C1-C6)alkynyl$, $(C=O)_aO_b(C3-C8)cycloalkyl$, $(C=O)_aO_b-aryl$, $(C=O)_aO_b-heteroaryl$, $(C=O)_aO_b-heterocyclyl$, halo, OH, CF_3 , CO_2H , CN and NH_2 , said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from $(C1-C6)alkyl$, $(C1-C6)alkenyl$, $(C1-C6)alkynyl$, halo, OH, CF_3 , CO_2H , CN, phenyl and NH_2 ;
- 15 R^7 and R^8 are independently selected from H, $(C=O)_aO_b(C1-C6)alkyl$, $(C=O)_aO_b(C1-C6)alkenyl$, $(C=O)_aO_b(C1-C6)alkynyl$, $(C=O)_aO_b(C3-C8)cycloalkyl$, $(C=O)_aO_b-aryl$, $(C=O)_aO_b-heteroaryl$, $(C=O)_aO_b-heterocyclyl$, halo, OH, CF_3 , CO_2H , CN, $N(R^b)_2$, $(C=O)N(R^b)_2$, $(C=S)N(R^b)_2$, $S(O)_mN(R^b)_2$, $S(O)_m(C1-C6)alkyl$, $S(O)_m(C1-C6)alkenyl$, $S(O)_m(C1-C6)alkynyl$, $S(O)_m(C3-C8)cycloalkyl$, $S(O)_m-aryl$, $S(O)_m-heteroaryl$, and $S(O)_m-heterocyclyl$, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a} ; and
- 20 R^b is independently selected from H, $(C=O)_aO_b(C1-C6)alkyl$, $(C=O)_aO_b(C1-C6)alkenyl$, $(C=O)_aO_b(C1-C6)alkynyl$, $(C=O)_aO_b(C3-C8)cycloalkyl$, $(C=O)_aO_b-aryl$, $(C=O)_aO_b-heteroaryl$ and $(C=O)_aO_b-heterocyclyl$, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from $(C1-C6)alkyl$, $(C1-C6)alkenyl$, $(C1-C6)alkynyl$, halo, OH, CF_3 , CO_2H , CN, phenyl and NH_2 ;
- 25 or a tautomer thereof;
- 30 or a pharmaceutically acceptable salt or a stereoisomer thereof.

3. A compound according to Formula B of Claim 1, wherein,



is selected from



a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; n is 0, 1, 2 or 3;

5

R¹ is independently selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_mNR⁷R⁸, S(O)_m(C₁-C₄)alkyl, S(O)_m(C₁-C₄)alkenyl, S(O)_m(C₁-C₄)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R⁶;

10

R³ and R^{3'} are independently selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl and phenyl are optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂, or R³ and R^{3'} can be taken together to form a (C₃-C₅)cycloalkyl, said cycloalkyl is optionally substituted with one or more substituents selected from H, (C₁-C₄)alkyl, (C₁-C₄)alkenyl, (C₁-C₄)alkynyl, (C₃-C₅)cycloalkyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;

20

R⁶ is selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_mNR⁷R⁸, S(O)_m(C₁-C₄)alkyl, S(O)_m(C₁-C₄)alkenyl, S(O)_m(C₁-C₄)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a};

30

R^{6a} is selected from (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN and NH₂, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;

35



R⁷ and R⁸ are independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, N(R^b)₂, (C=O)N(R^b)₂, (C=S)N(R^b)₂, S(O)_mN(R^b)₂, S(O)_m(C₁-C₆)alkyl, S(O)_m(C₁-C₆)alkenyl, S(O)_m(C₁-C₆)alkynyl, S(O)_m(C₃-C₈)cycloalkyl, S(O)_m-aryl, S(O)_m-heteroaryl, and S(O)_m-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a}; and

R^b is independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_aO_b(C₁-C₆)alkenyl, (C=O)_aO_b(C₁-C₆)alkynyl, (C=O)_aO_b(C₃-C₈)cycloalkyl, (C=O)_aO_b-aryl, (C=O)_aO_b-heteroaryl and (C=O)_aO_b-heterocyclyl, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from (C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₁-C₆)alkynyl, halo, OH, CF₃, CO₂H, CN, phenyl and NH₂;

or a tautomer thereof;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

4. A compound according to Formula B of Claim 1, wherein,



a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; n is 0, 1 or 2;

R¹ is independently selected from (C=O)_aO_b(C₁-C₆)alkyl, (C₁-C₆)alkenyl, (C₃-C₈)cycloalkyl, O_b-aryl, O_b-heteroaryl, (C=O)_aO_b-heterocyclyl, halo, OH, CF₃, CO₂H, CN, NR⁷R⁸, (C=O)NR⁷R⁸, S(O)_m(C₁-C₄)alkyl, said alkyl, alkenyl, cycloalkyl, aryl, heteroaryl and heterocyclyl are optionally substituted with one or more substituents selected from R⁶;

R³ and R^{3'} are H, or R³ and R^{3'} can be taken together to form cyclobutyl, said cyclobutyl is optionally substituted with one or more substituents selected from H, OH, methyl and cyclopropyl;

R⁶ is selected from O_b(C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, O_b-aryl, halo, OH, CF₃, NR⁷R⁸, said alkyl and cycloalkyl are optionally substituted with one or more substituents selected from R^{6a};

- 5 R^{6a} is selected from O_b(C₁-C₆)alkyl, aryl, OH and NH₂, said alkyl and aryl are optionally substituted with one or more substituents selected from halo;

- R⁷ and R⁸ are independently selected from H, (C=O)_aO_b(C₁-C₆)alkyl, (C=O)_a(C₃-C₈)cycloalkyl, (C=O)_a-aryl, heterocyclyl, (C=O)N(R^b)₂, (C=S)N(R^b)₂, S(O)_m(C₁-C₆)alkyl, and S(O)_m-aryl, said alkyl, cycloalkyl, aryl and heterocyclyl are optionally substituted with one or more substituents selected from R^{6a}; and

R^b is independently selected from H, (C₁-C₆)alkyl and aryl;

- 15 or a tautomer thereof;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

5. A compound which is selected from:

- 20 1-[4-(6-phenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine;
 1-[4-(2-isopropyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine;
 1-[4-(2-methyl-6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine;
 1-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine;
 25 3-amino-3-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]-1-methylcyclobutanol;
 3-amino-1-methyl-3-[4-(2-methyl-3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanol;
 1-[4-(2-methyl-3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]cyclobutanamine;
 3-amino-1-methyl-3-[4-(2-methyl-6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol;
 30 3-amino-1-methyl-3-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol;
 1-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine;
 1-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanamine;
 3-amino-1-methyl-3-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]cyclobutanol;
 35 1-[4-(3,6-diphenylimidazo[1,2-*a*]pyrimidin-7-yl)phenyl]methanamine;
 1-[4-(6-phenylpyrazolo[1,5-*a*]pyrimidin-5-yl)phenyl]methanamine;
 1-[4-(2-methyl-6-phenyl[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)phenyl]methanamine;
 4-(6-Phenyl-[1,2,4]triazolo[1,5-*a*]pyrimidin-5-yl)-benzylamine;

- 3-Amino-1-cyclopropyl-3-[4-(6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutanol;
 4-(6-Phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-benzylamine;
 1-[4-(6-Phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-[4-(3-Bromo-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 5 1-[4-(6-Phenyl-3-vinyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-{4-[3-(2-Methoxy-phenyl)-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutylamine;
 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-3-carbonitrile;
 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-3-carboxylic acid amide;
 1-[4-(3-Chloro-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 10 1-[4-(3-Methyl-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-{4-[3-(2-Cyclopropyl-vinyl)-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-
 cyclobutylamine;
 1-[4-(2,6-Diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-[4-(3-Bromo-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 15 1-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-
 cyclobutylamine;
 1-[4-(2,6-Diphenyl-3-vinyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-[4-(2-Ethyl-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-[4-(6-Phenyl-2-trifluoromethyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 20 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-2-carboxylic acid ethyl
 ester;
 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-2-carboxylic acid;
 7-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-imidazo[1,2-a]pyrimidine-2-carboxylic acid methyl
 ester;
 25 1-{4-[2-(4-Fluoro-phenyl)-6-phenyl-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-cyclobutylamine;
 1-[4-(6-Phenyl-2-p-tolyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-[4-(3-Bromo-6-phenyl-2-trifluoromethyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-
 cyclobutylamine;
 1-[4-(3-Methyl-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine
 30 1-[4-(2,3,6-Triphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-[4-(3-Cyclopropyl-2,6-diphenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 1-{4-[2,6-Diphenyl-3-(2,3,4-trifluoro-phenyl)-imidazo[1,2-a]pyrimidin-7-yl]-phenyl}-
 cyclobutylamine;
 1-[4-(2,6-Diphenyl-3-propenyl-imidazo[1,2-a]pyrimidin-7-yl)-phenyl]-cyclobutylamine;
 35 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-ylamine;
 N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-
 acetamide;
 1-[4-(2,6-Diphenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;

- 1-[4-(2-Phenoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
 1-[4-(2-Morpholin-4-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
 1-[4-(6-Phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
 1-[4-(2-Methyl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
 5 Ethanesulfonic acid {5-[4-(1-amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-amide;
 N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-benzenesulfonamide;
 N-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-
 10 benzamide;
 Cyclopropanecarboxylic acid {5-[4-(1-amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-amide;
 1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-3-phenyl-urea;
 15 1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-3-phenyl-thiourea;
 {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-carbamic acid ethyl ester;
 1-{4-[2-(6-Methoxy-pyridin-2-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-
 20 cyclobutylamine;
 1-[4-(6-Phenyl-2-pyrazin-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
 1-{4-[2-(2,3-Dimethyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
 1-{4-[2-(3,4-Difluoro-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-
 25 cyclobutylamine;
 1-{4-[2-(5-Fluoro-2-methyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
 1-[4-(6-Phenyl-2-pyridin-4-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
 4-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-
 30 phenylamine;
 1-[4-(6-Phenyl-2-p-tolyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
 1-{4-[2-(4-Methoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
 1-[4-(6-Phenyl-2-pyridin-3-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
 35 1-{4-[2-(2-Methoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
 1-{4-[2-(3-Chloro-4-fluoro-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;

- 1-{4-[2-(2-Fluoro-4-trifluoromethyl-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-{4-[6-Phenyl-2-(2,3,4-trifluoro-phenyl)-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 5 1-{4-[2-(2,4-Dimethoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-[4-(2-Furan-3-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(6-Phenyl-2-thiophen-3-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(6-Phenyl-2-thiophen-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 10 1-{4-[2-(6-Methoxy-naphthalen-2-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-[4-(2-Dibenzofuran-4-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 4-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-phenol;
- 15 1-[4-(6-Phenyl-2-m-tolyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-{4-[2-(4-Phenoxy-phenyl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-[4-(2-Biphenyl-2-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(2-Benzothiazol-2-yl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 20 1-[4-(6-Phenyl-2-pyridin-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(6-Phenyl-2-thiazol-2-yl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-{4-[6-Phenyl-2-(pyridin-4-yloxy)-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 25 1-[4-(2-Methoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(2-Phenoxy-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(2-Methylsulfanyl-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-{4-[2-(Naphthalen-2-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 30 1-{4-[2-(Biphenyl-2-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-{4-[2-(Isoquinolin-7-yloxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-{4-[2-(4-Phenoxy-phenoxy)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 35 1-{4-[2-(3,4-Dihydro-2H-quinolin-1-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;

- 1-{4-[2-(2,3-Dihydro-indol-1-yl)-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-methylamine;
- 5 {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-dimethylamine;
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-pyridin-3-ylamine;
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-diethylamine;
- 10 N1-{5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-ethane-1,2-diamine;
- 2-(5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl)-methylamino)-ethanol;
- 15 {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-benzylmethylamine;
- 1-[4-(2-{4-[Bis-(4-chloro-phenyl)-methyl]-piperazin-1-yl}-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-methylphenylamine;
- 20 {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-(2-methoxyethyl)-amine;
- 3-Amino-3-[4-(2,6-diphenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol;
- 3-Amino-1-methyl-3-[4-(6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutanol;
- 25 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methyl ester;
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid;
- 2-(5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-propan-2-ol;
- 30 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methylamide;
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-diphenylmethanol;
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid
- 35 amide;
- {5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-yl}-morpholin-4-yl-methanone;

- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid phenethyl-amide;
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid diethylamide;
- 5 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid phenylamide;
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid benzyl-methyl-amide;
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methoxy-methyl-amide;
- 10 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-[1,2,4]triazolo[1,5-a]pyrimidine-2-carboxylic acid methyl-phenyl-amide;
- 1-[4-(2,6-Diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(2,3,6-Triphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 15 1-[4-(2,3,6-Triphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(6-Phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(2-Methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 5-[4-(1-Amino-cyclobutyl)-phenyl]-6-phenyl-pyrazolo[1,5-a]pyrimidin-2-ol;
- 1-[4-(6-Phenyl-2-p-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 20 1-{4-[2-(4-Methoxy-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-[4-(6-Phenyl-2-o-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-{4-[2-(4-Chloro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-{4-[2-(3-Fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-{4-[6-Phenyl-2-(2,4,6-trimethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 25 1-{4-[2-(4-Fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-[4-(3-Chloro-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(3-Bromo-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(2,6-Diphenyl-3-pyridin-4-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 30 1-[4-(3-Methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-{4-[3-(2,4-Difluoro-phenyl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;
- 1-[4-(3-Methyl-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-[4-(3,6-Diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 35 1-[4-(6-Phenyl-3-pyrimidin-5-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
- 1-{4-[3-(4-Methoxy-phenyl)-2-methyl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-cyclobutylamine;

- 1-{4-[3-(2-Methoxy-pyrimidin-5-yl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-
cyclobutylamine;
1-{4-[6-Phenyl-3-(4-trifluoromethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-
cyclobutylamine;
5 1-[4-(3-Naphthalen-1-yl-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-cyclobutylamine;
1-{4-[3-(2-Fluoro-5-trifluoromethyl-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-
cyclobutylamine;
1-{4-[3-(3-Chloro-4-fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-
cyclobutylamine;
10 1-{4-[3-(3-Fluoro-5-isopropoxy-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-
cyclobutylamine;
1-{4-[6-Phenyl-3-(2,3,4-trifluoro-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-
cyclobutylamine;
3-Amino-3-[4-(2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-cyclobutanol;
15 3-Amino-3-[4-(3-bromo-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-
cyclobutanol;
3-Amino-1-methyl-3-[4-(3-methyl-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-
cyclobutanol;
3-Amino-1-methyl-3-[4-(6-phenyl-2-o-tolyl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-
20 cyclobutanol;
3-Amino-3-{4-[2-(4-fluoro-phenyl)-6-phenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-1-methyl-
cyclobutanol;
3-Amino-3-[4-(2,6-diphenyl-3-pyridin-3-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-
cyclobutanol;
25 3-Amino-3-{4-[3-(3-amino-phenyl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-phenyl}-1-
methyl-cyclobutanol;
3-Amino-3-[4-(2,6-diphenyl-3-pyrimidin-5-yl-pyrazolo[1,5-a]pyrimidin-5-yl)-phenyl]-1-methyl-
cyclobutanol; and
3-Amino-3-{4-[3-(2-amino-pyrimidin-5-yl)-2,6-diphenyl-pyrazolo[1,5-a]pyrimidin-5-yl]-
30 phenyl}-1-methyl-cyclobutanol;

or a tautomer thereof;

or a pharmaceutically acceptable salt or stereoisomer thereof.

35

6. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 1.

7. The use of the compound according to Claim 1 for the preparation of a medicament useful in the treatment or prevention of cancer in a mammal in need of such treatment.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 10/28989

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/44 (2010.01)

USPC - 514/300

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC: 514/300

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC: 514/303; 514/301, 546/113, 546/117, 546/118 (keywords below)Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
PubWEST (USPT, PGPB, EPAB, JPAB), Google Patents/Scholar: Akt, imidazol, pyrazolo, triazolo, pyrimidinyl, biphenyl

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2007/0083044 A1 (Paruch et al.) 12 April 2007 (12.04.2007) para [0017]- [0023], [0031], [0034], [0205], [0216], [0242]	1-7
Y	US 7,399,764 B2 (Duggan et al.) 15 July 2008 (15.07.2008) col 3, ln 21-45	1-7

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

19 May 2010 (19.05.2010)

Date of mailing of the international search report

01 JUN 2010

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774