




(11) (21) (C) **2,012,876**
(22) 1990/03/22
(43) 1990/09/22
(45) 2001/02/13

(72) Le Grazie, Cristina, IT

(73) KNOLL FARMACEUTICI S.P.A., IT

(51) Int.Cl.⁵ A61K 31/505

(30) 1989/03/22 (19867 a/89) IT

(54) **UTILISATION DE L'ACIDE 5-METHYLE-
TETRAHYDROFOLIQUE ET DE L'ACIDE
5- FORMYLE-TETRAHYDROFOLIQUE DANS LE
TRAITEMENT DES AGITATIONS MENTALES ORGANIQUES**

(54) **USE OF 5-METHYLTETRAHYDROFOLIC ACID AND
5-FORMYLTETRAHYDROFOLIC ACID IN THE
TREATMENT OF ORGANIC MENTAL DISTURBANCES**

(57) This invention relates to pharmaceutical compositions containing 5-methyltetrahydrofolic acid, 5-formyltetrahydrofolic acid and their pharmaceutical acceptable salts in controlled-release form which are active in the therapy of organic mental disturbances and in particular in the treatment of senile and presenile primary degenerative dementia of Alzheimer type and multiinfarctual demetia.

20 128 76

ABSTRACT

This invention relates to pharmaceutical compositions containing 5-methyltetrahydrofolic acid, 5-formyltetrahydrofolic acid and their pharmaceutical acceptable salts in controlled-release form which are active in the therapy of organic mental disturbances and in particular in the treatment of senile and presenile primary degenerative dementia of Alzheimer type and multiinfarctual demetia.

USE OF 5-METHYLTETRAHYDROFOLIC ACID AND 5-FORMYLTETRAHYDROFOLIC
ACID IN THE TREATMENT OF ORGANIC MENTAL DISTURBANCES

5 This invention relates to pharmaceutical compositions
containing 5-methyltetrahydrofolic acid (MTHF), 5-
formyltetrahydrofolic acid (FTHF) and their pharmaceutically
acceptable salts in the preparation of pharmaceutical
compositions in controlled-release form active in the therapy
10 of organic mental disturbances.

In particular, the pharmaceutical compositions according to the
present invention are active in the treatment of senile and
presenile primary degenerative dementia of Alzheimer type and in
the treatment of multiinfarctual dementia.
15

In the present text, for greater clarity and simplicity, the
expression 5-methyltetrahydrofolic acid and the initials MTHF
refer to compounds having the following complete chemical
denomination: (+)-L-5-methyl-5,6,7,8-tetrahydrofolic acid and
20 (-)-L-5-methyl-5,6,7,8-tetrahydrofolic acid and their salts,
whereas the expression 5-formyltetrahydrofolic acid and the
initials FTHF refer to compounds having the following complete
chemical denomination: (+)-L-5-formyl-5,6,7,8-tetrahydrofolic acid
and (-)-L-5-formyl-5,6,7,8-tetrahydrofolic acid and their salts.
25

No effective treatment is currently available in the therapy of

- 2 -

organic mental disturbances, and in the particular case of dementia of Alzheimer type the drugs currently under study are still far from being unequivocally considered active (Davis K.L., Mohs R.C.: Cholinergic drugs in Alzheimer's disease; New Engl. J. Med. 315: 1286-7, 1986).

With regard to the numerous drugs used to treat the so-called "senile brain", their effectiveness has up to the present time been evaluated more in terms of the modifications induced by the drug in the cerebral flow and in the electroencephalogram, rather than in terms of clinical response. It has therefore not been established whether these physiological variations correspond to a real improvement in the elderly patient.

5-methyltetrahydrofolic acid, 5-formyltetrahydrofolic acid and their salts are a group of substances pertaining to the vitamin B complex, structurally related to pteroylglutamic acid (folic acid). This acid, which is not synthesized by the cells of mammals, is of particular biological importance because it intervenes in a series of chemical reactions involving transfer of monocarbon groups and in particular in the synthesis of the purine ring and thymidylate and in the neogenesis of methyl groups. In the circulating blood the folate pool is mostly represented by MTHF, but also by FTHF. MTHF represents the main form of folate transport in the blood. At the choroid plexus level it passes from the blood to the body fluid and from here, by passive diffusion, into the tissue and nerve cells. In the central nervous system the folates and in particular MTHF participate in basic biochemical processes by intervening in the synthesis of S-adenosyl-L-methionine (SAdMe), in the metabolism of certain amino-acids (glycine, serine, glutamic acid), in the modulatory activity of monoaminergic transmission systems (noradrenaline, serotonin, dopamine), in nucleic acid synthesis and in ATP and GTP production.

35

The therapeutic use of folic acid and its cofactors has up to the present time been limited to the prevention and treatment of body

deficiencies of this vitamin, i.e. to the treatment of hypofolatememic subjects.

An object of the present invention is to allow effective therapy of organic mental disturbances by providing pharmaceutical compositions which possess demonstrated clinical effectiveness in the therapy of such disturbances and are free of side effects.

10 Accordingly, the present invention relates to a pharmaceutical composition in controlled-release form for use in the therapy of an organic mental disturbance, containing as active principle MTHF or FTHF, or one or more of their pharmaceutically acceptable salts.

We have also surprisingly found that for highest effectiveness the pharmaceutical compositions of controlled-release type have an active principle release time varying from 15 minutes to 8 hours, and preferably varying from 20 to 60 minutes, and contain from 5 to 200 mg and preferably from 10 to 50 mg of MTHF or FTHF or their pharmaceutically acceptable salts.

20

The organic mental disturbances are diagnosed on the basis of criteria contained in the Diagnostic and Statistical Manual of Mental Disorders, Third Edition Revised (DSM-III-R), published by the American Psychiatric Association in 1987. This manual describes organic mental disturbances according to the following codes:

290.21 Senile primary degenerative dementia of Alzheimer type with depression;

290.13 Presenile primary degenerative dementia of Alzheimer type with depression;

30 290.43 Multiinfarctual dementia with depression.

According to the definition given to it by Stone in the American Psychiatric Glossary (Am. Psych. Press, Washington, 6th Edition, 1988 p. 46), dementia is "an organic mental illness in which there is deterioration of previously acquired intellectual capacity, this deterioration being of sufficient severity to interfere with work or social activities. The most important symptom is memory disturbance. In addition abstract thought, capacity for judgement and control of impulses are compromised, and/or there is personality change.

40 Dementia can be progressive, stationary or

reversible depending on the morbid symptoms and on the availability of effective treatment". The diagnostic criteria for dementia are given in Table 1, taken from DMS-III-R.

5

TABLE 1

Diagnostic criteria for dementia

- A. Loss of intellectual faculties to such an extent as to interfere with social or professional activities.
- B. Memory deficit.
- 10 C. At least one of the following elements:
 - 1) deficit of abstract thought, encountered in literal interpretation of proverbs, in incapacity to recognise similarities and differences between related words, in difficulty in defining words and concepts, and in other similar tests;
 - 15 2) deficit of critical judgement;
 - 3) other disturbances of the higher cortical functions, such as aphasia (language disturbance related to cerebral malfunction), apraxia (inability to execute motor activities notwithstanding soundness of comprehension and motricity), agnosia (inability to
 - 20 recognise or identify objects notwithstanding soundness of sensitive functions), "constructive apraxia" (such as inability to copy three-dimensional figures and put blocks together, or to arrange sticks in predetermined designs);
 - 4) personality changes, such as alteration or accentuation of
 - 25 premorbid states.
- D. Absence of obnubilation of consciousness i.e. lack of response to the diagnostic criteria for Delirium or Intoxication, although these symptoms may mutually superimpose).
- E. One of the following elements:
 - 30 1) demonstration of a specific organic factor etiologically related to the disturbance, on the basis of anamnesis, clinical examination or laboratory examinations;
 - 2) in the absence of such demonstration, presumption of an
 - 35 organic factor necessary for development of the syndrome, when there are proper reasons for excluding situations other than organic mental disturbances, and when the behavioural alteration is represented by an intellectual deficit in different areas.

The characteristics and advantages of the present invention will be more apparent from the summary description of a significant clinical trial selected from those carried out using the compositions according to the present invention.

5

Clinical trial

The purpose of this trial was to verify the therapeutic effects of prolonged administration of controlled-release MTHF compared with the administration of MTHF not in controlled-release form and of a placebo to elderly subjects affected by organic mental disturbances with depression of mood in accordance with the definitions given in DSM-III-R. This was a multi-centre perspective randomized trial of double-blind controlled type. The trial was carried out in 7 centres, each of which involved 30 patients giving a total of 210 patients. The patients were divided randomly into three sub-groups, these receiving MTHF in controlled-release form, MTHF not in controlled-release form and placebo respectively.

On the basis of inclusion criteria patients of both sexes 65 years of age and over, institutionalised for at least 3 months, took part. The patients had to have shown evident clinical signs of cerebral deterioration for at least one year. In particular, the presence of at least 3 of the following symptoms was required: mental confusion with compromised cognition, absence or reduction of appetite, ease of exhaustion, fear, irritability, impulsiveness, absence of cooperation, emotive weakness, poor sociability, anxiety and depression. The criteria for admission were a point score of between 10 and 24 on the MMSE (Mini Mental State Examination) scale and a point score of 18 or more on the Hamilton depression scale (HAM-D).

The criteria for exclusion were represented by: age less than 65 years, MMSE point score less than 10 or greater than 24, HAM-D point score less than 18, presence of serious cardiovascular, renal, respiratory, hepatic, dismetabolic, hematological or neoplastic pathology. The following patients were also excluded:

those with anamnesis of epilepsy or convulsive manifestations, those with Parkinson's disease at stages III, IV or V, those with self-wounding tendencies, those with brain damage of traumatic or infective origin, and those with functional psychosis. Also
5 excluded were those patients under treatment with drugs which could interfere with the results of the present research. After a wash-out period of 15 days in which all pharmacological treatment for the pathology under examination was suspended, the patients taking part were assigned in a consecutive and random manner to
10 the treatment, which was carried out over the scheduled period of 90 days. The MTHF in controlled-release form (average release time 1 hour) was administered orally at a dose of 50 mg/day in a single administration, the MTHF not in controlled-release form was administered orally at a dose of 50 mg/day in a single
15 administration, and the placebo likewise.

To evaluate the effects of the treatment certain psychiatric scales particularly suitable for investigating behaviour, autonomy and depressed humour were used. The evaluation scales used were
20 as follows:

1) Geriatric Rating Scale (GRS) of Plutchik (Plutchik et al., J. Amer. Geriatric Soc., 1970, 18: 491) which evaluates the alterations in the physical, mental and social state which
25 influence daily living, i.e. the self-sufficiency and behaviour of the patient. This scale comprises 31 items graduated from 0 (normal) to 2 (serious). Mostly, these items are combined into symptom groups in relation to the aspects to be explored. Thus self-sufficiency combines 6 items, sleep disturbance 3, global
30 deficit 5, initiative 4, and aggressivity and sociability 6.

2) Dementia Rating Scale (DRS) of Gottfries (Gottfries et al., Clin. Neuropharmacol., 1984, 7/1: 12) with 26 items graduated from 0 to 6 indicating increasing severity of the syndrome. It is
35 divided into the following 4 sections, which evaluate:

a) psychiatric symptoms (confusion, irritability, anxiety, distress, depression of humour, restlessness;

- b) emotion;
- c) bodily movement,
- d) mental functions (memory, attention, vigilance).

- 5 3) Hamilton depression evaluation scale (HAM-D) with 21 items
 (Hamilton, British J. Med. Psychol., 1959, 32: 50).

Evaluation of the individual parameters of the aforesaid scales
was done before commencement of treatment (basal value) and after
10 21, 45 and 90 days of therapy (T21, T45 and T90).

The following psychometric tests were used to evaluate the
individual state of performance:

- 15 1) Wechsler Adult Intelligence Scale Test (W.A.I.S.) (Wechsler,
 "Wechsler Adult Intelligence Scale Manual", Psychological
 Corporation, New York, 1955), which consists of reconstructing
 human figures divided into 5 segments.
- 20 2) Randt Memory Test (Randt et al., Clin. Neuropsychol., 1980,
 2: 184) limited to the acquisition and repeating of 5 words; it
 allows evaluation both of any modifications in the various memory
 aspects and the effectiveness of pharmacological treatment on this
 function.
- 25 3) Toulouse-Pieron test (Andreoli et al., Il Fracastoro, 1975.
 68/Suppl. No.1-2: 71) for visual exploration.
- 30 4) Semantic Verbal Memory Test (Villardita, Intern.
 Neuropsychol. Soc. 7th Europ. Conference, Aachen, 13.06.1984)
 which consists of the immediate and deferred (20 minutes) recall
 of 15 frequently used words pertaining to 3 semantic categories.
- 35 5) Digit Span (Wechsler, "Wechsler Adult Intelligence Scale
 Manual", Psychological Corporation, New York, 1955), which
 consists of the forward and backward repeating of numbers of
 increasing length read to the patient by the examiner.

The psychometric tests were carried out before commencement of treatment (basal value) and after 21 and 90 days of treatment (T21 and T90). During the course (T21 and T45) and at the end (T90) of treatment the medical practitioner, with the cooperation of the patient's relatives and the paramedic personnel, expressed a global judgement on the effects of the therapy.

The variations in the individual parameters were evaluated during and at the end of the treatment, using variance analysis based on two evaluation criteria (drug and time), with multiple Tukey comparisons.

Having undergone the above selection procedure, 184 patients were available for evaluation divided as follows:

15	Group using MTHF in controlled-release form	60
	Group using MTHF not in controlled-release form	62
	Group using placebo	62

The demographic and clinical characteristics of the patients are shown in Table 2, from which the similarity between the groups can be seen. The values in parentheses represent standard deviation (S.D.).

TABLE 2

25	<u>Characteristics of the 184 patients evaluated</u>			
		MTHF controlled release	MTHF not controlled release	Placebo
		60	62	62
	Sex M/F	27/33	25/37	30/32
30	Average age (years)	74.6	75.3	73.9
	(S.D.)	(5.6)	(4.8)	(5.3)
	Average weight (kg)	63.6	62.7	59.8
35	(S.D.)	(10.3)	(11.2)	(9.4)
	Average illness duration (years)	4.5	5.1	4.7

(S.D.)	(7.2)	(8.2)	(8.4)
--------	-------	-------	-------

In the figures cited below, the MTHF in controlled-release form is indicated by MTHF R C and the MTHF not in controlled-release form is indicated by MTHF non R C.

Figure 1 shows the results obtained using the Plutchik geriatric evaluation scale (G.R.S.).

In Figures 1a to 1e the horizontal axis represents time in days and the vertical axis represents the average point score ($\pm$ standard error) obtained on the Plutchik G.R.S. for the 3 groups of patients under treatment. The individual graphs represent the following items: self-sufficiency (Figure 1a), sleep disturbance (Figure 1b), global deficit (Figure 1c), initiative (Figure 1d) and unadaptable behaviour (Figure 1e).

It can be seen that whereas the groups of patients treated with MTHF not in controlled-release form or with placebo show no significant difference from the basal values at T0 (b) and at the end of treatment (T90), the group of patients treated with MTHF in controlled-release form shows a statistically significant difference even at T21 for the global deficit (Figure 1c) and for initiative (Figure 1d), and at T45 for self-sufficiency. In contrast, sleep disturbance (Figure 1b) and unadaptable behaviour (Figure 1e) show no significant differences after the treatment period compared with the basal values. Analysis of the variance shows the existence of interaction between treatment and time ($p < 0.01$) for the above parameters with statistically significant difference.

Figure 2 shows the results obtained using the Gottfries Dementia Scale Rating (DRS) in which the horizontal axis represents time in days and the vertical axis represents the average point score ($\pm$ standard error), the results being clearly most apparent in the group receiving MTHF in controlled-release form. By T45 there is a significant difference compared with the treatments with placebo

and MTHF not in controlled-release form ($p < 0.05$), this difference increasing further by T90 ($p < 0.01$).

Figure 3 shows the results obtained using the Hamilton Rating Scale (HAM-D) for depression, the horizontal axis representing time in days and the vertical axis the average point score ($\pm$ standard error). Treatment with MTHF in controlled-release form acts on temperament to clearly improve depressive symptomatology from T21. At the end of treatment (T90) the average point score on the HAM-D scale, equal to 20.3 in the basal evaluation, passes to 8.3, corresponding to an improvement of 59.1%, which is decidedly better ($p < 0.01$) than that obtained with MTHF not in controlled-release form (23.0%) or with placebo (22.2%).

Figures 4 to 8 show the results obtained for all the psychometric tests used, namely: W.A.I.S test (Figure 4), Randt memory test for word acquisition (Figure 5a), Randt memory test for word repetition (Figure 5b), Toulouse-Pieron test (Figure 6), semantic verbal memory test (Figure 7), digit span for forward test (Figure 8a) and digit span for backward test (Figure 8b).

All the psychometric tests with the exception of the Toulouse-Pieron test (Figure 6) showed statistically significant improvements for treatment with MTHF in controlled-release form ($p < 0.05$) according to the present invention. In particular at T21 there was a statistically significant difference for the W.A.I.S. test (Figure 4), the Randt memory test (Figure 5), the semantic verbal memory test (Figure 7) and the digit span for the forward test (Figure 8a), and at T90 in the case of the digit span for the backward test (Figure 8b). In contrast, the group of patients treated with MTHF not in controlled-release form and with placebo showed no modifications of any kind.

Tolerance during the treatment was good for nearly all patients, with the exception of two patients of the MTHF controlled-release group, who complained of slight cephalgia during the initial days of treatment. This condition subsequently spontaneously

disappeared, although drug administration continued.

Table 3 shows the global judgement, expressed by the medical practitioner who performed the therapy, on the effectiveness of the treatment at T21, T45 and T90. In formulating this judgement the practitioner also consulted the patient's relatives and the paramedic personnel. The greater number of positive replies for treatment with MTHF in controlled-release form was statistically significant by T45 and further increased until T90 when 60% of the patients were recognised as and declared to be improved, against 24% of the patients in the group using MTHF not in controlled-release form and 26% in the placebo group.

TABLE 3

GLOBAL JUDGEMENT ON THE TREATMENT EFFECTIVENESS

Time 21 days					
	WORSENER	UNCHANGED	IMPROVED	MUCH	TOTAL
				IMPROVED	
MTHF in					
controlled-release					
form:	5	35	18	2	60
MTHF not in					
controlled-release					
form:	6	45	9	2	62
placebo:	10	40	10	2	62
				not	
				significant	
Time 45 days					
	WORSENER	UNCHANGED	IMPROVED	MUCH	TOTAL
				IMPROVED	
MTHF in					
controlled-release					
form:	4	29	22	5	60
MTHF not in					
controlled-release					
form:	6	44	10	2	62

2012876

- 12 -

placebo:	7	42	11	2	62
				P=0.05	

Time 90 days

5		WORSENER	UNCHANGED	IMPROVED	MUCH	TOTAL
					IMPROVED	

MTHF in

controlled-release

form:	2	22	28	8	60
-------	---	----	----	---	----

10 MTHF not in

controlled-release

form:	4	43	12	3	62
-------	---	----	----	---	----

placebo:	6	40	14	2	62
----------	---	----	----	---	----

P=0.001

15

Analogous experiments were conducted using pharmaceutical compositions according to the present invention containing different doses of MTHF and FTHF in controlled-release form, namely 5 mg (20 patients), 15 mg (20 patients), 20 mg (20 patients), 25 mg (15 patients), 40 mg (20 patients), 100 mg (20 patients) and 200 mg (20 patients), it being found that these doses also produce positive effects on the parameters analyzed.

25 The pharmaceutical compositions obtained using MTHF, FTHF and their salts, in raceme or optically active form, according to the invention, do not interfere with the sleeping/waking cycle, do not produce sedation, do not produce dependence or inurement and in general do not produce side effects.

30 Some examples of pharmaceutical compositions according to the present invention are given hereinafter for the sole purpose of non-limiting illustration.

EXAMPLE 1

35 Gastro-resistant controlled-release tablet containing 10 mg of MTHF; release time = 15 or 20 minutes.

1 tablet contains:

	MTHF calcium pentahydrate salt	12.6 mg
	(equivalent to 10 mg MTHF)	
	Pregelatinized starch	115.0 mg
	Lactose 100 mesh	72.7 mg
5	Hydroxypropylmethylcellulose	5.0 mg
	Magnesium stearate	1.0 mg
	Cellulose acetophthalate	7.5 mg
	Diethylphthalate	2.5 mg

10 EXAMPLE 2

Gastro-resistant controlled-release tablet containing 15 mg of MTHF; release time = 20 or 30 minutes.

1 tablet contains:

	MTHF calcium pentahydrate salt	18.8 mg
15	(equivalent to 15 mg MTHF)	
	Pregelatinized starch	115.0 mg
	Lactose 100 mesh	60.2 mg
	Hydroxypropylmethylcellulose	5.0 mg
	Magnesium stearate	1.0 mg
20	Cellulose acetophthalate	7.5 mg
	Diethylphthalate	2.5 mg

EXAMPLE 3

Gastro-resistant controlled-release tablet containing 20 mg of MTHF; release time = 30 or 35 minutes.

1 tablet contains:

	MTHF calcium pentahydrate salt	25.1 mg
	(equivalent to 20 mg MTHF)	
	Microcrystalline cellulose	80.0 mg
30	Lactose 100 mesh	79.9 mg
	Hydroxypropylmethylcellulose	10.0 mg
	Glyceryl behenate	5.0 mg
	Cellulose acetophthalate	7.5 mg
	Diethylphthalate	2.5 mg

35

EXAMPLE 4

Gastro-resistant controlled-release tablet containing 25 mg of

2012876

- 14 -

MTHF; release time = 35 or 40 minutes.

1 tablet contains:

	MTHF calcium pentahydrate salt	31.6 mg
	(equivalent to 25 mg MTHF)	
5	Microcrystalline cellulose	80.0 mg
	Lactose 100 mesh	73.4 mg
	Hydroxypropylmethylcellulose	10.0 mg
	Glyceryl behenate	5.0 mg
	Cellulose acetophthalate	7.5 mg
10	Diethylphthalate	2.5 mg

EXAMPLE 5

Gastro-resistant controlled-release tablet containing 40 mg of

MTHF; release time = 50 or 60 minutes.

15 1 tablet contains:

	MTHF calcium pentahydrate salt	50.6 mg
	(equivalent to 40 mg MTHF)	
	Microcrystalline cellulose	80.0 mg
	Lactose 100 mesh	54.4 mg
20	Hydroxypropylmethylcellulose	10.0 mg
	Glyceryl behenate	5.0 mg
	Cellulose acetophthalate	7.5 mg
	Diethylphthalate	2.5 mg

25 EXAMPLE 6

Gastro-resistant controlled-release tablet containing 50 mg of

MTHF; release time = 60 minutes.

1 tablet contains:

	MTHF calcium pentahydrate salt	63.2 mg
30	(equivalent to 50 mg MTHF)	
	Microcrystalline cellulose	80.0 mg
	Lactose 100 mesh	41.7 mg
	Hydroxypropylmethylcellulose	10.0 mg
	Glyceryl behenate	5.0 mg
35	Cellulose acetophthalate	7.5 mg
	Diethylphthalate	2.5 mg

EXAMPLE 7

Gastro-resistant controlled-release tablet containing 50 mg of FTHF; release time = 60 minutes.

1 tablet contains:

5	FTHF calcium pentahydrate salt (equivalent to 50 mg FTHF)	66.7 mg
	Carboxyvinylpolymer	20.0 mg
	Microcrystalline cellulose	112.3 mg
	Magnesium stearate	1.0 mg
10	Cellulose acetophthalate	7.5 mg
	Diethylphthalate	2.5 mg

EXAMPLE 8

Gastro-resistant controlled-release tablet containing 100 mg of MTHF; release time = 90 or 120 minutes.

1 tablet contains:

15	MTHF calcium pentahydrate salt (equivalent to 100 mg MTHF)	126.5 mg
	Dibasic calcium phosphate	90.0 mg
20	Lactose 100 mesh	163.0 mg
	Hydroxypropylmethylcellulose	15.0 mg
	Magnesium stearate	5.5 mg
	Cellulose acetophthalate	15.5 mg
	Diethylphthalate	5.0 mg

25

EXAMPLE 9

Gastro-resistant controlled-release tablet containing 200 mg of MTHF; release time = 180 or 210 or 240 minutes.

1 tablet contains:

30	MTHF calcium pentahydrate salt (equivalent to 200 mg MTHF)	251.0 mg
	Hydroxypropylmethylcellulose	30.0 mg
	Lactose 100 mesh	149.0 mg
	Glyceryl behenate	5.5 mg
35	Cellulose acetophthalate	15.0 mg
	Diethylphthalate	5.0 mg

EXAMPLE 10

Gastro-resistant controlled-release tablet containing 200 mg of FTHF; release time = 4 hours.

1 tablet contains:

5	FTHF calcium pentahydrate salt	266.7 mg
	(equivalent to 200 mg MTHF)	
	Microcrystalline cellulose	63.3 mg
	Hydrogenated castor oil	100.0 mg
	Glyceryl behenate	20.0 mg
10	Cellulose acetophthalate	15.0 mg
	Diethylphthalate	5.0 mg

EXAMPLE 11

Controlled-release suppository containing 50 mg of MTHF.

15 1 suppository contains:

	MTHF calcium pentahydrate salt	63.2 mg
	(equivalent to 50 mg MTHF)	
	Hydroxypropylmethylcellulose	50.0 mg
	Semisynthetic glycerides	1886.8 mg

20

EXAMPLE 12

Controlled-release injectable form containing 50 mg of MTHF.

1 vial contains:

	MTHF calcium pentahydrate salt	63.2 mg
25	(equivalent to 50 mg MTHF)	
	Glutathione	10.0 mg
	Citric acid	30.0 mg
	Hydroxyethylcellulose	10.0 mg
	Mannitol	160.0 mg
30	Methyl p-hydroxybenzoate	1.0 mg
	Sodium hydroxide	17.7 mg
	Water for injectable preparations	to make up to 3 ml

EXAMPLE 13

35 Controlled-release transdermic system containing 20 mg of MTHF

1 transdermic system contains:

	MTHF calcium pentahydrate salt	25.1 mg
--	--------------------------------	---------

(equivalent to 20 mg MTHF)

Fluid silicone	174.8 mg
Precipitated silica	15.2 mg

5 EXAMPLE 14

Controlled-release transdermic system containing 50 mg of MTHF

1 transdermic system contains:

	MTHF calcium pentahydrate salt	63.3 mg
	(equivalent to 50 mg MTHF)	
10	Glycerin	135.0 mg
	Polyvinyl alcohol	7.5 mg
	Polyvinylpyrrolidone	5.0 mg
	Citric acid	2.5 mg
	Purified water	100.0 mg

15

The present invention is susceptible to numerous modifications all falling within the inventive concept, and in addition all details can be replaced by others technically equivalent.

20 128 76

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A pharmaceutical composition in controlled-release form for use in the therapy of an organic mental disturbance, containing as active principle MTHF or FTHF or one or more of their pharmaceutically-acceptable salts.

2. The pharmaceutical composition as claimed in claim 1, wherein the organic mental disturbance is selected from the group consisting of senile primary degenerative dementia of Alzheimer type with depression, presenile primary degenerative dementia of Alzheimer type with depression, and multiinfarctual dementia with depression.

3. The pharmaceutical composition according to claim 1 or 2, wherein the active principle is released within a time period of between 15 minutes and 8 hours.

4. The pharmaceutical composition according to claim 1 or 2, wherein the active principle is released within a time period of between 20 and 60 minutes.

5. The pharmaceutical composition according to any one of claims 1 to 4, wherein the content of MTHF or FTHF or one or more of their salts in controlled-release form is between 5 and 200 mg.

6. The pharmaceutical composition according to any one of claims 1 to 4, wherein the content of MTHF or FTHF or their salts is between 10 and 50 mg.

B

20 12876

7. The pharmaceutical composition according to any one of claims 1 to 6, in gastrosoluble form suitable for oral administration.

8. The pharmaceutical composition according to any one of claims 1 to 6, in enterosoluble form suitable for oral administration.

9. The pharmaceutical composition according to any one of claims 1 to 6, in injectable form.

10. The pharmaceutical composition according to any one of claims 1 to 6, in the form of suppositories for rectal administration.

11. The pharmaceutical composition according to any one of claims 1 to 6, in the form of a transdermic system.

12. Use of a pharmaceutical composition in controlled-release form comprising MTHF or FTHF or one of their pharmaceutically-acceptable salts as active principle in the treatment of an organic mental disturbance in normofolatemmic or hypofolatemmic individuals.

13. The use according to claim 12, wherein the organic mental disturbance is selected from the group consisting of senile primary degenerative dementia of Alzheimer type with depression, presenile primary degenerative dementia of Alzheimer type with depression, and multiinfarctual dementia with depression.

B

20 128 76

14. The use according to claim 12 or 13, wherein the active principle is releasable within a time period of between 15 minutes and 8 hours.

15. The use according to claim 12 or 13, wherein the active principle is releasable within a time period of between 20 and 60 minutes.

16. The use according to any one of claims 12 to 15, wherein the content of the active principle is between 5 and 200 mg.

17. The use according to any one of claims 12 to 15, wherein the content of the active principle is between 10 and 50 mg.

18. The use according to any one of claims 12 to 17, wherein the pharmaceutical composition is in gastrosoluble or enterosoluble form suitable for oral administration.


19. The use according to any one of claims 12 to 17, wherein the pharmaceutical composition is in injectable form.

20. The use according to any one of claims 12 to 17, wherein the pharmaceutical composition is in the form of suppositories for rectal administration.

21. The use according to any one of claims 12 to 17, wherein the pharmaceutical composition is in the form of a transdermic system.


B

FIG. 1



Marks & Clerk

FIG. 2




* comparison in the group vs basal $p < 0.01$

❖ comparison vs placebo and vs MTHF not C R $p < 0.01$

❖❖ comparison vs placebo and vs MTHF not C R $p < 0.05$

Handwritten signature: *Handwritten signature*

FIG. 3




* comparison in the group vs basal $p < 0.01$

❖ comparison vs placebo and vs MTHF not C R $p < 0.01$

Marks & Clerk

FIG. 4



* comparison in the group vs basal $p < 0.01$

❖ comparison vs placebo and vs MTHF not C R $p < 0.01$

Hearts & Clark

FIG. 5

FIG. 5a






FIG. 5b




* comparison in the group vs basal $p < 0.01$

♦ comparison vs placebo and vs MTHF not C R
 $p < 0.01$

♦♦ comparison vs placebo and vs MTHF not C R
 $p < 0.05$


Marks & Clerk

FIG. 6



Marks & Clerk

FIG. 7



* comparison in the group vs basal p 0.01

❖ comparison vs placebo and vs MTHF not C R p < 0.01

❖❖ comparison vs placebo and vs MTHF not C R p < 0.05

Marks & Clerk

FIG. 8

FIG. 8a






FIG. 8b



* comparison in the group vs basal $p < 0.01$

❖ comparison vs placebo and vs MTHF not C R
 $p < 0.01$

❖❖ comparison vs placebo and vs MTHF not C R
 $p < 0.05$

Marks & Clerk