



(86) Date de dépôt PCT/PCT Filing Date: 2017/09/07  
 (87) Date publication PCT/PCT Publication Date: 2018/03/15  
 (45) Date de délivrance/Issue Date: 2022/06/28  
 (85) Entrée phase nationale/National Entry: 2019/02/27  
 (86) N° demande PCT/PCT Application No.: US 2017/050489  
 (87) N° publication PCT/PCT Publication No.: 2018/049036  
 (30) Priorité/Priority: 2016/09/07 (US62/384,433)

(51) Cl.Int./Int.Cl. *C11D 3/36* (2006.01),  
*C11D 17/00* (2006.01), *C11D 3/10* (2006.01),  
*C11D 3/386* (2006.01), *C11D 7/12* (2006.01),  
*C11D 7/36* (2006.01), *C11D 7/42* (2006.01)  
 (72) Inventeurs/Inventors:  
 LO, WENDY, US;  
 OLSON, ERIK C., US  
 (73) Propriétaire/Owner:  
 ECOLAB USA INC., US  
 (74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : COMPOSITIONS DE DETERGENT CONTENANT UNE ENZYME STABILISEE PAR DES PHOSPHONATES  
 (54) Title: DETERGENT COMPOSITIONS CONTAINING AN ENZYME STABILIZED BY PHOSPHONATES

(57) **Abrégé/Abstract:**

Detergent compositions that contain an enzyme, alkaline source, and phosphonate or amine phosphonate salt are described here. A use solution of the detergent compositions containing disclosed phosphonates can retain its enzyme activity for an extended period of time. Specifically, one specific type of phosphonates and another specific type of amine phosphonate salts were discovered to stabilize enzymes in detergent compositions. Solid detergent compositions that contain disclosed phosphonate or amine phosphonate salts are more effective to remove soils and can save production and use costs.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization

International Bureau

(43) International Publication Date  
15 March 2018 (15.03.2018)(10) International Publication Number  
**WO 2018/049036 A1**

## (51) International Patent Classification:

*C11D 3/386* (2006.01) *C11D 17/00* (2006.01)*C11D 3/36* (2006.01) *C11D 17/06* (2006.01)*C11D 3/30* (2006.01)

## (21) International Application Number:

PCT/US2017/050489

## (22) International Filing Date:

07 September 2017 (07.09.2017)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

62/384,433 07 September 2016 (07.09.2016) US

(71) Applicant: **ECOLAB USA INC.** [US/US]; 1 Ecolab Place, Saint Paul, Minnesota 55102 (US).(72) Inventors: **LO, Wendy**; c/o Ecolab USA Inc., 1 Ecolab Place, Saint Paul, Minnesota 55102 (US). **OLSON, Erik C.**; c/o Ecolab USA Inc., 1 Ecolab Place, Saint Paul, Minnesota 55102 (US).(74) Agent: **LINK, Jill N.** et al.; McKee, Voorhees & Sease, P.L.C., 801 Grand Avenue, Suite 3200, Des Moines, Iowa 50309-2721 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,

DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

## Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

## Published:

— with international search report (Art. 21(3))  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: DETERGENT COMPOSITIONS CONTAINING A STABILIZED ENZYME BY PHOSPHONATES




Figure 1A

(57) Abstract: Detergent compositions that contain an enzyme, alkaline source, and phosphonate or amine phosphonate salt are described here. A use solution of the detergent compositions containing disclosed phosphonates can retain its enzyme activity for an extended period of time. Specifically, one specific type of phosphonates and another specific type of amine phosphonate salts were discovered to stabilize enzymes in detergent compositions. Solid detergent compositions that contain disclosed phosphonate or amine phosphonate salts are more effective to remove soils and can save production and use costs.

  
 WO 2018/049036 A1

**TITLE: DETERGENT COMPOSITIONS CONTAINING AN ENZYME STABILIZED BY PHOSPHONATES**

5

**FIELD OF THE INVENTION**

10           The present disclosure relates generally to the field of cleaning compositions. In particular, solid detergent compositions that comprise an enzyme, a phosphonate, an alkaline source, and optionally other ingredients. The enzyme in these detergent compositions shows a superior stability after dissolving into use solutions of the compositions. Because of the enzyme's stability, the enzyme can function longer and  
15 therefore provide more effective removing/preventing re-deposition of soils. Conversely, also because of the superior stability of the enzyme, the amount of enzyme and other ingredients for increasing enzyme's effectiveness in these disclosed compositions can be reduced to achieve a similar cleaning effect.

20           **BACKGROUND OF THE INVENTION**

          Detergency is defined as the ability to wet, emulsify, suspend, penetrate, and disperse soils. Conventional detergents used in the warewashing and laundering industries include alkaline detergents. Alkaline detergent formulations employing alkali metal carbonates and/or alkali metal hydroxides, intended for both institutional and consumer  
25 use, are known to provide effective detergency.

          Enzymes have been employed in cleaning compositions since early 20<sup>th</sup> century. However, it was not until the mid-1960's when enzymes were commercially available with both the pH stability and soil reactivity for detergent applications. Enzymes are known as effective chemicals for use with detergents and other cleaning agents to break down soils.  
30 Enzymes break down soils, make them more soluble, and enable surfactants to remove them from a surface to provide enhanced cleaning of a substrate.

Specifically, enzymes can provide desirable activity for removal of, for example, protein-based, carbohydrate-based, or triglyceride-based stains from substrates. As a result, enzymes have been used for various cleaning applications in order to digest or degrade soils such as grease, oils (*e.g.*, vegetable oils or animal fat), protein, carbohydrate, or the like. For example, enzymes may be added as a component of a composition for laundry, textiles, ware washing, cleaning-in-place, cleaning drains, floors, carpets, medical or dental instruments, meat cutting tools, hard surfaces, personal care, or the like. Although products containing enzymes have evolved from simple powders containing alkaline protease to more complex granular compositions containing multiple enzymes and still further to liquid compositions, there remains a need for alternative cleaning applications employing stabilized enzymes. Numerous mechanisms for improving stabilization of enzymes for storage in detergent compositions have been used. However, there remains a need for improvement such that use solutions of detergent compositions retain detergency and cleaning performance when exposed to high temperatures and pH for extended periods of time as in actual cleaning applications.

Accordingly, it is an objective to develop a detergent composition with an enzyme and stabilizing agent such that its use solution is able to retain suitable enzyme stability under an elevated temperature and pH condition of use for a much longer period of time.

It is a further objective to develop multi-use solid detergent compositions that have not only storage, shelf, and dimensional stability but also provide a superior enzyme stability within the solid compositions and in their use solutions, under an elevated temperature and pH condition to provide improved detergency. The enzymatic activity in these compositions or use solutions thereof is retained under elevated temperature and pH conditions for a longer period of time.

It is an objective to develop methods for use of the stabilized enzymes in either detergent compositions themselves and in use solutions for improved detergency.

It is a further objective to develop methods to stabilize an enzyme in a solid detergent composition and its use solution.

Beneficially, such objectives overcome significant limitations of the state of the art of enzyme stability in detergent compositions, namely wherein un-stabilized enzyme significantly decreases its activity over time, including within short time periods of as little as 5-20 minutes, in the use solutions of these detergent compositions.

A further object is to develop multi-use compositions and methods for employing the same, to improve protein removal and anti-redeposition properties of detergent compositions, in particular non-caustic detergents compositions.


5 These and other objects, advantages and features of the present disclosure will become apparent from the following specification .

### BRIEF SUMMARY OF THE INVENTION

10 An advantage of the present disclosure is that an enzyme in a solid detergent composition can retain its activity for an extended period of time not only during the solid composition's storage but also in a use solution of the composition.

It is surprisingly discovered that adding a specific type of phosphonates or its salts, or another specific type of amine phosphonate salts in a detergent composition containing an enzyme can maintain the enzyme's activity for an extended period of time in a use  
15 solution of the composition during the actual use of the composition for cleaning purposes. It is also surprisingly discovered that enzymes in the use solutions of the detergent compositions of the present disclosure have superior stability. This discovery leads to a more effective composition due to a prolonged enzyme activity for removing soils. Beneficially, this improvement further allows the elimination or reduced use of enzymes,  
20 stabilizers, or some other ingredients commonly found in detergent compositions.


In one aspect, provided is a composition that comprises an enzyme, a phosphonate



represented by a formula of or salt thereof, and an alkaline source; wherein the enzyme is a protease, amylase, lipase, cellulase, peroxidase, gluconase, or mixture thereof; the alkaline source is a metal carbonate, metal bicarbonate, metal silicate,  
25 or mixture thereof; and R<sup>10</sup> and R<sup>11</sup> are independently hydrogen, a substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, phosphonate,


phosphonate ester, or derivative thereof, with a proviso that R<sup>10</sup> and R<sup>11</sup> are both -CH<sub>2</sub>-PO(OH)<sub>2</sub> groups.

In other aspect, provided is a composition that comprises an enzyme, an alkaline source, and an amine phosphonate salt; wherein the amine phosphonate salt is a product of



5 a phosphonate represented by a formula of and an amine, the enzyme is a protease, amylase, lipase, cellulase, peroxidase, gluconase, or mixture thereof; the alkaline source comprises a metal carbonate, metal bicarbonate, metal silicate, or mixture thereof; and R<sup>12</sup>, R<sup>13</sup>, and R<sup>14</sup> are independently hydroxyl, methyl, -PO(OH)<sub>2</sub>, -CH<sub>2</sub>COOH, a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof.


10 In yet another aspect, provided is a solid detergent composition that comprises an alkaline source, a phosphonate, and an enzyme; wherein the alkaline source comprises a metal carbonate, metal bicarbonate, metal silicate, or mixture thereof; the enzyme is a protease, amylase, lipase, cellulase, peroxidase, gluconase, or mixture thereof; the



15 phosphonate is represented by a formula of or salt thereof, wherein R<sup>10</sup> and R<sup>11</sup> are independently hydrogen, a substituted carboxylic acid, phosphonate, ethanol, diglyco, substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, or phosphonate-methyl; with a proviso that R<sup>10</sup> and R<sup>11</sup> are both -CH<sub>2</sub>-PO(OH)<sub>2</sub> groups, the ingredients of the composition is mixed and used to produce a solid detergent.

20 In another aspect, provided is a solid detergent composition that comprises an alkaline source, an enzyme, and an amine phosphonate salt; wherein the alkaline source comprises a metal carbonate, metal bicarbonate, metal silicate, or mixture thereof; the


enzyme is a protease, amylase, lipase, cellulase, peroxidase, gluconase, or mixture thereof; the amine phosphonate salt is a product of a phosphonate represented by a formula of



and an amine, wherein  $\text{R}^{12}$ ,  $\text{R}^{13}$ , and  $\text{R}^{14}$  are independently hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof; and the ingredients of the composition is mixed and used to produce a solid detergent.


In yet another aspect, provided is a method of cleaning, sanitizing and/or bleaching that comprises generating a use solution of a composition disclosed herein, and contacting a surface or object in need of cleaning and sanitizing with the use solution.

In other aspect, provided is a method of stabilizing an enzyme in a solid detergent composition. The method comprises adding a phosphonate of formula



or salt thereof, or an amine phosphonate salt in a detergent composition containing an enzyme, wherein  $\text{R}^{10}$  and  $\text{R}^{11}$  are independently hydrogen, a substituted carboxylic acid, phosphonate, ethanol, diglyco, substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, or phosphonate-methyl with a proviso that  $\text{R}^{10}$  and  $\text{R}^{11}$  are both  $-\text{CH}_2-\text{PO}(\text{OH})_2$  groups; the amine

phosphonate salt is a product of a phosphonate of formula



and an amine,

and R<sup>12</sup>, R<sup>13</sup>, and R<sup>14</sup> are independently hydroxyl, methyl, -PO(OH)<sub>2</sub>, -CH<sub>2</sub>COOH, a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof.

While multiple embodiments are disclosed, still other embodiments of the present disclosure will become apparent to those skilled in the art from the following detailed description, which shows and describes illustrative embodiments of the invention.  
5 Accordingly, the examples, figures, drawings, and detailed description are to be regarded as illustrative in nature and not restrictive.

### BRIEF DESCRIPTION OF THE DRAWINGS

10 **Figure 1A-Figure 1C** show the protease activities at different time points in the use solutions of the various base formula detergent compositions containing a different phosphonate at different levels at 120 °F. **Figure 1A** shows the protease activities at a level of 0.3 wt-% phosphonate. **Figure 1B** shows the protease activities at a level of 0.6 wt-% phosphonate. **Figure 1C** shows the protease activities at a level of 0.1 wt-%  
15 elemental phosphorus.

**Figures 2A-Figure 2D** show the protease activities at different time points in the use solutions of the base or all ash formula detergent compositions containing a phosphonate with or without an alkanolamine at 120 °F. **Figure 2A** shows the protease activities with phosphonebutane tricarboxylic acid (PBTC) alone or together with  
20 alkanolamines in the base formula detergent composition. **Figure 2B** shows the protease activities with 1-hydroxy ethylidene-1,1-diphosphonic acid (HEDP) alone or together with alkanolamines in the base formula detergent composition. **Figure 2C** shows the protease activities with PSO alone or together with alkanolamines in the base formula detergent composition. **Figure 2D** shows the protease activities in the all ash detergent  
25 compositions, *e.g.*, no bicarbonate in the detergent compositions, with various phosphonates.

**Figure 3A-Figure 3C** show the amylase activities at different time points in the use solutions of the various base formula detergent compositions containing a phosphonate at different levels at 120 °F. **Figure 3A** shows the amylase activities at a level of 0.3 wt-% of  
30 phosphonate. **Figure 3B** shows the amylase activities at a level of 0.6 wt-% of phosphonate. **Figure 3C** shows the amylase activities at a level of 0.1 wt-% elemental phosphorus.

**Figures 4A-Figure 4D** show the amylase activities at different time points in the use solutions of the base or all ash formula detergent compositions containing a phosphonate with or without an alkanolamine at 120 °F. **Figure 4A** shows the amylase activities with PBTC alone or together with alkanolamines in the base formula detergent compositions. **Figure 4B** shows the amylase activities with HEDP alone or together with alkanolamines in the base formula detergent compositions. **Figure 4C** shows the amylase activities with PSO alone or together with alkanolamines in the base formula detergent compositions. **Figure 4D** shows the amylase activity in the all ash formula detergent compositions with various phosphonates, respectively.

**Figure 5A-Figure 5C** show the lipase activities at different time points in the use solutions of the various base formula detergent compositions containing a phosphonate at different levels at 120 °F. **Figure 5A** shows the lipase activities at a level of 0.3 wt-% phosphonate. **Figure 5B** shows the lipase activities at a level of 0.6 wt-% phosphonate. **Figure 5C** shows the lipase activities at a level of 0.1 wt-% elemental phosphorus.

**Figures 6A-Figure 6G** show the lipase activities at different time points in a use solution of the base or all ash formula detergent compositions containing a phosphonate with or without an alkanolamine at 120 °F or at room temperature. **Figure 6A** shows the lipase activities with PBTC alone or together with alkanolamines in the base formula detergent compositions at 120 °F. **Figure 6B** shows the lipase activities with HEDP alone or together with alkanolamines in the base formula detergent compositions at 120 °F. **Figure 6C** shows the lipase activities with PSO alone or together with alkanolamines in the base formula detergent compositions at 120 °F. **Figure 6D** shows the lipase activities with PBTC at room temperature alone or together with alkanolamines in the base formula detergent compositions. **Figure 6E** shows the lipase activities with HEDP at room temperature, or together with alkanolamines in the base formula detergent compositions. **Figure 6F** shows the lipase activities with PSO at room temperature, alone or together with alkanolamines in the base formula detergent compositions. **Figure 6G** shows the lipase activities in the all ash formula detergent compositions with various phosphonates at 120 °F, respectively.

Various embodiments of the present disclosure will be described in detail with reference to the examples, figures, and drawings, wherein like reference numerals represent like parts throughout the several views. Reference to various embodiments does not limit

the scope of the disclosure. Figures represented herein are not limitations to the various embodiments according to the disclosure and are presented for exemplary illustration of the disclosure.

## 5 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The present disclosure relates to a detergent composition that contains an enzyme and has a superior enzyme stability in its use solution. The use solution of such a composition retains its enzyme activity for a long period of time. Especially, in a use solution produced from a detergent composition disclosed here, enzymes have such a superior stability that they can be effective to remove soil, protein, and starch for a long period time.

The embodiments of this disclosure are not limited to particular compositions and methods of use, which can vary and are understood by skilled artisans. It is further to be understood that all terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting in any manner or scope. For example, as used in this specification and the appended claims, the singular forms "a," "an" and "the" can include plural referents unless the content clearly indicates otherwise. Further, all units, prefixes, and symbols may be denoted in its SI accepted form.

Numeric ranges recited within the specification are inclusive of the numbers within the defined range. Throughout this disclosure, various aspects of this disclosure are presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the disclosure. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub-ranges as well as individual numerical values within that range (*e.g.* 1 to 5 includes 1, 1.5, 2, 2.75, 3, 3.80, 4, and 5).

So that the present disclosure may be more readily understood, certain terms are first defined. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which embodiments of the disclosure pertain. Many methods and materials similar, modified, or equivalent to those described herein can be used in the practice of the embodiments of the present disclosure without undue experimentation, the preferred materials and methods are

described herein. In describing and claiming the embodiments of the present disclosure, the following terminology will be used in accordance with the definitions set out below.

The term "about," as used herein, refers to variation in the numerical quantity that can occur, for example, through typical measuring and liquid handling procedures used for making concentrates or use solutions in the real world; through inadvertent error in these  
5 procedures; through differences in the manufacture, source, or purity of the ingredients used to make the compositions or carry out the methods; and the like. The term "about" also encompasses amounts that differ due to different equilibrium conditions for a composition resulting from a particular initial mixture. Whether or not modified by the  
10 term "about", the claims include equivalents to the quantities.

The term "actives" or "percent actives" or "percent by weight actives" or "actives concentration" are used interchangeably herein and refers to the concentration of those ingredients involved in cleaning expressed as a percentage minus inert ingredients such as water or salts.

As used herein, "substituted" refers to an organic group as defined below (*e.g.*, an alkyl group) in which one or more bonds to a hydrogen atom contained therein are replaced by a bond to non-hydrogen or non-carbon atoms. Substituted groups also include groups in which one or more bonds to carbon(s) or hydrogen(s) atom replaced by one or more bonds, including double or triple bonds, to a heteroatom. Thus, a substituted group is substituted  
15 with one or more substituents, unless otherwise specified. A substituted group can be substituted with 1, 2, 3, 4, 5, or 6 substituents.

Substituted ring groups include rings and ring systems in which a bond to a hydrogen atom is replaced with a bond to a carbon atom. Therefore, substituted cycloalkyl, aryl, heterocyclyl, and heteroaryl groups may also be substituted with  
25 substituted or unsubstituted alkyl, alkenyl, and alkynyl groups are defined herein.

As used herein, the term "alkyl" or "alkyl groups" refers to saturated hydrocarbons having one or more carbon atoms, including straight-chain alkyl groups (*e.g.*, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, etc.), cyclic alkyl groups (or "cycloalkyl" or "alicyclic" or "carbocyclic" groups) (*e.g.*, cyclopropyl, cyclopentyl,  
30 cyclohexyl, cycloheptyl, cyclooctyl, etc.), branched-chain alkyl groups (*e.g.*, isopropyl, tert-butyl, sec-butyl, isobutyl, etc.), and alkyl-substituted alkyl groups (*e.g.*, alkyl-substituted cycloalkyl groups and cycloalkyl-substituted alkyl groups).

Unless otherwise specified, the term "alkyl" includes both "unsubstituted alkyls" and "substituted alkyls." As used herein, the term "substituted alkyls" refers to alkyl groups having substituents replacing one or more hydrogens on one or more carbons of the hydrocarbon backbone. Such substituents may include, for example, alkenyl, alkynyl, halogeno, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonato, phosphinato, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonates, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclic, alkylaryl, or aromatic (including heteroaromatic) groups.

In some embodiments, substituted alkyls can include a heterocyclic group. As used herein, the term "heterocyclic group" includes closed ring structures analogous to carbocyclic groups in which one or more of the carbon atoms in the ring is an element other than carbon, for example, nitrogen, sulfur or oxygen. Heterocyclic groups may be saturated or unsaturated. Exemplary heterocyclic groups include, but are not limited to, aziridine, ethylene oxide (epoxides, oxiranes), thiirane (episulfides), dioxirane, azetidine, oxetane, thietane, dioxetane, dithietane, dithiete, azolidine, pyrrolidine, pyrroline, oxolane, dihydrofuran, and furan.

Alkenyl groups or alkenes are straight chain, branched, or cyclic alkyl groups having two to about 30 carbon atoms, and further including at least one double bond. In some embodiments alkenyl groups have from 2 to about carbon, or typically, from 2 to 10 carbons atoms. Alkenyl groups may be substituted or unsubstituted. Alkenyl groups may be substituted similarly to alkyl groups.

As used herein, the terms "alkylene", "cycloalkylene", and "alkenylene", alone or as part of another substituent, refer to a divalent radical derived from an alkyl, cycloalkyl, or alkenyl group, respectively, as exemplified by  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ . For alkylene, cycloalkylene, and alkenylene groups, no orientation of the linking group is implied.

The term "ester" as used herein refers to  $-\text{R}^{30}\text{COOR}^{31}$  group.  $\text{R}^{30}$  is absent, a substituted or unsubstituted alkylene, cycloalkylene, alkenylene, alkynylene, arylene,

aralkylene, heterocyclalkylene, or heterocyclene group as defined herein.  $R^{31}$  is a substituted or unsubstituted alkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, heterocyclalkyl, or heterocycl group as defined herein.

5 The term "amine" (or "amino") as used herein refers to  $-R^{32}NR^{33}R^{34}$  groups.  $R^{32}$  is absent, a substituted or unsubstituted alkylene, cycloalkylene, alkenylene, alkynylene, arylene, aralkylene, heterocyclalkylene, or heterocyclene group as defined herein.  $R^{33}$  and  $R^{34}$  are independently hydrogen, or a substituted or unsubstituted alkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, heterocyclalkyl, or heterocycl group as defined herein.

10 The term "amine" as used herein also refers to an independent compound. When an amine is a compound, it can be represented by a formula of  $R^{32'}NR^{33'}R^{34'}$  groups, wherein  $R^{32'}$ ,  $R^{33'}$ , and  $R^{34'}$  are independently hydrogen, or a substituted or unsubstituted alkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, heterocyclalkyl, or heterocycl group as defined herein.

15 The term "alcohol" as used herein refers to  $-R^{35}OH$  groups.  $R^{35}$  is absent, a substituted or unsubstituted alkylene, cycloalkylene, alkenylene, alkynylene, arylene, aralkylene, heterocyclalkylene, or heterocyclene group as defined herein.

The term "carboxylic acid" as used herein refers to  $-R^{36}COOH$  groups.  $R^{36}$  is absent, a substituted or unsubstituted alkylene, cycloalkylene, alkenylene, alkynylene, arylene, aralkylene, heterocyclalkylene, or heterocyclene group as defined herein.

20 The term "ether" as used herein refers to  $-R^{37}OR^{38}$  groups.  $R^{37}$  is absent, a substituted or unsubstituted alkylene, cycloalkylene, alkenylene, alkynylene, arylene, aralkylene, heterocyclalkylene, or heterocyclene group as defined herein.  $R^{38}$  is a substituted or unsubstituted alkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, heterocyclalkyl, or heterocycl group as defined herein.

25 An "antiredeposition agent" refers to a compound that helps keep suspended in water instead of redepositing onto the object being cleaned. Antiredeposition agents are useful in the present disclosure to assist in reducing redepositing of the removed soils onto the surface being cleaned.

30 As used herein, the term "cleaning" refers to perform, facilitate, or aid in soil removal, bleaching, microbial population reduction, and any combination thereof. As used herein, the term "microorganism" refers to any noncellular or unicellular (including colonial) organism. Microorganisms include all prokaryotes. Microorganisms include

bacteria (including cyanobacteria), spores, lichens, fungi, protozoa, virinos, viroids, viruses, phages, and some algae. As used herein, the term "microbe" is synonymous with microorganism.

As used herein, the term "disinfectant" refers to an agent that kills all vegetative  
5 cells including most recognized pathogenic microorganisms, using the procedure described in *A.O.A.C. Use Dilution Methods*, Official Methods of Analysis of the Association of Official Analytical Chemists, paragraph 955.14 and applicable sections, 15th Edition, 1990 (EPA Guideline 91-2). As used herein, the term "high level disinfection" or "high level disinfectant" refers to a compound or composition that kills substantially all organisms,  
10 except high levels of bacterial spores, and is effected with a chemical germicide cleared for marketing as a sterilant by the Food and Drug Administration. As used herein, the term "intermediate-level disinfection" or "intermediate level disinfectant" refers to a compound or composition that kills mycobacteria, most viruses, and bacteria with a chemical germicide registered as a tuberculocide by the Environmental Protection Agency (EPA).  
15 As used herein, the term "low-level disinfection" or "low level disinfectant" refers to a compound or composition that kills some viruses and bacteria with a chemical germicide registered as a hospital disinfectant by the EPA.

As used herein, the phrase "food processing surface" refers to a surface of a tool, a machine, equipment, a structure, a building, or the like that is employed as part of a food  
20 processing, preparation, or storage activity. Examples of food processing surfaces include surfaces of food processing or preparation equipment (*e.g.*, slicing, canning, or transport equipment, including flumes), of food processing wares (*e.g.*, utensils, dishware, wash ware, and bar glasses), and of floors, walls, or fixtures of structures in which food processing occurs. Food processing surfaces are found and employed in food anti-spoilage  
25 air circulation systems, aseptic packaging sanitizing, food refrigeration and cooler cleaners and sanitizers, ware washing sanitizing, blancher cleaning and sanitizing, food packaging materials, cutting board additives, third-sink sanitizing, beverage chillers and warmers, meat chilling or scalding waters, autodish sanitizers, sanitizing gels, cooling towers, food processing antimicrobial garment sprays, and non-to-low-aqueous food preparation  
30 lubricants, oils, and rinse additives.

As used herein, the phrase "food product" includes any food substance that might require treatment with an antimicrobial agent or composition and that is edible with or

without further preparation. Food products include meat (*e.g.*, red meat and pork), seafood, poultry, produce (*e.g.*, fruits and vegetables), eggs, living eggs, egg products, ready to eat food, wheat, seeds, roots, tubers, leafs, stems, corns, flowers, sprouts, seasonings, or a combination thereof. The term "produce" refers to food products such as  
5 fruits and vegetables and plants or plant-derived materials that are typically sold uncooked and, often, unpackaged, and that can sometimes be eaten raw.

The term "hard surface" refers to a solid, substantially non-flexible surface such as a counter top, tile, floor, wall, panel, window, plumbing fixture, kitchen and bathroom furniture, appliance, engine, circuit board, and dish. Hard surfaces may include for  
10 example, health care surfaces and food processing surfaces.

As used herein, the phrase "health care surface" refers to a surface of an instrument, a device, a cart, a cage, furniture, a structure, a building, or the like that is employed as part of a health care activity. Examples of health care surfaces include surfaces of medical or dental instruments, of medical or dental devices, of electronic apparatus employed for  
15 monitoring patient health, and of floors, walls, or fixtures of structures in which health care occurs. Health care surfaces are found in hospital, surgical, infirmity, birthing, mortuary, and clinical diagnosis rooms. These surfaces can be those typified as "hard surfaces" (such as walls, floors, bed-pans, etc.), or fabric surfaces, *e.g.*, knit, woven, and non-woven surfaces (such as surgical garments, draperies, bed linens, bandages, etc.), or patient-care  
20 equipment (such as respirators, diagnostic equipment, shunts, body scopes, wheel chairs, beds, etc.), or surgical and diagnostic equipment. Health care surfaces include articles and surfaces employed in animal health care.

As used herein, the term "instrument" refers to the various medical or dental instruments or devices that can benefit from cleaning with a composition according to the  
25 present disclosure.

The term "laundry" refers to items or articles that are cleaned in a laundry washing machine. In general, laundry refers to any item or article made from or including textile materials, woven fabrics, non-woven fabrics, and knitted fabrics. The textile materials can include natural or synthetic fibers such as silk fibers, linen fibers, cotton fibers, polyester  
30 fibers, polyamide fibers such as nylon, acrylic fibers, acetate fibers, and blends thereof including cotton and polyester blends. The fibers can be treated or untreated. Exemplary treated fibers include those treated for flame retardancy. It should be understood that the

term "linen" is often used to describe certain types of laundry items including bed sheets, pillow cases, towels, table linen, table cloth, bar mops and uniforms. The disclosure additionally provides a composition and method for treating non-laundry articles and surfaces including hard surfaces such as dishes, glasses, and other ware.

5           As used herein, the phrases "medical instrument," "dental instrument," "medical device," "dental device," "medical equipment," or "dental equipment" refer to instruments, devices, tools, appliances, apparatus, and equipment used in medicine or dentistry. Such instruments, devices, and equipment can be cold sterilized, soaked or washed and then heat sterilized, or otherwise benefit from cleaning in a composition of the present disclosure.

10       These various instruments, devices and equipment include, but are not limited to: diagnostic instruments, trays, pans, holders, racks, forceps, scissors, shears, saws (*e.g.*, bone saws and their blades), hemostats, knives, chisels, rongeurs, files, nippers, drills, drill bits, rasps, burrs, spreaders, breakers, elevators, clamps, needle holders, carriers, clips, hooks, gouges, curettes, retractors, straightener, punches, extractors, scoops, keratomes,

15       spatulas, expressors, trocars, dilators, cages, glassware, tubing, catheters, cannulas, plugs, stents, scopes (*e.g.*, endoscopes, stethoscopes, and arthoscopes) and related equipment, and the like, or combinations thereof.

          As used herein, the term "polymer" generally includes, but is not limited to, homopolymers, copolymers, such as for example, block, graft, random and alternating

20       copolymers, terpolymers, and higher "x"mers, further including their derivatives, combinations, and blends thereof. Furthermore, unless otherwise specifically limited, the term "polymer" shall include all possible isomeric configurations of the molecule, including, but are not limited to isotactic, syndiotactic and random symmetries, and combinations thereof. Furthermore, unless otherwise specifically limited, the term

25       "polymer" shall include all possible geometrical configurations of the molecule.

          For the purpose of this patent application, successful microbial reduction is achieved when the microbial populations are reduced by at least about 50%, or by significantly more than is achieved by a wash with water. Larger reductions in microbial population provide greater levels of protection.

30       As used herein, the term "sanitizer" refers to an agent that reduces the number of bacterial contaminants to safe levels as judged by public health requirements. In an embodiment, sanitizers for use in this disclosure will provide at least a 3 log reduction and

more preferably a 5-log order reduction. These reductions can be evaluated using a procedure set out in *Germicidal and Detergent Sanitizing Action of Disinfectants*, Official Methods of Analysis of the Association of Official Analytical Chemists, paragraph 960.09 and applicable sections, 15th Edition, 1990 (EPA Guideline 91-2). According to this  
5 reference a sanitizer should provide a 99.999% reduction (5-log order reduction) within 30 seconds at room temperature, 25±2°C, against several test organisms. Criteria for sanitizers and disinfectants may be different, depending on applications and regions.

As used herein, the term "soil" or "stain" refers to a non-polar oily substance which may or may not contain particulate matter such as mineral clays, sand, natural mineral  
10 matter, carbon black, graphite, kaolin, environmental dust, etc.

As used in this disclosure, the term "sporicide" refers to a physical or chemical agent or process having the ability to cause greater than a 90% reduction (1-log order reduction) in the population of spores of *Bacillus cereus* or *Bacillus subtilis* within 10  
15 seconds at 60° C. In certain embodiments, the sporicidal compositions of the disclosure provide greater than a 99% reduction (2-log order reduction), greater than a 99.99% reduction (4-log order reduction), or greater than a 99.999% reduction (5-log order reduction) in such population within 10 seconds at 60° C.

Differentiation of antimicrobial "-cidal" or "-static" activity, the definitions which describe the degree of efficacy, and the official laboratory protocols for measuring this  
20 efficacy are considerations for understanding the relevance of antimicrobial agents and compositions. Antimicrobial compositions can affect two kinds of microbial cell damage. The first is a lethal, irreversible action resulting in complete microbial cell destruction or incapacitation. The second type of cell damage is reversible, such that if the organism is rendered free of the agent, it can again multiply. The former is termed microbiocidal and  
25 the later, microbistatic. A sanitizer and a disinfectant are, by definition, agents which provide antimicrobial or microbiocidal activity. In contrast, a preservative is generally described as an inhibitor or microbistatic composition

As used herein, the term "substantially free of" or "free of" refers to compositions completely lacking the component or having such a small amount of the component that  
30 the component does not affect the performance of the composition. The component may be present as an impurity or as a contaminant and shall be less than 0.5 wt-%. In another

embodiment, the amount of the component is less than 0.1 wt-% and in yet another embodiment, the amount of component is less than 0.01 wt-%.

The term "substantially similar cleaning performance" refers generally to achievement by a substitute cleaning product or substitute cleaning system of generally the same degree (or at least not a significantly lesser degree) of cleanliness or with generally the same expenditure (or at least not a significantly lesser expenditure) of effort, or both.

As used herein, the term "ware" refers to items such as eating and cooking utensils, dishes, and other hard surfaces such as showers, sinks, toilets, bathtubs, countertops, windows, mirrors, transportation vehicles, and floors. As used herein, the term "warewashing" refers to washing, cleaning, or rinsing ware. Ware also refers to items made of plastic. Types of plastics that can be cleaned with the compositions according to the disclosure include but are not limited to, those that include polypropylene polymers (PP), polycarbonate polymers (PC), melamine formaldehyde resins or melamine resin (melamine), acrylonitrile-butadiene-styrene polymers (ABS), and polysulfone polymers (PS). Other exemplary plastics that can be cleaned using the compounds and compositions of the disclosure include polyethylene terephthalate (PET) polystyrene polyamide.

As used herein, the term "waters" includes food process or transport waters. Food process or transport waters include produce transport waters (*e.g.*, as found in flumes, pipe transports, cutters, slicers, blanchers, retort systems, washers, and the like), belt sprays for food transport lines, boot and hand-wash dip-pans, third-sink rinse waters, and the like. Waters also include domestic and recreational waters such as pools, spas, recreational flumes and water slides, fountains, and the like.

As used herein, the phrase "water soluble" means that the material is soluble in water in the present composition. In general, the material should be soluble at 25°C at a concentration of about 0.1 wt.% of the water, alternatively at about 1 wt.%, alternatively at about 5 wt.%, and alternatively at about 15 wt.%.

As used here, "an essentially similar composition" is referred to a composition in which everything else is the same except the addition of a different amount of the first solid, or of which the weight percent of alkaline compounds is within 10% of one for the reference composition. The compared blocks have identical shapes and dimensions.

The term "weight percent," "wt-%," "percent by weight," "% by weight," and variations thereof, as used herein, refer to the concentration of a substance as the weight of

that substance divided by the total weight of the composition and multiplied by 100. It is understood that, as used here, "percent," "%," and the like are intended to be synonymous with "weight percent," "wt-%," etc.

5 The methods and compositions of the present disclosure may comprise, consist essentially of, or consist of the components and ingredients of the present disclosure as well as other ingredients described herein. As used herein, "consisting essentially of" means that the methods and compositions may include additional steps, components or ingredients, but only if the additional steps, components or ingredients do not materially alter the basic and novel characteristics of the claimed methods and compositions.

10 It should also be noted that, as used in this specification and the appended claims, the term "configured" describes a system, apparatus, or other structure that is constructed or configured to perform a particular task or adopt a particular configuration. The term "configured" can be used interchangeably with other similar phrases such as arranged and configured, constructed and arranged, adapted and configured, adapted, constructed,  
15 manufactured and arranged, and the like.

#### *Detergent Compositions and Detergent Products*

As used herein, the term "composition" refers to chemical ingredients of a product or article. A product or article can be in a liquid, solid, powder form, or mixture thereof. It  
20 is possible that the same or similar composition can lead to different products or articles, due to the different process, arrangement, or amount in which each ingredient of the composition is put together in the product or article.

As used herein, the term "detergent composition" refers to chemical ingredients of a detergent product or detergent. A detergent product or detergent is usually used for  
25 cleaning purpose, by the detergent or detergent product itself or by a use solution thereof. A detergent or detergent product can be in a liquid, solid, powder form, or mixture thereof. A detergent product or detergent can be supplied in one package or separate packages. It is possible that the same or similar detergent composition can lead to different detergent products, due to the different process or amount in which each ingredient of the  
30 composition is put together in the detergent product. In this disclosure, the terms of "detergent product" and "detergent" are used interchangeably.

*Solid Detergents*

As used herein, the term "solid" refers to a state of matter known to those of skill in the art. A solid may be of crystalline, amorphous form, or a mixture thereof. In a solid can be a single compound or a mixture of compounds. A solid may be a mixture of two or  
5 more different solids. A solid may be aggregates of particles each of which has a size of a few, a few tens, a few hundreds of micrometers or nanometers. A solid may be a powder of one or more compounds.

As used herein, a solid detergent or cleaning composition refers to a detergent or cleaning composition in the form of a solid such as a powder, a flake, a granule, a pellet, a  
10 tablet, a lozenge, a puck, a briquette, a brick, a block, or another solid form known to those of skill in the art. Although the term solid block is often referred to herein, it is understood that the solid compositions can take various forms. In a preferred aspect, a pressed solid block is employed. It should be understood that the term "solid detergent" refers to the state of the detergent composition under the expected  
15 conditions of storage and use of the solid detergent composition. In general, it is expected that the detergent composition will remain a solid when provided at a temperature of a room temperature up to about 120°F.

A solid detergent composition can be provided as a pressed solid block, a cast solid block, an extruded pellet or block, or a tablet so that one or a plurality of the  
20 solids will be available in a package having a size of between about 1 grams and about 11,000 grams.

A solid detergent composition may be provided in the form of a unit dose. A unit dose refers to a solid detergent composition unit sized so that the entire unit is used during a single washing cycle. When the solid detergent composition is provided as a  
25 unit dose, it is preferably provided as a pressed solid, a cast solid, an extruded pellet, or a tablet having a size of between about 1 gram and about 50 grams. Alternatively, a pressed solid, a cast solid, an extruded pellet, or a tablet may have a size of between 50 grams up through 250 grams. An extruded, cast, or press solid may also have a weight of about 100 grams or greater.

30 A solid detergent composition may also be provided in the form of a multiple use (*e.g.*, multi-use) solid, such as, a block or a plurality of pellets, and can be repeatedly used to generate aqueous use solutions of the detergent composition for multiple cycles or

a predetermined number of dispensing cycles. A multiple use solid detergent composition can be repeatedly used to generate an aqueous detergent composition, *e.g.*, use solution, for multiple washing cycles. A multiple use solid detergent composition can have a mass of about 1 kilogram to about 10 kilograms or greater.

5           Typically, the solid detergent composition as disclosed herein dissolves quickly and completely upon contact with an aqueous solution into a stable use solution. In some aspects of the disclosure, the amount and type of anionic surfactants employed in the solid detergent composition provides a desired dissolution rate for a particular dispense rate. A stable use solution does not contain  
10 any solids upon visual inspection.

          Pressed solid detergent blocks are made suitable to provide stability such that reactive components in the compositions do not react with each other until a point of dilution and/or use. In some aspects, the order of introducing the components to form the solid are non-limiting as there is minimal and/or no water introduced into the solid  
15 compositions. However, in some aspects, pressed solid detergent blocks are made by using a binding system to minimize any damage to the coated granules which may be employed.

          Beneficially, a pressing process to make the pressed solid detergent blocks generates a pressed solid detergent block and prevents the reaction or mix of the components. In an aspect of the disclosure, the solid detergent composition remains  
20 unreacted or unmixed until a point of use, *e.g.* dilution.

          In a pressed solid process, a flowable solid, such as granular solids or other particle solids including binding agents are combined under pressure. In a pressed solid process, flowable solids of the compositions are placed into a form (*e.g.*, a mold or container). The method can include gently pressing the flowable solid in the form to produce the solid  
25 cleaning composition.

          The method can further include a curing step to produce the solid cleaning composition. As referred to herein, an uncured composition including the flowable solid is compressed to provide sufficient surface contact between particles making up the flowable solid that the uncured composition will solidify into a stable solid cleaning composition. A  
30 sufficient quantity of particles (*e.g.* granules) in contact with one another provides binding of particles to one another effective for making a stable solid composition. Inclusion of a curing step may include allowing the pressed solid to solidify for a period of time, such as

a few hours, or about 1 day (or longer). In additional aspects, the methods could include vibrating the flowable solid in the form or mold, such as the methods disclosed in U.S. Patent No. 8,889,048.

5 The use of pressed solids provides numerous benefits over conventional solid block or tablet compositions requiring high pressure in a tablet press, or casting requiring the melting of a composition consuming significant amounts of energy, and/or by extrusion requiring expensive equipment and advanced technical know-how. Pressed solids overcome such various limitations of other solid blocks, therefore there is a need for making new pressed solid cleaning compositions. Moreover, pressed solid blocks have  
10 more consistent and attractive appearance than extruded ones, therefore pressed solid detergent blocks can form solid blocks of distinct shapes for identification and control of use. They can retain its shape under conditions in which the blocks may be stored or handled. In general, it is expected that the detergent composition will remain a solid when provided at a temperature of up to about 120°F.

15 In some situations, the methods of making pressed blocks reduce or eliminate water from the system prior to solidification. Preferably, the compositions are formed using components in an anhydrous form. In some other situations, compositions have a water content of less than about 20% by weight, less than about 15% by weight, less than about 12% by weight, 10% by weight, less than about 5% by weight, less than about 1% by  
20 weight, less than about 0.1% by weight, less than about 0.05% by weight, and most preferably free of water (*e.g.* dried). In an aspect, the dried composition may be in the form of granules. On contrast, cast or extruded solid detergent blocks can have from about 20 to about 40 wt-% water. Therefore, pressed solid blocks are preferred due to the removal or reduction of water from the compositions and ash hydration is not employed as  
25 a solidification mechanism.

The particulate components of the disclosure can be in the form of granules and/or flakes, but is preferably presented in the form of regular small granules. Thereafter, the granules are used to form solid detergent blocks. The solidification process may last from a few seconds to several hours, depending on factors including, but not limited to the size  
30 of the formed or cast composition, the ingredients of the composition, and the temperature of the composition.

The solid detergent compositions may be formed using a batch or continuous mixing system. To make extruded blocks, powders and liquids of a detergent composition are blended to form a mixture, then the blended mixture is pressed through a mold to form a product, then the product hardens with time to an extruded solid block.

5 A single- or twin-screw extruder is used to combine and mix one or more cleaning agents at high shear to form a homogeneous mixture to make extruded blocks. To make pressed solid blocks, solid powders and/or other liquid ingredients of a detergent composition are mixed to form a blended power, then the blended power is poured into a mold and pressed into a solid detergent block. Generally, a solid detergent block  
10 processed according to the method of the disclosure is substantially homogeneous with regard to the distribution of ingredients throughout its mass and is dimensionally stable.

In some embodiments, the solid detergent composition of the present disclosure is provided as a pressed solid block having a mass of between about 5 grams and 10 kilograms. In certain embodiments, a pressed solid detergent block  
15 has a mass between about 1 and about 10 kilograms. In further embodiments, a block of the solid detergent composition has a mass of between about 5 kilograms and about 8 kilograms. In other embodiments, a block of the solid detergent composition has a mass of between about 5 grams and about 1 kilogram, or between about 5 grams and about 500 grams.

20 In some embodiments, the pressed solid detergent block produced from the disclosed composition has a water content of less than about 20 wt-%, 15 wt-%, 12 wt-%, 10 wt-%, 9 wt-%, 8 wt-%, 7 wt-%, 6 wt-%, 5 wt-%, 4 wt-%, 3 wt-%, 2 wt-%, 1 wt-%, 0.7 wt-%, 0.5 wt-%, 0.3 wt-%, 0.1 wt-%, or 0.05 wt-%. In some other embodiments, the pressed solid detergent block produced from the disclosed composition has a water  
25 content of between about 0.1 and about 15 wt-%, between about 0.1 and about 5 wt-%, between about 0.1 and about 3 wt-%, between about 1 and about 8 wt-%, between about 5 and about 10 wt-%, between about 5 and about 15 wt-%, or between about 5 and about 15 wt-%. In an aspect, the dried composition may be in the form of granules. On  
30 contrast, cast or extruded solid detergent blocks can have from about 20 to about 40 wt-% water.


*Phosphonate*

The detergent compositions disclosed here contains a specific type of phosphonates or salts thereof or amine salt of another specific type of phosphonates. Applicant unexpectedly discovered that these specific types of phosphonates or salts thereof disclosed here stabilize enzymes in detergent compositions.

The term “phosphonate” as used herein refers to an independent compound with a formula of  $R^{40'}PO(OH)_2$  groups, wherein  $R^{40'}$  is a substituted or unsubstituted alkyl, cycloalkyl, alkenyl, alkynyl, aryl, aralkyl, heterocyclalkyl, or heterocycl group as defined herein.


The term “phosphonate” as used herein may also refer to  $-R^{40}PO(OH)_2$  groups.  $R^{40}$  is a substituted or unsubstituted alkylene, cycloalkylene, alkenylene, alkynylene, arylene, aralkylene, heterocyclalkylene, or heterocyclylene group as defined herein.

One type phosphonate compound to stabilize enzyme(s) in a detergent composition




is a type of phosphonate represented by formula  $\text{R}^{10}\text{N}(\text{R}^{11})\text{CH}_2\text{PO}(\text{OH})_2$ , wherein  $\text{R}^{10}$  and  $\text{R}^{11}$  are independently hydrogen, a phosphonate, unsubstituted alkyl, or substituted alkyl, with an exception that  $\text{R}^{10}$  and  $\text{R}^{11}$  are both  $-\text{CH}_2\text{PO}(\text{OH})_2$  groups. This type of phosphonate can be added into a detergent composition in its acid form, or in a salt form after being neutralized by a base. Since a use solution of the detergent compositions disclosed here has a pH of from about 8 to about 12, the two  $-\text{OH}$  group of the phosphonate group are in their salt forms, *e.g.*, neutralized when the phosphonate is in the use solution.

The other type of phosphonate compound to stabilize enzyme(s) in a detergent composition is an amine phosphonate salt that is a product of a phosphonate compound



represented by a formula and an amine, and  $R^{12}$ ,  $R^{13}$ , and  $R^{14}$  are independently hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof. This amine phosphonate salt is added to a detergent composition in its salt form, produced usually by reacting the phosphonate with an amine to neutralize all or part of its OH groups. The amine can be an alkanolamine, monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof. In some other embodiments, the amine of the amine phosphonate salt is a  $\text{C}_{2-6}$  alkylamine or mixture thereof. The alkylamine of course can be a mono-, di-, or tri-amine.



In some embodiments, the disclosed detergent compositions contains a



phosphonate of formula , wherein  $R^{10}$  and  $R^{11}$  are independently hydrogen, a unsubstituted alkyl, substituted alkyl, a substituted carboxylic acid, phosphonate, ethanol, diglyco, 2-(EO) $_n$ -biphosphonateamine-ethyl, 2-(PO) $_n$ -biphosphonateamine-isopropyl, or phosphonate-methyl. In other embodiments,  $R^{10}$  is hydrogen and  $R^{11}$  is a unsubstituted alkyl, substituted alkyl, a substituted carboxylic acid, phosphonate, ethanol, diglyco, 2-(EO) $_n$ -biphosphonateamine-ethyl, 2-(PO) $_n$ -biphosphonateamine-isopropyl, or phosphonate-methyl. In some other embodiments,  $R^{10}$  is hydrogen and  $R^{11}$  is a substituted carboxylic acid, phosphonate, ethanol, diglyco, 2-(EO) $_n$ -biphosphonateamine-ethyl, 2-(PO) $_n$ -biphosphonateamine-isopropyl, or phosphonate-methyl.

As used here, EO refers to  $-\text{CH}_2\text{CH}_2\text{O}-$ , *e.g.*, ethylene oxide group, and PO to  $-\text{CH}_2\text{CH}(\text{CH}_3)\text{O}-$  group. When (EO) $_n$  or (PO) $_n$  is used,  $n$  is an integer of 1-30.


In some embodiments,  $R^{11}$  is  $-\text{CH}_2\text{-PO}(\text{OH})_2$  group. In some other embodiments,  $R^{11}$  is  $-\text{CH}_2\text{-PO}(\text{OH})_2$  group and  $R^{10}$  is ethanoyl, diglyco, substituted alkyl, isopropyl-2-(EO) $_n$ -biphosphonateamine, or methyl-phosphonate. In yet some other embodiments, the



5

, aminomethyl phosphonic acid, a mixture thereof, or a salt thereof.

In some embodiments, the amine phosphonate salt is an amine salt of a



phosphonate of a formula  $\text{R}^{12}\text{R}^{13}\text{R}^{14}\text{C-P}(\text{OH})_3$ , wherein  $\text{R}^{12}$ ,  $\text{R}^{13}$ , and  $\text{R}^{14}$  are independently hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof. In some other embodiments, one of  $\text{R}^{12}$ ,  $\text{R}^{13}$ , and  $\text{R}^{14}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , ester thereof, salt thereof, or derivative thereof.

In some other embodiments, the phosphonate of the amine phosphonate salt is  $\text{PO}(\text{OH})_2(\text{C}(\text{CH}_2\text{COOH})_3)$ , phosphonobutane tricarboxylic acid (PBTC). In yet some other embodiments, the phosphonate of the amine phosphonate is  $\text{C}(\text{CH}_3)(\text{OH})(\text{PO}(\text{OH})_2)_2$ , 1-hydroxy ethylidene-1,1-diphosphonic acid (HEDP).

In some embodiments, the amine of the amine phosphonate salt is an alkanolamine, monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof. In some other embodiments, the amine of the amine phosphonate salt is a  $\text{C}_{2-6}$  alkylamine or mixture thereof. The alkylamine can be a mono-, di-, or tri-amine.

In some embodiments, the detergent composition disclosed here has from about 0.1 wt-% to about 35 wt-%, 0.1 wt-% to about 30 wt-%, 0.1 wt-% to about 25 wt-%, 0.1 wt-% to about 15 wt-%, 0.1 wt-% to about 10 wt-%, 0.1 wt-% to about 5 wt-%, about 0.5 wt-% to about 5 wt-%, about 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 10 wt-%, 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 5 wt-%, 5 wt-% to about 10 wt-%, 10 wt-% to about 15 wt-%, about 15 wt-% to about 20 wt-%, 20 wt-% to about 25 wt-%, 25 wt-% to about 30 wt-%, 30 wt-% to about 35 wt-%, about 0.1 wt-%, about 0.5 wt-%, about 1 wt-%, about 2 wt-%, about 3 wt-%, about 4 wt-%, about 5 wt-%, about 6 wt-%, about 7 wt-%, about 8 wt-%, about 9 wt-%, about 10 wt-%, about 15 wt-%, about 20 wt-%, about 25 wt-%, about 30 wt-%, about 32 wt-%, or about 35 wt-% of the phosphonate, salt thereof, or amine phosphonate salt.

### *Enzymes*

The disclosure disclosed here related to a detergent composition that comprises an enzyme. In some embodiments, the enzyme is supplied in a liquid or solid form and mixed with the other components of the detergent composition, by spraying or mixing.

Enzymes that can be used according to the disclosure include enzymes that provide desirable activity for removal of protein-based, carbohydrate-based, or triglyceride-based stains from substrates; for cleaning, destaining, and sanitizing presoaks, such as presoaks for medical and dental instruments, devices, and equipment; presoaks for flatware, cooking ware, and table ware; or presoaks for meat cutting equipment; for machine warewashing; for laundry and textile cleaning and destaining; for carpet cleaning and destaining; for cleaning-in-place (CIP) and destaining-in-place; for cleaning and destaining food processing surfaces and equipment; for drain cleaning; presoaks for cleaning; and the like.

Although not limiting to the present disclosure, enzymes suitable for the detergent compositions can act by degrading or altering one or more types of soil residues encountered on an instrument or device thus removing the soil or making the soil more removable by a surfactant or other component of the cleaning composition. Both degradation and alteration of soil residues can improve detergency by reducing the physicochemical forces that bind the soil to the instrument or device being cleaned, *e.g.*, the soil becomes more water soluble. For example, one or more

proteases can cleave complex, macro molecular protein structures present in soil residues into simpler short chain molecules which are, of themselves, more readily desorbed from surfaces, solubilized or otherwise more easily removed by detergent solutions containing said proteases.

5            Suitable enzymes include a protease, an amylase, a lipase, a gluconase, a cellulase, a peroxidase, or a mixture thereof of any suitable origin, such as vegetable, animal, bacterial, fungal or yeast origin. Preferred selections are influenced by factors such as pH-activity and/or stability optima, thermostability, and stability to active detergents, builders and the like. In this respect, bacterial or fungal enzymes are preferred, such as bacterial amylases and proteases, and fungal cellulases. Preferably  
10            the enzyme is a protease, a lipase, an amylase, or a combination thereof.

          "Detergent enzyme", as used herein, means an enzyme having a cleaning, destaining or otherwise beneficial effect as a component of a solid detergent composition for instruments, devices, or equipment, such as medical or dental  
15            instruments, devices, or equipment; or for laundry, textiles, warewashing, cleaning-in-place, drains, carpets, meat cutting tools, hard surfaces, personal care, or the like. Preferred detergent enzymes include a hydrolase such as a protease, an amylase, a lipase, or a combination thereof. Preferred enzymes in solid detergent compositions for cleaning medical or dental devices or instruments include a protease, an amylase,  
20            a cellulase, a lipase, or a combination thereof. Preferred enzymes in solid detergent compositions for food processing surfaces and equipment include a protease, a lipase, an amylase, a gluconase, or a combination thereof. Preferred enzymes in solid detergent compositions for laundry or textiles include a protease, a cellulase, a lipase, a peroxidase, or a combination thereof. Preferred enzymes in solid detergent  
25            compositions for carpets include a protease, an amylase, or a combination thereof. Preferred enzymes in solid detergent compositions for meat cutting tools include a protease, a lipase, or a combination thereof. Preferred enzymes in solid detergent compositions for hard surfaces include a protease, a lipase, an amylase, or a combination thereof. Preferred enzymes in solid detergent compositions for drains  
30            include a protease, a lipase, an amylase, or a combination thereof.

          Enzymes are normally incorporated into a solid detergent composition according to the disclosure in an amount sufficient to yield effective cleaning during a

washing or presoaking procedure. An amount effective for cleaning refers to an amount that produces a clean, sanitary, and, preferably, corrosion free appearance to the material cleaned, particularly for medical or dental devices or instruments. An amount effective for cleaning also can refer to an amount that produces a cleaning, stain removal, soil removal, whitening deodorizing, or freshness improving effect on substrates such as medical or dental devices or instruments and the like. Such a cleaning effect can be achieved with amounts of enzyme as low as about 0.1 wt-% of the detergent composition.

In detergent compositions of the present disclosure, suitable cleaning can typically be achieved when an enzyme is also preferably present at about 1 to about 35 wt-%; preferably about 2 to about 15 wt-%; preferably about 3 to about 10 wt-%; preferably about 4 to about 8 wt-%; preferably about 4, about 5, about 6, about 7, or about 8 wt-%. The higher enzyme levels are typically desirable in highly concentrated cleaning or presoak formulations. A presoak is preferably formulated for use upon a dilution of about 1:500, or to a formulation concentration of about 2000 to about 4000 ppm, which puts the use concentration of the enzyme at about 20 to about 40 ppm.

Commercial enzymes, such as alkaline proteases, are obtainable in liquid or dried form, are sold as raw aqueous solutions or in assorted purified, processed and compounded forms, and include about 0.1% to about 80% by weight active enzyme generally in combination with stabilizers, buffers, cofactors, impurities and inert vehicles. The actual active enzyme content depends upon the method of manufacture and is not critical; assuming the solid detergent composition has the desired enzymatic activity. The particular enzyme chosen for use in the process and products of this disclosure depends upon the conditions of final utility, including the physical product form, use pH, use temperature, and soil types to be degraded or altered. The enzyme can be chosen to provide optimum activity and stability for any given set of utility conditions.

A valuable reference on enzymes is "Industrial Enzymes", Scott, D., in Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Edition, (editors Grayson, M. and Eckroth, D.) Vol. 9, pp. 173 224, John Wiley & Sons, New York, 1980.

In some other embodiments, the enzyme in the detergent composition is a single enzyme. In some other embodiments, the enzyme in the detergent composition is a mixture of two or more enzymes. In some other embodiments, the enzyme in the composition is a protease, amylase, lipase, hydrolase, cellulase, gluconase, peroxidase, 5 mannanase, or a mixture thereof. In some other embodiments, the enzyme is a protease, amylase, lipase, cellulose, peroxidase, gluconase, or mixture thereof. In some other embodiments, the enzyme in the detergent compositions disclosed here is a protease, amylase, lipase, or mixture thereof. In some other embodiments, the enzyme is a protease, amylase, or mixture thereof. In some other embodiments, the enzyme is a protease, lipase, 10 or mixture thereof. In some other embodiments, the enzyme is an amylase, lipase, or mixture thereof. In some other embodiments, the enzyme is a protease. In some other embodiments, the enzyme is an amylase. In yet some other embodiments, the enzyme is a lipase.

In some embodiments, the detergent composition disclosed here has from about 15 0.1 wt-% to about 35 wt-%, 0.1 wt-% to about 30 wt-%, 0.1 wt-% to about 25 wt-%, 0.1 wt-% to about 15 wt-%, 0.1 wt-% to about 10 wt-%, 0.1 wt-% to about 5 wt-%, about 0.5 wt-% to about 5 wt-%, about 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 10 wt-%, 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 5 wt-%, 5 wt-% to about 10 wt-%, 10 wt-% to about 15 wt-%, about 15 wt-% to about 20 wt-%, 20 wt-% to about 25 wt-%, 25 20 wt-% to about 30 wt-%, 30 wt-% to about 35 wt-%, about 0.1 wt-%, about 0.5 wt-%, about 1 wt-%, about 2 wt-%, about 3 wt-%, about 4 wt-%, about 5 wt-%, about 6 wt-%, about 7 wt-%, about 8 wt-%, about 9 wt-%, about 10 wt-%, about 15 wt-%, about 20 wt-%, about 25 wt-%, about 30 wt-%, about 32 wt-%, or about 35 wt-% of the enzyme.

The detergent composition of the current disclosure had further been found, 25 surprisingly, to have a significantly stabilized enzyme, especially, protease, lipase and/or amylase, activity toward digesting proteins and enhancing soil removal in their use solution. Applicant surprisingly discovered that through using a specific type of phosphonates alone or another specific amine phosphonate salts in a detergent composition, the enzymes in a use solution made from a detergent composition of the 30 present disclosure stay active much longer than those from the detergent compositions that do not contain phosphonates disclosed here. As shown in this disclosure, a use solution produced from the solid detergent of the present disclosure, both protease and

lipase stay active for a much longer time. More stable the enzymes are, the longer they are effective in removing soil, protein, or starch (and fats if lipases are included). As a result, the detergent composition disclosed here is also more effective.

5 Because of the superior stability of enzymes in the detergent composition of the present disclosure, it is possible for the composition to use less enzymes and to be free of other stabilizers or other ingredients commonly found in existing detergent compositions. Some stabilizers could be liquid and difficult to be included in a solid detergent composition, or could lead to undesirable reactions with other ingredients. Some stabilizers raise health/safety/labeling concerns in a concentrated composition (e.g. GHS  
10 label icon warnings that are not desired). At a minimum stabilizers add complexity to a formula and take up “formulation space” for other functional ingredients. It is an extra advantage of using the disclosed disclosure that no or a reduced amount of other stabilizers is used to produce the detergent disclosed here.


#### 15 *Alkaline Source*

The detergent compositions and methods, according to the present disclosure includes an effective amount of alkaline source. The alkaline source in turn comprises one or more alkaline compounds. In general, an effective amount of the alkaline source should be considered as an amount that provides a use solution having  
20 a pH of at least about 8. When the use solution has a pH of between about 8 and about 10, it can be considered mildly alkaline, and when the pH is greater than about 12, the use solution can be considered caustic. In general, it is desirable to provide the use solution as a mildly alkaline cleaning composition because it is considered to be safer than the caustic based use compositions.

25 The alkaline source can include an alkali metal carbonate, an alkali metal hydroxide, alkaline metal silicate, or a mixture thereof. Suitable metal carbonates that can be used include, for example, sodium or potassium carbonate, bicarbonate, sesquicarbonate, or a mixture thereof. Suitable alkali metal hydroxides that can also be used include, for example, sodium, lithium, or potassium hydroxide. Examples of  
30 useful alkaline metal silicates include sodium or potassium silicate (with  $M_2O:SiO_2$  ratio of 2.4 to 5:1, M representing an alkali metal) or metasilicate. The alkaline

source may also include a metal borate such as sodium or potassium borate, and the like.

The alkaline source may also include ethanolamines, urea sulfate, amines, amine salts, and quaternary ammonium. The simplest cationic amines, amine salts and quaternary ammonium compounds can be schematically drawn thus:



in which, R represents a long alkyl chain, R', R'', and R''' may be either long alkyl chains or smaller alkyl or aryl groups or hydrogen and X represents an anion.

The alkaline source can be added to the composition in the form of solid. For example, alkali metal hydroxides are commercially available as a solid in the form of prilled solids or beads having a mix of particle sizes ranging from 25 about 12-100 U.S. mesh. For example, an alkali metal hydroxide may be added to the solid detergent composition in a variety of solid forms, including for example in the form of solid beads. Alkali metal hydroxides are commercially available.

The alkaline source is preferably in an amount to enhance the cleaning of a substrate and improve soil removal performance of the composition. In general, it is expected that the concentrate will include the alkaline source in an amount of at least about 5 wt-%, at least about 10 wt-%, or at least about 15 wt-%. The pressed solid detergent composition can include between about 10 wt-% and about 95 wt-%, preferably between about 15 wt-% and about 70 wt-%, between about 20 wt-% and about 60 wt-%, and even more preferably between about 70 wt-% and about 95 wt-% of the alkaline source.

In some embodiments, the detergent compositions disclosed here contains a metal carbonate, metal bicarbonate, metal silicate, or mixture thereof as their alkaline source. In some other embodiments, the detergent compositions disclosed here contains a metal carbonate, metal bicarbonate, or mixture thereof as their alkaline source. In some embodiments, the alkaline source in the detergent compositions disclosed here is an alkali metal carbonate, alkali metal bicarbonate solid, alkali metal silicate, or mixture thereof. In


some other embodiments, the alkaline source in the detergent compositions disclosed here is an alkali metal carbonate, alkali metal bicarbonate, or a mixture thereof. In some other embodiments, the alkaline source in the detergent compositions disclosed here is a mixture of an alkali metal carbonate and alkali metal bicarbonate. In some other embodiments, the alkaline source in the detergent compositions disclosed here is just an alkali metal carbonate (*e.g.* all ash). In some embodiments, the alkaline source in the detergent compositions disclosed here is sodium carbonate, sodium bicarbonate, sodium metal silicate, or a mixture thereof. In some embodiments, the alkaline source in the detergent compositions disclosed here is sodium carbonate and sodium bicarbonate. In some  
5  
10  
15  
20  
25  
30  
embodiments, the alkaline source in the detergent compositions disclosed here is just sodium carbonate.

In some embodiments, the detergent produced from the disclosed detergent compositions or method has about 1 wt% to about 90 wt%, 5 wt% to about 85 wt%, 15 wt% to about 80 wt%, 20 wt% to about 75 wt%, 25 wt% to about 70 wt%, 30 wt% to about 65 wt%, 35 wt% to about 60 wt%, 40 wt% to about 55 wt%, or 45 wt% to about 50 wt% of the alkaline source. In some other embodiments, the detergent produced from the disclosed detergent compositions or method has about 80 wt% to about 90 wt%, about 70 wt% to about 80 wt%, about 60 wt% to about 70 wt%, about 50 wt% to about 60 wt%, about 40 wt% to about 50 wt%, about 30 wt% to about 40 wt%, about 20 wt% to about 30 wt%, about 10 wt% to about 10 wt%, about 1 wt% to about 10 wt%, or about 0.1 wt% to about 1 wt% of the alkaline source. In some embodiments, the detergent produced from the disclosed detergent compositions or method has about 90 wt%, about 85 wt%, about 80 wt%, about 75 wt%, about 70 wt%, about 65 wt%, about 60 wt%, about 55 wt%, about 50 wt%, about 45 wt%, about 40 wt%, about 35 wt%, about 30 wt%, about 25 wt%, about 20 wt%, about 15 wt%, about 10 wt%, about 5 wt%, about 1 wt%, or about 0.5 wt% of the alkaline source. In some other embodiments, the detergent produced from the disclosed compositions and methods has about 10 wt% to about 90 wt%, 20 wt% to about 90 wt%, 30 wt% to about 90 wt%, 40 wt% to about 90 wt%, 50 wt% to about 90 wt%, 60 wt% to about 90 wt%, 70 wt% to about 90 wt%, about 85 wt%, 75 wt%, about 65 wt%, about 55 wt%, about 45 wt%, about 35 wt%, about 25 wt%, about 15 wt%, or about 5 wt% of the alkaline source.

In some embodiments, the detergent compositions include a sufficient amount of the alkaline source to provide the use composition with a pH of from about 8 to about 12. In some other embodiment, the detergent compositions include a sufficient amount of the alkaline source to provide the use composition with a pH of from about 8 to about 11, from about 8 to about 9, about 9 to about 12, about 9 to about 11, about 9 to about 10, about 8, about 9, about 10, about 11, about 12, about 8.5, about 9.5, about 10.5, or about 11.5.


In some embodiments, the detergent compositions disclosed here may include additional alkaline compounds, such as alkali metal sesquicarbonate, alkali hydroxide, metasilicate, urea sulfate, amine, amine salt, quaternary ammonia, hydrate thereof, or a mixture of two or more thereof, as additional alkaline source.

In one aspect, provided herein is a composition that comprises an enzyme, a




phosphonate represented by a formula of  $\text{R}^{10}\text{N}(\text{R}^{11})\text{CH}_2\text{PO}(\text{OH})_2$  or salt thereof, and an alkaline source; wherein the enzyme is a protease, amylase, lipase, cellulase, peroxidase, gluconase, or mixture thereof; the alkaline source is a metal carbonate, metal bicarbonate, metal silicate, or mixture thereof;  $\text{R}^{10}$  is hydrogen, a substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, phosphonate, phosphonate ester, or derivative thereof, and  $\text{R}^{11}$  is hydrogen, a substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, phosphonate, phosphonate ester, or derivative thereof; with a proviso that  $\text{R}^{10}$  and  $\text{R}^{11}$  are both  $-\text{CH}_2-\text{PO}(\text{OH})_2$  groups.

In some other embodiments,  $\text{R}^{11}$  is  $-\text{CH}_2-\text{PO}(\text{OH})_2$  group. In yet some other embodiments,  $\text{R}^{11}$  is  $-\text{CH}_2-\text{PO}(\text{OH})_2$  group and  $\text{R}^{10}$  is a substituted alkyl. In some embodiments,  $\text{R}^{11}$  is  $-\text{CH}_2-\text{PO}(\text{OH})_2$  group and  $\text{R}^{10}$  is an phosphonate, phosphonate ester, or derivative thereof.



In some embodiments, the phosphonate is



phosphonic acid, or a mixture thereof, wherein n is an integer of 1-30.

In some embodiments, the phosphonate is aminotrimethylene phosphonic acid  
 5 (ATMP). In some other embodiments, the phosphonate is diglycolamine phosphonate  
 (DGAP).

In some embodiments, the phosphonate is a fully neutralized salt of  
 phosphonebutane tricarboxylic acid (PBTC) by an alkanolamine. In some other  
 embodiments, the phosphonate is a fully neutralized salt of 1-hydroxy ethylidene-1,1-  
 10 diposphonic acid (HEDP) by an alkanolamine. The alkanolamine can be  
 monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof.


In some embodiments, the alkaline source is a metal carbonate and metal  
 bicarbonate. In some other embodiments, the molar ratio of the metal carbonate and the  
 metal bicarbonate is from about 0.25:1 to about 1:0.25, from 0.5:1 to 1:0.5, or from 0.75:1  
 15 to 1:0.75. In some other embodiments, the alkaline source is a metal carbonate (*e.g.* all  
 ash). In some embodiments, the alkaline source is an alkali metal carbonate and alkali  
 metal bicarbonate. In some other embodiments, the molar ratio of the alkali metal  
 carbonate and the alkali metal bicarbonate is from about 0.25:1 to about 1:0.25, from 0.5:1  
 to 1:0.5, or from 0.75:1 to 1:0.75. In some other embodiments, the alkaline source is an  
 20 alkali metal carbonate (all ash).


In some embodiments, the enzyme is a protease, amylase, lipase, or mixture  
 thereof. In some embodiments, the enzyme is a protease, amylase, or mixture thereof. In  
 some embodiments, the enzyme is a protease, lipase, or mixture thereof. In some  
 embodiments, the enzyme is a protease. In some other embodiments, the enzyme is an

amylase, lipase, or mixture thereof. In some other embodiments, the enzyme is an amylase. In yet some other embodiments, the enzyme is a lipase.


In some embodiments, in a use solution of the composition, the enzyme retains at least 15% of its activity at 120 °F for at least 4 hours. In some other embodiments, in a use solution of the composition, the enzyme retains at least 20% of its activity at 120 °F for at least 4 hours. In some embodiments, in a use solution of the detergent composition, the enzyme retains at least 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% of its activity at 120 °F for at least 240 minutes.

In some embodiments, the composition comprises from about 0.1 wt-% to about 5 wt-%, from 0.5 wt-% to about 3 wt-%, from about 1 wt-% to about 1.5 wt-% of an enzyme, from about 0.01 wt-% to about 2 wt-%, from 0.05 wt-% to about 1.5 wt-%, or from 0.1 wt-



% to about 1 wt-% of a phosphonate represented by a formula of  or salt thereof, and from about 50 wt-% to about 95 wt-%, from about 50 wt-% to 90 wt-%, from about 60 wt-% to about 90 wt-%, from about 70 wt-% to about 90 wt-%, or from about 80 wt-% to about 90 wt% of an alkaline source. In some of these embodiments, the phosphonate is aminotrimethylene phosphonic acid (ATMP). In some others of these embodiments, the phosphonate is diglycolamine phosphonate (DGAP). In some of these embodiments, the alkaline source is a mixture of alkali metal carbonate and alkali metal bicarbonate with a ratio of from 0.25:1 to 1:0.5, from 0.5:1 to 1:0.5, or from 0.75:1 to 1:0.75. In some others of these embodiments, the alkaline source is alkali metal carbonate (all ash).


In some embodiments, the composition comprises an enzyme, a phosphonate



represented by a formula of  $\text{R}^{10}$  or salt thereof, an alkaline source and an amine. In some embodiments, the amine is about 0.1 wt-% to about 35 wt-%, 0.1 wt-% to about 30 wt-%, 0.1 wt-% to about 25 wt-%, 0.1 wt-% to about 15 wt-%, 0.1 wt-% to about 10 wt-%, 0.1 wt-% to about 5 wt-%, about 0.5 wt-% to about 5 wt-%, about 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 10 wt-%, 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 5 wt-%, 5 wt-% to about 10 wt-%, 10 wt-% to about 15 wt-%, about 15 wt-% to about 20 wt-%, 20 wt-% to about 25 wt-%, 25 wt-% to about 30 wt-%, 30 wt-% to about 35 wt-%, about 0.1 wt-%, about 0.5 wt-%, about 1 wt-%, about 2 wt-%, about 3 wt-%, about 4 wt-%, about 5 wt-%, about 6 wt-%, about 7 wt-%, about 8 wt-%, about 9 wt-%, about 10 wt-%, about 15 wt-%, about 20 wt-%, about 25 wt-%, about 30 wt-%, about 32 wt-%, or about 35 wt-% of the about 0.1-35 wt-% of the composition.


In some other embodiments, the amine is an alkanolamine or a mixture thereof. In some other embodiments, the amine is monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof. In some other embodiments, the amine of the amine phosphonate salt is a C<sub>2-6</sub> alkylamine or mixture thereof. The alkylamine of course can be a mono-, di-, or tri-amine. In some embodiments, the composition has a molar ratio of the phosphonate to the amine is from about 0.5:1 to 1:0.5.

In some embodiments, the composition comprises an enzyme, a phosphonate




represented by a formula of  $\text{R}^{10}$  or salt thereof, an alkaline source, and a metal hydroxide, tripoly phosphate, or mixture thereof. In some embodiments, the

composition comprises an enzyme, a phosphonate represented by a formula of




or salt thereof, an alkaline source, an amine, and a metal hydroxide, tripoly phosphate, or mixture thereof.

In some other embodiments, the composition comprises an enzyme, a phosphonate



5 represented by a formula of or salt thereof, an alkaline source, a metal hydroxide, tripoly phosphate, or mixture thereof, and one or more additional functional ingredients comprising an oxidizer, builder or water conditioner/water conditioning agent, peroxyacid and its initializer, chelant, threshold agent, crystal modifier; sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, 10 rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance and/or dye, rheology modifier or thickener, nonionic surfactant, cationic surfactant, or zwitterionic surfactant, hydrotrope or coupler, or combination thereof.

In some other embodiments, the composition comprises an enzyme, a phosphonate




15 represented by a formula of or salt thereof, an alkaline source, an amine, a metal hydroxide, tripoly phosphate, or mixture thereof, and one or more

additional functional ingredients comprising an oxidizer, builder or water conditioner/water conditioning agent, peroxyacid and its initializer, chelant, threshold agent, crystal modifier; sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance and/or dye, rheology modifier or thickener, nonionic surfactant, cationic surfactant, or zwitterionic surfactant, hydrotrope or coupler, or combination thereof.

In some embodiments, the composition comprises from about 2 wt-% to about 15 wt-% or from about 5 wt-% to 10 wt-% of a water conditioning agent. In some other embodiments, the composition comprises from about 0.1 wt-% to about 5 wt-%, from about 0.5 wt-% to about 4 wt-%, or from about 1 wt-% to about 3 wt-% of a surfactant.

In other aspect, the disclosure is a composition that comprises an enzyme, an alkaline source, and an amine phosphonate salt; wherein the amine salt is a product of a



phosphonate represented by a formula of and an amine, the enzyme is a protease, amylase, lipase, cellulase, peroxidase, gluconase, or mixture thereof; the alkaline source comprises a metal carbonate, metal bicarbonate, metal silicate, or mixture thereof;  $\text{R}^{12}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof;  $\text{R}^{13}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof; and  $\text{R}^{14}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof.

In some other embodiments, one of  $\text{R}^{12}$ ,  $\text{R}^{13}$ , and  $\text{R}^{14}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , ester thereof, salt thereof, or derivative thereof.


In some embodiments, the phosphonate of the amine phosphonate salt is  $\text{PO}(\text{OH})_2(\text{C}(\text{CH}_2\text{COOH})_3)$ , PBTC. In other embodiments, the phosphonate of the amine phosphonate salt is  $\text{C}(\text{CH}_3)(\text{OH})(\text{PO}(\text{OH})_2)_2$ , HEDP.

In some embodiments, the amine of the amine phosphonate salt is an alkanolamine or a mixture thereof. In other embodiments, the amine of the amine phosphonate salt is monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof. In some other embodiments, the amine of the amine phosphonate salt is a C<sub>2-6</sub> alkylamine or mixture thereof. The alkylamine of course can be a mono-, di-, or tri-amine. In some  
5                   embodiments, the composition has a molar ratio of the phosphonate to the amine is from about 0.5:1 to 1:0.5. In some embodiments, the amine phosphonate salt is only partially neutralized. In some other embodiments, the amine phosphonate salt is fully neutralized by the amine.

10                   In some embodiments, the enzyme is protease, amylase, lipase, or mixture thereof. In some other embodiments, the enzyme is protease, amylase, or mixture thereof. In some embodiments, the enzyme is a protease, lipase, or mixture thereof. In some embodiments, the enzyme is a protease. In some other embodiments, the enzyme is an amylase, lipase, or mixture thereof. In some other embodiments, the enzyme is an amylase. In yet some  
15                   other embodiments, the enzyme is a lipase.

In some embodiments, the alkaline source of the composition is a metal carbonate and metal bicarbonate. In some other embodiments, a molar ratio of the metal carbonate to the metal bicarbonate is from about 0.5:1 to about 1:0.5, from 0.5:1 to 1:0.5, or from 0.75:1 to 1:0.75. In some embodiments, the alkaline source is a metal carbonate. In some  
20                   embodiments, the alkaline source of the composition is an alkali metal carbonate and alkali metal bicarbonate. In some other embodiments, a molar ratio of the alkali metal carbonate to the alkali metal bicarbonate is from about 0.5:1 to about 1:0.5, from 0.5:1 to 1:0.5, or from 0.75:1 to 1:0.75. In some embodiments, the alkaline source is an alkali metal carbonate. In some embodiments, the alkaline source of the composition is sodium  
25                   carbonate and sodium bicarbonate. In some other embodiments, a molar ratio of sodium carbonate to sodium bicarbonate is from about 0.5:1 to about 1:0.5, from 0.5:1 to 1:0.5, or from 0.75:1 to 1:0.75. In some embodiments, the alkaline source is sodium carbonate.

In some embodiments, the composition comprises from about 0.1 wt-% to about 5 wt-%, from 0.5 wt-% to about 3 wt-%, from about 1 wt-% to about 1.5 wt-% of an enzyme,  
30                   from about 0.01 wt-% to about 2 wt-%, from 0.05 wt-% to about 1.5 wt-%, or from 0.1 wt-% to about 1 wt-% of an amine phosphonate salt; wherein the amine salt is a product of a



phosphonate represented by a formula of  $\begin{array}{c} \text{O} \\ \parallel \\ \text{HO}-\text{P}-\text{OH} \\ | \\ \text{R}^{12} \quad \text{R}^{14} \\ | \\ \text{R}^{13} \end{array}$  and an amine, and from about 50 wt-% to about 95 wt-%, from about 50 wt-% to 90 wt-%, from about 60 wt-% to about 90 wt-%, from about 70 wt-% to about 90 wt-%, or from about 80 wt-% to about 90 wt% of an alkaline source. In some of these embodiments, the phosphonate is a fully  
 5 neutralized salt of phosphonebutane tricarboxylic acid (PBTC) by an alkanolamine. In some others of these embodiments, the phosphonate is a fully neutralized salt of 1-hydroxy ethylidene-1,1-diphosphonic acid (HEDP) by an alkanolamine. The alkanolamine can be monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof. In some of these embodiments, the alkaline source is a mixture of alkali metal carbonate  
 10 and alkali metal bicarbonate with a ratio of from 0.25:1 to 1:0.5, from 0.5:1 to 1:0.5, or from 0.75:1 to 1:0.75. In some others of these embodiments, the alkaline source is alkali metal carbonate (all ash).

In some embodiments, in a use solution of the composition, the enzyme retains at least 15% of its activity at 120 °F for at least 40 minutes. In some other embodiments, in a  
 15 use solution of the detergent composition, the enzyme retains at least 20% of its activity at 120 °F for at least 4 hours. In some embodiments, in a use solution of the detergent composition, the enzyme retains at least 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% of its activity at 120 °F for at least 240 minutes.

In some other embodiments, the amine phosphonate salt is about 0.1 wt-% to about  
 20 35 wt-%, 0.1 wt-% to about 30 wt-%, 0.1 wt-% to about 25 wt-%, 0.1 wt-% to about 15 wt-%, 0.1 wt-% to about 10 wt-%, 0.1 wt-% to about 5 wt-%, about 0.5 wt-% to about 5 wt-%, about 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 10 wt-%, 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 5 wt-%, 5 wt-% to about 10 wt-%, 10 wt-% to about 15 wt-%, about 15 wt-% to about 20 wt-%, 20 wt-% to about 25 wt-%, 25 wt-% to about 30 wt-%,  
 25 30 wt-% to about 35 wt-%, about 0.1 wt-%, about 0.5 wt-%, about 1 wt-%, about 2 wt-%, about 3 wt-%, about 4 wt-%, about 5 wt-%, about 6 wt-%, about 7 wt-%, about 8 wt-%,


about 9 wt-%, about 10 wt-%, about 15 wt-%, about 20 wt-%, about 25 wt-%, about 30 wt-%, about 32 wt-%, or about 35 wt-% of the about 0.1-35 wt-% of the composition.

In some embodiments, the composition further comprises a metal hydroxide, tripoly phosphate, or mixture thereof. In some other embodiments, the composition further  
5 comprises one or more additional functional ingredients comprising an oxidizer, builder or water conditioner/water conditioning agent, peroxyacid and its initializer, chelant, threshold agent, crystal modifier; sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance  
10 and/or dye, rheology modifier or thickener, nonionic surfactant, cationic surfactant, or zwitterionic surfactant, hydrotrope or coupler, or combination thereof.

In some embodiments, the composition further comprises a metal hydroxide, tripoly phosphate, or mixture thereof and one or more additional functional ingredients. The additional functional can be an oxidizer, builder or water conditioner/water  
15 conditioning agent, peroxyacid and its initializer, chelant, threshold agent, crystal modifier; sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance and/or dye, rheology modifier or thickener, nonionic surfactant, cationic surfactant, or zwitterionic surfactant, hydrotrope or  
20 coupler, or combination thereof.

In some embodiments, the composition comprises from about 2 wt-% to about 15 wt-% or from about 5 wt-% to 10 wt-% of a water conditioning agent. In some other embodiments, the composition comprises from about 0.1 wt-% to about 5 wt-%, from about 0.5 wt-% to about 4 wt-%, or from about 1 wt-% to about 3 wt-% of a surfactant.

25 In yet another aspect, the disclosure is a solid detergent composition comprising: an alkaline source, a phosphonate, and an enzyme; wherein the alkaline source comprises a metal carbonate, metal bicarbonate, metal silicate, or mixture thereof; the enzyme is a protease, amylase, lipase, cellulase, peroxidase, gluconase, or mixture thereof; the




phosphonate is represented by a formula of  $\text{R}^{10}$  or salt thereof, wherein  $\text{R}^{10}$  is hydrogen, a substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, phosphonate, phosphonate ester, or derivative thereof, and  $\text{R}^{11}$  is hydrogen, a substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, phosphonate, phosphonate ester, or derivative thereof; with  
 5 a proviso that  $\text{R}^{10}$  and  $\text{R}^{11}$  are both  $-\text{CH}_2-\text{PO}(\text{OH})_2$  groups in the molecule, the composition is mixed and used to produce a solid detergent.

In some embodiments, the solid detergent is produced by a cast, extrude, or press process. In other embodiments, the solid detergent is produced by a press process. In  
 10 some embodiments, the solid detergent is a block, tablet, or particulate. In some other embodiments, the solid detergent is a multi-use solid detergent.

In some embodiments, the solid detergent has a dimensional stability and has a growth exponent of less than 3% if heated at a temperature of 122°F. In some other  
 15 embodiments, the solid detergent has a dimensional stability and has a growth exponent of less than 2% if heated at a temperature of 122°F.

In some other embodiments,  $\text{R}^{11}$  is  $-\text{CH}_2-\text{PO}(\text{OH})_2$  group. In yet some other  
 embodiments,  $\text{R}^{11}$  is  $-\text{CH}_2-\text{PO}(\text{OH})_2$  group and  $\text{R}^{10}$  is a substituted alkyl. In some  
 embodiments,  $\text{R}^{11}$  is  $-\text{CH}_2-\text{PO}(\text{OH})_2$  group and  $\text{R}^{10}$  is an phosphonate, phosphonate ester,  
 or derivative thereof.




phosphonic acid, or a mixture thereof, wherein n is an integer of 1-30.

In some embodiments, the alkaline source is a metal carbonate and metal  
 5 bicarbonate. In some other embodiments, the molar ratio of the metal carbonate and the  
 metal bicarbonate is from about 0.25:1 to about 1:0.25. In some embodiments, the alkaline  
 source is a metal carbonate. In some embodiments, the alkaline source of the composition  
 is an alkali metal carbonate and alkali metal bicarbonate. In some other embodiments, a  
 molar ratio of the alkali metal carbonate to the alkali metal bicarbonate is from about 0.5:1  
 10 to about 1:0.5. In some embodiments, the alkaline source is an alkali metal carbonate. In  
 some embodiments, the alkaline source of the composition is sodium carbonate and sodium  
 bicarbonate. In some other embodiments, a molar ratio of sodium carbonate to sodium  
 bicarbonate is from about 0.5:1 to about 1:0.5. In some embodiments, the alkaline source  
 is sodium carbonate.

15 In some embodiments, the enzyme is a protease, amylase, lipase, or mixture  
 thereof. In some embodiments, the enzyme is a protease, amylase, or mixture thereof. In  
 some embodiments, the enzyme is a protease, lipase, or mixture thereof. In some  
 embodiments, the enzyme is a protease. In some other embodiments, the enzyme is an  
 amylase, lipase, or mixture thereof. In some other embodiments, the enzyme is an  
 20 amylase. In yet some other embodiments, the enzyme is a lipase.

In some embodiments, the composition comprises from about 0.1 wt-% to about 5  
 wt-%, from 0.5 wt-% to about 3 wt-%, from about 1 wt-% to about 1.5 wt-% of an enzyme,  
 from about 0.01 wt-% to about 2 wt-%, from 0.05 wt-% to about 1.5 wt-%, or from 0.1 wt-%



5 % to about 1 wt-% of a phosphonate represented by a formula of  $\text{R}^{10}\text{N}(\text{R}^{11})\text{CH}_2\text{P}(\text{OH})_2\text{O}$  or salt thereof, and from about 50 wt-% to about 95 wt-%, from about 50 wt-% to 90 wt-%, from about 60 wt-% to about 90 wt-%, from about 70 wt-% to about 90 wt-%, or from about 80 wt-% to about 90 wt% of an alkaline source. In some of these embodiments, the phosphonate is aminotrimethylene phosphonic acid (ATMP). In some others of these

10 (all ash).

In some embodiments, in a use solution of the solid detergent composition, the enzyme retains at least 15% of its activity at 120 °F for at least 4 hours. In some other

15 solutions of the solid detergent composition, the enzyme retains at least 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% of its activity at 120 °F for at least 240 minutes.

In some embodiments, the solid detergent composition further comprises an amine. In some embodiments, the amine is about 0.1 wt-% to about 35 wt-%, 0.1 wt-% to about 30

20 wt-%, 0.1 wt-% to about 25 wt-%, 0.1 wt-% to about 15 wt-%, 0.1 wt-% to about 10 wt-%, 0.1 wt-% to about 5 wt-%, about 0.5 wt-% to about 5 wt-%, about 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 10 wt-%, 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 5 wt-%, 5 wt-% to about 10 wt-%, 10 wt-% to about 15 wt-%, about 15 wt-% to about 20 wt-%, 20 wt-% to about 25 wt-%, 25 wt-% to about 30 wt-%, 30 wt-% to about 35 wt-%, about 0.1 wt-%, about 0.5 wt-%, about 1 wt-%, about 2 wt-%, about 3 wt-%, about 4 wt-%, about

25 5 wt-%, about 6 wt-%, about 7 wt-%, about 8 wt-%, about 9 wt-%, about 10 wt-%, about 15 wt-%, about 20 wt-%, about 25 wt-%, about 30 wt-%, about 32 wt-%, or about 35 wt-% of the about 0.1-35 wt-% of the solid detergent composition.

In some embodiments, the composition has a molar ratio of the phosphonate to the amine is from about 0.5:1 to 1:0.5. In some other embodiments, the amine is monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof. In some other embodiments, the amine of the amine phosphonate salt is a C<sub>2-6</sub> alkylamine or mixture thereof. The alkylamine of course can be a mono-, di-, or tri-amine.


In some embodiments, the solid detergent composition further comprises a metal hydroxide, tripoly phosphate, or mixture thereof. In some other embodiments, the composition comprises one or more additional functional ingredients comprising an oxidizer, builder or water conditioner/water conditioning agent, peroxyacid and its  
10 initializer, chelant, threshold agent, crystal modifier; sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance and/or dye, rheology modifier or thickener, nonionic surfactant, cationic surfactant, or zwitterionic surfactant, hydrotrope or coupler, and combination thereof.

In some embodiments, the composition further comprises a metal hydroxide, tripoly phosphate, or mixture thereof, an amine and one or more additional functional ingredients. In some other embodiments, the composition further comprises a metal hydroxide, tripoly phosphate, or mixture thereof and an amine. In yet some other  
15 embodiments, the composition further comprises an amine and one or more additional functional ingredients. The additional ingredient can be an oxidizer, builder or water conditioner/water conditioning agent, peroxyacid and its initializer, chelant, threshold agent, crystal modifier; sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance and/or  
20 dye, rheology modifier or thickener, nonionic surfactant, cationic surfactant, or zwitterionic surfactant, hydrotrope or coupler, and combination thereof.

In some embodiments, the composition comprises from about 2 wt-% to about 15 wt-% or from about 5 wt-% to 10 wt-% of a water conditioning agent. In some other  
25 embodiments, the composition comprises from about 0.1 wt-% to about 5 wt-%, from about 0.5 wt-% to about 4 wt-%, or from about 1 wt-% to about 3 wt-% of a surfactant.

In another aspect, the disclosure is a solid detergent composition that comprises an alkaline source, an enzyme, and an amine phosphonate salt; wherein the alkaline source

comprises a metal carbonate, metal bicarbonate, metal silicate, or mixture thereof; the enzyme is a protease, amylase, lipase, cellulase, peroxidase, gluconase, or mixture thereof; the amine phosphonate salt is a product of a phosphonate represented by a formula of



and an amine, wherein  $\text{R}^{12}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof;  $\text{R}^{13}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof; and  $\text{R}^{14}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof; and the composition is mixed and used to produce a solid detergent.

In some embodiments, the solid detergent is produced by a cast, extrude, or press process. In other embodiments, the solid detergent is produced by a press process. In some embodiments, the solid detergent is a block, tablet, or particulate. In some other embodiments, the solid detergent is a multi-use solid detergent.

In some embodiments, the solid detergent has a dimensional stability and has a growth exponent of less than 3% if heated at a temperature of 122°F. In some other embodiments, the solid detergent has a dimensional stability and has a growth exponent of less than 2% if heated at a temperature of 122°F.

In some other embodiments, one of  $\text{R}^{12}$ ,  $\text{R}^{13}$ , and  $\text{R}^{14}$  is hydroxyl, methyl,  $-\text{PO}(\text{OH})_2$ ,  $-\text{CH}_2\text{COOH}$ , ester thereof, salt thereof, or derivative thereof.


In some embodiments, the phosphonate of the amine phosphonate salt is  $\text{PO}(\text{OH})_2(\text{C}(\text{CH}_2\text{COOH})_3)$ , PBTC. In other embodiments, the phosphonate of the amine phosphonate salt is  $\text{C}(\text{CH}_3)(\text{OH})(\text{PO}(\text{OH})_2)_2$ , HEDP. In yet another embodiments, the amine of the amine phosphonate salt is an alkanolamine, monoethanolamine, diethanolamine, triethanolamine, ethanolamine, isopropylamine, or a mixture thereof. In some other embodiments, the amine of the amine phosphonate salt is a  $\text{C}_{2-6}$  alkylamine or mixture thereof. The alkylamine of course can be a mono-, di-, or tri-amine. In some

embodiments, the composition has a molar ratio of the phosphonate to the amine is from about 0.5:1 to 1:0.5.

In some embodiments, the enzyme is protease, amylase, lipase, or mixture thereof. In some other embodiments, the enzyme is protease, amylase, or mixture thereof. In some  
 5 embodiments, the enzyme is a protease, lipase, or mixture thereof. In some embodiments, the enzyme is a protease. In some other embodiments, the enzyme is an amylase, lipase, or mixture thereof. In some other embodiments, the enzyme is an amylase. In yet some other embodiments, the enzyme is a lipase.

In some embodiments, the alkaline source of the composition is a metal carbonate  
 10 and metal bicarbonate. In some other embodiments, a molar ratio of the metal carbonate to the metal bicarbonate is from about 0.25:1 to about 1:0.25, from 0.5:1 to 1:0.5, or from 0.75:1 to 1:0.75. In some other embodiments, the alkaline source is a metal carbonate. In some embodiments, the alkaline source of the composition is an alkali metal carbonate and alkali metal bicarbonate. In some other embodiments, a molar ratio of the alkali metal  
 15 carbonate to the alkali metal bicarbonate is from about 0.5:1 to about 1:0.5. In some embodiments, the alkaline source is an alkali metal carbonate. In some embodiments, the alkaline source of the composition is sodium carbonate and sodium bicarbonate. In some other embodiments, a molar ratio of sodium carbonate to sodium bicarbonate is from about 0.5:1 to about 1:0.5. In some embodiments, the alkaline source is sodium carbonate.

20 In some embodiments, the composition comprises from about 0.1 wt-% to about 5 wt-%, from 0.5 wt-% to about 3 wt-%, from about 1 wt-% to about 1.5 wt-% of an enzyme, from about 0.01 wt-% to about 2 wt-%, from 0.05 wt-% to about 1.5 wt-%, or from 0.1 wt-% to about 1 wt-% of an amine phosphonate salt; wherein the amine salt is a product of a



phosphonate represented by a formula of  
 25 50 wt-% to about 95 wt-%, from about 50 wt-% to 90 wt-%, from about 60 wt-% to about 90 wt-%, from about 70 wt-% to about 90 wt-%, or from about 80 wt-% to about 90 wt% of an alkaline source. In some of these embodiments, the phosphonate is a fully

neutralized salt of phosphonobutane tricarboxylic acid (PBTC) by an alkanolamine. In some others of these embodiments, the phosphonate is a fully neutralized salt of 1-hydroxy ethylidene-1,1-diphosphonic acid (HEDP) by an alkanolamine. The alkanolamine can be monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof.

5 In some of these embodiments, the alkaline source is a mixture of alkali metal carbonate and alkali metal bicarbonate with a ratio of from 0.25:1 to 1:0.5, from 0.5:1 to 1:0.5, or from 0.75:1 to 1:0.75. In some others of these embodiments, the alkaline source is alkali metal carbonate (all ash). In some embodiments, the amine phosphonate salt is only partially neutralized. In some other embodiments, the amine phosphonate salt is fully  
10 neutralized by the amine.

In some embodiments, in a use solution of the solid detergent composition, the enzyme retains at least 15% of its activity at 120 °F for at least 4 hours. In some other embodiments, in a use solution of the solid detergent composition, the enzyme retains at least 50% of its activity at 120 °F for at least 4 hours. In some embodiments, in a use  
15 solution of the solid detergent composition, the enzyme retains at least 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% of its activity at 120 °F for at least 240 minutes.

In some other embodiments, the amine phosphonate salt is about 0.1 wt-% to about 35 wt-%, 0.1 wt-% to about 30 wt-%, 0.1 wt-% to about 25 wt-%, 0.1 wt-% to about 15 wt-%, 0.1 wt-% to about 10 wt-%, 0.1 wt-% to about 5 wt-%, about 0.5 wt-% to about 5 wt-%,  
20 about 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 10 wt-%, 0.1 wt-% to about 1 wt-%, about 1 wt-% to about 5 wt-%, 5 wt-% to about 10 wt-%, 10 wt-% to about 15 wt-%, about 15 wt-% to about 20 wt-%, 20 wt-% to about 25 wt-%, 25 wt-% to about 30 wt-%, 30 wt-% to about 35 wt-%, about 0.1 wt-%, about 0.5 wt-%, about 1 wt-%, about 2 wt-%, about 3 wt-%, about 4 wt-%, about 5 wt-%, about 6 wt-%, about 7 wt-%, about 8 wt-%,  
25 about 9 wt-%, about 10 wt-%, about 15 wt-%, about 20 wt-%, about 25 wt-%, about 30 wt-%, about 32 wt-%, or about 35 wt-% of the about 0.1-35 wt-% of the composition.

In some embodiments, the composition further comprises a metal hydroxide, tripoly phosphate, or mixture thereof. In some other embodiments, the composition further comprises one or more additional functional ingredients comprising an oxidizer,  
30 builder or water conditioner/water conditioning agent, peroxyacid and its initializer, chelant, threshold agent, crystal modifier; sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, rinse aid, polymer,

metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance and/or dye, rheology modifier or thickener, nonionic surfactant, cationic surfactant, or zwitterionic surfactant, hydrotrope or coupler, and combination thereof.

In some embodiments, the composition further comprises a metal hydroxide,  
5 tripoly phosphate, or mixture thereof and one or more additional functional ingredients. The additional ingredient can be an oxidizer, builder or water conditioner/water conditioning agent, peroxyacid and its initializer, chelant, threshold agent, crystal modifier; sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility  
10 modifier, dispersant, rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance and/or dye, rheology modifier or thickener, nonionic surfactant, cationic surfactant, or zwitterionic surfactant, hydrotrope or coupler, and combination thereof.

In some embodiments, the composition comprises from about 2 wt-% to about 15 wt-% or from about 5 wt-% to 10 wt-% of a water conditioning agent. In some other  
15 embodiments, the composition comprises from about 0.1 wt-% to about 5 wt-%, from about 0.5 wt-% to about 4 wt-%, or from about 1 wt-% to about 3 wt-% of a surfactant.

#### *Additional Functional Ingredients*

In some embodiments, the disclosed compositions contain additional ingredients.  
20 These ingredients can be in solid or liquid form and therefore be mixed with other components of the disclosed compositions.

The functional ingredients provide desired properties and functionalities to the detergent composition. For the purpose of this application, the term "functional ingredients" includes an ingredient that when dispersed or dissolved in a use and/or  
25 concentrate, such as an aqueous solution, provides a beneficial property in a particular use. Some particular examples of functional ingredients are discussed in more detail below, although the particular materials discussed are given by way of example only, and that a broad variety of other functional ingredients may be used. For example, many of the functional ingredients discussed below relate to materials used in cleaning applications.  
30 However, other embodiments may include functional ingredients for use in other applications.

Exemplary additional functional ingredients include for example: builders or water conditioners/water conditioning agents, including detergent builders; chelants; threshold agents; crystal modifiers; hardening agents; bleaching agents; fillers; defoaming agents; anti-redeposition agents; stabilizing agents; dispersants; glass and metal corrosion inhibitors; fragrances and dyes; thickeners; etc. Further description of suitable additional functional ingredients is set forth in U.S. Patent Application Serial No. 12/977,340.

In some embodiments, the blocks produced from the disclosed method, process, or composition further comprises additional functional ingredient comprising an oxidizer, peroxyacid and its initializer, sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, threshold agent, crystal modifier, phosphonate, binding agent, rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant and/or chelating agent, fragrance and/or dye, rheology modifier or thickener, anionic surfactant, nonionic surfactant, cationic surfactant, amphoteric surfactant, zwitterionic surfactant, hydrotrope or coupler, and combination thereof.

#### *Anionic Surfactants*

The method of adjusting dispense rate of a solid detergent block of a detergent composition, the process to produce a solid detergent block with a predetermined dispense rate, or the press solid composition according to this disclosure includes a first solid comprising an effective amount of one or more anionic surfactants.


Anionic surfactants are surface active substances in which the charge on the hydrophobe is negative; or surfactants in which the hydrophobic section of the molecule carries no charge unless the pH is elevated to neutrality or above (*e.g.*, carboxylic acids). Carboxylate, sulfonate, sulfate and phosphate are the polar (hydrophilic) solubilizing groups found in anionic surfactants. Of the cations (counter ions) associated with these polar groups, sodium, lithium and potassium impart water solubility; ammonium and substituted ammonium ions provide both water and oil solubility; and, calcium, barium, and magnesium promote oil solubility. As those skilled in the art understand, anionics are excellent deterative surfactants and are therefore favored additions to heavy duty detergent compositions.


Anionic sulfate surfactants suitable for use in the present compositions include alkyl ether sulfates, alkyl sulfates, the linear and branched primary and secondary alkyl sulfates, alkyl ethoxysulfates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, the C<sub>5</sub>-C<sub>17</sub> acyl-N-(C<sub>1</sub>-C<sub>4</sub> alkyl) and -N-(C<sub>1</sub>-C<sub>2</sub> hydroxyalkyl) glucamine sulfates, and sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside, and the like. Also included are the alkyl sulfates, alkyl poly(ethyleneoxy) ether sulfates and aromatic poly(ethyleneoxy) sulfates such as the sulfates or condensation products of ethylene oxide and nonyl phenol (usually having 1 to 6 oxyethylene groups per molecule).


Anionic sulfonate surfactants suitable for use in the present compositions also include alkyl sulfonates, the linear and branched primary and secondary alkyl sulfonates, and the aromatic sulfonates with or without substituents.


Anionic carboxylate surfactants suitable for use in the present compositions include carboxylic acids (and salts), such as alkanolic acids (and alkanooates), ester carboxylic acids (*e.g.*, alkyl succinates), ether carboxylic acids, sulfonated fatty acids, such as sulfonated oleic acid, and the like. Such carboxylates include alkyl ethoxy carboxylates, alkyl aryl ethoxy carboxylates, alkyl polyethoxy polycarboxylate surfactants and soaps (*e.g.*, alkyl carboxyls). Secondary carboxylates useful in the present compositions include those which contain a carboxyl unit connected to a secondary carbon. The secondary carbon can be in a ring structure, *e.g.*, as in *p*-octyl benzoic acid, or as in alkyl-substituted cyclohexyl carboxylates. The secondary carboxylate surfactants typically contain no ether linkages, no ester linkages and no hydroxyl groups. Further, they typically lack nitrogen atoms in the head-group (amphiphilic portion). Suitable secondary soap surfactants typically contain 11-13 total carbon atoms, although more carbons atoms (*e.g.*, up to 16) can be present. Suitable carboxylates also include acylamino acids (and salts), such as acylgluamates, acyl peptides, sarcosinates (*e.g.*, N-acyl sarcosinates), taurates (*e.g.*, N-acyl taurates and fatty acid amides of methyl tauride), and the like.


Suitable anionic surfactants include alkyl or alkylaryl ethoxy carboxylates of the following formula:





in which R is a C<sub>8</sub> to C<sub>22</sub> alkyl group or , in which R<sup>1</sup> is a C<sub>4</sub>-C<sub>16</sub> alkyl group; n is an integer of 1-20; m is an integer of 1-3; and X is a counter ion, such as hydrogen, sodium, potassium, lithium, ammonium, or an amine salt such as monoethanolamine, diethanolamine or triethanolamine. In some embodiments, n is an integer of 4 to 10 and m is 1. In some embodiments, R is a C<sub>8</sub>-C<sub>16</sub> alkyl group. In some  
 5       embodiments, R is a C<sub>12</sub>-C<sub>14</sub> alkyl group, n is 4, and m is 1.



In other embodiments, R is  and R<sup>1</sup> is a C<sub>6</sub>-C<sub>12</sub> alkyl group. In still yet other embodiments, R<sup>1</sup> is a C<sub>9</sub> alkyl group, n is 10 and m is 1.

Such alkyl and alkylaryl ethoxy carboxylates are commercially available. These  
 10       ethoxy carboxylates are typically available as the acid forms, which can be readily converted to the anionic or salt form. Commercially available carboxylates include, Neodox 23-4, a C<sub>12-13</sub> alkyl polyethoxy (4) carboxylic acid (Shell Chemical), and Emcol CNP-110, a C<sub>9</sub> alkylaryl polyethoxy (10) carboxylic acid (Witco Chemical). Carboxylates are also available from Clariant, *e.g.*, the product Sandopan<sup>®</sup> DTC, a C<sub>13</sub> alkyl polyethoxy  
 15       (7) carboxylic acid.

#### *Nonionic Surfactants*

Useful nonionic surfactants are generally characterized by the presence of an organic hydrophobic group and an organic hydrophilic group and are typically produced by  
 20       the condensation of an organic aliphatic, alkyl aromatic or polyoxyalkylene hydrophobic compound with a hydrophilic alkaline oxide moiety which in common practice is ethylene oxide or a polyhydration product thereof, polyethylene glycol. Practically any hydrophobic compound having a hydroxyl, carboxyl, amino, or amido group with a reactive hydrogen atom can be condensed with ethylene oxide, or its polyhydration adducts, or its mixtures  
 25       with alkoxylenes such as propylene oxide to form a nonionic surface-active agent. The length of the hydrophilic polyoxyalkylene moiety which is condensed with any particular hydrophobic compound can be readily adjusted to yield a water dispersible or water soluble compound having the desired degree of balance between hydrophilic and hydrophobic properties. Useful nonionic surfactants include:

Block polyoxypropylene-polyoxyethylene polymeric compounds based upon propylene glycol, ethylene glycol, glycerol, trimethylolpropane, and ethylenediamine as the initiator reactive hydrogen compound. Examples of polymeric compounds made from a sequential propoxylation and ethoxylation of initiator are commercially available from

5 BASF Corp. One class of compounds are difunctional (two reactive hydrogens) compounds formed by condensing ethylene oxide with a hydrophobic base formed by the addition of propylene oxide to the two hydroxyl groups of propylene glycol. This hydrophobic portion of the molecule weighs from about 1,000 to about 4,000. Ethylene oxide is then added to sandwich this hydrophobe between hydrophilic groups, controlled

10 by length to constitute from about 10% by weight to about 80% by weight of the final molecule. Another class of compounds are tetra-functional block copolymers derived from the sequential addition of propylene oxide and ethylene oxide to ethylenediamine. The molecular weight of the propylene oxide hydrotype ranges from about 500 to about 7,000; and, the hydrophile, ethylene oxide, is added to constitute from about 10% by weight to

15 about 80% by weight of the molecule.

Condensation products of one mole of alkyl phenol wherein the alkyl chain, of straight chain or branched chain configuration, or of single or dual alkyl constituent, contains from about 8 to about 18 carbon atoms with from about 3 to about 50 moles of ethylene oxide. The alkyl group can, for example, be represented by diisobutylene, di-

20 amyl, polymerized propylene, iso-octyl, nonyl, and di-nonyl. These surfactants can be polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols. Examples of commercial compounds of this chemistry are available on the market under the trade names Igepal<sup>®</sup> manufactured by Rhone-Poulenc and Triton<sup>®</sup> manufactured by Union Carbide.

25 Condensation products of one mole of a saturated or unsaturated, straight or branched chain alcohol having from about 6 to about 24 carbon atoms with from about 3 to about 50 moles of ethylene oxide. The alcohol moiety can consist of mixtures of alcohols in the above delineated carbon range or it can consist of an alcohol having a specific number of carbon atoms within this range. Examples of like commercial surfactant are

30 available under the trade names Lutensol<sup>™</sup>, Dehydol<sup>™</sup> manufactured by BASF, Neodol<sup>™</sup> manufactured by Shell Chemical Co. and Alfonic<sup>™</sup> manufactured by Vista Chemical Co.

Condensation products of one mole of saturated or unsaturated, straight or branched chain carboxylic acid having from about 8 to about 18 carbon atoms with from about 6 to about 50 moles of ethylene oxide. The acid moiety can consist of mixtures of acids in the above defined carbon atoms range or it can consist of an acid having a specific number of carbon atoms within the range. Examples of commercial compounds of this chemistry are available on the market under the trade names Disponil or Agnique manufactured by BASF and Lipopeg<sup>TM</sup> manufactured by Lipo Chemicals, Inc.

In addition to ethoxylated carboxylic acids, commonly called polyethylene glycol esters, other alkanolic acid esters formed by reaction with glycerides, glycerin, and polyhydric (saccharide or sorbitan/sorbitol) alcohols have application in this disclosure for specialized embodiments, particularly indirect food additive applications. All of these ester moieties have one or more reactive hydrogen sites on their molecule which can undergo further acylation or ethylene oxide (alkoxide) addition to control the hydrophilicity of these substances. Care must be exercised when adding these fatty ester or acylated carbohydrates to compositions of the present disclosure containing amylase and/or lipase enzymes because of potential incompatibility.

Examples of nonionic low foaming surfactants include:


Compounds from (1) which are modified, essentially reversed, by adding ethylene oxide to ethylene glycol to provide a hydrophile of designated molecular weight; and, then adding propylene oxide to obtain hydrophobic blocks on the outside (ends) of the molecule. The hydrophobic portion of the molecule weighs from about 1,000 to about 3,100 with the central hydrophile including 10% by weight to about 80% by weight of the final molecule. These reverse Pluronics<sup>TM</sup> are manufactured by BASF Corporation under the trade name Pluronic<sup>TM</sup> R surfactants. Likewise, the Tetronic<sup>TM</sup> R surfactants are produced by BASF Corporation by the sequential addition of ethylene oxide and propylene oxide to ethylenediamine. The hydrophobic portion of the molecule weighs from about 2,100 to about 6,700 with the central hydrophile including 10% by weight to 80% by weight of the final molecule.

Compounds from groups (1), (2), (3) and (4) which are modified by "capping" or "end blocking" the terminal hydroxy group or groups (of multi-functional moieties) to reduce foaming by reaction with a small hydrophobic molecule such as propylene oxide, butylene oxide, benzyl chloride; and, short chain fatty acids, alcohols or alkyl halides

containing from 1 to about 5 carbon atoms; and mixtures thereof. Also included are reactants such as thionyl chloride which convert terminal hydroxy groups to a chloride group. Such modifications to the terminal hydroxy group may lead to all-block, block-heteric, heteric-block or all-heteric nonionics.

5 Additional examples of effective low foaming nonionics include:

The alkylphenoxypolyethoxyalkanols of U.S. Pat. No. 2,903,486 issued Sep. 8, 1959 to Brown et al. and represented by the formula



10 in which R is an alkyl group of 8 to 9 carbon atoms, A is an alkylene chain of 3 to 4 carbon atoms, n is an integer of 7 to 16, and m is an integer of 1 to 10.

The polyalkylene glycol condensates of U.S. Pat. No. 3,048,548 issued Aug. 7, 1962 to Martin et al. having alternating hydrophilic oxyethylene chains and hydrophobic oxypropylene chains where the weight of the terminal hydrophobic chains, the weight of the middle hydrophobic unit and the weight of the linking hydrophilic units each represent  
15 about one-third of the condensate.

The defoaming nonionic surfactants disclosed in U.S. Pat. No. 3,382,178 issued May 7, 1968 to Lissant et al. having the general formula  $Z[(\text{OR})_n\text{OH}]_z$  wherein Z is alkoxylatable material, R is a radical derived from an alkylene oxide which can be ethylene and propylene and n is an integer from, for example, 10 to 2,000 or more and z is an  
20 integer determined by the number of reactive oxyalkylatable groups.

The conjugated polyoxyalkylene compounds described in U.S. Pat. No. 2,677,700, issued May 4, 1954 to Jackson et al. corresponding to the formula  $\text{Y}(\text{C}_3\text{H}_6\text{O})_n(\text{C}_2\text{H}_4\text{O})_m\text{H}$  wherein Y is the residue of organic compound having from about 1 to 6 carbon atoms and one reactive hydrogen atom, n has an average value of at least about 6.4, as determined by  
25 hydroxyl number and m has a value such that the oxyethylene portion constitutes about 10% to about 90% by weight of the molecule.

The conjugated polyoxyalkylene compounds described in U.S. Pat. No. 2,674,619, issued Apr. 6, 1954 to Lundsted et al. having the formula  $Y[(C_3H_6O)_n(C_2H_4O)_mH]_x$  wherein Y is the residue of an organic compound having from about 2 to 6 carbon atoms and containing x reactive hydrogen atoms in which x has a value of at least about 2, n has a value such that the molecular weight of the polyoxypropylene hydrophobic base is at least about 900 and m has value such that the oxyethylene content of the molecule is from about 10% to about 90% by weight. Compounds falling within the scope of the definition for Y include, for example, propylene glycol, glycerine, pentaerythritol, trimethylolpropane, ethylenediamine and the like. The oxypropylene chains optionally, but advantageously, contain small amounts of ethylene oxide and the oxyethylene chains also optionally, but advantageously, contain small amounts of propylene oxide.

Additional conjugated polyoxyalkylene surface-active agents which are advantageously used in the compositions of this disclosure correspond to the formula:  $P[(C_3H_6O)_n(C_2H_4O)_mH]_x$  wherein P is the residue of an organic compound having from about 8 to 18 carbon atoms and containing x reactive hydrogen atoms in which x has a value of 1 or 2, n has a value such that the molecular weight of the polyoxyethylene portion is at least about 44 and m has a value such that the oxypropylene content of the molecule is from about 10% to about 90% by weight. In either case the oxypropylene chains may contain optionally, but advantageously, small amounts of ethylene oxide and the oxyethylene chains may contain also optionally, but advantageously, small amounts of propylene oxide.

Polyhydroxy fatty acid amide surfactants suitable for use in the present compositions include those having the structural formula  $R_2CONR_1Z$  in which: R<sub>1</sub> is H, C<sub>1</sub>-C<sub>4</sub> hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl, ethoxy, propoxy group, or a mixture thereof; R<sub>2</sub> is a C<sub>5</sub>-C<sub>31</sub> hydrocarbyl, which can be straight-chain; and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative (preferably ethoxyated or propoxyated) thereof. Z can be derived from a reducing sugar in a reductive amination reaction; such as a glycityl moiety.

The alkyl ethoxylate condensation products of aliphatic alcohols with from about 0 to about 25 moles of ethylene oxide are suitable for use in the present compositions. The

alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from 6 to 22 carbon atoms.

The ethoxylated C<sub>6</sub>-C<sub>18</sub> fatty alcohols and C<sub>6</sub>-C<sub>18</sub> mixed ethoxylated and propoxylated fatty alcohols are suitable surfactants for use in the present compositions, particularly those that are water soluble. Suitable ethoxylated fatty alcohols include the C<sub>6</sub>-C<sub>18</sub> ethoxylated fatty alcohols with a degree of ethoxylation of from 3 to 50.

Suitable nonionic alkylpolysaccharide surfactants, particularly for use in the present compositions include those disclosed in U.S. Pat. No. 4,565,647, Llenado, issued Jan. 21, 1986. These surfactants include a hydrophobic group containing from about 6 to about 30 carbon atoms and a polysaccharide, *e.g.*, a polyglycoside, hydrophilic group containing from about 1.3 to about 10 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, *e.g.*, glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties. (Optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside.) The intersaccharide bonds can be, *e.g.*, between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6-positions on the preceding saccharide units.

Fatty acid amide surfactants suitable for use the present compositions include those having the formula: R<sub>6</sub>CON(R<sub>7</sub>)<sub>2</sub> in which R<sub>6</sub> is an alkyl group containing from 7 to 21 carbon atoms and each R<sub>7</sub> is independently hydrogen, C<sub>1</sub>- C<sub>4</sub> alkyl, C<sub>1</sub>- C<sub>4</sub> hydroxyalkyl, or --(C<sub>2</sub>H<sub>4</sub>O)<sub>x</sub>H, where x is in the range of from 1 to 3.

A useful class of non-ionic surfactants include the class defined as alkoxyated amines or, most particularly, alcohol alkoxyated/aminated/alkoxyated surfactants. These non-ionic surfactants may be at least in part represented by the general formulae: R<sup>20</sup>--(PO)<sub>s</sub>N--(EO)<sub>t</sub>H, R<sup>20</sup>--(PO)<sub>s</sub>N--(EO)<sub>t</sub>H(EO)<sub>u</sub>H, and R<sup>20</sup>--N(EO)<sub>t</sub>H; in which R<sup>20</sup> is an alkyl, alkenyl or other aliphatic group, or an alkyl-aryl group of from 8 to 20, preferably 12 to 14 carbon atoms, EO is oxyethylene, PO is oxypropylene, s is 1 to 20, preferably 2-5, t is 1-10, preferably 2-5, and u is 1-10, preferably 2-5. Other variations on the scope of these compounds may be represented by the alternative formula: R<sup>20</sup>--(PO)<sub>v</sub>--N[(EO)<sub>w</sub>H][(EO)<sub>z</sub>H] in which R<sup>20</sup> is as defined above, v is 1 to 20 (*e.g.*, 1, 2, 3, or 4 (preferably 2)), and w and z are independently 1-10, preferably 2-5. These compounds are represented commercially by a line of products sold by Huntsman Chemicals as nonionic surfactants. A


preferred chemical of this class includes Surfonic™ PEA 25 Amine Alkoxylate. Preferred nonionic surfactants for the compositions of the disclosure include alcohol alkoxylates, EO/PO block copolymers, alkylphenol alkoxylates, and the like.

The treatise *Nonionic Surfactants*, edited by Schick, M. J., Vol. 1 of the Surfactant Science Series, Marcel Dekker, Inc., New York, 1983 is an excellent reference on the wide variety of nonionic compounds generally employed in the practice of the present disclosure. A typical listing of nonionic classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975. Further examples are given in "Surface Active Agents and detergents" (Vol. I and II by Schwartz, Perry and Berch).

#### *Semi-Polar Nonionic Surfactants*

The semi-polar type of nonionic surface active agents are another class of nonionic surfactant useful in compositions of the present disclosure. Generally, semi-polar nonionics are high foamers and foam stabilizers, which can limit their application in CIP systems. However, within compositional embodiments of this disclosure designed for high foam cleaning methodology, semi-polar nonionics would have immediate utility. The semi-polar nonionic surfactants include the amine oxides, phosphine oxides, sulfoxides and their alkoxylated derivatives.

Amine oxides are tertiary amine oxides corresponding to the general formula:




wherein the arrow is a conventional representation of a semi-polar bond; and, R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> may be aliphatic, aromatic, heterocyclic, alicyclic, or combinations thereof. Generally, for amine oxides of detergent interest, R<sup>1</sup> is an alkyl radical of from about 8 to about 24 carbon atoms; R<sup>2</sup> and R<sup>3</sup> are alkyl or hydroxyalkyl of 1-3 carbon atoms or a mixture thereof; R<sup>2</sup> and R<sup>3</sup> can be attached to each other, *e.g.* through an oxygen or nitrogen atom, to form a ring structure; R<sup>4</sup> is an alkylene or a hydroxyalkylene group containing 2 to 3 carbon atoms; and n ranges from 0 to about 20.

Useful water soluble amine oxide surfactants are selected from the coconut or

tallow alkyl di-(lower alkyl) amine oxides, specific examples of which are dodecyl dimethylamine oxide, tridecyl dimethylamine oxide, e-tradecyl dimethylamine oxide, pentadecyl dimethylamine oxide, hexadecyl dimethylamine oxide, heptadecyl dimethylamine oxide, octadecyl dimethylamine oxide, dodecyl dipropylamine oxide, tetradecyl dipropylamine oxide, hexadecyl dipropylamine oxide, tetradecyl dibutylamine oxide, octadecyl dibutylamine oxide, bis(2-hydroxyethyl) dodecylamine oxide, bis(2-hydroxyethyl)-3-dodecoxy-1-hydroxypropylamine oxide, dimethyl-(2-hydroxydodecyl)amine oxide, 3,6,9-trioctadecyl dimethylamine oxide and 3-dodecoxy-2-hydroxypropyl di-(2-hydroxyethyl)amine oxide.


Useful semi-polar nonionic surfactants also include the water soluble phosphine oxides having the following structure:



wherein the arrow is a conventional representation of a semi-polar bond; and,  $\text{R}^1$  is an alkyl, alkenyl or hydroxyalkyl moiety ranging from 10 to about 24 carbon atoms in chain length; and,  $\text{R}^2$  and  $\text{R}^3$  are each alkyl moieties separately selected from alkyl or hydroxyalkyl groups containing 1 to 3 carbon atoms.

Examples of useful phosphine oxides include dimethyldecylphosphine oxide, dimethyltetradecylphosphine oxide, methylethyltetradecylphosphine oxide, dimethylhexadecylphosphine oxide, diethyl-2-hydroxyoctyldecylphosphine oxide, bis(2-hydroxyethyl)dodecylphosphine oxide, and bis(hydroxymethyl)tetradecylphosphine oxide.

Semi-polar nonionic surfactants useful herein also include the water soluble sulfoxide compounds which have the structure:



wherein the arrow is a conventional representation of a semi-polar bond; and,  $\text{R}^1$  is an alkyl or hydroxyalkyl moiety of about 8 to about 28 carbon atoms, from 0 to about 5

ether linkages and from 0 to about 2 hydroxyl substituents; and R<sup>2</sup> is an alkyl moiety consisting of alkyl and hydroxyalkyl groups having 1 to 3 carbon atoms.

Useful examples of these sulfoxides include dodecyl methyl sulfoxide; 3-hydroxy tridecyl methyl sulfoxide; 3-methoxy tridecyl methyl sulfoxide; and 3-hydroxy-4-dodecoxybutyl methyl sulfoxide.

Semi-polar nonionic surfactants for the compositions of the disclosure include dimethyl amine oxides, such as lauryl dimethyl amine oxide, myristyl dimethyl amine oxide, cetyl dimethyl amine oxide, combinations thereof, and the like. Useful water soluble amine oxide surfactants are selected from the octyl, decyl, dodecyl, isododecyl, coconut, or tallow alkyl di-(lower alkyl) amine oxides, specific examples of which are octyldimethylamine oxide, nonyldimethylamine oxide, decyldimethylamine oxide, undecyldimethylamine oxide, dodecyldimethylamine oxide, iso-dodecyldimethylamine oxide, tridecyldimethylamine oxide, tetradecyldimethylamine oxide, pentadecyldimethylamine oxide, hexadecyldimethylamine oxide, heptadecyldimethylamine oxide, octadecyldimethylamine oxide, dodecyldipropylamine oxide, tetradecyldipropylamine oxide, hexadecyldipropylamine oxide, tetradecyldibutylamine oxide, octadecyldibutylamine oxide, bis(2-hydroxyethyl)dodecylamine oxide, bis(2-hydroxyethyl)-3-dodecoxy-1-hydroxypropylamine oxide, dimethyl-(2-hydroxydodecyl)amine oxide, 3,6,9-trioctadecyldimethylamine oxide and 3-dodecoxy-2-hydroxypropyl-di-(2-hydroxyethyl)amine oxide.

Suitable nonionic surfactants suitable for use with the compositions of the present disclosure include alkoxyated surfactants. Suitable alkoxyated surfactants include EO/PO copolymers, capped EO/PO copolymers, alcohol alkoxyates, capped alcohol alkoxyates, mixtures thereof, or the like. Suitable alkoxyated surfactants for use as solvents include EO/PO block copolymers, such as the Pluronic and reverse Pluronic surfactants; alcohol alkoxyates, such as Dehypon LS-54 (R-(EO)<sub>5</sub>(PO)<sub>4</sub>) and Dehypon LS-36 (R-(EO)<sub>3</sub>(PO)<sub>6</sub>); and capped alcohol alkoxyates, such as Plurafac LF221 and Tegoten EC11; mixtures thereof, or the like.

### *Cationic Surfactants*


Surface active substances are classified as cationic if the charge on the hydrotrope

portion of the molecule is positive. Surfactants in which the hydrotrope carries no charge unless the pH is lowered close to neutrality or lower, but which are then cationic (e.g. alkyl amines), are also included in this group. In theory, cationic surfactants may be synthesized from any combination of elements containing an "onium" structure  $R_nX^+Y^-$  and could include compounds other than nitrogen (ammonium) such as phosphorus (phosphonium) and sulfur (sulfonium). In practice, the cationic surfactant field is dominated by nitrogen containing compounds, probably because synthetic routes to nitrogenous cationics are simple and straightforward and give high yields of product, which can make them less expensive.

Cationic surfactants preferably include, more preferably refer to, compounds containing at least one long carbon chain hydrophobic group and at least one positively charged nitrogen. The long carbon chain group may be attached directly to the nitrogen atom by simple substitution; or more preferably indirectly by a bridging functional group or groups in so-called interrupted alkylamines and amido amines. Such functional groups can make the molecule more hydrophilic and/or more water dispersible, more easily water solubilized by co-surfactant mixtures, and/or water soluble. For increased water solubility, additional primary, secondary or tertiary amino groups can be introduced or the amino nitrogen can be quaternized with low molecular weight alkyl groups. Further, the nitrogen can be a part of branched or straight chain moiety of varying degrees of unsaturation or of a saturated or unsaturated heterocyclic ring. In addition, cationic surfactants may contain complex linkages having more than one cationic nitrogen atom.

The surfactant compounds classified as amine oxides, amphoteric and zwitterions are themselves typically cationic in near neutral to acidic pH solutions and can overlap surfactant classifications. Polyoxyethylated cationic surfactants generally behave like nonionic surfactants in alkaline solution and like cationic surfactants in acidic solution.

The simplest cationic amines, amine salts and quaternary ammonium compounds can be schematically drawn thus:



in which, R represents an alkyl chain, R', R'', and R''' may be either alkyl chains or aryl groups or hydrogen and X represents an anion. The amine salts and quaternary ammonium compounds are preferred for practical use in this disclosure due to their high degree of water solubility.


5           The majority of large volume commercial cationic surfactants can be subdivided into four major classes and additional sub-groups known to those of skill in the art and described in "Surfactant Encyclopedia", *Cosmetics & Toiletries*, Vol. 104 (2) 86-96 (1989). The first class includes alkylamines and their salts. The second class includes alkyl imidazolines. The third class includes ethoxylated amines. The fourth class includes  
10           quaternaries, such as alkylbenzyltrimethylammonium salts, alkyl benzene salts, heterocyclic ammonium salts, tetra alkylammonium salts, and the like. Cationic surfactants are known to have a variety of properties that can be beneficial in the present compositions. These desirable properties can include detergency in compositions of or below neutral pH, antimicrobial efficacy, thickening or gelling in cooperation with other agents, and the like.

15           Cationic surfactants useful in the compositions of the present disclosure include those having the formula  $R^1_m R^2_x Y_L Z$  wherein each  $R^1$  is an organic group containing a straight or branched alkyl or alkenyl group optionally substituted with up to three phenyl or hydroxy groups and optionally interrupted by up to four of the following structures:



20           or an isomer or mixture of these structures, and which contains from about 8 to 22 carbon atoms. The  $R^1$  groups can additionally contain up to 12 ethoxy groups.  $m$  is a number from 1 to 3. Preferably, no more than one  $R^1$  group in a molecule has 16 or more carbon atoms when  $m$  is 2 or more than 12 carbon atoms when  $m$  is 3. Each  $R^2$  is an alkyl or  
25           hydroxyalkyl group containing from 1 to 4 carbon atoms or a benzyl group with no more than one  $R^2$  in a molecule being benzyl, and  $x$  is a number from 0 to 11, preferably from 0 to 6. The remainder of any carbon atom positions on the Y group are filled by hydrogens.

Y is can be a group including, but not limited to:



- or a mixture thereof. Preferably, L is 1 or 2, with the Y groups being separated by a moiety selected from R<sup>1</sup> and R<sup>2</sup> analogs (preferably alkylene or alkenylene) having from 1 to about 22 carbon atoms and two free carbon single bonds when L is 2. Z is a water soluble anion, such as a halide, sulfate, methylsulfate, hydroxide, or nitrate anion, particularly preferred being chloride, bromide, iodide, sulfate or methyl sulfate anions, in a number to give electrical neutrality of the cationic component.

#### 10 *Amphoteric Surfactants*

Amphoteric, or ampholytic, surfactants contain both a basic and an acidic hydrophilic group and an organic hydrophobic group. These ionic entities may be any of anionic or cationic groups described herein for other types of surfactants. A basic nitrogen and an acidic carboxylate group are the typical functional groups employed as the basic

and acidic hydrophilic groups. In a few surfactants, sulfonate, sulfate, phosphonate or phosphate provide the negative charge.


Amphoteric surfactants can be broadly described as derivatives of aliphatic secondary and tertiary amines, in which the aliphatic radical may be straight chain or branched and wherein one of the aliphatic substituents contains from about 8 to 18 carbon atoms and one contains an anionic water solubilizing group, *e.g.*, carboxy, sulfo, sulfato, phosphato, or phosphono. Amphoteric surfactants are subdivided into two major classes known to those of skill in the art and described in "Surfactant Encyclopedia" Cosmetics & Toiletries, Vol. 104 (2) 69-71 (1989).

The first class includes acyl/dialkyl ethylenediamine derivatives (*e.g.* 2-alkyl hydroxyethyl imidazoline derivatives) and their salts. The second class includes N-alkylamino acids and their salts. Some amphoteric surfactants can be envisioned as fitting into both classes.


Amphoteric surfactants can be synthesized by methods known to those of skill in the art. For example, 2-alkyl hydroxyethyl imidazoline is synthesized by condensation and ring closure of a long chain carboxylic acid (or a derivative) with dialkyl ethylenediamine. Commercial amphoteric surfactants are derivatized by subsequent hydrolysis and ring-opening of the imidazoline ring by alkylation -- for example with chloroacetic acid or ethyl acetate. During alkylation, one or two carboxy-alkyl groups react to form a tertiary amine and an ether linkage with differing alkylating agents yielding different tertiary amines.

Long chain imidazole derivatives having application in the present disclosure generally have the general formula:

(MONO)ACETATE




(DI)PROPIONATE



Neutral pH Zwitternion

AMPHOTERIC SULFONATE



wherein R is an acyclic hydrophobic group containing from about 8 to 18 carbon atoms and M is a cation to neutralize the charge of the anion, generally sodium.

Commercially prominent imidazoline-derived amphoteric surfactants that can be employed in the present compositions include for example: Cocoamphopropionate, Cocoamphocarboxy-propionate, Cocoamphoglycinate, Cocoamphocarboxy-glycinate, Cocoamphopropyl-sulfonate, and Cocoamphocarboxy-propionic acid. Amphocarboxylic acids can be produced from fatty imidazolines in which the dicarboxylic acid functionality of the amphodicarboxylic acid is diacetic acid and/or dipropionic acid.

The carboxymethylated compounds (glycinates) described herein above frequently are called betaines. Betaines are a special class of amphoteric discussed herein below in the section entitled, Zwitterion Surfactants.

Long chain N-alkylamino acids are readily prepared by reaction  $\text{RNH}_2$ , in which  $\text{R}=\text{C}_8\text{-C}_{18}$  straight or branched chain alkyl, fatty amines with halogenated carboxylic acids. Alkylation of the primary amino groups of an amino acid leads to secondary and tertiary amines. Alkyl substituents may have additional amino groups that provide more than one reactive nitrogen center. Most commercial N-alkylamine acids are alkyl derivatives of beta-alanine or beta-N(2-carboxyethyl) alanine. Examples of commercial N-alkylamino acid ampholytes having application in this disclosure include alkyl beta-amino dipropionates,  $\text{RN}(\text{C}_2\text{H}_4\text{COOM})_2$  and  $\text{RNHC}_2\text{H}_4\text{COOM}$ . In an embodiment, R can be an acyclic hydrophobic group containing from about 8 to about 18 carbon atoms, and M is a cation to neutralize the charge of the anion.

Suitable amphoteric surfactants include those derived from coconut products such as coconut oil or coconut fatty acid. Additional suitable coconut derived surfactants include as part of their structure an ethylenediamine moiety, an alkanolamide moiety, an amino acid moiety, *e.g.*, glycine, or a combination thereof; and an aliphatic substituent of from about 8 to 18 (*e.g.*, 12) carbon atoms. Such a surfactant can also be considered an alkyl amphodicarboxylic acid. These amphoteric surfactants can include chemical


structures represented as:  $C_{12}\text{-alkyl-C(O)-NH-CH}_2\text{-CH}_2\text{-N}^+(\text{CH}_2\text{-CH}_2\text{-CO}_2\text{Na})_2\text{-CH}_2\text{-CH}_2\text{-OH}$  or  $C_{12}\text{-alkyl-C(O)-N(H)-CH}_2\text{-CH}_2\text{-N}^+(\text{CH}_2\text{-CO}_2\text{Na})_2\text{-CH}_2\text{-CH}_2\text{-OH}$ . Disodium cocoampho dipropionate is one suitable amphoteric surfactant and is commercially available under the tradename Miranol™ FBS from Rhodia Inc., Cranbury, N.J. Another suitable coconut derived amphoteric surfactant with the chemical name disodium cocoampho diacetate is sold under the tradename Mirataine™ JCHA, also from Rhodia Inc., Cranbury, N.J.

A typical listing of amphoteric classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

#### *Zwitterionic Surfactants*

Zwitterionic surfactants can be thought of as a subset of the amphoteric surfactants and can include an anionic charge. Zwitterionic surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. Typically, a zwitterionic surfactant includes a positive charged quaternary ammonium or, in some cases, a sulfonium or phosphonium ion; a negative charged carboxyl group; and an alkyl group. Zwitterionics generally contain cationic and anionic groups which ionize to a nearly equal degree in the isoelectric region of the molecule and which can develop strong "inner-salt" attraction between positive-negative charge centers. Examples of such zwitterionic synthetic surfactants include derivatives of aliphatic quaternary ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight chain or branched, and wherein one of the aliphatic substituents contains from 8 to 18 carbon atoms and one contains an anionic water solubilizing group, *e.g.*, carboxy, sulfonate, sulfate, phosphate, or phosphonate.


Betaine and sultaine surfactants are exemplary zwitterionic surfactants for use herein. A general formula for these compounds is:



wherein R<sup>1</sup> contains an alkyl, alkenyl, or hydroxyalkyl radical of from 8 to 18 carbon atoms having from 0 to 10 ethylene oxide moieties and from 0 to 1 glyceryl moiety; Y is selected from the group consisting of nitrogen, phosphorus, and sulfur atoms; R<sup>2</sup> is an alkyl or monohydroxy alkyl group containing 1 to 3 carbon atoms; x is 1 when Y is a sulfur atom and 2 when Y is a nitrogen or phosphorus atom, R<sup>3</sup> is an alkylene or hydroxy alkylene or hydroxy alkylene of from 1 to 4 carbon atoms and Z is a radical selected from the group consisting of carboxylate, sulfonate, sulfate, phosphonate, and phosphate groups.

Examples of zwitterionic surfactants having the structures listed above include: 4- [N,N-di(2-hydroxyethyl)-N-octadecylammonio]-butane-1-carboxylate; 5-[S-3-hydroxypropyl-S-hexadecylsulfonio]-3-hydroxy pentane-1-sulfate; 3-[P,P-diethyl-P-3,6,9-trioxatetracosanephosphonio]-2-hydroxypropane-1-phosphate; 3-[N,N-dipropyl-N-3-dodecoxy-2-hydroxypropyl-ammonio]-propane-1-phosphonate; 3-(N,N-dimethyl-N-hexadecylammonio)-propane-1-sulfonate; 3-(N,N-dimethyl-N-hexadecylammonio)-2-hydroxy-propane-1-sulfonate; 4-[N,N-di(2(2-hydroxyethyl)-N(2-hydroxydodecyl)ammonio)-butane-1-carboxylate; 3-[S-ethyl-S-(3-dodecoxy-2-hydroxypropyl)sulfonio]-propane-1-phosphate; 3-[P,P-dimethyl-P-dodecylphosphonio]-propane-1-phosphonate; and S[N,N-di(3-hydroxypropyl)-N-hexadecylammonio]-2-hydroxy-pentane-1-sulfate. The alkyl groups contained in said detergent surfactants can be straight or branched and saturated or unsaturated.

The zwitterionic surfactant suitable for use in the present compositions includes a betaine of the general structure:



These surfactant betaines typically do not exhibit strong cationic or anionic characters at pH extremes nor do they show reduced water solubility in their isoelectric range. Unlike "external" quaternary ammonium salts, betaines are compatible with anionics. Examples of suitable betaines include coconut acylamidopropyldimethyl betaine; hexadecyl dimethyl

betaine; C<sub>12-14</sub> acylamidopropylbetaine; C<sub>8-14</sub> acylamidohexyldiethyl betaine; 4-C<sub>14-16</sub> acylmethyldiethylammonio-1-carboxybutane; C<sub>16-18</sub> acylamidodimethylbetaine; C<sub>12-16</sub> acylamidopentanedethylbetaine; and C<sub>12-16</sub> acylmethyldimethylbetaine.

5 Sultaines useful in the present disclosure include those compounds having the formula  $(R(R^1)_2 N^+ R^2 SO_3^-)$ , in which R is a C<sub>6</sub>-C<sub>18</sub> hydrocarbyl group, each R<sup>1</sup> is typically independently C<sub>1</sub>-C<sub>3</sub> alkyl, *e.g.*, methyl, and R<sup>2</sup> is a C<sub>1</sub>-C<sub>6</sub> hydrocarbyl group, *e.g.*, a C<sub>1</sub>-C<sub>3</sub> alkylene or hydroxyalkylene group.

A typical listing of zwitterionic classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975. Further  
10 examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

#### *Defoaming Agent*

15 A defoaming agent for reducing the stability of foam may also be included in the warewashing composition. Examples of defoaming agents include, but are not limited to: ethylene oxide/propylene block copolymers such as those available under the name Pluronic N-3; silicone compounds such as silica dispersed in polydimethylsiloxane, polydimethylsiloxane, and functionalized polydimethylsiloxane  
20 such as those available under the name Abil B9952; fatty amides, hydrocarbon waxes, fatty acids, fatty esters, fatty alcohols, fatty acid soaps, ethoxylates, mineral oils, polyethylene glycol esters, and alkyl phosphate esters such as monostearyl phosphate. A discussion of defoaming agents may be found, for example, in U.S. Patent No. 3,048,548 to Martin et al., U.S. Patent No. 3,334,147 to Brunelle et al., and U.S.  
25 Patent No. 3,442,242 to Rue et al.

When the concentrate includes a defoaming agent, the defoaming agent can be provided in an amount of between approximately 0.0001% and approximately 10% by weight, between approximately 0.001% and approximately 5% by weight, or between approximately 0.01% and approximately 1.0% by weight.

30

*Concentrate and Use Solutions for Methods of Use*

The detergent compositions as provided in a block are concentrate compositions. In general, a concentrate refers to a composition that is intended to be diluted with water to provide a use solution that contacts an object to provide the desired cleaning, rinsing, or the like.

A use solution may be prepared from the concentrate by diluting the concentrate with water at a dilution ratio that provides a use solution having desired deterative properties. The water that is used to dilute the concentrate to form the use composition can be referred to as water of dilution or a diluent, and can vary from one location to another. The typical dilution factor is between approximately 1 and approximately 10,000 but will depend on factors including water hardness, the amount of soil to be removed and the like. A concentrate may be diluted at a ratio of between about 1:10 and about 1:10,000 concentrate to water. Particularly, A concentrate is diluted at a ratio of between about 1:100 and about 1:5,000 concentrate to water. More particularly, a concentrate may be diluted at a ratio of between about 1:250 and about 1:2,000 concentrate to water.

In an aspect of the disclosure, a use solution of the detergent compositions has between about 10 ppm to about 6000 ppm alkaline source. In a preferred aspect of the disclosure, a use solution of the detergent composition has between about 500 ppm to about 4000 ppm alkaline source. In a still further preferred aspect of the disclosure, a use solution of the detergent composition has between 2500 ppm to about 3500 ppm alkaline source. In addition, without being limited according to the disclosure, all ranges recited are inclusive of the numbers defining the range and include each integer within the defined range.

In an aspect of the disclosure, the detergent composition preferably provides efficacious cleaning at low use dilutions, *e.g.*, require less volume to clean effectively. In an aspect, the detergent composition may be diluted in water prior to use at dilutions ranging from about 1/16 oz./gal. to about 2 oz./gal. or more. A detergent composition that requires less volume to achieve the same or better cleaning efficacy and provides hardness scale control and/or other benefits at low use dilutions is desirable.

In some aspects, the detergent compositions are contacted by a diluent, such as water to generate a concentrate and/or use solution for the various applications of use.

In some aspects, the present disclosure provides methods for removing soils from a surface, *e.g.*, a hard surface, and/or bleaching a surface. In some embodiments, the method comprises contacting a use solution of the detergent compositions with a surface, and removing the composition from the surface after an amount of time sufficient to facilitate  
5 soil removal and/or bleaching. The contacting step can last for any suitable time. In some embodiments, the contacting step lasts for at least 10 seconds, 20 seconds, 30 seconds, 40 seconds, 50 seconds, 1 minute, 10 minutes, 30 minutes, 1 hour, 2 hours, 4 hours, 8 hours, 16 hours, 1 day, 3 days, 1 week, or longer. The detergent composition can contact the surface (or target for soil removal and/or bleaching) in any suitable manner. In some  
10 embodiments, the detergent composition is applied by means of a spray, a foam, soaking or the like.

The methods can be used to achieve any suitable removal of soil (*e.g.*, cleaning), sanitizing, disinfecting, bleaching and/or reduction of the microbial population in and/or on the surface or target. In some embodiments, the methods can be used to reduce the  
15 microbial population by at least one log<sub>10</sub>. In other embodiments, the present methods can be used to reduce the microbial population in and/or on the target or the treated target composition by at least two log<sub>10</sub>. In still other embodiments, the present methods can be used to reduce the microbial population in and/or on the target or the treated target composition by at least three log<sub>10</sub>.

20 In some embodiments, the method further comprises rinsing the surface. In some embodiments, the method further comprises a mechanical application of force, agitation and/or pressure to assist in removing the soils and/or bleaching the surface.

The methods of the present disclosure can be used to remove a variety of soils from a variety of surfaces and/or bleaching a variety of surfaces. For example, surfaces suitable  
25 for cleaning using the methods of the present disclosure include, but are not limited to, walls, floors, ware, dishes, flatware, pots and pans, heat exchange coils, ovens, fryers, smoke houses, sewer drain lines, and the like.

In some embodiments, the methods of the present disclosure are followed by only a rinse step. In other embodiments, the methods of the present disclosure are followed by a  
30 conventional CIP method suitable for the surface to be cleaned. In still yet other embodiments, the methods of the present disclosure are followed by a CIP method such as those described in U.S. Patent Nos. 8,398,781 and 8,114,222 entitled "Methods for

Cleaning Industrial Equipment with Pre-treatment," .


*Methods of Use*

5            In another aspect, disclosed here is a method of cleaning, sanitizing and/or bleaching comprising generating a use solution of the disclosed compositions that comprise an alkaline source, an enzyme, and a specific type of phosphonate or amine phosphonate salt.


10           In yet another aspect, disclosed here is a method of cleaning, sanitizing and/or bleaching comprising generating a use solution of the disclosed compositions that comprise an alkaline source, an enzyme, and a specific type of phosphonate or amine phosphonate salt, and contacting a surface or object in need of cleaning and sanitizing with the use solution.

15           In some embodiments, the use solution of the disclosed detergent compositions has maintained at least 15% of its enzyme activity after 240 minutes of its generation. In some other embodiments, the use solution of the disclosed detergent composition has maintained at least 20% of its enzyme activity after 120 minutes of its generation.

             In yet another aspect, the disclosure is a method of stabilizing an enzyme in a solid



20           detergent. The method comprises adding a phosphonate of formula or salt thereof, or an amine phosphonate salt in an existing detergent composition containing an enzyme, wherein R<sup>10</sup> and R<sup>11</sup> are independently hydrogen, a substituted carboxylic acid, phosphonate, ethanol, diglyco, substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, or phosphonate-methyl; the amine phosphonate salt is a product of a phosphonate of formula



and an amine, and R<sup>12</sup>, R<sup>13</sup>, and R<sup>14</sup> are independently hydroxyl, methyl, -PO(OH)<sub>2</sub>, -CH<sub>2</sub>COOH, a substituted alkyl, phosphonate, ester thereof, salt thereof, or derivative thereof.

5 All publications and patent applications in this specification are indicative of the level of ordinary skill in the art to which this disclosure pertains.

## 10 **EXAMPLES**

Embodiments of the present disclosure are further defined in the following non-limiting Examples. It should be understood that these Examples, while indicating certain embodiments of the disclosure, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential  
 15 characteristics of this disclosure, and without departing from the spirit and scope thereof, can make various changes and modifications of the embodiments of the disclosure to adapt it to various usages and conditions. Thus, various modifications of the embodiments of the disclosure, in addition to those shown and described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to  
 20 fall within the scope of the appended claims.

The following materials are used in the Examples:

Bio-terge® AS-90 - 90% active C<sub>14</sub>-C<sub>16</sub> alpha olefin sulfonate (AOS);

Ufaryl DL90C - 90% C<sub>10</sub>-C<sub>13</sub> active linear alkylbenzene sulfonate (LAS), drum dried powder;

25 Belclene 200 - 50% active 500-100 MW polymaleic acid;

Acusol™ 445N - 45% active polyacrylic acid (4500 – 10000 MW);

Acusol™ 445ND - 45% active polyacrylic acid (4500 – 10,000MW), spray dried;

- Acusol™ 820 - a Hydrophobically modified Alkali Soluble acrylic polymer Emulsion (HASE) with unusually high aqueous thickening and stabilising efficiency;
- Acusol™ 929 - 46% active polyacrylic acid (~10,000 – 15,000 MW);
- Dense Ash – Sodium Carbonate;
- 5 Light Ash – Sodium Carbonate;
- Sodium Bicarbonate, granular;
- PEG 8000 – Polyethylene glycol with an average molecular weight of 8,000;
- Powder Bicarb – Sodium bicarbonate, in powder;
- CMC-7LT – carboxymethylcellulose;
- 10 LAE 24-7 - Linear alcohol ethoxylate (7 moles EO);
- ATMP - Aminotri (methylene phosphonic acid);
- AMPA – Aminomethyl phosphonic acid;
- PBTC - Phosphonobutane tricarboxylic acid; Bayhibit AM;
- STPP - sodium tripolyphosphate;
- 15 HEDP - 1-hydroxy ethylidene-1, 1-diphosphonic acid, Dequest 2010;
- DGAP – Diglycolamine phosphonate, Scale inhibitor 2588;
- MEAP – Monoethanolamine phosphonate, Scale inhibitor 2670;
- PAPEMP – Polyamino Polyether Methylene Phosphonic Acid, Kemguard 8010;
- 20 PSO – Phosphinosuccinic Mix from Nalco, Nalco TX15712SQ.

#### EXAMPLE 1

- The enzyme activities in the detergents with different phosphonate levels were tested at 4000 ppm in 5 grains per gallon (GPG) water at 120 °F with various time points collected up to 4 hours, after a use solution was generated from the detergent composition.
- 25 During the tests, the samples at different time points were collected and immediately frozen in an acetone/dry ice bath and stored in -80 °C before the activity of the enzyme was evaluated. For the use of such an assay, t=0 min was the reference point for 100% enzyme activity.

- 30 Assays of enzyme activity in formulations (% retention) were conducted to simulate a presoak condition in a beaker using the chemistry, temperature, and pH conditions relevant to manual warewash or presoak applications. Enzyme activity is an

indicator of the stability of the enzyme in the detergent, specifically in an aqueous use solution within a sump (which is under conditions of high pH, temperature and dilution).

The analysis by protease assay was conducted as follows. For the assays, a detergent composition was used to generate an aqueous use solution evaluated herein. The components in the tested detergent compositions are listed in **Table 1**.

Enzyme activity under presoak or manual warewash conditions was traced quantitatively using a standard protease assay. Samples were prepared under bench top conditions, whereby the use solution from a detergent composition or detergent was obtained and maintained at a presoak or manual warewash temperature in a stirring water bath. After the time course for assessing enzyme stability is initiated, aliquots were taken at various time points and flash-frozen. A time = 0 sample was prepared for each series by dissolving the detergent formulation at room temperature, mixing thoroughly, and flash freezing. Samples were thawed and diluted as necessary in an assay buffer usually for use in the protease assay. A glycine buffer at pH 9.0 is used here. The assay monitored the direct reaction of the protease on a small, commercially available peptidyl substrate, with liberation of the product providing correlation to the active enzyme content. The product was detected using a plate reader with an appreciable dynamic range (upper absorbance limit of the instrument >3.5). Enzyme activity levels were assessed relative to a calibration curve with average values for replicate tests used to map protease stability under presoak or manual warewash use conditions. Enzyme retention at each time point was calculated as the % enzyme activity relative to the time = 0 sample.

The analysis by lipase and amylase assay was conducted similarly, except with a different substrate and buffers. For lipase activity, the substrate is p-nitrophenyl valerate, and for amylase, the substrate is an ethylidene substrate (EPS). The buffer used in lipase assay is TRIS (Tris(hydroxymethyl)aminomethane) buffer at pH 8.0, and in amylase assay HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) buffer at pH 8.0.

The phosphonates and their concentration levels evaluated are listed in **Table 2**. The phosphonate structures and their overall effects on enzyme activity are listed in **Table 3**. The relative enzyme activities in a use solution of the tested detergent compositions containing a different phosphonate at different concentrations over a period time are listed in **Table 4A – Table 4E, Table 5A – Table 5E, and Table 6A – Table 6F**.

**Table 1.** List of the Components and their weight percentage in the Tested Detergent Compositions

Raw Material	Base Formula <sup>a</sup>	All Ash Formula <sup>a</sup>
Sodium carbonate	47.47 (35-55)	85.09 (50-90)
Sodium bicarbonate	37.62 (30-45)	
Alpha olefin sulfonate	4 (0.1-10)	4 (0.1-10)
Enzyme	1.3 (0.1-5)	1.3 (0.1-5)
Water Conditioning Agents	7.61 (2-15)	7.61 (2-15)
Triethanolamine	Added to neutralize HEDP and PBTC (see results)	
Surfactant	2 (0.1-5)	2 (0.1-5)







<sup>a</sup>The values in brackets are exemplary preferred ranges for the respective ingredient in the tested compositions.



**Table 2.** List of Phosphonates and their Concentration Levels that were Evaluated for Their Effect on Enzyme Activity.

Name	Levels tested
ATMP – aminotris (methylenephosphonic acid)	0.3% active 0.6% active 0.1% elemental phosphorus
PBTC – Phosphono butanetricarboxylic acid	0.3% active 0.6% active 0.1% elemental phosphorus 0.3% active neutralized with triethanolamine
HEDP – Etidronic acid	0.3% active 0.6% active 0.1% elemental phosphorus 0.3% active neutralized with triethanolamine
DGAP – Diglycolamine phosphonate	0.3% active 0.6% active 0.1% elemental phosphorus
MEAP – Monoethanolamine phosphonate	0.3% active 0.6% active 0.1% elemental phosphorus
PAPEMP – Polyamino Polyether Methylene Phosphonic Acid	0.3% active 0.6% active 0.1% elemental phosphorus
PSO	0.3% active 0.6% active 0.1% elemental phosphorus

AMPA – Aminomethyl phosphoric acid	0.3% active 0.6% active 0.1% elemental phosphorus
------------------------------------	---

**Table 3.** List of Phosponates, Their Structures, and Effects on Enzyme Activity

Name	Structure	Stability
ATMP		Yes
PBTC		No
HEDP		No
DGAP		Yes
PAPEMP	$\begin{array}{c} \text{H}_2\text{O}_3\text{P}-\text{CH}_2 \quad \text{CH}_3 \quad \text{CH}_3 \quad \text{CH}_2-\text{PO}_3\text{H}_2 \\   \quad \quad \quad   \quad \quad \quad   \\ \text{N}-\text{CH}_2-\text{CH}-(\text{O}-\text{CH}_2-\text{CH})_n-\text{N} \\   \quad \quad \quad   \\ \text{H}_2\text{O}_3\text{P}-\text{CH}_2 \quad \quad \quad \text{CH}_2-\text{PO}_3\text{H}_2 \end{array}$	Yes
PSO (mix)		No
MEAP		Yes

AMPA		Yes
Triethanolamine		Yes, when neutralized with a phosphonate

**Table 4A.** Relative Protease Enzyme Activity in a Use Solution of the Detergent Compositions Containing a Protease and Various Phosphonates at a Level of 0.3 wt-% at 120 °F.

time (min)	No Phosphorous	ATMP	PBTC	HEDP
Blank	-2.79%	-1.41%	-0.82%	-0.90%
0	43.38%	100.00%	100.00%	100.00%
5	100.00%	94.95%	52.04%	52.21%
10	75.23%	94.68%	30.98%	32.95%
20	53.12%	89.76%	14.75%	15.91%
40	34.98%	87.06%	7.33%	8.68%
60	24.31%	83.93%	5.15%	5.52%
90	19.75%	80.79%	4.01%	4.10%
120	13.39%	76.31%	2.91%	3.10%
150	12.48%	73.96%	2.21%	2.39%
180	12.46%	70.37%	1.52%	1.65%
210	10.72%	65.72%	1.01%	1.36%
240	9.69%	63.75%	0.63%	1.20%

5

**Table 4A - Continued**

DGAP	MEAP	PAPEMP	PSO	AMPA
-0.34%	-0.66%	-0.07%	-0.19%	-1.80%
100.00%	100.00%	100.00%	100.00%	85.83%
100.00%	103.02%	100.00%	52.04%	100.00%
98.30%	98.29%	107.31%	36.88%	86.30%
92.44%	96.59%	109.98%	19.71%	78.58%
91.75%	90.19%	105.51%	10.45%	63.30%
89.16%	82.48%	100.17%	7.19%	51.37%
79.16%	79.14%	89.50%	5.10%	44.31%
84.87%	72.58%	96.21%	3.55%	37.90%

82.96%	68.59%	99.48%	3.35%	31.17%
82.44%	63.56%	93.74%	3.21%	28.70%
75.52%	58.70%	86.60%	2.75%	25.64%
76.14%	60.60%	92.79%	2.27%	22.72%

**Table 4B.** Relative Protease Enzyme Activity in a Use Solution of the Detergent Compositions Containing a Protease and Various Phosphonates at a Level of 0.6 wt-% at 120 °F.

time (min)	ATMP	PBTC	HEDP	DGAP
Blank	0.31%	0.05%	-2.35%	-1.75%
0	100.00%	100.00%	100.00%	100.00%
5	91.70%	60.27%	69.69%	105.28%
10	102.58%	35.81%	51.90%	100.03%
20	100.03%	16.27%	27.67%	101.24%
40	97.16%	7.44%	12.41%	100.88%
60	92.31%	4.84%	7.62%	---
90	89.99%	2.96%	4.70%	98.76%
120	91.60%	1.76%	2.94%	93.04%
150	86.21%	1.32%	1.79%	88.27%
180	89.08%	0.91%	0.92%	86.23%
210	89.86%	0.70%	0.71%	88.24%
240	82.33%	0.39%	0.45%	89.57%

5 **Table 4B - Continued**

MEAP	PAPEMP	PSO	AMPA
-0.43%	0.48%	0.16%	-2.15%
100.00%	100.00%	100.00%	100.00%
106.85%	98.49%	59.03%	109.27%
101.06%	96.69%	34.62%	94.52%
104.08%	93.71%	14.30%	84.97%
95.75%	92.60%	6.47%	77.92%
95.13%	92.60%	4.30%	75.86%
86.12%	86.28%	3.15%	62.50%
85.48%	88.30%	1.60%	54.43%
80.22%	89.66%	1.15%	51.19%
82.71%	86.90%	0.86%	47.47%
70.64%	87.49%	0.70%	43.42%
72.77%	86.53%	0.74%	40.00%

**Table 4C.** Relative Protease Enzyme Activity in a Use Solution of the Detergent Compositions Containing a Protease and Various Phosphonates at a Level of 0.1% elemental phosphorus at 120 °F.

time (min)	ATMP	PBTC	HEDP	DGAP
Blank	-0.33%	-0.15%	-0.67%	-0.51%
0	100.00%	100.00%	100.00%	100.00%
5	122.55%	86.68%	77.74%	98.29%
10	116.65%	55.40%	53.71%	103.47%
20	115.14%	23.91%	23.58%	101.47%
40	107.73%	8.20%	9.12%	88.60%
60	102.70%	4.68%	5.36%	86.17%
90	111.18%	2.99%	3.76%	85.84%
120	106.19%	2.04%	2.44%	83.06%
150	96.80%	1.54%	1.45%	81.53%
180	93.35%	1.01%	0.74%	76.41%
210	90.51%	0.73%	0.64%	74.23%
240	88.60%	0.42%	0.45%	72.01%

5 **Table 4C - Continued**

MEAP	PAPEMP	PSO	AMPA
-0.97%	-1.75%	-1.49%	-1.85%
100.00%	100.00%	100.00%	107.11%
119.22%	95.74%	53.94%	100.00%
118.42%	95.47%	28.39%	99.75%
113.68%	92.78%	10.95%	84.62%
108.32%	86.55%	2.62%	70.58%
105.97%	87.49%	1.03%	62.19%
95.99%	86.61%	0.29%	49.17%
91.30%	86.34%	0.76%	43.34%
82.74%	84.01%	1.05%	38.26%
81.94%	71.92%	1.20%	34.55%
79.85%	80.74%	1.32%	32.27%
75.73%	79.08%	1.52%	28.25%

**Table 4D.** Relative Protease Enzyme Activity in a Use Solution of the Detergent Compositions Containing a Protease and Various Phosphonates Neutralized with an Alkanolamine at a Level of 0.3 wt-% at 120 °F.

PBTC neutralized with TEA	HEDP neutralized with TEA	PSO neutralized with TEA	PBTC neutralized with MEA	HEDP neutralized with MEA
-0.84%	-0.67%	-0.89%	-2.17%	-3.15%
100.00%	100.00%	100.00%	100.00%	100.00%
100.00%	100.00%	97.66%	87.92%	91.36%
89.46%	92.11%	76.32%	65.16%	74.15%
73.58%	77.81%	67.03%	48.64%	60.15%
52.71%	55.84%	48.91%	36.58%	41.89%
39.31%	43.76%	38.11%	27.37%	32.68%
28.74%	30.12%	28.75%	20.02%	24.63%
23.33%	22.99%	24.12%	14.33%	19.44%
16.83%	19.37%	20.99%	10.42%	15.29%
15.04%	17.13%	17.03%	12.17%	13.78%
13.21%	14.12%	15.22%	10.85%	11.80%
10.57%	13.15%	13.69%	9.37%	11.33%

Table 4D - Continued

PSO neutralized with MEA	PBTC neutralized with IPA	HEDP neutralized with IPA	PSO neutralized with IPA
-3.24%	-2.66%	-3.00%	-2.13%
100.00%	100.00%	100.00%	100.00%
91.94%	73.74%	102.03%	94.42%
70.31%	52.09%	85.18%	68.24%
48.71%	38.16%	65.55%	45.69%
32.24%	31.66%	48.32%	37.11%
23.13%	25.45%	38.50%	26.83%
17.02%	18.30%	28.67%	19.73%
13.58%	13.92%	22.10%	15.94%
11.49%	12.46%	16.09%	13.39%
10.56%	10.39%	16.19%	11.79%
8.01%	8.14%	14.38%	8.28%
7.33%	7.14%	12.80%	7.09%

- 5 **Table 4E.** Relative Protease Enzyme Activity in a Use Solution of the All Ash Detergent Compositions Containing a Protease and Various Phosphonates at a Level of 0.3 wt-% at 120 °F.

time (min)	ATMP	DGAP	PBTC	HEDP
Blank	1.16%	0.80%	-1.26%	-1.19%
0	100.00%	100.00%	100.00%	100.00%
5	84.60%	81.79%	11.99%	8.63%
10	75.67%	75.39%	3.18%	3.25%
20	53.32%	59.47%	-0.81%	0.61%
40	33.78%	39.98%	-1.79%	-1.32%
60	22.22%	27.49%	-1.94%	-1.66%
90	15.40%	18.95%	-1.57%	-0.97%
120	11.13%	12.48%	-1.11%	-1.23%
150	9.39%	10.56%	-1.04%	-0.85%
180	7.41%	9.69%	-2.62%	-0.93%
210	6.17%	7.02%	-3.07%	-0.42%
240	5.58%	6.34%	-2.02%	-0.42%

**Table 5A.** Relative Amylase Enzyme Activity in a Use Solution of the Detergent Compositions Containing a Protease and Various Phosphonates at a Level of 0.3 wt-% at 120 °F.

time (min)	No Phosphorous	ATMP	PBTC	HEDP	DGAP
Blank	-4.42%	-1.41%	-0.83%	-0.92%	-5.35%
0	26.50%	100.00%	100.00%	100.00%	39.01%
5	100.00%	89.47%	96.90%	81.99%	100.00%
10	96.04%	82.94%	94.03%	90.32%	113.48%
20	92.60%	79.67%	86.69%	95.23%	112.07%
40	94.11%	90.79%	83.68%	84.92%	110.36%
60	100.70%	89.94%	86.51%	72.27%	109.99%
90	90.23%	94.47%	84.89%	79.95%	118.04%
120	93.38%	95.08%	85.85%	84.46%	110.39%
150	93.75%	94.00%	76.25%	79.54%	111.92%
180	99.53%	90.87%	81.01%	84.18%	112.24%
210	102.48%	94.61%	81.99%	69.49%	115.96%
240	96.48%	96.54%	81.88%	88.63%	110.74%

**Table 5A - Continued**

MEAP	PAPEMP	PSO	AMPA
4.36%	-4.91%	3.80%	-0.50%
100.00%	45.92%	100.00%	82.13%

100.65%	100.00%	92.97%	100.00%
102.68%	101.33%	83.56%	95.55%
102.07%	101.28%	90.12%	101.29%
91.36%	89.91%	91.75%	80.17%
93.57%	93.07%	91.41%	80.10%
94.85%	89.58%	90.27%	87.43%
102.84%	89.74%	90.00%	92.65%
93.01%	92.85%	91.32%	93.49%
96.46%	87.45%	88.35%	102.37%
100.65%	93.95%	78.96%	103.78%
103.40%	108.16%	86.96%	102.93%

**Table 5B.** Relative Amylase Enzyme Activity in a Use Solution of the Detergent Compositions Containing a Protease and Various Phosphonates at a Level of 0.6 wt-% at 120 °F.

time (min)	ATMP	PBTC	HEDP	DGAP
Blank	0.13%	-5.98%	-1.11%	-0.53%
0	100.00%	100.00%	100.00%	100.00%
5	87.10%	107.86%	96.09%	102.43%
10	92.19%	111.05%	99.71%	95.89%
20	80.12%	106.52%	91.20%	99.40%
40	77.64%	73.11%	94.89%	97.26%
60	91.13%	79.10%	97.28%	100.23%
90	92.19%	71.11%	94.67%	95.37%
120	93.02%	64.98%	88.12%	102.17%
150	101.54%	74.97%	90.33%	102.37%
180	80.83%	85.22%	86.96%	101.89%
210	92.78%	94.67%	96.92%	102.69%
240	86.75%	97.07%	92.50%	105.89%

**Table 5B - Continued**

MEAP	PAPEMP	PSO	AMPA
-1.01%	-4.64%	-5.39%	-3.26%
100.00%	100.00%	100.00%	100.00%
95.91%	91.71%	98.30%	91.80%
99.98%	99.05%	97.66%	92.54%
90.83%	94.55%	94.69%	85.05%
91.75%	97.22%	84.89%	94.06%
97.99%	97.94%	78.56%	86.14%
99.08%	101.28%	83.38%	90.32%
97.36%	99.25%	84.70%	93.67%

101.28%	105.34%	79.34%	92.05%
94.70%	98.44%	89.95%	81.43%
99.20%	101.64%	88.55%	94.06%
99.15%	96.02%	94.02%	81.87%

**Table 5C.** Relative Amylase Enzyme Activity in a Use Solution of the Detergent Compositions Containing a Protease and Various Phosphonates at a 0.1% elemental phosphorus level at 120 °F.

time (min)	ATMP	PBTC	HEDP	DGAP
Blank	-1.95%	-1.50%	-3.71%	-4.45%
0	82.19%	90.07%	---	100.00%
5	100.00%	100.00%	100.00%	98.51%
10	107.32%	98.00%	97.58%	108.49%
20	106.35%	89.44%	90.30%	---
40	100.35%	90.60%	81.48%	97.68%
60	106.09%	86.05%	83.84%	97.62%
90	101.79%	90.47%	85.26%	99.36%
120	104.38%	89.11%	82.59%	92.67%
150	101.75%	88.36%	---	98.24%
180	103.89%	82.80%	75.47%	93.89%
210	100.38%	84.45%	85.77%	107.33%
240	107.10%	91.39%	81.23%	103.75%

**Table 5C - Continued**

MEAP	PAPEMP	PSO	AMPA
5.88%	-3.90%	-4.72%	-3.90%
100.00%	100.00%	100.00%	97.43%
95.47%	85.10%	87.63%	100.00%
112.93%	95.50%	86.00%	102.34%
117.67%	96.02%	82.60%	105.43%
122.41%	91.46%	68.15%	92.18%
125.44%	97.42%	72.63%	92.23%
127.99%	89.60%	71.93%	96.88%
123.91%	99.26%	65.82%	92.96%
120.88%	101.88%	69.91%	103.38%
113.29%	91.85%	75.23%	102.05%
112.78%	99.00%	74.61%	91.32%
118.72%	96.59%	76.21%	99.25%

**Table 5D.** Relative Amylase Enzyme Activity in a Use Solution of the Detergent Compositions Containing a Protease and Various Phosphonates Neutralized with an Alkanolamine at a Level of 0.3 wt-% at 120 °F.

PBTC neutralized with TEA	HEDP neutralized with TEA	PSO neutralized with TEA	PBTC neutralized with MEA	HEDP neutralized with MEA
1.45%	1.09%	6.53%	1.66%	3.10%
78.26%	97.09%	100.00%	100.00%	88.65%
100.00%	100.00%	79.64%	74.46%	100.00%
100.23%	102.11%	90.67%	76.22%	95.28%
98.16%	97.01%	99.47%	74.00%	103.84%
95.30%	92.21%	107.22%	74.18%	89.21%
95.53%	92.69%	102.80%	82.04%	84.15%
100.38%	97.65%	116.70%	83.37%	84.46%
100.48%	97.26%	117.67%	84.41%	90.08%
97.55%	98.42%	114.79%	93.20%	97.93%
95.18%	100.68%	98.22%	74.68%	95.62%
96.12%	102.40%	92.27%	78.79%	97.86%
94.33%	101.82%	98.73%	74.22%	108.86%

**Table 5D - Continued**

PSO neutralized with MEA	PBTC neutralized with IPA	HEDP neutralized with IPA	PSO neutralized with IPA
3.18%	-4.90%	-0.52%	-0.27%
78.22%	100.00%	95.84%	91.22%
100.00%	85.98%	100.00%	100.00%
99.59%	88.34%	96.93%	102.42%
100.02%	84.63%	95.39%	100.25%
81.36%	82.64%	90.50%	102.59%
83.13%	86.12%	89.15%	101.65%
84.18%	84.53%	97.78%	105.89%
85.55%	86.52%	93.19%	109.47%
98.17%	90.97%	94.21%	107.16%
98.34%	86.45%	96.62%	100.22%
103.65%	89.86%	98.72%	103.00%
99.44%	91.26%	109.38%	107.60%

**Table 5E.** Relative Amylase Enzyme Activity in a Use Solution of the All Ash Detergent Compositions Containing a Protease and Various Phosphonates at a Level of 0.3 wt-% at 120 °F.

time (min)	ATMP	DGAP	PBTC	HEDP
Blank	-2.61%	0.85%	0.39%	1.29%
0	98.49%	100.00%	98.88%	92.90%
5	100.00%	89.94%	100.00%	100.00%
10	98.40%	87.50%	94.76%	97.04%
20	97.90%	88.87%	90.04%	87.85%
40	91.03%	86.87%	63.67%	70.85%
60	86.91%	82.06%	58.65%	55.60%
90	87.68%	81.79%	54.03%	63.30%
120	80.93%	75.74%	45.38%	58.18%
150	83.80%	79.71%	44.40%	54.50%
180	80.91%	65.75%	41.15%	45.78%
210	84.34%	71.27%	41.56%	44.79%
240	83.21%	70.12%	40.40%	41.67%

**Table 6A.** Relative Lipase Enzyme Activity in a Use Solution of the Detergent Composition Containing a Lipase and Various Phosphonates at a Level of 0.3 wt-% at 120 °F.

time (min)	No Phosphorous	ATMP	PBTC	HEDP	DGAP	MEAP	PAPEMP	PSO
Blank	-7.58%	-5.56%	-8.69%	-2.91%	-4.86%	-3.99%	-3.64%	-7.44%
0	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	49.99%	48.83%
5	72.61%	79.77%	73.63%	47.15%	83.54%	84.99%	100.00%	100.00%
10	76.29%	71.10%	66.55%	44.35%	88.25%	78.55%	92.85%	84.45%
20	62.07%	62.47%	48.22%	34.94%	77.15%	58.36%	77.81%	64.78%
40	38.96%	39.28%	26.23%	23.40%	62.99%	39.07%	54.24%	29.73%
60	25.66%	34.01%	15.51%	17.75%	52.84%	31.84%	51.08%	19.72%
90	14.12%	26.70%	7.72%	11.06%	43.46%	26.97%	45.80%	11.86%
120	8.54%	22.13%	3.90%	6.92%	35.40%	22.23%	36.29%	6.82%
150	5.72%	19.07%	2.70%	5.38%	31.10%	20.19%	33.14%	5.20%
180	-0.84%	17.48%	-0.75%	2.90%	22.42%	19.50%	28.25%	3.31%

210	1.12%	15.49%	-1.86%	2.62%	24.78%	16.46%	26.56%	2.38%
240	0.88%	14.68%	-2.20%	2.32%	22.08%	15.76%	24.03%	2.08%

**Table 6B.** Relative Lipase Enzyme Activity in a Use Solution of the Detergent Composition Containing a Lipase and Various Phosphonates at a Level of 0.6 wt-% at 120 °F.

time (min)	ATMP	PBTC	HEDP	DGAP
Blank	-2.73%	-2.06%	-1.35%	-2.99%
0	100.00%	100.00%	100.00%	100.00%
5	93.26%	61.52%	62.99%	95.36%
10	87.80%	62.20%	59.92%	100.58%
20	81.36%	44.99%	48.34%	94.94%
40	60.03%	31.89%	33.30%	93.85%
60	56.12%	23.30%	23.22%	86.18%
90	49.58%	16.36%	17.91%	76.99%
120	42.20%	12.11%	14.72%	71.87%
150	37.91%	9.68%	12.63%	72.77%
180	33.77%	6.06%	7.01%	64.17%
210	32.19%	6.80%	8.08%	63.68%
240	29.80%	5.96%	5.20%	56.85%

5

**Table 6B - Continued**

MEAP	PAPEMP	PSO	AMPA
-2.42%	-3.29%	-3.68%	-1.32%
100.00%	100.00%	100.00%	100.00%
95.37%	86.04%	67.75%	80.17%
90.69%	83.42%	48.43%	64.48%
83.63%	73.22%	32.89%	49.45%
72.89%	64.86%	14.07%	37.77%
69.90%	59.26%	10.48%	26.82%
65.92%	47.33%	8.07%	19.59%
59.18%	42.05%	5.40%	15.84%
53.85%	38.58%	4.42%	15.01%
51.72%	29.04%	2.99%	13.23%
47.03%	31.37%	5.01%	12.05%
42.96%	29.77%	2.12%	11.16%

**Table 6C.** Relative Lipase Enzyme Activity in a Use Solution of the Detergent Composition Containing a Lipase and Various Phosphonates at a Level of 0.1% elemental phosphorus at 120 °F or at room temperature.

time (min)	ATMP	PBTC	HEDP	DGAP
Blank	-3.96%	-1.76%	-1.60%	-3.52%
0	100.00%	100.00%	100.00%	100.00%
5	82.14%	87.75%	79.12%	88.13%
10	78.45%	73.69%	67.70%	80.42%
20	64.87%	56.53%	51.16%	74.76%
40	45.84%	25.32%	25.69%	49.43%
60	37.66%	19.97%	19.99%	46.05%
90	30.16%	15.95%	16.10%	39.53%
120	25.91%	12.52%	13.04%	34.69%
150	21.51%	11.11%	11.14%	28.57%
180	15.43%	10.01%	9.12%	25.49%
210	17.94%	9.16%	7.99%	24.18%
240	15.64%	9.12%	7.70%	22.78%

5 **Table 6C - Continued**

MEAP	PAPEMP	PSO	AMPA
-3.13%	-2.56%	-1.04%	-3.14%
100.00%	100.00%	100.00%	100.00%
81.36%	91.09%	58.47%	79.03%
84.98%	83.66%	52.95%	77.51%
71.62%	75.51%	43.79%	62.55%
57.03%	58.49%	27.97%	53.26%
47.80%	54.87%	18.63%	44.19%
41.02%	48.29%	13.24%	33.99%
31.46%	42.46%	11.36%	28.67%
28.32%	38.07%	9.72%	23.85%
20.93%	33.57%	3.32%	16.98%
23.04%	31.15%	5.29%	17.74%
20.87%	31.01%	4.75%	16.10%

**Table 6D.** Relative Lipase Enzyme Activity in a Use Solution of the Detergent Composition Containing a Lipase and Various Phosphonates Neutralize With a Alkanolamine at a Level of 0.3 wt at 120 °F or at room temperature.

PBTC neutralized with TEA	HEDP neutralized with TEA	PSO neutralized with TEA	PBTC neutralized with MEA	HEDP neutralized with MEA
-1.43%	-1.96%	-5.25%	-0.97%	-5.48%
100.00%	100.00%	100.00%	100.00%	100.00%
65.46%	68.45%	75.32%	48.91%	74.09%
64.43%	54.84%	71.42%	46.95%	64.35%
47.72%	42.36%	53.53%	34.80%	46.99%
31.53%	22.75%	35.00%	20.43%	24.45%
23.15%	17.11%	24.95%	13.60%	16.43%
16.58%	13.43%	17.45%	10.75%	12.53%
12.91%	10.58%	13.62%	8.59%	9.01%
10.68%	8.60%	11.19%	7.84%	6.81%
6.55%	7.53%	6.65%	2.43%	5.52%
6.64%	6.32%	6.99%	3.12%	4.36%
6.49%	8.02%	6.52%	3.04%	3.98%

**Table 6D - Continued**

PSO neutralized with MEA	PBTC neutralized with IPA	HEDP neutralized with IPA	PSO neutralized with IPA
-5.72%	1.00%	1.04%	-1.11%
100.00%	100.00%	100.00%	100.00%
68.36%	81.40%	--	69.16%
51.37%	64.90%	52.89%	62.12%
41.11%	51.13%	46.32%	43.17%
21.86%	--	32.93%	18.78%
15.13%	--	25.04%	13.42%
8.55%	14.75%	18.40%	10.38%
7.00%	12.55%	14.58%	8.29%
4.90%	11.70%	14.58%	7.65%
4.01%	10.22%	--	7.05%
1.16%	10.12%	6.03%	6.56%
0.64%	9.27%	7.48%	6.44%

- 5 **Table 6E.** Relative Lipase Enzyme Activity in a Use Solution of the Detergent Composition Containing a Lipase and Various Phosphonates at a Level of 0.3 wt-% at room temperature.

time (min)	PBTC at RT	HEDP at RT	PSO at RT	PBTC neutralized with TEA at RT	HEDP neutralized with TEA at RT
Blank	-2.78%	-2.51%	1.88%	6.08%	4.65%
0	100.00%	100.00%	100.00%	100.00%	100.00%
5	84.82%	101.22%	78.10%	83.27%	104.63%
10	101.70%	102.97%	92.68%	94.42%	108.56%
20	98.72%	98.65%	91.22%	99.52%	113.36%
40	109.31%	81.12%	89.22%	102.52%	81.20%
60	103.07%	89.82%	100.34%	102.36%	84.32%
90	104.82%	104.91%	92.88%	103.86%	97.74%
120	102.36%	100.99%	97.71%	112.18%	100.05%
150	105.67%	100.18%	98.98%	115.98%	102.42%
180	81.94%	100.45%	72.79%	78.60%	107.49%
210	101.51%	98.87%	84.35%	95.28%	114.06%
240	110.35%	99.95%	90.49%	94.05%	110.61%

Table 6E - Continued

PSO neutralized with TEA at RT	PBTC neutralized with MEA at RT	HEDP neutralized with MEA at RT	PSO neutralized with MEA at RT
-1.66%	-2.12%	-1.96%	-2.59%
100.00%	100.00%	88.97%	100.00%
94.34%	91.42%	100.00%	104.04%
99.51%	95.13%	111.81%	105.40%
91.53%	92.96%	107.42%	100.56%
97.66%	86.20%	112.52%	98.21%
101.47%	89.72%	112.16%	102.41%
97.52%	92.34%	110.32%	104.19%
95.18%	96.23%	106.74%	104.32%
97.39%	96.43%	105.87%	106.85%
91.23%	94.51%	98.16%	95.81%
100.11%	95.96%	108.81%	103.98%
105.34%	93.66%	106.77%	101.63%

5 Table 6E - Continued

PBTC neutralized with IPA at RT	HEDP neutralized with IPA at RT	PSO neutralized with IPA at RT
1.82%	-3.60%	-2.60%
75.36%	100.00%	100.00%

100.00%	79.69%	87.26%
96.16%	97.28%	89.70%
95.53%	102.58%	87.14%
75.16%	103.94%	70.95%
82.79%	97.13%	73.31%
96.16%	102.38%	84.42%
89.94%	100.00%	84.38%
95.29%	98.30%	84.50%
91.79%	81.44%	85.60%
92.90%	94.02%	80.61%
89.89%	103.60%	86.09%

**Table 6F.** Relative Lipase Enzyme Activity in a Use Solution of the All Ash Detergent Composition Containing a Lipase and Various Phosphonates at a Level of 0.3 wt-% at 120 °F.

time (min)	ATMP	DGAP	PBTC	HEDP
Blank	1.74%	0.45%	-0.27%	-2.79%
0	100.00%	100.00%	100.00%	100.00%
5	61.64%	87.31%	44.27%	39.01%
10	63.66%	76.38%	26.34%	21.35%
20	54.08%	57.74%	13.39%	7.45%
40	41.20%	40.00%	3.62%	0.88%
60	36.19%	33.56%	1.82%	-0.91%
90	30.66%	25.59%	0.36%	-1.68%
120	19.17%	21.03%	0.52%	-1.38%
150	23.54%	17.27%	2.28%	-1.94%
180	14.43%	14.06%	2.33%	-4.15%
210	13.63%	12.18%	3.25%	-4.41%
240	12.97%	10.79%	3.58%	-3.60%

5

The protease, amylase, and lipase activities at different time points in a use solution of the various detergent compositions containing a phosphonate were plotted in **Figures 1A-Figure 6G**, respectively. The data in **Table 4A – Table 4E** and **Figure 1A-Figure 2C** shows that protease in the base formula is more stable in the detergent composition that contains ATMP, DGAP, PAPERMP, and MEAP, which can retain at least 50% protease activity even after its use solution was generated for 4 hours. AMPA also improves protease stability, but to a lesser extent. These phosphonates share a common structure feature of the NR'R''-PO(OH)<sub>2</sub>. On the other hand, a detergent composition that contains

10

no phosphonate or other type of phosphonates, such as PBTC and HEDP, show a very short enzyme activity time frame and lose more than 70% of its enzyme activity after merely 20 minutes after the use solution is generated. However, for at least some other phosphonates, such as PBTC and HEDP that share common structure feature of

5 CR'R'R'''-PO(OH)<sub>2</sub>, if the phosphonate is neutralized with an amine, such as triethanolamine, a detergent composition that contains an amine salt of such a phosphonate shows a much improved enzyme activity than the corresponding detergent composition in which the phosphonate is not neutralized by an amine. **Figure 2D** shows that protease in the all ash formula is more stable in a use solution of the detergent composition that

10 contains ATMP or DGAP than in a use solution of the same detergent composition that contains PBTC or HEDP without amine(s).

The data in **Table 5A – Table 5E** and **Figure 3A-Figure 4C** shows that amylase activity in the base formula is also affected differently by different types of phosphonates, in a similar manner as the protease activity, although the effect of phosphonates on

15 amylase activity is not as significant as on protease activity. **Figure 4D** shows, however, that amylase in the all ash formula is more stable in a use solution of the detergent composition that contains ATMP or DGAP than in a use solution of the same detergent composition that contains PBTC or HEDP without amine(s).

The data in **Table 6A – Table 6F** and **Figure 5A-Figure 6F** shows that lipase is

20 also more stable in the detergent composition that contains ATMP, DGAP, PAPERMP, and MEAP, which can retain at least about 15% lipase activity even after its use solution was generated for 4 hours. **Figure 6G** shows, however, that lipase in the all ash formula is more stable in a use solution of the detergent composition that contains ATMP or DGAP than in a use solution of the same detergent composition that contains PBTC or HEDP

25 without amine(s).

Applicant unexpectedly discovered that some phosphonates, some of which are commonly used in detergent compositions, actually destabilize protease, amylase, or lipase as shown in **Figures 1A-Figure 6G**.

Although phosphonates, as scale inhibitors, sequestrants, or antiscalants, are used in

30 many consumer or HI&I detergents to combat hard water use in order to boost the performance of other actives, their general ability as enzyme stabilizers in detergent compositions has not been recognized or investigated, let alone for the specific types of

phosphonates as disclosed here. Applicant unexpectedly discovered that these two specific types of phosphonates, alone or in their amine salt forms, can stabilize enzymes in a detergent composition after the generation of its use solution during the cleaning application. This discovery led to increased effectiveness of the detergent compositions and new way to produce or formulate new detergent compositions that are more efficient to remove soils and cost effective.



The inventions being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the disclosures and all such modifications are intended to be included within the scope of the following claims.

The above specification provides a description of the manufacture and use of the disclosed compositions and methods.

15

**CLAIMS:**

1. A composition comprising:  
 an enzyme, wherein the enzyme is a protease, amylase, lipase, cellulose, peroxidase, gluconase, or a mixture thereof;



or salt thereof; and

from about 50 wt-% to about 95 wt-% of a metal carbonate, metal bicarbonate, or a mixture thereof;

wherein R<sup>10</sup> and R<sup>11</sup> are independently hydrogen, a substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, an ethanol, a diglycol, -CH<sub>2</sub>-PO(OH)<sub>2</sub>, or phosphonate ester, or wherein R<sup>10</sup> and R<sup>11</sup> are both -CH<sub>2</sub>-PO(OH)<sub>2</sub> groups;

wherein (EO) is ethylene oxide, (PO) is propylene oxide, and n is an integer of 1-30; and


wherein the composition is a pressed solid block.

2. The composition of claim 1, wherein R<sup>11</sup> is the -CH<sub>2</sub>-PO(OH)<sub>2</sub> group.

3. The composition of claim 1, wherein R<sup>11</sup> is the -CH<sub>2</sub>-PO(OH)<sub>2</sub> group and R<sup>10</sup> is the substituted alkyl.

4. The composition of claim 1, wherein R<sup>11</sup> is the -CH<sub>2</sub>-PO(OH)<sub>2</sub> group and R<sup>10</sup> is the phosphonate ester.

5. The composition of claim 1, wherein the phosphonate is



or aminomethyl phosphonic acid, and wherein n is an integer of 1-30.

6. The composition of any one of claims 1-5, wherein the composition comprises the metal carbonate and the metal bicarbonate.

7. The composition of claim 6, wherein the molar ratio of the metal carbonate and the metal bicarbonate is from about 0.25:1 to about 1:0.25.

8. The composition of any one of claims 1-7, wherein the enzyme is the protease, amylase, lipase, or mixture thereof.

9. The composition of any one of claims 1-8, wherein the phosphonate is about 0.1-35 wt-% of the composition.

10. The composition of any one of claims 1-9, wherein the enzyme is about 0.1-35 wt-% of the composition.

11. The composition of any one of claims 1-10, wherein the metal carbonate, metal bicarbonate, or mixture thereof is about 60-90 wt-% of the composition.
12. The composition of any one of claims 1-11, wherein a use solution of the composition has a pH of from about 8 to about 12.
13. The composition of any one of claims 1-12, wherein in a use solution of the composition, the enzyme retains at least 15% of its activity at 120 °F for at least 4 hours.
14. The composition of any one of claims 1-12, wherein in a use solution of the composition, the enzyme retains at least 20% of its activity at 120 °F for at least 4 hours.
15. The composition of any one of claims 1-14, further comprising an amine.
16. The composition of claim 15, wherein the amine is about 0.1-35 wt-% of the composition.
17. The composition of claim 15, wherein a molar ratio of the phosphonate to the amine is from about 0.5:1 to 1:0.5.
18. The composition of any one of claims 15-17, wherein the amine is an alkanolamine, monoethanolamine, diethanolamine, triethanolamine, isopropylamine, or a mixture thereof.
19. The composition of any one of claims 1-18, wherein the composition further comprises a metal hydroxide, tripolyphosphate, or mixture thereof.
20. The composition of any one of claims 1-19, further comprising one or more of an oxidizer, builder, water conditioning agent, peroxyacid, precursor of a peroxyacid, chelant, threshold agent, crystal modifier, sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, rinse aid, polymer, metal protecting agent,

stabilizing agent, corrosion inhibitor, sequestrant, chelating agent, fragrance, dye, rheology modifier, thickener, nonionic surfactant, cationic surfactant, zwitterionic surfactant, hydrotrope, coupler, or any combination thereof.


21. A solid detergent composition comprising:

from about 50 wt-% to about 95 wt-% of a metal carbonate, metal bicarbonate, or mixture thereof;

a phosphonate; and

an enzyme, wherein the enzyme is a protease, amylase, lipase, cellulose, peroxidase, gluconase, or a mixture thereof;

wherein the phosphonate is represented by a formula of



independently hydrogen, a substituted carboxylic acid, -CH<sub>2</sub>-PO(OH)<sub>2</sub>, phosphonate ester, an ethanol, a diglycol, substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, or methyl phosphonate, or wherein R<sup>10</sup> and R<sup>11</sup> are both -CH<sub>2</sub>-PO(OH)<sub>2</sub> groups;

wherein (EO) is ethylene oxide, (PO) is propylene oxide, and n is an integer of 1-30; and

wherein the solid detergent composition is a pressed solid block.

22. The solid detergent composition of claim 21, wherein the solid detergent is produced by a press process.


23. The solid detergent composition of claim 21 or claim 22, wherein the solid detergent has a mass of about 1 kilogram to about 10 kilograms.

24. The solid detergent composition of any one of claims 21-23, wherein the solid detergent is a multi-use solid detergent.

25. The solid detergent composition of any one of claims 21-24, wherein R<sup>11</sup> is the -CH<sub>2</sub>-PO(OH)<sub>2</sub> group.

26. The solid detergent composition of any one of claims 21-24, wherein R<sup>11</sup> is the -CH<sub>2</sub>-PO(OH)<sub>2</sub> group and R<sup>10</sup> is the diglycol, substituted alkyl, isopropyl-2-(EO)<sub>n</sub>-biphosphonateamine, or methyl-phosphonate.

27. The solid detergent composition of any one of claims 21-24, wherein the phosphonate is






aminomethyl phosphonic acid, or a salt thereof.

28. The solid detergent composition of any one of claims 21-27, wherein the solid detergent composition comprises the metal carbonate and the metal bicarbonate.

29. The solid detergent composition of claim 28, wherein the mole ratio of the metal carbonate and the metal bicarbonate is from about 0.25:1 to about 1:0.25.
30. The solid detergent composition of any one of claims 21-29, wherein the enzyme is the protease, amylase, lipase, or mixture thereof.
31. The solid detergent composition of any one of claims 21-30, further comprising an amine or salt thereof.
32. The solid detergent composition of claim 31, wherein the amine is fully neutralized.
33. The solid detergent composition of claim 31 or claim 32, wherein the amine comprises an alkanolamine, monoethanolamine, diethanolamine, triethanolamine, isopropylamine, a salt thereof, or any mixture thereof.
34. The solid detergent composition of any one of claims 21-33, further comprising a metal hydroxide, tripolyphosphate, or mixture thereof.
35. The solid detergent composition of any one of claims 21-34, further comprising one or more of an oxidizer, builder, water conditioning agent, peroxyacid, precursor of a peroxyacid, chelant, threshold agent, crystal modifier, sanitizing agent, defoaming agent, anti-redeposition agent, bleaching agent, solubility modifier, dispersant, rinse aid, polymer, metal protecting agent, stabilizing agent, corrosion inhibitor, sequestrant, chelating agent, fragrance, dye, rheology modifier, thickener, nonionic surfactant, cationic surfactant, zwitterionic surfactant, hydrotrope, coupler, or any combination thereof.
36. The solid detergent composition of any one of claims 21-35, wherein a use solution of the composition has a pH of from about 8 to about 12.

37. The solid detergent composition of any one of claims 21-36, wherein in a use solution of the composition, the enzyme retains at least 15% of its activity at 120 °F for at least 4 hours.
38. The solid detergent composition of any one of claims 21-37, wherein the phosphonate comprises about 0.1-35 wt-% of the composition.
39. The solid detergent composition of any one of claims 21-38, wherein the enzyme comprises about 0.1-35 wt-% of the composition.
40. The solid detergent composition of any one of claims 21-39, wherein the alkaline source comprises about 0.1-90 wt-% of the composition.
41. The solid detergent composition of any one of claims 21-40, wherein the solid detergent has a dimensional stability and has a growth exponent of less than 3% if heated at a temperature of 122°F.
42. A method of cleaning, sanitizing and/or bleaching comprising generating a use solution of the composition as defined by any one of claims 1-41, and contacting a surface or object in need of cleaning and sanitizing with the use solution.
43. A method of stabilizing an enzyme in a solid detergent, the method comprising:

adding a phosphonate of formula



, or salt thereof in an existing

detergent composition containing an enzyme and from about 50 wt-% to about 95 wt-% of a metal carbonate, metal bicarbonate, or mixture thereof, wherein  $R^{10}$  and  $R^{11}$  are independently hydrogen, a substituted carboxylic acid,  $-\text{CH}_2\text{-PO}(\text{OH})_2$ , phosphonate ester, an ethanol, a diglycol, substituted alkyl, 2-(EO)<sub>n</sub>-biphosphonateamine-ethyl, 2-(PO)<sub>n</sub>-biphosphonateamine-isopropyl, or methyl phosphonate, or wherein  $R^{10}$  and  $R^{11}$  are both  $-\text{CH}_2\text{-PO}(\text{OH})_2$  groups;

wherein (EO) is ethylene oxide, (PO) is propylene oxide, and n is an integer of 1-30; and wherein the solid detergent is a pressed solid block.




Figure 1A




Figure 1B

2/13




Figure 1C




Figure 2A




Figure 2B

4/13




Figure 2C




Figure 2D

5/13




Figure 3A




Figure 3B

6/13




Figure 3C

7/13




Figure 4A




Figure 4B

8/13




Figure 4C




Figure 4D

9/13




Figure 5A




Figure 5B

10/13




Figure 5C

11/13




Figure 6A




Figure 6B








Figure 6C





**Figure 6G**