



US 20040052866A1

(19) **United States**

(12) **Patent Application Publication**
Otterbein et al.

(10) **Pub. No.: US 2004/0052866 A1**

(43) **Pub. Date: Mar. 18, 2004**

(54) **METHODS OF TREATING HEPATITIS**

Related U.S. Application Data

(76) Inventors: **Leo E. Otterbein**, New Kensington, PA (US); **Augustine M. K. Choi**, Pittsburgh, PA (US); **Brian Zuckerbraun**, Pittsburgh, PA (US)

(60) Provisional application No. 60/381,527, filed on May 17, 2002.

Publication Classification

Correspondence Address:
FISH & RICHARDSON PC
225 FRANKLIN ST
BOSTON, MA 02110 (US)

(51) **Int. Cl.⁷** **A61K 33/00**
(52) **U.S. Cl.** **424/699**

(57) **ABSTRACT**

(21) Appl. No.: **10/439,632**

The present invention relates to a method of treating hepatitis in a patient, which includes administering a pharmaceutical composition that includes carbon monoxide to the patient.

(22) Filed: **May 16, 2003**




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6A-6C




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11A-11H




Fig. 12




Fig. 13A-13D




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21

METHODS OF TREATING HEPATITIS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No. 60/381,527 filed May 17, 2002, which is incorporated herein by reference in its entirety.

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

[0002] This invention was made with Government support under National Institutes of Health Grant Nos. R01-GM-44100, HL 58688, HL55330, HL60234, and AI42365. The Government has certain rights in this invention.

TECHNICAL FIELD

[0003] This invention relates to the treatment of hepatitis.

BACKGROUND

[0004] Carbon monoxide gas is poisonous in high concentrations. However, it is now recognized as an important signaling molecule (Verma et al., *Science* 259:381-384, 1993). It has also been suggested that carbon monoxide acts as a neuronal messenger molecule in the brain (Id.) and as a neuro-endocrine modulator in the hypothalamus (Pozzoli et al., *Endocrinology* 735:2314-2317, 1994). Like nitric oxide (NO), carbon monoxide is a smooth muscle relaxant (Utz et al., *Biochem Pharmacol.* 47:195-201, 1991; Christodoulides et al., *Circulation* 97:2306-9, 1995) and inhibits platelet aggregation (Mansouri et al., *Thromb Haemost.* 48:286-8, 1982). Inhalation of low levels of carbon monoxide (CO) has been shown to have anti-inflammatory effects in some models.

[0005] Hepatitis is a disease characterized by inflammation of the liver. The inflammation can be characterized by diffuse or patchy necrosis affecting acini. Causative agents of hepatitis include, for example, viruses, e.g., specific hepatitis viruses, e.g., hepatitis A, B, C, D, E, and G viruses; alcohol; and other drugs (e.g., isoniazid, methyl dopa, acetaminophen, amiodarone, and nitrofurantoin) (see *The Merck Manual of Diagnosis and Therapy*, 17th Edition, Section 4, Chapter 42).

SUMMARY

[0006] The present invention is based, in part, on the discovery that administration of CO can protect against the development of hepatitis.

[0007] Accordingly, the present invention features a method of treating, preventing, or reducing the risk of, hepatitis in a patient. The method includes identifying a patient diagnosed as suffering from or at risk for hepatitis (e.g., a patient diagnosed as suffering from or at risk for hepatitis), and administering to the patient a pharmaceutical composition comprising an amount of carbon monoxide effective to treat hepatitis in the patient.

[0008] The pharmaceutical composition can be administered to the patient by any method known in the art for administering gases and/or liquids to patients, e.g., via inhalation, insufflation, infusion, injection, and/or ingestion. In one embodiment of the present invention, the pharmaceutical composition is administered to the patient by inha-

lation. In another embodiment, the pharmaceutical composition is administered to the patient orally. In still another embodiment, the pharmaceutical composition is administered directly to the abdominal cavity of the patient. In yet another embodiment, the pharmaceutical composition is administered by an extracorporeal membrane gas exchange device or an artificial lung. In another embodiment, the patient is an alcoholic.

[0009] The patient can be an animal, human or non-human. For example, the patient can be any mammal, e.g., humans, other primates, pigs, rodents such as mice and rats, rabbits, guinea pigs, hamsters, cows, horses, cats, dogs, sheep and goats. The hepatitis can be the result of, or a person may be considered at risk for hepatitis because of, any of a number of factors, e.g., infections, e.g., viral infections, e.g., infection with hepatitis A, B, C, D, E and/or G virus; alcohol use (e.g., alcoholism); drug use (e.g., one or more drugs described herein, e.g., acetaminophen, anesthetics, anti-tuberculous drugs, antifungal agents, antidiabetic drugs, neuroleptic agents, and drugs used to treat HIV infection and AIDS); autoimmune conditions (e.g., autoimmune hepatitis); and/or surgical procedures. The pharmaceutical composition can be in any form, e.g., gaseous or liquid form.

[0010] In another embodiment, the method further includes administering to the patient at least one of the following treatments: inducing HO-1 or ferritin in the patient; expressing recombinant HO-1 or ferritin in the patient; and administering a pharmaceutical composition comprising HO-1, bilirubin, biliverdin, ferritin, or apoferritin, iron, desferoxamine, or iron dextran to the patient. Also contemplated is use of CO and any of the above-listed agents in the preparation of a medicament for treatment or prevention of hepatitis.

[0011] In another embodiment, the hepatitis (or the risk for hepatitis) is not caused by surgery (e.g., abdominal or transplant surgery), bacterial endotoxin, septic shock, and/or systemic inflammation.

[0012] In another aspect, the invention features a method of treating or preventing hepatitis in a patient, which includes identifying a patient suffering from or at risk for hepatitis (e.g., a patient diagnosed as suffering from or at risk for hepatitis), providing a vessel containing a pressurized gas comprising carbon monoxide gas, releasing the pressurized gas from the vessel to form an atmosphere comprising carbon monoxide gas, and exposing the patient to the atmosphere, wherein the amount of carbon monoxide in the atmosphere is sufficient to treat hepatitis in the patient.

[0013] In still another aspect, the invention features a method of performing abdominal surgery, e.g., liver transplantation, on a patient, which includes identifying a patient in need of abdominal surgery, wherein hepatitis is a risk of the abdominal surgery; performing abdominal surgery on the patient, and before, during, or after the performing step, causing the patient to inhale an amount of carbon monoxide gas sufficient to reduce the risk of hepatitis in the patient. Also contemplated is use of CO in the preparation of a medicament, e.g., a gaseous or liquid medicament, for use in the treatment or prevention of hepatitis.

[0014] The invention also features a method of treating hepatitis in a patient suffering from or at risk for hepatitis not

caused by surgery and/or endotoxin, e.g., hepatitis caused by any factor described herein other than surgery and/or endotoxin. The method includes identifying a patient suffering from or at risk for hepatitis not caused by surgery and/or endotoxin and administering to the patient a pharmaceutical composition comprising an amount of carbon monoxide effective to treat hepatitis in the patient.

[0015] Also within the invention is a method of administering a hepatitis-inducing drug (i.e., a hepatotoxic drug, e.g., isoniazid, methyl dopa, acetaminophen, amiodarone, or nitrofurantoin) to a patient. The method includes administering the drug to the patient, and before, during, and/or after administering the drug, administering to the patient a pharmaceutical composition comprising carbon monoxide in an amount effective to treat hepatitis in the patient.

[0016] In another aspect, the invention provides a vessel comprising medical grade compressed CO gas. The vessel can bear a label indicating that the gas can be used to treat hepatitis in a patient. Alternatively or in addition, the vessel can bear a label indicating that the gas can be administered to a patient in conjunction with administration of a hepatitis-inducing drug (i.e., a hepatotoxic drug), e.g., acetaminophen. The CO gas can be in an admixture with nitrogen gas, with nitric oxide and nitrogen gas, or with an oxygen-containing gas. The CO gas can be present in the admixture at a concentration of at least about 0.025%, e.g., at least about 0.05%, 0.10%, 0.50%, 1.0%, 2.0%, 10%, 50%, or 90%.

[0017] Also within the invention is the use of CO in the manufacture of a medicament for treatment or prevention of hepatitis. The medicament can be used in a method for treating hepatitis in a patient suffering from or at risk for hepatitis in accordance with the methods described herein. The medicament can be in any form described herein, e.g., a liquid or gaseous CO composition.

[0018] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Suitable methods and materials are described below, although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. The materials, methods, and examples are illustrative only and not intended to be limiting.

[0019] The details of one or more embodiments of the invention are set forth in the description below. Other features, objects, and advantages of the invention will be apparent from the description and the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] FIG. 1 is a bar graph illustrating that induction of HO-1 protects mouse hepatocytes from TNF- α /D-gal-induced cell death. CoPP=cobalt protoporphyrin; ALT=serum alanine aminotransferase; TNF=tumor necrosis factor alpha. Results are the mean \pm SD of 6-8 mice/group *p<0.005.

[0021] FIG. 2 is a bar graph illustrating that exogenous CO protects hepatocytes against TNF-A induced cell death in a cGMP/p38 pathway-independent and an NF- κ B activation-dependent manner. CO=carbon monoxide; Air=room air; TNF=tumor necrosis factor alpha; BAY=BAY 11-7082 (inhibits NF- κ B activation); I κ B=I κ B α (prevents NF- κ B activation); ODQ=1H-[1,2,4]Oxadiazolo[4,3-a]quinoxalin-1-one (a selective guanylyl cyclase inhibitor); Lac-Z=pIEP-Lac-Z (adenoviral control). Results shown are the mean \pm SD of triplicate wells from four independent experiments (*p<0.01).

[0022] FIG. 3 is a bar graph illustrating that exogenous CO protects human hepatocytes against TNF- α /ActD-induced cell death. CO=carbon monoxide; Air=room air; TNF=TNF- α /ActD. Results are mean \pm SD of triplicate wells from 3 independent experiments. *p<0.05.

[0023] FIG. 4 is a bar graph illustrating that exogenous CO causes an increase in NF- κ B activation in hepatocytes. CO=carbon monoxide; Air=room air; BAY=BAY 11-7082; CM=cytokine mixture (TNF- α (500 U/ml), IL-1 β (100 U/ml), and IFN- δ (100 U/ml)). Results shown are the mean \pm SE of triplicate wells from three independent experiments. *p<0.001 versus Air.

[0024] FIG. 5 is a picture of a polyacrylamide gel illustrating that exogenous CO induces an increase in NF- κ B nuclear translocation and DNA binding as measured by electrophoretic mobility shift assay (EMSA). FP=free probe (no nuclear protein, thus no DNA binding); TOTAL=Nf κ B bands without antibody supershifting.

[0025] FIGS. 6A-6C are photomicrographs of primary hepatocytes immunostained to detect nuclear p65 localization, illustrating that exogenous CO causes an increase in NF- κ B activation in hepatocytes. FIG. 6A: air-exposed hepatocytes. FIG. 6B: hepatocytes exposed to cytokine mixture (TNF- α (500 U/ml), IL-1 (100 U/ml), and IFN- δ (100 U/ml)). FIG. 6C: CO-exposed hepatocytes. Images are representative of 6 different fields. Bar represents 10 μ m.

[0026] FIG. 7 is a bar graph illustrating that exogenous CO-induced protection of hepatocytes involves NF- κ B-dependent iNOS expression. CO=carbon monoxide; Air=room air; BAY=BAY 11-7082; CM=cytokine mixture. Results shown are the mean \pm SE of triplicate wells from four independent experiments. *p<0.001 versus air and air/BAY-treated cells.

[0027] FIG. 8 is a picture of a Western blot illustrating that iNOS protein expression in hepatocytes is markedly increased by exposure to TNF-a in the presence of CO as compared to exposure to TNF- α alone. iNOS=inducible nitric oxide synthase; CO=carbon monoxide; Air=room air; TNF=TNF- α /ActD; β -Actin=control protein. The immunoblot is representative of 3 independent experiments.

[0028] FIG. 9 is a bar graph illustrating that iNOS activity-deficient mouse (inos^{-/-}) hepatocytes are not protected by CO against TNF-a-induced cell death. CO=carbon monoxide; Air=room air; TNF=TNF- α /ActD; inos^{-/-}=iNOS knockout mice; L-NIO=L-N5-(1-iminoethyl)-ornithine-2HCl. Results shown are the mean \pm SE of triplicate wells from four independent experiments. *p<0.01 versus non-TNF/ActD and CO/TNF/ActD-treated cells.

[0029] FIG. 10 is a bar graph illustrating that exogenously-administered CO prevents TNF- α /D-Gal-induced liver injury in mice. ALT=serum alanine aminotransferase;

CO=carbon monoxide; Air=room air. Results presented as mean \pm SD of 18-20 mice. * p <0.001 versus air-treated.

[0030] FIGS. 11A-11H are photomicrographs of liver samples illustrating that exogenously-administered CO prevents TNF- α /D-Gal-induced liver injury in mice. FIGS. 11A and 11B: liver samples from mice exposed to room air and CO, respectively, and stained with hematoxylin & eosin (H & E). FIGS. 11C and 11D: liver samples from TNF-WD-Gal-treated mice exposed to room air and CO, respectively, and stained with H & E. FIGS. 11E and 11F: liver samples from TNF- α /D-Gal-treated mice exposed to room air and CO, respectively, and stained to detect activated caspase-3. FIGS. 11G and 11H: liver samples from TNF- α /D-Gal-treated mice exposed to room air and CO, respectively, and stained using terminal deoxynucleotidyl transferase mediated dUTP nick end labeling (TUNEL). Images are representative sections from 15-20 sections/liver from 3-4 individual mice/group. Bar represents 20 μ m.

[0031] FIG. 12 is a picture of a Western blot illustrating that livers of mice exposed to TNF- α /D-Gal and treated with inhaled CO display increased iNOS protein levels. Wild type=wild type mice; iNOS $^{-/-}$ =iNOS deficient mice; CO=carbon monoxide; Air=room air; TNF=TNF- α /D-Gal; β -Actin=control protein.

[0032] FIGS. 13A-13D are photomicrographs of liver samples illustrating that the livers of mice exposed to TNF- α /D-Gal and treated with inhaled CO display increased iNOS protein levels. FIG. 13A: liver sample from room air-exposed mouse. FIG. 13B: liver sample from CO-exposed mouse. FIG. 13C: liver sample from mouse exposed to TNF- α /D-Gal and room air. FIG. 13D: liver sample from mouse exposed to TNF- α /D-Gal and CO. Images are representative of 6 separate animals and 6-10 different sections/liver sample. Bar represents 20 μ m.

[0033] FIG. 14 is a bar graph illustrating that CO does not protect against liver damage in the absence of iNOS function/expression. L-NIL=L-N6-(1-iminoethyl)-lysine-dihydrochloride (a selective inhibitor of iNOS); CO=carbon monoxide; Air=room air; TNF=TNF- α /D-Gal. Results are mean \pm SD of 6-8 animals/group. * p <0.01 versus CO/TNF- α /D-gal and air and CO controls.

[0034] FIG. 15 is a picture of a Western blot illustrating that the livers of CO-treated mice displayed increased expression of HO-1 in both the presence and absence of TNF- α /D-Gal. CO=carbon monoxide; Air=room air; TNF=TNF-60 /D-Gal; β -Actin=control protein. Blot is representative of 2 independent experiments.

[0035] FIG. 16 is a picture of a Western blot illustrating that the livers of CO-treated mice do not display increased expression of HO-1 in the presence or absence of TNF- α /D-Gal if iNOS is inhibited using L-NIL. CO=carbon monoxide; Air=room air; TNF=TNF- α /D-Gal; β -Actin=control protein; L-NIL=L-N6-(1-iminoethyl)-lysine-dihydrochloride (a selective inhibitor of iNOS). Blot is representative of 2 independent experiments.

[0036] FIG. 17 is a bar graph illustrating that CO-induced HO-1 is protective against TNF- α -induced liver damage in mice. ALT=serum alanine aminotransferase; Air=room air; TNF=TNF- α /D-Gal; Sn=tin protoporphyrin (an inhibitor of HO-1); VP=V-PYRRO (a nitric oxide donor). Results are expressed as mean \pm SD of 8-10 mice/group. * p <0.05 versus CO/TNF/D-gal-treated mice.

[0037] FIG. 18 is a bar graph illustrating that induction of HO-1 is protective against TNF- α -induced liver injury independent of iNOS activity. ALT=serum alanine aminotransferase; Air=room air; TNF=TNF- α /D-Gal; L-NIL=L-N6-(1-iminoethyl)-lysine-dihydrochloride (a selective inhibitor of iNOS); CoPP=cobalt protoporphyrin (an inducer of HO-1); iNOS $^{-/-}$ =iNOS deficient mice. Results are mean \pm SD of 6-8 mice/group. * p <0.001 versus Air/TNF and L-NIL/TNF.

[0038] FIG. 19 is bar graph illustrating that HO-1 expression is required for CO-induced protection of mouse hepatocytes from TNF- α /ActD-induced cell death. Wild type (black bars)=hepatocytes isolated from wild type C57BL/6J mice; hmox-1 $^{-/-}$ (white bars)=hepatocytes isolated from HO-1 null mice; CO=carbon monoxide; Air=room air; TNF- α =TNF- α /ActD. * p <0.01 versus non-TNF- α /ActD treated cells and versus TNF- α /ActD-treated cells that were also treated with CO.

[0039] FIG. 20 is bar graph illustrating that HO-1 expression is required for NO-induced protection of mouse hepatocytes from TNF- α /ActD-induced cell death. Wild type (black bars)=hepatocytes isolated from wild type C57BL/6J mice; hmox-1 $^{-/-}$ (white bars)=hepatocytes isolated from HO-1 null mice; SNAP=s-nitroso-N-acetyl-penicillamine (an NO donor); Air=room air; TNF-A=TNF- α /ActD. * p <0.01 versus non-TNF- α /ActD treated cells and versus TNF- α /ActD-treated cells that were also treated with NO.

[0040] FIG. 21 is a bar graph illustrating that CO-exposed mice were protected from acetaminophen-induced liver injury. ALT=serum alanine aminotransferase; Air=room air; APAP=acetaminophen. Results are mean \pm SD of 4-8 mice/group.

DETAILED DESCRIPTION

[0041] The term "carbon monoxide" (or "CO") as used herein describes molecular carbon monoxide in its gaseous state, compressed into liquid form, or dissolved in aqueous solution. The terms "carbon monoxide composition" and "pharmaceutical composition comprising carbon monoxide" is used throughout the specification to describe a gaseous or liquid composition containing carbon monoxide that can be administered to a patient and/or an organ, e.g., the liver. The skilled practitioner will recognize which form of the pharmaceutical composition, e.g., gaseous, liquid, or both gaseous and liquid forms, is preferred for a given application.

[0042] The terms "effective amount" and "effective to treat," as used herein, refer to an amount or concentration of carbon monoxide utilized for period of time (including acute or chronic administration and periodic or continuous administration) that is effective within the context of its administration for causing an intended effect or physiological outcome. Effective amounts of carbon monoxide for use in the present invention include, for example, amounts that prevent hepatitis, reduce the risk of hepatitis, reduce the symptoms of hepatitis, or improve the outcome of other hepatitis treatments.

[0043] For gases, effective amounts of carbon monoxide generally fall within the range of about 0.0000001% to about 0.3% by weight, e.g., 0.0001% to about 0.25% by weight, preferably at least about 0.001%, e.g., at least 0.005%, 0.010%, 0.02%, 0.025%, 0.03%, 0.04%, 0.05%, 0.06%, 0.08%, 0.10%, 0.15%, 0.20%, 0.22%, or 0.24% by weight

carbon monoxide. Preferred ranges include, e.g., 0.001% to about 0.24%, about 0.005% to about 0.22%, about 0.005% to about 0.05%, about 0.010% to about 0.20%, about 0.02% to about 0.15%, about 0.025% to about 0.10%, or about 0.03% to about 0.08%, or about 0.04% to about 0.06%. For liquid solutions of CO, effective amounts generally fall within the range of about 0.0001 to about 0.0044 g CO/100 g liquid, e.g., at least 0.0001, 0.0002, 0.0004, 0.0006, 0.0008, 0.0010, 0.0013, 0.0014, 0.0015, 0.0016, 0.0018, 0.0020, 0.0021, 0.0022, 0.0024, 0.0026, 0.0028, 0.0030, 0.0032, 0.0035, 0.0037, 0.0040, or 0.0042 g CO/100 g aqueous solution. Preferred ranges include, e.g., about 0.0010 to about 0.0030 g CO/100 g liquid, about 0.0015 to about 0.0026 g CO/100 g liquid, or about 0.0018 to about 0.0024 g CO/100 g liquid. A skilled practitioner will appreciate that amounts outside of these ranges may be used, depending upon the application.

[0044] The term “patient” is used throughout the specification to describe an animal, human or non-human, to whom treatment according to the methods of the present invention is provided. Veterinary applications are contemplated by the present invention. The term includes but is not limited to mammals, e.g., humans, other primates, pigs, rodents such as mice and rats, rabbits, guinea pigs, hamsters, cows, horses, cats, dogs, sheep and goats. The term “treat(ment),” is used herein to describe delaying the onset of, inhibiting, or alleviating the effects of a condition, e.g., hepatitis, in a patient.

[0045] The term “hepatitis” is an art-recognized term and is used herein to refer to a disease of patients characterized in part by inflammation of the liver. Causative agents of hepatitis include, for example, infections, e.g., infection with specific hepatitis viruses, e.g., hepatitis A, B, C, D, E, and G viruses; or hepatotoxic agents, e.g., hepatotoxic drugs (e.g., isoniazid, methyl dopa, acetaminophen, amiodarone, and nitrofurantoin), and toxins (e.g., endotoxin or environmental toxins). Hepatitis may occur postoperatively in liver transplantation patients. Further examples of drugs and toxins that may cause hepatitis (i.e., hepatotoxic agents) are described in Feldman: Sleisenger & Fordtran’s Gastrointestinal and Liver Disease, 7th ed., Chapter 17 (Liver Disease Caused by Drugs, Anesthetics, and Toxins), the contents of which are expressly incorporated herein by reference in their entirety. Such examples include, but are not limited to, methyl dopa and phenytoin, barbiturates, e.g., phenobarbital; sulfonamides (e.g., in combination drugs such as co-trimoxazole (sulfamethoxazole and trimethoprim); sulfasalazine; salicylates; disulfiram; β -adrenergic blocking agents e.g., acebutolol, labetalol, and metoprolol); calcium channel blockers, e.g., nifedipine, verapamil, and diltiazem; synthetic retinoids, e.g., etretinate; gastric acid suppression drugs e.g., oxmetidine, ebrotidine, cimetidine, ranitidine, omeprazole and famotidine; leukotriene receptor antagonists, e.g., zafirlukast; anti-tuberculous drugs, e.g., rifampicin and pyrazinamide; antifungal agents, e.g., ketoconazole, terbinafine, fluconazole, and itraconazole; antidiabetic drugs, e.g., thiazolidinediones, e.g., troglitazone and rosiglitazone; drugs used in neurologic disorders, e.g., neuroleptic agents, antidepressants (e.g., fluoxetine, paroxetine, venlafaxine, trazodone, tolcapon, and nefazodone), hypnotics (e.g., alpidem, zolpidem, and bentazepam), and other drugs, e.g., tacrine, dantrolene, riluzole, tizanidine, and alverine; nonsteroidal anti-inflammatory drugs, e.g., bromfenac; COX-2 inhibitors; cyproterone acetate; leflunomide; anti-

ral agents, e.g., flaluridine, didanosine, zalcitabine, stavudine, lamivudine, zidovudine, abacavir; anticancer drugs, e.g., tamoxifen and methotrexate; recreational drugs, e.g., cocaine, phencyclidine, and 5-methoxy-3,4-methylenedioxymethamphetamine; L-asparaginase; amodiaquine; hycanthone; anesthetic agents; e.g., halothane, enflurane, and isoflurane; vitamins e.g., vitamin A; and dietary and/or environmental toxins, e.g., pyrrolizidine alkaloids, toxin from *Amanita phalloides* or other toxic mushrooms, aflatoxin, arsenic, Bordeaux mixture (copper salts and lime), vinyl chloride monomer; carbon tetrachloride, beryllium, dimethylformamide, dimethylnitrosamine, methylenedianiline, phosphorus, chlordecone (Kepone), 2,3,7,8-tetrachloro-dibenzo p-dioxin (TCDD), tetrachloroethane, tetrachloroethylene, 2,4,5-trinitrotoluene, 1,1,1-trichloroethane, toluene, and xylene, and known “herbal remedies,” e.g., ephedrine and eugenol.

[0046] Symptoms of hepatitis can include fatigue, loss of appetite, stomach discomfort, and/or jaundice (yellowing of the skin and/or eyes). More detailed descriptions of hepatitis are provided, for example, in the *The Merck Manual of Diagnosis and Therapy*, 17th Edition, Section 4, Chapter 42, Section 4, Chapter 44, and Section 4, Chapter 40, the contents of which are expressly incorporated herein by reference in their entirety.

[0047] Skilled practitioners will appreciate that a patient can be diagnosed by a physician as suffering from hepatitis by any method known in the art, e.g., by assessing liver function, e.g., using blood tests for serum alanine aminotransferase (ALT) levels, alkaline phosphatase (AP), or bilirubin levels.

[0048] Individuals considered at risk for developing hepatitis may benefit particularly from the invention, primarily because prophylactic treatment can begin before there is any evidence of hepatitis. Individuals “at risk” include, e.g., patients infected with hepatitis viruses, or individuals suffering from any of the conditions or having the risk factors described herein (e.g., patients exposed to hepatotoxic agents). The skilled practitioner will appreciate that a patient can be determined to be at risk for hepatitis by a physician’s diagnosis.

[0049] Amounts of CO effective to treat hepatitis can be administered to a patient on the day the patient is diagnosed as suffering from hepatitis or any condition associated with hepatitis, or as having any risk factor associated with an increased likelihood that the patient will develop hepatitis (e.g., that the patient has recently been, is being, or will be exposed to a hepatotoxic agent, e.g., a hepatotoxic drug such as acetaminophen). Patients can inhale CO at concentrations ranging from 10 ppm to 1000 ppm, e.g., about 100 ppm to about 800 ppm, about 150 ppm to about 600 ppm, or about 200 ppm to about 500 ppm. Preferred concentrations include, e.g., about 30 ppm, 50 ppm, 75 ppm, 100 ppm, 125 ppm, 200 ppm, 250 ppm, 500 ppm, 750 ppm, or about 1000 ppm. CO can be administered to the patient intermittently or continuously. CO can be administered for about 1, 2, 4, 6, 8, 10, 12, 14, 18, or 20 days, or greater than 20 days, e.g., 1, 2, 3, 5, or 6 months, or until the patient no longer exhibits symptoms of hepatitis, or until the patient is diagnosed as no longer being at risk for hepatitis. In a given day, CO can be administered continuously for the entire day, or intermittently, e.g., a single whiff of CO per day (where a high

concentration is used), or for up to 23 hours per day, e.g., up to 20, 15, 12, 10, 6, 3, or 2 hours per day, or up to 1 hour per day.

[0050] If the patient needs to be treated with a hepatotoxic drug (e.g., because prescribed by a physician), the patient can be treated with CO (e.g., a gaseous CO composition) before, during, and/or after administration of the drug. For example, CO can be administered to the patient, intermittently or continuously, starting 0 to 20 days before the drug is administered (and where multiple doses are given, before each individual dose), e.g., starting at least about 30 minutes, e.g., about 1, 2, 3, 5, 7, or 10 hours, or about 1, 2, 4, 6, 8, 10, 12, 14, 18, or 20 days, or greater than 20 days, before the administration. Alternatively or in addition, CO can be administered to the patient concurrent with administration of the drug. Alternatively or in addition, CO can be administered to the patient after administration of the drug, e.g., starting immediately after administration, and continuing intermittently or continuously for about 1, 2, 3, 5, 7, or 10 hours, or about 1, 2, 5, 8, 10, 20, 30, 50, or 60 days, indefinitely, or until a physician determines that administration of CO is no longer necessary (e.g., after the hepatotoxic drug is eliminated from the body or can no longer cause damage to the liver).

[0051] Preparation of Gaseous Compositions

[0052] A carbon monoxide composition may be a gaseous carbon monoxide composition. Compressed or pressurized gas useful in the methods of the invention can be obtained from any commercial source and in any type of vessel appropriate for storing compressed gas. For example, compressed or pressurized gases can be obtained from any source that supplies compressed gases, such as oxygen, for medical use. The term "medical grade" gas, as used herein, refers to gas suitable for administration to patients as defined herein. The pressurized gas including CO used in the methods of the present invention can be provided such that all gases of the desired final composition (e.g., CO, He, NO, CO₂, O₂, N₂) are in the same vessel, except that NO and O₂ cannot be stored together. Optionally, the methods of the present invention can be performed using multiple vessels containing individual gases. For example, a single vessel can be provided that contains carbon monoxide, with or without other gases, the contents of which can be optionally mixed with room air or with the contents of other vessels, e.g., vessels containing oxygen, nitrogen, carbon dioxide, compressed air, or any other suitable gas or mixtures thereof.

[0053] Gaseous compositions administered to a patient according to the present invention typically contain 0% to about 79% by weight nitrogen, about 21% to about 100% by weight oxygen and about 0.000001% to about 0.3% by weight (corresponding to about 1 ppb or 0.001 ppm to about 3,000 ppm) carbon monoxide. Preferably, the amount of nitrogen in the gaseous composition is about 79% by weight, the amount of oxygen is about 21% by weight and the amount of carbon monoxide is about 0.0001% to about 0.25% by weight, preferably at least about 0.001%, e.g., at least about 0.005%, 0.010%, 0.02%, 0.025%, 0.03%, 0.04%, 0.05%, 0.06%, 0.08%, 0.10%, 0.15%, 0.20%, 0.22%, or 0.24% by weight. Preferred ranges of carbon monoxide include about 0.005% to about 0.24%, about 0.01% to about 0.22%, about 0.015% to about 0.20%, about 0.08% to about 0.20%, and about 0.025% to about 0.1% by weight. It is

noted that gaseous carbon monoxide compositions having concentrations of carbon monoxide greater than 0.3% (such as 1% or greater) may be used for short periods (e.g., one or a few breaths), depending upon the application.

[0054] A gaseous carbon monoxide composition may be used to create an atmosphere that comprises carbon monoxide gas. An atmosphere that includes appropriate levels of carbon monoxide gas can be created, for example, by providing a vessel containing a pressurized gas comprising carbon monoxide gas, and releasing the pressurized gas from the vessel into a chamber or space to form an atmosphere that includes the carbon monoxide gas inside the chamber or space. Alternatively, the gases can be released into an apparatus that culminates in a breathing mask or breathing tube, thereby creating an atmosphere comprising carbon monoxide gas in the breathing mask or breathing tube, ensuring the patient is the only person in the room exposed to significant levels of carbon monoxide.

[0055] Carbon monoxide levels in an atmosphere can be measured or monitored using any method known in the art. Such methods include electrochemical detection, gas chromatography, radioisotope counting, infrared absorption, colorimetry, and electrochemical methods based on selective membranes (see, e.g., Sunderman et al., *Clin. Chem.* 28:2026-2032, 1982; Ingi et al., *Neuron* 16:835-842, 1996). Sub-parts per million carbon monoxide levels can be detected by, e.g., gas chromatography and radioisotope counting. Further, it is known in the art that carbon monoxide levels in the sub-ppm range can be measured in biological tissue by a midinfrared gas sensor (see, e.g., Morimoto et al., *Am. J. Physiol. Heart. Circ. Physiol.* 280:H482-H488, 2001). Carbon monoxide sensors and gas detection devices are widely available from many commercial sources.

[0056] Preparation of Liquid Compositions

[0057] A carbon monoxide composition may also be a liquid carbon monoxide composition. A liquid can be made into a carbon monoxide composition by any method known in the art for causing gases to become dissolved in liquids. For example, the liquid can be placed in a so-called "CO₂ incubator" and exposed to a continuous flow of carbon monoxide, preferably balanced with carbon dioxide, until a desired concentration of carbon monoxide is reached in the liquid. As another example, carbon monoxide gas can be "bubbled" directly into the liquid until the desired concentration of carbon monoxide in the liquid is reached. The amount of carbon monoxide that can be dissolved in a given aqueous solution increases with decreasing temperature. As still another example, an appropriate liquid may be passed through tubing that allows gas diffusion, where the tubing runs through an atmosphere comprising carbon monoxide (e.g., utilizing a device such as an extracorporeal membrane oxygenator). The carbon monoxide diffuses into the liquid to create a liquid carbon monoxide composition.

[0058] It is likely that such a liquid composition intended to be introduced into a living animal will be at or about 37° C. at the time it is introduced into the animal.

[0059] The liquid can be any liquid known to those of skill in the art to be suitable for administration to patients (see, for example, *Oxford Textbook of Surgery*, Morris and Malt, Eds., Oxford University Press (1994)). In general, the liquid

will be an aqueous solution. Examples of solutions include Phosphate Buffered Saline (PBS), Celsior™, Perfadex™, Collins solution, citrate solution, and University of Wisconsin (UW) solution (Oxford Textbook of Surgery, Morris and Malt, Eds., Oxford University Press (1994)). In one embodiment of the present invention, the liquid is Ringer's Solution, e.g., lactated Ringer's Solution, or any other liquid that can be used infused into a patient. In another embodiment, the liquid includes blood, e.g., whole blood.

[0060] Any suitable liquid can be saturated to a set concentration of carbon monoxide via gas diffusers. Alternatively, pre-made solutions that have been quality controlled to contain set levels of carbon monoxide can be used. Accurate control of dose can be achieved via measurements with a gas permeable, liquid impermeable membrane connected to a carbon monoxide analyzer. Solutions can be saturated to desired effective concentrations and maintained at these levels.

[0061] Treatment of Patients with Carbon Monoxide Compositions

[0062] A patient can be treated with a carbon monoxide composition by any method known in the art of administering gases and/or liquids to patients. Carbon monoxide compositions can be administered to a patient diagnosed with, or determined to be at risk for, hepatitis. The present invention contemplates the systemic administration of liquid or gaseous carbon monoxide compositions to patients (e.g., by inhalation and/or ingestion), and the topical administration of the compositions to the patient's liver (e.g., by introduction into the abdominal cavity).

[0063] Systemic Delivery of Carbon Monoxide

[0064] Gaseous carbon monoxide compositions can be delivered systemically to a patient, e.g., a patient diagnosed with, or determined to be at risk for hepatitis. Gaseous carbon monoxide compositions are typically administered by inhalation through the mouth or nasal passages to the lungs, where the carbon monoxide is readily absorbed into the patient's bloodstream. The concentration of active compound (CO) utilized in the therapeutic gaseous composition will depend on absorption, distribution, inactivation, and excretion (generally, through respiration) rates of the carbon monoxide as well as other factors known to those of skill in the art. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition. Treatments can be monitored and CO dosages can be adjusted to ensure optimal treatment of the patient. Acute, sub-acute and chronic administration of carbon monoxide are contemplated by the present invention, depending upon, e.g., the severity or persistence of hepatitis in the patient. Carbon monoxide can be delivered to the patient for a time (including indefinitely) sufficient to treat the condition and exert the intended pharmacological or biological effect.

[0065] The following are examples of some methods and devices that can be utilized to administer gaseous carbon monoxide compositions to patients.

[0066] Ventilators

[0067] Medical grade carbon monoxide (concentrations can vary) can be purchased mixed with air or another oxygen-containing gas in a standard tank of compressed gas (e.g., 21% O₂, 79% N₂). It is non-reactive, and the concentrations that are required for the methods of the present invention are well below the combustible range (10% in air). In a hospital setting, the gas presumably will be delivered to the bedside where it will be mixed with oxygen or house air in a blender to a desired concentration in ppm (parts per million). The patient will inhale the gas mixture through a ventilator, which will be set to a flow rate based on patient comfort and needs. This is determined by pulmonary graphics (i.e., respiratory rate, tidal volumes etc.). Fail-safe mechanism(s) to prevent the patient from unnecessarily receiving greater than desired amounts of carbon monoxide can be designed into the delivery system. The patient's carbon monoxide level can be monitored by studying (1) carboxyhemoglobin (COHb), which can be measured in venous blood, and (2) exhaled carbon monoxide collected from a side port of the ventilator. Carbon monoxide exposure can be adjusted based upon the patient's health status and on the basis of the markers. If necessary, carbon monoxide can be washed out of the patient by switching to 100% O₂ inhalation. Carbon monoxide is not metabolized; thus, whatever is inhaled will ultimately be exhaled except for a very small percentage that is converted to CO₂. Carbon monoxide can also be mixed with any level of O₂ to provide therapeutic delivery of carbon monoxide without consequential hypoxic conditions.

[0068] Face Mask and Tent A carbon monoxide-containing gas mixture is prepared as above to allow passive inhalation by the patient using a facemask or tent. The concentration inhaled can be changed and can be washed out by simply switching over to 100% O₂. Monitoring of carbon monoxide levels would occur at or near the mask or tent with a fail-safe mechanism that would prevent too high of a concentration of carbon monoxide from being inhaled.

[0069] Portable Inhaler

[0070] Compressed carbon monoxide can be packaged into a portable inhaler device and inhaled in a metered dose, for example, to permit intermittent treatment of a recipient who is not in a hospital setting. Different concentrations of carbon monoxide could be packaged in the containers. The device could be as simple as a small tank (e.g., under 5 kg) of appropriately diluted CO with an on-off valve and a tube from which the patient takes a whiff of CO according to a standard regimen or as needed.

[0071] Intravenous Artificial Lung

[0072] An artificial lung (a catheter device for gas exchange in the blood) designed for O₂ delivery and CO₂ removal can be used for carbon monoxide delivery. The catheter, when implanted, resides in one of the large veins and would be able to deliver carbon monoxide at given concentrations either for systemic delivery or at a local site. The delivery can be a local delivery of a high concentration of carbon monoxide for a short period of time at the site of the procedure, e.g., in proximity to the liver (this high concentration would rapidly be diluted out in the bloodstream), or a relatively longer exposure to a lower concentration of carbon monoxide (see, e.g., Hattler et al., *Artif. Organs* 18(11):806-812 (1994); and Golob et al., *ASAIO J.*, 47(5):432-437 (2001)).

[0073] Normobaric Chamber

[0074] In certain instances, it would be desirable to expose the whole patient to carbon monoxide. The patient would be inside an airtight chamber that would be flooded with carbon monoxide (at a level that does not endanger the patient, or at a level that poses an acceptable risk without the risk of bystanders being exposed. Upon completion of the exposure, the chamber could be flushed with air (e.g., 21% O₂, 79% N₂) and samples could be analyzed by carbon monoxide analyzers to ensure no carbon monoxide remains before allowing the patient to exit the exposure system.

[0075] Systemic Delivery of Liquid CO Compositions

[0076] The present invention further contemplates that aqueous solutions comprising carbon monoxide can be created for systemic delivery to a patient, e.g., for oral delivery and/or by infusion into the patient, e.g., intravenously, intra-arterially, intraperitoneally, and/or subcutaneously. For example, liquid CO compositions, such as CO-saturated Ringer's Solution, can be infused into a patient suffering from or at risk for hepatitis. Alternatively or in addition, CO-partially or completely saturated whole (or partial) blood can be infused into the patient.

[0077] The present invention also contemplates that agents capable of delivering doses of gaseous CO compositions or liquid CO compositions can be utilized (e.g., CO-releasing gums, creams, ointments, lozenges, or patches).

[0078] Topical Treatment of Organs with Carbon Monoxide

[0079] Alternatively or in addition, carbon monoxide compositions can be applied directly to the liver, e.g., to the entire liver, or to any portion thereof. A gaseous composition can be directly applied to the liver of a patient by any method known in the art for insufflating gases into a patient. For example, gases, e.g., carbon dioxide, are often insufflated into the abdominal cavity of patients to facilitate examination during laproscopic procedures (see, e.g., Oxford Textbook of Surgery, Morris and Malt, Eds., Oxford University Press (1994)). The skilled practitioner will appreciate that similar procedures could be used to administer carbon monoxide compositions directly to the liver of a patient.

[0080] Aqueous carbon monoxide compositions can also be administered topically to the liver of a patient. Aqueous forms of the compositions can be administered by any method known in the art for administering liquids to patients. As with gaseous compositions, aqueous compositions can be applied directly to the liver. For example, liquids, e.g., saline solutions containing dissolved CO, can be injected into the abdominal cavity of patients during laproscopic procedures. The skilled practitioner will appreciate that similar procedures could be used to administer liquid carbon monoxide compositions directly to the liver of a patient. Further, an in situ exposure can be carried out by flushing the liver or a portion thereof with a liquid carbon monoxide composition (see Oxford Textbook of Surgery, Morris and Malt, Eds., Oxford University Press (1994)).

[0081] Use of Hemoxygenase-1, Other Compounds, and Other Treatments for Hepatitis

[0082] Also contemplated by the present invention is the induction or expression of hemoxygenase-1 (HO-1) in

conjunction with administration of CO. For example, HO-1 can be induced in a patient suffering from or at risk for hepatitis. As used herein, the term "induce(d)" means to cause increased production of a protein, e.g., HO-1, in isolated cells or the cells of a tissue, organ or animal using the cells' own endogenous (e.g., non-recombinant) gene that encodes the protein.

[0083] HO-1 can be induced in a patient by any method known in the art. For example, production of HO-1 can be induced by hemin, by iron protoporphyrin, or by cobalt protoporphyrin. A variety of non-heme agents including heavy metals, cytokines, hormones, NO, COCl₂, endotoxin and heat shock are also strong inducers of HO-1 expression (Choi et al., *Am. J. Respir. Cell Mol. Biol.* 15:9-19, 1996; Maines, *Annu. Rev. Pharmacol. Toxicol.* 37:517-554, 1997; and Tenhunen et al., *J. Lab. Clin. Med.* 75:410-421, 1970). HO-1 is also highly induced by a variety of agents causing oxidative stress, including hydrogen peroxide, glutathione depletors, UV irradiation, endotoxin and hyperoxia (Choi et al., *Am. J. Respir. Cell Mol. Biol.* 15:9-19, 1996; Maines, *Annu. Rev. Pharmacol. Toxicol.* 37:517-554, 1997; and Keyse et al., *Proc. Natl. Acad. Sci. USA* 86:99-103, 1989). A "pharmaceutical composition comprising an inducer of HO-1" means a pharmaceutical composition containing any agent capable of inducing HO-1 in a patient, e.g., any of the agents described above, e.g., NO, hemin, iron protoporphyrin, and/or cobalt protoporphyrin.

[0084] HO-1 expression in a cell can be increased via gene transfer. As used herein, the term "express(ed)" means to cause increased production of a protein, e.g., HO-1 or ferritin, in isolated cells or the cells of a tissue, organ or animal using an exogenously administered gene (e.g., a recombinant gene). The HO-1 or ferritin is preferably of the same species (e.g., human, mouse, rat, etc.) as the recipient, in order to minimize any immune reaction. Expression could be driven by a constitutive promoter (e.g., cytomegalovirus promoters) or a tissue-specific promoter (e.g., milk whey promoter for mammary cells or albumin promoter for liver cells). An appropriate gene therapy vector (e.g., retrovirus, adenovirus, adeno associated virus (AAV), pox (e.g., vaccinia) virus, human immunodeficiency virus (HIV), the minute virus of mice, hepatitis B virus, influenza virus, Herpes Simplex Virus-1, and lentivirus) encoding HO-1 or ferritin would be administered to a patient suffering from or at risk for hepatitis, by mouth, by inhalation, or by injection into the liver. Similarly, plasmid vectors encoding HO-1 or apoferritin can be administered, e.g., as naked DNA, in liposomes, or in microparticles.

[0085] Further, exogenous HO-1 protein can be directly administered to a patient by any method known in the art. Exogenous HO-1 can be directly administered in addition, or as an alternative, to the induction or expression of HO-1 in the patient as described above. The HO-1 protein can be delivered to a patient, for example, in liposomes, and/or as a fusion protein, e.g., as a TAT-fusion protein (see, e.g., Becker-Hapak et al., *Methods* 24:247-256, 2001).

[0086] Alternatively or in addition, any of the products of metabolism by HO-1, e.g., bilirubin, biliverdin, iron, and/or ferritin, can be administered to a patient in conjunction with CO in order to prevent or treat hepatitis. Further, the present invention contemplates that iron-binding molecules other than ferritin, e.g., desferoxamine (DFO), iron dextran, and/

or apoferritin, can be administered to the patient. Further still, the present invention contemplates that enzymes (e.g., biliverdin reductase) that catalyze the breakdown any of these products can be inhibited to create/enhance the desired effect. Any of the above can be administered, e.g., orally, intravenously, intraperitoneally, or by direct administration to the liver.

[0087] The present invention contemplates that compounds that release CO into the body after administration of the compound (e.g., CO-releasing compounds, e.g., photo-activatable CO-releasing compounds), e.g., dimanganese decacarbonyl, tricarbonyldichlororuthenium (II) dimer, and methylene chloride (e.g., at a dose of between 400 to 600 mg/kg, e.g., about 500 mg/kg), can also be used in the methods of the present invention, as can carboxyhemoglobin and CO-donating hemoglobin substitutes.

[0088] The above can be administered to a patient in any way, e.g., by oral, intraperitoneal, intravenous, or intraarterial administration. Any of the above compounds can be administered to the patient locally and/or systemically, and in any combination.

[0089] The present invention further contemplates treating/preventing hepatitis by administering CO to the patient in combination with any other known methods or compounds for treating hepatitis, e.g., cessation or reducing administration of causative drugs; administering corticosteroids and/or α -interferon or other antiviral agents to the patient; and/or performing surgery on the patient, e.g., liver transplantation.

[0090] The invention is illustrated in part by the following examples, which are not to be taken as limiting the invention in any way.

EXAMPLE 1

Carbon Monoxide Attenuates Liver Injury

[0091] Animals

[0092] Male C57BL/6J (Charles Rivers Laboratories, Bar Harbor, Me.), 8-12-wk-old *inos^{-/-}* mice and wild type littermates (bred/maintained at the University of Pittsburgh) were used for in vivo experiments.

[0093] Acute Hepatic Injury Models

[0094] Groups of mice were administered TNF- α /D-gal (0.3 μ g/8 mg/mouse, i.p., respectively). Depending on the experimental condition, some mice received CO (250 ppm), the selective NO donor O₂-vinyl 1-(pyrrolidin-1-yl) diazen-1-ium-1,2-diolate (V-PYRRO; 10 mg/kg subcutaneously (s.c.), Alexis Biochem., San Diego, Calif.) or cobalt protoporphyrin (CoPP, 5 mg/kg, intraperitoneally (i.p.), Frontier Scientific, Logan, Utah). Additionally, the selective inhibitor of iNOS L-N⁶-(1-iminoethyl)-lysine-dihydrochloride (L-NIL; 5 mg/kg, i.p., Alexis Biochemicals) or the HO-1 inhibitor tin protoporphyrin (SnPP; 50 μ mol/kg, i.p., Frontier Scientific) was administered when specified. Where indicated, acetaminophen (Sigma Chem. Co.; St Louis, Mo.) was administered (500 mg/kg, i.p.).

[0095] Hepatocyte Cell Culture.

[0096] Mouse primary hepatocytes were harvested from C57BL/6J, *mkk3^{-/-}*, *inos^{-/-}* (in-house breeding colony), or

hmox-1^{-/-} mice as described in Kim et al. (J. Biol. Chem. 272: 1402-1411 (1997)). Hepatocytes were used on days 1-3 following harvest.

[0097] Induction of Hepatocyte Death/Apoptosis

[0098] Cells were treated with TNF- α (10 ng/ml) and actinomycin-D (Act-D; 200 ng/ml, Sigma Chemical Co. St. Louis, Mo.) to induce cell death. TNF- α /ActD treatment has been demonstrated to induce cell death, specifically apoptosis, in primary hepatocytes (see, e.g., Kim et al. (J. Biol. Chem. 272: 1402-1411 (1997)). Hepatocytes were treated with CO, the NO donor *s*-nitroso-N-acetyl-penicillamine (SNAP; 250-750 μ M), and/or additional pharmacologic agents where indicated. Twelve hours after TNF- α /ActD treatment, cells were washed and stained with crystal violet to determine viability as previously described (Id.). Where indicated, the selective in vitro inhibitor of iNOS, L-N⁵-(1-iminoethyl)-ornithine-2HCl (LNIO; 1-2 MM; Calbiochem, San Diego, Calif.) was administered.

[0099] Gene Transfer/Plasmids.

[0100] In some experiments, gene transfer of an I κ B α superrepressor (Hellerbrand et al., Hepatology 27:1285-1295 (1998)) or β -galactosidase using adenoviral vectors (10 pfu/cell) was performed 12 hours prior to TNF- α /ActD treatment. NF- κ B activation was evaluated using a luciferase reporter assay as described in Chow et al. (J. Biol. Chem. 274: 10689-10692 (1999)). Briefly, hepatocytes were co-transfected with NF- κ B reporter constructs (pGL3-kappa β luciferase, 100 ng/well; and pIEP-Lac-z 0.5 μ g/well) using Lipofectin™ (Invitrogen, Carlsbad, Calif.) as instructed by the manufacturer. Evaluation of iNOS expression was performed using a luciferase reporter assay as described in Lowenstein et al. (Proc. Natl. Acad. Sci. U.S.A 90: 9730-9734 (1993)). Briefly, hepatocytes were co-transfected with iNOS promoter reporter constructs (pXP2; 1 μ g/well) and pIEP-LacZ (0.5 μ g/well) as described above.

[0101] Luciferase Reporter Assays

[0102] Hepatocytes were transfected with plasmids as described above and treated with various stimuli 24 hours after transfection. Luciferase activity (reported as arbitrary units; A.U.) was assayed 6 hours after initiation of treatment, using a luciferase assay kit (Promega, Madison, Wis.) and a Berthold Luminometer. Results were corrected for transfection efficiency and protein concentration.

[0103] Electrophoretic Mobility Shift Assay

[0104] Nuclei were extracted from hepatocytes following treatment. A double-stranded DNA NF- κ B consensus sequence (GGGGACTTTCCTCC (SEQ ID NO: 1)); Santa Cruz Biotechnology, Santa Cruz, Calif.) was labelled with [δ -³²P]-ATP and incubated with 5 mg of total nuclear protein. Some incubations were performed in the presence of antibodies against p65/RelA or p50 (Santa Cruz Biotech) to evaluate for supershift. Electrophoretic mobility shift assay (EMSA) were performed as described in Taylor et al. (J. Biol. Chem. 273:15148-15156 (1998)).

[0105] Immunoblot Analysis

[0106] Western blot analysis was performed on primary hepatocytes in culture or from liver homogenates with antibodies to iNOS (Transduction Laboratories, Lexington, Ky.; 1:1000), HO-1 (Calbiochem; 1:2000), or β -actin

(Sigma Chemical; 1:5000). Thirty μg protein in cell culture experiments or 100 μg protein from liver homogenates was loaded per well for SDS-PAGE.

[0107] Histology/Immunohistochemistry

[0108] For histology and immunohistochemistry, livers were fixed in 2% paraformaldehyde and then snap frozen in liquid nitrogen. Livers were then sectioned (7 microns thick) and stained with hematoxylin and eosin (H&E). Liver sections were also stained for TUNEL and activated caspase-3 using kits according to the manufacturer's instructions (Promega). Sections for iNOS immunocytochemistry were blocked with 5% goat serum containing 0.2% bovine serum albumin. Thereafter, sections were incubated for 1 hour at room temperature with anti-iNOS antibody (Transduction Laboratories; 1:300), then washed and probed with a secondary antibody conjugated to Alexa-488 (Molecular Probes, Eugene, Oreg.). Nuclei were stained with Hoechst dye. Images were acquired using an Olympus Provus microscope. Hepatocytes in culture were plated on gelatinized coverslips, stimulated as indicated, and then fixed in 2% paraformaldehyde containing 0.1% Triton X-100. Blocking and staining was similar to liver sections except anti-p65/RelA antibody (Santa Cruz Biotechnology; 1:350) was utilized.

[0109] CO Exposure

[0110] The animals were exposed to CO at a concentration of 250 ppm. Briefly, 1% CO in air was mixed with air (21% oxygen) in a stainless steel mixing cylinder and then directed into a 3.70 ft³ glass exposure chamber at a flow rate of 12 L/min. A CO analyzer (InterScan, Chatsworth, Calif.) was used to measure CO levels continuously in the chamber. CO concentrations were maintained at 250 ppm at all times. Mice were placed in the exposure chamber as required.

[0111] HO-1 Protects Against Liver Injury.

[0112] Whether HO-1 is protective against acute hepatic failure was investigated. The results are presented in FIG. 1. Cobalt protoporphyrin (5 mg/kg, i.p.) was administered to male C57BL/6J mice. Twenty-four hours later, TNF- α /D-gal (0.3 μg /8 mg/mouse, i.p., respectively) was administered to the mice. Serum alanine aminotransferase (ALT) levels in the mice were measured 8 hours after administration of TNF- α /D-gal. Induction of HO-1 prevented liver injury as measured by serum ALT levels.

[0113] Exogenous CO Protects Hepatocytes

[0114] Whether exogenous CO is protective against hepatocyte cell death in vitro was investigated. The results are presented in FIGS. 2 and 3. To generate the data presented in FIG. 2, mouse hepatocytes were pre-incubated with CO (250 ppm) for 1 hr (standard pre-treatment time for all experiments) prior to addition of TNF- α /Act-D (10 ng/200 ng/ml respectively). Cells were maintained in CO for the duration of the experiment. Twelve hours afterward, cell viability was measured as described in Kim et al. (J. Biol. Chem. 272: 1402-1411 (1997)). Adenoviral experiments involved incubating hepatocytes overnight with 10 pfu/cell of the adenovirus prior to addition of TNF- α /ActD, and then assaying for viability using crystal violet. The roles of signaling molecules guanylyl cyclase and p38 MAPK were also investigated in this model. To evaluate the role of cGMP and confirm the role of NF- κ B, hepatocytes were treated

separately with the soluble guanylate cyclase (sGC) inhibitor 1H-[1,2,4]Oxadiazolo[4,3-a]quinoxalin-1-one (ODQ; Calbiochem; 2-10 μM) or the NF- κ B inhibitor BAY 11-7082, (10 μM). Cells were treated with the inhibitors for 1 hour prior to the 1 hour pretreatment with CO. TNF- α /ActD was then added and the cells tested for viability 12 hours later. NF- κ B activation was critical to the protection elicited by CO while cGMP was not involved. Exposure to CO led to significantly less cell death (*p<0.01) than without CO.

[0115] To generate the data presented in FIG. 3, human primary hepatocytes obtained from a donor liver resection were treated with CO and TNF- α /ActD as described above.

[0116] Exposure of primary mouse, rat, and human hepatocytes to CO inhibited TNF- α induced apoptosis. Inhibition of hepatocyte apoptosis was independent of cGMP generation, as the selective guanylyl cyclase inhibitor ODQ did not reverse the protection provided by CO (FIG. 2). Additionally, CO treatment inhibited cell death both in the presence of SB203580 (3-30 μM , Calbiochem), a selective inhibitor of p38 MAPK activation, and in hepatocytes from mkk3^{-/-} mice, the dominant upstream kinase for p38 (data not shown). Thus, the effects of CO were independent of the cGMP/p38 MAPK pathway. In these experiments, hepatocytes were pre-treated with CO for one hour prior to addition of TNF- α /ActD to the medium. If CO treatment was initiated after addition of TNF- α , less protection was observed (data not shown).

[0117] The Role of NF- κ B in CO Protection

[0118] Whether CO-induced protection of hepatocytes depends upon NF- κ B was investigated. FIGS. 4, 5, and 6A-6C present data illustrating that CO induced an increase in NF- κ B nuclear translocation and DNA binding in mouse hepatocytes as measured by NF- κ B luciferase reporter assay activity, EMSA, and immunostaining for RelA/p65 nuclear translocation, respectively.

[0119] To generate the data presented in FIG. 4, evaluation of NF- κ B activation was performed using a luciferase reporter assay as described in Chow et al. (J. Biol. Chem. 274: 10689-10692 (1999)). Briefly, hepatocytes were co-transfected with NF- κ B reporter constructs and pIEP-Lac-z 24 hr prior to addition of BAY 11-7082 (10 μM) or vehicle. Cells were incubated for 1 hr prior to CO (250 ppm). Luciferase activity (reported as arbitrary units; A.U.) was assayed 6 hr after exposure to CO or a cytokine mixture (CM) composed of TNF- α (500 U/ml), IL-1 β (100 U/ml), and IFN- δ (100 U/ml), which was used as a positive control for NF- κ B activation. Results were corrected for transfection efficiency and protein concentration.

[0120] To generate the data in FIG. 5, NF- κ B DNA binding was evaluated using EMSA in hepatocytes treated with CO (250 ppm). Note the time-dependent increase in NF- κ B binding (total) with expression peaking at one hr (Lanes 1, 4, 7). Extracts were then supershifted to identify the different NF- κ B dimers using antibodies against p50 (Lanes 2, 5, 8) and p65 (Lanes 3, 6, 9).

[0121] To generate the data in FIGS. 6A-6C, primary hepatocytes were immunostained for nuclear p65 localization following exposure to 1 hr CO (250 ppm). Images depict nuclear translocation of NF- κ B (arrows pointing to green nuclei that depict the translocation of NF- κ B) in both

CM (used as a positive control) and CO-treated cells versus no localization in air treated cells (arrows pointing to blue nuclei).

[0122] NF- κ B luciferase reporter assay activity peaked one hour after placing cells in the CO atmosphere. A cytokine mixture (CM) was included in the treatment groups as a positive signal as well as a standard for maximum reporter activity by which to evaluate the effects of CO. Transfection efficiency in primary hepatocytes is difficult, but the reporter activity was very significant (* $p < 0.001$ versus control). These data combined with the positive immunostaining and EMSA results support the notion that CO induces a moderate increase in NF- κ B that in itself may in part result in selective gene expression. To evaluate whether NF- κ B activity is needed for protection mediated by CO, adenoviral gene transfer of I κ B α was utilized to prevent NF- κ B translocation and BAY 11-7082 (1-10 mM, Calbiochem) was used to inhibit NF- κ B activation. The protective effects of CO were abrogated by inhibition of NF- κ B activation.

[0123] The Role of NF- κ B-Dependent iNOS Expression in CO Protection

[0124] Whether CO-mediated protection of hepatocytes requires expression of iNOS and generation of NO was investigated. The results are presented in FIGS. 7, 8, and 9.

[0125] To generate the data in FIG. 7, evaluation of iNOS expression was performed using a luciferase reporter assay as described in Lowenstein et al. (Proc. Natl. Acad. Sci. U.S.A 90: 9730-9734 (1993)). Briefly, hepatocytes were co-transfected with an iNOS promoter reporter construct and pIEP-LacZ 24 hr prior to exposure to BAY 11-7082 (10 μ M) or vehicle. Cells were incubated with BAY 1 hr prior to exposure to CO (250 ppm). Luciferase activity (reported as arbitrary units; A.U.) was assayed as above. Cytokine mixture (CM; see above) was used as a positive control to induce iNOS expression, and results were corrected for transfection efficiency and protein concentration.

[0126] To generate the data in FIG. 8, expression of iNOS protein was evaluated using immunoblotting techniques. Briefly, cell extracts from hepatocytes were treated with TNF- α /ActD for 6-8 hr in the presence and absence of CO (250 ppm). Control cells received air or CO alone. Note in FIG. 8 that TNF- α induces iNOS expression minimally, while those cells treated with TNF- α in the presence of CO show a significantly greater induction in iNOS protein.

[0127] To generate the data presented in FIG. 9, mouse hepatocytes were isolated from *inos^{-/-}* or from wild type C57BL/6J mice, which were then pre-treated for 1 hr with L-NIO (1 mM) to inhibit iNOS prior to CO administration. Those groups exposed to CO received a one-hour pretreatment prior to addition of TNF- α /ActD and were then returned to CO exposure. CO did not provide protection against cell death, as evaluated via crystal violet exclusion 12 hr later, in cells where iNOS expression was absent or inhibited.

[0128] Exposure of hepatocytes to CO produced a highly significant increase in activity in an iNOS luciferase reporter assay (FIG. 7). Again, a cytokine mixture was used as both a positive control in these low efficiency transfections and as a standard by which to evaluate the effects of CO. Consistent with the NF- κ B dependence of iNOS expression, decreased

reporter activity was observed in hepatocytes treated with BAY 11-7082 (FIG. 7). Additionally, iNOS protein was markedly increased in response to TNF- α in the presence of CO compared to TNF- α alone (FIG. 8). Using hepatocytes from iNOS knockout mice (*inos^{-/-}*) and wild type hepatocytes treated with the selective iNOS inhibitor L-NIO (1 mM, Calbiochem), applicants investigated whether CO could protect against TNF- α -induced death in the absence of iNOS activity. Hepatocytes lacking iNOS activity were not protected by CO from TNF- α -induced cell death while wild type hepatocytes were protected (FIG. 9). Taken together, these data show that CO requires NF- κ B activation and iNOS expression to protect hepatocytes from cell death in vitro.

[0129] Inhaled CO is Protective Against Liver Failure

[0130] Whether inhaled CO protects mice against liver injury in a TNF- α /D-gal model of fulminant hepatic failure was investigated. The results are presented in FIGS. 10 and 11A-11H.

[0131] To generate the data presented in FIG. 10, mice were pre-treated with CO (250 ppm) for one hour prior to receiving TNF- α /D-gal (0.3 μ g/8 mg/mouse; i.p., respectively). After receiving TNF- α /D-gal, mice were returned to the CO exposure chamber and their serum was analyzed for ALT levels 6-8 hr later. Without exposure to CO, liver failure occurred in 6-8 hr driven primarily by apoptosis of hepatocytes as in the in vitro model described above. Serum ALT in mice treated with CO was 74% lower than in air-exposed mice.

[0132] To generate the data presented in FIGS. 11A-11H, liver samples from mice treated with TNF- α /D-gal in the presence and absence of CO (250 ppm) for 8 hr were sectioned and stained for hematoxylin & eosin (H&E), activated caspase 3 (as indicated by an increase in red intensity), and for TUNEL positive cells (as demarcated by the increased green cellular staining; a marker of cell death). Nuclei stained blue. Exposure to CO markedly reduced TNF- α /D-gal-induced liver damage as assessed by H&E staining. Livers from mice exposed to CO also displayed fewer TUNEL positive cells, displayed less staining of activated caspase-3, and had normal architecture. Air-exposed control mice that received TNF- α /D-gal showed marked hepatic inflammation, edema, hemorrhage and loss of architecture.

[0133] Results discussed above were confirmed using lipopolysaccharide (LPS, also referred to as endotoxin) in place of TNF. In these confirmatory studies, LPS/D-Gal administration resulted in an increase in serum ALT levels from a control level of 20+/-5 IU/ml to >1000 IU/ml, as measured 8 hours following LPS/D-Gal administration. In mice pretreated with 250 ppm CO, the increase in ALT was reduced by >75%, to 250+/-75 IU/ml. To further characterize the effects observed with CO in this model, serum interleukin-6 was measured, and found to be reduced 65% in animals breathing CO vs air-breathing controls (data not shown). Tissue histopathology of the livers from these mice was similar to that demonstrated using TNF/D-Gal. Untreated and CO-treated mice (no LPS/D-Gal) had no signs of injury while those treated with air and LPS/D-Gal showed marked injury including edema, hemorrhage, neutrophil infiltration and an overall destruction of normal morphology and architecture. In contrast, livers from mice

treated with CO and LPS/D-Gal were protected to the same extent as mice treated with CO and TNF/D-Gal. Few changes in the markers of inflammation (edema, hemorrhage, neutrophil infiltration) were observed. Architecture was maintained and appeared grossly similar to untreated and CO (in the absence of LPS/D-Gal)-treated mice. Overall, the use of LPS/D-Gal to induce acute hepatitis paralleled and confirmed data generated using TNF/D-Gal treatment.

[0134] The Role of iNOS in CO Protection Against Liver Damage

[0135] Whether hepatic iNOS protein levels were increased in the livers of CO-exposed mice after treatment with TNF- α /D-gal was investigated using immunoblotting techniques and immunohistochemistry. Further, whether CO would protect *inos^{-/-}* mice or wild type mice treated with the selective iNOS inhibitor L-NIL (10 mg/kg, i.p.; dosed every 2 hours) was investigated to determine whether iNOS expression has a functional role. The results are provided in FIGS. 12, 13A-13D, and 14.

[0136] To generate the data presented in FIG. 12, male C57BL/6J mice were treated with air or CO (250 ppm) 1 hr prior to TNF- α /D-gal (0.3 μ g/8 mg/mouse, i.p., respectively) administration. Six hours later, livers were harvested to evaluate iNOS expression by immunoblotting. Results show that iNOS expression was increased modestly in air/TNF- α /D-gal-treated mice, but was markedly increased in mice treated with TNF- α /D-gal and CO. As expected, *inos^{-/-}* mice showed no expression of iNOS protein.

[0137] To generate the data in FIGS. 13A-13D, mouse liver sections were immunostained for iNOS expression. The liver sections were obtained from mice treated with TNF- α /D-gal in the presence or absence of CO, and from air and CO controls that received no TNF- α /D-gal. Livers from mice exposed to CO and not receiving TNF- α /D-gal displayed a modest increase in iNOS expression. However, a significantly greater increase in expression (indicated by an increase in green-stained cells) was observed in livers from mice that were exposed to CO and received TNF- α /D-gal. The increased expression appeared to be localized around blood vessels.

[0138] To generate the data in FIG. 14, the efficacy of CO-induced protection was tested in the absence of iNOS activity using *inos^{-/-}* and wild type mice that were treated with L-NIL, the selective inhibitor of iNOS (L-NIL; 5 mg/kg, i.p. dosed every two hours). L-NIL was administered 2 hr prior to CO. CO-treated animals were then pre-treated (250 ppm) for 1 hr prior to TNF- α /D-gal. In the absence of iNOS function/expression, CO is unable to protect against liver damage as assessed by serum ALT levels and histopathology (data not shown).

[0139] Thus, it appears the protective effect of inhaled CO in TNF- α -induced liver failure is dependent upon iNOS activity.

[0140] The Role of HO-1 in CO Protection Against Acute Liver Failure

[0141] Whether CO and NO exert protection against acute liver failure through an HO-1-dependent mechanism was investigated. The data are presented in FIGS. 15, 16, 17, and 18.

[0142] To generate the data presented in FIG. 15, immunoblotting was performed to observe HO-1 expression in the livers of mice that received TNF- α /D-gal in the presence and absence of CO (250 ppm). CO-treated mice showed a significant increase in HO-1 expression in both the presence and absence of TNF- α /D-gal.

[0143] To assess the role of iNOS on TNF- α /D-gal-induced HO-1 expression in the liver (data presented in FIG. 16), mice were administered L-NIL (5 mg/kg, i.p.) 2 hr prior to pre-treatment with CO (250 ppm) and every 2 hr thereafter. Control mice received L-NIL and remained in room air. Note in FIG. 16 that CO increased HO-1 expression in vehicle-treated mice, but was unable to induce expression when iNOS was inhibited. L-NIL treatment alone had a minimal effect on HO-1 expression.

[0144] To test the protective role of CO-induced HO-1 (data presented in FIG. 17), mice were given SnPP (50 μ mol/kg, s.c.), the selective inhibitor of HO-1, 5 hr prior to CO. Alternatively, the mice were given VPYRRO (VP), an NO donor (10 mg/kg, s.c.). VP was selectively designed to deliver NO directly to the liver. One hour after the initial VP dose, the animals were exposed to CO for 1 hr prior to administration of TNF- α /D-gal (see above). Serum ALT levels were determined 6-8 hr later. Note that CO was not able to provide protection in animals where HO-1 activity was blocked. VP, when administered 2 hr prior and then every 2 hr thereafter, provided protection against injury as determined 8 hour later by serum ALT measurements.

[0145] To generate the data presented in FIG. 18, wild type C57BL/6J mice were pretreated for 24 hr with L-NIL in the drinking water (4.5 mM) as described in Stenger et al. (J. Exp. Med. 183: 1501-1514 (1996)). These mice and *inos^{-/-}* mice were then administered CoPP. L-NIL was maintained in the water throughout the experiment. Control and *inos^{-/-}* mice received normal drinking water. Twenty-four hr after administration of CoPP, TNF- α /D-gal was administered and serum ALT determined 6-8 hr later. Note in FIG. 18 that induction of HO-1 provides protection regardless of the presence of iNOS.

[0146] Immunoblotting of liver extracts from mice treated with CO in the presence or absence of TNF- α /D-gal showed up-regulation of HO-1 (FIG. 15). The addition of the iNOS inhibitor L-NIL to these above groups, which abrogated the protection (FIG. 17), also prevented up-regulation of HO-1 (FIG. 16). To determine whether HO-1 was central to CO-elicited hepatoprotection, tin protoporphyrin-IX (SnPP, 50 μ mol/kg, s.c., Frontier Scientific) was used as a selective inhibitor of HO-1 activity. SnPP significantly diminished the protective effects of CO in this model (FIG. 17). SnPP administration in the absence of TNF- α /D-gal had no deleterious or protective effects (data not shown). These results suggest that up-regulation of HO-1 is important to the protective effects of CO.

[0147] To determine if up-regulation of HO-1 would also be needed if protection was initiated by NO, mice were treated with the pharmacological NO donor V-PYRRO/NO. This agent is metabolized by the liver, resulting in release of NO by hepatocytes. V-PYRRO/NO also provides protection following LPS/D-gal or TNF- α /D-gal administration. Mice were randomized and treated with TNF- α /D-gal with or without SnPP to evaluate the role of HO-1. V-PYRRO/NO was protective, as assayed by serum ALT. However, SnPP

abrogated the ability of this NO donor to protect against liver damage (FIG. 17). Thus, it appears that CO- or NO-initiated hepatoprotection is at least partially dependent on HO-1.

[0148] Because these data suggest that CO and NO require HO-1 activity to protect against TNF- α -induced hepatocyte death, whether protection mediated by HO-1 requires iNOS activity was investigated. Using *inos^{-/-}* mice, HO-1 was induced via administration of CoPP. TNF- α /D-gal was injected 24 hr thereafter, at the peak of HO-1 expression, and liver damage was assessed 6-8 hr later. The results show that induction of HO-1 was able to significantly prevent liver injury independently of iNOS activity with a >50% reduction in serum ALT (FIG. 18). These results were confirmed using L-NIL. Mice were pre-treated with drinking water containing L-NIL (4.5 mM) for 24 hours. This method effectively inhibits NOS activity. Control mice received normal water. Subsequently, CoPP was administered to induce HO-1 expression and 24 hours thereafter mice were challenged with TNF- α /D-gal. L-NIL treatment alone did not change the severity of injury induced in this model. All animals receiving CoPP (with and without L-NIL) were protected from liver injury (FIG. 18).

[0149] Whether HO-1 expression is required for CO- or NO-induced protection from TNF- α /ActD-induced hepatocyte cell death was investigated. The data are presented in FIGS. 19 and 20.

[0150] To generate the data presented in FIG. 19, mouse hepatocytes were isolated from HO-1 null mice (*hmox-1^{-/-}*) and wild type (C57BL/6J) littermates, pretreated for 1 hour with CO (250 ppm), and treated with TNF- α /ActD. Viability was assayed as described above. CO significantly protected wild type hepatocytes, but was unable to protect hepatocytes isolated from *hmox-1^{-/-}* mice.

[0151] To generate the data presented in FIG. 20, mouse hepatocytes were isolated from HO-1 null mice (*hmox-1^{-/-}*) and wild type (C57BL/6J) littermates, pretreated with the NO donor SNAP (500 μ M), and then treated with TNF- α /ActD 1 hour later. SNAP has been demonstrated to protect hepatocytes in this model. SNAP significantly protected against cell death in wild type hepatocytes but did not provide significant protection against cell death in hepatocytes isolated from *hmox-1^{-/-}* mice.

[0152] As discussed above, air-treated wild type and *hmox-1^{-/-}* cells exposed to TNF- α /ActD underwent cell death as expected, while CO- or NO-treated wild type cells were protected in the presence of TNF- α /ActD (FIGS. 19 and 20). The protection conferred by CO and NO was lost in cells lacking functional HO-1 (*hmox-1^{-/-}*). Thus, it appears that HO-1 can provide protection in this model without the involvement of iNOS, suggesting that HO-1 or one or more of its catalytic products can, in part, exert cytoprotective effects in this model.

[0153] Inhaled CO is Protective Against Acetaminophen-Induced Hepatitis

[0154] Whether inhaled CO is protective against acetaminophen (APAP)-induced hepatitis was investigated. The data are presented in FIG. 21.

[0155] To generate the data in FIG. 21, Male C57BL/6J mice were exposed to CO (250 ppm) either 1 hr prior or 4 hr post administration of APAP (500 mg/kg, i.p.). The mice

were then maintained in CO for the duration of the experiment. Serum ALT levels were determined 20 hr after APAP administration. Control mice received APAP and were maintained in air. This protocol was designed to allow hepatitis to develop for four hours before administering CO. CO significantly reduced damage to the liver as assessed by serum ALT (622 \pm 44 vs 175 \pm 137, $p < 0.01$ as compared to controls). This protection was similar to that observed in a separate group of animals that had been pre-treated with CO prior to APAP. These data support the therapeutic use of CO in a clinically relevant situation where treatment would begin after the initiation of hepatitis.

[0156] The results discussed in this Example demonstrate that a low concentration of CO can protect against TNF- α /D-gal-induced fulminant hepatitis and illustrate a unique and previously unrecognized dependence on both HO-1 and iNOS in the CO-induced protection of livers from damage by TNF- α /D-gal.

[0157] Without intending to be bound by theory, it is possible that CO mediated protection operates by activating NF- κ B, which in the presence of an inflammatory stimulus leads to the up-regulation of iNOS with the consequent production of NO. In addition to the induction of iNOS, other NF- κ B dependent antiapoptotic/protective genes may be induced. During the 1 hour pre-treatment with CO and before the cells are treated with TNF- α , significant activation of NF- κ B was present, which could be part of the priming of the cellular apparatus discussed above. The activation of NF- κ B by CO may in part result from a mild increase in reactive oxygen species generation originating from the mitochondria (preliminary observations). One hour might also permit time for expression of NF- κ B-dependent anti-apoptotic genes. The next step in such a hypothetical model might lead to NO production following the up-regulation of iNOS. NO leads to up-regulation of HO-1, the activity of which confers protective effects. The protective effect of HO-1 could be due to removal of heme or to any one or more of its three products: CO, biliverdin/bilirubin or iron/ferritin. Given that exogenous CO was administered throughout the duration of the experiments, it appears unlikely that endogenously-produced CO alone mediates HO-1 protection. However, the combination of CO with other products of HO-1 or these other products acting individually might be involved.

[0158] In a study described above, CO was administered in a clinically-relevant model of acetaminophen (APAP)-induced hepatitis that has a time course that is similar to the development of acute hepatitis in humans. The data demonstrate that exposure to CO 4 hours after administration of APAP (500 mg/kg, i.p.) resulted in a 62% reduction in liver injury (FIG. 21). In this model of APAP-induced liver injury, mice show signs of hepatitis as early as 2-4 hours after APAP administration and lethality occurs by 24-48 hours. Thus, CO was administered after the initiation of liver injury. Consistent with the data in the APAP model are the results in a murine model of hemorrhagic shock where the therapeutic initiation of inhaled CO during resuscitation following a 2.5 hour shock phase resulted in protection against liver injury (>65% reduction in serum ALT at 24 hr $p < 0.01$; $n = 6-10$ /group).

[0159] In summary, employing a model of liver injury driven principally by TNF- α -induced apoptosis, the follow-

ing was demonstrated: first, inhaled CO can prevent hepatitis in this model; second, protection by CO requires generation of a second gaseous molecule, NO; third, NO exerts its beneficial effects, at least in part, via upregulation of HO-1; and fourth, up-regulation of HO-1 is protective without a need for iNOS/NO activity, i.e., without an obligate continuation of the cycle.

EXAMPLE 2

Protocol for the Treatment of Hepatitis

[0160] The following example illustrates protocols for use in treating a patient diagnosed as suffering from hepatitis. The example also illustrates protocols for treating patients before, during, and/or after surgical procedures, e.g., a surgical procedure to transplant a liver. Skilled practitioners will appreciate that any protocol described herein can be adapted based on a patient's individual needs, and can be adapted to be used in conjunction with any other treatment for hepatitis.

[0161] Treatment of Patients

[0162] Treatment of a patient with CO can begin on the day the patient is diagnosed as suffering from hepatitis, for example, hepatitis caused by viral infection and/or alcohol abuse. The patient can be diagnosed by a physician using any art-known method. For example, a physician may make such a diagnosis using data obtained from blood tests, e.g., tests to determine serum ALT levels and tests to determine whether a patient is infected with a particular virus (e.g., any known hepatitis virus). Further, a physician may consider a patient's medical history in making such a diagnosis (e.g., by considering whether a patient is an alcoholic or a chronic drug user). The patient can inhale CO at concentration of about 250 to 500 ppm for one hour per day. This treatment can continue for about 30 days, or until the patient is diagnosed as no longer having or being at risk for hepatitis.

[0163] Liver Transplant Procedures

[0164] Treatment of a Liver Donor

[0165] Prior to harvesting a liver or portion thereof, the donor can be treated with inhaled carbon monoxide (250 ppm) for one hour. Treatment can be administered at doses varying from 10 ppm to 1000 ppm for times varying from one hour to six hours, or for the entire period from the moment when it becomes possible to treat a brain-dead (cadaver) donor to the time the organ is removed. For a human donor, treatment should start as soon as possible following the declaration that brain death is present. In some applications, it may be desirable to begin treatment before brain death.

[0166] For non-human animals (e.g., pigs) to be used as xenotransplantation donors, the live donor animal can be treated with relatively high levels of inhaled carbon monoxide, as desired, so long as the carboxyhemoglobin so produced does not compromise the viability and function of the organ to be transplanted. For example, one could use levels greater than 500 ppm (e.g., 1000 ppm or higher, and up to 10,000 ppm, particularly for brief times).

[0167] Treatment of the Liver In Situ

[0168] Before a liver is harvested from a donor, it can be flushed or perfused with a solution, e.g., a buffer or medium,

while it is still in the donor. The intent is to flush the liver with a solution saturated with carbon monoxide and maintained in a carbon monoxide atmosphere so that the carbon monoxide content remains at saturation. Flushing can take place for a time period of at least 10 minutes, e.g., 1 hour, several hours, or longer. The solution should ideally deliver the highest concentration of carbon monoxide possible to the cells of the liver (or portion thereof).

[0169] Treatment of the Liver Ex Vivo

[0170] A liver can be preserved in a medium that includes carbon monoxide from the time it is removed from the donor to the time it is transplanted to the recipient. This can be performed by maintaining the liver in the medium comprising CO, or by perfusing it with such a medium. Since this occurs ex vivo rather than in an animal, very high concentrations of CO gas can be used (e.g., 10,000 ppm) to keep the medium saturated with CO.

[0171] Treatment of a Liver Recipient

[0172] Treatment of the recipient with CO can begin on the day of transplantation at least 30 minutes before surgery begins. Alternatively, it could begin at least 30 minutes before reperfusion of the organ in the recipient. It can be continued for at least 30 minutes, e.g., 1 hour. Carbon monoxide doses between 10 ppm and 3000 ppm can be delivered for varying times, e.g., minutes or hours, and can be administered on the day of and on days following transplantation. For example, the patient can inhale a concentration of carbon monoxide, e.g., 3000 ppm, for three consecutive 10 second breath holds. Alternatively, a lower concentration of the gas can be delivered intermittently or constantly, for a longer period of time, with regular breathing rather than breath holding. Carboxyhemoglobin concentrations can be utilized as a guide for appropriate administration of carbon monoxide to a patient. Usually, treatments for recipients should not raise carboxyhemoglobin levels above those considered to pose an acceptable risk for a patient in need of a transplant.

[0173] A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, other embodiments are within the scope of the following claims.

What is claimed is:

1. A method of treating hepatitis in a patient, comprising:
 - identifying a patient diagnosed as suffering from hepatitis; and
 - administering to the patient a pharmaceutical composition comprising an amount of carbon monoxide effective to treat hepatitis in the patient.
2. The method of claim 1, wherein the pharmaceutical composition is in gaseous form and is administered to the patient by inhalation.
3. The method of claim 1, wherein the pharmaceutical composition is in liquid form and is administered to the patient orally.
4. The method of claim 1, wherein the pharmaceutical composition is administered directly to the abdominal cavity of the patient.
5. The method of claim 1, wherein the patient is infected with a virus selected from the group consisting of: hepatitis

A virus; hepatitis B virus; hepatitis C virus; hepatitis D virus; hepatitis E virus; and hepatitis G virus.

6. The method of claim 1, wherein the patient is an alcoholic.

7. The method of claim 1, further comprising administering to the patient a treatment selected from the group consisting of: withholding or reducing administration of hepatitis-inducing drugs; and administering corticosteroids or antiviral agents to the patient.

8. The method of claim 1, wherein the pharmaceutical composition is administered by artificial lung.

9. The method of claim 1, wherein the pharmaceutical composition is administered by an extracorporeal membrane gas exchange device.

10. The method of claim 1, wherein the hepatitis is caused by exposure to a hepatotoxic agent.

11. A method of treating hepatitis in a patient, comprising:

- (a) identifying a patient suffering from or at risk for hepatitis;
- (b) providing a vessel containing a pressurized gas comprising carbon monoxide gas;
- (c) releasing the pressurized gas from the vessel, to form an atmosphere comprising carbon monoxide gas; and
- (d) exposing the patient to the atmosphere, wherein the amount of carbon monoxide in the atmosphere is sufficient to treat hepatitis in the patient.

12. A method of administering a hepatotoxic drug to a patient, comprising:

- (a) administering the hepatotoxic drug to the patient; and
- (b) before, during, or after step (a), administering to the patient a pharmaceutical composition comprising carbon monoxide in an amount effective to treat hepatitis in the patient.

13. The method of claim 12, wherein carbon monoxide is administered before step (a).

14. The method of claim 12, wherein carbon monoxide is administered during step (a).

15. The method of claim 12, wherein carbon monoxide is administered after step (a).

16. The method of claim 12, wherein the hepatotoxic drug is selected from the group consisting of: isoniazid, methyl-dopa, acetaminophen, amiodarone, and nitrofurantoin.

17. A vessel comprising medical grade compressed carbon monoxide gas, the vessel bearing a label indicating that the gas can be used to treat hepatitis in a patient.

18. The vessel of claim 17, wherein the carbon monoxide gas is in admixture with an oxygen-containing gas.

19. The vessel of claim 17, wherein the carbon monoxide gas is present in the admixture at a concentration of at least about 0.025%.

20. The vessel of claim 17, wherein the carbon monoxide gas is present in the admixture at a concentration of at least about 0.05%.

21. The vessel of claim 17, wherein the carbon monoxide gas is present in the admixture at a concentration of at least about 0.10%.

22. The vessel of claim 17, wherein the carbon monoxide gas is present in the admixture at a concentration of at least about 1.0%.

23. The vessel of claim 17, wherein the carbon monoxide gas is present in the admixture at a concentration of at least about 2.0%.

24. The vessel of claim 17, wherein the label further indicates that the gas can be administered to a patient in conjunction with administration of a hepatotoxic drug.

25. A vessel comprising medical grade compressed carbon monoxide gas, the vessel bearing a label indicating that the gas can be administered to a patient in conjunction with administration of a hepatotoxic drug.

26. A method of treating hepatitis in a patient, the method comprising:

identifying a patient suffering from or at risk for hepatitis not caused by surgery or endotoxin; and

administering to the patient a pharmaceutical composition comprising an amount of carbon monoxide effective to treat hepatitis in the patient.

* * * * *