




(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) **Date de dépôt PCT/PCT Filing Date:** 2022/03/21
(87) **Date publication PCT/PCT Publication Date:** 2022/09/29
(85) **Entrée phase nationale/National Entry:** 2023/08/03
(86) **N° demande PCT/PCT Application No.:** US 2022/021080
(87) **N° publication PCT/PCT Publication No.:** 2022/203984
(30) **Priorité/Priority:** 2021/03/22 (US63/164,135)

(51) **Cl.Int./Int.Cl.** **C07D 471/04** (2006.01),
A61K 31/4745 (2006.01)
(71) **Demandeur/Applicant:**
NEUROCRINE BIOSCIENCES, INC., US
(72) **Inventeurs/Inventors:**
HARRIOTT, NICOLE, US;
PAGANO, NICHOLAS, US;
BOON, BYRON A., US
(74) **Agent:** GOWLING WLG (CANADA) LLP

(54) **Titre : INHIBITEURS DE VMAT2 ET PROCEDES D'UTILISATION**
(54) **Title: VMAT2 INHIBITORS AND METHODS OF USE**



(57) **Abrégé/Abstract:**

This disclosure relates generally to VMAT2 inhibitor compounds of formula (I), compositions, and methods related thereto.

Date Submitted: 2023/08/03

CA App. No.: 3207452

Abstract:

This disclosure relates generally to VMAT2 inhibitor compounds of formula (I), compositions, and methods related thereto.

VMAT2 INHIBITORS AND METHODS OF USE

BACKGROUND

Technical Field

5 This disclosure relates generally to VMAT2 inhibitor compounds, compositions, and methods related thereto.

Description of the Related Technology

10 Dysregulation of dopaminergic systems is integral to several central nervous system (CNS) disorders, including neurological and psychiatric diseases and disorders. These neurological and psychiatric diseases and disorders include hyperkinetic movement disorders, and conditions, such as, schizophrenia and mood disorders. The transporter protein vesicular monoamine transporter-2 (VMAT2) plays an important role in presynaptic dopamine release and regulates monoamine uptake from the cytoplasm to the synaptic vesicle for storage and release.

15 (±)-Tetrabenazine ((±)-TBZ), has been used as a drug for decades. (±)-TBZ is reported as a potent, reversible inhibitor of catecholamine uptake by VMAT2 ($IC_{50} = 3.2$ nM) (see, *e.g.*, Scherman *et al.*, Proc. Natl. Acad. Sci. USA, (1983) 80:584-8) and is currently used in the treatment of various hyperkinetic disorders. Inhibition of VMAT2 by (±)-TBZ results in depletion of brain monoamines *in vivo* (see, *e.g.*, Pettibone *et al.*, Eur. J. Pharmacol. (1984) 102:431-6). (±)-TBZ also inhibits
20 presynaptic and postsynaptic dopamine receptors in rat brain (see, *e.g.*, Login *et al.*, (1982) Ann. Neurology 12:257-62; Reches *et al.*, J. Pharmacol. Exp. Ther. (1983) 225:515-521). (±)-TBZ exhibits extensive first pass metabolism following oral administration to humans with little or no (±)-TBZ observed in systemic circulation. The pharmacological activity of (±)-TBZ is therefore thought to be mediated primarily by active metabolites. (±)-TBZ has two chiral centers and is a racemic mixture of
25 two stereoisomers. (±)-TBZ has been determined to be rapidly and extensively metabolized *in vivo* by carbonyl reductase to four metabolic stereoisomers of 3-isobutyl-9,10-dimethoxy-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-ol, also known as dihydrotetrabenazine (DHTBZ). The inhibitory constants of these four metabolites for VMAT2 have been reported, such as, in WO2008/058261, Example 7. As shown, only two of the four DHTBZ isomers ([+]-alpha-DHTBZ
30 and [+]-beta-DHTBZ) show significant potency as inhibitors of VMAT2.

(±)-TBZ Metabolite	pKi	Ki
[+]-alpha-DHTBZ (2R,3R,11bR)	8.7 ± 0.2 (n=6)	1.9 nM
[+]-beta-DHTBZ (2S,3R,11bR)	7.9 ± 0.1 (n=5)	13 nM
[-]-alpha-DHTBZ (2S,3S,11bS)	6.7 ± 0.1 (n=3)	202 nM
[-]-beta-DHTBZ (2R,3S,11bS)	6.1 ± 0.1 (n=4)	714 nM

In patients administered (±)-TBZ, [-]-alpha-DHTBZ (2S,3S,11bS-DHTBZ) and [+]-beta-DHTBZ (2S,3R,11bR-DHTBZ) were the most abundant DHTBZ isomers, while [-]-beta-DHTBZ

(2*R*,3*S*,11*bS*-DHTBZ) and [+-]-alpha-DHTBZ (2*R*,3*R*,11*bR*-DHTBZ) were present as minor metabolites. The [+-]-alpha-DHTBZ (2*R*,3*R*,11*bR*-DHTBZ) isomer was determined to be present in the least amount of all four isomers. Thus, [+-]-beta-DHTBZ (2*S*,3*R*,11*bR*-DHTBZ) appears to be the major DHTBZ isomer contributing to the pharmacological activity of (±)-TBZ. Once formed, the half-life of [+-]-beta-DHTBZ (2*S*,3*R*,11*bR*-DHTBZ) is relatively short (approximately 5 hours) which requires (±)-TBZ to have a sub-optimal (TID) dosing regimen.

(±)-TBZ has a narrow therapeutic window and its clinical use requires careful dose titration. Side effects associated with (±)-TBZ and/or its metabolites include neuroleptic malignant syndrome, drowsiness, fatigue, nervousness, anxiety, insomnia, agitation, confusion, orthostatic hypotension, nausea, dizziness, sedation, depression, akathisia, and Parkinsonism. Generally speaking, the probability of observing side effects is a function of the achieved plasma concentrations from a given dosing regimen. Compounds with a longer half-life ($t_{1/2}$) and lower clearance will have lower peak-to-trough fluctuations in plasma exposure given an equivalent dosing interval. These longer half-life compounds may exhibit improved tolerability by maintaining drug concentrations at levels needed for efficacy but below levels that may elicit side effects.


A fundamental and effective strategy to improve drug half-life is to reduce clearance. The term clearance describes the process of drug elimination from the body or from a single organ, defined as the volume of fluid cleared of drug from the body per unit of time. Clearance is a fundamental pharmacokinetic parameter and is commonly measured in drug research and development as this parameter impacts drug attributes such as half-life and, ultimately, the dosing regimen. When compound and dose selection is optimized, the benefits of small plasma-concentration fluctuations seen in compounds with low clearance include potentially reduced steady state peak concentrations, increased trough concentrations, and the prospect of improving medication adherence because of a possibly improved risk-benefit profile.

Despite the advances that have been made in this field, a need remains in the art for improved VMAT2 inhibitors, including compounds, compositions, and methods related thereto. The identification of long half-life/low clearance VMAT2 small molecules is advantageous for drug development, particularly when being developed for chronic administration. In certain disease populations where patient compliance and pill burden are an ongoing challenge, reduced dosing frequency is highly desirable and offers increased patient benefit.

The present disclosure fulfills these, such as, improved *in vitro* VMAT2 potency or improved pharmacokinetics, or both, and other needs, as evident in reference to the following disclosure.

SUMMARY

Some embodiments provide a compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

R¹ is **R^{2A}**, $-(CH_2)_m-R^{2B}$, or $-(CH_2)_n-OR^{2C}$;

R^{2A} is C₂-C₄alkenyl, C₁-C₆alkyl, C₃-C₈cycloalkyl, or 4-7 membered heterocyclyl, each
 5 unsubstituted or substituted with one or more **R^A**;

each **R^A** is independently selected from the group consisting of: C₁-C₄alkoxy, C₁-C₄alkyl, C₁-C₄alkylamino, C₁-C₄alkylsulfonyl, C₂-C₄dialkylamino, C₂-C₄dialkylsulfamoyl, halogen, haloC₁-C₄alkoxy, haloC₁-C₄alkyl, haloC₂-C₄dialkylamino, 4-7 membered heterocyclyl, -CN, -OH, and oxo;

R^{2B} is aryl, C₃-C₈cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each
 10 unsubstituted or substituted with one or more **R^B**;

each **R^B** is independently selected from the group consisting of: C₁-C₄alkyl, C₁-C₄alkylsulfonyl, C₁-C₄alkylsulfonyloxy, -CN, -CH₂CN, halogen, haloC₁-C₄alkyl, haloC₁-C₄alkylsulfonyl, oxo, and -OH;

R^{2C} is C₁-C₄alkyl, C₃-C₈cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or
 15 substituted with one or more **R^C**;

each **R^C** is independently -CN or C₃-C₆cycloalkyl;

m is 0, 1, 2, 3, or 4; and

n is 1, 2, or 3.

Also provided herein is a pharmaceutical product comprising a compound of Formula **(I)**, or a
 20 pharmaceutically acceptable salt thereof.

Also provided herein is a pharmaceutical product selected from: a pharmaceutical composition, a formulation, a unit dosage form, and a kit; each comprising a compound of Formula **(I)**, or a pharmaceutically acceptable salt thereof.

Also provided herein is a pharmaceutical product comprising a compound of Formula **(I)**, or a
 25 pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient.

Also provided herein is a pharmaceutical product selected from: a pharmaceutical composition, a formulation, a unit dosage form, and a kit; each comprising a compound of Formula **(I)**, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient.

Also provided herein is a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient.

Also provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt thereof, for use in treating a neurological or psychiatric disease or disorder in a subject in need thereof.

Also provided herein is a method of inhibiting VMAT2 comprising contacting said VMAT2 with a compound of Formula (I).

Also provided herein is a method of reducing the level of monoamines in the central nervous system of a subject comprising administering to the subject an amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, in an amount sufficient to lower the level of monoamines in the central nervous system relative to the level prior to administration.

Also provided herein is a method of treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder in a subject in need thereof, comprising administering to the subject a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical product comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; or a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

Also provided herein is a method of treating a neurological or psychiatric disease or disorder in a subject in need thereof, comprising administering to the subject a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical product comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; or a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

Also provided herein is a method of treating a neurological or psychiatric disease or disorder in a subject comprising administering a compound of Formula (I), or a pharmaceutically acceptable salt thereof, to the subject.

Also provided herein is a method of treating a neurological or psychiatric disease or disorder in a subject in need thereof comprising administering a compound of Formula (I), or a pharmaceutically acceptable salt thereof, to the subject in combination with a further pharmaceutical agent selected from an antidepressant, an antipsychotic (typical or atypical), an antiepileptic, an antimicrobial, an antiarrhythmic, a mood stabilizer, and a gastrointestinal drug.

Also provided herein is a method of ameliorating one or more symptoms of a neurological or psychiatric disease or disorder, comprising administering to a subject in need thereof an effective amount of at least one compound of Formula (I), or a pharmaceutically acceptable salt thereof; wherein the neurological or psychiatric disease or disorder is selected from the group consisting of: hyperkinetic movement disorder, schizophrenia, schizoaffective disorder, a mood disorder, treatment-refractory obsessive-compulsive disorder, neurological dysfunction associated with Lesch-Nyhan

syndrome, agitation associated with Alzheimer's disease, Fragile X syndrome or Fragile X-associated tremor-ataxia syndrome, autism spectrum disorder, Rett syndrome, and chorea-acanthocytosis.

Also provided herein is the use of a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical product comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; in the manufacture of a medicament for treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder.

Also provided herein is the use of a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical product comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; in the manufacture of a medicament for treating a neurological or psychiatric disease or disorder.

Also provided herein is the use of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a neurological or psychiatric disease or disorder.

Also provided herein is the use of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, for treating a neurological or psychiatric disease or disorder, wherein the neurological or psychiatric disease or disorder is selected from the group consisting of: hyperkinetic movement disorder, schizophrenia, schizoaffective disorder, a mood disorder, treatment-refractory obsessive-compulsive disorder, neurological dysfunction associated with Lesch-Nyhan syndrome, agitation associated with Alzheimer's disease, Fragile X syndrome or Fragile X-associated tremor-ataxia syndrome, autism spectrum disorder, Rett syndrome, and chorea-acanthocytosis.

Also provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical product comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; for use in a method of treatment of the human or animal body by therapy.

Also provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt thereof, for use in a method of treatment of the human or animal body by therapy.

Also provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical product comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; for use in a method for treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder.

Also provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical product comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof; a pharmaceutical composition comprising a compound of Formula (I), or a

pharmaceutically acceptable salt thereof; for use in a method for treating a neurological or psychiatric disease or disorder.

Also provided herein is a compound of Formula (I), or a pharmaceutically acceptable salt thereof, for treating a neurological or psychiatric disease or disorder.

In some embodiments, the neurological or psychiatric disease or disorder is selected from the group consisting of: hyperkinetic movement disorder, schizophrenia, schizoaffective disorder, a mood disorder, treatment-refractory obsessive-compulsive disorder, neurological dysfunction associated with Lesch-Nyhan syndrome, agitation associated with Alzheimer's disease, Fragile X syndrome or Fragile X-associated tremor-ataxia syndrome, autism spectrum disorder, Rett syndrome, and chorea-acanthocytosis.

In some embodiments, the vesicular monoamine transporter-2 (VMAT2) disease or disorder is selected from: an ataxias or spinal muscular atrophy; a chorea; a congenital malformation, deformation, or abnormality; a dementia; an oral cavity, salivary gland, or jaw disease; a dyskinesia; a dystonia; an endocrine, nutritional, or metabolic disease; an epilepsy; a habit or impulse disorder; a Huntington's disease or related disorder; a mood or psychotic disorder; a neurotic, stress-related, and somatoform disorder; a degenerative disease of the basal ganglia; an extrapyramidal and movement disorder; a neurological or psychiatric disease or disorder; a nervous system or motor function disorder; a Parkinson's/parkinsonism disorder; a pediatric-onset behavioral and emotional disorder; a pervasive developmental disorder; and a substance abuse or dependence disorder.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A shows a representation of the single crystal structure of (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*S*,11*bS*)**).

FIG. 1B shows a representation of the single crystal structure of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*R*,11*bR*)**).

FIG. 2 shows a powder X-ray diffraction (PXRD) diffractogram of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-1**) free base crystalline Form I.

FIG. 3 shows a DSC and TGA thermogram of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-1**) free base crystalline Form I.

FIG. 4 shows a powder X-ray diffraction (PXRD) diffractogram of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-15**) free base crystalline Form I.

FIG. 5 shows a DSC and TGA thermogram of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-15**) free base crystalline Form I.

FIG. 6 shows a general synthetic scheme for the preparation of Compounds of Formula (I) and intermediates related thereto, wherein R¹ has the same meaning as described herein and X is a leaving group, such as, halo (*e.g.* I, Cl, and Br), or OS(O)₂R^z (*e.g.* mesylate (-OMs), triflate (-OTf), and tosylate (-OTs), wherein R^z has the same meaning as described herein). As shown in **FIG. 6**, intermediates, **Compound 1-B**, **Compound 1-C**, and **Compound 1-D** (including the corresponding stereoisomers thereof), are novel and useful in the preparation of Compounds of Formula (I), and pharmaceutical salts thereof, as described herein.

FIG. 7 shows a general synthetic scheme for the preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*R*,11*bR*)**) utilizing a resolution step of 9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 1-C**) with (2*S*,3*S*)-2,3-*bis*(4-methylbenzoyloxy)butanedioic acid (DPTTA) to provide (2*R*,3*R*,11*bR*)-9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 1-C (2*R*,3*R*,11*bR*)**) that is subsequently deprotected to provide **Compound 1-D (2*R*,3*R*,11*bR*)**.

FIG. 8 shows a representative reaction for the preparation of the salt of (2*R*,3*R*,11*bR*)-9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 1-C (2*R*,3*R*,11*bR*)**) and (2*S*,3*S*)-2,3-*bis*(4-methylbenzoyloxy)butanedioic acid (DPTTA), see **Example 1B**.

FIG. 9 shows a general synthetic scheme for the preparation of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*R*,11*bR*)**) and (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*S*,11*bS*)**) using chiral supercritical fluid chromatography (SFC), see **Example 2**.

FIG. 10 shows a representative reduction step as described in **Example 3** for the preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*R*,11*bR*)**) from (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*R*,11*bR*)**); and a representative reduction step for the preparation of (2*S*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*S*,3*S*,11*bS*)**) from (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*S*,11*bS*)**).

FIG. 11 shows a representative reduction step as described in **Example 4** for the preparation of (2*S*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*S*,3*R*,11*bR*)**) from (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*R*,11*bR*)**); and a representative reduction step for the preparation of (2*R*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*S*,11*bS*)**) from (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*S*,11*bS*)**).

FIG. 12 shows a representative reaction used in the preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-1**) using (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*R*,11*bR*)**) and 2,2,2-trifluoroethyl trifluoromethanesulfonate (CF₃CH₂OTf) in the presence of cesium carbonate and acetone, see **Example 5A**. Also shown in **FIG. 12** is a representative reaction used in the preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-55**) using (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*R*,11*bR*)**) and (*R*)-2-methyloxirane in the presence of cesium carbonate and DMF, see **Example 5F**.

FIG. 13 shows a representative reaction scheme used in the preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-1**), see **Example 7**; and a representative reaction scheme used in the preparation of (2*S*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-3**), see **Example 8**; both reactions utilize dimethyl sulfate as the methylating agent and sodium borohydride as the reducing agent.

FIG. 14 shows a representative reduction reaction used in the preparation of (2*S*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-2**), see **Example 9**; and a representative reduction reaction used in the preparation of (2*R*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-4**), see **Example 10**; both reactions utilize lithium tri-*sec*-butylborohydride as the reducing agent in THF.

FIG. 15 shows the VMAT2 K_i values (nM) between compounds provided herein (*i.e.*, **Compound 4-1**, **Compound 4-13**, and **Compound 4-14**) and comparator compounds (*i.e.*, **Compound 4-1C**, **Compound 4-13C**, and **Compound 4-14C**).

FIG. 16 shows the VMAT2 K_i values (nM) between compounds provided herein (*i.e.*, **Compound 4-15**, **Compound 4-17**, and **Compound 4-18**) and comparator compounds (*i.e.*, **Compound 4-15C**, **Compound 4-17C**, and **Compound 4-18C**).

FIG. 17 shows the VMAT2 K_i values (nM) between compounds provided herein (*i.e.*, **Compound 4-22**, **Compound 4-26**, and **Compound 4-28**) and comparator compounds (*i.e.*, **Compound 4-22C**, **Compound 4-26C**, and **Compound 4-28C**).

FIG. 18 shows the VMAT2 K_i values (nM) between compounds provided herein (*i.e.*, **Compound 4-30**, **Compound 4-32**, and **Compound 4-38**) and comparator compounds (*i.e.*, **Compound 4-30C**, **Compound 4-32C**, and **Compound 4-38C**).

FIG. 19 shows the VMAT2 K_i values (nM) between compounds provided herein (*i.e.*, **Compound 4-39** and **Compound 5-1**) and comparator compounds (*i.e.*, **Compound 4-39C** and **Compound 5-1C**).

FIG. 20 shows the VMAT2 K_i values (nM) between compounds provided herein (*i.e.*, **Compound 4-27**, **Compound 4-88**, and **Compound 4-119**) and comparator compounds (*i.e.*, **Compound 4-27C**, **Compound 4-88C**, and **Compound 4-119C**).

FIG. 21 shows the VMAT2 K_i values (nM) between compounds provided herein (*i.e.*, **Compound 4-120** and **Compound 4-121**) and comparator compounds (*i.e.*, **Compound 4-120C** and **Compound 4-121C**).

FIG. 22 shows *in vivo* pharmacology in Sprague Dawley rats for **Compound 4-1** and **Compound 4-15** in the Open Field Hypolocomotion model as described in **Example 12**.

FIG. 23 shows *in vivo* pharmacology in Sprague Dawley rats for **Compound 4-27**, **Compound 4-50**, and **Compound 4-55** in the Open Field Hypolocomotion model as described in **Example 12**.

FIG. 24 shows *in vivo* pharmacology in Sprague Dawley rats for **Compound 4-31** in the Open Field Hypolocomotion model as described in **Example 12**.

DETAILED DESCRIPTION

DEFINITIONS

For clarity and consistency, the following definitions will be used throughout this patent document.

As used herein, “about” means $\pm 20\%$ of the stated value, and includes more specifically values of $\pm 10\%$, $\pm 5\%$, $\pm 2\%$, and $\pm 1\%$ of the stated value.

As used herein, “administering” refers to providing a compound described herein or other therapy to a subject in a form that can be introduced into that subject’s body in a therapeutically useful form and therapeutically useful amount, including, but not limited to: oral dosage forms, such as, tablets, capsules, syrups, suspensions, and the like; injectable dosage forms, such as, IV, IM, IP, and the like; transdermal dosage forms, including creams, jellies, powders, and patches; buccal dosage forms; inhalation powders, sprays, suspensions, and the like; and rectal suppositories.

A health care practitioner can directly provide a compound described herein to a subject in the form of a sample or can indirectly provide a compound to a subject by providing an oral or written

prescription for the compound. Also, for example, a subject can obtain a compound by themselves without the involvement of a health care practitioner. When the compound is administered to the subject, the body is transformed by the compound in some way. When a compound described herein is provided in combination with one or more other agents, "administration" is understood to include the compound and other agents are administered at the same time or at different times. When the agents of a combination are administered at the same time, they can be administered together in a single composition, or they can be administered separately. The preferred method of administration can vary depending on various factors, *e.g.*, the components of the pharmaceutical formulation, the site of the disease, and the severity of the disease.

The term "composition" refers to a compound or crystalline form thereof, including but not limited to, salts, solvates, and hydrates of a compound described herein, in combination with at least one additional component, such as, a composition obtained/prepared during synthesis, preformulation, in-process testing (*e.g.*, TLC, HPLC, NMR samples), and the like.

The term, "compound" as used herein is meant to include all stereoisomers, geometric isomers, tautomers, and isotopic variants of the structures depicted. The term is also meant to refer to compounds described herein, regardless of how they are prepared, *e.g.*, synthetically, through biological process (*e.g.*, metabolism or enzyme conversion), or a combination thereof. All compounds, and pharmaceutically acceptable salts thereof, can be found together with other substances, such as, water and solvents (*e.g.*, hydrates and solvates) or can be isolated. When in the solid state, the compounds described herein and salts thereof can occur in various forms and can, *e.g.*, take the form of co-crystals or solvates, including hydrates. The compounds can be in any solid-state form, such as, a polymorph or solvate, so unless clearly indicated otherwise, reference in the specification to compounds and salts thereof should be understood as encompassing any solid-state form of the compound. In some embodiments, the compounds described herein, or salts thereof, are substantially isolated. By "substantially isolated" is meant that the compound is at least partially or substantially separated from the environment in which it was formed or detected. Partial separation can include, *e.g.*, a composition enriched in the compounds described herein. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compounds described herein, or salts thereof.

The term "solvate" as used herein refers to a solid-state form of a compound described herein, or a pharmaceutically acceptable salt thereof which includes a stoichiometric or non-stoichiometric amount of a solvent bound by non-covalent intermolecular forces. When the solvent is water, the solvate is a hydrate.

The term "hydrate" as used herein refers to a compound described herein or a salt thereof that further includes a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.

The term “in need of treatment” and the term “in need thereof” when referring to treatment are used interchangeably to mean a judgment made by a caregiver (*e.g.* physician, nurse, nurse practitioner, *etc.* in the case of humans; veterinarian in the case of animals, including non-human mammals) that a subject or animal requires or will benefit from treatment. This judgment is made
5 based on a variety of factors that are in the realm of a caregiver's expertise, but that includes the knowledge that the subject or animal is ill, or will become ill, as the result of a disease, condition or disorder that is treatable by the compound described herein. Accordingly, the compound described herein can be used in a protective or preventive manner; or compound described herein can be used to alleviate, inhibit, or ameliorate the disease, condition, or disorder.

10 The term “subject” refers to any animal, including mammals, such as, mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, primates, and humans. In the context of a clinical trial or screening or activity experiment the subject can be a healthy volunteer or healthy participant without an underlying VMAT2 mediated disorder or condition or a volunteer or participant that has received a diagnosis for a disorder or condition in need of medical treatment as determined by
15 a health care professional. In the context outside of a clinical trial a subject under the care of a health care professional who has received a diagnosis for a disorder or condition is typically described as a subject.

The term “pediatric subject” refers to a subject under the age of 21 years at the time of diagnosis or treatment. The term “pediatric” can be further divided into various subpopulations
20 including: neonates (from birth through the first month of life); infants (1 month up to two years of age); children (two years of age up to 12 years of age); and adolescents (12 years of age through 21 years of age (up to, but not including, the twenty-second birthday)) see *e.g.*, Berhman *et al.*, Textbook of Pediatrics, 15th Ed. Philadelphia: W.B. Saunders Company, 1996; Rudolph *et al.*, Rudolph's Pediatrics, 21st Ed. New York: McGraw-Hill, 2002; and Avery *et al.*, Pediatric Medicine, 2nd Ed.
25 Baltimore: Williams & Wilkins; 1994.

The phrase “pharmaceutically acceptable” refers to compounds (and salts thereof), compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk
30 ratio.

The term “pharmaceutical composition” refers to a specific composition comprising at least one active ingredient; including but not limited to, salts, solvates, and hydrates of compounds described herein, whereby the composition is amenable to investigation for a specified, efficacious outcome in a mammal (for example, without limitation, a human). Those of ordinary skill in the art
35 will understand and appreciate the techniques appropriate for determining whether an active ingredient has a desired efficacious outcome based upon the needs of the artisan.

The terms “prevent”, “preventing”, and “prevention” refer to the elimination or reduction of the occurrence or onset of one or more symptoms associated with a particular disorder. For example, the terms “prevent”, “preventing”, and “prevention” can refer to the administration of therapy on a prophylactic or preventative basis to a subject who may ultimately manifest at least one symptom of a disorder but who has not yet done so. Such subjects can be identified on the basis of risk factors that are known to correlate with the subsequent occurrence of the disease, such as, the presence of a biomarker. Alternatively, prevention therapy can be administered as a prophylactic measure without prior identification of a risk factor. Delaying the onset of the at least one episode and/or symptom of a disorder can also be considered prevention or prophylaxis. In some embodiments, the subject can be a pediatric subject.

The terms “treat”, “treating”, and “treatment” refer to medical management of a disease, disorder, or condition of a subject (*e.g.*, subject) (see, *e.g.*, Stedman’s Medical Dictionary). In general, an appropriate dose and treatment regimen provide the VMAT2 inhibitor in an amount sufficient to provide therapeutic benefit. Therapeutic benefit for subjects to whom the VMAT2 inhibitor compound(s) described herein are administered, includes, for example, an improved clinical outcome, wherein the object is to prevent or slow or retard (lessen) an undesired physiological change associated with the disease, or to prevent or slow or retard (lessen) the expansion or severity of such disease. The effectiveness of one or more VMAT2 inhibitors can include beneficial or desired clinical results that comprise, but are not limited to, abatement, lessening, or alleviation of symptoms that result from or are associated with the disease to be treated; decreased occurrence of symptoms; improved quality of life; longer disease-free status (*i.e.*, decreasing the likelihood or the propensity that a subject will present symptoms on the basis of which a diagnosis of a disease is made); diminishment of extent of disease; stabilized (*i.e.*, not worsening) state of disease; delay or slowing of disease progression; amelioration or palliation of the disease state; and remission (whether partial or total), whether detectable or undetectable; and/or overall survival. In some embodiments, the subject can be a pediatric subject.

The term “therapeutically effective amount” refers to the amount of the compound described herein, or a pharmaceutically acceptable salt thereof, or an amount of a pharmaceutical composition comprising the compound described herein or a pharmaceutically acceptable salt thereof, that elicits the biological or medicinal response in a tissue, system, animal, or human that is being sought by a subject, researcher, veterinarian, medical doctor, or other clinician or caregiver, which can include one or more of the following:

(1) preventing the disorder, for example, preventing a disease, condition, or disorder in a subject who can be predisposed to the disease, condition, or disorder but does not yet experience or display the relevant pathology or symptomatology;

(2) inhibiting the disorder, for example, inhibiting a disease, condition, or disorder in a subject who is experiencing or displaying the relevant pathology or symptomatology (*i.e.*, arresting further development of the pathology and/or symptomatology); and

(3) ameliorating the disorder, for example, ameliorating a disease, condition, or disorder in a subject who is experiencing or displaying the relevant pathology or symptomatology (*i.e.*, reversing the pathology and/or symptomatology).

As used herein, the term “contacting” refers to the bringing together of indicated moieties in an *in vitro* system or an *in vivo* system. For example, “contacting” the VMAT2 protein with a compound provided herein includes the administration of a compound provided herein (or a pharmaceutically acceptable salt thereof) to a subject, such as, a human, having a VMAT2 protein, as well as, for example, introducing a compound provided herein into a sample containing a cellular or purified preparation containing the VMAT2 protein.

CHEMICAL GROUPS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this disclosure belongs. All patents, applications, published applications, and other publications are incorporated by reference in their entirety. In the event that there is a plurality of definitions for a term herein, those in this section prevail unless stated otherwise.

The term “n-membered” where n is an integer typically describes the number of ring-forming atoms in a moiety where the number of ring-forming atoms is n. For example, piperidinyl is an example of a 6-membered heterocyclyl ring, pyrazolyl is an example of a 5-membered heteroaryl ring, pyridyl is an example of a 6-membered heteroaryl ring, and 1,2,3,4-tetrahydro-naphthalene is an example of a 10-membered cycloalkyl group.

For compounds of Formula (I), and pharmaceutically acceptable salts thereof, in which a variable appears more than once, each variable can be a different moiety independently selected from the group defining the variable. For example, where a structure is described having, for example, two R groups that are simultaneously present on the same compound, the two R groups can represent different moieties independently selected from the group defined for R, or the two R groups can be the same.

Whenever a group is described as being “substituted” that group can be substituted with one or more of the indicated substituents. Likewise, when a group is described as being “unsubstituted or substituted” if substituted, the substituent(s) can be selected from one or more of the indicated substituents. It is to be understood that substitution at a given atom is limited by valency.

As used herein, “C_a-C_b” in which “a” and “b” are integers refer to the number of carbon atoms in, for example, an alkyl, alkenyl, or alkynyl group, or the number of carbon atoms in the ring of a cycloalkyl, cycloalkenyl, or aryl group. That is, these groups can contain from “a” to “b”, inclusive,

carbon atoms. Thus, for example, a “C₁-C₄alkyl” group refers to all alkyl groups having from 1 to 4 carbons, that is, methyl (CH₃-), ethyl (CH₃CH₂-), *n*-propyl (CH₃CH₂CH₂-), *iso*-propyl ((CH₃)₂CH-), *n*-butyl (CH₃CH₂CH₂CH₂-), *iso*-butyl ((CH₃)₂CHCH₂-), *sec*-butyl (CH₃CH₂CH(CH₃)-), and *tert*-butyl ((CH₃)₃C-). If no “a” and “b” are designated with regard to an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, or aryl group, the broadest range described in these definitions is to be assumed.

In addition to the foregoing, as used in the specification and appended claims, unless specified to the contrary, the following terms have the meaning indicated:

The term “alkenyl” refers to an alkyl group that contains in the straight or branched hydrocarbon chain one or more double bonds. Examples of alkenyl groups include allenyl, vinylmethyl, and ethenyl. Examples of alkenyl groups include: vinyl (CH₂=CH-), prop-1-en-2-yl (CH₂=C(Me)-), prop-1-en-1-yl (MeCH=CH-), allyl (CH₂=CHCH₂-), but-2-en-2-yl (CH₃CH=C(Me)-), but-1-en-1-yl (MeCH₂CH=CH-), but-1-en-2-yl (CH₂=C(Et)-), 2-methylprop-1-en-1-yl (Me₂C=CH-), but-3-en-2-yl (CH₂=CHCH(Me)-), 2-methylallyl (CH₂=C(Me)CH₂-), but-2-en-1-yl (MeCH=CHCH₂-), but-3-en-1-yl (CH₂=CHCH₂CH₂-), and buta-1,3-dien-1-yl (CH₂=CHCH=CH-). In some embodiments, an alkenyl group can be unsubstituted or substituted. In some embodiments, the alkenyl group can have 2 to 6 carbon atoms. In some embodiments, the alkenyl group can have 2 to 4 carbon atoms. The alkenyl group of the compounds can be designated as “C₂-C₆alkenyl”, “C₂-C₄alkenyl”, or similar designations.

The term “alkoxy” refers to the formula —OR wherein R is an alkyl defined herein. A non-limiting list of alkoxys are methoxy, ethoxy, *n*-propoxy, 1-methylethoxy (*iso*-propoxy), *n*-butoxy, *iso*-butoxy, *sec*-butoxy, and *tert*-butoxy. The alkoxy group can be “C₁-C₆alkoxy”, “C₁-C₄alkoxy”, or similar designations. In some embodiments, an alkoxy can be substituted or unsubstituted.

The term “alkyl” refers to a fully saturated straight or branched hydrocarbon radical. The alkyl group can have 1 to 20 carbon atoms (whenever it appears herein, a numerical range, such as, “1 to 20” refers to each integer in the given range; *e.g.*, “1 to 20 carbon atoms” means that the alkyl group can consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc., up to and including 20 carbon atoms. In some embodiments, the alkyl group can have 1 to 6 carbons (*i.e.*, “C₁-C₆alkyl”). Some embodiments are 1 to 5 carbons (*i.e.*, C₁-C₅alkyl), some embodiments are 1 to 4 carbons (*i.e.*, C₁-C₄alkyl), some embodiments are 1 to 3 carbons (*i.e.*, C₁-C₃alkyl), and some embodiments are 1 or 2 carbons. By way of example only, “C₁-C₄alkyl” indicates that there are one to four carbon atoms in the alkyl chain, *i.e.*, the alkyl chain is selected from methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, and *tert*-butyl. Examples of an alkyl group include: methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, pentyl, *iso*-pentyl, *tert*-pentyl, *neo*-pentyl, 1-methylbutyl [*i.e.*, -CH(CH₃)CH₂CH₂CH₃], 2-methylbutyl [*i.e.*, -CH₂CH(CH₃)CH₂CH₃], *n*-hexyl and the like. When one or more substituents are present on the alkyl group, the substituent(s) can be bonded at any available carbon atom. In some embodiments, an alkyl group can be substituted or unsubstituted.

The term “alkylamino” refers to an amino group (-NH_2) where the nitrogen atom is substituted with one alkyl group. The term “alkyl” has the same definition as described herein. The “alkylamino” group can have 1 to 4 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_4$ alkylamino). Examples for an alkylamino group include,

- 5 $\text{-NH(CH}_3\text{)}$, $\text{-NH(CH}_2\text{CH}_3\text{)}$, $\text{-NH(CH}_2\text{CH}_2\text{CH}_3\text{)}$, and the like.

The term “alkynyl” refers to an alkyl group that contains in the straight or branched hydrocarbon chain one or more triple bonds. Examples of alkynyls include ethynyl and propynyl. An alkynyl group can be unsubstituted or substituted. In some embodiments, an alkynyl group can be unsubstituted or substituted. In some embodiments, the alkynyl group can have 2 to 6 carbon atoms.

- 10 The alkenyl group of the compounds can be designated as “ $\text{C}_2\text{-C}_6$ alkynyl” or similar designations.

The term “alkylsulfonyl” refers to a radical consisting of an alkyl radical bonded to the sulfur of a sulfone radical of the formula: $\text{-S(=O)}_2\text{-}$ wherein alkyl has the same definition as described herein. The “alkylsulfonyl” group can have 1 to 4 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_4$ alkylsulfonyl). Examples include methylsulfonyl, ethylsulfonyl, *n*-propylsulfonyl, *iso*-propylsulfonyl, *n*-butylsulfonyl, *sec*-butylsulfonyl, *iso*-butylsulfonyl, *t*-butylsulfonyl, and the like.

- 15 The term “alkylsulfonyloxy” refers to a radical consisting of an alkyl radical bonded to the sulfur of a sulfonyloxy radical of the formula: $\text{-S(=O)}_2\text{O-}$ wherein alkyl has the same definition as described herein. The “alkylsulfonyloxy” group can have 1 to 4 carbon atoms (*i.e.*, $\text{C}_1\text{-C}_4$ alkylsulfonyloxy). Examples include methylsulfonyloxy ($\text{CH}_3\text{S(=O)}_2\text{O-}$), ethylsulfonyloxy, *n*-propylsulfonyloxy, *iso*-propylsulfonyloxy, *n*-butylsulfonyloxy, *sec*-butylsulfonyloxy, *iso*-butylsulfonyloxy, *t*-butylsulfonyloxy, and the like.

The term “amino” refers to the group -NH_2 .

- The term “aryl” refers to an aromatic ring system containing 6, 10, or 14 carbon atoms that can contain a single ring, two fused rings or three fused rings, such as, phenyl, naphthalenyl, and phenanthrenyl. In some embodiments, the aryl group can have 6 or 10 carbon atoms (*i.e.*, C_6 or C_{10} aryl). In some embodiments, the aryl group is phenyl. When one or more substituents are present on the “aryl” ring, the substituent(s) can be bonded at any available ring carbon. In some embodiments, an aryl group can be substituted or unsubstituted.

- 30 The “C-amido” group refers to a “ $\text{—C(=O)N(R}^{\text{A}}\text{R}^{\text{B}}\text{)}$ ” group that is connected to the rest of the molecule via a carbon atom, and in which R^{A} and R^{B} can be independently hydrogen, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_3\text{-C}_7$ cycloalkyl, $\text{C}_5\text{-C}_8$ cycloalkenyl, C_6 or C_{10} aryl, heteroaryl, or heterocyclyl.

The term “carbonyl” refers to the group -C(=O)- .

- 35 The term “cycloalkenyl” refers to a mono- or multi-cyclic hydrocarbon ring system that contains one or more double bonds in at least one ring; although, if there is more than one, the double bonds cannot form a fully delocalized pi-electron system throughout all the rings (*i.e.*, an aromatic system), otherwise the group would be “aryl,” as defined herein. When composed of two or more

rings, the rings can be connected together in a fused, bridged, or spiro fashion. A cycloalkenyl can contain 3 to 12 atoms in the ring(s) or 3 to 8 atoms in the ring(s). In some embodiments, a cycloalkenyl group can be unsubstituted or substituted. In some embodiments, the cycloalkenyl group can have 4 to 8 carbon atoms (*i.e.*, “C₄-C₈cycloalkenyl”). An example is cyclohexenyl.

5 The term “cycloalkyl” refers to a fully saturated all carbon mono- or multi-cyclic ring system. In some embodiments, the cycloalkyl is a monocyclic ring containing 3 to 8 carbon atoms (*i.e.*, “C₃-C₈cycloalkyl”). In some embodiments, the cycloalkyl is a monocyclic ring containing 3 to 7 carbon atoms (*i.e.*, “C₃-C₇cycloalkyl”). In some embodiments, the cycloalkyl is a monocyclic or bicyclic ring containing 3 to 7 carbon atoms (*i.e.*, “C₃-C₇cycloalkyl”). Some embodiments contain 3 to 6 carbons
10 (*i.e.*, “C₃-C₆cycloalkyl”). Some embodiments contain 3 to 5 carbons. Some embodiments contain 5 to 7 carbons. Some embodiments contain 3 to 4 carbons. Examples include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. Examples include cyclopropyl, cyclobutyl, cyclopentyl, bicyclo[1.1.1]pentane, and cyclohexyl. When one or more substituents are present on the cycloalkyl group, the substituent(s) can be bonded at any available carbon atom. In some embodiments, a
15 cycloalkyl group can be substituted or unsubstituted.

 The term “dialkylamino” refers to an amino group (-NH₂) where the nitrogen is substituted with two alkyl groups. The two alkyl groups can be the same or different. The term “alkyl” has the same definition as described herein. The “dialkylamino” group can have 1 to 4 carbon atoms (*i.e.*, C₂-C₄dialkylamino) provided that the two alkyl groups do not exceed a total of 4 carbon atoms between
20 the two groups. Examples include, -N(CH₃)₂, -N(CH₃)(CH₂CH₃), -N(CH₃)(CH₂CH₂CH₃), and the like.

 The term “dialkylsulfamoyl” refers to a sulfonamide group (-S(=O)₂NH₂) where the nitrogen is substituted with two alkyl groups, the alkyl groups can be the same or different. The term “alkyl” has the same definition as described herein. The “dialkylsulfamoyl” group can have 1 to 4 carbon atoms (*i.e.*, C₂-C₄dialkylsulfamoyl) provided that two alkyl groups do not exceed a total of 4 carbon
25 atoms between the two alkyl groups. Examples include, -SO₂N(CH₃)₂, -SO₂N(CH₃)(CH₂CH₃), -SO₂N(CH₃)(CH₂CH₂CH₃), -SO₂N(CH₂CH₃)₂, -SO₂N(CH₂CH₃)(CH₂CH₂CH₃), and the like.

 The term “haloalkoxy” refers to an alkoxy group in which one or more of the hydrogen atoms are replaced by a halogen (*e.g.*, mono-haloalkoxy, di-haloalkoxy, and tri-haloalkoxy). Such groups
30 include but are not limited to, chloromethoxy, fluoromethoxy, difluoromethoxy, trifluoromethoxy, 1-chloro-2-fluoromethoxy, and 2-fluoroisobutoxy. In some embodiments, the haloalkoxy group can have 1 to 6 carbon atoms. The haloalkoxy group of the compounds can be designated as “haloC₁-C₆alkoxy”, “haloC₁-C₄alkoxy”, or similar designations.

 The term “haloalkyl” refers to an alkyl group, as defined herein, wherein one or more
35 hydrogen atoms of the alkyl group have been replaced by a halogen atom (*e.g.*, mono-haloalkyl, di-haloalkyl, and tri-haloalkyl). In some embodiments, the haloalkyl group can have 1 to 6 carbons (*i.e.*, “haloC₁-C₆alkyl”). The haloC₁-C₆alkyl can be fully substituted in which case it can be represented by

the formula C_nL_{2n+1} , wherein L is a halogen and “n” is 1, 2, 3, 4, 5, or 6. When more than one halogen is present then they can be the same or different and selected from: fluorine, chlorine, bromine, and iodine. In some embodiments, haloalkyl contains 1 to 5 carbons (*i.e.*, haloC₁-C₅alkyl). In some embodiments, haloalkyl contains 1 to 4 carbons (*i.e.*, haloC₁-C₄alkyl). In some embodiments, haloalkyl contains 1 to 3 carbons (*i.e.*, haloC₁-C₃alkyl). In some embodiments, haloalkyl contains 1 or 2 carbons. Examples of haloalkyl groups include fluoromethyl, difluoromethyl, trifluoromethyl, chlorodifluoromethyl, 1-fluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 4,4,4-trifluorobutyl, and the like.

The term “haloalkylsulfonyl” refers to a radical consisting of a haloalkyl radical bonded to the sulfur of a sulfone radical of the formula: $-S(=O)_2-$, wherein haloalkyl has the same definition as described herein. The “haloalkylsulfonyl” group can have 1 to 4 carbon atoms (*i.e.*, haloC₁-C₄alkylsulfonyl). Examples include fluoromethylsulfonyl, difluoromethylsulfonyl, trifluoromethylsulfonyl, chlorodifluoromethylsulfonyl, 1-fluoroethylsulfonyl, 2,2,2-trifluoroethylsulfonyl, pentafluoroethylsulfonyl, 4,4,4-trifluorobutylsulfonyl, and the like.

The term “halodialkylamino” refers to an amino group ($-NH_2$) where the nitrogen is substituted with either one alkyl group and one haloalkyl group, or two haloalkyl groups. When the nitrogen is substituted with two haloalkyl groups the groups can be the same or different. The terms “alkyl” and “haloalkyl” has the same definitions as described herein. The “halodialkylamino” group can have 1 to 4 carbon atoms (*i.e.*, haloC₂-C₄dialkylamino) provided that the two groups do not exceed a total of 4 carbon atoms between the two groups. Examples include, $-N(CH_3)(CF_3)$, $-N(CH_3)(CF_3)_2$, $-N(CH_3)(CH_2CF_3)$, $-N(CH_2CF_3)_2$, and the like.

The term “halogen” or “halo” refers to fluoro, chloro, bromo, or iodo group. In some embodiments, halogen or halo is fluoro, chloro, or bromo. In some embodiments, halogen or halo is fluoro or chloro. In some embodiments, halogen or halo is fluoro.

The term “heteroaryl” refers to a monocyclic or fused multicyclic aromatic ring system and having at least one heteroatom in the ring system, that is, an element other than carbon, including but not limited to, nitrogen, oxygen, and sulfur. Some embodiments are “5-6 membered heteroaryl” and refers to an aromatic ring containing 5 to 6 ring atoms in a single ring and having at least one heteroatom in the ring system. Examples of heteroaryl rings include, but are not limited to, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, isoindolyl, oxazolyl, benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, purinyl, carbazolyl, dibenzo[b,d]furan, dibenzo[b,d]thiophene, phenanthridinyl, benzimidazolyl, pyrrolyl, quinolinyl, isoquinolinyl, benzisoxazolyl, imidazo[1,2-b]thiazolyl, and the like. A heteroaryl group can be substituted or unsubstituted. In some embodiments, the heteroaryl group has 5 to 10 ring members or 5 to 7 ring members. The heteroaryl group can be designated as “5-7 membered heteroaryl,” “5-10 membered

heteroaryl,” or similar designations. In some embodiments, the heteroaryl can be a substituted or unsubstituted C₁-C₁₃ five-, six-, seven-, eight-, nine-, ten-, up to 14-membered monocyclic, bicyclic, or tricyclic ring system including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the heteroaryl can be a substituted or unsubstituted C₁-C₅ five- or six-membered monocyclic ring including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the heteroaryl can be a substituted or unsubstituted C₅-C₉ 8-, 9-, or 10-membered bicyclic ring system including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the heteroaryl is a substituted or unsubstituted C₅-C₉ 8-, 9- or 10-membered heteroaryl. In some embodiments, the C₅-C₉ 8-, 9-, or 10-membered bicyclic heteroaryl is imidazo[2,1-b]thiazolyl, 1*H*-indolyl, isoindolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzisoxazolyl, indazolyl, purinyl, quinolinyl, isoquinolinyl, quinoxaliny, pyrido[3,4-b]pyrazinyl or pyrido[4,3-d]pyrimidinyl. In some embodiments, the heteroaryl is a substituted or unsubstituted C₈-C₁₃ 13- or 14-membered tricyclic ring system including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the heteroaryl can be an azolyl, such as, imidazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, tetrazolyl, 1,2,4-thiadiazolyl, thiazolyl, isothiazolyl, oxazolyl, or isoxazolyl, each of which can be substituted or unsubstituted. In some embodiments, the heteroaryl is a C₁-C₄ 5-membered heteroaryl. In some embodiments, the C₁-C₄ 5-membered heteroaryl is furanyl, thienyl, 1,2,4-thiadiazolyl, 1,2,3-thiadiazolyl, isothiazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, oxazolyl, pyrrolyl, triazolyl, tetrazolyl. In some embodiments, the heteroaryl is a C₃-C₅ 6-membered heteroaryl. In some embodiments, the C₃-C₅ 6-membered heteroaryl is pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, or triazinyl. In some embodiments, “5-10 membered heteroaryl” refers to: furanyl, thienyl, pyrrolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, quinoxaliny, triazinyl, benzofuranyl, 1*H*-indolyl, benzo[b]thiophenyl, and the like. In some embodiments, “5-10 membered heteroaryl” refers to: pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, 1*H*-indolyl, quinoxaliny, thiadiazolyl, and the like. In some embodiments, “5-10 membered heteroaryl” refers to: pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, 1*H*-indolyl, quinoxaliny, thiadiazolyl, imidazo[4,5-c]pyridinyl, and the like. In some embodiments, a heteroaryl group can be substituted or unsubstituted.

The term “heterocyclyl” refers to a 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, up to 18-membered monocyclic, bicyclic, and tricyclic ring system wherein carbon atoms together with from 1 to 5 heteroatoms constitute said ring system and optionally containing one or more unsaturated bonds situated in such a way, however, that a fully delocalized pi-electron system (aromatic system) does not occur in the monocyclic ring or in at least one ring of the bicyclic or tricyclic ring system. The heteroatom(s) is an element other than carbon including, but not limited to, oxygen, sulfur, and nitrogen. When composed of two or more rings, the rings can be joined together in a fused, bridged, or spiro fashion where the heteroatom(s) can be present in either a non-aromatic or aromatic ring in

the ring system. In some embodiments, “4-7 membered heterocyclyl” refers to a saturated non-aromatic ring system containing 4 to 7 ring atoms, where at least one ring atom is a heteroatom. In some embodiments, “3-6 membered heterocyclyl” refers to a saturated non-aromatic ring radical containing 3 to 6 ring atoms, where at least one ring atom is a heteroatom. In some embodiments, “4-6 membered heterocyclyl” refers to a saturated non-aromatic ring radical containing 4 to 6 ring atoms, where at least one ring atom is a heteroatom. In some embodiments, the one or two heteroatoms in the ring system are selected independently from: O (oxygen) and N (nitrogen). In some embodiments, a heterocyclyl can include a carbonyl (C=O) group adjacent to a hetero atom, that is, be substituted with an oxo on a carbon adjacent to a hetero atom, where the substituted ring system is a lactam, lactone, cyclic imide, cyclic thioimide, or cyclic carbamate. Examples of unsubstituted or oxo substituted “heterocyclyl” groups include but are not limited to, aziridinyl, azetidiny, tetrahydrofuranyl, 1,3-dioxinyl, 1,3-dioxanyl, 1,4-dioxanyl, 1,2-dioxolanyl, 1,3-dioxolanyl, 1,4-dioxolanyl, 1,3-oxathianyl, 1,4-oxathiinyl, 1,3-oxathiolanyl, 1,3-dithiolyl, 1,3-dithiolanyl, 1,4-oxathianyl, tetrahydro-1,4-thiazinyl, 2*H*-1,2-oxazinyl, maleimidyl, succinimidyl, dioxopiperazinyl, hydantoinyl, imidazoliny, imidazolidinyl, isoxazoliny, isoxazolidinyl, isoindoliny, indoliny, oxazoliny, oxazolidinyl, oxazolidinonyl, thiazoliny, thiazolidinyl, morpholiny, oxiranyl, piperidinyl N-oxide, piperidinyl, piperazinyl, pyrrolidinyl, pyrrolidonyl, pyrrolidionyl, 4-piperidonyl, pyrazoliny, pyrazolidinyl, 2-oxopyrrolidinyl, tetrahydropyranyl, 4*H*-pyranyl, tetrahydrothiopyranyl, 1,4-diazabicyclo[2.2.2]octane, 1,4-diazabicyclo[3.1.1]heptane, 2-azaspiro[3,3]heptane, 2,6-diazaspiro[3,3]heptane, 2-oxa-6-azaspiro[3,3]heptane, and their benzo-fused analogs (*e.g.*, benzimidazolidinonyl, tetrahydroquinoliny, and 3,4-methylenedioxyphenyl). The heterocyclyl group can be designated as “3-10 membered heterocyclyl” or similar designations. In some embodiments, the heterocyclyl can be a C₂-C₁₂ 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, up to 13-membered monocyclic, bicyclic, or tricyclic ring system including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the heterocyclyl can be a substituted or unsubstituted C₂-C₆ 3-, 4-, 5-, 6-, or 7-membered monocyclic ring including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the heterocyclyl can be a substituted or unsubstituted C₂-C₁₀ 4-, 5-, 6-, 7-, 8-, 9-, 10-, or 11-membered bicyclic ring system including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the heterocyclyl can be a substituted or unsubstituted C₇-C₁₂ 12- or 13-membered tricyclic ring system including 1 to 5 heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the heteroatom(s) of six membered monocyclic heterocyclyls are selected from one up to three of O (oxygen), N (nitrogen), or S (sulfur), and the heteroatom(s) of five membered monocyclic heterocyclyls are selected from one or two heteroatoms selected from O (oxygen), N (nitrogen), or S (sulfur). In some embodiments, the heterocyclyl can be aziridinyl, azetidiny, tetrahydrofuranyl, 1,3-dioxinyl, 1,3-dioxanyl, 1,4-dioxanyl, 1,2-dioxolanyl, 1,3-dioxolanyl, 1,3-oxathianyl, 1,4-oxathianyl, 1,3-oxathiolanyl, 1,3-dithiolyl, 1,3-dithiolanyl, 1,4-oxathianyl, tetrahydro-1,4-thiazinyl, imidazoliny, imidazolidinyl, isoxazoliny, isoxazolidinyl, isoindoliny, indoliny,

oxazolinyl, oxazolidinyl, thiazolinyl, thiazolidinyl, morpholinyl, oxiranyl, piperidinyl, piperazinyl, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, 1,4-diazabicyclo[2.2.2]octane, 1,4-diazabicyclo[3.1.1]heptane, 2-azaspiro[3,3]heptane, 2,6-diazaspiro[3,3]heptane, tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydro-2,6-naphthyridinyl, 1,2,3,4-tetrahydro-2,7-naphthyridinyl, 1,2,3,4-tetrahydro-1,7-naphthyridinyl, 1,2,3,4-tetrahydro-1,6-naphthyridinyl, 5,6,7,8-tetrahydropyrido[2,3-d]pyrimidinyl, 5,6,7,8-tetrahydropyrido[3,4-d]pyrimidinyl, [1,3]dioxolo[4,5-c]pyridinyl, [1,3]dioxolo[4,5-b]pyridinyl, [1,3]dioxolo[4,5-d]pyrimidinyl, or 3,4-methylenedioxyphenyl. In some embodiments, the unsubstituted or substituted heterocyclyl can be selected from aziridinyl, azetidiny, piperidinyl, morpholinyl, oxetanyl, piperazinyl, pyrrolidinyl, thiomorpholinyl, 2-piperidone, 1,1-dioxidothiomorpholinyl, oxolanyl (tetrahydrofuranly), and oxanyl (tetrahydropyranyl). When one or more substituents are present on the heterocyclyl group, the substituent(s) can be bonded at any available carbon atom and/or heteroatom. In some embodiments, a heterocyclyl group can be substituted or unsubstituted.

The term “oxo” refers to the =O substituent.

The term “nitro” refers to a -NO₂ group.

The term “sulfamoyl” refers to the group -S(=O)₂NH₂.

As used herein, an “excipient” refers to a substance that is added to a composition to provide, without limitation, bulk, consistency, stability, binding ability, lubrication, disintegrating ability, *etc.*, to the composition. A “diluent” is a type of excipient and refers to an ingredient in a pharmaceutical composition that lacks pharmacological activity but can be pharmaceutically necessary or desirable. For example, a diluent can be used to increase the bulk of a potent drug whose mass is too small for manufacture and/or administration. It can also be a liquid for the dissolution of a drug to be administered by injection, ingestion, or inhalation. A pharmaceutically acceptable excipient is a physiologically and pharmaceutically suitable non-toxic and inactive material or ingredient that does not interfere with the activity of the drug substance. Pharmaceutically acceptable excipients are well known in the pharmaceutical art and described, for example, in Rowe *et al.*, Handbook of Pharmaceutical Excipients: A Comprehensive Guide to Uses, Properties, and Safety, 5th Ed., 2006, and in Remington: The Science and Practice of Pharmacy (Gennaro, 21st Ed. Mack Pub. Co., Easton, PA (2005)). Preservatives, stabilizers, dyes, buffers, and the like can be provided in the pharmaceutical composition. One skilled in this art can further formulate a compound as disclosed and described herein in an appropriate manner, and in accordance with accepted practices, such as, those disclosed in Remington, *supra*.

As used herein, a “dose” or “dosage” refers to the measured quantity of drug substance to be taken at one time by a subject. In certain embodiments, wherein the drug substance is not a free base or free acid, the quantity is the molar equivalent to the corresponding amount of free base or free acid.

As used herein, a “pharmaceutically acceptable salt” refers to salts of a compound having an acidic or basic moiety which are not biologically or otherwise undesirable for use in a pharmaceutical. In many cases, the compounds disclosed herein are capable of forming acid and/or base salts by virtue of the presence of an acidic or basic moiety (*e.g.* amino and/or carboxyl groups or groups similar thereto). Pharmaceutically acceptable acid addition salts can be formed by combining a compound having a basic moiety with inorganic acids and organic acids. Pharmaceutically acceptable base addition salts can be formed by combining a compound having an acidic moiety with inorganic and organic bases.

The compounds described herein can have one or more stereocenters. All stereoisomers, such as, enantiomers and diastereomers, are intended unless otherwise indicated. It is understood that, in any compound described herein having one or more chiral centers, if an absolute stereochemistry is not expressly indicated, then each center can independently be the (R)-configuration, or the (S)-configuration, or a mixture thereof. Thus, the compounds provided herein can be enantiomerically pure, enantiomerically enriched, a racemic mixture, diastereomerically pure, diastereomerically enriched, or a stereoisomeric mixture. Preparation of enantiomerically pure or enantiomerically enriched forms can be accomplished by resolution of racemic mixtures or by using enantiomerically pure or enriched starting materials or by stereoselective or stereospecific synthesis. Stereochemical definitions are available in E.L. Eliel, S.H. Wilen & L.N. Mander, *Stereochemistry of Organic Compounds*, John Wiley & Sons, Inc., New York, NY, 1994 which is incorporated herein by reference in its entirety. In some embodiments, where the compound described herein is chiral or otherwise includes one or more stereocenters, the compound can be prepared with an enantiomeric excess or diastereomeric excess of greater than about 75%, greater than about 80%, greater than about 85%, greater than about 90%, greater than about 95%, greater than about 99%, greater than about 99.5%, or greater than about 99.9%.

Resolution of racemic mixtures of compounds can be carried out by any of numerous methods known in the art. An example method includes fractional recrystallization using a chiral resolving organic acid with a racemic compound containing a basic group. Other chiral resolving agents suitable for fractional crystallization methods include stereoisomerically pure forms of methylbenzylamine (*e.g.*, S and R forms, or diastereomerically pure forms), 2-phenylglycinol, norephedrine, ephedrine, *N*-methylephedrine, cyclohexylethylamine, 1,2-diaminocyclohexane, and the like. Similarly, fractional recrystallization using a chiral resolving base can be utilized with a racemic compound containing a basic group.

Resolution of racemic mixtures can also be carried out by elution on a chiral column. A suitable elution solvent composition can be determined by one skilled in the art.

In some embodiments, a compound described herein can be prepared having at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at

least about 99%, at least about 99.5%, or at least about 99.9% enantiomeric excess, or an enantiomeric excess within a range defined by any of the preceding numbers. In some embodiments, a compound described herein can be prepared having at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, at least 99.5%, or at least 99.9% enantiomeric excess, or an enantiomeric excess within a range defined by any of the preceding numbers.

In some embodiments, a compound described herein can be prepared having at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 99%, at least about 99.5%, or at least about 99.9% diastereomeric excess, or a diastereomeric excess within a range defined by any of the preceding numbers. In some embodiments, a compound described herein can be prepared having at least 5%, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 99%, at least 99.5%, or at least 99.9% diastereomeric excess, or a diastereomeric excess within a range defined by any of the preceding numbers.

In addition, it is understood that, when a compound described herein contain one or more double bond(s) (*e.g.*, C=C, C=N, and the like) or other centers of geometric asymmetry, and unless specified otherwise, it is understood that the compound includes both E and Z geometric isomers (*e.g.*, *cis* or *trans*). *Cis* and *trans* geometric isomers of the compounds described herein can be isolated as a mixture of isomers or as separated isomeric form.

The compounds described herein also include tautomeric forms. Tautomeric forms result from the swapping of a single bond with an adjacent double bond together with the concomitant migration of a proton. Tautomeric forms include prototropic tautomers which are isomeric protonation states having the same empirical formula and total charge. Example prototropic tautomers include ketone – enol pairs, amide – imidic acid pairs, lactam – lactim pairs, enamine – imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, for example, 1*H*- and 3*H*-imidazole, 1*H*-, 2*H*-, and 4*H*-1,2,4-triazole, 1*H*- and 2*H*-isoindole, and 1*H*- and 2*H*-pyrazole. Tautomeric forms can be in equilibrium or sterically locked into one form by appropriate substitution.

The compounds described herein, and their pharmaceutically acceptable salts, can be found together with other substances, such as, water and solvents, for example, in the form of hydrates or solvates. When in the solid-state, the compounds described herein and salts thereof can occur in various forms and can, *e.g.*, take the form of solvates, including hydrates. The compounds can be in any solid-state form, such as, a crystalline form, amorphous form, solvated form, *etc.* and unless clearly indicated otherwise, reference in the specification to compounds and salts thereof should be understood as reading on any solid-state form of the compound.

The compounds described herein can be used in a neutral form, such as, a free acid or free base form. Alternatively, the compounds can be used in the form of pharmaceutically acceptable salts, such as, pharmaceutically acceptable addition salts of acids or bases.

In some embodiments, the compounds described herein, or salts thereof, are substantially isolated. The phrase "substantially isolated" refers to the compound that is at least partially or substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the compound described herein. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compound described herein, or salt thereof.

ISOTOPES

The compounds disclosed and described herein allow and embrace atoms at each position of the compound independently to have: 1) an isotopic distribution for a chemical element in proportional amounts to those usually found in nature or 2) an isotopic distribution in proportional amounts different to those usually found in nature unless the context clearly dictates otherwise. A particular chemical element has an atomic number defined by the number of protons within the atom's nucleus. Each atomic number identifies a specific chemical element, but not the isotope; an atom of a given element can have a wide range in its number of neutrons. The number of both protons and neutrons in the nucleus is the atom's mass number, and each isotope of a given element has a different mass number. Compounds wherein one or more atoms have an isotopic distribution for a chemical element in proportional amounts different to those usually found in nature is commonly referred to as being isotopically-labeled. Each chemical element as represented in a compound structure can include any isotopic distribution of said element. For example, in a compound structure a hydrogen atom can be explicitly disclosed or understood to be present in the compound. At any position of the compound that a hydrogen atom can be present, the hydrogen atom can be an isotopic distribution of hydrogen, including but not limited to protium (^1H) and deuterium (^2H) in proportional amounts to those usually found in nature and in proportional amounts different to those usually found in nature. Thus, reference herein to a compound encompasses all potential isotopic distributions for each atom unless the context clearly dictates otherwise. Examples of isotopes include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine, chlorine, bromine, and iodine. As one of skill in the art would appreciate, any of the compounds as disclosed and described herein can include radioactive isotopes. Accordingly, also contemplated is use of compounds as disclosed and described herein, wherein one or more atoms have an isotopic distribution different to those usually found in nature, such as, having ^2H or ^3H in greater proportion, or ^{11}C , ^{13}C , or ^{14}C in greater proportion than found in nature. By way of general example, and without limitation, isotopes of hydrogen include protium (^1H), deuterium

(²H), and tritium (³H). Isotopes of carbon include carbon-11 (¹¹C), carbon-12 (¹²C), carbon-13 (¹³C), and carbon-14 (¹⁴C). Isotopes of nitrogen include nitrogen-13 (¹³N), nitrogen-14 (¹⁴N), and nitrogen-15 (¹⁵N). Isotopes of oxygen include oxygen-14 (¹⁴O), oxygen-15 (¹⁵O), oxygen-16 (¹⁶O), oxygen-17 (¹⁷O), and oxygen-18 (¹⁸O). Isotope of fluorine include fluorine-17 (¹⁷F), fluorine-18 (¹⁸F), and fluorine-19 (¹⁹F). Isotopes of phosphorous include phosphorus-31 (³¹P), phosphorus-32 (³²P), phosphorus-33 (³³P), phosphorus-34 (³⁴P), phosphorus-35 (³⁵P), and phosphorus-36 (³⁶P). Isotopes of sulfur include sulfur-32 (³²S), sulfur-33 (³³S), sulfur-34 (³⁴S), sulfur-35 (³⁵S), sulfur-36 (³⁶S), and sulfur-38 (³⁸S). Isotopes of chlorine include chlorine-35 (³⁵Cl), chlorine-36 (³⁶Cl), and chlorine-37 (³⁷Cl). Isotopes of bromine include bromine-75 (⁷⁵Br), bromine-76 (⁷⁶Br), bromine-77 (⁷⁷Br), bromine-79 (⁷⁹Br), bromine-81 (⁸¹Br), and bromine-82 (⁸²Br). Isotopes of iodine include iodine-123 (¹²³I), iodine-124 (¹²⁴I), iodine-125 (¹²⁵I), iodine-131 (¹³¹I), and iodine-135 (¹³⁵I). In some embodiments, atoms at every position of the compound have an isotopic distribution for each chemical element in proportional amounts to those usually found in nature. In some embodiments, an atom in one position of the compound has an isotopic distribution for a chemical element in proportional amounts different to those usually found in nature (remainder atoms having an isotopic distribution for a chemical element in proportional amounts to those usually found in nature). In some embodiments, atoms in at least two positions of the compound independently have an isotopic distribution for a chemical element in proportional amounts different to those usually found in nature (remainder atoms having an isotopic distribution for a chemical element in proportional amounts to those usually found in nature). In some embodiments, atoms in at least three positions of the compound independently have an isotopic distribution for a chemical element in proportional amounts different to those usually found in nature (remainder atoms having an isotopic distribution for a chemical element in proportional amounts to those usually found in nature). In some embodiments, atoms in at least four positions of the compound independently have an isotopic distribution for a chemical element in proportional amounts different to those usually found in nature (remainder atoms having an isotopic distribution for a chemical element in proportional amounts to those usually found in nature). In some embodiments, atoms in at least five positions of the compound independently have an isotopic distribution for a chemical element in proportional amounts different to those usually found in nature (remainder atoms having an isotopic distribution for a chemical element in proportional amounts to those usually found in nature). In some embodiments, atoms in at least six positions of the compound independently have an isotopic distribution for a chemical element in proportional amounts different to those usually found in nature (remainder atoms having an isotopic distribution for a chemical element in proportional amounts to those usually found in nature).

Certain compounds, for example those having incorporated radioactive isotopes, such as, ³H and ¹⁴C, are also useful in drug or substrate tissue distribution assays. Tritium (³H) and carbon-14 (¹⁴C) isotopes are particularly preferred for their ease of preparation and detectability. Compounds with isotopes, such as, deuterium (²H), in proportional amounts greater than usually found in nature

can afford certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements. Isotopically-labeled compounds can generally be prepared by performing procedures routinely practiced in the chemical art. Methods are readily available to measure such isotope perturbations or enrichments, such as, mass

spectrometry, and for isotopes that are radio-isotopes additional methods are available, such as, radio-detectors used in connection with HPLC or GC.

As used herein, "isotopic variant" means a compound that contains an unnatural proportion of an isotope at one or more of the atoms that constitute such a compound. In certain embodiments, an "isotopic variant" of a compound contains unnatural proportions of one or more isotopes, including, but not limited to, protium (^1H), deuterium (^2H), tritium (^3H), carbon-11 (^{11}C), carbon-12 (^{12}C), carbon-13 (^{13}C), carbon-14 (^{14}C), nitrogen-13 (^{13}N), nitrogen-14 (^{14}N), nitrogen-15 (^{15}N), oxygen-14 (^{14}O), oxygen-15 (^{15}O), oxygen-16 (^{16}O), oxygen-17 (^{17}O), oxygen-18 (^{18}O), fluorine-17 (^{17}F), fluorine-18 (^{18}F), phosphorus-31 (^{31}P), phosphorus-32 (^{32}P), phosphorus-33 (^{33}P), sulfur-32 (^{32}S), sulfur-33 (^{33}S), sulfur-34 (^{34}S), sulfur-35 (^{35}S), sulfur-36 (^{36}S), chlorine-35 (^{35}Cl), chlorine-36 (^{36}Cl), chlorine-37 (^{37}Cl), bromine-79 (^{79}Br), bromine-81 (^{81}Br), iodine-123 (^{123}I), iodine-125 (^{125}I), iodine-127 (^{127}I), iodine-129 (^{129}I), and iodine-131 (^{131}I). In certain embodiments, an "isotopic variant" of a compound is in a stable form, that is, non-radioactive. In certain embodiments, an "isotopic variant" of a compound contains unnatural proportions of one or more isotopes, including, but not limited to, hydrogen (^1H), deuterium (^2H), carbon-12 (^{12}C), carbon-13 (^{13}C), nitrogen-14 (^{14}N), nitrogen-15 (^{15}N), oxygen-16 (^{16}O), oxygen-17 (^{17}O), and oxygen-18 (^{18}O). In certain embodiments, an "isotopic variant" of a compound is in an unstable form, that is, radioactive. In certain embodiments, an "isotopic variant" of a compound described herein contains unnatural proportions of one or more isotopes, including, but not limited to, tritium (^3H), carbon-11 (^{11}C), carbon-14 (^{14}C), nitrogen-13 (^{13}N), oxygen-14 (^{14}O), and oxygen-15 (^{15}O). It will be understood that, in a compound as provided herein, any hydrogen can include ^2H as the major isotopic form, as example, or any carbon include be ^{13}C as the major isotopic form, as example, or any nitrogen can include ^{15}N as the major isotopic form, as example, and any oxygen can include ^{18}O as the major isotopic form, as example. In certain embodiments, an "isotopic variant" of a compound contains an unnatural proportion of deuterium (^2H).

With regard to the compounds provided herein, when a particular atomic position is designated as having deuterium or "D" or "d", it is understood that the abundance of deuterium at that position is substantially greater than the natural abundance of deuterium, which is about 0.015%. A position designated as having deuterium typically has a minimum isotopic enrichment factor of, in certain embodiments, at least 3500 (52.5% deuterium incorporation), at least 4000 (60% deuterium incorporation), at least 4500 (67.5% deuterium incorporation), at least 5000 (75% deuterium incorporation), at least 5500 (82.5% deuterium incorporation), at least 6000 (90% deuterium incorporation), at least 6333.3 (95% deuterium incorporation), at least 6466.7 (97% deuterium

incorporation), at least 6600 (99% deuterium incorporation), or at least 6633.3 (99.5% deuterium incorporation) at each designated deuterium position. Similarly, with regard to the compounds provided herein when any atomic position is designated as a specific isotope, it is understood that the abundance of the specific isotope at that position is substantially greater than the natural abundance of that isotope. A position designated as having a specific isotope typically has a minimum isotopic enrichment factor of, in certain embodiments, at least 52.5%, at least 60%, at least 67.5%, at least 75%, at least 82.5%, at least 90%, at least 95%, at least 97%, at least 99%, or at least 99.5% incorporation of the isotope at each designated position.

Synthetic methods for incorporating radio-isotopes into organic compounds are applicable to compound described herein and are well known in the art. These synthetic methods, for example, incorporating activity levels of tritium into target molecules, are as follows:

A. Catalytic Reduction with Tritium Gas: This procedure normally yields high specific activity products and requires halogenated or unsaturated precursors.

B. Reduction with Sodium Borohydride [^3H]: This procedure is rather inexpensive and requires precursors containing reducible functional groups, such as, aldehydes, ketones, lactones, esters, and the like.

C. Reduction with Lithium Aluminum Hydride [^3H]: This procedure offers products at almost theoretical specific activities. It also requires precursors containing reducible functional groups, such as, aldehydes, ketones, lactones, esters, and the like.

D. Tritium Gas Exposure Labeling: This procedure involves exposing precursors containing exchangeable protons to tritium gas in the presence of a suitable catalyst.

E. *N*-Methylation using Methyl Iodide [^3H]: This procedure is usually employed to prepare *O*-methyl or *N*-methyl (^3H) products by treating appropriate precursors with high specific activity methyl iodide (^3H). This method in general allows for higher specific activity, such as, for example, about 70-90 Ci/mmol.

Synthetic methods for incorporating activity levels of ^{125}I into target molecules include:

A. Sandmeyer and like reactions: This procedure transforms an aryl amine or a heteroaryl amine into a diazonium salt, such as, a diazonium tetrafluoroborate salt and subsequently to ^{125}I labeled compound using Na^{125}I . A representative procedure was reported by Zhu, G-D. and co-workers in *J. Org. Chem.*, 2002, 67, 943-948.

B. Ortho ^{125}I iodination of phenols: This procedure allows for the incorporation of ^{125}I at the ortho position of a phenol as reported by Collier, T. L. and co-workers in *J. Labelled Compd. Radiopharm.*, 1999, 42, S264-S266.


C. Aryl and heteroaryl bromide exchange with ^{125}I : This method is generally a two-step process. The first step is the conversion of the aryl or heteroaryl bromide to the corresponding tri-alkyltin intermediate using for example, a Pd catalyzed reaction [*i.e.* $\text{Pd}(\text{Ph}_3\text{P})_4$] or through an aryl or heteroaryl lithium, in the presence of a tri-alkyltinhalide or hexaalkylditin [*e.g.*, $(\text{CH}_3)_3\text{SnSn}(\text{CH}_3)_3$].

A representative procedure was reported by Le Bas, M.-D. and co-workers in J. Labelled Compd. Radiopharm., 2001, 44, S280-S282.

A radiolabeled form of a compound described herein can be used in a screening assay to identify/evaluate compounds. In general terms, a newly synthesized or identified compound (*i.e.*, test compound) can be evaluated for its ability to reduce binding of a radiolabeled form of a compound disclosed herein to VMAT2. The ability of a test compound to compete with a radiolabeled form of a compound described herein for the binding to VMAT2 correlates to its binding affinity.

COMPOUNDS

Some embodiments provide a compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

R^1 is R^{2A} , $-(CH_2)_m-R^{2B}$, or $-(CH_2)_n-OR^{2C}$;

R^{2A} is C_2 - C_4 alkenyl, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more R^A ;

each R^A is independently selected from the group consisting of: C_1 - C_4 alkoxy, C_1 - C_4 alkyl, C_1 - C_4 alkylamino, C_1 - C_4 alkylsulfonyl, C_2 - C_4 dialkylamino, C_2 - C_4 dialkylsulfamoyl, halogen, halo C_1 - C_4 alkoxy, halo C_1 - C_4 alkyl, halo C_2 - C_4 dialkylamino, 4-7 membered heterocyclyl, $-CN$, $-OH$, and oxo;

R^{2B} is aryl, C_3 - C_8 cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more R^B ;

each R^B is independently selected from the group consisting of: C_1 - C_4 alkyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 alkylsulfonyloxy, $-CN$, $-CH_2CN$, halogen, halo C_1 - C_4 alkyl, halo C_1 - C_4 alkylsulfonyl, oxo, and $-OH$;


R^{2C} is C_1 - C_4 alkyl, C_3 - C_8 cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more R^C ;

each R^C is independently $-CN$ or C_3 - C_6 cycloalkyl;

m is 0, 1, 2, 3, or 4; and

n is 1, 2, or 3.

Some embodiments provide a compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

R^1 is R^{2A} , $-(CH_2)_m-R^{2B}$, or $-(CH_2)_n-OR^{2C}$;

R^{2A} is C_1 - C_4 alkyl, unsubstituted or substituted with one or more R^A ;

each R^A is independently selected from the group consisting of: fluoro and $-OH$;

R^{2B} is C_3 - C_8 cycloalkyl or 4-6 membered heterocyclyl, each unsubstituted or substituted with one or more R^B ;

each R^B is independently selected from the group consisting of: fluoro, $-CN$, $-CH_2CN$, and $-OH$;


R^{2C} is C_1 - C_4 alkyl or C_3 - C_8 cycloalkyl, each unsubstituted or substituted with one or more R^C ;

each R^C is independently $-CN$ or C_3 - C_6 cycloalkyl;

m is 0, 1, 2, or 3; and


n is 1, 2, or 3.

In some embodiments, the compound of Formula (I) is a compound of Formula (Ia):




or a pharmaceutically acceptable salt thereof. The stereochemistry for a compound of Formula (Ia) is $2R,3R,11bR$.

In some embodiments, the compound of Formula (I) is a compound of Formula (Ic):




or a pharmaceutically acceptable salt thereof. The stereochemistry for a compound of Formula (Ic) is $2S,3R,11bR$.

In some embodiments, the compound of Formula (I) is a compound of Formula (Ie):



or a pharmaceutically acceptable salt thereof. The stereochemistry for a compound of Formula (Ie) is 2*R*,3*S*,11*bS*.

5 In some embodiments, the compound of Formula (I) is a compound of Formula (Ig):



or a pharmaceutically acceptable salt thereof. The stereochemistry for a compound of Formula (Ig) is 2*S*,3*S*,11*bS*.

In some embodiments, **R**¹ is **R**^{2A}.

10 In some embodiments, **R**^{2A} is C₂-C₄alkenyl, C₁-C₆alkyl, C₃-C₈cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more **R**^A, wherein **R**^A is as described herein *supra*, and *infra*. In some embodiments, **R**^{2A} is C₂-C₄alkenyl, C₁-C₆alkyl, C₃-C₈cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more **R**^A, each **R**^A is independently selected from the group consisting of: C₁-C₄alkoxy, C₁-C₄alkyl, C₁-C₄alkylamino, C₁-C₄alkylsulfonyl, C₂-C₄dialkylamino, C₂-C₄dialkylsulfamoyl, halogen, haloC₁-C₄alkoxy, haloC₁-C₄alkyl, haloC₂-C₄dialkylamino, 4-7 membered heterocyclyl, -CN, -OH, and oxo. It is understood that when more than one **R**^A is present, they may be the same or different. In some embodiments, when more than one **R**^A is present, they are the same. In some embodiments, when more than one **R**^A is present, they are different.

20 In some embodiments, **R**^{2A} is C₂-C₄alkenyl, C₁-C₆alkyl, C₃-C₈cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one, two, three, or four **R**^A. In some embodiments, **R**^{2A} is C₂-C₄alkenyl, C₁-C₆alkyl, C₃-C₈cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one **R**^A. In some embodiments, **R**^{2A} is C₂-C₄alkenyl, C₁-C₆alkyl, C₃-C₈cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with two **R**^A. In some embodiments, **R**^{2A} is C₂-C₄alkenyl, C₁-C₆alkyl, C₃-C₈cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with three **R**^A.

In some embodiments, R^{2A} is C₂-C₄alkenyl, unsubstituted or substituted with one or more R^A , wherein R^A is as described herein *supra*, and *infra*.

In some embodiments, R^{2A} is methylallyl, unsubstituted or substituted with one or more R^A .

In some embodiments, R^{2A} is prop-2-en-1-yl, unsubstituted or substituted with one or more methyl. In some embodiments, R^{2A} is methylallyl.

In some embodiments, R^{2A} is C₁-C₆alkyl, unsubstituted or substituted with one or more R^A , wherein R^A is as described herein *supra*, and *infra*.

In some embodiments, R^{2A} is C₁-C₆alkyl, unsubstituted or substituted with one or more R^A .

In some embodiments, R^{2A} is *n*-butyl, ethyl, 2-ethylbutyl, *n*-propyl, propan-2-yl, 2-methylpropyl, methyl, or *n*-pentyl, each unsubstituted or substituted with one or more R^A .

In some embodiments, each R^A is independently selected from the group consisting of: C₁-C₄alkoxy, C₁-C₄alkylamino, C₁-C₄alkylsulfonyl, C₂-C₄dialkylamino, C₂-C₄dialkylsulfamoyl, halogen, haloC₁-C₄alkoxy, haloC₂-C₄dialkylamino, 4-7 membered heterocyclyl, -CN, -OH, and oxo. In some embodiments, each R^A is independently selected from the group consisting of: methoxy, methylamino, methylsulfonyl, dimethylamino, *N,N*-dimethylsulfamoyl, chloro, fluoro, trifluoromethoxy, methyl(2,2,2-trifluoroethyl)amino, methyl(trifluoromethyl)amino, morpholino, -CN, -OH, and oxo.

In some embodiments, R^{2A} is *n*-butyl, ethyl, 2-ethylbutyl, *n*-propyl, propan-2-yl, 2-methylpropyl, methyl, or *n*-pentyl, each unsubstituted or substituted with one or more R^A independently selected from the group consisting of: C₁-C₄alkoxy, C₁-C₄alkylamino, C₁-C₄alkylsulfonyl, C₂-C₄dialkylamino, C₂-C₄dialkylsulfamoyl, halogen, haloC₁-C₄alkoxy, haloC₂-C₄dialkylamino, 4-7 membered heterocyclyl, -CN, -OH, and oxo.

In some embodiments, R^{2A} is 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, ethyl, 2-fluoroethyl, 3-fluoropropyl, propan-2-yl, 1,1,1-trifluoropropan-2-yl, 2-hydroxy-2-methylpropyl, 3,3,3-trifluoro-2-hydroxypropyl, 3,3-difluoro-2-hydroxypropyl, methyl, 2-hydroxypropyl, 2-hydroxybutyl, 4,4,4-trifluoro-2-hydroxybutyl, 3,3,3-trifluoro-2-hydroxy-2-methylpropyl, 2,3-dihydroxypropyl, 3-hydroxypropyl, 2-hydroxyethyl, cyanomethyl, 2-(dimethylamino)-2-oxoethyl, 2-(methylamino)-2-oxoethyl, 2-hydroxy-3-methoxypropyl, 2-chloro-2,2-difluoroethyl, 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl, 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl, 3-(methylsulfonyl)propyl, (*N,N*-dimethylsulfamoyl)methyl, 2-methoxypropyl, 2-ethyl-2-hydroxybutyl, 3-chloro-3,3-difluoropropyl, 2-hydroxy-3-morpholinopropyl, 5,5,5-trifluoropentyl, or 4,4,4-trifluorobutyl.

In some embodiments, R^{2A} is C₁-C₄alkyl, unsubstituted or substituted with one two, three, or four R^A ; and each R^A is independently selected from the group consisting of: fluoro and -OH. In some embodiments, R^{2A} is C₁-C₄alkyl, unsubstituted or substituted with one, two, or three R^A ; and each R^A is independently selected from the group consisting of: fluoro and -OH. In some embodiments, R^{2A} is C₂-C₄alkyl. In some embodiments, R^{2A} is -CH₂CH₃, -CH₂CH₂CH₃, or -CH(CH₃)₂. In some

embodiments, R^{2A} is C_1 - C_4 alkyl substituted with one, two, or three fluoro. In some embodiments, R^{2A} is C_1 - C_4 alkyl substituted with one, two, or three groups each independently selected from fluoro and a C-amido group. In some embodiments, R^{2A} is C_1 - C_4 alkyl substituted with one, two, or three fluoro. In some embodiments, R^{2A} is $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2CH_2CHF_2$, or $-CH_2CH_2CF_3$. In

5 some embodiments, R^{2A} is $-CH_2CH_2CHF_2$ or $-CH_2CH_2CF_3$.

In some embodiments, R^{2A} is C_3 - C_8 cycloalkyl, unsubstituted or substituted with one or more R^A , wherein R^A is as described herein *supra*, and *infra*.

10 In some embodiments, R^{2A} is C_3 - C_8 cycloalkyl, unsubstituted or substituted with one or more R^A .

In some embodiments, R^{2A} is cyclobutyl, cyclopropyl, cyclopentyl, or cyclohexyl, each unsubstituted or substituted with one or more R^A .

In some embodiments, each R^A is independently selected from the group consisting of: halogen and halo C_1 - C_4 alkyl. In some embodiments, each R^A is independently selected from the group consisting of: fluoro and trifluoromethyl.

15 In some embodiments, R^{2A} is cyclobutyl, cyclopropyl, cyclopentyl, or cyclohexyl, each unsubstituted or substituted with one or more R^A independently selected from the group consisting of: fluoro and trifluoromethyl. In some embodiments, R^{2A} is cyclopropyl, cyclobutyl, 4-(trifluoromethyl)cyclohexyl, 2-fluorocyclopentyl, or 3,3-difluorocyclopentyl.

20 In some embodiments, R^{2A} is 4-7 membered heterocyclyl, unsubstituted or substituted with one or more R^A , wherein R^A is as described herein *supra*, and *infra*.

In some embodiments, R^{2A} is azetidin-3-yl, oxetan-3-yl, or pyrrolidin-3-yl, each unsubstituted or substituted with one or more R^A .

25 In some embodiments, each R^A is independently selected from the group consisting of: C_1 - C_6 alkyl, halo C_1 - C_4 alkyl, and oxo. In some embodiments, each R^A is independently selected from the group consisting of: methyl, 2,2,2-trifluoroethyl, and oxo.

30 In some embodiments, R^{2A} is azetidin-3-yl, oxetan-3-yl, or pyrrolidin-3-yl, each unsubstituted or substituted with one or more R^A independently selected from the group consisting of: methyl, 2,2,2-trifluoroethyl, and oxo. In some embodiments, R^{2A} is oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl, or 1-methylazetidin-3-yl.

35 In some embodiments, R^{2A} is methylallyl, *n*-butyl, ethyl, 2-ethylbutyl, *n*-propyl, propan-2-yl, 2-methylpropyl (isobutyl), methyl, *n*-pentyl, cyclobutyl, cyclopropyl, cyclopentyl, cyclohexyl, azetidin-3-yl, oxetan-3-yl, or pyrrolidin-3-yl; each unsubstituted or substituted with one or more R^A independently selected from the group consisting of: methoxy, methylamino, methylsulfonyl, dimethylamino, *N,N*-dimethylsulfamoyl, chloro, fluoro, trifluoromethoxy, methyl(2,2,2-trifluoroethyl)amino, methyl(trifluoromethyl)amino, morpholino, -CN, -OH, trifluoromethyl, methyl, 2,2,2-trifluoroethyl, and oxo.


In some embodiments, R^{2A} is methylallyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, ethyl, 2-fluoroethyl, 3-fluoropropyl, propan-2-yl, 1,1,1-trifluoropropan-2-yl, 2-hydroxy-2-methylpropyl, 3,3,3-trifluoro-2-hydroxypropyl, 3,3-difluoro-2-hydroxypropyl, methyl, 2-hydroxypropyl, 2-hydroxybutyl, 4,4,4-trifluoro-2-hydroxybutyl, 3,3,3-trifluoro-2-hydroxy-2-methylpropyl, 2,3-dihydroxypropyl, 3-hydroxypropyl, 2-hydroxyethyl, cyanomethyl, 2-(dimethylamino)-2-oxoethyl, 2-(methylamino)-2-oxoethyl, 2-hydroxy-3-methoxypropyl, 2-chloro-2,2-difluoroethyl, 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl, 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl, 3-(methylsulfonyl)propyl, (*N,N*-dimethylsulfamoyl)methyl, 2-methoxypropyl, 2-ethyl-2-hydroxybutyl, 3-chloro-3,3-difluoropropyl, 2-hydroxy-3-morpholinopropyl, cyclopropyl, cyclobutyl, 4-(trifluoromethyl)cyclohexyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl, 1-methylazetidin-3-yl, 5,5,5-trifluoropentyl, or 4,4,4-trifluorobutyl.

In some embodiments, R^{2A} is methylallyl. In some embodiments, R^{2A} is 2,2,2-trifluoroethyl. In some embodiments, R^{2A} is 2,2-difluoroethyl. In some embodiments, R^{2A} is 2,2-difluoropropyl. In some embodiments, R^{2A} is 3,3,3-trifluoropropyl. In some embodiments, R^{2A} is ethyl. In some embodiments, R^{2A} is 2-fluoroethyl. In some embodiments, R^{2A} is 3-fluoropropyl. In some embodiments, R^{2A} is propan-2-yl. In some embodiments, R^{2A} is 1,1,1-trifluoropropan-2-yl. In some embodiments, R^{2A} is 2-hydroxy-2-methylpropyl. In some embodiments, R^{2A} is 3,3,3-trifluoro-2-hydroxypropyl. In some embodiments, R^{2A} is 3,3-difluoro-2-hydroxypropyl. In some embodiments, R^{2A} is methyl. In some embodiments, R^{2A} is 2-hydroxypropyl. In some embodiments, R^{2A} is 2-hydroxybutyl. In some embodiments, R^{2A} is 4,4,4-trifluoro-2-hydroxybutyl. In some embodiments, R^{2A} is 3,3,3-trifluoro-2-hydroxy-2-methylpropyl. In some embodiments, R^{2A} is 2,3-dihydroxypropyl. In some embodiments, R^{2A} is 3-hydroxypropyl. In some embodiments, R^{2A} is 2-hydroxyethyl. In some embodiments, R^{2A} is cyanomethyl. In some embodiments, R^{2A} is 2-(dimethylamino)-2-oxoethyl. In some embodiments, R^{2A} is 2-(methylamino)-2-oxoethyl. In some embodiments, R^{2A} is 2-hydroxy-3-methoxypropyl. In some embodiments, R^{2A} is 2-chloro-2,2-difluoroethyl. In some embodiments, R^{2A} is 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl. In some embodiments, R^{2A} is 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl. In some embodiments, R^{2A} is 3-(methylsulfonyl)propyl. In some embodiments, R^{2A} is (*N,N*-dimethylsulfamoyl)methyl. In some embodiments, R^{2A} is 2-methoxypropyl. In some embodiments, R^{2A} is 2-ethyl-2-hydroxybutyl. In some embodiments, R^{2A} is 3-chloro-3,3-difluoropropyl. In some embodiments, R^{2A} is 2-hydroxy-3-morpholinopropyl. In some embodiments, R^{2A} is cyclopropyl. In some embodiments, R^{2A} is cyclobutyl. In some embodiments, R^{2A} is 4-(trifluoromethyl)cyclohexyl. In some embodiments, R^{2A} is 2-fluorocyclopentyl. In some embodiments, R^{2A} is 3,3-difluorocyclopentyl. In some embodiments, R^{2A} is oxetan-3-yl. In some embodiments, R^{2A} is oxolan-3-yl (tetrahydrofuran-3-yl). In some embodiments, R^{2A} is 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl. In some embodiments, R^{2A} is 1-

methylazetidin-3-yl. In some embodiments, R^{2A} is 5,5,5-trifluoropentyl. In some embodiments, R^{2A} is 4,4,4-trifluorobutyl.





In some embodiments, R^1 is $-(CH_2)_m-R^{2B}$.

In some embodiments, the compound of Formula (I) is a compound of Formula (III):



or a pharmaceutically acceptable salt thereof, wherein: m and R^{2B} have the same definitions as described herein, and each can be selected independently from any of the embodiments as described herein, *supra*, and *infra*.

In some embodiments, the compound of Formula (I) is a compound selected from Formula (IIIa), Formula (IIIc), Formula (IIIe), or Formula (IIIg):



or a pharmaceutically acceptable salt thereof, wherein: m and R^{2B} have the same definitions as described herein, and each can be selected independently from any of the embodiments as described herein, *supra*, and *infra*.

In some embodiments, the compound of Formula (I) is a compound of Formula (IIIa). In some embodiments, the compound of Formula (I) is a compound of Formula (IIIc). In some embodiments, the compound of Formula (I) is a compound of Formula (IIIe). In some embodiments, the compound of Formula (I) is a compound of Formula (IIIg).

In some embodiments, m is 0, 1, 2, 3, or 4.

In some embodiments, m is 0, 1, 2, or 3.

In some embodiments, m is 0, 1, or 2.

In some embodiments, m is 0 or 1.

In some embodiments, **m** is 0.

In some embodiments, **m** is 1.

In some embodiments, **m** is 2.

In some embodiments, **m** is 3.

5 In some embodiments, **m** is 4.

In some embodiments, **R^{2B}** is aryl, C₃-C₈cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more **R^B**. It is understood that when more than one **R^B** is present, they may be the same or different. In some embodiments, when more than one **R^B** is present, they are the same. In some embodiments, when more than one **R^B** is present, they are different.

In some embodiments, **R^{2B}** is aryl, C₃-C₈cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one, two, three, or four **R^B**.

In some embodiments, **R^{2B}** is aryl, C₃-C₈cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one **R^B**. In some embodiments, **R^{2B}** is aryl, C₃-C₈cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with two **R^B**. In some embodiments, **R^{2B}** is aryl, C₃-C₈cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with three **R^B**. In some embodiments, **R^{2B}** is aryl, C₃-C₈cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with four **R^B**.

20 In some embodiments, **R^{2B}** is aryl, unsubstituted or substituted with one or more **R^B**, wherein **R^B** is as described herein *supra*, and *infra*.

In some embodiments, **R^{2B}** is phenyl, unsubstituted or substituted with one or more **R^B**.

In some embodiments, each **R^B** is independently selected from the group consisting of: halogen, haloC₁-C₄alkyl, haloC₁-C₄alkylsulfonyl, and -CN. In some embodiments, each **R^B** is independently selected from the group consisting of: fluoro, trifluoromethyl, (difluoromethyl)sulfonyl, and -CN.

In some embodiments, **R^{2B}** is phenyl, unsubstituted or substituted with one or more **R^B** independently selected from the group consisting of: fluoro, trifluoromethyl, (difluoromethyl)sulfonyl, and -CN. In some embodiments, **R^{2B}** is phenyl, 4-fluorophenyl, 3-fluorophenyl, 4-(trifluoromethyl)phenyl, 4-((difluoromethyl)sulfonyl)phenyl, or 3-cyanophenyl.

In some embodiments, **R^{2B}** is C₃-C₈cycloalkyl, unsubstituted or substituted with one or more **R^B**, wherein **R^B** is as described herein *supra*, and *infra*.

In some embodiments, **R^{2B}** is cyclobutyl, cyclopropyl, cyclohexyl, cyclopentyl, or bicyclo[1.1.1]pentan-1-yl, each unsubstituted or substituted with one or more **R^B**.

35 In some embodiments, each **R^B** is independently selected from the group consisting of: C₁-C₄alkyl, C₁-C₄alkylsulfonyl, C₁-C₄alkylsulfonyloxy, -CN, -CH₂CN, halogen, haloC₁-C₄alkyl, and -OH.

In some embodiments, each R^B is independently selected from the group consisting of: ethyl, methylsulfonyl, methylsulfonyloxy, -CN, -CH₂CN, fluoro, difluoromethyl, trifluoromethyl, and -OH.

In some embodiments, R^{2B} is cyclobutyl, cyclopropyl, cyclohexyl, cyclopentyl, or bicyclo[1.1.1]pentan-1-yl, each unsubstituted or substituted with one or more R^B independently selected from the group consisting of: ethyl, methylsulfonyl, methylsulfonyloxy, -CN, -CH₂CN, fluoro, difluoromethyl, trifluoromethyl, and -OH.

In some embodiments, R^{2B} is 1-hydroxycyclobutyl, cyclopropyl, 1-cyanocyclopropyl, 1-fluorocyclopropyl, cyclobutyl, 1-cyanocyclobutyl, 1-(cyanomethyl)cyclopropyl, 4,4-difluoro-1-hydroxycyclohexyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, 3-fluorobicyclo[1.1.1]pentan-1-yl, 1-(difluoromethyl)cyclopropyl, 3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl, 3-hydroxy-3-(trifluoromethyl)cyclobutyl, 1-(trifluoromethyl)cyclopentyl, 2-ethylcyclopropyl, 1-((methylsulfonyl)oxy)cyclopropyl, 1-(methylsulfonyl)cyclopropyl, or 2,2-difluorocyclopropyl.

In some embodiments, R^{2B} is C₃-C₆cycloalkyl, unsubstituted or substituted with one or two R^B ; and each R^B is independently selected from the group consisting of: fluoro, CN, -CH₂CN, and -OH. In some embodiments, R^{2B} is cyclopropyl, cyclobutyl, or cyclopentyl, each unsubstituted or substituted with one or two fluoro.

In some embodiments, R^{2B} is 5-10 membered heteroaryl, unsubstituted or substituted with one or more R^B , wherein R^B is as described herein *supra*, and *infra*.

In some embodiments, R^{2B} is 1,2,4-oxadiazol-5-yl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 1H-imidazol-2-yl, 1H-pyrazol-5-yl, 1H-tetrazol-5-yl, furan-2-yl, imidazol-1-yl, isoxazol-3-yl, isoxazol-5-yl, pyrazin-2-yl, pyridin-2-yl, pyridin-3-yl, pyrimidin-2-yl, or thiazol-4-yl, each unsubstituted or substituted with one or more R^B .

In some embodiments, each R^B is independently selected from the group consisting of: C₁-C₆alkyl and -CN. In some embodiments, each R^B is independently selected from the group consisting of: methyl and -CN.

In some embodiments, R^{2B} is 1,2,4-oxadiazol-5-yl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 1H-imidazol-2-yl, 1H-pyrazol-5-yl, 1H-tetrazol-5-yl, furan-2-yl, imidazol-1-yl, isoxazol-3-yl, isoxazol-5-yl, pyrazin-2-yl, pyridin-2-yl, pyridin-3-yl, pyrimidin-2-yl, or thiazol-4-yl, each unsubstituted or substituted with one or more R^B independently selected from the group consisting of: methyl and -CN. In some embodiments, R^{2B} is imidazol-1-yl, 5-methyl-1,3-oxazol-2-yl, 5-cyanofuran-2-yl, isoxazol-3-yl, oxazol-4-yl, 6-cyanopyridin-2-yl, 5-cyanopyridin-2-yl, 1,2,4-oxadiazol-5-yl, pyridin-3-yl, pyrimidin-2-yl, isoxazol-5-yl, 1H-imidazol-2-yl, or 1H-tetrazol-5-yl.

In some embodiments, R^{2B} is 4-7 membered heterocyclyl, unsubstituted or substituted with one or more R^B , wherein R^B is as described herein *supra*, and *infra*.

In some embodiments, R^{2B} is oxetan-3-yl, oxetan-2-yl, oxolan-3-yl (tetrahydrofuran-3-yl), pyrrolidin-1-yl, piperidin-4-yl, tetrahydro-2H-pyran-4-yl, azetidin-3-yl, 1,4-dioxepan-6-yl, or tetrahydro-2H-thiopyran-4-yl, unsubstituted or substituted with one or more R^B .

In some embodiments, each R^B is independently selected from the group consisting of: C_1 - C_4 alkyl, halogen, oxo, and -CN. In some embodiments, each R^B is independently selected from the group consisting of: methyl, fluoro, oxo, and -CN.

In some embodiments, R^{2B} is oxetan-3-yl, oxetan-2-yl, oxolan-3-yl (tetrahydrofuran-3-yl), pyrrolidin-1-yl, piperidin-4-yl, tetrahydro-2*H*-pyran-4-yl, azetidin-3-yl, 1,4-dioxepan-6-yl, or tetrahydro-2*H*-thiopyran-4-yl, each unsubstituted or substituted with one or more R^B independently selected from the group consisting of: methyl, fluoro, oxo, and -CN. In some embodiments, R^{2B} is 3-fluorooxetan-3-yl, oxetan-2-yl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 3,3-difluoropyrrolidin-1-yl, 1-cyanopiperidin-4-yl, 1-methylazetidin-3-yl, 6-fluoro-1,4-dioxepan-6-yl, or 1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl.

In some embodiments, R^{2B} is a 4-5 membered monocyclic heterocyclyl having one ring forming heteroatom selected from oxygen and sulfur, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or more R^B ; and each R^B is independently selected from the group consisting of: fluoro and -OH. In some embodiments, R^{2B} is a C_3 - C_4 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two R^B ; and each R^B is independently selected from the group consisting of: fluoro and -OH. In some embodiments, R^{2B} is a 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two fluoro. In some embodiments, R^{2B} is a C_3 - C_4 4-5 membered monocyclic heterocyclyl having one ring forming heteroatom selected from oxygen and sulfur, wherein the C_3 - C_4 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or more R^B ; and each R^B is independently selected from the group consisting of: fluoro and -OH. In some embodiments, R^{2B} is a 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two R^B ; and each R^B is independently selected from the group consisting of: fluoro and -OH. In some embodiments, R^{2B} is a C_3 - C_4 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the C_3 - C_4 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two R^B ; and each R^B is independently selected from the group consisting of: fluoro and -OH. In some embodiments, R^{2B} is a 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two fluoro. In some embodiments, R^{2B} is a C_3 - C_4 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the C_3 - C_4 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two fluoro. In some embodiments, R^{2B} is oxetan-3-yl or oxolan-3-yl, each unsubstituted or substituted with one fluoro.

In some embodiments, R^{2B} is heterocyclyl, unsubstituted or substituted with one or more R^B ; and each R^B is independently selected from the group consisting of: fluoro, chloro, bromo, -CN, - CH_2CN ,

-CH₂F, -CHF₂, -CF₃, -OH, -OCF₃, -OCH₃, and -OCH₂CH₃. In some embodiments, **R^{2B}** is a 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, 12-, or 13-membered monocyclic, bicyclic, or tricyclic heterocyclyl ring system having 1 to 5 ring forming heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the heterocyclyl is unsubstituted or substituted with one or more **R^B**; and each **R^B** is independently selected from the group consisting of: fluoro, chloro, bromo, -CN, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -OH, -OCF₃, -OCH₃, and -OCH₂CH₃. In some embodiments, **R^{2B}** is a 3-, 4-, 5-, 6-, or 7-membered monocyclic heterocyclyl ring having 1 to 5 ring forming heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the heterocyclyl is unsubstituted or substituted with one or more **R^B**, and each **R^B** is independently selected from the group consisting of: fluoro, chloro, bromo, -CN, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -OH, -OCF₃, -OCH₃, and -OCH₂CH₃. In some embodiments, **R^{2B}** is a 4-, 5-, 6-, 7-, 8-, 9-, 10-, or 11-membered bicyclic heterocyclyl ring system having 1 to 5 ring forming heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the heterocyclyl is unsubstituted or substituted with one or more **R^B**; and each **R^B** is independently selected from the group consisting of: fluoro, chloro, bromo, -CN, -CHF₂, -CF₃, -OH, and -OCH₃. In some embodiments, **R^{2B}** is a 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two **R^B**; and each **R^B** is independently selected from the group consisting of: fluoro and -OH. In some embodiments, **R^{2B}** is a C₂-C₁₂ 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, up to 13-membered monocyclic, bicyclic, or tricyclic heterocyclyl ring system having 1 to 5 ring forming heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the heterocyclyl is unsubstituted or substituted with one or more **R^B**; and each **R^B** is independently selected from the group consisting of: fluoro, chloro, bromo, -CN, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -OH, -OCF₃, -OCH₃, and -OCH₂CH₃. In some embodiments, **R^{2B}** is a C₂-C₆ 3-, 4-, 5-, 6-, or 7-membered monocyclic heterocyclyl ring having 1 to 5 ring forming heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the heterocyclyl is unsubstituted or substituted with one or more **R^B**; and each **R^B** is independently selected from the group consisting of: fluoro, chloro, bromo, -CN, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -OH, -OCF₃, -OCH₃, and -OCH₂CH₃. In some embodiments, **R^{2B}** is a C₂-C₁₀ 4-, 5-, 6-, 7-, 8-, 9-, 10-, or 11-membered bicyclic heterocyclyl ring system having 1 to 5 ring forming heteroatoms selected from nitrogen, oxygen, and sulfur, wherein the heterocyclyl is unsubstituted or substituted with one or more **R^B**; and each **R^B** is independently selected from the group consisting of: fluoro, chloro, -CN, -CHF₂, -CF₃, -OH, and -OCH₃. In some embodiments, **R^{2B}** is a C₃-C₄ 4-5 membered monocyclic heterocyclyl having one ring forming heteroatom selected from oxygen and sulfur, wherein the C₃-C₄ 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or more **R^B**; and each **R^B** is independently selected from the group consisting of: fluoro and -OH. In some embodiments, **R^{2B}** is a C₃-C₄ 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the C₃-C₄ 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two **R^B**; and each **R^B** is

independently selected from the group consisting of: fluoro and -OH. In some embodiments, **R**^{2B} is oxetan-3-yl or oxolan-3-yl, each unsubstituted or substituted with one fluoro.

In some embodiments, **R**^{2B} is: 1,2,4-oxadiazol-5-yl, 1,3-oxazol-2-yl, 1,4-dioxepan-6-yl, 1*H*-imidazol-2-yl, 1*H*-pyrazol-5-yl, 1*H*-tetrazol-5-yl, azetidin-3-yl, bicyclo[1.1.1]pentan-1-yl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopropyl, furan-2-yl, imidazol-1-yl, isoxazol-3-yl, isoxazol-5-yl, oxazol-4-yl, oxetan-2-yl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), phenyl, piperidin-4-yl, pyrazin-2-yl, pyridin-2-yl, pyridin-3-yl, pyrimidin-2-yl, pyrrolidin-1-yl, tetrahydro-2*H*-pyran-4-yl, tetrahydro-2*H*-thiopyran-4-yl, or thiazol-4-yl; each unsubstituted or substituted with one or more groups independently selected from the group consisting of: -CH₂CN, -CN, difluoromethyl, ethyl, fluoro, methyl, methylsulfonyl, methylsulfonyloxy, -OH, oxo, phenyl, and trifluoromethyl.


In some embodiments, **R**^{2B} is 1-((methylsulfonyl)oxy)cyclopropyl, 1-(cyanomethyl)cyclopropyl, 1-(difluoromethyl)cyclopropyl, 1-(methylsulfonyl)cyclopropyl, 1-(trifluoromethyl)cyclopentyl, 1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl, 1,2,4-oxadiazol-5-yl, 1-cyanocyclobutyl, 1-cyanocyclopropyl, 1-cyanopiperidin-4-yl, 1-fluorocyclobutyl, 1-fluorocyclopropyl, 1*H*-imidazol-2-yl, 1*H*-tetrazol-5-yl, 1-hydroxycyclobutyl, 1-methylazetidin-3-yl, 2-ethylcyclopropyl, 2-fluorocyclopentyl, 3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl, 3,3-difluorocyclopentyl, 3,3-difluoropyrrolidin-1-yl, 3-fluorobicyclo[1.1.1]pentan-1-yl, 3-fluorooxetan-3-yl, 3-fluorophenyl, 3-hydroxy-3-(trifluoromethyl)cyclobutyl, 4-((difluoromethyl)sulfonyl)phenyl, 4-(trifluoromethyl)phenyl, 4,4-difluoro-1-hydroxycyclohexyl, 4-fluorophenyl, 5-cyanofuran-2-yl, 5-cyanopyridin-2-yl, 5-methyl-1,3-oxazol-2-yl, 6-cyanopyridin-2-yl, 6-fluoro-1,4-dioxepan-6-yl, cyclobutyl, cyclopropyl, imidazol-1-yl, isoxazol-3-yl, isoxazol-5-yl, oxazol-4-yl, oxetan-2-yl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), phenyl, pyridin-3-yl, pyrimidin-2-yl, 3-cyanophenyl, or 2,2-difluorocyclopropyl.

In some embodiments, **R**^{2B} is 1-((methylsulfonyl)oxy)cyclopropyl. In some embodiments, **R**^{2B} is 1-(cyanomethyl)cyclopropyl. In some embodiments, **R**^{2B} is 1-(difluoromethyl)cyclopropyl. In some embodiments, **R**^{2B} is 1-(methylsulfonyl)cyclopropyl. In some embodiments, **R**^{2B} is 1-(trifluoromethyl)cyclopentyl. In some embodiments, **R**^{2B} is 1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl. In some embodiments, **R**^{2B} is 1,2,4-oxadiazol-5-yl. In some embodiments, **R**^{2B} is 1-cyanocyclobutyl. In some embodiments, **R**^{2B} is 1-cyanocyclopropyl. In some embodiments, **R**^{2B} is 1-cyanopiperidin-4-yl. In some embodiments, **R**^{2B} is 1-fluorocyclobutyl. In some embodiments, **R**^{2B} is 1-fluorocyclopropyl. In some embodiments, **R**^{2B} is 1*H*-imidazol-2-yl. In some embodiments, **R**^{2B} is 1*H*-tetrazol-5-yl. In some embodiments, **R**^{2B} is 1-hydroxycyclobutyl. In some embodiments, **R**^{2B} is 1-methylazetidin-3-yl. In some embodiments, **R**^{2B} is 2-ethylcyclopropyl. In some embodiments, **R**^{2B} is 2-fluorocyclopentyl. In some embodiments, **R**^{2B} is 2-methylthiazol-4-yl. In some embodiments, **R**^{2B} is 3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl. In some embodiments, **R**^{2B} is 3,3-difluorocyclopentyl. In some embodiments, **R**^{2B} is 3,3-difluoropyrrolidin-1-yl. In some embodiments, **R**^{2B} is 3-fluorobicyclo[1.1.1]pentan-1-yl. In some embodiments, **R**^{2B} is 3-fluorooxetan-3-yl. In some

embodiments, R^{2B} is 3-fluorophenyl. In some embodiments, R^{2B} is 3-hydroxy-3-(trifluoromethyl)cyclobutyl. In some embodiments, R^{2B} is 4-((difluoromethyl)sulfonyl)phenyl. In some embodiments, R^{2B} is 4-(trifluoromethyl)phenyl. In some embodiments, R^{2B} is 4,4-difluoro-1-hydroxycyclohexyl. In some embodiments, R^{2B} is 4-fluorophenyl. In some embodiments, R^{2B} is 5-cyanofuran-2-yl. In some embodiments, R^{2B} is 5-cyanopyridin-2-yl. In some embodiments, R^{2B} is 5-methyl-1,3-oxazol-2-yl. In some embodiments, R^{2B} is 6-cyanopyridin-2-yl. In some embodiments, R^{2B} is 6-fluoro-1,4-dioxepan-6-yl. In some embodiments, R^{2B} is cyclobutyl. In some embodiments, R^{2B} is cyclopropyl. In some embodiments, R^{2B} is imidazol-1-yl. In some embodiments, R^{2B} is isoxazol-3-yl. In some embodiments, R^{2B} is isoxazol-4-yl. In some embodiments, R^{2B} is isoxazol-5-yl. In some embodiments, R^{2B} is oxazol-4-yl. In some embodiments, R^{2B} is oxetan-2-yl. In some embodiments, R^{2B} is oxetan-3-yl. In some embodiments, R^{2B} is oxolan-3-yl (tetrahydrofuran-3-yl). In some embodiments, R^{2B} is phenyl. In some embodiments, R^{2B} is pyridin-3-yl. In some embodiments, R^{2B} is pyrimidin-2-yl. In some embodiments, R^{2B} is 3-cyanophenyl. In some embodiments, R^{2B} is 2,2-difluorocyclopropyl.


In some embodiments, R^1 is $-(CH_2)_n-OR^{2C}$.


In some embodiments, the compound is a compound of Formula (V):



or a pharmaceutically acceptable salt thereof, wherein: n and R^{2C} have the same definitions as described herein, and each can be selected independently from any of the embodiments as described herein, *supra*, and *infra*.

In some embodiments, the compound is a compound selected from Formula (Va), Formula (Vc), Formula (Ve), or Formula (Vg):





or a pharmaceutically acceptable salt thereof, wherein: **n** and **R^{2C}** have the same definitions as described herein, and each can be selected independently from any of the embodiments as described herein, *supra*, and *infra*.

In some embodiments, the compound of Formula (I) is a compound of Formula (Va). In some embodiments, the compound of Formula (I) is a compound of Formula (Vc). In some embodiments, the compound of Formula (I) is a compound of Formula (Ve). In some embodiments, the compound of Formula (I) is a compound of Formula (Vg).

In some embodiments, **n** is 1, 2, or 3.

In some embodiments, **n** is 1 or 2.

In some embodiments, **n** is 1.

In some embodiments, **n** is 2.

In some embodiments, **n** is 3.

In some embodiments, **R^{2C}** is C₁-C₄alkyl, C₃-C₈cycloalkyl, haloC₁-C₄alkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more **R^C**. It is understood that when more than one **R^C** is present, they may be the same or different. In some embodiments, when more than one **R^C** is present, they are the same. In some embodiments, when more than one **R^C** is present, they are different.

In some embodiments, **R^{2C}** is C₁-C₄alkyl, C₃-C₈cycloalkyl, haloC₁-C₄alkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one, two, three, or four **R^C**. In some embodiments, **R^{2C}** is C₁-C₄alkyl, C₃-C₈cycloalkyl, haloC₁-C₄alkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one **R^C**. In some embodiments, **R^{2C}** is C₁-C₄alkyl, C₃-C₈cycloalkyl, haloC₁-C₄alkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with two **R^C**. In some embodiments, **R^{2C}** is C₁-C₄alkyl, C₃-C₈cycloalkyl, haloC₁-C₄alkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with three **R^C**. In some embodiments, **R^{2C}** is C₁-C₄alkyl, C₃-C₈cycloalkyl, haloC₁-C₄alkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with four **R^C**.

In some embodiments, **R^{2C}** is C₁-C₄alkyl, unsubstituted or substituted with one or more **R^C**, wherein **R^C** is as described herein *supra*, and *infra*. In some embodiments, **R^{2C}** is ethyl, isopropyl, or methyl, each unsubstituted or substituted with one or more **R^C**.

In some embodiments, each **R^C** is independently -CN or C₃-C₆cycloalkyl. In some embodiments, each **R^C** is independently selected from the group consisting of: cyclopropyl and -CN.

In some embodiments, R^{2C} is ethyl, isopropyl, or methyl, each unsubstituted or substituted with one or more R^C independently selected from the group consisting of: cyclopropyl and -CN. In some embodiments, R^{2C} is methyl, cyano(cyclopropyl)methyl, 2-cyanopropan-2-yl, cyanomethyl, or 1-cyanoethyl.

5 In some embodiments, R^{2C} is C_3 - C_8 cycloalkyl, unsubstituted or substituted with one or more R^C , wherein R^C is as described herein *supra*, and *infra*.

In some embodiments, R^{2C} is cyclobutyl, unsubstituted or substituted with one or more R^C .

In some embodiments, each R^C is -CN.

In some embodiments, R^{2C} is cyclobutyl, unsubstituted or substituted with one or more -CN.

10 In some embodiments, R^{2C} is 1-cyanocyclobutyl.

In some embodiments, R^{2C} is 4-7 membered heterocyclyl, unsubstituted or substituted with one or more R^C , wherein R^C is as described herein *supra*, and *infra*.

In some embodiments, R^{2C} is oxetan-3-yl or tetrahydro-2H-pyran-4-yl, each unsubstituted or substituted with one or more R^C . In some embodiments, R^C is -CN.

15 In some embodiments, R^{2C} is oxetan-3-yl or tetrahydro-2H-pyran-4-yl, each unsubstituted or substituted with one or more -CN. In some embodiments, R^{2C} is oxetan-3-yl or 4-cyanotetrahydro-2H-pyran-4-yl.

In some embodiments, R^{2C} is C_1 - C_4 alkyl or C_3 - C_5 cycloalkyl, each substituted with one, two, three, or four R^C ; and each R^C is independently -CN or C_3 - C_6 cycloalkyl. In some embodiments, R^{2C} is C_1 - C_4 alkyl or C_3 - C_5 cycloalkyl, each substituted with one or more R^C ; and each R^C is independently -CN or C_3 - C_6 cycloalkyl. In some embodiments, R^{2C} is cyclobutyl substituted with one -CN. In some
20
embodiments, R^{2C} is C_1 - C_3 alkyl substituted with one -CN and one C_3 - C_5 cycloalkyl. In some
embodiments, R^{2C} is C_1 - C_2 alkyl substituted with one -CN and one C_3 - C_4 cycloalkyl. In some
embodiments, R^{2C} is C_1 - C_2 alkyl substituted with one -CN and one cyclopropyl. In some
25
embodiments, R^{2C} is C_1 - C_3 alkyl substituted with one -CN.

In some embodiments, R^{2C} is ethyl, isopropyl, methyl, cyclobutyl, oxetan-3-yl, or tetrahydro-2H-pyran-4-yl, each unsubstituted or substituted with one or more groups independently selected from the group consisting of: cyclopropyl and -CN.

In some embodiments, R^{2C} is methyl, cyano(cyclopropyl)methyl, 2-cyanopropan-2-yl, cyanomethyl, 1-cyanoethyl, 1-cyanocyclobutyl, oxetan-3-yl, or 4-cyanotetrahydro-2H-pyran-4-yl.
30

In some embodiments, R^{2C} is methyl. In some embodiments, R^{2C} is cyano(cyclopropyl)methyl. In some embodiments, R^{2C} is 2-cyanopropan-2-yl. In some embodiments, R^{2C} is cyanomethyl. In some embodiments, R^{2C} is 1-cyanoethyl. In some embodiments, R^{2C} is 1-cyanocyclobutyl. In some embodiments, R^{2C} is oxetan-3-yl. In some embodiments, R^{2C} is 4-cyanotetrahydro-2H-pyran-4-yl.
35

R¹ Groups

In some embodiments, **R¹** is methylallyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, ethyl, 2-fluoroethyl, 3-fluoropropyl, propan-2-yl, 1,1,1-trifluoropropan-2-yl, 2-hydroxy-2-methylpropyl, 3,3,3-trifluoro-2-hydroxypropyl, 3,3-difluoro-2-hydroxypropyl, methyl, 2-hydroxypropyl, 2-hydroxybutyl, 4,4,4-trifluoro-2-hydroxybutyl, 3,3,3-trifluoro-2-hydroxy-2-methylpropyl, 2,3-dihydroxypropyl, 3-hydroxypropyl, 2-hydroxyethyl, cyanomethyl, 2-(dimethylamino)-2-oxoethyl, 2-(methylamino)-2-oxoethyl, 2-hydroxy-3-methoxypropyl, 2-chloro-2,2-difluoroethyl, 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl, 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl, 3-(methylsulfonyl)propyl, (*N,N*-dimethylsulfamoyl)methyl, 2-methoxypropyl, 2-ethyl-2-hydroxybutyl, 3-chloro-3,3-difluoropropyl, 2-hydroxy-3-morpholinopropyl, cyclopropyl, cyclobutyl, 4-(trifluoromethyl)cyclohexyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl, 1-methylazetidin-3-yl, 1-((methylsulfonyl)oxy)cyclopropyl)methyl, (1-(cyanomethyl)cyclopropyl)methyl, (1-(difluoromethyl)cyclopropyl)methyl, (1-(methylsulfonyl)cyclopropyl)methyl, (1-(trifluoromethyl)cyclopentyl)methyl, (1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl)methyl, (1,2,4-oxadiazol-5-yl)methyl, (1-cyanocyclobutyl)methyl, (1-cyanocyclopropyl)methyl, (1-cyanopiperidin-4-yl)methyl, (1-fluorocyclobutyl)methyl, (1-fluorocyclopropyl)methyl, (1*H*-imidazol-2-yl)methyl, (1*H*-tetrazol-5-yl)methyl, (1-hydroxycyclobutyl)methyl, (1-methylazetidin-3-yl)methyl, (2-ethylcyclopropyl)methyl, (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)methyl, (3-fluorobicyclo[1.1.1]pentan-1-yl)methyl, (3-fluorooxetan-3-yl)methyl, (3-hydroxy-3-(trifluoromethyl)cyclobutyl)methyl, (4,4-difluoro-1-hydroxycyclohexyl)methyl, (5-cyanofuran-2-yl)methyl, (5-cyanopyridin-2-yl)methyl, (5-methyl-1,3-oxazol-2-yl)methyl, (6-cyanopyridin-2-yl)methyl, (6-fluoro-1,4-dioxepan-6-yl)methyl, (isoxazol-3-yl)methyl, (isoxazol-5-yl)methyl, (oxazol-4-yl)methyl, (oxetan-2-yl)methyl, (oxetan-3-yl)methyl, (pyridin-3-yl)methyl, 2-(1-cyanocyclopropyl)ethyl, 2-(1*H*-imidazol-1-yl)ethyl, 2-(3,3-difluoropyrrolidin-1-yl)ethyl, 3-fluorobenzyl, 4-((difluoromethyl)sulfonyl)benzyl, 4-(trifluoromethyl)benzyl, 4-(trifluoromethyl)phenethyl, 4-fluorobenzyl, benzyl, cyclopropylmethyl, pyrimidin-2-ylmethyl, (2-cyanopropan-2-yl)oxy)ethyl, 2-((4-cyanotetrahydro-2*H*-pyran-4-yl)oxy)ethyl, 2-(1-cyanocyclobutoxy)ethyl, 2-(1-cyanoethoxy)ethyl, 2-(cyano(cyclopropyl)methoxy)ethyl, 2-(cyanomethoxy)ethyl, 2-(oxetan-3-yloxy)ethyl, 2-methoxyethyl, 5,5,5-trifluoropentyl, 3-cyanobenzyl, 4-fluorophenethyl, cyclobutylmethyl, 4,4,4-trifluorobutyl, or (2,2-difluorocyclopropyl)methyl.

In some embodiments, **R¹** is methylallyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, ethyl, 2-fluoroethyl, 3-fluoropropyl, propan-2-yl, 1,1,1-trifluoropropan-2-yl, 2-hydroxy-2-methylpropyl, 3,3,3-trifluoro-2-hydroxypropyl, 3,3-difluoro-2-hydroxypropyl, methyl, 2-hydroxypropyl, 2-hydroxybutyl, 4,4,4-trifluoro-2-hydroxybutyl, 3,3,3-trifluoro-2-hydroxy-2-methylpropyl, 2,3-dihydroxypropyl, 3-hydroxypropyl, 2-hydroxyethyl,

cyanomethyl, 2-(dimethylamino)-2-oxoethyl, 2-(methylamino)-2-oxoethyl, 2-hydroxy-3-methoxypropyl, 2-chloro-2,2-difluoroethyl, 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl, 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl, 3-(methylsulfonyl)propyl, (*N,N*-dimethylsulfamoyl)methyl, 2-methoxypropyl, 2-ethyl-2-hydroxybutyl, 3-chloro-3,3-difluoropropyl, 2-hydroxy-3-morpholinopropyl, cyclopropyl, cyclobutyl, 4-(trifluoromethyl)cyclohexyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl, 1-methylazetidin-3-yl, 5,5,5-trifluoropentyl, or 4,4,4-trifluorobutyl.

In some embodiments, R^1 is (1-((methylsulfonyl)oxy)cyclopropyl)methyl, (1-(cyanomethyl)cyclopropyl)methyl, (1-(difluoromethyl)cyclopropyl)methyl, (1-(methylsulfonyl)cyclopropyl)methyl, (1-(trifluoromethyl)cyclopentyl)methyl, (1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl)methyl, (1,2,4-oxadiazol-5-yl)methyl, (1-cyanocyclobutyl)methyl, (1-cyanocyclopropyl)methyl, (1-cyanopiperidin-4-yl)methyl, (1-fluorocyclobutyl)methyl, (1-fluorocyclopropyl)methyl, (1*H*-imidazol-2-yl)methyl, (1*H*-tetrazol-5-yl)methyl, (1-hydroxycyclobutyl)methyl, (1-methylazetidin-3-yl)methyl, (2-ethylcyclopropyl)methyl, (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)methyl, (3-fluorobicyclo[1.1.1]pentan-1-yl)methyl, (3-fluorooxetan-3-yl)methyl, (3-hydroxy-3-(trifluoromethyl)cyclobutyl)methyl, (4,4-difluoro-1-hydroxycyclohexyl)methyl, (5-cyanofuran-2-yl)methyl, (5-cyanopyridin-2-yl)methyl, (5-methyl-1,3-oxazol-2-yl)methyl, (6-cyanopyridin-2-yl)methyl, (6-fluoro-1,4-dioxepan-6-yl)methyl, (isoxazol-3-yl)methyl, (isoxazol-5-yl)methyl, (oxazol-4-yl)methyl, (oxetan-2-yl)methyl, (oxetan-3-yl)methyl, (pyridin-3-yl)methyl, 2-(1-cyanocyclopropyl)ethyl, 2-(1*H*-imidazol-1-yl)ethyl, 2-(3,3-difluoropyrrolidin-1-yl)ethyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, 3-fluorobenzyl, 4-((difluoromethyl)sulfonyl)benzyl, 4-(trifluoromethyl)benzyl, 4-(trifluoromethyl)phenethyl, 4-fluorobenzyl, benzyl, cyclobutyl, cyclopropyl, cyclopropylmethyl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), pyrimidin-2-ylmethyl, 3-cyanobenzyl, 4-fluorophenethyl, cyclobutylmethyl, or (2,2-difluorocyclopropyl)methyl.

In some embodiments, R^1 is (2-cyanopropan-2-yl)oxyethyl, 2-((4-cyanotetrahydro-2*H*-pyran-4-yl)oxy)ethyl, 2-(1-cyanocyclobutoxy)ethyl, 2-(1-cyanoethoxy)ethyl, 2-(cyano(cyclopropyl)methoxy)ethyl, 2-(cyanomethoxy)ethyl, 2-(oxetan-3-yl)oxyethyl, or 2-methoxyethyl.

In some embodiments, R^1 is methylallyl. In some embodiments, R^1 is 2,2,2-trifluoroethyl. In some embodiments, R^1 is 2,2-difluoroethyl. In some embodiments, R^1 is 2,2-difluoropropyl. In some embodiments, R^1 is 3,3,3-trifluoropropyl. In some embodiments, R^1 is ethyl. In some embodiments, R^1 is 2-fluoroethyl. In some embodiments, R^1 is 3-fluoropropyl. In some embodiments, R^1 is propan-2-yl. In some embodiments, R^1 is 1,1,1-trifluoropropan-2-yl. In some embodiments, R^1 is 2-hydroxy-2-methylpropyl. In some embodiments, R^1 is 3,3,3-trifluoro-2-hydroxypropyl. In some embodiments, R^1 is 3,3-difluoro-2-hydroxypropyl. In some embodiments, R^1 is methyl. In some embodiments, R^1 is

2-hydroxypropyl. In some embodiments, R^1 is 2-hydroxybutyl. In some embodiments, R^1 is 4,4,4-trifluoro-2-hydroxybutyl. In some embodiments, R^1 is 3,3,3-trifluoro-2-hydroxy-2-methylpropyl. In some embodiments, R^1 is 2,3-dihydroxypropyl. In some embodiments, R^1 is 3-hydroxypropyl. In some embodiments, R^1 is 2-hydroxyethyl. In some embodiments, R^1 is cyanomethyl. In some
5 embodiments, R^1 is 2-(dimethylamino)-2-oxoethyl. In some embodiments, R^1 is 2-(methylamino)-2-oxoethyl. In some embodiments, R^1 is 2-hydroxy-3-methoxypropyl. In some embodiments, R^1 is 2-chloro-2,2-difluoroethyl. In some embodiments, R^1 is 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl. In some embodiments, R^1 is 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl. In some embodiments, R^1 is 3-(methylsulfonyl)propyl. In some embodiments, R^1 is (*N,N*-dimethylsulfamoyl)methyl. In some
10 embodiments, R^1 is 2-methoxypropyl. In some embodiments, R^1 is 2-ethyl-2-hydroxybutyl. In some embodiments, R^1 is 3-chloro-3,3-difluoropropyl. In some embodiments, R^1 is 2-hydroxy-3-morpholinopropyl. In some embodiments, R^1 is cyclopropyl. In some embodiments, R^1 is cyclobutyl. In some embodiments, R^1 is 4-(trifluoromethyl)cyclohexyl. In some embodiments, R^1 is 2-fluorocyclopentyl. In some embodiments, R^1 is 3,3-difluorocyclopentyl. In some embodiments, R^1 is
15 oxetan-3-yl. In some embodiments, R^1 is oxolan-3-yl (tetrahydrofuran-3-yl). In some embodiments, R^1 is 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl. In some embodiments, R^1 is 1-methylazetidin-3-yl. In some embodiments, R^1 is (1-((methylsulfonyl)oxy)cyclopropyl)methyl. In some embodiments, R^1 is (1-(cyanomethyl)cyclopropyl)methyl. In some embodiments, R^1 is (1-(difluoromethyl)cyclopropyl)methyl. In some embodiments, R^1 is (1-(methylsulfonyl)cyclopropyl)methyl. In some embodiments, R^1 is (1-(trifluoromethyl)cyclopentyl)methyl. In some embodiments, R^1 is (1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl)methyl. In some embodiments, R^1 is (1,2,4-oxadiazol-5-yl)methyl. In some
20 embodiments, R^1 is (1-cyanocyclobutyl)methyl. In some embodiments, R^1 is (1-cyanocyclopropyl)methyl. In some embodiments, R^1 is (1-cyanopiperidin-4-yl)methyl. In some
25 embodiments, R^1 is (1-fluorocyclobutyl)methyl. In some embodiments, R^1 is (1-fluorocyclopropyl)methyl. In some embodiments, R^1 is (1*H*-imidazol-2-yl)methyl. In some
embodiments, R^1 is (1*H*-tetrazol-5-yl)methyl. In some embodiments, R^1 is (1-hydroxycyclobutyl)methyl. In some embodiments, R^1 is (1-methylazetidin-3-yl)methyl. In some
embodiments, R^1 is (2-ethylcyclopropyl)methyl. In some embodiments, R^1 is (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)methyl. In some embodiments, R^1 is (3-fluorobicyclo[1.1.1]pentan-1-yl)methyl. In some embodiments, R^1 is (3-fluorooxetan-3-yl)methyl. In
30 some embodiments, R^1 is (3-hydroxy-3-(trifluoromethyl)cyclobutyl)methyl. In some embodiments, R^1 is (4,4-difluoro-1-hydroxycyclohexyl)methyl. In some embodiments, R^1 is (5-cyanofuran-2-yl)methyl. In some embodiments, R^1 is (5-cyanopyridin-2-yl)methyl. In some embodiments, R^1 is (5-methyl-1,3-oxazol-2-yl)methyl. In some embodiments, R^1 is (6-cyanopyridin-2-yl)methyl. In some
35 embodiments, R^1 is (6-fluoro-1,4-dioxepan-6-yl)methyl. In some embodiments, R^1 is (isoxazol-3-yl)methyl. In some embodiments, R^1 is (isoxazol-5-yl)methyl. In some embodiments, R^1 is (oxazol-4-

yl)methyl. In some embodiments, R^1 is (oxetan-2-yl)methyl. In some embodiments, R^1 is (oxetan-3-yl)methyl. In some embodiments, R^1 is (pyridin-3-yl)methyl. In some embodiments, R^1 is 2-(1-cyanocyclopropyl)ethyl. In some embodiments, R^1 is 2-(1*H*-imidazol-1-yl)ethyl. In some embodiments, R^1 is 2-(3,3-difluoropyrrolidin-1-yl)ethyl. In some embodiments, R^1 is 3-fluorobenzyl.

5 In some embodiments, R^1 is 4-((difluoromethyl)sulfonyl)benzyl. In some embodiments, R^1 is 4-(trifluoromethyl)benzyl. In some embodiments, R^1 is 4-(trifluoromethyl)phenethyl. In some embodiments, R^1 is 4-fluorobenzyl. In some embodiments, R^1 is benzyl. In some embodiments, R^1 is cyclopropylmethyl. In some embodiments, R^1 is pyrimidin-2-ylmethyl. In some embodiments, R^1 is (2-cyanopropan-2-yl)oxy)ethyl. In some embodiments, R^1 is 2-((4-cyanotetrahydro-2*H*-pyran-4-yl)oxy)ethyl. In some embodiments, R^1 is 2-(1-cyanocyclobutoxy)ethyl. In some embodiments, R^1 is 2-(1-cyanoethoxy)ethyl. In some embodiments, R^1 is 2-(cyano(cyclopropyl)methoxy)ethyl. In some

10 embodiments, R^1 is 2-(cyanomethoxy)ethyl. In some embodiments, R^1 is 2-(oxetan-3-yloxy)ethyl. In some embodiments, R^1 is 2-methoxyethyl. In some embodiments, R^1 is 5,5,5-trifluoropentyl. In some embodiments, R^1 is 3-cyanobenzyl. In some embodiments, R^1 is 4-fluorophenethyl. In some


15 embodiments, R^1 is cyclobutylmethyl. In some embodiments, R^1 is 4,4,4-trifluorobutyl. In some embodiments, R^1 is (2,2-difluorocyclopropyl)methyl. It is understood that when the R^1 group has a chiral carbon, the chiral carbon can be (*R*) or (*S*). In some embodiments, when the R^1 group has a chiral carbon, the chiral carbon is (*R*). In some embodiments, when the R^1 group has a chiral carbon, the chiral carbon is (*S*). Further, it is understood that when the R^1 group has two chiral carbons, the


20 chiral carbons can be independently (*R*) or (*S*). In some embodiments, when the R^1 group has two chiral carbons, both chiral carbons are (*R*). In some embodiments, when the R^1 group has two chiral carbons, both chiral carbons are (*S*). In some embodiments, when the R^1 group has two chiral carbons, one chiral carbon is (*R*) and the other chiral carbon is (*S*).

In some embodiments, R^1 is $-CH_3$. In some embodiments, R^1 is $-CD_3$. In some embodiments, R^1 is $-CH_2CH_3$. In some embodiments, R^1 is $-CD_2CD_3$. In some embodiments, R^1 is $-CH_2CH_2OCH_3$, $-CH_2CH(OH)CH_3$, $-CH_2CH(OH)CF_3$, $-CH_2CF_3$, $-CH_2CF_2H$, $-CH_2CH_2F$, $-CH_2CH_2CF_3$, $-CH_2CH_2CHF_2$, $-CH_2CH_2CH_2F$, or $-CH(CH_3)(CF_3)$. In some embodiments, R^1 is $-CH_2CF_3$, $-CH_2CF_2H$, $-CH_2CH_2CF_3$, or


30 $-CH_2CH_2CHF_2$.

In some embodiments, R^1 is $-CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CN$, $-CH_2CF_3$, $-CH(CH_3)(CF_3)$, $-CH_2CF_2H$, $-CH_2CF_2H$, $-CH_2CF_2CH_3$, $-CH_2CH_2CF_3$, $-CH_2CH_2CH_2H$, $-CH_2CH_2OCH_3$,






5 In some embodiments, R¹ is -CH₂CH₃, -CH(CH₃)₂, -CH₂CF₃, -CH(CH₃)(CF₃), -CH₂CF₂H, -CH₂CF₂H, -CH₂CF₂CH₃, -CH₂CH₂CF₃, -CH₂CH₂CH₂H, -CH₂CH₂OCH₃,









10 In some embodiments, R¹ is -CH₃, -CH₂CH₃, -CH(CH₃)₂, -CH₂CH(CH₃)₂, -CH₂CN,















Some embodiments include every combination of one or more compounds and pharmaceutically acceptable salts thereof selected from the following group shown in **Table A**. Some
 15 embodiments include every combination of one or more compounds selected from the following group shown in **Table A**.

Table A







Cmpd No.	Chemical Structure	Chemical Name
4-1		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-2		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-3		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-4		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-5		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-6		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol






Cmpd No.	Chemical Structure	Chemical Name
4-7		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-8		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-9		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-10		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-11		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-12		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-13		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-14		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-15		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-16		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-17		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-ethoxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-18		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(2-fluoroethoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-19		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(3-fluorooxetan-3-yl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-20		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[2-(oxetan-3-yloxy)ethoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-21		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>S</i>)-oxetan-2-ylmethoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-22		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-23		1-(2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethoxy)cyclobutane-1-carbonitrile
4-24		2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}acetonitrile







Cmpd No.	Chemical Structure	Chemical Name
4-25		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-hydroxycyclobutyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-26		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2-methoxyethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-27		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-cyclopropoxy-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-28		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(3-fluoropropoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-29		1-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)cyclopropane-1-carbonitrile
4-30		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-yloxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-31		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclopropyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-32		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-cyclobutoxy-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-33		2-([(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}- <i>N,N</i> -dimethylacetamide
4-34		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[2-(1 <i>H</i> -imidazol-1-yl)ethoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-35		2-([(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}- <i>N</i> -methylacetamide
4-36		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(5-methyl-1,3-oxazol-2-yl)methoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-37		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1-methylazetidin-3-yl)methoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-38		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(propan-2-yloxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-39		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1,1,1-trifluoropropan-2-yl)oxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-40		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(ethoxy- <i>d</i> ₅)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol or (2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(² H ₅)ethoxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-41		4-(2-([(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethoxy)oxanc-4-carbonitrile
4-42		1-([(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)cyclobutane-1-carbonitrile







Cmpd No.	Chemical Structure	Chemical Name
4-43		2-(2-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethoxy)-2-cyclopropylacetonitrile
4-44		2-(2-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethoxy)-2-methylpropanenitrile
4-45		1-(2-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethyl)cyclopropane-1-carbonitrile
4-46		2-[1-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)cyclopropyl]acetonitrile
4-47		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(2-hydroxy-2-methylpropoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-48		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>S</i>)-2-hydroxy-3-methoxypropoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol






Cmpd No.	Chemical Structure	Chemical Name
4-49		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>R</i>)-3,3,3-trifluoro-2-hydroxypropoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-50		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>S</i>)-3,3,3-trifluoro-2-hydroxypropoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-51		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(3,3-difluoro-2-hydroxypropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-52		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(methoxy- <i>d</i> ₃)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol or (2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(² H ₃)methoxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-53		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3 <i>R</i>)-oxolan-3-yloxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-54		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3 <i>S</i>)-oxolan-3-yloxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-55		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>R</i>)-2-hydroxypropoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-56		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2-methylprop-2-en-1-yl)oxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-57		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-{[1-(difluoromethyl)cyclopropyl]methoxy}-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-58		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>R</i>)-2-hydroxybutoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-59		5-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)furan-2-carbonitrile
4-60		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-61		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclobutyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-62		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(4,4,4-trifluoro-2-hydroxybutoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-63		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(4-fluorophenyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-64		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(3-fluorophenyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-65		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2-chloro-2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-66		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoro-2-hydroxy-2-methylpropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-67		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{{3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl}methoxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-68		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>S</i>)-2-hydroxybutoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-69		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2-oxazol-3-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-70		2-(2-{{(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl}oxy}ethoxy)acetonitrile
4-71		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[2-(3,3-difluoropyrrolidin-1-yl)ethoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-72		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(6-fluoro-1,4-dioxepan-6-yl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-73		4-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)piperidine-1-carbonitrile
4-74		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{2-[4-(trifluoromethyl)phenyl]ethoxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-75		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,3-oxazol-4-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-76		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[(4,4-difluoro-1-hydroxycyclohexyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-77		6-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)pyridine-2-carbonitrile
4-78		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-{3-(3-hydroxy-3-(trifluoromethyl)cyclobutyl)methoxy}-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol







Cmpd No.	Chemical Structure	Chemical Name
4-79		((2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[2-hydroxy-3-(2,2,2-trifluoroethoxy)propoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-80		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[1-(trifluoromethyl)cyclopentyl]methoxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-81		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[4-(trifluoromethyl)phenyl]methoxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-82		6-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)pyridine-3-carbonitrile
4-83		(2 <i>R</i>)-3-{{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}propane-1,2-diol







Cmpd No.	Chemical Structure	Chemical Name
4-84		2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}- <i>N</i> -methyl- <i>N</i> -(2,2,2-trifluoroethyl)acetamide
4-85		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(3-methanesulfonylpropoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-86		1-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}- <i>N,N</i> -dimethylmethanesulfonamide
4-87		3-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}-1-(2,2,2-trifluoroethyl)pyrrolidin-2-one
4-88		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(cyclopropylmethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-89		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(3-hydroxypropoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol

Cmpd No.	Chemical Structure	Chemical Name
4-90		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-([(1 <i>R</i> ,2 <i>R</i>)-2-ethylcyclopropyl]methoxy}-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-91		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>R</i>)-2-methoxypropoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-92		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>S</i>)-2-methoxypropoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-93		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(2-ethyl-2-hydroxybutoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-94		1-([(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy)methyl)cyclopropyl methanesulfonate
4-95		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2,4-oxadiazol-5-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol

Cmpd No.	Chemical Structure	Chemical Name
4-96		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-methanesulfonylcyclopropyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-97		4-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)-1 <i>λ</i> ⁶ -thiane-1,1-dione
4-98		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(2-hydroxyethoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-99		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[4-(trifluoromethyl)cyclohexyl]oxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-100		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2-fluorocyclopentyl)oxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-101		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[(3,3-difluorocyclopentyl)oxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol

Cmpd No.	Chemical Structure	Chemical Name
4-102		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[(4-difluoromethanesulfonylphenyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-103		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(3-chloro-3,3-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-104		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1-methylazetidin-3-yl)oxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-105		2-(2-[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethoxy)propanenitrile
4-106		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-({3-fluorobicyclo[1.1.1]pentan-1-yl}methoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-107		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(pyridin-3-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol

Cmpd No.	Chemical Structure	Chemical Name
4-108		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(pyrimidin-2-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-109		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2-oxazol-5-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-110		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(1 <i>H</i> -imidazol-2-ylmethoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-111		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1 <i>H</i> -1,2,3,4-tetrazol-5-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-112		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>R</i>)-2-hydroxy-3-methoxypropoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-113		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[2-hydroxy-3-(morpholin-4-yl)propoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol

Cmpd No.	Chemical Structure	Chemical Name
4-114		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>S</i>)-2-hydroxypropoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-115		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(5,5,5-trifluoropentyl)oxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-116		3-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)benzonitrile
4-117		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[2-(4-fluorophenyl)ethoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-118		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclopropyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-119		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(cyclobutylmethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol

Cmpd No.	Chemical Structure	Chemical Name
4-120		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(4,4,4-trifluorobutoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
4-121		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[(2,2-difluorocyclopropyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
5-1		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
5-2		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
5-3		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol
5-4		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol


Each and every compound of Formula (I), method, composition, or use described herein also optionally includes the limitation that the compound of Formula (I) is not one or more compounds selected from the group consisting of: **Compound 4-33, Compound 4-34, Compound 4-35,**

5 **Compound 4-37, Compound 4-95, Compound 4-96, Compound 4-97, Compound 4-100, Compound 4-101, Compound 4-102, Compound 4-103, Compound 4-104, Compound 4-105,**


Compound 4-106, Compound 4-107, Compound 4-108, Compound 4-109, Compound 4-110, Compound 4-111, Compound 4-112, Compound 4-113, and Compound 5-4.

Some embodiments provide a compound of Table 10A and/or Table 10B, or a pharmaceutically acceptable salt thereof. Some embodiments provide a compound having the

5 structure:




10

**Compound 4-39**

or a pharmaceutically acceptable salt thereof.


Some embodiments provide a compound having the structure:



5

or a pharmaceutically acceptable salt thereof.


Some embodiments provide a compound having the structure:



or a pharmaceutically acceptable salt thereof.


10

Some embodiments provide a compound having the structure:




or a pharmaceutically acceptable salt thereof.

Some embodiments provide a compound having the structure:




15 or a pharmaceutically acceptable salt thereof.

Some embodiments provide a compound having the structure:



or a pharmaceutically acceptable salt thereof.


Some embodiments provide a compound having the structure:



5

or a pharmaceutically acceptable salt thereof.


Some embodiments provide a compound having the structure:



or a pharmaceutically acceptable salt thereof.


10

Some embodiments provide a compound having the structure:



or a pharmaceutically acceptable salt thereof.

Some embodiments provide a compound having the structure:



15


or a pharmaceutically acceptable salt thereof.

INTERMEDIATES USEFUL IN PREPARING COMPOUNDS OF FORMULA (I)


Certain intermediates as described herein, *supra*, and *infra*, are novel and useful in the preparation of compounds of Formula (I).

In some embodiments, a compound is selected from the group consisting of: 9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-one (**Compound 1-B**); 9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-ol (**Compound 1-C**); 10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinoline-2,9-diol (**Compound 1-D**); and 9-hydroxy-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-one (**Compound 2-B**), or a salt thereof.

In some embodiments, a compound is: 9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-one (**Compound 1-B**), or a salt thereof. In some embodiments, a compound is selected from the group consisting of: (3*R*,11*bR*)-9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-one (**Compound 1-B (3*R*,11*bR*)**); and (3*S*,11*bS*)-9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-one (**Compound 1-B (3*S*,11*bS*)**), or a salt thereof. In some embodiments, a compound is: (3*R*,11*bR*)-9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-one (**Compound 1-B (3*R*,11*bR*)**), or a salt thereof. In some embodiments, a compound is:



In some embodiments, a compound is: 9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-ol (**Compound 1-C**), or a salt thereof. In some embodiments, the salt is DPTTA. In some embodiments, a compound is: (2*R*,3*R*,11*bR*)-9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinolin-2-ol (**Compound 1-C (2*R*,3*R*,11*bR*)**), or a salt thereof. In some embodiments, the salt is DPTTA. In some embodiments, a compound is:



In some embodiments, a compound is: 10-methoxy-3-neopentyl-1,3,4,6,7,11b-hexahydro-2H-pyrido[2,1-a]isoquinoline-2,9-diol (**Compound 1-D**), or a salt thereof. In some embodiments, a


compound is selected from the group consisting of: (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D**

(**2*R*,3*R*,11*bR***); (2*S*,3*S*,11*bS*)-10-methoxy-3-neopentyl-1,3,4,6,7,11*b*-hexahydro-2*H*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D** (**2*S*,3*S*,11*bS***)); (2*S*,3*R*,11*bR*)-10-methoxy-3-neopentyl-

1,3,4,6,7,11*b*-hexahydro-2*H*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D** (**2*S*,3*R*,11*bR***)); and (2*R*,3*S*,11*bS*)-10-methoxy-3-neopentyl-1,3,4,6,7,11*b*-hexahydro-2*H*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D** (**2*R*,3*S*,11*bS***)), or a salt thereof. In some embodiments, a compound is:

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D** (**2*R*,3*R*,11*bR***)), or a salt thereof. In some embodiments, a

compound is:




In some embodiments, a compound is: 9-hydroxy-10-methoxy-3-neopentyl-1,3,4,6,7,11*b*-hexahydro-2*H*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B**), or a salt thereof. In some

embodiments, a compound is selected from the group consisting of: (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B** (**3*R*,11*bR***)); and (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B** (**3*S*,11*bS***)), or a salt thereof. In some embodiments, a compound is: (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B** (**3*R*,11*bR***)), or

a salt thereof. In some embodiments, a compound is:



It is further appreciated that certain features, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely,

various features which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

PHARMACEUTICAL COMPOSITIONS, FORMULATION, AND DOSAGE FORMS

The present disclosure further provides for pharmaceutical products, such as, pharmaceutical compositions, formulations, unit dosage forms, and kits; each comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof.

5 The present disclosure further provides for pharmaceutical compositions comprising any one of the VMAT2 inhibitor compounds described herein (*e.g.*, a compound of Formula (I), including specific compounds described herein) and at least one pharmaceutically acceptable excipient for use in the methods described here, such as, for treating hyperkinetic movement disorders. A pharmaceutically acceptable excipient is a physiologically and pharmaceutically suitable non-toxic
10 and inactive material or ingredient that does not interfere with the activity of the drug substance; an excipient also can be called a carrier. The formulation methods and excipients described herein are exemplary and are in no way limiting. Pharmaceutically acceptable excipients are well known in the pharmaceutical art and described, for example, in Rowe *et al.*, Handbook of Pharmaceutical Excipients: A Comprehensive Guide to Uses, Properties, and Safety, 5th Ed., 2006, and in Remington:
15 The Science and Practice of Pharmacy (Gennaro, 21st Ed. Mack Pub. Co., Easton, PA (2005)).

The compositions can also be formulated as pills, capsules, granules, or tablets which contain, in addition to an VMAT2 inhibitor, diluents, dispersing and surface active agents, binders, and lubricants. One skilled in this art can further formulate the VMAT2 inhibitor in an appropriate manner, and in accordance with accepted practices, such as, those disclosed in Remington, *supra*.

20 Methods of administration include systemic administration of a VMAT2 inhibitor described herein, preferably in the form of a pharmaceutical composition as discussed above. As used herein, systemic administration includes oral and parenteral methods of administration. For oral administration, suitable pharmaceutical compositions include powders, granules, pills, tablets, and capsules as well as liquids, syrups, suspensions, and emulsions.

25 Pharmaceutical preparations for oral administration can be obtained by any suitable method, typically by uniformly mixing the compound(s) with liquids or finely divided solid carriers, or both, in the required proportions and then, if necessary, processing the mixture, after adding suitable auxiliaries, if desired, forming the resulting mixture into a desired shape to obtain tablets or dragee cores.

30 Conventional excipients, such as, binding agents, fillers, adjuvant, carrier, acceptable wetting agents, tableting lubricants, and disintegrants can be used in tablets and capsules for oral administration. Liquid preparations for oral administration can be in the form of solutions, emulsions, aqueous or oily suspensions, and syrups. Alternatively, the oral preparations can be in the form of dry powder that can be reconstituted with water or another suitable liquid vehicle before use. Parenteral
35 dosage forms can be prepared by dissolving the compound in a suitable liquid vehicle and filter sterilizing the solution before lyophilization, or simply filling and sealing an appropriate vial or ampule.

Some embodiments provide methods for preparing a pharmaceutical composition comprising the step of admixing a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

Formulations suitable for vaginal administration can be presented as pessaries, tampons, 5 creams, gels, pastes, foams, or sprays containing in addition to the drug substance such carriers as are known in the art to be appropriate.

In making pharmaceutical compositions, the drug substance is typically mixed (*i.e.*, admixed) with an excipient, diluted by an excipient or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be 10 a solid, semi-solid, or liquid material, which acts as a vehicle, carrier, or medium for the drug substance. Thus, the compositions can be in the form of tablets, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

For preparing solid form pharmaceutical compositions, such as, powders, tablets, capsules, 15 cachets, suppositories, and dispersible granules an excipient can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These 20 preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

For preparing suppositories, a low melting wax, such as, an admixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by 25 stirring. The molten homogenous mixture is then poured into convenient sized molds, allowed to cool and thereby to solidify.

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water-propylene glycol solutions. Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or 30 wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent.

The pharmaceutical compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles and can contain formulatory agents, such as, suspending, stabilizing and/or dispersing agents. Alternatively, the pharmaceutical compositions can be in powder form, obtained by 35 aseptic isolation of sterile solid or by lyophilization from solution, for constitution with a suitable vehicle, *e.g.* sterile, pyrogen-free water, before use.

The pharmaceutical compositions can be formulated as an aqueous solution, an aqua-alcoholic solution, a solid suspension, an emulsion, a liposomal suspension, or a freeze-dried powder for reconstitution. Such pharmaceutical compositions can be administered directly or as an admixture for further dilution/reconstitution. Route of administration includes intravenous bolus, intravenous
5 infusion, irrigation, and instillation.

Aqueous formulations suitable for oral use can be prepared by dissolving or suspending the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents, as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided drug
10 substance in water with viscous material.

For topical administration to the epidermis the compounds described herein, or pharmaceutically acceptable salts thereof can be formulated as gels, ointments, creams, or lotions, or as a transdermal patch. Also, formulations suitable for topical administration in the mouth include lozenges comprising drug substance in a flavored base.

15 Solutions or suspensions can be applied directly to the nasal cavity by conventional means, for example with a dropper, pipette, or spray. The formulations can be provided in single or multi-dose form. In the latter case of a dropper or pipette, this can be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this can be achieved for example by means of a metering atomizing spray pump.

20 Administration to the respiratory tract can also be achieved by means of an aerosol formulation provided in a pressurized pack with a suitable propellant. If the compounds described herein, or pharmaceutically acceptable salts thereof or pharmaceutical compositions comprising them are administered as aerosols, for example as nasal aerosols or by inhalation, this can be carried out, for example, using a spray, a nebulizer, a pump nebulizer, an inhalation apparatus, a metered inhaler,
25 or a dry powder inhaler.

Alternatively, the pharmaceutical composition can be provided in the form of a dry powder, for example, a powder mix of the compound in a suitable, powder base, such as, lactose, starch, starch derivatives. Conveniently the powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form for example in capsules or cartridges of, *e.g.*, gelatin,
30 or blister packs from which the powder may be administered by means of an inhaler.

The compounds Formula (I), or pharmaceutically acceptable salts thereof can also be administered via a rapid dissolving or a slow-release composition, wherein the composition includes a biodegradable rapid dissolving or slow-release carrier.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the
35 preparation is subdivided into unit doses containing appropriate quantities of the drug substance. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as, packeted tablets, capsules, and powders in vials or ampoules. Also, the unit

dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form. In some embodiments, the pharmaceutical preparation is a tablet or capsule for oral administration. In some embodiments, the pharmaceutical preparation is a liquid formulated for intravenous administration.

5 The compositions can be formulated in a unit dosage form, each dosage containing the drug substance or equivalent mass of the drug substance. The term “unit dosage forms” refers to physically discrete units of a formulation suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable excipient, as described herein.

10 The liquid forms including the drug substance can be incorporated for administration orally or by injection include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils.

 The pharmaceutical compositions described herein can be sterilized by conventional sterilization techniques, or can be sterile filtered. Aqueous solutions can be packaged for use as is, or
15 lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration.

 Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable excipients as described herein. In some
20 embodiments, the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions can be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device, or the nebulizing device can be attached to a face mask tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions can be administered orally or nasally from devices which deliver the formulation in an
25 appropriate manner.

 The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more-unit dosage forms containing the drug substance. The pack may for example comprise metal or plastic foil, such as, a blister pack. The pack or dispenser device can be accompanied by instructions for administration. The pack or dispenser may also be accompanied with
30 a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, can be the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions that can include a compound described herein formulated in a
35 compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

For preparing solid compositions, such as, tablets, the drug substance can be mixed with an excipient to form a solid preformulation composition containing a homogeneous mixture of components. When referring to these preformulation compositions as homogeneous, the drug substance is typically dispersed evenly throughout the composition so that the composition can be readily subdivided into equally effective unit dosage forms, such as, tablets and capsules.

Kits with unit doses of one or more of the compounds described herein, or a pharmaceutically acceptable salt thereof, usually in oral or injectable doses, are provided. Such kits can include a container containing the unit dose, an informational package insert describing the use and attendant benefits of the drugs in treating pathological condition of interest, and optionally an appliance or device for delivery of the composition.

The compounds described herein, or a pharmaceutically acceptable salt thereof, can be effective over a wide dosage range and is generally administered in a therapeutically effective amount. It will be understood, however, that the amount of the compound actually administered will usually be determined by a physician, according to the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual subject, the severity of the subject's symptoms, and the like.

The amount of compound or composition administered to a subject will also vary depending upon what is being administered, the purpose of the administration, such as, prophylaxis or therapy, the state of the subject, the manner of administration, and the like. In therapeutic applications, compositions can be administered to a subject already suffering from a disease in an amount sufficient to cure or at least partially arrest the symptomology and/or pathology of the disease and its complications. Therapeutically effective doses will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors, such as, the severity of the disease, the age, weight, and general condition of the subject, and the like.

The desired dose may conveniently be presented in a single dose or presented as divided doses administered at appropriate intervals, for example, as two, three, four, or more sub-doses per day. The sub-dose itself can be further divided, *e.g.*, into a number of discrete loosely spaced administrations. The daily dose can be divided, especially when relatively large amounts are administered as deemed appropriate, into several, for example two, three, or four-part administrations. If appropriate, depending on individual behavior, it can be necessary to deviate upward or downward from the daily dose indicated.

It will be apparent to those skilled in the art that the dosage forms described herein can comprise a compound described herein or pharmaceutically acceptable salt thereof.

PREPARATIONS

As used herein, a "preparation" is the product of a process used to make or isolate a compound as disclosed and described herein, wherein the preparation contains at least one other

component in addition to the compound. In some embodiments, the preparation comprises a chemical entity.

As used herein, a “chemical entity” defined in the context of a “preparation,” refers to a compound as disclosed and described herein and at least one other component in addition to the compound. For example, a chemical entity can be a co-crystal or salt of a compound as disclosed and described herein.

Some embodiments provide a preparation comprising a compound as disclosed and described herein. In some embodiments, the compound is a component of a chemical entity. In some embodiments, the chemical entity is a salt of a compound as disclosed and described herein. In some embodiments, the chemical entity is a (2*S*,3*S*)-2,3-bis(4-methylbenzoyloxy)butanedioic acid (DPTTA) salt of a compound as disclosed and described herein.

In some embodiments, the compound of the preparation is in at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 99.5%, or 99.9% enantiomeric excess, or an enantiomeric excess within a range defined by any of the preceding numbers. In some embodiments, the compound of the preparation is in at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, or 99.9% enantiomeric excess, or an enantiomeric excess within a range defined by any of the preceding numbers.

In some embodiments, the compound of the preparation is in at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, or 99.9% diastereomeric excess, or a diastereomeric excess within a range defined by any of the preceding numbers. In some embodiments, the compound of the preparation is in at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, or 99.9% diastereomeric excess, or a diastereomeric excess within a range defined by any of the preceding numbers.

In some embodiments, the preparation comprises at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 93% by weight of the compound, or a % by weight within a range defined by any of the preceding numbers. In some embodiments, the preparation comprises at least 50%, 60%, 70%, 80%, 90%, or 93% by weight of the compound, or a % by weight within a range defined by any of the preceding numbers. In some embodiments, the preparation comprises at most 50%, 60%, 70%, 80%, 90%, 93%, or 95% by weight of the compound, or a % by weight within a range defined by any of the preceding numbers.

In some embodiments, the preparation comprises at least 50% by weight of the compound. In some embodiments, the preparation is in the form of a solid, *i.e.*, a solid preparation. In some embodiments, the preparation is used to prepare a pharmaceutical composition.

METHODS OF USE

The compounds of Formula (I) as disclosed and described herein are inhibitors of VMAT2. Accordingly, the present disclosure includes a method of inhibiting VMAT2 (*i.e.*, decreasing at least

one function of VMAT2 or decreasing expression of VMAT2) by contacting the VMAT2 with a compound as disclosed and described herein, or a pharmaceutically acceptable salt thereof. In some embodiments, the contacting can occur *in vitro*, such as, where the VMAT2 is located in a purified preparation or in a cell outside of a living organism (*e.g.*, in a tissue sample or a cellular preparation).

5 In some embodiments, the contacting can occur *in vivo*, such as, where the VMAT2 is located in a living organism.

The VMAT2 inhibitors described herein can reduce the level of monoamines in the central nervous system. Accordingly, the present disclosure includes a method of reducing the level of monoamines in the central nervous system of a subject comprising administering to the subject
10 amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, sufficient to lower the level of monoamines relative to the level prior to administration.

The VMAT2 inhibitors as disclosed and described herein are believed to have utility over a wide range of therapeutic applications, and may be used to treat or prevent a variety of disorders which are caused by or linked to inhibition of the human vesicular monoamine transporter isoform 2.
15 These disorders include neurological and psychiatric disorders, for example, hyperkinetic movement disorders, schizophrenia, and mood disorders. A compound of Formula (I), or a pharmaceutically acceptable salt thereof, can be used in any of the therapeutic methods disclosed and described herein. Also, (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-
a]isoquinolin-2-one (**Compound 5-A (3*R*,11*bR*)**), or a pharmaceutically acceptable salt thereof, or
20 (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-
a]isoquinolin-2-one (**Compound 5-A (3*S*,11*bS*)**) or a pharmaceutically acceptable salt thereof, or an isotopic variant of a **Compound 5-A (3*R*,11*bR*)** or **Compound 5-A (3*S*,11*bS*)** can be used in any of the therapeutic methods disclosed and described herein. In some embodiments, the isotopic variant of
25 a **Compound 5-A (3*R*,11*bR*)** or **Compound 5-A (3*S*,11*bS*)** can have a deuterium in three or more positions in an isotopic distribution in proportional amounts different to those usually found in nature. In some embodiments, a **Compound 5-A (3*R*,11*bR*)** or **Compound 5-A (3*S*,11*bS*)** can have at least 95%, 97%, 99%, 99.5%, or 99.9% deuterium incorporation in three or more positions, or deuterium incorporation in three or more positions within a range defined by any of the preceding numbers.

Accordingly, in various embodiments as disclosed herein, methods are provided for treating
30 or preventing a neurological and/or psychiatric disease or disorder in a subject in need thereof by administering to the subject a pharmaceutically effective amount of a VMAT2 inhibitor as described herein, or a pharmaceutically acceptable salt thereof. The neurological and/or psychiatric disease or disorder can be, for example, a hyperkinetic movement disorder, schizophrenia, schizoaffective disorder, a mood disorder, treatment-refractory obsessive-compulsive disorder, neurological
35 dysfunction associated with Lesch-Nyhan syndrome, agitation associated with Alzheimer's disease, Fragile X syndrome or Fragile X-associated tremor-ataxia syndrome, autism spectrum disorder (*e.g.*,

restricted and repetitive behaviors associated with Autism spectrum disorder (ASD)), Rett syndrome, or chorca-acanthocytosis.

In various other embodiments as disclosed herein, methods are provided for treating or preventing a vesicular monoamine transporter-2 (VMAT2) disease or disorder in a subject in need thereof by administering to the subject a pharmaceutically effective amount of a VMAT2 inhibitor as described herein, or a pharmaceutically acceptable salt thereof. The VMAT2 disease or disorder can be, for example, an ataxias or spinal muscular atrophy; a chorea; a congenital malformation, deformation, or abnormality; a dementia; an oral cavity, salivary gland, or jaw disease; a dyskinesia; a dystonia; an endocrine, nutritional, or metabolic disease; an epilepsy; a habit or impulse disorder; a Huntington's disease or related disorder; a mood or psychotic disorder; a neurotic, stress-related, and somatoform disorder; a degenerative disease of the basal ganglia; an extrapyramidal and movement disorder; a neurological or psychiatric disease or disorder; a nervous system or motor function disorder; a Parkinson's/parkinsonism disorder; a pediatric-onset behavioral and emotional disorder; a pervasive developmental disorder; and a substance abuse or dependence disorder.

Accordingly, in various embodiments as disclosed herein, methods are provided for treating or preventing a hyperkinetic movement disorder in a subject in need thereof by administering to the subject in need thereof a pharmaceutically effective amount of a VMAT2 inhibitor described herein, or a pharmaceutically acceptable salt thereof. In some embodiments, the hyperkinetic movement disorder is tardive dyskinesia, Tourette's syndrome, Huntington's disease, chorca associated with Huntington's disease, or tics. In other embodiments, the hyperkinetic movement disorder is ataxia, chorea, dystonia, hemifacial spasm, myoclonus, restless leg syndrome, or tremors.

In some embodiments, methods are provided for treating or preventing a mood disorder in a subject in need thereof by administering to the subject in need thereof a pharmaceutically effective amount of a VMAT2 inhibitor described herein, or a pharmaceutically acceptable salt thereof. In some embodiments, the mood disorder is bipolar disorder, major depressive disorder, mania in a mood disorder, or depression in a mood disorder.

In some embodiments as disclosed herein, methods are provided for treating or preventing schizophrenia or schizoaffective disorder in a subject in need thereof by administering to the subject in need thereof a pharmaceutically effective amount of a VMAT2 inhibitor described herein, or a pharmaceutically acceptable salt thereof.

In some embodiments, the neurological or psychiatric disease or disorder is a hyperkinetic movement disorder.

In some embodiments, the hyperkinetic movement disorder is tardive dyskinesia.

In some embodiments, the hyperkinetic movement disorder is Tourette's syndrome.

In some embodiments, the hyperkinetic movement disorder is Huntington's disease.

In some embodiments, the hyperkinetic movement disorder is tics.

In some embodiments, the hyperkinetic movement disorder is chorea associated with Huntington's disease.

In some embodiments, the hyperkinetic movement disorder is ataxia, chorea, dystonia, hemifacial spasm, Huntington's disease, myoclonus, restless leg syndrome, or tremors.

5 In some embodiments, the neurological or psychiatric disease or disorder is restricted and repetitive behaviors associated with Autism spectrum disorder (ASD).

In some embodiments, the neurological or psychiatric disease or disorder is obsessions and compulsions in partial and non-responders (or completely refractory) with obsessive-compulsive disorder (OCD). In some embodiments, the neurological or psychiatric disease or disorder is
10 obsessions and compulsions in partial and non-responders (or completely refractory) with obsessive-compulsive disorder (OCD) and the compounds described herein are administered as adjunctive therapy. In some embodiments, the compounds described here are administered as adjunctive therapy with the primary therapy being treatment with antidepressants.

In some embodiments, the neurological or psychiatric disease or disorder is Bipolar I
15 Disorder. In some embodiments, the compound described herein is administered as monotherapy for the treatment of Bipolar I Disorder. In some embodiments, the compound described herein is administered as maintenance therapy for the treatment of Bipolar I Disorder. In some embodiments, the compound described herein is administered as monotherapy maintenance therapy for the treatment of Bipolar I Disorder.

20 In some embodiments, the VMAT2-inhibiting compound of Formula (I), or a pharmaceutically acceptable salt thereof, is administered to the patient to treat or prevent a disease or disorder selected from:

ataxias or spinal muscular atrophies, such as, spinocerebellar ataxia type 17 (SCA17) / HDL4, ataxia, spinal muscular atrophy, amyotrophic lateral sclerosis, familial amyotrophic lateral sclerosis,
25 bulbo-spinal muscular atrophy congenital, dentatorubral-pallidoluysian atrophy, hereditary motor neuron disease, and hereditary spastic paraplegia;

chorea, such as, benign hereditary chorea, chorea, chorea associated with mitochondrial disease/causes, chorea associated with Wilson's disease, chorea gravidarum, chorea-acanthocytosis, drug-induced chorea, hemiballism, rheumatic/Sydenham's chorea, and thyrotoxic chorea/hyperthyroid
30 chorea;

congenital malformations, deformations, or abnormalities, such as, Angelman syndrome, congenital neurological disorder, Aicardi's syndrome, neurofibromatosis, congenital facial nerve hypoplasia, Moebius II syndrome, Cockayne's syndrome, Sjogren-Larsson syndrome, Laurence-Moon-Bardet-Biedl syndrome, Fragile X syndrome, and Prader-Willi syndrome;

35 dementia, such as, AIDS-related dementia, Alzheimer's disease, congenital neurological degeneration, Lewy body dementia, micro-infarct dementia, pre-senile dementia, senile dementia, and vascular dementia;

diseases of oral cavity, salivary glands, and jaws, such as, glossodynia / burning mouth syndrome and temporomandibular joint disorder;

dyskinesia, such as, pharyngeal dyskinesia, dyskinesia, dyskinesia (neonatal), dyskinesia (oesophageal), levodopa-induced dyskinesia, paroxysmal kinesigenic dyskinesias, paroxysmal nonkinesigenic dyskinesias, and respiratory dyskinesia;

dystonia, such as, blepharospasm, buccoglossal syndrome, drug-induced acute dystonia, dystonia, early onset primary dystonia, genetic torsion dystonia, hand dystonia/writer's cramp, idiopathic nonfamilial dystonia, idiopathic orofacial dystonia/Meige's disease, laryngeal dystonia, oromandibular dystonia, and spasmodic torticollis/cervical dystonia;

endocrine, nutritional, and metabolic diseases, such as, Wilson's Disease, diabetes mellitus, obesity, syndrome X, and Lesch-Nyhan syndromes;

epilepsy, such as, Baltic myoclonic epilepsy, benign familial neonatal convulsions, epilepsy, epilepsy congenital, Lafora's myoclonic epilepsy, severe myoclonic epilepsy of infancy, and convulsions;

habit and impulse disorders, such as, binge eating disorder, kleptomania, impulse control disorders, trichotillomania, intermittent explosive disorder, pathological gambling, and pyromania;

Huntington's disease or related disorders, such as, Huntington's disease, Huntington's disease-like syndromes 1-3, Huntington's chorea, and X-linked McLeod Neuroacanthocytosis syndrome;

mood or psychotic disorders, such as, schizophrenia, psychosis, mania, bipolar disorder, depression, and mood disorders;

other diseases or disorders, such as, fumbling, hypokinesia, hypokinesia (neonatal), movement disorder, rabbit syndrome, spasticity, up and down phenomenon, asthma, cancer, congenital nystagmus, familial hemiplegic migraine, fetal movement disorder, and rheumatoid arthritis;

neurotic, stress-related, and somatoform disorders, such as, social anxiety disorder, panic disorder, generalized anxiety disorder, obsessive-compulsive disorder, post-traumatic stress disorder, and psychogenic movement disorder;

other degenerative diseases of basal ganglia, such as, pantothenate kinase-associated neurodegeneration, progressive supranuclear palsy, multiple system atrophy, dyslexia, basal ganglion degeneration, and neuroferritinopathy;

other extrapyramidal and movement disorders, such as, hemiballismus, extrapyramidal disorder, essential tremor, geniospasm, hyperexplexia, akathisia, ballismus / hemiballism, myoclonus, and restless legs syndrome / Willis-Ekbom's syndrome;

other nervous system or motor function, such as, sleep-related bruxism, abnormal involuntary movement disorders, alien limb syndrome, Alzheimer's disease (agitation), clumsiness, clonic hemifacial spasm, olfactory nerve agenesis, congenital cranial nerve paralysis, exercise ataxia

syndrome, familial periodic paralysis, congenital hemiparesis, fine motor delay, fine motor skill dysfunction, gross motor delay, multiple sclerosis, congenital flaccid paralysis, congenital Horner's syndrome, alternating hemiplegia of childhood, motor developmental delay, cerebral palsy, athetoid cerebral palsy, posturing, pseudoparalysis, psychomotor hyperactivity, bradykinesia, synkinesis, akinesia, Riley-Day syndrome, and athetosis;

Parkinson's/parkinsonism, such as, parkinsonism, drug-induced parkinsonism, micrographia, and Parkinson's disease;

pediatric-onset behavioral and emotional disorders, such as, attention deficit hyperactivity disorder, attention deficit disorder, hyperkinesia, hyperkinesia (neonatal), oppositional defiant disorder, provisional tic disorder, persistent (chronic) motor or vocal tic disorder, stereotypic movement disorder, stereotypy, and Tourette's syndrome;

pervasive developmental disorders, such as, autism spectrum disorders, Rett's syndrome, Asperger's syndrome, pervasive developmental disorder NOS, and dyslexia; and

substance abuse or dependence, such as, addiction disorders, alcoholism, cocaine dependence, illegal drug abuse, methamphetamine abuse, methamphetamine addiction/dependence, methamphetamine use disorder, morphine abuse, morphine-analogue abuse, nicotine dependence, polysubstance abuse, and prescription drug abuse.

In some embodiments, the patient which is treated has been determined to have 22q11.2 deletion syndrome. In some embodiments, the patient is predisposed to developing a psychiatric disorder due to the patient having 22q11.2 deletion syndrome. In some embodiments, the patient has been determined to have COMT haploinsufficiency. In some embodiments, the patient is predisposed to developing a psychiatric disorder due to the patient having COMT haploinsufficiency.

In another embodiment, the VMAT2 inhibitors described herein may be hydrolyzed in the body of a mammal to compounds that may inhibit the human vesicular monoamine transporter isoform 2. As such, these VMAT2 inhibitors may have additional utility in altering the *in vivo* properties of the metabolite in a mammal, such as, the maximum concentration or duration of action.

Characterizing any of the VMAT2 inhibitors described herein may be determined using methods described herein, and those in the art. For example, dopamine depletion may be determined using the locomotor activity (LMA) assay. Another *in vivo* animal model includes the conditioned avoidance response (CAR) test, which has been shown to be an effective and reliable preclinical model for assessing the antipsychotic activity of compounds.

COMINATION THERAPY

The compounds of the present disclosure, or their pharmaceutically acceptable salts, can be used as monotherapy, or in combination with one or more other pharmaceutical agents. In some embodiments, a compound of Formula (I), or its pharmaceutically acceptable salt, is administered together with (simultaneously or sequentially) one or more pharmaceutical agents selected from

antidepressants, antipsychotics (typical or atypical), antiepileptics, antimicrobials, antiarrhythmics, mood stabilizers, and gastrointestinal drugs. In some embodiments, a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is used in adjunctive therapy which refers to a treatment that is used in conjunction with a primary treatment and its purpose is to assist the primary treatment.

- 5 Adjunctive therapies are typically co-administered therapies. As an example of adjunctive therapy, if obsessive-compulsive disorder is being treated, the primary therapy may be, *e.g.*, an antidepressant, and the co-administration of a compound described herein would be considered an adjunctive therapy.

COMPOUND SYNTHESIS

- 10 Detailed compound synthesis methods are described herein in the Examples. In general, starting components are commercially available chemicals and may be obtained from commercial sources or may be made according to organic synthesis techniques known to those skilled in this art, starting from commercially available chemicals and/or from compounds described in the chemical literature.

- 15 In general, the compounds used in the reactions described herein may be made according to organic synthesis techniques known to those skilled in this art, starting from commercially available chemicals and/or from compounds described in the chemical literature. Methods known to one of ordinary skill in the art may be identified through various reference books and databases. Suitable reference books and treatise that detail the synthesis of reactants useful in the preparation of
20 compounds of the present disclosure, or provide references to articles that describe the preparation, include for example, "Synthetic Organic Chemistry," John Wiley & Sons, Inc., New York; S. R. Sandler *et al.*, "Organic Functional Group Preparations," 2nd Ed., Academic Press, New York, 1983; H. O. House, "Modern Synthetic Reactions", 2nd Ed., W. A. Benjamin, Inc. Menlo Park, Calif. 1972; T. L. Gilchrist, "Heterocyclic Chemistry", 2nd Ed., John Wiley & Sons, New York, 1992; J. March,
25 "Advanced Organic Chemistry: Reactions, Mechanisms and Structure," 4th Ed., Wiley-Interscience, New York, 1992.

- Specific and analogous reactants may also be identified through the indices of known chemicals prepared by the Chemical Abstract Service of the American Chemical Society, which are available in most public and university libraries, as well as through on-line databases (the American
30 Chemical Society, Washington, D.C., may be contacted for more details). Chemicals that are known but not commercially available in catalogs may be prepared by custom chemical synthesis houses according to known methods, where many of the standard chemical supply houses (*e.g.*, those listed above) provide custom synthesis services.

- The term "reducing agent" refers to a compound that contributes a hydride to an electrophilic
35 position of a reactant compound, such as, an unsaturated carbon (*e.g.* carbon of a carbonyl moiety), such as, converting a ketone containing reactant compound to an alcohol product compound or converting an ester containing reactant compound to an alcohol product compound. The reducing

agent can be a hydride reducing agent. Example hydride reducing agents include, but are not limited to, diborane, borane (*e.g.* borane tetrahydrofuran complex), 9-borabicyclo[3.3.1]nonane, lithium aluminum hydride, diisobutylaluminum hydride, lithium diisobutyl-*tert*-butoxyaluminum hydride, lithium tri-*tert*-butoxyaluminum hydride, lithium tris[(3-ethyl-3-pentyl)oxy]aluminumhydride, sodium bis(2-methoxyethoxy)aluminum dihydride, sodium aluminum hydride, calcium borohydride, lithium borohydride, magnesium borohydride, potassium borohydride, tetrabutylammonium borohydride, tetraethylammonium borohydride, tetramethylammonium borohydride, bis(triphenylphosphine)copper(I) borohydride, lithium 9-borabicyclo[3.3.1]nonane hydride, sodium triacetoxymborohydride, potassium tri-*sec*-butylborohydride, sodium tri-*sec*-butylborohydride, potassium trisiamylborohydride, lithium triethylborohydride, potassium triethylborohydride, sodium triethylborohydride, potassium triphenylborohydride, lithium dimethylaminoborohydride, lithium pyrrolidinoborohydride, sodium cyanoborohydride, sodium trimethoxyborohydride, sodium borohydride, and the like.

The term “halogenating agent” refers to a compound that contributes a halogen atom to a reactant compound, such as, converting an alcohol reactant compound to an alkyl halide product compound. Examples of halogenating agents include, but not limited to, thionyl chloride, oxalyl chloride, phosphorus oxychloride, phosphorus pentachloride, phosphorus trichloride, methanesulfonyl chloride and NaI, *p*-toluenesulfonyl chloride and NaI, phosphorus tribromide, triphenylphosphine dibromide, phosphorus pentabromide, or thionyl bromide, and the like.

The term “base” refers to a compound that is an electron pair donor in an acid-base reaction. The base can be an inorganic base or an organic base.

The term “organic base” refers to a base including at least one C-H bond (*e.g.* an amine base). In some embodiments, the amine base can be a primary, secondary, or tertiary amine.

The term “inorganic base” refers to a base that does not include at least one C-H bond and includes at least one alkali metal or alkaline earth metal.

The term “acid” refers to a compound that is an electron pair acceptor in an acid-base reaction.

The acid can be an inorganic acid or organic acid.

The term “inorganic acid” refers to an acid that does not include a carbon bond. Inorganic acids can be a strong acid or a weak acid.

The term “organic acid” refers to an acid including at least one C-H bond, C-F bond, or C-C bond.

CERTAIN ABBREVIATIONS

The specification includes numerous abbreviations, whose definitions are listed in the following table:

Abbreviation	Definition
ACN	Acetonitrile
aq.	Aqueous
EtOAc	Ethyl acetate
DCM	Dichloromethane or methylene chloride
d	Day
de	Diastereomeric excess
DPTTA	(2 <i>S</i> ,3 <i>S</i>)-2,3-bis(4-methylbenzoyloxy)butanedioic acid
ee	Enantiomeric excess
EtOH	Ethyl alcohol
h	Hour
HPLC	High-performance liquid chromatography
<i>i</i> -PrOH	Isopropyl alcohol
min.	Minute(s)
MeOH	Methyl alcohol
MTBE	Methyl <i>tert</i> -butyl ether
RT	Room temperature
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran

EXAMPLES

The following examples are included to demonstrate embodiments of the disclosure. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the disclosure.

Analytical HPLC analyses were performed on an LC-MS system with a UV Detector (DionexTM UVD 170u UV/VIS Detector), Corona array detector (ThermoTM VeoTM RS), and mass spectrometer (Dionex MSQ PlusTM). Reverse-phase preparative HPLC purifications were performed on a liquid chromatography-mass spectrometry (LCMS) system C18 Kinetix 5 μ 100 Å 150 x 21.2 mm column by Phenomenex using ACN/water gradient containing 0.05% TFA run at 1.1 mL/min and 50 °C. Supercritical fluid chromatography purification (SFC) was performed using a WatersTM Prep 100qTM system, equipped with a UV Detector (WatersTM 2998 Photodiode Array DetectorTM) and mass spectrometer (WatersTM Acquity QDa DetectorTM). A WatersTM ViridisTM BEH 2-Ethylpyridine 130 Å 5 μ m, 30 mm x 100 mm column was used with a CO₂ and 0.3% NH₄OH in MeOH gradient, run at 100 mL/min, 40 °C, and 105 bar back pressure regulator. All final compounds were analyzed by analytical HPLC-MS, and peaks were monitored at 220, 212, 270, and 254 nm in the UV, by mass spectrometer, and through the charged aerosol detector for purity. ¹H was recorded in an appropriate NMR solvent, such as, dimethylsulfoxide-*d*₆ (DMSO-*d*₆), on a Bruker 400 MHz spectrometer equipped with a Broad Band NMR probe, or on a Bruker 500 MHz spectrometer equipped with a 5

mm QNP probe with Z gradient. The ^1H chemical signals are given in parts per million (ppm) with the residual solvent signal used as reference. The chemical shifts are expressed in ppm (δ) and coupling constants (J) are reported in hertz (Hz). Reactions were performed under an atmosphere of dry nitrogen unless otherwise stated.

5

I. The general synthetic scheme for the preparation of compounds of Formula (I) is shown in FIG. 6, wherein X is halo (*e.g.* I, Cl, or Br), $\text{OS}(\text{O})_2\text{OR}^1$, or $\text{OS}(\text{O})_2\text{R}^2$ wherein R^2 is $\text{C}_1\text{-C}_6$ alkyl, halo $\text{C}_1\text{-C}_6$ alkyl, or phenyl, said phenyl unsubstituted or substituted with one or more halo, nitro, $\text{C}_1\text{-C}_6$ alkyl, halo $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, or halo $\text{C}_1\text{-C}_6$ alkoxy. Alternatively, $\text{R}^1\text{-X}$, where X is I, Cl, or Br can be prepared by reacting commercially available HO-R^1 with a halogenating agent.

10

According to FIG. 6, compounds of general Formula (I), as described herein, can be synthesized in several steps from **Compound 1-A**. In one instance, a sequence for the formation of compounds of general Formula (I) includes condensation of **Compound 1-A** with 3-[(dimethylamino)methyl]-5,5-dimethylhexan-2-one, optionally in the presence of an acid and an additive, such as, NaI , in a solvent or a solvent mixture, such as, *i*-PrOH/water and the like to afford **Compound 1-B**. Reduction of **Compound 1-B** in the presence of a reducing agent, such as, sodium aluminum hydride, calcium borohydride, lithium borohydride, magnesium borohydride, potassium borohydride, tetrabutylammonium borohydride, tetraethylammonium borohydride, tetramethylammonium borohydride, bis(triphenylphosphine)copper(I) borohydride, lithium 9-borabicyclo[3.3.1]nonane hydride, sodium triacetoxyborohydride, lithium tri-*sec*-butylborohydride, potassium tri-*sec*-butylborohydride, sodium tri-*sec*-butylborohydride, potassium trisiamylborohydride, lithium triethylborohydride, potassium triethylborohydride, sodium triethylborohydride, potassium triphenylborohydride, lithium dimethylaminoborohydride, lithium pyrrolidinoborohydride, sodium cyanoborohydride, sodium trimethoxyborohydride, sodium borohydride, and the like, in a solvent or a solvent mixture, such as, methanol, MTBE, and the like, optionally in the presence of an additive, such as, acetic acid, affords **Compound 1-C**. Preparation of **Compound 1-D** can be accomplished by hydrogenolysis or solvolysis, for example, treating **Compound 1-C** with an acid in a solvent, such as, methanol, and the like. Finally, **Compound 1-D** can be treated with $\text{R}^1\text{-X}$, wherein X is a leaving group, such as, I, Br, Cl, triflate, and the like; or an epoxide, in the presence of a base in a solvent, such as, DMF, acetone, and the like, to afford compounds of general Formula (I).


15

20

25

30

EXAMPLE 1A: Preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (Compound 1-D (2*R*,3*R*,11*bR*)), see FIG. 7.



Step 1: Synthesis of 9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-one (Compound 1-B).

A suspension of 6-(benzyloxy)-7-methoxy-3,4-dihydroisoquinoline (**Compound 1-A**, 25 g, 93.5 mmol, 1.2 eq) in water (63 mL) was prepared and cooled to 0 °C. Then, aq. HCl (6.7 mL, 12.1N, 1.05 eq) followed by sodium iodide (5.83 g, 38.9 mmol, 0.5 eq) were added and the resulting mixture was warmed to RT. Lastly, a solution of 3-[(dimethylamino)methyl]-5,5-dimethylhexan-2-one (14.4 g, 77.9 mmol, 1.0 eq) in *i*-PrOH (22 mL) was added and the reaction mixture was stirred vigorously at 45 °C for 3d. The resulting suspension was cooled, the precipitate collected, and the filtrate extracted into DCM. The collected precipitate was dissolved in DCM and added to the combined organic layers, dried over MgSO₄, filtered to remove solid and concentrated *in vacuo*. Silica gel columns (220g x 2, 120g x 1) were loaded using DCM and run with an increasing gradient of EtOAc (0-40% over 20 minutes) in hexanes to afford 9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-one (**Compound 1-B**, 22.6 g, 55.5 mmol, 71%).

Step 2: Synthesis of 9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-ol (Compound 1-C).

To a solution of 9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-one (**Compound 1-B**, 22.6 g, 55.5 mmol, 1.0 eq) in MTBE:MeOH (4:1, 113 mL) and acetic acid (3.2 mL, 55.5 mmol, 1.0 eq) was carefully added solid sodium borohydride (2.3 g, 61.0 mmol, 1.1 eq) portionwise. Reaction progression was monitored hourly by LC/MS. Subsequently, 4:1 MTBE:MeOH (56 mL), 0.5 eq of acetic acid and an additional 2.0 eq of sodium borohydride were added to the stirring mixture. After stirring overnight at RT, 1N NaOH (180 mL) was added, and the resulting mixture heated to 50 °C for 3h. The mixture was cooled to RT and allowed to stand for 1h. The resulting precipitate was collected by vacuum filtration, rinsed with excess water (~ 600 mL) until pH < 10 and lastly washed with MTBE. The filter cake was dried to provide 9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-ol (**Compound 1-C**, 18.7 g, 42.1 mmol, 76%).

Step 3: Synthesis of (2R,3R,11bR)-9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-ol (Compound 1-C (2R,3R,11bR) also referred to as Compound 4-60 in Table A).


To a suspension of 9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 1-C**, 18.7 g, 42.1 mmol, 1.0 eq) in MeOH (150 mL) was added (2*S*,3*S*)-2,3-*bis*(4-methylbenzoyloxy)butanedioic acid (DPTTA, 17.9 g, 46.3 mmol, 1.1 eq) portion wise. Additional MeOH (100 mL) and heat (65 °C) were required to dissolve solids. The resulting solution was cooled to 60 °C, seeded with previously prepared **Compound 1-C (2*R*,3*R*,11*bR*)-(DPTTA)** (138 mg, for example see **EXAMPLE 1B**) and stirred at 60 °C for 30 min. Then, the mixture was cooled to RT at a rate of 4 °C/h then stirred overnight at RT. The resulting precipitate was collected by vacuum filtration and dried to provide **Compound 1-C (2*R*,3*R*,11*bR*) (DPTTA)** (14.7 g, 18.5 mmol, 44%) as a white solid. An optical purity of 100% ee as a single diastereomer was determined by chiral supercritical fluid chromatography (SFC) analysis. Chiral purity analysis was performed using a WatersTM Ultra-Performance Convergence Chromatography (UPC2)TM supercritical fluid chromatography (SFC) system, equipped with a UV Detector (WatersTM Acquity UPC2 PDA DetectorTM) and mass spectrometer (WatersTM Acquity QDa DetectorTM). A Chiral Technologies IncTM ChiralPakTM IC/SFC 1.6 µm, 2.1 mm x 50 mm column was used with an isocratic gradient at 85% CO₂ and 15% 0.5% DMEA in MeOH, at 1.5 mL/min, 55 °C, and 1500 psi back pressure regulator. An extracted wavelength of 285 nm was used for %cc quantification and analysis.

The chiral salt pair was suspended in DCM and water was added. The mixture was basified with saturated aq. NH₄OH until pH ~ 10. The mixture was extracted three times with DCM. The organic extracts were dried over MgSO₄, filtered, and concentrated *in vacuo* to give (2*R*,3*R*,11*bR*)-9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 1-C (2*R*,3*R*,11*bR*)**, 7.41 g, 18.1 mmol, 43%).

Step 4: Synthesis of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (Compound 1-D (2*R*,3*R*,11*bR*)**).**

To a solution of (2*R*,3*R*,11*bR*)-9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 1-C (2*R*,3*R*,11*bR*)**, 6.9 g, 16.8 mmol, 1.0 eq) in MeOH (68 mL) cooled to 0 °C was added 37% aq. HCl (34 mL) dropwise over 30 min. The resulting mixture was stirred at 0 °C for 30 min, warmed to RT and stirred for 30 min and finally stirred at 80 °C overnight. The mixture was transferred to a 2 L flask, diluted with water (60 mL) and DCM (100 mL) and cooled to 0 °C. Then, saturated aq. NaHCO₃ was added dropwise until reaching a pH of ~ 8, adding additional DCM as needed to aid stirring. The resulting precipitate was collected by vacuum filtration, rinsed with water and MTBE, dried and set aside. The filtrate was extracted with 5:1 DCM:*i*-PrOH. The combined organic layers were dried over MgSO₄, filtered to remove solid and concentrated *in vacuo*. The resulting solid was combined with the filter cake to provide (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*R*,11*bR*)**, 5.5 g, 17.2 mmol) in near quantitative yield.

EXAMPLE 1B: Preparation of the salt of (2*R*,3*R*,11*bR*)-9-(benzyloxy)-10-methoxy-3-neopentyl-1,3,4,6,7,11*b*-hexahydro-2*H*-pyrido[2,1-*a*]isoquinolin-2-ol and (2*S*,3*S*)-2,3-bis(4-methylbenzoyloxy)butanedioic acid (Compound 1-*C*·(2*R*,3*R*,11*bR*) (DPTTA)), see FIG. 8.



To a suspension of 9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 1-*C***, 506 mg, 1.24 mmol, 1.0 eq) in MeOH (7 mL) was added (2*S*,3*S*)-2,3-bis(4-methylbenzoyloxy)butanedioic acid (DPTTA, 494 mg, 1.24 mmol, 1.0 eq) portion wise. The resulting mixture was heated to reflux with stirring to achieve a clear solution. The mixture was cooled to RT without stirring and let stand 16 h at RT. The resulting precipitate was collected by vacuum filtration, rinsed with methanol (1 mL) and dried to provide **Compound 1-*C*·(2*R*,3*R*,11*bR*) (DPTTA)** (365 mg, 37%) as a white solid. An optical purity of 90% ee was determined by chiral SFC analysis. Chiral purity analysis was performed using a WatersTM Ultra-Performance Convergence Chromatography (UPC2)TM supercritical fluid chromatography (SFC) system, equipped with a UV Detector (WatersTM Acquity UPC2 PDA DetectorTM) and mass spectrometer (WatersTM Acquity QDa DetectorTM). A Chiral Technologies IncTM ChiralPakTM IC/SFC 1.6 μ m, 2.1 mm x 50 mm column was used with an isocratic gradient at 85% CO₂ and 15% 0.5% DMEA in MeOH, at 1.5 mL/min, 55 °C, and 1500 psi back pressure regulator. An extracted wavelength of 285 nm was used for %ee quantification and analysis. A 20 mg sample of the solid was set aside for further recrystallizations and the mother liquors and remaining white solids were recombined and concentrated *in vacuo*. This mixture was suspended in MeOH (5 mL) and heated to reflux to achieve a clear solution. Stirring was stopped and the mixture cooled to 60 °C and seeded with 2 mg of the prior 90% ee **Compound 1-*C*·(2*R*,3*R*,11*bR*) (DPTTA)** solid. The mixture was allowed to stand for 1 h at 60 °C then reduced the temperature by 5 °C each hour until at RT, then let stand for 16 h. The resulting precipitate was collected by vacuum filtration, rinsed with methanol (1 mL), and dried to provide **Compound 1-*C*·(2*R*,3*R*,11*bR*) (DPTTA)** (200 mg, 20%). An optical purity of 100% ee as a single diastereomer was determined by chiral SFC analysis.

EXAMPLE 2: Preparation of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 2-*B* (3*R*,11*bR*)) and (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 2-*B* (3*S*,11*bS*)), see FIG. 9.

Step 1: Synthesis of 3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 2-B).

To a solution of 7-methoxy-3,4-dihydroisoquinolin-6-ol (**Compound 2-A**, 7.2 g, 40.7 mmol, 1.2 eq) in EtOH (66 mL) was added [2-(2,2-dimethylpropyl)-3-oxobutyl]trimethylazanium iodide (11.1 g, 33.9 mmol, 1.0 eq) and the resulting mixture was stirred at reflux for 3d. The mixture was cooled, diluted with aq. NaHCO₃, and extracted with DCM. The combined organic layers were dried over MgSO₄, filtered, and concentrated *in vacuo*. The crude material was dry loaded on celite and chromatographed on a silica gel column (220 g) with an increasing gradient of EtOAc (0-60% over 20 minutes) in hexanes. Due to a co-eluting impurity, the material was dry loaded on celite and re-chromatographed on a silica gel column (80 g) using a shallower gradient of EtOAc (0-45% over 20 minutes) in hexanes to provide 3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B**, 4.2 g, 13.3 mmol, 39%) as a 7:1 mixture of racemic diastereomers.

Step 2: Isolation of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 2-B (3*R*,11*bR*)) and (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 2-B (3*S*,11*bS*)).

Separation of the 3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B**) enantiomers was achieved by preparing a solution in a minimal volume of MeOH and subjecting to preparative chiral SFC. Chiral compound separation was performed using a Pic SolutionTM SFC-PICLAB-Prep 200TM supercritical fluid chromatography (SFC) system, equipped with a UV Detector and an AdvionTM mass spectrometer. A Chiral Technologies IncTM ChiralPakTM IG/SFC 5 μ m, 20 mm x 250 mm column was used with an isocratic gradient at 80% CO₂ and 20% 0.5% DMEA in MeOH, at 150 mL/min, 55 °C, and 100 bar back pressure regulator. Following purification, 1.6 g each of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*R*,11*bR*))** (99% ee) and (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*S*,11*bS*))** (100% ee) were isolated.

Suitable crystals were generated by slow evaporation of a solution of **Compound 2-B (3*S*,11*bS*)** in acetone. Absolute stereochemistry was confirmed by obtaining a small molecule crystal structure of **Compound 2-B (3*S*,11*bS*)**.


Single Crystal X-Ray Structure of Compound 2-B (3*S*,11*bS*).

The single crystal structure of (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*S*,11*bS*))** was determined. The absolute configuration parameter (Flack parameter) is near zero when the configuration assignment is correct, and near +1 when the configuration is wrong. For this data set,

the Flack parameter is 0.06(3) for the assignment shown in **FIG. 1A**. Crystallographic information is shown in **Table 1**.

Table 1: Crystal Data for Compound 2-B (3<i>S</i>,11<i>bS</i>)		
Molecular Formula	C ₁₉ H ₂₇ NO ₃	
Formula weight (g/mol)	317.41	
Wavelength	1.54178 Å	
Radiation (Å)	Cu Kα (λ = 1.54178 Å)	
Instrument	Bruker Microstar APEX II CCD	
Crystal system	Monoclinic	
space group	P 1 2 ₁ 1	
Temperature (K)	100.0	
Unit cell dimensions		
	a = 6.8023(2) Å	α = 90°
	b = 8.9511(3) Å	β = 94.837(3)°
	c = 13.9008(5) Å	γ = 90°
Volume	843.38(5) Å ³	
Z, Z'	2	
Density (calculated)	1.250 Mg/m ³	
F(000)	344	
Crystal size	0.18 x 0.12 x 0.02 mm ³	

- 5 **EXAMPLE 3: Alternative Preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (Compound 1-D (2*R*,3*R*,11*bR*)) and preparation of (2*S*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (Compound 1-D (2*S*,3*S*,11*bS*)), see FIG. 10.**



Synthesis of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (Compound 1-D (2*R*,3*R*,11*bR*)).

To a solution of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*R*,11*bR*)**), 733 mg, 2.31 mmol, 1.0 eq) in MeOH (5 mL) was added sodium borohydride (131 mg, 3.46 mmol, 1.5 eq). The resulting mixture was stirred 16 h at RT at which point additional sodium borohydride was added (44 mg, 1.16 mmol, 0.5 eq). The mixture was stirred an additional 3 h at RT. The excess reagent was

quenched with water, and the resulting mixture extracted three times with 5:1 DCM:*i*-PrOH. The combined organic layers were dried over MgSO₄, filtered to remove solid and concentrated *in vacuo*. The resultant solid was suspended in MTBE (5.9 mL) and MeOH (1.5 mL) and the resulting mixture was heated to 50 °C for 1 h with stirring. The suspension was cooled to RT and the solids collected by filtration. The filter cake was rinsed with MTBE (2.2 mL) and the solids were collected to give (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*R*,11*bR*)**), 567 mg, 1.77 mmol, 35:1 dr, 77%).

Single Crystal X-Ray Structure of Compound 1-D (2*R*,3*R*,11*bR*).

Crystals of **Compound 1-D (2*R*,3*R*,11*bR*)** were grown using a CrystalBreeder system. Approximately 5 mg of **Compound 1-D (2*R*,3*R*,11*bR*)** in 0.25 mL of DMF (~20 mg/mL) was slow thermo-cycled several times from 5 °C to 80 °C. Multiple vials were utilized. After the thermo-cycling, large needles formed on the sides of the vials. The solvent was decanted, and the crystals were analyzed. A suitable crystal was selected for single crystal X-ray structure determination. A similar preparation using ACN also provided crystals, but the crystals grown from DMF provided better crystals that were more suitable for further analysis.

The crystal structure of **Compound 1-D (2*R*,3*R*,11*bR*)** was solved. The structure was determined to be anhydrous and unsolvated; the three chiral centers at C2, C3, and C11b were all determined to have the *R* configuration. See **FIG. 1B** for the asymmetric unit for the crystal structure. The crystal data, unit system and space group for **Compound 1-D (2*R*,3*R*,11*bR*)** are shown in **Table 2**. Data collection and refinement parameters for **Compound 1-D (2*R*,3*R*,11*bR*)** are shown in **Table 3**.


Table 2: Crystal Data for Compound 1-D (2<i>R</i>,3<i>R</i>,11<i>bR</i>)	
Chemical formula	C ₁₉ H ₂₉ NO ₃
<i>Mr</i>	319.43
Crystal system, Space group	Orthorhombic, P2 ₁ 2 ₁ 2 ₁
Data collection temperature (K)	150
<i>a</i> (Å)	9.1345 (3)
<i>b</i> (Å)	12.3990 (4)
<i>c</i> (Å)	14.9842 (6)
volume (Å ³)	1697.09 (10)
<i>Z</i>	4
Radiation type	Cu Kα
Crystal shape	Plate
Color	Colorless
Crystal size (mm)	0.13 × 0.11 × 0.04

Table 3: Data collection and Refinement Parameters for Compound 1-D (2R,3R,11bR)	
Diffractionmeter	Bruker AXS D8 Quest diffractometer with PhotonIII_C14 charge-integrating and photon counting pixel array detector
Radiation source	I-mu-S microsource X-ray tube
Absolute structure parameter	0.03 (10)

Synthesis of (2S,3S,11bS)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol (Compound 1-D (2S,3S,11bS)).

(2S,3S,11bS)-3-(2,2-Dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol (**Compound 1-D (2S,3S,11bS)**) was made in an analogous manner to prepare **Compound 1-D (2R,3R,11bR)** using (3S,11bS)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-one (**Compound 2-B (3S,11bS)**) as the starting material and sodium borohydride as the reducing agent.

EXAMPLE 4: Preparation of (2S,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol (Compound 1-D (2S,3R,11bR)) and (2R,3S,11bS)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol (Compound 1-D (2R,3S,11bS)), see FIG. 11.




Synthesis of (2S,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol (Compound 1-D (2S,3R,11bR)).

A solution of (3R,11bR)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-one (**Compound 2-B (3R,11bR)**), 48 mg, 0.15 mmol, 1.0 eq) in THF (0.75 mL) was cooled to 0 °C. Then, a solution of lithium tri-*sec*-butylborohydride (1M THF, 0.30 mL, 0.30 mmol, 2.0 eq) was added dropwise. The resulting mixture was stirred at 0 °C for 30 min, warmed to RT and stirred for 3h. The mixture was concentrated *in vacuo*, diluted with water, and extracted with DCM. The combined organic layers were dried over MgSO₄, filtered to remove solids, and concentrated *in vacuo*. Silica gel column (4 g) was loaded using DCM and run with an increasing gradient of EtOAc (0-60% over 30 minutes) in hexanes. Fractions were split between pure product (**Compound 1-D (2S,3R,11bR)**, 16 mg) and product with the minor diastereomer (28 mg, 10:1 diastereomeric ratio between **Compound 1-D (2S,3R,11bR)** : **Compound 1-D (2R,3S,11bR)**). Taken together, (2S,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol was isolated in a combined yield (**Compound 1-D (2S,3R,11bR)**, 44 mg) of 91%.

Synthesis of (2*R*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (Compound 1-D (2*R*,3*S*,11*bS*)).


(2*R*,3*S*,11*bS*)-3-(2,2-Dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (Compound 1-D (2*R*,3*S*,11*bS*)) was made in an analogous manner to prepare
 5 **Compound 1-D (2*S*,3*R*,11*bR*)** using (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 2-B (3*S*,11*bS*)) as the starting material and lithium tri-*sec*-butylborohydride as the reducing agent.

EXAMPLE 5A: Preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 4-1), see FIG. 12.




To a solution of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (Compound 1-D (2*R*,3*R*,11*bR*), 0.20
 15 g, 0.63 mmol, 1.0 eq) in acetone (6.3 mL) was added cesium carbonate (0.61 g, 1.9 mmol, 3.0 eq) and the resulting mixture stirred at RT for 10 min. Then, 2,2,2-trifluoroethyl trifluoromethanesulfonate (0.19 g, 0.82 mmol, 1.3 eq) was added and the mixture stirred at RT overnight. The mixture was diluted with water and extracted with 5:1 DCM:*i*-PrOH. The combined organic layers were dried over MgSO₄, filtered to remove solids, and concentrated *in vacuo*. Silica gel column (40 g) was loaded
 20 using DCM and run with an increasing gradient of EtOAc (0-50% over 25 min) in hexanes to afford (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 4-1, 0.19 g, 0.47 mmol, 75%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ (ppm) 6.81 (s, 1H), 6.79 (s, 1H), 4.63 (q, *J* = 8.9 Hz, 2H), 4.56 (d, *J* = 6.4 Hz, 1H), 3.76 (s, 3H), 3.13 – 3.00 (m, 2H), 2.99 – 2.83 (m, 3H), 2.57 – 2.44
 25 (m, 2H), 2.32 – 2.26 (m, 1H), 1.94 (t, *J* = 11.2 Hz, 1H), 1.70 (d, *J* = 13.6 Hz, 1H), 1.48 – 1.40 (m, 1H), 1.27 (q, *J* = 11.5 Hz, 1H), 0.91 (s, 9H), 0.81 (dd, *J* = 14.0, 7.6 Hz, 1H).

EXAMPLE 5B: Preparation of (2*R*,3*R*,11*bR*)-9-cyclopropoxy-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 4-27).




To a solution of (2R,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol (**Compound 1-D (2R,3R,11bR)**), 0.80 g, 2.50 mmol, 1.0 eq) in DMF (40 mL) was added cesium carbonate (2.44 g, 7.50 mmol, 3.0 eq) at 50 °C. A solution of cyclopropyl trifluoromethanesulfonate (0.60 g, 3.13 mmol, 1.25 eq) in DMF (3 mL) was added via syringe pump over 6 h. The resulting mixture was stirred at 50 °C an additional 10 h. The mixture was diluted with water and extracted three times with EtOAc. The combined organic extracts were washed five times with water, dried over MgSO₄, filtered, and concentrated *in vacuo*. A silica gel column (24 g) was loaded using DCM and run with an increasing gradient of EtOAc (0-100% over 25 min) in hexanes to afford impure (2R,3R,11bR)-9-cyclopropoxy-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-ol (**Compound 4-27**, 0.83 g, 2.31 mmol) as an off white solid. To remove trace impurities, crude **Compound 4-27** (0.83 g, 2.31 mmol, 1.0 eq) was dissolved in anhydrous THF (14 mL) under an atmosphere of nitrogen at 0 °C with stirring. A 2M solution of lithium aluminum hydride in THF (2.3 mL, 4.62 mmol, 2.0 eq) was added dropwise over 10 min. The temperature was increased to 60 °C and stirred 16 h. The mixture was cooled to 0 °C then sodium sulfate dodecahydrate (0.65 g, 2.02 mmol, 0.9 eq) was added over the course of 10 min then allowed to stir 20 min. The mixture was filtered over celite, and the filter cake was rinsed with acetone, MeOH, and DCM. The filtrate was concentrated *in vacuo* then redissolved in 1:1 DCM/EtOAc and washed three times with 1M aq. NaOH. The organic layer was dried with MgSO₄, filtered, and concentrated *in vacuo*. The crude material was submitted to three chromatographic separations: a silica gel column (12 g) loaded using DCM run with an increasing gradient of EtOAc (0-75% over 25 min) in hexanes, a basic alumina column (24 g) run with an increasing gradient of EtOAc (0-75% over 25 min) in hexanes and a reversed-phase C18 column (30 g) loaded using DMSO run with an increasing gradient of MeCN (10-70% over 20 min) in water to afford (2R,3R,11bR)-9-cyclopropoxy-3-(2,2-dimethylpropyl)-10-methoxy-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-ol (**Compound 4-27**, 0.40 g, 1.11 mmol, 44% over two steps) as a white solid. ¹H NMR (500 MHz, DMSO-*d*₆): δ (ppm) 6.89 (s, 1H), 6.71 (s, 1H), 4.55 (d, *J* = 6.3 Hz, 1H), 3.75 (tt, *J* = 6.0, 3.0 Hz, 1H), 3.69 (s, 3H), 3.11 – 3.04 (m, 1H), 3.00 (br d, *J* = 11.2 Hz, 1H), 2.94 (dd, *J* = 11.8, 4.4 Hz, 1H), 2.92 – 2.86 (m, 2H), 2.55 (dd, *J* = 14.3, 3.8 Hz, 1H), 2.48 – 2.42 (m, 1H), 2.33 – 2.25 (m, 1H), 1.92 (t, *J* = 11.5 Hz, 1H), 1.69 (d, *J* = 12.9 Hz, 1H), 1.48 – 1.39 (m, 1H), 1.25 (q, *J* = 11.5 Hz, 1H), 0.90 (s, 9H), 0.80 (dd, *J* = 13.8, 7.5 Hz, 1H), 0.75 – 0.70 (m, 2H), 0.64 – 0.60 (m, 2H).

EXAMPLE 5C: Preparation of (2R,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-yloxy)-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-ol (Compound 4-30).



To a solution of (2R,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-
 5 1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol (**Compound 1-D (2R,3R,11bR)**), 0.20 g, 0.63 mmol, 1.0 eq) in DMF (6 mL) was added cesium carbonate (0.61 g, 1.88 mmol, 3.0 eq) and the resulting mixture stirred at RT for 10 min. Then, 3-bromooxetane (0.12 mL, 1.56 mmol, 2.5 eq) was added and the mixture stirred at 60 °C for 6 h. The mixture was cooled to RT, diluted with water, and extracted three times with EtOAc. The combined organic extracts were washed five times with
 10 water, dried over MgSO₄, filtered, and concentrated *in vacuo*. The crude material was submitted to two chromatographic separations: a silica gel column (4 g) loaded using DCM run with an increasing gradient of EtOAc (0-100% over 30 min) in hexanes and a reversed-phase C18 column (16 g) loaded using DMSO run with an increasing gradient of MeCN (10-100% over 25 min) in water to afford (2R,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-yloxy)-1H,2H,3H,4H,6H,7H,11bH-
 15 pyrido[2,1-a]isoquinolin-2-ol (**Compound 4-30**), 0.15 g, 0.41 mmol, 65%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ (ppm) 6.77 (s, 1H), 6.31 (s, 1H), 5.15 (quin, *J* = 5.5 Hz, 1H), 4.88 (t, *J* = 6.7 Hz, 2H), 4.57 – 4.51 (m, 3H), 3.75 (s, 3H), 3.12 – 3.04 (m, 1H), 3.04 – 2.91 (m, 2H), 2.90 – 2.79 (m, 2H), 2.46 – 2.42 (m, 1H), 2.31 – 2.23 (m, 1H), 1.92 (br t, *J* = 11.3 Hz, 1H), 1.70 (d, *J* = 13.6 Hz, 1H), 1.47 – 1.38 (m, 1H), 1.25 (q, *J* = 1.0 Hz, 1H), 0.90 (s, 9H), 0.81 (dd, *J* = 13.8, 7.5 Hz, 1H).


EXAMPLE 5D: Preparation of (2R,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2S)-3,3,3-trifluoro-2-hydroxypropoxy]-1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinolin-2-ol (Compound 4-50).



To a solution of (2R,3R,11bR)-3-(2,2-dimethylpropyl)-10-methoxy-
 25 1H,2H,3H,4H,6H,7H,11bH-pyrido[2,1-a]isoquinoline-2,9-diol (**Compound 1-D (2R,3R,11bR)**), 0.20 g, 0.63 mmol, 1.0 eq) in DMF (6 mL) was added cesium carbonate (0.61 g, 1.88 mmol, 3.0 eq) and the resulting mixture stirred at RT for 10 min. Then, (2S)-2-(trifluoromethyl)oxirane (72 μL, 0.94 mmol, 1.5 eq) was added and the mixture stirred at RT for 6 h. An additional portion of (2S)-2-

(trifluoromethyl)oxirane (24 μ L, 0.31 mmol, 0.5 eq) was added and the mixture stirred at RT for an additional hour. The mixture was diluted with water and extracted three times with EtOAc. The combined organic extracts were washed five times with water, dried over MgSO_4 , filtered, and concentrated *in vacuo*. The crude material was submitted to two chromatographic separations: a silica gel column (12 g) loaded using DCM run with an increasing gradient of EtOAc (0-65% over 20 min) in hexanes and a reversed-phase C18 column (16 g) loaded using DMSO run with an increasing gradient of MeCN (10-40% over 25 min, then 40%-90% over 15 min) in water to afford (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*S*)-3,3,3-trifluoro-2-hydroxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-50**, 0.13 g, 0.31 mmol, 50%) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ (ppm) 6.76 (s, 1H), 6.72 (s, 1H), 6.60 (d, J = 6.2 Hz, 1H), 4.55 (d, J = 6.5 Hz, 1H), 4.41 – 4.30 (m, 1H), 4.12 (dd, J = 10.5, 4.0 Hz, 1H), 3.98 (dd, J = 10.3, 6.7 Hz, 1H), 3.74 (s, 3H), 3.12 – 2.83 (m, 5H), 2.59 – 2.54 (m, 1H), 2.46 – 2.43 (m, 1H), 2.32 – 2.23 (m, 1H), 1.93 (br t, J = 11.2 Hz, 1H), 1.70 (bd, J = 14.1 Hz, 1H), 1.48 – 1.38 (m, 1H), 1.26 (q, J = 11.5 Hz, 1H), 0.91 (s, 9H), 0.81 (br dd, J = 13.8, 7.5 Hz, 1H).


EXAMPLE 5E: Preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3*R*)-oxolan-3-yloxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 4-53**).**



To a solution of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D** (2*R*,3*R*,11*bR*), 0.30 g, 0.94 mmol, 1.0 eq) in DMF (3 mL) was added cesium carbonate (0.92 g, 2.82 mmol, 3.0 eq) and the resulting mixture stirred at RT for 10 min. Then, (3*S*)-3-iodooxolane (0.15 mL, 1.41 mmol, 1.5 eq) was added and the mixture stirred at RT overnight. The mixture was diluted with water and extracted three times with EtOAc. The combined organic extracts were washed five times with water, dried over MgSO_4 , filtered, and concentrated *in vacuo*. The crude material was submitted to four chromatographic separations: a dry loaded (celite) basic alumina column (24 g) run with an increasing gradient of MeOH (0-5% over 10 min, then 100% for 10 min) in DCM, a dry loaded (celite) silica column (4 g) run with an increasing gradient of EtOAc (0-100% over 20 min) in hexanes, a reversed-phase C18 column (30 g) loaded with DMSO run with an increasing gradient of MeCN (10-50% over 20 min) in water and a dry loaded (celite) silica column (4 g) run with an increasing gradient of EtOAc (0-100% over 20 min) in hexanes to afford (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3*R*)-oxolan-3-yloxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-53**, 0.14 g, 0.35 mmol, 38%) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ (ppm) 6.74 (s, 1H), 6.62

(s, 1H), 4.95 – 4.89 (m, 1H), 4.55 (d, $J = 6.5$ Hz, 1H), 3.86 – 3.73 (m, 4H), 3.72 (s, 3H), 3.13 – 3.04 (m, 1H), 3.01 (br d, $J = 11.5$ Hz, 1H), 2.95 (dd, $J = 11.7, 4.2$ Hz, 1H), 2.92 – 2.82 (m, 2H), 2.58 – 2.53 (m, 1H), 2.49 – 2.44 (m, 1H), 2.32 – 2.24 (m, 1H), 2.19 – 2.08 (m, 1H), 1.99 – 1.88 (m, 2H), 1.70 (d, $J = 13.1$ Hz, 1H), 1.48 – 1.39 (m, 1H), 1.26 (q, $J = 11.4$ Hz, 1H), 0.91 (s, 9H), 0.81 (dd, $J = 13.9, 7.4$ Hz, 1H).

EXAMPLE 5F: Preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 4-55), see FIG. 12.





































To a solution of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinoline-2,9-diol (**Compound 1-D (2*R*,3*R*,11*bR*)**, 1.00 g, 3.13 mmol, 1.0 eq) in DMF (28 mL) was added cesium carbonate (3.06 g, 9.39 mmol, 3.0 eq) and the resulting mixture stirred at RT for 10 min. Then, (*R*)-2-methyloxirane (0.31 mL, 6.26 mmol, 2 eq) was added and the mixture stirred at 80 °C for 16 h. An additional portion of (*R*)-2-methyloxirane (0.16 mL, 3.13 mmol, 1 eq) was added and the mixture stirred at 80 °C for an additional 4 h. A third addition of (*R*)-2-methyloxirane (39 μ L, 0.78 mmol, 0.25 eq) was added and the mixture stirred at 80 °C for an additional 3 h. The mixture was cooled to RT, diluted with water, and extracted three times with 5:1 DCM:*i*-PrOH. The combined organic extracts were washed five times with water, dried over MgSO_4 , filtered, and concentrated *in vacuo*. A silica gel column (24 g) was loaded using DCM and run with an increasing gradient of EtOAc (0-100% over 20 min) in hexanes to afford impure (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-55**, 0.95 g) as an off white solid. The crude white solid was recrystallized from 1:1 hexanes/EtOAc to afford crude 4-55 (0.70 g) as an off white solid. A reversed-phase C18 column (100 g) was loaded using DMSO and run with an increasing gradient of MeCN (10-70% over 25 min) in water to afford (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-55**, 0.59 g, 1.55 mmol, 50%) as a white solid. ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ (ppm) 6.71 (s, 1H), 6.63 (s, 1H), 4.77 (d, $J = 4.4$ Hz, 1H), 4.54 (d, $J = 6.6$ Hz, 1H), 3.96 – 3.87 (m, 1H), 3.77 (dd, $J = 9.3, 6.3$ Hz, 1H), 3.72 (s, 3H), 3.68 (dd, $J = 9.5, 5.5$ Hz, 1H), 3.11 – 3.05 (m, 1H), 2.99 (br d, $J = 11.0$ Hz, 1H), 2.94 (dd, $J = 11.8, 4.1$ Hz, 1H), 2.90 – 2.83 (m, 2H), 2.55 – 2.51 (m, 1H), 2.47 – 2.42 (m, 1H), 2.30 – 2.24 (m, 1H), 1.91 (t, $J = 11.2$ Hz, 1H), 1.69 (d, $J = 12.6$

Hz, 1H), 1.47 – 1.40 (m, 1H), 1.25 (q, $J = 11.5$ Hz, 1H), 1.13 (d, $J = 6.0$ Hz, 3H), 0.90 (s, 9H), 0.80 (dd, $J = 13.7, 7.7$ Hz, 1H).

Representative reagents (*i.e.*, R^1-X) used to synthesize compounds in Table 10A and Table 10B, include those shown in Table 8.

Table 8

R^1-X	R^1-X	R^1-X	R^1-X
			
			
			
			
			
			
			
			
			
			

Representative epoxide reagents used to synthesize compounds shown in **Table 10A** and **Table 10B**, include those shown in **Table 9**.

Table 9
































Epoxide	Epoxide	Epoxide
		
		

Table 10A and **Table 10B**, shown below, provide the observed (Obs) ion m/z ratio for **Compound 4-1**, **Compound 4-27**, **Compound 4-30**, **Compound 4-50**, **Compound 4-55**, and other representative compounds that were made analogously to the procedure described in the above examples. In many cases, a dialkylsulfate, alkyl bromide, alkyl chloride, or alkyl iodide shown in **Table 8** was used in place of a triflate. In other cases, an epoxide shown in **Table 9** was used in place of a triflate. Also frequently, final compounds were submitted directly to purification by preparative HPLC.

Table 10A

Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-1		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	402.2
4-2		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	384.2
4-3		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	398.3
4-4		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	402.2
4-5		(2 <i>R</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	416.2
4-6		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	384.2

Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-7		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	398.3
4-8		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	402.2
4-9		(2 <i>S</i> ,3 <i>S</i> ,11 <i>bS</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	416.2
4-10		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	384.2
4-11		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	398.3
4-12		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	416.2
4-13		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	384.2
4-14		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	398.3
4-15		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	416.2
4-16		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	402.2
4-17		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-ethoxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	348.3
4-18		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(2-fluoroethoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	366.2
4-19		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(3-fluorooxetan-3-yl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	408.3












Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-20		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[2-(oxetan-3-yloxy)ethoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	420.3
4-21		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>S</i>)-oxetan-2-ylmethoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	390.3
4-22		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	390.3
4-23		1-(2-((2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl)oxy)cyclobutane-1-carbonitrile	443.3
4-24		2-((2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl)oxy)acetonitrile	359.2
4-25		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-hydroxycyclobutyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	404.3
4-26		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2-methoxyethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	378.3
4-27		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-cyclopropoxy-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	360.3
4-28		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(3-fluoropropoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	380.3
4-29		1-((2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl)methylcyclopropane-1-carbonitrile	399.3
4-30		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-yloxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	376.3
4-31		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclopropyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	392.3
4-32		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-cyclobutoxy-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	374.3












Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-33		2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}- <i>N,N</i> -dimethylacetamide	405.3
4-34		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[2-(1 <i>H</i> -imidazol-1-yl)ethoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	414.3
4-35		2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}- <i>N</i> -methylacetamide	391.3
4-36		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(5-methyl-1,3-oxazol-2-yl)methoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	415.3
4-37		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1-methylazetidin-3-yl)methoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	403.3
4-38		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(propan-2-yloxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	362.3
4-39		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1,1,1-trifluoropropan-2-yl)oxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	416.2
4-40		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(ethoxy- <i>d</i> ₅)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol or (2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(² H ₃)ethoxy-10-methoxy-	353.3
4-41		4-(2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethoxy)oxane-4-carbonitrile	473.3
4-42		1-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)cyclobutane-1-carbonitrile	413.3
4-43		2-(2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethoxy)-2-cyclopropylacetonitrile	443.3
4-44		2-(2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethoxy)-2-methylpropanenitrile	431.3
4-45		1-(2-{[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}ethyl)cyclopropane-1-carbonitrile	413.3












Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-46		2-[1-({(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl}oxy)methyl)cyclopropyl]acetonitrile	413.3
4-47		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(2-hydroxy-2-methylpropoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	392.3
4-48		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>S</i>)-2-hydroxy-3-methoxypropoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	408.3
4-49		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>R</i>)-3,3,3-trifluoro-2-hydroxypropoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	432.2
4-50		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>S</i>)-3,3,3-trifluoro-2-hydroxypropoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	432.2
4-51		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(3,3-difluoro-2-hydroxypropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	414.5
4-52		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(methoxy- <i>d</i> ₃)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol or (2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(² H ₃)methoxy-10-methoxy-	337.3
4-53		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3 <i>R</i>)-oxolan-3-yloxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	390.3
4-54		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3 <i>S</i>)-oxolan-3-yloxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	390.3
4-55		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>R</i>)-2-hydroxypropoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	378.3












Table 10B











Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-56		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2-methylprop-2-en-1-yl)oxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	374.3

Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-57		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9- $\{[1-(\text{difluoromethyl})\text{cyclopropyl}]\text{methoxy}\}$ -3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	424.3
4-58		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>R</i>)-2-hydroxybutoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	392.3
4-59		5- $\{[(2R,3R,11bR)-3-(2,2\text{-dimethylpropyl})-2\text{-hydroxy-10-methoxy-1}H,2H,3H,4H,6H,7H,11bH\text{-pyrido[2,1-}a\text{]isoquinolin-9-yl)oxy}\}$ methyl)furan-2-carbonitrile	425.2
4-60		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	410.3
4-61		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclobutyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	406.3
4-62		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(4,4,4-trifluoro-2-hydroxybutoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	446.3
4-63		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(4-fluorophenyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	428.3
4-64		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(3-fluorophenyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	428.3
4-65		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(2-chloro-2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	418.2
4-66		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoro-2-hydroxy-2-methylpropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	446.2
4-67		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9- $\{[3-(\text{trifluoromethyl})\text{bicyclo[1.1.1]pentan-1-yl}]\text{methoxy}\}$ -1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	468.2

Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-68		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>S</i>)-2-hydroxybutoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	392.3
4-69		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2-oxazol-3-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	401.3
4-70		2-(2-{{(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl}oxy}ethoxy)acetonitrile	403.3
4-71		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[2-(3,3-difluoropyrrolidin-1-yl)ethoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	453.3
4-72		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(6-fluoro-1,4-dioxepan-6-yl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	452.3
4-73		4-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl}oxy)methyl)piperidine-1-carbonitrile	442.3
4-74		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{2-[4-(trifluoromethyl)phenyl]ethoxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	492.3
4-75		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,3-oxazol-4-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	401.3
4-76		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[(4,4-difluoro-1-hydroxycyclohexyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	468.3
4-77		6-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl}oxy)methyl}pyridine-2-carbonitrile	436.3
4-78		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-{[3-hydroxy-3-(trifluoromethyl)cyclobutyl]methoxy}-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	472.3

Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-79		((2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[2-hydroxy-3-(2,2,2-trifluoroethoxy)propoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	476.3
4-80		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[1-(trifluoromethyl)cyclopentyl]methoxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	470.3
4-81		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[4-(trifluoromethyl)phenyl]methoxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	478.3
4-82		6-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)pyridine-3-carbonitrile	436.3
4-83		(2 <i>R</i>)-3-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}propane-1,2-diol	394.3
4-84		2-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}- <i>N</i> -methyl- <i>N</i> -(2,2,2-trifluoroethyl)acetamide	473.2
4-85		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(3-methanesulfonylpropoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	440.2
4-86		1-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}- <i>N,N</i> -dimethylmethanesulfonamide	441.3
4-87		3-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}-1-(2,2,2-trifluoroethyl)pyrrolidin-2-one	485.2
4-88		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(cyclopropylmethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	374.3
4-89		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(3-hydroxypropoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	378.3

Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-90		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-{[(1 <i>R</i> ,2 <i>R</i>)-2-ethylcyclopropyl]methoxy}-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	402.3
4-91		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>R</i>)-2-methoxypropoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	392.3
4-92		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2 <i>S</i>)-2-methoxypropoxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	392.3
4-93		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(2-ethyl-2-hydroxybutoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	420.3
4-94		1-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)cyclopropyl methanesulfonate	468.3
4-95		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2,4-oxadiazol-5-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	402.3
4-96		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-methanesulfonylcyclopropyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	452.3
4-97		4-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)-1 <i>λ</i> ⁶ -thiane-1,1-dione	466.3
4-98		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(2-hydroxyethoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	364.3
4-99		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-{{4-(trifluoromethyl)cyclohexyl}oxy}-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	470.3
4-100		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2-fluorocyclopentyl)oxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	405.3



Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-101		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[(3,3-difluorocyclopentyl)oxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	424.3
4-102		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[(4-difluoromethanesulfonylphenyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	524.2
4-103		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(3-chloro-3-(difluoropropoxy))-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	431.2
4-104		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1-methylazetidin-3-yl)oxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	389.3
4-105		2-(2-({(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl}oxy)ethoxy)propanenitrile	417.3
4-106		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-({3-fluorobicyclo[1.1.1]pentan-1-yl}methoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	418.3
4-107		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(pyridin-3-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	411.3
4-108		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(pyrimidin-2-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	412.3
4-109		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2-oxazol-5-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	401.3
4-110		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-(1 <i>H</i> -imidazol-2-ylmethoxy)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	400.3
4-111		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1 <i>H</i> -1,2,3,4-tetrazol-5-ylmethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	402.3

Cmpd. No.	-R ¹	Chemical Name	Obs. Ion (m/z)
4-112		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>R</i>)-2-hydroxy-3-methoxypropoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	407.3
4-113		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[2-hydroxy-3-(morpholin-4-yl)propoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	462.3
4-114		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(2 <i>S</i>)-2-hydroxypropoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	378.3
4-115		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(5,5,5-trifluoropentyl)oxy]-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	444.4
4-116		3-({[(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-9-yl]oxy}methyl)benzonitrile	435.3
4-117		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[2-(4-fluorophenyl)ethoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	442.3
4-118		(2 <i>S</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclopropyl)methoxy]-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	392.3
4-119		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-(cyclobutylmethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	388.4
4-120		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-dimethylpropyl)-10-methoxy-9-(4,4,4-trifluorobutoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	430.3
4-121		(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-9-[(2,2-difluorocyclopropyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol	410.3

A summary of representative physical properties for **Compound 4-1** prepared according to the methods disclosed herein are provided in **Table 11** and **Table 12**, and a summary of representative

physical properties for **Compound 4-15** prepared according to the methods disclosed herein are provided in **Table 13** and **Table 14**.

EXAMPLE 6: Powder X-Ray Diffraction (PXRD), Differential Scanning Calorimetry (DSC), and Thermal Gravimetric (TGA) Analyses for (2*R*,3*R*,11*bR*)-3-(2,2-Dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 4-1) and (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 4-15).



POWDER X-RAY DIFFRACTION (PXRD)

The powder X-ray powder diffraction (PXRD) analysis was performed on a Rigaku Powder X-Ray Diffractometer Miniflex 600 Serial Number BD66000190-01. For analysis, approximately 0.5-1 mg of sample was added to a zero-background sample holder. The powder was pressed down gently with a piece of weigh paper, and the sample holder was placed in the sample changer. Run Parameters: Miniflex Counter Detector, K α Filter (x2), Scan Axis Theta/2-Theta, Mode Continuous, Start (deg) 2.0, Stop (deg) 45.0, Step (deg) 0.020, Speed (deg/min) 10.0, Spin-yes, Voltage (kV) 40, Current (mA) 15. The reported 2 θ values can vary by plus or minus 0.2° (*i.e.*, $\pm 0.2^\circ$).

It is understood that peak intensities can vary from one diffractogram to another for the same crystalline form based on any number of factors that are known to those skilled in the art, such as, preferred orientation effects, preparation technique, the sample mounting procedure, the instrument employed, sample purity, etc. One skilled in the art would understand that despite the differences in the peak intensities, a particular crystal form can be identified by its characteristic peak(s).

DIFFERENTIAL SCANNING CALORIMETRY (DSC) AND THERMAL GRAVIMETRIC ANALYSIS (TGA):

The DSC and TGA analyses were performed on TA Instruments Discovery 2500 calorimeter with serial number: 2500-00547 (DSC) and Discovery 5500 with serial number: 5500-0126 (TGA). For the DSC analysis, obtained and recorded the weight of a Tzero pan and a Tzero lid. ~1-3 mg of material was weighed into the Tzero Pan and the Tzero lid was pressed on with tweezers. The pan was transferred to the DSC autosampler for analysis. The method for analysis was a ramp at 10 °C/min to 275 °C. The reference pan was prepared with the same procedures, absent of material.

For TGA analysis, a standard aluminum sample pan was placed into the platinum TGA pan and the blank was tared with the instrument. Approximately 1-5 mg of material was added to the standard aluminum pan and analyzed at 10 °C/min up to ~400 °C. For TGA and DSC, the temperature features reported herein may vary by plus or minus 3 °C (*i.e.*, ± 3 °C).

5 **Table 11** provides select powder X-ray diffraction (PXRD) diffractogram peak positions at 2 θ , DSC and TGA data for (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-1**) free base crystalline Form I.

Table 11	(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-Dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol (Compound 4-1)
PXRD	FIG. 2: Peaks at $10.2 \pm 0.2^\circ$, $11.7 \pm 0.2^\circ$, $15.2 \pm 0.2^\circ$ and $17.0 \pm 0.2^\circ$ 2 θ
DSC	FIG. 3: Apparent onset temperature: about 154.7 °C
TGA	FIG. 3: Weight percent loss: about 4.0%

10 **Table 12** provides powder X-ray diffraction (PXRD) diffractogram peak positions at 2 θ for (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-1**) free base crystalline Form I.

15 **Table 12**

Angle 2 θ (2-theta)	Height (cps)
$6.6 \pm 0.2^\circ$	2546 (146)
$10.2 \pm 0.2^\circ$	17569 (383)
$11.7 \pm 0.2^\circ$	6874 (239)
$13.2 \pm 0.2^\circ$	3069 (160)
$13.9 \pm 0.2^\circ$	2883 (155)
$14.1 \pm 0.2^\circ$	5957 (223)
$15.2 \pm 0.2^\circ$	8497 (266)
$16.4 \pm 0.2^\circ$	1461 (110)
$17.0 \pm 0.2^\circ$	10114 (290)
$19.1 \pm 0.2^\circ$	15522 (360)
$19.5 \pm 0.2^\circ$	5227 (209)
$19.7 \pm 0.2^\circ$	4430 (192)
$20.6 \pm 0.2^\circ$	1622 (116)
$21.6 \pm 0.2^\circ$	14740 (350)

Angle 2 θ (2-theta)	Height (cps)
$21.9 \pm 0.2^\circ$	9609 (283)
$22.8 \pm 0.2^\circ$	13213 (332)
$23.7 \pm 0.2^\circ$	3139 (162)
$24.0 \pm 0.2^\circ$	8568 (267)
$25.4 \pm 0.2^\circ$	2646 (148)
$25.6 \pm 0.2^\circ$	5154 (207)
$27.2 \pm 0.2^\circ$	7410 (248)
$27.6 \pm 0.2^\circ$	6166 (227)
$27.9 \pm 0.2^\circ$	1120 (97)
$29.8 \pm 0.2^\circ$	1596 (115)
$30.3 \pm 0.2^\circ$	642 (73)
$31.2 \pm 0.2^\circ$	1408 (108)
$31.9 \pm 0.2^\circ$	1120 (97)
$32.3 \pm 0.2^\circ$	1013 (92)

Angle 2θ (2-theta)	Height (cps)
$34.0 \pm 0.2^\circ$	380 (56)
$34.9 \pm 0.2^\circ$	1948 (127)
$35.4 \pm 0.2^\circ$	750 (79)
$35.8 \pm 0.2^\circ$	617 (72)
$37.6 \pm 0.2^\circ$	431 (60)
$39.3 \pm 0.2^\circ$	566 (69)

Angle 2θ (2-theta)	Height (cps)
$40.2 \pm 0.2^\circ$	251 (46)
$41.2 \pm 0.2^\circ$	1987 (129)
$41.6 \pm 0.2^\circ$	1642 (117)
$42.0 \pm 0.2^\circ$	550 (68)
$43.1 \pm 0.2^\circ$	1155 (98)
$44.4 \pm 0.2^\circ$	507 (65)

Table 13 provides select powder X-ray diffraction (PXRD) diffractogram peak positions at 20, DSC and TGA data for (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-15**) free base crystalline Form I.

Table 13	(2 <i>R</i> ,3 <i>R</i> ,11 <i>bR</i>)-3-(2,2-Dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1 <i>H</i> ,2 <i>H</i> ,3 <i>H</i> ,4 <i>H</i> ,6 <i>H</i> ,7 <i>H</i> ,11 <i>bH</i> -pyrido[2,1- <i>a</i>]isoquinolin-2-ol (Compound 4-15)
PXRD	FIG. 4: Peaks at $11.4 \pm 0.2^\circ$, $14.1 \pm 0.2^\circ$, $15.7 \pm 0.2^\circ$ and $18.9 \pm 0.2^\circ$ 2θ
DSC	FIG. 5: Apparent onset temperature: about 146.2 °C, and about 153.4 °C
TGA	FIG. 5: Weight percent loss: about 0.17%

Table 14 provides powder X-ray diffraction (PXRD) diffractogram peak positions at 2θ for (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-15**) free base crystalline Form I.

Table 14


Angle 2θ (2-theta)	Height (cps)
$5.6 \pm 0.2^\circ$	1248 (102)
$6.7 \pm 0.2^\circ$	1744 (121)
$7.1 \pm 0.2^\circ$	1151 (98)
$8.4 \pm 0.2^\circ$	2630 (148)
$9.2 \pm 0.2^\circ$	5583 (216)
$10.6 \pm 0.2^\circ$	8271 (263)
$11.4 \pm 0.2^\circ$	13856 (340)
$12.3 \pm 0.2^\circ$	1809 (123)
$12.8 \pm 0.2^\circ$	1234 (101)

Angle 2θ (2-theta)	Height (cps)
$13.3 \pm 0.2^\circ$	4742 (199)
$14.1 \pm 0.2^\circ$	12229 (319)
$14.6 \pm 0.2^\circ$	489 (64)
$15.7 \pm 0.2^\circ$	24533 (452)
$16.0 \pm 0.2^\circ$	4993 (204)
$16.9 \pm 0.2^\circ$	8830 (271)
$17.3 \pm 0.2^\circ$	2584 (147)
$17.6 \pm 0.2^\circ$	8835 (271)
$17.9 \pm 0.2^\circ$	8446 (265)

Angle 2θ (2-theta)	Height (cps)
$18.5 \pm 0.2^\circ$	5690 (218)
$18.9 \pm 0.2^\circ$	16374 (369)
$19.6 \pm 0.2^\circ$	11400 (308)
$20.1 \pm 0.2^\circ$	13479 (335)
$20.4 \pm 0.2^\circ$	6903 (240)
$21.2 \pm 0.2^\circ$	10829 (300)
$21.9 \pm 0.2^\circ$	6742 (237)
$22.4 \pm 0.2^\circ$	1919 (126)
$22.9 \pm 0.2^\circ$	15091 (355)
$23.1 \pm 0.2^\circ$	19640 (405)
$23.7 \pm 0.2^\circ$	3951 (181)
$24.6 \pm 0.2^\circ$	3618 (174)
$24.9 \pm 0.2^\circ$	3464 (170)
$25.2 \pm 0.2^\circ$	957 (89)
$26.1 \pm 0.2^\circ$	2206 (136)
$27.1 \pm 0.2^\circ$	6631 (235)
$27.9 \pm 0.2^\circ$	3317 (166)
$28.4 \pm 0.2^\circ$	2722 (151)

Angle 2θ (2-theta)	Height (cps)
$29.9 \pm 0.2^\circ$	3566 (172)
$30.9 \pm 0.2^\circ$	794 (81)
$31.7 \pm 0.2^\circ$	2852 (154)
$32.6 \pm 0.2^\circ$	1383 (107)
$33.1 \pm 0.2^\circ$	1610 (116)
$33.8 \pm 0.2^\circ$	558 (68)
$34.3 \pm 0.2^\circ$	1942 (127)
$35.2 \pm 0.2^\circ$	418 (59)
$36.6 \pm 0.2^\circ$	1069 (94)
$37.1 \pm 0.2^\circ$	921 (88)
$37.7 \pm 0.2^\circ$	481 (63)
$38.4 \pm 0.2^\circ$	1013 (92)
$40.0 \pm 0.2^\circ$	675 (75)
$40.3 \pm 0.2^\circ$	989 (91)
$41.0 \pm 0.2^\circ$	563 (68)
$42.1 \pm 0.2^\circ$	854 (84)
$44.2 \pm 0.2^\circ$	1678 (118)

EXAMPLE 7: Preparation of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 5-1), see FIG. 13.



Step 1: Synthesis of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 5-A (3*R*,11*bR*)).


To a solution of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 2-B (3*R*,11*bR*)**), 95 mg, 0.30 mmol, 1.0 eq) in acetone (1.5 mL) was added cesium carbonate (0.19 g, 0.60 mmol, 2.0 eq) and the resulting mixture stirred at RT for 10 min. Then, dimethyl sulfate (17 μ L, 0.18 mmol, 0.6 eq) was added and the resulting mixture stirred at RT for 1h. An additional 0.1 eq of dimethyl sulfate was added and the mixture was stirred an additional 30 min. The resulting suspension was filtered to remove solids and concentrated *in vacuo*. The crude material was re-dissolved in DCM, quenched with 1M aq. NaOH and extracted with DCM. The combined organic layers were dried over MgSO_4 ,

filtered to remove solids, and concentrated *in vacuo*. Silica gel column (12 g) was loaded using DCM and run with an increasing gradient of EtOAc (0-40% over 25 min) in hexanes to provide (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 5-A (3*R*,11*bR*), 75 mg, 0.23 mmol, 77%).

Step 2: Synthesis of (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 5-1).

To a solution of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 5-A (3*R*,11*bR*), 33 mg, 0.10 mmol, 1.0 eq) in EtOH (0.6 mL) cooled to 0 °C was added sodium borohydride (7.4 mg, 0.20 mmol, 2.0 eq). The resulting mixture was stirred at 0 °C for 30 min, warmed to RT and stirred for 1h. The mixture was diluted with DCM, washed twice with sat. aq. K₂CO₃ and the aqueous layer was further extracted with DCM. The combined organic layers were dried over MgSO₄, filtered to remove solid and concentrated *in vacuo*. Silica gel column (12 g) was loaded using DCM and run with an increasing gradient of EtOAc (0-60%) in hexanes over 15 min to provide (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 5-1, 20 mg, 0.06 mmol, 60%) as a white solid. 10 mg was dissolved in MeOH (1 mL) and submitted for additional purification by preparative HPLC. ¹H NMR (400 MHz, DMSO-*d*₆): δ (ppm) 6.72 (s, 1H), 6.64 (s, 1H), 4.55 (d, *J* = 6.4 Hz, 1H), 3.72 (s, 3H), 3.71 (s, 3H), 3.13 – 3.05 (m, 1H), 3.01 (d, *J* = 11.2 Hz, 1H), 2.95 (dd, *J* = 11.8, 4.2 Hz, 1H), 2.91 – 2.82 (m, 2H), 2.58 – 2.54 (m, 1H), 2.48 – 2.43 (m, 1H), 2.33 – 2.25 (m, 1H), 1.93 (t, *J* = 11.4 Hz, 1H), 1.70 (d, *J* = 12.8 Hz, 1H), 1.48 – 1.39 (m, 1H), 1.25 (q, *J* = 12.8 Hz, 1H), 0.91 (s, 9H), 0.81 (dd, *J* = 13.6, 7.6 Hz, 1H). The observed (Obs) ion *m/z* ratio for Compound 5-1 was 334.2.

EXAMPLE 8: Preparation of (2*S*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 5-3), see FIG. 13.




Step 1: Synthesis of (3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (Compound 5-A (3*S*,11*bS*)).

Compound 5-A (3*S*,11*bS*) was prepared according to the same procedure used to prepare Compound 5-A (3*R*,11*bR*) (see Example 7, Step 1) using Compound 2-B (3*S*,11*bS*) in place of Compound 2-B (3*R*,11*bR*), see FIG. 13.

Step 2: Synthesis of (2*S*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 5-3).


Compound 5-3 was prepared according to the same procedure used to prepare **Compound 5-1** (see **Example 7, Step 2**) using **Compound 5-A (3*S*,11*bS*)** in place of **Compound 5-A (3*R*,11*bR*)**. The observed (Obs) ion *m/z* ratio for **Compound 5-3** was 334.2.

EXAMPLE 9: Preparation of (2*S*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 5-2), see FIG. 14.



A solution of (3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-one (**Compound 5-A (3*R*,11*bR*)**, 27 mg, 0.08 mmol, 1.0 eq) in THF (0.6 mL) was prepared and cooled to 0 °C. Then, lithium tri-*sec*-butylborohydride (1M THF, 0.16 mL, 0.16 mmol, 2.0 eq) was added dropwise. The excess reagent was quenched with water. The resulting aqueous mixture was extracted with DCM. The combined organic layers were dried over MgSO₄, filtered to removed solids and concentrated *in vacuo*. Silica gel column (4g) was loaded using DCM and run with an increasing gradient of EtOAc in hexanes to provide (2*S*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-2**, 17 mg, 0.05 mmol, 63%). ¹H NMR (400 MHz, DMSO-*d*₆): δ (ppm) 6.64 (s, 1H), 6.63 (s, 1H), 4.46 (d, *J* = 4.0 Hz, 1H), 3.84 – 3.80 (m, 1H), 3.70 (s, 6H), 3.38 (d, *J* = 11.2 Hz, 1H), 2.94 – 2.83 (m, 2H), 2.56 – 2.46 (m, 2H), 2.41 – 2.26 (m, 3H), 1.72 – 1.65 (m, 1H), 1.50 – 1.43 (m, 2H), 0.90 (s, 9H), 0.84 (dd, *J* = 14.2, 4.6 Hz, 1H). The observed (Obs) ion *m/z* ratio for **Compound 5-2** was 334.2.

EXAMPLE 10: Preparation of (2*R*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (Compound 5-4), see FIG. 14.



(2*R*,3*S*,11*bS*)-3-(2,2-Dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-4**) was prepared according to the same procedure to prepare **Compound 5-2** (see **Example 9**) using **Compound 5-A (3*S*,11*bS*)** in place of **Compound 5-A (3*R*,11*bR*)**. The observed (Obs) ion *m/z* ratio for **Compound 5-4** was 334.2.

EXAMPLE 11: Methods for Determining VMAT2 Inhibitory Activity of a Compound.

Examples of techniques for determining the capability of a compound to inhibit VMAT2 are provided below. The procedure was adapted from that described previously (see, *e.g.*, Near, (1986), Mol. Pharmacol. 30: 252-57; Teng, *et al.*, J. Neurochem. 71, 258-65, 1998). Homogenates from human platelets were prepared by homogenization and then washed by centrifugation as described previously (see, *e.g.*, Hoare *et al.*, (2003) Peptides 24:1881-97).

The human VMAT2 K_i values for the compounds listed in Table 15A and Table 15B were determined using the following procedures. Compound dilution series in DMSO were generated from either powder stocks by hand, or from DMSO stocks using serial dilution on a Vantage (Hamilton), or direct dilution using an Echo 655 (Beckman). In a total volume of 0.145 to 0.150 mL in low-binding 96-well plates (Corning #3605), twelve concentrations of test compound were competed against 10 nM ^3H -dihydrotetrabenzine (American Radiolabeled Chemicals) on human platelet homogenate (30 μg membrane protein per well) in VMAT2 binding buffer (Dulbecco's phosphate buffered saline, 1 mM EDTA, pH 7.4). Following incubation at 25 °C for 90 minutes, bound radioligand was collected by rapid filtration onto either GF/B or GF/C glass fiber filters, pretreated with 0.1% polyethylenimine, using either a Unifilter-96 Harvester (PerkinElmer) or Microlab Star (Hamilton). Following harvesting the filter plates were washed with 0.8 mL VMAT2 binding buffer, and bound radioligand was quantified by scintillation counting using a Topcount NXT or Microplate Counter Microbeta (PerkinElmer). Data from 12-point concentration response curves were analyzed to calculate an IC_{50} using a four-parameter logistic regression algorithm, where top was constrained to 100 and bottom was constrained to 0. The K_i value for each compound was calculated using the Cheng-Prusoff equation, utilizing a K_d of 2.5 nM or 4.06 nM for ^3H -dihydrotetrabenzine, depending on the batch of platelets and radiolabel used in each experiment.

Compound K_i (nM) values are provided in Table 15A and Table 15B.

Table 15A

Cmpd. No.	VMAT2 K_i (nM)	Cmpd. No.	VMAT2 K_i (nM)	Cmpd. No.	VMAT2 K_i (nM)
4-1	2.4	4-21	28	4-41	238
4-2	467	4-22	14	4-42	35
4-3	152	4-23	0.41	4-43	4.8
4-4	170	4-24	35	4-44	2.4
4-5	135	4-25	5.5	4-45	7.3
4-6	860	4-26	14	4-46	25
4-7	539	4-27	1.0	4-47	15
4-8	597	4-28	0.48	4-48	313
4-9	277	4-29	83	4-49	20
4-10	15	4-30	12	4-50	1.4
4-11	14	4-31	1.6	4-51	9.4
4-12	3.4	4-32	0.59	4-52	2.5
4-13	2.2	4-33	> 10,000	4-53	12

Cmpd. No.	VMAT2 K _i (nM)
4-14	2.6
4-15	1.1
4-16	22
4-17	1.1
4-18	5.5
4-19	8.1
4-20	336

Cmpd. No.	VMAT2 K _i (nM)
4-34	> 10,000
4-35	> 10,000
4-36	51
4-37	> 10,000
4-38	3.7
4-39	12
4-40	1.4

Cmpd. No.	VMAT2 K _i (nM)
4-54	12
4-55	4.8
5-1	6.6
5-2	50
5-3	700
5-4	> 10,000

Table 15B

Cmpd. No.	VMAT2 K _i (nM)
4-56	0.52
4-57	0.93
4-58	1.1
4-59	1.5
4-60	3.1
4-61	3.5
4-62	3.7
4-63	4.2
4-64	5.5
4-65	8.7
4-66	10
4-67	14
4-68	29
4-69	32
4-70	36
4-71	37
4-72	51
4-73	51
4-74	65
4-75	76
4-76	82
4-77	85

Cmpd. No.	VMAT2 K _i (nM)
4-78	92
4-79	99
4-80	133
4-81	177
4-82	198
4-83	207
4-84	272
4-85	276
4-86	287
4-87	650
4-88	0.68
4-89	140
4-90	0.29
4-91	28
4-92	51
4-93	281
4-94	387
4-95	> 10,000
4-96	> 10,000
4-97	> 10,000
4-98	NT
4-99	NT

Cmpd. No.	VMAT2 K _i (nM)
4-100	NT
4-101	NT
4-102	NT
4-103	NT
4-104	NT
4-105	NT
4-106	NT
4-107	NT
4-108	NT
4-109	NT
4-110	NT
4-111	NT
4-112	NT
4-113	NT
4-114	70
4-115	6.1
4-116	56
4-117	2.2
4-118	13
4-119	0.29
4-120	1.5
4-121	0.82

NT = Not Tested

5 **FIG. 15 to FIG. 21** provide VMAT2 K_i (nM) values for representative compounds of Formula (I) and comparator compounds. The VMAT2 K_i values were determined according to **Example 11**. The comparator compounds were prepared as described in Intl. Pub. No. WO 2018/195121 A1, which is incorporated herein by reference in its entirety. Representative compounds of Formula (I) are more potent than the comparator compounds in terms of VMAT2 K_i values (*e.g.*, 10 4.1-times to 20.9-times).

EXAMPLE 12: VMAT2 Inhibitor-Induced Reduction of Open-Field Hypolocomotor Activity.

The effect of VMAT2 inhibitors on dopamine depletion was measured using the locomotor activity (LMA) assay. Following a pre-treatment time of 60 or 120 minutes, male Sprague-Dawley rats (200-250 g) were placed in a clear cage surrounded by photocell detectors (San Diego Instruments). Rat locomotor activity was detected by breaks in the photocell beams and activity is defined as the number of beam breaks in 30 min. Data was analyzed by one-way analysis of variance (ANOVA; SigmaStat version 3.0.1, SPSS, Chicago, IL) followed by the Student Newman Keuls *post-hoc* test for significance. The ED₅₀ was calculated for **Compound 4-1** (0.24 mg/kg, see **FIG. 22**), **Compound 4-15** (0.18 mg/kg, see **FIG. 22**), **Compound 4-27** (0.14 mg/kg, see **FIG. 23**), **Compound 4-31** (0.31 mg/kg, see **FIG. 24**), **Compound 4-50** (0.21 mg/kg, see **FIG. 23**), and **Compound 4-55** (0.29 mg/kg, see **FIG. 23**).

EXAMPLE 13A: Conditioned Avoidance Response Assay of Antipsychotic Activity.

The conditioned avoidance response (CAR) test has been shown to be an effective and reliable preclinical model for assessing the antipsychotic activity of compounds. In the CAR paradigm, a rat is trained in a two-chamber shuttle box to respond to a conditioned stimulus (auditory) by negative reinforcement. If the animal fails to move to the other chamber upon presentation of an auditory stimulus, a mild foot shock is applied to the side where the rat is located. The rat learns to avoid the mild foot shock by moving to the other chamber upon initiation of the auditory signal, termed a conditioned avoidance response. Crossing to the other chamber during administration of the shock is termed an escape response. If a rat fails to move to the other chamber even upon administration of the foot shock, the rat is considered to have an escape failure. Numerous studies have shown that typical and atypical antipsychotic drugs selectively suppress CAR, thus making it an ideal assay to screen potential antipsychotic compounds (see, *e.g.*, Wadenberg *et al.*, *Biobehav. Rev.* (1999) 23: 851-62).

Male Wistar rats are trained every day for 3 to 4 weeks. In the training session, rats are placed in the CAR two-way shuttle box and the training period of 20 trials ensued. A trial consisted of a 10-sec presentation of an 80 dB white noise followed by a scrambled 0.6 mA foot shock lasting up to 20 sec. The inter-trial interval ranged from 20-60 sec. The rat learns to avoid shock by moving from one compartment to the other when the conditioned stimulus is presented (a conditioned avoidance response). A rat is deemed sufficiently trained if it avoided the shock when presented with the conditioned stimulus at least 19 times out of the 20 trials. Rats that do not pass these criteria are not used.

On test day, trained animals are acclimated in the test room for 30 minutes prior to testing. They were then dosed with compound and are placed in the CAR two-way shuttle box. In the test, 20 trials are performed on each rat. In each trial the conditioned stimulus is applied (10-sec presentation of 80 dB white noise), followed by the foot shock (a scrambled 0.6 mA foot shock lasting up to 20

sec). If the animal moves to the other chamber on presentation of the conditioned stimulus, it is scored as a conditioned avoidance response. If it moved upon presentation of the foot shock, it is scored as an escape. If it fails to move upon presentation of the foot shock, it is scored as an escape failure.

Antipsychotic efficacy is evident by an increase in the number of escapes. Data are analyzed by analysis of variance (ANOVA) followed by post-hoc comparisons with the Bonferroni Test when appropriate. An effect is considered significant if $p < 0.05$. Outliers defined as two standard deviations above or below the mean are detected and are removed from all analysis.

EXAMPLE 13B: Conditioned Avoidance Response (CAR) Assay of Antipsychotic Activity.

Material and Methods

Animals: Adult male Wistar rats from Envigo (Indianapolis, IN) were used in this study. Rats were received at approximately 150g, assigned unique identification numbers and group housed 2-3 per cage in ventilated cages. All rats were examined, handled, and weighed prior to initiation of the study to assure adequate health and suitability. During the course of the study, 12/12 light/dark cycles were maintained. The RT was maintained between 20 °C and 23 °C with a relative humidity around 50%. Chow and water were provided *ad libitum* for the duration of the study. All testing was conducted during the light cycle of the rats.

Compound Formulation

Compound 4-1 (0.03, 0.1, 0.3, and 1 mg/kg) was formulated in 0.25% methyl cellulose and administered orally at a dose volume of 3 mL/kg 120 minutes prior to test.

Compound 4-15 (0.03, 0.1, 0.3, and 1 mg/kg) was formulated in 0.25% methyl cellulose and administered orally at a dose volume of 3 mL/kg 120 minutes prior to test.

Behavioral Testing -- Conditioned Avoidance Response

The conditioned avoidance response (CAR) test has been shown to be a very reliable animal model for screening antipsychotic drugs (see Wadenberg & Hicks, Neuroscience and Biobehavioral Reviews, (1999) 23:851–862). In the CAR paradigm, an animal is trained to respond to a conditioned stimulus (auditory and visual) by negative reinforcement (foot shock). Numerous studies have shown that typical and atypical antipsychotic drugs selectively suppress CAR, thus making it an ideal assay to screen potential antipsychotic compounds (Wadenberg & Hicks, Neuroscience and Biobehavioral Reviews, (1999) 23:851–862).

Rats were placed in the CAR two-way shuttle box and the training period of 20 trials ensued. A trial consisted of a 10-sec presentation of an 80 dB white noise followed by a scrambled 0.6 mA foot shock lasting up to 20 sec. The inter-trial interval ranged from 20-60 sec. The rat learned to avoid shock by moving from one compartment to the other when the cue is presented. If it failed to do so, it received the foot-shock, during which it most likely crossed compartments to escape the shock. Finally, if the rat failed to leave the compartment, it endures the full 20 s foot-shock. The following measures were captured from the assay:

Avoidance response: If the rat moved from one compartment to the other during the cued-stimulus (CS) presentation. Decreased avoidance responding is the typical signature of an efficacious dose of an antipsychotic.

Escape Failure: If the rat failed to move into the other compartment during the 20-sec foot-shock.

Data Analysis

Avoidance response data were expressed as the number of avoidance responses as well as percent of avoidance responses based on the three baseline responses prior to drug test. Escape failures were expressed as the total number of failures during the test session.

Body weight and avoidance data were analyzed by analysis of variance (ANOVA) followed by Dunnett's post-hoc comparisons when appropriate. Escape failures were analyzed by Kruskal-Wallis nonparametric analysis followed by Dunn's post hoc comparisons when appropriate. Results are reported as mean $\pm$ SEM. An effect was considered statistically significant if $p < 0.05$.

The ED₅₀ was calculated for **Compound 4-1** (0.55 mg/kg). The ED₅₀ (mg/kg) was also calculated for **Compound 4-15** (0.44 mg/kg). In addition, rats treated with either **Compound 4-1** or **Compound 4-15** were observed to display a reduction in avoidance without significant increase in escape failures at 0.3 mg/kg.

EXAMPLE 14: Method to Determine Stability of Compounds in Human Liver Microsomes.

A test compound (0.5 μ M) was incubated with pooled mixed gender liver microsomes from humans (0.5mg/mL total protein) at 37 °C in the presence of an NADPH-generating system containing 50 mM, pH 7.4 potassium phosphate buffer, 3 mM magnesium chloride, 1 mM EDTA, 1 mM NADP, 5 mM glucose-6-phosphate, and 1 Unit/mL glucose-6-phosphate dehydrogenase. All concentrations were relative to the final incubation volume of 125 μ L. Incubations were conducted at 37 °C for 0, 5, 10, 20, 40, and 60 minutes in a water bath and terminated by rapid mixing with 150 μ L of ice-cold acetonitrile containing internal standard. Precipitated proteins were removed by centrifugation prior to LC-MS/MS analysis. Aliquots of the resulting supernatant fractions were analyzed by LC-MS/MS monitoring for depletion of parent compound. The resultant peak area ratio versus time data was fitted to a non-linear regression using XLfit Scientific Curve Fitting Software (IDBS Ltd., Surrey, UK) and the elimination half-life ($t_{1/2}$, min) was calculated from the slope. Pharmacokinetic parameters were predicted using the method described by Obach *et al.* (J. Pharmacol. Exp. Ther. 1997; 283: 46.58). Briefly, the values for intrinsic clearance were calculated from the elimination half-life data and were then scaled to represent the clearance expected in the entire animal, see **Table 16** (human). Additional values calculated included predicted extraction ratio and predicted maximum bioavailability.

For very stable compounds the *in vitro* half-life calculated from the HLM method is maxed at 420 minutes. Accordingly, for these stable compounds the *in vitro* half-life is at least 420 minutes but

could be greater. Further, for these stable compounds the predicted systemic clearance and scaled intrinsic clearance is at least 2.59 and 2.97 respectively but could be lower, and the predicted maximum bioavailability (%F) is at least 87 but could be higher.

Table 16 (Human LM)

Cmpd. No.	HLM Elimination half-life (t _{1/2} , min)	HLM Predicted Systemic Clearance (ml/min/kg)	HLM Scaled Intrinsic Clearance (ml/min/kg)	Predicted Max. Bioavailability (%F)
4-1	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-9	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-10	342	3.09	3.65	85
4-11	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-12	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-13	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-14	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-15	308	3.37	4.05	83
4-16	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-17	38	12.44	32.9	38
4-18	155	5.74	8.05	71
4-19	88	8.3	14.18	59
4-22	130	6.49	9.6	68
4-23	21	14.9	58.49	25
4-24	117	6.97	10.7	65
4-25	109	7.26	11.41	64
4-26	88	8.32	14.24	58
4-27	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-28	39	12.28	31.79	39
4-29	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-30	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-31	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-32	19	15.38	66.53	23
4-36	93	8.02	13.39	60
4-38	58	10.35	21.46	48
4-40	52	10.88	23.86	46
4-41	28	13.87	45.21	31
4-42	51	10.98	24.33	45
4-43	18	15.5	68.8	23
4-44	46	11.47	26.9	43
4-45	129	6.53	9.7	67
4-46	28	13.86	45.1	31
4-47	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-49	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-50	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-51	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-52	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-53	146	6	8.57	70

Cmpd. No.	HLM Elimination half-life ($t_{1/2}$, min)	HLM Predicted Systemic Clearance (ml/min/kg)	HLM Scaled Intrinsic Clearance (ml/min/kg)	Predicted Max. Bioavailability (%F)
4-54	164	5.5	7.59	73
4-55	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-56	10	17.16	120.93	14
4-57	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-58	208	4.61	5.99	77
4-59	50	11.08	24.82	45
4-60	37	12.58	33.93	37
4-61	30	13.55	42.02	32
4-62	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-63	29	13.58	42.33	32
4-64	33	13.1	37.96	35
4-65	227	4.31	5.5	78
4-66	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-67	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-68	196	4.83	6.37	76
4-69	80	8.74	15.51	56
4-70	110	7.25	11.37	64
4-71	33	13.15	38.38	34
4-72	80	8.77	15.62	56
4-73	119	6.88	10.48	66
4-74	36	12.68	34.62	37
4-75	104	7.48	11.94	63
4-76	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-77	156	5.72	8.01	71
4-78	195	4.86	6.41	76
4-79	212	4.54	5.88	77
4-88	44	11.68	28.05	42
4-90	8	17.72	155.72	11
4-91	63	9.92	19.7	50
4-92	49	11.2	25.46	44
4-114	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-115	133	6.39	9.38	68
4-116	57	10.44	21.84	48
4-117	11	16.98	112.26	15
4-118	189	4.96	6.6	75
4-119	9	17.4	133.88	13.00
4-120	≥ 420	≤ 2.59	≤ 2.97	≥ 87
4-121	280	3.64	4.46	82
5-1	80	8.77	15.63	56
5-2	38	12.39	32.56	38

The various embodiments described above can be combined to provide further embodiments.

All U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents,


foreign patent applications, and non-patent publications referred to in this specification and/or listed in the Application Data Sheet are incorporated herein by reference in their entirety. Aspects of the embodiments can be modified, if necessary, to employ concepts of the various patents, applications, and publications to provide yet further embodiments.

- 5 These and other changes can be made to the embodiments in light of the above-detailed description. In general, in the following claims, the terms used should not be construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

CLAIMS

What is claimed is:

1. A compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein:

R^1 is R^{2A} , $-(CH_2)_m-R^{2B}$, or $-(CH_2)_n-OR^{2C}$;

R^{2A} is C_2 - C_4 alkenyl, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more R^A ;

each R^A is independently selected from the group consisting of: C_1 - C_4 alkoxy, C_1 - C_4 alkyl, C_1 - C_4 alkylamino, C_1 - C_4 alkylsulfonyl, C_2 - C_4 dialkylamino, C_2 - C_4 dialkylsulfamoyl, halogen, halo C_1 - C_4 alkoxy, halo C_1 - C_4 alkyl, halo C_2 - C_4 dialkylamino, 4-7 membered heterocyclyl, -CN, -OH, and oxo;

R^{2B} is aryl, C_3 - C_8 cycloalkyl, 5-10 membered heteroaryl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more R^B ;

each R^B is independently selected from the group consisting of: C_1 - C_4 alkyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 alkylsulfonyloxy, -CN, -CH₂CN, halogen, halo C_1 - C_4 alkyl, halo C_1 - C_4 alkylsulfonyl, oxo, and -OH;


R^{2C} is C_1 - C_4 alkyl, C_3 - C_8 cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more R^C ;

each R^C is independently -CN or C_3 - C_6 cycloalkyl;

m is 0, 1, 2, 3, or 4; and


n is 1, 2, or 3.

2. The compound of claim 1, wherein the compound has Formula (Ia):




or a pharmaceutically acceptable salt thereof.

3. The compound of claim 1, wherein the compound has Formula (Ic):




or a pharmaceutically acceptable salt thereof.

4. The compound of claim 1, wherein the compound has Formula (Ie):



or a pharmaceutically acceptable salt thereof.

5. The compound of claim 1, wherein the compound has Formula (Ig):



or a pharmaceutically acceptable salt thereof.

6. The compound according to any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein R^1 is R^{2A} .

In some embodiments, R^{2A} is C_2 - C_4 alkenyl, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, or 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more R^A .

7. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is C_2 - C_4 alkenyl, unsubstituted or substituted with one or more R^A .
8. The compound according to claim 7, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is methylallyl, unsubstituted or substituted with one or more R^A .
9. The compound according to claim 7 or 8, or a pharmaceutically acceptable salt thereof, wherein R^A is halogen.
10. The compound according to any one of claims 7 to 9, or a pharmaceutically acceptable salt thereof, wherein R^A is fluoro.
11. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is prop-2-en-1-yl, unsubstituted or substituted with one or more methyl.
12. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is methylallyl.
13. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is C_1 - C_6 alkyl, unsubstituted or substituted with one or more R^A .
14. The compound according to claim 13, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is *n*-butyl, ethyl, 2-ethylbutyl, *n*-propyl, propan-2-yl, 2-methylpropyl, methyl, or *n*-pentyl, each unsubstituted or substituted with one or more R^A .
15. The compound according to claim 13 or 14, or a pharmaceutically acceptable salt thereof, wherein each R^A is independently selected from the group consisting of: C_1 - C_4 alkoxy, C_1 - C_4 alkylamino, C_1 - C_4 alkylsulfonyl, C_2 - C_4 dialkylamino, C_2 - C_4 dialkylsulfamoyl, halogen, halo C_1 - C_4 alkoxy, halo C_2 - C_4 dialkylamino, 4-7 membered heterocyclyl, -CN, -OH, and oxo.

16. The compound according to any one of claims 13 or 15, or a pharmaceutically acceptable salt thereof, wherein each R^A is independently selected from the group consisting of: methoxy, methylamino, methylsulfonyl, dimethylamino, *N,N*-dimethylsulfamoyl, chloro, fluoro, trifluoromethoxy, methyl(2,2,2-trifluoroethyl)amino, methyl(trifluoromethyl)amino, morpholino, -CN, -OH, and oxo.
17. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is *n*-butyl, ethyl, 2-ethylbutyl, *n*-propyl, propan-2-yl, 2-methylpropyl, methyl, or *n*-pentyl, each unsubstituted or substituted with one or more R^A independently selected from the group consisting of: C₁-C₄alkoxy, C₁-C₄alkylamino, C₁-C₄alkylsulfonyl, C₂-C₄dialkylamino, C₂-C₄dialkylsulfamoyl, halogen, haloC₁-C₄alkoxy, haloC₂-C₄dialkylamino, 4-7 membered heterocyclyl, -CN, -OH, and oxo.
18. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, ethyl, 2-fluoroethyl, 3-fluoropropyl, propan-2-yl, 1,1,1-trifluoropropan-2-yl, 2-hydroxy-2-methylpropyl, 3,3,3-trifluoro-2-hydroxypropyl, 3,3-difluoro-2-hydroxypropyl, methyl, 2-hydroxypropyl, 2-hydroxybutyl, 4,4,4-trifluoro-2-hydroxybutyl, 3,3,3-trifluoro-2-hydroxy-2-methylpropyl, 2,3-dihydroxypropyl, 3-hydroxypropyl, 2-hydroxyethyl, cyanomethyl, 2-(dimethylamino)-2-oxoethyl, 2-(methylamino)-2-oxoethyl, 2-hydroxy-3-methoxypropyl, 2-chloro-2,2-difluoroethyl, 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl, 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl, 3-(methylsulfonyl)propyl, (*N,N*-dimethylsulfamoyl)methyl, 2-methoxypropyl, 2-ethyl-2-hydroxybutyl, 3-chloro-3,3-difluoropropyl, 2-hydroxy-3-morpholinopropyl, 5,5,5-trifluoropentyl, or 4,4,4-trifluorobutyl.
19. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is C₃-C₈cycloalkyl, unsubstituted or substituted with one or more R^A .
20. The compound according to claim 19, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is cyclobutyl, cyclopropyl, cyclopentyl, or cyclohexyl, each unsubstituted or substituted with one or more R^A .
21. The compound according to claim 19 or 20, or a pharmaceutically acceptable salt thereof, wherein each R^A is independently selected from the group consisting of: halogen and haloC₁-C₄alkyl.

22. The compound according to any one of claims 19 to 21, or a pharmaceutically acceptable salt thereof, wherein each R^A is independently selected from the group consisting of: fluoro and trifluoromethyl.
23. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is cyclobutyl, cyclopropyl, cyclopentyl, or cyclohexyl, each unsubstituted or substituted with one or more R^A independently selected from the group consisting of: fluoro and trifluoromethyl.
24. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is cyclopropyl, cyclobutyl, 4-(trifluoromethyl)cyclohexyl, 2-fluorocyclopentyl, or 3,3-difluorocyclopentyl.
25. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is 4-7 membered heterocyclyl, unsubstituted or substituted with one or more R^A .
26. The compound according to claim 25, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is azetidin-3-yl, oxetan-3-yl, or pyrrolidin-3-yl, each unsubstituted or substituted with one or more R^A .
27. The compound according to claim 25 or 26, or a pharmaceutically acceptable salt thereof, wherein each R^A is independently selected from the group consisting of: C_1 - C_6 alkyl, halo C_1 - C_4 alkyl, and oxo.
28. The compound according to any one of claims 25 to 27, or a pharmaceutically acceptable salt thereof, wherein R^A is independently selected from the group consisting of: methyl, 2,2,2-trifluoroethyl, and oxo.
29. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is azetidin-3-yl, oxetan-3-yl, or pyrrolidin-3-yl, each unsubstituted or substituted with one or more R^A independently selected from the group consisting of: methyl, 2,2,2-trifluoroethyl, and oxo.
30. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl, or 1-methylazetidin-3-yl.

31. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein **R^{2A}** is methylallyl, *n*-butyl, ethyl, 2-ethylbutyl, *n*-propyl, propan-2-yl, 2-methylpropyl (isobutyl), methyl, *n*-pentyl, cyclobutyl, cyclopropyl, cyclopentyl, cyclohexyl, azetidin-3-yl, oxetan-3-yl, or pyrrolidin-3-yl; each unsubstituted or substituted with one or more **R^A** independently selected from the group consisting of: methoxy, methylamino, methylsulfonyl, dimethylamino, *N,N*-dimethylsulfamoyl, chloro, fluoro, trifluoromethoxy, methyl(2,2,2-trifluoroethyl)amino, methyl(trifluoromethyl)amino, morpholino, -CN, -OH, trifluoromethyl, methyl, 2,2,2-trifluoroethyl, and oxo.

32. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein **R^{2A}** is methylallyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, ethyl, 2-fluoroethyl, 3-fluoropropyl, propan-2-yl, 1,1,1-trifluoropropan-2-yl, 2-hydroxy-2-methylpropyl, 3,3,3-trifluoro-2-hydroxypropyl, 3,3-difluoro-2-hydroxypropyl, methyl, 2-hydroxypropyl, 2-hydroxybutyl, 4,4,4-trifluoro-2-hydroxybutyl, 3,3,3-trifluoro-2-hydroxy-2-methylpropyl, 2,3-dihydroxypropyl, 3-hydroxypropyl, 2-hydroxyethyl, cyanomethyl, 2-(dimethylamino)-2-oxoethyl, 2-(methylamino)-2-oxoethyl, 2-hydroxy-3-methoxypropyl, 2-chloro-2,2-difluoroethyl, 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl, 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl, 3-(methylsulfonyl)propyl, (*N,N*-dimethylsulfamoyl)methyl, 2-methoxypropyl, 2-ethyl-2-hydroxybutyl, 3-chloro-3,3-difluoropropyl, 2-hydroxy-3-morpholinopropyl, cyclopropyl, cyclobutyl, 4-(trifluoromethyl)cyclohexyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl, 1-methylazetidin-3-yl, 5,5,5-trifluoropentyl, or 4,4,4-trifluorobutyl.

33. The compound according to any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein **R¹** is $-(CH_2)_m-R^{2B}$.

34. The compound according to claim 33, or a pharmaceutically acceptable salt thereof, wherein **m** is 0.

35. The compound according to claim 33, or a pharmaceutically acceptable salt thereof, wherein **m** is 1.

36. The compound according to claim 33, or a pharmaceutically acceptable salt thereof, wherein **m** is 2.

37. The compound according to claim 33, or a pharmaceutically acceptable salt thereof, wherein **m** is 3.

38. The compound according to claim 33, or a pharmaceutically acceptable salt thereof, wherein **m** is 4.

39. The compound according to any one of claims 33 to 38, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is aryl, unsubstituted or substituted with one or more R^B .
40. The compound according to claim 39, wherein R^{2B} is phenyl, unsubstituted or substituted with one or more R^B .
41. The compound according to claim 39 or 40, or a pharmaceutically acceptable salt thereof, wherein each R^B is independently selected from the group consisting of: halogen, haloC₁-C₄alkyl, haloC₁-C₄alkylsulfonyl, and -CN.
42. The compound according to any one of claims 39 to 41, or a pharmaceutically acceptable salt thereof, wherein each R^B is independently selected from the group consisting of: fluoro, trifluoromethyl, (difluoromethyl)sulfonyl, and -CN.
43. The compound according to any one of claims 33 to 38, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is phenyl, unsubstituted or substituted with one or more R^B independently selected from the group consisting of: fluoro, trifluoromethyl, (difluoromethyl)sulfonyl, and -CN.
44. The compound according to any one of claims 33 to 38, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is phenyl, 4-fluorophenyl, 3-fluorophenyl, 4-(trifluoromethyl)phenyl, 4-((difluoromethyl)sulfonyl)phenyl, or 3-cyanophenyl.
45. The compound according to any one of claims 33 to 38, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is C₃-C₈cycloalkyl, unsubstituted or substituted with one or more R^B .
46. The compound according to claim 45, wherein R^{2B} is cyclobutyl, cyclopropyl, cyclohexyl, cyclopentyl, or bicyclo[1.1.1]pentan-1-yl, each unsubstituted or substituted with one or more R^B .
47. The compound according to claim 45 or 46, or a pharmaceutically acceptable salt thereof, wherein each R^B is independently selected from the group consisting of: C₁-C₄alkyl, C₁-C₄alkylsulfonyl, C₁-C₄alkylsulfonyloxy, -CN, -CH₂CN, halogen, haloC₁-C₄alkyl, and -OH.
48. The compound according to any one of claims 45 to 47, or a pharmaceutically acceptable salt thereof, wherein each R^B is independently selected from the group consisting of: ethyl,

methylsulfonyl, methylsulfonyloxy, -CN, -CH₂CN, fluoro, difluoromethyl, trifluoromethyl, and -OH.

49. The compound according to claim 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is cyclobutyl, cyclopropyl, cyclohexyl, cyclopentyl, or bicyclo[1.1.1]pentan-1-yl, each unsubstituted or substituted with one or more **R^B** independently selected from the group consisting of: ethyl, methylsulfonyl, methylsulfonyloxy, -CN, -CH₂CN, fluoro, difluoromethyl, trifluoromethyl, and -OH.
50. The compound according to claim 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is 1-hydroxycyclobutyl, cyclopropyl, 1-cyanocyclopropyl, 1-fluorocyclopropyl, cyclobutyl, 1-cyanocyclobutyl, 1-(cyanomethyl)cyclopropyl, 4,4-difluoro-1-hydroxycyclohexyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, 3-fluorobicyclo[1.1.1]pentan-1-yl, 1-(difluoromethyl)cyclopropyl, 3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl, 3-hydroxy-3-(trifluoromethyl)cyclobutyl, 1-(trifluoromethyl)cyclopentyl, 2-ethylcyclopropyl, 1-((methylsulfonyl)oxy)cyclopropyl, 1-(methylsulfonyl)cyclopropyl, or 2,2-difluorocyclopropyl.
51. The compound according to any one of claims 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is 5-10 membered heteroaryl, unsubstituted or substituted with one or more **R^B**.
52. The compound according to claim 51, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is 1,2,4-oxadiazol-5-yl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 1*H*-imidazol-2-yl, 1*H*-pyrazol-5-yl, 1*H*-tetrazol-5-yl, furan-2-yl, imidazol-1-yl, isoxazol-3-yl, isoxazol-5-yl, pyrazin-2-yl, pyridin-2-yl, pyridin-3-yl, pyrimidin-2-yl, or thiazol-4-yl, each unsubstituted or substituted with one or more **R^B**.
53. The compound according to claim 51 or 52, or a pharmaceutically acceptable salt thereof, wherein each **R^B** is independently selected from the group consisting of: C₁-C₄alkyl and -CN.
54. The compound according to any one of claims 51 to 53, or a pharmaceutically acceptable salt thereof, wherein each **R^B** is independently selected from the group consisting of: methyl and -CN.

55. The compound according to claim 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is 1,2,4-oxadiazol-5-yl, 1,3-oxazol-2-yl, 1,3-oxazol-4-yl, 1*H*-imidazol-2-yl, 1*H*-pyrazol-5-yl, 1*H*-tetrazol-5-yl, furan-2-yl, imidazol-1-yl, isoxazol-3-yl, isoxazol-5-yl, pyrazin-2-yl, pyridin-2-yl, pyridin-3-yl, pyrimidin-2-yl, or thiazol-4-yl, each unsubstituted or substituted with one or more **R^B** independently selected from the group consisting of: methyl and -CN.
56. The compound according to claim 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is imidazol-1-yl, 5-methyl-1,3-oxazol-2-yl, 5-cyanofuran-2-yl, isoxazol-3-yl, oxazol-4-yl, 6-cyanopyridin-2-yl, 5-cyanopyridin-2-yl, 1,2,4-oxadiazol-5-yl, pyridin-3-yl, pyrimidin-2-yl, isoxazol-5-yl, 1*H*-imidazol-2-yl, or 1*H*-tetrazol-5-yl.
57. The compound according to any one of claims 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is 4-7 membered heterocyclyl, unsubstituted or substituted with one or more **R^B**.
58. The compound according to claim 57, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is oxetan-3-yl, oxetan-2-yl, oxolan-3-yl (tetrahydrofuran-3-yl), pyrrolidin-1-yl, piperidin-4-yl, tetrahydro-2*H*-pyran-4-yl, azetidin-3-yl, 1,4-dioxepan-6-yl, or tetrahydro-2*H*-thiopyran-4-yl, unsubstituted or substituted with one or more **R^B**.
59. The compound according to claim 57 or 58, or a pharmaceutically acceptable salt thereof, wherein each **R^B** is independently selected from the group consisting of: C₁-C₄alkyl, halogen, oxo, and -CN.
60. The compound according to any one of claims 57 to 59, or a pharmaceutically acceptable salt thereof, wherein each **R^B** is independently selected from the group consisting of: methyl, fluoro, oxo, and -CN.
61. The compound according to claim 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is oxetan-3-yl, oxetan-2-yl, oxolan-3-yl (tetrahydrofuran-3-yl), pyrrolidin-1-yl, piperidin-4-yl, tetrahydro-2*H*-pyran-4-yl, azetidin-3-yl, 1,4-dioxepan-6-yl, or tetrahydro-2*H*-thiopyran-4-yl, each unsubstituted or substituted with one or more **R^B** independently selected from the group consisting of: methyl, fluoro, oxo, and -CN.
62. The compound according to claim 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is 3-fluorooxetan-3-yl, oxetan-2-yl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-

- yl), 3,3-difluoropyrrolidin-1-yl, 1-cyanopiperidin-4-yl, 1-methylazetidin-3-yl, 6-fluoro-1,4-dioxepan-6-yl, or 1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl.
63. The compound according to any one of claims 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is 1,2,4-oxadiazol-5-yl, 1,3-oxazol-2-yl, 1,4-dioxepan-6-yl, 1*H*-imidazol-2-yl, 1*H*-pyrazol-5-yl, 1*H*-tetrazol-5-yl, azetidin-3-yl, bicyclo[1.1.1]pentan-1-yl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopropyl, furan-2-yl, imidazol-1-yl, isoxazol-3-yl, isoxazol-5-yl, oxazol-4-yl, oxetan-2-yl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), phenyl, piperidin-4-yl, pyrazin-2-yl, pyridin-2-yl, pyridin-3-yl, pyrimidin-2-yl, pyrrolidin-1-yl, tetrahydro-2*H*-pyran-4-yl, tetrahydro-2*H*-thiopyran-4-yl, or thiazol-4-yl; each unsubstituted or substituted with one or more groups independently selected from the group consisting of: -CH₂CN, -CN, difluoromethyl, ethyl, fluoro, methyl, methylsulfonyl, methylsulfonyloxy, -OH, oxo, phenyl, and trifluoromethyl.
64. The compound according to any one of claims 33 to 38, or a pharmaceutically acceptable salt thereof, wherein **R^{2B}** is 1-((methylsulfonyl)oxy)cyclopropyl, 1-(cyanomethyl)cyclopropyl, 1-(difluoromethyl)cyclopropyl, 1-(methylsulfonyl)cyclopropyl, 1-(trifluoromethyl)cyclopentyl, 1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl, 1,2,4-oxadiazol-5-yl, 1-cyanocyclobutyl, 1-cyanocyclopropyl, 1-cyanopiperidin-4-yl, 1-fluorocyclobutyl, 1-fluorocyclopropyl, 1*H*-imidazol-2-yl, 1*H*-tetrazol-5-yl, 1-hydroxycyclobutyl, 1-methylazetidin-3-yl, 2-ethylcyclopropyl, 2-fluorocyclopentyl, 3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl, 3,3-difluorocyclopentyl, 3,3-difluoropyrrolidin-1-yl, 3-fluorobicyclo[1.1.1]pentan-1-yl, 3-fluorooxetan-3-yl, 3-fluorophenyl, 3-hydroxy-3-(trifluoromethyl)cyclobutyl, 4-((difluoromethyl)sulfonyl)phenyl, 4-(trifluoromethyl)phenyl, 4,4-difluoro-1-hydroxycyclohexyl, 4-fluorophenyl, 5-cyanofuran-2-yl, 5-cyanopyridin-2-yl, 5-methyl-1,3-oxazol-2-yl, 6-cyanopyridin-2-yl, 6-fluoro-1,4-dioxepan-6-yl, cyclobutyl, cyclopropyl, imidazol-1-yl, isoxazol-3-yl, isoxazol-5-yl, oxazol-4-yl, oxetan-2-yl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), phenyl, pyridin-3-yl, pyrimidin-2-yl, or 3-cyanophenyl.
65. The compound according to any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein **R¹** is -(CH₂)_n-OR^{2C}.
66. The compound according to claim 65, wherein **n** is 1.
67. The compound according to claim 65, wherein **n** is 2.
68. The compound according to claim 65, wherein **n** is 3.

In some embodiments, R^{2C} is C_1 - C_4 alkyl, C_3 - C_8 cycloalkyl, or 4-7 membered heterocycl, each unsubstituted or substituted with one or more R^C .

69. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is C_1 - C_4 alkyl, unsubstituted or substituted with one or more R^C .
70. The compound according to claim 69, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is ethyl, isopropyl, or methyl, each unsubstituted or substituted with one or more R^C .
71. The compound according to claim 69 or 70, or a pharmaceutically acceptable salt thereof, wherein each R^C is independently -CN or C_3 - C_6 cycloalkyl.
72. The compound according to any one of claims 69 to 71, or a pharmaceutically acceptable salt thereof, wherein each R^C is independently selected from the group consisting of: cyclopropyl and -CN.
73. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is ethyl, isopropyl, or methyl, each unsubstituted or substituted with one or more R^C independently selected from the group consisting of: cyclopropyl and -CN.
74. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is methyl, cyano(cyclopropyl)methyl, 2-cyanopropan-2-yl, cyanomethyl, or 1-cyanoethyl.
75. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is C_3 - C_8 cycloalkyl, unsubstituted or substituted with one or more R^C .
76. The compound according to claim 75, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is cyclobutyl, unsubstituted or substituted with one or more R^C .
77. The compound according to claim 75 or 76, wherein each R^C is -CN.
78. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is cyclobutyl, unsubstituted or substituted with one or more -CN.
79. The compound according to any one of claims 65 to 68, wherein R^{2C} is 1-cyanocyclobutyl.

80. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is 4-7 membered heterocyclyl, each unsubstituted or substituted with one or more R^C .
81. The compound according to claim 80, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is oxetan-3-yl or tetrahydro-2*H*-pyran-4-yl, each unsubstituted or substituted with one or more R^C .
82. The compound according to claim 80 or 81, or a pharmaceutically acceptable salt thereof, wherein R^C is -CN.
83. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is oxetan-3-yl or tetrahydro-2*H*-pyran-4-yl, each unsubstituted or substituted with one or more -CN.
84. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is oxetan-3-yl or 4-cyanotetrahydro-2*H*-pyran-4-yl.
85. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is ethyl, isopropyl, methyl, cyclobutyl, oxetan-3-yl, or tetrahydro-2*H*-pyran-4-yl, each unsubstituted or substituted with one or more groups independently selected from the group consisting of: cyclopropyl and -CN.
86. The compound according to any one of claims 65 to 68, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is methyl, cyano(cyclopropyl)methyl, 2-cyanopropan-2-yl, cyanomethyl, 1-cyanoethyl, 1-cyanocyclobutyl, oxetan-3-yl, or 4-cyanotetrahydro-2*H*-pyran-4-yl.
87. The compound according to any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein R^1 is methylallyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, ethyl, 2-fluoroethyl, 3-fluoropropyl, propan-2-yl, 1,1,1-trifluoropropan-2-yl, 2-hydroxy-2-methylpropyl, 3,3,3-trifluoro-2-hydroxypropyl, 3,3-difluoro-2-hydroxypropyl, methyl, 2-hydroxypropyl, 2-hydroxybutyl, 4,4,4-trifluoro-2-hydroxybutyl, 3,3,3-trifluoro-2-hydroxy-2-methylpropyl, 2,3-dihydroxypropyl, 3-hydroxypropyl, 2-hydroxyethyl, cyanomethyl, 2-(dimethylamino)-2-oxoethyl, 2-(methylamino)-2-oxoethyl, 2-hydroxy-3-methoxypropyl, 2-chloro-2,2-difluoroethyl, 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl, 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl, 3-

(methylsulfonyl)propyl, (*N,N*-dimethylsulfamoyl)methyl, 2-methoxypropyl, 2-ethyl-2-hydroxybutyl, 3-chloro-3,3-difluoropropyl, 2-hydroxy-3-morpholinopropyl, cyclopropyl, cyclobutyl, 4-(trifluoromethyl)cyclohexyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl, 1-methylazetidin-3-yl, (1-((methylsulfonyl)oxy)cyclopropyl)methyl, (1-(cyanomethyl)cyclopropyl)methyl, (1-(difluoromethyl)cyclopropyl)methyl, (1-(methylsulfonyl)cyclopropyl)methyl, (1-(trifluoromethyl)cyclopentyl)methyl, (1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl)methyl, (1,2,4-oxadiazol-5-yl)methyl, (1-cyanocyclobutyl)methyl, (1-cyanocyclopropyl)methyl, (1-cyanopiperidin-4-yl)methyl, (1-fluorocyclobutyl)methyl, (1-fluorocyclopropyl)methyl, (1*H*-imidazol-2-yl)methyl, (1*H*-tetrazol-5-yl)methyl, (1-hydroxycyclobutyl)methyl, (1-methylazetidin-3-yl)methyl, (2-ethylcyclopropyl)methyl, (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)methyl, (3-fluorobicyclo[1.1.1]pentan-1-yl)methyl, (3-fluorooxetan-3-yl)methyl, (3-hydroxy-3-(trifluoromethyl)cyclobutyl)methyl, (4,4-difluoro-1-hydroxycyclohexyl)methyl, (5-cyanofuran-2-yl)methyl, (5-cyanopyridin-2-yl)methyl, (5-methyl-1,3-oxazol-2-yl)methyl, (6-cyanopyridin-2-yl)methyl, (6-fluoro-1,4-dioxepan-6-yl)methyl, (isoxazol-3-yl)methyl, (isoxazol-5-yl)methyl, (oxazol-4-yl)methyl, (oxetan-2-yl)methyl, (oxetan-3-yl)methyl, (pyridin-3-yl)methyl, 2-(1-cyanocyclopropyl)ethyl, 2-(1*H*-imidazol-1-yl)ethyl, 2-(3,3-difluoropyrrolidin-1-yl)ethyl, 3-fluorobenzyl, 4-((difluoromethyl)sulfonyl)benzyl, 4-(trifluoromethyl)benzyl, 4-(trifluoromethyl)phenethyl, 4-fluorobenzyl, benzyl, cyclopropylmethyl, pyrimidin-2-ylmethyl, (2-cyanopropan-2-yl)oxy)ethyl, 2-((4-cyanotetrahydro-2*H*-pyran-4-yl)oxy)ethyl, 2-(1-cyanocyclobutoxy)ethyl, 2-(1-cyanoethoxy)ethyl, 2-(cyano(cyclopropyl)methoxy)ethyl, 2-(cyanomethoxy)ethyl, 2-(oxetan-3-yloxy)ethyl, 2-methoxyethyl, 5,5,5-trifluoropentyl, 3-cyanobenzyl, 4-fluorophenethyl, cyclobutylmethyl, 4,4,4-trifluorobutyl, or (2,2-difluorocyclopropyl)methyl.

88. The compound according to any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein **R¹** is methylallyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, ethyl, 2-fluoroethyl, 3-fluoropropyl, propan-2-yl, 1,1,1-trifluoropropan-2-yl, 2-hydroxy-2-methylpropyl, 3,3,3-trifluoro-2-hydroxypropyl, 3,3-difluoro-2-hydroxypropyl, methyl, 2-hydroxypropyl, 2-hydroxybutyl, 4,4,4-trifluoro-2-hydroxybutyl, 3,3,3-trifluoro-2-hydroxy-2-methylpropyl, 2,3-dihydroxypropyl, 3-hydroxypropyl, 2-hydroxyethyl, cyanomethyl, 2-(dimethylamino)-2-oxoethyl, 2-(methylamino)-2-oxoethyl, 2-hydroxy-3-methoxypropyl, 2-chloro-2,2-difluoroethyl, 2-hydroxy-3-(2,2,2-trifluoroethoxy)propyl, 2-(methyl(2,2,2-trifluoroethyl)amino)-2-oxoethyl, 3-(methylsulfonyl)propyl, (*N,N*-dimethylsulfamoyl)methyl, 2-methoxypropyl, 2-ethyl-2-hydroxybutyl, 3-chloro-3,3-difluoropropyl, 2-hydroxy-3-morpholinopropyl, cyclopropyl,

cyclobutyl, 4-(trifluoromethyl)cyclohexyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), 2-oxo-1-(2,2,2-trifluoroethyl)pyrrolidin-3-yl, 1-methylazetidin-3-yl, 5,5,5-trifluoropentyl, or 4,4,4-trifluorobutyl.

89. The compound according to any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein **R¹** is (1-((methylsulfonyl)oxy)cyclopropyl)methyl, (1-(cyanomethyl)cyclopropyl)methyl, (1-(difluoromethyl)cyclopropyl)methyl, (1-(methylsulfonyl)cyclopropyl)methyl, (1-(trifluoromethyl)cyclopentyl)methyl, (1,1-dioxidotetrahydro-2*H*-thiopyran-4-yl)methyl, (1,2,4-oxadiazol-5-yl)methyl, (1-cyanocyclobutyl)methyl, (1-cyanocyclopropyl)methyl, (1-cyanopiperidin-4-yl)methyl, (1-fluorocyclobutyl)methyl, (1-fluorocyclopropyl)methyl, (1*H*-imidazol-2-yl)methyl, (1*H*-tetrazol-5-yl)methyl, (1-hydroxycyclobutyl)methyl, (1-methylazetidin-3-yl)methyl, (2-ethylcyclopropyl)methyl, (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)methyl, (3-fluorobicyclo[1.1.1]pentan-1-yl)methyl, (3-fluorooxetan-3-yl)methyl, (3-hydroxy-3-(trifluoromethyl)cyclobutyl)methyl, (4,4-difluoro-1-hydroxycyclohexyl)methyl, (5-cyanofuran-2-yl)methyl, (5-cyanopyridin-2-yl)methyl, (5-methyl-1,3-oxazol-2-yl)methyl, (6-cyanopyridin-2-yl)methyl, (6-fluoro-1,4-dioxepan-6-yl)methyl, (isoxazol-3-yl)methyl, (isoxazol-5-yl)methyl, (oxazol-4-yl)methyl, (oxetan-2-yl)methyl, (oxetan-3-yl)methyl, (pyridin-3-yl)methyl, 2-(1-cyanocyclopropyl)ethyl, 2-(1*H*-imidazol-1-yl)ethyl, 2-(3,3-difluoropyrrolidin-1-yl)ethyl, 2-fluorocyclopentyl, 3,3-difluorocyclopentyl, 3-fluorobenzyl, 4-((difluoromethyl)sulfonyl)benzyl, 4-(trifluoromethyl)benzyl, 4-(trifluoromethyl)phenethyl, 4-fluorobenzyl, benzyl, cyclobutyl, cyclopropyl, cyclopropylmethyl, oxetan-3-yl, oxolan-3-yl (tetrahydrofuran-3-yl), pyrimidin-2-ylmethyl, 3-cyanobenzyl, 4-fluorophenethyl, cyclobutylmethyl, or (2,2-difluorocyclopropyl)methyl.
90. The compound according to any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein **R¹** is (2-cyanopropan-2-yl)oxy)ethyl, 2-((4-cyanotetrahydro-2*H*-pyran-4-yl)oxy)ethyl, 2-(1-cyanocyclobutoxy)ethyl, 2-(1-cyanoethoxy)ethyl, 2-(cyano(cyclopropyl)methoxy)ethyl, 2-(cyanomethoxy)ethyl, 2-(oxetan-3-yl)oxy)ethyl, or 2-methoxyethyl.
91. The compound according to claim 1, selected from the following compounds:
 (2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-1**);
 (2*R*,3*S*,11*bS*)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-2**);

(2*R*,3*S*,11*bS*)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-3**);

(2*R*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-4**);

(2*R*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-5**);

(2*S*,3*S*,11*bS*)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-6**);

(2*S*,3*S*,11*bS*)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-7**);

(2*S*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-8**);

(2*S*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-9**);

(2*S*,3*R*,11*bR*)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-10**);

(2*S*,3*R*,11*bR*)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-11**);

(2*S*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-12**);

(2*R*,3*R*,11*bR*)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-13**);

(2*R*,3*R*,11*bR*)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-14**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-15**);

(2*S*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-16**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-ethoxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-17**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(2-fluoroethoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-18**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(3-fluorooxetan-3-yl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-19**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[2-(oxetan-3-yloxy)ethoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-20**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*S*)-oxetan-2-ylmethoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-21**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-22**);

1-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)cyclobutane-1-carbonitrile (**Compound 4-23**);

2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}acetonitrile (**Compound 4-24**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-hydroxycyclobutyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-25**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2-methoxyethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-26**);

(2*R*,3*R*,11*bR*)-9-cyclopropoxy-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-27**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(3-fluoropropoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-28**);

1-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)cyclopropane-1-carbonitrile (**Compound 4-29**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-yl)oxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-30**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclopropyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-31**);

(2*R*,3*R*,11*bR*)-9-cyclobutoxy-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-32**);

2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}-*N,N*-dimethylacetamide (**Compound 4-33**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[2-(1*H*-imidazol-1-yl)ethoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-34**);

2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}-*N*-methylacetamide (**Compound 4-35**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(5-methyl-1,3-oxazol-2-yl)methoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-36**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1-methylazetidin-3-yl)methoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-37**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(propan-2-yloxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-38**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1,1,1-trifluoropropan-2-yl)oxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-39**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(ethoxy-*d*5)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-40**);

4-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)oxane-4-carbonitrile (**Compound 4-41**);

1-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)cyclobutane-1-carbonitrile (**Compound 4-42**);

2-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)-2-cyclopropylacetonitrile (**Compound 4-43**);

2-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)-2-methylpropanenitrile (**Compound 4-44**);

1-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethyl)cyclopropane-1-carbonitrile (**Compound 4-45**);

2-[1-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)cyclopropyl]acetonitrile (**Compound 4-46**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(2-hydroxy-2-methylpropoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-47**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*S*)-2-hydroxy-3-methoxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-48**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*R*)-3,3,3-trifluoro-2-hydroxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-49**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*S*)-3,3,3-trifluoro-2-hydroxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-50**);

(2*R*,3*R*,11*bR*)-9-(3,3-difluoro-2-hydroxypropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-51**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(methoxy-*d*3)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-52**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3*R*)-oxolan-3-yloxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-53**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3*S*)-oxolan-3-yloxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-54**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-55**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2-methylprop-2-en-1-yl)oxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-56**);

(2*R*,3*R*,11*bR*)-9-{[1-(difluoromethyl)cyclopropyl]methoxy}-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-57**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxybutoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-58**);

5-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)furan-2-carbonitrile (**Compound 4-59**);

(2*R*,3*R*,11*bR*)-9-(benzyloxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-60**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclobutyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-61**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(4,4,4-trifluoro-2-hydroxybutoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-62**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(4-fluorophenyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-63**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(3-fluorophenyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-64**);

(2*R*,3*R*,11*bR*)-9-(2-chloro-2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-65**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoro-2-hydroxy-2-methylpropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-66**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl]methoxy}-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-67**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*S*)-2-hydroxybutoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-68**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2-oxazol-3-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-69**);

2-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)acetonitrile (**Compound 4-70**);

(2*R*,3*R*,11*bR*)-9-[2-(3,3-difluoropyrrolidin-1-yl)ethoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-71**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(6-fluoro-1,4-dioxepan-6-yl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-72**);

4-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)piperidine-1-carbonitrile (**Compound 4-73**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-{2-[4-(trifluoromethyl)phenyl]ethoxy}-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-74**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,3-oxazol-4-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-75**);

(2*R*,3*R*,11*bR*)-9-[(4,4-difluoro-1-hydroxycyclohexyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-76**);

6-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)pyridine-2-carbonitrile (**Compound 4-77**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-{[3-hydroxy-3-(trifluoromethyl)cyclobutyl]methoxy}-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-78**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[2-hydroxy-3-(2,2,2-trifluoroethoxy)propoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-79**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[1-(trifluoromethyl)cyclopentyl]methoxy}-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-80**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[4-(trifluoromethyl)phenyl]methoxy}-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-81**);

6-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)pyridine-3-carbonitrile (**Compound 4-82**);

(2*R*)-3-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}propane-1,2-diol (**Compound 4-83**);

2-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}-*N*-methyl-*N*-(2,2,2-trifluoroethyl)acetamide (**Compound 4-84**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(3-methanesulfonylpropoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-85**);

1-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}-*N,N*-dimethylmethanesulfonamide (**Compound 4-86**);

3-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}-1-(2,2,2-trifluoroethyl)pyrrolidin-2-one (**Compound 4-87**);

(2*R*,3*R*,11*bR*)-9-(cyclopropylmethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-88**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(3-hydroxypropoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-89**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-({[(1*R*,2*R*)-2-ethylcyclopropyl]methoxy}-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-90**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*R*)-2-methoxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-91**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*S*)-2-methoxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-92**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(2-ethyl-2-hydroxybutoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-93**);

1-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)cyclopropyl methanesulfonate (**Compound 4-94**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2,4-oxadiazol-5-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-95**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-methanesulfonylcyclopropyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-96**);

4-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)-1*λ*⁶-thiane-1,1-dione (**Compound 4-97**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(2-hydroxyethoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-98**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-{[4-(trifluoromethyl)cyclohexyl]oxy}-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-99**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2-fluorocyclopentyl)oxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-100**);

(2*R*,3*R*,11*bR*)-9-[(3,3-difluorocyclopentyl)oxy]-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-101**);

(2*R*,3*R*,11*bR*)-9-[(4-difluoromethanesulfonylphenyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-102**);

(2*R*,3*R*,11*bR*)-9-(3-chloro-3,3-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-103**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1-methylazetidin-3-yl)oxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-104**);

2-(2-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)propanenitrile (**Compound 4-105**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-({3-fluorobicyclo[1.1.1]pentan-1-yl}methoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-106**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(pyridin-3-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-107**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(pyrimidin-2-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-108**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1,2-oxazol-5-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-109**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(1*H*-imidazol-2-ylmethoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-110**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(1*H*-1,2,3,4-tetrazol-5-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-111**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxy-3-methoxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-112**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[2-hydroxy-3-(morpholin-4-yl)propoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-113**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*S*)-2-hydroxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-114**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(5,5,5-trifluoropentyl)oxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-115**);

3-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)benzonitrile (**Compound 4-116**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[2-(4-fluorophenyl)ethoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-117**);

(2*S*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclopropyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-118**);

(2*R*,3*R*,11*bR*)-9-(cyclobutylmethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-119**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(4,4,4-trifluorobutoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-120**);

(2*R*,3*R*,11*bR*)-9-[(2,2-difluorocyclopropyl)methoxy]-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-121**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-1**);

(2*S*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-2**);

(2*S*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-3**); and

(2*R*,3*S*,11*bS*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-4**);

or a pharmaceutically acceptable salt thereof.

92. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein:

R^1 is R^{2A} , $-(CH_2)_m-R^{2B}$, or $-(CH_2)_n-OR^{2C}$;

R^{2A} is C_1 - C_4 alkyl, unsubstituted or substituted with one or more R^A ;

each R^A is independently selected from the group consisting of: fluoro and -OH;

R^{2B} is C₃-C₈cycloalkyl or 4-6 membered heterocyclyl, each unsubstituted or substituted with one or more R^B ;

each R^B is independently selected from the group consisting of: fluoro, -CN, -CH₂CN, and -OH;


R^{2C} is C₁-C₄alkyl or C₃-C₈cycloalkyl, each unsubstituted or substituted with one or more R^C ;

each R^C is independently -CN or C₃-C₆cycloalkyl;

m is 0, 1, 2, or 3; and

n is 1, 2, or 3.

93. The compound according to claim 92, wherein the compound has Formula (Ia):



or a pharmaceutically acceptable salt thereof.

94. The compound according to claim 92 or 93, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is C₁-C₄alkyl, unsubstituted or substituted with one two, three, or four R^A ; and each R^A is independently selected from the group consisting of: fluoro and -OH.
95. The compound according to claim 94, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is C₁-C₄alkyl, unsubstituted or substituted with one, two, or three R^A ; and each R^A is independently selected from the group consisting of: fluoro and -OH.
96. The compound according to claim 94, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is C₂-C₄alkyl.
97. The compound according to claim 94, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is -CH₂CH₃, -CH₂CH₂CH₃, or -CH(CH₃)₂.
98. The compound according to claim 94, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is C₁-C₄alkyl substituted with one, two, or three fluoro.

99. The compound according to claim 94, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2CH_2CHF_2$, or $-CH_2CH_2CF_3$.
100. The compound according to claim 94, or a pharmaceutically acceptable salt thereof, wherein R^{2A} is $-CH_2CH_2CHF_2$ or $-CH_2CH_2CF_3$.
101. The compound according to claim 92 or 93, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is C_3 - C_6 cycloalkyl, unsubstituted or substituted with one or two R^B ; and each R^B is independently selected from the group consisting of: fluoro, CN, $-CH_2CN$, and $-OH$.
102. The compound according to claim 101, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is cyclopropyl, cyclobutyl, or cyclopentyl, each unsubstituted or substituted with one or two fluoro.
103. The compound according to claim 92 or 93, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is a 4-5 membered monocyclic heterocyclyl having one ring forming heteroatom selected from oxygen and sulfur, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or more R^B ; and each R^B is independently selected from the group consisting of: fluoro and $-OH$.
104. The compound according to claim 103, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is a C_3 - C_4 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two R^B ; and each R^B is independently selected from the group consisting of: fluoro and $-OH$.
105. The compound according to claim 104, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is a 4-5 membered monocyclic heterocyclyl having one ring forming oxygen, wherein the 4-5 membered monocyclic heterocyclyl is unsubstituted or substituted with one or two fluoro.
106. The compound according to claim 105, or a pharmaceutically acceptable salt thereof, wherein R^{2B} is oxetan-3-yl or oxolan-3-yl, each unsubstituted or substituted with one fluoro.
107. The compound according to claim 92 or 93, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is C_1 - C_4 alkyl or C_3 - C_5 cycloalkyl, each substituted with one, two, three, or four R^C ; and each R^C is independently $-CN$ or C_3 - C_6 cycloalkyl.

108. The compound according to claim 107, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is cyclobutyl substituted with one -CN.
109. The compound according to claim 107, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is C_1 - C_3 alkyl substituted with one -CN and one C_3 - C_5 cycloalkyl.
110. The compound according to claim 109, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is C_1 - C_2 alkyl substituted with one -CN and one C_3 - C_4 cycloalkyl.
111. The compound according to claim 110, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is C_1 - C_2 alkyl substituted with one -CN and one cyclopropyl.
112. The compound according to claim 109, or a pharmaceutically acceptable salt thereof, wherein R^{2C} is C_1 - C_3 alkyl substituted with one -CN.
113. The compound according to any one of claims 92, 93, and 101 to 106, or a pharmaceutically acceptable salt thereof, wherein m is 0, 1, or 2.
114. The compound according to any one of claims 92, 93, and 101 to 106, or a pharmaceutically acceptable salt thereof, wherein m is 0 or 1.
115. The compound according to any one of claims 92, 93, and 101 to 106, or a pharmaceutically acceptable salt thereof, wherein m is 1.
116. The compound according to any one of claims 92, 93, and 101 to 106, or a pharmaceutically acceptable salt thereof, wherein m is 0.
117. The compound according to any one of claims 92, 93, and 107-112, or a pharmaceutically acceptable salt thereof, wherein n is 1 or 2.
118. The compound according to any one of claims 92, 93, and 107-112, or a pharmaceutically acceptable salt thereof, wherein n is 1.
119. The compound according to claim 92, selected from the following compounds:
(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-1**);

(2*R*,3*R*,11*bR*)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-13**);

(2*R*,3*R*,11*bR*)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-14**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-15**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*S*)-3,3,3-trifluoro-2-hydroxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-50**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*R*)-3,3,3-trifluoro-2-hydroxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-49**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(2-fluoroethoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-18**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1,1,1-trifluoropropan-2-yl)oxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-39**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-1**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-ethoxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-17**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(propan-2-yloxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-38**);

(2*R*,3*R*,11*bR*)-9-(3,3-difluoro-2-hydroxypropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-51**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-55**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(2-hydroxy-2-methylpropoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-47**);

1-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethyl)cyclopropane-1-carbonitrile (**Compound 4-45**);

2-[1-({[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)cyclopropyl]acetonitrile (**Compound 4-46**);

(2*R*,3*R*,11*bR*)-9-cyclopropoxy-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-27**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-hydroxycyclobutyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-25**);

(2*R*,3*R*,11*bR*)-9-cyclobutoxy-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-32**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-yloxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-30**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-ylmethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-22**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(3-fluorooxetan-3-yl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-19**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3*R*)-oxolan-3-yloxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-53**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3*S*)-oxolan-3-yloxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-54**);

1-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)cyclobutane-1-carbonitrile (**Compound 4-23**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclopropyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-31**);

2-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)-2-cyclopropylacetonitrile (**Compound 4-43**);

2-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)-2-methylpropanenitrile (**Compound 4-44**); and

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2-methoxyethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-26**);

or a pharmaceutically acceptable salt thereof.

120. The compound according to claim 92, selected from the following compounds:

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2,2,2-trifluoroethoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-1**);

(2*R*,3*R*,11*bR*)-9-(2,2-difluoroethoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-13**);

(2*R*,3*R*,11*bR*)-9-(2,2-difluoropropoxy)-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-14**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(3,3,3-trifluoropropoxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-15**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*S*)-3,3,3-trifluoro-2-hydroxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-50**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(2*R*)-3,3,3-trifluoro-2-hydroxypropoxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-49**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(2-fluoroethoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-18**); and

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(1,1,1-trifluoropropan-2-yl)oxy]-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-39**);

or a pharmaceutically acceptable salt thereof.

121. The compound according to claim 92, selected from the following compounds:

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9,10-dimethoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 5-1**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-ethoxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-17**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(propan-2-yloxy)-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-38**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(2*R*)-2-hydroxypropoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-55**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-(2-hydroxy-2-methylpropoxy)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-47**);

1-(2-([(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethyl)cyclopropane-1-carbonitrile (**Compound 4-45**);

2-[1-([(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}methyl)cyclopropyl]acetonitrile (**Compound 4-46**);

(2*R*,3*R*,11*bR*)-9-cyclopropoxy-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-27**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-hydroxycyclobutyl)methoxy]-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-25**); and

(2*R*,3*R*,11*bR*)-9-cyclobutoxy-3-(2,2-dimethylpropyl)-10-methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-32**);

or a pharmaceutically acceptable salt thereof.

122. The compound according to claim 92, selected from the following compounds:

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-yloxy)-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-30**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(oxetan-3-ylmethoxy)-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-22**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(3-fluorooxetan-3-yl)methoxy]-10-methoxy-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-19**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3*R*)-oxolan-3-yloxy]-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-53**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-[(3*S*)-oxolan-3-yloxy]-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-54**);

1-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)cyclobutane-1-carbonitrile (**Compound 4-23**);

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-9-[(1-fluorocyclopropyl)methoxy]-10-

methoxy-1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-31**);

2-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)-2-cyclopropylacetonitrile (**Compound 4-43**);

2-(2-{[(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-2-hydroxy-10-methoxy-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-9-yl]oxy}ethoxy)-2-methylpropanenitrile (**Compound 4-44**); and

(2*R*,3*R*,11*bR*)-3-(2,2-dimethylpropyl)-10-methoxy-9-(2-methoxyethoxy)-

1*H*,2*H*,3*H*,4*H*,6*H*,7*H*,11*bH*-pyrido[2,1-*a*]isoquinolin-2-ol (**Compound 4-26**);

or a pharmaceutically acceptable salt thereof.

123. A pharmaceutical product selected from: a pharmaceutical composition, a formulation, a unit dosage form, and a kit; each comprising the compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof.

124. A pharmaceutical product selected from: a pharmaceutical composition, a formulation, a unit dosage form, and a kit; each comprising the compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient.

125. A pharmaceutical composition comprising the compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient.
126. A method of treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder in a subject in need thereof, comprising administering to the subject a compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125.
127. A method of treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder in a subject in need thereof, comprising administering to the subject a compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; wherein the VMAT2 disease or disorder is selected from: an ataxias or spinal muscular atrophy; a chorea; a congenital malformation, deformation, or abnormality; a dementia; an oral cavity, salivary gland, or jaw disease; a dyskinesia; a dystonia; an endocrine, nutritional, or metabolic disease; an epilepsy; a habit or impulse disorder; a Huntington's disease or related disorder; a mood or psychotic disorder; a neurotic, stress-related, and somatoform disorder; a degenerative disease of the basal ganglia; an extrapyramidal and movement disorder; a neurological or psychiatric disease or disorder; a nervous system or motor function disorder; a Parkinson's/parkinsonism disorder; a pediatric-onset behavioral and emotional disorder; a pervasive developmental disorder; and a substance abuse or dependence disorder.
128. A method of treating a neurological or psychiatric disease or disorder in a subject in need thereof, comprising administering to the subject a compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125.
129. A method of treating a neurological or psychiatric disease or disorder in a subject in need thereof, comprising administering to the subject a compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; wherein the neurological or psychiatric disease or disorder is selected from the group consisting of: hyperkinetic movement disorder, schizophrenia, schizoaffective disorder, a mood disorder, treatment-refractory obsessive-compulsive disorder, neurological dysfunction associated with Lesch-Nyhan syndrome, agitation associated with Alzheimer's disease, Fragile X syndrome

- or Fragile X-associated tremor-ataxia syndrome, autism spectrum disorder, Rett syndrome, and chorea-acanthocytosis.
130. Use of a compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; in the manufacture of a medicament for treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder.
131. Use of a compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; in the manufacture of a medicament for treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder selected from: an ataxias or spinal muscular atrophy; a chorea; a congenital malformation, deformation, or abnormality; a dementia; an oral cavity, salivary gland, or jaw disease; a dyskinesia; a dystonia; an endocrine, nutritional, or metabolic disease; an epilepsy; a habit or impulse disorder; a Huntington's disease or related disorder; a mood or psychotic disorder; a neurotic, stress-related, and somatoform disorder; a degenerative disease of the basal ganglia; an extrapyramidal and movement disorder; a neurological or psychiatric disease or disorder; a nervous system or motor function disorder; a Parkinson's/parkinsonism disorder; a pediatric-onset behavioral and emotional disorder; a pervasive developmental disorder; and a substance abuse or dependence disorder.
132. Use of a compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; in the manufacture of a medicament for treating a neurological or psychiatric disease or disorder.
133. Use of a compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; in the manufacture of a medicament for treating a neurological or psychiatric disease or disorder selected from the group consisting of: hyperkinetic movement disorder, schizophrenia, schizoaffective disorder, a mood disorder, treatment-refractory obsessive-compulsive disorder, neurological dysfunction associated with Lesch-Nyhan syndrome, agitation associated with Alzheimer's disease, Fragile X syndrome or Fragile X-associated tremor-ataxia syndrome, autism spectrum disorder, Rett syndrome, and chorea-acanthocytosis.

134. A compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; for use in a method of treatment of the human or animal body by therapy.
135. A compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; for use in a method for treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder.
136. A compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; for use in a method for treating a vesicular monoamine transporter-2 (VMAT2) disease or disorder selected from: an ataxias or spinal muscular atrophy; a chorea; a congenital malformation, deformation, or abnormality; a dementia; an oral cavity, salivary gland, or jaw disease; a dyskinesia; a dystonia; an endocrine, nutritional, or metabolic disease; an epilepsy; a habit or impulse disorder; a Huntington's disease or related disorder; a mood or psychotic disorder; a neurotic, stress-related, and somatoform disorder; a degenerative disease of the basal ganglia; an extrapyramidal and movement disorder; a neurological or psychiatric disease or disorder; a nervous system or motor function disorder; a Parkinson's/parkinsonism disorder; a pediatric-onset behavioral and emotional disorder; a pervasive developmental disorder; and a substance abuse or dependence disorder.
137. A compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; for use in a method for treating a neurological or psychiatric disease or disorder.
138. A compound according to any one of claims 1 to 122, or a pharmaceutically acceptable salt thereof; a pharmaceutical product according to claim 123 or 124; or a pharmaceutical composition according to claim 125; for use in a method for treating a neurological or psychiatric disease or disorder selected from the group consisting of: hyperkinetic movement disorder, schizophrenia, schizoaffective disorder, a mood disorder, treatment-refractory obsessive-compulsive disorder, neurological dysfunction associated with Lesch-Nyhan syndrome, agitation associated with Alzheimer's disease, Fragile X syndrome or Fragile X-associated tremor-ataxia syndrome, autism spectrum disorder, Rett syndrome, and chorea-acanthocytosis.



Representation of the crystal structure of Compound 2-B (3*S*,11*bS*)

FIG. 1A



Representation of the crystal structure of Compound 1-D (2*R*,3*R*,11*bR*)

FIG. 1B

PXRD Diffractogram of Compound 4-1, Form I




FIG. 2

DSC and TGA Thermogram of Compound 4-1, Form I




FIG. 3

PXRD Diffractogram of Compound 4-15, Form I




FIG. 4

DSC and TGA Thermogram of Compound 4-15, Form I




FIG. 5

General synthetic scheme for preparing Compounds of Formula (I)




FIG. 6

Example 1A: Preparation of Compound 1-D (2R,3R,11bR)





FIG. 7

Example 1B: Preparation of Compound 1-C (2*R*,3*R*,11*bR*)•DPTTA Salt**FIG. 8**

Example 2: Compound 2-B (3*R*,11*bR*) and Compound 2-B (3*S*,11*bS*)









FIG. 9

Example 3: Preparation of Compound 1-D (2*R*,3*R*,11*bR*)**Example 3: Preparation of Compound 1-D (2*S*,3*S*,11*bS*)****FIG. 10**

Example 4: Preparation of Compound 1-D (2*S*,3*R*,11*bR*)**Example 4: Preparation of Compound 1-D (2*R*,3*S*,11*bS*)****FIG. 11**

Example 5A: Preparation of Compound 4-1



Example 5F: Preparation of Compound 4-55








FIG. 12

Example 7: Preparation of Compound 5-1



Example 8: Preparation of Compound 5-3

**FIG. 13**

Example 9: Preparation of Compound 5-2**Example 10: Preparation of Compound 5-4****FIG. 14**

VMAT2 Ki values (nM) comparison




Cmpd of Formula (I) vs Comparator Cmpd	Cmpd No.	Z group	VMAT Ki (nM)	Potency Improvement
	4-1		2.4	6.7x
	4-1C		16	
	4-13		2.2	12.3x
	4-13C		27	
	4-14		2.6	5.8x
	4-14C		15	

FIG. 15

VMAT2 Ki values (nM) comparison




Cmpd of Formula (I) vs Comparator Cmpd	Cmpd No.	Z group	VMAT Ki (nM)	Potency Improvement
	4-15		1.1	10.0x
	4-15C		11	
	4-17		1.1	8.1x
	4-17C		8.9	
	4-18		5.5	5.3x
	4-18C		29	

FIG. 16

VMAT2 Ki values (nM) comparison




Cmpd of Formula (I) vs Comparator Cmpd	Cmpd No.	Z group	VMAT Ki (nM)	Potency Improvement
	4-22		14	6.6x
	4-22C		93	
	4-26		14	12.1x
	4-26C		169	
	4-28		0.48	7.1x
	4-28C		3.4	

FIG. 17

VMAT2 Ki values (nM) comparison







Compd of Formula (I) vs Comparator Compd	Compd No.	Z group	VMAT Ki (nM)	Potency Improvement
	4-30		12	20.9x
	4-30C		251	
	4-32		0.59	5.8x
	4-32C		3.4	
	4-38		3.7	7.0x
	4-38C		26	

FIG. 18

VMAT2 Ki values (nM) comparison



Cmpd of Formula (I) vs Comparator Cmpd	Cmpd No.	Z group	VMAT Ki (nM)	Potency Improvement
	4-39		12	15.1x
	4-39C		181	
	5-1		6.6	4.4x
	5-1C		29	

FIG. 19

VMAT2 Ki values (nM) comparison







Cmpd of Formula (I) vs Comparator Cmpd	Cmpd No.	Z group	VMAT Ki (nM)	Potency Improvement
	4-27		1.0	4.4x
	4-27C		4.4	
	4-88		0.68	4.1x
	4-88C		2.8	
	4-119		0.29	9.7x
	4-119C		2.8	

FIG. 20

VMAT2 Ki values (nM) comparison





Compd of Formula (I) vs Comparator Compd	Compd No.	Z group	VMAT Ki (nM)	Potency Improvement
	4-120		1.5	13.3x
	4-120C		20	
	4-121		0.82	6.0x
	4-121C		4.9	

FIG. 21




FIG. 22





FIG. 23


**FIG. 24**


INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2022/021 080

Box No. IV Text of the abstract (Continuation of item 5 of the first sheet)

This disclosure relates generally to VMAT2 inhibitor compounds of formula (I), compositions, and methods related thereto.





(I)

(I)