





(86) Date de dépôt PCT/PCT Filing Date: 2000/03/30
(87) Date publication PCT/PCT Publication Date: 2000/10/19
(45) Date de délivrance/Issue Date: 2005/01/11
(85) Entrée phase nationale/National Entry: 2001/10/05
(86) N° demande PCT/PCT Application No.: US 2000/008337
(87) N° publication PCT/PCT Publication No.: 2000/061136
(30) Priorités/Priorities: 1999/04/08 (60/128,400) US;
2000/03/30 (09/538,182) US

(51) Cl.Int.⁷/Int.Cl.⁷ A61K 31/255, A61K 31/7048,
A61K 31/35, A61P 9/12
(72) Inventeur/Inventor:
COTTRELL, SANDRA C., US
(73) Propriétaire/Owner:
ORTHO-MCNEIL PHARMACEUTICAL, INC., US
(74) Agent: OGILVY RENAULT

(54) Titre : DERIVES ANTICONVULSIVANTS UTILES POUR ABAISSER LA PRESSION SANGUINE
(54) Title: ANTICONVULSANT DERIVATIVES USEFUL IN LOWERING BLOOD PRESSURE



(57) **Abrégé/Abstract:**


Use of anticonvulsant derivatives of formula (I) for lowering blood pressure particularly in overweight individuals, wherein X is CH₂ or oxygen; R₁ is hydrogen or alkyl; and R₂, R₃, R₄ and R₅ are independently hydrogen or alkyl and, when X is CH₂, R₄ and R₅ may be alkene groups joined to form a benzene ring and, when X is oxygen, R₂ and R₃, and/or R₄ and R₅ together may be a methylenedioxy group of formula (II), wherein R₆ and R₇ are the same or different and are hydrogen, or alkyl and are joined to form a cyclopentyl or cyclohexyl ring.




**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/255, 31/35, 31/7048, A61P 9/12	A1	(11) International Publication Number: WO 00/61136 (43) International Publication Date: 19 October 2000 (19.10.00)
(21) International Application Number: PCT/US00/08337 (22) International Filing Date: 30 March 2000 (30.03.00) (30) Priority Data: 60/128,400 8 April 1999 (08.04.99) US 09/538,182 30 March 2000 (30.03.00) US (71) Applicant: ORTHO-MCNEIL PHARMACEUTICAL, INC. [US/US]; U.S. Route #202, P.O. Box 300, Raritan, NJ 08869-0602 (US). (72) Inventor: COTTRELL, Sandra, C.; 2 Granite Circle, Doylestown, PA 18901 (US). (74) Agents: CIAMPORCERO, Audley, A., Jr. et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 (US).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: ANTICONVULSANT DERIVATIVES USEFUL IN LOWERING BLOOD PRESSURE

(I)



(II)

(57) Abstract


Use of anticonvulsant derivatives of formula (I) for lowering blood pressure particularly in overweight individuals, wherein X is CH₂ or oxygen; R₁ is hydrogen or alkyl; and R₂, R₃, R₄ and R₅ are independently hydrogen or alkyl and, when X is CH₂, R₄ and R₅ may be alkene groups joined to form a benzene ring and, when X is oxygen, R₂ and R₃, and/or R₄ and R₅ together may be a methylenedioxy group of formula (II), wherein R₆ and R₇ are the same or different and are hydrogen, or alkyl and are joined to form a cyclopentyl or cyclohexyl ring.

ANTICONVULSANT DERIVATIVES USEFUL IN LOWERING BLOOD PRESSURE

BACKGROUND OF THE INVENTION

5

Compounds of Formula I:



are structurally novel antiepileptic compounds that are highly effective anticonvulsants in animal tests (Maryanoff, B.E, Northey, S.O., Gardocki, J.F., Shank, R.P. and Dodgson, S.P. *J. Med. Chem.* 30, 880-887, 1987; Maryanoff, B.E., Costanzo, M.J., Shank, R.P., Schupsky, J.J., Ortegon, M.E., and Vaught J.L. *Bioorganic & Medicinal Chemistry Letters* 3, 2653-2656, 1993). These compounds are covered by US Patent No.4,513,006. One of these compounds 2,3:4,5-bis-O-(1-methylethylidene)-β-D-fructopyranose sulfamate known as topiramate has been demonstrated in clinical trials of human epilepsy to be effective as adjunctive therapy or as monotherapy in treating simple and complex partial seizures and secondarily generalized seizures (E. FAUGHT, B.J. WILDER, R.E. RAMSEY, R.A. REIFE, L D. KRAMER, G.W. PLEDGER, R.M. KARIM et. al., *Epilepsia* 36 (S4) 33, 1995; S.K. SACHDEO, R.C. SACHDEO, R.A. REIFE, P. LIM and G. PLEDGER, *Epilepsia* 36 (S4) 33, 1995), and is currently marketed for the treatment of simple and complex partial seizure epilepsy with or without secondary generalized seizures in approximately twenty countries including the United States, and applications for regulatory approval are presently pending in several additional countries throughout the world.

Compounds of Formula I were initially found to possess anticonvulsant activity in the traditional maximal electroshock seizure (MES) test in mice (SHANK, R.P., GARDOCKI, J.F., VAUGHT, J.L., DAVIS, C.B., SCHUPSKY, J.J., RAFFA, R.B., DODGSON, S.J., NORTEY, S.O., and MARYANOFF, B.E., *Epilepsia* 35 450-460, 1994). Subsequent studies revealed that Compounds of Formula I were also highly


effective in the MES test in rats. More recently topiramate was found to effectively block seizures in several rodent models of epilepsy (J. NAKAMURA, S. TAMURA, T. KANDA, A. ISHII, K. ISHIHARA, T. SERIKAWA, J. YAMADA, and M. SASA, Eur. J. Pharmacol. 254 83-89, 1994), and in an animal model of kindled epilepsy (A. WAUQUIER and S. ZHOU, Epilepsy Res. 24, 73-77, 1996).

Clinical studies on topiramate have revealed previously unrecognized pharmacological properties which suggest that topiramate will be effective in lowering blood pressure, both systolic and diastolic, in humans, particularly in overweight individuals.

10

DISCLOSURE OF THE INVENTION

Accordingly, it has been found that compounds of the following formula I:




15

wherein X is O or CH₂, and R₁, R₂, R₃, R₄ and R₅ are as defined hereinafter are useful in lowering blood pressure.

20 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The sulfamates of the invention are of the following formula (I):



wherein

25

X is CH₂ or oxygen;


WO 00/61136

PCT/US00/08337

R₁ is hydrogen or alkyl; and

R₂, R₃, R₄ and R₅ are independently hydrogen or alkyl and, when X is CH₂, R₄ and R₅ may be alkene groups joined to form a benzene ring and, when X is oxygen, R₂ and R₃ and/or R₄ and R₅ together may be a methylenedioxy group of the following

5 formula (II):



wherein


10 R₆ and R₇ are the same or different and are hydrogen, lower alkyl or are alkyl and are joined to form a cyclopentyl or cyclohexyl ring.

R₁ in particular is hydrogen or alkyl of about 1 to 4 carbons, such as methyl, ethyl and iso-propyl. Alkyl throughout this specification includes straight and branched chain alkyl. Alkyl groups for R₂, R₃, R₄, R₅, R₆ and R₇ are of about 1 to 3 carbons and include methyl, ethyl, iso-propyl and n-propyl. When X is CH₂, R₄ and R₅ may 15 combine to form a benzene ring fused to the 6-membered X-containing ring, i.e., R₄ and R₅ are defined by the alkatrienyl group =C-CH=CH-CH=.

A particular group of compounds of formula (I) is that wherein X is oxygen and both R₂ and R₃ and R₄ and R₅ together are methylenedioxy groups of the formula (II), wherein R₆ and R₇ are both hydrogen both alkyl or combine to form a spiro 20 cyclopentyl or cyclohexyl ring, in particular where R₆ and R₇ are both alkyl such as methyl. A second group of compounds is that wherein X is CH₂ and R₄ and R₅ are joined to form a benzene ring. A third group of compounds of formula (I) is that wherein both R₂ and R₃ are hydrogen.

The compounds of formula (I) may be synthesized by the following methods:

25 (a) Reaction of an alcohol of the formula RCH₂OH with a chlorosulfamate of the formula ClSO₂NH₂ or ClSO₂NHR₁ in the presence of a base such as potassium *n*-butoxide or sodium hydride at a temperature of about -20° to 25° C and in a solvent such as toluene, THF or dimethylformamide wherein R is a moiety of the following formula (III):



(b) Reaction of an alcohol of the formula RCH_2OH with sulfurylchloride of the
 5 formula SO_2Cl_2 in the presence of a base such as triethylamine or pyridine at a
 temperature of about -40° to 25° C in a solvent such as diethyl ether or methylene
 chloride to produce a chlorosulfate of the formula RCH_2OSO_2Cl .

The chlorosulfate of the formula RCH_2OSO_2Cl may then be reacted with an
 amine of the formula R_1NH_2 at a temperature of about -40° to 25° C in a solvent such
 10 as methylene chloride or acetonitrile to produce a compound of formula (I). The
 reaction conditions for (b) are also described by T. Tsuchiya et al. in Tet. Letters, No.
 36, p. 3365 to 3368 (1978).

(c) Reaction of the chlorosulfate RCH_2OSO_2Cl with a metal azide such as sodium
 azide in a solvent such as methylene chloride or acetonitrile yields an azidosulfate of
 15 the formula $RCH_2OSO_2N_3$ as described by M. Hedayatullah in Tet. Lett. p. 2455-2458
 (1975). The azidosulfate is then reduced to a compound of formula (I) wherein R_1 is
 hydrogen by catalytic hydrogenation, e.g. with a noble metal and H_2 or by heating
 with copper metal in a solvent such as methanol.

The starting materials of the formula RCH_2OH may be obtained commercially
 20 or as known in the art. For example, starting materials of the formula RCH_2OH
 wherein both R_2 and R_3 and R_4 and R_5 are identical and are of the formula (II) may
 be obtained by the method of R.F. Brady in Carbohydrate Research, Vol. 14, p. 35 to
 40 (1970) or by reaction of the trimethylsilyl enol ether of a R_6COR_7 ketone or
 aldehyde with fructose at a temperature of about 25° C, in a solvent such as a
 25 halocarbon, e.g. methylene chloride in the presence of a protic acid such as
 hydrochloric acid or a Lewis Acid such as zinc chloride. The trimethylsilyl enol ether
 reaction is described by G.L. Larson et al in J. Org. Chem. Vol. 38, No. 22, p. 3935
 (1973).

Further, carboxylic acids and aldehydes of the formulae RCOOH and RCHO may be reduced to compounds of the formula RCH₂OH by standard reduction techniques, e.g. reaction with lithium aluminum hydride, sodium borohydride or borane-THF complex in an inert solvent such as diglyme, THF or toluene at a temperature of about 0° to 100° C, e.g. as described by H.O. House in "Modern Synthetic Reactions", 2nd Ed., pages 45 to 144 (1972).

The compounds of formula I may also be made by the known process disclosed in United States Patent No. 5,387,700.

The compounds of formula I include the various individual isomers as well as the racemates thereof, e.g., the various alpha and beta attachments, i.e., below and above the plane of the drawing, of R₂, R₃, R₄ and R₅ on the 6-membered ring. Preferably, the oxygens of the methylenedioxy group (II) are attached on the same side of the 6-membered ring.

In a retrospective analysis of the topiramate safety database, mean decreases in systolic and diastolic blood pressure of 3.8 mmHg and 2.6 mmHg, respectively, were observed in all subjects (n=1160). Those subjects with weight at baseline in excess of 100 kg (n=114) had greater decreases of 4.0 mmHg and 3.5 mmHg, respectively. In subjects with both abnormally high blood pressure (defined as ≥140 mmHg systolic or ≥90 mmHg diastolic) and weight in excess of 100 kg (n=39), systolic and diastolic blood pressure decreased by 11.6 mmHg and 8.5 mmHg, respectively.

For lowering blood pressure, a compound of formula (I) may be employed at a daily dose in the range of about 100 mg to 400 mg, usually in two daily divided doses, for an average adult human. A unit dose would contain about 15 to 200 mg of the active ingredient.

To prepare the pharmaceutical compositions of this invention, one or more sulfamate compounds of formula (I) are intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral, by suppository, or parenteral. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be

employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs, and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules and tablets, suitable carriers
5 and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid

pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. Suppositories may be prepared, in which case cocoa butter could be used as the carrier. For parenterals, the carrier will usually comprise sterile water, though other ingredients, for example, for purposes such as

5 aiding solubility or for preservation, may be included. Injectable solutions may also be prepared in which case appropriate stabilizing agents may be employed. Topiramate is currently available for oral administration in round tablets containing 25 mg, 100 mg or 200 mg of active agent. The tablets contain the following inactive ingredients: lactose hydrous, pregelatinized starch, microcrystalline cellulose, sodium starch glycolate,

10 magnesium stearate, purified water, carnauba wax, hydroxypropyl methylcellulose, titanium dioxide, polyethylene glycol, synthetic iron oxide, and polysorbate 80.


The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder injection, teaspoonful, suppository and the like from about 25 to about 200 mg of the active ingredient.

15

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. The use of a compound of formula (I):

5




wherein

X is CH₂ or oxygen;

- 10 R₁ is hydrogen or C₁-C₄ alkyl where alkyl includes straight and branched chain alkyl; and

R₂, R₃, R₄ and R₅ are independently hydrogen or C₁-C₃ alkyl and, when X is CH₂, R₄ and R₅ may be alkene groups joined to form a benzene ring and, when X is oxygen, R₂ and R₃ and/or R₄ and R₅ together may be a methylenedioxy group of formula (II):

15




- 20 wherein R₆ and R₇ are the same or different and are hydrogen or C₁-C₃ alkyl, or R₆ and R₇ together with the carbon atom to which they are attached form a cyclopentyl or cyclohexyl ring; in a therapeutically effective amount for lowering blood pressure in mammals.

2. The use of claim 1, wherein the compound of formula (I) is topiramate.
3. The use of claim 1 or 2, wherein the therapeutically effective amount is from about 100 mg to about 400 mg per day.

4. The use of claim 1 or 2, wherein the therapeutically effective amount is from about 15 mg to about 200 mg per unit dose.

5. The use of a therapeutically effective amount of a compound of formula (I):



wherein

X is CH₂ or oxygen;

10 R₁ is hydrogen or C₁-C₄ alkyl where alkyl includes straight and branched chain alkyl;
and

R₂, R₃, R₄ and R₅ are independently hydrogen or C₁-C₃ alkyl and, when X is CH₂, R₄ and R₅ may be alkene groups joined to form a benzene ring and, when X is oxygen, R₂ and R₃ and/or R₄ and R₅ together may be a methylenedioxy group of formula (II):




20 wherein R₆ and R₇ are the same or different and are hydrogen or C₁-C₃ alkyl, or R₆ and R₇ together with the carbon atom to which they are attached form a cyclopentyl or cyclohexyl ring; in the preparation of a medicament for lowering blood pressure in mammals.


6. The use of claim 5, wherein the compound of formula (I) is topiramate.

7. The use of claim 5 or 6, wherein the therapeutically effective amount is from about 100 mg to about 400 mg per day.

8. The use of claim 5 or 6, wherein the therapeutically effective amount is from about 15 mg to about 200 mg per unit dose.
9. The use of any of claims 5 to 8, wherein the medicament comprises a pharmaceutical composition, comprising a therapeutically effective amount of a
5 compound of formula (I) together with a pharmaceutically acceptable carrier.



(I)



(II)