



(86) **Date de dépôt PCT/PCT Filing Date:** 2008/11/21
(87) **Date publication PCT/PCT Publication Date:** 2009/05/28
(45) **Date de délivrance/Issue Date:** 2016/01/12
(85) **Entrée phase nationale/National Entry:** 2010/04/21
(86) **N° demande PCT/PCT Application No.:** US 2008/084427
(87) **N° publication PCT/PCT Publication No.:** 2009/067706
(30) **Priorité/Priority:** 2007/11/21 (US60/989,786)

(51) **Cl.Int./Int.Cl. A61K 31/6615** (2006.01),
A61K 31/52 (2006.01), **A61K 31/675** (2006.01),
A61K 31/7048 (2006.01), **A61K 31/7068** (2006.01),
A61P 35/00 (2006.01)
(72) **Inventeurs/Inventors:**
CHAPLIN, DAVID, GB;
SIIM, BRONWYN G., GB
(73) **Propriétaire/Owner:**
OXIGENE, INC., US
(74) **Agent:** RIDOUT & MAYBEE LLP

(54) **Titre : PROCÉDE POUR TRAITER DES NEOPLASMES HEMATOPOIETIQUES**
(54) **Title: METHOD FOR TREATING HEMATOPOIETIC NEOPLASMS**

(57) **Abrégé/Abstract:**

This invention relates to methods for treating a hematopoietic neoplasm comprising administering a therapeutically effective amount of a combretastatin compound (CA1dP), or a pharmaceutically acceptable salt thereof, to a subject having a hematological malignancy, wherein the combretastatin compound comprises a catechol or quinone moiety and is capable of forming a reactive oxygen species (ROS) in vivo. The method may further comprise co-administering a second chemotherapeutic agent.



ABSTRACT

This invention relates to methods for treating a hematopoietic neoplasm comprising administering a therapeutically effective amount of a combretastatin compound (CA1dP), or a pharmaceutically acceptable salt thereof, to a subject having a hematological malignancy, wherein the combretastatin compound comprises a catechol or quinone moiety and is capable of forming a reactive oxygen species (ROS) *in vivo*. The method may further comprise co-administering a second chemotherapeutic agent.

METHOD FOR TREATING HEMATOPOIETIC NEOPLASMS

I. Introduction

5 Cancer is a leading cause of death in the industrialized world and despite years of research, many types of cancer lack an effective therapeutic treatment. Although chemotherapy induces remission in the majority of adult patients with acute myeloid leukemia (AML), only a small percent are cured with conventional chemotherapy. Relapse of leukemias is in part due to the persistence of minimal residual leukemias
10 that remain viable within specialized niches, such as vascular niches. Hence, novel treatment strategies are urgently needed to block the interaction of hematopoietic neoplasms with activated vascular cells, interfering with the establishment of pro-leukemic niches in various organs and to eradicate resistant disease.

Adhesion of leukemic cells to stromal cells has been shown to confer increased
15 resistance to chemotherapeutic agents and diminish the rate of apoptosis of the leukemic cells. This process, named cell adhesion-mediated drug resistance (CAM-DR), depends on the interaction of integrins with their ligands. Adhesion of VLA4 (very late antigen 4, $\alpha 4\beta 1$) integrin-positive myeloid cells, to VCAM-1+ stromal cells is an important mediator of CAM-DR. Indeed, expression of VLA4 by leukemic cells portends
20 a poor prognosis and a decreased five-year survival rate. Therefore, identification of novel anti-leukemic agents that inhibit interaction of leukemic cells with vascular cells provides novel strategies to target organ-infiltrating, angiogenesis-dependent leukemias.

Combretastatin A-1, a novel tubulin-destabilizing agent, was isolated from the South African tree *Combretum caffrum*. Combretastatin A-1 binds to tubulin at the same
25 site as colchicine, but with higher affinity. Its phosphate pro-drug, combretastatin A-1 phosphate (CA1dP) induces rapid microtubule depolymerization and vascular shutdown in subcutaneous solid tumors causing tumor necrosis at concentrations well below the maximum tolerated dose. Combretastatin A-1 comprises a single ortho-catechol moiety and is known to be capable of generating an enhanced antitumor response by forming
30 reactive oxygen species (ROS) in the locality of the tumor in addition to selectively reducing the flow of blood to at least a portion of a tumor. This results in both direct inhibition of the proliferation of tumor cells, and tumor selective induction of hypoxia and subsequent necrosis in a portion of the tumor tissue without substantial necrosis of adjoining non-tumor tissue. Other catechol- and quinone-containing vascular disrupting
35 agents have been shown to have enhanced therapeutic activity, relative to vascular

disrupting agents that act solely through interaction with tubulin, particularly against solid tumors (see co-pending USSN 10/790,662, U.S. Patent Publication No. 2004-024696).

In this report, we show that combretastatin compounds that comprise a catechol or quinone moiety induce rapid cell death of non-adherent leukemic cells, at low, non-toxic doses. We also demonstrate that single-agent treatment with a catechol- or quinone-containing combretastatin compound is effective in eradicating both circulating, and vascular-adherent leukemic cells in subcutaneous mouse models of AML, without affecting normal hematopoiesis. CA1dP-treated mice had significantly prolonged survival and significantly decreased tumor load. Co-administration of an additional chemotherapeutic agent, e.g. AraC, decreases tumor load even further. Therefore, catechol- or quinone-containing combretastatin compounds delivered alone or in combination with chemotherapeutic agents represent a promising novel therapeutic approach to eradicate hematopoietic neoplasms.

15 II. SUMMARY OF THE INVENTION

One aspect of the invention provides methods of treating a hematopoietic neoplasm comprising administering a therapeutically effective amount of a combretastatin compound, or a pharmaceutically acceptable salt thereof, to a subject having a hematological malignancy, wherein the combretastatin a quinone or catechol moiety and is capable of forming a reactive oxygen species (ROS) *in vivo*. Preferably, the combretastatin compound is an ortho-quinone, ortho-quinone prodrug, ortho-catechol or ortho-catechol prodrug. More preferably, the compound is combretastatin A1 (CA1) or combretastatin A-1 diphosphate (CA1dP), or a pharmaceutically acceptable salt thereof.

25 Another aspect of the invention provides the use of a combretastatin capable of forming a reactive oxygen species *in vivo*, particularly combretastatin A-1 or a prodrug and/or salt thereof, for the treatment of a hematopoietic neoplasm. The invention also contemplates use of a combretastatin capable of forming a reactive oxygen species *in vivo* in the preparation of a medicament for use in treating a hematopoietic neoplasm.

30 Yet another aspect of the invention provides methods of treating a non-solid tumor comprising administering, to a subject suffering from non-solid tumor, a therapeutically effective amount of combretastatin compound or a pharmaceutically acceptable salt thereof, wherein the combretastatin compound comprises a catechol or quinone moiety and is capable of forming a reactive oxygen species *in vivo*.

III. BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates average tumor volume (\pm SD) as a function of time for subcutaneous HL60 tumors treated with CA4P, CA1dP and/or AraC.

5 Figures 2 and 3 provide individual tumor volumes for selected experiments summarized in Figure 1.

Figure 4 illustrates the dose response activity of CA1dP (\pm AraC) against leukemia cells.

IV. DETAILED DESCRIPTION

A. Definitions

10 As used herein, a "therapeutically effective amount" of combretastatin A-4 phosphate (CA4P), or a therapeutically acceptable salt thereof, according to the present invention is intended to mean that amount of the CA4P that will inhibit the growth of, or retard cancer, or kill malignant cells, and cause the regression and palliation of cancer, *i.e.*, reduce the proliferation rate and/or the number of malignant cells within the body.
15 Other desired anti-tumor effects include, without limitation, the modulation of neoplasm growth rates, the enhancement of necrosis or hypoxia in malignant cells, reduced retention of CEPs and other pro-angiogenic cells, amelioration or minimization of the clinical impairment or symptoms of hematopoietic neoplasms, extending the survival of the subject beyond that which would otherwise be expected in the absence of such
20 treatment, and the prevention of neoplastic growth in an animal lacking any neoplasm formation prior to administration, *i.e.*, prophylactic administration.

As used herein, the terms "modulate", "modulating" or "modulation" refer to changing the rate at which a particular process occurs, inhibiting a particular process, reversing a particular process, and/or preventing the initiation of a particular process.
25 Accordingly, if the particular process is neoplastic growth or metastasis, the term "modulation" includes, without limitation, decreasing the rate at which neoplastic growth and/or metastasis occurs; inhibiting neoplastic growth and/or metastasis, including tumor re-growth following treatment with an anticancer agent; reversing neoplastic growth and/or metastasis (including tumor shrinkage and/or eradication) and/or
30 preventing neoplastic growth and/or metastasis.

"Catechol" is any group of optionally substituted compounds with aryl functionality and containing at least two OH groups at ortho or para positions on the aryl ring, wherein a conjugated system is formed with at least one C=C bond. The preferred

catechol of the present invention is an ortho-benzocatechol. The term "catechol" also encompasses catechols in prodrug form, wherein one or both hydroxyl groups are substituted with a moiety that is metabolically converted to hydroxyl *in vivo*.

"Quinone" is any group of optionally substituted aromatic polyketone compounds derived from a compound with an aryl moiety. At least two C=O groups are in the ortho or para position on the aryl ring, and form a conjugated system with at least one C=C bond. The preferred quinone of the present invention is an ortho-benzoquinone. Quinones can be synthesized in a number of ways by oxidation of a phenolic precursor such as ortho-catechol. The oxidant reagents used in the reaction can include Jones reagent (Chromate salts), Fremy's salt ((KSO₃)₂NO), and the like. Such oxidation also can occur *in vivo* by virtue of chemical or enzymatic activity. The preferred oxidant is o-iodoxybenzoic acid.

"Alkyl" refers to monovalent saturated aliphatic hydrocarbyl groups having from 1 to 10 carbon atoms and preferably 1 to 6 carbon atoms. This term includes, by way of example, linear and branched hydrocarbyl groups such as methyl (CH₃-), ethyl (CH₃CH₂-), n-propyl (CH₃CH₂CH₂-), isopropyl ((CH₃)₂CH-), n-butyl (CH₃CH₂CH₂CH₂-), isobutyl ((CH₃)₂CHCH₂-), sec-butyl ((CH₃)(CH₃CH₂)CH-), t-butyl ((CH₃)₃C-), n-pentyl (CH₃CH₂CH₂CH₂CH₂-), and neopentyl ((CH₃)₃CCH₂-).

"Alkoxy" refers to the group -O-alkyl, wherein alkyl is as defined herein. Alkoxy includes, by way of example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, t-butoxy, sec-butoxy, n-pentoxy, and the like.

"Aryl" refers to a monovalent aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl or anthryl) which condensed rings may or may not be aromatic (e.g., 2-benzoxazolinone, 2H-1,4-benzoxazin-3(4H)-one-7-yl, and the like), provided that the point of attachment is through an atom of the aromatic aryl group. Preferred aryl groups include phenyl and naphthyl.

"Halo" or "halogen" refers to fluoro, chloro, bromo, and iodo and is preferably fluoro or chloro.


"Hydroxy" or "hydroxyl" refers to the group -OH.

As used herein, the term "prodrug" refers to a precursor form of the drug which is metabolically converted *in vivo* to produce the active drug. Thus, for example,

combretastatin phosphate prodrug salts administered to an animal in accordance with the present invention undergo metabolic activation and regenerate combretastatin A-1 *in vivo*, e.g., following dissociation and exposure to endogenous non-specific phosphatases in the body, the drug which is metabolically converted *in vivo* to produce the active drug.

5 Preferred prodrugs of the present invention include the phosphate, phosphoramidate, or amino acid acyl groups as defined herein. The phosphate ester salt moiety may also include $(-OP(O)(O\text{-alkyl})_2)$ or $(-OP(O)(O^-\text{NH}_4^+)_2)$. In preferred embodiments, a prodrug of the invention comprises a substitution of a phenolic moiety or amine moiety of the active drug with a phosphate, phosphoramidate, or amino acid acyl group. A wide variety of
10 methods for the preparation of prodrugs are known to those skilled in the art (see, for example, Pettit and Lippert, *Anti-Cancer Drug Design*, (2000), 15, 203-216).

Catechol prodrugs of the present invention are precursor forms of catechols that are metabolically converted *in vivo* to produce corresponding catechols. In a preferred implementation, the catechol prodrug of the invention comprises at least one phosphate
15 replacing a hydroxyl moiety of the catechol. typically, the phosphate moiety has the general formula:



wherein

each Y independently is OR or NHR, S^- , NH^- or O^- , wherein each R

20 independently is selected from hydrogen, C_{1-6} alkyl, a mono or divalent metal cation, an ammonium cation; and

Z is O or S.

“Hematopoietic neoplasm” refers to a cell proliferative disorder arising from cells of the hematopoietic lineage. Generally, hematopoiesis is the physiological process
25 whereby undifferentiated cells or stem cells develop into various cells found in the peripheral blood. In the initial phase of development, hematopoietic stem cells, typically found in the bone marrow, undergo a series of cell divisions to form multipotent progenitor cells that commit to two main developmental pathways: the lymphoid lineage and the myeloid lineage. The committed progenitor cells of the myeloid lineage
30 differentiate into three major sub-branches comprised of the erythroid, megakaryocyte, and granulocyte/monocyte developmental pathways. An additional pathway leads to formation of dendritic cells, which are involved in antigen presentation. The erythroid

lineage gives rise to red blood cells while the megakaryocytic lineage gives rise to blood platelets. Committed cells of the granulocyte/monocyte lineage split into granulocyte or monocyte developmental pathways, the former pathway leading to formation of neutrophils, eosinophils, and basophils and the latter pathway giving rise to blood
5 monocytes and macrophages.

Neoplasms of hematopoietic cells can involve cells of any phase of hematopoiesis, including hematopoietic stem cells, multipotent progenitor cells, oligopotent committed progenitor cells, precursor cells, and mature differentiated cells. The categories of hematopoietic neoplasms can generally follow the descriptions and
10 diagnostic criteria employed by those of skill in the art (see, *e.g.*, International Classification of Disease and Related Health Problems (ICD 10), World Health Organization (2003)). Hematopoietic neoplasms can also be characterized based on the molecular features, such as cell surface markers and gene expression profiles, cell
15 phenotype exhibited by the aberrant cells, and/or chromosomal aberrations (*e.g.*, deletions, translocations, insertions, etc.) characteristic of certain hematopoietic neoplasms, such as the Philadelphia chromosome found in chronic myelogenous leukemia. Other classifications include National Cancer Institute Working Formulation (Cancer, 1982, 49:2112-2135) and Revised European-American Lymphoma Classification (REAL).

20 The term "hematopoietic neoplasm" includes, but is not limited to, acute lymphoblastic leukemia (ALL), acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL), hairy cell leukemia, Hodgkin's disease, non-Hodgkin's lymphoma, multiple myeloma, and myeloplastic syndrome.

25 "Myeloid neoplasm" refers to proliferative disorder of cells of the myeloid lineage of hematopoiesis. Neoplasms can arise from hematopoietic stem cells, myeloid committed progenitor cells, precursor cells, and terminally differentiated cells. Myeloid neoplasms can be subdivided based on the phenotypic attributes of the aberrant cells or the differentiated state from which the abnormal cells arise. Subdivisions include,
30 among others, myeloproliferative diseases, myelodysplastic/myeloproliferative diseases, myelodysplastic syndromes, acute myeloid leukemia, and acute biphenotypic leukemia.

"Lymphoid neoplasm" refers a proliferative disorder involving cells of the lymphoid lineage of hematopoiesis. Lymphoid neoplasms can arise from hematopoietic stem cells as well as lymphoid committed progenitor cells, precursor cells, and

terminally differentiated cells. These neoplasms can be subdivided based on the phenotypic attributes of the aberrant cells or the differentiated state from which the abnormal cells arise. Subdivisions include, among others, B cell neoplasms, T cell neoplasms, NK cell neoplasms, and Hodgkin's lymphoma. Committed progenitor cells of the lymphoid lineage develop into the B cell pathway, T cell pathway, or the non-T/B cell pathway. Similar to the myeloid lineage, an additional lymphoid pathway appears to give rise to dendritic cells involved in antigen presentation. The B cell progenitor cell develops into a precursor B cell (pre-B), which differentiates into B cells responsible for producing immunoglobulins. Progenitor cells of the T cell lineage differentiate into precursor T cells (pre-T) that, based on the influence of certain cytokines, develop into cytotoxic or helper/suppressor T cells involved in cell mediated immunity. Non-T/B cell pathway leads to generation of natural killer (NK) cells.

The term "hematopoietic neoplasm" includes, but is not limited to, acute lymphoblastic leukemia (ALL), acute myelogenous leukemia (AML), chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL), hairy cell leukemia, Hodgkin's disease, non-Hodgkin's lymphoma, multiple myeloma, and myeloplastic syndrome.

As used herein, the term "pharmaceutically acceptable salt" includes salts that are physiologically tolerated by a subject. Such salts are typically prepared from an inorganic and/or organic acid. Examples of suitable inorganic acids include, but are not limited to, hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, and phosphoric acid. Organic acids may be aliphatic, aromatic, carboxylic, and/or sulfonic acids. Suitable organic acids include, but are not limited to, formic, acetic, propionic, succinic, camphorsulfonic, citric, fumaric, gluconic, lactic, malic, mucic, tartaric, para-toluenesulfonic, glycolic, glucuronic, maleic, furoic, glutamic, benzoic, anthranilic, salicylic, phenylacetic, mandelic, pamoic, methanesulfonic, ethanesulfonic, pantothenic, benzenesulfonic (besylate), stearic, sulfanilic, alginic, galacturonic, and the like. Other pharmaceutically acceptable salts include alkali metal cations such as Na, K, Li; alkali earth metal salts such as Mg or Ca; or organic amine salts such as those disclosed in PCT International Application Nos. WO02/22626 or WO00/48606 and US Patent Nos. 6,855,702 and 6,670,344.

Particularly preferred salts include organic amine salts such tromethamine (TRIS) and amino acid salts such as histidine. Other exemplary salts that can be synthesized using the methods of the invention include those described in US Patent No. 7,018,987.

B. Methods of Treating Hematopoietic Neoplasms

Adhesion of leukemic cells to vascular cells may confer resistance to chemotherapeutic agents. Therefore, disruption of leukemic cell cytoskeletal stability and interference with vascular cell interactions should promote leukemic cell death. Indeed, as disclosed in greater detail below, low and non-toxic doses of the catechol compound combretastatin A-1 diphosphate (CA1dP) inhibit leukemic cell proliferation *in vitro* and induce mitotic arrest and cell death. In addition, CA1dP rapidly increases intracellular reactive oxygen species (ROS), and antioxidant treatment provides partial protection from cell death. As such, combretastatin compounds comprising a catechol or quinone moiety, such as CA1dP, provide for an effective means to treat refractory organ-infiltrating leukemias.

Accordingly, one aspect of the present invention provides a method of treating a hematopoietic neoplasm, the method comprising administering, to a mammal suffering from a hematopoietic neoplasm, a therapeutically effective amount of a combretastatin compound comprising a catechol or quinone moiety. Preferably the combretastatin compound is combretastatin A1, a combretastatin A-1 prodrug (such as combretastatin A-1 diphosphate), or a pharmaceutically acceptable salt thereof.

Derived from the South African tree *Combretum caffrum*, combretastatins such as combretastatin A-4 (CA-4) were initially identified in the 1980's as a potent inhibitors of tubulin polymerization. CA-4, and other combretastatins (*e.g.* CA-1) have been shown to bind at or near the colchicine binding site on tubulin with high affinity. *In vitro* studies clearly demonstrated that combretastatins are potent cytotoxic agents against a diverse spectrum of tumor cell types in culture. CA4P and CA1P, respective phosphate prodrugs of CA-4 and CA-1, were subsequently developed to combat problems with aqueous insolubility. Surprisingly, CA1P has also been shown to cause a rapid and acute shutdown of the blood flow to tumor tissue that is separate and distinct from the anti-proliferative effects of the agents on tumor cells themselves. A number of studies have shown that combretastatins cause extensive shut-down of blood flow within the tumor microvasculature, leading to secondary tumor cell death (Dark *et al.*, *Cancer Res.*, 57: 1829-34, (1997); Chaplin *et al.*, *Anticancer Res.*, 19: 189-96, (1999); Hill *et al.*, *Anticancer Res.*, 22(3):1453-8 (2002); Holwell *et al.*, *Anticancer Res.*, 22(2A):707-11, (2002). Blood flow to normal tissues is generally far less affected by CA1P than blood flow to tumors, although blood flow to some organs, such as spleen, skin, skeletal muscle and brain, can be transiently inhibited.

As used herein, the term "combretastatin" or "combretastatin compound" denotes at least one of the combretastatin family of compounds, derivatives or analogs thereof, their prodrugs (preferably phosphate prodrugs) and derivatives thereof, and salts of these compounds. Combretastatins include those anti-cancer compounds isolated from the South African tree *Combretum caffrum*, including without limitation, Combretastatins A-1, A-2, A-3, A-4, B-1, B-2, B-3, B-4, D-1, and D-2, and various prodrugs thereof, exemplified by Combretastatin A-4 phosphate (CA4P) compounds, Combretastatin A-1 diphosphate (CA1dP) compounds and salts thereof (see for example Pettit *et al.*, *Can. J. Chem.*, (1982); Pettit *et al.*, *J. Org. Chem.*, 1985; Pettit *et al.*, *J. Nat. Prod.*, 1987; Lin *et al.*, *Biochemistry*, (1989); Pettit *et al.*, *J. Med. Chem.*, 1995; Pettit *et al.*, *Anticancer Drug Design*, (2000); Pettit *et al.*, *Anticancer Drug Design*, 16(4-5): 185-93 (2001)).


Combretastatin and combretastatin salts contemplated for use in the methods of the invention are described in WO 99/35150; WO 01/81355; WO 02/022626; US Patent Nos. 4,996,237; 5,409,953; 5,561,122; 5,569,786; 6,538,038; 6,670,344; 6,855,702; 7,018,987; 7,078,552; and 7,279,466. Derivatives or analogs of combretastatins also are described in Singh *et al.*, *J. Org. Chem.*, 1989; Cushman *et al.*, *J. Med. Chem.*, 1991; Getahun *et al.*, *J. Med. Chem.*, 1992; Andres *et al.*, *Bioorg. Med. Chem. Lett.*, 1993; Manila, *et al.*, *Liebigs. Ann. Chem.*, 1993; Shirai *et al.*, *Bioorg. Med. Chem. Lett.*, 1994; Medarde *et al.*, *Bioorg. Med. Chem. Lett.*, 1995; Wood *et al.*, *Br. J. Cancer*, 1995; Bedford *et al.*, *Bioorg. Med. Chem. Lett.*, 1996; Dorr *et al.*, *Invest. New Drugs*, 1996; Jonnalagadda *et al.*, *Bioorg. Med. Chem. Lett.*, 1996; Shirai *et al.*, *Heterocycles*, 1997; Aleksandrak, *et al.*, *Anticancer Drugs*, 1998; Chen *et al.*, *Biochem. Pharmacol.*, 1998; Ducki *et al.*, *Bioorg. Med. Chem. Lett.*, 1998; Hatanaka *et al.*, *Bioorg. Med. Chem. Lett.*, 1998; Medarde *et al.*, *Eur. J. Med. Chem.*, 1998; Medina *et al.*, *Bioorg. Med. Chem. Lett.*, 1998; Ohsumi *et al.*, *Bioorg. Med. Chem. Lett.*, 1998; Ohsumi *et al.*, *J. Med. Chem.*, 1998; Pettit, *et al.*, *J. Med. Chem.*, 1998; Shirai *et al.*, *Bioorg. Med. Chem. Lett.*, 1998; Banwell *et al.*, *Aust. J. Chem.*, 1999; Medarde *et al.*, *Bioorg. Med. Chem. Lett.*, 1999; Shan *et al.*, *PNAS*, 1999; Combeau *et al.*, *Mol. Pharmacol.*, 2000; Pettit *et al.*, *J. Med. Chem.*, 2000; Pinney *et al.*, *Bioorg. Med. Chem. Lett.*, 2000; Flynn *et al.*, *Bioorg. Med. Chem. Lett.*, 2001; Gwaltney *et al.*, *Bioorg. Med. Chem. Lett.*, 2001; Lawrence *et al.*, 2001; Nguyen-Hai *et al.*, *Bioorg. Med. Chem. Lett.*, 2001; Xia *et al.*, *J. Med. Chem.*, 2001; Tahir *et al.*, *Cancer Res.*, 2001; Wu-Wong *et al.*, *Cancer Res.*, 2001; Janik *et al.*, *Bioorg. Med. Chem. Lett.*, 2002; Kim *et al.*, *Bioorg Med Chem Lett.*, 2002; Li *et al.*, *Bioorg. Med. Chem. Lett.*, 2002; Nam *et al.*, *Bioorg. Med. Chem. Lett.*, 2002; Wang *et al.*, *J. Med. Chem.* 2002; Hsieh *et al.*, *Bioorg. Med. Chem. Lett.*, 2003; Hadimani *et al.*,

Bioorg. Med. Chem. Lett., 2003; Mu *et al.*, *J. Med. Chem.*, 2003; Nam *et al.*, *Curr. Med. Chem.*, 2003; Pettit *et al.*, *J. Med. Chem.*, 2003; Gaukroger *et al.*, *Org Biomol Chem.* 2003; Bailly *et al.*, *J Med Chem.* 2003; Sun *et al.*, *Anticancer Res.* 2004; Sun *et al.*, *Bioorg Med Chem Lett.* 2004; Liou *et al.*, *J Med Chem.* 2004; Perez-Melero *et al.*, *Bioorg Med Chem Lett.* 2004; Liou *et al.*, *J Med Chem.* 2004; Mamane *et al.*, *Chemistry.* 2004 ; De Martini *et al.*, *J Med Chem.* 2004; Ducki *et al.*, *J Med Chem.* 2005; Maya *et al.*, *J Med Chem.* 2005; Medarde *et al.*, *J Enzyme Inhib Med Chem.* 2004; Simoni *et al.*, *J Med Chem.* 2005; Sanchez *et al.*, *Bioorg Med Chem.* 2005; Vongvanich *et al.*, *Planta Med.* 2005; Tron *et al.*, *J Med Chem.* 2005; Borrel *et al.*, *Bioorg Med Chem.* 2005; Hsieh *et al.*, *Curr Pharm Des.* 2005 ; Lawrence *et al.*, *Curr Pharm Des.* 2005; Hadfield *et al.*, *Eur J Med Chem.* 2005; Pettit *et al.*, *J Med Chem.* 2005; Coggioloa *et al.*, *Bioorg Med Chem Lett.* 2005; Kaffy *et al.*, *Org Biomol Chem.* 2005; Mateo *et al.*, *J Org Chem.* 2005; LeBlanc *et al.*, *Bioorg Med Chem.* 2005; Srivistava *et al.*, *Bioorg Med Chem.* 2005; Nguyen *et al.*, *J Med Chem.* 2005; Kong *et al.*, *Chem Biol.* 2005; Li *et al.*, *Bioorg Med Chem Lett.* 2005; Pettit *et al.*, *J Nat Prod.* 2005; Nicholson *et al.*, *Anticancer Drugs.* 2006; Monk *et al.*, *Bioorg Med Chem.* 2006; De Martino *et al.*, *J Med Chem.* 2006; Peifer *et al.*, *J Med Chem.* 2006; Kaffy *et al.*, *Bioorg Med Chem.* 2006; Banwell *et al.*, *Bioorg Med Chem.* 2006; Dupeyre *et al.*, *Bioorg Med Chem.* 2006 Simoni *et al.*, *J Med Chem.* 2006 ; Tron *et al.*, *J Med Chem.* 2006; Romagnoli *et al.*, *J Med Chem.* 2006; Pandit *et al.*, *Bioorg Med Chem.* 2006; Nakamura *et al.*, *ChemMedChem.* 2006; Pirali *et al.*, *J Med Chem.* 2006; Bellina *et al.*, *Bioorg Med Chem Lett.* 2006; Hu *et al.*, *J Med Chem.* 2006; Chang *et al.*, *J Med Chem.* 2006; Thomson *et al.*, *Mol Cancer Ther.* 2006; Fortin *et al.*, *Bioorg Med Chem Lett.*, 2007; Duan *et al.*, *J Med Chem.*, 2007; Zhang *et al.*, *J Med Chem.* 2007; Wu *et al.*, *Bioorg Med Chem Lett.* 2007; Sun *et al.*, *Bioorg Med Chem Lett.* 2007, WO 06/138427; WO 036743; WO 05/007635, WO 03/040077, WO 03/035008, WO 02/50007, WO 02/14329; WO 01/12579, WO 01/09103, WO 01/81288, WO 01/84929, WO 00/48590, WO 00/73264, WO 00/06556, WO 00/35865, WO 99/34788, WO 99/48495, WO 92/16486, US Patent Nos. 7,125,906; 7,105,695; 7,105,501; 7,087,627; 7,030,123; 7,078,552; 7,030,123; 7,018,987; 6,992,106; 6,919,324; 6,846,192, 6,855,702; 6,849,656; 6,794,384; 6,787,672, 6,777,578, 6,723,858, 6,720,323, 6,433,012, 6,423,753, 6,201,001, 6,150,407, 6,169,104, 5,731,353, 5,674,906, 5,430,062, 5,525,632, 4,996,237 and 4,940,726. Any combretastatin compound having a catechol or quinone moiety is suitable for use in the methods of the present invention.

In a preferred implementation, the combretastatin comprises a quinone or catechol, preferable an orthoquinone or ortho-catechol. Oxidation of an ortho-catechol,

such as CA1, to an ortho-quinone can result in oxidative damage to neoplastic cells via redox cycling. This is a process in which the quinone is reduced to a radical (ie. semiquinone), which in turn reduces oxygen to superoxide radicals with the quinone being reformed or cycled. The generation of a quinone derivative from CA1 has been demonstrated *in vivo* and the quinone was found to react rapidly with the reducing agents glutathione and ascorbate. In addition, rapid consumption of oxygen in the presence of ascorbate confirmed the formation of CA1 quinone. Furthermore, redox-cycling, confirming the formation of CA1 semiquinone, was observed with CA1.

In a particular embodiment, the combretastatin compound is of the Formula I:



10

or a pharmaceutically acceptable salt thereof, wherein


each R^1 independently is selected from C_{1-6} alkoxy, halogen and halo-substituted C_{1-6} alkyl;

each R^2 independently is selected from C_{1-6} alkoxy, halogen and halo-substituted C_{1-6} alkyl;

p is 1, 2, 3, or 4;

q is 0, 1 or 2;

R^3 and R^4 each independently is selected from hydroxyl or



20


wherein

each Y independently is OR^5 or NHR^5 , or O^- , wherein each R^5 independently is selected from hydrogen and C_{1-6} alkyl;

Z is O or S.


In a preferred implementation, the compound is a pharmaceutically acceptable salt of the compound of Formula I, wherein each of R^3 and R^4 is a phosphate:

25



wherein the cation of the salt can be either a divalent cation or two monovalent cations. Preferably, the divalent cation is a divalent metal cation. In some implementations, the monovalent cation is an alkali metal, an aliphatic amine or an ammonium. In other
 5 implementations, the monovalent cation is selected from the group consisting of sodium, TRIS, histidine, ethanolamine, diethanolamine, ethylenediamine, diethylamine, triethanolamine, glucamine, N-methylglucamine, ethylenediamine, 2-(4-imidazolyl)-ethylamine, choline, and hydrabamine.

In another embodiment, Formula I is represented by a compound of Formula II:



10


wherein

each R¹ independently is selected from C₁₋₆ alkoxy, halogen and halo-substituted C₁₋₆ alkyl;

R² is selected from C₁₋₆ alkoxy, halogen and halo-substituted C₁₋₆ alkyl;

15

R³ and R⁴ each independently is selected from hydroxyl or




wherein

each Y independently is OR⁵ or NHR⁵, or O⁻, wherein each R⁵ independently is selected from hydrogen and C₁₋₆ alkyl;

20

Z is O or S.

In a particularly preferred implementation, the compound is combretastatin A-1 diphosphate (CA1dP):



or a pharmaceutically acceptable salt thereof.

The method of the invention can further comprise co-administering a chemotherapeutic agent, such a cytosine arabinoside (Ara-C), to the subject. "Co-administration" or "co-administering" can be in the form of a single formulation (combining, for example, CA1dP and Ara-C with pharmaceutically acceptable excipients, optionally segregating the two active ingredients in different excipient mixtures designed to independently control their respective release rates and durations) or by independent administration of separate formulations containing the active agents.

"Co-administration" further includes concurrent administration (e.g. administration of CA1dP and a Ara-C at the same time) and time varied administration (administration of CA1dP at a time different from that of the Ara-C), as long as both the combretastatin A-1 diphosphate, or a pharmaceutically acceptable salt thereof,, and the chemotherapeutic agent, e.g., Ara-C, are present in the body in therapeutically effective concentrations during at least partially overlapping times. In preferred implementations the chemotherapeutic agent is Ara-C, etoposide, thioguanine or cyclophosphamide.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and CYTOXAN® (cyclophosphamide); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN®), CPT-11 (irinotecan, CAMPTOSAR®), acetylcamptothecin, scopolectin, and 9-aminocamptothecin); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide; cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin;

duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin;
 pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil,
 chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine,
 mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine,
 5 prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine,
 chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as
 the enediyne antibiotics (e. g., calicheamicin, especially calicheamicin gamma1I and
 calicheamicin omega1 (see, e.g., Agnew, Chem Intl. Ed. Engl., 33: 183-186 (1994));
 dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin
 10 chromophore and related chromoprotein enediyne antiobiotic chromophores),
 aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin,
 carabycin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin,
 detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (including ADRIAMYCIN®,
 morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin,
 15 doxorubicin HCl liposome injection (DOXIL®) and deoxydoxorubicin), epirubicin,
 esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic
 acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin,
 rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-
 metabolites such as methotrexate, gemcitabine (GEMZAR®), tegafur (UFTORAL®),
 20 capecitabine (XELODA®), an epothilone, and 5-fluorouracil (5-FU); folic acid analogues
 such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as
 fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as
 ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine (Ara-C), dideoxyuridine,
 doxifluridine, encitabine, floxuridine; androgens such as calusterone, dromostanolone
 25 propionate, epitiostanol, mepitiothane, testolactone; anti-adrenals such as
 aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid;
 aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine;
 bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine;
 elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine;
 30 maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone;
 mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-
 ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products,
 Eugene, Oreg.); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid;
 triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin
 35 A, roridin A and anguidine); urethan; vindesine (ELDISINE®, FILDESIN®); dacarbazine;
 mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C");
 thiotepa; taxoids, e.g., paclitaxel (TAXOL®), albumin-engineered nanoparticle

formulation of paclitaxel (ABRAXANE™), and doxorubicin (TAXOTERE®); chlorambucil; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine (VELBAN®); platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN®); oxaliplatin; leucovorin; vinorelbine
5 (NAVELBINE®); novantrone; edatrexate; daunomycin; aminopterin; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone,
10 and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATIN™) combined with 5-FU and leucovorin.

As is well-known in the art, solid tumors are quite distinct from non-solid tumors, such as those found in hematopoietic-related cancers. Examples of non-solid tumors include leukemias, such as myeloid leukemias and lymphoid leukemias, myelomas, and
15 lymphomas. In some implementations, the non-solid tumor is a hematopoietic neoplasm, which is aberrant growth of cells of the hematopoietic system.

Hematopoietic malignancies can have their origins in pluripotent stem cells, multipotent progenitor cells, oligopotent committed progenitor cells, precursor cells, and terminally differentiated cells involved in hematopoiesis. Some hematological malignancies are
20 believed to arise from hematopoietic stem cells, which have the ability for self renewal. For instance, cells capable of developing specific subtypes of acute myeloid leukemia (AML) upon transplantation display the cell surface markers of hematopoietic stem cells, implicating hematopoietic stem cells as the source of leukemic cells. Although hematopoietic neoplasms often originate from stem cells, committed progenitor cells or
25 more terminally differentiated cells of a developmental lineage can also be the source of some leukemias. For example, forced expression of the fusion protein Bcr/Abl (associated with chronic myelogenous leukemia) in common myeloid progenitor or granulocyte/macrophage progenitor cells produces a leukemic-like condition.

Hematopoietic neoplasms differ from solid tumors in being capable of circulating
30 and having access to various organs through interaction with activated vascular cells. Indeed, some hematopoietic neoplastic cells may adhere to vascular cells, establishing perivascular infiltrates, and as such may be endowed with a unique mechanism of resistance to chemotherapy. Both circulating and vascular-adherent hematopoietic neoplasms require cytoskeletal stability to maintain mitochondrial and cellular function
35 and avoid cell death. Low and non-toxic doses of combretastatins and combretastatin

compounds comprising a catechol or quinone moiety, can selectively induce apoptosis of circulating and vascular-bound leukemic cells, leading to cell death. This induction of apoptosis occurs by a caspase-dependent mechanism as well as ROS-mediated mitochondrial damage. Thus, combretastatin compounds comprising a catechol or quinone moiety are effective in treating hematopoietic neoplasms, as demonstrated by their ability to target hematopoietic neoplasms *in vitro* and *in vivo* and to eradicate circulating, marrow- and organ-resident vascular-adherent hematopoietic neoplasms.

In some implementations, the hematopoietic neoplasm treated is a lymphoid neoplasm, where the abnormal cells are derived from and/or display the characteristic phenotype of cells of the lymphoid lineage. Lymphoid neoplasms can be subdivided into B-cell neoplasms, T and NK -cell neoplasms, and Hodgkin's lymphoma. B-cell neoplasms can be further subdivided into precursor B-cell neoplasm and mature/peripheral B-cell neoplasm. Exemplary B-cell neoplasms are precursor B-lymphoblastic leukemia/lymphoma (precursor B-cell acute lymphoblastic leukemia) while exemplary mature/peripheral B-cell neoplasms are B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma, B-cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, splenic marginal zone B-cell lymphoma, hairy cell leukemia, plasma cell myeloma/plasmacytoma, extranodal marginal zone B-cell lymphoma of MALT type, nodal marginal zone B-cell lymphoma, follicular lymphoma, mantle-cell lymphoma, diffuse large B-cell lymphoma, mediastinal large B-cell lymphoma, primary effusion lymphoma, and Burkitt's lymphoma/Burkitt cell leukemia. T-cell and NK-cell neoplasms are further subdivided into precursor T-cell neoplasm and mature (peripheral) T-cell neoplasms. Exemplary precursor T-cell neoplasm is precursor T-lymphoblastic lymphoma/leukemia (precursor T-cell acute lymphoblastic leukemia) while exemplary mature (peripheral) T-cell neoplasms are T-cell prolymphocytic leukemia T-cell granular lymphocytic leukemia, aggressive NK-cell leukemia, adult T-cell lymphoma/leukemia (HTLV-1), extranodal NK/T-cell lymphoma, nasal type, enteropathy-type T-cell lymphoma, hepatosplenic gamma-delta T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, Mycosis fungoides/Sezary syndrome, Anaplastic large-cell lymphoma, T/null cell, primary cutaneous type, Peripheral T-cell lymphoma, not otherwise characterized, Angioimmunoblastic T-cell lymphoma, Anaplastic large-cell lymphoma, T/null cell, primary systemic type. The third member of lymphoid neoplasms is Hodgkin's lymphoma, also referred to as Hodgkin's disease. Exemplary diagnosis of this class that can be treated with the compounds include, among others, nodular lymphocyte-predominant Hodgkin's lymphoma, and various classical forms of Hodgkin's disease, exemplary members of which are Nodular

sclerosis Hodgkin's lymphoma (grades 1 and 2), Lymphocyte-rich classical Hodgkin's lymphoma, Mixed cellularity Hodgkin's lymphoma, and Lymphocyte depletion Hodgkin's lymphoma. In various implementations, any of the lymphoid neoplasms can be treated with the combretastatin compounds of the invention.

5 In some implementations, the hematopoietic neoplasm treated is a myeloid neoplasm. This group comprises a large class of cell proliferative disorders involving or displaying the characteristic phenotype of the cells of the myeloid lineage. Myeloid neoplasms can be subdivided into myeloproliferative diseases, myelodysplastic/myeloproliferative diseases, myelodysplastic syndromes, and acute
10 myeloid leukemias. Exemplary myeloproliferative diseases are chronic myelogenous leukemia (*e.g.*, Philadelphia chromosome positive (t(9;22)(qq34;q11)), chronic neutrophilic leukemia, chronic eosinophilic leukemia/hypereosinophilic syndrome, chronic idiopathic myelofibrosis, polycythemia vera, and essential thrombocythemia. Exemplary myelodysplastic/myeloproliferative diseases are chronic myelomonocytic
15 leukemia, atypical chronic myelogenous leukemia, and juvenile myelomonocytic leukemia. Exemplary myelodysplastic syndromes are refractory anemia, with ringed sideroblasts and without ringed sideroblasts, refractory cytopenia (myelodysplastic syndrome) with multilineage dysplasia, refractory anemia (myelodysplastic syndrome) with excess blasts, 5q- syndrome, and myelodysplastic syndrome. In various
20 implementations, any of the myeloid neoplasms can be treated with the combretastatin compounds of the invention.

In some implementations, combretastatin compounds comprising a quinone or catechol, can be used to treat acute myeloid leukemias (AML), which represent a large class of myeloid neoplasms having its own subdivision of disorders. These subdivisions
25 include, among others, AMLs with recurrent cytogenetic translocations, AML with multilineage dysplasia, and other AML not otherwise categorized. Exemplary AMLs with recurrent cytogenetic translocations include, among others, AML with t(8;21)(q22;q22), AML1(CBF-alpha)/ETO, Acute promyelocytic leukemia (AML with t(15;17)(q22;q11-12) and variants, PML/RAR-alpha), AML with abnormal bone marrow eosinophils
30 (inv(16)(p13q22) or t(16;16)(p13;q11), CBFb/MYH11X), and AML with 11q23 (MLL) abnormalities. Exemplary AML with multilineage dysplasia are those that are associated with or without prior myelodysplastic syndrome. Other acute myeloid leukemias not classified within any definable group include, AML minimally differentiated, AML without maturation, AML with maturation, acute myelomonocytic

leukemia, acute monocytic leukemia, acute erythroid leukemia, acute megakaryocytic leukemia, acute basophilic leukemia, and acute panmyelosis with myelofibrosis.

One aspect of the invention is a pharmaceutical composition useful for treating a hematopoietic neoplasm in a mammal, which composition comprises combretastatin
5 compound comprising a quinone or catechol, in combination with a pharmaceutically acceptable excipient. The composition is prepared in accordance with known formulation techniques to provide a composition suitable for oral, topical, transdermal, rectal, by inhalation, parenteral (intravenous, intramuscular, or intraperitoneal) administration, and the like. Detailed guidance for preparing compositions of the
10 invention are found by reference to the 18th or 19th Edition of Remington's Pharmaceutical. Sciences, Published by the Mack Publishing Co., Easton, Pa. 18040. In certain implementations, the pharmaceutical composition further comprises a chemotherapeutic agent, such as Ara-C, etoposide, thioguanine or cyclophosphamide.

Unit doses or multiple dose forms are contemplated, each offering advantages in
15 certain clinical settings. The unit dose would contain a predetermined quantity of active compound calculated to produce the desired effect(s) in the setting of treating cancer. The multiple dose form may be particularly useful when multiples of single doses, or fractional doses, are required to achieve the desired ends. Either of these dosing forms may have specifications that are dictated by or directly dependent upon the unique
20 characteristic of the particular compound, the particular therapeutic effect to be achieved, and any limitations inherent in the art of preparing the particular compound for treatment of cancer.

A unit dose will contain a therapeutically effective amount sufficient to treat a hematopoietic neoplasm in a subject and may contain from about 1.0 to 1000 mg of
25 compound, for example about 50 to 500 mg.

The combretastatin compound comprising a catechol or quinone moiety preferably is administered parenterally, e.g., intravenously, intramuscularly, intravenously, subcutaneously, or intraperitoneally. The carrier or excipient or excipient mixture can be a solvent or a dispersive medium containing, for example, various polar
30 or non-polar solvents, suitable mixtures thereof, or oils. As used herein "carrier" or "excipient" means a pharmaceutically acceptable carrier or excipient and includes any and all solvents, dispersive agents or media, coating(s), antimicrobial agents, iso/hypo/hypertonic agents, absorption-modifying agents, and the like. The use of such substances and the agents for pharmaceutically active substances is well known in the

art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use in therapeutic compositions is contemplated. Moreover, other or supplementary active ingredients can also be incorporated into the final composition.

5 Solutions of the compound may be prepared in suitable diluents such as water, ethanol, glycerol, liquid polyethylene glycol(s), various oils, and/or mixtures thereof, and others known to those skilled in the art.

10 The pharmaceutical forms suitable for injectable use include sterile solutions, dispersions, emulsions, and sterile powders. The final form must be stable under conditions of manufacture and storage. Furthermore, the final pharmaceutical form must be protected against contamination and must, therefore, be able to inhibit the growth of microorganisms such as bacteria or fungi. A single intravenous or intraperitoneal dose can be administered. Alternatively, a slow long term infusion or multiple short term daily infusions may be utilized, typically lasting from 1 to 8 days. Alternate day or dosing once every several days may also be utilized.

15 Sterile, injectable solutions are prepared by incorporating a compound in the required amount into one or more appropriate solvents to which other ingredients, listed above or known to those skilled in the art, may be added as required. Sterile injectable solutions are prepared by incorporating the compound in the required amount in the appropriate solvent with various other ingredients as required. Sterilizing procedures, such as filtration, then follow. Typically, dispersions are made by incorporating the compound into a sterile vehicle which also contains the dispersion medium and the required other ingredients as indicated above. In the case of a sterile powder, the preferred methods include vacuum drying or freeze drying to which any required ingredients are added.

25 In all cases the final form, as noted, must be sterile and must also be able to pass readily through an injection device such as a hollow needle. The proper viscosity may be achieved and maintained by the proper choice of solvents or excipients. Moreover, the use of molecular or particulate coatings such as lecithin, the proper selection of particle size in dispersions, or the use of materials with surfactant properties may be utilized.

30 Prevention or inhibition of growth of microorganisms may be achieved through the addition of one or more antimicrobial agents such as chlorobutanol, ascorbic acid,

parabens, thimerosal, or the like. It may also be preferable to include agents that alter the tonicity such as sugars or salts.

The combretastatin compound comprising a catechol or quinone moiety also can be administered orally in a suitable formulation as an ingestible tablet, a buccal tablet, capsule, caplet, elixir, suspension, syrup, trouche, wafer, lozenge, and the like. Generally, the most straightforward formulation is a tablet or capsule (individually or collectively designated as an "oral dosage unit"). Suitable formulations are prepared in accordance with a standard formulating techniques available that match the characteristics of the compound to the excipients available for formulating an appropriate composition. A tablet or capsule will preferably contain about 50 to about 500 mg of such a combretastatin compound.

The form may deliver a compound rapidly or may be a sustained-release preparation. The compound may be enclosed in a hard or soft capsule, may be compressed into tablets, or may be incorporated with beverages, food or otherwise into the diet. The percentage of the final composition and the preparations may, of course, be varied and may conveniently range between 1 and 90% of the weight of the final form, e.g., tablet. The amount in such therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions according to the current invention are prepared so that an oral dosage unit form contains between about 5.0 to about 50% by weight (%w) in dosage units weighing between 5 and 1000 mg.

The suitable formulation of an oral dosage unit may also contain: a binder, such as gum tragacanth, acacia, corn starch, gelatin; sweetening agents such as lactose or sucrose; disintegrating agents such as corn starch, alginic acid and the like; a lubricant such as magnesium stearate; or flavoring such a peppermint, oil of wintergreen or the like. Various other materials may be present as coating or to otherwise modify the physical form of the oral dosage unit. The oral dosage unit may be coated with shellac, a sugar or both. Syrup or elixir may contain the compound, sucrose as a sweetening agent, methyl and propylparabens as a preservative, a dye and flavoring. Any material utilized should be pharmaceutically-acceptable and substantially non-toxic. Details of the types of excipients useful may be found in the nineteenth edition of "Remington: The Science and Practice of Pharmacy," Mack Printing Company, Easton, Pa. See particularly chapters 91-93 for a fuller discussion.

Another aspect of this invention is a method for treating a hematopoietic neoplasm in a warm-blooded animal, which method comprises administering a

therapeutically effective amount of a combretastatin compounds comprising a catechol or quinone moiety. A combretastatin compound useful in this invention is administered to an appropriate subject in need of these agents in a therapeutically effective dose by a medically acceptable route of administration such as orally, parentally (e.g.,
5 intramuscularly, intravenously, subcutaneously, interperitoneally), transdermally, rectally, by inhalation and the like.

With mammals, including humans, the effective amounts can be administered on the basis of body surface area. The interrelationship of dosages varies for animals of various sizes and species, and for humans (based on mg/m^2 of body surface) is
10 described by E. J. Freireich et al., *Cancer Chemother. Rep.*, 50(4) :219 (1966). Body surface area may be approximately determined from the height and weight of an individual (see, e.g., *Scientific Tables*, Geigy Pharmaceuticals, Ardsley, N.Y. pp. 537-538 (1970)). A suitable dose range is from 1 to 1000 mg of equivalent per m^2 body surface area of a compound of the invention, for instance from 50 to 500 mg/m^2 .

15 Another important feature of the method provided by the present invention relates to the relatively low apparent overall toxicity of the derivatives administered in accordance with the teachings herein. Overall toxicity can be judged using various criteria. For example, loss of body weight in a subject over 10% of the initially recorded body weight (i.e., before treatment) can be considered as one sign of toxicity. In
20 addition, loss of overall mobility and activity and signs of diarrhea or cystitis in a subject can also be interpreted as evidence of toxicity.

V. EXAMPLES

A. EXAMPLE 1: Subcutaneous *in vivo* leukemia model

HL60 (1×10^7 cells in 0.1 ml) were injected subcutaneously into the dorsa of 4-6
25 week old athymic nu/nu mice (Southern Research Institute). When palpable tumors were evident (average tumor volume 100mm^3 ; i.e. approximately 12 days after inoculation), 6 experimental groups of mice were randomized, each with 10 animals. Treatment was initiated according to the following regimens:

Group	Animals	Compound	Dose	Treatment Schedule
1	10	Control (PBS)	0	Q4D x 2/2 wks (SD) – Q3Hx3 ()
2	10	CA1dP	75 mg/kg/inj	Q7Dx2 (SD+1)
3	10	CA4P	75 mg/kg/inj	Q7Dx2 (SD+1)
4	10	Ara-C	20 mg/kg/inj	Q4Dx2/2 wks (SD) – Q3Hx3 ()
5	10	CA1dP Ara-C	75 mg/kg/inj 20 mg/kg/inj	Q7Dx2 (SD+1) Q4Dx2/2 wks (SD) – Q3Hx3 ()
6	10	CA4P Ara-C	75 mg/kg/inj 20 mg/kg/inj	Q7Dx2 (SD+1) Q4Dx2/2 wks (SD) – Q3Hx3 ()

Tumor volume was measured three times a week until the endpoint was reached (tumor volume >3000 mm³), or for 90 days post-treatment. Treatment with Ara-C alone showed little improvement over treatment with phosphate buffered saline (control).

- 5 Treatment with CA4P as a single agent showed delayed tumor growth and extended life span as compared to controls (See Figure 1). Treatment with an orthoquinone prodrug, CA1dP, demonstrated substantial decreases in tumor volume and five complete tumor regressions in a population of ten treated mice (Figure 2). Addition of Ara-C to CA1dP further decreased tumor volumes and led to six complete regressions out of ten treated mice (Figure 3).

A. EXAMPLE 2: Dose Response Activity

- HL60 (1 X 10⁷ cells in 0.1 ml) were injected subcutaneously into the dorsa of 4-6 week old athymic nu/nu mice (Southern Research Institute). When palpable tumors were evident (average tumor volume 100mm³; i.e. approximately 12 days after inoculation), ten experimental groups of mice were randomized, each with 10 animals. Treatment was initiated according to the following regimens:


Group	Compound	Dose	Treatment Schedule
1	Control	0 mg/kg/inj	
2	CA1dP	75 mg/kg/inj	day 3 and 10
3	CA1dP	25 mg/kg/inj	day 3 and 10
4	CA1dP	10 mg/kg/inj	day 3 and 10
5	CA1dP	2.5 mg/kg/inj	day 3 and 10
6	Ara-C	20 mg/kg/inj	tid day 1, 4, 8 and 12
7	Ara-C +CA1dP	20 mg/kg/inj 75 mg/kg/inj	tid day 1, 4, 8 and 12 day 3 and 10
8	Ara-C +CA1dP	20 mg/kg/inj 25 mg/kg/inj	tid day 1, 4, 8 and 12 day 3 and 10
9	Ara-C +CA1dP	20 mg/kg/inj 10 mg/kg/inj	tid day 1, 4, 8 and 12 day 3 and 10
10	Ara-C +CA1dP	20 mg/kg/inj 2.5 mg/kg/inj	tid day 1, 4, 8 and 12 day 3 and 10

Tumor volume was measured three times per week. The decrease in tumor volume showed a clear dose response relationship relative to the dose of CA1dP administered (See Figure 4). administration of doses of CA1dP above 2.5 mg/kg showed significant anti-tumor activity. Although the number of complete responses was not significantly altered by the combination with araC, tumors showed longer growth inhibition following treatment with CA1dP and araC as compared to tumors treated with CA1dP alone.

The scope of the claims should not be limited by the embodiments set forth in the examples, but should be given the broadest interpretation consistent with the description as a whole.

WE CLAIM:

1. A combretastatin compound, or a pharmaceutically acceptable salt thereof, for use in treating a myeloid neoplasm, wherein the combretastatin compound is combretastatin A-1 diphosphate (CA1dP):



or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, wherein the compound is a pharmaceutically acceptable salt, wherein the cation of the salt can be either a divalent cation or two monovalent cations.
3. The compound of claim 2, wherein, the divalent cation is a divalent metal cation.
4. The compound of claim 2, wherein, the monovalent cation is an alkali metal, an aliphatic amine or an ammonium.
5. The compound of claim 2, wherein, the monovalent cation is selected from the group consisting of sodium, TRIS, histidine, ethanolamine, diethanolamine, ethylenediamine, diethylamine, triethanolamine, glucamine, N-methylglucamine, ethylenediamine, 2-(4-imidazolyl)-ethylamine, choline, and hydrabamine.
6. The compound of claim 1, wherein the myeloid neoplasm is an acute myeloid leukemia (AML).

7. The compound of claim 1, wherein the myeloid neoplasm is a refractory organ-infiltrating leukemia.
8. The compound of claim 1, wherein the pharmaceutically acceptable salt is a potassium salt of combretastatin A-1 diphosphate.
9. The compound of claim 1 in combination with a chemotherapeutic agent.
10. The compound of claim 9, wherein the chemotherapeutic agent is Ara-C, etoposide, thioguanine or cyclophosphamide.




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4