



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2671531 C 2016/06/14

(11)(21) **2 671 531**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2007/12/20

(87) Date publication PCT/PCT Publication Date: 2008/07/03

(45) Date de délivrance/Issue Date: 2016/06/14

(85) Entrée phase nationale/National Entry: 2009/06/03

(86) N° demande PCT/PCT Application No.: US 2007/025983

(87) N° publication PCT/PCT Publication No.: 2008/079250

(30) Priorité/Priority: 2006/12/21 (US60/876,555)

(51) Cl.Int./Int.Cl. *C07C 5/03* (2006.01)

(72) Inventeurs/Inventors:

HUTCHENSON, KEITH W., US;
JACKSON, SCOTT CHRISTOPHER, US;
MANZER, LEO ERNEST, US;
SCIALDONE, MARK A., US;
SEAPAN, MAYIS, US

(73) Propriétaire/Owner:

E. I. DU PONT DE NEMOURS AND COMPANY, US


(74) Agent: TORYS LLP

(54) Titre : HYDROGENATION DE CARYOPHYLLENES

(54) Title: HYDROGENATION OF CARYOPHYLLENE

(57) Abrégé/Abstract:

A process for the hydrogenation of a mixture of caryophyllenes by contact with hydrogen in the presence of a supported catalyst. The resulting products are useful for the application to skin.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 July 2008 (03.07.2008)

PCT

(10) International Publication Number
WO 2008/079250 A3

(51) International Patent Classification:
C07C 5/03 (2006.01)

(74) Agent: **LANGWORTHY, John, A.**; E. I. du Pont de Nemours and Company, Legal Patent Records Center, 4417 Lancaster Pike, Wilmington, Delaware 19805 (US).

(21) International Application Number:
PCT/US2007/025983

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
20 December 2007 (20.12.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/876,555 21 December 2006 (21.12.2006) US

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **E. I. DU PONT DE NEMOURS AND COMPANY** [US/US]; 1007 Market Street, Wilmington, Delaware 19898 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **HUTCHENSON, Keith, W.** [US/US]; 311 Belmont Court, Lincoln University, Pennsylvania 19352 (US). **JACKSON, Scott, Christopher** [US/US]; 700 Kilburn Road, Wilmington, Delaware 19803 (US). **MANZER, Leo, Ernest** [US/US]; 714 Burnley Road, Wilmington, Delaware 19803 (US). **SCIALDONE, Mark, A.** [US/US]; 77 Allsmeer Drive, West Grove, Pennsylvania 19390 (US). **SEAPAN, Mayis** [US/US]; 912 Barley Court, Landenberg, Pennsylvania 19350-1347 (US).

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

(88) Date of publication of the international search report:
24 December 2008

(54) Title: HYDROGENATION OF CARYOPHYLLENE

(57) Abstract: A process for the hydrogenation of a mixture of caryophyllenes by contact with hydrogen in the presence of a supported catalyst. The resulting products are useful for the application to skin.



WO 2008/079250 A3

WO 2008/079250

PCT/US2007/025983

TITLE

Hydrogenation of Caryophyllene

5

10

Technical Field

This invention relates to a process for hydrogenating caryophyllenes. Mixtures comprising hydrogenated caryophyllene are useful for the purpose of providing
15 enhancements to skin.

Background

Caryophyllene is a sesquiterpenoid derivative that
20 can be obtained from the oils of a number of plants, such as clove and catmint.

The partial reduction of caryophyllene, or isomers thereof, to their corresponding dihydro forms has been
25 disclosed in several references. U.S. Patent No. 3,966,819 discloses the photoisomerization of caryophyllene to iso-caryophyllene, and discloses that iso-caryophyllene can be hydrogenated to iso-dihydrocaryophyllene in the presence of a Raney nickel
30 catalyst. Rao and Devaprabhakara [Tetrahedron (1978) 34:2223-2227] describe the partial reduction of

caryophyllene and isocaryophyllene with diimide to the dihydro products dihydrocaryophyllene and dihydroisocaryophyllene, respectively.

5 Several references describe the complete hydrogenation of caryophyllene to the tetrahydro derivative. Naves and Perrottet ["Volatile Plant Materials, XIV, Structure of Caryophyllene", *Helv. Chim. Acta*, (1941) 24:789-804] disclose the hydrogenation of
10 caryophyllene to tetrahydrocaryophyllene in the presence of PtO₂, which is an unsupported catalyst. Similarly, Sorm et al [Chemistry & Industry (1956) 154-155] disclose the preparation of tetrahydrocaryophyllene by catalytic hydrogenation of caryophyllene with PtO₂ in the presence
15 of glacial acetic acid.

A need nevertheless exists for an economically viable route to the synthesis of the hydrogenated forms of caryophyllene.

20

Summary

One embodiment of this invention involves a process for preparing a reaction product comprising at
25 least one member of the group consisting of dihydro- β -caryophyllene, α -humulane and β -caryophyllane; by

(a) contacting a mixture comprising α -humulene and β -caryophyllene with hydrogen in the presence of at least one supported catalyst to form the reaction
30 product; and

(b) optionally, separating the reaction product from the supported catalyst.

The catalytic metals useful for the process include those that may be selected from elements in group consisting of iron, ruthenium, rhenium, copper, osmium, cobalt, rhodium, iridium, nickel, palladium, platinum, and alloys or compounds thereof. A support for the catalytic metal includes those that may be selected from the group consisting of carbon, alumina, silica, silica-alumina, silica-titania, titania, titania-alumina, barium sulfate, calcium carbonate, strontium carbonate, compounds thereof, and combinations thereof. The weight of the catalytic metal in the supported catalyst typically constitutes from about 0.1% to about 70% of the combined weight of the catalytic metal plus the support.

The reaction may be run at a temperature in the range of from about 0°C to about 200°C. Hydrogen may be fed at a pressure in the range of from about 0.1 MPa to about 20.7 MPa. Solvents useful for the reaction include without limitation ethanol, isopropanol, hexane, cyclohexane, ethyl acetate, dioxane, tetrahydrofuran and diethyl ether.

Another embodiment of the processes hereof involves a process wherein there is an additional step of incorporating the reaction product into a composition of matter that contains, in addition to the reaction product, a carrier and/or a cosmetically-active adjuvant.

A further embodiment of this invention involves a composition of matter (such as a topical treatment for skin) or an article of manufacture that includes at least one member of the group consisting of dihydro- β -caryophyllene, α -humulane and β -caryophyllane. Such a composition may also contain a carrier, a cosmetically-active adjuvant, and/or a dihydronepetalactone.

Yet another embodiment of this invention involves a method for imparting fragrance to, or revitalizing, the skin, hide, hair, fur, feathers or other surface of a human or a domesticated animal by applying thereto a composition of matter that contains (a) at least one member of the group consisting of dihydro- β -caryophyllene, α -humulane and β -caryophyllane; and (b) a carrier and/or a cosmetically-active adjuvant.

The compositions of this invention are useful for the purpose of imparting a fragrance to, or otherwise enhancing or revitalizing, the skin, hide, hair, fur, feathers or other surface of a human or a domesticated, companion or sporting animal, such as a dog, cat or horse.

25

30


Detailed Description

This invention provides a process for the hydrogenation of a mixture comprising alpha- and beta-caryophyllene in the presence of a supported catalyst, as well as the reaction-product compounds obtained from such process, and compositions prepared from those compounds.

Definitions:

In the description of the inventions hereof, the following definitional structure is provided for certain terminology as employed in various locations in the specification:


The term "beta-caryophyllene" or " β -caryophyllene" or "(1R,4E,9S)-4,11,11-trimethyl-8-methylene-bicyclo[7.2.0]undec-4-ene", as used herein, refers to the compound having the structural formula [Molecular Weight (MW) 204]:




The term "dihydro- β -caryophyllene" or "4,8,11,11-tetramethyl-bicyclo[7.2.0]undec-4-ene", as used herein, refers to a single isomer or a mixture of isomers having the 8R and/or 8S isomeric configuration shown below,

wherein the 8R isomer is (1R,4E,8R,9R)-4,8,11,11-tetramethylbicyclo[7.2.0]undec-4-ene and the 8S isomer is (1R,4E,8S,9R)-4,8,11,11-tetramethylbicyclo[7.2.0]undec-4-ene (MW 206):

5



The term "alpha-humulene" or " α -humulene" or "alpha-caryophyllene" or "(1E,4E,8E)-2,6,6,9-tetramethyl-1,4,8-cycloundecatriene", as used herein, refers to the compound having the structural formula (MW 204):



15

The term "beta-caryophyllane" or " β -caryophyllane" or "2,6,10,10-tetramethyl-bicyclo[7.2.0]undecane", as used herein, refers to a single isomer or a mixture of isomers having the (i) 2R,6R, (ii) 2R,6S, (iii) 2S,6S and/or (iv) 2S,6R isomeric configuration shown below, wherein

the 2R,6R isomer is (1R,2R,6R,9R)-2,6,10,10-tetramethylbicyclo[7.2.0]undecane,


the 2R,6S isomer is (1R,2R,6S,9R)-2,6,10,10-tetramethylbicyclo[7.2.0]undecane,

25

the 2S,6S isomer is (1R,2S,6S,9R)-2,6,10,10-tetramethylbicyclo[7.2.0]undecane, and

the 2S,6R isomer is (1R,2S,6R,9R)-2,6,10,10-tetramethylbicyclo[7.2.0]undecane (MW 208):

5



10 The term "alpha-humulane" or "α-humulane" or "1,1,4,8-tetramethyl- (6CI,8CI,9CI) cycloundecane", as used herein, refers to a single isomer or a mixture of isomers having the (i) 4R,8R, (ii) 4R,8S, (iii) 4S,8S and/or (iv) 4S,8R isomeric configuration shown below,

15 wherein


the 4R,8R isomer is (4R,8R)-1,1,4,8-tetramethylcycloundecane,

the 4R,8S isomer is (4R,8S)-1,1,4,8-tetramethylcycloundecane,

20 the 4S,8S isomer is (4S,8S)-1,1,4,8-tetramethylcycloundecane, and

the 4S,8R isomer is (4S,8R)-1,1,4,8-tetramethylcycloundecane (MW 210);

25




A "catalyst" is a substance that affects the rate of
 5 the reaction but not the reaction equilibrium, and emerges
 from the process chemically unchanged.

More specifically, this invention provides a process
 for preparing a reaction product comprising at least one
 10 member of the group consisting of dihydro- β -caryophyllene,
 α -humulane and β -caryophyllane; by

(a) contacting a mixture comprising α -humulene
 and β -caryophyllene with hydrogen in the presence of
 at least one supported catalyst to form the reaction
 15 product; and

(b) optionally, separating the reaction product
 from the supported catalyst.

According to the processes hereof, the mixture
 20 comprising α -humulene and β -caryophyllene can be partially
 reduced, wherein one or more of the double bonds of the
 caryophyllene isomers are hydrogenated, or the mixture may
 be completely reduced, such that all of the double bonds
 are hydrogenated. A scheme depicting several possible
 25 reaction sequences for the partial and complete
 hydrogenation of caryophyllenes is shown below:



5

In various embodiments, it may be found that the exocyclic methylene group of beta-caryophyllene is hydrogenated more readily than the endocyclic double bonds of α -humulene or β -caryophyllene.

10

Caryophyllenes useful as starting materials in the reaction may be obtained commercially from a supplier such as Sigma-Aldrich (St. Louis MO). Caryophyllenes are sesquiterpenoids found in the essential oils of many plants, such as clove and catmint; the essential oils may be obtained from plant sources by known methods, including steam distillation such as described in Regnier et al [Phytochem. (1967) 6:1271-1280].

20

A supported catalyst, as used herein, at least one catalytic metal on a support. The catalytic metal is deposited on the support by any one of a number of methods, such as spraying, soaking or physical mixing, followed by drying, calcination, and if necessary, activation through methods such as reduction or

25

oxidation/reduction. A supported catalyst may also be made by co-precipitation or blending of the active components and the support followed by drying, calcination, and if necessary, activation through methods
5 such as reduction or oxidation/reduction. Materials frequently used as a catalyst support are porous solids with high total surface areas (external and internal) which can provide high concentrations of active sites per unit weight of catalyst.

10

In the processes hereof, supported catalysts are preferred to unsupported catalysts for a variety of reasons. The support allows greater efficiency in the use of the metal by increasing the active metal surface
15 and by facilitating metal recovery. Supported catalysts generally have a greater resistance to poisoning. In addition, the support provides for better control over selectivity.

20

The catalytic metal may be selected from elements in the group consisting of iron, ruthenium, rhenium, copper, osmium, cobalt, rhodium, iridium, nickel, palladium, platinum, alloys or compounds thereof, and combinations thereof. In a more specific embodiment, the catalytic
25 metal may be selected from elements in the group consisting of palladium, rhodium, iridium, alloys or compounds thereof, and combinations thereof. The catalytic metal may, however, be selected from any subgroup of the foregoing list formed by omitting other
30 members of the list to leave only the members of the subgroup, and the reaction in such instance is run in the

absence of the members of the list omitted to form the subgroup.

A support for the supported catalyst may be selected
5 from the group consisting of carbon, alumina, silica, silica-alumina, silica-titania, titania, titania-alumina, barium sulfate, calcium carbonate, strontium carbonate, compounds thereof, and combinations thereof. In a more specific embodiment, the support may be selected from the
10 group consisting of carbon, alumina, silica, silica-alumina, compounds thereof, and combinations thereof. In an even more specific embodiment, the support material may be carbon with a surface area greater than $100 \text{ m}^2/\text{g}$, or carbon with a surface area greater than $200 \text{ m}^2/\text{g}$.
15 Preferably, the carbon has an ash content that is less than 5% by weight of the catalyst support; the ash content is the inorganic residue (expressed as a percentage of the original weight of the carbon) which remains after incineration of the carbon.

20

Commercially available carbons which may be used herein as a support include those sold under the following trademarks: Bameby & SutcliffeTM, DarcoTM, NucharTM, Columbia JXNTM, Columbia LCKTM, Calgon PCBTM, Calgon BPLTM,
25 WestvacoTM, NoritTM and Bamaby Cheny NBTM. The carbon can also be commercially available carbon such as Calsicat C, Sibunit C, or Calgon C (commercially available under the registered trademark Centaur®).

The catalyst support can be in the form of powder, granules, pellets, extrudates, or the like.

In the process hereof, the content of the catalytic metal in a supported catalyst will depend on the choice of the catalyst and support. In one embodiment, the content of the catalytic metal in the supported catalyst may be from about 0.1% to about 70% of the supported catalyst based on catalytic metal weight plus the support weight.

Promoter metals may be used to enhance the physical or chemical function of a catalyst. Promoter metals are often selected from metals from Groups 3 through 8, 11 and 12 of the Periodic Table, and are typically used at up to about 2% by weight of the total metal of the promoter plus the catalytic metal. Promoter metals may be used to enhance the activity of the supported catalyst useful for the present invention.

Combinations of catalytic metal and support useful herein include at least one catalytic metal as described above with at least one support as described above. In a more specific embodiment, combinations of catalytic metal and support may be selected from the group consisting of palladium on carbon, palladium on alumina, palladium on silica, palladium on silica-alumina, rhodium on carbon, rhodium on alumina, rhodium on silica, rhodium on silica-alumina, iridium on carbon, iridium on alumina, iridium on silica, iridium on silica-alumina, and combinations thereof.

The reaction may be run at a temperature in the range of from about 0°C to about 200°C. In a more specific embodiment, the reaction may be run at a temperature in the range of from about 25°C to about 150°C. In an even
5 more specific embodiment, the reaction may be run at a temperature of from about 50°C to about 100°C.

The hydrogen feed useful for the reaction may be supplied at a pressure in the range of from about 0.1 MPa
10 to about 20.7 MPa. In one embodiment, the hydrogen is maintained at a pressure to achieve saturation levels of the hydrogen in the mixture at the temperature of the reaction.

15 Solvents may be used in the processes hereof, if desired, and when used are typically inert solvents. Suitable solvents may include without limitation alcohols such as ethanol or isopropanol; alkanes such as hexane or cyclohexane; esters such as ethyl acetate; and ethers
20 such as dioxane, tetrahydrofuran or diethyl ether.

The reaction product comprising the hydrogenated caryophyllenes may be separated from the at least one supported catalyst by any suitable method, such as
25 decantation. For those reactions in which a solvent is used, the reaction product comprising the hydrogenated caryophyllenes may additionally comprise the solvent, or alternatively, the solvent may be recovered from the reaction product using standard techniques known to those
30 skilled in the art, such as distillation or vacuum stripping.

The processes hereof may be carried out in batch in a single reactor, in sequential batch in a series of reactors, in reaction zones within one or more reactors, 5 or in continuous mode in any of the equipment customarily employed for continuous processes.

The reaction products will be determined by the conditions under which the reaction is carried out 10 inasmuch as temperature, hydrogen pressure, catalytic metal, support and time can affect the reaction kinetics, product yield and product selectivity. For example, with a given supported catalyst at a given pressure, β -caryophyllene may be hydrogenated to dihydro- β -caryophyllene at 50°C, whereas α -humulene is not 15 hydrogenated. Under the same conditions except using a temperature of 100°C, both β -caryophyllene and α -humulene may be completely hydrogenated to β -caryophyllane and α -humulane, respectively. The conditions required for 20 producing a specified mixture of products may be similarly determined.

This invention relates to the Hydrogenation Reaction Products ("HRP") as described above (dihydro- β -caryophyllene, α -humulane and/or β -caryophyllane), to 25 compositions comprising HRP, and to the use of HRP and the compositions thereof. The preparations of this invention, which include the HRP compounds described above, and the compositions, formulations and other 30 materials that may be prepared from such compounds according to this invention, and mixtures thereof, may all

be used for a multiplicity of purposes. These purposes include, for example, use as a fragrance compound itself or as an ingredient in a perfume composition, or use as a topical treatment for skin.

5

For example, the preparations hereof may be applied in a topical manner to the skin, hide, hair, fur, feathers or other surface of a mammal, such as a human or domesticated animal.

10

The preparations hereof may also be applied to the skin and/or hair of humans to impart a pleasant odor or aroma as a fragrance compound itself, or as an ingredient in a perfume composition; and the preparations hereof may also be used as a topical treatment for skin by application to the skin and/or hair of humans in the form of a body wash, rinse, conditioner, toner, lotion, splash, spray or other type of cosmetic product as applied personally by the user.

20

Alternatively, the active ingredient may be contained in microcapsules to control the rate of loss from a surface or an article; a precursor molecule, which slowly disintegrates on a surface or in an article, may be used to control the rate of release of the active ingredient; or a synergist may be used to continually stimulate the evaporation of the active from the formulated composition.

25

The release of an active ingredient that is intended for application to the skin or other surface of a user may be accomplished, for example, by sub-micron

30

encapsulation, in which the active ingredient is encapsulated or enveloped in a skin-nourishing protein. The protein may be used, for example, at about a 20 wt% concentration. An application of repellent contains many
5 of these protein capsules that may be suspended in either a water-based lotion, or water for spray application. After contact with skin, the protein capsules begin to break down, releasing the encapsulated active. The process continues as each microscopic capsule is depleted
10 then replaced in succession by a new capsule that contacts the surface and releases its active ingredient. The process may take up to 24 hours for one application. Because a protein adheres very effectively to skin, these formulations are very resistant to perspiration (sweat-
15 off) and dilution by water from other sources.

One of the distinct advantages of the preparations of this invention is that they are all characterized by a relative volatility that makes them
20 suitable for use to obtain a desirably high level of concentration of active ingredient on, above and around a surface of a user, as described above. One or more of these preparations may be used for such purpose as an active, or an active formulation, in a composition in
25 which the preparation is admixed with a carrier suitable for wet or dry application of the composition to a surface in the form, for example, of a liquid, aerosol, gel, aerogel, foam or powder (such as a sprayable powder or a dusting powder). Suitable carriers include any one of a
30 variety of commercially available organic and inorganic liquid, solid, or semi-solid carriers or carrier

formulations usable in formulating a variety of cosmetic products. When formulating a composition for application to the skin or other surface of a human, it is important to select a dermatologically acceptable carrier. A carrier suitable for use herein may include water, alcohol, silicone, petrolatum, lanolin; or may include an organic liquid carrier such as a liquid aliphatic hydrocarbon (e.g. pentane, hexane, heptane, nonane, decane and their analogs) or a liquid aromatic hydrocarbon.

Examples of other useful liquid hydrocarbons include oils produced by the distillation of coal and the distillation of various types and grades of petrochemical stocks, including kerosene oils that are obtained by fractional distillation of petroleum. Suitable petroleum oils include those generally referred to as agricultural spray oils (e.g. the so-called light and medium spray oils, consisting of middle fractions in the distillation of petroleum and which are only slightly volatile). Such oils are usually highly refined and may contain only minute amounts of unsaturated compounds. Such oils, moreover, are generally paraffin oils and accordingly can be emulsified with water and an emulsifier, diluted to lower concentrations, and used as sprays. Tall oils, obtained from sulfate digestion of wood pulp, like the paraffin oils, can similarly be used. Other organic liquid carriers can include liquid terpene hydrocarbons and terpene alcohols such as alpha-pinene, dipentene, terpineol, and the like.

Other suitable carriers include silicone, petrolatum, lanolin, liquid hydrocarbons, agricultural spray oils, paraffin oil, tall oils, liquid terpene hydrocarbons and terpene alcohols, aliphatic and aromatic alcohols, esters, aldehydes, ketones, mineral oil, higher alcohols, finely divided organic and inorganic solid materials. In addition to the above-mentioned liquid hydrocarbons, the carrier can contain conventional emulsifying agents which can be used for causing the active ingredient to be dispersed in, and diluted with, water for end-use application. Still other liquid carriers can include organic solvents such as aliphatic and aromatic alcohols, esters, aldehydes, and ketones. Aliphatic monohydric alcohols include methyl, ethyl, normal-propyl, isopropyl, normal-butyl, sec-butyl, and tert-butyl alcohols. Suitable alcohols include glycols (such as ethylene and propylene glycol) and pinacols. Suitable polyhydroxy alcohols include glycerol, arabitol, erythritol, sorbitol, and the like. Finally, suitable cyclic alcohols include cyclopentyl and cyclohexyl alcohols.

Conventional aromatic and aliphatic esters, aldehydes and ketones can be used as carriers, and occasionally are used in combination with the above-mentioned alcohols. Still other liquid carriers include relatively high-boiling petroleum products such as mineral oil and higher alcohols (such as cetyl alcohol). Additionally, conventional or so-called "stabilizers" (e.g. tert-butyl sulfinyl dimethyl dithiocarbonate) can be used in conjunction with, or as a component of, the

carrier or carriers used in a composition as made according to this invention.

Numerous clays having a layered structure with
5 interstices, and synthetic inorganic materials that resemble such clays in respect of chemical composition, crystallinity and layered morphology, are suitable for use herein as carriers. Suitable clays having a layered structure with interstices include smectite, kaolin,
10 muscovite, vermiculite, phlogopite, xanthophyllite, and chrysotile, and mixtures thereof. Preferred are smectite clays and kaolin clays. Smectite clays include montmorillonite, beidellite, nontronite, saponite, hectorite, sauconite, and others. Kaolin clays include
15 kaolinite, dickite, nacrite, antigorite, and others. Most preferred is montmorillonite. Average particle sizes range from 0.5 to 50 micrometers.


The application of a preparation hereof may be
20 accomplished by dispersing the preparation into the air, or by dispersing the preparation as a liquid mist or incorporated into a powder or dust, and this will permit the composition to fall on the desired surfaces of a user. It may also be desirable to combine a preparation hereof
25 with a fugitive vehicle for application in the form of a spray. Such a composition may be an aerosol, sprayable liquid or sprayable powder composition adapted to disperse the active ingredient into the atmosphere by means of a compressed gas, or a mechanical pump spray. Likewise,
30 directly spreading a preparation in liquid/semi-solid/solid form on a user in wet or dry form (as a

friable solid, for example) is a useful method of contacting a surface of the user.

Further, it may also be desirable to combine a
5 preparation hereof with one or more other compounds known
to have insect repellency in a composition. Suitable
insect repellent compounds combinable for such purpose
include nepetalactones, nepetalactams,
dihydronepetalactones and derivatives thereof,
10 dihydronepetalactams and derivatives thereof, benzil,
benzyl benzoate, 2,3,4,5-bis(butyl-2-ene)
tetrahydrofurfural, butoxypolypropylene glycol, *N*-
butylacetanilide, normal-butyl-6,6-dimethyl-5,6-dihydro-
1,4-pyrone-2-carboxylate, dibutyl adipate, dibutyl
15 phthalate, di-normal-butyl succinate, *N,N*-diethyl-meta-
toluamide, dimethyl carbate, dimethyl phthalate, 2-ethyl-
2-butyl-1,3-propanediol, 2-ethyl-1,3-hexanediol, di-
normal-propyl isocinchomeronate, 2-phenylcyclohexanol, *p*-
methane-3,8-diol, and normal-propyl *N,N*-
20 diethylsuccinamate.

Suitable dihydronepetalactones for use as a component in a composition hereof include the 9S dihydronepetalactones as shown below:

5



10

In addition, a preparations hereof may also include one or more essential oils and/or active ingredients of essential oils. An essential oil includes any type of volatile oil that is obtained from a plant and possesses the odor and other characteristic properties of the plant. Examples of useful essential oils include: almond bitter oil, anise oil, basil oil, bay oil, caraway oil, cardamom oil, cedar oil, celery oil, chamomile oil, cinnamon oil, citronella oil, clove oil, coriander oil, cumin oil, dill oil, eucalyptus oil, fennel oil, ginger oil, grapefruit oil, lemon oil, lime oil, mint oil, parsley oil, peppermint oil, pepper oil, rose oil, spearmint oil (menthol), sweet orange oil, thyme oil, turmeric oil, and oil of wintergreen. Examples of active ingredients in essential oils are: citronellal, methyl salicylate, ethyl salicylate, propyl salicylate, citronellol, safrole, and limonene.

20

In another embodiment, a preparation hereof may be used as a fragrance material or as an active in a fragrance composition, and be applied in a topical manner to human or animal skin or hair to impart a pleasing scent or aroma thereto, as in colognes or perfumes for humans or pets. Alternatively, the pleasing scent or aroma may be obtained by the use of a preparation hereof as an insect/arthropod repellent where the preparation has the dual attributes of simultaneously imparting both repellency as well as the pleasing scent or aroma.

30

As used herein, the terms "alter" and "modify" in their various forms refer to a means of supplying or imparting a fragrance, aroma character or note to otherwise bland substances, or augmenting existing aroma characteristics where natural aroma is deficient in some regard, or supplementing an existing aroma impression to modify its quality, character, or aroma.

The term "enhance" is intended to mean the intensification (without effecting a change in kind or quality of aroma) of one or more aroma nuances and their organoleptic impression of a fragrance, perfume composition, or one or more perfumed articles.

The term "fragrance note(s)" or "note(s)" refers to the three stages that most fragrances go through. The top note is the first impression of the fragrance. The middle note is the main character of a fragrance. These are stronger, mid-range notes that emerge after the top and linger longest, as the 'heart' of the fragrance. Finally, the base note is the final scent of the fragrance. These rich, heavy notes emerge slowly and definitely, echoing resonantly after the others die down. Bottom notes, by definition, linger behind and act as a fixative to stop the lighter oils from dispersing too quickly.

The terms "perfume composition" or "fragrance composition" or "aroma composition" are used herein to mean a mixture of organic compounds including, for example, alcohols, aldehydes, ketones, nitriles, esters, lactones, natural essential oils, synthetic essential

oils, and mercaptans, which are admixed so that the combined odors of the individual components produce a pleasant or desired fragrance. Such compositions usually contain: (1) the main note or the "bouquet" or foundation
5 stone of the composition; (2) modifiers which round off and accompany the main note; (3) fixatives which include odorous substances which lend a particular note to the perfume throughout all stages of evaporation and substances which retard evaporation; and (4) top notes
10 which are usually low-boiling, fresh-smelling materials.

A perfume is characterized by its uniquely pleasing fragrance or aroma. In perfume, fragrance or aroma compositions, the individual component will contribute its
15 particular olfactory characteristics, but the overall effect of the composition will be the sum of each of the effects of each of the ingredients. Thus, an HRP can be used to alter the aroma characteristics of such compositions, for example, by highlighting or moderating
20 the olfactory reaction contributed by another ingredient in the composition.

A "perfume composition" may contain a perfume compound, a fragrance compound or an aroma compound in
25 addition to other materials. Moreover, a "perfume composition" or "fragrance composition" can be used as components of an article such as a "perfumed article" or "fragrant article", or the composition may be added to perfumed or fragrant articles, wherein the term "perfumed
30 article" or "fragrant article" refers to an article of manufacture possessing a pleasing fragrance or aroma, or a

fragrance or aroma that is enhanced, altered, or augmented by the perfume composition.

The unique fragrance notes of an HRP thus make them
5 useful in imparting, altering, augmenting or enhancing the overall olfactory component of a perfume composition, for example, by utilizing or moderating the olfactory reaction contributed by one or more other ingredients in the composition. Specifically, the composition may be
10 utilized to either: (1) impart a characteristic perfume or aroma to a perfume or article; or (2) mask or modify the odor of one or more of the components thereof.

As will be appreciated, fragrant materials are
15 typically utilized in combinations that may include both natural and synthetic ingredients to achieve the desired overall perfume effect. Fragrance developers consider the scent of the compound, as well as its efficacy, degree of stability within the final formulation, activity during a
20 product's shelf life, and lack of adverse reaction with the product or its intended function as a perfumed article. Typically, fragrances are used to mask the odor contributed by other ingredients in the formulation of the final scented product, and/ or to enhance consumer appeal
25 of a product.

An HRP possesses unique fragrant notes and, therefore, are particularly useful individually and in combination with other fragrant chemicals in a variety of
30 perfume compositions. The overall effect of the perfume composition will be the sum of the effects of each of the

ingredients. Thus, an HRP as used herein or mixtures thereof with other perfumery materials can be used to alter the aroma characteristics of a perfume composition, for example, by highlighting or moderating the olfactory reaction contributed by another ingredient in the preparation, such as a fragrance compound that is not an HRP. The HRPs according to the present invention can be used in practically every field of modern perfumery. Embodiments of the invention therefore are suitable for applications in fine perfumery, i.e. in the preparations of perfumes and colognes in which new and original effects can be obtained.

In a further embodiment, the insect/arthropod repellency, fragrance and/or other properties of products directed to other fundamental purposes will be improved by the presence therein of a preparation of this invention. Those other products include, for example, a body wash, rinse, lotion, splash, tonic or toner, bath and shower gels, foam products (e.g. shaving foams), makeup, deodorants, shampoo, hair lacquers/hair rinses, personal soap compositions (e.g. hand soaps and bath/shower soaps) or other personal care treatments or palliatives, and cleaning agents such as detergents and solvents, and air fresheners and odor removers. Such products may be fabricated, for example, in the form of a sprayable liquid, an aerosol, a foam, a cream, an ointment, a gel, a paste, a powder or a friable solid. The process of fabricating such a product would thus include admixing a preparation hereof with suitable carriers or other inert ingredients to facilitate delivery in the physical form as

described, such as liquid carriers that are readily
sprayed; a propellant for an aerosol or a foam; viscous
carriers for a cream, an ointment, a gel or a paste; or
dry or semi-solid carriers for a powder or a friable
5 solid.

Any of the above described products may also
contain other therapeutically or cosmetically active
adjuvants or supplemental ingredients as are typical in
10 the personal care industry. Examples of these include
fungicides, sunscreens agents, sunblocking agents,
vitamins, tanning agents, plant extracts, anti-
inflammatory agents, anti-oxidants, radical scavenging
agents, retinoids, alpha-hydroxy acids, antiseptics,
15 antibiotics, antibacterial agents, antihistamines;
adjuvants such as thickeners, buffering agents, chelating
agents, preservatives, gelling agents, stabilizers,
surfactants, emollients, coloring agents, aloe vera, waxes,
and penetration enhancers; and mixtures of any two or
20 more thereof.

Articles into which a preparation hereof may be
incorporated to impart an improved fragrance include
articles or manufactured goods such as textile and fibrous
25 goods, clothing, sanitary goods, carpeting, linens,
outdoor or military equipment such as tents, tarpaulins,
backpacks or mosquito netting, candles, paper, paint, ink,
wood products such as furniture, plastics and other
polymers, and the like.

30

A preparation hereof may be formulated as or incorporated into a composition for application to an animate host by any of the same methods known in the cosmetics industry, such as dilution, mixing, thickening, emulsifying, bottling and pressurizing. A preparation hereof may be incorporated into an article by mixing during production or by post-production steps such as spraying or dipping.

10 A preparation hereof may be admixed in a composition with other components, such as a carrier, in an amount that is effective for usage for a particular purpose. The amount of a HRP as described herein, contained in a composition will generally not exceed about 80% by weight
15 based on the weight of the final product, however, greater amounts may be utilized in certain applications, and this amount is not limiting. More preferably, a suitable amount of a HRP will be at least about 0.001% by weight and preferably about 0.01% up to about 50% by weight; and
20 more preferably, from about 0.01% to about 20% weight percent, based on the total weight of the total composition or article. Specific compositions will depend on the intended use.

25 Other compositions, materials and methods relevant to the use of a HRP are as disclosed in US 2003/062,357; US 2003/079,786; US 2003/191,047; and US 2006/148,842.

30

The advantageous attributes and effects of the processes hereof may be seen in a series of examples, as described below. The embodiments of these processes on which the examples are based are representative only, and
5 the selection of those embodiments to illustrate the invention does not indicate that materials, conditions, arrangements, components, reactants, techniques or configurations not described in these examples are not suitable for practicing these processes, or that subject
10 matter not described in these examples is excluded from the scope of the appended claims and equivalents thereof.

15

EXAMPLES

The following abbreviations are used in the examples: GC is gas chromatograph(y); GC-MS is gas chromatography-mass spectrometry; NMR is nuclear magnetic resonance; C is Centigrade, MPa is mega Pascal; rpm is revolutions per
20 minute; mL is milliliter; MW is molecular weight; CMO is catmint oil; wt% is weight percent; TOS is time on stream; h is hour; conc. is concentration; conv. is conversion; temp. is temperature; °C is degrees Centigrade.

25 GC-MS Analysis:

Samples were analyzed using an HP6890 GC (Agilent Technologies, Palo Alto, CA) GC equipped with an HP-5973 mass detector and a DB column (JW Scientific (available through Fisher Scientific, Pittsburgh, PA)).

30

NMR Analysis:

NMR spectra were obtained on a Bruker DRX Advance (500 MHz ^1H , 125 MHz ^{13}C ; Bruker Biospin Corp., Billerica, MA) using deuterated solvents obtained from Cambridge Isotope Laboratories, Inc. (Andover, MA).

Example 1: Hydrogenation of caryophyllene (50°C)

An 18.5 wt% solution of caryophyllene (Sigma-Aldrich; St. Louis, MO; purchased as beta-caryophyllene) in absolute ethanol was prepared. GC-MS and NMR analysis of the caryophyllene indicated that it was a mixture of both the alpha- (13.3%) and beta- (86.7%) isomers (MW 204).

This solution was hydrogenated using ESCAT-142 (Engelhard, Iselin, NJ; 5% Pd/C powder catalyst) using 10% powdered catalyst by weight relative to the beta caryophellene; the reaction was carried out for 2 hours at 50°C and 3.45 MPa hydrogen pressure. GC-MS and NMR analysis of the hydrogenated solution indicated that alpha-caryophyllene (alpha-humulene) was not hydrogenated, and beta-caryophyllene was converted to dihydro- β -caryophyllene.

Example 2: Hydrogenation of caryophyllene (100 degrees C)

An 18.5% solution of caryophyllene (Sigma-Aldrich) in absolute ethanol was prepared as described in Example 1.

This solution was hydrogenated using ESCAT-142 using 10% powdered catalyst by weight relative to the beta

caryophellene; the reaction was carried out for 2 hours at 100 degrees C and 3.45 MPa hydrogen pressure. GC-MS and NMR analysis of the hydrogenated solution indicated that alpha-humulene and beta-caryophyllene were completely
5 hydrogenated to alpha-humulane and beta-caryophyllane, respectively.

Example 3: Hydrogenation of caryophyllene

10 A 50 mL stirred batch autoclave reactor was charged with 31.41 g of a 50 wt% solution of caryophyllene (prepared as described in Example 1) and 1.59 g of ESCAT 142. The reactor was equipped with a magnetically-coupled gas entrainment agitator which was rotated at about 1000
15 rpm during the reaction. The reactor was sealed and then flushed and evacuated with nitrogen several times to remove oxygen. These flushes were followed by two rapid flushes with hydrogen to minimize residual nitrogen in the reactor. The reactor was heated to approximately 50
20 degrees C via an external band heater and then charged to 83.9 bar (8.39 MPa) pressure with hydrogen. Hydrogen was continuously fed to the reactor during the course of the run to maintain this pressure as hydrogen was consumed by reaction.

25

Small samples of the reaction mixture were removed periodically via a dip tube equipped with a 2 micron sintered metal frit. The reactor was maintained at an average temperature of 54°C for a total reaction time of
30 3.0 hours. The reactor was then thermally quenched via an external cooling coil supplied with a chilled propylene

glycol/water mixture from a recirculating bath and vented. Product analysis by gas chromatography (GC-FID) using 1,2-dibromobenzene as an internal standard added post reaction showed 99.7% conversion of both caryophyllene isomers
 5 after 0.5 hours, and quantitative conversion by 1.10 hours reaction time. Additional analysis by mass spectroscopy (GC-MS) and C^{13} NMR identified the anticipated hydrogenation product isomers for both the alpha- and beta-caryophyllene (humulane and caryophyllane,
 10 respectively). Concentrations of the pertinent reactants and reaction products are summarized in Table 1, and Figure 1 plots the concentrations of the various isomers of each molecular weight. This example demonstrates complete and rapid hydrogenation of alpha- and beta-
 15 caryophyllene at commercially-viable reactor concentrations in the solvent (i.e. high reactor productivity).

Table 1

20

Time on Stream (hr)	Beta-Caryophyllene (MW 204) (wt%)	Alpha-Caryophyllene (MW 204) (wt%)	Sum of MW 206 Isomers (wt%)	Sum of MW 208 Isomers (wt%)	Sum of MW 210 Isomers (Humulane) (wt%)
0.00	84.50	7.82	0.21	0.02	0.07
0.17	1.56	1.91	75.13	5.53	0.79
0.50	0.08	0.20	41.73	39.06	5.75
1.10	0.04	0.02	10.89	68.03	9.69
1.50	0.03	0.02	3.16	75.48	10.51
2.10	0.02	0.02	0.83	80.06	11.92
2.50	0.02	0.02	0.60	81.42	11.75
3.00	0.02	0.02	0.56	81.48	11.66

Example 4: Hydrogenation of caryophyllene in catnip oil

A 50 mL stirred batch autoclave reactor was charged with 34.6 g of a 50 wt% solution of catnip oil in absolute ethanol and 1.73 g of ESCAT 142. The catnip oil was obtained by steam distillation of catnip plants (George Thacker Sons Farm, Alberta, Canada, 2003), and this as-received oil was azeotropically dried with isopropanol. The reactor was equipped with a magnetically-coupled gas entrainment agitator which was rotated at about 1000 rpm during the reaction. The reactor was sealed and then chilled to 15 degrees C via an external cooling coil supplied with a chilled propylene glycol/water mixture from a recirculating bath. The reactor was flushed and evacuated with nitrogen several times to remove oxygen, and these flushes were followed by two rapid flushes with hydrogen to minimize residual nitrogen in the reactor. The reactor was charged to 86 bar (8.6 MPa) pressure with hydrogen. Hydrogen was continuously fed to the reactor during the course of the run to maintain this pressure as hydrogen was consumed by reaction.

Small samples of the reaction mixture were removed periodically via a dip tube equipped with a 2 micron sintered metal frit. The reactor was maintained at an average temperature of 15 degrees C for a reaction time of 4 hours, then the reactor was heated to approximately 100 degrees C via an external band heater for an additional 2 hours. The reactor was then thermally quenched via the external cooling coil and vented. Product analysis by gas chromatography (GC-FID) using 1,2-dibromobenzene as an

internal standard added post reaction showed 97.6% conversion of both caryophyllene isomers after 1.75 hours at 15 degrees C, and quantitative conversion by the end of the reaction at 100 degrees C. Additional analysis by mass spectroscopy (GC-MS) identified the anticipated hydrogenation product isomers for both the alpha- and beta-caryophyllene (humulane and caryophyllane, respectively). Concentrations of the pertinent reactants and reaction products are summarized in Table 2. This example demonstrates complete and rapid hydrogenation of alpha- and beta-caryophyllene as minor components of a natural product.

Table 2

Time on Stream (hr)	Reactor Temp. (°C)	Beta-Caryophyllene (MW 204) (wt%)	Alpha-Caryophyllene (MW 204) (wt%)	Sum of MW 208 Isomers (Caryophyllane) (wt%)	Sum of MW 210 Isomers (Humulane) (wt%)
0.00	14.0	7.70	1.29	-	-
0.17	15.4	4.40	0.80	0.17	0.20
0.50	15.3	0.19	0.18	0.30	0.46
1.75	15.0	0.20	0.02	0.71	0.13
2.75	15.1	0.19	0.03	0.78	0.13
4.00	15.2	0.20	0.03	0.94	0.29
4.60	94.2	0.15	0.00	4.36	0.51
5.85	100.5	0.05	0.00	6.44	1.08

Where a range of numerical values is recited herein, the range includes the endpoints thereof and all the individual integers and fractions within the range, and also includes each of the narrower ranges therein formed
5 by all the various possible combinations of those endpoints and internal integers and fractions to form subgroups of the larger group of values within the stated range to the same extent as if each of those narrower ranges was explicitly recited. Where a range of
10 numerical values is stated herein as being greater than a stated value, the range is nevertheless finite and is bounded on its upper end by a value that is operable within the context of the invention as described herein. Where a range of numerical values is stated herein as
15 being less than a stated value, the range is nevertheless bounded on its lower end by a non-zero value.

In this specification, unless explicitly stated otherwise or indicated to the contrary by the context of
20 usage, amounts, sizes, ranges, formulations, parameters, and other quantities and characteristics recited herein, particularly when modified by the term "about", may but need not be exact, and may also be approximate and/or larger or smaller (as desired) than stated, reflecting
25 tolerances, conversion factors, rounding off, measurement error and the like, as well as the inclusion within a stated value of those values outside it that have, within the context of this invention, functional and/or operable equivalence to the stated value.

30

In this specification, unless explicitly stated otherwise or indicated to the contrary by the context of usage, where an embodiment of the subject matter hereof is stated or described as comprising, including, containing, 5 having, being composed of or being constituted by or of certain features or elements, one or more features or elements in addition to those explicitly stated or described may be present in the embodiment. An alternative embodiment of the subject matter hereof, 10 however, may be stated or described as consisting essentially of certain features or elements, in which embodiment features or elements that would materially alter the principle of operation or the distinguishing characteristics of the embodiment are not present therein. 15 A further alternative embodiment of the subject matter hereof may be stated or described as consisting of certain features or elements, in which embodiment, or in insubstantial variations thereof, only the features or elements specifically stated or described are present.

20

A catalyst suitable for use herein may be selected as any one or more or all of the members of the whole population of catalysts described by name or structure above. A suitable catalyst may, however, also 25 be selected as any one or more or all of the members of a subgroup of the whole population, where the subgroup may be any size (1, 2, 4 or 6, for example), and where the subgroup is formed by omitting any one or more of the members of the whole population as described above. As a 30 result, the catalyst may in such instance not only be selected as one or more or all of the members of any

subgroup of any size that may be formed from the whole population of catalysts as described above, but the catalyst may also be selected in the absence of the members that have been omitted from the whole population
5 to form the subgroup. For example, in certain embodiments, the catalyst useful herein may be selected as one or more or all of the members of a subgroup of catalysts that excludes from the whole population ruthenium supported on titania, with or without the
10 exclusion from the whole population of other catalysts too.

15

CLAIMS

What is claimed is:

1. A composition of matter comprising
 - 5 (a) dihydro- β -caryophyllene and at least one of α -humulane or β -caryophyllane;
 - (b) a dihydronepetalactone or a derivative thereof; and
 - (c) a sunscreensing or sunblocking agent.
2. The composition according to Claim 1 wherein a dihydronepetalactone
 - 10 comprises 1*S*,9*S*,5*R*,6*R*-5,9-dimethyl-3-oxabicyclo[4.3.0]nonan-2-one.
3. The composition according to Claim 1 or 2 that is fabricated in the form of a sprayable liquid, an aerosol, a foam, a cream, an ointment, a gel, a paste, a powder or a friable solid.
 - 15
4. The composition according to any one of Claims 1 to 3 further comprising one or more of adjuvants consisting of thickeners, chelating agents, preservatives, stabilizers, or surfactants.
5. The composition according to Claim 1 or 2 that further comprises one or more
 - 20 of adjuvants consisting of thickeners, chelating agents, preservatives, stabilizers, or surfactants, and that is formulated as a lotion.
6. A method for imparting fragrance to, or revitalizing, the skin, hide, hair, fur,
 - 25 feathers or other surface of a human or a domesticated animal comprising applying thereto a composition of matter according to any one of Claims 1 to 5.
7. An article that comprises
 - (a) dihydro- β -caryophyllene and at least one of α -humulane or β -caryophyllane;
 - 30 and
 - (b) a dihydronepetalactone or a derivative thereof; and

that is fabricated as a textile, fibrous goods, clothing, sanitary goods, carpeting, linens, outdoor equipment, military equipment, candles, paper, paint, ink, wood products, plastics, or polymers.

5 8. The article according to Claim 7 that is fabricated as mosquito netting.

 9. The article according to Claim 7 or 8 wherein a dihydronepetalactone comprises 1*S*,9*S*,5*R*,6*R*-5,9-dimethyl-3-oxabicyclo[4.3.0]nonan-2-one.