Innovation, Sciences et Développement économique Canada
Office de la Propriété Intellectuelle du Canada

Innovation, Science and Economic Development Canada
Canadian Intellectual Property Office

CA 2916481 C 2018/05/15

(11)(21) **2 916 481**

(12) BREVET CANADIEN / CANADIAN PATENT

(13) **C**

(22) **Date de dépôt/Filing Date:** 2009/12/14

(41) **Mise à la disp. pub./Open to Public Insp.:** 2010/06/24

(45) **Date de délivrance/Issue Date:** 2018/05/15

(62) **Demande originale/Original Application:** 2 744 624

(30) **Priorité/Priority:** 2008/12/16 (EP08021835.7)

(51) **Cl.Int./Int.Cl.** ***C07K 16/22*** (2006.01), ***A61K 39/395*** (2006.01), ***A61P 35/00*** (2006.01), ***A61P 9/10*** (2006.01), ***C12N 15/13*** (2006.01), ***C12N 15/63*** (2006.01)

(72) **Inventeurs/Inventors:**
BRINKMANN, ULRICH, DE;
GRIEP, REMKO ALBERT, NO;
KALUZA, KLAUS, DE;
KLEIN, CHRISTIAN, CH;
REGULA, JOERG THOMAS, DE;
SCHEUER, WERNER, DE;
KAVLIE, ANITA, NO

(73) **Propriétaire/Owner:**

(54) **Titre : ANTICORPS DIRIGES CONTRE L'ANGIOPOIETINE 2 HUMAINE**
(54) **Title: ANTIBODIES AGAINST HUMAN ANGIOPOIETIN 2**

(57) **Abrégé/Abstract:**
The present invention relates to antibodies against human Angiopoietin-2 (anti- ANG-2 antibodies), methods for their production, pharmaceutical compositions containing said antibodies, and uses thereof.

CA 2916481 C 2018/05/15

(11)(21) **2 916 481**

(13) **C**

(73) **Propriétaires(suite)/Owners(continued):**F. HOFFMANN-LA ROCHE AG, CH

(74) **Agent:** GOWLING WLG (CANADA) LLP

Abstract

The present invention relates to antibodies against human Angiopoietin-2 (anti- ANG-2 antibodies), methods for their production, pharmaceutical compositions containing said antibodies, and uses thereof.

Antibodies against human Angiopoietin 2

The present invention relates to antibodies against human Angiopoietin 2 (anti-ANG-2 antibodies), methods for their production, pharmaceutical compositions containing said antibodies, and uses thereof.

Background of the Invention

Angiogenesis is implicated in the pathogenesis of a variety of disorders which include solid tumors, intraocular neovascular syndromes such as proliferative retinopathies or age-related macular degeneration (AMD), rheumatoid arthritis, and psoriasis (Folkman, J., et al., J. Biol. Chem. 267 (1992) 10931-10934; Klagsbrun, M., et al., Annu. Rev. Physiol. 53 (1991) 217-239; and Garner, A., Vascular diseases, In: Pathobiology of ocular disease, A dynamic approach, Garner, A., and Klintworth, G. K. (eds.), 2nd edition, Marcel Dekker, New York (1994), pp 1625-1710). In the case of solid tumors, the neovascularization allows the tumor cells to acquire a growth advantage and proliferative autonomy compared to the normal cells. Accordingly, a correlation has been observed between density of microvessels in tumor sections and patient survival in breast cancer as well as in several other tumors (Weidner, N., et al., N. Engl. J. Med. 324 (1991) 1-8; Horak, E.R., et al., Lancet 340 (1992) 1120-1124; and Macchiarini, P., et al., Lancet 340 (1992) 145-146).

ANG-2 and anti-ANG-2 antibodies

Human angiopoietin-2 (ANG-2) (alternatively abbreviated with ANGPT2 or ANG2) (SEQ ID No: 107) is described in Maisonpierre, P.C., et al., Science 277 (1997) 55-60 and Cheung, A.H., et al, Genomics 48 (1998) 389-91. The angiopoietins-1 and -2 (ANG-1(SEQ ID No: 108) and ANG-2 (SEQ ID No: 107) were discovered as ligands for the Ties, a family of tyrosine kinases that is selectively expressed within the vascular endothelium. Yancopoulos, G.D., et al., Nature 407 (2000) 242-48. There are now four definitive members of the angiopoietin family. Angiopoietin-3 and -4 (Ang-3 and Ang-4) may represent widely diverged counterparts of the same gene locus in mouse and man. Kim, I., et al., FEBS Let, 443 (1999) 353-56; Kim, I., et al., J Biol Chem 274 (1999) 26523-28. ANG-1 and ANG-2 were originally identified in tissue culture experiments as agonist and antagonist, respectively (see for ANG-1: Davies, S., et al., Cell, 87 (1996) 1161-1169; and for ANG-2: Maisonpierre, P.C., et al., Science

277 (1997) 55-60). All of the known angiopoietins bind primarily to Tie2, and both Ang-1 and -2 bind to Tie2 with an affinity of 3 nM (Kd). Maisonpierre, P.C., et al., Science 277 (1997) 55-60. Ang-1 was shown to support EC survival and to promote endothelium integrity, Davis, S., et al., Cell, 87 (1996) 1161-1169; Kwak, H.J., et al., FEBS Lett 448 (1999) 249-53; Suri, C., et al., Science 282 (1998) 468-71; Thurston, G., et al., Science 286 (1999) 2511-14; Thurston, G., et al., Nat. Med. 6 (2000) 460-63, whereas ANG-2 had the opposite effect and promoted blood vessel destabilization and regression in the absence of the survival factors VEGF or basic fibroblast growth factor. Maisonpierre, P.C., et al., Science 277 (1997) 55-60. However, many studies of ANG-2 function have suggested a more complex situation. ANG-2 might be a complex regulator of vascular remodeling that plays a role in both vessel sprouting and vessel regression. Supporting such roles for ANG-2, expression analyses reveal that ANG-2 is rapidly induced, together with VEGF, in adult settings of angiogenic sprouting, whereas ANG-2 is induced in the absence of VEGF in settings of vascular regression. Holash, J., et al., Science 284 (1999) 1994-98; Holash, J., et al., Oncogene 18 (1999) 5356-62. Consistent with a context-dependent role, ANG-2 specifically binds to the same endothelial-specific receptor, Tie-2, which is activated by Ang-1, but has context-dependent effects on its activation. Maisonpierre, P.C., et al., Science 277 (1997) 55-60.

Corneal angiogenesis assays have shown that both ANG-1 and ANG-2 had similar effects, acting synergistically with VEGF to promote growth of new blood vessels. Asahara, T., et al., Circ. Res., 83, (1998) 233-40. The possibility that there was a dose-dependent endothelial response was raised by the observation that in vitro at high concentration, ANG-2 can also be pro-angiogenic. Kim, I., et al., Oncogene 19 (2000) 4549-52. At high concentration, ANG-2 acts as an apoptosis survival factor for endothelial cells during serum deprivation apoptosis through activation of Tie2 via PI-3 Kinase and Akt pathway. Kim, I., et al., Oncogene 19 (2000) 4549-52.

Other in vitro experiments suggested that during sustained exposure, the effects of ANG-2 may progressively shift from that of an antagonist to an agonist of Tie2, and at later time points, it may contribute directly to vascular tube formation and neovessel stabilization. Teichert-Kuliszewska, K., et al., Cardiovasc. Res. 49 (2001) 659-70. Furthermore, if ECs were cultivated on fibrin gel, activation of Tie2 with ANG-2 was also observed, perhaps suggesting that the action of ANG-2 could

depend on EC differentiation state. Teichert-Kuliszewska; K., et al., Cardiovasc. Res. 49 (2001) 659-70. In microvascular EC cultured in a three-dimensional collagen gel, ANG-2 can also induce Tie2 activation and promote formation of capillary-like structures. Mochizuki, Y., et al., J. Cell. Sci. 115 (2002) 175-83. Use of a 3-D spheroidal coculture as an in-vitro model of vessel maturation demonstrated that direct contact between ECs and mesenchymal cells abrogates responsiveness to VEGF, whereas the presence of VEGF and ANG-2 induced sprouting. Korff, T., et al., Faseb J. 15 (2001) 447-57. Etoh, T., et al. demonstrated that ECs that constitutively express Tie2, the expression of MMP-1, -9 and u-PA were strongly upregulated by ANG-2 in the presence of VEGF. Etoh, T., et al., Cancer Res. 61 (2001) 2145-53. With an in vivo pupillary membrane model, Lobov, I.B., et al. showed that ANG-2 in the presence of endogenous VEGF promotes a rapid increase in capillary diameter, remodeling of the basal lamina, proliferation and migration of endothelial cells, and stimulates sprouting of new blood vessels. Lobov, I.B., et al., Proc. Natl. Acad. Sci. USA 99 (2002) 11205-10. By contrast, ANG-2 promotes endothelial cell death and vessel regression without endogenous VEGF. Lobov, I.B., et al., Proc. Natl. Acad. Sci. USA 99 (2002) 11205-10. Similarly, with an in vivo tumor model, Vajkoczy, P., et al. demonstrated that multicellular aggregates initiate vascular growth by angiogenic sprouting via the simultaneous expression of VEGFR-2 and ANG-2 by host and tumor endothelium. Vajkoczy, P., et al., J. Clin. Invest. 109 (2002) 777-85. This model illustrated that the established microvasculature of growing tumors is characterized by a continuous remodeling, putatively mediated by the expression of VEGF and ANG-2. Vajkoczy, M.A., et al., J Clin. Invest. 09 (2002) 777-85.

Knock-out mouse studies of Tie-2 and Angiopoietin-1 show similar phenotypes and suggest that Angiopoietin-1 stimulated Tie-2 phosphorylation mediates remodeling and stabilization of developing vessel, promoting blood vessel maturation during angiogenesis and maintenance of endothelial cell-support cell adhesion (Dumont, D.J., et al., Genes & Development, 8 (1994) 1897-1909; Sato, T.N., Nature, 376 (1995) 70-74; (Thurston, G., et al., Nature Medicine 6 (2000) 460-463). The role of Angiopoietin-1 is thought to be conserved in the adult, where it is expressed widely and constitutively (Hanahan, D., Science, 277 (1997) 48-50; Zagzag, D., et al., Exp Neurology, 159 (1999) 391-400). In contrast, Angiopoietin-2 expression is primarily limited to sites of vascular remodeling where it is thought to block the constitutive stabilizing or maturing function of Angiopoietin-1, allowing vessels to revert to, and remain in, a plastic state which

may be more responsive to sprouting signals (Hanahan, D., 1997; Holash, J., et al., Orzcogerze 18 (1999) 5356-62; Maisonpierre, P.C., 1997). Studies of Angiopoietin-2 expression in pathological angiogenesis have found many tumor types to show vascular Angiopoietin-2 expression (Maisonpierre, P.C., et al., Science 277 (1997) 55-60). Functional studies suggest Angiopoietin-2 is involved in tumor angiogenesis and associate Angiopoietin-2 overexpression with increased tumor growth in a mouse xenograft model (Ahmad, S.A., et al., Cancer Res., 61 (2001)1255-1259). Other studies have associated Angiopoietin-2 overexpression with tumor hypervascularity (Etoh, T., et al., Cancer Res. 61 (2001) 2145-53; Tanaka, F., et al., Cancer Res. 62 (2002) 7124-29).

In recent years Angiopoietin-1, Angiopoietin-2 and/or Tie-2 have been proposed as possible anti-cancer therapeutic targets. For example US 6,166,185, US 5,650,490 and US 5,814,464 each disclose anti-Tie-2 ligand and receptor antibodies. Studies using soluble Tie-2 were reported to decrease the number and size of tumors in rodents (Lin, P, 1997; Lin, P., 1998). Siemester, G., et al. (1999) generated human melanoma cell lines expressing the extracellular domain of Tie-2, injected these into nude mice and reported soluble Tie-2 to result in significant inhibition of tumor growth and tumor angiogenesis. Given both Angiopoietin-1 and Angiopoietin-2 bind to Tie-2, it is unclear from these studies whether Angiopoietin-1, Angiopoietin-2 or Tie-2 would be an attractive target for anti-cancer therapy. However, effective anti-Angiopoietin-2 therapy is thought to be of benefit in treating diseases such as cancer, in which progression is dependant on aberrant angiogenesis where blocking the process can lead to prevention of disease advancement (Folkman, J., Nature Medicine. 1, (1995) 27-31.

In addition some groups have reported the use of antibodies and peptides that bind to Angiopoietin-2. See, for example, US 6,166,185 and US 2003/10124129. WO 03/030833, WO 2006/068953, WO 03/057134 or US 2006/0122370.

Study of the effect of focal expression of Angiopoietin-2 has shown that antagonizing the Angiopoietin-1/Tie-2 signal loosens the tight vascular structure thereby exposing ECs to activating signals from angiogenesis inducers, e.g. VEGF (Hanahan, 1997). This pro-angiogenic effect resulting from inhibition of Angiopoietin-1 indicates that anti-Angiopoietin-1 therapy would not be an effective anti-cancer treatment.

ANG-2 is expressed during development at sites where blood vessel remodeling is occurring. Maisonpierre, P.C., et al., Science 277 (1997) 55-60. In adult individuals, ANG-2 expression is restricted to sites of vascular remodeling as well as in highly vascularized tumors, including glioma, Osada, H., et al., Int. J. Oncol. 18 (2001) 305-09; Koga, K., et al., Cancer Res. 61 (2001) 6248-54, hepatocellular carcinoma, Tanaka, S., et al, J. Clin. Invest. 103 (1999) 341-45, gastric carcinoma, Etoh, T., et al., Cancer Res. 61 (2001) 2145-53; Lee, J.H., et al, Int. J. Oncol. 18 (2001) 355-61, thyroid tumor, Bunone, G., et al., Am J Pathol 155 (1999) 1967-76, non-small cell lung cancer, Wong, M.P., et al., Lung Cancer 29 (2000) 11-22, and cancer of colon, Ahmad, S.A., et al., Cancer 92 (2001) 1138-43, and prostate Wurmbach, J.H., et al., Anticancer Res. 20 (2000) 5217-20. Some tumor cells are found to express ANG-2. For example, Tanaka, S., et al., J. Clin. Invest. 103 (1999) 341-45 detected ANG-2 mRNA in 10 out of 12 specimens of human hepatocellular carcinoma (HCC). Ellis' group reported that ANG-2 is expressed ubiquitously in tumor epithelium. Ahmad, S.A., et al., Cancer 92 (2001) 1138-43. Other investigators reported similar findings. Chen, L., et al., J. Tongji Med. Univ. 21 (2001) 228-30, 235 (2001). By detecting ANG-2 mRNA levels in archived human breast cancer specimens, Sfilogoi, C,. et al., Int. J. Cancer 103 (2003) 466-74 reported that ANG-2 mRNA is significantly associated with auxiliary lymph node invasion, short disease-free time and poor overall survival. Tanaka, F., et al., Cancer Res. 62 (2002) 7124-29 reviewed a total of 236 patients of non-small cell lung cancer (NSCLC) with pathological stage-I to -IIIA, respectively. Using immunohistochemistry, they found that 16.9% of the NSCLC patients were ANG-2 positive. The microvessel density for ANG-2 positive tumor is significantly higher than that of ANG-2 negative. Such an angiogenic effect of ANG-2 was seen only when VEGF expression was high. Moreover, positive expression of ANG-2 was a significant factor to predict a poor postoperative survival. Tanaka, F., et al., Cancer Res. 62 (2002) 7124-29. However, they found no significant correlation between Ang-1 expression and the microvessel density. Tanaka, F., et al., Cancer Res. 62 (2002) 7124-29. These results suggest that ANG-2 is an indicator of poor prognosis patients with several types of cancer.

Recently, using an ANG-2 knockout mouse model, Yancopoulos' group reported that ANG-2 is required for postnatal angiogenesis. Gale, N.W., et al., Dev. Cell 3 (2002) 411-23. They showed that the developmentally programmed regression of the hyaloid vasculature in the eye does not occur in the ANG-2 knockout mice and their retinal blood vessels fail to sprout out from the central retinal artery. Gale,

N.W., et al., Dev. Cell 3 (2002) 411-23. They also found that deletion of ANG-2 results in profound defects in the patterning and function of the lymphatic vasculature. Gale, N.W., et al., Dev. Cell 3 (2002) 411-23. Genetic rescue with Ang-1 corrects the lymphatic, but not the angiogenesis defects. Gale, N.W., et al., Dev. Cell 3 (2002) 411-23.

Peters and his colleagues reported that soluble Tie2, when delivered either as recombinant protein or in a viral expression vector, inhibited in vivo growth of murine mammary carcinoma and melanoma in mouse models. Lin, P., et al., Proc. Natl. Acad. Sci. USA 95 (1998) 8829-34; Lin , P., et al., J. Clin. Invest. 100 (1997) 2072-78. Vascular densities in the tumor tissues so treated were greatly reduced. In addition, soluble Tie2 blocked angiogenesis in the rat corneal stimulated by tumor cell conditioned media. Lin, P., et al., J. Clin. Invest. 100 (1997) 2072-78. Furthermore, Isner and his team demonstrated that addition of ANG-2 to VEGF promoted significantly longer and more circumferential neovascularity than VEGF alone. Asahara, T., et al., Circ. Res., 83 (1998) 233-40. Excess soluble Tie2 receptor precluded modulation of VEGF-induced neovascularization by ANG-2. Asahara, T., et al., Circ. Res., 83, (1998) 233-40. Siemeister, G., et al., Cancer Res. 59 (1999) 3185-91 showed with nude mouse xenografts that overexpression of the extracellular ligand-binding domains of either Flt-1 or Tie2 in the xenografts results in significant inhibition of pathway could not be compensated by the other one, suggesting that the VEGF receptor pathway and the Tie2 pathway should be considered as two independent mediators essential for the process of in vivo angiogenesis. Siemeister, G., et al., Cancer Res. 59 (1999) 3185-91. This is proven by a more recent publication by White, R.R., et al., Proc. Natl. Acad. Sci. USA 100 (2003) 5028-33. In their study, it was demonstrated that a nuclease-resistant RNA aptamer that specifically binds and inhibits ANG-2 significantly inhibited neovascularization induced by bFGF in the rat corneal micropocket angiogenesis model.

Summary of the Invention

The invention comprises an antibody specifically binding to human angiopoietin-2 (ANG-2), characterized in comprising as heavy chain variable domain CDR3 region a CDR3 region of SEQ ID NO: 1, SEQ ID NO: 9, SEQ ID NO: 17, SEQ ID NO: 25, SEQ ID NO: 33, SEQ ID NO: 41, or SEQ ID NO: 49.

Preferably the antibody is characterized in that

a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, SEQ ID NO: 9, SEQ ID NO: 17, SEQ ID NO: 25, SEQ ID NO: 33, SEQ ID NO: 41, or SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 2, SEQ ID NO: 10, SEQ ID NO: 18, SEQ ID NO: 26, SEQ ID NO: 34, SEQ ID NO: 42, or SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 3, SEQ ID NO: 11, SEQ ID NO: 19, SEQ ID NO: 27, SEQ ID NO: 35, SEQ ID NO: 43, or SEQ ID NO: 51, and

b) the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, SEQ ID NO: 12, SEQ ID NO: 20, SEQ ID NO: 28, SEQ ID NO: 36, SEQ ID NO: 44, or SEQ ID NO: 52, a CDR2 region of SEQ ID NO: 5, SEQ ID NO: 13, SEQ ID NO: 21, SEQ ID NO: 29, SEQ ID NO: 37, SEQ ID NO: 45, or SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 6, SEQ ID NO: 14, SEQ ID NO: 22, SEQ ID NO: 30, SEQ ID NO: 38, SEQ ID NO: 46, or SEQ ID NO: 54.

Preferably the antibody is characterized in comprising

a) the heavy chain variable domain of SEQ ID NO: 7, SEQ ID NO: 15, SEQ ID NO: 23, SEQ ID NO: 31, SEQ ID NO: 39, SEQ ID NO: 47, or SEQ ID NO: 55; and

b) the light chain variable domain of SEQ ID NO: 8, SEQ ID NO: 16, SEQ ID NO: 24, SEQ ID NO: 32, SEQ ID NO: 40, SEQ ID NO: 48, or SEQ ID NO: 56.

Preferably the antibody is characterized in that the antibody is not specifically binding to Angiopoietin 1 (ANG-1).

A further embodiment of the invention is a pharmaceutical composition comprising an antibody according to the invention.

A further embodiment of the invention is the use of an antibody according to the invention for the manufacture of a pharmaceutical composition.

A further embodiment of the invention is the use of an antibody according to the invention for the prevention of metastasis.

A further embodiment of the invention is the use of an antibody according to the invention for the treatment of cancer.

A further embodiment of the invention is the use of an antibody according to the invention for the treatment of vascular diseases.

A further embodiment of the invention is the use of an antibody according to the invention for the treatment of retinopathy.

A further embodiment of the invention is a nucleic acid encoding a heavy chain variable domain and/or a light chain variable domain of an antibody according to the invention.

The invention further provides expression vectors containing nucleic acid according to the invention capable of expressing said nucleic acid in a prokaryotic or eukaryotic host cell, and host cells containing such vectors for the recombinant production of such an antibody.

The invention further comprises a prokaryotic or eukaryotic host cell comprising a vector according to the invention.

The invention further comprises a method for the production of a recombinant human or humanized antibody according to the invention, characterized by expressing a nucleic acid according to the invention in a prokaryotic or eukaryotic host cell and recovering said antibody from said cell or the cell culture supernatant. The invention further comprises the antibody obtainable by such a recombinant method.

The antibodies according to the invention are especially useful for the prevention of secondary tumors/metastasis or in the treatment of vascular diseases such as retinopathies.

Detailed Description of the Invention

The invention comprises an antibody specifically binding to human angiopoietin-2 (ANG-2), characterized in comprising as heavy chain variable domain CDR3 region a CDR3 region of SEQ ID NO: 1, SEQ ID NO: 9, SEQ ID NO: 17, SEQ ID NO: 25, SEQ ID NO: 33, SEQ ID NO: 41, or SEQ ID NO: 49.

In one embodiment of the invention the antibody is characterized in that

a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, SEQ ID NO: 9, SEQ ID NO: 17, SEQ ID NO: 25, SEQ ID NO: 33, SEQ

ID NO: 41, or SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 2, SEQ ID NO: 10, SEQ ID NO: 18, SEQ ID NO: 26, SEQ ID NO: 34, SEQ ID NO: 42, or SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 3, SEQ ID NO: 11, SEQ ID NO: 19, SEQ ID NO: 27, SEQ ID NO: 35, SEQ ID NO: 43, or SEQ ID NO: 51, and

b) the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, SEQ ID NO: 12, SEQ ID NO: 20, SEQ ID NO: 28, SEQ ID NO: 36, SEQ ID NO: 44, or SEQ ID NO: 52, a CDR2 region of SEQ ID NO: 5, SEQ ID NO: 13, SEQ ID NO: 21, SEQ ID NO: 29, SEQ ID NO: 37, SEQ ID NO: 45, or SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 6, SEQ ID NO: 14, SEQ ID NO: 22, SEQ ID NO: 30, SEQ ID NO: 38, SEQ ID NO: 46, or SEQ ID NO: 54.

Preferably the antibody is characterized in comprising

a) the heavy chain variable domain of SEQ ID NO: 7, SEQ ID NO: 15, SEQ ID NO: 23, SEQ ID NO: 31, SEQ ID NO: 39, SEQ ID NO: 47, or SEQ ID NO: 55; and

b) the light chain variable domain of SEQ ID NO: 8, SEQ ID NO: 16, SEQ ID NO: 24, SEQ ID NO: 32, SEQ ID NO: 40, SEQ ID NO: 48, or SEQ ID NO: 56.

Another embodiment of the invention is an antibody specifically binding to human ANG-2, which is characterized in that the antibody is not specifically binding to human Angiopoietin 1 (ANG-1). Typical antibodies which specifically bind to human ANG-2, but not to human ANG-1 are e.g. Ang2s_R3_LC03, Ang2s_LC09, Ang2i_LC06, Ang2i_LC07, or antibodies binding to the same epitope as Ang2s_R3_LC03, Ang2s_LC09, Ang2i_LC06, Ang2i_LC07, Ang2i_LC10, preferably binding to the same epitope as Ang2i_LC06. Therefore in one embodiment of the invention the antibody specifically binding to human angiopoietin-2 (ANG-2) but not to human ANG-1 binds to the same epitope as Ang2s_R3_LC03, Ang2s_LC09, Ang2i_LC06, Ang2i_LC07, Ang2i_LC10, preferably to the same epitope as Ang2i_LC06. Such antibodies specifically binding to ANG-2, but not to ANG-1 can have improved properties such as efficacy, less toxicity, pharmacokinetic properties compared to ANG-2 and ANG-1 specific antibodies.

Therefore in one embodiment of the invention the antibody specifically binding to human angiopoietin-2 (ANG-2) but not to human ANG-1 is characterized in that

a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, SEQ ID NO: 9, SEQ ID NO: 25, SEQ ID NO: 33, or SEQ ID NO: 49, a CDR2 region of SEQ ID NO: 2, SEQ ID NO: 10, SEQ ID NO: 26, SEQ ID NO: 34, or SEQ ID NO: 50, and a CDR1 region of SEQ ID NO: 3, SEQ ID NO: 11, SEQ ID NO: 27, SEQ ID NO: 35, or SEQ ID NO: 51, and

b) the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, SEQ ID NO: 12, SEQ ID NO: 28, SEQ ID NO: 36, or SEQ ID NO: 52, a CDR2 region of SEQ ID NO: 5, SEQ ID NO: 13, SEQ ID NO: 29, SEQ ID NO: 37, or SEQ ID NO: 53, and a CDR1 region of SEQ ID NO: 6, SEQ ID NO: 14, SEQ ID NO: 30, SEQ ID NO: 38, or SEQ ID NO: 54.

Preferably such antibody specifically binding to human angiopoietin-2 (ANG-2) but not to human ANG-1 is characterized in comprising

a) the heavy chain variable domain of SEQ ID NO: 7, SEQ ID NO: 15, SEQ ID NO: 31, SEQ ID NO: 39, or SEQ ID NO: 55; and

b) the light chain variable domain of SEQ ID NO: 8, SEQ ID NO: 16, SEQ ID NO: 32, SEQ ID NO: 40, or SEQ ID NO: 56.

In one embodiment said antibody according to the invention is characterized in that

a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, or SEQ ID NO: 9, a CDR2 region of SEQ ID NO: 2, or SEQ ID NO: 10, and a CDR1 region of SEQ ID NO: 3, or SEQ ID NO: 11, and

b) the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, or SEQ ID NO: 12, a CDR2 region of SEQ ID NO: 5, or SEQ ID NO: 13, and a CDR1 region of SEQ ID NO: 6, or SEQ ID NO: 14.

In one embodiment said antibody according to the invention is characterized in comprising

a) the heavy chain variable domain of SEQ ID NO: 7, or SEQ ID NO: 15; and

b) the light chain variable domain of SEQ ID NO: 8, or SEQ ID NO: 16.

In one embodiment said antibody according to the invention is characterized in that

a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 1, a CDR2 region of SEQ ID NO: 2, and a CDR1 region of SEQ ID NO: 3, and

b) the light chain variable domain comprises a CDR3 region of SEQ ID NO: 4, a CDR2 region of SEQ ID NO: 5, and a CDR1 region of SEQ ID NO: 6.

In one embodiment said antibody according to the invention is characterized in comprising

a) the heavy chain variable domain of SEQ ID NO: 7; and

b) the light chain variable domain of SEQ ID NO: 8.

In one embodiment said antibody according to the invention is characterized in that

a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 17, a CDR2 region of SEQ ID NO: 18, and a CDR1 region of SEQ ID NO: 19, and

b) the light chain variable domain comprises a CDR3 region of SEQ ID NO: 20, a CDR2 region of SEQ ID NO: 21, and a CDR1 region of SEQ ID NO: 22.

In one embodiment said antibody according to the invention is characterized in comprising

a) the heavy chain variable domain of SEQ ID NO: 23; and

b) the light chain variable domain of SEQ ID NO: 24.

Preferably the antibody according to the invention is characterized in that said antibody is of human IgG1 subclass or is of human IgG4 subclass.

The term "antibody" encompasses the various forms of antibody structures including but not being limited to whole antibodies and antibody fragments, The antibody according to the invention is preferably a humanized antibody, chimeric

antibody, or further genetically engineered antibody,as long as the characteristic properties according to the invention are retained.

"Antibody fragments" comprise a portion of a full length antibody, preferably the variable domain thereof, or at least the antigen binding site thereof. Examples of antibody fragments include diabodies, single-chain antibody molecules (scFv or scFab), and multispecific antibodies (e.g. bispecific) formed from antibody fragments. scFv antibodies are, e.g. described in Houston, J.S., Methods in Enzymol. 203 (1991) 46-88). In addition, antibody fragments comprise single chain polypeptides having the characteristics of a V_H domain, namely being able to assemble together with a V_L domain, or of a V_L domain binding to ANG-2, namely being able to assemble together with a V_H domain to a functional antigen binding site and thereby providing the property. ScFvs can be stabilized using e.g. a) disulfide stabilization (see e.g. in WO 94/029350, Rajagopal, V., et al., Prot. Engin. (1997) 1453-59; Kobayashi, H., et al., Nuclear Medicine & Biology, Vol. 25, (1998) 387-393; or Schmidt, M., et al., Oncogene (1999) 18 1711-1721.) or b) stabilized frameworks (e.g. by specific mutations of the see e.g. WO 2007/109254 specific stabilized frameworks see e.g. US7,258,985, Furrer, F., et al., Invest. Ophthalmol. Vis. Sci. 50 (2009), pp. 771–778 or Ottiger, M., et al., Invest. Ophthalmol. Vis. Sci. 50 (2009), pp. 779–786.

The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of a single amino acid composition.

The term "chimeric antibody" refers to an antibody comprising a variable region, i.e., binding region, from one source or species and at least a portion of a constant region derived from a different source or species, usually prepared by recombinant DNA techniques. Chimeric antibodies comprising a murine variable region and a human constant region are preferred. Other preferred forms of "chimeric antibodies" encompassed by the present invention are those in which the constant region has been modified or changed from that of the original antibody to generate the properties according to the invention, especially in regard to C1q binding and/or Fc receptor (FcR) binding. Such chimeric antibodies are also referred to as "class-switched antibodies.". Chimeric antibodies are the product of expressed immunoglobulin genes comprising DNA segments encoding immunoglobulin variable regions and DNA segments encoding immunoglobulin constant regions. Methods for producing chimeric antibodies involve conventional recombinant

DNA and gene transfection techniques are well known in the art. See e.g. Morrison, S.L., et al., Proc. Natl. Acad. Sci. USA 81 (1984) 6851-6855; US 5,202,238 and US 5,204,244.

The term "humanized antibody" refers to antibodies in which the framework or "complementarity determining regions" (CDR) have been modified to comprise the CDR of an immunoglobulin of different specificity as compared to that of the parent immunoglobulin. In a preferred embodiment, a murine CDR is grafted into the framework region of a human antibody to prepare the "humanized antibody." See e.g. Riechmann, L., et al., Nature 332 (1988) 323-327; and Neuberger, M.S., et al., Nature 314 (1985) 268-270. Particularly preferred CDRs correspond to those representing sequences recognizing the antigens noted above for chimeric antibodies. Other forms of "humanized antibodies" encompassed by the present invention are those in which the constant region has been additionally modified or changed from that of the original antibody to generate the properties according to the invention, especially in regard to C1q binding and/or Fc receptor (FcR) binding.

The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germ line immunoglobulin sequences. Human antibodies are well-known in the state of the art (van Dijk, M.A., and van de Winkel, J.G., Curr. Opin. Chem. Biol. 5 (2001) 368-374). Human antibodies can also be produced in transgenic animals (e.g., mice) that are capable, upon immunization, of producing a full repertoire or a selection of human antibodies in the absence of endogenous immunoglobulin production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge (see, e.g., Jakobovits, A., et al., Proc. Natl. Acad. Sci. USA 90 (1993) 2551-2555; Jakobovits, A., et al., Nature 362 (1993) 255-258; Brueggemann, M., et al., Year Immunol. 7 (1993) 33-40). Human antibodies can also be produced in phage display libraries (Hoogenboom, H.R., and Winter, G., J. Mol. Biol. 227 (1992) 381-388; Marks, J.D., et al., J. Mol. Biol. 222 (1991) 581-597). The techniques of Cole, S.P.C., et al., and Boerner, et al. are also available for the preparation of human monoclonal antibodies (Cole, S.P.C., et al., Monoclonal Antibodies and Cancer Therapy, Liss, A.R., (1985) 77-96; and Boerner, P., et al., J. Immunol. 147 (1991) 86-95). As already mentioned for chimeric and humanized antibodies according to the invention the term "human

antibody" as used herein also comprises such antibodies which are modified in the constant region to generate the properties according to the invention, especially in regard to C1q binding and/or FcR binding, e.g. by "class switching" i.e. change or mutation of Fc parts (e.g. from IgG1 to IgG4 and/or IgG1/IgG4 mutation.)

The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from a host cell such as a NS0 or CHO cell or from an animal (e.g. a mouse) that is transgenic for human immunoglobulin genes or antibodies expressed using a recombinant expression vector transfected into a host cell. Such recombinant human antibodies have variable and constant regions in a rearranged form. The recombinant human antibodies according to the invention have been subjected to in vivo somatic hypermutation. Thus, the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germ line VH and VL sequences, may not naturally exist within the human antibody germ line repertoire in vivo.

The "variable domain" (variable domain of a light chain (V_L), variable domain of a heavy chain (V_H)) as used herein denotes each of the pair of light and heavy chain domains which are involved directly in binding the antibody to the antigen. The variable light and heavy chain domains have the same general structure and each domain comprises four framework (FR) regions whose sequences are widely conserved, connected by three "hypervariable regions" (or complementary determining regions, CDRs). The framework regions adopt a β-sheet conformation and the CDRs may form loops connecting the β-sheet structure. The CDRs in each chain are held in their three-dimensional structure by the framework regions and form together with the CDRs from the other chain the antigen binding site. The antibody's heavy and light chain CDR3 regions play a particularly important role in the binding specificity/affinity of the antibodies according to the invention and therefore provide a further object of the invention.

The term "antigen-binding portion of an antibody" when used herein refer to the amino acid residues of an antibody which are responsible for antigen-binding. The antigen-binding portion of an antibody comprises amino acid residues from the "complementary determining regions" or "CDRs". "Framework" or "FR" regions are those variable domain regions other than the hypervariable region residues as herein defined. Therefore, the light and heavy chain variable domains of an antibody comprise from N- to C-terminus the domains FR1, CDR1, FR2, CDR2,

FR3, CDR3, and FR4. Especially, CDR3 of the heavy chain is the region which contributes most to antigen binding and defines the antibody's properties. CDR and FR regions are determined according to the standard definition of Kabat, E.A., et al., Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD (1991) and/or those residues from a "hypervariable loop."

The terms "nucleic acid" or "nucleic acid molecule", as used herein, are intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA.

The term "amino acid" as used within this application denotes the group of naturally occurring carboxy α-amino acids comprising alanine (three letter code: ala, one letter code: A), arginine (arg, R), asparagine (asn, N), aspartic acid (asp, D), cysteine (cys, C), glutamine (gln, Q), glutamic acid (glu, E), glycine (gly, G), histidine (his, H), isoleucine (ile, I), leucine (leu, L), lysine (lys, K), methionine (met, M), phenylalanine (phe, F), proline (pro, P), serine (ser, S), threonine (thr, T), tryptophan (trp, W), tyrosine (tyr, Y), and valine (val, V).

A nucleic acid is "operable linked" when it is placed into a functional relationship with another nucleic acid. For example, DNA for a presequence or secretory leader is operable linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operable linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operable linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operable linked" means that the DNA sequences being linked are colinear, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

As used herein, the expressions "cell", "cell line", and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to

deliberate or inadvertent mutations. Variant progeny that have the same function or biological activity as screened for in the originally transformed cell are included.

As used herein, the term "binding" or "specifically binding" refers to the binding of the antibody to an epitope of the antigen (ANG-2) in an in vitro assay, preferably in an plasmon resonance assay (BIAcore, GE-Healthcare Uppsala, Sweden) (Example 3) with purified wild-type ANG-2 antigen. The affinity of the binding is defined by the terms ka (rate constant for the association of the antibody from the antibody/antigen complex), k_D (dissociation constant), and K_D (k_D/ka). Binding or specifically binding means a binding affinity (K_D) of 10^{-8} mol/l or less, preferably 10^{-9} M to 10^{-13} mol/l.

Binding of the antibody to the FcγRIII can be investigated by a BIAcore assay (GE-Healthcare Uppsala, Sweden). The affinity of the binding is defined by the terms ka (rate constant for the association of the antibody from the antibody/antigen complex), k_D (dissociation constant), and K_D (k_D/ka).

As used herein, the term "not binding to ANG-1" or "not specifically binding to ANG-1" denotes that the antibody has an EC50-value above 8000 ng/ml in an in vitro ANG-1 binding ELISA assay (according to Example 2).

The term "epitope" includes any polypeptide determinant capable of specific binding to an antibody. In certain embodiments, epitope determinant include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three dimensional structural characteristics, and or specific charge characteristics. An epitope is a region of an antigen that is bound by an antibody.

The "Fc part" of an antibody is not involved directly in binding of an antibody to an antigen, but exhibit various effector functions. A "Fc part of an antibody" is a term well known to the skilled artisan and defined on the basis of papain cleavage of antibodies. Depending on the amino acid sequence of the constant region of their heavy chains, antibodies or immunoglobulins are divided in the classes: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g. IgG1, IgG2, IgG3, and IgG4, IgA1, and IgA2. According to the heavy chain constant regions the different classes of immunoglobulins are called α, δ, ε, γ, and μ,. respectively. The Fc part of an antibody is directly involved in ADCC (antibody-dependent cell-mediated cytotoxicity) and CDC (complement-

dependent cytotoxicity) based on complement activation, C1q binding and Fc receptor binding. Complement activation (CDC) is initiated by binding of complement factor C1q to the Fc part of most IgG antibody subclasses. While the influence of an antibody on the complement system is dependent on certain conditions, binding to C1q is caused by defined binding sites in the Fc part. Such binding sites are known in the state of the art and described e.g. by Boakle, R.J., et al., Nature 282 (1975) 742-743, Lukas, T.J., et al., J. Immunol. 127 (1981) 2555-2560, Brunhouse, R., and Cebra, J.J., Mol. Immunol. 16 (1979) 907-917, Burton, D.R., et al., Nature 288 (1980) 338-344, Thommesen, J.E., et al., Mol. Immunol. 37 (2000) 995-1004, Idusogie, E.E., et al., J. Immunol.164 (2000) 4178-4184, Hezareh, M., et al., J. Virology 75 (2001) 12161-12168, Morgan, A., et al., Immunology 86 (1995) 319-324, EP 0307434. Such binding sites are e.g. L234, L235, D270, N297, E318, K320, K322, P331 and P329 (numbering according to EU index of Kabat, see below). Antibodies of subclass IgG1, IgG2 and IgG3 usually show complement activation and C1q and C3 binding, whereas IgG4 do not activate the complement system and do not bind C1q and C3.

The antibody according to the invention preferably comprises a Fc part from human origin which is Fc part of a human antibody of the subclass IgG1.

The antibody according to the invention is characterized in that the constant chains are of human origin. Such constant chains are well known in the state of the art and e.g. described by Kabat, E.A. (see e.g. Johnson, G. and Wu, T.T., Nucleic Acids Res. 28 (2000) 214-218). For example, a useful human heavy chain constant region comprises an amino acid sequence of SEQ ID NO: 57 or of SEQ ID NO: 58. For example, a useful human light chain constant region comprises an amino acid sequence of a kappa-light chain constant region of SEQ ID NO: 59, or of a lambda-light chain constant region of SEQ ID NO: 60.

The term "constant region" as used within the current applications denotes the sum of the domains of an antibody other than the variable region. The constant region is not involved directly in binding of an antigen, but exhibit various effector functions. Depending on the amino acid sequence of the constant region of their heavy chains, antibodies are divided in the classes: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses, such as IgG1, IgG2, IgG3, and IgG4, IgA1 and IgA2. The heavy chain constant regions that correspond to the different classes of antibodies are called α, δ, ε, γ, and μ, respectively. The

light chain constant regions which can be found in all five antibody classes are called κ (kappa) and λ (lambda).

The term "constant region derived from human origin" as used in the current application denotes a constant heavy chain region of a human antibody of the subclass IgG1, IgG2, IgG3, or IgG4 and/or a constant light chain κ region. Such constant regions are well known in the state of the art and e.g. described by Kabat, E.A., (see e.g. Johnson, G. and Wu, T.T., Nucleic Acids Res. 28 (2000) 214-218; Kabat, E.A., et al., Proc. Natl. Acad. Sci. USA 72 (1975) 2785-2788).

While antibodies of the IgG4 subclass show reduced Fc receptor (FcγRIIIa) binding, antibodies of other IgG subclasses show strong binding. However Pro238, Asp265, Asp270, Asn297 (loss of Fc carbohydrate), Pro329, Leu234, Leu235, Gly236, Gly237, Ile253, Ser254, Lys288, Thr307, Gln311, Asn434, and His435 are residues which, if altered, provide also reduced Fc receptor binding (Shields, R.L., et al., J. Biol. Chem. 276 (2001) 6591-6604; Lund, J., et al., FASEB J. 9 (1995) 115-119; Morgan, A., et al., Immunology 86 (1995) 319-324; EP 0 307 434).

In one embodiment an antibody according to the invention has a reduced FcR binding compared to an IgG1 antibody and the monospecific bivalent parent antibody is in regard to FcR binding of IgG4 subclass or of IgG1 or IgG2 subclass with a mutation in S228, L234, L235 and/or D265, and/ or contains the PVA236 mutation. In one embodiment the mutations in the monospecific bivalent parent antibody are S228P, L234A, L235A, L235E and/or PVA236. In another embodiment the mutations in the monospecific bivalent parent antibody are in IgG4 S228P and in IgG1 L234A and L235A. Constant heavy chain regions shown in SEQ ID NO: 57 and 58. In one embodiment the constant heavy chain region of the monospecific bivalent parent antibody is of SEQ ID NO: 57 with mutations L234A and L235A. In another embodiment the constant heavy chain region of the monospecific bivalent parent antibody is of SEQ ID NO: 58 with mutation S228P. In another embodiment the constant light chain region of the monospecific bivalent parent antibody is a kappa light chain region of SEQ ID NO: 59, or a lambda light chain constant region of SEQ ID NO: 60. In one embodiment of the invention the constant heavy chain region of the monospecific bivalent parent antibody is of SEQ ID NO: 57 or of SEQ ID NO: 58 with mutation S228P.

The constant region of an antibody is directly involved in ADCC (antibody-dependent cell-mediated cytotoxicity) and CDC (complement-dependent

cytotoxicity). Complement activation (CDC) is initiated by binding of complement factor C1q to the constant region of most IgG antibody subclasses. Binding of C1q to an antibody is caused by defined protein-protein interactions at the so called binding site. Such constant region binding sites are known in the state of the art and described e.g. by Lukas, T.J., et al., J. Immunol. 127 (1981) 2555-2560; Brunhouse, R. and Cebra, J.J., Mol. Immunol. 16 (1979) 907-917; Burton, D.R., et al., Nature 288 (1980) 338-344; Thommesen, J.E., et al., Mol. Immunol. 37 (2000) 995-1004; Idusogie, E.E., et al., J. Immunol. 164 (2000) 4178-4184; Hezareh, M., et al., J. Virol. 75 (2001) 12161-12168; Morgan, A., et al., Immunology 86 (1995) 319-324; and EP 0 307 434. Such constant region binding sites are, e.g., characterized by the amino acids L234, L235, D270, N297, E318, K320, K322, P331, and P329 (numbering according to EU index of Kabat).

The term "antibody-dependent cellular cytotoxicity (ADCC)" refers to lysis of human target cells by an antibody according to the invention in the presence of effector cells. ADCC is measured preferably by the treatment of a preparation of CCR5 expressing cells with an antibody according to the invention in the presence of effector cells such as freshly isolated PBMC or purified effector cells from buffy coats, like monocytes or natural killer (NK) cells or a permanently growing NK cell line.

The term "complement-dependent cytotoxicity (CDC)" denotes a process initiated by binding of complement factor C1q to the Fc part of most IgG antibody subclasses. Binding of C1q to an antibody is caused by defined protein-protein interactions at the so called binding site. Such Fc part binding sites are known in the state of the art (see above). Such Fc part binding sites are, e.g., characterized by the amino acids L234, L235, D270, N297, E318, K320, K322, P331, and P329 (numbering according to EU index of Kabat). Antibodies of subclass IgG1, IgG2, and IgG3 usually show complement activation including C1q and C3 binding, whereas IgG4 does not activate the complement system and does not bind C1q and/or C3.

The antibody according to the invention is produced by recombinant means. Thus, one aspect of the current invention is a nucleic acid encoding the antibody according to the invention and a further aspect is a cell comprising said nucleic acid encoding an antibody according to the invention. Methods for recombinant production are widely known in the state of the art and comprise protein expression in prokaryotic and eukaryotic cells with subsequent isolation of the antibody and

usually purification to a pharmaceutically acceptable purity. For the expression of the antibodies as aforementioned in a host cell, nucleic acids encoding the respective modified light and heavy chains are inserted into expression vectors by standard methods. Expression is performed in appropriate prokaryotic or eukaryotic host cells like CHO cells, NS0 cells, SP2/0 cells, HEK293 cells, COS cells, PER.C6 cells, yeast, or E.coli cells, and the antibody is recovered from the cells (supernatant or cells after lysis). General methods for recombinant production of antibodies are well-known in the state of the art and described, for example, in the review articles of Makrides, S.C., Protein Expr. Purif. 17 (1999) 183-202; Geisse, S., et al., Protein Expr. Purif. 8 (1996) 271-282; Kaufman, R.J., Mol. Biotechnol. 16 (2000) 151-161; Werner, R.G., J. Drug Res. 48 (1998) 870-880.

The antibodies according to the invention are suitably separated from the culture medium by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography. DNA and RNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures. The hybridoma cells can serve as a source of such DNA and RNA. Once isolated, the DNA may be inserted into expression vectors, which are then transfected into host cells such as HEK 293 cells, CHO cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of recombinant monoclonal antibodies in the host cells.

Amino acid sequence variants (or mutants) of the antibody according to the invention are prepared by introducing appropriate nucleotide changes into the antibody DNA, or by nucleotide synthesis. Such modifications can be performed, however, only in a very limited range, e.g. as described above. For example, the modifications do not alter the above mentioned antibody characteristics such as the IgG isotype and antigen binding, but may improve the yield of the recombinant production, protein stability or facilitate the purification.

The term "host cell" as used in the current application denotes any kind of cellular system which can be engineered to generate the antibodies according to the current invention. In one embodiment HEK293 cells and CHO cells are used as host cells. As used herein, the expressions "cell," "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also

understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Variant progeny that have the same function or biological activity as screened for in the originally transformed cell are included.

Expression in NS0 cells is described by, e.g., Barnes, L.M., et al., Cytotechnology 32 (2000) 109-123; Barnes, L.M., et al., Biotech. Bioeng. 73 (2001) 261-270. Transient expression is described by, e.g., Durocher, Y., et al., Nucl. Acids. Res. 30 (2002) E9. Cloning of variable domains is described by Orlandi, R., et al., Proc. Natl. Acad. Sci. USA 86 (1989) 3833-3837; Carter, P., et al., Proc. Natl. Acad. Sci. USA 89 (1992) 4285-4289; and Norderhaug, L., et al., J. Immunol. Methods 204 (1997) 77-87. A preferred transient expression system (HEK 293) is described by Schlaeger, E.-J., and Christensen, K., in Cytotechnology 30 (1999) 71-83 and by Schlaeger, E.-J., in J. Immunol. Methods 194 (1996) 191-199.

The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, enhancers and polyadenylation signals.

A nucleic acid is "operably linked" when it is placed in a functional relationship with another nucleic acid sequence. For example, DNA for a pre-sequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a pre-protein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

Purification of antibodies is performed in order to eliminate cellular components or other contaminants, e.g. other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis, and others well known in the art. See Ausubel, F., et al., ed. Current Protocols in Molecular Biology, Greene Publishing and Wiley Interscience, New York (1987). Different methods are well established and widespread used for protein purification, such as affinity chromatography with

microbial proteins (e.g. protein A or protein G affinity chromatography), ion exchange chromatography (e.g. cation exchange (carboxymethyl resins), anion exchange (amino ethyl resins) and mixed-mode exchange), thiophilic adsorption (e.g. with beta-mercaptoethanol and other SH ligands), hydrophobic interaction or aromatic adsorption chromatography (e.g. with phenyl-sepharose, aza-arenophilic resins, or m-aminophenylboronic acid), metal chelate affinity chromatography (e.g. with Ni(II)- and Cu(II)-affinity material), size exclusion chromatography, and electrophoretical methods (such as gel electrophoresis, capillary electrophoresis) (Vijayalakshmi, M.A. Appl. Biochem. Biotech. 75 (1998) 93-102).

The invention comprises a method for the treatment of a patient in need of therapy, characterized by administering to the patient a therapeutically effective amount of an antibody according to the invention.

The invention comprises the use of an antibody according to the invention for therapy.

The invention comprises the use of an antibody according to the invention for the preparation of a medicament for the prevention of metastasis.

The invention comprises the use of an antibody according to the invention for the preparation of a medicament for the treatment of cancer.

One aspect of the invention is a pharmaceutical composition comprising an antibody according to the invention. Another aspect of the invention is the use of an antibody according to the invention for the manufacture of a pharmaceutical composition. A further aspect of the invention is a method for the manufacture of a pharmaceutical composition comprising an antibody according to the invention. In another aspect, the present invention provides a composition, e.g. a pharmaceutical composition, containing an antibody according to the present invention, formulated together with a pharmaceutical carrier.

Another aspect of the invention is said pharmaceutical composition for the prevention of metastasis.

Another aspect of the invention is an antibody according to the invention for the prevention of metastasis.

Another aspect of the invention is the use of an antibody according to the invention for the manufacture of a medicament for the prevention of metastasis.

Another aspect of the invention is a method of prevention metastasis in patient suffering from primary cancer by administering an antibody according to the invention to a patient in the need of such preventative treatment.

We could show higly efficient prevention of spotanouenes metastasis/secondary tumors in vivo in a orthotopic and a subcutanoues cancer model (see Example 9) (in contrast to experimental model where the tumor cells are injected i.v. This is similar to the clinical situation wherein cells disseminate from a primary tumor and metastase to secondary organ like lung or liver (where secondary tumors).

The term "metastasis" according to the invention refers to the transmission of cancerous cells from the primary tumor to one or more sites elsewhere in a patient where then secondary tumors develop. MetastasMeans to determine if a cancer has metastasized are known in the art and include bone scan, chest X-ray, CAT scan, MRI scan, and tumor marker tests.

The term "prevention of metastasis" or "prevention of secondary tumors" as used herein have the same meaning and refers a prophylactic agent against metastasis in patient suffering from relapsed HER2 positive cancer in this way inhibiting or reducing a further transmission of cancerous cells from the primary tumor to one or more sites elsewhere in a patient. This means that the metastasis of the primary, tumor or cancer is prevented, delayed, or reduced and thus the developement of secondary tunmors is is prevented, delayed, or reduced. Preferably the metastasis i.e secondary tumors of the lung are prevented or reduced, which means that metastatic transmission of cancerous cells from the primary tumor to the lung is prevented or reduced.

Another aspect of the invention is said pharmaceutical composition for the treatment of cancer.

Another aspect of the invention is an antibody according to the invention for the treatment of cancer.

Another aspect of the invention is the use of an antibody according to the invention for the manufacture of a medicament for the treatment of cancer.

Another aspect of the invention is method of treatment of patient suffering from cancer by administering an antibody according to the invention to a patient in the need of such treatment.

As used herein, "pharmaceutical carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g. by injection or infusion).

A composition of the present invention can be administered by a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. To administer a compound of the invention by certain routes of administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the compound may be administered to a subject in an appropriate carrier, for example, liposomes, or a diluent. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Pharmaceutical carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intra-arterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

The term cancer as used herein refers to proliferative diseases, such as lymphomas, lymphocytic leukemias, lung cancer, non small cell lung (NSCL) cancer, bronchioloalviolar cell lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the

vagina, carcinoma of the vulva, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, hepatocellular cancer, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwanomas, ependymonas, medulloblastomas, meningiomas, squamous cell carcinomas, pituitary adenoma and Ewings sarcoma, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers.

Another aspect of the invention is said pharmaceutical composition as anti-angiogenic agent. Such anti-angiogenic agent can be used for the treatment of cancer, especially solid tumors, and other vascular diseases.

Another aspect of the invention is the use of an antibody according to the invention for the manufacture of a medicament for the treatment of vascular diseases.

Another aspect of the invention is an antibody according to the invention for the treatment of vascular diseases.

A preferred embodiment is an antibody according to the invention for the treatment of retinopathy.

A preferred embodiment is the use of an antibody according to the invention for the manufacture of a medicament for the treatment of retinopathy

Another aspect of the invention is method of treatment of patient suffering from vascular diseases by administering an antibody according to the invention to a patient in the need of such treatment.

The term "vascular diseases" includes Cancer, Inflammatory diseases, Atherosclerosis, Ischemia, Trauma, Sepsis, COPD, Asthma, Diabetes, AMD, Retinopathy, Stroke, Adipositas, Acute lung injury, Hemorrhage, Vascular leak e.g. Cytokine induced, Allergy, Graves' Disease , Hashimoto's Autoimmune Thyroiditis, Idiopathic Thrombocytopenic Purpura, Giant Cell Arteritis, Rheumatoid Arthritis, Systemic Lupus Erythematosus (SLE), Lupus Nephritis, Crohn's Disease, Multiple Sclerosis, Ulcerative Colitis, especially to solid tumors,

intraocular neovascular syndromes (such as proliferative retinopathies or age-related macular degeneration (AMD)), rheumatoid arthritis, and psoriasis (Folkman, J., et al., J. Biol. Chem. 267 (1992) 10931- 10934; Klagsbrun, M., et al., Annu. Rev. Physiol. 53 (1991) 217-239; and Garner, A., Vascular diseases, In: Pathobiology of ocular disease, A dynamic approach, Garner, A., and Klintworth, G. K. (eds.), 2nd edition, Marcel Dekker, New York (1994), pp 1625-1710).

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of presence of microorganisms may be ensured both by sterilization procedures, supra, and by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

The composition must be sterile and fluid to the extent that the composition is deliverable by syringe. In addition to water, the carrier preferably is an isotonic buffered saline solution.

Proper fluidity can be maintained, for example, by use of coating such as lecithin, by maintenance of required particle size in the case of dispersion and by use of surfactants. In many cases, it is preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol or sorbitol, and sodium chloride in the composition.

As used herein, the expressions "cell," "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Variant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, it will be clear from the context.

The term “transformation” as used herein refers to process of transfer of a vectors/nucleic acid into a host cell. If cells without formidable cell wall barriers are used as host cells, transfection is carried out e.g. by the calcium phosphate precipitation method as described by Graham, F.L., and van der Eb, Virology 52 (1973) 456-467. However, other methods for introducing DNA into cells such as by nuclear injection or by protoplast fusion may also be used. If prokaryotic cells or cells which contain substantial cell wall constructions are used, e.g. one method of transfection is calcium treatment using calcium chloride as described by Cohen, F. N, et al, PNAS. 69 (1972) 7110ff.

As used herein, "expression" refers to the process by which a nucleic acid is transcribed into mRNA and/or to the process by which the transcribed mRNA (also referred to as transcript) is subsequently being translated into peptides, polypeptides, or proteins. The transcripts and the encoded polypeptides are collectively referred to as gene product. If the polynucleotide is derived from genomic DNA, expression in a eukaryotic cell may include splicing of the mRNA.

A "vector" is a nucleic acid molecule, in particular self-replicating, which transfers an inserted nucleic acid molecule into and/or between host cells. The term includes

vectors that function primarily for insertion of DNA or RNA into a cell (e.g., chromosomal integration), replication of vectors that function primarily for the replication of DNA or RNA, and expression vectors that function for transcription and/or translation of the DNA or RNA. Also included are vectors that provide more than one of the functions as described.

An "expression vector" is a polynucleotide which, when introduced into an appropriate host cell, can be transcribed and translated into a polypeptide. An "expression system" usually refers to a suitable host cell comprised of an expression vector that can function to yield a desired expression product.

The following examples, sequence listing and figures are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

Description of the Amino Acid Sequences

SEQ ID NO: 1	heavy chain CDR3, < ANG-2>Ang2i_LC06
SEQ ID NO: 2	heavy chain CDR2, <ANG-2> Ang2i_LC06
SEQ ID NO: 3	heavy chain CDR1, <ANG-2>Ang2i_LC06
SEQ ID NO: 4	light chain CDR3, <ANG-2>Ang2i_LC06
SEQ ID NO: 5	light chain CDR2, <ANG-2>Ang2i_LC06
SEQ ID NO: 6	light chain CDR1, <ANG-2>Ang2i_LC06
SEQ ID NO: 7	heavy chain variable domain, <ANG-2>Ang2i_LC06
SEQ ID NO: 8	light chain variable domain, <ANG-2>Ang2i_LC06
SEQ ID NO: 9	heavy chain CDR3, <ANG-2>Ang2i_LC07
SEQ ID NO: 10	heavy chain CDR2, <ANG-2>Ang2i_LC07
SEQ ID NO: 11	heavy chain CDR1, <ANG-2>Ang2i_LC07
SEQ ID NO: 12	light chain CDR3, <ANG-2>Ang2i_LC07
SEQ ID NO: 13	light chain CDR2, <ANG-2>Ang2i_LC07
SEQ ID NO: 14	light chain CDR1, <ANG-2>Ang2i_LC07
SEQ ID NO: 15	heavy chain variable domain, <ANG-2>Ang2i_LC07
SEQ ID NO: 16	light chain variable domain, <ANG-2>Ang2i_LC07
SEQ ID NO: 17	heavy chain CDR3, < ANG-2>Ang2k_LC08
SEQ ID NO: 18	heavy chain CDR2, <ANG-2> Ang2k_LC08
SEQ ID NO: 19	heavy chain CDR1, <ANG-2> Ang2k_LC08
SEQ ID NO: 20	light chain CDR3, <ANG-2> Ang2k_LC08

SEQ ID NO: 21	light chain CDR2, <ANG-2> Ang2k_LC08
SEQ ID NO: 22	light chain CDR1, <ANG-2> Ang2k_LC08
SEQ ID NO: 23	heavy chain variable domain, <ANG-2> Ang2k_LC08
SEQ ID NO: 24	light chain variable domain, <ANG-2> Ang2k_LC08
SEQ ID NO: 25	heavy chain CDR3, <ANG-2> Ang2s_LC09
SEQ ID NO: 26	heavy chain CDR2, <ANG-2> Ang2s_LC09
SEQ ID NO: 27	heavy chain CDR1, <ANG-2> Ang2s_LC09
SEQ ID NO: 28	light chain CDR3, <ANG-2> Ang2s_LC09
SEQ ID NO: 29	light chain CDR2, <ANG-2> Ang2s_LC09
SEQ ID NO: 30	light chain CDR1, <ANG-2> Ang2s_LC09
SEQ ID NO: 31	heavy chain variable domain, <ANG-2> Ang2s_LC09
SEQ ID NO: 32	light chain variable domain, <ANG-2> Ang2s_LC09
SEQ ID NO: 33	heavy chain CDR3, <ANG-2> Ang2i_LC10
SEQ ID NO: 34	heavy chain CDR2, <ANG-2> Ang2i_LC10
SEQ ID NO: 35	heavy chain CDR1, <ANG-2> Ang2i_LC10
SEQ ID NO: 36	light chain CDR3, <ANG-2> Ang2i_LC10
SEQ ID NO: 37	light chain CDR2, <ANG-2> Ang2i_LC10
SEQ ID NO: 38	light chain CDR1, <ANG-2> Ang2i_LC10
SEQ ID NO: 39	heavy chain variable domain, <ANG-2> Ang2i_LC10
SEQ ID NO: 40	light chain variable domain, <ANG-2> Ang2i_LC10
SEQ ID NO: 41	heavy chain CDR3, <ANG-2> Ang2k_LC11
SEQ ID NO: 42	heavy chain CDR2, <ANG-2> Ang2k_LC11
SEQ ID NO: 43	heavy chain CDR1, <ANG-2> Ang2k_LC11
SEQ ID NO: 44	light chain CDR3, <ANG-2> Ang2k_LC11
SEQ ID NO: 45	light chain CDR2, <ANG-2> Ang2k_LC11
SEQ ID NO: 46	light chain CDR1, <ANG-2> Ang2k_LC11
SEQ ID NO: 47	heavy chain variable domain, <ANG-2> Ang2k_LC11
SEQ ID NO: 48	light chain variable domain, <ANG-2> Ang2k_LC11
SEQ ID NO: 49	heavy chain CDR3, <ANG-2> Ang2s_R3_LC03
SEQ ID NO: 50	heavy chain CDR2, <ANG-2> Ang2s_R3_LC03
SEQ ID NO: 51	heavy chain CDR1, <ANG-2> Ang2s_R3_LC03
SEQ ID NO: 52	light chain CDR3, <ANG-2> Ang2s_R3_LC03
SEQ ID NO: 53	light chain CDR2, <ANG-2> Ang2s_R3_LC03
SEQ ID NO: 54	light chain CDR1, <ANG-2> Ang2s_R3_LC03
SEQ ID NO: 55	heavy chain variable domain, <ANG-2> Ang2s_R3_LC03
SEQ ID NO: 56	light chain variable domain, <ANG-2> Ang2s_R3_LC03
SEQ ID NO: 57	human heavy chain constant region derived from IgG1

SEQ ID NO: 58	human heavy chain constant region derived from IgG4
SEQ ID NO: 59	kappa light chain constant region
SEQ ID NO: 60	lambda light chain constant region
SEQ ID NO: 61	Human Tie-2 receptor
SEQ ID NO: 62	Human angiopoietin-2 (ANG-2) with leader and His-tag
SEQ ID NO: 63	Human angiopoietin-1 (ANG-1) with leader and His-tag

Description of the Figures

Figure 1	Cloning of IgGs for transient expressions into expression vectors transient expressions A) Ang2i-LC06 **(Fig 1A)** B.) Ang2i-LC06 **(Fig 1B)**
Figure 2	SDS-PAGE Gel of purified anti ANG-2 antibodies Ang2i-LC06, Ang2i-LC07 and Ang2k-LC08
Figure 3	Angiopoietin-Tie2 interaction ELISA
Figure 4	Inhibition of ANG-2 binding to Tie2 by Ang2i-LC06 and Ang2k-LC08
Figure 5	Inhibition of ANG-1 binding to Tie2 by Ang2i-LC06 and Ang2k-LC08
Figure 6	Colo205 xenograft model to test in vivo efficacy of anti ANG-2 antibodies
Figure 7	KPL-4 xenograft model to test in vivo efficacy of anti ANG-2 antibodies.
Figure 8	ANG-1 binding via Biacore sensogramm.
Figure 9	Prevention of lung metastasis/secondary tumors by the antibodies according to the invention in primary colon tumor xenograft (9A) and primary breast xenograft (9B)
Figure10	Inhibition of tethinopaty by the antibodies according to the invention.

Experimental Procedure 1

Materials & general methods

General information regarding the nucleotide sequences of human immunoglobulins light and heavy chains is given in: Kabat, E.A., et al., Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD (1991). Amino acids of antibody chains are

numbered and referred to according to EU numbering (Edelman, G.M., et al., Proc. Natl. Acad. Sci. USA 63 (1969) 78-85; Kabat, E.A., et al., Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD, (1991)).

Recombinant DNA techniques

Standard methods were used to manipulate DNA as described in Sambrook, J. et al., Molecular cloning: A laboratory manual; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989. The molecular biological reagents were used according to the manufacturer's instructions.

Gene synthesis

Desired gene segments were prepared from oligonucleotides made by chemical synthesis. The gene segments, which are flanked by singular restriction endonuclease cleavage sites, were assembled by annealing and ligation of oligonucleotides including PCR amplification and subsequently cloned via the indicated restriction sites e.g. KpnI/ SacI or AscI/PacI into a pPCRScript (Stratagene) based pGA4 cloning vector. The DNA sequences of the subcloned gene fragments were confirmed by DNA sequencing. Gene synthesis fragments were ordered according to given specifications at Geneart (Regensburg, Germany).

DNA sequence determination

DNA sequences were determined by double strand sequencing performed at MediGenomix GmbH (Martinsried, Germany) or Sequiserve GmbH (Vaterstetten, Germany).

DNA and protein sequence analysis and sequence data management

The GCG's (Genetics Computer Group, Madison, Wisconsin) software package version 10.2 and Infomax's Vector NT1 Advance suite version 8.0 was used for sequence creation, mapping, analysis, annotation and illustration.

Expression vectors

For the expression of the described antibodies variants of expression plasmids for transient expression (e.g. in HEK293 EBNA or HEK293-F cells) or for stable expression (e.g. in CHO cells) based either on a cDNA organization with a CMV-

Intron A promoter or on a genomic organization with a CMV promoter (e.g. Figure 1) were applied.

Beside the antibody expression cassette the vectors contained:

- an origin of replication which allows replication of this plasmid in E. coli, and
- a ß-lactamase gene which confers ampicillin resistance in E. coli.

The transcription unit of the antibody gene is composed of the following elements:

- unique restriction site(s) at the 5' end
- the immediate early enhancer and promoter from the human cytomegalovirus,
- followed by the Intron A sequence in the case of the cDNA organization,
- a 5'-untranslated region of a human antibody gene,
- a immunoglobulin heavy chain signal sequence,
- the human antibody chain (heavy chain, modified heavy chain or light chain) either as cDNA or as genomic organization with an the immunoglobulin exon-intron organization
- a 3' untranslated region with a polyadenylation signal sequence, and
- unique restriction site(s) at the 3' end.

The fusion genes comprising the heavy chain sequences of the selected antibody as described below were generated by PCR and/or gene synthesis and assembled with known recombinant methods and techniques by connection of the according nucleic acid segments e.g. using unique NsiI and EcoRI sites in the genomic heavy chain vectors. The subcloned nucleic acid sequences were verified by DNA sequencing. For transient and stable transfections larger quantities of the plasmids were prepared by plasmid preparation from transformed E. coli cultures (Nucleobond AX, Macherey-Nagel).

Cell culture techniques

Standard cell culture techniques were used as described in Current Protocols in Cell Biology (2000), Bonifacino, J.S., Dasso, M., Harford, J.B., Lippincott-Schwartz, J. and Yamada, K.M. (eds.), John Wiley & Sons, Inc.

Transient transfections in HEK293-F system

Antibodies were generated by transient transfection of the two plasmids encoding the heavy or modified heavy chain, respectively and the corresponding light chain using the HEK293-F system (Invitrogen) according to the manufacturer's instruction. Briefly, HEK293-F cells (Invitrogen) growing in suspension either in a shake flask or in a stirred fermenter in serumfree FreeStyle 293 expression medium (Invitrogen) were transfected with a mix of the two respective expression plasmids and 293fectin or fectin (Invitrogen). For e.g. 2 L shake flask (Corning) HEK293-F cells were seeded at a density of 1.0E*6 cells/mL in 600 mL and incubated at 120 rpm, 8% CO2. The day after the cells were transfected at a cell density of ca. 1.5E*6 cells/mL with ca. 42 mL mix of A) 20 mL Opti-MEM (Invitrogen) with 600 µg total plasmid DNA (1 µg/mL) encoding the heavy or modified heavy chain, respectively and the corresponding light chain in an equimolar ratio and B) 20 ml Opti-MEM + 1.2 mL 293 fectin or fectin (2 µl/mL). According to the glucose consumption glucose solution was added during the course of the fermentation. The supernatant containing the secreted antibody was harvested after 5-10 days and antibodies were either directly purified from the supernatant or the supernatant was frozen and stored.

Protein determination

The protein concentration of purified antibodies and derivatives was determined by determining the optical density (OD) at 280 nm, using the molar extinction coefficient calculated on the basis of the amino acid sequence according to Pace, C.N., et. al., Protein Science, 4 (1995), 2411-1423.

Antibody concentration determination in supernatants

The concentration of antibodies and derivatives in cell culture supernatants was estimated by immunoprecipitation with Protein A Agarose-beads (Roche). 60 µL Protein A Agarose beads are washed three times in TBS-NP40 (50 mM Tris, pH 7.5, 150 mM NaCl, 1% Nonidet-P40). Subsequently, 1 -15 mL cell culture

supernatant are applied to the Protein A Agarose beads pre-equilibrated in TBS-NP40. After incubation for at 1 h at room temperature the beads are washed on an Ultrafree-MC-filter column (Amicon) once with 0.5 mL TBS-NP40, twice with 0.5 mL 2x phosphate buffered saline (2xPBS, Roche) and briefly four times with 0.5 mL 100 mM Na-citrate pH 5,0. Bound antibody is eluted by addition of 35 µl NuPAGE® LDS Sample Buffer (Invitrogen). Half of the sample is combined with NuPAGE® Sample Reducing Agent or left unreduced, respectively, and heated for 10 min at 70°C. Consequently, 20 µl are applied to an 4-12% NuPAGE® Bis-Tris SDS-PAGE (Invitrogen) (with MOPS buffer for non-reduced SDS-PAGE and MES buffer with NuPAGE® Antioxidant running buffer additive (Invitrogen) for reduced SDS-PAGE) and stained with Coomassie Blue.

The concentration of antibodies and derivatives in cell culture supernatants was measured by Protein A-HPLC chromatography. Briefly, cell culture supernatants containing antibodies and derivatives that bind to Protein A were applied to a HiTrap Protein A column (GE Healthcare) in 50 mM K2HPO4, 300 mM NaCl, pH 7.3 and eluted from the matrix with 50 mM acetic acid, pH 2.5 on a Dionex HPLC-System. The eluted protein was quantified by UV absorbance and integration of peak areas. A purified standard IgG1 antibody served as a standard.

Alternatively, the concentration of antibodies and derivatives in cell culture supernatants was measured by Sandwich-IgG-ELISA. Briefly, StreptaWell High Bind Strepatavidin A-96 well microtiter plates (Roche) were coated with 100 µL/well biotinylated anti-human IgG capture molecule F(ab')2<h-Fcγ> BI (Dianova) at 0.1 µg/mL for 1 h at room temperature or alternatively over night at 4°C and subsequently washed three times with 200 µL/well PBS, 0.05% Tween (PBST, Sigma). 100 µL/well of a dilution series in PBS (Sigma) of the respective antibody containing cell culture supernatants was added to the wells and incubated for 1-2 h on a microtiterplate shaker at room temperature. The wells were washed three times with 200 µL/well PBST and bound antibody was detected with 100 µl F(ab')2<hFcgamma>POD (Dianova) at 0.1 µg/mL as detection antibody for 1-2 h on a microtiterplate shaker at room temperature. Unbound detection antibody was washed away three times with 200 µL/well PBST and the bound detection antibody was detected by addition of 100 µL ABTS/well. Determination of absorbance was performed on a Tecan Fluor Spectrometer at a measurement wavelength of 405 nm (reference wavelength 492 nm).

Protein purification

Proteins were purified from filtered cell culture supernatants referring to standard protocols. In brief, antibodies were applied to a Protein A Sepharose column (GE Healthcare) and washed with PBS. Elution of antibodies was achieved at acidic pH followed by immediate neutralization of the sample. Aggregated protein was separated from monomeric antibodies by size exclusion chromatography (Superdex 200, GE Healthcare) in 20 mM Histidine, 140 mM NaCl pH 6.0. Monomeric antibody fractions were pooled, concentrated if required using e.g. a MILLIPORE Amicon Ultra (30 MWCO) centrifugal concentrator and stored at -80 °C. Part of the samples were provided for subsequent protein analytics and analytical characterization e.g. by SDS-PAGE, size exclusion chromatography, mass spectrometry and Endotoxin determination (see Figure 2).

SDS-PAGE

The NuPAGE® Pre-Cast gel system (Invitrogen) was used according to the manufacturer's instruction. In particular, 4-20% NuPAGE® Novex® TRIS-Glycine Pre-Cast gels and a Novex® TRIS-Glycine SDS running buffer were used. (see e.g. Figure 1). Reducing of samples was achieved by adding NuPAGE® sample reducing agent prior to running the gel.

Analytical size exclusion chromatography

Size exclusion chromatography for the determination of the aggregation and oligomeric state of antibodies was performed by HPLC chromatography. Briefly, Protein A purified antibodies were applied to a Tosoh TSKgel G3000SW column in 300 mM NaCl, 50 mM KH2PO4/K2HPO4, pH 7.5 on an Dionex HPLC system or to a Superdex 200 column (GE Healthcare) in 2 x PBS on a Dionex HPLC-System. The eluted protein was quantified by UV absorbance and integration of peak areas. BioRad Gel Filtration Standard 151–1901 served as a standard.

Mass spectrometry

The total deglycosylated mass of antibodies was determined and confirmed via electrospray ionization mass spectrometry (ESI-MS). Briefly, 100 µg purified antibodies were deglycosylated with 50 mU N-Glycosidase F (PNGaseF, ProZyme) in 100 mM KH2PO4/K2HPO4, pH 7 at 37°C for 12-24 h at a protein concentration of up to 2 mg/ml and subsequently desalted via HPLC on a Sephadex G25 column (GE Healthcare). The mass of the respective heavy and light chains was determined by ESI-MS after deglycosylation and reduction. In brief, 50 µg antibody in 115 µl were incubated with 60 µl 1M TCEP and 50 µl 8 M Guanidinium-hydrochloride subsequently desalted. The total mass and the mass of the reduced heavy and light chains was determined via ESI-MS on a Q-Star Elite MS system equipped with a NanoMate source.

ANG-1 and ANG-2 binding ELISA

The binding properties of antibodies directed against ANGPTs (Angiopoietin 1 or 2) were evaluated in an ELISA assay with full-length Angiopoietin-2-His protein (R&D Systems #623-AN/CF or in house produced material) or Angiopoietin-1-His (R&D systems #923-AN). Therefore 96 well plates (Falcon polystyrene clear enhanced microtiter plates or Nunc Maxisorb) were coated with 100 µl 1 µg/mL recombinant human Angiopoietin-1 or Angiopoietin-2 (carrier-free) in PBS (Sigma) for 2 h at room temperature or over night at 4°C. The wells were washed three times with 300µl PBST (0,2% Tween 20) and blocked with 200 µl 2% BSA 0,1% Tween 20 for 30 min at room temperature and subsequently washed three times with 300µl PBST. 100 µL/well of a dilution series (40pM-0.01 pM) of purified test antibody against <ANG-2> and as a reference Mab536 (Oliner, J., et al., Cancer Cell. Nov 6 (2004) 507-16, US 2006/0122370) in PBS was added to the wells and incubated for 1 h on a microtiterplate shaker at room temperature. The wells were washed three times with 300µl PBST (0,2% Tween 20) and bound antibody was detected with 100 µL/well 0.1 µg/ml F(ab') <hk>POD (Biozol Cat.No. 206005) in 2% BSA 0,1% Tween 20 as detection antibody for 1 h on a microtiterplate shaker at room temperature. Unbound detection antibody was washed away three times with 300 µL/well PBST and the bound detection antibody was detected by addition of 100 µL ABTS/well. Determination of absorbance was performed on a Tecan Fluor Spectrometer at a measurement wavelength of 405 nm (reference wavelength 492 nm).

ANG-2 binding BIACORE

Binding of the antibodies to the antigen e.g. human ANG-2 were investigated by surface plasmon resonance using a BIACORE T100 instrument (GE Healthcare Biosciences AB, Uppsala, Sweden). Briefly, for affinity measurements goat<hIgG-Fcgamma> polyclonal antibodies were immobilized on a CM4 chip via amine coupling for presentation of the antibodies against human ANG-2. Binding was measured in HBS buffer (HBS-P (10 mM HEPES, 150 mM NaCl, 0.05% Tween 20, ph 7.4), 25°C. Purified ANG-2-His (R&D systems or in house purified) was added in various concentrations between 0,41 nM and 200 nM in solution. Association was measured by an ANG-2-injection of 3 minutes; dissociation was measured by washing the chip surface with HBS buffer for 5 minutes and a KD value was estimated using a 1:1 Langmuir binding model. Due to heterogenity of the ANG-2 preparation no 1:1 binding could be observed; KD values are thus only relative estimations. Negative control data (e.g. buffer curves) were subtracted from sample curves for correction of system intrinsic baseline drift and for noise signal reduction. Biacore T100 Evaluation Software version 1.1.1 was used for analysis of sensorgrams and for calculation of affinity data. Alternatively, Ang-2 could be captured with a capture level of 2000-1700 RU via a PentaHisAntibody (PentaHis-Ab BSA-free, Qiagen No. 34660) that was immobilized on a CM5 chip via amine coupling (BSA-free) (see below).

Inhibition of huANG-2 binding to Tie-2 (ELISA)

The interaction ELISA was performed on 384 well microtiter plates (MicroCoat, DE, Cat.No. 464718) at RT. After each incubation step plates were washed 3 times with PBST. ELISA plates were coated with 0.5 µg/ml Tie-2 protein (R&D Systems, UK, Cat.No.313-TI) for at least 2 hours (h).Thereafter the wells were blocked with PBS supplemented with 0.2% Tween-20 and 2% BSA (Roche Diagnostics GmbH, DE) for 1 h. Dilutions of purified antibodies in PBS were incubated together with 0.2 µg/ml huAngiopoietin-2 (R&D Systems, UK, Cat.No. 623-AN) for 1 h at RT. After washing a mixture of 0.5 µg/ml biotinylated anti-Angiopoietin-2 clone BAM0981 (R&D Systems,UK) and 1:3000 diluted streptavidin HRP (Roche Diagnostics GmbH, DE, Cat.No.11089153001) was added for 1 h. Thereafter the plates were washed 6 times with PBST. Plates were developed with freshly prepared ABTS reagent (Roche Diagnostics GmbH, DE, buffer #204 530 001, tablets #11 112 422 001) for 30 minutes at RT. Absorbance was measured at 405 nm.

Inhibition of huANG-1 binding to Tie-2 (ELISA)

The interaction ELISA was performed on 384 well microtiter plates (MaxiSorb Nunc#442768) at RT. After each incubation step plates were washed 3 times with PBST. ELISA plates were coated with 0.5 µg/ml Tie-2 protein (R&D Systems, UK, Cat.No.313-TI or in house produced material) for at least 2 hours (h).Thereafter the wells were blocked with PBS supplemented with 0.2% Tween-20 and 2% BSA (Roche Diagnostics GmbH, DE) for 1 h. Dilutions of purified antibodies in PBS were incubated together with 0.2 µg/ml huAngiopoietin-1 (R&D Systems #923-AN/CF or in house produced material) for 1 h at RT. After washing a mixture of 0.5 µg/ml biotinylated anti-Angiopoietin-1 clone (R&D Systems #BAF923) and 1:3000 diluted streptavidin HRP (Roche Diagnostics GmbH, DE, Cat.No.11089153001) was added for 1 h. Thereafter the plates were washed 6 times with PBST. Plates were developed with freshly prepared ABTS reagent (Roche Diagnostics GmbH, DE, buffer #204 530 001, tablets #11 112 422 001) for 30 minutes at RT. Absorbance was measured at 405 nm.

Generation of HEK293-Tie2 cell line

In order to determine the interference of Angiopoietin-2 antibodies with ANGPT2 stimulated Tie2 phosphorylation and binding of ANGPT2 to Tie2 on cells a recombinant HEK293-Tie cell line was generated. Briefly, a pcDNA3 based plasmid (RB22-pcDNA3 Topo hTie2) coding for full-length human Tie2 (SEQ ID 61) under control of a CMV promoter and a Neomycin resistance marker was transfected using Fugene (Roche Applied Science) as transfection reagent into HEK293 cells (ATCC) and resistant cells were selected in DMEM 10% FCS, 500µg/ml G418. Individual clones were isolated via a cloning cylinder, and subsequently analyzed for Tie2 expression by FACS. Clone 22 was identified as clone with high and stable Tie2 expression even in the absence of G418 (HEK293-Tie2 clone22). HEK293-Tie2 clone22 was subsequently used for cellular assays: ANGPT2 induced Tie2 phosphorylation and ANGPT2 cellular ligand binding assay.

ANGPT2 induced Tie2 phosphorylation assay

Inhibition of ANGPT2 induced Tie2 phosphorylation by ANGPT2 antibodies was measured according to the following assay principle. HEK293-Tie2 clone22 was stimulated with ANGPT2 for 5 minutes in the absence or presence of ANGPT2

antibody and P-Tie2 was quantified by a sandwich ELISA. Briefly, 2x105 HEK293-Tie2 clone 22 cells per well were grown over night on a Poly-D-Lysine coated 96 well- microtiter plate in 100µl DMEM, 10% FCS, 500 µg/ml Geneticin. The next day a titration row of ANGPT2 antibodies was prepared in a microtiter plate (4-fold concentrated, 75µl final volume/well, duplicates) and mixed with 75µl of an ANGPT2 (R&D systems # 623-AN] dilution (3.2 µg/ml as 4-fold concentrated solution). Antibodies and ANGPT2 were pre-incubated for 15 min at room temperature. 100 µl of the mix were added to the HEK293-Tie2 clone 22 cells (pre-incubated for 5 min with 1 mM NaV3O4, Sigma #S6508) and incubated for 5 min at 37°C. Subsequently, cells were washed with 200µl ice-cold PBS + 1mM NaV3O4 per well and lysed by addition of 120µl lysis buffer (20 mM Tris, pH 8.0, 137 mM NaCl, 1% NP-40, 10% glycerol, 2mM EDTA, 1 mM NaV3O4, 1 mM PMSF and 10 µg/ml Aprotinin) per well on ice. Cells were lysed for 30 min at 4°C on a microtiter plate shaker and 100 µl lysate were transferred directly into a p-Tie2 ELISA microtiter plate (R&D Systems, R&D #DY990) without previous centrifugation and without total protein determination. P-Tie2 amounts were quantified according to the manufacturer's instructions and IC50 values for inhibition were determined using XLfit4 analysis plug-in for Excel (Dose-response one site, model 205). IC50 values can be compared within on experiment but might vary from experiment to experiment.

ANGPT1 induced Tie2 phosphorylation assay

Inhibition of ANGPT1 induced Tie2 phosphorylation by ANGPT1 antibodies was measured according to the following assay principle. HEK293-Tie2 clone22 was stimulated with ANGPT1 for 5 minutes in the absence or presence of ANGPT1 antibody and P-Tie2 was quantified by a sandwich ELISA. Briefly, 2x105 HEK293-Tie2 clone 22 cells per well were grown over night on a Poly-D-Lysine coated 96 well- microtiter plate in 100µl DMEM, 10% FCS, 500 µg/ml Geneticin. The next day a titration row of ANGPT1 antibodies was prepared in a microtiter plate (4-fold concentrated, 75µl final volume/well, duplicates) and mixed with 75µl of an ANGPT1 (R&D systems # 923-AN] dilution (0.8 µg/ml as 4-fold concentrated solution). Antibodies and ANGPT1 were pre-incubated for 15 min at room temperature. 100 µl of the mix were added to the HEK293-Tie2 clone 22 cells (pre-incubated for 5 min with 1 mM NaV3O4, Sigma #S6508) and incubated for 5 min at 37°C. Subsequently, cells were washed with 200µl ice-cold PBS + 1mM NaV3O4 per well and lysed by addition of 120µl lysis buffer (20 mM Tris,

pH 8.0, 137 mM NaCl, 1% NP-40, 10% glycerol, 2mM EDTA, 1 mM NaV3O4, 1 mM PMSF and 10 µg/ml Aprotinin) per well on ice. Cells were lysed for 30 min at 4°C on a microtiter plate shaker and 100 µl lysate were transferred directly into a p-Tie2 ELISA microtiter plate (R&D Systems, R&D #DY990) without previous centrifugation and without total protein determination. P-Tie2 amounts were quantified according to the manufacturer's instructions and IC50 values for inhibition were determined using XLfit4 analysis plug-in for Excel (Dose-response one site, model 205). IC50 values can be compared within on experiment but might vary from experiment to experiment.

Example 1

Expression & purification of monoclonal <ANG-2> antibodies Ang2i-LC06, Ang2i-LC07 and Ang2k-LC08

Light and heavy chains of the corresponding antibodies Ang2i-LC06, Ang2i-LC07 and Ang2k-LC08 were constructed in expression vectors as described above. The heavy chain and the kappa light was cloned in a genomic expression cassette, whereas the lambda light chain was cloned as cDNA with intron A (Fig 1B). The plasmids were amplified in E. coli, purified, and subsequently transfected for transient expression of recombinant proteins in HEK293-F cells (utilizing Invitrogen's FreeStyle 293 system). After 7 days, HEK 293-F cell supernatants were harvested, filtered and the antibodies were purified by protein A and size exclusion chromatography. Homogeneity of all antibodies was confirmed by SDS-PAGE under non reducing and reducing conditions and analytical size exclusion chromatography. Under reducing conditions (Figure 1), polypeptide heavy chains of <ANG-2> antibodies showed upon SDS-PAGE apparent molecular sizes of ca. 50 kDa analogous to the calculated molecular weights, polypeptide light chains showed apparent molecular masses of 25 kDa according to their predicted size. Mass spectrometry confirmed the identity of the purified antibodies. Expression levels of all constructs were analyzed by Protein A HPLC.

Size exclusion chromatography analysis of the purified. All antibodies were prepared and analytically characterized analogously to the procedure described. The SEC data of the corresponding antibodies were summarized in the table below.

	Antibody chain	Theoretical mass (Da)	Experimental mass (Da)	SEC (%) main peak
<ANG-2>Ang-2i_LC07	HC	50343	50325 (pyro-Glu)	99.7%
	LC	22738	22720 (pyro-Glu)	
<ANG-2>Ang-2i_LC06	HC	50343	50325 (pyro-Glu)	99.8%
	LC	22620	22605 (pyro-Glu)	
<ANG-2>Ang-2k_LC08	HC	49544	49527 (pyro-Glu)	99.8%
	LC	22685	22667 (pyro-Glu)	

Example 2

ELISA Binding assay to human ANG-1 and to human ANG-2

The binding of <ANG-2> antibodies Ang2i-LC06, Ang2i-LC07 and Ang2k-LC08 to human ANG-1 and human ANG-2 was determined in an ANG-1 or ANG-2 binding ELISA as described above. Briefly, the ELISA-type assay is based on the immobilization of human wild-type Angiopoieti-1 or -2 in a microtiter plate. Binding of an antibody directed against the immobilized ANG-1 or ANG-2 is measured via an <human Fc> (anti-IgG) antibody with a POD conjugate. A dilution series of the <ANG-2>antibody allows determining an EC50 concentration. As a reference the human anti-ANG-2 antibody <ANG-2> antibody Mab536 (Oliner et al., Cancer Cell. Nov 6 (2004) 507-16, US 2006/0122370) was used. The determined EC50 concentrations are summarized in the table below.

Antibody	hANG-1 binding EC50	hANG-2 binding EC50
<ANG-2>MAb536	2538 ng/mL	133 ng/mL
<ANG-2>Ang2i-LC06	> 8000 ng/mL	84 ng/mL
<ANG-2>Ang2i-LC07	> 8000 ng/mL	3006 ng/mL
<ANG-2>Ang2i-LC08	4044 ng/mL	105 ng/mL

All antibodies are specifically binding to ANG-2. MAb536 and Ang2k-LC08 show also specific binding towards ANG-1, whereas Ang2i-LC06 and Ang2i-LC07 are not specifically binding to ANG-1 as they have an EC50-value of above 8000 ng/ml (detection limit).

Example 3:

Binding to ANG-2 via Biacore

The affinity for binding to human ANGPT2 was examined with a Biacore assay as describes above. Briefly, is this assay a capturing antibody (anti-Fc) is immobilized to the surface of the Biacore chip, which captures and presents the corresponding antibody (for example Ang2i-LC06). The ligand (here ANGPT2) is captured from solution. The affinity for this interaction is determined with the assumption of a 1:1 interaction. Details of this experiment can be found in the general methods section. The affinities determined for ANGPT2-binding (KD) are summarized in the table below.

hAng-2	**Experiment 1**			**Experiment 2**			**Average (from 1+2)**	
	KD (pM)	**kd (1/s)**	**$t_{(1/2)}$diss (min)**	**KD (pM)**	**kd (1/s)**	**$t_{(1/2)}$diss (min)**	**KD (pM)**	**$t_{(1/2)}$diss (min)**
Ang2i-LC06	11	7.16E-05	161	21	1.14E-04	102	16	132
Ang2k-LC08	16	1.61E-04	72	27	2.28E-04	51	22	61
MAb536	29	1.44E-04	80	29	1.25E-04	92	29	86

The antibodies Ang2i-LC06 and Ang2k bind with high affinity to ANGPT2.

Example 4:

Neutralization of ANGPT1/2-Tie2 interaction (human)

Blocking of human ANGPT1/2/human Tie2 interaction was shown by receptor interaction ELISA. 384-well Maxisorp plates (Nunc) were coated with 0.5 µg/ml human Tie2 (R&D Systems, UK, Cat.No.313-TI or in house produced material) for 2 h at room temperature and blocked with PBS supplemented with 0.2% Tween-20 and 2% BSA (Roche Diagnostics GmbH, DE) for 1 h at room temperature under shaking. In the meantime, Dilutions of purified antibodies in PBS were incubated together with 0.2 µg/ml huAngiopoietin-1/2 (R&D Systems #923-AN/CF, R&D Systems,UK, Cat.No. 623-AN or in house produced material) for 1 h at RT. After washing a mixture of 0.5 µg/ml biotinylated anti-Angiopoietin-1/2 clone (R&D Systems #BAF923, BAM0981 R&D Systems,UK) and 1:3000 diluted streptavidin HRP (Roche Diagnostics GmbH, DE, Cat.No.11089153001) was added for 1 h. Thereafter the plates were washed 6 times with PBST. Plates were developed with freshly prepared ABTS reagent (Roche Diagnostics GmbH, DE, buffer #204 530 001, tablets #11 112 422 001) for 30 minutes at RT. Absorbance was measured at 405 nm.

The obtained inhibitory concentrations are summarized in the following table.

Antibody	**ANGPT1/Tie2 interaction ELISA**	**ANGPT2/Tie2 interaction ELISA**
Ang2i-LC06	> 100 nM	0.1 nM
Ang2k-LC08	11 nM	0.17 nM
MAb536	n.d.	0.15 nM

The table above shows different selectivity profiles for the two antibodies Ang2i-LC06 and Ang2k-LC08. Ang2i-LC06 is ANGPT2 selective, whereas Ang2k-LC08 is ANGPT1/2 cross reactive in inhibition for ANGPT1/2 Tie2 interaction.

Example 5:

Tie2 phosphorylation

The ability of the identified ANGPT2 antibodies to interfere with ANGPT2 and ANGPT1 mediated Tie2 phosphorylation was examined in the ANGPT2 and

ANGPT1 induced Tie2 phosphorylation assays as described above. A schematic representation of the assay setup is depicted in Figure 3.

Both antibodies Ang2i-LC06 and Ang2k-LC08 showed a dose-dependent interference with ANGPT2 stimulated Tie2 phosphorylation as shown in Figure 4 with comparable IC50 values. Ang2i-LC06 interfered with ANGPT2 stimulated Tie2 phosphorylation with a IC50 value of approx. 508 ng/ml and Ang2k-LC08 interfered with ANGPT2 stimulated Tie2 phosphorylation with a IC50 value of approx. 499 ng/ml. In contrast, only Ang2k-LC08 interfered with ANGPT1 stimulated Tie2 phosphorylation with a IC50 value of approx. 391 ng/ml whereas Ang2i-LC06 did not interfere with ANGPT2 stimulated Tie2 phosphorylation in the same tested concentration range (Figure 5).

Example 6: In vivo efficacy

Effect of anti ANGPT antibodies on Colo205 xenograft growth

In vivo efficacy of <ANGPT2> antibodies Ang2i-LC06 and Ang2k-LC08 in comparison to <ANGPT2> Mab536 in staged subcutaneous Colo205 xenograft model

The purified Ang2i-LC06 and Ang2k-LC08 antibodies were compared to the antibody Mab536 in the staged subcutaneous Colo205 xenograft model (Ang2_PZ_Colo205_006) in female Scid beige mice.

Antibodies: Mab536 was provided as frozen stock solution (c = 4.5 mg/mL), Ang2i-LC06 and Ang2k-LC08 were provided as frozen stock solution (c = 1 mg/mL) in 20 mM Histidine, 140 mM NaCl, pH 6.0. Antibody solution was diluted appropriately in PBS from stock prior injections where required and PBS was applied as vehicle. The humanized IgG1 anti-IgE antibody Xolair (Omalizumab) served as positive control and was bought from a pharmacy.

Cell lines and culture conditions: Colo205 human colorectal cancer cells were originally obtained from ATCC and after expansion deposited in the Roche Penzberg internal cell bank. Tumor cell line was routinely cultured in RPMI 1640 medium (PAA, Laboratories, Austria) supplemented with 10% fetal bovine serum (PAA Laboratories, Austria) and 2 mM L-glutamine, at 37 °C in a water-saturated atmosphere at 5% CO2. Passage 3 was used for transplantation.

Animals: Female SCID beige mice (purchased from Charles River Germany) were maintained under specific-pathogen-free condition with daily cycles of 12 h light /12 h darkness according to committed guidelines (GV-Solas; Felasa; TierschG). Experimental study protocol was reviewed and approved by local government. After arrival animals were maintained in the quarantine part of the animal facility for one week to get accustomed to new environment and for observation. Continuous health monitoring was carried out on regular basis. Diet food (Provimi Kliba 3337) and water (acidified pH 2.5-3) were provided ad libitum. Age of mice at start of the study was about 12-14 weeks.

Monitoring: Animals were controlled daily for clinical symptoms and detection of adverse effects. For monitoring throughout the experiment body weight of animals was documented and tumor volume was measured by caliper after staging.

Tumor cell injection: At day of injection Colo205 cells were centrifuged, washed once and resuspended in PBS. After an additional washing with PBS cell concentration and cell size were determined using a cell counter and analyzer system (Vi-CELL, Beckman Coulter). For injection of Colo205 cells, the final titer was adjusted to 5.0 x 10E7 cells/ml, viability ca. 90%. Subsequently 100 µl of this suspension corresponding to 2.5*106 cells per animal was injected s.c. into the right flank of the mice.

Treatment of animals: Animal treatment started at day of randomisation, 16 days after cell transplantation (study Ang2_PZ_Colo205_006) at a mean tumor volume of 178 mm3.

Dose schedule of study Ang2_PZ_Colo205_006:

Group	No of animals	Compound	Dose (mg/kg)	Route/Mode of administration	No of treatments	Cumulative dose (mg/kg)
1	10	Vehicle		i.p. once weekly	5	
2	10	Xolair	10	i.p. once weekly	5	50
3	10	Ang2i-LC06	10	i.p. once weekly	5	50
5	10	Ang2k-LC08	10	i.p. once weekly	5	50
6	10	MAB536	10	i.p. once weekly	5	50

Tumor growth inhibition until Day 50 is shown in Figure 6. The data show that the ANGPT2 selective antibody Ang2i-LC06 was the most active antibody (Tumor control ration (TCR) value 0.39). Ang2i-LC06 was more efficacious in inhibiting tumor growth than antibody MAb536 (TCR value 0.47) and the ANGPT2 selective, ANGPT1 cross-reactive antibody Ang2k-LC08 (TCR value 0.46).

Effect of anti ANGPT antibodies on KPL-4 xenograft growth

In vivo efficacy of <ANGPT2> antibodies Ang2i-LC06 and Ang2k-LC08 in comparison to <ANGPT2> Mab536 in staged orthotopic KPL-4 xenograft model

The purified Ang2i-LC06 and Ang2k-LC08 antibodies were compared to the antibody Mab536 in the staged orthotopic KPL-4 xenograft model (Ang2_PZ_KPL-4_002) in female Scid beige mice.

Antibodies: Mab536 was provided as frozen stock solution (c = 4.5 mg/mL), Ang2i-LC06 and Ang2k-LC08 were provided as frozen stock solution (c = 1 mg/mL) in 20 mM Histidine, 140 mM NaCl, pH 6.0. Antibody solution was diluted appropriately in PBS from stock prior injections where required and PBS was applied as vehicle.

Cell lines and culture conditions: KPL-4 human breast cancer cells were originally established from the malignant pleural effusion of a breast cancer patient with an inflammatory skin metastasis. KPL-4 cells were kindly provided by Prof. J. Kurebayashi (Kawasaki Medical School, Kurashiki, Japan). Tumor cells were routinely cultured in DMEM medium (PAN Biotech, Germany) supplemented with 10% fetal bovine serum (PAN Biotech, Germany) and 2 mM L-glutamine (PAN Biotech, Germany) at 37 °C in a water-saturated atmosphere at 5% CO2. Culture passage was performed with trypsin / EDTA 1x (PAN) splitting three times / week.

Animals: Female SCID beige mice (purchased from Charles River Germany) were maintained under specific-pathogen-free condition with daily cycles of 12 h light /12 h darkness according to committed guidelines (GV-Solas; Felasa; TierschG). Experimental study protocol was reviewed and approved by local government. After arrival animals were maintained in the quarantine part of the animal facility for one week to get accustomed to new environment and for observation. Continuous health monitoring was carried out on regular basis. Diet food (Provimi Kliba 3337) and water (acidified pH 2.5-3) were provided ad libitum. Age of mice at start of the study was about 12 weeks.

Monitoring: Animals were controlled daily for clinical symptoms and detection of adverse effects. For monitoring throughout the experiment body weight of animals was documented and tumor volume was measured by caliper after staging.

Tumor cell injection: At the day of injection tumor cells were harvested (trypsin-EDTA) from culture flasks (Greiner TriFlask) and transferred into 50 ml culture medium, washed once and resuspended in PBS. After an additional washing step with PBS and filtration (cell strainer; Falcon™; 100µm) the final cell titer was adjusted to 1.5 x 108 / ml. Tumor cell suspension was carefully mixed with transfer pipette to avoid cell aggregation. Anesthesia was performed using a Stephens's inhalation unit for small animals with preincubation chamber (plexiglas), individual mouse nose-mask (silicon) and not flammable or explosive anesthesia compound Isoflurane (Pharmacia-Upjohn, Germany) in a closed circulation system. Two days before injection coat of the animals were shaved. For i.m.f.p. injection cells were injected orthotopically at a volume of 20 µl (3*106 / animal) into the right penultimate inguinal mammary fat pad of each anesthetized mouse. For the orthotopic implantation, the cell suspension was injected through the skin under the nipple using a using a Hamilton microliter syringe and a 30Gx1/2" needle.

Treatment of animals started at day of randomization with tumors ranging from 60-180 mm 3,35 days after cell transplantation (study Ang2_PZ_KPL-4_002) at a mean tumor volume of ca. 90 mm3.

Dose schedule of study Ang2_PZ_KPL-4_002:

Group	No of animals	Compound	Dose (mg/kg)	Route/Mode of administration	No of treatments	Cumulative dose (mg/kg)
1	10	Vehicle		i.p. once weekly	5	
2	10	Xolair	10	i.p. once weekly	5	50
3	10	Ang2i-LC06	10	i.p. once weekly	5	50
5	10	Ang2k-LC08	10	i.p. once weekly	5	50
6	10	MAB536	10	i.p. once weekly	5	50

Tumor growth inhibition until day 64 is shown in Figure 7. The data show that the ANGPT2 selective antibody Ang2i-LC06 was the most active antibody (TCR value 0.55) in the KPL-4 model. Ang2i-LC06 was more efficacious in inhibiting tumor growth than antibody MAb536 (TCR value 0.57) and the ANGPT2 selective, ANGPT1 cross-reactive antibody Ang2k-LC08 (TCR value 0.57).

Example 7:

Binding to ANG-1 via Biacore

The affinity for binding to human ANG-1 was examined with a Biacore assay: huAng-1 was immobilized on a CM5 biosensorchip using amine-coupling chemistry. The protein was injected for 20 min in sodium acetate pH 4.5 at 10 µg/ml at a flow rate of 5 µl/min. This resulted in a surface density of appr. 20000 RU. On the reference flow cell BSA was immobilized under the same conditions. The antibodies were diluted in HBS-P to 100 nM and injected for 3 min (association phase). After washing with running buffer for 3 min (dissociation phase), the surface was regenerated by injecting 10 mM sodium hydroxide for 1 min at 5 µl/min. Results are shown in Fig. 8: Ang2k_LC08 had a halftime of complex dissociation of approximately 50s, Ang2i_LC06 of appr. 5s and Ang2i_LC10 showed no binding to ANG-1.

Example 8: Prevention of metastasis/secondary tumors in vivo in bearing primary tumors

a) Prevention of metastasis/ secondary in mice xenografted with primary Colo205 tumors

Cell lines and culture conditions:

Colo205 human colorectal cancer cells were originally obtained from ATCC and after expansion deposited in the Roche Penzberg internal cell bank. Tumor cell line was routinely cultured in RPMI 1640 medium (PAA, Laboratories, Austria) supplemented with 10 % fetal bovine serum (PAA Laboratories, Austria) and 2 mM L-glutamine, at 37 °C in a water-saturated atmosphere at 5 % CO_2. Passage 3 was used for transplantation.

Animals:

Female SCID beige mice; age 4–5 weeks at arrival (purchased from Charles River Germanyd) were maintained under specific-pathogen-free condition with daily cycles of 12 h light /12 h darkness according to committed guidelines (GV-Solas; Felasa; TierschG). Experimental study protocol was reviewed and approved by local government. After arrival animals were maintained in the quarantine part of the animal facility for one week to get accustomed to new environment and for observation. Continuous health monitoring was carried out on regular basis. Diet food (Provimi Kliba 3337) and water (acidified pH 2.5-3) were provided ad libitum. Age of mice at start of the study was about 10 weeks.

Tumor cell injection:

At the day of injection, Colo205 tumor cells were harvested (trypsin-EDTA) from culture flasks (Greiner) and transferred into 50 ml culture medium, washed once and resuspended in PBS. After an additional washing step with PBS and filtration (cell strainer; Falcon ⌀ 100μm) the final cell titer was adjusted to 2.5 x 10^7 / ml. Tumor cell suspension was carefully mixed with transfer pipette to avoid cell aggregation. After this, cell suspension was filled into a 1.0 ml tuberculin syringe (Braun Melsungen) using a wide needle (1.10 x 40 mm); for injection needle size was changed (0.45 x 25 mm) and for every injection a new needle was used. Anesthesia was performed using a Stephens inhalation unit for small animals with

preincubation chamber (plexiglas), individual mouse nose-mask (silicon) and not flammable or explosive anesthesia compound Isoflurane (cp-pharma) in a closed circulation system. Two days before injection coat of the animals were shaved and for cell injection skin of anaesthetized animals was carefully lifted up with an anatomic forceps and 100 µl cell suspension (= 2.5 x 10^6 cells) was injected subcutaneously in the right flank of the animals. Tumor growth of the primary tumors was monitored (data not shown)

Monitoring of secondary tumors e.g. in the lung by Quantification of human Alu sequences

At study termination (day 103) lungs were collected from animals of all groups. Briefly, samples are transferred immediately into fluid nitrogen. In a further step total DNA was isolated from the samples with MagNA Pure LC Instrument according to manufacturer's instructions. Human Alu specific primers were chosen for selective amplification of Alu sequences by quantitative PCR (LightCycler instrument). (T. Schneider et. al., Clin. Exp. Metas. 2002; 19: 571-582).

Treatment of animals

Treatment of animals with Avastin (10 mg/kg i.p. once weekly) was started 14 days after cell transplantation (study Ang2_PZ_Colo205_008) at a mean tumor volume of 340 mm^3. After 7 weeks mice were randomized for subsequent secondary treatment starting at day 51 with compounds listed in table below. Secondary treatment starting at day 51 of Study Ang2_PZ_Colo205_008.

Group	No of animals	Compound	Dose (mg/kg)	Route/Mode of administration	No of treatments	Cumulative dose (mg/kg)
1	10	Avastin	10 mg/kg	i.p. once weekly	11	110
2	10	LC06 + Avastin	10 mg/kg 10 mg/kg	i.p. once weekly i.p. once weekly	6 11	60 110
3	10	LC06	10 mg/kg	i.p. once weekly	6	60

Results of prevention of metastasis/ secondary tumors (in the lung) are listed in the table below and shown in Figure 9A

Table 1 : Quantification of human ALU DNA in the lungs of mice originally bearing primary Colo205 tumors , after treatment with different antibodies

	Avastin		Avastin + Ang2i-LC06		Ang2i_LC06	
	101	0,0264	201	0,0042	301	0,0047
	102	5,6740	202	0,0044	302	0,0055
	103	0,0307	203	0,0065	303	0,0050
	104	0,0203	204	0,0081	304	0,0064
	105	0,0215	205	0,0063	305	0,0062
	106	0,0338	206	0,0061	306	0,0066
	107	0,0075	207	0,0053	307	0,0250
	108	0,0113	208	0,0506	308	0,0062
	109	0,0087	209	0,0065	309	0,0067
	110	0,0587	210	0,0160	310	0,0064
mean		0,5893		0,0114		0,0079
median		**0,0240**		**0,0064**		**0,0063**

Results show a clearly improved preventionof secondary tumors/metastasis by ANG2i-LC06 compared with Avastin

b) Prevention of metastasis/ secondary in mice xenografted with primary KPL-4 tumors

Tumor cell line

The human breast cancer cell line KPL-4 (kindly provided by Prof. J. Kurebayashi) has been established from the malignant pleural effusion of a breast cancer patient with an inflammatory skin metastasis. Tumor cells are routinely cultured in DMEM medium (PAN Biotech, Germany) supplemented with 10% fetal bovine serum (PAN Biotech, Germany) and 2 mM L-glutamine (PAN Biotech, Germany) at 37°C in a water-saturated atmosphere at 5% CO_2. Culture passage is performed with trypsin / EDTA 1x (PAN) splitting three times / week.

Mice

After arrival, female SCID beige mice (age 10-12 weeks; body weight 18-20 g) Charles River, Sulzfeld, Germany) were maintained in the quarantine part of the animal facility for one week to get them accustomed to the new environment and for observation. Continuous health monitoring was carried out. The mice were kept under SPF-conditions according to the international guidelines (GV-Solas; Felasa;

TierschG) with daily cycles of 12 h light /12 h darkness. Diet food (Kliba Provimi 3347) and water (filtered) were provided ad libitum. Experimental study protocol was reviewed and approved by the local government (Regierung von Oberbayern; registration no. 211.2531.2-22/2003).

Tumor cell injection

At the day of injection tumor cells were harvested (trypsin-EDTA) from culture flasks (Greiner TriFlask) and transferred into 50 ml culture medium, washed once and resuspended in PBS. After an additional washing step with PBS and filtration (cell strainer; Falcon ∅ 100µm) the final cell titer was adjusted to 1.5×10^8 / ml. Tumor cell suspension was carefully mixed with transfer pipette to avoid cell aggregation. Anesthesia is performed using a Stephens inhalation unit for small animals with preincubation chamber (plexiglas), individual mouse nose-mask (silicon) and not flammable or explosive anesthesia compound Isoflurane (Pharmacia-Upjohn, Germany) in a closed circulation system. Two days before injection coat of the animals were shaved. For i.m.f.p. injection cells were injected orthotopically at a volume of 20 µl into the right penultimate inguinal mammary fat pad of each anesthetized mouse. For the orthotopic implantation, the cell suspension was injected through the skin under the nipple using a using a Hamilton microliter syringe and a 30Gx1/2" needle. Tumor growth of the primary tumors was monitored (data not shown)

Monitoring of secondary tumors e.g. in the lung by Quantification of human Alu sequences

At study termination (day 103) lungs were collected from animals of all groups. Briefly, samples are transferred immediately into fluid nitrogen. In a further step total DNA was isolated from the samples with MagNA Pure LC Instrument according to manufacturer's instructions. Human Alu specific primers were chosen for selective amplification of Alu sequences by quantitative PCR (LightCycler instrument). (T. Schneider et. al., Clin. Exp. Metas. 2002; 19: 571-582)

Treatment of animals

Treatment of animals was started 35 days after cell transplantation at a mean tumor volume of 60-160 mm^3. Compounds and dose schedule is listed in the table below.

Group	No of animals	Compound	Dose (mg/kg)	Route/Mode of administration	No of treatments	Cumulative dose (mg/kg)
4	10	Vehicle		i.p. twice weekly	5	
5	10	Xolair	10	i.p. twice weekly	5	50
6	10	Ang2i_LC06	10	i.p. once weekly	4	40
7	10	Ang2i_LC07	10	i.p. once weekly	4	40
8	10	Ang2k_LC08	10	i.p. once weekly	4	40

Results of prevention of metastasis/ secondary tumors (in the lung) are listed in the table below and shown in Figure 9B

Table 2: Quantification of human ALU DNA in the lungs of mice originally bearing primary KPL4 tumors , after treatment with different antibodies

	Vehicle		Xolair		Ang2i_LC06		Ang2i_LC07		Ang2i_LC08	
	101	0,0098	201	0,0157			401	0,0273	501	0,0069
	102	0,0090	202	0,0516	302	0,0076	402	0,0060	502	0,0261
	103	0,0119	203	0,0108	303	0,0413	403	0,0046	503	0,0067
	104	0,0405	204	0,0148	304	0,0042	404	0,0164	504	0,0044
			205	0,0020	305	0,0041	405	0,0040	505	0,0039
	106	0,0381	206	0,0340	306	0,0093	406	0,0044	506	0,0051
	107	0,0281	207	0,0141	307	0,0038	407	0,0060	507	0,0037
			208	0,0422	308	0,0044	408	0,0174	508	0,0037
	109	0,0121	209	0,0227	309	0,0036	409	0,0314	509	0,0051
	110	0,0143	210	0,0383	310	0,0094	410	0,0083	540	0,0200
median		**0,0132**		**0,0192**		**0,0044**		**0,0072**		**0,0051**
mean		0,0205		0,0246		0,0098		0,0126		0,0086

Results show a very efficient prevention of secondary tumors/metastasis by ANG2i-LC06, ANG2i-LC07, ANG2k-LC08.

Example 9: Effects in the treatment of retinopathy

Methods

C57/Bl6 pups are cross fostered to CD1 nursing dams and are exposed to 75% oxygen from P7 to P12 (PRO-OX 110 chamber oxygen controller, Biospherix Ltd, Redfield, NY) which induces vessel obliteration and cessation of capillaries in the centre of the retina. The pups and nursing dams are placed in normal air leading to relative hypoxia and the induction of neovascularisation. On P13, pups were anaesthetised using isofluorane (5% induction, 3% maintenance combined with 1.5% oxygen) and the eye was exposed and 1 µl intraocular injections using a Nanofil syringe fitted with a 35 gauge needle (WPI, Sarasota, FL) into the left eye was performed. On P17, both eyes were dissected, fixed in 4% paraformaldehyde for 4 h at 4°C and retinas were dissected. Retinas were permeabilised in PBS containing 0.5% Triton X-100 and 1% bovine serum albumin, stained with 20 µg/ml biotinylated isolectin B4 (Sigma Aldrich, Gillingham, UK) in PBS pH 6.8, 1% Triton-X100, 0.1 mM $CaCl_2$, 0.1 mM $MgCl_2$, followed by 20 µg/ml ALEXA 488-streptavidin (Molecular Probes, Eugene, OR) and flat mounted in Vectashield (Vector Laboratories, Burlingame, CA). Retinas were imaged using a Nikon epi-fluorescence microscope at 4x magnification. Quantification of neovascular and ischaemic areas were performed in a blinded fashion using Photoshop CS3 along with Image J (NIH) and expressed as percentage of total retinal area (=normal+ischaemic+neovascular).

Results

Figure 10A show representative flat mounted retinas with the retinal vasculature visualised by isolectin staining. The centre ischemic areas induce neovascularisation and re-growth of the retinal vessels by upregulation of angiogenic inducers. The neovascular front is hyperproliferative leading to tortuous vessels in an irregular vessel pattern. The most outer areas contain the normal unaffected vessels. Quantification of retinal flat mounts showed that inhibition of VEGF with Avastin reduced retinal neovascularisation (see fig 10B , uninjected 36.7±1.8% to injected 22.4±3.0%) as expected. Inhibition of Ang2 using antibodies LC06 or LC08 also led to a reduction in neovascularisation (31.5±1.1% to 18.8±1.3% and 34.0±3.1% to 25.4±3.4%). Control injection of human Ig G had no effect on neovascularisation (see fig 10B , 38.3±1.1% to 38.3±0.8%).

What is claimed is:

1. An antibody or antibody fragment specifically binding to human angiopoietin-2 (ANG-2), wherein

a) the heavy chain variable domain comprises a CDR3 region of SEQ ID NO: 33, a CDR2 region of SEQ ID NO: 34, and a CDR1 region of SEQ ID NO: 35; and

b) the light chain variable domain comprises a CDR3 region of SEQ ID NO: 36, a CDR2 region of SEQ ID NO: 37, and a CDR1 region of SEQ ID NO: 38.

2. The antibody or antibody fragment according to claim 1, comprising

a) the heavy chain variable domain of SEQ ID NO: 39; and

b) the light chain variable domain of SEQ ID NO: 40.

3. The antibody or antibody fragment according to any one of claims 1 to 2, wherein

the antibody is not binding to human Angiopoietin 1 (ANG-1).

4. The antibody or antibody fragment according to any one of claims 1 to 3, wherein said antibody is of human IgG4 subclass or is of human IgG1 subclass.

5. A pharmaceutical composition comprising an antibody or fragment according to any one of claims 1 to 3, and a pharmaceutically acceptable carrier.

6. Use of the antibody or antibody fragment according to any one of claims 1 to 3 for the manufacture of a medicament for the prevention of metastasis.

7. Use of the antibody or antibody fragment according to any one of claims 1 to 3 for the manufacture of a medicament for the treatment of cancer.

8. Use of the antibody or antibody fragment according to any one of claims 1 to 3 for the manufacture of a medicament for the treatment of cancer, inflammatory diseases, atherosclerosis, ischemia, trauma, sepsis, COPD, asthma, diabetes, AMD, retinopathy, stroke, adipositas, acute lung injury, hemorrhage, vascular leak, allergy, Graves' disease, Hashimoto's autoimmune thyroiditis, idiopathic thrombocytopenic purpura, giant cell arteritis, rheumatoid arthritis, systemic lupus erythematosus (SLE), lupus nephritis, Crohn's disease, multiple sclerosis, or ulcerative colitis.

9. Use of the antibody or antibody fragment according to any one of claims 1 to 3 for the prevention of metastasis.

10. Use of the antibody or antibody fragment according to any one of claims 1 to 3 for the treatment of cancer.

11. Use of the antibody or antibody fragment according to any one of claims 1 to 3 for the treatment of cancer, inflammatory diseases, atherosclerosis, ischemia, trauma, sepsis, COPD, asthma, diabetes, AMD, retinopathy, stroke, adipositas, acute lung injury, hemorrhage, vascular leak, allergy, Graves' disease, Hashimoto's autoimmune thyroiditis, idiopathic thrombocytopenic purpura, giant cell arteritis, rheumatoid arthritis, systemic lupus erythematosus (SLE), lupus nephritis, Crohn's disease, multiple sclerosis, or ulcerative colitis.

12. A nucleic acid encoding a heavy chain of an antibody specifically binding to human angiopoietin-2 (ANG-2), wherein said antibody comprises a heavy chain variable domain and a light chain variable domain according to claim 1.

13. An expression vector comprising a nucleic acid according to claim 12 for the expression of an antibody specifically binding to human angiopoietin-2 (ANG-2) in a prokaryotic or eukaryotic host cell.

14. Prokaryotic or eukaryotic host cell comprising a vector according to claim 13.

Fig. 1A


pAng2i LC06 LC-lambda
8277 bp
BamHI (1)
Leader
Ang2i LC06 VL
CL lambda
XbaI (722)
BGH pA
P-SV40
HygR
pUC origin
AmpR
P-CMV
ClaI (6715)
IntronA
pAng2i LC06 HC
5483 bp
HindIII (1)
Leader + Leader intron
Ang2i LC06 VH
XhoI (587)
intron
CH1-IgG1
intron
Hinge
intron
CH2
intron
CH3
EagI (2571)
SV40 pA
AvrII (2723)
pAng-2i LC06 LC-lambda
5483 bp
P-CMV
ClaI (4797)
AmpR
pUC origin

Fig. 1B


pAng2k LC08 LC-lambda
8283 bp
Bam HI (1)
Leader
Ang2k LC08 VL
CL lambda
Xba I (728)
BGH pA
P-SV40
HygR
pUC origin
AmpR
P-CMV
Cla I (6721)
Intron A


pAng2k LC08 HC
5468 bp
HindIII (1)
Leader + Leader intron
Ang2k LC08 VH
Xho I (572)
intron
CH1-IgG1
intron
Hinge
intron
CH2
intron
CH3
SV40 pA
Aur II (2708)
pUC origin
AmpR
P-CMV
Cla I (4782)

Fig. 2


Reduced
Not reduced
Angpt2-LC06
Angpt2-LC07
Angpt2k_LC08
Angpt2-LC06
Angpt2-LC07
Angpt2k_LC08
250 kD
150 kD
100 kD
75 kD
50 kD
37 kD
25 kD
20 kD
15 kD
10 kD

Fig. 3


SA-POD
biot <hAng-2>
Antagonistic Ab
hAng1
Tie-2
Tie-2
h-Fc
h-Fc
plate

Fig. 4


Ang2k_LC08
1,2
1
0,8
0,6
0,4
0,2
0
1
0,1
0,01
Ang2i_LC06
1,2
1
0,8
0,6
0,4
0,2
0
1
0,1
0,01

Fig. 5


Ang2k_LC08
0,75
0,65
0,55
0,45
0,35
0,25
0,15
0,05
1
0,1
0,01
Ang2i_LC06
0,99
0,97
0,95
0,93
0,91
0,89
0,87
0,85
0,83
0,81
1
0,1
0,01

Fig. 6


ANG2_PZ_COLO205_006
Vehicle
ANG2i-LC06 10 mg/kg
Mab536 10 mg/kg
Xolair
Ang2k-LC08 10 mg/kg
Tumor volume [mm3]
Study Day
0
100
200
300
400
500
600
700
15
20
25
30
35
40
45
50
55

Fig. 7


ANG2_PZ_KPL-4_002
Vehicle
ANG2i-LC06 10 mg/kg
Ang2k-LC08 10 mg/kg
MAB536 10 mg/kg
Median tumor volume [mm3]
0
100
200
300
400
500
600
34
39
44
49
54
59
64
Study Day

Fig. 8



Fig. 9A


Ang2_Pz_Colo205_008
Median values
Ang2I_LC06
Avastin + Ang2i-LC06
Avastin
0,035
0,030
0,025
0,020
0,015
0,010
0,005
0,000
human DNA [pg/µl]

Fig. 9B


Metastasis in lung
Ang2_Pz_KPL-4_002
Ang2i_LC08
Ang2i_LC07
Ang2i_LC06
Xolair
Vehicle
0,025
0,020
0,015
0,010
0,005
0,000
human DNA [ng/ml]

Fig. 10A



Fig. 10B


i
% neovascularisation
% ischemia
NS
**
Uninjected
Injected
Human Ig G
ANG2i-LC06
ANG2i-LC08
Avastin
Relative change (inj/uninj)
Neovascular
Ischemia
Normal
ii
iii
iv
***