



(86) Date de dépôt PCT/PCT Filing Date: 2005/05/14
(87) Date publication PCT/PCT Publication Date: 2005/11/24
(85) Entrée phase nationale/National Entry: 2006/11/14
(86) N° demande PCT/PCT Application No.: IB 2005/002383
(87) N° publication PCT/PCT Publication No.: 2005/111211
(30) Priorités/Priorities: 2004/05/14 (US10/709,572);
2004/05/14 (US10/709,577); 2004/10/03 (US60/522,452);
2004/10/03 (US60/522,449); 2004/10/04 (US60/522,457);
2004/11/15 (US60/522,860); 2004/12/08 (US60/593,081);
2005/01/06 (US60/593,329); 2005/03/17 (US60/662,742);
2005/03/25 (US60/665,094); 2005/03/30 (US60/666,340)

(51) Cl.Int./Int.Cl. *C12Q 1/68* (2006.01)
(71) Demandeur/Applicant:
ROSETTA GENOMICS LTD., IL
(72) Inventeurs/Inventors:
BENTWICH, ITZHAK, IL;
AVNIEL, AMIR, IL;
KAROV, Yael, IL;
AHARONOV, RANIT, IL
(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : MICRO-ARN ET UTILISATIONS DE CEUX-CI
(54) Title: MICRONAS AND USES THEREOF

(57) **Abrégé/Abstract:**

Described herein are novel polynucleotides associated with prostate and lung cancer. The polynucleotides are miRNAs and miRNA precursors. Related methods and compositions that can be used for diagnosis, prognosis, and treatment of those medical conditions are disclosed. Also described herein are methods that can be used to identify modulators of prostate and lung cancer.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 November 2005 (24.11.2005)

PCT

(10) International Publication Number
WO 2005/111211 A3

(51) International Patent Classification:
C12Q 1/68 (2006.01)

KAROV, Yael [IL/IL]; 10 Plaut Street, 76706 Rehovot (IL). **AHARONOV, Ranit** [IL/IL]; 10 Plaut Street, 76706 Rehovot (IL).

(21) International Application Number:
PCT/IB2005/002383

(22) International Filing Date: 14 May 2005 (14.05.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

10/709,572	14 May 2004 (14.05.2004)	US
10/709,577	14 May 2004 (14.05.2004)	US
60/522,452	3 October 2004 (03.10.2004)	US
60/522,449	3 October 2004 (03.10.2004)	US
60/522,457	4 October 2004 (04.10.2004)	US
60/522,860	15 November 2004 (15.11.2004)	US
60/593,081	8 December 2004 (08.12.2004)	US
60/593,329	6 January 2005 (06.01.2005)	US
60/662,742	17 March 2005 (17.03.2005)	US
60/665,094	25 March 2005 (25.03.2005)	US
60/666,340	30 March 2005 (30.03.2005)	US

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
13 July 2006

(71) Applicant (*for all designated States except US*): **ROSETTA GENOMICS LTD.** [IL/IL]; 10 Plaut Street, 76706 Rehovot (IL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **BENTWICH, Itzhak** [IL/IL]; 76802 Kvutzat Shiler (IL). **AVNIEL, Amir** [IL/IL]; 10 Plaut Street, 76706 Rehovot (IL).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MICRONAS AND USES THEREOF

(57) Abstract: Described herein are novel polynucleotides associated with prostate and lung cancer. The polynucleotides are miRNAs and miRNA precursors. Related methods and compositions that can be used for diagnosis, prognosis, and treatment of those medical conditions are disclosed. Also described herein are methods that can be used to identify modulators of prostate and lung cancer.



WO 2005/111211 A3

MICRORNAS AND USES THEREOF

FIELD OF THE INVENTION

[0001] The invention relates in general to microRNA molecules as well as various nucleic acid molecules relating thereto or derived therefrom.

BACKGROUND OF THE INVENTION

[0002] MicroRNAs (miRNAs) are short RNA oligonucleotides of approximately 22 nucleotides that are involved in gene regulation. MicroRNAs regulate gene expression by targeting mRNAs for cleavage or translational repression. Although miRNAs are present in a wide range of species including *C. elegans*, *Drosophilla* and humans, they have only recently been identified. More importantly, the role of miRNAs in the development and progression of disease has only recently become appreciated.

[0003] As a result of their small size, miRNAs have been difficult to identify using standard methodologies. A limited number of miRNAs have been identified by extracting large quantities of RNA. MiRNAs have also been identified that contribute to the presentation of visibly discernable phenotypes. Expression array data shows that miRNAs are expressed in different developmental stages or in different tissues. The restriction of miRNAs to certain tissues or at limited developmental stages indicates that the miRNAs identified to date are likely only a small fraction of the total miRNAs.

[0004] Computational approaches have recently been developed to identify the remainder of miRNAs in the genome. Tools such as MiRscan and MiRseeker have identified miRNAs that were later experimentally confirmed. Based on these computational tools, it has been estimated that the human genome contains 200-255 miRNA genes. These estimates are based on an assumption, however, that the miRNAs remaining to be identified will have the same properties as those miRNAs already identified. Based on the fundamental importance of miRNAs in mammalian biology and disease, the art needs to identify unknown miRNAs. The present invention satisfies this need and provides a significant number of miRNAs and uses therefore.

SUMMARY OF THE INVENTION

[0005] The present invention is related to an isolated nucleic acid comprising a sequence of a pri-miRNA, pre-miRNA, miRNA, miRNA*, anti-miRNA, or a miRNA binding site, or a variant thereof. The nucleic acid may comprise the sequence of a hairpin referred to in Table 1; the sequence of sequence identifiers 6757248-6894882 of Table 10; the sequence of sequence identifiers 1-6318 or 18728-18960 of Table 17; the sequence of a miRNA referred to in Table 1; the sequence of sequence identifiers 1-117750 or 6894883-10068177 of Table 10; the sequence of sequence identifiers 6319-18727 or 18961-19401 of Table 17; the sequence of a target gene binding site referred to in Table 4; the sequence of sequence identifiers 117751-6757247 of Table 10; a complement thereof; or a sequence comprising at least 12 contiguous nucleotides at least 60% identical thereto. The isolated nucleic acid may be from 5-250 nucleotides in length.

[0006] The present invention is also related to a probe comprising the nucleic acid. The probe may comprise at least 8-22 contiguous nucleotides complementary to a miRNA referred to in Table 2 as differentially expressed in prostate cancer or lung cancer.

[0007] The present invention is also related to a plurality of the probes. The plurality of probes may comprise at least one probe complementary to each miRNA referred to in Table 2 as differentially expressed in prostate cancer. The plurality of probes may also comprise at least one probe complementary to each miRNA referred to in Table 2 as differentially expressed in lung cancer.

[0008] The present invention is also related to a composition comprising a probe or plurality of probes.

[0009] The present invention is also related to a biochip comprising a solid substrate, said substrate comprising a plurality of probes. Each of the probes may be attached to the substrate at a spatially defined address. The biochip may comprise probes that are complementary to a miRNA referred to in Table 2 as differentially expressed in prostate cancer. The biochip may also comprise probes that are complementary to a miRNA referred to in Table 2 as differentially expressed in lung cancer.

sequence identifiers 6319-18727 or 18961-19401 of Table 17; or variants thereof. A difference in the level of the nucleic acid compared to a control is indicative of differential expression.

[0011] The present invention is also related to a method of identifying a compound that modulates a pathological condition. A cell may be provided that is capable of expressing a nucleic acid at least 70% identical to a sequence of a miRNA referred to in Table 1; the sequence of sequence identifiers 1-117750 or 6894883-10068177 of Table 10; the sequence of sequence identifiers 6319-18727 or 18961-19401 of Table 17; or variants thereof. The cell may be contacted with a candidate modulator and then measuring the level of expression of the nucleic acid. A difference in the level of the nucleic acid compared to a control identifies the compound as a modulator of a pathological condition associated with the nucleic acid.

[0012] The present invention is also related to a method of inhibiting expression of a target gene in a cell. Into the cell, a nucleic acid may be introduced in an amount sufficient to inhibit expression of the target gene. The target gene may comprise a binding site substantially identical to a binding site referred to in Table 4; a sequence of sequence identifiers 117751-6757247 of Table 10; or a variant thereof. The nucleic acid may comprise a sequence of SEQ ID NOS: 1-760616; a sequence of sequence identifiers 1-117750 and 6757248-10068177 of Table 10; a sequence set forth on Table 17; or a variant thereof. Expression of the target gene may be inhibited in vitro or in vivo.

[0013] The present invention is also related to a method of increasing expression of a target gene in a cell. Into the cell, a nucleic acid may be introduced in an amount sufficient to inhibit expression of the target gene. The target gene may comprise a binding site substantially identical to a binding site referred to in Table 4; a sequence of sequence identifiers 117751-6757247 of Table 10; or a variant thereof. The nucleic acid may comprise a sequence substantially complementary to SEQ ID NOS: 1-760616; a sequence of sequence identifiers 1-117750 and 6757248-10068177 of Table 10; a sequence set forth on Table 17; or a variant thereof. Expression of the target gene may be inhibited in vitro or in vivo. Expression of the target gene may be increased in vitro or in vivo.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] Figure 1 demonstrates a model of maturation for miRNAs.

[0016] Figure 2 shows a schematic illustration of the MC19cluster on 19q13.42. Panel A shows the ~500,000bp region of chromosome 19, from 58,580,001 to 59,080,000 (according to the May 2004 USCS assembly), in which the cluster is located including the neighboring protein-coding genes. The MC19-1 cluster is indicated by a rectangle. Mir-371, mir-372, and mir-373 are indicated by lines. Protein coding genes flanking the cluster are represented by large arrow-heads. Panel B shows a detailed structure of the MC19-1 miRNA cluster. A region of ~102,000bp, from 58,860,001 to 58,962,000 (according to the May 2004 USCS assembly), is presented. MiRNA precursors are represented by a black bars. It should be noted that all miRNAs are at the same orientation from left to right. Shaded areas around miRNA precursors represent repeating units in which the precursor is embedded. The location of mir-371, mir-372, and mir-373, is also presented.

[0017] Figure 3 is a graphical representation of multiple sequence alignment of 35 human repeat units at distinct size of ~690nt (A) and 26 chimpanzees repeat units (B). The graph was generated by calculating a similarity score for each position in the alignment with an averaging sliding window of 10nt (Maximum score -1, minimum score-0). The repeat unit sequences were aligned by ClustalW program. Each position of the resulting alignment was assigned a score which represented the degree of similarity at this position. The region containing the miRNA precursors is bordered by vertical lines. The exact location of the mature miRNAs derived from the 5' stems (5p) and 3' stems (3p) of the precursors is indicated by vertical lines.

[0018] Figure 4 shows sequence alignments of the 43 A-type pre-miRNAs of the MC19-1 cluster. Panel A shows the multiple sequence alignment with the Position of the mature miRNAs marked by a frame. The consensus sequence is shown at the bottom. Conserved nucleotides are colored as follows: black-100%, dark grey- 80% to 99%, and clear grey- 60

mir-98 and ethidium bromide staining of the tRNA band served as control. Panel B shows RT-PCR analysis of the mRNA transcript containing the A-type miRNA precursors. Reverse transcription of 5 μ g total RNA from placenta was performed using oligo-dT. This was followed by PCR using the denoted primers (indicated by horizontal arrows). The region examined is illustrated at the top. Vertical black bars represent the pre-miRNA; shaded areas around the pre-miRNAs represent the repeating units; the location of four ESTs is indicated at the right side; the poly-A site, as found in the ESTs and located downstream to a AATAAA consensus, is indicated by a vertical arrow. The fragments expected from RT-PCR using three primer combinations are indicated below the illustration of the cluster region. The results of the RT-PCR analysis are presented below the expected fragments. Panel C shows the sequencing strategy of the FR2 fragment. The fragment was cloned into the pTZ57R/T vector and sequenced using external and internal primers.

DETAILED DESCRIPTION

[0020] The present invention provides nucleotide sequences of miRNAs, precursors thereto, targets thereof and related sequences. Such nucleic acids are useful for diagnostic purposes, and also for modifying target gene expression. Other aspects of the invention will become apparent to the skilled artisan by the following description of the invention.

1. Definitions

[0021] Before the present compounds, products and compositions and methods are disclosed and described, it is to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise.

a. animal

[0022] "Animal" as used herein may mean fish, amphibians, reptiles, birds, and mammals, such as mice, rats, rabbits, goats, cats, dogs, cows, apes and humans.

b. attached

[0023

covalent. Covalent bonds may be formed directly between the probe and the solid support or may be formed by a cross linker or by inclusion of a specific reactive group on either the solid support or the probe or both molecules. Non-covalent binding may be one or more of electrostatic, hydrophilic, and hydrophobic interactions. Included in non-covalent binding is the covalent attachment of a molecule, such as streptavidin, to the support and the non-covalent binding of a biotinylated probe to the streptavidin. Immobilization may also involve a combination of covalent and non-covalent interactions.

c. biological sample

[0024] "Biological sample" as used herein may mean a sample of biological tissue or fluid that comprises nucleic acids. Such samples include, but are not limited to, tissue isolated from animals. Biological samples may also include sections of tissues such as biopsy and autopsy samples, frozen sections taken for histologic purposes, blood, plasma, serum, sputum, stool, tears, mucus, hair, and skin. Biological samples also include explants and primary and/or transformed cell cultures derived from patient tissues. A biological sample may be provided by removing a sample of cells from an animal, but can also be accomplished by using previously isolated cells (e.g., isolated by another person, at another time, and/or for another purpose), or by performing the methods of the invention in vivo. Archival tissues, such as those having treatment or outcome history, may also be used.

d. complement

[0025] "Complement" or "complementary" as used herein may mean Watson-Crick or Hoogsteen base pairing between nucleotides or nucleotide analogs of nucleic acid molecules.

e. differential expression

[0026] "Differential expression" may mean qualitative or quantitative differences in the temporal and/or cellular gene expression patterns within and among cells and tissue. Thus, a differentially expressed gene can qualitatively have its expression altered, including an activation or inactivation, in, e.g., normal versus disease tissue. Genes may be turned on or turned off in a particular state, relative to another state thus permitting comparison of two or more states. A qualitatively regulated gene will exhibit an expression pattern within a state or cell type which may be detectable by standard techniques. Some genes will be expressed in one state or cell type, but not in both. Alternatively, the difference in expression may be quantitative, e.g., in that expression is mod

down-regulated, resulting in a decreased amount of transcript. The degree to which expression differs need only be large enough to quantify via standard characterization techniques such as expression arrays, quantitative reverse transcriptase PCR, northern analysis, and RNase protection.

f. gene

[0027] "Gene" used herein may be a genomic gene comprising transcriptional and/or translational regulatory sequences and/or a coding region and/or non-translated sequences (e.g., introns, 5'- and 3'-untranslated sequences). The coding region of a gene may be a nucleotide sequence coding for an amino acid sequence or a functional RNA, such as tRNA, rRNA, catalytic RNA, siRNA, miRNA and antisense RNA. A gene may also be an mRNA or cDNA corresponding to the coding regions (e.g., exons and miRNA) optionally comprising 5'- or 3'-untranslated sequences linked thereto. A gene may also be an amplified nucleic acid molecule produced in vitro comprising all or a part of the coding region and/or 5'- or 3'-untranslated sequences linked thereto.

g. host cell

[0028] "Host cell" used herein may be a naturally occurring cell or a transformed cell that contains a vector and supports the replication of the vector. Host cells may be cultured cells, explants, cells in vivo, and the like. Host cells may be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells, such as CHO, HeLa.

h. identity

[0029] "Identical" or "identity" as used herein in the context of two or more nucleic acids or polypeptide sequences, may mean that the sequences have a specified percentage of nucleotides or amino acids that are the same over a specified region. The percentage may be calculated by comparing optimally aligning the two sequences, comparing the two sequences over the specified region, determining the number of positions at which the identical residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the specified region, and multiplying the result by 100 to yield the percentage of sequence identity. In cases where the two sequences are of different lengths or the alignment produces staggered end and the specified region of comparison includes only a single sequence, the residues of single sequence

uracil (U) are considered equivalent. Identity may be performed manually or by using computer sequence algorithm such as BLAST or BLAST 2.0.

i. label

[0030] "Label" as used herein may mean a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, chemical, or other physical means. For example, useful labels include ^{32}P , fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, digoxigenin, or haptens and other entities which can be made detectable. A label may be incorporated into nucleic acids and proteins at any position.

j. nucleic acid

[0031] "Nucleic acid" or "oligonucleotide" or "polynucleotide" used herein may mean at least two nucleotides covalently linked together. As will be appreciated by those in the art, the depiction of a single strand also defines the sequence of the complementary strand. Thus, a nucleic acid also encompasses the complementary strand of a depicted single strand. As will also be appreciated by those in the art, many variants of a nucleic acid may be used for the same purpose as a given nucleic acid. Thus, a nucleic acid also encompasses substantially identical nucleic acids and complements thereof. As will also be appreciated by those in the art, a single strand provides a probe for a probe that may hybridize to the target sequence under stringent hybridization conditions. Thus, a nucleic acid also encompasses a probe that hybridizes under stringent hybridization conditions.

[0032] Nucleic acids may be single stranded or double stranded, or may contain portions of both double stranded and single stranded sequence. The nucleic acid may be DNA, both genomic and cDNA, RNA, or a hybrid, where the nucleic acid may contain combinations of deoxyribo- and ribo-nucleotides, and combinations of bases including uracil, adenine, thymine, cytosine, guanine, inosine, xanthine hypoxanthine, isocytosine and isoguanine. Nucleic acids may be obtained by chemical synthesis methods or by recombinant methods.

[0033] A nucleic acid will generally contain phosphodiester bonds, although nucleic acid analogs may be included that may have at least one different linkage

containing one or more non-naturally occurring or modified nucleotides are also included within one definition of nucleic acids. The modified nucleotide analog may be located for example at the 5'-end and/or the 3'-end of the nucleic acid molecule. Representative examples of nucleotide analogs may be selected from sugar- or backbone-modified ribonucleotides. It should be noted, however, that also nucleobase-modified ribonucleotides, i.e. ribonucleotides, containing a non-naturally occurring nucleobase instead of a naturally occurring nucleobase such as uridines or cytidines modified at the 5-position, e.g. 5-(2-amino)propyl uridine, 5-bromo uridine; adenosines and guanosines modified at the 8-position, e.g. 8-bromo guanosine; deaza nucleotides, e.g. 7-deaza-adenosine; O- and N-alkylated nucleotides, e.g. N6-methyl adenosine are suitable. The 2'-OH-group may be replaced by a group selected from H, OR, R, halo, SH, SR, NH₂, NHR, NR₂ or CN, wherein R is C₁-C₆ alkyl, alkenyl or alkynyl and halo is F, Cl, Br or I. Modifications of the ribose-phosphate backbone may be done for a variety of reasons, e.g., to increase the stability and half-life of such molecules in physiological environments or as probes on a biochip. Mixtures of naturally occurring nucleic acids and analogs may be made; alternatively, mixtures of different nucleic acid analogs, and mixtures of naturally occurring nucleic acids and analogs may be made.

k. operably linked

[0034] "Operably linked" used herein may mean that expression of a gene is under the control of a promoter with which it is spatially connected. A promoter may be positioned 5' (upstream) or 3' (downstream) of the gene under its control. The distance between the promoter and the gene may be approximately the same as the distance between that promoter and the gene it controls in the gene from which the promoter is derived. As is known in the art, variation in this distance can be accommodated without loss of promoter function.

l. probe

[0035] "Probe" as used herein may mean an oligonucleotide capable of binding to a target nucleic acid of complementary sequence through one or more types of chemical bonds, usually through complementary base pairing, usually through hydrogen bond formation. Probes may bind

that no hybridization can occur under even the least stringent of hybridization conditions, the sequence is not a complementary target sequence. A probe may be single stranded or partially single and partially double stranded. The strandedness of the probe is dictated by the structure, composition, and properties of the target sequence. Probes may be directly labeled or indirectly labeled such as with biotin to which a streptavidin complex may later bind.

m. promoter

[0036] "Promoter" as used herein may mean a synthetic or naturally-derived molecule which is capable of conferring, activating or enhancing expression of a nucleic acid in a cell. A promoter may comprise one or more specific regulatory elements to further enhance expression and/or to alter the spatial expression and/or temporal expression of same. A promoter may also comprise distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A promoter may be derived from sources including viral, bacterial, fungal, plants, insects, and animals. A promoter may regulate the expression of a gene component constitutively, or differentially with respect to cell, the tissue or organ in which expression occurs or, with respect to the developmental stage at which expression occurs, or in response to external stimuli such as physiological stresses, pathogens, metal ions, or inducing agents. Representative examples of promoters include the bacteriophage T7 promoter, bacteriophage T3 promoter, SP6 promoter, lac operator-promoter, tac promoter, SV40 late promoter, SV40 early promoter, RSV-LTR promoter, CMV IE promoter, SV40 early promoter or SV40 late promoter and the CMV IE promoter.

n. selectable marker

[0037] "Selectable marker" used herein may mean any gene which confers a phenotype on a cell in which it is expressed to facilitate the identification and/or selection of cells which are transfected or transformed with a genetic construct. Representative examples of selectable markers include the ampicillin-resistance gene (Amp^r), tetracycline-resistance gene (Tc^r), bacterial kanamycin-resistance gene (Kan^r), zeocin resistance gene, the AURI-C gene which confers resistance to the antibiotic aureobasidin A, phosphinothricin-resistance gene, neomycin phosphotransferase gene (nptII), hygromycin-resistance gene, beta-glucuronidase (GUS) gene, chloramphenicol acetyltransferase (CAT) gene, green fluorescent protein-encoding gene and luciferase gene.

o. stringent hybridization conditions

[0038] "Stringent hybridization conditions" used herein may mean conditions under which a first nucleic acid sequence (e.g., probe) will hybridize to a second nucleic acid sequence (e.g., target), such as in a complex mixture of nucleic acids, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5-10° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. The T_m may be the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T_m , 50% of the probes are occupied at equilibrium). Stringent conditions may be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01-1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., about 10-50 nucleotides) and at least about 60°C for long probes (e.g., greater than about 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal may be at least 2 to 10 times background hybridization. Exemplary stringent hybridization conditions include the following: 50% formamide, 5x SSC, and 1% SDS, incubating at 42°C, or, 5x SSC, 1% SDS, incubating at 65°C, with wash in 0.2x SSC, and 0.1% SDS at 65°C.

p. substantially complementary

[0039] "Substantially complementary" used herein may mean that a first sequence is at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% identical to the complement of a second sequence over a region of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50 or more nucleotides, or that the two sequences hybrid

r. target

[0041] "Target" as used herein may mean a polynucleotide that may be bound by one or more probes under stringent hybridization conditions.

s. terminator

[0042] "Terminator" used herein may mean a sequence at the end of a transcriptional unit which signals termination of transcription. A terminator may be a 3'-non-translated DNA sequence containing a polyadenylation signal, which may facilitate the addition of polyadenylate sequences to the 3'-end of a primary transcript. A terminator may be derived from sources including viral, bacterial, fungal, plants, insects, and animals. Representative examples of terminators include the SV40 polyadenylation signal, HSV TK polyadenylation signal, CYC1 terminator, ADH terminator, SPA terminator, nopaline synthase (NOS) gene terminator of *Agrobacterium tumefaciens*, the terminator of the Cauliflower mosaic virus (CaMV) 35S gene, the zein gene terminator from *Zea mays*, the Rubisco small subunit gene (SSU) gene terminator sequences, subclover stunt virus (SCSV) gene sequence terminators, rho-independent *E. coli* terminators, and the lacZ alpha terminator.

t. Vector

[0043] "Vector" used herein may mean a nucleic acid sequence containing an origin of replication. A vector may be a plasmid, bacteriophage, bacterial artificial chromosome or yeast artificial chromosome. A vector may be a DNA or RNA vector. A vector may be either a self-replicating extrachromosomal vector or a vector which integrate into a host genome.

2. MicroRNA

[0044] While not being bound by theory, the current model for the maturation of mammalian miRNAs is shown in Figure 1. A gene coding for a miRNA may be transcribed leading to production of an miRNA precursor known as the pri-miRNA. The pri-miRNA may be part of a polycistronic RNA comprising multiple pri-miRNAs.

overhang. Approximately one helical turn of stem (~10 nucleotides) extending beyond the Drosha cleavage site may be essential for efficient processing. The pre-miRNA may then be actively transported from the nucleus to the cytoplasm by Ran-GTP and the export receptor Exportin-5.

[0046] The pre-miRNA may be recognized by Dicer, which is also an RNase III endonuclease. Dicer may recognize the double-stranded stem of the pre-miRNA. Dicer may also recognize the 5' phosphate and 3' overhang at the base of the stem loop. Dicer may cleave off the terminal loop two helical turns away from the base of the stem loop leaving an additional 5' phosphate and ~2 nucleotide 3' overhang. The resulting siRNA-like duplex, which may comprise mismatches, comprises the mature miRNA and a similar-sized fragment known as the miRNA*. The miRNA and miRNA* may be derived from opposing arms of the pri-miRNA and pre-miRNA. MiRNA* sequences may be found in libraries of cloned miRNAs but typically at lower frequency than the miRNAs.

[0047] Although initially present as a double-stranded species with miRNA*, the miRNA may eventually become incorporated as single-stranded RNAs into a ribonucleoprotein complex known as the RNA-induced silencing complex (RISC). Various proteins can form the RISC, which can lead to variability in specificity for miRNA/miRNA* duplexes, binding site of the target gene, activity of miRNA (repress or activate), which strand of the miRNA/miRNA* duplex is loaded in to the RISC.

[0048] When the miRNA strand of the miRNA:miRNA* duplex is loaded into the RISC, the miRNA* may be removed and degraded. The strand of the miRNA:miRNA* duplex that is loaded into the RISC may be the strand whose 5' end is less tightly paired. In cases where both ends of the miRNA:miRNA* have roughly equivalent 5' pairing, both miRNA and miRNA* may have gene silencing activity.

[0049] The RISC may identify target nucleic acids based on high levels of complementarity between the miRNA and the mRNA, especially by nucleotides 2-8 of the miRNA. Only one case has been reported in animals where the interaction between the miRNA and its target was along the entire length of the miRNA. This was shown for mir-196 and Hox B8 and it was further shown that mir-196 mediates the cleavage of the Hox B8 mRNA (

[0050] A number of studies have looked at the base-pairing requirement between miRNA and its mRNA target for achieving efficient inhibition of translation (reviewed by Bartel 2004, Cell 116-281). In mammalian cells, the first 8 nucleotides of the miRNA may be important (Doench & Sharp 2004 GenesDev 2004-504). However, other parts of the microRNA may also participate in mRNA binding. Moreover, sufficient base pairing at the 3' can compensate for insufficient pairing at the 5' (Brennecke et al, 2005 PLoS 3-e85). Computation studies, analyzing miRNA binding on whole genomes have suggested a specific role for bases 2-7 at the 5' of the miRNA in target binding but the role of the first nucleotide, found usually to be "A" was also recognized (Lewis et al 2005 Cell 120-15). Similarly, nucleotides 1-7 or 2-8 were used to identify and validate targets by Krek et al (2005, Nat Genet 37-495).

[0051] The target sites in the mRNA may be in the 5' UTR, the 3' UTR or in the coding region. Interestingly, multiple miRNAs may regulate the same mRNA target by recognizing the same or multiple sites. The presence of multiple miRNA complementarity sites in most genetically identified targets may indicate that the cooperative action of multiple RISCs provides the most efficient translational inhibition.

[0052] MiRNAs may direct the RISC to downregulate gene expression by either of two mechanisms: mRNA cleavage or translational repression. The miRNA may specify cleavage of the mRNA if the mRNA has a certain degree of complementarity to the miRNA. When a miRNA guides cleavage, the cut may be between the nucleotides pairing to residues 10 and 11 of the miRNA. Alternatively, the miRNA may repress translation if the miRNA does not have the requisite degree of complementarity to the miRNA. Translational repression may be more prevalent in animals since animals may have a lower degree of complementarity.

[0053] It should be noted that there may be variability in the 5' and 3' ends of any pair of miRNA and miRNA*. This variability may be due to variability in the enzymatic processing of Drosha and Dicer with respect to the site of cleavage. Variability at the 5' and 3' ends of miRNA and miRNA* may also be due to mismatches in the stem structures of the pri-miRNA and pre-miRNA. The mismatches of the stem strands may lead to a population of different hairpin structures. Variability in the stem structures may also lead to variability in

3. Nucleic Acid

[0054] The present invention relates to an isolated nucleic acid comprising a nucleotide sequence referred to in SEQ ID NOS: 1-760616, the sequences set forth on Table 10, and the sequences set forth on Table 17, or variants thereof. The variant may be a complement of the referenced nucleotide sequence. The variant may also be a nucleotide sequence that is substantially identical to the referenced nucleotide sequence or the complement thereof. The variant may also be a nucleotide sequence which hybridizes under stringent conditions to the referenced nucleotide sequence, complements thereof, or nucleotide sequences substantially identical thereto.

[0055] The nucleic acid may have a length of from 10 to 100 nucleotides. The nucleic acid may have a length of at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 60, 70, 80 or 90 nucleotides. The nucleic acid may be synthesized or expressed in a cell (in vitro or in vivo) using a synthetic gene described below. The nucleic acid may be synthesized as a single strand molecule and hybridized to a substantially complementary nucleic acid to form a duplex, which is considered a nucleic acid of the invention. The nucleic acid may be introduced to a cell, tissue or organ in a single- or double-stranded form or capable of being expressed by a synthetic gene using methods well known to those skilled in the art, including as described in U.S. Patent No. 6,506,559 which is incorporated by reference.

a. Pri-miRNA

[0056] The nucleic acid of the invention may comprise a sequence of a pri-miRNA or a variant thereof. The pri-miRNA sequence may comprise from 45-250, 55-200, 70-150 or 80-100 nucleotides. The sequence of the pri-miRNA may comprise a pre-miRNA, miRNA and miRNA* as set forth below. The pri-miRNA may also comprise a miRNA or miRNA* and the complement thereof, and variants thereof. The pri-miRNA may comprise at least 19% adenosine nucleotides, at least

described in Hofacker et al., Monatshefte f. Chemie 125: 167-188 (1994), the contents of which are incorporated herein. The hairpin may comprise a terminal loop of 4-20, 8-12 or 10 nucleotides.

[0058] The sequence of the pri-miRNA may comprise the sequence of a hairpin referred to in Table 1, the sequence of sequence identifiers 6757248-6894882 of Table 10, the sequence of sequence identifiers 1-6318 or 18728-18960 of Table 17, or variants thereof.

b. Pre-miRNA

[0059] The nucleic acid of the invention may also comprise a sequence of a pre-miRNA or a variant thereof. The pre-miRNA sequence may comprise from 45-90, 60-80 or 60-70 nucleotides. The sequence of the pre-miRNA may comprise a miRNA and a miRNA* as set forth below. The pre-miRNA may also comprise a miRNA or miRNA* and the complement thereof, and variants thereof. The sequence of the pre-miRNA may also be that of a pri-miRNA excluding from 0-160 nucleotides from the 5' and 3' ends of the pri-miRNA.

[0060] The sequence of the pre-miRNA may comprise the sequence of a hairpin referred to in Table 1, the sequence of sequence identifiers 6757248-6894882 of Table 10, the sequence of sequence identifiers 1-6318 or 18728-18960 of Table 17, or variants thereof.

c. MiRNA

[0061] The nucleic acid of the invention may also comprise a sequence of a miRNA, miRNA* or a variant thereof. The miRNA sequence may comprise from 13-33, 18-24 or 21-23 nucleotides. The sequence of the miRNA may be the first 13-33 nucleotides of the pre-miRNA. The sequence of the miRNA may be the last 13-33 nucleotides of the pre-miRNA.

regions of the target site from the 5' end of said miRNA, or (b) at least 5-12 nucleotides that are substantially identical to the 3' of a miRNA and at least 5 nucleotide that are substantially complementary to the flanking region of the target site from the 3' end of said miRNA.

[0064] The sequence of the anti-miRNA may comprise the complement of a sequence of a miRNA referred to in Table 1, the sequence of sequence identifiers 1-117750 or 6894883-10068177 of Table 10, the sequence of sequence identifiers 6319-18727 or 18961-19401 of Table 17, or variants thereof.

e. Binding Site of Target

[0065] The nucleic acid of the invention may also comprise a sequence of a target miRNA binding site, or a variant thereof. The target site sequence may comprise a total of 5-100 or 10-60 nucleotides. The target site sequence may comprise at least 5 nucleotides of the sequence of a target gene binding site referred to in Table 4, the sequence of sequence identifiers 117751-6757247 of Table 10, or variants thereof.

4. Synthetic Gene

[0066] The present invention also relates to a synthetic gene comprising a nucleic acid of the invention operably linked to a transcriptional and/or translational regulatory sequences. The synthetic gene may be capable of modifying the expression of a target gene with a binding site for the nucleic acid of the invention. Expression of the target gene may be modified in a cell, tissue or organ. The synthetic gene may be synthesized or derived from naturally-occurring genes by standard recombinant techniques. The synthetic gene may also comprise terminators at the 3'-end of the transcriptional unit of the synthetic gene sequence. The synthetic gene may also comprise a selectable marker.

5. Vector

[0067] The present invention also relates to a vector comprising a synthetic gene of the invention. The vector may be an expression vector. An expression vector may comprise additional elements. For example,

homologous sequence for inclusion in the vector. The vector may also comprise a selectable marker gene to allow the selection of transformed host cells.

6. Host Cell

[0068] The present invention also relates to a host cell comprising a vector of the invention. The cell may be a bacterial, fungal, plant, insect or animal cell.

7. Probes

[0069] The present invention also relates to a probe comprising a nucleic acid of the invention. Probes may be used for screening and diagnostic methods, as outlined below. The probe may be attached or immobilized to a solid substrate, such as a biochip.

[0070] The probe may have a length of from 8 to 500, 10 to 100 or 20 to 60 nucleotides. The probe may also have a length of at least 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 220, 240, 260, 280 or 300 nucleotides. The probe may further comprise a linker sequence of from 10-60 nucleotides.

8. Biochip

[0071] The present invention also relates to a biochip. The biochip may comprise a solid substrate comprising an attached probe or plurality of probes of the invention. The probes may be capable of hybridizing to a target sequence under stringent hybridization conditions. The probes may be attached at spatially defined address on the substrate. More than one probe per target sequence may be used, with either overlapping probes or probes to different sections of a particular target sequence. The probes may be

CLAIMS

1. An isolated nucleic acid comprising a sequence selected from the group consisting of:

- (a) the sequence of a hairpin referred to in Table 1;
- (b) the sequence of sequence identifiers 6757248-6894882 of Table 10;
- (c) the sequence of sequence identifiers 1-6318 or 18728-18960 of Table 17;
- (d) the sequence of a miRNA referred to in Table 1;
- (e) the sequence of sequence identifiers 1-117750 or 6894883-10068177 of Table 10;
- (f) the sequence of sequence identifiers 6319-18727 or 18961-19401 of Table 17;
- (g) the sequence of a target gene binding site referred to in Table 4;
- (h) the sequence of sequence identifiers 117751-6757247 of Table 10;
- (i) complement of (a)-(h);
- (j) nucleotide sequence comprising at least 12 contiguous nucleotides at least 60% identical to (a)-(h);

wherein the nucleic acid is from 5-250 nucleotides in length.

2. A probe comprising the nucleic acid of claim 1.

3. The probe of claim 2 wherein the nucleic acid comprises at least 8-22 contiguous nucleotides complementary to a miRNA referred to in Table 2 as differentially expressed in prostate cancer or lung cancer.

4. A plurality of probes of claim 3.

5. The plurality of probes of claim 4 comprising at least one probe complementary to each miRNA referred to in Table 2 as differentially expressed in prostate cancer.

6. The plurality of probes of claim 4 comprising at least one probe complementary

9. The biochip of claim 8 wherein the probes are complementary to a miRNA referred to in Table 2 as differentially expressed in prostate cancer.

10. The biochip of claim 8 wherein the probes are complementary to a miRNA referred to in Table 2 as differentially expressed in lung cancer.

11. A method for detecting differential expression of a disease-associated miRNA comprising:

- (a) providing a biological sample; and
- (b) measuring the level of a nucleic acid at least 70% identical to (i) a sequence of a miRNA referred to in Table 1, (ii) the sequence of sequence identifiers 1-117750 or 6894883-10068177 of Table 10, (iii) the sequence of sequence identifiers 6319-18727 or 18961-19401 of Table 17, or (iv) variants of (i)-(iii),

wherein a difference in the level of the nucleic acid compared to a control is indicative of differential expression.

12. A method for identifying a compound that modulates a pathological condition comprising:

- (a) providing a cell is capable of expressing a nucleic acid at least 70% identical to (i) a sequence of a miRNA referred to in Table 1, (ii) the sequence of sequence identifiers 1-117750 or 6894883-10068177 of Table 10, (iii) the sequence of sequence identifiers 6319-18727 or 18961-19401 of Table 17, or (iv) variants of (i)-(iii);
- (b) contacting the cell with a candidate modulator;
- (c) measuring the level of expression of the nucleic acid,

wherein a difference in the level of the nucleic acid compared to a control identifies the compound

(b) sequence of sequence identifiers 1-117750 and 6757248-10068177 of Table 10, (c) set forth on Table 17, or (d) a variant of (a)-(c).


14. The method of claim 12 wherein expression is inhibited in vitro or in vivo.

15. A method of increasing expression of a target gene in a cell comprising introducing a nucleic acid into the cell in an amount sufficient to increase expression of the target gene, wherein the target gene comprises a (i) binding site substantially identical to a binding site referred to in Table 4, (ii) sequence of sequence identifiers 117751-6757247 of Table 10, or (iii) a variant of (i) or (ii); wherein the nucleic acid comprises a sequence substantially complementary to a sequence (a) of SEQ ID NOS: 1-760616, (b) set forth on Table 10, (c) set forth on Table 17, or (d) a variant of (a)-(c).

16. The method of claim 15 wherein expression is inhibited in vitro or in vivo.


17. A method of treating a patient with a disorder set forth on Table 6 comprising administering to a patient in need thereof a nucleic acid comprising a sequence (a) of SEQ ID NOS: 1-760616, (b) set forth on Table 10, (c) set forth on Table 17, or (d) a variant of (a)-(c).

FIGURE 1



SHEET 2/5

FIGURE 2



SHEET 3/5

FIGURE 3




FIGURE 5

