

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3004875 T3**

(12) **Oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **G 01 N 33/50 (2006.01)** **G 01 N 33/574 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-06-08**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-03-04**
- (86) Europæisk ansøgning nr.: **14725114.4**
- (86) Europæisk indleveringsdag: **2014-05-15**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-04-13**
- (86) International ansøgning nr.: **EP2014059995**
- (87) Internationalt publikationsnr.: **WO2014191222**
- (30) Prioritet: **2013-05-30 GB 201309657**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Mologen AG, Fabeckstrasse 30, 14195 Berlin, Tyskland**
- (72) Opfinder: **SCHROFF, Matthias, Schützallee 136, 14196 Berlin, Tyskland**
KAPP, Kerstin, Dolziger Str. 7, 10247 Berlin, Tyskland
WITTIG, Burghardt, Salzachstr. 33, 14129 Berlin, Tyskland
SCHMIDT, Manuel, Tölzer Str. 31, 14199 Berlin, Tyskland
- (74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Bredgade 30, 1260 København K, Danmark**
- (54) Benævnelse: **PRÆDIKTIV BIOMARKØR TIL CANCERTERAPI**
- (56) Fremdragne publikationer:
WO-A1-2010/055340
HANS-JOACHIM SCHMOLL ET AL: "Maintenance treatment with the immunomodulator MGN1703, a Toll-like receptor 9 (TLR9) agonist, in patients with metastatic colorectal carcinoma and disease control after chemotherapy: a randomised, double-blind, placebo-controlled trial", JOURNAL OF CANCER RESEARCH AND CLINICAL ONCOLOGY, 10 May 2014 (2014-05-10), XP055131369, ISSN: 0171-5216, DOI: 10.1007/s00432-014-1682-7
H.-J. SCHMOLL ET AL: "O-0012 * UPDATED RESULTS OF THE RANDOMIZED PHASE 2 IMPACT TRIAL: MAINTENANCE WITH TLR-9 AGONIST MGN1703 VS PLACEBO IN PATIENTS WITH METASTATIC COLORECTAL CARCINOMA (mCRC)", ANNALS OF ONCOLOGY, vol. 24, no. suppl 4, 1 June 2013 (2013-06-01), pages iv16-iv17, XP055131314, ISSN: 0923-7534, DOI: 10.1093/annonc/mdt201.12
Anonymous: "Maintenance with the TLR-9 agonist MGN1703 versus placebo in patients with advanced colorectal carcinoma (mCRC): A randomized phase II trial (IMPACT). -- Riera-Knorrenschild et al. 31 (15): 3643 -- ASCO Meeting Abstracts", , 20 May 2013 (2013-05-20), XP055131313, Retrieved from the Internet: URL:http://meeting.ascopubs.org/cgi/content/abstract/31/15_suppl/3643 [retrieved on 2014-07-23]
Anonymous: "A novel iNKT-mediated vaccine maneuver overcomes PD-1-mediated T cell exhaustion in cancer -- Varghese et al. 188 (1001): 165.13 -- The Journal of Immunology", , 1 January 2012 (2012-01-01),

Fortsættes ...

XP055131252, Retrieved from the Internet: URL:http://www.jimmunol.org/cgi/content/meeting_abstract/188/1_MeetingAbstracts/165. 13 [retrieved on 2014-07-23]

MONTOYA C J ET AL: "Increased IFN-gamma production by NK and CD3+/CD56+ cells in sexually HIV-1-exposed but uninfected individuals", *CLINICAL IMMUNOLOGY*, ACADEMIC PRESS, US, vol. 120, no. 2, 1 August 2006 (2006-08-01), pages 138-146, XP024941265, ISSN: 1521-6616, DOI: 10.1016/J.CLIM.2006.02.008 [retrieved on 2006-08-01]

J. VAS ET AL: "Regulatory Roles for NKT Cell Ligands in Environmentally Induced Autoimmunity", *THE JOURNAL OF IMMUNOLOGY*, vol. 181, no. 10, 3 November 2008 (2008-11-03), pages 6779-6788, XP055131246, ISSN: 0022-1767, DOI: 10.4049/jimmunol.181.10.6779

Lan Wu ET AL: "Natural Killer T Cells and Autoimmune Disease", *CURRENT MOLECULAR MEDICINE*, vol. 9, no. 1, 1 February 2009 (2009-02-01), pages 4-14, XP055307021, NL ISSN: 1566-5240, DOI: 10.2174/156652409787314534

OREN A ET AL: "A comparative study of immunomagnetic methods used for separation of human natural killer cells from peripheral blood", *JOURNAL OF IMMUNOLOGICAL METHODS*, ELSEVIER SCIENCE PUBLISHERS B.V., AMSTERDAM, NL, vol. 303, no. 1-2, 1 August 2005 (2005-08-01), pages 1-10, XP027659286, ISSN: 0022-1759 [retrieved on 2005-08-01]

Gitte S. Jensen ET AL: "Immunomodulation by SanPharma Fungal Metabolic Products", *JOURNAL OF ALTERNATIVE AND COMPLEMENTARY MEDICINE*, vol. 12, no. 4, 24 May 2006 (2006-05-24), pages 409-416, XP055307016, US ISSN: 1075-5535, DOI: 10.1089/acm.2006.12.409

DESCRIPTION

FIELD OF THE INVENTION

[0001] The present invention relates generally to the identification of cancer patients as to whether they will respond to specific therapies. More particularly the invention relates to a method and DNA construct for identifying responders to a therapy involving immune activating DNA constructs, as well as to a DNA construct intended to treat cancer patients individually characterized as potential responders prior to the start of a treatment with said construct.

BACKGROUND OF THE INVENTION

[0002] Biomarkers are substances found in blood or other body fluids, in tissues or as receptors on cells that signal the presence or absence of a condition or disease, like cancer for instance. Biomarkers can be differentiated into predictive, prognostic and pharmacodynamics biomarkers.

[0003] Predictive biomarkers are used to assess the probability that a patient will respond to or benefit from a particular treatment. Prognostic biomarkers allow for the classification of tumours with regard to their aggressive potential to guide a decision of whom to treat or how aggressively to treat. Pharmacodynamic biomarkers measure the near-term treatment effects of a drug on the tumour and might be used to guide dose selection during clinical development of a new anti-cancer drug.

[0004] Targeted therapies are treatments that work on a molecular level to stop a cancer from growing or spreading. In order to avoid unnecessary approaches, because such therapies are usually related to the use of very aggressive drugs, there is a need for tools allowing to predict whether a patient will respond to a targeted therapy or not.

[0005] With regards to the fact that therapies become more and more target specific, the role of biomarkers will increase. They will help to individualise therapeutic approaches and to open the way to individualised oncology. Even the possibility to distinguish a responder to a specific therapy from a non-responder will help to prevent that patients will be subject to unnecessary treatments.

[0006] A commonly known and well-approved treatment of cancer is chemotherapy. One major disadvantage of chemotherapy is the use of very aggressive pharmaceuticals causing severe side effects. In order to minimize these side effects many attempts have been undertaken to minimize dosage of chemotherapeutics like for instance combining chemotherapy with immune activating agents. One approach is the use of immune activating DNA.

[0007] So-called unmethylated CG sequences have been shown to activate the immune system very effectively (Krieg AM, Yi AK, Matson S, Waldschmidt TJ, Bishop GA, Teasdale R, Koretzky GA, Klinman DM; CpG motifs in bacterial DNA trigger direct B-cell activation; Nature 1995 Apr 6 374:6522 546-9). Those sequences are derived from bacteria. EP 1 196 178 discloses a covalently closed circular DNA with partially self-complementary sequences resulting in a DNA construct having a double stranded stem with single stranded loops at both ends comprising the unmethylated CG motifs.

[0008] The combination of the dumbbell-shaped DNA constructs of EP 1 196 178 with chemotherapeutics to treat cancerous diseases has been proposed by EP 1 776 124. Patients who were treated with the DNA constructs of EP 1 196 178 received subsequently chemotherapeutic. It turned out that the amount of chemotherapeutic could be reduced by the disclosed combination. But this document does not disclose information about tools for identifying whether a patient will respond to the application of the combination therapy at all.

[0009] It is an object of the present invention to provide a method for predicting whether a cancer patient will respond to the treatment with immune activating DNA. Another object of the invention is the immune activating DNA itself, which can be used in such a method, i.e. to treat a patient identified as responder prior to the start of treatment.

[0010] Riera-Knorrenschild et al. (31 (15): 3643 - ASCO Meeting Abstracts, 20 May 2013) disclose that the synthetic DNA-based TLR-9 agonist MGN1703 prolongs progression free survival (PFS) in a subgroup of advanced colorectal carcinoma patients with disease control after induction chemotherapy vs placebo. The document does not disclose a predictive biomarker.

[0011] Varghese et al. (188(1001): 165.13 - The Journal of Immunology, (2012)) disclose that invariant NKT (iNKT) cell stimulation using the ligand alpha-galactosylceramide, combined with TLR9 stimulation using the ligand CpG can induce long-lasting therapeutic immune responses in mice with large advanced (>1cm²) tumors. The document refers to the iNKT cell subgroup only.

[0012] Document WO 2010/055340 discloses a method of protecting a mammalian subject against, or treating, a disease, wherein the mammalian subject has elevated numbers and/or activities of myeloid-derived suppressor cell (MDSC) comprising administering to the subject a pharmaceutically acceptable amount of an iNKT agonist, such as alpha galactosylceramide or an analogue thereof, or a TLR agonist, or a combination thereof. The document refers to the iNKT cell subgroup only

[0013] Montoya C. et al. (Clinical Immunology, Academic press, US, vol. 120(2):138-146 (2006)) discloses CpG ODN induced CD86 and CD69 up-regulation on innate cells.

BRIEF SUMMARY OF THE INVENTION

[0014] The present invention provides a method for predicting or monitoring whether a patient suffering from cancer or an autoimmune disease will respond or responds to treatment with a TLR-9 agonist by determining the frequency of CD3+/CD56+/CD69+ activated natural killer T (NKT) cells, wherein the TLR-9 agonist is a DNA construct comprising at least one sequence motif $N^1N^2CGN^3N^4$, wherein N^1N^2 and N^3N^4 is any combination of A, C, T, and G, and C is deoxycytidine, G is deoxyguanosine, A is deoxyadenosine and T is deoxythymidine. The method is not carried out in vivo. The ratio of activated NKT cells of the whole NKT cell population may also be determined in order to assess the probability whether a patient will respond to a treatment with a TLR-9 agonist.

[0015] A responder to the treatment with the TLR-9 agonist shall have a frequency of at least 3 % of activated NKT cells of the whole NKT cell population.

[0016] It is further intended that previously an induction therapy with a non-DNA drug may be performed. Such a therapy comprises chemotherapy with or without angiogenesis inhibitor or the use of antibodies.

[0017] A further object of the present disclosure is a TLR-9 agonist for use in a method of treatment of cancer, agonist comprises at least a DNA construct comprising at least one sequence motif $N^1N^2CGN^3N^4$, wherein N^1N^2 and N^3N^4 is any combination of A, C, T, and G, and C is deoxycytidine, G is deoxyguanosine, A is deoxyadenosine and T is deoxythymidine and characterised in that the patient has elevated levels of CD3+/CD56+/CD69 activated NKT cells.

[0018] The agonist for use in the method of treatment of cancer may further be characterised by N^1N^2 being an element selected from the group comprising GT, GG, GA, AT or AA, N^3N^4 being an element selected from the group comprising CT or TT.

[0019] The agonist for use in the method of treatment of cancer may comprise DNA that is a linear open-chained DNA construct comprising single or double-stranded DNA or is a linear double-stranded DNA construct, which comprises at least one end with a single stranded loop. Instead of using loops to protect the ends of the linear double strands at least one L-DNA nucleotide may be used as part of the DNA double strand.

[0020] The sequence motif $N^1N^2CGN^3N^4$ shall be located within a single-stranded and/or a double-stranded region of the DNA sequence. It is further intended that at least one nucleotide of a DNA TLR-9 agonist is modified with a functional group selected from the group comprising carboxyl, amine, amide, aldimine, ketal, acetal, ester, ether, disulfide, thiol and aldehyde groups.

[0021] The agonist for use in the method of treatment of cancer comprising a DNA construct can be linked to a compound selected from the group comprising peptides, proteins,

carbohydrates, antibodies, lipids, micelles, vesicles, synthetic molecules, polymers, micro projectiles, metal particles, nanoparticles, or a solid phase.

[0022] It is further intended that the agonist for use in the method of treatment of cancer comprising a DNA construct can be part of a pharmaceutical composition, wherein the pharmaceutical composition can be a vaccine.

[0023] Another object of the present disclosure is the use of a kit for predicting or monitoring whether a patient suffering from cancer or auto immune disease will respond or responds to treatment with a TLR-9 agonist comprising means for detecting and quantifying the frequency of CD3+ CD56+ CD69+ activated NKT cells of the whole NKT cell population, and wherein the TLR-9 agonist comprises a DNA construct comprising at least one sequence motif $N^1N^2CGN^3N^4$, wherein N^1N^2 and N^3N^4 is any combination of A, C, T, and G, and C is deoxycytidine, G is deoxyguanosine, A is deoxyadenosine and T is deoxythymidine. The kit is not used in vivo.

BRIEF SUMMARY OF THE FIGURES

[0024]

Figure 1:

ROC curves in verum patients for activated NKT cells.

Figure 2:

Activated NKT cells versus sensitivity and specificity.

Figure 3:

a: Kaplan-Meier plot in biomarker positive patients for verum and placebo; b: Kaplan-Meier curve for the biomarker negative patients in both the verum and placebo arm.

Figure 4:

a: Kaplan-Meier plot of biomarker positive patients versus biomarker negative patients in the verum arm; b: Kaplan-Meier plot of the placebo group for biomarker positive and negative patients.

DETAILED DESCRIPTION OF THE INVENTION

[0025] Within the meaning of the present invention MGN1703 designates a DNA construct of a covalently closed partially self-complementary DNA chain having a double stranded stem and single stranded terminal loops bearing unmethylated CG motifs.

[0026] A "stem" according to the present disclosure shall be understood as a DNA double strand formed by base pairing either within the same DNA molecule (which is then partially

self-complementary) or within different DNA molecules (which are partially or completely complementary). Intramolecular base pairing designates base pairing within the same molecules, and base pairing between different DNA molecules is termed as intermolecular base-pairing.

[0027] A "loop" within the meaning of the present disclosure shall be understood as an unpaired, single-stranded region either within or at the end of a stem structure. A "hairpin" is a distinct combination of a stem and a loop, which occurs when two self-complementary regions of the same DNA molecule hybridize to form a stem with an unpaired loop.

[0028] A "dumbbell-shape" describes a linear DNA construct with hairpins at both ends flanking a stem region. Thus, a "linear DNA construct" within the context of the present disclosure describes a linear dumbbell-shaped DNA construct comprising single stranded loops at both ends of a double stranded DNA stem.

[0029] Immunomodulation according to the present disclosure refers to immune activation and immunosuppression. Immune activation means preferentially that effector cells of the immune system are activated in order to proliferate, migrate, differentiate or become active in any other form. B cell proliferation for instance can be induced without co-stimulatory signals by immune activating DNA molecules, which normally require a co-stimulatory signal from helper T-cells.

[0030] Immunosuppression on the other hand shall be understood as reducing the activation or efficacy of the immune system. Immunosuppression is generally deliberately induced to prevent for instance the rejection of a transplanted organ, to treat graft-versus-host disease after a bone marrow transplant, or for the treatment of autoimmune diseases such as, for example, rheumatoid arthritis or Crohn's disease.

[0031] In this context, immunomodulation may also refer to influencing the nature or the character of an immune reaction, either by affecting an immune reaction, which is still developing or maturing or by modulating the character of an established immune reaction.

[0032] The term "vaccination" used in this disclosure refers to the administration of antigenic material (a vaccine) to produce immunity to a disease. Vaccines can prevent or ameliorate the effects of infection by many pathogens such as viruses, fungi, protozoan parasites, bacteria but also of allergic diseases and asthma, as well as of tumors. Vaccines typically contain one or more adjuvants, e.g. immune activating nucleic acids, used to boost the immune response. Vaccination is generally considered to be the most effective and cost-effective method of preventing infectious and other diseases.

[0033] The material administered can, for example, be live but weakened forms of pathogens (bacteria or viruses), killed or inactivated forms of these pathogens, purified material such as proteins, nucleic acids encoding antigens, or cells such as tumor cells or dendritic cells. In particular, DNA vaccination has recently been developed. DNA vaccination works by insertion (and expression, triggering immune system recognition) of DNA encoding antigens into human

or animal cells. Some cells of the immune system that recognize the proteins expressed will mount an attack against these proteins and against cells expressing them. One advantage of DNA vaccines is that they are very easy to produce and store. In addition, DNA vaccines have a number of advantages over conventional vaccines, including the ability to induce a wider range of immune response types.

[0034] Vaccination can be used as a prophylactic approach, leading to immunity against the antigen in the vaccinated, healthy individual upon exposure to the antigen. Alternatively, a therapeutic vaccination can cause an improved response of the immune system of the vaccinated, diseased individual, by guiding the immune system of the individual towards the antigens. Both prophylactic and therapeutic vaccination can be applied to humans as well as animals.

[0035] The term "cancer" comprises cancerous diseases or a tumor being treated or prevented that is selected from the group comprising, but not limited to, mammary carcinomas, melanoma, skin neoplasms, lymphoma, leukemia, gastrointestinal tumors, including colon carcinomas, stomach carcinomas, pancreas carcinomas, colon cancer, small intestine cancer, ovarian carcinomas, cervical carcinomas, lung cancer, prostate cancer, kidney cell carcinomas and/or liver metastases.

[0036] Autoimmune diseases according to the present disclosure comprise rheumatoid arthritis, Crohn's disease, systemic lupus (SLE), autoimmune thyroiditis, Hashimoto's thyroiditis, multiple sclerosis, Graves' disease, myasthenia gravis, celiac disease and Addison's disease.

[0037] During experiments with immune activating dumbbell-shaped DNA constructs with unmethylated CG sequences in the single-stranded terminal loops it turned out that elevated frequencies of certain cell types of the immune activation and effector pathway were related to a successful therapy with the DNA constructs.

[0038] In vertebrates, the so-called "toll-like receptors" (TLRs) are part of the innate immune system. TLRs are a family of specialized immune receptors that induce protective immune responses when they detect highly conserved pathogen-related molecular patterns, such as proteins, lipid structures, saccharidic structures, and certain nucleic acids. Synthetic agonists for several TLRs, including TLR-3, TLR-4, TLR-7, TLR-8, and TLR-9, have been or are being developed for the treatment of cancer, generally with the intention to activate the immune system in the presence of tumours. TLR-9 recognizes the presence of unmethylated CG-containing DNA sequences, which are typically found in bacteria, but practically never in human genomic DNA. Thus, unmethylated CG-containing DNA sequences have been designed as artificial TLR-9 agonists. The effect of such unmethylated CG-containing DNA constructs depends on their interaction with TLR-9, and DNA-protein interaction depends on the conformation of both DNA and protein. Experimental data demonstrate that dumbbell-shaped DNA molecules are surprisingly suitable for the induction of an immune response.

[0039] In order to identify a tool for predicting whether a cancer patient will respond to the application of a TLR-9 agonist, a dumbbell-shape DNA construct having unmethylated CG motifs in the single stranded terminal loops was used.

[0040] In total, 46 patients suffering from metastatic colorectal cancer who had previously been treated for 4.5 to 6 month in a standard first-line combination therapy with or without a human monoclonal antibody inhibiting vascular endothelial growth factor A were selected for the study. After a treatment-free interval of ca. 1-6 weeks, a randomization of the patients was performed, so that 32 patients received 60 mg per dose of the DNA construct MGN1703 and 14 patients received a placebo, each patient twice weekly by subcutaneous injection.

[0041] After 12 weeks of treatment, all patients were examined for tumour-progression. Based on the presence or absence of tumour progression, patients were divided into two groups, designated as "PFS groups". Patients whose tumour had not progressed were designated as progression-free patients and labelled "PFS1", while patients with tumour progression were labelled as "PFS0". Obviously, a lack of tumour progression (PFS1) indicates a possible response to the treatment, while tumour growth (PFS2) indicates a lack of response. Treatment was continued for each patient until tumour progression was found.

[0042] Table 1 summarizes the results. It is obvious that nearly all progression-free patients received the DNA construct.

Table 1: Designation of patients

Treatment arms	Group designation		
	PFS0	PFS1	Total
MGN1703	19	13	32
Placebo	13	1	14
Total	32	14	46

[0043] Prior to the first application of MGN1703 or placebo (at baseline) blood samples of all patients were collected. The distribution of the following immunovables, e.g. part of whole PBMC population and relation of activated and non-activated sub populations was determined. A correlation of the respective PFS group designation after 12 weeks and all immunovables at baseline was investigated. The following immunovables were determined: monocytes , activated monocytes 1 ,activated monocytes 2, B-Cells, activated B-cells, T-cells, activated T-cells, natural killer (NK)-cells, activated NK-cells, NKT-cells, activated NKT-cells, plasmacytoid dendritic cells (pDCs), activated pDCs, myeloid dendritic cells (mDCs), and activated mDCs. Table 2 summarizes the context of cell types, CDs and the determined frequencies.

Table 2: Relation of cell types, CDs and determined frequencies

Cell Type	Identified by	Frequency as percentage of
monocytes	CD14+	all PBMCs

Cell Type	Identified by	Frequency as percentage of
activated monocytes 1	CD14+ / CD86+	all monocytes
activated monocytes 2	CD14+ / CD169+	all monocytes
B-cells	CD19+	all PBMCs
activated B-cells	CD19+ / CD86+	all B-cells
T-cells	CD3+ / CD56-	all PBMCs
activated T-cells	CD3+ / CD56- / CD69+	all T-cells
natural killer (NK-)cells	CD3- / CD56+	all PBMCs
activated NK-cells	CD3- / CD56+ / CD69+	all natural killer cells
NKT cells	CD3+ / CD56+	all PBMCs
activated NKT cells	CD3+ / CD56+ / CD69+	all NKT cells
plasmacytoid dendritic cells (pDCs)	Lin1- / CD123+ / HLA-DR+	all PBMCs
activated pDCs	Lin1- / CD123+ / HLA-DR+ / CD40+	all pDCs
myeloid dendritic cells (mDCs)	Lin1- / CD11c+ / HLA-DR+	all PBMCs
activated mDCs	Lin1- / CD11c+ / HLA-DR+ / CD86+	all mDCs

[0044] To assess whether one of the immunovariabiles may serve as a proper biomarker, a so-called Cox regression was calculated for each immunovariable. A Cox regression allows estimating the effect of parameter(s) without any consideration of the hazard function. The resulting hazard ratio from such a calculation should be below 1 and related to a significant p value being below 0.05. Otherwise the observed effect would not be related to the applied TLR-9 agonist. Those criteria applied only to activated NKT cells having a hazard ratio of about 0.933 and a p value of 0.0309.

[0045] Surprisingly, the percentage of activated NKT cells could be used to predict treatment success within the verum group. The relationship between the percentage of activated NKT cells and the PFS group status was studied using advanced, well-established statistical analyses within the verum group.

[0046] A receiver operating characteristic (ROC) curve, or simply ROC curve, shows the performance of a binary classifier system as its discrimination threshold is varied. ROC curves are created by plotting the fraction of true positives out of the positives versus the fraction of false positives out of the negatives at various threshold settings. The true positives are also designated as sensitivity and the false positives is one minus the specificity or true negative rate.

[0047] ROC analysis is used in medicine, radiology, biometrics, and other areas for many decades and is increasingly used in machine learning and data mining research. In biomarkers, it can be used to study whether a potential biomarker can have clinical validity, i.e., whether it can be used for predictive purposes. A successful diagnostic test or biomarker will result in a curve that bends above the diagonal while an unsuccessful test will mirror the diagonal, or fall below it. Thus, a ROC curve provides information whether a diagnostic test is successful or not.

[0048] ROC curves in verum patients for activated NKT cells were established (figure 1). The area under the curve was determined to be 0.71, which is a clear indication of the reliability of the biomarker. The Youden-index, which can be used to determine an optimum cut-off value for the test's readout, shows an optimum at cut-off at 3.08% activated NKT cells. Figure 2 shows activated NKT cells versus sensitivity and specificity.

[0049] All patients were now sorted into groups depending on their level of activated NKT cells. Patients with cell levels above the cut-off of 3.08% activated NKT cells were designated as biomarker positive, while patients with levels below the cut-off were labelled biomarker negative.

[0050] Figure 3a shows a Kaplan-Meier plot in biomarker positive patients for verum and placebo (solid line: patients treated with MGN1703; dotted line: patients treated with placebo). It is obvious that the survival probability within the biomarker positive group is surprisingly related to the application of the TLR-9 agonist.

[0051] Figure 3b shows a Kaplan-Meier curve for the biomarker negative patients in both the verum and placebo arm. It is obvious that the time of progression free survival is clearly shorter within this verum group as compared to the biomarker positive verum patients (comp. fig. 3a).

[0052] Figure 4a shows a Kaplan-Meier plot of biomarker positive patients versus biomarker negative patients, only from the verum treatment arm (solid line: biomarker positive patients; dotted line: biomarker negative patients). Clearly, the biomarker positive patients have a significant advantage in survival probability compared to the biomarker negative patients, even when both groups received the verum treatment MGN1703.

[0053] Figure 4b shows a Kaplan-Meier plot of the placebo group for biomarker positive and negative patients (solid line: biomarker positive patients; dotted line: biomarker negative patients). It is obvious that both groups show a more or less identical progression free survival as they were only treated with placebo. Clearly, the biomarker is suitable for assessing whether a patient is a responder to the treatment with a TLR-9 agonist or not. Further, the placebo group shows that the biomarker is not related to effects caused by the overall health status of a patient.

[0054] The present invention provides new predictive biomarker for responder to a cancer

treatment with a TLR-9 agonist, especially a covalently closed partially self-complementary DNA chain having a double stranded stem and single stranded terminal loops bearing unmethylated CG motifs. The determination of the frequencies of activated NKT cells (CD3+/CD56+/Cd69+) at baseline allows assessing the probability whether a patient is a responder to treatment with the DNA construct or not. Importantly, in the placebo arm, the patients with responder-like characteristics, behave the same way as non-responders did, showing that the prolonged progression-free survival time of the biomarker positive verum patients is in fact due to the applicability of the biomarkers for the selected therapy, not just a better overall health or any non-specific effect.

MATERIAL AND METHODS

Sample handling

[0055] Whole blood (10 mL) for FACS was collected in Streck Cyto-Chex® BCT tubes. Within 2 hours after sampling the blood samples were shipped to the analytical laboratory. According to established Protocol, samples were stored at room temperature. The frequency and activation status of plasmacytoid dendritic cells (pDC), myeloid dendritic cells (mDC), monocytes, natural killer (NK) cells, NKT cells, B cells, T cells and other cell populations were evaluated.

Analytical Methods

[0056] Fluorescence activated cell sorting (FACS) were performed according to established principles. Whole blood samples were stained with fluorescencelabeled antibodies and incubated. Phenotypical analysis of the immune cells was performed with a FACScalibur (Becton Dickinson) flow cytometer. Frequencies of the respective analyzed cell populations were documented for each sample of the patients.

[0057] Analysis of human PBMC for the activation of specific cell populations

CD40 expression of plasmacytoid dendritic cells (pDC)

Cells were stained with the following combination of monoclonal antibodies: Anti-Lineage marker-FITC, (antibody cocktail containing antibodies directed against CD3, CD14, CD16, CD19, CD20, CD56); Anti-CD123-PE; Anti-HLA-DR-PerCP; Anti-CD40-APC; PDC were gated as: lineage negative, HLA-DR positive, CD123 positive cells. Within the PDC population CD40 was used as activation marker.

[0058] Activation of NK-, *NK-T* and T cells using the activation marker CD69

Cells were stained with the following combination of monoclonal antibodies: Anti CD3-FITC; Anti CD56-PE; Anti CD69-APC

NK cells were gated as: CD3 negative, CD56 positive cells

NK-T cells were gated as: CD3 positive, CD56 positive cells
T cells were gated as: CD3 positive, CD56 negative
CD69 was used as activation marker for all 3 populations.

[0059] CD86 expression of myeloid dendritic cells (MDC):

Cells were stained with the following combination of monoclonal antibodies: Anti-Lineage marker-FITC, (antibody cocktail containing antibodies directed against CD3, CD14, CD16, CD19, CD20, CD56); Anti-CD11c-PE; Anti-HLA-DR-PerCP ; Anti-CD86-APC

MDC were gated as: lineage negative, HLA-DR positive, CD11c positive cells.

Within the MDC population CD86 was used as activation marker.

[0060] CD86 expression of B cells and monocytes; CD169 expression of monocytes

Cells were stained with the following combination of monoclonal antibodies: Anti CD14-FITC; Anti CD169-PE; Anti CD19-PerCP; Anti CD86-APC

B cells were gated as CD19 positive cells. Within the B cell population CD86 was used as activation marker.

[0061] Monocytes were gated as CD14 positive cells. Within the monocyte population CD86 and CD169 were used as activation markers.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [EP1196178A \[0007\] \[0008\] \[0008\]](#)
- [EP1776124A \[0008\]](#)
- [WO2010055340A \[0012\]](#)

Non-patent literature cited in the description

- **KRIEG AMYI AKMATSON SWALDSCHMIDT TJBISHOP GATEASDALE RKORETZKY GAKLINMAN DM**CpG motifs in bacterial DNA trigger direct B-cell activation *Nature*, 1995, vol. 374, 6522546-9 [0007]
- **RIERA-KNORRENSCHILD et al.** ASCO Meeting Abstracts, 2013, vol. 31, 153643- [0010]
- **VARGHESE et al.** *The Journal of Immunology*, 2012, vol. 188, 1001165.13- [0011]
- **MONTOYA C. et al.** *Clinical Immunology Academic press* 20060000 vol. 120, 138-146 [0013]

PATENTKRAV

1. Fremgangsmåde til forudsigelse eller monitorering af, om en patient, der lider af cancer, vil reagere eller reagerer på behandling med en TLR-9-agonist ved bestemmelse af frekvensen af CD3+ CD56+ CD69+ aktiverede naturlige dræber-T- (NKT) celler, hvor TLR-9-agonisten er et DNA-konstrukt omfattende mindst ét sekvensmønster $N^1N^2CGN^3N^4$, hvor N^1N^2 og N^3N^4 er en hvilken som helst kombination af A, C, T og G, og C er deoxycytidin, G er deoxyguanosin, A er deoxyadenosin og T er deoxythymidin, hvor fremgangsmåden ikke udføres in vivo.
2. Fremgangsmåde ifølge krav 1, hvor forholdet mellem CD3+ CD56+ CD69+ aktiverede NKT-celler fra hele NKT-cellepopulationen bestemmes.
3. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 2, hvor en responder på behandlingen med TLR-9-agonisten har en frekvens af mindst 3 % CD3+ CD56+ CD69+ aktiverede NKT-celler fra hele NKT-cellepopulationen.
4. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 3, hvor der forinden er foretaget en induktionsterapi med et ikke-DNA-lægemiddel.
5. TLR-9-agonist til anvendelse i en fremgangsmåde til behandling af cancer omfattende mindst et DNA-konstrukt, der omfatter mindst ét sekvensmønster $N^1N^2CGN^3N^4$, hvor N^1N^2 og N^3N^4 er en hvilken som helst kombination af A, C, T og G, og C er deoxycytidin, G er deoxyguanosin, A er deoxyadenosin og T er deoxythymidin, og **kendetegnet ved, at** patienten har forhøjede niveauer af aktiveret CD3+ CD56+ CD69+ NKT-celler.
6. Agonist til anvendelse i fremgangsmåden til behandling af cancer ifølge krav 5, hvor N^1N^2 er et element udvalgt fra gruppen omfattende GT, GG, GA, AT eller AA, N^3N^4 er et element udvalgt fra gruppen omfattende CT eller TT.
7. Agonist til anvendelse i fremgangsmåden til behandling af cancer ifølge krav 5 eller 6, hvor DNA'et er et lineært åbenkædet DNA-konstrukt, der omfatter enkelt- eller dobbeltstrenget DNA, eller er et lineært dobbeltstrenget DNA-konstrukt, der omfatter mindst én ende med en enkeltstrenget sløjfe.

8. Agonist til anvendelse i fremgangsmåden til behandling af cancer ifølge et hvilket som helst af kravene 5 til 7, hvor sekvensmønstret $N^1N^2CGN^3N^4$ er placeret inde i et enkeltstrenget og/eller et dobbeltstrenget område af DNA-sekvensen.

5 9. Agonist til anvendelse i fremgangsmåden til behandling af cancer ifølge et hvilket som helst af kravene 5 til 6, der omfatter mindst ét L-DNA-nukleotid.

10 10. Agonist til anvendelse i fremgangsmåden til behandling af cancer ifølge et hvilket som helst af kravene 5 til 9, hvor mindst ét nukleotid er modificeret med en funktionel gruppe udvalgt fra gruppen omfattende carboxyl-, amin-, amid-, aldimin-, ketal-, acetal-, ester-, ether-, disulfid-, thiol- og aldehydgrupper.

15 11. Agonist til anvendelse i fremgangsmåden til behandling af cancer ifølge et hvilket som helst af kravene 5 til 9, hvor DNA-konstruktet er bundet til en forbindelse udvalgt fra gruppen omfattende peptider, proteiner, carbohydrater, antistoffer, lipider, miceller, vesikler, syntetiske molekyler, polymerer, mikroprojektiler, metalpartikler, nanopartikler eller en fast fase.

20 12. Agonist til anvendelse i fremgangsmåden til behandling af cancer ifølge et hvilket som helst af kravene 5 til 10, hvor DNA-konstruktet er en del af en farmaceutisk sammensætning.

13. Agonist til anvendelse i fremgangsmåden til behandling af cancer ifølge krav 12, hvor den farmaceutiske sammensætning er en vaccine.

25 14. Anvendelse af et kit til forudsigelse eller monitorering af, om en patient, der lider af cancer, vil reagere eller reagerer på behandling med en TLR-9-agonist omfattende midler til detektering og kvantificering af frekvensen af $CD3^+ CD56^+ CD69^+$ aktiverede NKT-celler fra hele NKT-cellepopulationen, og hvor TLR-9-agonisten omfatter et DNA-konstrukt, der omfatter mindst ét sekvensmønster $N^1N^2CGN^3N^4$, hvor N^1N^2 og N^3N^4 er en hvilken som helst
30 kombination af A, C, T og G, og C er deoxycytidin, G er deoxyguanosin, A er deoxyadenosin og T er deoxythymidin, hvor kittet ikke anvendes in vivo.

DRAWINGS

Figure 1




Figure 2




Figure 3a




Figure 3b




Figure 4a




Figure 4b

