

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
20 October 2005 (20.10.2005)

PCT

(10) International Publication Number
WO 2005/098049 A2

(51) International Patent Classification⁷: **C12Q 1/68**

(21) International Application Number:
PCT/US2005/009471

(22) International Filing Date: 22 March 2005 (22.03.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/556,147 25 March 2004 (25.03.2004) US

(71) Applicant (for all designated States except US): **CALIFORNIA INSTITUTE OF TECHNOLOGY** [US/US];
Office of Technology Transfer, 1200 E. California Blvd.,
MC 201-85, Pasadena, CA 91125 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DIRKS, Robert**
[US/US]; 183 N. Chester Ave., Apt. 8, Pasadena, CA
91106 (US). **PIERCE, Niles, A.** [US/US]; 800 N. Michi-
gan Avenue, Pasadena, CA 91104 (US).

(74) Agent: **HART, Daniel**; Knobbe Martens Olson & Bear
LLP, 2040 Main Street, 14th Floor, Irvine, CA 92614 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,
ZM, ZW.


(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: HYBRIDIZATION CHAIN REACTION



(57) **Abstract:** The present invention relates to the use of nucleic acid probes to identify analytes in a sample. In the preferred embodiments, metastable nucleic acid monomers are provided that associate in the presence of an initiator nucleic acid. Upon exposure to the initiator, the monomers self-assemble in a hybridization chain reaction. The initiator nucleic acid may be, for example, a portion of an analyte to be detected or may be part of an initiation trigger such that it is made available in the presence of a target analyte.

HYBRIDIZATION CHAIN REACTION

Reference to Related Applications

[0001] The present application claims priority to U.S. Provisional Application Number 60/556,147, filed March 25, 2004

Background of the Invention

Field of the Invention

[0002] The present invention relates generally to the use of nucleic acid probes to identify analytes in a sample and more particularly to the use of metastable nucleic acid monomers that self-assemble upon exposure to a target analyte.

Description of the Related Art

[0003] Single stranded DNA is a versatile material that can be programmed to self-assemble into complex structures driven by the free energy of base pair formation. Synthetic DNA machines can be powered by strand displacement interactions initiated by the sequential introduction of auxiliary DNA fuel strands. Typically, various DNA strands begin to associate as soon as they are mixed together. Catalytic fuel delivery proves a conceptual approach to powering autonomous DNA machines by storing potential energy in loops that are difficult to access kinetically except in the presence of a catalyst strand.

Summary of the Invention

[0004] Metastable nucleic acid monomers can be made that self-assemble upon exposure to an initiator, such as a target analyte. In one aspect of the invention, methods are provided for using such monomers to detect an analyte in a sample. The sample is contacted with a first metastable monomer comprising an initiator complement region and a second metastable monomer comprising a region that is complementary to a portion of the first monomer. The monomers may be, for example, hairpin nucleic acid structures comprising a loop region and a duplex region.

[0005] The first and second monomers polymerize in the presence of an initiator. Preferably, hybridization of the initiator to the initiator complement region of

the first monomer initiates polymerization. Polymerization continues until the supply of one of the monomers is exhausted. The identification of polymers comprising the first and second monomers is indicative of the presence of the analyte in the sample. Polymers may be identified, for example, by gel electrophoresis

[0006] The initiator is preferably a nucleic acid. In some embodiments, the analyte comprises the initiator. In other embodiments the sample is additionally contacted with an initiation trigger. The initiation trigger preferably comprises the initiator and a binding molecule, such as an aptamer, that is able to specifically recognize the analyte of interest. The initiator is able to hybridize to the first monomer, triggering polymerization, when the binding molecule is bound by the analyte.

[0007] In one embodiment the analyte is a nucleic acid that is associated with a pathogen, such as HIV. The sample may be a biological sample from a patient.

[0008] According to another aspect of the invention, methods are provided for forming a structure comprising hybridized nucleic acid monomers. An initiator comprising a nucleic acid initiation region is provided. The initiator may be a single stranded nucleic acid. In some embodiments, the initiator comprises a recognition molecule, such as an aptamer. In one embodiment the initiator is an analyte to be detected in a sample, such as a biological sample obtained from a patient.

[0009] A first metastable nucleic acid hairpin monomer is provided comprising a duplex region, a first loop region and an initiator complement region that is substantially complementary to the initiation region. The loop region of the first monomer preferably comprises from about 3 to about 30 nucleotides.

[0010] A second metastable nucleic acid hairpin monomer is provided that comprises a second duplex region and a propagation region that is substantially complementary to the initiator complement region of the first monomer. The propagation region may be, for example, a second loop region.

[0011] The first and second monomers are able to hybridize in a chain reaction to form a polymer structure upon hybridization of the initiator to the first monomer. The polymer structure may be detected, for example, by a method selected from the group consisting of gel electrophoresis, capillary electrophoresis, mass spectrometry, light scattering spectroscopy, colorimetry and fluorescent spectroscopy.

[0012] In some embodiment, a third and fourth metastable nucleic acid monomer are provided. The third monomer preferably comprises a region that is

complementary to a portion of the first and/or second monomer and the fourth monomer preferably comprises a region complementary to a portion of the third monomer.

[0013] In a further aspect of the invention, a kit is provided for detecting an analyte in a sample. The kit preferably comprises a first metastable nucleic acid monomer comprising an initiator complement region and a second metastable nucleic acid monomer comprising a propagation region. In one embodiment the monomers are hairpin nucleic acid monomers. The kit may also comprise an initiation trigger. The initiation trigger preferably comprises a region that is substantially complementary to the initiator complement region of the first monomer. The initiation trigger may also comprise a target recognition molecule, such as an aptamer that is able to specifically bind the analyte to be detected.

Brief Description of the Drawings

[0014] Figure 1 schematically illustrates one embodiment of an HCR system. Each letter represents a segment of nucleic acids. Letters marked with a * are complementary to the corresponding unmarked letter.

[0015] Figure 1A shows two hairpins, labeled H1 and H2, that are metastable in the absence of initiator I. The hairpins comprise sticky ends 'a' and 'c*', respectively. Potential energy is stored in the hairpin loops.

[0016] Figure 1B shows how a single initiator strand 'I' can nucleate or bind to the sticky end of H1 and displace one arm to open the hairpin. This frees up the bases that were trapped in the hairpin, allowing them to perform a similar displacement reaction on H2.

[0017] As illustrated in Figure 1C, the newly exposed c region of H1 nucleates at the sticky end of H2 and opens the hairpin to expose a region on H2 (a*) that is identical in sequence to the initiator I. As a result, each copy of I can propagate a chain reaction of hybridization events between alternating H1 and H2 hairpins to form a nicked double helix, thereby amplifying the signal of initiator binding. The process can continue until the monomers (H1 and H2) are exhausted. At each step, energy is gained from the hybridization of 'a' or 'c'. The reactions diagrammed in Figure 1 have been successfully carried out and are summarized in Figure 1D.

[0018] Figure 1D illustrates the results of an HCR reaction and the effect of initiator concentration on amplification. Lanes 2-7: Six different concentrations of

initiator were used (0.00, 10.00, 3.20, 1.00, 0.32 and 0.10 μM , respectively) in a 1 μM mixture of H1 and H2. Lanes 1 and 8 were DNA markers with 100-bp and 500-bp increments respectively.

[0019] Figure 1E illustrates HCR kinetics monitored by substituting a fluorescently labeled base in the sticky end of an HCR monomer. Here 2-aminopurine (2AP) was substituted for A (base 3) in the sticky end of H1. The hairpin monomers H1 and H2 did not hybridize prior to triggering by initiator ($(\text{H1}^{2\text{AP}} + 1.2\text{x H2 for 24 hours} + 0.5\text{x initiator})$, red). The same quenched baseline was achieved without HCR by adding excess initiator to $\text{H1}^{2\text{AP}}$ in the absence of H2 ($\text{H1}^{2\text{AP}} + 4.0\text{x initiator}$, green). Addition of insufficient initiator to $\text{H1}^{2\text{AP}}$ provided only partial quenching ($\text{H12AP} + 0.5\text{x initiator}$ (blue), demonstrating that HCR, and not initiator alone, was responsible for exhausting the supply of H12AP monomer.

[0020] Figure 2 illustrates an embodiment in which HCR utilizing two monomer pairs produces quadratic signal amplification. Figure 2A illustrates two hairpin monomers Q1 and Q2 that are metastable in the absence of initiator IQ. As shown in Figure 2B, binding of IQ leads to a strand displacement interaction that exposes sticky end f^*b^* . This single stranded region then nucleates at the f^* sticky end of Q2, and a subsequent branch migration exposes segments cb^*d^* and e^* as shown in Figure 2C. The d^*e^* region initiates the next Q1 molecule, leading to amplification in one direction, while the exposed cb^* region initiates a second HCR reaction involving monomers H1 and H2 (Figure 1). As illustrated in Figure 2D, the resulting branched polymer has a Q1/Q2 main chain with H1/H2 side chains branching off at each Q2 segment.

[0021] Figure 3 illustrates a pair of monomers (E1 and E2) that can be used in combination with at least one other pairs of monomers to achieve exponential amplification by HCR. In the presence of cb^* , E1 and E2 form a linear chain that includes periodic single stranded d^*e^* regions. The initiator sequence for E1/E2 matches the periodic single stranded region produced by Q1/Q2 and vice versa. Consequently, a mixture of Q1, Q2, E1 and E2 plus either initiator (cb^* or d^*e^*) will lead to the formation of a structure in which each branch of the polymer is itself a branched polymer. Sustained growth will ultimately decrease to cubic amplification.

[0022] Figures 4A-E illustrate another embodiment for HCR with exponential growth. Eight different strands are used in this embodiment. Strand one (1) is the 'hub' of the system and has an exposed, single-stranded region joining two hairpins (Figure

4A). When the initiator (2) binds, it creates a long helix with one sticky end on each side. The two sticky ends generated by the initiated 'hub' bind with strands (3) and (6), respectively (Figure 4B and C). Next, auxiliary strands (4) and (7) bind to previously protected bulge loops (Figures 4D and E), and expose two hairpin regions. These hairpins then bind to strands (5) and (8), respectively, to generate sticky ends similar to the initiator molecule (2). Thus, each initiator produces two new initiators, leading to exponential growth. As a side note, two subsets of strands (1,2,3,4,5) and (1,2,6,7,8) produce linear systems in the absence of the other strands.

[0023] Figure 5A illustrates an aptamer HCR trigger mechanism for the detection of ATP. Binding of the DNA aptamer to ATP induces a conformation change that exposes a sticky end.

[0024] Figure 5B shows an agarose gel demonstrating ATP detection via HCR.

[0025] Figure 6 illustrates a self-complementary hairpin with an interior loop that is a double helix (dotted lines). DNA hairpin can also exist as a dimer with an interior loop. One possible concern is that the interior loops may be easier to invade than the corresponding hairpins. To prevent this side reaction, self-complementary hairpins would convert the interior loop to a simple double helix (dotted lines). However, this added precaution may not be necessary.

Detailed Description of the Preferred Embodiment

[0026] Hybridization Chain Reaction (HCR) is a novel method for the triggered hybridization of nucleic acid molecules starting from metastable monomer hairpins or other metastable nucleic acid structures. Dirks, R. and Pierce, N. Proc. Natl. Acad. Sci. USA 101(43): 15275-15278 (2004), incorporated herein by reference in its entirety. HCR does not require any enzymes and can operate isothermally.

[0027] In one embodiment of HCR, two or more metastable monomer hairpins are used. The hairpins preferably comprise loops that are protected by long stems. The loops are thus resistant to invasion by complementary single-stranded nucleic acids. This stability allows for the storage of potential energy in the loops. Potential energy is released when a triggered conformational change allows the single-stranded bases in the loops to hybridize with a complementary strand, preferably in a second hairpin monomer.

[0028] Each monomer is caught in a kinetic trap, preventing the system from rapidly equilibrating. That is, pairs of monomers are unable to hybridize with each other in the absence of an initiator. Introduction of an initiator strand causes the monomers to undergo a chain reaction of hybridization events to form a nicked helix (see Figures 1A-C). HCR can be used, for example, to detect the presence of an analyte of interest in a sample. This and other applications are discussed in more detail below.

Definitions

[0029] "Nucleic Acids" as used herein means oligomers of DNA or RNA. Nucleic acids may also include analogs of DNA or RNA having modifications to either the bases or the backbone. For example, nucleic acid, as used herein, includes the use of peptide nucleic acids (PNA). The term "nucleic acids" also includes chimeric molecules.

[0030] The term "sticky end" refers to a nucleic acid sequence that is available to hybridize with a complementary nucleic acid sequence. The secondary structure of the "sticky end" is such that the sticky end is available to hybridize with a complementary nucleic acid under the appropriate reaction conditions without undergoing a conformational change. Typically the sticky end is a single stranded nucleic acid.

[0031] "Monomers" are individual nucleic acid oligomers. Typically, at least two monomers are used in hybridization chain reactions, although three, four, five, six or more monomers may be used. In some embodiments more than two monomers are utilized, such as in the HCR systems displaying quadratic and exponential growth discussed below. Typically each monomer comprises at least one region that is complementary to at least one other monomer being used for the HCR reaction.

[0032] A first monomer in a monomer pair preferably comprises an initiator complement region that is complementary to a portion of an initiator molecule. The initiator complement region is preferably a sticky end. Binding of the initiator to the initiator complement region begins an HCR reaction.

[0033] In addition, the second monomer preferably comprises a propagation region that is able to hybridize to the initiator complement region of another monomer, preferably another copy of the first monomer, to continue the HCR reaction begun by the initiator. The propagation region may be, for example, the loop region of a hairpin monomer as described below. In one embodiment the propagation region on the second

monomer is identical to the portion of the initiator that is complementary to the initiator complement region of the first monomer.

[0034] The propagation region on the second monomer is preferably only available to interact with the initiator complement region of the first monomer when an HCR reaction has been started by the initiator. That is, the propagation region becomes available to hybridize to the initiator complement region of another monomer when one copy of the first monomer has already hybridized to a second monomer, as discussed in more detail below.

[0035] Preferred monomers are “metastable.” That is, in the absence of an initiator they are kinetically disfavored from associating with other monomers comprising complementary regions. “HCR” monomers are monomers that are able to assemble upon exposure to an initiator nucleic acid to form a polymer.

[0036] As used herein, “polymerization” refers to the association of two or more monomers to form a polymer. The “polymer” may comprise covalent bonds, non-covalent bonds or both. For example, in some embodiments two species of monomers are able to hybridize in an alternating pattern to form a polymer comprising a nicked double helix. The polymers are also referred to herein as “HCR products.”

[0037] An “initiator” is a molecule that is able to initiate the polymerization of monomers. Preferred initiators comprise a nucleic acid region that is complementary to the initiator complement region of an HCR monomer.

Monomers

[0038] Two or more distinct species of nucleic acid monomers are preferably utilized in an HCR reaction. Each monomer species typically comprises at least one region that is complementary to a portion of another monomer species. However, the monomers are designed such that they are kinetically trapped and the system is unable to equilibrate in the absence of an initiator molecule that can disrupt the secondary structure of one of the monomers. Thus, the monomers are unable to polymerize in the absence of the initiator. Introduction of an initiator species triggers a chain reaction of alternating kinetic escapes by the two or more monomer species resulting in formation of a polymer. In the examples below, the two hairpin monomers polymerize in the presence of an initiator to form a nicked, double helix.

[0039] In a preferred embodiment, two or more monomer species are employed that have a hairpin structure. The hairpin monomers preferably comprise loops protected by long stems. In other embodiments, monomers with a different secondary structure are provided. However, the secondary structure is preferably such that the monomers are metastable under the reaction conditions in the absence of an initiator nucleic acid. In the presence of an initiator, the secondary structure of a first monomer changes such that it is able to hybridize to a sticky end of a second monomer species. This in turn leads to a change in the secondary structure of the second monomer, which is then able to hybridize to another first monomer and continue the process. In this way, once a single copy of the first monomer interacts with a single copy of the initiator, a chain reaction is produced such that the monomers are able to assemble into a polymer comprising alternating monomer species.

[0040] A number of criteria can be used to design the monomers to achieve the desired properties. These include, for example and without limitation, sequence symmetry minimization, the probability of adopting the target secondary structure at equilibrium, the average number of incorrect nucleotides at equilibrium relative to the target structure, and hybridization kinetics.

[0041] Monomers can be synthesized using standard methods, including commercially available nucleic acid synthesizers or obtained from commercial sources such as Integrated DNA Technologies (Coralville, IA).

[0042] In some embodiments, monomers are derivitized with a compound or molecule to increase the molecular weight of the polymer resulting from HCR. Preferably they are derivitized at a location that does not interfere with their ability to hybridize. In other embodiments monomers comprise a fluorophore or colorimetric compound that allows the resulting polymers to be visualized.

[0043] In preferred embodiments, at least two hairpin monomers are utilized as illustrated in Figure 1A. The monomers each preferably comprise a sticky end (a and c^* , respectively), a first complementary segment (b and b^* , respectively), a loop segment (c and a^* , respectively), and a second complementary segment (b and b^* , respectively). The first and second complementary segments are also referred to as “stems” and together form a duplex region.

[0044] The first monomer (H1) preferably comprises a sticky end a that is complementary to a first nucleic acid portion a^* of an initiator (I; Figure 1B). This sticky

end is referred to herein as the “initiator complement region.” The initiator may be, for example, an analyte of interest, or a nucleic acid that is able to contact the first monomer only in the presence of an analyte of interest, as discussed in more detail below.

[0045] The second monomer (H2) preferably comprises a sticky end c^* that is complementary to a portion of the first monomer that becomes accessible upon initiator binding. Preferably the sticky end c^* is complementary to the loop segment c of the first monomer (Figure 1A). The loop segment c of the first monomer is preferably not available to hybridize with sticky end c^* of the second monomer in the absence of initiator.

[0046] The first and second complementary segments (b and b^*) in the first and second monomers are typically substantially identical. That is, the first complementary segment b of the first monomer (H1) is able to hybridize to the second complementary segment b^* of the second monomer (H2).

[0047] The first complementary segment of each monomer is also able to hybridize to the second complementary segment of the same monomer to form the hairpin structure. For example, as shown in Figure 1A, the first monomer (H1) comprises a first complementary segment b that is able to hybridize to the second complementary segment b^* . In the absence of an initiator, the first and second complementary segments of each monomer are generally hybridized to form a duplex region of the metastable monomer.

[0048] Preferably, the first complementary segment b of the first monomer is also complementary to a portion b^* of the initiator, such that upon hybridization of the initiator region a^* to the sticky end a (the initiator complement region) of the first monomer H1, one arm of the hairpin structure is displaced. This opens the hairpin and allows binding of the first complementary segment b to the second portion b^* of the initiator strand (Figure 1B).

[0049] The loop segment c of the first monomer is also exposed by the opening of the hairpin and is able to bind to the sticky end c^* of the second monomer H2, as illustrated in Figure 1C. This opens the second monomer hairpin H2 and the second complementary segment b^* of the first monomer is able to hybridize to the first complementary segment b of the second monomer H2.

[0050] This leaves the loop region a^* and first complementary region b^* of the second monomer H2 exposed (Figure 1C). The sticky end a of another first monomer (H1) species is able to bind to the exposed loop region a^* of the second monomer H2,

thus opening the H1 hairpin and continuing the process described above. Because the loop region *a* of the second monomer acts as an initiator on a second H1 monomer and allows the process to continue in the absence of further initiator, it is referred to as the propagation region.

[0051] At each step, energy is gained from the hybridization of the the sticky end of the monomer. The result is a nicked, double helix polymer comprising alternating H1 and H2 fragments. This process preferably continues in a chain reaction until all of one or both of the monomer species is used up, or the reaction is stopped by some other mechanism. If desired, the nicks in the nucleic acid polymer structures that result from HCR can be ligated (for example, using T4 DNA ligase).

[0052] Because of the self-propagating nature of the reaction, each copy of the initiator species can begin the chain reaction. Further, as long as there is a fixed supply of monomers the average molecular weight of the resulting polymers is inversely related to the initiator concentration, as can be seen in Figure 1D.

[0053] The length of the loop, stem and sticky ends of the monomers can be adjusted, for example to ensure kinetic stability in particular reaction conditions and to adjust the rate of polymerization in the presence of initiator. In one preferred embodiment the length of the sticky ends is the same as the length of the loops. In other embodiments the sticky ends are longer or shorter than the loops. However, if the loops are longer than the sticky ends, the loops preferably comprise a region that is complementary to the sticky end of a monomer.

[0054] In some preferred embodiments the length of the loops is short relative to the stems. For example, the stems may be two or three times as long as the loops.

[0055] The loop regions are preferably between about 1 and about 100 nucleotides, more preferably between about 3 and about 30 nucleotides and even more preferably between about 4 and about 7 nucleotides. In one embodiment the loops and sticky ends of a pair of hairpin monomers are about 6 nucleotides in length and the stems are about 18 nucleotides long.

[0056] Other refinements to the system stabilize the monomer hairpins to help prevent HCR in the absence of an initiator. This can be achieved, for example, via super-stable hairpin loop sequences (Nakano et al. Biochemistry 41:14281-14292 (2002)), with ostensible structural features that could further inhibit direct hybridization to the hairpin. In other embodiments hairpin loops are made to be self-complementary at their ends.

This self-complementarity “pinches” the hairpin loops, making them shorter. However, if the reactive sticky ends of each monomer are complementary to the loop regions on the opposite monomer, as described above, they will have a slight propensity to close up, thereby slowing down the reaction. This feature can be utilized if a slower reaction is desired. Completely self-complementary hairpins can also be used, for example if the monomer hairpins are forming dimers with interior loops that are more easily invaded than their hairpin counterparts. Figure 6 illustrates a self-complementary hairpin with an interior loop that is a double helix.

[0057] Reaction conditions are preferably selected such that hybridization is able to occur, both between the initiator and the sticky end of a first monomer, and between the complementary regions of the monomers themselves. The reaction temperature does not need to be changed to facilitate the hybridization chain reaction. That is, the HCR reactions are isothermic. They also do not require the presence of any enzymes.

Variations

[0058] There are many possible variations to HCR that may improve its speed, stability and ability to amplify chemical signals. The system illustrated in Figure 1 and discussed above exhibits linear growth in response to initiator. However, increasing the rate of polymer growth can enhance the ability to detect the presence of low copy number targets, such as a single target molecule in a large test volume. For example, monomers can be designed to undergo triggered self-assembly into branched structures exhibiting quadratic growth or dendritic structures exhibiting exponential growth. The exponential growth is limited by the available space such that it decreases to cubic amplification as the volume around the initiator fills. However, if chain reactions products are able to dissociate, exponential growth can be maintained until the supply of monomers is exhausted.

[0059] In order to achieve non-linear growth, 3 or more HCR monomers can be used. In preferred embodiments at least 4 HCR monomers are used. In some embodiments, at least one monomer in a primary monomer pair incorporate a trigger nucleic acid segment that is complementary to the exposed sticky end of one of the monomers from a secondary set of HCR monomers. Upon exposure to the nucleic acid that is to be detected, the set of primary monomers undergoes HCR to form a polymer

with a periodic single stranded trigger region. Thus the trigger nucleic acid is exposed, leading to a polymerization chain reaction in the secondary set of monomers. In other embodiments, both the primary and secondary set of monomers includes a trigger segment, such that exponential growth is achieved. Exemplary schemes are presented in Figures 2 and 3 for achieving quadratic and exponential growth, respectively.

[0060] In one embodiment, one of a first pair of monomers comprises a bulge loop. Upon polymerization, a single stranded region results from the presence of the bulge loop. The bulge loop segment is preferably complementary to the sticky end of one of a second pair of HCR monomers. Thus, upon exposure to the initiator, the first pair of monomers undergoes HCR to form a polymer with a single stranded region that acts to trigger polymerization of the second pair of monomers. Figures 2A-C depict such a quadratic amplification scheme. Monomers Q1 and Q2 interact with hairpin monomers H1 and H2 (Figure 1) after initiation by I_Q to form the branched polymer schematically illustrated in Figure 2D.

[0061] Q1 and Q2 (Fig. 2a) are metastable in the absence of the initiator I_Q . I_Q binds to Q1 and a subsequent strand displacement interaction exposes segments f, e* and b* as shown in Figure 2B. This single-stranded region contacts sticky end f* of Q2 and a subsequent branch migration exposes segments c, b*, d* and e*. Segment d* then interacts with another copy of Q1 at sticky end d, causing the hairpin to open up such that e* can also hybridize. At the same time, the exposed c segment initiates a linear HCR reaction with hairpins H1 and H2 (not shown). The resulting branched polymer has a main chain comprising alternating Q1 and Q2 segments and H1/H2 side chains branching off at each Q2 segment.

[0062] In a further embodiment, exponential growth is achieved in response to an initiator by combining two or more pairs of monomers. For example, monomer pair Q1 and Q2 (Figure 2) can be used in conjunction with monomers E1 and E2 (Figure 3) to obtain exponential growth in response to an initiator. In the presence of nucleic acid segment cb*, E1 and E2 form a linear chain that includes periodic single stranded d*e* regions. By design, the initiator sequence for E1/E2 matches the periodic single stranded region produced by Q1/Q2 and vice versa. Consequently, a mixture of Q1, Q2, E1 and E2 monomers in the presence of initiator will form a structure in which each branch of the polymer is itself a branched polymer. Either initiator cb*, corresponding to the sticky end

and first complementary region of E1, or d^*e^* , corresponding the sticky end and first complementary region of Q1, will activate the chain reaction.

[0063] While non-linear amplification systems provide enhanced sensitivity over a linear system, they may also have an increased chance for spurious initiation of HCR and a resultant increase in false-positive signals. Several methods may be used to decrease the possibility for initiation of the system in the absence of the initiator. In systems utilizing hairpin monomers, these may include helix clamping, helix elongation and loop entropy ratchets.

[0064] The quadratic and exponential growth HCR schemes illustrated in Figures 2 and 3 include long single-stranded regions (b^* and e^* respectively). These long regions could potentially function as weak initiators. Several methods are available to reduce spurious monomer polymerization in the absence of initiator for both higher order growth schemes and linear growth schemes. These include helix clamping, helix lengthening and loop entropy ratchets. In helix clamping, the single stranded regions in one or more of the monomers are truncated at each end so that the helices that they could potentially invade in other monomers are effectively clamped at the ends by bases that are not present in the single stranded (b^* and e^*) regions. Experiments have shown that this can eliminate any spurious initiation. The amount of truncation that is effective to decrease or eliminate spurious initiation can be determined by routine experimentation. For example, control experiments can be performed using fluorescent gel electrophoresis time courses to monitor strand exchange between single stranded DNA and duplex DNA (e.g., strand b^* invading duplex bb^*) for different clamp lengths. Using spectrally distinct dyes for the initially single stranded DNA and for the two DNA species in the duplex allows independent monitoring of all species as strand exchange proceeds. These controls can provide a systematic basis for selection of clamp dimensions.

[0065] The length of the helices in the linear HCR scheme illustrated in Figure 1 contributes directly to the height of the kinetic barrier that prevents spurious polymerization between the two hairpin species. Interactions between H1 and H2 are sterically impeded by the loop size. However, the long helices (bb^*) in each hairpin provide a more fundamental kinetic barrier; the length of the helices has a direct effect on the height of the kinetic barrier that impedes spurious HCR. An increase in the length of the helices will increase the initial kinetic barrier in the uninitiated system. Thus, in some embodiments utilizing hairpin monomers, for example if spurious initiation is observed,

the length of the duplex region can be increased to reduce the background noise. The helix length necessary to reduce polymerization in the absence of initiator to an acceptable level can be readily determined by routine experimentation. In some embodiments helix lengthening is combined with helix clamping.

[0066] In still other embodiments utilizing hairpin monomers, loop entropy ratchets are used to reduce HCR in the absence of initiator. An initiator opens an HCR hairpin via a three-way branch migration. This reaction is reversible because the displaced strand is tethered in the proximity of the new helix. However, by increasing the length of the single-stranded loop, the entropy penalty associated with closing the loop increases. As a result, a longer loop will bias the reaction to proceed forward rather than returning to the uninitiated state. However, larger loops are more susceptible to strand invasion. To counter this effect and allow the use of larger loops, mismatches can be introduced between the loop sequences and the complementary regions of the other monomers. Again, the loop length and amount of mismatch that produces the desired reduction in non-specific HCR can be determined by the skilled artisan through routine experimentation.

Initiator

[0067] The initiator is preferably a nucleic acid molecule. The initiator comprises an initiator region that is complementary to a portion of an HCR monomer, preferably a portion of the monomer that is available for hybridization with the initiator while the monomer is in its kinetically stable state. The initiator also preferably comprises a sequence that is complementary to a portion of the monomer adjacent to the sticky end such that hybridization of the initiator to the sticky end causes a conformational change in the monomer and begins the HCR chain reaction. For example, the initiator may comprise a region that is complementary to the first complementary region of the HCR monomer, as described above.

[0068] In the preferred embodiments, the sequence of the initiator is complementary the sticky end (initiator complementary region) and first complementary region of a first monomer. As described above, in some embodiments this will also influence the sequence of the second complementary region and the loop of the second monomer species.

[0069] In some embodiments the initiator is a nucleic acid that is to be detected in a sample or a portion of a nucleic acid that is to be detected. In this case, the sequence of the target nucleic acid is taken into consideration in designing the HCR monomers. For example, the initiator complement region, preferably a sticky end, of one monomer is designed to be complementary to a portion of the target nucleic acid sequence. Similarly, a region adjacent to the sticky end of the same monomer can be designed to be complementary to a second region of the target sequence as well. Because the second monomer will hybridize to the first monomer, the sequence of the second monomer will also reflect at least a portion of the sequence of the target nucleic acid.

[0070] In other embodiments, the initiator comprises at least a portion of a nucleic acid that is part of a “initiation trigger” such that the initiator is made available when a predetermined physical event occurs. In the preferred embodiments that predetermined event is the presence of an analyte of interest. However, in other embodiments the predetermined event may be any physical process that exposes the initiator. For example, and without limitation, the initiator may be exposed as a result of a change in temperature, pH, the magnetic field, or conductivity. In each of these embodiments the initiator is preferably associated with a molecule that is responsive to the physical process. Thus, the initiator and the associated molecule together form the initiation trigger. For example, the initiator may be associated with a molecule that undergoes a conformational change in response to the physical process. The conformational change would expose the initiator and thereby stimulate polymerization of the HCR monomers. In other embodiments, however, the initiation trigger comprises a single nucleic acid. The initiator region of the nucleic acid is made available in response to a physical change. For example, the conformation of the initiation trigger may change in response to pH to expose the initiator region.

[0071] The structure of the trigger is preferably such that when the analyte of interest is not present (or the other physical event has not occurred), the initiator is not available to hybridize with the sticky end of a monomer. Analyte frees the initiator such that it can interact with a metastable monomer, triggering the HCR polymerization reactions described above. In some embodiments analyte causes a conformational change in the trigger that allows the initiator to interact with the monomer.

[0072] The initiator may be part of a trigger comprising a nucleic acid that is linked to or associated with a recognition molecule, such as an aptamer, that is capable of

interacting with an analyte of interest. The trigger is designed such that when the analyte of interest interacts with the recognition molecule, the initiator is able to stimulate HCR. Preferably, the recognition molecule is one that is capable of binding the analyte of interest.

[0073] Recognition molecules include, without limitation, polypeptides, such as antibodies and antibody fragments, nucleic acids, such as aptamers, and small molecules. The use of an initiator bound to an aptamer is described in more detail below.

[0074] In some particular embodiments, amplification of diverse recognition events is achieved by coupling HCR to nucleic acid aptamer triggers. An aptamer is identified that is able to specifically bind an analyte of interest. The analyte is not limited to a nucleic acid but may be, for example, a polypeptide or small molecule. The aptamer is linked to a nucleic acid comprising an initiator region in such a way that the initiator is unavailable to stimulate HCR in the absence of analyte binding to the aptamer.

[0075] Preferably, conformational changes in the aptamer secondary structure expose the initiator segment. In one embodiment, such an aptamer trigger is a hairpin nucleic acid that comprises an initiator segment that is complementary to the initiator complement region or sticky end of an HCR monomer. The aptamer trigger also comprises a complementary region that is complementary to a region of the HCR monomer adjacent to the sticky end, a loop region and an aptamer sequence. The hairpin aptamer trigger may also comprise a region that enhances the stability of the hairpin in the absence of aptamer binding to the analyte, such as a nucleic acid region in one arm of the hairpin that is complementary to a region of the other arm.

[0076] Figure 5A depicts a scheme for HCR amplification of ATP binding using an aptamer construct that exposes an initiator strand upon ATP binding. The sticky end can act as a trigger for the HCR mechanism of Figure 1 by opening hairpin H2. The region x is introduced to help stabilize the trigger in the absence of analyte. The region b* includes both the hairpin loop and the portion of the stem complementary to x. This trigger mechanism is based on conformational changes in the aptamer secondary structure (Yingfu Li (2003) Journal of the American Chemical Society 125:4771-4778) that make the initiator strand available to stimulate HCR. Figure 5B illustrates successful detection of ATP, as well as specificity in differentiating ATP from GTP, as discussed in more detail in the Examples below.

Detecting HCR

[0077] The products of HCR are readily detectable by methods known to one of skill in the art for the detection of nucleic acids, including, for example, agarose gel electrophoresis, polyacrylamide gel electrophoresis, capillary electrophoresis, and gel-filled capillary electrophoresis. As the polymers comprise nucleic acids, they can be visualized by standard techniques, such as staining with ethidium bromide. Other methods also may be suitable including light scattering spectroscopy, such as dynamic light scattering (DLS), viscosity measurement, colorimetric systems and fluorescence spectroscopy.

[0078] In some embodiments HCR is monitored by fluorescence resonance energy transfer (FRET). Certain monomers are labeled with fluorescent dyes so that conformational changes resulting from HCR can be monitored by detecting changes in fluorescence. In one embodiment, one of a pair of hairpin molecules is labeled with a fluorophore at the junction of the region complementary to the initiator strand and the duplex region and labeled at the opposing side of the duplex region with a quencher molecule. Upon polymerization, the fluorophore and quencher are separated spatially in the aggregated nucleic acid structure, providing relief of the fluorescence quenching. In this case, the presence of a single initiator is amplified by the chain of fluorescent events caused by HCR.

[0079] Because the size of the HCR products is inversely related to the amount of the target analyte in a sample, HCR can be used to determine analyte concentration. The average molecular weight of the HCR products is obtained by standard measurements. In some embodiments the average molecular weight of HCR products obtained from one sample is compared to the average molecular weight of HCR products from one or more other samples with an unknown analyte concentration. In this way, the relative concentration of analyte in each sample can be determined.

[0080] In other embodiments, the concentration of analyte is determined by comparing the average molecular weight from a sample with unknown concentration to the average molecular weight of HCR products from HCR reactions in one or more control samples with a known concentration of the analyte. In this way the concentration of analyte in the unknown samples can be determined to be the same as one of the control samples, greater than one of the control samples, less than one of the control samples, or in the range of concentration between two control samples. Thus, the number of control

reactions can be adjusted based on the particular circumstances to provide more or less sensitive determination of analyte concentration. For example, if a relatively exact analyte concentration is necessary, the number of control samples can be increased. On the other hand, if only a broad idea of analyte concentration is necessary, fewer control samples can be used.

Applications

[0081] HCR may be used to produce and amplifying a signal indicating the presence of a target molecule. The initial signal can be the presence of a target molecule comprising an initiator nucleic acid region. In other embodiments the initial signal can be generated by any event that exposes an initiator nucleic acid strand (for example, an aptamer binding to its target ligand so as to expose an initiator strand, as discussed above). In both cases, hybridization of the initiator strand to the initiator complementary region on an HCR monomer triggers polymerization of the HCR monomers. For some amplification applications in which detection is the primary objective, HCR may serve as an attractive alternative to PCR (polymerase chain reaction).

[0082] HCR can be used to detect a physical change in a sample. For example, as discussed above an initiator trigger may be used to trigger polymerization of two or more HCR monomers in response to a physical change in the sample. This may be, for example, and without limitation, a change in pH, temperature, magnetic field or conductivity in the sample.

[0083] In other embodiments HCR is used to identify a physical characteristic of a sample. An initiator trigger may be used that is activated only when the sample has a particular characteristic, such as a particular pH, temperature or conductivity. In one embodiment an initiator trigger is utilized that exposes the initiator and triggers HCR only when the pH is above a predetermined level. A sample can then be contacted with HCR monomers and the initiator trigger. After incubation, the sample is analyzed for the presence of HCR reaction products. The presence of such products indicates a pH above the predetermined level.

[0084] HCR may be used to construct a biosensor. In one embodiment the sensor would comprise two or more HCR monomers and would detect the presence of initiator in a sample. Whenever initiator is present, HCR will occur, resulting in the creation of a polymer from the individual monomer species.

[0085] HCR can be used to identify the presence of nucleic acids of interest in a sample. For example, nucleic acids associated with a pathogen can be identified in a biological sample obtained from a patient. In some embodiments, a sample is obtained from a patient and tested for the presence of nucleic acids associated with a virus, such as HIV. One or more pairs of HCR monomers are contacted with the sample, where at least one of the monomers comprises an initiator complementary region that is complementary

to a portion of the nucleic acid to be detected. After incubation, the sample is tested for the presence of polymerized HCR monomers, such as by gel electrophoresis.

[0086] Furthermore, the average molecular weight of the resulting polymers will be inversely related to the initiator concentration. As a result, HCR can be used for quantitative sensing. In one embodiment the concentration of analyte in a sample is determined by comparing HCR products in the sample with HCR products obtained from control reactions using known concentrations of analyte.

[0087] For some applications, it may be useful to employ HCR for both amplification and capture of a target. Thus, the HCR mechanism can be designed to capture a target molecule, leaving the target tethered to the resulting polymer. The separation step is facilitated by the increased size of the HCR product with respect to the analyte of interest. Thus, even small, rare nucleic acid targets can be readily isolated from a sample. In one embodiment the HCR reactions are run on a gel under conditions that do not disrupt binding between the target and the HCR monomer. The HCR products are then isolated from the gel and the target is purified.

[0088] More sophisticated biosensors can be created by the incorporation of DNA or RNA aptamers or other mechanisms that expose HCR initiator strands only in the presence of a particular analyte of interest or a particular physical change in the system. Figure 5A shows how ATP can be used to trigger HCR. Figure 5B shows illustrative results demonstrating sensitivity to initiator concentration and specificity in detecting ATP relative to GTP.

[0089] Thus in some embodiments an initiator region is linked to a recognition molecule that is specific for an analyte to be detected in a sample to form an initiator trigger. The initiator is only made available to stimulate HCR upon analyte binding to the recognition molecule. Preferably, the recognition molecule is an aptamer that is specific for an analyte of interest, such as a polypeptide associated with a disease or disorder. The polypeptide may be, for example, a cancer antigen or a polypeptide associated with a pathogen or infectious agent.

[0090] In one embodiment a biological sample, such as blood, urine, cerebrospinal fluid, or tissue, is obtained from a patient suspected of suffering from a disease or disorder. The sample is incubated with one or more pairs of HCR monomers and an initiator trigger molecule comprising an initiator and an aptamer specific for a polypeptide associated with the disease or disorder. Following incubation the sample is

analyzed for the presence of polymers comprising the HCR monomers. The presence of the polymers indicates that the analyte associated with the disease or disorder is present in the sample.

Examples

[0091] HCR monomers comprising DNA sequences were designed using a combination of criteria (Dirks et al. Nucleic Acids Research 32:1392-1403 (2004)). These included sequence symmetry minimization (Seeman N. C. J. Theor. Biol. 99:237-247 (1982)), the probability of adopting the target secondary structure at equilibrium (Hofacker et al. Monatsh. Chem. 125:167-188 (1994)), the average number of incorrect nucleotides at equilibrium relative to the target structure (Dirks et al. Nucleic Acids Research 32:1392-1403 (2004)) and hybridization kinetics (Flamm et al. RNA 6:325-338 (2000)). The sequences of the monomers and initiator for the basic HCR system illustrated in Figure 1 and the aptamer trigger HCR system illustrated in Figure 5 are shown in Table 1. The aptamer system included new sequences to ensure compatibility with the fixed sequence of the aptamer. DNA was synthesized and purified by Integrated DNA Technologies (Coralville, IA).

Table 1

System	Strand	Sequence*
	d	
Basic	H1	5'- <i>TTA ACC</i> CAC GCC GAA TCC TAG ACT CAA AGT AGT CTA GGA TTC GGC GTG - 3' (SEQ ID NO: 1)
	H2	5'- AGT CTA GGA TTC GGC GTG GGT TAA CAC GCC GAA TCC TAG ACT <i>ACT TTG</i> - 3' (SEQ ID NO:2)
	I	5'- AGT CTA GGA TTC GGC GTG GGT TAA - 3' (SEQ ID NO: 3)
Aptamer †	H1	5'- <i>CAT CTC</i> GGT TTG GCT TTC TTG TTA CCC AGG TAA CAA GAA AGC CAA ACC - 3' (SEQ ID NO: 4)
	H2	5'- TAA CAA GAA AGC CAA ACC GAG ATG GGT TTG GCT TTC TTG TTA <i>CCT GGG</i> - 3' (SEQ ID NO: 5)
	I ^{ATP}	5'- CCC AGG TAA CAA GAA AGC CAA ACC TCT TGT <i>TAC CTG GGG GAG TAT TGC GGA GGA AGG T</i> - 3' (SEQ ID NO: 6)
	I	5' - CCC AGG TAA CAA GAA AGC CAA ACC - 3' (SEQ ID NO: 7)

*In the hairpin sequences, loops are in bold and sticky ends are italicized.

†Aptamer nucleotides are italicized and bolded.

[0092] For the basic HCR system illustrated in Figure 1, concentrated DNA stock solutions were prepared in buffer that was later diluted to reaction conditions. The buffer comprised 50 mM Na₂HPO₄/0.5M NaCl (pH 6.8).

[0093] Monomers H1 and H2 (Figure 1B) were mixed at various concentrations in the presence and absence of initiator. Stock solutions of I, H1 and H2 were diluted in reaction buffer to three times their final concentration and 9 µl of each species was combined, in that order to give a 27 µl reaction volume. Six different concentrations of initiator were used (0.00, 10.00, 3.20, 1.00, 0.32 and 0.10 µM) in a 1 µM mixture of H1 and H2. Reactions were incubated at room temperature for 24 hours before running 24 µl of each product on a gel. Samples were loaded on 1% agarose gels containing 0.5 µg EtBr per ml of gel volume. The gels were prepared using 1X SB buffer (10 mM NaOH, pH adjusted to 8.5 with boric acid). The agarose gels were run at 150 V for 60 minutes and visualized under UV light.

[0094] Figure 1D illustrates the results of the HCR reactions and the effect of initiator concentration on amplification. Lanes 2-7 of the gel shown in Figure 1D are the results of the HCR reactions at the various initiator concentrations, respectively. Lanes 1 and 8 are DNA markers with 100-bp and 500-bp increments respectively. As illustrated by Figure 1D, introduction of an initiator strand triggers a chain reaction of hybridization that results in the production of polymers of various sizes. The average molecular weight of the polymers is inversely related to the initiator concentration (Fig. 1(d)). The inverse relationship follows from the fixed supply of monomer hairpins, but the phenomenon was observed after 10 minutes, when the supply of monomers had not been exhausted.

[0095] These results confirmed an earlier experiment in which 1 µM of H1 and H2 were reacted overnight in 0.5M NaCl, 50mM Na₂HPO₄ at pH 6.5 with initiator at concentrations of 0, 1, 0.1, 0.01, 0.001 and 0.0001 µM. With no initiator HCR reactions were not observed. In addition, no visible polymer growth was observed at initiator concentrations of 0.001 and 0.0001µM. At the other initiator concentrations an inverse relationship was observed between the initiator concentration and the average molecular weight of the resulting polymers.

[0096] In another set of experiments, the aptamer trigger illustrated in Figure 5A was utilized. The aptamer trigger (I^{ATP}) comprised an initiator region corresponding to the initiator used in the experiments described above, linked to an aptamer that is able

to specifically interact with ATP (Huizenga et al. Biochemistry 34:656-665 (1995)). In addition, the aptamer trigger comprises a stabilizing region designed to stabilize the trigger in the absence of ATP. The aptamer trigger is designed such that in the presence of ATP the hairpin is opened and the initiator region exposed, thereby triggering polymerization of the H1 and H2 monomers. The sequence of I^{ATP} is provided in Table 1, above.

[0097] Reactions were carried out with various combinations of H1, H2, I, I^{ATP} , ATP and GTP. Concentrated stock solutions of the constituents were diluted to reaction conditions: 5 mM MgCl₂/0.3 M NaCl/20 mM Tris (pH 7.6). Reactions were performed with 1.4 mM ATP and/or GTP. DNA species were combined to yield 1 μ M concentrations in 27 μ l of reaction buffer, with additions made in the following order: buffer and/or initiator I or aptamer trigger I^{ATP} , H1, and then H2 (I and I^{ATP} interact with H2 rather than H1). In this case, 1 μ l of 40 mM ATP, 40mM GTP or water was added to each reaction, as appropriate, for a total reaction volume of 28 μ l.

[0098] Reactions were incubated at room temperature for one hour and run on agarose gels (as described above) or native polyacrylamide gels. Native polyacrylamide gels were 10% precast gels made with 1X TBE buffer (90mM Tris, 89mM boric acid, 2.0mM EDTA, pH 8.0). The polyacrylamide gels were run at 150V for 40 minutes in 1X TBE and stained for 30 minutes in a solution containing 5 μ g EtBr per ml.

[0099] Figure 5B shows a representative agarose gel illustrating that the aptamer trigger I^{ATP} can initiate polymerization of H1 and H2 in the presence of ATP and that ATP can be distinguished from GTP. Hairpins H1 and H2 did not polymerize when mixed in the absence of initiator (H1 + H2; Lane 1), but did polymerize when the initiator I was added (H1 + H2 + I; Lane 2). ATP alone was unable to trigger the polymerization of the hairpin monomers (H1 + H2 + ATP; Lane 3) and no polymers were observed from the combination of aptamer trigger with ATP in the absence of hairpin monomers (I^{ATP} + ATP; Lane 4). Some weak spurious HCR was observed in the absence of ATP from the combination of monomers and aptamer trigger (H1 + H2 + I^{ATP} ; Lanes 5) or in the presence of GTP (H1 + H2 + I^{ATP} + GTP; Lane 7), respectively. Strong HCR amplification of ATP recognition was seen when the monomers were combined with the aptamer trigger in the presence of ATP (H1 + H2 + I^{ATP} + ATP; Lane 6). A DNA ladder is shown in Lane 8 (100-1000 bp in 100 bp increments).

[0100] The kinetics of HCR reactions were explored using fluorescence quenching. The adenine analog 2-aminopurine (2AP) fluoresces when single stranded but is significantly quenched when in a stacked double-helical conformation (Rachofsky et al. *Biochemistry* 40: 996-956 (2001)). Monomer usage was monitored as polymerization occurred by replacing H1 with the labeled hairpin H1^{2AP}. H1^{2AP} was prepared by substituting 2AP for the third base (A) in the sticky end of H1 (see Table 1). Monitoring 2AP fluorescence was used rather than standard end-labeled strands because the local environment of quenched 2AP was the same regardless of whether initiator (I) or monomer (H2) performs the quenching.

[0101] Fluorescence data were obtained using a fluorometer from Photon Technology International (Lawrenceville, NJ), with the temperature controller set to 22°C. Excitation and emission wavelengths were 303 and 365 nm, respectively, with 4 nm bandwidths. Stock solutions of 0.40 μ M H12AP and 0.48 μ M H2 were prepared in reaction buffer as described above, heated to 90°C for 90 seconds and allowed to cool to room temperature for 1 hour before use. For each experiment, 250 μ l of H12AP was added to either 250 μ l of H2 or 250 μ l of reaction buffer. These 0.20 μ M H12AP solutions were allowed to sit at room temperature for at least 24 hours before fluorescence measurements were taken. The initial signal was obtained after rapidly pipetting the sample in the cuvette to obtain a stable fluorescence baseline. After acquiring at least 2,000 seconds of the baseline, runs were paused for about 1 minute to add 20 μ l of initiator (either 20 μ M or 2.5 μ M) and allow mixing by rapid pipetting. The final reaction volume was 520 μ l for all experiments. The variation in initial fluorescence intensities was about 10% across three experiments.

[0102] As evidenced by the results presented in Figure 1E the hairpin monomers H1 and H2 do not hybridize in the absence of initiator. Addition of initiator (I) to the hairpin mixture led to fluorescence quenching via HCR (bottom band from 0 to 2000 seconds, then dropping to middle band from 2000 seconds on)

[0103] The same quenched baseline was achieved without HCR by combining H1^{2AP} with excess initiator in the absence of H2 (Fig. 1(E), middle band from 0 to 2000 seconds, then dropping to bottom band from 2000 seconds on). In this case, each initiator molecule caused one fluorescent signaling event by binding to H1^{2AP}. With H2 present, HCR performed fluorescent amplification, allowing each initiator molecule to alter the fluorescence of multiple hairpins.

[0104] Addition of insufficient initiator to $\text{H1}^{2\text{AP}}$ provided only partial quenching (Fig. 1(E), top band), demonstrating that HCR, and not initiator alone, was responsible for exhausting the supply of $\text{H1}^{2\text{AP}}$ monomer.

[0105] Although the foregoing invention has been described in terms of certain preferred embodiments, other embodiments will be apparent to those of ordinary skill in the art. Additionally, other combinations, omissions, substitutions and modification will be apparent to the skilled artisan, in view of the disclosure herein. Accordingly, the present invention is not intended to be limited by the recitation of the preferred embodiments, but is instead to be defined by reference to the appended claims. All references cited herein are incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. A method for detecting an analyte in a sample by hybridization chain reaction, the method comprising:
 - contacting the sample with a first metastable monomer comprising an initiator complement region;
 - contacting the sample with a second metastable monomer comprising a region complementary to a portion of the first monomer; and
 - identifying polymers comprising the first and second monomers, wherein the first and second monomers polymerize in the presence of an initiator.
2. The method of Claim 1, wherein the initiator is a nucleic acid.
3. The method of Claim 1, wherein the analyte comprises the initiator.
4. The method of Claim 1, wherein polymerization of the first and second monomers is initiated by hybridization of the initiator to the initiator complement region of the first monomer.
5. The method of Claim 1, wherein the initiator is able to hybridize with the initiator complement region when analyte is present in the sample.
6. The method of Claim 1, wherein the sample is additionally contacted with an initiation trigger.
7. The method of Claim 6, wherein the initiation trigger comprises an aptamer.
8. The method of Claim 7, wherein the aptamer is able to specifically bind the analyte.
9. The method of Claim 7, wherein the initiation trigger comprises the initiator.
10. The method of Claim 9, wherein the initiator is able to hybridize to the first monomer when the aptamer is bound by the analyte.
11. The method of Claim 1, wherein identifying polymers comprises gel electrophoresis.
12. The method of Claim 1, wherein at least one of the first and second monomers is fluorescently labeled.
13. The method of Claim 1, wherein the analyte is a nucleic acid associated with a pathogen.

14. The method of Claim 1, wherein the sample is obtained from a patient.
15. The method of Claim 1, additionally comprising contacting the sample with a third and fourth metastable monomers.
16. The method of Claim 1, wherein the first and second metastable monomers are hairpin monomers.
17. A method of forming a structure comprising hybridized nucleic acid monomers, the method comprising:
 - a) providing an initiator comprising a nucleic acid initiation region;
 - b) providing a first nucleic acid hairpin monomer comprising a duplex region, a first loop region, and an initiator complement region that is substantially complementary to the initiation region;
 - (c) providing a second nucleic acid hairpin monomer comprising a second duplex region, and a propagation region that is substantially complementary to the initiator complement region of the first monomer;wherein the first and second monomers hybridize in a chain reaction upon binding of the initiator to the first monomer.
18. The method of claim 17, wherein the first loop region comprises from about 3 to about 30 nucleotides.
19. The method of Claim 17, wherein the propagation region of the second monomer is a second loop region.
20. The method of claim 17, wherein the initiator is an analyte to be detected in a sample.
21. The method of claim 17, wherein the initiator is a single stranded nucleic acid.
22. The method of Claim 17, wherein the initiator additionally comprises a recognition molecule.
23. The method of Claim 22, wherein the recognition molecule is an aptamer.
24. The method of claim 17, wherein the initiator undergoes a conformational change in the presence of an analyte.
25. The method of claim 25, wherein the analyte comprises a polypeptide
26. The method of Claim 17, wherein the structure comprising hybridized monomers is detected by a method selected from the group consisting of gel

electrophoresis, capillary electrophoresis, mass spectrometry, light scattering spectroscopy, colorimetry and fluorescent spectroscopy.

27. The method of claim 17, further comprising:

providing a third nucleic acid monomer comprising a first complementary region that is complementary to a portion of the first and/or second monomer; and
providing a fourth nucleic acid monomer comprising a second complementary region that is complementary to a portion of the third monomer.

28. The method of Claim 27, wherein the first monomer comprises a bulge region that is complementary to a portion of the third monomer.

29. An kit for the detection of an analyte in a sample, the kit comprising:

a) a first metastable nucleic acid monomer comprising an initiator complement region; and

b) a second metastable nucleic acid monomer comprising a propagation region that is substantially complementary to a portion of the first nucleic acid.

30. The kit of Claim 29, wherein the first and second monomers are hairpin monomers.

31. The kit of Claim 29, additionally comprising a third and fourth metastable nucleic acid monomer.

32. The kit of Claim 29, wherein the initiation complement region is substantially complementary to a portion of the analyte to be detected.

33. The kit of Claim 29, additionally comprising an initiation trigger.

34. The kit of Claim 33, wherein a portion of the initiation trigger is substantially complementary to the initiator complement region of the first monomer.

35. The kit of Claim 31, wherein the initiation trigger comprises an aptamer.

36. The kit of Claim 35, wherein the aptamer is specific for the analyte to be identified.

FIG. 1A




FIG. 1B





FIG. 1C



2/8




FIG. 1E






FIG. 1D

FIG. 2A




Q1 +

FIG. 2B




$I_Q + Q1$

FIG. 2C



$I_Q \cdot Q1 + Q2$

3/8



$I_Q \cdot Q1 \cdot Q2$

FIG. 2D




FIG. 3




FIG. 4A





FIG. 4B



5/8

FIG. 4C




FIG. 4D




FIG. 4E




FIG. 5A




FIG. 5B





FIG. 6



SEQUENCE LISTING

<110> Dirks, Robert M.
Pierce, Niles A.

<120> HYBRIDIZATION CHAIN REACTION

<130> CALTE.020VPC

<160> 7

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> DNA Oligomer

<400> 1

ttaacccacg ccgaatccta gactcaaagt agtctaggat tcggcgtg

48

<210> 2

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> DNA Oligomer

<400> 2

agtctaggat tcggcgtggg ttaacacgcc gaatcctaga ctactttg

48

<210> 3

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> DNA Oligomer

<400> 3

agtctaggat tcggcgtggg ttaa

24

<210> 4

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> DNA Oligomer

<400> 4

catctcgggtt tggctttctt gttacccagg taacaagaaa gccaaacc

48

<210> 5

<211> 48
<212> DNA
<213> Artificial Sequence

<220>
<223> DNA Oligomer

<400> 5
taacaagaaa gccaaaccga gatgggtttg gctttcttgt tacctggg 48

<210> 6
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> DNA Oligomer

<400> 6
cccaggtaac aagaaagcca aacctcttgt tacctggggg agtattgcgg aggaaggt 58

<210> 7
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> DNA Oligomer

<400> 7
cccaggtaac aagaaagcca aacc 24