



(51) International Patent Classification:

C11D 1/02 (2006.01) *C11D 17/06* (2006.01)
C11D 3/20 (2006.01) *C11D 3/42* (2006.01)
C11D 11/00 (2006.01) *C11D 3/40* (2006.01)

(21) International Application Number:

PCT/US2017/054818

(22) International Filing Date:

03 October 2017 (03.10.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

16192053.3 03 October 2016 (03.10.2016) EP

(71) Applicant: **THE PROCTER & GAMBLE COMPANY**
 [US/US]; One Procter & Gamble Plaza, Cincinnati, Ohio
 45202 (US).


(72) Inventors: **CHIEFFI, Andre**; Procter & Gamble Technical Centres Limited, Whitley Road, Longbenton, Newcastle-upon-Tyne NE12 9TS (GB). **BROOKER, Alan, Thomas**; Procter & Gamble Technical Centres Limited, Whitley Road, Longbenton, Newcastle-upon-Tyne NE12 9TS (GB). **WILKINSON, Craig, Adam**; Procter & Gamble Technical Centres Limited, Whitley Road, Longbenton, Newcastle-upon-Tyne NE12 9TS (GB). **FULLER, Linsey**,

Sarah; Procter & Gamble Technical Centres Limited, Whitley Road, Longbenton, Newcastle-upon-Tyne NE12 9TS (GB). **GOULD, Paul, Anthony**; Procter & Gamble Technical Centres Limited, Whitley Road, Longbenton, Newcastle-upon-Tyne NE12 9TS (GB). **TANTAWY, Hossam, Hassan**; Procter & Gamble Technical Centres Limited, Whitley Road, Longbenton, Newcastle-upon-Tyne NE12 9TS (GB). **HADDRELL, Benjamin, Mark**; Procter & Gamble Technical Centres Limited, Whitley Road, Longbenton, Newcastle-upon-Tyne NE12 9TS (GB).

(74) Agent: **KREBS, Jay A.**; c/o THE PROCTER & GAMBLE COMPANY, Global IP Services, One Procter & Gamble Plaza, C9, Cincinnati, Ohio 45202 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(54) Title: LOW PH LAUNDRY DETERGENT COMPOSITION



(57) Abstract: The present invention relates to a solid free flowing particulate laundry detergent composition comprising: (a) anionic deterative surfactant; (b) from 0wt% to 8wt% zeolite builder; (c) from 0wt% to 4wt% phosphate builder; (d) from 0wt% to 8wt% sodium carbonate; (e) from 0wt% to 8wt% sodium silicate; (f) from 4wt% to 20wt% organic acid; and (g) hueing agent having the following structure: wherein: R1 and R2 are independently selected from the group consisting of: H; alkyl; alkoxy; alkyleneoxy; alkyl capped alkyleneoxy; urea; and amido; R3 is a substituted aryl group; X is a substituted group comprising sulfonamide moiety and optionally an alkyl and/or aryl moiety, and wherein the substituent group comprises at least one alkyleneoxy chain that comprises an average molar distribution of at least four alkyleneoxy moieties, wherein the composition at 1wt% dilution in deionized water at 20°C, has an equilibrium pH in the range of from 6.5 to 9.0, wherein the composition comprises from 30wt% to 90wt% base detergent particle, wherein the base detergent particle comprising (by weight of the base detergent particle): (a) from 4wt% to 35wt% anionic deterative surfactant; (b) optionally, from 1wt% to 8wt% zeolite builder; (c) from 0wt% to 4wt% phosphate builder; (d) from 0wt% to 8wt% sodium carbonate; (e) from 0wt% to 8wt% sodium silicate; (f) from 1wt% to 10wt% organic acid; and (g) optionally, from 1wt% to 10wt% magnesium sulphate.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

LOW PH LAUNDRY DETERGENT COMPOSITION

FIELD OF THE INVENTION

The present invention relates to a solid free flowing particulate laundry detergent
5 composition having a low pH profile. The compositions of the present invention provide good solubility profile, good cleaning profile, good stability profile and good fabric care profile.

BACKGROUND OF THE INVENTION

Laundry detergent powder manufacturers seek to provide solid free-flowing particulate
10 laundry detergent compositions that have good solubility profile, good cleaning profile, good stability profile and good fabric care profile. Typically, a performance balance is required between the chosen formulation to ensure that these profile requirements are met.

The pH profile of a typical laundry detergent powder is quite high, around pH 10.5 and sometimes even higher. This pH profile ensures the good performance of historic cleaning
15 mechanisms: such as grease saponification mechanisms and/or fabric fibre swelling mechanisms. However, this high pH profile also means that the detergent formulators are having to address problems with improving the fabric care profile, and ensuring fabric appearance performance and/or fabric shape retention performance is still adequate.

The inventors have found that an alternative approach to this historic dichotomy of
20 formulating high pH detergent powders to ensure good cleaning performance whilst needing to balance the formulation so as to also provide good fabric care performance, is to formulate the solid detergent powder at a lower pH and then to balance the formulation so as to also provide good cleaning performance.

This low pH laundry detergent powder formulation approach ensures good fabric
25 appearance and good fabric care profiles, but careful attention is needed to ensure good cleaning performance, and especially to address any undesirable cleaning performance skews that result due to the low pH profile.

The inventors have found that the cleaning performance of low pH laundry detergent
30 powders can be improved by careful formulation of specific technologies, specific formulation features and particle architecture as defined by the present invention.

In particular, the inventors have found that a good cleaning performance is achieved by the combination of a low pH solid laundry detergent powder when formulated using a specific base detergent particle, specific formulation features and a specific hueing dye. This combination of

technologies provide good whiteness performance, whilst also minimizing/negating any undesirable build up of hueing dye on the fabric.


WO00/18856 relates to detergent compositions. However, the compositions disclosed by WO00/18856 differ from the composition required by the present invention. In particular, example composition E of WO00/18856 has a calculated pH of 9.7. This is higher (more alkaline) than the pH profile required by the present invention. Data in the application shows the benefit of combining the reduced pH profile with the specific hueing dye and other formulation features required by the present invention (c.f. invention example 7 compared to comparative example 9).

WO03/038028 relates to detergent compositions. However, the compositions disclosed by WO03/038028 differ from the compositions required by the present invention. In particular, example E of WO03/18856 comprises high levels of carbonate in excess of the levels required by the present invention. Data in the application shows the benefit of formulating at lower sodium carbonate levels when formulated in combination with the specific hueing dye and other formulation features required by the present invention (c.f. invention example 7 compared to comparative example 8).

SUMMARY OF THE INVENTION

The present invention relates to a solid free flowing particulate laundry detergent composition comprising:

- (a) anionic deterative surfactant;
- (b) from 0wt% to 8wt% zeolite builder;
- (c) from 0wt% to 4wt% phosphate builder;
- (d) from 0wt% to 8wt% sodium carbonate;
- (e) from 0wt% to 8wt% sodium silicate;
- (f) from 4wt% to 20wt% organic acid; and
- (g) hueing agent having the following structure:



wherein:

R1 and R2 are independently selected from the group consisting of: H; alkyl; alkoxy; alkyleneoxy; alkyl capped alkyleneoxy; urea; and amido;

5 R3 is a substituted aryl group;

X is a substituted group comprising sulfonamide moiety and optionally an alkyl and/or aryl moiety, and wherein the substituent group comprises at least one alkyleneoxy chain that comprises an average molar distribution of at least four alkyleneoxy moieties,

10 wherein the composition at 1wt% dilution in deionized water at 20°C, has an equilibrium pH in the range of from 6.5 to 9.0, preferably from 6.5 to 8.0,

wherein the composition comprises from 30wt% to 90wt% base detergent particle, wherein the base detergent particle comprising (by weight of the base detergent particle):

- (a) from 4wt% to 35wt% anionic deterative surfactant;
- (b) optionally, from 1wt% to 8wt% zeolite builder;
- 15 (c) from 0wt% to 4wt% phosphate builder;
- (d) from 0wt% to 8wt% sodium carbonate;
- (e) from 0wt% to 8wt% sodium silicate;
- (f) from 1wt% to 10wt% organic acid; and
- (g) optionally, from 1wt% to 10wt% magnesium sulphate.


20

DETAILED DESCRIPTION OF THE INVENTION

The solid free flowing particulate laundry detergent composition comprises:

- (a) anionic deterative surfactant;
- (b) from 0wt% to 8wt% zeolite builder;
- 25 (c) from 0wt% to 4wt% phosphate builder;
- (d) from 0wt% to 8wt% sodium carbonate;
- (e) from 0wt% to 8wt% sodium silicate;
- (f) from 4wt% to 20wt% organic acid; and
- (g) hueing agent having the following structure:

30



wherein:

R1 and R2 are independently selected from the group consisting of: H; alkyl; alkoxy;

5 alkyleneoxy; alkyl capped alkyleneoxy; urea; and amido;

R3 is a substituted aryl group;

X is a substituted group comprising sulfonamide moiety and optionally an alkyl and/or aryl moiety, and wherein the substituent group comprises at least one alkyleneoxy chain that comprises an average molar distribution of at least four alkyleneoxy moieties,

10

wherein the composition at 1wt% dilution in deionized water at 20°C, has an equilibrium pH in the range of from 6.5 to 9.0, preferably from 6.5 to 8.0,

wherein the composition comprises from 30wt% to 90wt% base detergent particle, wherein the base detergent particle comprising (by weight of the base detergent particle):

15

(a) from 4wt% to 35wt% anionic deterative surfactant;

(b) optionally, from 1wt% to 8wt% zeolite builder;

(c) from 0wt% to 4wt% phosphate builder;

(d) from 0wt% to 8wt% sodium carbonate;

(e) from 0wt% to 8wt% sodium silicate;

20

(f) from 1wt% to 10wt% organic acid; and

(g) optionally, from 1wt% to 10wt% magnesium sulphate.

Solid free-flowing particulate laundry detergent composition: Typically, the solid free-flowing particulate laundry detergent composition is a fully formulated laundry detergent composition, not a portion thereof such as a spray-dried, extruded or agglomerate particle that only forms part of the laundry detergent composition. Typically, the solid composition comprises a plurality of chemically different particles, such as spray-dried base detergent particles and/or agglomerated base detergent particles and/or extruded base detergent particles, in combination with one or more, typically two or more, or five or more, or even ten or more particles selected

25

from: surfactant particles, including surfactant agglomerates, surfactant extrudates, surfactant needles, surfactant noodles, surfactant flakes; phosphate particles; zeolite particles; polymer particles such as carboxylate polymer particles, cellulosic polymer particles, starch particles, polyester particles, polyamine particles, terephthalate polymer particles, polyethylene glycol particles; aesthetic particles such as coloured noodles, needles, lamellae particles and ring particles; enzyme particles such as protease granulates, amylase granulates, lipase granulates, cellulase granulates, mannanase granulates, pectate lyase granulates, xyloglucanase granulates, bleaching enzyme granulates and co- granulates of any of these enzymes, preferably these enzyme granulates comprise sodium sulphate; bleach particles, such as percarbonate particles, especially coated percarbonate particles, such as percarbonate coated with carbonate salt, sulphate salt, silicate salt, borosilicate salt, or any combination thereof, perborate particles, bleach activator particles such as tetra acetyl ethylene diamine particles and/or alkyl oxybenzene sulphonate particles, bleach catalyst particles such as transition metal catalyst particles, and/or isoquinolinium bleach catalyst particles, pre-formed peracid particles, especially coated pre-formed peracid particles; filler particles such as sulphate salt particles and chloride particles; clay particles such as montmorillonite particles and particles of clay and silicone; flocculant particles such as polyethylene oxide particles; wax particles such as wax agglomerates; silicone particles, brightener particles; dye transfer inhibition particles; dye fixative particles; perfume particles such as perfume microcapsules and starch encapsulated perfume accord particles, or pro-perfume particles such as Schiff base reaction product particles; hueing dye particles; chelant particles such as chelant agglomerates; and any combination thereof.

Typically, the solid free flowing particulate laundry detergent composition comprises:

- (a) anionic deterative surfactant;
- (b) from 0wt% to 8wt% zeolite builder;
- (c) from 0wt% to 4wt% phosphate builder;
- (d) from 0wt% to 8wt% sodium carbonate;
- (e) from 0wt% to 8wt% sodium silicate; and (f) from 4wt% to 20wt% organic acid.

Typically, the composition at 1wt% dilution in deionized water at 20°C, has an equilibrium pH in the range of from 6.5 to 9.0, preferably from 6.5 to 8.5, more preferably from 7.0 to 8.0.

Typically, the composition at 1wt% dilution in deionized water at 20°C, has a reserve alkalinity to pH 7.0 of less than 4.0gNaOH/100g, preferably less than 3.0gNaOH/100g, or even less than 2.0gNaOH/100g.

As used herein, the term “reserve alkalinity” is a measure of the buffering capacity of the detergent composition (g/NaOH/100g detergent composition) determined by titrating a 1% (w/v) solution of detergent composition with hydrochloric acid to pH 7.0 i.e. in order to calculate Reserve Alkalinity as defined herein:

5

$$\text{Reserve Alkalinity (to pH 7.0) as \% alkali in g NaOH/100 g product} = \frac{T \times M \times 40 \times \text{Vol}}{10 \times \text{Wt} \times \text{Aliquot}}$$

10

T = titre (ml) to pH 7.0
 M = Molarity of HCl = 0.2
 40 = Molecular weight of NaOH
 Vol = Total volume (ie. 1000 ml)
 W = Weight of product (10 g)
 Aliquot = (100 ml)

15

Obtain a 10g sample accurately weighed to two decimal places, of fully formulated detergent composition. The sample should be obtained using a Pascall sampler in a dust cabinet. Add the 10g sample to a plastic beaker and add 200 ml of carbon dioxide-free de-ionised water. Agitate using a magnetic stirrer on a stirring plate at 150 rpm until fully dissolved and for at least 15 minutes. Transfer the contents of the beaker to a 1 litre volumetric flask and make up to 1 litre with deionised water. Mix well and take a 100 mls $\pm$ 1 ml aliquot using a 100 mls pipette immediately. Measure and record the pH and temperature of the sample using a pH meter capable of reading to ± 0.01 pH units, with stirring, ensuring temperature is 21°C $\pm$ 2°C. Titrate whilst stirring with 0.2M hydrochloric acid until pH measures exactly 7.0. Note the millilitres of hydrochloric acid used. Take the average titre of three identical repeats. Carry out the calculation described above to calculate the reserve alkalinity to pH 7.0.

Typically, the composition comprises from 30wt% to 90wt% base detergent particle, wherein the base detergent particle comprising (by weight of the base detergent particle): (a) from 4wt% to 35wt% anionic deterative surfactant; (b) optionally, from 1wt% to 8wt% zeolite builder; (c) from 0wt% to 4wt% phosphate builder; (d) from 0wt% to 8wt%, preferably from 0wt% to 4wt%, sodium carbonate; (e) from 0wt% to 8wt%, preferably from 0wt% to 4wt%, sodium silicate; (f) from 1wt% to 10wt% organic acid; and (g) optionally, from 1wt% to 10wt%


magnesium sulphate. Typically, the base detergent particle is in the form of a spray-dried particle.

Typically, the organic acid comprises citric acid and the base detergent particle comprises from 1wt% to 10wt% citric acid.

5 The organic acid may be at least partially coated, or even completely coated, by a water-dispersible material. Water-dispersible material also typically includes water-soluble material. A suitable water-dispersible material is wax. A suitable water-soluble material is citrate.


Typically, the anionic deterative surfactant comprises alkyl benzene sulphonate and wherein the base detergent particle comprises from 4wt% to 35wt% alkyl benzene sulphonate.

10 Typically, the base detergent particle comprises from 0.5wt% to 5wt% carboxylate co-polymer, wherein the carboxylate co-polymer comprises: (i) from 50 to less than 98 wt% structural units derived from one or more monomers comprising carboxyl groups; (ii) from 1 to less than 49 wt% structural units derived from one or more monomers comprising sulfonate moieties; and (iii) from 1 to 49 wt% structural units derived from one or more types of
15 monomers selected from ether bond-containing monomers represented by formulas (I) and (II):
formula (I):



wherein in formula (I), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, X represents a number 0-5 provided X represents a
20 number 1-5 when R is a single bond, and R_1 is a hydrogen atom or C_1 to C_{20} organic group;

formula (II)



wherein in formula (II), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, x represents a number 0-5, and R_1 is a hydrogen atom or C_1 to C_{20} organic group.

Typically, the base detergent particle comprises from 30wt% to 70wt% sodium sulphate.

Typically, the composition comprises from 1wt% to 20wt% co-surfactant particle, wherein the co-surfactant particle comprises: (a) from 25wt% to 60wt% co-surfactant; (b) from 10wt% to 50wt% carbonate salt; and (c) from 1wt% to 30wt% silica. Typically, the co-surfactant particle is in the form of an agglomerate.

Typically, the co-surfactant comprises alkyl ethoxylated sulphate having an average degree of ethoxylation of from 0.5 to 2.5, and wherein the co-surfactant particle comprises from 25wt% to 60wt% alkyl ethoxylated sulphate having an average degree of ethoxylation of from 0.5 to 2.5.

The co-surfactant particle may comprise linear alkyl benzene sulphonate and alkyl ethoxylated sulphate having an average degree of ethoxylation of from 0.5 to 2.5.

The composition at 1wt% dilution in deionized water at 20°C, may have an equilibrium pH in the range of from 6.5 to 8.5.

The composition may have a reserve alkalinity to pH 7.5 of less than 3.0gNaOH/100g.

The composition may comprise from 0wt% to 6wt%, preferably from 0wt% to 4wt%, sodium bicarbonate.

The composition may comprise from 0wt% to 4wt% sodium carbonate.

The composition may comprise from 0wt% to 4wt% sodium silicate.

The composition may comprise from 0wt% to 4wt% phosphate builder.

The composition is preferably substantially free of phosphate builder.

The composition may be substantially free of sodium carbonate.

The composition may be substantially free of sodium bicarbonate.

The composition may be substantially free of sodium silicate.

By “substantially free” it is typically meant herein to mean: “comprises no deliberately added”.

The composition may comprise the combination of lipase enzyme and soil release polymer.

5 Preferably, the composition comprises alkyl benzene sulphonate, wherein the alkyl benzene sulphonate comprises at least 25wt% of the 2-phenyl isomer. A suitable alkyl benzene sulphonate having this feature is obtained by DETAL synthesis.


The composition may comprises alkyl amine oxide.

10 The composition may comprises from 0.5wt% to 8wt% carboxylate co-polymer, wherein the carboxylate co-polymer comprises: (i) from 50 to less than 98 wt% structural units derived from one or more monomers comprising carboxyl groups;

(ii) from 1 to less than 49 wt% structural units derived from one or more monomers comprising sulfonate moieties; and (iii) from 1 to 49 wt% structural units derived from one or more types of monomers selected from ether bond-containing monomers represented by formulas (I) and (II):


15

formula (I):



20 wherein in formula (I), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, X represents a number 0-5 provided X represents a number 1-5 when R is a single bond, and R_1 is a hydrogen atom or C_1 to C_{20} organic group;


formula (II)



- 5 wherein in formula (II), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, X represents a number 0-5, and R_1 is a hydrogen atom or C_1 to C_{20} organic group.

The composition may comprise polyethylene glycol polymer, wherein the polyethylene glycol polymer comprises a polyethylene glycol backbone with grafted polyvinyl acetate side chains.

- 10 The composition may comprise a polyester soil release polymer having the structure:



wherein n is from 1 to 10; m is from 1 to 15 ;




- 15 X is H or SO_3Me ;

wherein Me is H, Na^+ , Li^+ , K^+ , Mg^{2+} , Ca^{2+} , Al^{3+} , ammonium, mono-, di-, tri-, or tetra-alkylammonium; wherein the alkyl groups are C_1 - C_{18} alkyl or C_2 - C_{10} hydroxyalkyl, or any mixture thereof;

R_1 are independently selected from H or C_1 - C_{18} n- or iso-alkyl.

- 20 The composition may comprise a polyester soil release polymer consisting of structure units (1) to (3):

11



wherein:

a, b and c are from 1 to 10;

5 x, y is from 1 to 10;

z is from 0.1 to 10;

Me is H, Na⁺, Li⁺, K⁺, Mg²⁺, Ca²⁺, Al³⁺, ammonium, mono-, di-, tri-, or tetra-alkylammonium wherein the alkyl groups are C₁-C₁₈ alkyl or C₂-C₁₀ hydroxyalkyl, or any mixture thereof;

10 R₁, are independently selected from H or C₁-C₁₈ n- or iso-alkyl;

R₂ is a linear or branched C₁-C₁₈ alkyl, or a linear or branched C₂-C₃₀ alkenyl, or a cycloalkyl group with 5 to 9 carbon atoms, or a C₆-C₃₀ aryl group, or a C₆-C₃₀ arylalkyl group.

The composition may comprise carboxymethyl cellulose having a degree of substitution greater than 0.65 and a degree of blockiness greater than 0.45.

15 The composition may comprise an alkoxyated polyalkyleneimine, wherein said alkoxyated polyalkyleneimine has a polyalkyleneimine core with one or more side chains bonded to at least one nitrogen atom in the polyalkyleneimine core, wherein said alkoxyated polyalkyleneimine has an empirical formula (I) of (PEI)_a-(EO)_b-R₁, wherein a is the average number-average molecular weight (MW_{PEI}) of the polyalkyleneimine core of the alkoxyated polyalkyleneimine and is in the range of from 100 to 100,000 Daltons, wherein b is the average degree of ethoxylation in said one or more side chains of the alkoxyated polyalkyleneimine and is in the range of from 5 to 40, and wherein R₁ is independently selected from the group consisting of hydrogen, C₁-C₄ alkyls, and combinations thereof.

20

The composition may comprise an alkoxyated polyalkyleneimine, wherein said alkoxyated polyalkyleneimine has a polyalkyleneimine core with one or more side chains bonded to at least one nitrogen atom in the polyalkyleneimine core, wherein the alkoxyated polyalkyleneimine has an empirical formula (II) of $(PEI)_o-(EO)_m(PO)_n-R_2$ or $(PEI)_o-(PO)_n(EO)_m-R_2$, wherein o is the average number-average molecular weight (MW_{PEI}) of the polyalkyleneimine core of the alkoxyated polyalkyleneimine and is in the range of from 100 to 100,000 Daltons, wherein m is the average degree of ethoxylation in said one or more side chains of the alkoxyated polyalkyleneimine which ranges from 10 to 50, wherein n is the average degree of propoxylation in said one or more side chains of the alkoxyated polyalkyleneimine which ranges from 1 to 50, and wherein R_2 is independently selected from the group consisting of hydrogen, C_1 - C_4 alkyls, and combinations thereof.

The composition may comprise the combination of a non-ionic soil release polymer and an anionic soil release polymer.

Highly preferably, the composition is substantially free of pre-formed peracid.

The composition may comprise:

- (a) from 1wt% to 20wt% sodium percarbonate;
- (b) from 0.5wt% to 5wt% bleach activator; and
- (c) from 0.5wt% to 5wt% chelant.

The bleach activator may comprise sodium tetraacetythylenediamine, and wherein the composition may comprise from 0.5wt% to 5wt% sodium tetraacetythylenediamine.

The chelant may comprise sodium salt of methylglycine diacetic acid (MGDA), and wherein the composition may comprise from 0.5wt% to 5wt% sodium salt of methylglycine diacetic acid (MGDA).


The chelant may comprise ethylenediamine disuccinic acid (EDDS), and wherein the composition may comprise from 0.5wt% to 5wt% ethylenediamine disuccinic acid (EDDS).

The chelant may comprise disodium 4,5-dihydroxy-1,3-benzenedisulfonate, and wherein the composition may comprise from 0.5wt% to 5wt% disodium 4,5-dihydroxy-1,3-benzenedisulfonate.

The composition may comprises 4,4'-bis-(triazinylamino)-stilbene-2,2'-disulfonic acid brightener and/or 4,4'-distyryl biphenyl brightener.

The composition may comprises an acyl hydrazone bleach catalyst, wherein the acyl hydrazone bleach catalyst has the formula I:

13




wherein, R¹ is selected from the groups comprising CF₃, C₁₋₂₈ alkyl, C₂₋₂₈ alkenyl, C₂₋₂₂ alkynyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkenyl, phenyl, naphthyl, C₇₋₉ aralkyl, C₃₋
 5 20 heteroalkyl, C₃₋₁₂ cycloheteroalkyl or a mixture thereof;

R² and R³ are independently selected from the group comprising hydrogen, substituted C₁₋₂₈ alkyl, C₂₋₂₈ alkenyl, C₂₋₂₂ alkynyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkenyl, C₇₋₉ aralkyl, C₃₋₂₈ heteroalkyl, C₃₋₁₂ cycloheteroalkyl, C₅₋₁₆ heteroaralkyl, phenyl, naphthyl, heteroaryl or a mixture thereof;

10 or R² and R³ are linked to form a substituted 5-, 6-, 7-, 8- or 9-membered ring that optionally comprises heteroatoms;

and R⁴ is selected from the groups comprising hydrogen, C₁₋₂₈ alkyl, C₂₋₂₈ alkenyl, C₂₋₂₂ alkynyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkenyl, C₇₋₉ aralkyl, C₃₋₂₀ heteroalkyl, C₃₋₁₂ cycloheteroalkyl, C₅₋₁₆ heteroaralkyl, substituted phenyl, naphthyl, heteroaryl or a
 15 mixture thereof.

The composition may comprise a hueing agent having the following structure:




20 wherein:

R₁ and R₂ are independently selected from the group consisting of: H; alkyl; alkoxy; alkyleneoxy; alkyl capped alkyleneoxy; urea; and amido;

R₃ is a substituted aryl group;

X is a substituted group comprising sulfonamide moiety and optionally an alkyl and/or aryl
 25 moiety, and wherein the substituent group comprises at least one alkyleneoxy chain that comprises an average molar distribution of at least four alkyleneoxy moieties.

The composition may comprise a hueing agent having the following structure:



5 wherein the index values x and y are independently selected from 1 to 10.

The composition may comprise a hueing agent selected from Acid Violet 50, Direct Violet 9, 66 and 99, Solvent Violet 13 and any combination thereof.

The composition may comprise a protease having at least 90% identity to the amino acid sequence of *Bacillus amyloliquefaciens* as shown in SEQ ID NO:9

10 The composition may comprise a protease having at least 90% identity to the amino acid sequence of *Bacillus amyloliquefaciens* BPN' as shown in SEQ ID NO:10, and which comprises one or more mutations selected from group consisting of V4I, S9R, A15T, S24G, S33T, S53G, V68A, N76D, S78N, S101M/N, Y167F, and Y217Q.

15 The composition may comprise a protease having at least 90% identity to the amino acid sequence of *Bacillus thermoproteolyticus* as shown in SEQ ID NO:11.

The composition may comprise a protease having at least 90% identity to the amino acid sequence of *Bacillus lentus* as shown in SEQ IS NO:12, and which comprises one or mutations selected from the group consisting of S3T, V4I, A194P, V199M, V205I, and L217D.

20 The composition may comprise a protease having at least 90% identity to the amino acid sequence of *Bacillus sp. TY145* as shown in SEQ ID NO:13.

The composition may comprises a protease having at least 90% identity to the amino acid sequence of *Bacillus sp. KSM-KP43* as shown in SEQ ID NO:14.

25 The composition may comprise a variant of the wild-type amylase from *Bacillus sp.* which has at least 90% identity for amino acid sequence SEQ ID NO:5, and which comprises one or more mutations at positions N195, G477, G304, W140, W189, D134, V206, Y243, E260, F262, W284, W347, W439, W469 and/or G476, and optionally which comprises the deletions of D183* and/or G184*.

30 The composition may comprise a variant of the wild-type amylase from *Bacillus sp.* which has at least 90% identity for amino acid sequence SEQ ID NO:6, and which comprises one or more mutations at positions 9, 26, 30, 33, 82, 37, 106, 118, 128, 133, 149, 150, 160, 178, 182,

186, 193, 195, 202, 214, 231, 256, 257, 258, 269, 270, 272, 283, 295, 296, 298, 299, 303, 304, 305, 311, 314, 315, 318, 319, 320, 323, 339, 345, 361, 378, 383, 419, 421, 437, 441, 444, 445, 446, 447, 450, 458, 461, 471, 482 and/or 484, preferably that also contain the deletions of D183* and G184*.

5 The composition may comprise a variant of the wild-type amylase from *Bacillus sp. KSM-K38* which has at least 90% identity for amino acid sequence SEQ ID NO:7.

 The composition may comprise a variant of the wild-type amylase from *Cytophaga sp.* which has at least 60% identity for amino acid sequence SEQ ID NO:8.

10 The composition may comprise a variant of the wild-type lipase from *Thermomyces lanuginosus* which has at least 90% identity for amino acid sequence SEQ ID NO:1.

 The composition may comprise a variant of the wild-type lipase from *Thermomyces lanuginosus* which has at least 90% identity for amino acid sequence SEQ ID NO:1, and which comprises T231R and/or N233R mutations.

15 The composition may comprise a variant of the wild-type lipase from *Thermomyces lanuginosus* which has at least 90% identity for amino acid sequence SEQ ID NO:1, and which comprises G91A, D96G, G225R, T231R and/or N233R mutations.

 the composition may comprise a cellulase that is a wild-type or variant of a microbially-derived endoglucanase endogenous to *Bacillus sp.* exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4) which has at least 90% identity to the amino acid sequence SEQ ID NO:2.


20 The composition may comprise cellulase that is a wild-type or variant of a microbially-derived endoglucanase endogenous to *Paenibacillus polymyxa* exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4) which has at least 90% identity to amino acid sequence SEQ ID NO:3.

25 The composition may comprise a cellulase that is a hybrid fusion endoglucanase comprising a Glycosyl Hydrolase Family 45 catalytic domain that is a wild-type or variant of a microbially-derived endoglucanase endogenous to *Melanocarpus albomyces*, and a carbohydrate binding module that is a wild-type or variant of a carbohydrate binding module endogenous to *Trichoderma reesei*, and which has at least 90% identity to amino acid sequence SEQ ID NO:4.

30 The composition may comprise an enzyme selected from mannanase, pectate lyase, laccase, polyesterase, galactanase, acyltransferase, and any combination thereof.


 The composition may comprise a perfume, wherein the perfume comprises from 60wt% to 85wt% ester perfume raw materials having the structure:

16



wherein R1 and R2 are independently selected from C1 to C30 linear or branched, cyclic or non-cyclic, aromatic or non-aromatic, saturated or un-saturated, substituted or unsubstituted alkyl.

The composition may comprise: (a) alkyl ethoxylated sulphate having an average degree of ethoxylation of from 0.5 to 2.0; (b) perfume, wherein the perfume comprises from 60wt% to 85wt% ester perfume raw materials having the structure:



wherein R1 and R2 are independently selected from C1 to C30 linear or branched, cyclic or non-cyclic, aromatic or non-aromatic, saturated or un-saturated, substituted or unsubstituted alkyl.

The composition may comprise polyvinyl N oxide polymer.

The composition may comprise: silicate salt particles, especially sodium silicate particles; and/or carbonate salt particles, especially sodium bicarbonate particles. However it may be preferred for the composition to be free of silicate salt particles, especially free of sodium silicate particles. It may also be preferred for the composition to be free of carbonate salt particles, especially free of sodium carbonate particles.

Preferably, the composition comprises from 1wt% to 10wt% dry-added acid particles, preferably from 2wt% to 8wt% dry-added acid particles. A suitable dry-added acid is an organic acid, preferably a carboxylic acid, preferably citric acid.

Base detergent particle: The solid free-flowing particulate laundry detergent composition typically comprises a base detergent particle. The base detergent particle may be in the form of spray-dried particle, or an agglomerate, preferably the base particle is in the form of a spray-dried particle. Typically, the composition comprises from 30wt% to 90wt% base detergent particle, preferably from 40wt% to 80wt%, more preferably from 50wt% to 70wt% base detergent particle.

The base detergent particle typically comprises from 1wt% to 10wt% organic acid, preferably from 2wt% to 8wt%, or from 3wt% to 7wt% organic acid. A preferred organic acid is a carboxylic acid, preferably citric acid.


5 The base detergent particle typically comprises from 1wt% to 10wt% magnesium sulphate, preferably from 2wt% to 8wt%, or from 3wt% to 6wt% magnesium sulphate.

The base detergent particle typically comprises from 1wt% to 8wt%, preferably from 2wt% to 6wt% or from 2wt% to 4wt% zeolite. A preferred zeolite is zeolite A, especially zeolite 4A.

10 The base detergent particle typically comprises from 5wt% to 40wt%, preferably from 10wt% to 30wt% anionic deterative surfactant. A preferred anionic deterative surfactant is alkyl benzene sulphonate.

The base detergent particle typically comprises from 0.5wt% to 5wt% polymer, preferably from 1wt% to 3wt% polymer. A preferred polymer is a carboxylate polymer, more preferably a co-polymer that comprises: (i) from 50 to less than 98 wt% structural units derived from one or more monomers comprising carboxyl groups; (ii) from 1 to less than 49 wt% structural units
15 derived from one or more monomers comprising sulfonate moieties; and (iii) from 1 to 49 wt% structural units derived from one or more types of monomers selected from ether bond-containing monomers represented by formulas (I) and (II):

formula (I):




20

wherein in formula (I), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, X represents a number 0-5 provided X represents a number 1-5 when R is a single bond, and R_1 is a hydrogen atom or C_1 to C_{20} organic group;

25

formula (II)



5 wherein in formula (II), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, x represents a number 0-5, and R_1 is a hydrogen atom or C_1 to C_{20} organic group.

It may be preferred that the polymer has a weight average molecular weight of at least 50kDa, or even at least 70kDa.

10 Typically, the base detergent particle comprises from 30wt% to 70wt%, or from 40wt% to 70wt% sodium sulphate.

Co-surfactant particle: Typically, the detergent composition comprises a co-surfactant particle. Typically, the composition comprises from 1wt% to 20wt%, or from 2wt% to 15wt%, or from 3wt% to 10wt% co-surfactant particle. Typically, the co-surfactant particle is in the form of
 15 an agglomerate, extrudate, needle, noodle, flake or any combination thereof. Preferably, the co-surfactant particle is in the form of an agglomerate.

The co-surfactant particle typically comprises from 25wt% to 60wt% co-surfactant, preferably from 30wt% to 50wt% co-surfactant. A preferred co-surfactant is alkyl alkoxy sulphate, preferably a C_{10} - C_{20} alkyl ethoxylated sulphate having an average degree of
 20 ethoxylation of from 0.5 to 2.0.

Typically, the co-surfactant particle comprises from 10wt% to 50wt% carbonate salt. A preferred carbonate salt is sodium carbonate and/or sodium bicarbonate. However, it may be preferred for the co-surfactant particle to be free of carbonate salt, especially free of sodium carbonate.

25 Typically, the co-surfactant particle comprises from 1wt% to 30wt% silica, preferably from 5wt% to 20wt% silica.

Detergent Ingredients: Suitable laundry detergent compositions comprise a detergent ingredient selected from: deterative surfactant, such as anionic deterative surfactants, non-ionic deterative surfactants, cationic deterative surfactants, zwitterionic deterative surfactants and amphoteric deterative surfactants; polymers, such as carboxylate polymers, soil release polymer, anti-redeposition polymers, cellulosic polymers and care polymers; bleach, such as sources of hydrogen peroxide, bleach activators, bleach catalysts and pre-formed peracids; photobleach, such as such as zinc and/or aluminium sulphonated phthalocyanine; enzymes, such as proteases, amylases, cellulases, lipases; zeolite builder; phosphate builder; co-builders, such as citric acid and citrate; sulphate salt, such as sodium sulphate; chloride salt, such as sodium chloride; brighteners; chelants; hueing agents; dye transfer inhibitors; dye fixative agents; perfume; silicone; fabric softening agents, such as clay; flocculants, such as polyethyleneoxide; suds suppressors; and any combination thereof.

The composition may comprise: silicate salt, especially sodium silicate; and/or carbonate salt, especially sodium bicarbonate and/or sodium carbonate. However it may be preferred for the composition to be free of silicate salt, especially free of sodium silicate. It may also be preferred for the composition to be free of carbonate salt, especially free of sodium carbonate and/or sodium bicarbonate.

The composition may have a pH profile such that upon dilution in de-ionized water at a concentration of 1g/L at a temperature of 20°C, the composition has a pH in the range of from 6.5 to 8.5, preferably from 7.0 to 8.0.

Suitable laundry detergent compositions may have a low buffering capacity. Such laundry detergent compositions typically have a reserve alkalinity to pH 7.5 of less than 5.0gNaOH/100g, preferably less than 3.0gNaOH/100g.

The composition is preferably substantially free of pre-formed peracid. The composition is preferably substantially free of phthalimido-peroxycaproic acid. Substantially free means no deliberately added.

Deterative Surfactant: Suitable deterative surfactants include anionic deterative surfactants, non-ionic deterative surfactant, cationic deterative surfactants, zwitterionic deterative surfactants and amphoteric deterative surfactants. Suitable deterative surfactants may be linear or branched, substituted or un-substituted, and may be derived from petrochemical material or biomaterial.

Anionic deterative surfactant: Suitable anionic deterative surfactants include sulphonate and sulphate deterative surfactants.

Suitable sulphonate deterative surfactants include methyl ester sulphonates, alpha olefin sulphonates, alkyl benzene sulphonates, especially alkyl benzene sulphonates, preferably C₁₀₋₁₃ alkyl benzene sulphonate. Suitable alkyl benzene sulphonate (LAS) is obtainable, preferably obtained, by sulphonating commercially available linear alkyl benzene (LAB); suitable LAB includes low 2-phenyl LAB, other suitable LAB include high 2-phenyl LAB, such as those supplied by Sasol under the tradename Hyblene®.

Suitable sulphate deterative surfactants include alkyl sulphate, preferably C₈₋₁₈ alkyl sulphate, or predominantly C₁₂ alkyl sulphate.

A preferred sulphate deterative surfactant is alkyl alkoxyated sulphate, preferably alkyl ethoxyated sulphate, preferably a C₈₋₁₈ alkyl alkoxyated sulphate, preferably a C₈₋₁₈ alkyl ethoxyated sulphate, preferably the alkyl alkoxyated sulphate has an average degree of alkoxylation of from 0.5 to 20, preferably from 0.5 to 10, preferably the alkyl alkoxyated sulphate is a C₈₋₁₈ alkyl ethoxyated sulphate having an average degree of ethoxylation of from 0.5 to 10, preferably from 0.5 to 5, more preferably from 0.5 to 3 and most preferably from 0.5 to 1.5.

The alkyl sulphate, alkyl alkoxyated sulphate and alkyl benzene sulphonates may be linear or branched, substituted or un-substituted, and may be derived from petrochemical material or biomaterial.

Other suitable anionic deterative surfactants include alkyl ether carboxylates.

Suitable anionic deterative surfactants may be in salt form, suitable counter-ions include sodium, calcium, magnesium, amino alcohols, and any combination thereof. A preferred counter-ion is sodium.

Non-ionic deterative surfactant: Suitable non-ionic deterative surfactants are selected from the group consisting of: C₈-C₁₈ alkyl ethoxylates, such as, NEODOL® non-ionic surfactants from Shell; C₆-C₁₂ alkyl phenol alkoxyates wherein preferably the alkoxyate units are ethyleneoxy units, propyleneoxy units or a mixture thereof; C₁₂-C₁₈ alcohol and C₆-C₁₂ alkyl phenol condensates with ethylene oxide/propylene oxide block polymers such as Pluronic® from BASF; alkylpolysaccharides, preferably alkylpolyglycosides; methyl ester ethoxylates; polyhydroxy fatty acid amides; ether capped poly(oxyalkylated) alcohol surfactants; and mixtures thereof.


Suitable non-ionic deterative surfactants are alkylpolyglucoside and/or an alkyl alkoxyated alcohol.

Suitable non-ionic deterative surfactants include alkyl alkoxyated alcohols, preferably C₈₋₁₈ alkyl alkoxyated alcohol, preferably a C₈₋₁₈ alkyl ethoxyated alcohol, preferably the alkyl alkoxyated alcohol has an average degree of alkoxylation of from 1 to 50, preferably from 1 to 30, or from 1 to 20, or from 1 to 10, preferably the alkyl alkoxyated alcohol is a C₈₋₁₈ alkyl ethoxyated alcohol having an average degree of ethoxylation of from 1 to 10, preferably from 1 to 7, more preferably from 1 to 5 and most preferably from 3 to 7. The alkyl alkoxyated alcohol can be linear or branched, and substituted or un-substituted.

Suitable nonionic deterative surfactants include secondary alcohol-based deterative surfactants.

Cationic deterative surfactant: Suitable cationic deterative surfactants include alkyl pyridinium compounds, alkyl quaternary ammonium compounds, alkyl quaternary phosphonium compounds, alkyl ternary sulphonium compounds, and mixtures thereof.

Preferred cationic deterative surfactants are quaternary ammonium compounds having the general formula:



wherein, R is a linear or branched, substituted or unsubstituted C₆₋₁₈ alkyl or alkenyl moiety, R₁ and R₂ are independently selected from methyl or ethyl moieties, R₃ is a hydroxyl, hydroxymethyl or a hydroxyethyl moiety, X is an anion which provides charge neutrality, preferred anions include: halides, preferably chloride; sulphate; and sulphonate.

Zwitterionic deterative surfactant: Suitable zwitterionic deterative surfactants include amine oxides and/or betaines.


Polymer: Suitable polymers include carboxylate polymers, soil release polymers, anti-redeposition polymers, cellulosic polymers, care polymers and any combination thereof.

Carboxylate polymer: The composition may comprise a carboxylate polymer, such as a maleate/acrylate random copolymer or polyacrylate homopolymer. Suitable carboxylate polymers include: polyacrylate homopolymers having a molecular weight of from 4,000 Da to 9,000 Da; maleate/acrylate random copolymers having a molecular weight of from 50,000 Da to 100,000 Da, or from 60,000 Da to 80,000 Da.

Another suitable carboxylate polymer is a co-polymer that comprises: (i) from 50 to less than 98 wt% structural units derived from one or more monomers comprising carboxyl groups; (ii) from 1 to less than 49 wt% structural units derived from one or more monomers comprising


sulfonate moieties; and (iii) from 1 to 49 wt% structural units derived from one or more types of monomers selected from ether bond-containing monomers represented by formulas (I) and (II):

formula (I):



wherein in formula (I), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, X represents a number 0-5 provided X represents a number 1-5 when R is a single bond, and R_1 is a hydrogen atom or C_1 to C_{20} organic group;

formula (II)



wherein in formula (II), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, X represents a number 0-5, and R_1 is a hydrogen atom or C_1 to C_{20} organic group.

It may be preferred that the polymer has a weight average molecular weight of at least 50kDa, or even at least 70kDa.

Soil release polymer: The composition may comprise a soil release polymer. A suitable soil release polymer has a structure as defined by one of the following structures (I), (II) or (III):



wherein:

a, b and c are from 1 to 200;

d, e and f are from 1 to 50;

10 Ar is a 1,4-substituted phenylene;

sAr is 1,3-substituted phenylene substituted in position 5 with SO₃Me;

Me is Li, K, Mg/2, Ca/2, Al/3, ammonium, mono-, di-, tri-, or tetraalkylammonium wherein the alkyl groups are C₁-C₁₈ alkyl or C₂-C₁₀ hydroxyalkyl, or mixtures thereof;

R¹, R², R³, R⁴, R⁵ and R⁶ are independently selected from H or C₁-C₁₈ n- or iso-alkyl; and

15 R⁷ is a linear or branched C₁-C₁₈ alkyl, or a linear or branched C₂-C₃₀ alkenyl, or a cycloalkyl group with 5 to 9 carbon atoms, or a C₈-C₃₀ aryl group, or a C₆-C₃₀ arylalkyl group.

Suitable soil release polymers are sold by Clariant under the TexCare® series of polymers, e.g. TexCare® SRN240 and TexCare® SRA300. Other suitable soil release polymers are sold by Solvay under the Repel-o-Tex® series of polymers, e.g. Repel-o-Tex® SF2 and Repel-o-Tex®
20 Crystal.

Anti-redeposition polymer: Suitable anti-redeposition polymers include polyethylene glycol polymers and/or polyethyleneimine polymers.

Suitable polyethylene glycol polymers include random graft co-polymers comprising: (i) hydrophilic backbone comprising polyethylene glycol; and (ii) hydrophobic side chain(s)
25 selected from the group consisting of: C₄-C₂₅ alkyl group, polypropylene, polybutylene, vinyl ester of a saturated C₁-C₆ mono-carboxylic acid, C₁-C₆ alkyl ester of acrylic or methacrylic acid, and mixtures thereof. Suitable polyethylene glycol polymers have a polyethylene glycol backbone with random grafted polyvinyl acetate side chains. The average molecular weight of the polyethylene glycol backbone can be in the range of from 2,000 Da to 20,000 Da, or from
30 4,000 Da to 8,000 Da. The molecular weight ratio of the polyethylene glycol backbone to the polyvinyl acetate side chains can be in the range of from 1:1 to 1:5, or from 1:1.2 to 1:2. The average number of graft sites per ethylene oxide units can be less than 1, or less than 0.8, the average number of graft sites per ethylene oxide units can be in the range of from 0.5 to 0.9, or

the average number of graft sites per ethylene oxide units can be in the range of from 0.1 to 0.5, or from 0.2 to 0.4. A suitable polyethylene glycol polymer is Sokalan HP22. Suitable polyethylene glycol polymers are described in WO08/007320.

Cellulosic polymer: Suitable cellulosic polymers are selected from alkyl cellulose, alkyl alkoxyalkyl cellulose, carboxyalkyl cellulose, alkyl carboxyalkyl cellulose, sulphoalkyl cellulose, more preferably selected from carboxymethyl cellulose, methyl cellulose, methyl hydroxyethyl cellulose, methyl carboxymethyl cellulose, and mixtures thereof.

Suitable carboxymethyl celluloses have a degree of carboxymethyl substitution from 0.5 to 0.9 and a molecular weight from 100,000 Da to 300,000 Da.

Suitable carboxymethyl celluloses have a degree of substitution greater than 0.65 and a degree of blockiness greater than 0.45, e.g. as described in WO09/154933.

Care polymers: Suitable care polymers include cellulosic polymers that are cationically modified or hydrophobically modified. Such modified cellulosic polymers can provide anti-abrasion benefits and dye lock benefits to fabric during the laundering cycle. Suitable cellulosic polymers include cationically modified hydroxyethyl cellulose.

Other suitable care polymers include dye lock polymers, for example the condensation oligomer produced by the condensation of imidazole and epichlorhydrin, preferably in ratio of 1:4:1. A suitable commercially available dye lock polymer is Polyquart® FDI (Cognis).

Other suitable care polymers include amino-silicone, which can provide fabric feel benefits and fabric shape retention benefits.


Bleach: Suitable bleach includes sources of hydrogen peroxide, bleach activators, bleach catalysts, pre-formed peracids and any combination thereof. A particularly suitable bleach includes a combination of a source of hydrogen peroxide with a bleach activator and/or a bleach catalyst.

Source of hydrogen peroxide: Suitable sources of hydrogen peroxide include sodium perborate and/or sodium percarbonate.

Bleach activator: Suitable bleach activators include tetra acetyl ethylene diamine and/or alkyl oxybenzene sulphonate.

Bleach catalyst: The composition may comprise a bleach catalyst. Suitable bleach catalysts include oxaziridinium bleach catalysts, transition metal bleach catalysts, especially manganese and iron bleach catalysts. A suitable bleach catalyst has a structure corresponding to general formula below:

25



wherein R^{13} is selected from the group consisting of 2-ethylhexyl, 2-propylheptyl, 2-butyloctyl, 2-pentylonyl, 2-hexyldecyl, n-dodecyl, n-tetradecyl, n-hexadecyl, n-octadecyl, iso-nonyl, iso-decyl, iso-tridecyl and iso-pentadecyl.

Pre-formed peracid: Suitable pre-form peracids include phthalimido-peroxycaproic acid. However, it is preferred that the composition is substantially free of pre-formed peracid. By: “substantially free” it is meant: “no deliberately added”.

Enzymes: Suitable enzymes include lipases, proteases, cellulases, amylases and any combination thereof.

Protease: Suitable proteases include metalloproteases and/or serine proteases. Examples of suitable neutral or alkaline proteases include: subtilisins (EC 3.4.21.62); trypsin-type or chymotrypsin-type proteases; and metalloproteases. The suitable proteases include chemically or genetically modified mutants of the aforementioned suitable proteases.

Suitable commercially available protease enzymes include those sold under the trade names Alcalase®, Savinase®, Primase®, Durazym®, Polarzyme®, Kannase®, Liquanase®, Liquanase Ultra®, Savinase Ultra®, Ovozyme®, Neutrase®, Everlase® and Esperase® by Novozymes A/S (Denmark), those sold under the tradename Maxatase®, Maxacal®, Maxapem®, Preferenz P® series of proteases including Preferenz® P280, Preferenz® P281, Preferenz® P2018-C, Preferenz® P2081-WE, Preferenz® P2082-EE and Preferenz® P2083-A/J, Properase®, Purafect®, Purafect Prime®, Purafect Ox®, FN3®, FN4®, Excellase® and Purafect OXP® by DuPont, those sold under the tradename Opticlean® and Optimase® by Solvay Enzymes, those available from Henkel/ Kemira, namely BLAP (sequence shown in Figure 29 of US 5,352,604 with the following mutations S99D + S101 R + S103A + V104I + G159S, hereinafter referred to as BLAP), BLAP R (BLAP with S3T + V4I + V199M + V205I + L217D), BLAP X (BLAP with S3T + V4I + V205I) and BLAP F49 (BLAP with S3T + V4I + A194P + V199M + V205I + L217D) - all from Henkel/Kemira; and KAP (Bacillus alkalophilus subtilisin with mutations A230V + S256G + S259N) from Kao.

A suitable protease is described in WO11/140316 and WO11/072117.

Amylase: Suitable amylases are derived from AA560 alpha amylase endogenous to *Bacillus* sp. DSM 12649, preferably having the following mutations: R118K, D183*, G184*, N195F, R320K, and/or R458K. Suitable commercially available amylases include Stainzyme®, Stainzyme® Plus, Natalase, Termamyl®, Termamyl® Ultra, Liquezyme® SZ, Duramyl®, Everest® (all Novozymes) and Spezyme® AA, Preferenz S® series of amylases, Purastar® and Purastar® Ox Am, Optisize® HT Plus (all Du Pont).

A suitable amylase is described in WO06/002643.

Cellulase: Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are also suitable. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g., the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum*.

Commercially available cellulases include Celluzyme®, Carezyme®, and Carezyme® Premium, Celluclean® and Whitezyme® (Novozymes A/S), Revitalenz® series of enzymes (Du Pont), and Biotouch® series of enzymes (AB Enzymes). Suitable commercially available cellulases include Carezyme® Premium, Celluclean® Classic. Suitable cellulases are described in WO07/144857 and WO10/056652.

Lipase: Suitable lipases include those of bacterial, fungal or synthetic origin, and variants thereof. Chemically modified or protein engineered mutants are also suitable. Examples of suitable lipases include lipases from *Humicola* (synonym *Thermomyces*), e.g., from *H. lanuginosa* (*T. lanuginosus*).

The lipase may be a “first cycle lipase”, e.g. such as those described in WO06/090335 and WO13/116261. In one aspect, the lipase is a first-wash lipase, preferably a variant of the wild-type lipase from *Thermomyces lanuginosus* comprising T231R and/or N233R mutations. Preferred lipases include those sold under the tradenames Lipex®, Lipolex® and Lipoclean® by Novozymes, Bagsvaerd, Denmark.

Other suitable lipases include: Liprl 139, e.g. as described in WO2013/171241; and TfuLip2, e.g. as described in WO2011/084412 and WO2013/033318.

Other enzymes: Other suitable enzymes are bleaching enzymes, such as peroxidases/oxidases, which include those of plant, bacterial or fungal origin and variants thereof. Commercially available peroxidases include Guardzyme® (Novozymes A/S). Other suitable enzymes include choline oxidases and perhydrolases such as those used in Gentle Power Bleach™.

Other suitable enzymes include pectate lyases sold under the tradenames X-Pect®, Pectaway® (from Novozymes A/S, Bagsvaerd, Denmark) and PrimaGreen® (DuPont) and mannanases sold under the tradenames Mannaway® (Novozymes A/S, Bagsvaerd, Denmark), and Mannastar® (Du Pont).

5 Identity: When used herein identity or sequence identity refers to the relatedness between two amino acid sequences.

For purposes of the present invention, the degree of sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *J. Mol. Biol.* 48: 443-453) as implemented in the Needle program of the EMBOSS
10 package (EMBOSS: The European Molecular Biology Open Software Suite, Rice *et al.*, 2000, *Trends Genet.* 16: 276-277), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled “longest identity” (obtained using the –nobrief option) is used as the percent identity and is calculated as follows:

15
$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

Zeolite builder: The composition may comprise zeolite builder. The composition may comprise from 0wt% to 5wt% zeolite builder, or 3wt% zeolite builder. The composition may even be substantially free of zeolite builder; substantially free means “no deliberately added”. Typical zeolite builders include zeolite A, zeolite P and zeolite MAP.

20 Phosphate builder: The composition may comprise phosphate builder. The composition may comprise from 0wt% to 5wt% phosphate builder, or to 3wt%, phosphate builder. The composition may even be substantially free of phosphate builder; substantially free means “no deliberately added”. A typical phosphate builder is sodium tri-polyphosphate.

Carbonate salt: The composition may comprise carbonate salt. The composition may
25 comprise from 0wt% to 5wt% carbonate salt. The composition may even be substantially free of carbonate salt; substantially free means “no deliberately added”. Suitable carbonate salts include sodium carbonate and sodium bicarbonate.

Silicate salt: The composition may comprise silicate salt. The composition may comprise from 0wt% to 5wt% silicate salt. The composition may even be substantially free of silicate salt;
30 substantially free means “no deliberately added”. A preferred silicate salt is sodium silicate, especially preferred are sodium silicates having a Na₂O:SiO₂ ratio of from 1.0 to 2.8, preferably from 1.6 to 2.0.

Sulphate salt: A suitable sulphate salt is sodium sulphate.

Brightener: Suitable fluorescent brighteners include: di-styryl biphenyl compounds, e.g. Tinopal® CBS-X, di-amino stilbene di-sulfonic acid compounds, e.g. Tinopal® DMS pure Xtra and Blankophor® HRH, and Pyrazoline compounds, e.g. Blankophor® SN, and coumarin compounds, e.g. Tinopal® SWN.

- 5 Preferred brighteners are: sodium 2 (4-styryl-3-sulfophenyl)-2H-naphthol[1,2-d]triazole, disodium 4,4'-bis{[(4-anilino-6-(N methyl-N-2 hydroxyethyl)amino 1,3,5- triazin-2-yl)]amino}stilbene-2-2' disulfonate, disodium 4,4'-bis{[(4-anilino-6-morpholino-1,3,5-triazin-2-yl)]amino} stilbene-2-2' disulfonate, and disodium 4,4'- bis(2-sulfoxyethyl)biphenyl. A suitable fluorescent brightener is C.I. Fluorescent Brightener 260, which may be used in its beta or alpha crystalline forms, or a mixture of these forms.

- Chelant: The composition may also comprise a chelant selected from: diethylene triamine pentaacetate, diethylene triamine penta(methyl phosphonic acid), ethylene diamine-N'N'-disuccinic acid, ethylene diamine tetraacetate, ethylene diamine tetra(methylene phosphonic acid) and hydroxyethane di(methylene phosphonic acid). A preferred chelant is ethylene diamine-N'N'-disuccinic acid (EDDS) and/or hydroxyethane diphosphonic acid (HEDP). The composition preferably comprises ethylene diamine-N'N'- disuccinic acid or salt thereof. Preferably the ethylene diamine-N'N'-disuccinic acid is in S,S enantiomeric form. Preferably the composition comprises 4,5-dihydroxy-m-benzenedisulfonic acid disodium salt. Preferred chelants may also function as calcium carbonate crystal growth inhibitors such as: 1-hydroxyethanediphosphonic acid (HEDP) and salt thereof; N,N-dicarboxymethyl-2-aminopentane-1,5-dioic acid and salt thereof; 2-phosphonobutane-1,2,4-tricarboxylic acid and salt thereof; and combination thereof.

- Hueing agent: Suitable hueing agents include small molecule dyes, typically falling into the Colour Index (C.I.) classifications of Acid, Direct, Basic, Reactive (including hydrolysed forms thereof) or Solvent or Disperse dyes, for example classified as Blue, Violet, Red, Green or Black, and provide the desired shade either alone or in combination. Preferred such hueing agents include Acid Violet 50, Direct Violet 9, 66 and 99, Solvent Violet 13 and any combination thereof.

- Many hueing agents are known and described in the art which may be suitable for the present invention, such as hueing agents described in WO2014/089386.

Suitable hueing agents include phthalocyanine and azo dye conjugates, such as described in WO2009/069077.

Suitable hueing agents may be alkoxyated. Such alkoxyated compounds may be produced by organic synthesis that may produce a mixture of molecules having different degrees of alkoxylation. Such mixtures may be used directly to provide the hueing agent, or may undergo a purification step to increase the proportion of the target molecule. Suitable hueing agents include alkoxyated bis-azo dyes, such as described in WO2012/054835, and/or alkoxyated thiophene azo dyes, such as described in WO2008/087497 and WO2012/166768.

The hueing agent may be incorporated into the detergent composition as part of a reaction mixture which is the result of the organic synthesis for a dye molecule, with optional purification step(s). Such reaction mixtures generally comprise the dye molecule itself and in addition may comprise un-reacted starting materials and/or by-products of the organic synthesis route. Suitable hueing agents can be incorporated into hueing dye particles, such as described in WO 2009/069077.

Dye transfer inhibitors: Suitable dye transfer inhibitors include polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone, polyvinylloxazolidone, polyvinylimidazole and mixtures thereof. Preferred are poly(vinyl pyrrolidone), poly(vinylpyridine betaine), poly(vinylpyridine N-oxide), poly(vinyl pyrrolidone-vinyl imidazole) and mixtures thereof. Suitable commercially available dye transfer inhibitors include PVP-K15 and K30 (Ashland), Sokalan® HP165, HP50, HP53, HP59, HP56K, HP56, HP66 (BASF), Chromabond® S-400, S403E and S-100 (Ashland).

Perfume: Suitable perfumes comprise perfume materials selected from the group: (a) perfume materials having a ClogP of less than 3.0 and a boiling point of less than 250°C (quadrant 1 perfume materials); (b) perfume materials having a ClogP of less than 3.0 and a boiling point of 250°C or greater (quadrant 2 perfume materials); (c) perfume materials having a ClogP of 3.0 or greater and a boiling point of less than 250°C (quadrant 3 perfume materials); (d) perfume materials having a ClogP of 3.0 or greater and a boiling point of 250°C or greater (quadrant 4 perfume materials); and (e) mixtures thereof.

It may be preferred for the perfume to be in the form of a perfume delivery technology. Such delivery technologies further stabilize and enhance the deposition and release of perfume materials from the laundered fabric. Such perfume delivery technologies can also be used to further increase the longevity of perfume release from the laundered fabric. Suitable perfume delivery technologies include: perfume microcapsules, pro-perfumes, polymer assisted deliveries, molecule assisted deliveries, fiber assisted deliveries, amine assisted deliveries,

cyclodextrin, starch encapsulated accord, zeolite and other inorganic carriers, and any mixture thereof. A suitable perfume microcapsule is described in WO2009/101593.

Silicone: Suitable silicones include polydimethylsiloxane and amino-silicones. Suitable silicones are described in WO05075616.

- 5 Process for making the solid composition: Typically, the particles of the composition can be prepared by any suitable method. For example: spray-drying, agglomeration, extrusion and any combination thereof.

Typically, a suitable spray-drying process comprises the step of forming an aqueous slurry mixture, transferring it through at least one pump, preferably two pumps, to a pressure nozzle.
10 Atomizing the aqueous slurry mixture into a spray-drying tower and drying the aqueous slurry mixture to form spray-dried particles. Preferably, the spray-drying tower is a counter-current spray-drying tower, although a co-current spray-drying tower may also be suitable.

Typically, the spray-dried powder is subjected to cooling, for example an air lift. Typically, the spray-drying powder is subjected to particle size classification, for example a sieve, to obtain
15 the desired particle size distribution. Preferably, the spray-dried powder has a particle size distribution such that weight average particle size is in the range of from 300 micrometers to 500 micrometers, and less than 10wt% of the spray-dried particles have a particle size greater than 2360 micrometers.

It may be preferred to heat the aqueous slurry mixture to elevated temperatures prior to
20 atomization into the spray-drying tower, such as described in WO2009/158162.

It may be preferred for anionic surfactant, such as linear alkyl benzene sulphonate, to be introduced into the spray-drying process after the step of forming the aqueous slurry mixture: for example, introducing an acid precursor to the aqueous slurry mixture after the pump, such as described in WO 09/158449.

- 25 It may be preferred for a gas, such as air, to be introduced into the spray-drying process after the step of forming the aqueous slurry, such as described in WO2013/181205.

It may be preferred for any inorganic ingredients, such as sodium sulphate and sodium carbonate, if present in the aqueous slurry mixture, to be micronized to a small particle size such as described in WO2012/134969.

- 30 Typically, a suitable agglomeration process comprises the step of contacting a deterative ingredient, such as a deterative surfactant, e.g. linear alkyl benzene sulphonate (LAS) and/or alkyl alkoxylated sulphate, with an inorganic material, such as sodium carbonate and/or silica, in a mixer. The agglomeration process may also be an in-situ neutralization agglomeration process

wherein an acid precursor of a deterative surfactant, such as LAS, is contacted with an alkaline material, such as carbonate and/or sodium hydroxide, in a mixer, and wherein the acid precursor of a deterative surfactant is neutralized by the alkaline material to form a deterative surfactant during the agglomeration process.

5 Other suitable detergent ingredients that may be agglomerated include polymers, chelants, bleach activators, silicones and any combination thereof.

 The agglomeration process may be a high, medium or low shear agglomeration process, wherein a high shear, medium shear or low shear mixer is used accordingly. The agglomeration process may be a multi-step agglomeration process wherein two or more mixers are used, such as
10 a high shear mixer in combination with a medium or low shear mixer. The agglomeration process can be a continuous process or a batch process.

 It may be preferred for the agglomerates to be subjected to a drying step, for example to a fluid bed drying step. It may also be preferred for the agglomerates to be subjected to a cooling step, for example a fluid bed cooling step.

15 Typically, the agglomerates are subjected to particle size classification, for example a fluid bed elutriation and/or a sieve, to obtain the desired particle size distribution. Preferably, the agglomerates have a particle size distribution such that weight average particle size is in the range of from 300 micrometers to 800 micrometers, and less than 10wt% of the agglomerates have a particle size less than 150 micrometers and less than 10wt% of the agglomerates have a
20 particle size greater than 1200 micrometers.

 It may be preferred for fines and over-sized agglomerates to be recycled back into the agglomeration process. Typically, over-sized particles are subjected to a size reduction step, such as grinding, and recycled back into an appropriate place in the agglomeration process, such as the mixer. Typically, fines are recycled back into an appropriate place in the agglomeration
25 process, such as the mixer.

 It may be preferred for ingredients such as polymer and/or non-ionic deterative surfactant and/or perfume to be sprayed onto base detergent particles, such as spray-dried base detergent particles and/or agglomerated base detergent particles. Typically, this spray-on step is carried out in a tumbling drum mixer.

30 Method of laundering fabric: The method of laundering fabric comprises the step of contacting the solid composition to water to form a wash liquor, and laundering fabric in said wash liquor. Typically, the wash liquor has a temperature of above 0°C to 90°C, or to 60°C, or to 40°C, or to 30°C, or to 20°C. The fabric may be contacted to the water prior to, or after, or

simultaneous with, contacting the solid composition with water. Typically, the wash liquor is formed by contacting the laundry detergent to water in such an amount so that the concentration of laundry detergent composition in the wash liquor is from 0.2g/l to 20g/l, or from 0.5g/l to 10g/l, or to 5.0g/l. The method of laundering fabric can be carried out in a front-loading automatic washing machine, top loading automatic washing machines, including high efficiency automatic washing machines, or suitable hand-wash vessels. Typically, the wash liquor comprises 90 litres or less, or 60 litres or less, or 15 litres or less, or 10 litres or less of water. Typically, 200g or less, or 150g or less, or 100g or less, or 50g or less of laundry detergent composition is contacted to water to form the wash liquor.

Solid free-flowing particulate laundry detergent composition illustrative examples:

<u>Ingredient</u>	<u>Amount (in wt%)</u>
<u>Anionic deterative surfactant</u> (such as alkyl benzene sulphonate, alkyl ethoxylated sulphate and mixtures thereof)	from 8wt% to 15wt%
<u>Non-ionic deterative surfactant</u> (such as alkyl ethoxylated alcohol)	from 0.1wt% to 4wt%
<u>Cationic deterative surfactant</u> (such as quaternary ammonium compounds)	from 0wt% to 4wt%
<u>Other deterative surfactant</u> (such as zwitterionic deterative surfactants, amphoteric surfactants and mixtures thereof)	from 0wt% to 4wt%
<u>Carboxylate polymer</u> (such as co-polymers of maleic acid and acrylic acid and/or carboxylate polymers comprising ether moieties and sulfonate moieties)	from 0.1wt% to 4wt%
<u>Polyethylene glycol polymer</u> (such as a polyethylene glycol polymer comprising polyvinyl acetate side chains)	from 0wt% to 4wt%
<u>Polyester soil release polymer</u> (such as Repel-o-tex and/or Texcare polymers)	from 0wt% to 2wt%
<u>Cellulosic polymer</u> (such as carboxymethyl cellulose, methyl cellulose and combinations thereof)	from 0.5wt% to 2wt%
<u>Other polymer</u> (such as care polymers)	from 0wt% to 4wt%
<u>Zeolite builder and phosphate builder</u> (such as zeolite 4A and/or sodium tripolyphosphate)	from 0wt% to 4wt%

<u>Other co-builder</u> (such as sodium citrate and/or citric acid)	from 0wt% to 3wt%
<u>Citric Acid</u>	from 4wt% to 16wt%
<u>Magnesium Sulphate</u>	from 1wt% to 4wt%
<u>Carbonate salt</u> (such as sodium carbonate and/or sodium bicarbonate)	from 0wt% to 4wt%
<u>Silicate salt</u> (such as sodium silicate)	from 0wt% to 4wt%
<u>Filler</u> (such as sodium sulphate and/or bio-fillers)	from 10wt% to 70wt%
<u>Source of hydrogen peroxide</u> (such as sodium percarbonate)	from 0wt% to 20wt%
<u>Bleach activator</u> (such as tetraacetylene diamine (TAED) and/or nonanoyloxybenzenesulphonate (NOBS))	from 0wt% to 8wt%
<u>Bleach catalyst</u> (such as oxaziridinium-based bleach catalyst and/or transition metal bleach catalyst)	from 0wt% to 0.1wt%
<u>Other bleach</u> (such as reducing bleach and/or pre-formed peracid)	from 0wt% to 10wt%
<u>Photobleach</u> (such as zinc and/or aluminium sulphonated phthalocyanine)	from 0wt% to 0.1wt%
<u>Chelant</u> (such as ethylenediamine-N'N'-disuccinic acid (EDDS) and/or hydroxyethane diphosphonic acid (HEDP))	from 0.2wt% to 1wt%
<u>Hueing agent</u> (such as direct violet 9, 66, 99, acid red 50, solvent violet 13 and any combination thereof)	from 0wt% to 1wt%
<u>Brightener</u> (C.I. fluorescent brightener 260 or C.I. fluorescent brightener 351)	from 0.1wt% to 0.4wt%
<u>Protease</u> (such as Savinase, Savinase Ultra, Purafect, FN3, FN4 and any combination thereof)	from 0.1wt% to 0.4wt%
<u>Amylase</u> (such as Termamyl, Termamyl ultra, Natalase, Optisize, Stainzyme, Stainzyme Plus and any combination thereof)	from 0wt% to 0.2wt%
<u>Cellulase</u> (such as Carezyme and/or Celluclean)	from 0wt% to 0.2wt%
<u>Lipase</u> (such as Lipex, Lipolex, Lipoclean and any combination thereof)	from 0wt% to 1wt%
<u>Other enzyme</u> (such as xyloglucanase, cutinase, pectate lyase, mannanase, bleaching enzyme)	from 0wt% to 2wt%

<u>Fabric softener</u> (such as montmorillonite clay and/or polydimethylsiloxane (PDMS))	from 0wt% to 15wt%
<u>Flocculant</u> (such as polyethylene oxide)	from 0wt% to 1wt%
<u>Suds suppressor</u> (such as silicone and/or fatty acid)	from 0wt% to 4wt%
<u>Perfume</u> (such as perfume microcapsule, spray-on perfume, starch encapsulated perfume accords, perfume loaded zeolite, and any combination thereof)	from 0.1wt% to 1wt%
<u>Aesthetics</u> (such as coloured soap rings and/or coloured speckles/noodles)	from 0wt% to 1wt%
<u>Miscellaneous</u>	balance to 100wt%

EXAMPLES

Example 1 – Low pH formulation with hueing agent (embodiment of the present invention)

- 5 A low pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic detergent surfactant	18.22
Sodium sulphate	75.23
Citric acid	6.54
Water & miscellaneous	to 100wt%

- 10 143g Sodium sulphate, 18g sodium carbonate, 18g sodium silicate, and 2.5g hueing dye particle (dye having a structure in accordance with claim 1) were added to the 321g base powder to form 502.5g of solid free-flowing particulate laundry detergent composition (in accordance with the present invention) having the following formulation:

<u>Ingredient</u>	<u>Amount (wt% of composition)</u>
Alkyl benzene sulphonate anionic deterative surfactant	11.6
Sodium sulphate	76.4
Citric acid	4.2
Sodium carbonate	3.6
Sodium silicate	3.6
Hueing dye particle	0.5
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 7.0.

The composition had a reserve alkalinity to pH 7 at 1wt% dilution in deionized water at 20°C of 2.0.

5

Example 2 – Low pH formulation without hueing dye (comparative example)

A low pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

10

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.22
Sodium sulphate	75.23
Citric acid	6.45
Water & miscellaneous	to 100wt%

143g Sodium sulphate, 18g sodium carbonate, and 18g sodium silicate added to the 321g base powder to form 500g of solid free-flowing particulate laundry detergent composition (comparative example) having the following formulation:

15

<u>Ingredient</u>	<u>Amount (wt% of composition)</u>
Alkyl benzene sulphonate anionic deterative surfactant	11.7
Sodium sulphate	76.9
Citric acid	4.2
Sodium carbonate	3.6
Sodium silicate	3.6
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 7.0.

The composition had a reserve alkalinity to pH 7 at 1wt% dilution in deionized water at 20°C of 2.0.

5

Example 3 – High pH formulation with hueing dye (comparative example)

A high pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

10

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.22
Sodium sulphate	75.23
Citric acid	6.54
Water & miscellaneous	to 100wt%

25g Sodium sulphate, 100g sodium carbonate, 50g sodium silicate, 3g citric acid and 2.5 g hueing dye particle (dye having a structure in accordance with claim 1) were added to the 321g base powder to form 502.5g of solid free-flowing particulate laundry detergent composition (comparative example) having the following formulation:

15

<u>Ingredient</u>	<u>Amount (wt% of composition)</u>
Alkyl benzene sulphonate anionic deterative surfactant	11.6
Sodium sulphate	52.8
Citric acid	5
Sodium carbonate	20
Sodium silicate	10.0
Hueing dye particle	0.5
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 10.5.

The composition had a reserve alkalinity to pH 7 at 1wt% dilution in deionized water at 20°C of 9.6.

5

Example 4 – High pH formulation without hueing dye (comparative example)

A high pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.22
Sodium sulphate	75.23
Citric acid	6.54
Water & miscellaneous	to 100wt%

10

25g Sodium sulphate, 100g sodium carbonate, and 50g sodium silicate and 4g citric acid were added to the 321g base powder to form 500g of solid free-flowing particulate laundry detergent composition (comparative example) having the following formulation:

15

<u>Ingredient</u>	<u>Amount (wt% of composition)</u>
Alkyl benzene sulphonate anionic deterative surfactant	11.7
Sodium sulphate	53.3
Citric acid	5
Sodium carbonate	20
Sodium silicate	10
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 10.5.

The composition had a reserve alkalinity to pH 7 at 1wt% dilution in deionized water at 20°C of 9.6.

5

Washing and whiteness measure method: The following method demonstrates the ability of Samples 1-4 to improve whiteness of fabrics while reducing unwanted hue build-up. The above samples were added separately into the pots of a tergotometer (quantity of sample = 1% of the bulk preparation as described in the Examples, sampled-down uniformly to give a representative sample). The volume of each pot was 1 L. The wash temperature was set to 30 °C. Throughout the procedure, 8.1 gpg water was used. The products were agitated for 2 minutes before addition of fabrics (5x5 cm swatches of unbrightened cotton (10 internal replicates) with additional unbrightened cotton ballast (35 g)). Once the fabrics were added, the wash solution was agitated for 30 minutes. The wash solutions were then drained and the fabrics were subject to a 5 minute

10

rinse step before being drained and spun dry. The washed fabrics were then dried in a gas dryer before being analysed to measure the whiteness of the fabric. The procedure was repeated a further three times to build-up a four-cycle history on the fabrics, alternating tergotometer pots after each cycle to avoid apparatus bias.

15

Whiteness Analysis: The fabrics were analysed using commercially available ColorEye software for L, a, b values. CIE whiteness (WCIE) values were obtained from the L, a, b values using the Color Slide Rule by Axiphos. The higher the WCIE, the greater the whiteness. The higher the a value, the more red the hue.

20

	WCIE	a value
Sample 1: low pH with hue (invention)	32.22	-0.11
Sample 2: low pH nil hue (comparative)	29.46	+0.34
Sample 3: high pH with hue (comparative)	29.34	-0.14
Sample 4: high pH nil hue (comparative)	26.27	+0.49

With the low pH laundry detergent powder, the hueing agent provided greater whiteness performance (higher WCIE) and less unwanted dye build up (lower a value) compared to the high pH laundry detergent powder.

5

Example 5 – pH 8.4 formulation with 4% Sodium Carbonate with hueing agent (embodiment of the present invention)

A low pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

10

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.48
Sodium sulphate	76.3
Citric acid	5.21
Water & miscellaneous	to 100wt%

137g Sodium sulphate, 20g sodium carbonate, 18g sodium silicate, 5g zeolite builder, 3.5g citric acid and 2.5 g hueing dye particle (dye having a structure in accordance with claim 1) were added to the 316.5g base powder to form 502.5g of solid free-flowing particulate laundry detergent composition (in accordance with the present invention) having the following formulation:

15

Ingredient	Amount (wt% of composition)
Alkyl benzene sulphonate anionic deterative surfactant	11.7
Sodium sulphate	75.2
Citric acid	4
Sodium carbonate	4
Sodium silicate	3.6
Zeolite Builder	1
Hueing Dye Particle	0.5
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 8.5.

Example 6 – pH 8.4 formulation with 10% Sodium Carbonate with hueing agent (Comparative Example)

A low pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.48
Sodium sulphate	76.30
Citric acid	5.21
Water & miscellaneous	to 100wt%

88.5g Sodium sulphate, 50g sodium carbonate, 18g sodium silicate, 5g zeolite builder, 22g citric acid and 2.5 g hueing dye particle were added to the 316.5g base powder to form 502.5g of solid free-flowing particulate laundry detergent composition (comparative example) having the following formulation:

Ingredient	Amount (wt% of composition)
Alkyl benzene sulphonate anionic deterative surfactant	11.7
Sodium sulphate	65.5
Citric acid	7.7
Sodium carbonate	10
Sodium silicate	3.6
Zeolite Builder	1
Hueing Dye Particle	0.5
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 8.5.

Example 7 – pH 9.7 formulation with hueing agent (comparative example)

5

A high pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.22
Sodium sulphate	76.30
Citric acid	5.21
Water & miscellaneous	to 100wt%

- 10 140.5g Sodium sulphate, 20g sodium carbonate, 18g sodium silicate, 5g zeolite builder and 2.5 g hueing dye particle were added to the 321g base powder to form 502.5g of solid free-flowing particulate laundry detergent composition (comparative example) having the following formulation:

15

<u>Ingredient</u>	<u>Amount (wt% of composition)</u>
Alkyl benzene sulphonate anionic deterative surfactant	11.7
Sodium sulphate	75.9
Citric acid	3.3
Sodium carbonate	4
Sodium silicate	3.6
Zeolite Builder	1
Hueing Dye Particle	0.5
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 9.7

Example 8 – pH 8.4 formulation with 4% Sodium Carbonate without hueing agent (Comparative Example)

5

A low pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.48
Sodium sulphate	76.30
Citric acid	5.21
Water & miscellaneous	to 100wt%

- 10 137g Sodium sulphate, 20g sodium carbonate, 18g sodium silicate, 5g zeolite builder and 3.5g citric acid were added to the 316.5g base powder to form 500g of solid free-flowing particulate laundry detergent composition (comparative example) having the following formulation:

Ingredient	Amount (wt% of composition)
Alkyl benzene sulphonate anionic deterative surfactant	11.7
Sodium sulphate	75.7
Citric acid	4
Sodium carbonate	4
Sodium silicate	3.6
Zeolite Builder	1
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 8.5.

Example 9 – pH 8.4 formulation with 10% Sodium Carbonate without hueing agent
 5 (Comparative Example)

A low pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.48
Sodium sulphate	76.30
Citric acid	5.21
Water & miscellaneous	to 100wt%

10 88.5g Sodium sulphate, 50g sodium carbonate, 18g sodium silicate, 5g zeolite builder and were added to the 316.5g base powder to form 500g of solid free-flowing particulate laundry detergent composition (comparative example) having the following formulation:

Ingredient	Amount (wt% of composition)
Alkyl benzene sulphonate anionic deterative surfactant	11.7
Sodium sulphate	66
Citric acid	7.7
Sodium carbonate	10
Sodium silicate	3.6
Zeolite Builder	1
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 8.5.

Example 10 – pH 9.7 formulation without hueing agent (comparative example)

5

A high pH base powder was prepared by mixing the ingredients together. The composition of the base powder was:

<u>Ingredient</u>	<u>Amount (wt% of base powder)</u>
Alkyl benzene sulphonate anionic deterative surfactant	18.48
Sodium sulphate	76.30
Citric acid	5.21
Water & miscellaneous	to 100wt%

- 10 140.5g Sodium sulphate, 20g sodium carbonate, 18g sodium silicate, 5g zeolite builder and were added to the 316.5g base powder to form 500g of solid free-flowing particulate laundry detergent composition (comparative example) having the following formulation:

<u>Ingredient</u>	<u>Amount (wt% of composition)</u>
Alkyl benzene sulphonate anionic deterative surfactant	11.7
Sodium sulphate	76.4
Citric acid	3.3

Sodium carbonate	4
Sodium silicate	3.6
Zeolite Builder	1
Water & miscellaneous	to 100wt%

The composition had an equilibrium pH at 1wt% dilution in deionized water at 20°C of 9.7.

Washing and whiteness measure method: The following method demonstrates the ability of Samples 5-10 to improve whiteness of fabrics while reducing unwanted hue build-up. The above samples were added separately into the pots of a tergotometer (quantity of sample = 1% of the bulk preparation as described in the Examples, sampled-down uniformly to give a representative sample). The volume of each pot was 1 L. The wash temperature was set to 20 °C. Throughout the procedure, 0.05 water was used. The products were agitated for 2 minutes before addition of fabrics (5x5 cm swatches of knitted cotton and polyester (10 internal replicates) with additional knitted cotton ballast (35 g)). Once the fabrics were added, the wash solution was agitated for 20 minutes. The wash solutions were then drained and the fabrics were subject to a 5 minute rinse step before being drained and spun dry. The washed fabrics were then dried in a gas dryer before being analysed to measure the whiteness of the fabric. The procedure was repeated a further three times to build-up a four-cycle history on the fabrics, alternating tergotometer pots after each cycle to avoid apparatus bias.

Whiteness Analysis: The fabrics were analysed using commercially available ColorEye software for L, a, b values. CIE whiteness (WCIE) values were obtained from the L, a, b values using the Color Slide Rule by Axiphos. The higher the WCIE, the greater the whiteness.

Sample	Delta WCIE Vs Nil Hueing Agent
Example 7: pH 8.5 with 4% Sodium Carbonate with hueing agent (in accordance with the present invention)	6.97
Example 8: pH 8.5 with 10% Sodium Carbonate with hueing agent (Comparative Example)	2.51
Example 9: pH 9.7 with agent (Comparative Example)	2.67

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as “40 mm” is intended to mean


5 “about 40 mm”.

CLAIMS

What is claimed is:

1. A solid free flowing particulate laundry detergent composition comprising:

- (a) anionic deterative surfactant;
- (b) from 0wt% to 8wt% zeolite builder;
- (c) from 0wt% to 4wt% phosphate builder;
- (d) from 0wt% to 8wt% sodium carbonate;
- (e) from 0wt% to 8wt% sodium silicate;
- (f) from 4wt% to 20wt% organic acid; and
- (g) hueing agent having the following structure:



wherein:

R1 and R2 are independently selected from the group consisting of: H; alkyl; alkoxy; alkyleneoxy; alkyl capped alkyleneoxy; urea; and amido;

R3 is a substituted aryl group;

X is a substituted group comprising sulfonamide moiety and optionally an alkyl and/or aryl moiety, and wherein the substituent group comprises at least one alkyleneoxy chain that comprises an average molar distribution of at least four alkyleneoxy moieties, wherein the composition at 1wt% dilution in deionized water at 20°C, has an equilibrium pH in the range of from 6.5 to 9.0,


wherein the composition comprises from 30wt% to 90wt% base detergent particle,

wherein the base detergent particle comprising (by weight of the base detergent particle):

- (a) from 4wt% to 35wt% anionic deterative surfactant;
- (b) optionally, from 1wt% to 8wt% zeolite builder;
- (c) from 0wt% to 4wt% phosphate builder;
- (d) from 0wt% to 8wt% sodium carbonate;
- (e) from 0wt% to 8wt% sodium silicate;

- (f) from 1wt% to 10wt% organic acid; and
 - (g) optionally, from 1wt% to 10wt% magnesium sulphate.
2. A composition according to claim 1, wherein the composition at 1wt% dilution in deionized water at 20°C, has an equilibrium pH in the range of from 6.5 to 8.5.
 3. A composition according to any preceding claim, wherein the organic acid comprises citric acid, and wherein the base detergent particle comprises from 1wt% to 10wt% citric acid, and wherein optionally the organic acid is at least partially coated with a water-dispersible material.
 4. A composition according to any preceding claim, wherein:
 - (a) the anionic deterative surfactant comprises alkyl benzene sulphonate and wherein the base detergent particle comprises from 4wt% to 35wt% alkyl benzene sulphonate; and/or
 - (b) the base detergent particle comprises from 0.5wt% to 5wt% carboxylate co-polymer, wherein the carboxylate co-polymer comprises:
 - (i) from 50 to less than 98 wt% structural units derived from one or more monomers comprising carboxyl groups;
 - (ii) from 1 to less than 49 wt% structural units derived from one or more monomers comprising sulfonate moieties; and
 - (iii) from 1 to 49 wt% structural units derived from one or more types of monomers selected from ether bond-containing monomers represented by formulas (I) and (II):


formula (I):



wherein in formula (I), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, x represents a number 0-5 provided x represents a

number 1-5 when R is a single bond, and R₁ is a hydrogen atom or C₁ to C₂₀ organic group;

formula (II)



wherein in formula (II), R₀ represents a hydrogen atom or CH₃ group, R represents a CH₂ group, CH₂CH₂ group or single bond, X represents a number 0-5, and R₁ is a hydrogen atom or C₁ to C₂₀ organic group; and/or

(c) wherein the base detergent particle comprises from 30wt% to 70wt% sodium sulphate.

5. A composition according to any preceding claim, wherein the composition comprises from 1wt% to 20wt% co-surfactant particle, wherein the co-surfactant particle comprises:


- (a) from 25wt% to 60wt% co-surfactant;
- (b) from 10wt% to 50wt% carbonate salt; and
- (c) from 1wt% to 30wt% silica,

and wherein optionally:

- (a) the co-surfactant particle is in the form of an agglomerate; and/or
- (b) the co-surfactant comprises alkyl ethoxylated sulphate having an average degree of ethoxylation of from 0.5 to 2.5, and wherein the co-surfactant particle comprises from 25wt% to 60wt% alkyl ethoxylated sulphate having an average degree of ethoxylation of from 0.5 to 2.5; and/or
- (c) the co-surfactant particle comprises linear alkyl benzene sulphonate and alkyl ethoxylated sulphate having an average degree of ethoxylation of from 0.5 to 2.5.


6. A composition according to any preceding claim wherein the composition at 1wt% dilution in deionized water at 20°C, has an equilibrium pH in the range of from 6.5 to 8.5, and wherein optionally the composition has a reserve alkalinity to pH 7.0 of less than 3.0gNaOH/100g.
7. A composition according to any preceding claim, wherein the composition comprises:
 - (a) from 0wt% to 6wt% sodium bicarbonate;
 - (b) from 0wt% to 4wt% sodium carbonate;
 - (c) from 0wt% to 4wt% sodium silicate; and
 - (d) from 0wt% to 4wt% phosphate builder,and optionally wherein the composition is substantially free of phosphate builder, and optionally wherein the composition is substantially free of sodium carbonate, and optionally wherein the composition is substantially free of sodium bicarbonate, and optionally wherein the composition is substantially free of sodium silicate.
8. A composition according to any preceding claim, wherein the composition comprises the combination of a lipase enzyme and soil release polymer.
9. A composition according to any preceding claim wherein the composition comprises:
 - (a) alkyl benzene sulphonate, wherein the alkyl benzene sulphonate comprises at least 25wt% of the combined total of 2-phenyl isomer and 3-phenyl isomer; and/or
 - (b) alkyl amine oxide.
10. A composition according to any preceding claim, wherein the composition comprises:
 - (a) from 0.5wt% to 8wt% carboxylate co-polymer, wherein the carboxylate co-polymer comprises:
 - (i) from 50 to less than 98 wt% structural units derived from one or more monomers comprising carboxyl groups;
 - (ii) from 1 to less than 49 wt% structural units derived from one or more monomers comprising sulfonate moieties; and
 - (iii) from 1 to 49 wt% structural units derived from one or more types of monomers selected from ether bond-containing monomers represented by formulas (I) and (II):

formula (I):



wherein in formula (I), R₀ represents a hydrogen atom or CH₃ group, R represents a CH₂ group, CH₂CH₂ group or single bond, X represents a number 0-5 provided X represents a number 1-5 when R is a single bond, and R₁ is a hydrogen atom or C₁ to C₂₀ organic group;


formula (II)



wherein in formula (II), R₀ represents a hydrogen atom or CH₃ group, R represents a CH₂ group, CH₂CH₂ group or single bond, X represents a number 0-5, and R₁ is a hydrogen atom or C₁ to C₂₀ organic group; and/or

- (b) polyethylene glycol polymer, wherein the polyethylene glycol polymer comprises a polyethylene glycol backbone with grafted polyvinyl acetate side chains; and/or
- (c) polyester soil release polymer having the structure:

52






wherein n is from 1 to 10; m is from 1 to 15 ;

X is H or SO₃Me;

wherein Me is H, Na⁺, Li⁺, K⁺, Mg²⁺, Ca²⁺, Al³⁺, ammonium, mono-, di-, tri-, or tetra-alkylammonium; wherein the alkyl groups are C₁-C₁₈ alkyl or C₂-C₁₀ hydroxyalkyl, or any mixture thereof;

R1 are independently selected from H or C₁-C₁₈ n- or iso-alkyl; and/or

(d) polyester soil release polymer consisting of structure units (1) to (3):



wherein:

a, b and c are from 1 to 10;

x, y is from 1 to 10;

z is from 0.1 to 10;


Me is H, Na⁺, Li⁺, K⁺, Mg²⁺, Ca²⁺, Al³⁺, ammonium, mono-, di-, tri-, or tetra-alkylammonium wherein the alkyl groups are C₁-C₁₈ alkyl or C₂-C₁₀ hydroxyalkyl, or any mixture thereof;

R₁, are independently selected from H or C₁-C₁₈ n- or iso-alkyl;

R₂ is a linear or branched C₁-C₁₈ alkyl, or a linear or branched C₂-C₃₀ alkenyl, or a cycloalkyl group with 5 to 9 carbon atoms, or a C₆-C₃₀ aryl group, or a C₆-C₃₀ arylalkyl group; and/or

- (e) carboxymethyl cellulose having a degree of substitution greater than 0.65 and a degree of blockiness greater than 0.45; and/or
- (f) alkoxyated polyalkyleneimine, wherein said alkoxyated polyalkyleneimine has a polyalkyleneimine core with one or more side chains bonded to at least one nitrogen atom in the polyalkyleneimine core, wherein said alkoxyated polyalkyleneimine has an empirical formula (I) of (PEI)_a-(EO)_b-R₁, wherein a is the average number-average molecular weight (MW_{PEI}) of the polyalkyleneimine core of the alkoxyated polyalkyleneimine and is in the range of from 100 to 100,000 Daltons, wherein b is the average degree of ethoxylation in said one or more side chains of the alkoxyated polyalkyleneimine and is in the range of from 5 to 40, and wherein R₁ is independently selected from the group consisting of hydrogen, C₁-C₄ alkyls, and combinations thereof; and/or
- (g) alkoxyated polyalkyleneimine, wherein said alkoxyated polyalkyleneimine has a polyalkyleneimine core with one or more side chains bonded to at least one nitrogen atom in the polyalkyleneimine core, wherein the alkoxyated polyalkyleneimine has an empirical formula (II) of (PEI)_o-(EO)_m(PO)_n-R₂ or (PEI)_o-(PO)_n(EO)_m-R₂, wherein o is the average number-average molecular weight (MW_{PEI}) of the polyalkyleneimine core of the alkoxyated polyalkyleneimine and is in the range of from 100 to 100,000 Daltons, wherein m is the average degree of ethoxylation in said one or more side chains of the alkoxyated polyalkyleneimine which ranges from 10 to 50, wherein n is the average degree of propoxylation in said one or more side chains of the alkoxyated polyalkyleneimine which ranges from 1 to 50, and wherein R₂ is independently selected from the group consisting of hydrogen, C₁-C₄ alkyls, and combinations thereof; and/or
- (h) the combination of a non-ionic soil release polymer and an anionic soil release polymer.

11. A composition according to any preceding claim, wherein the composition is substantially free of pre-formed peracid.
12. A composition according to any preceding claim, wherein the composition comprises:
 - (a) from 1wt% to 20wt% sodium percarbonate;
 - (b) from 0.5wt% to 5wt% bleach activator; and
 - (c) from 0.5wt% to 5wt% chelant.
13. A composition according to any preceding claim, wherein the composition comprises from 0.5wt% to 5wt% sodium tetraacetythylenediamine.
14. A composition according to any preceding claim, wherein the composition comprises:
 - (a) from 0.5wt% to 5wt% tri sodium salt of methylglycine diacetic acid (MGDA); and/or
 - (b) from 0.5wt% to 5wt% ethylenediamine disuccinic acid (EDDS).
15. A composition according to any preceding claim, wherein the composition comprises 4,4'-bis-(triazinylamino)-stilbene-2,2'-disulfonic acid brightener and/or 4,4'-distyryl biphenyl brightener.
16. A composition according to any preceding claim, wherein the composition comprises from 0.5wt% to 4wt% disodium 4,5-dihydroxy-1,3-benzenedisulfonate.
17. A composition according to any preceding claim, wherein the composition comprises acyl hydrazone bleach catalyst, wherein the acyl hydrazone bleach catalyst has the formula I:



wherein, R¹ is selected from the groups comprising CF₃, C₁₋₂₈ alkyl, C₂₋₂₈ alkenyl, C₂₋₂₂ alkynyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkenyl, phenyl, naphthyl, C₇₋₉ aralkyl, C₃₋

20 heteroalkyl, C₃₋₁₂ cycloheteroalkyl or a mixture thereof;


R² and R³ are independently selected from the group comprising hydrogen, substituted C₁₋₂₈ alkyl, C₂₋₂₈ alkenyl, C₂₋₂₂ alkynyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkenyl, C₇₋₉ aralkyl, C₃₋₂₈ heteroalkyl, C₃₋₁₂ cycloheteroalkyl, C₅₋₁₆ heteroaralkyl, phenyl, naphthyl, heteroaryl or a mixture thereof;

or R² and R³ are linked to form a substituted 5-, 6-, 7-, 8- or 9-membered ring that optionally comprises heteroatoms;

and R⁴ is selected from the groups comprising hydrogen, C₁₋₂₈ alkyl, C₂₋₂₈ alkenyl, C₂₋₂₂ alkynyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkenyl, C₇₋₉ aralkyl, C₃₋₂₀ heteroalkyl, C₃₋₁₂ cycloheteroalkyl, C₅₋₁₆ heteroaralkyl, substituted phenyl, naphthyl, heteroaryl or a mixture thereof.

18. A composition according to any preceding claim, wherein the composition comprises:

(a) hueing agent having the following structure:



wherein the index values x and y are independently selected from 1 to 10; and/or


(b) hueing agent selected from Acid Violet 50, Direct Violet 9, 66 and 99, Solvent Violet 13 and any combination thereof.

19. A composition according to any preceding claim, wherein the composition comprises an enzyme selected from:

- (a) protease having at least 90% identity to the amino acid sequence of *Bacillus amyloliquefaciens* as shown in SEQ ID NO:9;
- (b) protease having at least 90% identity to the amino acid sequence of *Bacillus amyloliquefaciens BPN'* as shown in SEQ ID NO:10, and which comprises one or more mutations selected from group consisting of V4I, S9R, A15T, S24G, S33T, S53G, V68A, N76D, S78N, S101M/N, Y167F, and Y217Q;
- (c) protease having at least 90% identity to the amino acid sequence of *Bacillus thermoproteolyticus* as shown in SEQ ID NO:11;
- (d) protease having at least 90% identity to the amino acid sequence of *Bacillus lentus* as shown in SEQ IS NO:12, and which comprises one or mutations selected from the group consisting of S3T, V4I, A194P, V199M, V205I, and L217D;

- (e) protease having at least 90% identity to the amino acid sequence of *Bacillus sp. TY145* as shown in SEQ ID NO:13;
- (f) protease having at least 90% identity to the amino acid sequence of *Bacillus sp. KSM-KP43* as shown in SEQ ID NO:14;
- (g) variant of the wild-type amylase from *Bacillus sp.* which has at least 90% identity for amino acid sequence SEQ ID NO:5, and which comprises one or more mutations at positions N195, G477, G304, W140, W189, D134, V206, Y243, E260, F262, W284, W347, W439, W469 and/or G476, and optionally which comprises the deletions of D183* and/or G184*;
- (h) variant of the wild-type amylase from *Bacillus sp.* which has at least 90% identity for amino acid sequence SEQ ID NO:6, and which comprises one or more mutations at positions 9, 26, 30, 33, 82, 37, 106, 118, 128, 133, 149, 150, 160, 178, 182, 186, 193, 195, 202, 214, 231, 256, 257, 258, 269, 270, 272, 283, 295, 296, 298, 299, 303, 304, 305, 311, 314, 315, 318, 319, 320, 323, 339, 345, 361, 378, 383, 419, 421, 437, 441, 444, 445, 446, 447, 450, 458, 461, 471, 482 and/or 484, preferably that also contain the deletions of D183* and G184*;
- (i) variant of the wild-type amylase from *Bacillus sp. KSM-K38* which has at least 90% identity for amino acid sequence SEQ ID NO:7;
- (j) variant of the wild-type amylase from *Cytophaga sp.* which has at least 60% identity for amino acid sequence SEQ ID NO:8;
- (k) a variant of the wild-type lipase from *Thermomyces lanuginosus* which has at least 90% identity for amino acid sequence SEQ ID NO:1;
- (l) variant of the wild-type lipase from *Thermomyces lanuginosus* which has at least 90% identity for amino acid sequence SEQ ID NO:1, and which comprises T231R and/or N233R mutations;
- (m) variant of the wild-type lipase from *Thermomyces lanuginosus* which has at least 90% identity for amino acid sequence SEQ ID NO:1, and which comprises G91A, D96G, G225R, T231R and/or N233R mutations;
- (n) cellulase that is a wild-type or variant of a microbially-derived endoglucanase endogenous to *Bacillus sp.* exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4) which has at least 90% identity to the amino acid sequence SEQ ID NO:2;

- (o) cellulase that is a wild-type or variant of a microbially-derived endoglucanase endogenous to *Paenibacillus polymyxa* exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4) which has at least 90% identity to amino acid sequence SEQ ID NO:3;
 - (p) cellulase that is a hybrid fusion endoglucanase comprising a Glycosyl Hydrolase Family 45 catalytic domain that is a wild-type or variant of a microbially-derived endoglucanase endogenous to *Melanocarpus albomyces*, and a carbohydrate binding module that is a wild-type or variant of a carbohydrate binding module endogenous to *Trichoderma reesei*, and which has at least 90% identity to amino acid sequence SEQ ID NO:4;
 - (q) an enzyme selected from mannanase, pectate lyase, laccase, polyesterase, galactanase, acyltransferase, and any combination thereof; and
 - (r) any combination thereof.
20. A composition according to any preceding claim, wherein the composition comprises a perfume, wherein the perfume comprises from 60wt% to 85wt% ester perfume raw materials having the structure:



- wherein R1 and R2 are independently selected from C1 to C30 linear or branched, cyclic or non-cyclic, aromatic or non-aromatic, saturated or un-saturated, substituted or unsubstituted alkyl,
- and optionally wherein the composition comprises alkyl ethoxylated sulphate having an average degree of ethoxylation of from 0.5 to 2.0.
21. A composition according to any preceding claim, wherein the composition comprises polyvinyl N oxide polymer.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/054818

A. CLASSIFICATION OF SUBJECT MATTER		
INV.	C11D1/02 C11D3/40	C11D3/20 C11D11/00 C11D17/06 C11D3/42
ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) C11D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	WO 03/038028 A2 (HENKEL KGAA) 8 May 2003 (2003-05-08) page 7, paragraph 1 - pages 10-11; claims 1,3-10,12-20; example 1; table 1 page 1, paragraph 4 - page 2, paragraph 3 pages 8-9 - page 12, paragraph 1 page 8, paragraph 2 - page 9, paragraph 2 page 9, paragraph 3 page 11, paragraph 3 page 12, paragraph 2 - page 13, paragraph 1 page 14, paragraph 3 page 32, paragraph 4 - page 33, paragraph 1 page 33, paragraph 3 page 22, paragraph 2 - page 23, paragraph 1 page 32, paragraph 2 -/--	1-4,6,7, 9-15, 19-21 16,17
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 15 December 2017		Date of mailing of the international search report 07/02/2018
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Kanbier, Titia

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/054818

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	page 23, paragraph 4 - page 30, paragraph 3 page 16, paragraph 1-2 - page 18, paragraph 2 page 19, paragraph 3 page 23, paragraph 2 page 21, paragraph 5 page 27, paragraph 5 - page 28, paragraph 3 -----	
X	WO 95/12658 A1 (PROCTER & GAMBLE) 11 May 1995 (1995-05-11) pages 5-6 - page 20, paragraph 1; example 1 page 30, lines 12-16,3-5, paragraphs 3,4 - page 31, paragraph 2; claims 1,6-9 page 3, paragraph 5 page 11, lines 1-4, paragraph 4 - page 17, lines 1-3, paragraph 6 -----	1-14,16,19
X	WO 91/17232 A1 (PROCTER & GAMBLE) 14 November 1991 (1991-11-14) page 1, lines 7-16 page 3, lines 28-32 page 4, lines 13-15 - page 6, lines 4-5 page 7, lines 15-16 page 11, lines 16-17,29-30 page 12, line 15 - page 13, line 25 page 14, lines 9-14,19-22 page 15, line 8 - page 16, line 16; claims 1,2 page 17, lines 25-28; claims 7-10; examples I,II,III -----	1-4,6-12,15,19,20
X	WO 2013/036662 A1 (SUN PRODUCTS CORP) 14 March 2013 (2013-03-14)	1-3,5,18
A	pages 28-30; claims 1,2,6-8; examples 4,6 page 2, lines 28-29 page 7, lines 4-9 -----	15
A	WO 2015/169851 A1 (BASF SE; HENKEL AG & CO KGAA) 12 November 2015 (2015-11-12) page 1, lines 21-43; claims 1,12,13 page 5, line 31 - page 6, line 17 page 8, line 23 - page 9, line 22 page 10, lines 5-23 page 10, line 40 - page 11, line 9 page 11, lines 27-39; example C; table 1 ----- -/--	1,12,13,15,17

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/054818

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	WO 2013/184981 A2 (PROCTER & GAMBLE) 12 December 2013 (2013-12-12) paragraphs [0034] - [0040], [0079]; claims 1,6,18-23 paragraphs [0063] - [0066], [0075] - [0077], [0079], [0080], [0082] - [0085] pages 31-33, paragraphs 89-92,95,97-99; example 2 paragraphs [0101] - [0103], [0110] - [0114], [0116] paragraphs [0111], [0112], [0114], [0116], [0117] paragraphs [0121] - [0125] paragraphs [0126] - [155;159;161] -----	1,3-5, 7-21 2
X A	WO 2005/095572 A1 (UNILEVER NV) 13 October 2005 (2005-10-13) page 29, lines 16-23,33 - page 30, line 1 page 28, line 22 - page 29, line 5 page 2, lines 28-29; claims 1,2,6-8; examples 4,6 -----	1,5,15 10
X A	US 2011/082066 A1 (WRUBBEL N ET AL) 7 April 2011 (2011-04-07) paragraphs [0003], [0008] - [10;26], [0099], [0100]; table 2 paragraphs [0041], [0095] - [0097], [0104] paragraphs [0130] - [0132], [0142], [0151], [0172] paragraphs [0133], [0134] - [0138], [0142], [0143], [0146] - [0150], [0153], [0156] - [0159], [0169], [0170] -----	1,4,5,9, 10,12, 13,18-21 14,16,17
A	WO 2013/043855 A2 (PROCTER & GAMBLE) 28 March 2013 (2013-03-28) page 26, paragraph 2 - page 27, paragraph 2; claims 1,3,5,6,8,12; examples 1,2 page 31, paragraph 1 page 32, lines 1-3, paragraph 2 page 22, paragraph 3 - page 24, paragraph 3 -----	13-16
X A	WO 2012/054835 A1 (PROCTER & GAMBLE) 26 April 2012 (2012-04-26) page 52, lines 10-22; claims 1,9,12,13; table 5B page 41, line 12 - page 42, line 10 page 59, lines 5-15 page 42, line 12 - page 45, line 22 page 46, line 3 - page 51, line 27 page 36, line 20 - page 37, line 14 page 28, line 11 - page 31, line 13 -----	1-4,6,7, 9-15, 19-21 5,8,16
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/054818

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2011/011799 A2 (PROCTER & GAMBLE) 27 January 2011 (2011-01-27)	18
A	page 59, paragraph 2; claims 1,6-8 page 56, paragraph 2; claim 12 page 61, paragraph 3 - page 65, paragraph 1 page 67, paragraph 1 - page 68, paragraph 1 page 73, paragraph 4 - page 74, paragraph 4 page 76, paragraph 3 - page 77, paragraph 1	1
A	----- JP 2009 013291 A (SUMITOMO CHEMICAL CO) 22 January 2009 (2009-01-22) paragraphs [0002], [0038] - [0040], [0075], [0087], [0091] - [0093]; claims 1,3; examples 2-4; table 1 -----	1,18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/054818

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 03038028	A2	08-05-2003	AT 368726 T 15-08-2007 DE 10153551 A1 22-05-2003 EP 1440141 A2 28-07-2004 ES 2289163 T3 01-02-2008 WO 03038028 A2 08-05-2003
WO 9512658	A1	11-05-1995	AU 8125494 A 23-05-1995 EP 0651053 A1 03-05-1995 WO 9512658 A1 11-05-1995
WO 9117232	A1	14-11-1991	AR 244332 A1 29-10-1993 AT 141639 T 15-09-1996 AU 7856791 A 27-11-1991 CA 2081324 A1 09-11-1991 CN 1057292 A 25-12-1991 DE 69121460 D1 26-09-1996 EP 0456315 A2 13-11-1991 ES 2090229 T3 16-10-1996 JP H05507744 A 04-11-1993 MX 172329 B 13-12-1993 TW 200525 B 21-02-1993 WO 9117232 A1 14-11-1991
WO 2013036662	A1	14-03-2013	CA 2847546 A1 14-03-2013 CN 103946360 A 23-07-2014 EP 2751248 A1 09-07-2014 JP 2014529693 A 13-11-2014 KR 20140068141 A 05-06-2014 US 2013095717 A1 18-04-2013 US 2016160156 A1 09-06-2016 US 2016160157 A1 09-06-2016 WO 2013036662 A1 14-03-2013
WO 2015169851	A1	12-11-2015	EP 3140382 A1 15-03-2017 KR 20170003922 A 10-01-2017 US 2017175048 A1 22-06-2017 WO 2015169851 A1 12-11-2015
WO 2013184981	A2	12-12-2013	CN 104471050 A 25-03-2015 CN 104769094 A 08-07-2015 EP 2859082 A2 15-04-2015 EP 2859083 A2 15-04-2015 JP 6188794 B2 30-08-2017 JP 2015518920 A 06-07-2015 JP 2015524015 A 20-08-2015 RU 2014144338 A 27-05-2016 RU 2014151212 A 10-08-2016 US 2013345108 A1 26-12-2013 US 2015126428 A1 07-05-2015 WO 2013184981 A2 12-12-2013 WO 2013184987 A2 12-12-2013
WO 2005095572	A1	13-10-2005	AR 047984 A1 15-03-2006 AT 393207 T 15-05-2008 BR PI0507264 A 26-06-2007 CN 101076583 A 21-11-2007 DE 602005006244 T2 25-06-2009 EP 1720968 A1 15-11-2006

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/054818

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		ES 2306092 T3	01-11-2008
		US 2005197275 A1	08-09-2005
		WO 2005095572 A1	13-10-2005

US 2011082066 A1	07-04-2011	EP 2291505 A1	09-03-2011
		ES 2398026 T3	13-03-2013
		US 2011082066 A1	07-04-2011
		US 2017081612 A1	23-03-2017
		WO 2010000636 A1	07-01-2010

WO 2013043855 A2	28-03-2013	AR 088757 A1	02-07-2014
		US 2013072416 A1	21-03-2013
		WO 2013043855 A2	28-03-2013

WO 2012054835 A1	26-04-2012	AR 083503 A1	27-02-2013
		BR 112013009791 A2	09-08-2016
		BR 112013009792 A2	26-07-2016
		BR 112013009793 A2	26-07-2016
		BR 112013009794 A2	26-07-2016
		BR 112013009798 A2	26-07-2016
		CA 2815479 A1	26-04-2012
		CN 103168097 A	19-06-2013
		CN 103180424 A	26-06-2013
		CN 103180425 A	26-06-2013
		CN 103201370 A	10-07-2013
		CN 103339245 A	02-10-2013
		CN 105695160 A	22-06-2016
		EP 2630225 A1	28-08-2013
		EP 2630226 A1	28-08-2013
		EP 2630227 A1	28-08-2013
		EP 2630228 A1	28-08-2013
		EP 2630229 A1	28-08-2013
		JP 5833130 B2	16-12-2015
		JP 6049679 B2	21-12-2016
		JP 2013544916 A	19-12-2013
		JP 2015120906 A	02-07-2015
		US 2012101018 A1	26-04-2012
		US 2012122750 A1	17-05-2012
		US 2012122751 A1	17-05-2012
		US 2012129751 A1	24-05-2012
		US 2012129752 A1	24-05-2012
		US 2012129753 A1	24-05-2012
		US 2015291918 A1	15-10-2015
		WO 2012054820 A1	26-04-2012
		WO 2012054821 A1	26-04-2012
		WO 2012054823 A1	26-04-2012
		WO 2012054827 A1	26-04-2012
		WO 2012054835 A1	26-04-2012
		ZA 201302890 B	29-10-2014

WO 2011011799 A2	27-01-2011	BR 112013011851 A2	16-08-2016
		CA 2817718 A1	27-01-2011
		CN 103210073 A	17-07-2013
		EP 2638142 A2	18-09-2013
		JP 5833133 B2	16-12-2015
		JP 2014501802 A	23-01-2014
		WO 2011011799 A2	27-01-2011

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/054818

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 2009013291 A	22-01-2009	CN 101337942 A	07-01-2009
		JP 2009013291 A	22-01-2009
		KR 20090004759 A	12-01-2009
		TW 200911928 A	16-03-2009
