



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2312208 C 2011/01/25

(11)(21) **2 312 208**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1998/12/04
(87) Date publication PCT/PCT Publication Date: 1999/06/17
(45) Date de délivrance/Issue Date: 2011/01/25
(85) Entrée phase nationale/National Entry: 2000/06/05
(86) N° demande PCT/PCT Application No.: US 1998/025828
(87) N° publication PCT/PCT Publication No.: 1999/029888
(30) Priorité/Priority: 1997/12/05 (US08/986,016)

(51) Cl.Int./Int.Cl. *C12P 21/08* (2006.01),
C07K 16/28 (2006.01), *C07K 16/46* (2006.01)
(72) Inventeurs/Inventors:
BARBAS, CARLOS F., III, US;
RADER, CHRISTOPH, US
(73) Propriétaire/Owner:
THE SCRIPPS RESEARCH INSTITUTE, US
(74) Agent: CASSAN MACLEAN

(54) Titre : HUMANISATION D'ANTICORPS MURINS
(54) Title: HUMANIZATION OF MURINE ANTIBODY

(57) **Abrégé/Abstract:**

A humanized murine antibody is provided. The amino acid sequences of a light chain complementarity determining region from a mouse antibody are grafted onto a human light chain, and a heavy chain complementarity determining region from a mouse antibody are grafted onto a human antibody heavy chain to produce libraries from which a humanized murine antibody having the desired specificity is selected.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12P 21/08	A1	(11) International Publication Number: WO 99/29888 (43) International Publication Date: 17 June 1999 (17.06.99)
(21) International Application Number: PCT/US98/25828 (22) International Filing Date: 4 December 1998 (04.12.98) (30) Priority Data: 08/986,016 5 December 1997 (05.12.97) US (71) Applicant: THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10550 North Torrey Pines Road, Mail Drop TPC-8, La Jolla, CA 92037 (US). (72) Inventors: BARBAS, Carlos, F., III; 7081 Weller Street, San Diego, CA 92122 (US). RADER, Christoph; 3623 Marlesta Drive, San Diego, CA 92111 (US). (74) Agents: CEPURITIS, Talivaldis; Olson & Hierl, Ltd., 36th floor, 20 North Wacker Drive, Chicago, IL 60606 (US) et al.		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>With an indication in relation to deposited biological</i> <i>material furnished under Rule 13bis separately from the</i> <i>description.</i>
(54) Title: HUMANIZATION OF MURINE ANTIBODY (57) Abstract A humanized murine antibody is provided. The amino acid sequences of a light chain complementarity determining region from a mouse antibody are grafted onto a human light chain, and a heavy chain complementarity determining region from a mouse antibody are grafted onto a human antibody heavy chain to produce libraries from which a humanized murine antibody having the desired specificity is selected.		

HUMANIZATION OF MURINE ANTIBODY

Technical Field

This invention relates to humanization of murine antibodies.

5 Background of the Invention

Antibodies typically comprise two heavy chains linked together by disulfide bonds and two light chains linked to a respective heavy chain by a disulfide bond. Beginning at one end of each heavy chain there is a variable domain followed by several constant domains. Similarly, each light chain has a variable domain at one
10 end, but only a single constant domain at its other end. There are two types of light chain, which are termed lambda (λ) and kappa (κ) chains. No functional difference has been found between antibodies having λ or κ light chains. The ratio of the two types of light chain varies from species to species, however. In mice, the κ : λ ratio is 20:1, whereas in humans it is 2:1.

15 The variable domains of the light and heavy chains are aligned, as are the constant domain of the light chain and the first constant domain of the heavy chain. The constant domains in the light and heavy chains are not involved directly in binding the antibody to antigen.

It is the variable domains that form the antigen binding site of
20 antibodies. The general structure of each light and heavy chain domain comprises a framework of four regions, whose sequences are relatively conserved, connected by three complementarity determining regions (CDRs). The four framework regions employ a beta-sheet conformation and the CDRs form loops connecting, and in some cases forming part of, the beta-sheet structure. The CDRs are held in close
25 proximity by the framework regions and, with the CDRs from the other domain, contribute to the formation of the antigen binding site.

While cell surface antigens of tumor cells are the traditional targets for antibody-guided cancer therapy, one of the major limitations for the therapy of solid tumors is the low accessibility of tumor antigens to antibodies circulating in the
30 blood stream. The dense packing of tumor cells and the elevated interstitial pressure in the tumor core present formidable physical barriers.

A solution to the problem of poor penetration of antibodies into solid tumors would be to attack the endothelial cells lining the blood vessels of the

tumor rather than the tumor cells themselves. While it may be difficult to target the mature tumor vasculature specifically, i.e., without destroying healthy tissue, promising strategies aim at the inhibition of neovascularization.

Neovascularization, also termed angiogenesis, is induced by
5 cytokines that are secreted from tumor cells and depends on vascular cell migration and invasion, processes regulated by cell adhesion molecules (CAM) and protease. These molecules are currently considered potential targets for angiogenic inhibitors. In this regard, the vascular integrin $\alpha_v\beta_3$ has recently been identified as a marker of angiogenic blood vessels. See Brooks, P.C., et al. (1994), REQUIREMENT OF
10 INTEGRIN $\alpha_v\beta_3$ FOR ANGIOGENESIS, *Science* **264**, 569-571. Moreover, it was shown that the mouse monoclonal antibody (Mab) LM609 directed to integrin $\alpha_v\beta_3$ was able to suppress angiogenesis, indicating that integrin $\alpha_v\beta_3$ has a critical role in angiogenesis.

It has been further demonstrated that LM609 selectively promotes
15 apoptosis of vascular cells that have been stimulated to undergo angiogenesis. See Brooks, P.C., et al. (1994), INTEGRIN $\alpha_v\beta_3$ ANTAGONISTS PROMOTE TUMOR REGRESSION BY INDUCING APOPTOSIS OF ANGIOGENIC BLOOD VESSELS, *Cell* **79**, 1157-1164. These findings suggest that integrin $\alpha_v\beta_3$ may be a target and LM609 a tool for cancer diagnosis and therapy.

Indeed, LM609 not only prevented the growth of histologically
20 distinct human tumors implanted on the chorioallantoic membranes of chicken embryos, but also induced their regression. See, *Cell* **79**, 1157-1164. Using a more clinically relevant model of tumor growth, it was found that LM609 blocked human breast cancer growth in a SCID mouse/human chimeric model. Importantly, not only
25 did LM609 block tumor growth, but it also inhibited metastasis of the breast carcinomas examined. See Brooks, et al. (1995) ANTI-INTEGRIN $\alpha_v\beta_3$ BLOCKS HUMAN BREAST CANCER GROWTH AND ANGIOGENESIS IN HUMAN SKIN, *J. Clin. Invest.* **96**, 1815-1822.

The Brooks et al. results are consistent with previous studies that
30 have suggested that angiogenesis contributes to the metastatic spread of breast tumor cells. See Weidner, N., et al. (1991) TUMOR ANGIOGENESIS AND METASTASIS: CORRELATION IN INVASIVE BREAST CARCINOMA, *N.Engl. J. Med.* **324**, 1-8; and Weidner, N., et al. (1992) TUMOR ANGIOGENESIS: A NEW SIGNIFICANT AND

INDEPENDENT PROGNOSTIC INDICATOR IN EARLY-STAGE BREAST CARCINOMA, *J.Natl. Cancer Inst.* **84**, 1875-1887.

Within the last few years evidence has been presented that two cytokine-dependent pathways of angiogenesis exist and that these are defined by their dependency on distinct vascular integrins. See Friedlander, M., et al. (1995) DEFINITION OF TWO ANGIOGENIC PATHWAYS BY DISTINCT AV INTEGRINS, *Science* **270**, 1500-1502. The results of the Friedlander et al. studies show that anti- $\alpha_v\beta_3$ antibody LM609 blocked angiogenesis in response to bFGF and $\text{TNF}\alpha$, yet have little effect on angiogenesis induced by VEGF, $\text{TGF}\alpha$, or phorbol ester PMA. In contrast, the anti- $\alpha_v\beta_5$ antibody P1F6 blocks angiogenesis induced by VEGF, $\text{TGF}\alpha$, and phorbol ester PMA, while having minimal effects on that induced by bFGF or $\text{TNF}\alpha$.

It is conceivable, thus, that tumors showing less susceptibility to anti- $\alpha_v\beta_3$ antibodies might secrete cytokines that promote angiogenesis in an $\alpha_v\beta_5$ -dependent manner. Taken together, both anti- $\alpha_v\beta_3$ and anti- $\alpha_v\beta_5$ antibodies are promising tools for diagnosis and therapy of cancer.

Mouse monoclonal antibodies such as LM609, however, are highly immunogenic in humans, thus limiting their potential use for cancer therapy, especially when repeated administration is necessary. To reduce the immunogenicity of mouse monoclonal antibodies, chimeric monoclonal antibodies were generated, with the variable Ig domains of a mouse monoclonal antibody being fused to human constant Ig domains. See Morrison, S.L., et al. (1984) CHIMERIC HUMAN ANTIBODY MOLECULES; MOUSE ANTIGEN-BINDING DOMAINS WITH HUMAN CONSTANT REGION DOMAINS, *Proc. Natl. Acad. Sci. USA* **81**, 6841-6855; and, Boulianne, G.L., et al. (1984) PRODUCTION OF A FUNCTIONAL CHIMAERIC MOUSE/HUMAN ANTIBODY, *Nature* **312**, 643-646. This process is commonly referred to as "humanization" of an antibody.

In general, the chimeric monoclonal antibodies retain the binding specificity of the mouse monoclonal antibody and exhibit improved interactions with human effector cells. This results in an improved antibody-dependent cellular cytotoxicity which is presumed to be one of the ways of eliminating tumor cells using monoclonal antibodies. See Morrison, S.L. (1992) *IN VITRO* ANTIBODIES: STRATEGIES FOR PRODUCTION AND APPLICATION, *Ann. Rev. Immunol.* **10**, 239-265. Though some chimeric monoclonal antibodies have proved less immunogenic in humans, the

mouse variable Ig domains can still lead to a significant human anti-mouse response. See Bruggemann, M., et al. (1989) THE IMMUNOGENICITY OF CHIMERIC ANTIBODIES, *J.Exp. Med.* 170, 2153-2157. Therefore, for therapeutic purposes it may be necessary to fully humanize a murine monoclonal antibody by altering both the
5 variable and the constant Ig domains.

Full humanization is feasible by introducing the six CDRs from the mouse heavy and light chain variable Ig domains into the appropriate framework regions of human variable Ig domains. This CDR grafting technique (Riechmann, L., et al. (1988) RESHAPING HUMAN ANTIBODIES FOR THERAPY, *Nature* 332, 323)
10 takes advantage of the conserved structure of the variable Ig domains, with the four framework regions (FR1-FR4) serving as a scaffold to support the CDR loops which are the primary contacts with antigen. U.S. Patent No. 5,502,167 to Waldmann, et al. discloses a "humanised antibody" having the CDR loops LCDR1 through LCDR3 and HCDR1 through HCDR3 from YTH 655(5)6, a rat IgG2b monoclonal antibody,
15 grafted onto a human T cell antibody.

A drawback, however, of the CDR grafting technique is the fact that amino acids of the framework regions can contribute to antigen binding, as well as amino acids of the CDR loops can influence the association of the two variable Ig domains. To maintain the affinity of the humanized monoclonal antibody, the CDR
20 grafting technique relies on the proper choice of the human framework regions and site-directed mutagenesis of single amino acids aided by computer modeling of the antigen binding site (e.g., Co, M.S., et al. (1994) A HUMANIZED ANTIBODY SPECIFIC FOR THE PLATELET INTEGRIN gpIIb/IIIa, *J.Immunol.* 152, 2968-2976). A number of successful humanizations of mouse monoclonal antibodies by rational design have
25 been reported. Among them are several monoclonal antibodies that are directed to human integrins and have potential clinical application. See, *J.Immunol.* 152, 2968-2976; Hsiao, K.C., et al. (1994) HUMANIZATION OF 60.3, AN ANTI-CD18 ANTIBODY; IMPORTANCE OF THE L2 LOOP, *Protein Eng.* 7, 815-822; and, Poul, M.A., et al. (1995) INHIBITION OF T CELL ACTIVATION WITH A HUMANIZED ANTI-BETA 1
30 INTEGRIN CHAIN mAb, *Mol. Immunol.* 32, 101-116.

Human immunoglobulin transgenic mice provide a promising alternative to the humanization of mouse monoclonal antibodies. See, e.g., Fishwild, D.M., et al. (1996) HIGH-AVIDITY HUMAN IgG κ MONOCLONAL ANTIBODIES FROM A

NOVEL STRAIN OF MINILOCUS TRANSGENIC MICE, *Nature Biotechnology* 14, 845-851. In response to immunization, these mice express human monoclonal antibodies, which can be accessed by conventional hybridoma technology.

Rational design strategies in protein engineering have been
5 challenged by *in vitro* selection strategies that are mainly based on phage display libraries. See Clackson, T., and Wells, J.A. (1994) *IN VITRO SELECTION FROM PROTEIN AND PEPTIDE LIBRARIES*, *TIBTECH* 12, 173-184. In particular, *in vitro* selection and evolution of antibodies derived from phage display libraries has become a powerful tool. See Burton, D.R., and Barbas III, C.F. (1994) *HUMAN ANTIBODIES*
10 *FROM COMBINATORIAL LIBRARIES*, *Adv. Immunol.* 57, 191-280; and, Winter, G., et al. (1994) *MAKING ANTIBODIES BY PHAGE DISPLAY TECHNOLOGY*, *Annu. Rev. Immunol.* 12, 433-455.

The development of technologies for making repertoires of human antibody genes, and the display of the encoded antibody fragments on the surface of
15 filamentous bacteriophage, has provided a means for making human antibodies directly. The antibodies produced by phage technology are produced as antigen binding fragments—usually Fv or Fab fragments—in bacteria and thus lack effector functions. Effector functions can be introduced by one of two strategies: The fragments can be engineered either into complete antibodies for expression in
20 mammalian cells, or into bispecific antibody fragments with a second binding site capable of triggering an effector function.

Typically, the Fd fragment (V_H - C_{H1}) and light chain (V_L - C_L) of antibodies are separately cloned by PCR and recombined randomly in combinatorial phage display libraries, which can then be selected for binding to a particular antigen.
25 The Fab fragments are expressed on the phage surface, i.e., physically linked to the genes that encode them. Thus, selection of Fab by antigen binding co-selects for the Fab encoding sequences, which can be amplified subsequently. By several rounds of antigen binding and reamplification, a procedure termed panning, Fab specific for the antigen are enriched and finally isolated.

30 In 1994, an approach for the humanization of antibodies, called "guided selection", was described. Guided selection utilizes the power of the phage display technique for the humanization of mouse monoclonal antibody. See Jespers, L.S., et al. (1994) *GUIDING THE SELECTION OF HUMAN ANTIBODIES FROM PHAGE*

DISPLAY REPERTOIRES TO A SINGLE EPITOPE OF AN ANTIGEN, *Bio/Technology* 12, 899-903. For this, the Fd fragment of the mouse monoclonal antibody can be displayed in combination with a human light chain library, and the resulting hybrid Fab library may then be selected with antigen. The mouse Fd fragment thereby
5 provides a template to guide the selection.

Subsequently, the selected human light chains are combined with a human Fd fragment library. Selection of the resulting library yields entirely human Fab.

For the full humanization of murine monoclonal antibodies, the
10 present invention uses a unique combination of CDR grafting and guided selection. The anti-integrin antibody generated is useful for cancer diagnosis and therapy.

Summary of the Invention

Humanization of a mouse monoclonal antibody is achieved by a combination of guided selection and CDR grafting. The term "humanized" as used
15 herein and in the appended claims means that at least one chain of a mouse monoclonal antibody includes a region of a human monoclonal antibody.

A humanized mouse monoclonal antibody is produced by constructing a library of human antibody heavy chains or light chains in which each such chain includes a variable domain and has at least one complementarity
20 determining region (CDR) amino acid sequence which is that of a corresponding mouse heavy or light antibody chain, and then combining the library so constructed with a complementary chain from an antibody which binds a preselected antigen. In this manner, the complementary chain together with a human chain present in the constructed library forms a heavy and light chain pair in a resulting library of
25 humanized chain pairs. Thereafter a particular humanized heavy and light chain pair is selected from the humanized pair library using the aforementioned complementary chain.

In a particular embodiment, a mouse monoclonal antibody can be humanized by constructing a human light chain library in which each light chain
30 includes at least the variable domain thereof and at least one CDR amino acid sequence of a mouse light chain, and a human heavy chain library in which each such heavy chain includes at least the variable domain thereof and at least one CDR amino

acid sequence of a mouse heavy chain. The heavy chain usually is no more than about 200 amino acid residues in size.

A human light chain having a mouse CDR is selected from the constructed human light chain library using a heavy chain from an antibody which binds a preselected antigen. The constructed heavy chain library is combined with the selected human light having a mouse CDR to produce a humanized library of heavy and light chain pairs, each containing at least one mouse CDR. Thereafter, a heavy and light chain pair with mouse CDR is selected from the aforesaid humanized library using the selected human light chain with mouse CDR. The sequence of
10 aforementioned library construction is not critical.

Preferably, only the light chain complementarity determining region three (LCDR3) loop of the monoclonal antibody is grafted onto the human light chain. Similarly, it is preferable that only the HCDR3 loop be grafted onto the human heavy chain (HC) fragment. The selection of either the human light chain or
15 human heavy chain having the grafted mouse CDR is preferably made by using a chimeric mouse/human complementary chain as a template.

In CDR grafting onto a human light chain, the human light chain is cloned, then the clones are randomly recombined to form a library such as a combinatorial phage display library. The same method can be followed for grafting
20 onto the human heavy chain.

Brief Description of the Drawings

FIGURE 1 is a schematic illustration showing the sequence of the steps in the combined CDR grafting technique and guided selection technique to form the humanized Fab fragment.

FIGURES 2a and 2b show the amino acid sequences of V_L and V_H , respectively, of mouse monoclonal antibody LM609. The N-terminal two amino acids (Leu)(Glu) of V_L and (Glu)(Leu) of V_H encoded by the vector cloning sites CTCGAG (XhoI) and GAGCTC (SacI), respectively, are artificial. The CDR loops are underlined.
25

FIGURE 3a through FIGURE 3e show the amino acid sequence alignment of mouse LM609 V_H (top full line of each sequence grouping) and six selected human V_H 's (#1-6). Framework regions (FR1-3) and CDR (CDR1-2) loops are separated. Lines (-) indicate identical amino acids. Note that due to the LCDR3
30

grafting parts of FR3, entire CDR3 and entire FR4 are identical in mouse LM609 V_{κ} and the selected human V_{κ} 's. Therefore, these two sequences are not shown.

FIGURE 4 shows a comparison of three selected human fragment sequences and four unselected human fragment sequences to the original sequences of the mouse LCDR1 and LCDR2 loops.

FIGURE 5a through FIGURE 5f are line graphs which show the binding of human integrin $\alpha_v\beta_3$ on the cell surface by humanized LM609 clones 2, 4, 7, 11, 24, and control antibody, respectively. Line A indicates untransfected CS-1 hamster cells; Line B indicates human β_3 cDNA transfected CS-1 hamster cells (essentially the same line as line A in FIGURE 5f); and line C indicates human β_3 cDNA transfected CS-1 hamster cells.

FIGURE 6 is a bar graph showing the cross-reactivity of the LM609 antibody, and clones 2, 4, 7, 11, 24, and control antibody, respectively. Columns represent the mean of triplicates, with the left columns indicating binding to human integrin $\alpha_v\beta_3$, the central columns indicating binding to human integrin $\alpha_{IIb}\beta_3$, and the right columns indicating background binding. Error bars indicate standard deviations.

FIGURE 7 is a schematic illustration of a stretch of four amino acids in a light chain complementarity determining region three (LCDR3) and a heavy chain complementarity determining region three (HCDR3) being optimized.

FIGURES 8a and 8b are fragmented illustrations of the V_L amino acid sequences of a mouse antibody compared to the amino acid sequences of five versions of humanized clones represented by group letters A (clones 10, 11, and 37), and B (clones 7, 8, and 22), C (clones 4, 31, and 36), D (clones 24, 34, 35, and 40), and E (clone 2) which are combined.

FIGURES 8c through 8e are fragmented illustrations of the V_H amino acid sequences of a mouse antibody compared to the amino acid sequences of five versions of humanized clones represented by group letters A (clones 10, 11, and 37), B (clones 7, 8, and 22), C (clones 4, 31, and 36), D (clones 24, 34, 35, and 40), and E (clone 2).

Description of a Preferred Embodiment

While the present invention is susceptible to embodiments in many different forms, a preferred embodiment of the invention is described below. It

- 9 -

should be understood, however, that the present disclosure is to be considered as an exemplification of the principles of the invention and is not intended to limit the invention to the specific embodiments illustrated.

cDNA Cloning of mouse monoclonal antibody LM609

5

Beginning with a LM609 hybridoma cell line (ATCC Designation HB 9537), total RNA was prepared from 10^8 LM609 hybridoma cells using an RNA Isolation Kit (Stratagene, La Jolla, CA). Reverse transcription and polymerase chain reaction (PCR) amplification of Fd fragment and κ chain encoding sequences were performed essentially as described in "Combinatorial immunoglobulin libraries in phage 1", (*Methods* 2, 119 (1991)) by A.S. Kang, et al.

10

Fd fragment and κ chain encoding PCR products were cut with XhoI/SpeI and SacI/XbaI, respectively, and ligated sequentially into the appropriately digested phagemid vector pComb3H. The ligation products were introduced into *E.coli* strain XL1-BlueTM by electrotransformation and subsequent steps were as described in "Assembly of combinatorial antibody libraries on phage surfaces: the gene III site", (*Proc. Natl. Acad. Sci. USA* 88, 7978-7982) by C.F. Barbas III, et al., to produce phage displaying Fab on their surface. Phage were selected by panning against immobilized integrin $\alpha_v\beta_3$. After two panning rounds single clones were analyzed for LM609 Fab expression. Supernatant from IPTG-induced cultures was tested for binding to immobilized integrin $\alpha_v\beta_3$ by enzyme-linked immunosorbent assay (ELISA) using goat anti-mouse F(ab')₂ conjugated to alkaline phosphatase as a secondary antibody. The sequence of each Fd fragment and each κ chain encoding sequence of positive clones was determined by DNA sequencing.

15

20

25

Amplification of human light chain and Fd fragment sequences

Total RNA was prepared from the bone marrow of five donors (Poietic Technologies; Germantown, MD) using TRI Reagent (Molecular Research Center; Cincinnati, OH) and was further purified by lithium chloride precipitation. See Sambrook, J., et al. (1989) MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY. First-strand cDNA was synthesized using the 'SUPERScriptTM Preamplification System for First Strand

30

cDNA Synthesis' kit with oligo (dT) priming (Life Technologies; Gaithersburg, MD). The five generated first-strand cDNAs were subjected to separate PCR amplifications.

V_{κ} sequences of each of the first-strand cDNAs were amplified in eight separate reactions by combining four sense and two antisense primers (see list below). V_{λ} sequences were amplified in nine separate reactions using nine sense and one antisense primer (see list below). The V_{λ} and V_{κ} amino acid sequences, including the underlining of the CDR loops, are shown in FIGURES 2a and 2b, respectively (See also, SEQ ID NO:44 and SEQ ID NO:45, respectively).

V_H sequences (See SEQ ID NO:56) were amplified in four reactions using four sense and one antisense primer (see list below). All amplifications were performed under standard PCR conditions using Taq polymerase (Pharmacia; Uppsala, Sweden). While the sense primers hybridize to sequences that encode the N-terminal amino acids of the various V_{κ} , V_{λ} , and V_H families, the antisense primers hybridize to a sequence that encodes the C-terminal amino acids of FR3 of V_{κ} , V_{λ} , or V_H , respectively, which are highly conserved.

The list of primers used for the amplification of human antibody sequences include:

V_{κ} sense primers:

	HSCk1 - F	SEQ ID NO:3	HSCk6	SEQ ID
20	HSCk24 - F	SEQ ID NO:4	NO:14	
	HSCk3 - F	SEQ ID NO:5	HSCk70	SEQ ID
	HSCk5 - 5	SEQ ID NO:6	NO:15	

V_{κ} antisense primers:

	BKFR3UN	SEQ ID NO:7	NO:16	
25	BK2FR3UN	SEQ ID NO:8	HSCk9	SEQ ID
			NO:17	

V_{λ} sense primers:

	HSCk1a	SEQ ID NO:9	V_{λ} antisense primer:	
	HSCk1b	SEQ ID NO:10	BLFR3UN	SEQ ID
	HSCk2	SEQ ID NO:11	NO:18	
30	HSCk3	SEQ ID NO:12	V_H sense primers:	
	HSCk4	SEQ ID NO:13	HFVH1-F	SEQ ID
			NO:19	

HFVH2-F SEQ ID
 NO:20
 HFVH35-F SEQ ID NO:21
 HFVH4-F SEQ ID NO:22
 5 V_H antisense primers:
 BFR3UN SEQ ID NO:23

Construction of a chimeric mouse/human Fd fragment by fusing V_H of LM609 to human C_H1

10

The phagemid vector pComb3H containing the LM609 Fab sequences was used as a template for amplification of the sequence encoding the N-terminal FR1 through FR3 fragment of LM609's V_H by the PCR primer pair PELSEQ (SEQ ID NO:24) / BFR3UN (SEQ ID NO:25). The sense primer PELSEQ
 15 hybridizes to the pe1B leader sequence upstream of the Fd fragment encoding sequence in pComb3H. The antisense primer BFR3UN hybridizes to a sequence that encodes eight C-terminal amino acids of FR3 of V_H, which are highly conserved (SEQ ID NO:26), and differ in one amino acid from the corresponding amino acid sequence of LM609's V_H (SEQ ID NO:27).

20

By overlap extension PCR (*See* McArn Horton, R., and Readington Pease, L. (1991) RECOMBINATION AND MUTAGENESIS OF DNA SEQUENCES USING PCR IN *DIRECTED MUTAGENESIS: A PRACTICAL APPROACH*, ed. M.J. McPherson, IRL Press, Oxford, UK, pp. 217-247), the PELSEQ/BFR3UN product was fused to a PCR fragment encoding the HCDR3 (SEQ ID NO:1) of LM609 coupled to FR4 of
 25 V_H and the entire C_H1 domain of the human anti-gp120 antibody b8. This fragment had been amplified by the PCR primer pair CR501 (SEQ ID NO:28) / CR301 (SEQ ID NO:29). The sense primer CR501 encodes a synthetic link of the nine C-terminal amino acids of FR3, the eight amino acids forming the HCDR3 (SEQ ID NO:1) of LM609, and the six N-terminal amino acids of FR4 of b8. The FR4 of b8 is a
 30 preferred choice here because it is identical to FR4 of LM609's V_H with the exception of the C-terminal amino acid, which is A for LM609 and S for b8. The 24-bp overlap of CR501 and BFR3UN allowed to fuse the corresponding PCT products by overlap extension PCR.

The sense primer CR301 hybridizes to a sequence that encodes the C terminus of C_H1 and introduces a SpeI site that allows the PCR product to link to the gene III ORF on pComb3H. The product of the overlap extension PCR was cut with XhoI/SpeI, ligated into the appropriately digested phagemid vector pComb3H, and the correct sequence was confirmed by DNA sequencing.

Substitution of the LM609 light chain by a human light chain that contains the LCDR3 of LM609

Using overlap extension PCR, the amplified human sequences encoding the N-terminal FR1 through FR3 fragment of V_κ and V_λ were fused to PCR fragments encoding the LCDR3 (SEQ ID NO:2) of LM609 coupled to FR4 of human V_κ or V_λ and the human C_κ or C_λ domain. Two κ fragments were generated by the PCR primer pairs CR503 (SEQ ID NO:30) / T7B (SEQ ID NO:31) and CR508 (SEQ ID NO:32) / T7B using the sequence of the anti-gp120 antibody b11 in pComb3 as a template.

The sense primers CR503 and CR508 encode a synthetic link of eight C-terminal amino acids of FR3 of human V_κ (SEQ ID NO:33 or SEQ ID NO:34), the nine amino acids forming the LCDR3 (SEQ ID NO:2) of LM609, and the seven N-terminal amino acids of FR4 of b11. FR4 of b11 is the preferred choice because it is identical to FR4 of LM609's V_κ with the exception of the third N-terminal and C-terminal amino acid, which are G and T in LM609 versus Q and A in b11. The 23-bp overlap of CR503 with BKFR3UN and CR508 with BK2FR3UN allowed to fuse the corresponding PCR products by overlap extension PCR.

The backward primer T7B hybridizes to a pComb3 sequence downstream of the light chain encoding sequence. A λ fragment was generated by the PCR primer pair CR510 (SEQ ID NO:35) / CLext (SEQ ID NO:36) using CLext primed first strand cDNA from human bone marrow as a template.

The sense primer CR510 encodes a synthetic link of seven C-terminal amino acids of FR3 of human V_λ (SEQ ID NO:37), the nine amino acids forming the LCDR3 of LM609, and the seven N-terminal amino acids of FR4 of human V_λ (SEQ ID NO:38). The 21-bp overlap of CR510 with BLFR3UN allowed to fuse the corresponding PCR products by overlap extension PCR. The backward

- 13 -

primer CLext hybridizes to the 3' end of the human C_λ encoding sequence and introduces a XbaI site.

The generated light chain encoding sequences were cut with SacI/XbaI and ligated into the appropriately digested phagemid vector pComb3H that contained the chimeric mouse/human Fd fragment. Electrotransformation of the ligation products into *E. coli* strain ER 2537 (New England Biolabs; Beverly, MA) resulted in a light chain library consisting of 1.5 x 10⁸ independent transformants. DNA sequencing revealed the correct assembly of the fused fragments.

Four rounds of panning against immobilized human integrin α_vβ₃ were carried out essentially as described in "High-affinity self-reactive human antibodies by design and selection: targeting the integrin ligand binding site", (*Proc. Natl. Acad. Sci. USA* 90, 10003-10007 (1993)) by C.F. Barbas, III, et al. using 200 ng protein in 25 μl metal buffer (25 mM Tris-HCl, pH 7.5; 137 mM NaCl; 1 mM KCl; 1 mM MgCl₂; 1 mM CaCl₂; 1 mM MnCl₂) for coating, 0.05 percent TweenTM 20 in TBS for washing, and 10 mg/ml trypsin in TBS for elution. The washing steps were increased from 5 in the first round, to 10 in the second round and 15 in the third and fourth rounds. The output phage pool of each round was monitored by phage ELISA.

After the fourth round phage were produced from single clones and tested for binding to immobilized integrin α_vβ₃ by phage ELISA. Light chain encoding sequences of positive clones were analyzed by DNA sequencing using the primer OMPSEQ (SEQ ID NO:39) that hybridizes to the ompA leader sequence upstream of the light chain encoding sequence in pComb3H.

25 Substitution of the LM609 Fd fragment by a human Fd fragment that contains the heavy complementarity determining region three (HCDR3) of LM609

Three PCR fragments were fused in one step by overlap extension PCR. Using the selected phagemids from the light chain panning as a template, Fragment 1 was amplified with the PCR primer pair RSC-F (SEQ ID NO:40) / lead-B (SEQ ID NO:41). While the sense primer RSC-F hybridizes to a sequence upstream of the light chain encoding sequence, the antisense primer lead-B hybridizes to a sequence upstream of the Fd fragment encoding sequence. The amplified human

sequences encoding the FR1 through FR3 fragment of VH (see above) were used as fragment 2.

Fragment 3 was amplified with the PCR primer pair CR501 / HIgG1-B (SEQ ID 42) using the hybrid mouse/human Fd fragment (see above) as a template. The antisense primer HIgG1-B hybridizes to the 3' end of the C_H1 encoding sequence. Using the 21-bp overlap of lead-B with the HFVH-F primers and the 24-bp overlap of BFR3UN with CR501, the three fragments were fused and amplified with the PCR primer pair RSC-F / RSC-B (SEQ ID NO:43). The antisense primer RSC-B overlaps with HIgG1-B. RSC-F and RSC-B introduce two asymmetric SfiI sites.

To maintain high complexity, separate PCR reactions were performed for each selected phagemid from the light chain panning (Fragment 1) and for each of the five V_H fragment pools derived from the five first strand cDNA sources (Fragment 2). The generated fragments encoding the selected human light chains linked to human Fd fragments were cut with SfiI and ligated into the appropriately digested phagemid vector pComb3H generating a library of 3 x 10⁷ independent transformants.

DNA sequencing revealed the correct assembly of the fused DNA fragments. Four rounds of panning against immobilized human integrin $\alpha_v\beta_3$ were carried out exactly as described for the light chain panning. The output phage pool of each round was monitored by phage ELISA. After the fourth round, light chain and Fd fragment encoding sequences were isolated from the selected phagemids by SfiI digestion and subcloned into the compatible expression vector pPhoA-H6HA.

Lysates of individual clones grown in phosphate-deprived medium were analyzed for binding to immobilized integrin $\alpha_v\beta_3$ by ELISA using goat anti-human F(ab')₂ conjugated to alkaline phosphatase (Pierce) as secondary antibody. Light chain and Fd fragment encoding sequences of positive clones were analyzed by DNA sequencing using the primers OMPSEQ and PELSEQ, respectively.

Results

cDNA Cloning of LM609

Starting from LM609 expressing hybridoma cells, cDNAs encoding λ chain Fd fragments and entire κ chains were cloned by PCR. The PCR products

were cloned into the phagemid pComb3H, which is derived from pComb3, and engineered to facilitate the expression of Fab on the surface of M13 filamentous phage. Phage displaying LM609 Fab were selected by panning against integrin $\alpha_v\beta_3$ and the corresponding cDNA sequences were determined. Soluble LM609 Fab
5 purified from *E.coli* was analyzed and found to bind specifically to integrin $\alpha_v\beta_3$ by ELISA.

The approach for the sequential humanization of LM609 by a combination of guided selection and CDR grafting is illustrated in FIGURE 1. For the human light chain selection, the mouse Fd fragment is substituted by a chimeric
10 Fd fragment composed of mouse V_H linked to human C_H1 to stabilize the hybrid Fab of the first selection step by the interaction of two matching human constant domains, C_κ and C_H1 . A stabilization of the hybrid Fab also stabilizes the antigen binding site.

The guided selection is started by substituting the κ light chain of LM609 with a human κ and λ light chain library that contained the grafted LCDR3
15 loop of LM609. The corresponding phage libraries displaying hybrid Fab are then selected by four rounds of panning against immobilized integrin $\alpha_v\beta_3$. Though the output number does not increase from round to round, analysis of the output phage pool from each round for binding to integrin $\alpha_v\beta_3$ by phage ELISA reveals an increasing signal. After the fourth round, phage are produced from clones and tested
20 for binding to integrin $\alpha_v\beta_3$ by phage ELISA.

While the majority of these clones give signals that reveal some binding above background, in the present case, six clones gave very strong signals (See FIGURES 3a through 3e). DNA sequence analysis of these clones revealed 3 different light chain sequences. Two light chain sequences found in five out of six
25 positive clones differ only in four amino acids, i.e., are 96 percent identical. The third light chain sequence shares about 80 percent identity with the other two. However, this sequence had two parts, each of which could be aligned to germ-lines of different V_κ families, and, thus is deemed likely to have arisen from PCR cross-over, an artifact that has been reported to occur frequently in the amplification
30 antibody sequences.

Referring to FIGURES 3a through 3e, analysis of the six human V_κ revealed two groups of highly related κ chain sequences. In addition, the CDR1 loops of all six selected human V_κ , which are believed to play a role in the assembly

of V_{κ} and V_{λ} , resemble the corresponding region of LM609 V_{κ} . This indicates that the template V_{λ} of LM609 together with the LM609 antigen, the human integrin $\alpha_v\beta_3$, selected for human κ chains that are related to the LM609 κ chain. The fact that no repeated sequences were found may indicate that the grafted LCDR3 loop of LM609, which is identical in all selected human κ chains, is mainly responsible for the contribution of the LM609 κ chain to antigen binding.

This supposition is supported by two additional observations. First, the initial humanization approach was based on the original human κ chain libraries. Four rounds of panning selected a repeated human κ chain with a sequence related to the LM609 κ chain. However, the corresponding hybrid Fab appeared to bind only weakly to human integrin $\alpha_v\beta_3$. Therefore, the LCDR3 loop of LM609 was grafted in the human κ chain libraries, and, though only roughly estimated from ELISA, the binding of the corresponding selected hybrid Fab to human integrin $\alpha_v\beta_3$ improved.

Second, the selected phage were selected by two further rounds of panning against immobilized human integrin $\alpha_v\beta_3$. Again, soluble ones derived from selected phagemids were analyzed for binding to immobilized human integrin $\alpha_v\beta_3$ by ELISA. This time, all of the analyzed 20 clones were found to bind specifically. However, sequencing of 16 clones revealed no repeated sequences. It appears, thus, that a number of different human κ chain sequences that contain the LCDR3 loop of LM609 can substitute the LM609 κ chain without much difference in binding to human integrin $\alpha_v\beta_3$. This finding is believed to be of importance for the therapeutic application of humanized LM609.

Due to allotypic sequence variability, humanized antibodies can be neutralized by the patient's immune system after repeated injections. This problem is avoided by using humanized antibodies with identical antigen binding properties but different amino acid sequences for repeated administrations.

As with the original LM609 light chain, the selected light chains are each κ light chains. Moreover, database screening revealed that they were derived from the same germ-line, namely DPK-26, belonging to the $V_{\kappa}6$ family. This speaks in favor of a strong selection towards these sequences because the $V_{\kappa}6$ family is not frequently found in human antibodies. An obvious reason for this strong selection is a relatively high sequence similarity of the selected human light chains with the original mouse light chain.

Referring to FIGURE 4, for comparison, four clones from the unselected library were picked randomly and their light chain sequences determined. Three selected human light chains used in the comparison consist of eleven LCDR1 amino acids, the length of the original mouse LCDR1, while only one out of four
5 unselected human light chains shared the same LCDR1 length with the original mouse sequence (see FIGURE 4).

Moreover, both LCDR1 and LCDR2 of the selected human light chains are highly similar to the corresponding mouse sequence. The C-terminal amino acid of framework region 2 of the original mouse light chain sequence, a
10 lysine (Lys) shown in brackets in FIGURE 4, is an unusual amino acid at this position and, thus may be involved in the formation of the antigen binding site. Interestingly, all the selected human light chain sequences contain a lysine at this position, while all the unselected sequences contain a tyrosine (Tyr) instead. As a matter of fact, the $V_{\kappa}6$ family is the only human V_{κ} family that contains a lysine at
15 that position.

Taken together, this evidence shows that the mouse V_H template and the antigen selected for unbiased human V_{κ} sequences are related to the original mouse V_{κ} sequence. Three clones from the light chain selection, revealing weaker binding to integrin $\alpha_v\beta_3$ than the six clones discussed above but still having
20 significant binding above background, were also analyzed by DNA sequencing. These analyses revealed three unrelated V_{λ} sequences, together with selected V_{κ} sequences—except the one that stemmed from the PCR cross-over artefact— were used as templates in the humanization of the heavy chain of LM609.

Based on the aforescribed humanization strategy, five humanized
25 versions of the anti-human integrin $\alpha_v\beta_3$ monoclonal antibody LM609 were generated. The five version were revealed through the sequence analysis of 14 humanized clones that bind to $\alpha_v\beta_3$. Referring to FIGURES 8a and 8b, the amino acid sequences of a mouse V_L (SEQ ID NO:45) are compared to the amino acid sequences of the version A, and the combined versions (or groups) B, C, D, and E.

30 Similarly FIGURES 8c through 8e compare the amino acid sequences of a mouse V_H (SEQ ID NO:56) to the amino acid sequences of the five versions (or groups) A-E. Four of these versions, represented by clones 7, 8, and 22 (group B); 4, 31, and 36 (group C); 24, 34, and 40 (group D); and 2 (group E), are

highly related in amino acid sequence. The sequence group BCDE in FIGURES 8a and 8b represent the four versions that share an identical V_L domain (SEQ ID NO:49). The amino acid sequence identity of their V_H domains (SEQ ID NOS:50-53), which are all derived from the germ-line DP-65 or the highly related DP-78, is
5 at least 90% for each version.

In contrast, version five includes clones 10, 11, and 37 (A), and represents a humanized version with a V_H domain (SEQ ID NO:54) that is derived from a different germ-line family. This humanized version also contains a different V_L domain (SEQ ID NO:55), which is, however, 95% identical and derived from the
10 same germ-line. Germ-lines were determined by nucleic acid sequence alignment using DNAPLOT software provided by the VBASE Directory of Human V Gene Sequences from the MRC Centre for Protein Engineering.

By preserving the original complementarity determining region sequences such as the LCDR3 (SEQ ID NO:2) and HCDR3 (SEQ ID NO:1)
15 sequences of LM609, the disclosed humanization strategy ensures epitope conservation. Epitope conservation is a critical demand in the humanization of antibodies, especially in the case of LM609. The function-blocking anti-human integrin $\alpha_v\beta_3$ mouse monoclonal antibody LM609 binds to a yet unidentified nonlinear epitope that involves both the α_v and β_3 polypeptide chains. Importantly,
20 by binding to this epitope LM609 induces apoptosis in $\alpha_v\beta_3$ expressing vascular cells, a unique feature among a number of anti-human integrin $\alpha_v\beta_3$ mouse monoclonal antibodies. LM609 does not recognize the related human integrin $\alpha_{IIb}\beta_3$. Any cross-reactivity with human $\alpha_{IIb}\beta_3$, which is expressed on platelets, precludes the use of LM609 as a tool in cancer therapy.

25 The five humanized versions of LM609, clones 2, 4, 7, 11, and 24, which had been selected by binding to immobilized, thus potentially denatured human integrin $\alpha_v\beta_3$, were tested for binding to native human integrin $\alpha_v\beta_3$ expressed on the cell surface. For this, binding of humanized LM609 to untransfected CS-1 hamster cells and CS-1 hamster cells transfected with either human β_3 or β_5 cDNA was
30 analyzed by flow cytometry. Like mouse LM609, but in contrast to a control antibody, all five humanized versions of LM609 revealed specific binding to CS-1 hamster cells transfected with human β_3 cDNA (See FIGURES 5a-5f).

Potential cross-reactivity of humanized LM609 with human integrin $\alpha_{\text{IIb}}\beta_3$ was analyzed by ELISA. While antibody Fab-9 with known cross-reactivity bound to both immobilized human integrin $\alpha_v\beta_3$ and $\alpha_{\text{IIb}}\beta_3$, cross-reactivity was detected neither for mouse LM609 nor its five humanized versions (See FIGURE 6).

5 Thus, all evidence speaks in favor of epitope conservation through the process of humanization of LM609.

Affinity maturation is a highly relevant step in engineering antibodies for therapeutic applications. By increasing the target affinity, the *in vivo* concentration of an antibody that must be reached to be effective for therapy is
10 lowered. In addition to reducing the costs of antibody therapy, low effective *in vivo* concentrations will help to reduce the chance of immune response.

The CDR walking strategy for the affinity maturation of antibodies has been described elsewhere, and is known in the relevant art. For the affinity maturation of humanized LM609 a sequential optimization of LCDR3 and HCDR3
15 was chosen (See FIGURE 7). The randomized region in both CDRs was confined to a stretch of four amino acids that revealed highest variability in human antibody sequences. See Kabat, E.A. et al. (1991) SEQUENCES OF PROTEINS OF IMMUNOLOGICAL INTEREST, U.S. Dept of Health and Human Services, Washington, D.C. Using NNK doping (Barbas, C.F. et al., (1994) *IN VITRO* EVOLUTION OF A
20 NEUTRALIZING HUMAN ANTIBODY TO HIV-1 TO ENHANCE AFFINITY AND BROADEN STRAIN CROSS-REACTIVITY, *Proc. Natl. Acad. Sci. USA* 91, 3809-3813) the randomization of four codons results in 32^4 or 1×10^6 different nucleotide sequences. Based on this and assuming a Poisson distribution (Clackson, T., and Wells, J.A. (1994) *IN VITRO* SELECTION FROM PROTEIN AND PEPTIDE LIBRARIES, *Trends in*
25 *Biotechnol.* 12, 173-184), 5×10^6 independent transformants are required for a complete library with 99 percent confidence. For each of the five humanized versions of LM609 independent LCDR3 randomized libraries were generated. The number of independent transformants in each library lay well above 5×10^6 (see Experimental Procedures, below). Ten completely different LCDR3 sequences were
30 obtained when two clones of each of the five libraries were analyzed by DNA sequencing. The three libraries that were based on humanized LM609 clones 2, 4, and 24, which contain identical light chains and highly related heavy chains derived from germ-line DP-65, were pooled. Corresponding to their underlying germ-line

sequences the three remaining libraries were named DPK-26ranLCDR3/DP-10, DPK-26ranLCDR3/DP-65, and DPK-26ranLCDR3/DP-78. These three libraries were selected in parallel.

To improve the selection of higher affinity LCDR3 mutants, a solid phase off-rate selection (Yang, W.P. et al. (1995) CDR WALKING MUTAGENESIS FOR THE AFFINITY MATURATION OF A POTENT HUMAN ANTI-HIV-1 ANTIBODY INTO THE PICOMOLAR RANGE, *J.Mol Biol.* **254**, 392-403) was used. In five subsequent cycles of selection 20 μ g LM609 IgG was added to the well with 200 or 50 ng immobilized human integrin $\alpha_v\beta_3$ following phage incubation and washing. After 24 hours at room temperature, the well was washed again and bound phage eluted with trypsin. This off-rate selection step is discussed in the following paragraph.

Protein interactions are characterized by thermodynamic and kinetic parameters. While the affinity constant ($K_a = k_{on}/k_{off}$) is an equilibrium constant, the association (k_{on}) and dissociation (k_{off}) rate constants are more relevant to *in vivo* processes which are beyond equilibrium. See Williams, A.F. (1991) OUT OF EQUILIBRIUM, *Nature* **352**, 473-474. In fact, to occur *in vivo*, interactions with high affinity, i.e., high K_a values, still rely on rapid association, i.e., high k_{on} values. Antibodies are subject to kinetic selection based on binding target antigens rapidly, in parallel with thermodynamic selection for high affinity binding in order to allow sufficient time for antigen clearance. See Foote, J., and Milstein, C. (1991) KINETIC MATURATION OF AN IMMUNE RESPONSE, *Nature* **352**, 530-532. A typical antibody/antigen interaction with a K_a value in the range of 10^9 M^{-1} associates rapidly with a k_{on} value in the range of 10^5 to 10^6 $M^{-1}s^{-1}$ and dissociates slowly with a k_{off} value in the range of 10^{-3} to 10^{-4} s^{-1} . An off-rate selection for affinity maturation, i.e., decreasing k_{off} , requires consideration of the half-life of the antibody/antigen interactions that is given by $t_{1/2} = \ln 2/k_{off}$. An antibody/antigen interaction with $k_{off} = 1 \times 10^{-4}$ s^{-1} has a half-life of about 2 hours. A tenfold lower dissociation constant, i.e., $k_{off} = 1 \times 10^{-5}$ s^{-1} , results in a tenfold longer half-life, i.e., about 20 hours. These long half-lives limit the off-rate selection in our affinity maturation protocol. Using a reasonable time frame, antibodies with dissociation constants below 1×10^{-6} s^{-1} can not be enriched even after multiple selection cycles. However, using a similar protocol, an antibody was selected against gp120 with a k_{off} value in the range of 10^{-6} s^{-1} . The

corresponding affinity constant was in the range of 10^{11} M⁻¹, a more than 400-fold improvement of the parental antibody.

Eight clones from each of the three independently selected libraries were analyzed by DNA sequencing. The LCDR3 sequences are shown in Table 1, below. A strong selection towards a consensus sequence that is highly related to the original sequence took place. All 24 analyzed clones contain a serine (Ser) in Position 91 and a glutamine (Gln) in Position 92 of the randomized region. The corresponding amino acids in the parental LCDR3 are serine (Ser) and asparagine (Asn), respectively. Interestingly, all three serine codons of the NNK genetic code (TCT, TCG, and ACT) are found in Position 91. Position 94, a tryptophane (Trp) in the parental LCDR3, was re-selected in 22 out of 24 clones. Two clones contain a histidine (His) instead. Only Position 93 reveals greater diversity. The original serine (Ser) is substituted by an aromatic or hydrophobic amino acid, either phenylalanine (Phe-13/24), tryptophane (Trp-6/24), valine (Val-3/24), tyrosine (Tyr-1/24) or histidine (His-1/24). Analysis of the heavy chain sequences revealed that no cross-contamination between the three independently selected libraries took place. All eight clones selected from library DPK-26ranLCDR3/DP-65, which contained the pool of the three highly related heavy chains encoding sequences derived from germ-line DP-65, were identical and derived from humanized LM609 clone 24.

The conserved LCDR3 sequence speaks in favor of a highly defined epitope on human integrin $\alpha_v\beta_3$. Though binding to native human integrin $\alpha_v\beta_3$ on the cell surface needs to be proved yet, an epitope shift towards denatured human integrin $\alpha_v\beta_3$ is unlikely. The selected phage pools were analyzed for binding to human integrin $\alpha_v\beta_3$ by phage ELISA and in competition with LM609 IgG. These analyses suggest a significantly lower dissociation constant of the selected clones in comparison with LM609 as well as humanized LM609. The substitution of the original serine in Position 3 by an aromatic residue may give rise to a new hydrophobic interaction with a strong impact on the overall affinity.

TABLE 1

Selected LCDR3 Mutants

Kabat position ¹	91	92	93	94
LM609	Ser	Asn	Ser	Trp
5 Library DPK- 26ranLCDR3/DP-10 ²	Ser Ser Ser Ser Ser Ser Ser Ser	Gln Gln Gln Gln Gln Gln Gln Gln	Trp Trp Trp Trp Val Phe Phe Phe	Trp Trp Trp Trp Trp Trp His His
Library DPK- 26ranLCDR3/DP-65 ³	Ser Ser Ser Ser Ser Ser Ser Ser	Gln Gln Gln Gln Gln Gln Gln Gln	Phe Phe Phe Phe Phe Phe Phe Phe	Trp Trp Trp Trp Trp Trp Trp Trp
10 Library DPK- 26ranLCDR3/DP-78 ⁴	Ser Ser Ser Ser Ser Ser Ser Ser	Gln Gln Gln Gln Gln Gln Gln Gln	Trp Trp Val Val Phe Phe Tyr His	Trp Trp Trp Trp Trp Trp Trp Trp

¹ cf. Kabat, et al. (1991)² based on humanized LM609 clone 11³ based on humanized LM609 clones 2, 4, and 24⁴ based on humanized LM609 clone 7Experimental Procedures**Materials**

LM609 IgG purified from hybridoma cultures was provided by Dr. David A. Cheresh. Human integrin $\alpha_v\beta_3$ and human integrin $\alpha_{IIb}\beta_3$ were from sources described in HIGH-AFFINITY SELF-REACTIVE HUMAN ANTIBODIES BY DESIGN AND SELECTION: TARGETING THE INTEGRIN LIGAND BINDING SITE, Barbas III, C.F., et al. (1993) *Proc. Natl. Acad. Sci. USA*, **90**, 10003-10007. Untransfected CS-1

hamster cells and CS-1 hamster cells transfected with either human β_3 or β_5 cDNA were obtained from Dr. David A. Cheresh (Filardo, E.J., et al. (1995)

REQUIREMENT OF THE NPXY MOTIF IN THE INTEGRIN β_3 SUBUNIT CYTOPLASMIC TAIL FOR MELANOMA CELL MIGRATION *IN VITRO* AND *IN VIVO*, *J. Cell Biol.* **130**, 441-450)

5 and maintained in RPMI medium supplemented with 5% fetal calf serum (FCS) at 37°C in 7% CO₂.

***E. coli* expression of soluble humanized LM609 Fab**

Following phage library panning, the SfiI insert of the selected
 10 humanized LM609 phagemid pool was cloned into the *E. coli* expression vector pPhoA-H6HA (See Rader, C., and Barbas III, C.F. (1997) PHAGE DISPLAY OF COMBINATORIAL ANTIBODY LIBRARIES, *Curr. Opin. Biotechnol.* **8**, 503-508) for detection of $\alpha_v\beta_3$ binders. Sequence determination of 14 clones revealed five different humanized LM609 versions, represented by clones 2, 4, 7, 11, and 24.
 15 cDNAs of these clones were cut out by SacI/SpeI digestion and ligated into SacI/NheI cut pComb3H, thereby removing the gene III fragment encoding cDNA of pComb3H and allowing for production of soluble Fab (See Rader, C. (1997) *Curr. Opin. Biotechnol.* **8**, 503-508). The ligation products were electrotransformed into *E. coli* strain XL1-Blue. Fab production was induced by addition of isopropyl β -D-thiogalactopyranoside as described (Barbas III, C.F., et al. (1991) ASSEMBLY OF
 20 COMBINATORIAL ANTIBODY LIBRARIES ON PHAGE SURFACES: THE GENE III SITE, *Proc. Natl. Acad. Sci. USA* **88**, 7978-7982).

ELISA

25 Human integrins $\alpha_v\beta_3$ and $\alpha_{Iib}\beta_3$ were coated for 90 minutes at 37°C on a 96-well plate (Costar #3690) at a concentration of 60 ng/25 μ l metal buffer (25 mM Tris-HCl, pH 7.5; 137 mM NaCl; 1 mM KCl; 1 mM MgCl₂; 1 mM CaCl₂; 1 mM MnCl₂) per well. Following blocking with 150 μ l 3% BSA/TBS for 1 hour at 37°C, 25 μ l crude supernatants from overnight cultures of *E. coli* strain XL1-Blue
 30 expressing soluble LM609 or humanized LM609 Fab were added to the well and incubated for 2 hours at 37°C. Binding of each of the supernatants to wells coated with human integrin $\alpha_v\beta_3$ and $\alpha_{Iib}\beta_3$ as well as to uncoated but blocked wells was analyzed in triplicates. As a positive control, 25 μ l of 50 ng/ μ l purified Fab-9, an

- 24 -

antibody binding to both human integrin $\alpha_v\beta_3$ and $\alpha_{IIb}\beta_3$ (Barbas III, et al. (1993) *Proc. Natl. Acad. Sci. USA* 90, 10003-10007), and as a negative control, 25 μ l of plain bacterial culture medium were used. After extensive washing with tap water, 25 μ l of a 1:2,000 dilution of goat anti-mouse F(ab')₂ or goat anti-human F(ab')₂ conjugated with alkaline phosphatase (Pierce #31324 or 31312, respectively) in 1% BSA/TBS was added to the well and incubated for 1 hour at 37°C. Following another extensive washing with tap water, 50 μ l of alkaline phosphatase substrate (5 mg disodium p-nitrophenyl phosphate hexahydrate (Sigma #104-105) dissolved in 5 ml 10% diethanolamine, 1 mM MgCl₂, 3 mM NaN₃, pH 9.8) were added to the well. The plate was analyzed with an ELISA reader (Molecular Devices) after 15 minutes incubation at room temperature.

Flow cytometry

Flow cytometry was performed using a Becton Dickinson FACScanTM instrument. For each determination, 5x10³ untransfected hamster CS-1 cells or hamster CS-1 cells expressing either human β_3 or β_5 , were analyzed. Indirect immunofluorescence staining was achieved with crude lysates of *E.coli* strain XL1-Blue expressing soluble humanized LM609 Fab or, as a negative control, an unrelated human Fab. A 1:100 dilution of FITC-conjugated goat anti-human F(ab')₂ (Jackson #109-096-097) was used for detection.

Construction of LCDR3 libraries

Humanized LM609 clones 2, 4, 7, 11, and 24 in pPhoA-H6HA were separately utilized as templates for overlap extension PCR mutagenesis as described (Barbas III, et al. (1994) *IN VITRO EVOLUTION OF A NEUTRALIZING HUMAN ANTIBODY TO HIV-1 TO ENHANCE AFFINITY AND BROADEN STRAIN CROSS-REACTIVITY*, *Proc. Natl. Acad. Sci. USA* 91, 3809-3813). The two fragments required for this procedure were obtained with the PCR primer pairs OMPSEQ (SEQ ID NO:39) / CR320 (SEQ ID NO:46) and CR520 (SEQ ID NO:47) / DPSEQ (SEQ ID NO:48), respectively. The resulting five cDNAs with randomized LCDR3 were cut with SfiI, ligated into the appropriately digested phagemid vector pComb3H, and electrotransformed into *E.coli* strain ER 2537. Two clones of each of the five libraries were analyzed by DNA sequencing and revealed correct assembly as well as

10 different LCDR3 sequences. Prior to selection, libraries based on clones 2, 4, and 24 (V_L germ-line DP-26; V_H germ-line DP-65) were combined to give a complexity of 6×10^7 independent transformants. Libraries based on clone 11 (V_L germ-line DP-26; V_H germ-line DP-10) and 7 (V_L germ-line DP-26; V_H germ-line DP-78) were kept separate with a complexity of 3×10^7 and 4×10^7 independent transformants, respectively.

Selection of LCDR3 libraries

The three LCDR3 libraries were separately selected by panning against immobilized integrin $\alpha_v\beta_3$ for six cycles. Panning was performed substantially as described hereinabove for the LM609 humanization. The concentration of human integrin $\alpha_v\beta_3$ for coating was 200 ng/25 μ l in the first through fourth cycles and 50 ng/25 μ l in the fifth and sixth cycles. Also, the input number of phage, in the range of 10^{12} in the first through fourth cycles as usual, was decreased by a factor of 10 in the fifth cycle and by a factor of 100 in the sixth cycle. In the second through the sixth cycles of selection 20 μ g LM609 IgG in 50 μ l metal buffer (25 mM Tris-HCl, pH 7.5; 137 mM NaCl; 1 mM KCl; 1 mM $MgCl_2$; 1 mM $CaCl_2$; 1 mM $MnCl_2$) was added to the well following removal of unbound phage by five to ten washing steps. The plate was then incubated for 24 hours at room temperature (off-rate selection) prior to five additional washing steps and trypsin elution as described. After the sixth cycle phage were produced from single clones and tested for binding to immobilized human integrin $\alpha_v\beta_3$ by phage ELISA using a sheep anti-M13 antibody conjugated with horseradish peroxidase (Pharmacia #27-9411-01) as a secondary antibody. Light chain and heavy chain encoding sequences of positive clones were analyzed by DNA sequencing using the primers OMPSEQ (SEQ ID NO:39) and PELSEQ (SEQ ID NO:24), respectively.

Optimization of LM609 by CDR randomization

In addition to the humanization, LM609 can be optimized in two respects: First, by increasing the affinity to $\alpha_v\beta_3$ and, second, by broadening the species cross-reactivity. Increased affinity of engineered LM609 increases the potency and decreases the cost of a potential cancer therapy.

The original mouse monoclonal antibody LM609 already has a broad species cross-reactivity. It binds to human, dog, cat, bovine, rabbit, and chick but not mouse $\alpha_v\beta_3$. The fact that LM609 does not recognize host $\alpha_v\beta_3$ in the mouse models of human cancer is a major concern for the therapeutic applicability of LM609. Engineered LM609 binding to both human and mouse $\alpha_v\beta_3$ would be an important tool towards clinical trials. *In vitro* methods for the improvement of monoclonal antibody affinity include *chain shuffling* (See Marks, J.D., et al. (1992) BY-PASSING IMMUNIZATION: BUILDING HIGH HUMAN ANTIBODIES BY CHAIN SHUFFLING, *Bio/Technology* 10, 779-783). Binding to $\alpha_v\beta_3$ can be further improved by subsequent CDR randomization, an approach termed *CDR walking* (See Barbas III, C.F., et al. (1994) *IN VITRO* EVOLUTION OF A NEUTRALIZING HUMAN ANTIBODY TO HIV-1 TO ENHANCE AFFINITY AND BROADEN STRAIN CROSS-REACTIVITY, *Proc. Natl. Acad. Sci. USA* 91, 3809-3813). The *in vitro* strategies for humanization and affinity improvement of LM609 are likely to generate cross-reactivity with mouse $\alpha_v\beta_3$ concurrently. Directed selection for mouse $\alpha_v\beta_3$ recognition is complicated by the fact that mouse $\alpha_v\beta_3$ has not been purified yet. However, several mouse cell lines, e.g., NIH/3T3, are known to express $\alpha_v\beta_3$ and, thus, may be included in the screening procedure.

The foregoing discussion and the accompanying examples are presented as illustrative, and are not to be taken as limiting. Still other variations within the spirit and scope of this invention are possible and will readily present themselves to those skilled in the art.

SEQUENCE LISTING

<110> The Scripps Research Institute

<120> HUMANIZATION OF MURINE ANTIBODY

<130> TSRI 598.0

<140> PCT/US98/25828

<141> 1998-12-04

<150> 08/986,016

<151> 1997-12-05

<160> 56

<170> PatentIn Ver. 2.0

<210> 1

<211> 8

<212> PRT

<213> Murine hybridoma LM609

<400> 1

His Asn Tyr Gly Ser Phe Ala Tyr

1

5

<210> 2

<211> 9

<212> PRT

<213> Murine hybridoma LM609

<400> 2

Gln Gln Ser Asn Ser Trp Pro His Thr

1

5

<210> 3

<211> 37

<212> DNA

<213> Murine hybridoma LM609

<400> 3

gggccccaggc ggccgagctc cagatgaccc agtctcc

37

<210> 4

<211> 37

<212> DNA

<213> Murine hybridoma LM609

<400> 4

gggccccaggc ggccgagctc gtgatgacgc agtctcc

37

<210> 5

<211> 37

<212> DNA

<213> Murine hybridoma LM609

<400> 5

gggccccaggc ggccgagctc gtgwtgacgc agtctcc

37

<210> 6
<211> 37
<212> DNA
<213> Murine hybridoma LM609

<400> 6

gggcccaggc ggccgagctc acactcacgc agtctcc

37

<210> 7
<211> 23
<212> DNA
<213> Murine hybridoma LM609

<400> 7

cagtaataca ctgcaaaatc ttc

23

<210> 8
<211> 23
<212> DNA
<213> Murine hybridoma LM609

<400> 8

cagtaataaaa cccaacatc ctc

23

<210> 9
<211> 40
<212> DNA
<213> Murine hybridoma LM609

<400> 9

gggcccaggc ggccgagctc gtgbtgacgc agccgccctc

40

<210> 10
<211> 40
<212> DNA
<213> Murine hybridoma LM609

<400> 10

gggcccaggc ggccgagctc gtgctgactc agccaccctc

40

<210> 11
<211> 43
<212> DNA
<213> Murine hybridoma LM609

<400> 11

gggcccaggc ggccgagctc gccctgactc agcctccctc cgt

43

<210> 12
<211> 46
<212> DNA
<213> Murine hybridoma LM609

<400> 12

gggcccaggc ggccgagctc gagctgactc agccaccctc agtgctc 46

<210> 13
<211> 40
<212> DNA
<213> Murine hybridoma LM609

<400> 13

gggcccaggc ggccgagctc gtgctgactc aatcgccctc 40

<210> 14
<211> 40
<212> DNA
<213> Murine hybridoma LM609

<400> 14

gggcccaggc ggccgagctc atgctgactc agccccactc 40

<210> 15
<211> 40
<212> DNA
<213> Murine hybridoma LM609

<400> 15

gggcccaggc ggccgagctc gggcagactc agcagctctc 40

<210> 16
<211> 40
<212> DNA
<213> Murine hybridoma LM609

<400> 16

gggcccaggc ggccgagctc gtggtgacyc aggagccmtc 40

<210> 17
<211> 40
<212> DNA
<213> Murine hybridoma LM609

<400> 17

gggcccaggc ggccgagctc gtgctgactc agccaccttc 40

<210> 18
<211> 21
<212> DNA
<213> Murine hybridoma LM609

<400> 18

gcagtaataa tcagcctcrt c 21

<210> 19
<211> 44
<212> DNA

<213> Murine hybridoma LM609

<400> 19

gctgccaac cagccatggc ccaggtgcag ctggtgcagt ctgg

44

<210> 20

<211> 44

<212> DNA

<213> Murine hybridoma LM609

<400> 20

gctgccaac cagccatggc ccagatcacc ttgaaggagt ctgg

44

<210> 21

<211> 44

<212> DNA

<213> Murine hybridoma LM609

<400> 21

gctgccaac cagccatggc cgaggtgcag ctggtgsagt ctgg

44

<210> 22

<211> 44

<212> DNA

<213> Murine hybridoma LM609

<400> 22

gctgccaac cagccatggc ccaggtgcag ctgcaggagt cggg

44

<210> 23

<211> 24

<212> DNA

<213> Murine hybridoma LM609

<400> 23

cgcacagtaa tacacggccg tgtc

24

<210> 24

<211> 21

<212> DNA

<213> Murine hybridoma LM609

<400> 24

acctattgcc tacggcagcc g

21

<210> 25

<211> 24

<212> DNA

<213> Murine hybridoma LM609

<400> 25

cgcacagtaa tacacggccg tgtc

24

<210> 26
<211> 8
<212> PRT
<213> Murine hybridoma LM609

<400> 26

Asp Thr Ala Val Tyr Tyr Cys Ala

1 5

<210> 27
<211> 8
<212> PRT
<213> Murine hybridoma LM609

<400> 27

Asp Thr Ala Met Tyr Tyr Cys Ala

1 5

<210> 28
<211> 69
<212> DNA
<213> Murine hybridoma LM609

<400> 28

gacacggccg tgtattactg tgcgcgtcat aactacggca gttttgctta ctggggccag 60
ggaaccctg 69

<210> 29
<211> 42
<212> DNA
<213> Murine hybridoma LM609

<400> 29

gaggaggagg aggagactag ttttgtcaca agatttgggc tc 42

<210> 30
<211> 73
<212> DNA
<213> Murine hybridoma LM609

<400> 30

gaagattttg cagtgtatta ctgcccaaca gagtaacagc tggcctcaca cgtttggcca 60
ggggaccaag ctg 73

<210> 31
<211> 21
<212> DNA
<213> Murine hybridoma LM609

<400> 31

aatacgactc actatagggc g

21

<210> 32

<211> 72

<212> DNA

<213> Murine hybridoma LM609

<400> 32

gaggatgttg gggtttatta ctgccaacag agtaacagct ggcctcacac gtttggccag 60

gggaccaagc tg

72

<210> 33

<211> 8

<212> PRT

<213> Murine hybridoma LM609

<400> 33

Glu Asp Phe Ala Val Tyr Tyr Cys

1

5

<210> 34

<211> 8

<212> PRT

<213> Murine hybridoma LM609

<400> 34

Glu Asp Val Gly Val Tyr Tyr Cys

1

5

<210> 35

<211> 69

<212> DNA

<213> Murine hybridoma LM609

<400> 35

gaygaggctg attattactg ccaacagagt aacagctggc ctcacacgtt cggcggaggg 60

accaagctg

69

<210> 36

<211> 50

<212> DNA

<213> Murine hybridoma LM609

<400> 36

agagagagag agagagagag cgccgtctag aattatgaac attctgtagg

50

<210> 37

<211> 7

<212> PRT

<213> Murine hybridoma LM609

<400> 37

Asp Glu Ala Asp Tyr Tyr Cys

1

5

<210> 38

<211> 7

<212> PRT

<213> Murine hybridoma LM609

<400> 38

Phe Gly Gly Gly Thr Lys Leu

1

5

<210> 39

<211> 22

<212> DNA

<213> Murine hybridoma LM609

<400> 39

aagacagcta tcgcgattgc ag

22

<210> 40

<211> 41

<212> DNA

<213> Murine hybridoma LM609

<400> 40

gaggaggagg aggaggaggc ggggcccagg cggccgagct c

41

<210> 41

<211> 21

<212> DNA

<213> Murine hybridoma LM609

<400> 41

ggccatggct ggttgggcag c

21

<210> 42

<211> 42

<212> DNA

<213> Murine hybridoma LM609

<400> 42

gcagagccca aatcttgtga cactagtggc caggccggcc ag

42

<210> 43

<211> 41

<212> DNA

<213> Murine hybridoma LM609

<400> 43

gaggaggagg aggaggagcc tggccggcct ggccactagt g

41

<210> 44

<211> 130

<212> PRT

<213> Murine hybridoma LM609

<400> 44

Leu Glu Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly Ser Leu Lys

1		5		10		15
Leu	Ser	Cys	Ala	Ala	Ser	Gly

	20		25		30
Trp	Val	Arg	Gln	Ile	Pro

	35		40		45
Ser	Ser	Gly	Gly	Gly	Ser

	50		55		60
Phe	Thr	Ile	Ser	Arg	Asp

65		70		75		80
Ser	Ser	Leu	Asn	Ser	Glu	Asp

	85		90		95
Asn	Tyr	Gly	Ser	Phe	Ala

	100		105		110
Ser	Ala	Ala	Lys	Thr	Thr

115		120		125
-----	--	-----	--	-----

Ser Ala

130

<210> 45

<211> 109

<212> PRT

<213> Murine hybridoma LM609

<400> 45

Glu Leu Val Met Thr Gln Thr Pro Ala Thr Leu Ser Val Thr Pro Gly

1		5		10		15
Asp	Ser	Val	Ser	Leu	Ser	Cys

	20		25		30
--	----	--	----	--	----

Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile

	35		40		45
--	----	--	----	--	----

Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly

	50		55		60
--	----	--	----	--	----

Ser Gly Ser Gly Thr Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Thr

65		70		75		80
----	--	----	--	----	--	----

9/10

Glu Asp Phe Gly Met Tyr Phe Cys Gln Gln Ser Asn Ser Trp Pro His

85

90

95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala

100

105

<210> 46

<211> 57

<212> DNA

<213> Murine hybridoma LM609

<220>

<221> misc_feature

<222> (26)..(36)

<223> N=A, C, G or T

<400> 46

ggccccctgg ccaaactgtg gaggmnnmnn mnnmnnctgt tggcagtaat acactgc 57

<210> 47

<211> 23

<212> DNA

<213> Murine hybridoma LM609

<400> 47

cctcaccgtt tggccagggg acc 23

<210> 48

<211> 21

<212> DNA

<213> Murine hybridoma LM609

<400> 48

agaagcgtag tccggaacgt c 21

<210> 49

<211> 109

<212> PRT

<213> Murine hybridoma LM609

<400> 49

Glu Leu Val Met Thr Gln Ser Pro Glu Phe Gln Ser Val Thr Pro Lys

1

5

10

15

Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Gly Thr Ser

20

25

30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile

35

40

45

Lys Tyr Ala Ser Gln Pro Val Phe Gly Val Pro Ser Arg Phe Arg Gly

50

55

60

10/10

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Tyr Ser Leu Glu Ala
 65 70 75 80
 Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Asn Ser Trp Pro His
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr
 100 105

<210> 50
 <211> 118
 <212> PRT
 <213> Murine hybridoma LM609

<400> 50

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Ala Ser Ile Ser Arg Gly
 20 25 30
 Gly Tyr Tyr Trp Ser Trp Ile Arg Gln Tyr Pro Gly Lys Gly Leu Glu
 35 40 45
 Trp Ile Gly Tyr Ile His His Ser Gly Ser Thr Tyr Tyr Asn Pro Ser
 50 55 60
 Leu Lys Ser Arg Val Thr Ile Ala Ile Asp Thr Ser Lys Asn Gln Leu
 65 70 75 80
 Ser Leu Arg Leu Thr Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
 85 90 95
 Cys Ala Arg His Asn Tyr Gly Ser Phe Ala Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ser
 115

<210> 51
 <211> 118
 <212> PRT
 <213> Murine hybridoma LM609

<400> 51

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15
 Thr Leu Phe Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly
 20 25 30
 Gly Tyr Tyr Trp Ser Trp Ile Arg His His Pro Gly Lys Gly Leu Glu

11/10

35 40 45
 Trp Ile Gly Tyr Ile His His Arg Ala Ala Pro Tyr Tyr Asn Pro Ser
 50 55 60
 Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Arg Asn Gln Ile
 65 70 75 80
 Ser Leu Lys Leu Arg Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
 85 90 95
 Cys Ala Arg His Asn Tyr Gly Ser Phe Ala Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ser
 115

<210> 52
 <211> 118
 <212> PRT
 <213> Murine hybridoma LM609

<400> 52

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly
 20 25 30
 Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu
 35 40 45
 Trp Ile Gly Tyr Ile His His Ser Ala Gly Thr Tyr Tyr Asn Pro Ser
 50 55 60
 Leu Lys Ser Arg Val Thr Met Ser Val Asp Thr Ser Lys Asn Gln Leu
 65 70 75 80
 Ser Leu Lys Leu Thr Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
 85 90 95
 Cys Ala Arg His Asn Tyr Gly Ser Phe Ala Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ser
 115

<210> 53
 <211> 118
 <212> PRT
 <213> Murine hybridoma LM609

<400> 53

12/10

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu

1 5 10 15
 Thr Leu Ser Leu Thr Cys Ser Val Ser Gly Gly Ser Ile Ser Ser Gly

20 25 30

Gly Tyr Tyr Trp Ser Trp Ile Arg His His Pro Gly Lys Gly Leu Glu

35 40 45

Trp Ile Gly Tyr Ile His His Ser Ala Gly Thr Tyr Tyr Asn Pro Ser

50 55 60

Leu Lys Ser Arg Val Thr Met Ser Ala Asp Thr Ser Lys Asn Gln Leu

65 70 75 80

Ser Leu Lys Leu Ala Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr

85 90 95

Cys Ala Arg His Asn Tyr Gly Ser Phe Ala Tyr Trp Gly Gln Gly Thr

100 105 110

Leu Val Thr Val Ser Ser

115

<210> 54

<211> 117

<212> PRT

<213> Murine hybridoma LM609

<400> 54

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Arg Lys Pro Gly Ser

1 5 10 15
 Ser Val Arg Val Ser Cys Lys Ala Ser Gly Gly Thr Phe Ser Gly Phe

20 25 30

Ala Val Ser Trp Val Arg Gln Ala Pro Gly Gln Arg Phe Glu Trp Leu

35 40 45

Gly Gly Ile Val Ala Ser Leu Gly Ser Thr Asp Tyr Ala Gln Lys Phe

50 55 60

Gln Asp Lys Leu Thr Ile Thr Val Asp Glu Ser Thr Ala Thr Val Tyr

65 70 75 80

Met Glu Met Arg Asn Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys

85 90 95

Ala Arg His Asn Tyr Gly Ser Phe Ala Tyr Trp Gly Gln Gly Thr Leu

100 105 110

Val Thr Val Ser Ser

115

13/10

<210> 55
 <211> 109
 <212> PRT
 <213> Murine hybridoma LM609

<400> 55

Glu Leu Val Met Thr Gln Ser Pro Glu Phe Gln Ser Val Thr Pro Lys

1 5 10 15
 Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Gly Asn Ser

20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile

35 40 45

Lys Tyr Ala Ser Gln Pro Val Phe Gly Val Pro Ser Arg Phe Arg Gly

50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu Pro

65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Asn Ser Trp Pro His

85 90 95

Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr

100 105

<210> 56
 <211> 117
 <212> PRT
 <213> Murine hybridoma LM609

<400> 56

Glu Val Gln Leu Glu Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly

1 5 10 15
 Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Ala Phe Ser Ser Tyr

20 25 30

Asp Met Ser Trp Val Arg Gln Ile Pro Glu Lys Arg Leu Glu Trp Val

35 40 45

Ala Lys Val Ser Ser Gly Gly Gly Ser Thr Tyr Tyr Leu Asp Thr Val

50 55 60

Gln Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr

65 70 75 80

Leu Gln Met Ser Ser Leu Asn Ser Glu Asp Thr Ala Met Tyr Tyr Cys

85 90 95

Ala Arg His Asn Tyr Gly Ser Phe Ala Tyr Trp Gly Gln Gly Thr Leu

14/10

100

105

110

Val Thr Val Ser Ala

115

- 27 -

WHAT IS CLAIMED IS:

1. A method of producing a humanized mouse monoclonal antibody which consists of the steps:

- (a) constructing a first human antibody heavy or light chain library consisting of antibody heavy chains or antibody light chains in which each chain in the first human antibody light or heavy chain library has three complementarity determining region (CDR) loops in which the complementarity determining region three (CDR3) loop in each light or heavy chain has the amino acid sequence of a corresponding mouse antibody heavy or light chain CDR3 loop and is flanked by unaltered human framework residues;
- (b) constructing a second human antibody heavy or light chain library consisting of antibody heavy chains or antibody light chains, in which each chain in the second human antibody heavy or light chain library has three complementarity determining region (CDR) loops in which the complementarity determining region three (CDR3) loop in each light or heavy chain has the amino acid sequence of a corresponding mouse antibody heavy or light chain CDR3 loop and is flanked by unaltered human framework residues; wherein the chains of said second human antibody heavy or light chain library are the complementary heavy or light chain of said first human antibody heavy or light chain library chains, such that one library is a light chain library and the other library is a heavy chain library;
- (c) creating a library of heavy and light chain pairs by combining chains from said first human antibody heavy or light chain library of step (a) with a complementary chain from an antibody which binds a preselected antigen forming a library of heavy and light chain pairs;
- (d) isolating antigen binding chains of step (a) by selecting from the library of step (c), a heavy and light chain pair which binds to said preselected antigen, isolating from the antigen binding heavy and light chain pair, the antigen binding chains of step (a);
- (e) creating a humanized pair library by combining isolated antigen binding chains of step (d), with chains of said library of step (b), so that a first isolated chain

- 28 -

- of step (d), together with a complementary second chain in said library of step (b) combine to form a heavy and light chain humanized pair library;
- (f) selecting from the humanized pair library of step (e) a humanized heavy and light chain pair that binds to said preselected antigen; and
- (g) combining two of said selected humanized heavy and light chain pair of step (f) to form a whole antibody;

wherein said preselected antigen is the same antigen recognized by the mouse monoclonal antibody.

2. The method of claim 1 wherein the first human antibody heavy or light chain library contains only light chains and the complementary chain is a heavy chain.

3. The method of claim 1 wherein the first human antibody heavy or light chain library contains only heavy chains and the complementary chain is a light chain.

4. The method of claim 1 wherein the heavy chain is a Fd fragment.

5. A method of producing a humanized mouse monoclonal antibody heavy and light chain pair which consists of the steps:

- (a) constructing a human light chain library wherein each light chain of said light chain library has three complementarity determining region (CDR) loops in which the complementarity determining region three (CDR3) loop in each light chain has the amino acid sequence of a corresponding mouse antibody light chain CDR3 loop and is flanked by unaltered human framework residues;
- (b) selecting a light chain from the light chain library of step (a), wherein the selection comprises combining a heavy chain from an antibody that binds a preselected antigen with a light chain from the library of step (a) to form a heavy and light chain pair library, screening said pair library for binding to said preselected antigen, and isolating light chains that bind to said preselected antigen;
- (c) constructing a library of human heavy chains wherein each heavy chain of said library of human heavy chains has three CDR loops in which the complementarity determining region three (CDR3) loop in each heavy chain

- 29 -

has the amino acid sequence of a corresponding mouse antibody heavy chain CDR3 loop and is flanked by unaltered human framework residues;

- (d) combining a heavy chain from the heavy chain library of step (c) with a selected light chain of step (b) to produce a second library of humanized mouse monoclonal antibody heavy and light chain pairs;
- (e) isolating an antigen binding humanized mouse monoclonal antibody heavy and light chain pair from the library of step (d) by screening said library of step (d) for binding with said preselected antigen and isolating those humanized mouse monoclonal antibody heavy and light chain pairs that bind said preselected antigen; wherein said preselected antigen is the same antigen recognized by the mouse monoclonal antibody.

6. The method of claim 5 further comprising converting the selected heavy and light chain pair to a whole antibody.

7. The method of claim 5 wherein the heavy chain of step (b) is a Fd fragment.

8. The method of claim 7 wherein the heavy chain Fd is a humanized mouse heavy chain fragment or a template mouse heavy chain fragment.

9. The method of claim 5 wherein in step (b) a humanized mouse heavy chain is used.

10. The method of claim 5 wherein the CDR3 loop in each light and heavy chain is a CDR3 loop from antibody LM609 produced from hybridoma cell line deposited with American Type Culture Collection under ATCC Accession No. HB 9537.

11. The method of claim 7 wherein the heavy chain Fd is a humanized mouse heavy chain.

12. The method of claim 7 wherein the heavy chain Fd is a template mouse heavy chain fragment.

1/14




FIG. 1

WO 99/29888

PCT/US98/25828

2/14

 V_1

Leu Glu Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly Ser Leu Lys Leu Ser Cys Ala Ala Ser
 5 10 15 20
 Gly Phe Ala Phe Ser Ser Tyr Asp Met Ser Trp Val Arg Gln Ile Pro Glu Lys Arg Leu Glu Trp
 25 30 35 40
 Val Ala Lys Val Ser Ser Gly Gly Gly Ser Thr Tyr Tyr Leu Asp Thr Val Gln Gly Arg Phe Thr
 45 50 55 60 65
 Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr Leu Gln Met Ser Ser Leu Asn Ser Glu Asp Thr
 70 75 80 85
 Ala Met Tyr Tyr Cys Ala Arg His Asn Tyr Gly Ser Phe Ala Tyr Trp Gly Gln Gly Thr Leu Val
 90 95 100 105 110
 Thr Val Ser Ala Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala Pro Gly Ser Ala
 115 120 125 130

FIGURE 2a

 V_k

Glu Leu Val Met Thr Gln Thr Thr Pro Ala Thr Leu Ser Val Thr Pro Gly Asp Ser Val Ser Leu Ser
 5 10 15 20
 Cys Arg Ala Ser Gln Ser Ile Ser Asn His Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro
 25 30 35 40
 Arg Leu Leu Ile Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly Ser Gly
 45 50 55 60 65
 Ser Gly Thr Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Thr Glu Asp Phe Gly Met Tyr Phe Cys
 70 75 80 85
 Gln Gln Ser Asn Ser Trp Pro His Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala
 90 95 100 105

FIGURE 2b

WO 99/29888

PCT/US98/25828

3/14

FR 1

V _K	..	Thr	Gln	Thr	Pro	Ala	Thr	Leu	Ser	Val	Thr	Pro	Gly	Asp	Ser	Val	Ser	Leu	Ser	Cys
#1	..	-	-	Ser	-	Gly	Thr	-	-	Leu	Ser	-	-	Glu	Arg	Ala	Thr	-	-	-
#2	..	-	-	Ser	-	Gly	Thr	-	-	Leu	Ser	-	-	Glu	Arg	Gly	Ser	-	-	-
#3	..	-	-	Ser	-	Gly	Thr	-	-	Leu	Ser	-	-	Glu	Arg	Ala	Thr	-	-	-
#4	..	-	-	Ser	-	Ser	Ser	-	-	Ala	Ser	Val	-	-	Arg	-	Thr	Ile	Thr	-
#5	..	-	-	Ser	-	Ser	Ser	-	-	Ala	Ser	Val	-	-	*	-	Thr	Ile	Thr	-
#6	..	-	-	Ser	-	Ser	Ser	-	-	Ala	Ser	Val	-	-	Arg	-	Thr	Ile	Thr	-

FIGURE 3a

CDR1

V _K	Arg	Ala	Ser	Gln	Ser	Ile	Ser	Asn	His	Leu	His	
#1	-	-	-	-	-	Val	-	Ser	Ser	Tyr	Leu	Ala
#2	-	-	-	-	-	Val	-	Ser	Ser	Phe	Leu	Ala
#3	-	-	-	-	-	Val	Thr	Ser	Ser	Tyr	Leu	Ala
#4	-	-	-	-	-	-	-	Thr	Phe	-	Asn	
#5	-	-	-	-	-	-	-	Ser	Tyr	-	Asn	
#6	-	-	-	-	-	-	-	Ser	Tyr	-	Asn	

FIGURE 3b

WO 99/29888

PCT/US98/25828

4/14

FR2															
V _K	Trp	Tyr	Gln	Gln	Lys	Ser	His	Glu	Ser	Pro	Arg	Leu	Leu	Ile	Lys
#1	-	-	-	-	-	Pro Gly	Gln	Ala	-	-	-	-	-	-	Tyr
#2	-	-	-	-	-	Pro Gly	Gln	Ala	-	-	-	-	-	-	Tyr
#3	-	-	-	-	-	Pro Gly	Gln	Ala	-	-	-	-	-	-	Tyr
#4	-	-	-	-	-	Pro Gly	Lys	Ala	-	-	Lys Phe	-	-	-	Tyr
#5	-	-	-	Arg	-	Pro Gly	Lys	Ala	-	-	Lys Leu	-	-	-	Tyr
#6	-	-	-	-	-	Pro Gly	Lys	Ala	-	-	Lys Leu	-	-	-	Tyr

FIGURE 3c

CDR2							
V _K	Tyr	Ala	Ser	Gln	Ser	Ile	Ser
#1	Gly	-	-	Ser-Arg	Ala-Thr		
#2	Gly	-	-	Ser-Arg	Ala-Thr		
#3	Gly	-	-	Ser-Arg	Ala-Thr		
#4	Ala	-	-	Thr	Leu	Gln	-
#5	Ala	-	-	Thr	Leu	Gln	-
#6	Ala	-	-	Thr	Leu	Gln	-

FIGURE 3d

WO 99/29888

PCT/US98/25828

5/14

FR3																						
V _x	Gly Val	Ile Glu	Pro Thr	Ser Glu	Arg Asp	Phe Phe	Ser Gly	Gly Met	Ser Tyr	Gly Phe	Ser Cys	Gly Thr	Asp Phe	Thr Leu	Ser Ile	Asn Ser						
#1	-	-	-	Asp	-	-	-	-	-	-	-	-	-	-	Thr	-	Ser Arg					
	Leu	-	Pro	-	-	-	Ala Val	-	-	Tyr	-	-	-	-	-	-						
#2	-	-	-	Asp	-	-	-	-	-	-	-	-	-	-	Thr	-	Ser Arg					
	Leu	-	Pro	-	-	-	Ala Val	-	-	Tyr	-	-	-	-	-	-						
#3	-	-	-	Asp	-	-	-	-	-	-	-	-	-	-	Ile Phe Thr	-	Ser Arg					
	Leu	-	Pro	-	-	-	Ala Val	-	-	Tyr	-	-	-	-	-	-						
#4	-	Val	-	-	-	-	-	-	-	-	-	-	-	-	Thr	-	Ser					
	Leu	Gln Pro	-	-	-	-	Ala Val	-	-	Tyr	-	-	-	-	-	-						
#5	-	Val	-	-	-	-	-	-	-	-	-	Ala	-	-	Thr	-	Ser					
	Leu	Gln Pro	-	-	-	-	Ala Val	-	-	Tyr	-	-	-	-	-	-						
#6	-	Val	-	-	-	-	-	-	-	-	-	-	-	-	Thr	-	Ser					
	Leu	Gln Pro	-	-	-	-	Ala Val	-	-	Tyr	-	-	-	-	-	-						

FIGURE 3e

WO 99/29888

PCT/US98/25828

6/14

	LCDR1	LCDR2
mouse	Arg Ala Ser Gln Ser Ile Ser Asn	His Leu His (Lys) Tyr Ala Ser Gln Ser Ile Ser
selected human		
3 x	Arg Ala Ser Gln Asp Ile Gly Thr	(Lys) Tyr Ala Ser Gln Pro Val Phe
2 x	Arg Ala Ser Gln Asp Ile Gly Asn	(Lys) Tyr Ala Ser Gln Pro Val Phe
1 x	Arg Ala Ser Gln Ser Ile Gly Trp	(Lys) Tyr Ala Ser Gln Ser Ile Ser
unselected human		
	Arg Ser Ser Gln Ser Ile Asn Ile	(Tyr) His Ala Ser Lys Arg Ala Ser
	Arg Ala Ser Gln Ser Val Ser Asn Asn	(Tyr) Arg Ala Ser Ser Arg Ala Thr
	Arg Ser Ser Gln Ser Leu Val Tyr Ser Asp Gly Asn Thr	(Tyr) Lys Val Ser Asn Arg Asp Ser
	Thr Ala Ser Gln Ser Leu Val Tyr Thr Asp Gly Asn Thr	(Tyr) Met Val Ser Asn Arg Asp Ser

FIGURE 4

WO 99/29888

PCT/US98/25828

7/14




FIG. 5a




FIG. 5b




FIG. 5c




FIG. 5d




FIG. 5e




FIG. 5f

WO 99/29888

PCT/US98/25828

8/14




FIGURE 6

WO 99/29888

PCT/US98/25828

9/14




FIGURE 7

10/14

FIGURE 8a

V_L amino acid sequences

FR1

mouse
 Glu Leu Val Met Thr Gln Thr Pro Ala Thr Leu Ser Val Thr Pro Gly Asp Ser Val Ser Leu Ser Cys
human (Group A)
 Glu Leu Val Met Thr Gln Ser Pro Glu Phe Gln Ser Val Thr Pro Lys Glu Thr Val Thr Ile Thr Cys
human (Groups BCDE)
 Glu Leu Val Met Thr Gln Ser Pro Glu Phe Gln Ser Val Thr Pro Lys Glu Thr Val Thr Ile Thr Cys

CDR1

mouse
 Arg Ala Ser Gln Ser Ile Ser Asn His Leu His
human (Group A)
 Arg Ala Ser Gln Asp Ile Gly Thr Ser Leu His
human (Groups BCDE)
 Arg Ala Ser Gln Asp Ile Gly Asn Ser Leu His

FR2

mouse
 Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile Lys
human (Group A)
 Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Lys
human (Groups BCDE)
 Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Lys

CDR2

mouse
 Tyr Ala Ser Gln Ser Ile Ser
human (Group A)
 Tyr Ala Ser Gln Pro Val Phe
human (Groups BCDE)
 Tyr Ala Ser Gln Pro Val Phe

V_L amino acid sequences

CDR3	FR4
<u>mouse</u>	<u>mouse</u>
Gln Gln Ser Asn Ser Trp Pro His Thr	Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala
<u>human (Group A)</u>	<u>human (Group A)</u>
Gln Gln Ser Asn Ser Trp Pro His Thr	Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr
<u>human (Group BCDE)</u>	<u>human (Group BCDE)</u>
Gln Gln Ser Asn Ser Trp Pro His Thr	Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg Thr

12/14

FIGURE 8c

V_H amino acid sequences

FR1

mouse
 Glu Val Gln Leu Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly Ser Leu Lys Leu Ser Cys
 Ala Ala Ser Gly Phe Ala Phe Ser
human (Group A)
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Arg Lys Pro Gly Ser Ser Val Arg Val Ser Cys
 Lys Ala Ser Gly Gly Thr Phe Ser
human (Group B)
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln Thr Leu Ser Leu Thr Cys
 Thr Val Ser Gly Ala Ser Ile Ser
human (Group C)
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln Thr Leu Phe Leu Thr Cys
 Thr Val Ser Gly Gly Ser Ile Ser
human (Group D)
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu Thr Leu Ser Leu Thr Cys
 Thr Val Ser Gly Gly Ser Ile Ser
human (Group E)
 Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu Thr Leu Ser Leu Thr Cys
 Ser Val Ser Gly Gly Ser Ile Ser

CDR1

mouse
 Ser Tyr Asp Met Ser
human (Group A)
 Gly Phe Ala Val Ser
human (Group B)
 Arg Gly Gly Tyr Tyr Trp Ser
human (Group C)
 Ser Gly Gly Tyr Tyr Trp Ser
human (Group D)
 Ser Gly Gly Tyr Tyr Trp Ser
human (Group E)
 Ser Gly Gly Tyr Tyr Trp Ser

FR2

mouse
 Trp Val Arg Gln Ile Pro Glu Lys Arg Leu Glu Trp Val Ala
human (Group A)
 Trp Val Arg Gln Ala Pro Gly Gln Arg Phe Glu Trp Leu Gly
human (Group B)
 Trp Ile Arg Gln Tyr Pro Gly Lys Gly Leu Glu Trp Ile Gly
human (Group C)
 Trp Ile Arg His His Pro Gly Lys Gly Leu Glu Trp Ile Gly
human (Group D)
 Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu Trp Ile Gly
human (Group E)
 Trp Ile Arg His His Pro Gly Lys Gly Leu Glu Trp Ile Gly

13/14

FIGURE 8d

V_H amino acid sequences

CDR2

mouse
 Lys Val Ser Ser Gly Gly Gly Ser Thr Tyr Tyr Leu Asp Thr Val Gln Gly
human (Group A)
 Gly Ile Val Ala Ser Leu Gly Ser Thr Asp Tyr Ala Gln Lys Phe Gln Asp
human (Group B)
 Tyr Ile His His Ser Gly Ser Thr Tyr Tyr Asn Pro Ser Leu Lys Ser
human (Group C)
 Tyr Ile His His Arg Ala Ala Pro Tyr Tyr Asn Pro Ser Leu Lys Ser
human (Group D)
 Tyr Ile His His Ser Ala Gly Thr Tyr Tyr Asn Pro Ser Leu Lys Ser
human (Group E)
 Tyr Ile His His Ser Ala Gly Thr Tyr Tyr Asn Pro Ser Leu Lys Ser

FR3

mouse
 Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr Leu Gln Met Ser Ser Leu Asn Ser
 Glu Asp Thr Ala Met Tyr Tyr Cys Ala Arg
human (Group A)
 Lys Leu Thr Ile Thr Val Asp Glu Ser Thr Ala Thr Val Tyr Met Glu Met Arg Asn Leu Arg Ser
 Asp Asp Thr Ala Val Tyr Tyr Cys Ala Ar
human (Group B)
 Arg Val Thr Ile Ala Ile Asp Thr Ser Lys Asn Gln Leu Ser Leu Arg Leu Thr Ser Val Thr Ala
 Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg
human (Group C)
 Arg Val Thr Ile Ser Val Asp Thr Ser Arg Asn Gln Ile Ser Leu Lys Leu Arg Ser Val Thr Ala
 Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg
human (Group D)
 Arg Val Thr Met Ser Val Asp Thr Ser Lys Asn Gln Leu Ser Leu Lys Leu Thr Ser Val Thr Ala
 Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg
human (Group E)
 Arg Val Thr Met Ser Ala Asp Thr Ser Lys Asn Gln Leu Ser Leu Lys Leu Ala Ser Val Thr Ala
 Ala Asp Thr Ala Val Tyr Tyr Cys Ala Arg

WO 99/29888

PCT/US98/25828

14/14

FIGURE 8e

 V_H amino acid sequences

CDR3	FR4
<u>mouse</u>	<u>mouse</u>
His Asn Tyr Gly Ser Phe Ala Tyr	Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
<u>human (Group A)</u>	<u>human (Group A)</u>
His Asn Tyr Gly Ser Phe Ala Tyr	Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
<u>human (Group B)</u>	<u>human (Group B)</u>
His Asn Tyr Gly Ser Phe Ala Tyr	Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
<u>human (Group C)</u>	<u>human (Group C)</u>
His Asn Tyr Gly Ser Phe Ala Tyr	Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
<u>human (Group D)</u>	<u>human (Group D)</u>
His Asn Tyr Gly Ser Phe Ala Tyr	Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
<u>human (Group E)</u>	<u>human (Group E)</u>
His Asn Tyr Gly Ser Phe Ala Tyr	Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser