



HU000035870T2

(19) **HU****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**(11) Lajstromszám: **E 035 870**(13) **T2**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 14 713844**(22) A bejelentés napja: **2014. 03. 28.**(51) Int. Cl.: **C07D487/04** (2006.01)**A61P 29/00** (2006.01)**A61K 31/4985** (2006.01)**A61P 25/00** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20140713844

(97) Az európai bejelentés közzétételi adatai:

EP 2948455 A1 **2014. 10. 02.**

(86) A nemzetközi (PCT) bejelentési szám:

PCT/EP 14/056367

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2948455 B1 **2017. 09. 27.**

(87) A nemzetközi közzétételi szám:

WO 14154895

(30) Elsőbbségi adatok:

13161863 **2013. 03. 29.** **EP****13193025** **2013. 11. 15.** **EP****14154303** **2014. 02. 07.** **EP**

(73) Jogosult(ak):

OGEDA S.A., 6041 Charleroi (FR)

(72) Feltaláló(k):

HOVEYDA, Hamid, B-1190 Bruxelles (BE)**DUTHEUIL, Guillaume, B-5020 Vedrin (BE)****FRASER, Graeme, B-1470 Bousval (BE)**

(74) Képviselő:

**Arinova Szabadalmi és Védjegy Iroda,
Budapest**

(54) **N-acil-(3- szubsztituált)-(8- szubsztituált)-5,6-dihidro-[1,2,4]triazolo[4,3-A]pirazinok mint szelektív NK-3 receptor antagonisták, gyógyászati készítmény, eljárások NK-3 receptor által közvetített rendellenességek során történő alkalmazásra**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmat az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 948 455 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
27.09.2017 Bulletin 2017/39

(51) Int Cl.:
C07D 487/04 ^(2006.01) **A61K 31/4985** ^(2006.01)
A61P 25/00 ^(2006.01) **A61P 29/00** ^(2006.01)

(21) Application number: **14713844.0**

(86) International application number:
PCT/EP2014/056367

(22) Date of filing: **28.03.2014**

(87) International publication number:
WO 2014/154895 (02.10.2014 Gazette 2014/40)

(54) **N-ACYL-(3-SUBSTITUTED)-(8-SUBSTITUTED)-5,6-DIHYDRO-
[1,2,4]TRIAZOLO[4,3-A]PYRAZINES AS SELECTIVE NK-3 RECEPTOR ANTAGONISTS,
PHARMACEUTICAL COMPOSITION, METHODS FOR USE IN NK-3 RECEPTOR-MEDIATED
DISORDERS**

N-ACYL-(3-SUBSTITUIERTE)-(8-SUBSTITUIERTE)-5,6-DIHYDRO-[1,2,4]TRIAZOLO[4,3-A]PYRAZINE ALS SELEKTIVE NK-3-REZEPTOR-ANTAGONISTEN, PHARMAZEUTISCHE ZUSAMMENSETZUNG, VERFAHREN ZUR VERWENDUNG IN NK-3-REZEPTORVERMITTELTEN ERKRANKUNGEN

N-ACYL-(3-SUBSTITUÉES)-(8-SUBSTITUÉES)-5,6-DIHYDRO-[1,2,4]TRIAZOLO[4,3-A]PYRAZINE S EN TANT QU'ANTAGONISTES SÉLECTIFS DU RÉCEPTEUR NK-3, COMPOSITION PHARMACEUTIQUE ET PROCÉDÉS POUR LEUR UTILISATION DANS LES MALADIES MÉDIÉES PAR LE RÉCEPTEUR NK-3

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **29.03.2013 EP 13161863**
15.11.2013 EP 13193025
07.02.2014 EP 14154303

(43) Date of publication of application:
02.12.2015 Bulletin 2015/49

(60) Divisional application:
17165576.4 / 3 219 715

(73) Proprietor: **OGEDA S.A.**
6041 Charleroi (FR)

(72) Inventors:
• **HOVEYDA, Hamid**
B-1190 Bruxelles (BE)
• **DUTHEUIL, Guillaume**
B-5020 Vedrin (BE)
• **FRASER, Graeme**
B-1470 Bousval (BE)

(74) Representative: **Icosa**
83 avenue Denfert-Rochereau
75014 Paris (FR)

(56) References cited:
WO-A1-2010/125102 WO-A1-2011/121137

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**FIELD OF INVENTION**

[0001] The present invention relates to novel N-acyl-(3-substituted)-(8-substituted)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazines including their pharmaceutically acceptable solvates that are selective antagonists to neurokinin-3 receptor (NK-3) and are useful as therapeutic compounds, particularly in the treatment and/or prevention of a broad array of CNS and peripheral diseases or disorders.

BACKGROUND OF INVENTION

[0002] Tachykinin receptors are the targets of a family of structurally related peptides which include substance P (SP), neurokinin A (NKA) and neurokinin B (NKB), named collectively "tachykinins". Tachykinins are synthesized in the central nervous system (CNS) and peripheral tissues, where they exert a variety of biological activities. Three tachykinin receptors are known which are named neurokinin-1 (NK-1), neurokinin-2 (NK-2) and neurokinin-3 (NK-3) receptors. Tachykinin receptors belong to the rhodopsin-like seven membrane G-protein coupled receptors. SP has the highest affinity and is believed to be the endogenous ligand of NK-1, NKA for NK-2 receptor and NKB for NK-3 receptor, although cross-reactivity amongst these ligands does exist. The NK-1, NK-2 and NK-3 receptors have been identified in different species. NK-1 and NK-2 receptors are expressed in a wide variety of peripheral tissues and NK-1 receptors are also expressed in the CNS; whereas NK-3 receptors are primarily expressed in the CNS.

[0003] The neurokinin receptors mediate a variety of tachykinin-stimulated biological effects that include transmission of excitatory neuronal signals in the CNS and periphery (e.g. pain), modulation of smooth muscle contractile activity, modulation of immune and inflammatory responses, induction of hypotensive effects via dilatation of the peripheral vasculature and stimulation of endocrine and exocrine gland secretions.

[0004] In the CNS, the NK-3 receptor is expressed in regions including the medial prefrontal cortex, the hippocampus, the thalamus and the amygdala. Moreover, NK-3 receptors are expressed on dopaminergic neurons. Activation of NK-3 receptors has been shown to modulate dopamine, acetylcholine and serotonin release suggesting a therapeutic utility for NK-3 receptor modulators for the treatment of a variety of disorders including psychotic disorders, anxiety, depression, schizophrenia as well as obesity, pain or inflammation (Giardina et al., *Exp. Opin. Ther. Patents*, 2000, 10(6), 939-960; *Current Opinion in Investigational Drugs*, 2001, 2(7), 950-956 and Dawson and Smith, *Current Pharmaceutical Design*, 2010, 16, 344-357).

[0005] Schizophrenia is classified into subgroups. The paranoid type is characterized by delusions and hallucinations and absence of thought disorder, disorganized behavior, and affective flattening. In the disorganized type, which is also named 'hebephrenic schizophrenia' in the International Classification of Diseases (ICD), thought disorder and flat affect are present together. In the catatonic type, prominent psychomotor disturbances are evident, and symptoms may include catatonic stupor and waxy flexibility. In the undifferentiated type, psychotic symptoms are present but the criteria for paranoid, disorganized, or catatonic types have not been met. The symptoms of schizophrenia normally manifest themselves in three broad categories, i.e. positive, negative and cognitive symptoms. Positive symptoms are those, which represent an "excess" of normal experiences, such as hallucinations and delusions. Negative symptoms are those where the patient suffers from a lack of normal experiences, such as anhedonia and lack of social interaction. The cognitive symptoms relate to cognitive impairment in schizophrenics, such as a lack of sustained attention and deficits in decision making. The current antipsychotic drugs (APDs) are fairly successful in treating the positive symptoms but fare less well for the negative and cognitive symptoms. Contrary to that, NK-3 antagonists have been shown clinically to improve on both positive and negative symptoms in schizophrenics (Meltzer et al, *Am. J. Psychiatry*, 2004, 161, 975-984) and ameliorate cognitive behavior of schizophrenics (*Curr. Opin. Invest. Drug*, 2005, 6, 717-721).

[0006] In rat, morphological studies provide evidence for putative interactions between NKB neurons and the hypothalamic reproductive axis (Krajewski et al, *J. Comp. Neurol.*, 2005, 489(3), 372-386). In arcuate nucleus neurons, NKB expression co-localizes with estrogen receptor α and dynorphin, implicated in progesterone feedback to Gonadotropin Releasing Hormone (GnRH) secretion (Burke et al., *J. Comp. Neurol.*, 2006, 498(5), 712-726; Goodman et al., *Endocrinology*, 2004, 145(6), 2959-2967). Moreover, NK-3 receptor is highly expressed in the hypothalamic arcuate nucleus in neurons which are involved in the regulation of GnRH release.

[0007] WO 00/43008 discloses a method of suppressing gonadotropin and/or androgen production with specific NK-3 receptor antagonists. More particularly, the WO 00/43008 application relates to lowering luteinizing hormone (LH) blood level by administering an NK-3 receptor antagonist. Concurrently or alternatively with gonadotropin suppression, WO 00/43008 also relates to suppression of androgen production with NK-3 receptor antagonists. Recently it has been postulated that NKB acts autocrinally on kisspeptin neurons in the arcuate nucleus to synchronize and shape the pulsatile secretion of kisspeptin and drive the release of GnRH from fibers in the median eminence (Navarro et al., *J. of Neuroscience*, 2009, 23(38), 11859-11866). All these observations suggest a therapeutic utility for NK-3 receptor mod-

ulators for sex hormone-dependent diseases.

[0008] NK-3 receptors are also found in the human myenteric and submucosal plexus of the sigmoid colon as well as in the gastric fundus (Dass et al., Gastroenterol., 2002, 122 (Suppl 1), Abstract M1033) with particular expression noted on myenteric intrinsic primary afferent neurons (IPANs) (Lomax and Furness, Cell Tissue Res, 2000, 302, 59-3). Intense stimulation of IPANs changes patterns of intestinal motility and intestinal sensitivity. Electrophysiology experiments have shown that activation of the NK-3 receptor changes the voltage threshold of action potentials in IPANs and promotes the generation of long-lasting plateau potentials (Copel et al., J Physiol, 2009, 587, 1461-1479) that may sensitize these neurons to mechanical and chemical stimuli leading to effects on gut motility and secretion. Similarly, Irritable Bowel Syndrome (IBS) is characterized by patient hypersensitivity to mechanical and chemical stimuli. Thus, NK-3 antagonists have been tested in preclinical models of IBS where they have been shown to be effective to reduce nociceptive behavior caused by colo-rectal distension (Fioramonti et al., Neurogastroenterol Motil, 2003, 15, 363-369; Shafon et al., Neurogastroenterol Motil, 2004, 16, 223-231) and, on this basis, NK-3 antagonists have been advanced into clinical development for the treatment of IBS (Houghton et al., Neurogastroenterol Motil, 2007, 19, 732-743; Dukes et al., Gastroenterol, 2007, 132, A60).

[0009] Non-peptide antagonists have been developed for each of the tachykinin receptors. Some of them have been described as dual modulators able to modulate both NK-2 and NK-3 receptors (WO 06/120478). However, known non-peptide NK-3 receptor antagonists suffer from a number of drawbacks, notably poor safety profile and limited CNS penetrability that may limit the success of these compounds in clinical development.


[0010] On this basis, new potent and selective antagonists of NK-3 receptor may be of therapeutic value for the preparation of drugs useful in the treatment and/or prevention of CNS and peripheral diseases or disorders in which NKB and the NK-3 receptors are involved.

[0011] Target potency alone, which may be demonstrated by competitive binding data, is not sufficient for drug development. Rather, efficacy *in vivo* is contingent upon achieving a relevant "free" drug concentration relative to the target potency at the physiological site of action. Drug molecules typically bind reversibly to proteins and lipids in plasma. The "free" fraction refers to the drug concentration that is unbound and therefore available to engage the biological target and elicit pharmacological activity. This free fraction is commonly determined using plasma protein binding (PPB) assays. The free drug fraction is relevant to not only achieving the desired pharmacological activity, but also potentially undesirable activities including rapid hepatic metabolism (leading to high first-pass clearance and thereby poor oral bioavailability) as well as possible off-target activities that can lead to safety concerns (for example, inhibition of hERG ion channel activity, a widely accepted marker of cardiovascular toxicity).

[0012] The invention thus encompasses compounds of general Formula I, their pharmaceutically acceptable solvates as well as methods of use of such compounds or compositions comprising such compounds as antagonists to the NK-3 receptor. Compounds of Formula I are *N*-acyl-(3-substituted)-(8-substituted)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazines. The compounds of the invention are generally disclosed in international patent application WO2011/121137 but none is specifically exemplified therein. On another hand, unsubstituted and thus non-chiral 5,6,7,8-tetrahydro[1,2,4]triazolo[4,3-*a*]pyrazines have been disclosed in WO2010/125102 as modulators of an unrelated target, namely P2X7.

SUMMARY

[0013] In a general aspect, the invention provides compounds of general Formula I:



and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or, nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

X^1 is N and X^2 is S or O; or X^1 is S and X^2 is N;

— represents a single or a double bond depending on X^1 and X^2 ;

* — stands for the (*R*)-enantiomer or for the racemate of compound of Formula I.

[0014] In another aspect, the present invention provides a pharmaceutical composition comprising at least one compound according to the invention or a pharmaceutically acceptable solvate thereof.

[0015] The invention also relates to the use of the above compounds or their pharmaceutically acceptable solvates as modulators of NK-3 receptors, preferably as antagonists of NK-3 receptors.

[0016] The invention also relates to the use of the above compounds or their pharmaceutically acceptable solvates as lowering agents of the circulating LH levels.

[0017] The invention further provides compounds for use in methods of treatment and/or prevention of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis comprising the administration of a therapeutically effective amount of a compound or pharmaceutically acceptable solvate of Formula I, to a patient in need thereof. The invention further provides methods of treatment and/or prevention of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis comprising the administration of a therapeutically effective amount of a compound or pharmaceutically acceptable solvate of Formula I, to a patient in need thereof. Preferably the patient is a warm-blooded animal, more preferably a human.

[0018] The invention further provides compounds for use in methods of treatment for gynecological disorders and infertility. In particular, the invention provides methods to lower and/or suppress the LH-surge in assisted conception comprising the administration of a therapeutically effective amount of a compound or pharmaceutically acceptable solvate of Formula I, to a patient in need thereof. Preferably the patient is a warm-blooded animal, more preferably a woman.


[0019] The invention further provides compounds for use in methods to affect androgen production to cause male castration and to inhibit the sex drive in male sexual offenders comprising the administration of a therapeutically effective amount of a compound or pharmaceutically acceptable solvate of Formula I, to a patient in need thereof. Preferably the patient is a warm-blooded animal, more preferably a man.

[0020] The invention also provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof as a medicament. Preferably, the medicament is used for the treatment and/or prevention of depression, anxiety,

psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. Preferably, the medicament is used for the treatment and/or prevention of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. The medicament may also be used for the treatment of gynecologic disorders, infertility and to affect androgen production to cause male castration.

DETAILED DESCRIPTION

[0021] As noted above, the invention relates to compounds of Formula I:



and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;

R^{2'} is H or F;

R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁴ is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

R⁵ is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably **R⁵** is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

X¹ is N and **X²** is S or O; or **X¹** is S and **X²** is N;

== represents a single or a double bond depending on **X¹** and **X²**;

* stands for the (*R*)-enantiomer or for the racemate of compound of Formula I.

[0022] In one specific embodiment of the invention, **R⁵** is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl. In another specific embodiment, **R⁵** is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl. In another specific embodiment, **R⁵** is 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl.

[0023] Preferred compounds of Formula I and pharmaceutically acceptable solvates thereof are those wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;

R^{2'} is H or F;

R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁴ is methyl, ethyl, *n*-propyl or hydroxyethyl;


R⁵ is methyl, ethyl, trifluoromethyl, difluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably

R⁵ is methyl, ethyl or trifluoromethyl;

X¹ is N and **X²** is S or O, preferably **X¹** is N and **X²** is S.


[0024] In an embodiment of the invention, compound of Formula I is the (*R*)-enantiomer. In another embodiment, compound of Formula I is the racemate.

[0025] In one embodiment, preferred compounds of Formula I are those of Formula I':




and pharmaceutically acceptable solvates thereof, wherein **R¹**, **R^{1'}**, **R²**, **R^{2'}**, **R³**, **R⁴**, **R⁵**, **X¹** and **X²** are as defined in Formula I and --- represents a single or a double bond depending on **X¹** and **X²**.

[0026] In one embodiment, preferred compounds of Formula I are those of Formula I'':



and pharmaceutically acceptable solvates thereof, wherein **R¹**, **R^{1'}**, **R²**, **R^{2'}**, **R³**, **R⁴**, **R⁵**, **X¹** and **X²** are as defined in Formula I and --- represents a single or a double bond depending on **X¹** and **X²**.

[0027] In one embodiment, preferred compounds of Formula I are those of Formula Ia:



and pharmaceutically acceptable solvates thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably

R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;


R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or

2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl,


preferably R^5 is methyl, ethyl, trifluoromethyl or difluoromethyl, preferably R^5 is methyl, ethyl or trifluoromethyl;

* _ _ _ stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia.

[0028] In one embodiment, preferred compounds of Formula Ia are those of Formula Ia' and Formula Ia'':



Ia'



Ia''

and pharmaceutically acceptable solvates thereof, wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , R^4 and R^5 are as defined in Formula Ia.

[0029] Preferred compounds of Formula Ia' and Ia'' and pharmaceutically acceptable solvates thereof are those wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;


$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;

R^5 is methyl, ethyl, trifluoromethyl or difluoromethyl, preferably R^5 is methyl, ethyl or trifluoromethyl.

[0030] In one embodiment, preferred compounds of Formula Ia are those of Formula Ia-1:



and pharmaceutically acceptable solvates thereof, wherein:

R³ is H, F, Cl, methyl, trifluoromethyl or nitrile, preferably **R³** is H, F or Cl;


R⁴ is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably **R⁴** is methyl, ethyl, *n*-propyl or hydroxyethyl;

R⁵ is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably

R⁵ is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably **R⁵** is methyl, ethyl, trifluoromethyl or difluoromethyl, preferably **R⁵** is methyl, ethyl or trifluoromethyl;


* _ _ stands for the (*R*)-enantiomer or for the racemate.

[0031] In one embodiment, preferred compounds of Formula Ia-1 are those of Formula Ia-1':



and pharmaceutically acceptable solvates thereof, wherein **R³**, **R⁴** and **R⁵** are as defined in Formula Ia-1.

[0032] In one embodiment, preferred compounds of Formula Ia are those of Formula Ia-2:



and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;


$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is methyl, ethyl, trifluoromethyl or difluoromethyl, preferably R^5 is methyl, ethyl or trifluoromethyl;


* _ _ stands for the (*R*)-enantiomer or for the racemate.

[0033] In one embodiment, preferred compounds of Formula Ia-2 are those of Formula Ia-2':



and pharmaceutically acceptable solvates thereof, wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 and R^5 are as defined in Formula Ia-2.

[0034] In one embodiment, preferred compounds of Formula Ia are those of Formula Ia-3:



and pharmaceutically acceptable solvates thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;


R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably

R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;


* _ _ stands for the (*R*)-enantiomer or for the racemate.

[0035] In one embodiment, preferred compounds of Formula Ia-3 are those of Formula Ia-3':



and pharmaceutically acceptable solvates thereof, wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 and R^4 are as defined in Formula Ia-3.

[0036] In one embodiment, preferred compounds of Formula I are those of Formula Ib:




and pharmaceutically acceptable solvates thereof, wherein:

R^3 is F;

R^5 is methyl, ethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is methyl, ethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl or ethyl;


* _ _ _ stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ib.

[0037] In one embodiment, preferred compounds of Formula Ib are those of Formula Ib':




and pharmaceutically acceptable solvates thereof, wherein R^3 and R^5 are defined as in Formula Ib.

[0038] In one embodiment, preferred compounds of Formula Ib are those of Formula Ib'':




and pharmaceutically acceptable solvates thereof, wherein R^3 and R^5 are defined as in Formula Ib.

[0039] In one embodiment, preferred compounds of Formula Ib are those of Formula Ib-1:




and pharmaceutically acceptable solvates thereof, wherein R^5 is methyl, ethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is methyl or ethyl.

[0040] In one embodiment, preferred compounds of Formula Ib are those of Formula Ib-2:



and pharmaceutically acceptable solvates thereof, wherein R^5 is ethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is ethyl.

[0041] In one embodiment, preferred compounds of Formula I are those of Formula Ic:



and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;

R^{2'} is H or F;


R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁴ is methyl, ethyl, *n*-propyl or hydroxyethyl;

R⁵ is methyl, ethyl or trifluoromethyl;

* _ _ stands for the (*R*)-enantiomer or for the racemate.

[0042] In one embodiment, preferred compounds of Formula Ic are those of Formula Ic':



and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl, preferably **R¹** is H;

R^{1'} is H;

R² is H, F, Cl or methoxy, preferably **R²** is H;

R^{2'} is H or F, preferably **R^{2'}** is H;




R³ is H, F, Cl, methyl, trifluoromethyl or nitrile, preferably **R³** is F;

R⁴ is methyl, ethyl, *n*-propyl or hydroxyethyl, preferably **R⁴** is methyl;



R⁵ is methyl, ethyl or trifluoromethyl, preferably **R⁵** is methyl.

[0043] Particularly preferred compounds of Formula I of the invention are those listed in Table 1 hereafter.



TABLE 1

Cpd n°	Structure	Chemical name	MW
1		(R)-3-(3-(4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	409.29
2		(R)-3-(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	372.42
3		(R)-3-(3-(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	374.85



(continued)

Cpd n°	Structure	Chemical name	MW
4		(R)-(4-chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	392.84
5		(R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	358.39



(continued)

Cpd n°	Structure	Chemical name	MW
6		(R)-(3-chloro-4-fluorophenyl)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	392.84
7		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)-(3,4,5-trifluorophenyl)methanone	394.37



(continued)

Cpd n°	Structure	Chemical name	MW
8		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanone	394.37
9		(R)-(3,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	376.38



(continued)

Cpd n°	Structure	Chemical name	MW
10		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanone	412.36
11		(R)-(4-fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	388.42



(continued)

Cpd n°	Structure	Chemical name	MW
12		(4-fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	388.42
13		(R)-(3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	356.35



(continued)

Cpd n°	Structure	Chemical name	MW
14		(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	358.39
15		(R)-(3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	358.39



(continued)

Cpd n°	Structure	Chemical name	MW
16		(R)-(3-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	374.85
17		(R)-(3,5-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	376.38



(continued)

Cpd n°	Structure	Chemical name	MW
18		(R)-(2,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	376.38
19		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(p-tolyl)methanone	354.43



(continued)

Cpd n°	Structure	Chemical name	MW
20		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-1,2,4-triazolo[4,3-a]pyrazin-7(8H)-yl)(phenyl)methanone	340.4
21		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-1,2,4-triazolo[4,3-a]pyrazin-7(8H)-yl)(4-(trifluoromethyl)phenyl)methanone	408.4



(continued)

Cpd n°	Structure	Chemical name	MW
22		(R)-(8-ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	372.42
23		(8-ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	372.42




(continued)

Cpd n°	Structure	Chemical name	MW
24		(R)-(4-fluorophenyl)(3-(3-methyl-1,2,4-thiadiazol-5-yl)-8-propyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	386.45
25		(R)-(4-fluoro-3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	388.42



(continued)

Cpd n°	Structure	Chemical name	MW
26		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(o-tolyl)methanone	354.43
27		(R)-(3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	370.43



(continued)

Cpd n°	Structure	Chemical name	MW
28		(R)-4-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	342.33
29		(R)-4-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazine-7-carbonyl)benzonitrile	365.41
31		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanone	412.36

(continued)



Cpd n°	Structure	Chemical name	MW
32		(3,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	376.38
33		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanone	394.37

(continued)

Cpd n°	Structure	Chemical name	MW
34		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanone	394.37
35		(3-chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	392.84

Cpd n°	Structure	Chemical name	MW
36		(4-chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	392.84
37		(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	374.85
38		(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	409.29



(continued)

Cpd n°	Structure	Chemical name	MW
39		(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	372.42
40		(3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	356.35


(continued)

Cpd n°	Structure	Chemical name	MW
41		(R)-(4-fluorophenyl)(8-methyl-3-(3-(trifluoromethyl)-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	412.36
42		(R)-(3-(3-(difluoromethyl)-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	394.37

(continued)

Cpd n°	Structure	Chemical name	MW
43		(R)-3-(3-(1,1-difluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	392.34
44		(R)-3-(3-(2,2,2-trifluoroethyl)-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	410.33

(continued)

Cpd n°	Structure	Chemical name	MW
45		((8R)-3-(3-(1-fluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	374.34


and pharmaceutically acceptable solvates thereof.

[0044] In Table 1, the term "Cpd" means compound.

[0045] The compounds of Table 1 were named using ChemBioDraw® Ultra version 12.0 (PerkinElmer).

[0046] The compounds of Formula I can be prepared by different ways with reactions known to a person skilled in the art.

[0047] The invention further relates to a process of manufacturing of compounds of Formula I:



and pharmaceutically acceptable solvates thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably

R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is methyl, ethyl or trifluoromethyl;


X^1 is N and X^2 is S or O; or X^1 is S and X^2 is N, preferably X^1 is N and X^2 is S or O, more preferably, X^1 is N and X^2 is S;

== represents a single or a double bond depending on X^1 and X^2 ;

* _ _ stands for the (*R*)-enantiomer or for the racemate of compound of Formula I;

characterized in that it comprises the following steps:

a) reacting a compound of Formula (i)



wherein:


PG represents a suitable protecting group such as for example DMB, PMB, Boc, allyl, diphenyl-phosphinamide (DPP), 2-trimethylsilyl ethanesulfonyl (SES), preferably **PG** is DMB;

$R^{4'}$ is R^4 as defined above or a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl, such as for example $-CH_2CO_2Alkyl$; where the term "reducible precursor of hydroxyethyl or consequently a further precursor of methoxyethyl" refers to any chemical group which, when reacting with reducing agents, such as for example $LiAlH_4$, is reduced to hydroxyethyl and then optionally further converted

to methoxyethyl;

* --- stands for the (*R*)-enantiomer or for the racemate;

with a compound of Formula (ii)



(ii)


wherein:

R^{5'} is R⁵ as defined above, H or 1-((tert-butyldiphenylsilyl)oxy)ethyl, preferably R^{5'} is R⁵ as defined above or H;

X¹ and X² are as defined above; and

== represents a single or a double bond depending on X¹ and X²;


so as to obtain a compound of Formula (iii)



(iii)

wherein PG, R^{4'}, R^{5'}, X¹ and X² are as defined above, *--- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on X¹ and X²;


b) deprotecting compound of Formula (iii) with a suitable deprotection agent to afford compound of Formula (iv)



(iv)


wherein R^4 , R^5 , X^1 and X^2 are as defined above, *--- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on X^1 and X^2 ;

c) when R^5 is H, introducing a trifluoromethyl or difluoromethyl group by direct C-H trifluoro- or difluoromethylation, leading to compound of Formula (v)




wherein R^4 , X^1 and X^2 are as defined above and R^5 is trifluoromethyl or * difluoromethyl, *--- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on X^1 and X^2 ;

d) N-acylating compound of Formula (iv) wherein R^5 is not H or compound of Formula (v), with a compound of Formula (vi)



wherein R^1 , R^1 , R^2 , R^2 and R^3 are as defined above; leading to compound of Formula (vii)



wherein R^1 , R^1 , R^2 , R^2 , R^3 , R^4 , X^1 and X^2 are as defined above, *--- stands for the (*R*)-enantiomer or for the racemate, == represents a single or a double bond depending on X^1 and X^2 ; and $R^{5''}$ is R^5 as defined in Formula I or 1-((tert-butyl)diphenylsilyl)oxy)ethyl;

e) optionally further conducting one or both of the two following steps:

e') when $R^{4'}$ is a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl, a step of reduction optionally followed by methyl ether formation;

e'') when R^5 is 1-((tert-butyldiphenylsilyl)oxy)ethyl, a step of alcohol deprotection and subsequent fluorination to form 1-fluoroethyl R^5 group; or a step of alcohol deprotection, followed by an oxidation step and a subsequent fluorination step to afford 1,1-difluoroethyl R^5 group;

to afford compound of Formula I.

[0048] In a preferred embodiment, the protecting group PG used in the process of the invention is DMB.

[0049] According to one embodiment, the introduction of a trifluoromethyl or difluoromethyl group at step c) may be performed by direct C-H trifluoro- or difluoromethylation as described by Ji Y. et al. in PNAS, 2011, 108(35), 14411-14415 or by Fujiwara Y. et al. in JACS, 2012, 134, 1494-1497.

[0050] According to one embodiment, the fluorination step to form 1-fluoroethyl or 1,1-difluoroethyl R^5 groups at step e'') may be performed by DAST fluorination. DAST fluorination may be performed as described in WO2004/103953, page 51.

[0051] Reaction schemes as described in the example section are illustrative only and should not be construed as limiting the invention in any way. According to one embodiment, compounds of Formula I can be prepared using the chiral synthesis of the invention detailed in the examples below.

[0052] The invention is further directed to the use of the compounds of the invention or pharmaceutically acceptable solvates thereof as antagonists to the NK-3 receptor.

[0053] Accordingly, in a particularly preferred embodiment, the invention relates to the use of compounds of Formula I and subformulae in particular those of Table 1 above, or pharmaceutically acceptable solvates thereof, as NK-3 receptor antagonists.

[0054] Accordingly, in another aspect, the invention relates to the use of these compounds or solvates thereof for the synthesis of pharmaceutical active ingredients, such as selective NK-3 receptor antagonists.

USES

[0055] The compounds of the invention are therefore useful as medicaments, in particular in the prevention and/or treatment of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. The compounds of the invention are therefore useful as medicaments, in particular in the prevention and/or treatment of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis.

[0056] The invention also provides for a method for delaying in patient the onset of depression, anxiety, psychosis,

schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis comprising the administration of a pharmaceutically effective amount of a compound of Formula I or pharmaceutically acceptable solvate thereof to a patient in need thereof. The invention also provides for a method for delaying in patient the onset of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis comprising the administration of a pharmaceutically effective amount of a compound of Formula I or pharmaceutically acceptable solvate thereof to a patient in need thereof.

[0057] Preferably, the patient is a warm-blooded animal, more preferably a human.

[0058] The compounds of the invention are especially useful in the treatment and/or prevention of sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. The compounds of the invention are especially useful in the treatment and/or prevention of sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis.

[0059] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of benign prostatic hyperplasia (BPH), endometriosis, uterine fibrosis, uterine fibroid tumor, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of benign prostatic hyperplasia (BPH), endometriosis, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of

luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis.

[0060] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of endometriosis, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, polycystic ovary syndrome (PCOS) and benign prostatic hyperplasia (BPH).

[0061] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of endometriosis.

[0062] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of uterine fibrosis.

[0063] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of uterine fibroid tumor.

[0064] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of uterine leiomyoma.

[0065] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of polycystic ovary syndrome (PCOS).

[0066] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of benign prostatic hyperplasia (BPH).

[0067] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of hot flashes also known as hot flushes.

[0068] In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of peri-menopausal conditions (i.e. 'hot flashes'), in vitro fertilization ('IVF'), male contraceptive, female contraceptive, castration of sex offenders.

[0069] The compounds of the invention are also useful in the treatment of gynecological disorders and infertility. In particular, the invention provides methods to lower and/or suppress the LH-surge in assisted conception.

[0070] The compounds of the invention are also useful to cause male castration and to inhibit the sex drive in men. This is of particular interest in the treatment of male sexual offenders.

[0071] The invention further provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament for treating and/or preventing depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis in a patient. The invention further provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament for treating and/or preventing depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis in a patient.

[0072] Preferably, the patient is a warm-blooded animal, more preferably a human.

[0073] The invention especially provides the use of a compound of Formula I or a pharmaceutically acceptable solvate

thereof for the manufacture of a medicament to treat and/or prevent sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. The invention especially provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament to treat and/or prevent sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis.

[0074] In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent endometriosis, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, polycystic ovary syndrome (PCOS) and benign prostatic hyperplasia (BPH).

[0075] In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent endometriosis.

[0076] In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent uterine fibrosis.

[0077] In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent uterine fibroid tumor.

[0078] In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent uterine leiomyoma.

[0079] In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent polycystic ovary syndrome (PCOS).

[0080] In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent benign prostatic hyperplasia (BPH).

[0081] In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent hot flashes also known as hot flushes.

[0082] The invention further provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament to lower and/or suppress the LH-surge in assisted conception in a patient. Preferably the patient is a warm-blooded animal, more preferably a woman.

[0083] The invention further provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament to cause male castration and to inhibit the sex drive in men. This is of particular interest in the treatment of male sexual offenders.

[0084] According to a further feature of the present invention there is provided a method for modulating NK-3 receptor activity, in a patient, preferably a warm blooded animal, and even more preferably a human, in need of such treatment, which comprises administering to said patient an effective amount of compound of the present invention, or a pharmaceutically acceptable solvate thereof.

[0085] According to one embodiment, the compounds of the invention, their pharmaceutical acceptable solvates may be administered as part of a combination therapy. Thus, are included within the scope of the present invention embodiments comprising coadministration of, and compositions and medicaments which contain, in addition to a compound of the present invention, a pharmaceutically acceptable solvate thereof as active ingredient, additional therapeutic agents and/or active ingredients. Such multiple drug regimens, often referred to as "combination therapy", may be used in the treatment and/or prevention of any of the diseases or conditions mediated by or associated with NK-3 receptor modulation. The use of such combinations of therapeutic agents is especially pertinent with respect to the treatment of the above-mentioned disorders within a patient in need of treatment or one at risk of becoming such a patient.

[0086] In addition to the requirement of therapeutic efficacy, which may necessitate the use of active agents in addition to the NK-3 receptor modulator compounds of Formula I or pharmaceutical acceptable solvates thereof, there may be additional rationales which compel or highly recommend the use of combinations of drugs involving active ingredients which represent adjunct therapy, i.e., which complement and supplement the function performed by the NK-3 receptor modulator compounds of the present invention. Suitable supplementary therapeutic agents used for the purpose of

auxiliary treatment include drugs which, instead of directly treating or preventing a disease or condition mediated by or associated with NK-3 receptor modulation, treat diseases or conditions which directly result from or indirectly accompany the basic or underlying NK-3 receptor modulated disease or condition.

[0087] According to a further feature of the present invention, the compound of Formula I, a pharmaceutically acceptable solvate thereof may be used in combination therapy with antipsychotic drugs (APD), to improve the efficacy and to minimize secondary effects associated to APD including but not limited to Dopamine 2/3 and 5-HT₂ receptors antagonists. More particular the compound of Formula I, a pharmaceutically acceptable solvate thereof may be used as an adjunct therapy in combination with an atypical antipsychotic drug, including but not limited to risperidone, clozapine, olanzapine, where the NK-3 receptor modulator may serve a role as dose-limiting for the atypical antipsychotic and therefore spare the patient from some of the side effect of those atypical antipsychotic drugs.

[0088] Thus, the methods of treatment and pharmaceutical compositions of the present invention may employ the compounds of Formula I or pharmaceutical acceptable solvates thereof in the form of monotherapy, but said methods and compositions may also be used in the form of multiple therapy in which one or more compounds of Formula I or their pharmaceutically acceptable solvates are coadministered in combination with one or more other therapeutic agents.

[0089] In the above-described embodiment combinations of the present invention, the compound of Formula I, a pharmaceutically acceptable solvate thereof and other therapeutic active agents may be administered in terms of dosage forms either separately or in conjunction with each other, and in terms of their time of administration, either serially or simultaneously. Thus, the administration of one component agent may be prior to, concurrent with, or subsequent to the administration of the other component agent(s).

[0090] The invention also provides pharmaceutical compositions comprising a compound of Formula I or a pharmaceutically acceptable solvate thereof and at least one pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant. As indicated above, the invention also covers pharmaceutical compositions which contain, in addition to a compound of the present invention, a pharmaceutically acceptable solvate thereof as active ingredient, additional therapeutic agents and/or active ingredients.

[0091] Another object of this invention is a medicament comprising at least one compound of the invention, or a pharmaceutically acceptable solvate thereof, as active ingredient.

[0092] According to a further feature of the present invention there is provided the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament for modulating NK-3 receptor activity in a patient, in need of such treatment, which comprises administering to said patient an effective amount of compound of the present invention, or a pharmaceutically acceptable solvate thereof.

[0093] Preferably, the patient is a warm-blooded animal, more preferably a human.

[0094] As set forth above, the compounds of the invention, their pharmaceutically acceptable solvates may be used in monotherapy or in combination therapy. Thus, according to one embodiment, the invention provides the use of a compound of the invention for the manufacture of a medicament for at least one of the purposes described above, wherein said medicament is administered to a patient in need thereof, preferably a warm-blooded animal, and even more preferably a human, in combination with at least one additional therapeutic agent and/or active ingredient. The benefits and advantages of such a multiple drug regimen, possible administration regimens as well as suitable additional therapeutic agents and/or active ingredients are those described above.

[0095] Generally, for pharmaceutical use, the compounds of the invention may be formulated as a pharmaceutical preparation comprising at least one compound of the invention and at least one pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant, and optionally one or more further pharmaceutically active compounds.

[0096] By means of non-limiting examples, such a formulation may be in a form suitable for oral administration, for parenteral administration (such as by intravenous, intramuscular or subcutaneous injection or intravenous infusion), for topical administration (including ocular), for administration by inhalation, by a skin patch, by an implant, by a suppository, etc. Such suitable administration forms - which may be solid, semi-solid or liquid, depending on the manner of administration - as well as methods and carriers, diluents and excipients for use in the preparation thereof, will be clear to the skilled person; reference is made to the latest edition of Remington's Pharmaceutical Sciences.

[0097] Some preferred, but non-limiting examples of such preparations include tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, ointments, cremes, lotions, soft and hard gelatin capsules, suppositories, drops, sterile injectable solutions and sterile packaged powders (which are usually reconstituted prior to use) for administration as a bolus and/or for continuous administration, which may be formulated with carriers, excipients, and diluents that are suitable per se for such formulations, such as lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, polyethylene glycol, cellulose, (sterile) water, methylcellulose, methyl- and propylhydroxybenzoates, talc, magnesium stearate, edible oils, vegetable oils and mineral oils or suitable mixtures thereof. The formulations can optionally contain other substances that are commonly used in pharmaceutical formulations, such as lubricating agents, wetting agents, emulsifying and suspending agents, dispersing agents, desintegrants, bulking agents, fillers, preserving agents, sweetening agents, flavoring agents, flow regulators, release agents, etc.. The com-

positions may also be formulated so as to provide rapid, sustained or delayed release of the active compound(s) contained therein.

[0098] The pharmaceutical preparations of the invention are preferably in a unit dosage form, and may be suitably packaged, for example in a box, blister, vial, bottle, sachet, ampoule or in any other suitable single-dose or multi-dose holder or container (which may be properly labeled); optionally with one or more leaflets containing product information and/or instructions for use. Generally, such unit dosages will contain between 0.05 and 1000 mg, and usually between 1 and 500 mg, preferably between 2 and 150 mg of at least one compound of the invention, e.g. about 2, 4, 8, 16, 32, 64 or 128 mg per unit dosage. According to another embodiment, such unit dosages will contain between 0.05 and 1000 mg, and usually between 1 and 500 mg, preferably between 2 and 400 mg, preferably between 2 and 200 mg of at least one compound of the invention per unit dosage.

[0099] Usually, depending on the condition to be prevented or treated and the route of administration, the active compound of the invention will usually be administered between 0.001 and 10 mg per kilogram body weight, more often between 0.01 and 4 mg per kilogram body weight, preferably between 0.02 and 1.5 mg per kilogram body weight, for example about 0.02, 0.04, 0.08, 0.16, 0.32, 0.64 or 1.28 mg, per kilogram body weight of the patient per day, which may be administered as a single daily dose, divided over one or more daily doses, or essentially continuously, e.g. using a drip infusion. According to another embodiment, the active compound of the invention will usually be administered between 0.001 and 10 mg per kilogram body weight, more often between 0.01 and 7 mg per kilogram body weight, preferably between 0.03 and 3.5 mg per kilogram body weight of the patient per day, which may be administered as a single daily dose, divided over one or more daily doses, or essentially continuously, e.g. using a drip infusion.

[0100] According to one embodiment, the active compound of the invention will be administered as a single daily dose, divided over one, two or more daily doses, or essentially continuously, e.g. using a drip infusion.


DEFINITIONS

[0101] The definitions and explanations below are for the terms as used throughout the entire application, including both the specification and the claims.

[0102] When describing the compounds of the invention, the terms used are to be construed in accordance with the following definitions, unless indicated otherwise.

[0103] The term "alkyl" refers to a hydrocarbyl radical of formula C_nH_{2n+1} wherein n is a number greater than or equal to 1. Generally, alkyl groups of this invention comprise from 1 to 4 carbon atoms, preferably from 1 to 3 carbon atoms. Alkyl groups may be linear or branched. Suitable alkyl groups include but are not limited to methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl and *t*-butyl.

[0104] The ring atoms of (3-substituted)-(8-substituted)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazines of the invention are numbered based on scheme below.




[0105] Bonds from an asymmetric carbon in compounds are generally depicted using a solid line (—), a solid wedge (▲), or a dotted wedge (▤). The use of either a solid or dotted wedge to depict bonds from an asymmetric carbon atom is meant to indicate that only the stereoisomer shown is meant to be included.

[0106] The compounds of Formula I and subformulae thereof contain a stereogenic carbon center at position 8 and thus may exist as (*R*)- and (*S*)-enantiomers. In an embodiment of the invention, compounds of Formula I are not pure (*S*)-enantiomers relative to the C8 position.


[0107] In the compounds of the invention, a dotted wedge (▤) carrying a substituent at the C8 position is used to depict the (*R*)-enantiomer, thus excluding racemic mixtures thereof.

[0108] In the compounds of the invention, a dotted line with a star next to the C8 position (* - -) is used to depict either a dotted wedge (▤) to represent the (*R*)-enantiomer or a solid line (—) to depict the racemic mixture of (*R*)- and (*S*)-enantiomer, which is called "racemate".

[0109] For instance, (*R*)-(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone (compound n°1) is depicted as:



[0110] The racemic mixture of this compound, (3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone (compound n°38) is depicted as:



[0111] The term "solvate" is used herein to describe a compound in this invention that contains stoichiometric or sub-stoichiometric amounts of one or more pharmaceutically acceptable solvent molecule such as ethanol. The term "hydrate" refers to when the said solvent is water.

[0112] All references to compounds of Formula I include references to solvates, multicomponent complexes and liquid crystals thereof.

[0113] The compounds of the invention include compounds of Formula I as hereinbefore defined, including all polymorphs and crystal habits thereof, prodrugs and prodrugs thereof and isotopically- labeled compounds of Formula I.

[0114] The invention also generally covers all pharmaceutically acceptable prodrugs and prodrugs of the compounds of Formula I.

[0115] The term "prodrug" as used herein means the pharmacologically acceptable derivatives of compounds of Formula I, such as for example esters, whose *in vivo* biotransformation product generates the biologically active drug. Prodrugs are generally characterized by increased bio-availability and are readily metabolized into biologically active compounds *in vivo*.

[0116] The term "prodrug", as used herein, means any compound that will be modified to form a drug species, wherein the modification may take place either inside or outside of the body, and either before or after the prodrug reaches the area of the body where administration of the drug is indicated.

[0117] The term "patient" refers to a warm-blooded animal, more preferably a human, who/which is awaiting the receipt of, or is receiving medical care or is/will be the object of a medical procedure.

[0118] The term "human" refers to a subject of both genders and at any stage of development (i.e. neonate, infant, juvenile, adolescent, adult).

[0119] The terms "treat", "treating" and "treatment, as used herein, are meant to include alleviating, attenuating or abrogating a condition or disease and/or its attendant symptoms.

[0120] The terms "prevent", "preventing" and "prevention", as used herein, refer to a method of delaying or precluding the onset of a condition or disease and/or its attendant symptoms, barring a patient from acquiring a condition or disease, or reducing a patient's risk of acquiring a condition or disease.

[0121] The term "therapeutically effective amount" (or more simply an "effective amount") as used herein means the

amount of active agent or active ingredient (e.g. NK-3 antagonist) that is sufficient to achieve the desired therapeutic or prophylactic effect in the patient to which/whom it is administered.

[0122] The term "administration", or a variant thereof (e.g. "administering"), means providing the active agent or active ingredient (e.g. a NK-3 antagonist), alone or as part of a pharmaceutically acceptable composition, to the patient in whom/which the condition, symptom, or disease is to be treated or prevented.

[0123] By "pharmaceutically acceptable" is meant that the ingredients of a pharmaceutical composition are compatible with each other and not deleterious to the patient thereof.

[0124] The term "antagonist" as used herein means a compound that competitively or non-competitively binds to a receptor at the same site as an agonist (for example, the endogenous ligand) and has reversible and competitive binding affinity to a receptor without direct modulation of receptor signaling, but that nonetheless occupies the binding site of an agonist (for example, the endogenous ligand) to thereby block agonist-mediated receptor signaling.

[0125] The term "sex hormone-dependent disease" as used herein means a disease which is exacerbated by, or caused by, excessive, inappropriate or unregulated sex hormone production and/or an extraordinary physiological response to sex hormones. Examples of such diseases in men include but are not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, androgen dependent acne, male pattern baldness and precocious puberty in boys. Examples of such diseases in women include but are not limited to endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers (ovarian cancer, breast cancer), androgen-producing tumor (virilizing ovarian or adrenal tumor), hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), menorrhagia and adenomyosis (abnormal endometrial growth within the muscle of the uterus).

[0126] The term "Psychotic disorders" as used herein means a group of illnesses that affect the mind. These illnesses alter a patient's ability to think clearly, make good judgments, respond emotionally, communicate effectively, understand reality, and behave appropriately. When symptoms are severe, patient with psychotic disorders have difficulty staying in touch with reality and are often unable to meet the ordinary demands of daily life. Psychotic disorders include but are not limited to, schizophrenia, schizophreniform disorder, schizo-affective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, psychotic disorder due to a general medical condition, substance-induced psychotic disorder or psychotic disorders not otherwise specified (Diagnostic and Statistical Manual of Mental Disorders, Ed. 4th, American Psychiatric Association, Washington, D.C. 1994).

[0127] The term "pharmaceutical vehicle" as used herein means a carrier or inert medium used as solvent or diluent in which the pharmaceutically active agent is formulated and/or administered. Non-limiting examples of pharmaceutical vehicles include creams, gels, lotions, solutions, and liposomes.

[0128] The present invention will be better understood with reference to the following examples. These examples are intended to be representative of specific embodiments of the invention, and are not intended as limiting the scope of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0129]

Figure 1 is a graph showing the plasma testosterone levels over time in intact male rats after oral administration of compound n°5 (3 mg/kg) or of a vehicle (0.5% methyl cellulose).

Figure 2 is a histogram showing the prostate weight in a rat model of Benign Prostate Hyperplasia (BPH) after oral administration of 3, 10 or 30 mg/kg of compound n°5.

Figure 3 is a histogram showing the estradiol levels in adult, female rats tracked over the duration of consecutive estrous cycles, after oral administration of compound n°5 (10 mg/kg) or of a vehicle (0.5% methyl cellulose).

EXAMPLES

CHEMISTRY EXAMPLES

[0130] All reported temperatures are expressed in degrees Celsius (°C); all reactions were carried out at room temperature (RT) unless otherwise stated.

[0131] All reactions were followed by thin layer chromatography (TLC) analysis (TLC plates, silica gel 60 F₂₅₄, Merck)

was used to monitor reactions, establish silica-gel flash chromatography conditions. All other TLC developing agents/visualization techniques, experimental set-up or purification procedures that were used in this invention, when not described in specific details, are assumed to be known to those conversant in the art and are described in such standard reference manuals as: i) Gordon, A. J.; Ford, R. A. "The Chemist's Companion - A Handbook of Practical Data, Techniques, and References", Wiley: New York, 1972; ii) Vogel's Textbook of Practical Organic Chemistry, Pearson Prentice Hall: London, 1989.

[0132] HPLC-MS spectra were typically obtained on an Agilent LCMS using electrospray ionization (ESI). The Agilent instrument includes an autosampler 1100, a binary pump 1100, an ultraviolet multi-wavelength detector 1100 and a 6100 single-quad mass-spectrometer. The chromatography column used was Sunfire 3.5 μ m, C18, 3.0 x 50 mm in dimensions.

[0133] Eluent typically used was a mixture of solution A (0.1% TFA in H₂O) and solution B (0.1 % TFA in MeCN).

[0134] Gradient was applied at a flow rate of 1.3 mL per minute as follows: gradient A (for analysis of final compounds and intermediates): held the initial conditions of 5% solution B for 0.2 min, increased linearly to 95% solution B in 6 min, held at 95% during 1.75 min, returned to initial conditions in 0.25 min and maintained for 2.0 min; gradient B (for analysis of crude samples and reactions mixtures): held the initial conditions of 5% solution B for 0.2 min, increased linearly to 95% in 2.0 min, held at 95% during 1.75 min, returned to initial conditions in 0.25 min and maintained for 2 min.

[0135] Determination of chiral purity was made using chiral HPLC that was performed on an Agilent 1100 (binary pump and a ultraviolet multi wavelength detector) with manual or automatic (Autosampler 1100) injection capabilities. Column used is CHIRALPAK IA 5 μ m, 4.6 x 250 mm 4.6 x 250 mm in isocratic mode. Choice of eluent was predicated on the specifics of each separation. Further details concerning the chiral HPLC methods used are provided below.

[0136] Method A: column CHIRALPAK IA 5 μ m, 4.6 x 250 mm, eluent: EtOAc plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm; column at RT, eluent was used as sample solvent.

[0137] Method B: column CHIRALPAK IA 5 μ m, 4.6 x 250 mm, eluent: EtOAc/hexane (50:50) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm; column at RT, eluent was used as sample solvent.

[0138] Method C: column CHIRALPAK IA 5 μ m 4.6 x 250mm, eluent: hexane/ethanol (80:20 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0139] Method D: column CHIRALPAK IA 5 μ m 4.6 x 250mm, eluent: hexane/ethanol (50:50 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0140] Method E: column CHIRALPAK ID 5 μ m 4.6 x 250mm, eluent: hexane/ethanol (80:20 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0141] Method F: column CHIRALPAK IA 5 μ m 4.6 x 250mm, eluent: DCM/ethanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0142] Method G: column CHIRALPAK IA 5 μ m 4.6 x 250mm, eluent: DCM/ethanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0143] Method H: column CHIRALPAK IB 5 μ m 4.6 x 250mm, eluent: TBME plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0144] Method I: column CHIRALPAK IC 5 μ m 4.6 x 250mm, eluent: TBME/ethanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0145] Method J: column CHIRALPAK ID 5 μ m 4.6 x 250mm, eluent: EtOAc/DCM/IPAethanol (3:1:1 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0146] Method K: column CHIRALPAK IC 5 μ m 4.6 x 250mm, eluent: TBME/methanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0147] Method L: column CHIRALPAK IB 5 μ m 4.6 x 250mm, eluent: TBME/methanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

[0148] Preparative HPLC purifications were typically carried out on an Agilent 1200 instrument (preparative pump 1200 and ultraviolet multi wavelength detector 1200) with manual injection. The chromatography column used was Waters Sunfire 5 μ m, C18, 19 x 100 mm, or XBridge 5 μ m, C18, 19 x 100mm depending on the type of eluent system employed, i.e. low pH or high pH conditions.

[0149] For high-pH HPLC purifications, eluent typically consisted of a mixture of solution A (0.04 M ammonium bicarbonate in H₂O plus 0.1% of conc. NH₄OH) and solution B was MeCN. The gradient was adapted depending on the impurity profile in each sample purified, thereby allowing sufficient separation between the impurities and the desired compound.

[0150] In rare cases when high-pH HPLC purification did not provide sufficient purity, low-pH HPLC was applied. For low-pH HPLC purifications, eluent typically consisted of a mixture of solution A (0.1% of TFA in H₂O) and solution B was MeCN. The gradient was adapted depending on the impurity profile in each sample purified, thereby allowing sufficient separation between the impurities and the desired compound. TFA was removed from evaporated fractions by liquid-liquid extraction.

[0151] Chiral preparative HPLC purifications were performed on an Agilent 1200 instrument (preparative pump 1200

and ultraviolet multi wavelength detector 1200) with manual injection. The chiral columns used are CHIRALPAK IA 5 μm , 20 x 250 mm or CHIRALPAK IA 5 μm , 10 x 250 mm. All chiral HPLC methods were employed in an isocratic mode. The eluent mixture was selected based on the analytical chiral HPLC experiment (see above) that provided the best chiral separation.

[0152] ^1H (300 MHz), ^{19}F (282 MHz) and ^{13}C NMR (75 MHz) spectra were recorded on a Bruker Avance DRX 300 instrument. Chemical shifts are expressed in parts per million, (ppm, δ units). Coupling constants are expressed in Hertz (Hz). Abbreviations for multiplicities observed in NMR spectra are as follows: s (singlet), d (doublet), t (triplet), q (quadruplet), m (multiplet), br (broad).

[0153] Solvents, reagents and starting materials were purchased and used as received from commercial vendors unless otherwise specified.

[0154] The following abbreviations are used:


Boc: *tert*-Butoxycarbonyl,
 Cpd: Compound,
 DAST: (Diethylamino)sulfur trifluoride,
 DCM: Dichloromethane,
 DEA: Diethylamine,
 DMB: 2,4-Dimethoxybenzyl,
 DMB-CHO: 2,4-Dimethoxybenzaldehyde,
 DPP: Diphenylphosphinamide,
 ee: Enantiomeric excess,
 eq.: Equivalent(s),
 EtOAc: Ethyl acetate,
 EtOH: Ethanol,
 g: Gram(s),
 h: Hour(s),
 IPA: *iso*-Propylalcohol,
 L: Liter(s),
 MeOH: Methanol,
 μL : Microliter(s),
 mg: Milligram(s),
 mL: Milliliter(s),
 mmol: Millimole(s),
 min: Minute(s),
 P: UV purity at 254 nm or 215 nm determined by HPLC-MS,
 PMB: 4-Methoxybenzyl,
 rt: Room temperature,
 SES: 2-Trimethylsilylethanesulfonyl,
*t*Bu: *tert*-Butyl,
 TBDPS: *tert*-Butyldiphenylsilyl,
 TBME: *tert*-Butyl methyl ether,
 TFA: Trifluoroacetic acid,
 TLC: Thin layer chromatography.

[0155] The intermediates and compounds described below were named using ChemBioDraw[®] Ultra version 12.0 (PerkinElmer).

I. Racemic synthesis

I.1. General Synthetic Scheme for racemic synthesis

[0156] Compounds of the invention may be synthesized using the methodology described in Scheme 1, which represents the racemic product synthesis. The racemic products may then be subjected to chiral HPLC for chiral separation.



Scheme 1: General racemic synthetic scheme for the preparation of the compounds of the invention.

[0157] The general synthetic scheme comprises the following steps:

Step 1: DMB-protected ketopiperazine **1.1** was converted to iminoether **1** by using the Meerwein reagent (Et_3OBF_4).


[0158] Step 2: Ester **2.2** was subsequently converted to acyl hydrazide **2**. Ester **2.2** may be obtained by esterification of acid **2.1**.

[0159] Step 3: Cyclodehydration between the acyl hydrazide **2** and the iminoether **1** furnished the protected triazolopiperazine **3.1**. Thereafter, **3.1** was subjected to acidolytic deprotection to obtain **3**. When applicable, R^5 was introduced from $\text{R}^{5'}$ affording **3'**.

[0160] Step 4: The thus obtained triazolopiperazine intermediate **3** (or **3'**) was acylated through reaction with the appropriate acid chloride **4.1** to obtain the racemic final target structure represented by the general Formula **4**. Optionally, R^4 may be transformed, for example by reduction of $\text{R}^{4'}$ when $\text{R}^{4'}$ contains a reducible group such as an ester group. The chiral compound **4'** was subsequently obtained by purification using preparative chiral HPLC.

I.2. Sten 1: Protection and conversion to iminoether 1**Method A: Conversion of DMB-protected ketopiperazine 1.1 to iminoether 1**


[0161] Method A is the procedure used for the synthesis of the iminoether intermediates **1** with a DMB protecting group and is detailed below:

**Scheme 2: Conversion to iminoether 1.**

[0162] Method A is illustrated by the synthesis of intermediate **1a** wherein **R^{4'}** is Me.

Synthesis of 1-(2,4-dimethoxybenzyl)-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine 1a


[0163]

**Scheme 3: Synthesis of 1-(2,4-dimethoxybenzyl)-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine 1a.**

[0164] Oven-dried (115°C) sodium carbonate (18.6 g, 98 mmol, 2.25 eq.) was placed in a 500 mL round-bottom flask. The round-bottom flask was backfilled with Ar and then capped with a rubber septum. A solution of 4-(2,4-dimethoxybenzyl)-3-methylpiperazin-2-one **1.1a** (20.6 g, 78 mmol, 1 eq.) in anhydrous DCM (250 mL) was added, followed by triethyloxonium tetrafluoroborate (18.6g, 98 mmol, 1.25 eq.) in one portion. Thereafter, the reaction mixture was stirred further at RT for 1 h whereupon the reaction mixture was diluted with water (250 mL). The aqueous layer was extracted with DCM (3 x 150 mL). The organic layers were combined, dried over MgSO₄, filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel (EtOAc) to afford the desired product **1a** as orange oil. Yield: 13.2 g, 58 %. LCMS: P = 93 %, retention time = 1.8 min, (M+H+H₂O)⁺: 311; ¹H-NMR (CDCl₃): δ 7.23 (d, J= 8.8 Hz, 1H), 6.48 (d, J= 8.8 Hz, 1H), 6.44 (s, 1H), 4.02 (m, 2H), 3.92 (s, 3H), 3.91 (s, 3H), 3.86 (d, J_{AB}= 14.0 Hz, 1H), 3.46 (d, J_{AB}= 14.0 Hz, 1H), 3.44 (m, 2H), 3.10 (m, 1H), 2.79 (m, 1H), 2.32 (m, 1H), 1.35 (d, J= 6.8 Hz, 3H), 1.24 (t, J= 6.0 Hz, 3H).

I.3. Sten 2: Formation of acyl hydrazide 2**Method B: Acyl hydrazide2**

[0165] Method B is the procedure used for the synthesis of the acyl hydrazides **2** and is detailed below:




Scheme 4: Formation of acylhydrazide 2.

[0166] In a round-bottom flask equipped with a condenser, ester **2.2** (1 eq.) is dissolved in anhydrous EtOH and treated with hydrazine hydrate (1.2 to 20 eq., preferably 1.5 to 10 eq.) using a temperature range from RT to reflux. After allowing the reaction mixture to come to RT, the solution is concentrated under reduced pressure. Co-evaporations using a mixture of commercial DCM:MeOH (1:1) may be performed to remove residual water. The residue is then recrystallized and/or precipitated or purified on a pad of silica to afford **2**.

I.4. Step 3: Cyclodehydration leading to triazolonerazine 3

Method C: Cyclodehydration and acyolysis

[0167] Method C is the procedure used for the synthesis of the triazolopiperazine **3** and is detailed below:



Scheme 5: Cyclodehydration leading to triazolopiperazine 3.

[0168] Step 1: In a round-bottom flask equipped with a condenser, imino-ether **1** (1 eq.) is dissolved in anhydrous MeOH, to which is added **2** (1 eq.) in one portion. The resulting solution is stirred at reflux overnight. Thereafter, the reaction mixture is brought to RT and the volatiles are removed under reduced pressure. The crude compound is then purified using silica gel chromatography to afford the desired product **3.1**.

[0169] Step 2: In a round-bottom flask containing DCM is added **3.1** (1 eq.). Then, TFA (5 to 75 eq.), is added to the reaction mixture at RT. After 30 min stirring, the mixture is concentrated. Then DCM is added to the residue thus obtained, and washed with saturated NaHCO_3 . The aqueous layer is extracted twice with DCM, the organic layers are washed with brine, dried over MgSO_4 , filtered and concentrated under reduced pressure to obtain crude **3**. The crude **3** may be directly used in the next step without further purification.


[0170] In one embodiment, alternative work-up equally used involves treatment of the dried residue obtained above with 4 M HCl/dioxane (20 eq.) at RT under stirring. After 5 min, Et_2O is added to help precipitation. This precipitate is filtered off under vacuum, washed with Et_2O and dried under high vacuum to furnish **3** as hydrochloride salt.

[0171] In another embodiment, HCl could be used for **Step 2**: HCl 4M solution in 1,4-dioxane (3 to 20 eq.) is added in one portion to a solution of **3.1** (1 eq.) in commercial isopropanol or ethanol. The reaction mixture is stirred at 60°C . After complete conversion monitored by HPLC-MS (1 to 10 h), the reaction mixture is allowed to cool to room temperature and then further cooled to 0°C with an ice bath. Thereupon, Et_2O is added. After 15-30 min stirring, the precipitate is filtered and dried *in vacuo* to afford **3** as hydrochloride salt.

[0172] Remark: When $\text{R}^{5'} \neq \text{R}^5 = \text{H}$, introduction of groups such as trifluoro- or difluoromethyl through direct trifluoromethylation or direct difluoromethylation (Ji Y. et al., PNAS, 2011, 108(35), 14411-14415; Fujiwara Y. et al., JACS, 2012, 134, 1494-1497) may be performed.

I.5. Step 4: Acylation leading to final produces**Method D: Acylation and chiral HPLC purification**

[0173] Method D is the procedure used for the synthesis of the racemic product **4** and its purification to obtain (R)-enantiomer **4'** compounds of general Formula I. Method D is detailed below:



Scheme 6: Acylation and chiral HPLC purification.


[0174] To a solution of crude **3** (1 eq.) in anhydrous DCM are added, at RT, **4.1** (1.17 to 1.3 eq.), followed by N-methylmorpholine (1 eq. to 3.5 eq.) dropwise over 15 sec. The reaction mixture is stirred at RT for 1 to 30 minutes and the milky suspension is poured into 1 M HCl solution or directly diluted with DCM. The aqueous phase is extracted with DCM. The organic phases are combined, optionally washed with 1 M NaOH, water, brine, dried over $MgSO_4$ and evaporated to dryness. The residue is solubilized in DCM and Et_2O and is slowly added to induce precipitation. The solid was filtered off, washed with Et_2O and dried under vacuum to afford **4**. Alternatively, the residue is preliminary purified on silica gel before precipitation or purified on silica gel only. Substituent R^4 may then be transformed, when applicable, into R^4 . One example of such transformation is illustrated by the synthesis of compound **12** wherein R^4 is hydroxyethyl group, obtained by reduction of $R^{4'} = -CH_2CO_2Alkyl$.

[0175] To a solution of **4** (1 eq.) in anhydrous THF is added, at $-40^\circ C$, LAH (1 eq.). The reaction mixture is stirred at $-40^\circ C$ for 5 to 30 minutes and the mixture is quenched with 1 M NaOH solution. The resulting mixture is extracted with DCM twice. The organic phases are combined, dried over $MgSO_4$ and evaporated to dryness. The residue **4''** is then purified on silica gel.

[0176] Compound **4** or **4''** may be purified by chiral preparative HPLC according to the abovementioned method to yield the corresponding chiral(R)-compound **4'**. Compounds **4**, **4''** and **4'** are compounds of Formula I of the invention.

II. Chiral synthesis**II.1. General Synthetic Scheme for chiral synthesis**

[0177] Chiral compounds of the invention may be synthesized using the chiral process of the invention described in Scheme 7.



Scheme 7: General synthetic scheme for the preparation of chiral compounds of the invention.

[0178] Chiral ketopiperazine **B** was protected with "PG" protecting group leading to PG-protected chiral ketopiperazine **C**. PG-protected chiral ketopiperazine **C** was converted to iminoether **D** by using the Meerwein reagent (Et_3OBF_4). Condensation reaction between the acyl hydrazide **E** and iminoether **D** was conducted under heating conditions in methanol to provide PG-protected piperazine **F** that was subsequently deprotected to yield compound of Formula **G**.

[0179] In one embodiment, when the protecting group PG is DMB, the DMB group deprotection step (from **F** to **G**) is carried out using TFA in DCM at rt, followed by either TFA salt exchange with HCl or extraction at high pH recovering free piperazine **G**.

[0180] When applicable, R^5 was introduced from $\text{R}^{5'}$ of **G**, affording **G'**.


[0181] Acylation of **G** or **G'** with the appropriate acid chloride **H** afforded the (R)-enantiomer of **I** typically in > 90% enantiomeric excess (chiral HPLC). When applicable, $\text{R}^{4'}$ of **I** was then modified to afford R^4 , furnishing **I'**.

[0182] When applicable, $\text{R}^{5'}$ of **I** or **I'** was then modified to afford R^5 , furnishing **I''** or **I'''** respectively.

II.2. Step 1: Protection of ketopiperazine **B**

II.2.1. Protection of ketopiperazine **B** with an allyl to afford protected ketopiperazine **C**₁

[0183]

Scheme 8: Alkyl protection of **B**.

[0184] Alkyl protection is illustrated by the synthesis of intermediate (*R*)-4-allyl-3-methylpiperazin-2-one (i.e. compound **C₁** wherein **R^{4'}** is Me).


[0185] To a solution of (*R*)-3-methylpiperazin-2-one (0.5 g, 4.38 mmol) in commercial anhydrous THF (44 mL) at rt was added K_2CO_3 (1.2 g, 8.76 mmol). 3-bromoprop-1-ene (0.41 mL, 4.82 mmol) was then added at once, and the reaction mixture was stirred under reflux for 14h.

[0186] The reaction mixture was allowed to cool to rt, concentrated and the residue was then solubilized with water (10 mL) and DCM (10 mL). The organic layer was separated, dried over MgSO_4 , filtered and concentrated to afford 460 mg of yellow oil. $^1\text{H-NMR}$ analysis shows that desired product was clearly the main product. Crude was used as-is in the following step.

LCMS: P > 90 %, retention time = 0.2 min, $(\text{M}+\text{H})^+$: 155. $^1\text{H-NMR}$ (CDCl_3): δ 6.2 (m, 1H), 5.8 (m, 1H), 5.3 (m, 2H), 3.4 (m, 3H), 3.3 (q, $J = 7.2 \text{ Hz}$, 1H), 3.1 (m, 2H), 2.6 (m, 1H).

II.2.2. Protection of ketopiperazine **B** with DPP to afford protected ketopiperazine **C₂**

[0187]

Scheme 9: DPP protection of **B**.


[0188] DPP protection is illustrated by the synthesis of intermediate (*R*)-4-(diphenylphosphoryl)-3-methylpiperazin-2-one (i.e. compound **C₂** wherein **R^{4'}** is Me).

[0189] To a solution of (*R*)-3-methylpiperazin-2-one (0.5 g, 4.38 mmol) in commercial anhydrous DCM (9 mL) under Ar atmosphere at rt was added diphenylphosphinic chloride (0.84 mL, 4.38 mmol) in one portion, followed by N-methylmorpholine (1.2 mL, 8.76 mmol) dropwise. The reaction mixture was stirred under reflux for 72h.

[0190] The reaction mixture was concentrated and the crude compound was then purified on silica gel (DCM/MeOH 99/1) to afford the desired product as colorless oil. Yield: 0.54 g, 88 %. LCMS: P = 98 %, retention time = 2.0 min, $(\text{M}+\text{H})^+$: 315; chiral HPLC retention time = 26.7 min, ee = 99.4 %; $^1\text{H-NMR}$ (CDCl_3): δ 7.9 (m, 4H), 7.5 (m, 6H), 6.1 (bs, 1H), 3.9 (m, 1H), 3.6 (m, 1H), 3.3 (m, 2H), 3.2 (m, 1H), 1.5 (m, 3H).

II.2.3. Protection of ketopiperazine **B** with Boc to afford protected ketopiperazine **C₃**

[0191]

Scheme 10: Boc protection of **B**.


[0192] Boc protection is illustrated by the synthesis of intermediate (*R*)-tert-butyl 2-methyl-3-oxopiperazine-1-carboxylate (i.e. compound **C₃** wherein **R^{4'}** is Me).

[0193] To a solution of (*R*)-3-methylpiperazin-2-one (0.33 g, 2.87 mmol) in commercial anhydrous DCM (10 mL) at 0°C was added Boc₂O (0.77 mL, 3.30 mmol) in one portion. The reaction mixture was allowed to reach rt and stirred for 1h.

[0194] The reaction mixture was concentrated under reduced pressure and the residue was taken up in DCM (100 mL) and washed with HCl 0.5M (90 mL) and brine (120 mL), dried over MgSO₄, filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel (DCM/MeOH 99/1) to afford the desired product as colorless oil. Yield: 0.45 g, 33 %. LCMS: P = 98 %, retention time = 1.9 min, (M+H)⁺: 215; ¹H-NMR (CDCl₃): δ 6.3 (bs, 1H), 4.6 (m, 1H), 4.1 (m, 1H), 3.5 (m, 1H), 3.3-3.1 (m, 2H), 1.5 (m, 3H), 1.4 (s, 9H).

II.2.4. Protection of ketopiperazine **B** with SES to afford protected ketopiperazine **C₄**

[0195]

Scheme 11: SES protection of **B**.

[0196] SES protection is illustrated by the synthesis of intermediate (*R*)-3-methyl-4-((2-(trimethylsilyl)ethyl)sulfonyl)piperazin-2-one (i.e. compound **C₄** wherein **R^{4'}** is Me).


[0197] To a solution of (*R*)-3-methylpiperazin-2-one (0.25 g, 2.19 mmol) in commercial anhydrous DCM (4.5 mL) under Ar atmosphere at rt was added 2-(trimethylsilyl)ethanesulfonyl chloride (0.44 mL, 2.30 mmol) in one portion, followed by N-methylmorpholine (0.45 mL g, 4.38 mmol) dropwise. The reaction mixture was stirred at rt for 16h.

[0198] The reaction mixture was diluted with water (10 mL) and DCM (10 mL). The organic layer was separated, dried over MgSO₄, filtered and concentrated. The crude compound was then purified on silica gel (DCM/MeOH 99/1) to afford the desired product as colorless oil. Yield: 0.08 g, 13 %. LCMS: P = 95 %, retention time = 2.1 min, (M+H)⁺: 279; chiral HPLC retention time = 7.2 min, ee = 99.6 %; ¹H-NMR (CDCl₃): δ 6.1 (bs, 1H), 4.5 (m, 1H), 3.8 (m, 1H), 3.6 (m, 1H), 3.4 (m, 1H), 3.3 (m, 1H), 2.9 (m, 2H), 1.6 (m, 3H), 1.0 (m, 2H), 0.1 (s, 9H).

II.3. Step 2: Conversion to iminoether **D**

Method E: Conversion to iminoether

[0199] General Method E is the procedure used for the synthesis of intermediates **D**.

Scheme 12: Conversion to iminoether **D**.

[0200] Method E is illustrated by the synthesis of intermediate (*R*)-1-(2,4-dimethoxybenzyl)-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine **D₅-1** (i.e. compound **D** wherein **PG** is DMB and **R^{4'}** is Me). The corresponding DMB-protected keto-piperazine C₅ is commercially available.

[0201] Oven dried (115°C) sodium carbonate (2.48 g, 23.40 mmol, 2.25 eq.) was placed in a round-bottom flask. The round-bottom flask was backfilled with Ar and then capped with a rubber septum. A solution of (*R*)-4-(2,4-dimethoxybenzyl)-3-methylpiperazin-2-one **C-1** (2.75 g, 10.40 mmol, 1 eq.) in anhydrous DCM (35 mL) was added, followed by freshly prepared triethyloxonium tetrafluoroborate (2.48 g, 13.05 mmol, 1.25 eq.) in one portion. Thereafter the reaction mixture was stirred further at rt for 45 min to 1 hour, whereupon the reaction mixture was diluted with saturated aqueous NaHCO₃ (100 mL). The aqueous layer was extracted with DCM (3 x 200 mL). The organic layers were combined, dried over MgSO₄, filtered and concentrated under reduced pressure to afford 3.1 g of yellow oil. The crude compound was then purified on silica gel (EtOAc/MeOH: 99/1) to afford the desired product **D-1** as a pale yellow oil. Yield: 1.44 g, 48 %.

LCMS: P = 95 %, retention time = 1.8 min, (M+H₂O+H)⁺: 311; chiral HPLC retention time = 12.3 min, ee > 97 %. **¹H-NMR** (CDCl₃): δ 7.23 (d, *J* = 8.8, 1H), 6.48 (d, *J* = 8.8, 1H), 6.44 (s, 1H), 4.02 (m, 2H), 3.92 (s, 6H), 3.86 (d, *J*_{AB} = 14.0, 1H), 3.46 (d, *J*_{AB} = 14.0, 1H), 3.44 (m, 2H), 3.10 (m, 1H), 2.79 (m, 1H), 2.32 (m, 1H), 1.35 (d, *J* = 6.8, 3H), 1.24 (t, *J* = 6.0, 3H).

[0202] The reaction mixture may alternatively be treated with brine. After stirring for about 20 min, additional water and DCM were added leading to phase separation. The organic layers were then dried over MgSO₄, filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel.

[0203] Method E is further illustrated by the synthesis of intermediate (*R*)-1-allyl-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine (i.e. compound **D₁** wherein **PG** is allyl and **R^{4'}** is Me).

[0204] To a solution of (*R*)-4-allyl-3-methylpiperazin-2-one (0.35 g, 2.27 mmol, 1 eq.) in DCM (7.6 mL) at 0°C was added sodium carbonate (0.54 g, 5.11 mmol, 2.25 eq.) in one portion, followed by commercial triethyloxonium tetrafluoroborate (0.54 g, 2.84 mmol, 1.25 eq.) in one portion. Thereafter the reaction mixture was stirred further at rt for 45 min, whereupon the reaction mixture was diluted with DCM (10 mL) and brine (10 mL). The layers were separated and the aqueous layer was further extracted with DCM (2 x 5 mL). The organic layers were combined, dried over MgSO₄, filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel (EtOAc) to afford the desired product as colorless oil. Yield: 0.19 g, 46 %. LCMS: P = 95 %, retention time = 1.5 min, (M+H)⁺: 183; **¹H-NMR** (CDCl₃): δ 5.9 (m, 1H), 5.2 (m, 2H), 4.0 (m, 2H), 3.5 (m, 2H), 3.3 (m, 1H), 3.1-3.0 (m, 2H), 2.8 (m, 1H), 2.4 (m, 1H), 1.3 (m, 6H).

[0205] The following intermediates were also prepared from the *ad hoc* reagents:

(*R*)-(3-ethoxy-2-methyl-5,6-dihydropyrazin-1(2H)-yl)diphenylphosphine oxide in 44 % yield. LCMS: P = 98 %, retention time = 2.0 min, (M+H₂O+H)⁺: 361; chiral HPLC retention time = 4.8 min, ee = 99.4 %; **¹H-NMR** (CDCl₃): δ 7.9 (m, 4H), 7.5 (m, 6H), 4.0 (m, 2H), 3.7 (m, 1H), 3.6 (m, 1H), 3.5 (m, 1H), 3.1 (m, 2H), 1.4 (m, 3H), 1.2 (m, 3H).


[0206] (*R*)-tert-butyl 3-ethoxy-2-methyl-5,6-dihydropyrazine-1(2H)-carboxylate in 68% yield. LCMS: P = 98 %, retention time = 1.8 min, (M+H₂O+H)⁺: 261; **¹H-NMR** (CDCl₃): δ 4.3 (m, 1H), 4.1 (m, 2H), 3.9 (m, 1H), 3.5 (m, 2H), 2.9 (m, 1H), 1.5 (s, 9H), 1.3 (d, *J* = 6.9 Hz, 3H), 1.2 (t, *J* = 7.0 Hz, 3H).

[0207] (*R*)-5-ethoxy-6-methyl-1-((2-(trimethylsilyl)ethyl)sulfonyl)-1,2,3,6-tetrahydropyrazine in 68 % yield. LCMS: P = 70 %, retention time = 2.0 min, (M+H₂O+H)⁺: 325; chiral HPLC retention time = 4.8 min, ee = 97.3 %; **¹H-NMR** (CDCl₃): δ 4.3 (m, 1H), 4.1 (m, 2H), 3.6 (m, 3H), 3.2 (m, 1H), 2.9 (m, 2H), 1.5 (m, 3H), 1.3 (m, 3H), 1.0 (m, 2H), 0.0 (s, 9H).

II.4. Step 3: Cyclodehydration leading to **F**

Method F: Cyclodehydration

[0208] General Method F is the general procedure used for the synthesis of chiral triazolopiperazine intermediates **F**.

Scheme 13: Formation of acylhydrazide **F**.

[0209] In a round-bottom flask equipped with a condenser, imino-ether **D** (1 eq.) was dissolved in anhydrous MeOH, to which was added **E** (1 eq.) in one portion. The resulting solution was stirred at a temperature ranging from 55°C to 70 °C for a period of time ranging from 6 hours to 8 hours. Completion of the reaction was monitored by HPLC analysis. The reaction mixture was cooled down to rt and the solvent was removed under reduced pressure. The crude compound was then purified by silica gel chromatography to afford the desired product **F**.

[0210] In an embodiment of the invention, the crude compound precipitates during cooling of the reaction mixture. In this case, the precipitate is stirred at rt in MeOH for about 5 hours before being filtered, washed with MeOH and oven dried.

[0211] Cyclodehydration is illustrated by the synthesis of intermediate (R)-5-(7-allyl-8-methyl-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazin-3-yl)-3-methyl-1,2,4-thiadiazole (i.e. compound **F**₁ wherein **PG** is allyl, **R**^{4'} is Me, **X**¹ is N, **X**² is S and **R**^{5'} is methyl).

[0212] To (R)-1-allyl-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine (0.14 g, 0.77 mmol) at rt was added 3-methyl-1,2,4-thiadiazole-5-carbohydrazide (0.12 g, 0.77 mmol) at once. The mixture was diluted with commercially anhydrous MeOH (0.77 mL) to allow complete solubilization and the resulting mixture was heated to 60 °C for 16h.

[0213] The reaction mixture was then allowed to reach rt whereupon the solvent was removed under reduced pressure (1-2 mbar). The crude residue was then dissolved in DCM (10 mL), and thus-obtained organic phase washed with NaOH (1 M, 10 mL). The organic layer was then dried over MgSO₄, filtered and concentrated under reduced pressure (1-2 mbar) the desired product as a yellow solid. Yield: 0.09 g, 42 %. LCMS: P = 95 %, retention time = 1.6 min, (M+H)⁺: 277; chiral HPLC retention time = 21.6 min, ee = 98.9 %; ¹H-NMR (CDCl₃): δ 5.9 (m, 1H), 5.3 (m, 2H), 4.5 (m, 1H), 4.4 (m, 1H), 4.1 (m, 1H), 3.5 (m, 1H), 3.3 (m, 1H), 3.1 (m, 1H), 2.8 (m, 1H), 2.7 (s, 3H), 1.6 (m, 3H).

[0214] The following intermediates were also prepared from the *ad hoc* reagents:

(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo [4,3-a]pyrazin-7(8H)-yl)diphenylphosphine oxide in 31 % yield (reaction time: 48h and silica gel purification (EtOAc)). LCMS: P = 96 %, retention time = 2.2 min, (M+H)⁺: 437; chiral HPLC retention time = 7.5 min, ee = 98.3 %; ¹H-NMR (CDCl₃): δ 7.9 (m, 4H), 7.5 (m, 6H), 4.9 (m, 1H), 4.8 (dd, J = 3.1, 13.6 Hz, 1H), 4.3 (dt, J = 4.9, 12.2 Hz, 1H), 3.6 (m, 1H), 3.5 (m, 1H), 2.7 (s, 3H), 1.6 (d, J = 6.9 Hz, 3H).

[0215] (R)-tert-butyl 8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo [4,3-a]pyrazine-7(8H)-carboxylate in 83% yield (reaction time: 48h). LCMS: P = 97 %, retention time = 2.3 min, (M+H)⁺: 337; chiral HPLC retention time = 19.4 min, ee = 95.1 %; ¹H-NMR (CDCl₃): δ 5.7 (m, 1H), 4.9 (m, 1H), 4.5 (m, 1H), 4.2 (m, 1H), 3.3 (m, 1H), 2.7 (s, 3H), 1.6 (d, J = 6.9 Hz, 3H), 1.5 (s, 9H).

[0216] (R)-3-methyl-5-(8-methyl-7-((2-(trimethylsilyl)ethyl)sulfonyl)-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazin-3-yl)-1,2,4-thiadiazole in 28 % yield (reaction time: 48h). LCMS: P = 40 %, retention time = 2.5 min, (M+H)⁺: 401; chiral HPLC retention time = 7.1 min, ee = 92.4 %; ¹H-NMR (CDCl₃): δ 4.9 (m, 1H), 4.3 (m, 1H), 4.1 (m, 1H), 3.6 (m, 1H), 3.0 (m, 1H), 2.7 (s, 3H), 1.6 (m, 2H), 1.4 (m, 3H), 1.0 (m, 2H), 0.0 (s, 9H).

II.5. Step 4: PG-deprotection


[0217] The methods of deprotection of above Protecting Groups (PGs) are known to those skill-in-the-art. As examples, one may refer to "Greene's Protective Groups in Organic Synthesis":

- Allyl: p. 806 of fourth edition;
- DPP: p. 844 of fourth edition;

- Boc: p. 725 of fourth edition;
- SES: p. 854 in fourth edition.

Method G: DMB deprotection - TFA/DCM

[0218]



Scheme 14: DMB deprotection.


[0219] Deprotection of DMB may be performed using TFA.

[0220] When crude or precipitated **F** was used (in opposition to purified **F** on silica gel), pre-washing was performed before deprotection as follow: **F** was dissolved in DCM and optionally washed with 1M NaOH in order to remove remaining **E**. The DCM extracts were then dried over magnesium sulphate, filtered and the filter cake washed with DCM.

[0221] **F** was diluted with DCM and TFA (7.6 eq.) was added to the DCM solution of **F** at RT. The mixture was stirred at rt for 2h-2h30. Completion of the deprotection was monitored by HPLC. Water was added, the mixture stirred for 30 minutes and filtered. The filter cake was washed with water and DCM. The filtrate layers were separated. The pH of the aqueous layer was adjusted to 12-13 by the addition of 4M NaOH. Sodium chloride was then added and the aqueous solution was extracted with DCM. The DCM extract comprising **G** was concentrated and was used in the next step without further purification.

II.6. Optional conversion of R^{5'} to R⁵ in triazolopiperazine G

[0222]



Scheme 15: conversion of R^{5'} to R⁵ in triazolopiperazine G leading to G'.

[0223] Substituent R⁵ may then be introduced, when applicable, from R^{5'} (especially when R^{5'} = H). One example of such transformation is illustrated by the synthesis of intermediate **G'** wherein **R⁵** is trifluoromethyl.

[0224] To a solution of **G** (1 eq.) in DCM/water (3/1) are added, at rt, sodium trifluoromethanesulfonate (3 eq.) and 2-hydroperoxy-2-methylpropane (5 eq.). The reaction mixture is *not* stirred and left at rt. Monitoring conversion by HPLC-MS, extra amount of each reagent can be added if required. The resulting mixture is diluted with DCM and quenched with 4 M NaOH saturated solution. Layers were separated and aqueous layer was extracted twice with EtOAc. The organic phases are combined, dried over MgSO₄ and evaporated to dryness. The residue is purified on silica gel or used crude in next step.

[0225] A further example of such transformation is illustrated by the synthesis of intermediate **G'** wherein **R⁵** is difluoromethyl:

To a suspension of **G** (**R⁵** = H) (*R*)-5-(8-methyl-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazin-3-yl)-1,2,4-thiadiazole (0.29 g, 1.13 mmol) and bis(difluoromethylsulfinyloxy)zinc (0.67 g, 2.26 mmol) in DCM (5 mL) and Water (2 mL), was added TFA (0.09 mL, 1.13 mmol), followed by slow addition of 2-hydroperoxy-2-methylpropane (0.77 mL, 5.64 mmol) with vigorous stirring.


[0226] When conversion was not increasing any more (HPLC-MS monitoring) bis(difluoromethylsulfinyloxy)zinc and 2-hydroperoxy-2-methylpropane were added at rt still with vigorous stirring (3 additional times (1.001 g, 3.39 mmol) and (0.773 mL, 5.64 mmol) respectively).

[0227] After 4 days in total, reaction mixture was diluted with EtOAc (50 mL) and carefully quenched with NaHCO₃ sat. solution (30 mL) and then NaHCO₃ solid until no bubbling was observed. Reaction mixture was filtered on Celite pad and phases of the filtrate were separated. Aqueous phase was filtered again on Celite pad then filtrate was extracted with EtOAc (2 x 50 mL). Organic phases were combined, dried over MgSO₄, filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel (DCM/MeOH 99/1) to afford the desired product as colorless oil. Yield: 0.03 g, 10 %. LCMS: P = 97 %, retention time = 1.8 min, (M+H)⁺: 273; ¹H-NMR (CDCl₃): δ 6.8 (t, J_{H-F} = 53.5 Hz, 1H), 4.7 (m, 1H), 4.3 (m, 2H), 3.5 (m, 1H), 3.3 (m, 1H), 1.7 (d, J = 6.7 Hz, 3H); ¹⁹F-NMR (CDCl₃): δ -113.5 (dd, J = 3.2, 53.5 Hz, 2F).

II.7. Step 5: Acylation leading to products I

Method H: Acylation NMM/DCM

[0228] General Method H is the general procedure used for the synthesis of (*R*)-enantiomer of **Formula I** of the invention.



Scheme 16: Acylation.

[0229] To a solution of crude **G** or **G'** (1 eq.) in anhydrous DCM were added at 0°C **H** (1.3 eq.), followed by N-methylmorpholine (2.2 eq.) dropwise over 15 sec. The reaction mixture was stirred at rt for 10 minutes and, the milky suspension was poured into 1 M HCl. The aqueous phase was extracted with DCM. The organic phases were combined, washed with 1 M NaOH, brine, dried over MgSO₄ and evaporated to dryness. The crude compound was purified by silica gel chromatography to afford the desired product (*R*)-**I**.

[0230] Measurement of %ee confirmed that no detectable racemization occurs during the acidolytic deprotection and N-acylation steps.

Method I: Acylation - biphasic conditions

[0231] Alternatively, the reaction may be performed under biphasic conditions.

[0232] In this case, saturated sodium hydrogen carbonate solution was added to the DCM slurry of **G** or **G'** (1 eq.) at rt. **H** (1 eq.) was added and the mixture stirred for a period of time ranging from about 20 minutes to overnight at rt. Completion of the reaction was monitored by HPLC. The layers were separated and the DCM phase washed with water. The DCM extracts were dried with magnesium sulphate and filtered, washing the filter cake with DCM. The DCM extracts were then concentrated. TBME was added and the resulting slurry stirred overnight at rt. The solid was collected by filtration, washed with TBME and pulled dry. The crude compound may be purified by silica gel chromatography or by

crystallisation.

[0233] Measurement of %ee confirmed that no detectable racemization occurs during the acidolytic deprotection and N-acylation steps.

[0234] Substituent R⁴ may then be transformed, when applicable, into R⁴ (see racemic synthesis).

II.8. Optional further transformation leading to products I''/I''' from II'

[0235] Compound 45: From compound I/I' wherein R⁵ = 1-((tert-butyldiphenylsilyl)oxy)ethyl, well known *tert*-butylammonium fluoride TBDPS deprotection of alkoxy was applied, followed by DAST fluorination of the latter alcohol, leading to racemic compound 45. Both diastereomers can be separated by purification on preparative HPLC to afford 45-1 and 45-2.

[0236] Compound 43: From compound I/I' wherein R⁵ = 1-((tert-butyldiphenylsilyl)oxy)ethyl, well known *tert*-butylammonium fluoride TBDPS deprotection of alkoxy was applied, followed by Dess-Martin oxidation, then followed by DAST fluorination of the latter ketone, leading to compound 43.

III. Chemical characterization

[0237] Compound 1: HPLC-MS: t_R = 4.1 min, (M+H)⁺ = 409; Chiral HPLC (Method C): %ee = 99.0; ¹H-NMR (CDCl₃): δ 7.6 (m, 2H), 7.3 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

[0238] Compound 2: HPLC-MS: t_R = 3.8 min, (M+H)⁺ = 373; Chiral HPLC (Method A): %ee = 98.0; ¹H-NMR (300 MHz, CDCl₃): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 3.1 (q, 2H), 1.8 (d, 3H), 1.4 (t, 3H); ¹⁹F-NMR (CDCl₃): δ -98.5.

[0239] Compound 3: HPLC-MS: t_R = 3.8 min, (M+H)⁺ = 375; Chiral HPLC (Method C): %ee > 99.8; ¹H-NMR (CDCl₃): δ 7.5 (m, 4H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

[0240] Compound 4: HPLC-MS: t_R = 3.9 min, (M+H)⁺ = 393; Chiral HPLC (Method C): %ee = 99.0; ¹H-NMR (CDCl₃): δ 7.6 (m, 1H), 7.3 (s, 1H), 7.2 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H); ¹⁹F-NMR (CDCl₃): δ -98.4.

[0241] Compound 5: HPLC-MS: t_R = 3.4 min, (M+H)⁺ = 359; Chiral HPLC (Method C): %ee = 99.0; ¹H-NMR (CDCl₃): δ 7.5 (m, 2H), 7.3 (m, 2H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.5 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H); ¹⁹F-NMR (CDCl₃): δ -98.4.

[0242] Compound 6: HPLC-MS: t_R = 3.8 min, (M+H)⁺ = 393; Chiral HPLC (Method C): %ee = 99.5; ¹H-NMR (CDCl₃): δ 7.6 (m, 1H), 7.3 (m, 2H), 5.7 (m, 1H), 4.9 (m, 1H), 4.5 (m, 1H), 4.3 (m, 1H), 3.5 (m, 1H), 2.7 (s, 3H), 1.8 (d, 3H); ¹⁹F-NMR (CDCl₃): δ -96.2.

[0243] Compound 7: HPLC-MS: t_R = 3.8 min, (M+H)⁺ = 395; Chiral HPLC (Method C): %ee = 98.9; ¹H-NMR (CDCl₃): δ 7.1 (m, 2H), 5.8 (m, 1H), 5.0 (m, 1H), 4.5 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.8 (d, 3H); ¹⁹F-NMR (CDCl₃): δ -75.8.

[0244] Compound 8: HPLC-MS: t_R = 3.7 min, (M+H)⁺ = 395; Chiral HPLC (Method C): %ee = 99.0; ¹H-NMR (CDCl₃): δ 7.1 (m, 2H), 6.2 (m, 1H), 5.3-5.0 (m, 2H), 4.3-3.6 (m, 2H), 2.7 (s, 3H), 1.8 (m, 3H); ¹⁹F-NMR (CDCl₃): δ -49.4, -72.0, -77.4.

[0245] Compound 9: HPLC-MS: t_R = 3.6 min, (M+H)⁺ = 377; Chiral HPLC (Method C): %ee = 99.4; ¹H-NMR (CDCl₃): δ 7.3 (m, 3H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H); ¹⁹F-NMR (CDCl₃): δ -72.1, -74.4.

[0246] Compound 10: HPLC-MS: t_R = 4.0 min, (M+H)⁺ = 413; Chiral HPLC (Method C): %ee = 99.0; ¹H-NMR (CDCl₃): δ 7.2 (m, 1H), 6.2 (m, 1H), 5.2-5.0 (m, 2H), 4.3 (m, 1H), 3.9-3.4 (m, 2H), 2.7 (s, 3H), 1.8 (m, 3H); ¹⁹F-NMR (CDCl₃): δ -54.2, -56.3, -67.1, -72.8.

[0247] Compound 11: HPLC-MS: t_R = 3.2 min, (M+H)⁺ = 389; Chiral HPLC (Method A): %ee > 99.8; ¹H-NMR (CDCl₃): δ 7.5 (m, 2H), 7.2 (m, 2H), 6.1 (m, 1H), 4.9 (dd, 1H), 4.3 (m, 2H), 3.9 (m, 2H), 3.6 (m, 1H), 2.7 (s, 3H), 2.4 (m, 1H), 2.2 (m, 1H); ¹⁹F-NMR (CDCl₃): δ -97.9.

[0248] Compound 12: is racemate of compound 11.

[0249] Compound 13: HPLC-MS: t_R = 4.1 min, (M+H)⁺ = 357; Chiral HPLC (Method B): %ee = 98.7; ¹H-NMR (CDCl₃): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.8 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.9 (q, 2H), 1.8 (d, 3H), 1.4 (t, 3H); ¹⁹F-NMR (CDCl₃): δ -98.7.

[0250] Compound 14: is racemate of compound 5.

[0251] Compound 15: HPLC-MS: t_R = 3.4 min, (M+H)⁺ = 359; Chiral HPLC (Method B): %ee > 99.8; ¹H-NMR (CDCl₃): δ 7.5 (m, 1H), 7.2 (m, 3H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H); ¹⁹F-NMR (CDCl₃): δ -96.2.

[0252] Compound 16: HPLC-MS: t_R = 3.7 min, (M+H)⁺ = 375; ¹H-NMR (CDCl₃): δ 7.5-7.3 (m, 4H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

[0253] Compound 17: HPLC-MS: t_R = 3.6 min, (M+H)⁺ = 377; ¹H-NMR (CDCl₃): δ 7.3 (m, 1H), 7.0 (m, 2H), 5.8 (m,

1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.8 (s, 3H), 1.8 (d, 3H); ^{19}F -NMR (CDCl_3): δ -101.2.

[0254] Compound 18: HPLC-MS: t_R = 3.5 min, $(\text{M}+\text{H})^+ = 377$; ^1H -NMR (CDCl_3): δ 7.5 (m, 1H), 7.0-6.9 (m, 2H), 6.2 (m, 1H), 5.2-4.9 (m, 2H), 4.3 (m, 1H), 4.0-3.7 (m, 1H), 2.7 (s, 3H), 1.7 (m, 3H); ^{19}F -NMR (CDCl_3): δ -95.7, -102.5.

[0255] Compound 19: HPLC-MS: t_R = 3.6 min, $(\text{M}+\text{H})^+ = 355$; ^1H -NMR (CDCl_3): δ 7.4 (m, 4H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 2.4 (s, 3H), 1.7 (d, 3H).

[0256] Compound 20: HPLC-MS: t_R = 3.3 min, $(\text{M}+\text{H})^+ = 341$; Chiral HPLC (Method C): %ee = 96.8; ^1H -NMR (CDCl_3): δ 7.5 (m, 5H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

[0257] Compound 21: HPLC-MS: t_R = 4.0 min, $(\text{M}+\text{H})^+ = 409$; ^1H -NMR (CDCl_3): δ 7.7 (d, 2H), 7.6 (d, 1H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H); ^{19}F -NMR (CDCl_3): δ -60.1.

[0258] Compound 22: HPLC-MS: t_R = 3.7 min, $(\text{M}+\text{H})^+ = 373$; Chiral HPLC (Method B): %ee > 99.7; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.3 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 2.2-2.0 (m, 2H), 1.1 (m, 3H); ^{19}F -NMR (CDCl_3): δ -98.4.

[0259] Compound 23: is racemate of compound 22.

[0260] Compound 24: HPLC-MS: t_R = 4.0 min, $(\text{M}+\text{H})^+ = 387$; Chiral HPLC (Method D): %ee = 95.5; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.2 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 2.1-2.0 (m, 2H), 1.6 (m, 2H), 1.0 (m, 3H); ^{19}F -NMR (CDCl_3): δ -98.2.

[0261] Compound 25: HPLC-MS: t_R = 3.5 min, $(\text{M}+\text{H})^+ = 389$; ^1H -NMR (CDCl_3): δ 7.2 (m, 2H), 7.0 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.9 (s, 3H), 3.5 (td, 1H), 2.7 (s, 3H), 1.8 (d, 3H); ^{19}F -NMR (CDCl_3): δ -76.3.

[0262] Compound 26: HPLC-MS: t_R = 3.5 min, $(\text{M}+\text{H})^+ = 355$; ^1H -NMR (CDCl_3): δ 7.4-7.2 (m, 4H), 6.3 (m, 1H), 5.3-4.8 (m, 2H), 4.3-3.8 (m, 1H), 3.5-3.4 (m, 1H), 2.7 (2s, 3H), 2.3 (s, 3H), 1.7 (2s, 3H).

[0263] Compound 27: HPLC-MS: t_R = 3.4 min, $(\text{M}+\text{H})^+ = 371$; ^1H -NMR (CDCl_3): δ 7.4 (m, 1H), 7.0 (m, 3H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.8 (s, 3H), 3.5 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

[0264] Compound 28: HPLC-MS: t_R = 3.1 min, $(\text{M}+\text{H})^+ = 343$; Chiral HPLC (Method C): %ee = 96.1; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.5 (td, 1H), 2.5 (s, 3H), 1.7 (d, 3H); ^{19}F -NMR (CDCl_3): δ -98.3.

[0265] Compound 29: HPLC-MS: t_R = 3.2 min, $(\text{M}+\text{H})^+ = 366$; Chiral HPLC (Method B): %ee = 99.0; ^1H -NMR (CDCl_3): δ 7.8 (d, 2H), 7.6 (d, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

[0266] Compound 30: HPLC-MS: t_R = 4.9 min, $(\text{M}+\text{H})^+ = 421$; Chiral HPLC (Method A): %ee = 98.2; ^1H -NMR (CDCl_3): δ 7.7 (d, 2H), 7.5 (d, 2H), 7.4 (m, 2H), 7.1 (m, 1H), 5.9 (m, 1H), 4.8 (dd, 1H), 4.7 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.9 (q, 2H), 1.8 (d, 3H), 1.4 (t, 3H).

[0267] Compound 31: is racemate of compound 10.

[0268] Compound 32: is racemate of compound 9.

[0269] Compound 33: is racemate of compound 8.

[0270] Compound 34: is racemate of compound 7.

[0271] Compound 35: is racemate of compound 6.

[0272] Compound 36: is racemate of compound 4.

[0273] Compound 37: is racemate of compound 3.

[0274] Compound 38: is racemate of compound 1.

[0275] Compound 39: is racemate of compound 2.

[0276] Compound 40: is racemate of compound 13.

[0277] Compound 41: HPLC-MS: t_R = 4.8 min, $(\text{M}+\text{H})^+ = 413$; Chiral HPLC (Method B): %ee = 99.7; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 1.8 (d, 3H); ^{19}F -NMR (CDCl_3): δ -62.9, -98.7.

[0278] Compound 42: HPLC-MS: t_R = 4.3 min, $(\text{M}+\text{H})^+ = 395$; Chiral HPLC (Method B): %ee = 97.4; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.1 (m, 2H), 6.8 (t, $J_{\text{HF}} = 53.5$ Hz, 1H), 5.8 (m, 1H), 4.9 (dd, $J = 3.1, 13.6$ Hz, 1H), 4.6 (m, 1H), 4.3 (dt, $J = 4.6, 13.3$ Hz, 1H), 3.6 (m, 1H), 1.8 (d, $J = 6.9$ Hz, 3H); ^{19}F -NMR (CDCl_3): δ -105.2 (s, 1F), -113.4 (dd, $J = 9.6, 53.4$ Hz, 2F).

[0279] Compound 43: HPLC-MS: t_R = 4.4 min, $(\text{M}+\text{H})^+ = 393$; Chiral HPLC (Method C): %ee = 96.3; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.9 (m, 1H), 4.8 (dd, $J = 3.3, 13.5$ Hz, 1H), 4.6 (m, 1H), 4. (dt, $J = 4.2, 12.7$ Hz, 1H), 3.6 (m, 1H), 2.2 (t, $J = 8.6$ Hz, 3H), 1.8 (d, $J = 6.9$ Hz, 3H); ^{19}F -NMR (CDCl_3): δ -88.3 (q, $J = 18.3$ Hz, 2F), -105.0 (s, 1F).

[0280] Compound 44: HPLC-MS: t_R = 4.4 min, $(\text{M}+\text{H})^+ = 411$; Chiral HPLC (Method C): %ee = 98.6; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.8 (dd, $J = 3.5, 13.6$ Hz, 1H), 4.6 (m, 1H), 4.3 (dt, $J = 4.0, 12.2$ Hz, 1H), 3.7 (q, $J = 10.0$ Hz, 2H), 3.6 (m, 1H), 1.8 (d, $J = 6.9$ Hz, 3H); ^{19}F -NMR (CDCl_3): δ -61.1 (t, $J = 9.6$ Hz, 1F), -105.0 (s, 1F).

[0281] Compound 45: HPLC-MS: t_R = 4.4 min, $(\text{M}+\text{H})^+ = 375$; Chiral HPLC (Method C): %ee = 98.5; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.9 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 1.9 (d, $J = 6.9$ Hz, 3H), 1.8 (m, 3H); ^{19}F -NMR (CDCl_3): δ -105.0 (s, 1F), -175.0 (m, 1F).

[0282] Compound 45-1: HPLC-MS: t_R = 4.4 min, $(\text{M}+\text{H})^+ = 375$; Chiral HPLC (Method C): %ee = 99.2; ^1H -NMR (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.9 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 1.9 (d, $J = 6.9$ Hz, 3H), 1.8 (m, 3H); ^{19}F -NMR (CDCl_3): δ -105.0 (s, 1F), -175.0 (m, 1F).

[0283] Compound 45-2: HPLC-MS: $t_R = 4.4$ min, $(M+H)^+ = 375$; Chiral HPLC (Method C): %ee = 91.7; 1H -NMR ($CDCl_3$): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.9 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 1.9 (d, $J = 6.9$ Hz, 3H), 1.8 (m, 3H); ^{19}F -NMR ($CDCl_3$): δ -105.0 (s, 1F), -175.0 (m, 1F).

BIOLOGY EXAMPLES

Functional Assay

Aequorin assay with human NK-3 receptor

[0284] Changes in intracellular calcium levels are a recognized indicator of G protein-coupled receptor activity. The efficacy of compounds of the invention to inhibit NKA-mediated NK-3 receptor activation was assessed by an *in vitro* Aequorin functional assay. Chinese Hamster Ovary recombinant cells expressing the human NK-3 receptor and a construct that encodes the photoprotein apoaequorin were used for this assay. In the presence of the cofactor coelenterazine, apoaequorin emits a measurable luminescence that is proportional to the amount of intracellular (cytoplasmic) free calcium.

Antagonist testing

[0285] The antagonist activity of compounds of the invention is measured following preincubation (3 minutes) of the compound (at various concentrations) with the cells, followed by addition of the reference agonist (NKA) at a final concentration equivalent to the EC_{80} (3 nM) and recording of emitted light (FDSS 6000 Hamamatsu) over the subsequent 90-second period. The intensity of the emitted light is integrated using the reader software. Compound antagonist activity is measured based on the concentration-dependent inhibition of the luminescence response to the addition of Neurokinin A. Inhibition curves are obtained for compounds of the invention and the concentrations of compounds which inhibit 50% of reference agonist response (IC_{50}) were determined (see results in table 2 below). The IC_{50} values shown in table 2 indicate that compounds of the invention are potent NK-3 antagonist compounds.

Competitive binding assays

[0286] The affinity of compounds of the invention for the human NK-3 receptor was determined by measuring the ability of compounds of the invention to competitively and reversibly displace a well-characterized NK-3 radioligand in a concentration-dependent manner.

3H -SB222200 binding competition assay with human NK-3 receptor

[0287] The ability of compounds of the invention to inhibit the binding of the NK-3 receptor selective antagonist 3H -SB222200 was assessed by an *in vitro* radioligand binding assay. Membranes were prepared from Chinese hamster ovary recombinant cells stably expressing the human NK-3 receptor. The membranes were incubated with 5 nM 3H -SB222200 (ARC) in a HEPES 25 mM/ NaCl 0.1 M/CaCl₂ 1 mM/MgCl₂ 5 mM/ BSA 0.5%/ Saponin 10 μ g/ml buffer at pH 7.4 and various concentrations of compounds of the invention. The amount of 3H -SB222200 bound to the receptor was determined after filtration by the quantification of membrane associated radioactivity using the TopCount-NXT reader (Packard). Competition curves were obtained for compounds of the invention and the concentration that displaced 50% of bound radioligand (IC_{50}) were determined by linear regression analysis and then the apparent inhibition constant (K_i) values were calculated by the following equation: $K_i = IC_{50}/(1+[L]/K_d)$ where [L] is the concentration of free radioligand and K_d is its dissociation constant at the receptor, derived from saturation binding experiments (Cheng and Prusoff, 1973) (see results in table 2 below).

[0288] Table 2 shows biological results obtained using the 3H -SB222200 binding competition assay with compounds of the invention. These results indicate that compounds of the invention display potent affinity for the human NK-3 receptor.

TABLE 2

Cpd n°	Functional assay: Aequorin assay with human NK-3 receptor hNK-3 - AEQ(antagonist IC_{50} , nM)	Competitive binding assay with human NK-3 receptor hNK-3 (K_i , nM)
1	16	11
2	12	15

EP 2 948 455 B1

(continued)

Cpd n°	Functional assay: Aequorin assay with human NK-3 receptor hNK-3 - AEQ(antagonist IC ₅₀ , nM)	Competitive binding assay with human NK-3 receptor hNK-3 (K _i , nM)
3	32	19
4	19	20
5	18	23
6	30	24
7	30	26
8	33	26
9	21	30
10	56	31
11	73	32
12	170	59
13	44	40
14	57	42
15	50	45
16	71	49
17	50	51
18	87	54
19	110	56
20	93	60
21	150	63
22	130	69
23	220	150
24	120	78
25	110	85
26	74	88
27	220	100
28	160	110
29	170	150
30	5	7
31	64	70
32	44	59
33	120	89
34	62	38
35	54	73
36	58	43
37	52	41
38	34	26

(continued)

Cpd n°	Functional assay: Aequorin assay with human NK-3 receptor hNK-3 - AEQ(antagonist IC ₅₀ , nM)	Competitive binding assay with human NK-3 receptor hNK-3 (K _i , nM)
39	32	28
40	130	83
41	7	11
42	39	48
43	136	59
44	204	45
45	244	101
45-1	285	100
45-2	148	83

Selectivity assay

[0289] Selectivity of the compounds of the invention was determined over the other human NK receptors, namely NK-1 and NK-2 receptors.

Human NK-1

[0290] The affinity of compounds of the invention for the NK-1 receptor was evaluated in CHO recombinant cells which express the human NK-1 receptor. Membrane suspensions were prepared from these cells. The following radioligand: [³H] substance P (PerkinElmer Cat#NET111520) was used in this assay. Binding assays were performed in a 50 mM Tris / 5 mM MnCl₂ / 150 mM NaCl / 0.1% BSA at pH 7.4. Binding assays consisted of 25 µl of membrane suspension (approximately 5 µg of protein/well in a 96 well plate), 50 µl of compound or reference ligand (Substance P) at increasing concentrations (diluted in assay buffer) and 2nM [³H] substance P. The plate was incubated 60 min at 25°C in a water bath and then filtered over GF/C filters (Perkin Elmer, 6005174, presoaked in 0.5% PEI for 2h at room temperature) with a Filtration unit (Perkin Elmer). The radioactivity retained on the filters was measured by using the TopCount-NXT reader (Packard). Competition curves were obtained for compounds of the invention and the concentrations of compounds which displaced 50% of bound radioligand (IC₅₀) were determined and then apparent inhibition constant K_i values were calculated by the following equation: $K_i = IC_{50} / (1 + [L] / K_D)$ where [L] is the concentration of free radioligand and K_D is its dissociation constant at the receptor, derived from saturation binding experiments (Cheng and Prusoff, 1973).

Human NK-2

[0291] The affinity of compounds of the invention for the NK-2 receptor was evaluated in CHO recombinant cells which express the human NK-2 receptor. Membrane suspensions were prepared from these cells. The following radioligand [¹²⁵I]-Neurokinin A (PerkinElmer Cat#NEX252) was used in this assay. Binding assays were performed in a 25 mM HEPES / 1 mM CaCl₂ / 5 mM MgCl₂ / 0.5% BSA / 10 µg/ml saponin, at pH 7.4. Binding assays consisted of 25 µl of membrane suspension (approximately 3.75 µg of protein/well in a 96 well plate), 50 µl of compound or reference ligand (Neurokinin A) at increasing concentrations (diluted in assay buffer) and 0.1 nM [¹²⁵I]-Neurokinin A. The plate was incubated 60 min at 25°C in a water bath and then filtered over GF/C filters (Perkin Elmer, 6005174, presoaked in assay buffer without saponine for 2h at room temperature) with a Filtration unit (Perkin Elmer). The radioactivity retained on the filters was measured by using the TopCount-NXT reader (Packard). Competition curves were obtained for compounds of the invention and the concentrations of compounds which displaced 50% of bound radioligand (IC₅₀) were determined and then apparent inhibition constant K_i values were calculated by the following equation: $K_i = IC_{50} / (1 + [L] / K_D)$ where [L] is the concentration of free radioligand and K_D is its dissociation constant at the receptor, derived from saturation binding experiments (Cheng and Prusoff, 1973).

[0292] The compounds of the invention, which were tested in the above NK-1 and NK-2 described assays, demonstrated a low affinity at the human NK-1 and human NK-2 receptors: more than 200 fold shift of the K_i compared to the human NK-3 receptor (table 3). Thus, compounds according to the invention have been shown to be selective over NK-1 and

NK-2 receptors.

TABLE 3

Cpd n°	hNK-3 (K_i , nM)	hNK-1 (K_i , nM)	hNK-2 (K_i , nM)
1	11	10300	7500
2	15	23800	>30000
3	19	>30000	>30000
4	20	19900	23000
5	23	>30000	>30000
6	24	22100	25000
7	26	>30000	36000
8	26	>30000	>30000
9	30	>30000	>30000
10	31	>30000	49000
11	32	22000	>30000
12	59	NA	NA
13	40	>30000	>30000
14	42	>30000	>30000
15	45	>30000	>30000
16	49	>30000	37000
17	51	>30000	>30000
18	54	>30000	>30000
19	56	>30000	>30000
20	60	>30000	>30000
21	63	>30000	>30000
22	69	>30000	>30000
23	150	NA	NA
24	78	>30000	>30000
25	85	>30000	>30000
26	88	>30000	>30000
27	100	>30000	>30000
28	110	>30000	>30000
29	150	>30000	>30000
30	7	40000	32000
31	70	>30000	>30000
32	59	>30000	>30000
33	89	>30000	>30000
34	38	>30000	>30000
35	73	>30000	30000
36	43	>30000	36000
37	41	32000	>30000

(continued)

Cpd n°	hNK-3 (K _i , nM)	hNK-1 (K _i , nM)	hNK-2 (K _i , nM)
38	26	21000	28000
39	28	>30000	>30000
40	83	>30000	>30000
41	11	NA	NA
42	48	>30000	>30000
43	59	>30000	>30000
44	45	>30000	>30000
45	101	>30000	>30000
45-1	100	>30000	>30000
45-2	83	>30000	>30000
NA: not available			

hERG inhibition Assay

[0293] The human ether-a-go-go related gene (hERG) encodes the inward rectifying voltage gated potassium channel in the heart (I_{Kr}) which is involved in cardiac repolarisation. I_{Kr} current inhibition has been shown to elongate the cardiac action potential, a phenomenon associated with increased risk of arrhythmia. I_{Kr} current inhibition accounts for the vast majority of known cases of drug-induced QT-prolongation. A number of drugs have been withdrawn from late stage clinical trials due to these cardiotoxic effects, therefore it is important to identify inhibitors early in drug discovery.

[0294] The hERG inhibition study aims at quantifying the *in vitro* effects of compounds of the invention on the potassium-selective I_{Kr} current generated in normoxic conditions in stably transfected HEK 293 cells with the human ether-a-go-go-related gene (hERG).

[0295] Whole-cell currents (acquisition by manual patch-clamp) elicited during a voltage pulse were recorded in baseline conditions and following application of tested compounds (5 minutes of exposure). The concentrations of tested compounds (0.3 μM; 3 μM; 10 μM; 30 μM) reflect a range believed to exceed the concentrations at expected efficacy doses in preclinical models.

[0296] The pulses protocol applied is described as follow: the holding potential (every 3 seconds) was stepped from -80 mV to a maximum value of +40 mV, starting with -40 mV, in eight increments of +10 mV, for a period of 1 second. The membrane potential was then returned to -55 mV, after each of these incremented steps, for 1 second and finally repolarized to -80 mV for 1 second.

[0297] The current density recorded were normalized against the baseline conditions and corrected for solvent effect and time-dependent current run-down using experimental design in test compound free conditions.

[0298] Inhibition curves were obtained for compounds and the concentrations which decreased 50% of the current density determined in the baseline conditions (IC₅₀) were determined. All compounds for which the IC₅₀ value is above 10 μM are not considered to be potent inhibitors of the hERG channel whereas compounds with IC₅₀ values below 1 μM are considered potent hERG channel inhibitors.

[0299] When tested in the hERG inhibition assay, compounds of the invention were determined to have IC₅₀ values as shown in Table 4.

Determination of plasma protein binding

[0300] The pharmacokinetic and pharmacodynamic properties of chemicals/drugs are largely a function of the reversible binding of chemicals to plasma or serum proteins. Generally, only the unbound or "free fraction" of a drug is available for diffusion or transport across cell membranes, and for interaction with a pharmacological/toxicological target. Consequently, the extent of the plasma protein binding (PPB) of a compound influences its action as well as its distribution and elimination.

[0301] The determination of plasma protein binding (PPB) of a compound is enabled by equilibrium dialysis, an accepted and standard method for reliable estimation of the non-bound drug fraction in plasma. RED (Rapid Equilibrium Dialysis) device insert is made of two side-by-side chambers separated by an O-ring-sealed vertical cylinder of dialysis membrane

EP 2 948 455 B1

(MWCO ~8,000). Plasma containing drug (at 5 μ M or blood concentrations otherwise corresponding to efficacious doses, if known) is added to one chamber while buffer is added to the second. After 4 hours incubation at 37°C under shaking, an aliquot is removed from each chamber and analyzed by a LC-MS/MS procedure enables the determination of both free and bound drug.

[0302] The percentages provided in Table 4 represent for the compounds of the invention the bound drug fraction to the plasma protein. The "free fraction" may be calculated as 100% - % rPPB (i.e. the complementary percentage of that disclosed in Table 4, corresponding to the drug concentration that is unbound and therefore available to engage biological target and elicit pharmacological activity).

TABLE 4

Cpd n°	Exposure (%rPPB)	CardioSafety (hERG IC ₅₀ , μ M)
1	67	42
2	47	32
3	42	66
4	40	70
5	22	70
6	53	45
7	26	70
8	29	70
9	22	70
10	30	70
11	24	50
12	20	NA
13	37	70
14	21	NA
15	20	70
16	36	70
17	24	46
18	23	70
19	51	NA
20	26	50
21	38	45
22	27	70
23	34	NA
24	48	61
25	19	NA
26	19	NA
27	24	70
28	12	NA
29	10	59
30	94	32
31	31	NA

(continued)

Cpd n°	Exposure (%rPPB)	CardioSafety (hERG IC ₅₀ , µM)
32	25	NA
33	29	NA
34	24	NA
35	52	NA
36	60	NA
37	53	NA
38	76	NA
39	43	NA
40	24	NA
41	55	NA
42	16	NA
43	47	NA
44	31	NA
45	31	NA
45-1	27	NA
45-2	33	NA
NA: not available		

In vivo assay to assess compound activity in rat (oral dosing)

Castrated male rat model to assess the effect of compound of invention on circulating levels of luteinizing hormone (LH)

[0303] The effect of compounds of the invention to inhibit luteinizing hormone (LH) secretion is determined by the following biological studies.

[0304] In humans and rodents, castration is well-precedented to permit heightened, persistent GnRH signaling and consequent elevation of circulating LH. Thus, a castrated rat model is used to provide a broad index for measurement of LH inhibition as a marker of test compound inhibition of the GnRH signaling pathway.

[0305] Castrated adult male Sprague-Dawley (SD) rats (150-175 g,) were purchased from Janvier (St Berthevin, France). All animals were housed 2 per cage in a temperature-controlled room ($22 \pm 2^\circ\text{C}$) and $50 \pm 5\%$ relative humidity with a 12 hour/12 hour light/dark cycles (lights off at 6h00 pm). The animals were allowed 3 weeks of postoperative recovery prior to study. Animals were handled on a daily basis. Standard diet and tap water were provided *ad libitum*. Animal cage litters were changed once a week. On the study day, animals were acclimated to the procedure room for a period of one hour prior to the initiation of the experiment.

[0306] Compounds of the invention were formulated in 0.5% methyl cellulose.

[0307] After basal sampling (TO) a single dose of compounds of the invention or vehicle was administrated orally to rats. Blood samples were then collected at several time points post dosing (45, 90, 150, 300 and 420 minutes). Blood samples were obtained via tail vein bleed, drawn into EDTA-containing tubes and centrifuged immediately. Plasma samples were collected and stored in a -80°C freezer until assayed. Serum LH levels were determined using radioimmunoassay kit from RIAZEN - Rat LH, Zentech (Liege, Belgium). Baseline was defined as the initial basal blood sample.

[0308] When tested in the castrated male rat model described above, compounds n° **1, 2, 4, 5, 8, 9, 11, 13, 20** and **30** of the invention significantly suppressed circulating LH levels (statistically significant, $p < 0.05$) at a dose less than or equal to 30 mg/kg).

Effect of compounds of the invention on plasma testosterone in gonad intact male rats

[0309] The study was designed to evaluate the effect of compounds of the invention on testosterone circulating levels

following oral administration at 3 mg/kg on SD gonad intact male rats.

[0310] Briefly the experimental methods used for this study were as follows:

Two groups of non-fasted rats (male, Sprague-Dawley, 200 to 225g; n = 4 rats /group) with jugular vein cannulation, were dosed via a single oral administration of compounds of the invention at 3 mg/kg. The control group was dosed with the vehicle. Compounds of the invention were prepared in a dose formulation of pyrogen-free water with 0.5% methylcellulose. Blood samples were collected via the catheter implanted in the jugular vein at pre-determined intervals using EDTA-3K as anti-coagulant. Samples were chilled and rapidly processed by centrifugation to obtain corresponding plasma samples. Testosterone hormone levels were determined by RIA performed on plasma samples collected for all the groups at 5 minutes before administration (basal time), and at 45, 90, 150, 300, 480 minutes and 24 hours after dosing.

[0311] When tested in the gonad intact male rats, compound n°5 significantly suppressed plasma testosterone level over the test period as compared to the vehicle treated group (Figure 1).

Effect of compounds of the invention on prostate weight reduction in a Benign Prostatic Hyperplasia (BPH) rat model

[0312] Briefly, adult male rats were injected daily for four weeks with testosterone to cause an enlargement of the prostate as per methods previously described in the literature (Scolnick et al., J. Andrology, 1994, 15(4), 287-297; Rick et al., J. Urol., 2012, 187, 1498-1504; see Figure 2, Ctrl Neg vs BHP). Rats were then treated daily for three weeks with compounds of the invention. After 21 days treatment with compounds of the invention at 3, 10 or 30 mg/kg (q.d.; PO administration), the ratio of prostate to body weight (g prostate/100g of body weight) was evaluated as an indicator of BPH. Treated groups were compared to the BPH group (Testosterone-induced BPH group followed 21 days of vehicle administration) or to the Control group (Corn oil injection for the induction phase followed by vehicle treatment rather than test compound). Comparison between groups was made by using One-Way ANOVA followed by Dunnett's test for statistical analysis.

[0313] When tested in the Benign Prostatic Hyperplasia rat model, compound n°5, demonstrated a concentration-response to reduce prostate weight to normal levels (i.e. levels in rats not exposed to exogenous testosterone; Figure 2).

Effect of compounds of the invention on Estradiol circulating level in female rats

[0314] The aim of this study was to evaluate the effect of compounds of the invention on plasma estradiol levels following oral administration at 10 mg/kg (b.i.d.) for a period of 10 days in female rats.


[0315] Briefly the experimental methods used for this study were as follows:

Two groups of adult, female rats (Sprague-Dawley, ~320g) were treated in-phase with their individual estrous cycles. Thus, treatment was started in the proestrus phase (coincident with peak estradiol levels, as shown on Day 1 in Figure 3) and rats were dosed twice daily (~9h30 and 17h30) by oral administration either with a compound of the invention at 10 mg/kg or with the vehicle for the control group. Compounds of the invention were prepared in a dose formulation of pyrogen-free water with 0.5% methylcellulose. Estradiol levels were determined for all groups by ELISA performed on plasma samples derived from blood collections taken at 30 minutes before the daily, 9h30 test article administration on all days presented in Figure 3.

[0316] In vehicle-treated, adult female rats, estradiol peaks are observed every 4-5 days consistent with the anticipated duration of the rat estrous cycle. Treatment with compound n°5, significantly decreased estradiol levels over the time-course tracked over two consecutive estrous cycles. This finding is most apparent in the proestrus phase (i.e. for vehicle group, on Day 5 and Day 9) where estradiol levels rise coincident with ovulation.

Claims

1. A compound of Formula I:



or a pharmaceutically acceptable solvate thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;


R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl;

X^1 is N and X^2 is S or O; or X^1 is S and X^2 is N;

== represents a single or a double bond depending on X^1 and X^2 ;

* stands for the (R)-enantiomer or for the racemate of compound of Formula I.


2. The compound according to claim 1, having Formula I' and I'':



or a pharmaceutically acceptable solvate thereof, wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , R^4 , R^5 , X^1 and X^2 are as defined in claim 1; and

== represents a single or a double bond depending on X^1 and X^2 .

3. The compound according to claim 1 or claim 2, having Formula Ia:



or a pharmaceutically acceptable solvate thereof, wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;

R^{2'} is H or F;


R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁴ is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

R⁵ is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;


* - - stands for the (R)-enantiomer or for the racemate of compound of Formula Ia.

4. The compound according to anyone of claims 1 to 3, selected from Formulae Ia' and Ia'':



or a pharmaceutically acceptable solvate thereof, wherein **R¹**, **R^{1'}**, **R²**, **R^{2'}**, **R³**, **R⁴** and **R⁵** are as defined in claim 3.

5. The compound according to anyone of claims 1 to 4, having Formula Ia-1:



or a pharmaceutically acceptable solvate thereof, wherein:


R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

*--- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia-1.

6. The compound according to anyone of claims 1 to 5, having Formula Ia-2:



or a pharmaceutically acceptable solvate thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;


$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

*--- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia-2.

7. The compound according to anyone of claims 1 to 6, having Formula Ia-3:



or a pharmaceutically acceptable solvate thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;


$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

*--- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia-3.

8. The compound according to anyone of claims 1 to 7, having Formula Ib:




or a pharmaceutically acceptable solvate thereof, wherein:

R^3 is F;

R^5 is methyl, ethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl;


* stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ib.

9. The compound according to anyone of claims 1 to 8 having Formula Ib':



or a pharmaceutically acceptable solvate thereof, wherein R^3 and R^5 are defined as in claim 8.

10. The compound according to anyone of claims 1 to 9, having Formula Ic:



or a pharmaceutically acceptable solvate thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;


R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;

R^5 is methyl, ethyl or trifluoromethyl;

* _ _ stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ic.

11. The compound according to anyone of claims 1 to 10, having Formula Ic':



or a pharmaceutically acceptable solvate thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;





R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;

R^5 is methyl, ethyl or trifluoromethyl.





12. The compound according to anyone of claims 1 to 11, selected from the group consisting of:

1		(<i>R</i>)-(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
2		(<i>R</i>)-(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone





(continued)

3		(R)-(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
4		(R)-(4-chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
5		(R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
6		(R)-(3-chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone





(continued)

7		(<i>R</i>)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanone
8		(<i>R</i>)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanone
9		(<i>R</i>)-(3,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
10		(<i>R</i>)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanone





(continued)

5	11		(<i>R</i>)-(4-fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
10	12		(4-fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
15	13		(<i>R</i>)-(3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
20	14		(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
25			
30			
35			
40			
45			
50			





(continued)

5	15		(R)-(3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
10			
15	16		(R)-(3-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
20			
25			
30	17		(R)-(3,5-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
35			
40			
45	18		(R)-(2,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
50			





(continued)

19		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(p-tolyl)methanone
20		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(phenyl)methanone
21		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-(trifluoromethyl)phenyl)methanone
22		(R)-(8-ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone





(continued)

23		(8-ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
24		(<i>R</i>)-(4-fluorophenyl)(3-(3-methyl-1,2,4-thiadiazol-5-yl)-8-propyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
25		(<i>R</i>)-(4-fluoro-3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
26		(<i>R</i>)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(o-tolyl)methanone





(continued)

27		(R)-(3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
28		(R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
29		(R)-4-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazine-7-carbonyl)benzonitrile
31		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanone





(continued)

32		(3,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
33		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanone
34		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanone
35		(3-chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone



(continued)

36		(4-chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
37		(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
38		(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
39		(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone

(continued)

40		(3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
41		(R)-(4-fluorophenyl)(8-methyl-3-(3-(trifluoromethyl)-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
42		(R)-(3-(3-(difluoromethyl)-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
43		(R)-(3-(3-(1,1-difluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone

(continued)

5 10 15	44 	(R)-(4-fluorophenyl)(8-methyl-3-(3-(2,2,2-trifluoroethyl)-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
20 25 30	45 	((8R)-3-(3-(1-fluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone

and pharmaceutically acceptable solvates thereof.

13. The compound according to anyone of claims **1** to **12**, which selected from the group consisting of:

(R)-(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone;
 (R)-(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone;
 (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone;
 (4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone; and
 (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone,
 and pharmaceutically acceptable solvates thereof.

14. The compound according to anyone of claims **1** to **13**, which is (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone, or a pharmaceutically acceptable solvate thereof.

15. A pharmaceutical composition comprising a compound according to anyone of claims **1** to **14**, or a pharmaceutically acceptable solvate thereof, and at least one pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant.

16. A compound according to anyone of claims **1** to **14** or a pharmaceutically acceptable solvate thereof for use as a medicament.

17. A compound according to anyone of claims **1** to **14** or a pharmaceutically acceptable solvate thereof for use in treating and/or preventing depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders,

cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian tumor or virilizing adrenal tumor), menorrhagia and adenomyosis.

18. The compound for use according to claim 17, wherein the compound is (*R*)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazin-7(8H)-yl)methanone or a pharmaceutically acceptable solvate thereof.

19. A compound according to anyone of claims 1 to 14 or a pharmaceutically acceptable solvate thereof for use in treating and/or preventing hot flashes.

20. The compound for use according to claim 19, wherein the compound is (*R*)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazin-7(8H)-yl)methanone or a pharmaceutically acceptable solvate thereof.

21. A compound according to anyone of claims 1 to 14 or a pharmaceutically acceptable solvate thereof for use as a lowering agent of the circulating LH levels.

22. The compound for use according to claim 21, wherein the compound is (*R*)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazin-7(8H)-yl)methanone or a pharmaceutically acceptable solvate thereof.

23. Process of manufacturing a compound according to anyone of claims 1 to 14 or a pharmaceutically acceptable solvate thereof, **characterized in that** it comprises the following steps:

a) reacting a compound of Formula (i)




wherein:

PG represents a suitable protecting group such as for example DMB, PMB, Boc, allyl, diphenyl-phosphinamide or 2-trimethylsilylethanesulfonyl;

R^{4'} is **R⁴** as defined in claim 1 or a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl;

*--- stands for the (*R*)-enantiomer or for the racemate;

with a compound of Formula (ii)




(ii)

wherein:

$R^{5'}$ is R^5 as defined in claim 1, H or 1-((tert-butylidiphenylsilyl)oxy)ethyl;

X^1 and X^2 are as defined in claim 1;


== represents a single or a double bond depending on X^1 and X^2 ; so as to obtain a compound of Formula (iii)



(iii)

wherein PG , R^4 , $R^{5'}$, X^1 and X^2 are as defined above, * _ _ stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on X^1 and X^2 ;


b) deprotecting compound of Formula (iii) with a suitable deprotection agent to afford compound of Formula (iv)




(iv)

wherein R^4 , $R^{5'}$, X^1 and X^2 are as defined above, * _ _ stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on X^1 and X^2 ;


c) when $R^{5'}$ is H, introducing a trifluoromethyl or difluoromethyl group by direct C-H trifluoro- or difluoromethylation, leading to compound of Formula (v)



wherein $R^{4'}$, X^1 and X^2 are as defined above and R^5 is trifluoromethyl or * difluoromethyl, * _ _ stands for the (R)-enantiomer or for the racemate and _ _ _ represents a single or a double bond depending on X^1 and X^2 ; d) N-acylating compound of Formula (iv) wherein $R^{5'}$ is not H or compound of Formula (v), with a compound of Formula (vi)



wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$ and R^3 are as defined in claim 1; leading to compound of Formula (vii)



wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , $R^{4'}$, X^1 and X^2 are as defined above, * _ _ stands for the (R)-enantiomer or for the racemate, _ _ _ represents a single or a double bond depending on X^1 and X^2 ; and $R^{5''}$ is R^5 as defined in claim 1 or 1-((tert-butyldiphenylsilyl)oxy)ethyl;

e) optionally further conducting one or both of the two following steps e') and e''):


e') when $R^{4'}$ is a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl,

a step of reduction optionally followed by methyl ether formation;
 e") when R^5 is 1-((tert-butyl)diphenylsilyl)oxy)ethyl, a step of alcohol deprotection and subsequent fluorination to form 1-fluoroethyl R^5 group; or a step of alcohol deprotection, followed by an oxidation step and a subsequent fluorination step to afford 1,1-difluoroethyl R^5 group;

to afford compound of Formula I according to anyone of claims 1 to 14.

Patentansprüche

1. Verbindung mit der Formel I:



oder ein pharmazeutisch annehmbares Solvat davon, wobei:

R^1 H, F oder Methyl ist;

$R^{1'}$ H ist;

R^2 H, F, Cl oder Methoxy ist;

$R^{2'}$ H oder F ist;

R^3 H, F, Cl, Methyl, Trifluoromethyl oder Nitril ist;

R^4 Methyl, Ethyl, *n*-Propyl, Hydroxyethyl, Methoxyethyl, Trifluoromethyl, Difluoromethyl oder Fluoromethyl ist;


R^5 Methyl, Ethyl, Methoxymethyl, Trifluoromethyl, Difluoromethyl, Fluoromethyl, 1-Fluoroethyl, 1,1-Difluoroethyl oder 2,2,2-Trifluoroethyl ist;

X^1 N ist und X^2 S oder O ist; oder X^1 S ist und X^2 N ist;

== eine Einzel- oder eine Doppelbindung in Abhängigkeit von X^1 und X^2 darstellt;

* - - für das (*R*)-Enantiomer oder für das Recemat der Verbindung mit der Formel I steht.


2. Verbindung nach Anspruch 1 mit der Formel I' und I'':



oder ein pharmazeutisch annehmbares Solvat davon, wobei R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , R^4 , R^5 , X^1 und X^2 wie in Anspruch 1 definiert sind; und

== eine Einzel- oder eine Doppelbindung in Abhängigkeit von X^1 und X^2 darstellt.

3. Verbindung nach Anspruch 1 oder Anspruch 2 mit der Formel Ia:



oder ein pharmazeutisch annehmbares Solvat davon, wobei:

R¹ H, F oder Methyl ist;

R^{1'} H ist;

R² H, F, Cl oder Methoxy ist;


R^{2'} H oder F ist;

R³ H, F, Cl, Methyl, Trifluoromethyl oder Nitril ist;

R⁴ Methyl, Ethyl, *n*-Propyl, Hydroxyethyl, Methoxyethyl, Trifluoromethyl, Difluoromethyl oder Fluoromethyl ist;


R⁵ Methyl, Ethyl, Methoxymethyl, Trifluoromethyl, Difluoromethyl oder Fluoromethyl ist; * _ _ _ für das (*R*)-Enantiomer oder für das Recemat der Verbindung mit der Formel Ia steht.

4. Verbindung nach einem der Ansprüche 1 bis 3, ausgewählt aus den Formeln Ia' und Ia'':



oder ein pharmazeutisch annehmbares Solvat davon, wobei **R¹**, **R^{1'}**, **R²**, **R^{2'}**, **R³**, **R⁴** und **R⁵** wie in Anspruch 3 definiert sind.

5. Verbindung nach einem der Ansprüche 1 bis 4 mit der Formel Ia-1:



oder ein pharmazeutisch annehmbares Solvat davon, wobei:


R³ H, F, Cl, Methyl, Trifluoromethyl oder Nitril ist;

R⁴ Methyl, Ethyl, *n*-Propyl, Hydroxyethyl, Methoxyethyl, Trifluoromethyl, Difluoromethyl oder Fluoromethyl ist;

R⁵ Methyl, Ethyl, Methoxymethyl, Trifluoromethyl, Difluoromethyl oder Fluoromethyl ist;

* _ _ für das (*R*)-Enantiomer oder für das Recemat der Verbindung mit der Formel Ia-1 steht.

6. Verbindung nach einem der Ansprüche 1 bis 5 mit der Formel Ia-2:



oder ein pharmazeutisch annehmbares Solvat davon, wobei:

R¹ H, F oder Methyl ist;

R^{1'} H ist;

R² H, F, Cl oder Methoxy ist;


R^{2'} H oder F ist;

R³ H, F, Cl, Methyl, Trifluoromethyl oder Nitril ist;

R⁵ Methyl, Ethyl, Methoxymethyl, Trifluoromethyl, Difluoromethyl oder Fluoromethyl ist;

* _ _ für das (*R*)-Enantiomer oder für das Recemat der Verbindung mit der Formel Ia-2 steht.

7. Verbindung nach einem der Ansprüche 1 bis 6 mit der Formel Ia-3:



oder ein pharmazeutisch annehmbares Solvat davon, wobei:

R¹ H, F oder Methyl ist;

R^{1'} H ist;

R² H, F, Cl oder Methoxy ist;


R^{2'} H oder F ist;

R³ H, F, Cl, Methyl, Trifluoromethyl oder Nitril ist;

R⁴ Methyl, Ethyl, *n*-Propyl, Hydroxyethyl, Methoxyethyl, Trifluoromethyl, Difluoromethyl oder Fluoromethyl ist;

* _ _ für das (*R*)-Enantiomer oder für das Recemat der Verbindung mit der Formel Ia-3 steht.

8. Verbindung nach einem der Ansprüche 1 bis 7 mit der Formel Ib:




oder ein pharmazeutisch annehmbares Solvat davon, wobei:

R³ F ist;

R⁵ Methyl, Ethyl, Trifluoromethyl, Difluoromethyl, Fluoromethyl, 1-Fluoroethyl, 1,1-Difluoroethyl oder 2,2,2-Trifluoroethyl ist;


* _ _ für das (*R*)-Enantiomer oder für das Recemat der Verbindung mit der Formel Ib steht.

9. Verbindung nach einem der Ansprüche 1 bis 8 mit der Formel Ib':



oder ein pharmazeutisch annehmbares Solvat davon, wobei **R³** und **R⁵** wie in Anspruch 8 definiert sind.

10. Verbindung nach einem der Ansprüche 1 bis 9 mit der Formel Ic:



oder ein pharmazeutisch annehmbares Solvat davon, wobei:

R¹ H, F oder Methyl ist;

R^{1'} H ist;

R² H, F, Cl oder Methoxy ist;

R^{2'} H oder F ist;


R³ H, F, Cl, Methyl, Trifluoromethyl oder Nitril ist;

R⁴ Methyl, Ethyl, *n*-Propyl oder Hydroxyethyl ist;

R⁵ Methyl, Ethyl oder Trifluoromethyl ist;

* _ _ für das (*R*)-Enantiomer oder für das Recemat der Verbindung mit der Formel Ic steht.

11. Verbindung nach einem der Ansprüche 1 bis 10 mit der Formel Ic':



oder ein pharmazeutisch annehmbares Solvat davon, wobei:

R¹ H, F oder Methyl ist;

R^{1'} H ist;

R² H, F, Cl oder Methoxy ist;

R^{2'} H oder F ist;

R³ H, F, Cl, Methyl, Trifluoromethyl oder Nitril ist;

R⁴ Methyl, Ethyl, *n*-Propyl oder Hydroxyethyl ist;

R⁵ Methyl, Ethyl oder Trifluoromethyl ist.






12. Verbindung nach einem der Ansprüche 1 bis 11, ausgewählt aus der Gruppe, bestehend aus:

1		(<i>R</i>)-(3,4-Dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
2		(<i>R</i>)-(3-(3-Ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon
3		(<i>R</i>)-(4-Chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon






(fortgesetzt)

4		(<i>R</i>)-(4-Chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4] triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
5		(<i>R</i>)-(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
6		(<i>R</i>)-(3-Chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
7		(<i>R</i>)-(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4] triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanon
8		(<i>R</i>)-(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4] triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanon






(fortgesetzt)

5	9		(R)-(3,4-Difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
10			
15	10		(R)-(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanon
20			
25	11		(R)-(4-Fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
30			
35	12		(4-Fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
40			
45	13		(R)-(3-(3-Ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon
50			






(fortgesetzt)

5	14		(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
10			
15	15		(<i>R</i>)-(3-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
20			
25	16		(<i>R</i>)-(3-Chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
30			
35	17		(<i>R</i>)-(3,5-Difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
40			
45	18		(<i>R</i>)-(2,4-Difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
50			
55			






(fortgesetzt)

5	19		(R)-(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4] triazolo[4,3-a]pyrazin-7(8H)-yl)(p-tolyl)methanon
10			
15	20		(R)-(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4] triazolo[4,3-a]pyrazin-7(8H)-yl)(phenyl)methanon
20			
25	21		(R)-(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4] triazolo[4,3-a]pyrazin-7(8H)-yl)(4-(trifluoromethyl)phenyl)methanon
30			
35	22		(R)-(8-Ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4] triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon
40			
45	23		(8-Ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4] triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon
50			
55			






(fortgesetzt)

24		(R)-(4-Fluorophenyl)(3-(3-methyl-1,2,4-thiadiazol-5-yl)-8-propyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
25		(R)-(4-Fluoro-3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
26		(R)-(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(o-tolyl)methanon
27		(R)-(3-Methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
28		(R)-(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon






(fortgesetzt)

5	29		(<i>R</i>)-4-(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazin-7-carbonyl)benzonitril
10			
15	31		(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanon
20			
25	32		(3,4-Difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
30			
35	33		(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanon
40			
45	34		(8-Methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanon
50			


(fortgesetzt)

35		(3-Chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
36		(4-Chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
37		(4-Chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
38		(3,4-Dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
39		(3-(3-Ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon

(fortgesetzt)

5	40		(3-(3-Ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon
15	41		(<i>R</i>)-(4-Fluorophenyl)(8-methyl-3-(3-(trifluoromethyl)-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon
25	42		(<i>R</i>)-(3-(3-(Difluoromethyl)-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon
35	43		(<i>R</i>)-(3-(3-(1,1-Difluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon
45	44		(<i>R</i>)-(4-Fluorophenyl)(8-methyl-3-(3-(2,2,2-trifluoroethyl)-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon

(fortgesetzt)

45		((8R)-3-(3-(1-fluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon
----	---	---

und pharmazeutisch annehmbare Solvate davon.

13. Verbindung nach einem der Ansprüche 1 bis 12, ausgewählt aus der Gruppe, bestehend aus:

(R)-(3-(3-Ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo [4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanon;
 (R)-(4-Chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon;
 (R)-(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon;
 (4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon; und
 (R)-(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon;
 und pharmazeutisch annehmbare Solvate davon.

14. Verbindung nach einem der Ansprüche 1 bis 13, wobei es sich um (R)-(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon oder ein pharmazeutisch annehmbares Solvat davon handelt.

15. Pharmazeutische Zusammensetzung, umfassend eine Verbindung nach einem der Ansprüche 1 bis 14, oder ein pharmazeutisch annehmbares Solvat davon, und mindestens einen pharmazeutisch annehmbaren Trägerstoff, Verdünner, Hilfsstoff und/oder Adjuvans.

16. Verbindung nach einem der Ansprüche 1 bis 14 oder ein pharmazeutisch annehmbares Solvat davon zur Verwendung als ein Arzneimittel.

17. Verbindung nach einem der Ansprüche 1 bis 14, oder ein pharmazeutisch annehmbares Solvat davon, zur Verwendung bei der Behandlung und/oder Prävention von Depression, Angst, Psychose, Schizophrenie, psychotischen Störungen, bipolaren Störungen, kognitiven Störungen, Parkinson'scher Erkrankung, Alzheimer'scher Erkrankung, Aufmerksamkeits- und Hyperaktivitätsstörung (ADHD), Schmerz, Konvulsion, Fettleibigkeit, Entzündungserkrankungen, darin eingeschlossen Reizdarmsyndrom (IBS) und Reizdarmstörungen, Erbrechen, Präeklampsie, Atemweg-bedingten Erkrankungen, darin eingeschlossen chronisch obstruktive Lungenerkrankung, Asthma, Atemwegs-Hyperaktivität, Bronchokonstriktion und Husten, Harninkontinenz, Fortpflanzungsstörungen, Kontrazeption und Sexualhormon-abhängigen Erkrankungen, darin eingeschlossen, jedoch nicht beschränkt auf gutartige Prostatahyperplasie (BPH), Prostatahyperplasie, metastatischen Prostatakrebs, Hodenkrebs, Brustkrebs, Eierstockkrebs, Androgen-abhängige Akne, männliches Haarausfallmuster, Endometriose, abnormale Pubertät, Uterusfibrose, Uterusmyom, Uterusleiomyom, Hormon-abhängige Krebserkrankungen, Hyperandrogenämie, Hirsutismus, Virilisierung, polycystisches Ovarialsyndrom (PCOS), prämenstruelles Verstimmungssyndrom (PMDD), HAIR-AN-Syndrom (Hyperandrogenismus, Insulinresistenz und Acanthosis nigricans), *Hyperthecosis ovarii* (HAIR-AN mit Hyperplasie von Luteinisierungs-Thekazellen im ovariellen Stroma), weitere Manifestationen hoher intraovarieller Androgenkonzentrationen (z. B. Follikelreifungshemmung, Atresie, Anovulation, Dysmenorrhoe, dysfunktionale Uterusblutung, Unfruchtbarkeit), Androgen-erzeugender Tumor (virilisierender Eierstockkrebs oder virilisierender Nebennierentumor).

mor), Menorrhagie und Adenomyose.

18. Verbindung zur Verwendung nach Anspruch 17, wobei die Verbindung (*R*)-(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon oder ein pharmazeutisch annehmbares Solvat davon ist.

19. Verbindung nach einem der Ansprüche 1 bis 14 oder ein pharmazeutisch annehmbares Solvat davon zur Verwendung bei der Behandlung und/oder Prävention von Hitzewallungen.


20. Verbindung zur Verwendung nach Anspruch 19, wobei die Verbindung (*R*)-(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon oder ein pharmazeutisch annehmbares Solvat davon ist.

21. Verbindung nach einem der Ansprüche 1 bis 14 oder ein pharmazeutisch annehmbares Solvat davon zur Verwendung als ein Senkungsmittel der zirkulierenden LH-Niveaus.

22. Verbindung zur Verwendung nach Anspruch 21, wobei die Verbindung (*R*)-(4-Fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanon oder ein pharmazeutisch annehmbares Solvat davon ist.

23. Verfahren zur Herstellung einer Verbindung nach einem der Ansprüche 1 bis 14 oder ein pharmazeutisch annehmbares Solvat davon, **dadurch gekennzeichnet, dass** es die folgenden Schritte umfasst:

a) Reagieren einer Verbindung mit der Formel (i)



(i)


wobei:

PG eine geeignete Schutzgruppe darstellt, wie z. B. DMB, PMB, Boc, Allyl, Diphenyl-phosphinamid oder 2-Trimethylsilylethansulfonyl;

R^{4'} **R⁴** ist, wie in Anspruch 1 definiert, oder ein reduzierbarer Vorläufer von Hydroxyethyl und folglich ein weiterer Vorläufer von Methoxyethyl;

***** für das (*R*)-Enantiomer oder für das Recemat steht;

mit einer Verbindung mit der Formel (ii):




(ii)

wobei:

R^{5'} **R⁵**, wie in Anspruch 1 definiert, H oder 1-((Tert-butyldiphenylsilyl)oxy)ethyl ist;


X¹ und **X²** wie in Anspruch 1 definiert sind;

== eine Einzel- oder eine Doppelbindung in Abhängigkeit von **X¹** und **X²** darstellt; um eine Verbindung der Formel (iii) zu erhalten




wobei **PG**, **R^{4'}**, **R^{5'}**, **X¹** und **X²** wie oben definiert sind, * _ _ für das (R)-Enantiomer oder für das Recemat steht und == eine Einzel- oder eine Doppelbindung in Abhängigkeit von **X¹** und **X²** darstellt;

b) Entschützen der Verbindung mit der Formel (iii) mit einem geeigneten Entschützungsmittel, um die Verbindung der Formel (iv) zu erhalten




wobei **R^{4'}**, **R^{5'}**, **X¹** und **X²** wie oben definiert sind, * _ _ für das (R)-Enantiomer oder für das Recemat steht und == eine Einzel- oder eine Doppelbindung in Abhängigkeit von **X¹** und **X²** darstellt;

c) wobei **R^{5'}** H ist, eine Trifluoromethyl- oder Difluoromethyl-Gruppe durch direkte C-H- Trifluoro- oder Difluoromethylierung einführt, die zur Verbindung der Formel (v) führt




wobei **R^{4'}**, **X¹** und **X²** wie oben definiert sind, und **R⁵** Trifluoromethyl oder Difluoromethyl ist; * _ _ für das (R)-Enantiomer oder für das Recemat steht; und == eine Einzel- oder eine Doppelbindung in Abhängigkeit von **X¹** und **X²** darstellt;

d) N-Acylieren der Verbindung mit der Formel (iv), wobei **R^{5'}** nicht H oder die Verbindung von Formel (v) ist, mit einer Verbindung mit der Formel (vi)



(vi)

wobei R^1 , $R^{1'}$, R^2 , $R^{2'}$ und R^3 wie in Anspruch 1 definiert sind; zu Verbindung der Formel (vii) führen



(vii)

wobei R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , $R^{4'}$, X^1 und X^2 wie oben definiert sind;

* _ _ _ für das (*R*)-Enantiomer oder für das Recemat steht;

== eine Einzel- oder eine Doppelbindung in Abhängigkeit von X^1 und X^2 darstellt; und

$R^{5''}$ R^5 wie in Anspruch 1 definiert oder 1-((Tert-butyl-diphenylsilyl)oxy)ethyl ist;

e) optional weiter Durchführen von einem oder von beiden der zwei folgenden Schritte e') und e''):


e') wenn $R^{4'}$ ein reduzierbarer Vorläufer von Hydroxyethyl und folglich ein weiterer Vorläufer von Methoxyethyl ist, einen Schritt des Reduzierens, optional gefolgt von der Methyletherbildung;

e'') wenn $R^{5''}$ gleich 1-((Tert-butyl-diphenylsilyl)oxy)ethyl ist, einen Schritt des Alkohol-Entschützens und der folgenden Fluorierung, um die 1-Fluoroethyl R^5 -Gruppe zu bilden; oder einen Schritt des Alkohol-Entschützens, gefolgt von einem Schritt des Oxidierens und einem folgenden Schritt des Fluorierens, um die 1,1-Difluoroethyl R^5 -Gruppe zu erhalten;

um die Verbindung mit der Formel I nach einem der Ansprüche 1 bis 14 zu erhalten.

Revendications

1. Un composé de formule I:



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle :

R^1 est H, F ou un méthyle ;

$R^{1'}$ est H ;

R^2 est H, F, Cl ou un méthoxy ;

$R^{2'}$ est H ou F ;

R^3 est H, F, Cl, un méthyle, un trifluorométhyle ou un nitrile ;

R^4 est un méthyle, un éthyle, un *n*-propyle, un hydroxyéthyle, un méthoxyéthyle, un trifluorométhyle, un difluorométhyle ou un fluorométhyle ;


R^5 est un méthyle, un éthyle, un méthoxyméthyle, un trifluorométhyle, un difluorométhyle, un fluorométhyle, un 1-fluoroéthyle, un 1,1-difluoroéthyle ou un 2,2,2-trifluoroéthyle ;

X^1 est N et X^2 est S ou O ; ou X^1 est S et X^2 est N ;

== représente une liaison simple ou une liaison double en fonction de X^1 et X^2 ;

== représente l'énantiomère (R) ou le racémique du composé de formule I.


2. Le composé selon la revendication 1, ayant pour formule I' et I'' :



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , R^4 , R^5 , X^1 et X^2 sont tels que définis dans la revendication 1 ; et

== représente une liaison simple ou une liaison double en fonction de X^1 et X^2 .

3. Le composé selon la revendication 1 ou la revendication 2, ayant pour formule Ia :



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle :

R^1 est H, F ou un méthyle ;

$R^{1'}$ est H ;

R^2 est H, F, Cl ou un méthoxy ;

$R^{2'}$ est H ou F ;

R^3 est H, F, Cl, un méthyle, un trifluorométhyle ou un nitrile ;


R^4 est un méthyle, un éthyle, un *n*-propyle, un hydroxyéthyle, un méthoxyéthyle, un trifluorométhyle, un difluo-

rométhyle ou un fluorométhyle ;


R⁵ est un méthyle, un éthyle, un méthoxyméthyle, un trifluorométhyle, un difluorométhyle ou un fluorométhyle ;

* _ _ _ représente l'énantiomère (R) ou le racémique du composé de formule la.

4. Le composé selon l'une quelconque des revendications 1 à 3, choisi parmi les formules la' et la'' :




la'



la''

ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle **R¹**, **R^{1'}**, **R²**, **R^{2'}**, **R³**, **R⁴** et **R⁵** sont tels que définis dans la revendication 3.

5. Le composé selon l'une quelconque des revendications 1 à 4, ayant pour formule la-1 :



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle :


R³ est H, F, Cl, un méthyle, un trifluorométhyle ou un nitrile ;

R⁴ est un méthyle, un éthyle, un *n*-propyle, un hydroxyéthyle, un méthoxyéthyle, un trifluorométhyle, un difluorométhyle ou un fluorométhyle ;

R⁵ est un méthyle, un éthyle, un méthoxyméthyle, un trifluorométhyle, un difluorométhyle ou un fluorométhyle ;

* _ _ _ représente l'énantiomère (R) ou le racémique du composé de formule la-1.

6. Le composé selon l'une quelconque des revendications 1 à 5, ayant pour formule la-2 :



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle :

R^1 est H, F ou un méthyle ;

$R^{1'}$ est H ;

R^2 est H, F, Cl ou un méthoxy ;


$R^{2'}$ est H ou F ;

R^3 est H, F, Cl, un méthyle, un trifluorométhyle ou un nitrile ;

R^5 est un méthyle, un éthyle, un méthoxyméthyle, un trifluorométhyle, un difluorométhyle ou un fluorométhyle ;

* _ _ représente l'énantiomère (R) ou le racémique du composé de formule 1a-2.

7. Le composé selon l'une quelconque des revendications 1 à 6, ayant pour formule 1a-3 :



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle :

R^1 est H, F ou un méthyle ;

$R^{1'}$ est H ;

R^2 est H, F, Cl ou un méthoxy ;


$R^{2'}$ est H ou F ;

R^3 est H, F, Cl, un méthyle, un trifluorométhyle ou un nitrile ;

R^4 est un méthyle, un éthyle, un *n*-propyle, un hydroxyéthyle, un méthoxyéthyle, un trifluorométhyle, un difluorométhyle ou un fluorométhyle ;

* _ _ représente l'énantiomère (R) ou le racémique du composé de formule 1a-3.

8. Le composé selon l'une quelconque des revendications 1 à 7, ayant pour formule 1b :




ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle :

R^3 est F ;

R^5 est un méthyle, un éthyle, un trifluorométhyle, un difluorométhyle, un fluorométhyle, un 1-fluoroéthyle, un 1,1-difluoroéthyle ou un 2,2,2-trifluoroéthyle ;


* _ _ _ représente l'énantiomère (R) ou le racémique du composé de formule 1b.

9. Le composé selon l'une quelconque des revendications 1 à 8 ayant pour formule 1b' :



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle R^3 et R^5 sont tels que définis dans la revendication 8.

10. Le composé selon l'une quelconque des revendications 1 à 9, ayant pour formule 1c :



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle :

R^1 est H, F ou un méthyle ;

$R^{1'}$ est H ;

R^2 est H, F, Cl ou un méthoxy ;

$R^{2'}$ est H ou F ;


R^3 est H, F, Cl, un méthyle, un trifluorométhyle ou un nitrile ;

R^4 est un méthyle, un éthyle, un *n*-propyle ou un hydroxyéthyle ;

R^5 est un méthyle, un éthyle ou un trifluorométhyle ;

* *--- représente l'énantiomère (R) ou le racémique du composé de formule Ic.

11. Le composé selon l'une quelconque des revendications 1 à 10, ayant pour formule Ic' :



ou un solvate pharmaceutiquement acceptable de celui-ci, dans laquelle :

R^1 est H, F ou un méthyle ;

$R^{1'}$ est H ;

R^2 est H, F, Cl ou un méthoxy ;

$R^{2'}$ est H ou F ;

R^3 est H, F, Cl, un méthyle, un trifluorométhyle ou un nitrile ;





R^4 est un méthyle, un éthyle, un *n*-propyle ou un hydroxyéthyle ;

R^5 est un méthyle, un éthyle ou un trifluorométhyle.





12. Le composé selon l'une quelconque des revendications 1 à 11, choisi dans le groupe constitué par :

1		(R)-(3,4-dichlorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
2		(R)-(3-(3-éthyl-1,2,4-thiadiazol-5-yl)-8-méthyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone





(suite)

3		(R)-(4-chlorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
4		(R)-(4-chloro-3-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
5		(R)-(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
6		(R)-(3-chloro-4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone





(suite)

5 10	7 	(<i>R</i>)-(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophényl)méthanone
15 20 25	8 	(<i>R</i>)-(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophényl)méthanone
30 35	9 	(<i>R</i>)-(3,4-difluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
40 45 50	10 	(<i>R</i>)-(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tétrafluorophényl)méthanone





(suite)

5	11		(R)-(4-fluorophényl)(8-(2-hydroxyéthyl)-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
10	12		(4-fluorophényl)(8-(2-hydroxyéthyl)-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
15	13		(R)-(3-(3-éthyl-1,2,4-oxadiazol-5-yl)-8-méthyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone
20	14		(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
25			
30			
35			
40			
45			
50			





(suite)

5	15		(R)-(3-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
10			
15	16		(R)-(3-chlorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
20			
25			
30	17		(R)-(3,5-difluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
35			
40			
45	18		(R)-(2,4-difluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
50			





(suite)

19		(R)-(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(p-tolyl)méthanone
20		(R)-(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(phényl)méthanone
21		(R)-(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-(trifluorométhyl)phényl)méthanone
22		(R)-(8-éthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone





(suite)

23		(8-éthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone
24		(<i>R</i>)-(4-fluorophényl)(3-(3-méthyl-1,2,4-thiadiazol-5-yl)-8-propyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
25		(<i>R</i>)-(4-fluoro-3-méthoxyphényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
26		(<i>R</i>)-(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(o-tolyl)méthanone





(suite)

27		(R)-(3-méthoxyphényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
28		(R)-(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
29		(R)-4-(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6,7,8-tétrahydro-[1,2,4]triazolo[4,3-a]pyrazine-7-carbonyl)benzonitrile
31		(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tétrafluorophényl)méthanone





(suite)

5 10	32 	(3,4-difluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
15 20 25	33 	(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophényl)méthanone
30 35	34 	(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophényl)méthanone
40 45 50	35 	(3-chloro-4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone



(suite)

5 10	36		(4-chloro-3-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
15 20 25	37		(4-chlorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
30 35	38		(3,4-dichlorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
40 45 50	39		(3-(3-éthyl-1,2,4-thiadiazol-5-yl)-8-méthyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone

(suite)

40		(3-(3-éthyl-1,2,4-oxadiazol-5-yl)-8-méthyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone
41		(<i>R</i>)-(4-fluorophényl)(8-méthyl-3-(3-(trifluorométhyl)-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
42		(<i>R</i>)-(3-(3-(difluorométhyl)-1,2,4-thiadiazol-5-yl)-8-méthyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone
43		(<i>R</i>)-(3-(3-(1,1-difluoroéthyl)-1,2,4-oxadiazol-5-yl)-8-méthyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone

(suite)

5 10 15	44 	(<i>R</i>)-(4-fluorophényl)(8-méthyl-3-(3-(2,2,2-trifluoroéthyl)-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone
20 25 30	45 	((8<i>R</i>)-3-(3-(1-fluoroéthyl)-1,2,4-oxadiazol-5-yl)-8-méthyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone

et les solvates pharmaceutiquement acceptables de ceux-ci.

13. Le composé selon l'une quelconque des revendications **1** à **12**, choisi dans le groupe constitué par :

(*R*)-(3-(3-éthyl-1,2,4-thiadiazol-5-yl)-8-méthyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophényl)méthanone ;
 (*R*)-(4-chlorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone ;
 (*R*)-(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone ;
 (4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone ; et
 (*R*)-(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone ;
 et les solvates pharmaceutiquement acceptables de ceux-ci.

14. Le composé selon l'une quelconque des revendications **1** à **13**, qui est le (*R*)-(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone, ou un solvate pharmaceutiquement acceptable de celui-ci.

15. Une composition pharmaceutique comprenant un composé selon l'une quelconque des revendications **1** à **14**, ou un solvate pharmaceutiquement acceptable de celui-ci, et au moins un véhicule, diluant, excipient et/ou adjuvant pharmaceutiquement acceptables.

16. Un composé selon l'une quelconque des revendications **1** à **14** ou un solvate pharmaceutiquement acceptable de celui-ci pour son utilisation en tant que médicament.

17. Un composé selon l'une quelconque des revendications **1** à **14** ou un solvate pharmaceutiquement acceptable de

celui-ci pour son utilisation dans le traitement et/ou la prévention de la dépression, de l'anxiété, de la psychose, de la schizophrénie, des troubles psychotiques, des troubles bipolaires, des troubles cognitifs, de la maladie de Parkinson, de la maladie d'Alzheimer, du trouble déficitaire de l'attention avec hyperactivité (TDAH), de la douleur, des convulsions, de l'obésité, des maladies inflammatoires telles que le syndrome du côlon irritable et les troubles intestinaux inflammatoires, des vomissements, de la pré-éclampsie, des maladies associées aux voies respiratoires telles que la bronchopneumopathie obstructive chronique, l'asthme, l'hypersensibilité des voies respiratoires, la broncoconstruction et la toux, l'incontinence urinaire, les troubles de la reproduction, de la contraception et des maladies dépendantes des hormones sexuelles telles que, mais sans s'y limiter, l'hypertrophie bénigne de la prostate (HBP), l'hypertrophie de la prostate, le carcinome métastatique de la prostate, le cancer des testicules, le cancer du sein, le cancer des ovaires, l'acné dépendant des androgènes, la calvitie masculine, l'endométriome, une puberté anormale, la fibrose utérine, une tumeur fibrogénique utérine, léiomyome utérin, les cancers hormono-dépendants, l'hyperandrogénie, l'hirsutisme, le virilisme, le syndrome des ovaires polykystiques (SOPK), le trouble dysphorique prémenstruel (TDPM), le syndrome HAIR-AN (hyperandrogénie, résistance insuline et acanthosis nigricans), l'hyperthécose ovarienne (HAIR-AN avec hyperplasie des cellules thécales lutéinisées du stroma ovarien), d'autres manifestations de concentrations intra-ovariennes élevées en androgènes (par exemple, l'arrêt de la maturation folliculaire, l'atrésie, l'anovulation, les dysménorrhées, des saignements utérins dysfonctionnels, la stérilité), des tumeurs produisant des androgènes (tumeur ovarienne virilisante ou tumeur surrénalienne), la ménorragie et l'adénomyose.

18. Le composé pour son utilisation selon la revendication **17**, dans laquelle le composé est le (*R*)-(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone ou un solvate pharmaceutiquement acceptable de celui-ci.

19. Un composé selon l'une quelconque des revendications **1** à **14** ou un solvate pharmaceutiquement acceptable de celui-ci pour son utilisation dans le traitement et/ou la prévention des bouffées de chaleur.


20. Le composé pour son utilisation selon la revendication **19**, dans laquelle le composé est le (*R*)-(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone ou un solvate pharmaceutiquement acceptable de celui-ci.

21. Un composé selon l'une quelconque des revendications **1** à **14** ou un solvate pharmaceutiquement acceptable de celui-ci pour son utilisation en tant qu'agent diminuant les taux de LH circulants.

22. Le composé pour son utilisation selon la revendication **21**, dans laquelle le composé est le (*R*)-(4-fluorophényl)(8-méthyl-3-(3-méthyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)méthanone ou un solvate pharmaceutiquement acceptable de celui-ci.

23. Procédé de préparation d'un composé selon l'une quelconque des revendications **1** à **14** ou un solvate pharmaceutiquement acceptable de celui-ci, ledit procédé comprenant les étapes suivantes :

a) la réaction d'un composé de formule (i)



(i)

dans laquelle :


PG représente un groupement protecteur approprié tel que par exemple DMB, PMB, Boc, allyle, diphenylphosphinamide ou 2-triméthylsilyléthansulfonyle ;

R^{4'} est **R⁴** tel que défini dans la revendication **1** ou un précurseur réductible d'hydroxyéthyle et par consé-

quent un précurseur de méthoxyéthyle ;

* --- représente l'énantiomère (*R*) ou le racémique ;

avec un composé de formule (ii)



(ii)


dans laquelle:

R^{5'} est **R⁵** tel que défini dans la revendication 1, H ou un 1-((tert-butyl)diphénylsilyl)oxy)éthyle ;

X¹ et **X²** sont tels que définis dans la revendication 1 ;

== représente une liaison simple ou une liaison double en fonction de **X¹** et **X²**;


pour conduire à un composé de formule (iii)



(iii)

dans laquelle **PG**, **R^{4'}**, **R^{5'}**, **X¹** et **X²** sont tels que définis ci-dessus, *-représente l'énantiomère (*R*) ou le racémique et == représente une liaison simple ou une liaison double en fonction de **X¹** et **X²** ;

b) la déprotection du composé de formule (iii) avec un réactif de déprotection approprié pour obtenir le composé de formule (iv)




(iv)

dans laquelle **R^{4'}**, **R^{5'}**, **X¹** et **X²** sont tels que définis ci-dessus, *-représente l'énantiomère (*R*) ou le racémique


et --- représente une liaison simple ou une liaison double en fonction de $\mathbf{X^1}$ et $\mathbf{X^2}$;

c) quand $\mathbf{R^{5'}}$ est H, l'introduction d'un groupe trifluorométhyle ou difluorométhyle par C-H trifluoro- ou difluoro-méthylation directe, conduisant au composé de formule (v)




dans laquelle $\mathbf{R^{4'}}$, $\mathbf{X^1}$ et $\mathbf{X^2}$ sont tels que définis ci-dessus, et $\mathbf{R^5}$ est un * trifluorométhyle ou un difluorométhyle, *--- représente l'énantiomère (*R*) ou le racémique et --- représente une liaison simple ou une liaison double en fonction de $\mathbf{X^1}$ et $\mathbf{X^2}$;

d) la N-acylation du compound de formule (iv) dans laquelle $\mathbf{R^{5'}}$ n'est pas H ou du composé de formule (v), avec un composé de formule (vi)



dans laquelle $\mathbf{R^1}$, $\mathbf{R^{1'}}$, $\mathbf{R^2}$, $\mathbf{R^{2'}}$ et $\mathbf{R^3}$ sont tels que définis dans la revendication 1 ; conduisant au composé de formule (vii)



dans laquelle $\mathbf{R^1}$, $\mathbf{R^{1'}}$, $\mathbf{R^2}$, $\mathbf{R^{2'}}$, $\mathbf{R^3}$, $\mathbf{R^{4'}}$, $\mathbf{X^1}$ et $\mathbf{X^2}$ sont tels que définis ci-dessus ;

*--- représente l'énantiomère (*R*) ou le racémique ;

--- représente une liaison simple ou une liaison double en fonction de $\mathbf{X^1}$ et $\mathbf{X^2}$; et

EP 2 948 455 B1

$R^{5''}$ est R^5 tel que défini dans la revendication 1 ou un 1-((tert-butyldiphénylsilyl)oxy)éthyle ;

e) optionnellement, conduire également l'une ou les deux des étapes suivantes e') et e'') :

- 5 e') quand $R^{4'}$ est un précurseur réductible d'hydroxyéthyle et par conséquent, un précurseur de méthoxyéthyle, une étape de réduction optionnellement suivie par la formation d'un éther méthylique ;
e'') quand $R^{5''}$ est un 1-((tert-butyldiphénylsilyl)oxy)éthyle, une étape de deprotection d'alcool et une fluorination subséquente pour former un groupe R^5 1-fluoroéthyl ; ou une étape de deprotection d'alcool, suivie
10 par une étape d'oxydation et une étape de fluorination subséquente pour former un groupe R^5 1,1-difluoroéthyl ;

pour conduire au composé de formule I selon l'une quelconque des revendications 1 à 14.

15

20

25

30

35

40

45

50

55





FIG. 1



* $p < 0.05$, ** $p < 0.01$ compared to BHP; one-way ANOVA followed by Dunnett's post-hoc test
N=8-10 rats/group

FIG. 2




FIG. 3

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 0043008 A [0007]
- WO 06120478 A [0009]
- WO 2011121137 A [0012]
- WO 2010125102 A [0012]
- WO 2004103953 A [0050]


Non-patent literature cited in the description

- **GIARDINA et al.** *Exp. Opin. Ther. Patents*, 2000, vol. 10 (6), 939-960 [0004]
- *Current Opinion in Investigational Drugs*, 2001, vol. 2 (7), 950-956 [0004]
- **DAWSON ; SMITH.** *Current Pharmaceutical Design*, 2010, vol. 16, 344-357 [0004]
- **MELTZER et al.** *Am. J. Psychiatry*, 2004, vol. 161, 975-984 [0005]
- *Curr. Opin. Invest. Drug*, 2005, vol. 6, 717-721 [0005]
- **KRAJEWSKI et al.** *J. Comp. Neurol.*, 2005, vol. 489 (3), 372-386 [0006]
- **BURKE et al.** *J. Comp. Neurol.*, 2006, vol. 498 (5), 712-726 [0006]
- **GOODMAN et al.** *Endocrinology*, 2004, vol. 145 (6), 2959-2967 [0006]
- **NAVARRO et al.** *J. of Neuroscience*, 2009, vol. 23 (38), 11859-11866 [0007]
- **DASS et al.** *Gastroenterol.*, 2002, vol. 122 (1 [0008]
- **LOMAX ; FURNESS.** *Cell Tissue Res*, 2000, vol. 302, 59-3 [0008]
- **COPEL et al.** *J Physiol*, 2009, vol. 587, 1461-1479 [0008]
- **FIORAMONTI et al.** *Neurogastroenterol Motil*, 2003, vol. 15, 363-369 [0008]
- **SHAFTON et al.** *Neurogastroenterol Motil*, 2004, vol. 16, 223-231 [0008]
- **HOUGHTON et al.** *Neurogastroenterol Motil*, 2007, vol. 19, 732-743 [0008]
- **DUKES et al.** *Gastroenterol*, 2007, vol. 132, A60 [0008]
- **JI Y et al.** *PNAS*, 2011, vol. 108 (35), 14411-14415 [0049] [0172]
- **FUJIWARA Y et al.** *JACS*, 2012, vol. 134, 1494-1497 [0049] [0172]
- Diagnostic and Statistical Manual of Mental Disorders. American Psychiatric Association, 1994 [0126]
- **GORDON, A. J. ; FORD, R. A.** *The Chemist's Companion - A Handbook of Practical Data, Techniques, and References.* Wiley, 1972 [0131]
- Vogel's Textbook of Practical Organic Chemistry. Pearson Prentice Hall, 1989 [0131]
- **SCOLNICK et al.** *J. Andrology*, 1994, vol. 15 (4), 287-297 [0312]
- **RICK et al.** *J. Urol.*, 2012, vol. 187, 1498-1504 [0312]

**N-ACIL-(3- SZUBSZTITUÁLT)-(8- SZUBSZTITUÁLT)-5,6-DIHDRO-
[1,2,4]TRIAZOLO[4,3-A]PIRAZINOK MINT SZELEKTÍV NK-3
RECEPTOR ANTAGONISTÁK, GYÓGYÁSZATI KÉSZÍTMÉNY,
ELJÁRÁSOK NK-3 RECEPTOR ÁLTAL KÖZVETÍTETT
RENDELLENESÉGEK SORÁN TÖRTÉNŐ ALKALMAZÁSRA**

Szabadalmi igénypontok

1. Vegyület, amely a következő I képletű:



vagy annak egy gyógyászatilag elfogadható szolvátja, ahol:

R^1 jelentése H, F vagy metil;

$R^{1'}$ jelentése H;

R^2 jelentése H, F, Cl vagy metoxi;

$R^{2'}$ jelentése H vagy F;

R^3 jelentése H, F, Cl, metil, trifluor-metil vagy nitril;

R^4 jelentése metil, etil, *n*-propil, hidroxil-etil, metoxi-etil, trifluor-metil, difluor-metil vagy fluor-metil;


R^5 jelentése metil, etil, metoxi-metil, trifluor-metil, difluor-metil, fluor-metil, 1-fluor-etil, 1,1-difluor-etil vagy 2,2,2-trifluor-etil;

X^1 jelentése N és X^2 jelentése S vagy O; vagy X^1 jelentése S és X^2 jelentése N;

== jelentése egy egyes vagy egy kettős kötés, X^1 és X^2 jelentésétől függően;

* - - jelentése az I képletű vegyületnek az (R)-enantiomerje vagy a racemátja.

2. Vegyület az 1. igénypont szerint, amely a következő I' és I'' képlettel rendelkezik:



vagy annak egy gyógyászatilag elfogadható szolvátja, ahol R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , R^4 , R^5 , X^1 és X^2 jelentése az 1. igénypontban meghatározottak szerinti; és

== jelentése egy egyes vagy egy kettős kötés, X^1 és X^2 jelentésétől függően.

vagy annak egy gyógyászati lag elfogadható szolvátja, ahol:


R^3 jelentése H, F, Cl, metil, trifluor-metil vagy nitril;

R^4 jelentése metil, etil, *n*-propil, hidroxil-etil, metoxil-etil, trifluor-metil, difluor-metil vagy fluor-metil;

R^5 jelentése metil, etil, metoxil-metil, trifluor-metil, difluor-metil vagy fluor-metil;

* - - - jelentése az Ia-1 képletű vegyületnek az (R)-enantiomerje vagy a racemátja.

6. Vegyület az 1.-től 5.-ig igénypontok bármelyike szerint, amely a következő Ia-2 képlettel rendelkezik:



vagy annak egy gyógyászati lag elfogadható szolvátja, ahol:

R^1 jelentése H, F vagy metil;

$R^{1'}$ jelentése H;

R^2 jelentése H, F, Cl vagy metoxil;


$R^{2'}$ jelentése H vagy F;

R^3 jelentése H, F, Cl, metil, trifluor-metil vagy nitril;

R^5 jelentése metil, etil, metoxil-metil, trifluor-metil, difluor-metil vagy fluor-metil;

* - - - jelentése az Ia-2 képletű vegyületnek az (R)-enantiomerje vagy a racemátja.

7. Vegyület az 1.-től 6.-ig igénypontok bármelyike szerint, amely a következő Ia-3 képlettel rendelkezik:



vagy annak egy gyógyászati lag elfogadható szolvátja, ahol:

R^1 jelentése H, F vagy metil;

$R^{1'}$ jelentése H;

R^2 jelentése H, F, Cl vagy metoxil;


$R^{2'}$ jelentése H vagy F;

R^3 jelentése H, F, Cl, metil, trifluor-metil vagy nitril;

R^4 jelentése metil, etil, *n*-propil, hidroxil-etil, metoxil-etil, trifluor-metil, difluor-metil vagy fluor-metil;

* - - - jelentése az Ia-3 képletű vegyületnek az (R)-enantiomerje vagy a racemátja.

8. Vegyület az 1.-től 7.-ig igénypontok bármelyike szerint, amely a következő Ib képlettel rendelkezik:




vagy annak egy gyógyászatilag elfogadható szolvátja, ahol:

R^3 jelentése F;

R^5 jelentése metil, etil, trifluor-metil, difluor-metil, fluor-metil, 1-fluor-etil, 1,1-difluor-etil vagy 2,2,2-trifluor-etil;


* - - jelentése az Ib képletű vegyületnek az (R)-enantiomerje vagy a racemátja.

9. Vegyület az 1.-től 8.-ig igénypontok bármelyike szerint, amely a következő Ib' képlettel rendelkezik:



vagy annak egy gyógyászatilag elfogadható szolvátja, ahol R^3 és R^5 jelentése a 8. igénypontban meghatározottak szerinti.

10. Vegyület az 1.-től 9.-ig igénypontok bármelyike szerint, amely a következő Ic képlettel rendelkezik:



vagy annak egy gyógyászatilag elfogadható szolvátja, ahol:

R^1 jelentése H, F vagy metil;

$R^{1'}$ jelentése H;

R^2 jelentése H, F, Cl vagy metoxi;

$R^{2'}$ jelentése H vagy F;


R^3 jelentése H, F, Cl, metil, trifluor-metil vagy nitril;

R^4 jelentése metil, etil, *n*-propil vagy hidroxetil;

R^5 jelentése metil, etil vagy trifluor-metil;

* - * jelentése az Ic képletű vegyületnek az (R)-enantiomerje vagy a racemátja.

11. Vegyület az 1.-től 10.-ig igénypontok bármelyike szerint, amely a következő Ic' képlettel rendelkezik:



vagy annak egy gyógyászatilag elfogadható szolvátja, ahol:

R^1 jelentése H, F vagy metil;

$R^{1'}$ jelentése H;

R^2 jelentése H, F, Cl vagy metoxi;

$R^{2'}$ jelentése H vagy F;

R^3 jelentése H, F, Cl, metil, trifluor-metil vagy nitril;






R^4 jelentése metil, etil, *n*-propil vagy hidroxetil;

R^5 jelentése metil, etil vagy trifluor-metil.






12. Vegyület az 1.-től 11.-ig igénypontok bármelyike szerint, amely a következőkből álló csoportból van választva:

1		(R)-(3,4-diklór-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
2		(R)-(3-(3-etil-1,2,4-tiadiazol-5-il)-8-metil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-fenil)metanon






(táblázat folytatása)

3		(R)-(4-klór-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
4		(R)-(4-klór-3-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
5		(R)-(4-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
6		(R)-(3-klór-4-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
7		(R)-(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(3,4,5-trifluor-fenil)metanon






(táblázat folytatása)

8		(R)-(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(2,3,4-trifluor-fenil)metanon
9		(R)-(3,4-difluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
10		(R)-(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(2,3,4,5-tetrafluor-fenil)metanon
11		(R)-(4-fluor-fenil)(8-(2-hidroxi-etil)-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
12		(4-fluor-fenil)(8-(2-hidroxi-etil)-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon






(táblázat folytatása)

13		(R)-(3-(3-etil-1,2,4-oxadiazol-5-il)-8-metil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-fenil)metanon
14		(4-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
15		(R)-(3-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
16		(R)-(3-klór-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
17		(R)-(3,5-difluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon






(táblázat folytatása)

18		(R)-(2,4-difluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
19		(R)-(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(p-tolil)metanon
20		(R)-(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(fenil)metanon
21		(R)-(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-(trifluor-metil)fenil)metanon
22		(R)-(8-etil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-fenil)metanon






(táblázat folytatása)

23		(8-etil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-fenil)metanon
24		(<i>R</i>)-(4-fluor-fenil)(3-(3-metil-1,2,4-tiadiazol-5-il)-8-propil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
25		(<i>R</i>)-(4-fluor-3-metoxi-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
26		(<i>R</i>)-(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(o-tolil)metanon
27		(<i>R</i>)-(3-metoxi-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon






(táblázat folytatása)

28		(R)-(4-fluor-phenil)(8-metil-3-(3-metil-1,2,4-oxadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
29		(R)-4-(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6,7,8-tetrahidro-[1,2,4]triazolo[4,3-a]pirazin-7-karbonil)benzonitril
31		(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(2,3,4,5-tetrafluor-phenil)metanon
32		(3,4-difluor-phenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
33		(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(2,3,4-trifluor-phenil)metanon



(táblázat folytatása)

34		(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(3,4,5-trifluor-fenil)metanon
35		(3-klór-4-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
36		(4-klór-3-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
37		(4-klór-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
38		(3,4-diklór-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon

(táblázat folytatása)

39		(3-(3-etil-1,2,4-tiadiazol-5-il)-8-metil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-fenil)metanon
40		(3-(3-etil-1,2,4-oxadiazol-5-il)-8-metil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-fenil)metanon
41		(<i>R</i>)-(4-fluor-fenil)(8-metil-3-(3-(trifluor-metil)-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
42		(<i>R</i>)-(3-(3-(difluor-metil)-1,2,4-tiadiazol-5-il)-8-metil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-fenil)metanon
43		(<i>R</i>)-(3-(3-(1,1-difluor-etil)-1,2,4-oxadiazol-5-il)-8-metil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-fenil)metanon

(táblázat folytatása)

44		(R)-(4-fluor-phenil)(8-metil-3-(3-(2,2,2-trifluor-etil)-1,2,4-oxadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon
45		((8R)-3-(3-(1-fluor-etil)-1,2,4-oxadiazol-5-il)-8-metil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-phenil)metanon

és azok gyógyászatilag elfogadható szolvátjai.

13. Vegyület az 1.-től 12.-ig igénypontok bármelyike szerint, amely a következőkből álló csoportból van választva:


(R)-(3-(3-etil-1,2,4-tiadiazol-5-il)-8-metil-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)(4-fluor-phenil)metanon;
 (R)-(4-klór-phenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon;
 (R)-(4-fluor-phenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon;
 (4-fluor-phenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon; és
 (R)-(4-fluor-phenil)(8-metil-3-(3-metil-1,2,4-oxadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon,

és azok gyógyászatilag elfogadható szolvátjai.

14. Vegyület az 1.-től 13.-ig igénypontok bármelyike szerint, amely a következő: (R)-(4-fluor-phenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon, vagy annak egy gyógyászatilag elfogadható szolvátja.
15. Gyógyászati készítmény, amely tartalmaz egy az 1.-től 14.-ig igénypontok bármelyike szerinti vegyületet, vagy annak egy gyógyászatilag elfogadható szolvátját, és legalább egy gyógyászatilag elfogadható hordozót, hígítószer, segédanyagot (excipienst) és/vagy adjuvánst.
16. Vegyület az 1.-től 14.-ig igénypontok bármelyike szerint vagy annak egy gyógyászatilag elfogadható szolvátja, alkalmazás céljából egy gyógyszerként.

17. Vegyület az 1.-től 14.-ig igénypontok bármelyike szerint vagy annak egy gyógyászatilag elfogadható szolvátja, alkalmazás céljából a következők kezelésére és/vagy megelőzésére: depresszió, szorongás, pszichózis, skizofrénia, pszichotikus rendellenességek, bipoláris rendellenességek, kognitív rendellenességek, Parkinson-kór, Alzheimer-kór, figyelemhiányos hiperaktivitás zavar (ADHD), fájdalom, görcsök, elhízás (obezitás), gyulladásos betegségek, beleértve a következőket: irritábilis bélszindróma (IBS) és gyulladásos bélzavarok, hányás (emesis), preeklampszia, légutakkal kapcsolatos betegségek, beleértve a következőket: krónikus obstruktív tüdőbetegség, asztma, légúti túlérzékenység, hörgőszűkület (bronchoconstrictio) és köhögés, vizelet-inkontinencia, reprodukciós zavarok, fogamzásgátlás és nemi hormon-függő betegségek, beleértve, de nem kizárólag, a következőket: jóindulatú prosztata hiperplázia (BPH), prosztata hiperplázia, metasztatikus prosztata karcinóma, hererák, emlőrák (mellrák), petefészekrák, androgén-függő akne, férfias típusú hajhullás, endometriózis, abnormalis pubertás (angolul: „abnormal puberty”), méhfibrózis, méh fibroid tumor (mióma), méh-leiomióma, hormon-függő rákok, hiperandrogenizmus, túlzott szőrnövekedés (hirsutizmus), virilizáció, policisztás petefészek szindróma (PCOS), premenstruációs diszfóriás zavar (PMDD), HAIR-AN szindróma (hiperandrogenizmus, inzulin rezisztencia és acanthosis nigricans), petefészek hyperthecosis (HAIR-AN együtt a luteinizált theca-sejtek hiperpláziájával a petefészek strómában), magas petefészekben belüli (intraovariális) androgén koncentrációk más megjelenési formái (például tüsző (follikuláris) érés gátlás, atrézia (atresia), anovuláció, fájdalmas menstruáció (dysmenorrhea), diszfunkcionális méhvérvzés, meddőség), androgén-termelő tumor (virilizáló petefészek tumor vagy virilizáló mellékvese tumor), menorrhagia vagy adenomiózis.
18. Vegyület alkalmazás céljából a 17. igénypont szerint, ahol a vegyület a következő: (R)-(4-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon vagy annak egy gyógyászatilag elfogadható szolvátja.
19. Vegyület az 1.-től 14.-ig igénypontok bármelyike szerint vagy annak egy gyógyászatilag elfogadható szolvátja, alkalmazás céljából hőhullámok kezelésére és/vagy megelőzésére.
20. Vegyület alkalmazás céljából a 19. igénypont szerint, ahol a vegyület a következő: (R)-(4-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon vagy annak egy gyógyászatilag elfogadható szolvátja.
21. Vegyület az 1.-től 14.-ig igénypontok bármelyike szerint vagy annak egy gyógyászatilag elfogadható szolvátja, alkalmazás céljából egy olyan szerként (ágensként), amely csökkenti a cirkuláló LH-szinteket (angolul: „circulating LH levels”).
22. Vegyület alkalmazás céljából a 21. igénypont szerint, ahol a vegyület a következő: (R)-(4-fluor-fenil)(8-metil-3-(3-metil-1,2,4-tiadiazol-5-il)-5,6-dihidro-[1,2,4]triazolo[4,3-a]pirazin-7(8H)-il)metanon vagy annak egy gyógyászatilag elfogadható szolvátja.
23. Eljárás egy az 1.-től 14.-ig igénypontok bármelyike szerinti vegyületnek vagy annak egy gyógyászatilag elfogadható szolvátjának az előállítására, azzal jellemezve, hogy az magában foglalja a következő lépéseket:

a) egy vegyület reagáltatása, amely a következő (i) képletű:



(i)


ahol:

PG jelentése egy alkalmas védőcsoport, mint amilyen például a DMB, PMB, Boc, allil, difenil-foszfinamid vagy 2-trimetil-szilil-etán-szulfonil;

$R^{4'}$ jelentése R^4 , az 1. igénypontban meghatározottak szerint, vagy a hidroxil-etil egy redukálható prekuzora, és következésképpen a metoxi-etil egy további prekuzora;

* - - jelentése az (R)-enantiomer vagy a racemát;

egy vegyülettel, amely a következő (ii) képletű:



(ii)


ahol:

$R^{5'}$ jelentése R^5 , az 1. igénypontban meghatározottak szerint, H vagy 1-((terc-butil-difenil-szilil)oxi)etil;

X^1 és X^2 jelentése az 1. igénypontban meghatározottak szerinti;

== jelentése egy egyes vagy egy kettős kötés, X^1 és X^2 jelentésétől függően;


azon célból, hogy legyen előállítva egy vegyület, amely a következő (iii) képletű:



(iii)

ahol PG, $R^{4'}$, $R^{5'}$, X^1 és X^2 jelentése a fennebb meghatározottak szerinti, * - - jelentése az (R)-enantiomer vagy a racemát és == jelentése egy egyes vagy egy kettős kötés, X^1 és X^2 jelentésétől függően;


b) a védőcsoport eltávolítása a (iii) képletű vegyületből egy alkalmas védőcsoport-eltávolító szerrel (ágenssel), hogy legyen előállítva a következő (iv) képletű vegyület:



(iv)


ahol $R^{4'}$, $R^{5'}$, X^1 és X^2 jelentése a fennebb meghatározottak szerinti, $*$ - - jelentése az (R)-enantiomer vagy a racemát és $\equiv$ - - jelentése egy egyes vagy egy kettős kötés, X^1 és X^2 jelentésétől függően;

c) amikor $R^{5'}$ jelentése H, egy trifluor-metil vagy difluor-metil csoport bevezetése direkt (közvetlen) C-H trifluor- vagy difluor-metilzés (-metiláció) révén, ami a következő (v) képletű vegyülethez vezet:




ahol $R^{4'}$, X^1 és X^2 jelentése a fennebb meghatározottak szerinti és $R^{5'}$ jelentése trifluor-metil vagy difluor-metil, $*$ - - jelentése az (R)-enantiomer vagy a racemát és $\equiv$ - - jelentése egy egyes vagy egy kettős kötés, X^1 és X^2 jelentésétől függően;

d) a (iv) képletű vegyületnek, ahol $R^{5'}$ jelentése nem H, vagy az (v) képletű vegyületnek az N-acilezése, egy olyan vegyülettel, amely a következő (vi) képletű:



ahol R^1 , $R^{1'}$, R^2 , $R^{2'}$ és R^3 jelentése az 1. igénypontban meghatározottak szerinti; ami a következő (vii) képletű vegyülethez vezet:



ahol R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , $R^{4'}$, X^1 és X^2 jelentése a fennebb meghatározottak szerinti; $*$ - - jelentése az (R)-enantiomer vagy a racemát; $\equiv$ - - jelentése egy egyes vagy egy kettős kötés, X^1 és X^2 jelentésétől függően; és $R^{5''}$ jelentése R^5 , az 1. igénypontban meghatározottak szerint, vagy 1-((terc-butildifenil-szillil)oxi)etil;

e) opcionálisan továbbá az egyik vagy mindkettő megvalósítása a két következő e') és e'') lépés közül:

e') amikor $R^{4'}$ jelentése a hidroxietil egy redukálható prekursora, és következésképpen a metoxietil egy további prekursora, egy olyan lépés, amely redukálásból áll, amit opcionálisan metil-éter képzés követ;

e'') amikor $R^{5''}$ jelentése 1-((terc-butil-difenil-szilil)oxi)etil, egy olyan lépés, amely alkohol védőcsoport-eltávolításból és azután következő fluorozásból áll, hogy egy 1-fluor-etil R^5 csoport legyen képezve; vagy egy olyan lépés, amely alkohol védőcsoport-eltávolításból áll, amit egy oxidálási lépés és egy azután következő fluorozási lépés követ, hogy legyen előállítva egy 1,1-difluor-etil R^5 csoport;

hogy legyen előállítva az 1.-től 14.-ig igénypontok bármelyike szerinti I képletű vegyület.