




(21) (A1) **2,235,323**
(86) 1996/11/05
(87) 1997/05/22

- (72) ZAGAR, Cyrill, DE
(72) HARREUS, Albrecht, DE
(72) VON DEM BUSSCHE-HÜNNEFELD, Christoph-Sweder, DE
(72) HAMPRECHT, Gerhard, DE
(72) HEISTRACHER, Elisabeth, DE
(72) SCHÄFER, Peter, DE
(72) MENKE, Olaf, DE
(72) WESTPHALEN, Karl-Otto, DE
(72) MISSLITZ, Ulf, DE
(72) WALTER, Helmut, DE
(71) BASF AKTIENGESELLSCHAFT, DE
(51) Int.Cl.⁶ C07D 231/18, A01N 43/56
(30) 1995/11/15 (195 42 520.0) DE
(54) **1-METHYL-3-PHENYLPYRAZOLES SUBSTITUES ET LEUR
UTILISATION COMME HERBICIDES ET POUR LA
DESSICCATION/DEFOLIATION DE PLANTES**
(54) **SUBSTITUTED 1-METHYL-3-PHENYLPYRAZOLES**



(57) L'invention concerne des 1-méthyl-3-phénylpyrazoles substitués de la formule (I) et leurs sels d'addition d'acide utilisables en agriculture. Dans cette formule (I), R¹ désigne halogénure d'alcoxy C₁-C₄ ou halogénure d'alkylthio C₁-C₄; R² désigne halogène; R³ désigne F ou Cl; R⁴ désigne CN, fluoroalkyle C₁-C₄, fluoroalcényle C₂-C₄ ou -C(R⁵)(X-R⁶)(Y-R⁷); R⁵ désigne H ou alkyle C₁-C₄; X et Y désignent -O- ou

(57) The invention relates to substituted 1-methyl-3-phenylpyrazoles (I) and the agriculturally useful acid addition salts thereof, in which R¹ is C₁-C₄ alkylhalide or C₁-C₄ alkylthiohalide; R² is halogen; R³ is F or Cl; R⁴ is CN, C₁-C₄ fluoroalkyl, C₂-C₄ fluoroalkenyl or -C(R⁵)(X-R⁶)(Y-R⁷); R⁵ is H or C₁-C₄ alkyl; X and Y are -O- or -S-; R⁶, R⁷ is C₁-C₄ alkyl, C₁-C₄ alkylhalide, C₃-C₄ alkenyl, C₃-C₄ alkenylhalide or C₃-C₄ alkynyl. It





(21) (A1) **2,235,323**
(86) 1996/11/05
(87) 1997/05/22

-S-, R⁶, R⁷ désignent alkyle C₁-C₄, halogénure d'alkyle C₁-C₄, alcényle C₃-C₄, halogénure d'alcényle C₃-C₄ ou alkinyle C₃-C₄. Ces composés s'utilisent comme herbicides, ainsi que pour la dessiccation/défoliation de plantes.

also relates to the use thereof as herbicides and for the desiccation or defoliation of plants.



PCT


WELTORGANISATION FÜR GEISTIGES EIGENTUM
Internationales BüroINTERNATIONALE ANMELDUNG VERÖFFENTLICHT NACH DEM VERTRAG ÜBER DIE
INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT)

<p>(51) Internationale Patentklassifikation ⁶ : C07D 231/20, 231/18, A01N 43/56</p>	<p>A1</p>	<p>(11) Internationale Veröffentlichungsnummer: WO 97/18195</p> <p>(43) Internationales Veröffentlichungsdatum: 22. Mai 1997 (22.05.97)</p>
<p>(21) Internationales Aktenzeichen: PCT/EP96/04819</p> <p>(22) Internationales Anmeldedatum: 5. November 1996 (05.11.96)</p> <p>(30) Prioritätsdaten: 195 42 520.0 15. November 1995 (15.11.95) DE</p> <p>(71) Anmelder (für alle Bestimmungsstaaten ausser US): BASF AKTIENGESELLSCHAFT [DE/DE]; D-67056 Ludwigshafen (DE).</p> <p>(72) Erfinder; und</p> <p>(75) Erfinder/Anmelder (nur für US): ZAGAR, Cyrill [DE/DE]; Georg-Herwegh-Strasse 31, D-67061 Ludwigshafen (DE). HARREUS, Albrecht [DE/DE]; Beuthener Strasse 10, D-67063 Ludwigshafen (DE). VON DEM BUSSCHE-HÜNNEFELD, Christoph, Sweder [DE/DE]; Mönchwörthstrasse 39, D-68199 Mannheim 39 (DE). HAMPRECHT, Gerhard [DE/DE]; Rote-Turm-Strasse 28, D-69469 Weinheim (DE). HEISTRACHER, Elisabeth [DE/DE]; D 3,4, D-68159 Mannheim (DE). SCHÄFER, Peter [DE/DE]; Römerstrasse 1, D-67308 Ottersheim (DE). MENKE, Olaf [DE/DE]; Lerchenweg 3, D-67317 Altleiningen (DE). WESTPHALEN, Karl-Otto [DE/DE]; Mausbergweg 58, D-67346 Speyer (DE). MISSLITZ, Ulf [DE/DE]; Am Herzel 40, D-67433 Neustadt (DE).</p>	<p>WALTER, Helmut [DE/DE]; Grünstadter Strasse 82, D-67283 Obrigheim (DE).</p> <p>(74) Gemeinsamer Vertreter: BASF AKTIENGESELLSCHAFT; D-67056 Ludwigshafen (DE).</p> <p>(81) Bestimmungsstaaten: AU, BG, BR, BY, CA, CN, CZ, GE, HU, IL, JP, KE, KR, KZ, LV, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, UZ, VN, ARIPO Patent (KE, LS, MW, SD, SZ, UG), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Veröffentlicht Mit internationalem Recherchenbericht.</p> <p>046360 030113</p>	
<p>(54) Title: SUBSTITUTED 1-METHYL-3-PHENYLPYRAZOLES AND THE USE THEREOF AS HERBICIDES AND FOR THE DESICCATION OR DEFOLIATION OF PLANTS</p>		
<p>(54) Bezeichnung: SUBSTITUIERTE 1-METHYL-3-PHENYLPYRAZOLE UND IHRE VERWENDUNG ALS HERBIZIDE UND ZUR DESIKKATION/DEFOLIATION VON PFLANZEN</p>		
<p>(57) Abstract</p>		
<p>The invention relates to substituted 1-methyl-3-phenylpyrazoles (I) and the agriculturally useful acid addition salts thereof, in which R¹ is C₁-C₄ alkylhalide or C₁-C₄ alkylthiohalide; R² is halogen; R³ is F or Cl; R⁴ is CN, C₁-C₄ fluoroalkyl, C₂-C₄ fluoroalkenyl or -C(R⁵) (X-R⁶) (Y-R⁷); R⁵ is H or C₁-C₄ alkyl; X and Y are -O- or -S-; R⁶, R⁷ is C₁-C₄ alkyl, C₁-C₄ alkylhalide, C₃-4 alkenyl, C₃-C₄ alkenylhalide or C₃-C₄ alkynyl. It also relates to the use thereof as herbicides and for the desiccation or defoliation of plants.</p>		
<p>(57) Zusammenfassung</p>		
<p>Substituierte 1-Methyl-3-phenylpyrazole der Formel (I) sowie deren landwirtschaftlich brauchbaren Säureadditionssalze, wobei R¹ = C₁-C₄-Halogenalkoxy oder C₁-C₄-Halogenalkylthio; R² = Halogen; R³ = F oder Cl; R⁴ = CN, C₁-C₄-Fluoralkyl, C₂-C₄-Fluoralkenyl oder -C(R⁵) (X-R⁶) (Y-R⁷); R⁵ = H oder C₁-C₄-Alkyl; X und Y = -O- oder -S-; R⁶, R⁷ = C₁-C₄-Alkyl, C₁-C₄-Halogenalkyl, C₃-C₄-Alkenyl, C₃-C₄-Halogenalkenyl oder C₃-C₄-Alkynyl. Verwendung: als Herbizide; zur Desikkation/Defoliation von Pflanzen.</p>		
<p style="text-align: right;">(I)</p>		

Vot

Substituted 1-methyl-3-phenylpyrazoles

The present invention relates to new substituted 1-methyl-3-phenylpyrazoles of the formula I



15

where the substituents have the following meanings:

R¹ is C₁-C₄-haloalkoxy or C₁-C₄-haloalkylthio;

20 R² is halogen;

R³ is fluorine or chlorine;

R⁴ is cyano, C₁-C₄-fluoroalkyl, C₂-C₄-fluoroalkenyl or
25 -C(R⁵)(X-R⁶)(Y-R⁷), where

R⁵ is hydrogen or C₁-C₄-alkyl,

X and Y independently of one another are oxygen or sulfur
30 and

R⁶ and R⁷ independently of one another are C₁-C₄-alkyl,
C₁-C₄-haloalkyl, C₃-C₄-alkenyl, C₃-C₄-haloalkenyl or
C₃-C₄-alkynyl;

35

and to the agriculturally useful acid addition salts of I.

Furthermore, the invention relates to


- 40 - the use of the compounds I as herbicides or for the desiccation/defoliation of plants,
- herbicidal compositions and compositions for the desiccation/defoliation of plants which comprise the compounds I as active ingredients,
45 - processes for the preparation of herbicidal compositions and compositions for the desiccation/defoliation of plants using the compounds I, and

2

- methods for controlling undesirable vegetation and for the desiccation/defoliation of plants using the compounds I.

In WO 95/24403, many heterocycles having the common radical
 5 -S(O)₀₋₂-CH₂CH₂CH=CF₂ - inter alia 4-cyano-5-(4,4-difluorobut-3-enylthio)-1-methyl-3-phenylpyrazole - are reported as having nematocidal, insecticidal and acaricidal properties.

Furthermore, JP-A 03/072 460 describes herbicidally active phe-
 10 nylpyrazole derivatives of the formula II




20

where R^a can be, inter alia, halogen-substituted lower alkoxy or lower alkylthio.

Furthermore, JP-A 03/151 367 discloses, as herbicides, 1-lower-
 25 alkyl-3-(2,4-dihalophenyl)-4-halo-5-(haloalkoxy)pyrazoles which have attached to them in the 5-position on the phenyl ring a 1,3-dioxolan-2-yl group, inter alia.

JP 06/199 805 discloses a process for the preparation of herbi-
 30 cidally active 1-H- and 1-lower-alkyl-3-(dihalophenyl)-4-halo- gen-5-(haloalkoxy/haloalkylthio)pyrazoles which can have attached to them in the 5-position on the phenyl ring halogen or lower alkoxy, inter alia. Moreover, there can be found in JP-A 03/163 063 herbicidally active phenylpyrazoles of the general formula
 35 III



45

3

where R^b can be, inter alia, formyl, alkoxy, alkylthio, haloalkoxy, haloalkylthio, lower alkenyloxy, lower alkenylthio, lower haloalkenyloxy, lower haloalkenylthio, lower alkynyloxy or lower alkynylthio.

5

WO 92/02509 relates to pyrazole derivatives which are substituted by phenyl and alkylsulfonyl and which are also reported as having a herbicidal action. Substances which are suitable for their preparation are, inter alia, compounds of type I.

10

Other herbicides of the type of the compounds I can also be seen in the earlier WO 95/34659.

Finally, some of the substituted 1-methyl-3-phenylpyrazoles I formally come under the general formulae of the 3-phenylpyrazoles in the earlier Patent Applications WO 96/01255 and WO 96/15115.

However, the herbicidal properties of the prior art phenylpyrazoles with a view to the harmful plants are not always entirely satisfactory. It was therefore the object of the present invention to provide novel herbicidally active 3-phenylpyrazoles which allow better control of undesirable plants. Another object was to provide novel compounds which have a desiccant or defoliant action.

25

We have found that this object is achieved by the present substituted 1-methyl-3-phenylpyrazoles of the formula I and their herbicidal action. We have furthermore found herbicidal compositions which comprise the compounds I and have a very good herbicidal activity. Moreover, we have found processes for the preparation of these compositions and methods for controlling undesirable vegetation using the compounds I.

We have furthermore found that the compounds I are also suitable for the desiccation/defoliation of parts of plants, suitable plants being crop plants such as cotton, potatoes, oilseed rape, sunflower, soybean or field beans, in particular cotton. Thus, we have found compositions for the desiccation and/or defoliation of plants, processes for the preparation of these compositions and methods for the desiccation and/or defoliation of plants using the compounds I.

Depending on the substitution pattern, the compounds of the formula I can contain one or more chiral centers, in which case they exist in the form of enantiomer or diastereomer mixtures. The in-

vention relates to the pure enantiomers or diastereomers and also to their mixtures.

Agriculturally useful acid addition salts of I are generally to be understood as meaning the salts of I with acids whose anions do not adversely affect the herbicidal action of the compounds I. Accordingly, anions which are mainly suitable are fluoride, chloride, bromide, hydrogen sulfate, sulfate, dihydrogen phosphate, hydrogen phosphate, phosphate, nitrate, hydrogen carbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate, oxalate, dodecylbenzenesulfonate, and the anions of C₁-C₄-alkanoic acids, preferably formate, acetate, propionate and butyrate.

The organic moieties mentioned for the substituents R¹ and R⁴ to R⁷ are collective terms for individual enumerations of the individual group members. All carbon chains, i.e. all alkyl, fluoroalkyl, haloalkyl, haloalkoxy, haloalkylthio, alkenyl, fluoroalkenyl, haloalkenyl and alkynyl moieties, can be straight-chain or branched. Halogenated substituents preferably have attached to them one to five identical or different halogen atoms. The meaning of halogen represents in each case fluorine, bromine, chlorine or iodine, in particular fluorine or chlorine.

Other examples of meanings are, for example:

25

- C₁-C₄-alkyl: methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1-methylpropyl, 2-methylpropyl or 1,1-dimethylethyl, in particular methyl or ethyl;

30 -

C₁-C₄-fluoroalkyl: a C₁-C₄-alkyl radical as mentioned above which is partially or fully substituted by fluorine, eg. CHF₂, CH₂F, CF₃, 2-fluoroethyl, 1,1-difluoroethyl, 2,2,2-trifluoroethyl, 1,1,2,2-tetrafluoroethyl, C₂F₅, 3-fluoropropyl, 1,1-difluoropropyl, 3,3,3-trifluoropropyl, heptafluoropropyl, heptafluoro-1-methylethyl, 4-fluorobutyl, 1,1-difluorobutyl, 4,4,4-trifluorobutyl or nonafluorobutyl, in particular CH₂F, CHF₂, CF₃, C₂F₅ or heptafluoropropyl;

35

40

- C₁-C₄-haloalkyl: a C₁-C₄-alkyl radical as mentioned above which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, eg. 2-fluoroethyl, 2,2,2-trifluoroethyl, 2-chloroethyl, 2,2-dichloroethyl, 2,2,2-trichloroethyl, 2-bromoethyl, 2-iodoethyl, 3-fluoropropyl, 2,3-difluoropropyl, 3,3,3-trifluoropropyl, 2,2,3,3-tetrafluoropropyl, 2,3-dichloro-2,3-difluoropropyl, 3-chloropropyl, 2,3-dichloropropyl, 3,3,3-trichloropropyl, 2,2,3,3-tetrachloropropyl, 3-bromopropyl, 2,3-dibromopropyl,

45

5

- 3-iodopropyl, 4-fluorobutyl, 2,3-difluorobutyl, 3,4-difluorobutyl, 4,4,4-trifluorobutyl, 4-chlorobutyl, 2,3-dichlorobutyl, 3,4-dichlorobutyl, 4,4,4-trichlorobutyl, 4-bromobutyl or 4-iodobutyl, in particular 2-fluoroethyl, 2,2,2-trifluoroethyl, 2-chloroethyl, 3-chloropropyl or 4-chlorobutyl;
- 5
- C₁-C₄-haloalkoxy: C₁-C₄-alkoxy, such as OCH₃, OC₂H₅, OCH₂-C₂H₅, OCH(CH₃)₂, n-butoxy, 1-methylpropoxy, 2-methylpropoxy or OC(CH₃)₃, in particular OCH₃, which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, eg.
- 10 OCH₂Cl, OCH₂F, OCHF₂, OCF₃, chlorodifluoromethoxy, 2-fluoroethoxy, 2-bromoethoxy, 2-iodoethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 2,2,2-trichloroethoxy, OC₂F₅, 3-fluoropropoxy, 3-chloropropoxy, 3-bromopropoxy,
- 15 3-iodopropoxy, 2,3-difluoropropoxy, 3,3,3-trifluoropropoxy, 2,2,3,3-tetrafluoropropoxy, heptafluoropropoxy, heptafluoro-1-methylethoxy, 4-fluorobutoxy or 4-chlorobutoxy, preferably difluoromethoxy, chlorodifluoromethoxy or OCF₃;
- 20 - C₁-C₄-haloalkylthio: C₁-C₄-alkylthio, such as SCH₃, SC₂H₅, SCH₂-C₂H₅, SCH(CH₃)₂, n-butylthio, 1-methylpropylthio, 2-methylpropylthio or SC(CH₃)₃, in particular SCH₃, which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, eg. SCH₂Cl, SCH₂F, SCHF₂, SCF₃,
- 25 chlorodifluoromethylthio, 2-fluoroethylthio, 2-bromoethylthio, 2-iodoethylthio, 2,2-difluoroethylthio, 2,2,2-trifluoroethylthio, 2,2,2-trichloroethylthio, SC₂F₅, 3-fluoropropylthio, 3-chloropropylthio, 3-bromopropylthio, 3-iodopropylthio, 2,3-difluoropropylthio,
- 30 3,3,3-trifluoropropylthio, 2,2,3,3-tetrafluoropropylthio, heptafluoropropylthio, heptafluoro-1-methylethylthio, 4-fluorobutylthio or 4-chlorobutylthio, preferably difluoromethylthio, chlorodifluoromethylthio or SCF₃;
- 35 - C₂-C₄-fluoroalkenyl: C₂-C₄-alkenyl, such as ethenyl, prop-1-en-1-yl, prop-2-en-1-yl, 1-methylethenyl, but-1-en-1-yl, but-2-en-1-yl, but-3-en-1-yl, but-1-en-2-yl, but-3-en-2-yl, 2-methylprop-2-en-1-yl or 2-methylprop-1-en-1-yl, which is partially or fully substituted by
- 40 fluorine, eg. 1-fluoroethenyl, 2-fluoroethenyl, 2,2-difluoroethenyl, 1,2-difluoroethenyl, trifluoroethenyl, 1-fluoroprop-1-en-1-yl, pentafluoroprop-1-en-1-yl, 3,3-difluoroprop-2-en-1-yl, pentafluoroprop-2-en-1-yl, pentafluoro-1-methylethenyl, 1-fluorobut-1-en-1-yl, 4,4,4-tri-
- 45 fluorobut-1-en-1-yl, heptafluorobut-1-en-1-yl, 4,4-difluorobut-3-en-1-yl or 3,3,3-trifluoro-2-methylprop-1-en-1-yl,

6

preferably 1-fluoroethenyl, trifluoroethenyl or pentafluoro-prop-1-en-1-yl;

- 5 - C₃-C₄-alkenyl: prop-2-en-1-yl, but-2-en-1-yl, but-3-en-1-yl, but-3-en-2-yl or 2-methylprop-2-en-1-yl;
- 10 - C₃-C₄-haloalkenyl: C₃-C₄-alkenyl as mentioned above which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, eg. 2-chloroprop-2-en-1-yl, 3-chloroprop-2-en-1-yl, 3,3-dichloroprop-2-en-1-yl, 3-bromoprop-2-en-1-yl, 3-fluoroprop-2-en-1-yl, 4,4-difluorobut-3-en-1-yl or 4,4-dichlorobut-3-en-1-yl;
- 15 - C₃-C₄-alkynyl: prop-2-yn-1-yl, but-2-yn-1-yl, but-3-yn-1-yl or but-3-yn-2-yl.

Preferred with a view to the use of the substituted 1-methyl-3-phenylpyrazoles I as herbicides are those compounds I where the substituents have the following meanings, in each case alone or
20 in combination:


- R¹ is difluoromethoxy, chlorodifluoromethoxy, OCF₃ or difluoromethylthio, in particular difluoromethoxy;
- 25 R² is chlorine or bromine, in particular chlorine;
- R³ is fluorine;
- 30 R⁴ is cyano, fluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, heptafluoropropyl, 1-fluoroethenyl, 2,2-difluoroethenyl, trifluoroethenyl, pentafluoroprop-1-en-1-yl or -C(R⁵)(X-R⁶)(Y-R⁷), in particular fluoromethyl, difluoromethyl, trifluoromethyl or -C(R⁵)(X-R⁶)(Y-R⁷);
- 35 R⁵ is hydrogen, methyl or ethyl, in particular hydrogen;
- X and Y are both oxygen or both sulfur;
- 40 R⁶ and R⁷ independently of one another are C₁-C₄-alkyl, C₁-C₄-haloalkyl or C₃-C₄-alkenyl, in particular methyl, ethyl, n-propyl or allyl.

Very especially preferred are the compounds Ia (\cong I where R¹ =
45 difluoromethoxy; R² and R³ = chlorine) which are listed in Table 1 below:

Table 1

5

10



Ia

No.	-R ⁴
15	Ia.001 -CH(OCH ₃) ₂
	Ia.002 -CH(OC ₂ H ₅) ₂
	Ia.003 -CH(OCH ₂ -CH ₂ F) ₂
	Ia.004 -CH(OCH ₂ -CH ₂ Cl) ₂
20	Ia.005 -CH(OC ₃ H ₇) ₂
	Ia.006 -CH(OC ₄ H ₉) ₂
	Ia.007 -CH(OCH ₂ -CH(CH ₃) ₂) ₂
	Ia.008 -CH(OCH ₂ -CH=CH ₂) ₂
	Ia.009 -CH(OCH ₂ -CH=CHCl) ₂
25	Ia.010 -CH(OCH ₂ -C≡CH) ₂
	Ia.011 -CH(SCH ₃) ₂
	Ia.012 -CH(SC ₂ H ₅) ₂
	Ia.013 -CH(SCH ₂ -CH ₂ F) ₂
30	Ia.014 -CH(SCH ₂ -CH ₂ Cl) ₂
	Ia.015 -CH(SC ₃ H ₇) ₂
	Ia.016 -CH(SC ₄ H ₉) ₂
	Ia.017 -CH(SCH ₂ -CH(CH ₃) ₂) ₂
35	Ia.018 -CH(SCH ₂ -CH=CH ₂) ₂
	Ia.019 -CH(SCH ₂ -CH=CHCl) ₂
	Ia.020 -CH(SCH ₂ -C≡CH) ₂
	Ia.021 -CH(OCH ₃)(SCH ₃)
	Ia.022 -CH(OCH ₃)(SC ₂ H ₅)
40	Ia.023 -CH(OC ₂ H ₅)(SCH ₃)
	Ia.024 -CH(OC ₂ H ₅)(SC ₂ H ₅)
	Ia.025 -C(CH ₃)(OCH ₃) ₂
	Ia.026 -C(CH ₃)(OC ₂ H ₅) ₂
45	Ia.027 -C(CH ₃)(OCH ₂ -CH ₂ F) ₂
	Ia.028 -C(CH ₃)(OCH ₂ -CH ₂ Cl) ₂

No.	-R ⁴
Ia.029	-C(CH ₃)(OC ₃ H ₇) ₂
Ia.030	-C(CH ₃)(OC ₄ H ₉) ₂
5 Ia.031	-C(CH ₃)(OCH ₂ -CH(CH ₃) ₂) ₂
Ia.032	-C(CH ₃)(OCH ₂ -CH=CH ₂) ₂
Ia.033	-C(CH ₃)(OCH ₂ -CH=CHCl) ₂
Ia.034	-C(CH ₃)(OCH ₂ -C≡CH) ₂
10 Ia.035	-C(CH ₃)(SCH ₃) ₂
Ia.036	-C(CH ₃)(SC ₂ H ₅) ₂
Ia.037	-C(CH ₃)(SCH ₂ -CH ₂ F) ₂
Ia.038	-C(CH ₃)(SCH ₂ -CH ₂ Cl) ₂
Ia.039	-C(CH ₃)(SC ₃ H ₇) ₂
15 Ia.040	-C(CH ₃)(SC ₄ H ₉) ₂
Ia.041	-C(CH ₃)(SCH ₂ -CH(CH ₃) ₂) ₂
Ia.042	-C(CH ₃)(SCH ₂ -CH=CH ₂) ₂
Ia.043	-C(CH ₃)(SCH ₂ -CH=CHCl) ₂
20 Ia.044	-C(CH ₃)(SCH ₂ -C≡CH) ₂
Ia.045	-C(CH ₃)(OCH ₃)(SCH ₃)
Ia.046	-C(CH ₃)(OCH ₃)(SC ₂ H ₅)
Ia.047	-C(CH ₃)(OC ₂ H ₅)(SCH ₃)
25 Ia.048	-C(CH ₃)(OC ₂ H ₅)(SC ₂ H ₅)
Ia.049	-CN
Ia.050	-CH ₂ F
Ia.051	-CHF ₂
Ia.052	-CF ₃
30 Ia.053	-CF ₂ -CH ₃
Ia.054	-CH ₂ -CF ₃
Ia.055	-CH ₂ -CH ₂ F
Ia.056	-CHF-CH ₂ F
35 Ia.057	-CF ₂ -CF ₃
Ia.058	-CF ₂ -CHF ₂
Ia.059	-CH ₂ -CH ₂ -CH ₂ F
Ia.060	-CH ₂ -CHF-CH ₂ F
40 Ia.061	-CH ₂ -CH ₂ -CF ₃
Ia.062	-CF ₂ -CH ₂ -CH ₃
Ia.063	-CF ₂ -CF ₂ -CF ₃
Ia.064	-CF(CF ₃) ₂
45 Ia.065	-CH ₂ -CH ₂ -CH ₂ -CH ₂ F
Ia.066	-CF ₂ -CH ₂ -CH ₂ -CH ₃
Ia.067	-CH ₂ -CH ₂ -CH ₂ -CF ₃


No.	-R ⁴
Ia.068	-CF ₂ -CF ₂ -CF ₂ -CF ₃
Ia.069	-CF=CH ₂
5 Ia.070	-CH=CHF
Ia.071	-CH=CF ₂
Ia.072	-CF=CHF
Ia.073	-CF=CF ₂
Ia.074	-CF=CH ₂ -CH ₃
10 Ia.075	-CF=CF ₂ -CF ₃

Other especially preferred 3-phenylpyrazoles of the formulae Ib to Id are in particular

15

the compounds Ib.001 - Ib.075, which differ from the corresponding compounds Ia.001 - Ia.075 only by the fact that R³ is fluorine:

20




Ib

25

30

the compounds Ic.001 - Ic.075, which differ from the corresponding compounds Ia.001 - Ia.075 only by the fact that R² is bromine:

35



Ic


40

the compounds Id.001 - Id.075, which differ from the corresponding compounds Ia.001 - Ia.075 only by the fact that R² is bromine and R³ is fluorine:

45

10

5




Id

10 The substituted 1-methyl-3-phenylpyrazoles of the formula I can be obtained in a variety of ways, in particular following one of the processes below:


A) Reaction of a β -keto-carboxylic acid derivative IV with methylhydrazine in an inert solvent (cf., for example, JP-A 04/225 937 and JP-A 03/072 460) and alkylation of the process product V:

20



25

30



35

L¹ is a customary leaving group, such as halogen, -O-SO₂CH₃, -O-SO₂CF₃, -O-SO₂C₄F₉ and -O-SO₂(p-CH₃-C₆H₄);

40 R⁸ is preferably halogen, C₁-C₄-alkoxy or (C₁-C₄-alkyl)-carbonyloxy;

R⁹ is hydrogen or halogen;

45 R¹⁰ is methyl, bromine or iodine.

The solvent can be aprotic or protic. Suitable examples are organic acids such as acetic acid, hydrocarbons, halogenated hydrocarbons, ethers such as ethylene glycol dimethyl ether, alcohols such as methanol and ethanol, and sulfoxides.

5 However, the process may also be carried out in the absence of a solvent.

10 The reaction temperature is mainly determined by the melting point of the solvent or the compound IV and the boiling point of the reaction mixture. The process is preferably carried out at from approximately 60 to 120°C.

15 In general, approximately 0.95 to 5 times the molar amount, preferably 1 to 1.4 times the amount, of methylhydrazine is employed based on the β -ketocarboxylic acid derivative IV.

20 The amount of alkylating agent L¹-(C₁-C₄-haloalkyl) is usually also 0.95 up to 5 times the molar amount based on the intermediate V.

25 The alkylation is normally carried out with the halide, preferably the chloride or bromide, or with the sulfate of an alkane or haloalkane, if desired in the presence of an organic base, eg. a trialkylamine, or of pyridine, or of an inorganic base, eg. of an alkali metal carbonate.

30 The alkylation is expediently carried out in an inert organic solvent, eg. in an aliphatic or cyclic ether such as 1,2-dimethoxyethane, tetrahydrofuran and dioxane, in an aliphatic ketone such as acetone, in an amide such as dimethylformamide, in a sulfoxide such as dimethyl sulfoxide or in a mixture of one of these solvents and water.


35 The reaction can generally be carried out at from 0°C to the boiling point of the reaction mixture. It is preferably carried out at from approximately 20 to 80°C.

B) Halogenation of the compounds VI where R⁹ = hydrogen:

40

45

12



10 The reaction can be carried out in an inert solvent/diluent or in the absence of a solvent.

15 Examples of suitable solvents are organic acids, inorganic acids, hydrocarbons, halogenated hydrocarbons, aromatic hydrocarbons, ethers, sulfides, sulfoxides and sulfones.

20 Examples of suitable halogenating agents are chlorine, bromine, N-bromosuccinimides, N-chlorosuccinimides or sulfonyl chloride. Depending on the starting compound and the halogenating agent, the addition of a free-radical initiator, for example an organic peroxide such as dibenzoyl peroxide or an azo compound such as azobisisobutyronitrile, or irradiation with light, may have an advantageous effect on the course of the reaction.


25 The amount of halogenating agent is not critical. Both sub-stoichiometric amounts and large excesses of halogenating agent based on the compound VI to be halogenated (where $R^9 =$ hydrogen) are possible.

30 When using a free-radical initiator, a catalytic amount is usually sufficient.

35 The reaction temperature is normally at from (-100) to 200°C , mainly at from 10 to 100°C or the boiling point of the reaction mixture.

40 C) Halogenation of compounds VII where $R^{10} =$ methyl and, if desired, conversion of the process products VIII into compounds I:

45



The halogenation of VII where $R^{10} = \text{CH}_3$ is preferably a chlorination or bromination. As regards suitable solvents, reaction conditions and ratios, reference is made to the information given under B).

For example, the fluorination of VII ($R^{10} = \text{CH}_3$) to compounds I where $R^4 = \text{CH}_2\text{F}, \text{CHF}_2$ or CF_3 can be carried out in a manner known per se by reaction with hydrogen fluoride and chlorine (see, for example, U.S. 4 825 014).

Those halogenation process products which have attached to them chloro-, bromo-, dichloro-, dibromo-, trichloro- or tribromomethyl instead of R^4 can be converted into compounds I where $R^4 = \text{CH}_2\text{F}, \text{CHF}_2$ or CF_3 by means of nucleophilic substitution with fluoride (see, for example, AU 526 817 or P.L. Coe, D. Oldfield, J.C. Tatlow, J. Fluorine Chem. 29 (1985) 341). Examples of fluorides suitable for this purpose are alkali metal fluorides, which are employed in stoichiometric amounts or in an excess (up to approximately five times the molar amount based on the compound to be fluorinated):




The fluorination is preferably carried out in a polar organic solvent such as dimethylformamide and sulfolane. A reaction temperature of from 50 to 300°C is usually sufficient.

Process products VIII which have attached to them a dihalomethyl, preferably a dichloro- or dibromomethyl, group instead of R^4 can furthermore be reacted with alcohols or thiols (preferably thiols) $R^6\text{-XH}$ in the presence of a strong base, such as sodium

14

hydride, or with a salt $R^6\text{-XM}$, M being, in particular, an alkali metal, to give (thio)acetals I $\{R^4 = -\text{CH}(\text{X-R}^6)_2\}$:




As regards suitable solvents, ratios and reaction conditions, the information given in the previous section preferably applies.

15

Compounds VIII which have a dihalomethyl or trihalomethyl radical can furthermore be converted into aldehydes or carboxylic acids IX by means of acid hydrolysis:

20



30

Particularly suitable substances are strong acids, such as sulfuric and hydrochloric acid; the reaction temperature usually being from 50 to 100°C.


35 The aldehydes IX' can then be converted into carboxylic acids IX'' or ketones X in a manner known per se (cf., in this context, for example Houben-Weyl, Methoden der Organischen Chemie [Methods of Organic Chemistry], Georg Thieme Verlag Stuttgart, 4th Edition, Vol. 6/1a, 1980, p. 946 et seq., Vol. 7/2a, 1973, p. 699 et seq. and Vol. 8, 1952, p. 404 et seq.):

40

45

1) In this case, $X=Y$ and $R^6=R^7$.


15



Furthermore, compounds I where R⁴ = cyano are also accessible from the aldehydes IX' in a manner known per se. To this end, IX is


30 first reacted with hydroxylamine to give the oxime, whereupon the latter is dehydrated, for example by means of reaction with di-phosgene (cf., for example, Houben-Weyl, Methoden der Organischen Chemie [Methods of Organic Chemistry], Georg Thieme Verlag Stuttgart, 4th Edition, Vol. 10/4, 1968, p. 55 et seq. and Vol.

35 8, 1952, p. 325 et seq.):




16

- D) Acetalization of the aldehydes IX' or the ketones X in a manner known per se in the presence of an acid catalyst (cf., for example, Houben-Weyl, Methoden der Organischen Chemie [Methods of Organic Chemistry], Georg Thieme Verlag Stuttgart, Vol. 6/3, 4th Edition 1965, p. 204 et seq., p. 221 et seq.; ditto Vol. 9, 4th Edition 1955, p. 199 et seq. and Vol. E14a/3, 1992, p. 415 et seq.):




Examples of suitable acid catalysts are toluenesulfonic acid, hydrochloric acid or sulfuric acid.

The process products can be subjected to a transacetalization in a manner known per se using other alcohols or thiols $\text{R}^7\text{-YH}$ in the presence of catalytic amounts of acid or of a Lewis acid (eg. boron trifluoride) (see, in this context, for example, Houben-Weyl, Methoden der Organischen Chemie [Methods of Organic Chemistry], Georg Thieme Verlag, Stuttgart, Vol. E14a/1, 1991, p. 803 et seq. and Vol. E14a/3, 1992, p. 414 et seq.):



- 30 E) C-C linkage between compounds VII where R^{10} = bromine or iodine and metal cyanides or fluorinated alkyl/alkenyl halides:



Compounds I where R^4 = cyano can be prepared, as a rule, in a manner known per se (see, for example, Houben-Weyl, Methoden der Organischen Chemie [Methods of Organic Chemistry], Georg Thieme Verlag, Stuttgart, Vol. E5, 1985, p. 1459 et seq.) by

2) In this case, thus, $\text{Y}=\text{X}$ and $\text{R}^6=\text{R}^7$.

4) In this case, thus, $\text{X}=\text{Y}$ and $\text{R}^6=\text{R}^7$.

17

reacting VII ($R^{10} = \text{Br}, \text{I}$) with an alkali metal cyanide or with copper cyanide.

5 The process is usually carried out in a polar organic solvent, eg. in dimethylformamide or an ether such as tetrahydrofuran.

10 The reaction is preferably carried out in the presence of a transition-metal catalyst, such as tetrakis(triphenylphosphine)palladium.

To prepare I where $R^4 = \text{fluoroalkyl}$ or fluoroalkenyl , VII ($R^{10} = \text{Br}$ or I) is preferably reacted in a manner known per se with alkali metal perfluoroalkanecarboxylic acid salts and copper(I) iodide {cf. U.S. 4 814 480; K. Matsui, Chem. Lett., 1719 (1981); J. N. Freskos, Synth. Commun. 18 (1988) 965}, with fluoroalkenyl halides, palladium acetate and a base {cf. DE-A 40 13 305; W. Heitz, A. Knebelkamp, Makromol. Chem.-Rap-
20 pid 12 (1991) 69} or with fluoroalkenyl copper or tin compounds in the presence of tetrakis(triphenylphosphine)-palladium {cf. P.L. Heinze, D.J. Burton, J. Fluorine Chem. 31 (1986) 115; D.J. Burton, S.W. Hansen, J. Am. Chem. Soc. 108 (1986) 4229}.

25 Unless otherwise indicated, the reactions described above are expediently carried out under atmospheric pressure or under the inherent pressure of the reaction mixture in question.

30 As a rule, the mixtures are worked up by known methods, for example by diluting the reaction solution with water and subsequently isolating the product by means of filtration, crystallization or solvent extraction, or by removing the solvent, partitioning the residue in a mixture of water and a suitable organic solvent and working up the organic phase to give the product of value.

35 The substituted 1-methyl-3-phenylpyrazoles I can be obtained from their preparation in the form of isomer mixtures, but, if desired, the latter can be resolved into the essentially pure isomers by the methods customary for such cases, such as crystal-
40 lization or chromatography, also on an optically active adsorbate. Pure optically active isomers can be prepared advantageously from suitable optically active starting materials.

45

Agriculturally useful salts of the compounds I can be formed in a manner known per se by means of reaction with an acid of the corresponding anion, preferably of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid or nitric acid.

5

The compounds I and their agriculturally useful salts are suitable, in the form of isomer mixtures and in the form of the pure isomers, as herbicides. The herbicidal compositions comprising I control vegetation on non-crop areas very efficiently, especially at high rates of application. They act against broad-leaved weeds and grass weeds in crops such as wheat, rice, maize, soya and cotton without causing any essential damage to the crop plants. This effect is mainly observed at low rates of application.

15 Depending on the application method in question, the compounds I, or herbicidal compositions comprising them, can additionally be employed in a further number of crop plants for eliminating undesirable plants. Examples of suitable crops are the following:

20 *Allium cepa*, *Ananas comosus*, *Arachis hypogaea*, *Asparagus officinalis*, *Beta vulgaris spec. altissima*, *Beta vulgaris spec. rapa*, *Brassica napus var. napus*, *Brassica napus var. napobrassica*, *Brassica rapa var. silvestris*, *Camellia sinensis*, *Carthamus tinctorius*, *Carya illinoensis*, *Citrus limon*, *Citrus sinensis*, *Coffea arabica (Coffea canephora, Coffea liberica)*,
 25 *Cucumis sativus*, *Cynodon dactylon*, *Daucus carota*, *Elaeis guineensis*, *Fragaria vesca*, *Glycine max*, *Gossypium hirsutum*, (*Gossypium arboreum*, *Gossypium herbaceum*, *Gossypium vitifolium*), *Helianthus annuus*, *Hevea brasiliensis*, *Hordeum vulgare*, *Humulus lupulus*, *Ipomoea batatas*, *Juglans regia*, *Lens culinaris*, *Linum usitatissimum*, *Lycopersicon lycopersicum*, *Malus spec.*, *Manihot esculenta*, *Medicago sativa*, *Musa spec.*, *Nicotiana tabacum (N.rustica)*, *Olea europaea*, *Oryza sativa*, *Phaseolus lunatus*, *Phaseolus vulgaris*, *Picea abies*, *Pinus spec.*, *Pisum sativum*, *Prunus avium*, *Prunus persica*, *Pyrus communis*, *Ribes sylvestre*,
 35 *Ricinus communis*, *Saccharum officinarum*, *Secale cereale*, *Solanum tuberosum*, *Sorghum bicolor (s. vulgare)*, *Theobroma cacao*, *Trifolium pratense*, *Triticum aestivum*, *Triticum durum*, *Vicia faba*, *Vitis vinifera* and *Zea mays*.

40 In addition, the compounds I may also be used in crops which are tolerant to the action of herbicides due to breeding, including genetic engineering methods.

Moreover, the substituted 1-methyl-3-phenylpyrazoles I are also
 45 suitable for the desiccation and/or defoliation of plants.

As desiccants, they are suitable, in particular, for desiccating the aerial parts of crop plants such as potatoes, oilseed rape, sunflowers and soybeans. This allows completely mechanical harvesting of these important crop plants.

5

Also of economic interest is to facilitate harvesting, which is made possible by concentrating, over a period of time, dehiscence, or reducing the adherence to the tree, in citrus fruit, olives or other species and varieties of pomaceous fruit, stone
10 fruit and nuts. The same mechanism, ie. promotion of the formation of abscission tissue between fruit or leaf and shoot of the plants, is also important for readily controllable defoliation of useful plants, in particular cotton.

15 Moreover, shortening the period within which the individual cotton plants mature results in improved fiber quality after harvesting.

The compounds I, or the compositions comprising them, can be used
20 for example in the form of ready-to-spray aqueous solutions, powders, suspensions, also highly-concentrated aqueous, oily or other suspensions or dispersions, emulsions, oil dispersions, pastes, dusts, materials for spreading, or granules, by means of spraying, atomizing, dusting, spreading or pouring. The use forms
25 depend on the intended aims; in any case, they should guarantee the finest possible distribution of the active ingredients according to the invention.

Suitable inert auxiliaries are essentially: mineral oil fractions
30 of medium to high boiling point, such as kerosene and diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, eg. paraffins, tetrahydronaphthalene, alkylated naphthalenes and their derivatives, alkylated benzenes and their derivatives, alcohols such as
35 methanol, ethanol, propanol, butanol and cyclohexanol, ketones such as cyclohexanone, strongly polar solvents, eg. amines such as N-methylpyrrolidone, and water.

Aqueous use forms can be prepared from emulsion concentrates,
40 suspensions, pastes, wettable powders or water-dispersible granules by adding water. To prepare emulsions, pastes or oil dispersions, the substances, either as such or dissolved in an oil or solvent, can be homogenized in water by means of a wetting agent, tackifier, dispersant or emulsifier. Alternatively, it is pos-
45 sible to prepare concentrates comprising active ingredient, wetting agent, tackifier, dispersant or emulsifier and, if

desired, solvent or oil, which are suitable for dilution with water.

Suitable surfactants are the alkali metal salts, alkaline earth
5 metal salts and ammonium salts of aromatic sulfonic acids, eg. ligno-, phenol-, naphthalene- and dibutyl-naphthalenesulfonic acid, and of fatty acids, alkyl- and alkylarylsulfonates, alkyl sulfates, lauryl ether sulfates and fatty alcohol sulfates, and salts of sulfated hexa-, hepta- and octadecanols, and also of
10 fatty alcohol glycol ethers, condensates of sulfonated naphthalene and its derivatives with formaldehyde, condensates of naphthalene, or of the naphthalenesulfonic acids with phenol and formaldehyde, polyoxyethylene octylphenol ether, ethoxylated iso-octyl-, octyl- or nonylphenol, alkylphenyl or tributylphenyl
15 polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers or polyoxypropylene alkyl ethers, lauryl alcohol polyglycol ether acetate, sorbitol esters, lignin-sulfite waste liquors or methylcellulose.

20

Powders, materials for spreading and dusts can be prepared by mixing or grinding the active ingredients together with a solid carrier.

25 Granules, eg coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active ingredients to solid carriers. Solid carriers are mineral earths, such as silicas, silica gels, silicates, talc, kaolin, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium
30 sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders, or other solid carriers.

35

The concentrations of the active ingredients I in the ready-to-use preparations can be varied within wide ranges. In general, the formulations comprise approximately from 0.001 to 98 % by weight, preferably 0.01 to 95 % by weight of at least one active
40 ingredient. The active ingredients are employed in a purity of from 90 % to 100 %, preferably 95 % to 100 % (according to NMR spectrum).

The formulation examples which follow illustrate the preparation
45 of such products:

21

- I. 20 parts by weight of the compound No. Ia.002 are dissolved in a mixture composed of 80 parts by weight of alkylated benzene, 10 parts by weight of the adduct of 8 to 10 mol of ethylene oxide to 1 mol of oleic acid N-monoethanolamide, 5 parts by weight of calcium dodecylbenzenesulfonate and 5 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02 % by weight of the active ingredient.
- II. 20 parts by weight of the compound No. Ib.011 are dissolved in a mixture composed of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the adduct of 7 mol of ethylene oxide to 1 mol of iso-octylphenol and 10 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02 % by weight of the active ingredient.
- III. 20 parts by weight of the active ingredient No. Ib.002 are dissolved in a mixture composed of 25 parts by weight of cyclohexanone, 65 parts by weight of a mineral oil fraction of boiling point 210 to 280°C and 10 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02 % by weight of the active ingredient.
- IV. 20 parts by weight of the active ingredient No. Ib.001 are mixed thoroughly with 3 parts by weight of sodium diisobutyl-naphthalene- α -sulfonate, 17 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 60 parts by weight of pulverulent silica gel, and the mixture is ground in a hammer mill. Finely distributing the mixture in 20,000 parts by weight of water gives a spray mixture which comprises 0.1 % by weight of the active ingredient.
- V. 3 parts by weight of the active ingredient No. Ia.049 are mixed with 97 parts by weight of finely divided kaolin. This gives a dust which comprises 3 % by weight of active ingredient.

22

- VI. 20 parts by weight of the active ingredient No. Ib.052 are mixed intimately with 2 parts by weight of calcium dodecylbenzenesulfonate, 8 parts by weight of fatty alcohol polyglycol ether, 2 parts by weight of the sodium salt of a phenol/urea/formaldehyde condensate and 68 parts by weight of a paraffinic mineral oil. This gives a stable oily dispersion.
- VII. 1 part by weight of the compound No. Ib.051 is dissolved in a mixture composed of 70 parts by weight of cyclohexanone, 20 parts by weight of ethoxylated isooctylphenol and 10 parts by weight of ethoxylated castor oil. The mixture can then be diluted with water to the desired concentration of active ingredient. This gives a stable emulsion concentrate.
- VIII. 1 part by weight of the compound No. Ib.049 is dissolved in a mixture composed of 80 parts by weight of cyclohexanone and 20 parts by weight of Wettol® EM 31 (= nonionic emulsifier based on ethoxylated castor oil). The mixture can then be diluted with water to the desired concentration of active ingredient. This gives a stable emulsion concentrate.
- 25 The active ingredients I or the herbicidal compositions can be applied pre- or post-emergence. If the active ingredients are less well tolerated by certain crop plants, application techniques may be used in which the herbicidal compositions are sprayed, with the aid of the spraying equipment, in such a way that they come into as little contact as possible, if any, with the leaves of the sensitive crop plants, while the active ingredients reach the leaves of undesirable plants growing underneath, or the bare soil surface (post-directed, lay-by).
- 35 The rates of application of active ingredient I are from 0.001 to 3.0, preferably 0.01 to 1.0, kg/ha of active substance (a.s.), depending on the control target, the season, the target plants and the growth stage.
- 40 To widen the spectrum of action and to achieve synergistic effects, the substituted 1-methyl-3-phenylpyrazoles I may be mixed with a large number of representatives of other herbicidal or growth-regulating active ingredients and then applied concomitantly. Suitable components for mixtures are 1,2,4-thiadiazole, 45 1,3,4-thiadiazole, amides, aminophosphoric acid and its derivatives, aminotriazoles, anilides, aryloxy-/hetaryl-oxyalkanoic acids and their derivatives, benzoic acid and its

23

derivatives, benzothiadiazinones, 2-(hetaroyl/aroyl)-1,3-cyclohexanediones, hetaryl aryl ketones, benzyloxazolidinones, meta-CF₃-phenyl derivatives, carbamates, quinolinecarboxylic acid and its derivates, chloroacetanilides, cyclohexane-1,3-dione

5 derivatives, diazines, dichloropropionic acid and its derivatives, dihydrobenzofurans, dihydrofuran-3-ones, dinitroanilines, dinitrophenols, diphenyl ethers, dipyridyls, halocarboxylic acids and their derivatives, ureas, 3-phenyluracils, imidazoles, imidazolinones, N-phenyl-3,4,5,6-tetrahydrophthalimides,

10 oxadiazoles, oxiranes, phenols, aryloxy- and hetaryloxyphenoxypropionic esters, phenylacetic acid and its derivatives, 2-phenylpropionic acid and its derivatives, pyrazoles, phenylpyrazoles, pyridazines, pyridinecarboxylic acid and its derivatives, pyrimidyl ethers, sulfonamides, sulfonylureas,

15 triazines, triazinones, triazolinones, triazolecarboxamides and uracils.

It may furthermore be advantageous to apply the compounds I, alone or in combination with other herbicides, in the form of a

20 mixture with other crop protection agents, for example together with agents for controlling pests or phytopathogenic fungi or bacteria. Also of interest is the miscibility with mineral salt solutions, which are employed for treating nutritional and trace element deficiencies. Non-phytotoxic oils and oil concentrates

25 may also be added.

30

35

40

45

Preparation Examples

Example 1

5 4-Chloro-3-(4-chloro-2-fluoro-5-trifluoromethylphenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole (No. Ib.052)

1 g (2.8 mmol) of 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorobenzoic acid was introduced
10 into a 500 ml autoclave made of Hastelloy C. This was subsequently made inert using nitrogen. The autoclave was placed into a cold bath of dry ice/acetone and charged first with 20 g (1 mol) of hydrogen fluoride and then with 5 g (0.046 mol) of sulfur tetrafluoride. The stirred mixture was subsequently
15 allowed to come to room temperature, and the temperature was then increased to 105°C. Stirring was continued for 10 hours at this internal temperature, during which process a maximum pressure rise to 6.4 bar was observed. After the experiment has ended, the batch was cooled to 25°C, whereupon the mixture was degassed in a
20 washing tower using potassium hydroxide solution. In the autoclave, the residue was then taken up in 50 ml of dichloromethane. The mixture was poured into 50 ml of ice-water. The organic phase was thereupon separated off, after 1 hour washed using 50 ml of water, dried over 4 g of a 1:1 mixture of potassium fluoride/magnesium sulfate and finally concentrated under reduced pressure.
25 The crude product obtained (0.8 g) was purified by chromatography on silica gel (eluent: cyclohexane/methyl tert-butyl ether, ratio 5:1). After the fractions comprising product of value had been combined, the solvent was removed. Yield: 0.45 g of the desired
30 product of value in the form of an amorphous solid.
¹H NMR (250 MHz; in CDCl₃): δ [ppm] = 3.85 (s,3H), 6.70 (t,1H), 7.35 (d,1H), 7.95 (d,1H).

Precursor 1.1

35 5-(4-Chloro-2-fluoro-5-methylphenyl)-1,2-dihydro-2-methyl-3H-pyrazol-3-one

34. 7 g (0.75 mol) of methylhydrazine were added to a solution of 177 g (0.68 mol) of ethyl 3-(4-chloro-2-fluoro-5-methyl-
40 phenyl)-3-oxopropionate in 500 ml of diethylene glycol. After 6 hours at 100°C, the mixture was poured into 4 l of water. The solids were subsequently separated off and dried. Yield: 133 g; m.p.: 155-156°C.

Precursor 1.2

3-(4-Chloro-2-fluoro-5-methylphenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole

5 Gaseous chlorodifluoromethane was passed at room temperature for 2 hours into a solution of 133 g (0.55 mol) of 5-(4-chloro-2-fluoro-5-methylphenyl)-1,2-dihydro-2-methyl-3H-pyrazol-3-one and 110 g (2.7 mol) of sodium hydroxide in 1 l of dioxane and 0.5 l of water. The reaction solution was then poured into 2 l of
10 water, whereupon the aqueous phase was extracted three times using ethyl acetate. The combined organic phases were dried over magnesium sulfate and then filtered and concentrated. The residue was purified by means of column chromatography on silica gel (eluent: cyclohexane/ethyl acetate = 9:1), followed by
15 preparative MPLC on silica gel (same eluent). Yield: 43 g.
 ^1H NMR (250 MHz; in CDCl_3): δ [ppm] = 2.37 (s, 3H), 3.80 (s, 3H), 6.30 (s, 1H), 6.57 (t, 1H), 7.14 (d, 1H), 7.82 (d, 1H).

Precursor 1.3

20 4-Chloro-3-(4-chloro-2-fluoro-5-methylphenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole

22 g (0.16 mol) of sulfuryl chloride were added dropwise to a solution of 43 g (0.15 mol) of 3-(4-chloro-2-fluoro-5-methyl-
25 phenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole in 250 ml of tetrachloromethane. After the reaction solution had subsequently been stirred for 16 hours, 200 ml of saturated aqueous sodium hydrogen carbonate solution were added dropwise. The organic phase was thereupon separated off, washed using saturated sodium
30 chloride solution, dried over magnesium sulfate, then filtered and subsequently concentrated. Yield: 46 g.
 ^1H NMR (250 MHz; in CDCl_3): δ [ppm] = 2.37 (s, 3H), 3.84 (s, 3H), 6.71 (t, 1H), 7.21 (d, 1H), 7.40 (d, 1H).

35 Precursor 1.4

4-Chloro-3-(4-chloro-5-dibromomethyl-2-fluorophenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole

A solution of 46 g (0.14 mol) of 4-chloro-3-(4-chloro-2-fluoro-
40 5-methylphenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole and 62.9 g (0.35 mol) of N-bromosuccinimide in 2 l of tetrachloromethane was irradiated for 3.5 hours with a 1500 watt high-pressure mercury vapor lamp and a UV lamp. The solids were subsequently filtered off and washed twice more using tetrachloromethane. The combined
45 filtrates were then concentrated. Yield: 68 g.

26

¹H NMR (270 MHz; in CDCl₃): δ [ppm] = 3.86 (s, 3H), 6.72 (t, 1H), 7.05 (s, 1H), 7.19 (d, 1H), 8.23 (d, 1H).

Precursor 1.5

- 5 2-Chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorobenzaldehyde

68 g (0.14 mol) of 4-chloro-3-(4-chloro-5-dibromomethyl-2-fluorophenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole were dissolved in
 10 200 ml of concentrated sulfuric acid, with ice-cooling, whereupon the mixture was heated at 100°C until the evolution of gas had ceased. It was then poured into 4 l of ice-water. The mixture was subsequently extracted three times using ethyl acetate. The combined organic phases were washed with water, then dried over
 15 magnesium sulfate and finally concentrated. The crude product was purified by means of column chromatography on silica gel (eluent: hexane/ethyl acetate = 8:1). Yield: 35 g; m.p.: 95-98°C.

Precursor 1.6

- 20 2-Chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorobenzoic acid

A solution of 2.8 g (18 mmol) of sodium dihydrogen phosphate dihydrate in 28 ml of water was added at 10-15°C to a solution of 24
 25 g (71 mmol) of 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorobenzaldehyde in 130 ml of acetonitrile. Then, 6.8 ml of 30 % strength aqueous hydrogen peroxide solution and (in the course of one hour) 9.8 g (108 mmol) of sodium chlorite were added in succession. After the
 30 mixture had been stirred for a further hour, it was heated to room temperature. After it had been stirred for a further 16 hours, it was acidified using dilute hydrochloric acid. The aqueous solution was then extracted twice using ethyl acetate, whereupon the combined organic phases were washed in succession
 35 with concentrated aqueous thiosulfate solution, water and saturated sodium chloride solution, dried over magnesium sulfate and finally concentrated. Yield: 23 g; m.p.: 137-140°C.

Example 2

- 40 4-Chloro-3-(4-chloro-2-fluoro-5-di(methylthio)methylphenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole (No. 1b.011)

1.1 g (15 mmol) of sodium thiomethoxide were added, with ice-cooling, to a solution of 1.5 g (3 mmol) of
 45 4-chloro-3-(4-chloro-5-dibromomethyl-2-fluorophenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole in 30 ml of dimethyl sulfoxide. After 16 hours at room temperature, the mixture was poured into

27

0.5 l of water. The aqueous phase was subsequently extracted twice using ethyl acetate, whereupon the combined organic phases were washed water and saturated aqueous sodium chloride solution, dried over magnesium sulfate and finally concentrated. The crude
5 product was purified by means of silica gel chromatography (eluent: hexane/ethyl acetate = 9:1).

Yield: 0.2 g.

^1H NMR (270 MHz; in CDCl_3): δ [ppm] = 2.14 (s, 6H), 3.84 (s, 3H), 5.28 (s, 1H), 6.70 (t, 1H), 7.25 (d, 1H), 7.86 (d, 1H).

10

Example 3

4-Chloro-3-(4-chloro-2-fluoro-5-di(ethoxy)methylphenyl)-
5-difluoromethoxy-1-methyl-1H-pyrazole (No. 1b.002)

15 20 ml of triethyl orthoformate and 0.1 g of toluenesulfonic acid were added to a solution of 3 g (8.8 mmol) of 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorobenzaldehyde in 50 ml of tetrahydrofuran. After the mixture had been refluxed for 2 hours, it was treated with 3 ml of
20 triethylamine. The mixture was subsequently concentrated. The residue was treated with 50 ml of water. The product was subsequently extracted from the aqueous phase using ethyl acetate. The organic phase was then washed using saturated
aqueous sodium hydrogen carbonate solution, dried over magnesium
25 sulfate and finally concentrated. The crude product was purified by means of column chromatography on silica gel (eluent: hexane/ethyl acetate = 9:1). Yield: 1 g.

^1H NMR (270 MHz; in CDCl_3): δ [ppm] = 1.25 (t, 6H), 3.64 (m, 4H), 3.84 (s, 3H), 5.71 (s, 1H), 6.70 (t, 1H), 7.22 (d, 1H), 7.87 (d, 1H).

30

Example 4

4-Chloro-3-(4-chloro-5-difluoromethyl-2-fluorophenyl)-5-difluoromethoxy-1-methyl-1H-pyrazole (No. Ib.051)

35 0.9 g (5.4 mmol) of diethylaminosulfur tetrafluoride (DAST) were added dropwise at 50°C to 0.9 g (2.7 mmol) of 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorobenzaldehyde. The reaction mixture was subsequently stirred at this temperature for a further 1.5 hours. The batch was then
40 poured onto 10 g of ice, whereupon 10 ml of dichloromethane were added. The organic phase was separated off and washed twice using water (10 ml), then dried and concentrated under reduced pressure. The resulting crude product was purified by
chromatography on 30 g of silica gel (eluent: cyclohexane/methyl
45 tert-butyl ether in a ratio of 4:1). Yield after removal of the solvent: 0.35 g of product of value in the form of a viscous oil.

28

^1H NMR (400 MHz; in CDCl_3): δ [ppm] = 3.83 (s, 3H), 6.72 (t, 1H), 6.94 (t, 1H), 7.28 (d, 1H), 7.90 (d, 1H).

Example 5

5 2-Chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorobenzonitrile (No. Ib.049)


1.2 g (18 mmol) of hydroxylamine hydrochloride and 1.8 g (23 mmol) of pyridine were added to 2 g (5.9 mmol) of
 10 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorobenzaldehyde, whereupon the mixture is dissolved in 100 ml of methanol. The mixture was subsequently stirred first for 16 hours at approximately 20°C and then for another 2 hours at reflux temperature. The mixture was then concentrated. The
 15 residue was treated with water. The reaction product was extracted from the aqueous phase using ethyl acetate. The organic phase was finally dried over magnesium sulfate and concentrated.

The resulting oxime was then first dissolved in 50 ml of
 20 acetonitrile, and 1.3 g (6.7 mmol) of diphosgene were then added to the solution. After the reaction mixture had been stirred for 30 minutes, it was treated with 100 ml of water and stirred for a further 16 hours. The product of value was then extracted using diethyl ether. The ether phase was washed with water, dried over
 25 magnesium sulfate and concentrated. The residue was purified by means of column chromatography on silica gel (eluent: hexane/ethyl acetate = 7:1).
 Yield: 0.6 g; m.p.: $93-95^\circ\text{C}$.

30 Other 1-methyl-3-phenylpyrazoles I which were, or can be, prepared in a similar manner are listed in Table 2 below, in addition to the compounds described above:

Table 2

35



45

No.	R ³	R ⁴	M.p. / ^1H NMR [ppm] / MS [m/z]
Ia.002	Cl	$-\text{CH}(\text{OC}_2\text{H}_5)_2$	28 [M] ⁺
Ia.049	Cl	-CN	84-86°C

29

No.	R ³	R ⁴	M.p. / ¹ H NMR [ppm] / MS [m/z]
Ib.001	F	-CH(OCH ₃) ₂	68-69°C
5 Ib.002	F	-CH(OC ₂ H ₅) ₂	1.25 (t,6H), 3.64 (m,4H), 3.84 (s,3H), 5.71 (s,1H), 6.70 (t,1H), 7.22 (d,1H), 7.87 (d,1H)
Ib.011	F	-CH(SCH ₃) ₂	2.14 (s,6H), 3.84 (s,3H), 5.28 (s,1H), 6.70 (t,1H), 7.25 (d,1H), 7.86 (d,1H)
10 Ib.049	F	-CN	3.85 (s,3H), 6.71 (t,1H), 7.39 (d,1H), 7.93 (d,1H)
Ib.051	F	-CHF ₂	3.83 (s,3H), 6.72 (t,1H), 6.94 (t,1H), 7.28 (d,1H), 7.90 (d,1H)
Ib.052	F	-CF ₃	3.85 (s,3H), 6.70 (t,1H), 7.35 (d,1H), 7.95 (d,1H)

15

Use examples (herbicidal activity)

The herbicidal activity of the substituted 1-methyl-3-phenylpyrazoles I was demonstrated by the following greenhouse experiments:
 20 the culture containers used were plastic flowerpots containing loamy soil with approximately 3.0 % of humus as the substrate. The seeds of the test plants were sown separately for each species.

25 For the pre-emergence treatment, the active ingredients, which had been suspended or emulsified in water, were applied directly after sowing by means of finely distributing nozzles. The containers were irrigated gently to promote germination and growth and subsequently covered with translucent plastic hoods until the
 30 plants had rooted. This cover causes uniform germination of the test plants, unless this was adversely affected by the active ingredients.

For the post-emergence treatment, the test plants were first
 35 grown to a plant height of 3 to 15 cm, depending on the plant habit, and only then treated with the active ingredients which had been suspended or emulsified in water. The test plants for this purpose were either sown directly and grown in the same containers, or they were first grown separately as seedlings and
 40 transplanted into the test containers a few days prior to the treatment. The rate of application for the post-emergence treatment was 0.5 kg/ha of a.s. (active substance).

45

30

Depending on the species, the plants were kept at from 10-25°C or 20-35°C. The test period extended over 2 to 4 weeks. During this time, the plants were tended, and their response to the individual treatments was evaluated.

5

Evaluation was carried out using a scale from 0 to 100. 100 means no emergence of the plants, or complete destruction of at least the aerial parts, and 0 means no damage, or normal course of growth.

10

The plants used in the greenhouse experiments belonged to the following species:

15	Scientific name	Common name
	Echinochloa crus-galli	barnyard grass
	Galium aparine	catchweed bedstraw
	Ipomoea subspecies	morningglory
20	Viola arvensis	field pansy; field violet

The compound No. Ib.052, applied post-emergence, showed a very good activity against the abovementioned weeds at a rate of application of 0.5 kg/ha of a.s.

25

30

35

40

45

Use Examples (desiccant/defoliant activity)

The test plants used were young cotton plants with 4 leaves (without cotyledons) which had been grown under greenhouse conditions (relative atmospheric humidity 50 to 70 %; day/night temperature 27/20°C).

The young cotton plants were subjected to foliar treatment to runoff point with aqueous preparations of the active ingredients (with an addition of 0.15 % by weight of the fatty alcohol alkoxide Plurafac® LF 700⁶⁾, based on the spray mixture). The amount of water applied was 1000 l/ha (converted). After 13 days, the number of leaves shed and the degree of defoliation in % were determined.

15

No leaves were shed in the untreated control plants.

20

25

30

35

40

45 ⁶⁾ a low-foaming nonionic surfactant by BASF AG


1

We claim:

1. A substituted 1-methyl-3-phenylpyrazole of the formula I

5

10



I

15

where the substituents have the following meanings:

- R¹ is C₁-C₄-haloalkoxy or C₁-C₄-haloalkylthio;
- R² is halogen;
- R³ is fluorine or chlorine;
- R⁴ is cyano, C₁-C₄-fluoroalkyl, C₂-C₄-fluoroalkenyl or
-C(R⁵)(X-R⁶)(Y-R⁷), where
- R⁵ is hydrogen or C₁-C₄-alkyl,

25

- X and Y independently of one another are oxygen or sulfur
and

30

- R⁶ and R⁷ independently of one another are C₁-C₄-alkyl,
C₁-C₄-haloalkyl, C₃-C₄-alkenyl, C₃-C₄-haloalkenyl or
C₃-C₄-alkynyl;

35

and to the agriculturally useful acid addition salts of I.

2. A substituted 1-methyl-3-phenylpyrazole of the formula I or
an acid addition salt thereof, as claimed in claim 1, where
R⁴ is cyano.

40

45

3. A substituted 1-methyl-3-phenylpyrazole of the formula I or an acid addition salt thereof, as claimed in claim 1, where R^4 is C_1-C_4 -fluoroalkyl or C_2-C_4 -fluoroalkenyl.
- 5
4. A substituted 1-methyl-3-phenylpyrazole of the formula I or an acid addition salt thereof, as claimed in claim 1, where R^4 is $-C(R^5)(X-R^6)(Y-R^7)$.
- 10 5. The use of a substituted 1-methyl-3-phenylpyrazole I or of an acid addition salt thereof, as claimed in claims 1 to 4, as herbicides or for the desiccation/defoliation of plants.
6. A herbicidal composition comprising a herbicidally active amount of at least one substituted 1-methyl-3-phenylpyrazole of the formula I or of an acid addition salt of I, as claimed in claims 1 to 4, and at least one liquid and/or solid carrier and, if desired, at least one surfactant.
- 15
7. A composition for the desiccation and/or defoliation of plants, comprising such an amount of at least one substituted 1-methyl-3-phenylpyrazole of the formula I or of an acid addition salt of I, as claimed in claims 1 to 4, that it acts as a desiccant and/or defoliant, and at least one inert liquid and/or solid carrier and, if desired, at least one surfactant.
- 20
- 25
8. A process for the preparation of herbicidally active compositions, which comprises mixing a herbicidally active amount of at least one substituted 1-methyl-3-phenylpyrazole of the formula I or of an acid addition salt of I, as claimed in claims 1 to 4, with at least one liquid and/or solid carrier and, if desired, at least one surfactant.
- 30
9. A process for the preparation of compositions which act as desiccants and/or defoliants, which comprises mixing such an amount of at least one substituted 1-methyl-3-phenylpyrazole of the formula I or of an acid addition salt of I, as claimed in claims 1 to 4, that it acts as a desiccant and/or defoliant, with at least one liquid and/or solid carrier and, if desired, at least one surfactant.
- 35
- 40
10. A method for controlling undesirable vegetation, which comprises allowing a herbicidally active amount of at least one substituted 1-methyl-3-phenylpyrazole of the formula I or of
- 45

3

an acid addition salt of I, as claimed in claims 1 to 4, to act on plants, their environment or on seed.

11. A method for the desiccation and/or defoliation of plants,
5 which comprises allowing such an amount of at least one substituted 1-methyl-3-phenylpyrazole of the formula I or of an acid addition salt of I, as claimed in claims 1 to 4, to act on plants that it has a desiccant/defoliant action.

10 12. A method as claimed in claim 11, wherein cotton is treated.

15

20

25

30

35

40

45

