



(86) Date de dépôt PCT/PCT Filing Date: 2004/04/13
 (87) Date publication PCT/PCT Publication Date: 2004/10/28
 (45) Date de délivrance/Issue Date: 2011/06/07
 (85) Entrée phase nationale/National Entry: 2005/10/06
 (86) N° demande PCT/PCT Application No.: EP 2004/003884
 (87) N° publication PCT/PCT Publication No.: 2004/092156
 (30) Priorité/Priority: 2003/04/16 (US60/463,200)

(51) Cl.Int./Int.Cl. *C07D 333/38* (2006.01),
A61K 31/381 (2006.01), *A61K 31/4365* (2006.01),
A61P 3/10 (2006.01), *C07D 211/00* (2006.01),
C07D 311/00 (2006.01), *C07D 333/00* (2006.01),
C07D 333/78 (2006.01), *C07D 495/04* (2006.01)


(72) Inventeurs/Inventors:
 ERICKSON, SHAWN DAVID, US;
 GILLESPIE, PAUL, US;
 GUERTIN, KEVIN RICHARD, US

(73) Propriétaire/Owner:
 F. HOFFMANN-LA ROCHE AG, CH

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : ACETAMIDES DE 3-CYANOTHIOPHENE SUBSTITUES EN TANT QU'ANTAGONISTES DE RECEPTEUR DE
 GLUCAGON

(54) Title: SUBSTITUTED 3-CYANOTHIOPHENE ACETAMIDES AS GLUCAGON RECEPTOR ANTAGONISTS



(57) Abrégé/Abstract:

The present invention relates to compounds of formula (I) wherein R1, R2, R3, R4 and n are as defined in the description and claims, and pharmaceutically acceptable salts thereof. The compounds are useful for the treatment and/or prophylaxis of diseases which are associated with the antagonism of the glucagon receptor, such as diabetes.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
28 October 2004 (28.10.2004)

PCT

(10) International Publication Number
WO 2004/092156 A1

- (51) International Patent Classification⁷: **C07D 333/38**, 333/78, 495/04, A61K 31/381, 31/4365, A61P 3/10 // (C07D 495/04, 333:00, 211:00) (C07D 495/04, 333:00, 311:00)
- (21) International Application Number: PCT/EP2004/003884
- (22) International Filing Date: 13 April 2004 (13.04.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/463,200 16 April 2003 (16.04.2003) US
- (71) Applicant (for all designated States except US): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **ERICKSON, Shawn, David** [US/US]; 122 Oaktree Place, Leonia, NJ 07605 (US). **GILLESPIE, Paul** [GB/US]; 739 Tamaques Way, Westfield, NJ 07090 (US). **GUERTIN, Kevin, Richard** [CA/US]; 14 Wedgewood Drive, Apt. 20, Verona, NJ 07044 (US).
- (74) Agent: **SALUD, Carlos**; Grenzacherstrasse 124, CH-4070 Basel (CH).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED 3-CYANOTHIOPHENE ACETAMIDES AS GLUCAGON RECEPTOR ANTAGONISTS



(57) Abstract: The present invention relates to compounds of formula (I) wherein RI, R2, R3, R4 and n are as defined in the description and claims, and pharmaceutically acceptable salts thereof. The compounds are useful for the treatment and/or prophylaxis of diseases which are associated with the antagonism of the glucagon receptor, such as diabetes.

WO 2004/092156 A1


SUBSTITUTED 3-CYANO THIOPHENE ACETAMIDES AS GLUCAGON RECEPTOR ANTAGONISTS

The present invention is concerned with novel substituted 3-cyanothiophene acetamides, their manufacture and their use as medicaments. The present invention further relates to pharmaceutically acceptable salts of these 3-cyanothiophene compounds and pharmaceutical compositions containing these compounds.

5 The glucagon receptor (GLUR) is a G-protein coupled 7-transmembrane domain receptor (GPCR) of the secretin family. When the natural hormonal ligand glucagon binds to the GLUR, there is an activation of adenylate cyclase and a concomitant increase in cAMP production. This increase in cAMP causes an activation of glycogen phosphorylase resulting in an increase in hepatic glucose production. The actions of glucagon are counter-regulatory to insulin and thus it is believed to play a central role in glucose homeostasis. Glucagon has been used clinically to rescue diabetic patients from hypoglycemia. Thus, a small molecule GLUR antagonists has considerable potential for the treatment of diabetes.

Briefly stated, novel substituted 3-cyanothiophene acetamides have been found to be glucagon receptor antagonists that inhibit glucagon stimulated increase in cAMP production in a functional cell based assay. Consequently, the compounds of the present invention are useful for the treatment and/or prophylaxis of diabetes, and/or impaired glucose tolerance, as well as other conditions wherein the antagonism of the glucagon receptor gives a therapeutic benefit.

20 According to one aspect of the present invention, there is provided a compound of formula (I)



wherein R1, R2, R3, R4 and n are as defined below.

According to another aspect of the present invention, there is provided a pharmaceutical composition comprising a compound of formula (I), or pharmaceutically acceptable salts thereof, and a pharmaceutically acceptable carrier and/or adjuvant.

According to a further aspect of the present invention, there is provided a method for
5 treating or preventing diseases which are associated with antagonism of the glucagon receptor, comprising administering to a patient in need thereof, a therapeutically effective amount of a compound of formula (I).

These and other features, aspects and advantages of the present invention will become better understood with reference to the following drawings, description and claims.

10 Unless otherwise indicated, the following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

In this specification the term "lower" is used to mean a group consisting of one to seven, preferably of one to four carbon atom(s).

The term "halogen" refers to fluorine, chlorine, bromine and iodine, preferably to fluorine
15 and chlorine.

The term "alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of one to twenty carbon atoms, preferably one to sixteen carbon atoms, more preferably one to ten carbon atoms.

Alkyl groups can optionally be substituted e.g. with halogen, hydroxy, lower-alkoxy, lower-
20 alkoxy-carbonyl, NH_2 , $\text{N}(\text{H}, \text{lower-alkyl})$ and/or $\text{N}(\text{lower-alkyl})_2$. Furthermore, substituents as described in this specification come into consideration. Unsubstituted alkyl groups are preferred.

The term "lower-alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent alkyl radical of one to seven carbon atoms, preferably one to
25 four carbon atoms. This term is further exemplified by such radicals as methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, t-butyl and the like. A lower-alkyl group may optionally have a substitution pattern as described earlier in connection with the term "alkyl". Unsubstituted lower-alkyl groups are preferred.

The term "alkoxy" refers to the group $\text{R}'\text{-O-}$, wherein R' is alkyl. The term "lower-alkoxy"
30 refers to the group $\text{R}'\text{-O-}$, wherein R' is lower-alkyl. Examples of lower-alkoxy groups are

e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy and hexyloxy. Alkoxy and lower-alkoxy groups may optionally have a substitution pattern as described earlier in connection with the term "alkyl". Unsubstituted alkoxy and lower-alkoxy groups are preferred.

- 5 The term "cycloalkyl" refers to a monovalent carbocyclic radical of 3 to 10 carbon atom(s), preferably 3 to 6 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. Cycloalkyl groups can optionally be substituted, e.g. by lower-alkyl, lower-alkoxy, halogen, cyano, trifluoromethyl, hydroxy, nitro, amino, alkylamino, dialkylamino, carboxy, aminocarbonyl, phenyl, benzyl, phenoxy and benzyloxy. Unsubstituted cycloalkyl groups
10 are preferred.

The term "aryl" relates to the phenyl or naphthyl group, preferably the phenyl group, which can optionally be mono- or multiply-substituted by lower-alkyl, lower-alkoxy, halogen, CN, CF₃, hydroxy, NO₂, NH₂, N(H, lower-alkyl), N(lower-alkyl)₂, carboxy, aminocarbonyl, phenyl, benzyl, phenoxy, and/or benzyloxy. Preferred substituents are
15 lower-alkyl, lower-alkoxy, halogen, CN, and/or CF₃. Furthermore, substituents as described in this specification come into consideration.


The term "aralkyl" refers to a group aryl-lower-alkyl.

The term "heterocycle" refers to a 4- to 8-membered ring which can comprise 1, 2 or 3 atoms selected from nitrogen, oxygen and/or sulfur such as tetrahydropyridine,
20 dihydrofuran, dihydropyran, furyl, pyrrolyl, pyridyl, 1,2-, 1,3- and 1,4-diazinyl, thienyl, oxazolyl, oxadiazolyl, isoxazolyl, thiazolyl, isothiazolyl or imidazolyl. A heterocycle group may be optionally substituted with an aryl group or have a substitution pattern as described earlier in connection with the term "aryl". A heterocyclic ring may comprise a double bond.

- 25 The term "pharmaceutically acceptable salts" embraces salts of the compounds of formula (I) with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, citric acid, formic acid, maleic acid, acetic acid, fumaric acid, succinic acid, tartaric acid, methanesulfonic acid, p-toluenesulfonic acid and the like, which are non toxic to living organisms. Preferred salts with acids are formates,
30 maleates, citrates, hydrochlorides, hydrobromides and methanesulfonic acid salts.

The term "leaving group" relates to a group which is removed or replaced during a reaction. Examples of leaving groups are halogen, mesylate and tosylate.

In detail, the present invention relates to compounds of formula (I)



wherein

R1 and R2 are independently selected from the group consisting of lower alkyl, lower alkoxy, aminoalkyl, aryl, aralkyl, substituted lower alkyl, substituted lower alkoxy, substituted lower aminoalkyl, substituted aryl and substituted aralkyl, wherein the substituent is selected from the group consisting of one or more of halogen, hydroxy, lower alkoxy, amino, alkylamino, dialkylamino, cyano and nitro; or

R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a 4 – 8 membered, substituted or unsubstituted, carbocyclic or heterocyclic ring, wherein any substituents are independently selected from the group consisting of halogen, hydroxy, lower alkyl, aryl, aralkyl, amino, alkylamino, dialkylamino, alkylsulfonyl, and alkoxy carbonyl;

R3 is selected from the group consisting of lower alkyl, substituted or unsubstituted aryl, substituted or unsubstituted aralkyl, and substituted or unsubstituted cycloalkyl, wherein any substituents are selected from the group consisting of lower alkyl, lower alkoxy, halogen, cyano, trifluoromethyl, hydroxy, nitro, amino, alkylamino, dialkylamino, carboxy, aminocarbonyl, phenyl, benzyl, phenoxy and benzyloxy; and

R4 is selected from the group consisting of lower alkyl, lower alkoxy, halogen, cyano, trifluoromethyl, hydroxy, nitro, amino, alkylamino, dialkylamino, alkylsulfonyl, and alkoxy carbonyl; and

n is 0, 1, 2, 3, 4 or 5;

and pharmaceutically acceptable salts thereof.

Compounds of formula (I) as defined above represent a preferred embodiment of the present invention and pharmaceutically acceptable salts of compounds of formula (I) individually also represent a preferred embodiment of the present invention. Compounds as defined above, which are not salts, are more preferred.

A preferred embodiment of the present invention relates to compounds as defined above, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a 4 – 8 membered, substituted or unsubstituted, carbocyclic or heterocyclic ring, wherein any substituents are selected from the group consisting of halogen, hydroxy, lower alkyl, amino, alkylamino, dialkylamino, 5 alkylsulfonyl, and alkoxy carbonyl.

Another preferred embodiment of the present invention relates to compounds as defined above, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a carbocyclic ring. Preferably, 10 R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a cyclopentenyl or a cyclohexenyl ring.

Furthermore, the present invention relates to compounds as described above, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a substituted or unsubstituted heterocyclic ring. 15 Preferably, R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a six-membered substituted or unsubstituted heterocyclic ring containing at least one heteroatom. More preferably, R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a six-membered substituted or unsubstituted 20 heterocyclic ring containing one heteroatom. Preferably, said heteroatom is a nitrogen. Compounds, wherein said heteroatom is an oxygen are also preferred. Compounds, wherein said heteroatom is nitrogen or oxygen are also preferred.

Such compounds as described above, wherein the heterocyclic ring is substituted with a ring substituent selected from the group consisting of lower alkyl, alkylsulfonyl, 25 alkoxy carbonyl, aryl and aralkyl, are preferred. Particularly preferred are compounds as described above, wherein the ring substituent is attached to the nitrogen atom. Also particularly preferred are compounds as described above, wherein the ring substituent is attached to the nitrogen atom and the ring substituent is lower alkyl, alkoxy carbonyl or alkylsulfonyl.

30 Other preferred compounds as described above are those, wherein R1 and R2 are independently selected from the group consisting of lower alkyl, lower alkoxy, aminoalkyl, aryl, aralkyl, substituted lower alkyl, substituted lower alkoxy, substituted lower

aminoalkyl, substituted aryl and substituted aralkyl. Preferably, R1 and R2 independently are lower alkyl, more preferably methyl or ethyl.

Another preferred embodiment of the present invention relates to compounds as described above, wherein R3 is a substituted or unsubstituted cycloalkyl. Unsubstituted cycloalkyl is
5 preferred. Preferably, R3 is cyclopentyl.

Other preferred compounds as described above are those, wherein R3 is substituted or unsubstituted aryl. Unsubstituted aryl is preferred, particularly phenyl. Furthermore, compounds wherein R3 is substituted or unsubstituted aryl or aralkyl are preferred. Unsubstituted aryl or aralkyl is preferred, particularly phenyl or benzyl.

10 Another preferred embodiment of the present invention relates to compounds of formula (I) as defined above, wherein R3 is a lower alkyl. Compounds as described above, wherein R3 is selected from the group consisting of methyl, ethyl, propyl, isopropyl and sec-butyl are also especially preferred.

In another preferred embodiment of the present invention, n is 0.

15 Preferred compounds of general formula (I) as defined above are those selected from the group consisting of

N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,2-diphenyl-acetamide,

N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,3-diphenyl-propionamide,

N-(3-Cyano-4-methyl-5-ethyl-thiophen-2-yl)-2,3-diphenyl-propionamide,

20 N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2,2-diphenyl-acetamide,

3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,

3-Cyano-2-(2-phenyl-propionyl-amino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,

25 3-Cyano-2-(3-methyl-2-phenyl-butyrylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,

3-Cyano-2-(3-methyl-2-phenyl-pentanoylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,

30 N-(3-Cyano-6-methanesulfonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide,

N-(3-Cyano-6-methyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide,

3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-

- carboxylic acid methyl ester,
 N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2-phenyl-butyramide,
 3-Methyl-2-phenyl-pentanoic acid (3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide,
 5 3-Methyl-2-(2-phenyl-propionylamino)-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide,
 N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2,2-diphenyl-acetamide,
 N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-cyclopentyl-2-phenyl-acetamide,
 10 N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2,2-diphenyl-acetamide, and
 N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-phenyl-propionamide,
 and pharmaceutically acceptable salts thereof.

Other preferred compounds of general formula (I) as defined above are those selected from the group consisting of

- 15 N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,2-diphenyl-acetamide,
 N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,3-diphenyl-propionamide, and
 N-(3-Cyano-4-methyl-5-ethyl-thiophen-2-yl)-2,3-diphenyl-propionamide,
 and pharmaceutically acceptable salts thereof.

Other preferred compounds of general formula (I) as defined above are those selected from the group consisting of

- 20 N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2,2-diphenyl-acetamide,
 3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,
 3-Cyano-2-(2-phenyl-propionylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,
 25 3-Cyano-2-(3-methyl-2-phenyl-butyrylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,
 3-Cyano-2-(3-methyl-2-phenyl-pentanoylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,
 30 N-(3-Cyano-6-methanesulfonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide,
 N-(3-Cyano-6-methyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide,
 3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid methyl ester, and
 35

N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2-phenyl-butylamide,
and pharmaceutically acceptable salts thereof.

Other preferred compounds of general formula (I) as defined above are those selected
from the group consisting of

5 3-Methyl-2-phenyl-pentanoic acid-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide,

3-Methyl-2-(2-phenyl-propionylamino)-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide,

N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2,2-diphenyl-acetamide,

10 N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-cyclopentyl-2-phenyl-acetamide,


N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2,2-diphenyl-acetamide, and

N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-phenyl-propionamide, and
pharmaceutically acceptable salts thereof.

15 Compounds of formula (I) can have one or more asymmetric carbon atoms and can exist
in the form of optically pure enantiomers or as racemates. The invention embraces all of
these forms.


It will be appreciated, that the compounds of general formula (I) in this invention may be
derivatized at functional groups to provide derivatives which are capable of conversion
20 back to the parent compound in vivo.

In a further embodiment, the present invention relates to a process for the manufacture of
compounds of general formula (I) as defined above, which process comprised reacting a
compound of formula (II)



(II)

25 with a compound of formula (III)



wherein R₁, R₂, R₃, R₄ and n are as defined above.

Reactions of compounds of formula (II) with compounds of formula (III) can e.g. be carried out by procedures well known in the art.

- 5 The invention further relates to compounds of formula (I) as defined above, when manufactured by a process as defined above.

As described above, the compounds of formula (I) of the present invention may be used as medicaments for the treatment and/or prophylaxis of diseases mediated by the antagonism of the glucagon receptor. Preferably, the compounds of the present invention may be used to treat diabetes.

- 5 The invention therefore also relates to pharmaceutical compositions comprising a compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

Furthermore, the present invention relates to compounds as defined above for use as therapeutic active substances, particularly for use as therapeutic active substances for the treatment and/or prophylaxis of diseases which are associated with the glucagon receptor,
10 preferably for the treatment and/or prophylaxis of diabetes.

In another embodiment, the present invention relates to a method for the treatment and/or prophylaxis of diseases which are associated with the glucagon receptor, preferably for the treatment and/or prophylaxis of diabetes, which method comprises administering a compound of general formula (I) as defined above to a human being or animal.

- 15 The invention further relates to the use of compounds as defined above for the treatment and/or prophylaxis of diseases which are associated with the glucagons receptor, preferably for the treatment and/or prophylaxis of diabetes.

In addition, the invention relates to the use of compounds as defined above for the preparation of medicaments for the treatment and/or prophylaxis of diseases which are
20 associated with the glucagons receptor, preferably for the treatment and/or prophylaxis of diabetes. Such medicaments comprise a compound as defined above.

The compounds of formula (I) can be manufactured by the methods given below, by the methods given in the examples or by analogous methods. Appropriate reaction conditions for the individual reaction steps are known to the person skilled in the art. Starting
25 materials are either commercially available or can be prepared by methods analogous to the methods given below or in the examples or by methods known in the art.

The following tests were carried out in order to determine the activity of the compounds of formula I.

In general, glucagon antagonists may be identified by their ability to inhibit or reduce stimulation of cAMP production, relative to the cAMP production in the presence of
5 native glucagon alone, as determined in an adenylate cyclase assay. Adenylate cyclase assays are described, for example, by Lin et al. (Biochemistry 14:1559-1563, 1975).

Biological responses via the inositol triphosphate pathway may be assessed by measuring inositol phosphate metabolism as generally described in Subers and Nathanson (J. Mol. Cell. Cardiol. 20:131-140, 1988)
10 or Pittner and Fain (ibid.)
or by measuring the intracellular calcium concentration as generally described by Gryniewicz et al. (J. Biol. Chem. 260:3440-3450, 1985).

In a preferred embodiment, glucagon antagonists, including the 3-cyanothiophene
15 acetamides of the present invention, may be identified through their ability to specifically inhibit the glucagon-induced adenylate cyclase response pathway. Glucagon receptors have been reported in a number of tissues, for example, liver, kidney, cardiac muscle and adipose tissue from a number of species including dog, pig, human and rat. In addition, host cells expressing recombinant glucagon receptors may also be used. Adenylate cyclase
20 activity assays may be carried out using, for example, the method described by Lin et al. (Biochemistry. 14:1559-1563, 1975). These methods measure the level of stimulation of cAMP production relative to native glucagon and generally involve exposing a membrane preparation from tissue containing glucagon receptors to a mixture of glucagon and the glucagon antagonist in the presence of ATP. Membrane preparations from rat liver are
25 generally used for adenylate cyclase activity assays, although other tissues containing glucagon receptors or host cells expressing a recombinant glucagon receptor may be used. Membranes may be prepared using the method described by Neville (Biochim. Biophys Acta 154:540-552, 1968) as modified by Pohl (Methods in Receptor Research, Ed. Blecher, M., New York, pp 160-164, 1976).

cAMP measurement in CHO/GluR12B cells

The glucagon antagonists described herein have been characterized using CHO-K1 cells overexpressing the full length human glucagon receptor (CHO/GluR 12B). Cells were plated in a 384well plate at a density of 10000 cells/well in medium containing DMEM (Gibco #21063-029), 10% dialyzed FBS (Gibco#26400-044), 1% L-Glutamine (Gibco #25030-081) and 1% Pen/Strep (Gibco # 15140-122). The cells were allowed to adhere overnight at 37°C. The media was removed and the cells were then pre-incubated with antagonist solubilized in DMSO diluted in DMEM containing 0.5mM IBMX (Calbiochem #410957), 1mg/ml BSA (Sigma #A-8806), 25mM HEPES for 1hr. Glucagon (0.1nM) diluted in the same medium is then added for 30 min at 37°C. The formation of cAMP is measured in the cell lysates using a protocol and reagents purchased from Applied Biosystems for their cAMP-Screen™ System. An IC₅₀ of a glucagon receptor antagonist is calculated by plotting the dose response of antagonist versus the percent of maximum cAMP generated by 0.1nM glucagon using Microsoft XLFit equation #205 (Sigmoidal Dose Response w/ variable slope). Provided the appropriate concentrations of antagonist are used to obtain saturation at both the high and low concentrations, the midpoint of the curve is extrapolated as the IC₅₀.

The IC₅₀ values for the inhibition of glucagon stimulated cAMP production as measured in the above described cell based assay for each of the compounds described in the examples below are ≤ 10.0 μM.

The following table shows the IC₅₀ values for the inhibition of glucagon stimulated cAMP production (cell based assay) for some of the compounds of the present invention.

Example	IC ₅₀ (μM)	Example	IC ₅₀ (μM)
Example 3	0.7	Example 11	2.5
Example 4	1.5	Example 13	2.5
Example 5	2.3	Example 14	1.0
Example 8	1.0	Example 17	5.0

The compounds of formula I and/or their pharmaceutically acceptable salts can be used as medicaments, e.g. in the form of pharmaceutical preparations for enteral, parenteral or topical administration. They can be administered, for example, perorally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions, rectally, e.g. in the form of suppositories, parenterally, e.g. in the form of injection solutions or infusion solutions, or topically, e.g. in the form of ointments, creams or oils. Oral administration is preferred.

The production of the pharmaceutical preparations can be effected in a manner which will be familiar to any person skilled in the art by bringing the described compounds of formula I and/or their pharmaceutically acceptable salts, optionally in combination with other therapeutically valuable substances, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

Suitable carrier materials are not only inorganic carrier materials, but also organic carrier materials. Thus, for example, lactose, corn starch or derivatives thereof, talc, stearic acid or its salts can be used as carrier materials for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carrier materials for soft gelatine capsules are, for example, vegetable oils, waxes, fats and semi-solid and liquid polyols (depending on the nature of the active ingredient no carriers might, however, be required in the case of soft gelatine capsules). Suitable carrier materials for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar and the like. Suitable carrier materials for injection solutions are, for example, water, alcohols, polyols, glycerol and vegetable oils. Suitable carrier materials for suppositories are, for example, natural or hardened oils, waxes, fats and semi-liquid or liquid polyols. Suitable carrier materials for topical preparations are glycerides, semi-synthetic and synthetic glycerides, hydrogenated oils, liquid waxes, liquid paraffins, liquid fatty alcohols, sterols, polyethylene glycols and cellulose derivatives.

Usual stabilizers, preservatives, wetting and emulsifying agents, consistency-improving agents, flavor-improving agents, salts for varying the osmotic pressure, buffer substances, solubilizers, colorants and masking agents and antioxidants come into consideration as pharmaceutical adjuvants.

The dosage of the compounds of formula I can vary within wide limits depending on the disease to be controlled, the age and the individual condition of the patient and the mode of administration, and will, of course, be fitted to the individual requirements in each

particular case. For adult patients a daily dosage of about 1 to 1000 mg, especially about 1 to 100 mg, comes into consideration. Depending on severity of the disease and the precise pharmacokinetic profile the compound could be administered with one or several daily dosage units, e.g. in 1 to 3 dosage units.

- 5 The pharmaceutical preparations conveniently contain about 1-500 mg, preferably 1-100 mg, of a compound of formula I.

The following Examples serve to illustrate the present invention in more detail. They are, however, not intended to limit its scope in any manner.


General Methods

The compounds of formula (I) can be manufactured by the methods given below, by the methods outlined in the examples or by analogous methods.

Scheme 1, below, generally describes the synthesis of compounds of formula (I).


- 5 Thiophene (b) may be formed by contacting ketone (a) with malononitrile and sulfur in an alcoholic solvent in the presence of a secondary amine. The compound of formula (I) may then be formed by contacting acetyl chloride (c) with thiophene (b).

Scheme 1:



- 10 Derivatization of compounds of formula (I) may produce additional compounds within the scope of the present invention. Scheme 2, below, shows one example of various compounds produced from a t-BOC-tetrahydrothienopyridine compound (d). Removal of the t-BOC group yields the amine (e). Amine (e) may be functionalized by any method known to one of skill in the art. For example, amine (e) may be converted into an alkyl
- 15 amine (f), carbamate (g) or sulfonate (h).


Scheme 2:



Examples

Example 1

N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,2-diphenyl-acetamide



5 Step 1: Preparation of 2-Amino-3-cyano-4,5-dimethylthiophene.


To a stirred mixture of methylethyl ketone (4.51 mL; 50.0 mmol), malononitrile (3.34 g; 50.0 mmol) and sulfur (1.60 g; 50.0 mmol) in absolute ethanol (20 mL) under nitrogen at 0°C was added diethylamine (5.00 mL) dropwise. The mixture was then warmed to 45 - 50°C and stirred for 4 hours. The mixture was then allowed to cool to room temperature,
10 filtered to remove insoluble material and concentrated *in vacuo*. The crude product was chromatographed (Merck Silica gel 60, 230-400 mesh, eluent: 20% ethylacetate / hexanes) to provide 2.35 g (31%) of 2-Amino-3-cyano-4,5-dimethylthiophene as a light brown foam.

Step 2: Preparation of N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,2-diphenyl-acetamide

15 To a stirred solution of 2-Amino-3-cyano-4,5-dimethylthiophene (152 mg; 1.0 mmol) and diphenylacetyl chloride (300 mg; 1.3 mmol) in dry methylene chloride (10 mL) under nitrogen at room temperature was added triethylamine (0.42 mL; 3.0 mmol) dropwise. After 1.5 h, the mixture was washed with 1N HCl. The organic layer was dried over sodium sulfate, filtered and concentrated *in vacuo*. The crude product was triturated with
20 cold 30% ethylacetate/hexanes and filtered to give 162 mg (47%) of N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,2-diphenyl-acetamide as a white solid. ES-HRMS m/e calcd for C₂₁H₁₈N₂OS (M+H⁺) 347.1213, found 347.1217.

Example 2

N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,3-diphenyl-propionamide




EDCI (61 mg; 0.32 mmol) was added to a solution of 2-Amino-3-cyano-4,5-dimethylthiophene (prepared in Example 1, Step 1; 24 mg; 0.148 mmol) and 2,3-diphenylpropionic acid (72 mg; 0.32 mmol). After stirring for 19 h, the reaction mixture was applied directly to a silica gel column (Merck Silica gel 60, 230-400 mesh, eluent: 10% - 33% ethylacetate / hexanes) to provide N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,3-diphenyl-propionamas a white solid (17.0 mg; 39%) .

ES-HRMS m/e calcd for C₂₂H₂₀N₂OS (M+H⁺) 361.1369, found 361.1374.

10 Example 3


N-(3-Cyano-4-methyl-5-ethyl-thiophen-2-yl)-2,3-diphenyl-propionamide



By replacing the methylethylketone of Example 1 with methylpropylketone, the compound of Example may be prepared.

15 Example 4

N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2,2-diphenyl-acetamide



Step 1: Preparation of 2-Amino-3-cyano-4,7-dihydro-5H-thieno[2,3-c]pyran.


To a stirred mixture of dihydro-4H-pyran-4-one (5.00g; 50.0 mmol), malononitrile (3.34g; 50.0 mmol) and sulfur (1.60g; 50.0 mmol) in absolute ethanol (20 mL) under nitrogen at 0°C was added diethylamine (5.00 mL) dropwise. The mixture was then warmed to 45 - 50
 5 °C and stirred for 4.5 hours. The mixture was then allowed to cool to room temperature. The solid was filtered, washed several times with cold ethanol to provide 6.09g (68%) of 2-Amino-3-cyano-4,7-dihydro-5H-thieno[2,3-c]pyran as a tan colored solid.

Step 2: Preparation of of N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2,2-diphenyl-acetamide.

10 To a stirred solution of 2-Amino-3-cyano-4,7-dihydro-5H-thieno[2,3-c]pyran (400mg; 2.22 mmol) and triethylamine (0.93 mL; 6.66 mmol) in dry methylene chloride (10 mL) under nitrogen at room temperature was added diphenylacetyl chloride (768 mg; 3.33 mmol). After 72 h, the mixture was diluted with methylene chloride and washed with 1N HCl. The organic layer was dried over sodium sulfate, filtered and concentrated *in vacuo*.
 15 The crude product was triturated with cold diethyl ether and filtered to give 310 mg (37%) of N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2,2-diphenyl-acetamide as a white solid. ES-HRMS m/e calcd for C₂₂H₁₈N₂O₂S (M+H⁺) 375.1162, found 375.1165.

Example 5

20 3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester



Step 1: Preparation of 2-Amino-3-Cyano-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester.

25 To a stirred mixture of tert-butyl 4-oxo-1-piperidinecarboxylate (15.96g; 80.0 mmol), malononitrile (5.28g; 80.0 mmol) and sulfur (2.56g; 80.0 mmol) in absolute ethanol (40


mL) under nitrogen at 0°C was added diethylamine (8.00 mL) dropwise. The mixture was then warmed to room temperature for 1 hour and then further heated to 45 - 50 °C for 1 h. The mixture was then allowed to cool to room temperature and an additional 10 mL of ethanol was added. The resulting slurry cooled to 0°C and then filtered to collect the
5 product. The product was washed three times with 10 mL of cold ethanol to provide 15.75g (71 %) of 2-amino-3-cyano-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester as a light orange colored solid.

Step 2: Preparation of 3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-
10 c]pyridine-6-carboxylic acid tert-butyl ester.

To a stirred solution of 2-Amino-3-Cyano-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-
carboxylic acid tert-butyl ester (8.00g; 28.7 mmol) and triethylamine (10.01 mL; 71.8
mmol) in dry methylene chloride (50 mL) under nitrogen at room temperature was added
diphenylacetyl chloride (9.92 g; 43.0 mmol). After 18 h, the mixture was diluted with
15 methylene chloride and washed with 1N HCl. The organic layer was dried over sodium
sulfate, filtered and concentrated *in vacuo*. The crude product was chromatographed
(eluent: 35% ethyl acetate/hexanes to give 13.60 g (100%) of 3-Cyano-2-
diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-
butyl ester as a foam. ES-HRMS m/e calcd for C₂₇H₂₇N₃O₃S (M+H⁺) 474.1846, found
20 474.1849.

Example 6

3-Cyano-2-(2-phenyl-propionyl-amino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-
carboxylic acid tert-butyl ester



25

Step 1: Preparation of 2-phenylpropionyl chloride


To a stirred solution of 2-phenylpropionic acid (0.36 mL; 2.6 mmol) and a catalytic amount of dry dimethylformamide (ca. 10 μ L) in 3.0 mL of dry methylene chloride under nitrogen at room temperature was added oxalyl chloride (0.34 mL; 3.9 mmol) dropwise. After 30 minutes, the mixture was concentrated *in vacuo* and used immediately for Step 2.

5 Step 2: Preparation of 3-Cyano-2-(2-phenyl-propionylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester

To a stirred solution of 2-Amino-3-Cyano-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester (418 mg; 1.5 mmol; preparation outlined in Example 4, Step 1) and potassium carbonate (1106 mg; 8.0 mmol) in 10 mL of dry diethyl ether under
 10 nitrogen at room temperature was added a solution of 2-phenylpropionyl chloride (Prepared in Example 5, step 1) in 3.0 mL of dry methylene chloride. The mixture was allowed to stir for 22 h and then concentrated *in vacuo*. The residue was taken up into methylene chloride and washed with 10% potassium carbonate solution. The organic layer was dried over sodium sulfate, filtered and concentrated *in vacuo* to give 813 mg of
 15 product. Trituration with cold anhydrous diethyl ether provided 353 mg (54%) of 3-Cyano-2-(2-phenyl-propionylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester as an off white solid. ES-HRMS m/e calcd for $C_{22}H_{35}N_3O_3SNa$ ($M+Na^+$) 434.1509, found 434.1509.

20 Example 7

3-Cyano-2-(3-methyl-2-phenyl-butyrylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester



Step 1: Preparation of 3-methyl-2-phenylbutyryl chloride

25 To a stirred solution of 3-methyl-2-phenylbutyric acid (891 mg; 5.0 mmol) and a catalytic amount of dry dimethylformamide (ca. 10 μ L) in 10 mL of dry methylene chloride under


nitrogen at room temperature was added oxalyl chloride (0.65 mL; 7.5 mmol) dropwise. After 1 hour, the mixture was concentrated *in vacuo* and used immediately for Step 2.

Step 2: Preparation of 3-Cyano-2-(3-methyl-2-phenyl-butyrylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester

- 5 Reaction of 3-methyl-2-phenylbutyryl chloride (Prepared in Example 6, Step1) and of 2-Amino-3-Cyano-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester (836 mg; 3.0 mmol) according to the procedure outlined in Example 5 for 24 h at room temperature provided after workup, crude product which was subjected to flash chromatography (Merck Silica gel 60, 230-400 mesh; Eluent: gradient 20 to 30% ethyl acetate / hexanes) to provide 513 mg of 3-Cyano-2-(3-methyl-2-phenyl-butyrylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester as a white solid and 444 mg of recovered starting material. ES-HRMS m/e calcd for C₂₄H₃₉N₃O₃SNa (M+Na⁺) 462.1822, found 462.1825.

15 Example 8

3-Cyano-2-(3-methyl-2-phenyl-pentanoylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester



Step 1: Preparation of 3-methyl-2-phenylvaleryl chloride

- 20 To a stirred solution of 3-methyl-2-phenylvaleric acid (1.23g; 6.2 mmol) and a catalytic amount of dry dimethylformamide (ca. 20 μ L) in 5 mL of dry methylene chloride under nitrogen at room temperature was added oxalyl chloride (4.8 mL of 2.0 M solution in methylene chloride) dropwise. After 45 minutes, the mixture was concentrated *in vacuo* and used immediately for Step 2.


- 25 Step 2: Preparation of 3-Cyano-2-(3-methyl-2-phenyl-pentanoylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester

Reaction of 3-methyl-2-phenylvaleryl chloride (Prepared in Example 7, Step1) and of 2-Amino-3-Cyano-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester (869 mg; 3.1 mmol) according to the procedure outlined in Example 5 for 15 h at room temperature provided after workup, crude which was subjected to flash chromatography (Biotage 40M; eluent: gradient 5 to 10% ethyl acetate / hexanes) to provide 597 mg of 3-Cyano-2-(3-methyl-2-phenyl-pentanoylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester as a mixture of diastereomers and 209 mg of recovered starting material. ES-HRMS m/e calcd for $C_{25}H_{31}N_3O_3SNa$ ($M+Na^+$) 476.1978, found 476.1984.

10

Example 9

N-(3-Cyano-6-methanesulfonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide



15 Step 1: Preparation of N-(3-Cyano-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide

Trifluoroacetic acid (20 mL) was added to a stirred solution of 3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester (prepared in Example 4, Step 2; 13.60g) in dry methylene chloride under nitrogen at 0°C. After 5 minutes, the mixture was allowed to warm to room temperature and stirred for 1 hr. The mixture was concentrated in vacuo and the oily residue was triturated with dry diethyl ether and filtered. The filter cake was washed with dry ether (3 X 100 mL) and allowed to air dry to give 14.50g of N-(3-Cyano-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide as the trifluoacetate salt.


25 Recrystallization from tetrahydrofuran/ether provided 8.75g of of N-(3-Cyano-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide as the trifluoacetate salt

Step 2: Preparation of N-(3-Cyano-6-methanesulfonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide

Methanesulfonic anhydride (90 mg; 0.517 mmol) was added dropwise to a solution of N-(3-Cyano-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide (140 mg; 0.342 mmol) and Et₃N (200 μL; 1.43 mmol) in CH₂Cl₂ (6 mL). After stirring for 4.5 h, the reaction mixture was diluted with ethyl acetate and washed once with water and twice with brine. The combined aqueous layers were extracted twice with ethyl acetate. The combined organic layers were dried over Na₂SO₄. Filtration followed by removal of all volatiles *in vacuo* yielded a waxy solid from which N-(3-Cyano-6-methanesulfonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide (100 mg; 65%) was isolated by flash chromatography (40%-66% ethyl acetate in hexane). ES-HRMS m/e calcd for C₂₃H₂₁N₃O₃S₂ (M+H⁺) 452.1097, found 452.1096.

Example 10


N-(3-Cyano-6-methyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide



To a stirred solution of N-(3-Cyano-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide (prepared in Example 8, step 1; 75 mg; 0.15 mmol) and formaldehyde (0.1 mL of 37% solution) in dry methanol at room temperature was added sodium cyanoborohydride (14.5 mg; 0.23 mmol). The mixture was allowed to stir for 1 hour at room temperature then concentrated in vacuo. The residue was taken up into methylene chloride and washed with water. The organic layer was dried (sodium sulfate) and concentrated in vacuo to give 50 mg (85%) of N-(3-Cyano-6-methyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide. ES-HRMS m/e calcd for C₂₃H₂₁N₃OS (M+H⁺) 388.1478, found 388.1481.

Example 11


3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid methyl ester



Methylchloroformate (5 μ L; 0.04 mmol) was added dropwise to a solution of N-(3-Cyano-
 5 4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide (20 mg; 0.06 mmol)
 and Et₃N (200 μ L; 1.43 mmol) in CH₂Cl₂ (6 mL). After stirring for 3.5 h, the mixture was
 concentrated in vacuo and the crude product purified by by flash chromatography (Merck
 Silica gel 60, 230-400 mesh; Eluent: gradient 50% to 66% ethyl acetate / hexanes) to
 provide 11 mg (64%) as an oil. ES-HRMS m/e calcd for C₂₄H₂₁N₃O₃Sa (M+H⁺) 432.1377,
 10 found 432.1381.

Example 12

3-Methyl-2-phenyl-pentanoic acid (3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide



15

Step 1: Preparation of 2-Amino-3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophene.

Diethylamine (2.50 mL) was added dropwise to a stirred solution of cyclopentanone (2.21
 mL; 25.0 mmol), malononitrile (1.67g; 25.0 mmol) and sulfur (0.80g; 25.0 mmol) in
 absolute ethanol (10 mL) under nitrogen at 0°C. After 10 min., the mixture was allowed
 20 to warm to room temperature and stirred for 3h. The mixture was then concentrated in
 vacuo and the crude residue was chromatographed (Merck Silica gel 60, 230-400 mesh;

Eluent: 30% ethyl acetate / hexanes) to provide 1.18g of 2-Amino-3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophene as a light brown foam.

Step 2: Preparation of 3-Methyl-2-phenyl-pentanoic acid (3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide

- 5 Reaction of 3-methyl-2-phenylvaleryl chloride (Prepared in Example 7, Step1) and 2-Amino-3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophene (164 mg; 1.0 mmol) according to the procedure outlined in Example 5 for 20 h at room temperature provided after workup, crude which triturated with diethylether to give 94 mg of crude 3-Methyl-2-phenyl-pentanoic acid (3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide
- 10 which was further purified by dissolving in methylene chloride and passing through a short plug of Merck Silica gel 60 to provide 82 mg (24%) of 3-Methyl-2-phenyl-pentanoic acid (3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide as a mixture of diastereomers. ES-HRMS m/e calcd for C₂₆H₃₈N₂O₂ (M+H⁺) 411.3006, found 411.3011.

15 **Example 13**

3-Methyl-2-(2-phenyl-propionylamino)-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide

By replacing the 3-methyl-2-phenylvaleryl chloride of Example 12 with 2-phenylpropionyl chloride, the compound of Example may be prepared.

20


Example 14

N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2,2-diphenyl-acetamide

By replacing the cyclopentanone of Example 12 with cyclohexanone, the compound of Example may be prepared

25

Example 15


N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-cyclopentyl-2-phenyl-acetamide5 Step 1: Preparation of α -phenylcyclopentaneacetyl chloride.

To a stirred solution of α -phenylcyclopentaneacetic acid (422 mg; 2.0 mmol) and a catalytic amount of dry dimethylformamide (ca. 20 μ L) in 5 mL of dry methylene chloride under nitrogen at room temperature was added oxalyl chloride (1.5 mL of 2.0 M solution in methylene chloride) dropwise. After 45 minutes, the mixture was concentrated *in vacuo* to give crude α -phenylcyclopentaneacetyl chloride which was used immediately for Step 2.

Step 2: Preparation of N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-cyclopentyl-2-phenyl-acetamide


Reaction of α -phenylcyclopentaneacetyl chloride (Prepared in Example 12, Step1) and 2-Amino-3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophene (164 mg; 1.0 mmol) according to the procedure outlined in Example 5 for 20 h at room temperature provided after workup, crude product which triturated with diethylether to give 111 mg of which was further purified by recrystallization from ethylacetate / hexanes to provide 44 mg (%) of N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-cyclopentyl-2-phenyl-acetamide as a light brown solid. EI-HRMS *m/e* calcd for $C_{21}H_{32}N_2OS$ (M^+) 350.1453, found 350.1456.

Example 16

N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2,3-diphenyl-propionamide


EDCI (59 mg; 0.308 mmol) was added to a solution of 2-Amino-3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophene (25 mg; 0.152 mmol) and 2,3-diphenylpropionic acid (70 mg; 0.309 mmol). After stirring for 19 h, the reaction mixture was applied directly to a silica gel column (Merck Silica gel 60; eluent: 10% - 33% ethyl acetate / hexanes) to provide 13.0 mg (23%) of N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2,3-diphenylpropionamide as a white solid. EI-HRMS m/e calcd for $C_{22}H_{18}N_2OS$ ($M+H^+$) 358.1140, found 358.1140.

Example 17

N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-phenyl-propionamide

EDCI (42 mg; 0.219 mmol) was added to a solution of 2-Amino-3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophene (18 mg; 0.11 mmol) and 2-phenylpropionic acid (30 μ L; 0.22 mmol). After stirring for 19 h, the reaction mixture was applied directly to a silica gel column (Merck Silica gel 60; eluent: 12.5% - 33% ethyl acetate / hexanes) to provide 3.5 mg (11%) of N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-phenylpropionamide as a white solid. ES-HRMS m/e calcd for $C_{17}H_{16}N_2OS$ ($M+H^+$) 297.1056, found 297.1058.

Example 18

N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2-phenyl-butamide

To a stirred suspension of 2-Amino-3-cyano-4,7-dihydro-5H-thieno[2,3-c]pyran (361 mg;
5 2.0 mmol; prepared as described in Example 2, Step 1) and potassium carbonate (1.40g;
10.1 mmol) in diethylether (6.0 mL) was added 2-phenylbutyryl chloride dropwise. The
resulting mixture was stirred at room temperature for 24h. The solvent was then removed
in vacuo, the residue taken up into methylene chloride and washed with water. The
organic layer was dried (sodium sulfate) filtered and concentrated *in vacuo* to give crude
10 product which was triturated with diethyl ether to provide 428 mg (66%) of N-(3-Cyano-
4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2-phenyl-butamide as a beige colored solid.
EI-HRMS m/e calcd for C₁₈H₁₈N₂O₂S (M⁺) 326.1809, found 326.1809.

Examples**Example A**

Film coated tablets containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per tablet</u>	
Kernel:		
Compound of formula (I)	10.0 mg	200.0 mg
Microcrystalline cellulose	23.5 mg	43.5 mg
Lactose hydrous	60.0 mg	70.0 mg
Povidone K30	12.5 mg	15.0 mg
Sodium starch glycolate	12.5 mg	17.0 mg
Magnesium stearate	1.5 mg	4.5 mg
(Kernel Weight)	120.0 mg	350.0 mg
Film Coat:		
Hydroxypropyl methyl cellulose	3.5 mg	7.0 mg
Polyethylene glycol 6000	0.8 mg	1.6 mg
Talc	1.3 mg	2.6 mg
Iron oxyde (yellow)	0.8 mg	1.6 mg
Titan dioxide	0.8 mg	1.6 mg

5

The active ingredient is sieved and mixed with microcrystalline cellulose and the mixture is granulated with a solution of polyvinylpyrrolidone in water. The granulate is mixed with sodium starch glycolate and magnesium stearate and compressed to yield kernels of 120 or 350 mg respectively. The kernels are lacquered with an aqueous solution / suspension of the above mentioned film coat.

10

Example B

Capsules containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per capsule</u>
Compound of formula (I)	25.0 mg
Lactose	150.0 mg

WO 2004/092156

PCT/EP2004/003884

Maize starch	20.0 mg
Talc	5.0 mg

The components are sieved and mixed and filled into capsules of size 2.

Example C

5 Injection solutions can have the following composition:

Compound of formula (I)	3.0 mg
Polyethylene Glycol 400	150.0 mg
Acetic Acid	q.s. ad pH 5.0
Water for injection solutions	ad 1.0 ml

The active ingredient is dissolved in a mixture of Polyethylene Glycol 400 and water for injection (part). The pH is adjusted to 5.0 by Acetic Acid. The volume is adjusted to 1.0 ml by addition of the residual amount of water. The solution is filtered, filled into vials using
10 an appropriate overage and sterilized.

Example D

Soft gelatin capsules containing the following ingredients can be manufactured in a conventional manner:

Capsule contents

Compound of formula (I)	5.0 mg
Yellow wax	8.0 mg
Hydrogenated Soya bean oil	8.0 mg
Partially hydrogenated plant oils	34.0 mg
Soya bean oil	110.0 mg
Weight of capsule contents	165.0 mg
Gelatin capsule	
Gelatin	75.0 mg
Glycerol 85 %	32.0 mg
Karion 83	8.0 mg (dry matter)
Titan dioxide	0.4 mg
Iron oxide yellow	1.1 mg

- 5 The active ingredient is dissolved in a warm melting of the other ingredients and the mixture is filled into soft gelatin capsules of appropriate size. The filled soft gelatin capsules are treated according to the usual procedures.

Example E

- 10 Sachets containing the following ingredients can be manufactured in a conventional manner:

Compound of formula (I)	50.0 mg
Lactose, fine powder	1015.0 mg
Microcrystalline cellulose (AVICEL PH 102)	1400.0 mg
Sodium carboxymethyl cellulose	14.0 mg
Polyvinylpyrrolidon K 30	10.0 mg
Magnesiumstearate	10.0 mg
Flavoring additives	1.0 mg


The active ingredient is mixed with lactose, microcrystalline cellulose and sodium carboxymethyl cellulose and granulated with a mixture of polyvinylpyrrolidone in water. The granulate is mixed with magnesium stearate and the flavouring additives and filled into sachets.

- 5 The amount of active compound of formula (I) in the above examples is for illustrative purposes. The actual amount of the compound of formula (I) may vary depending on the patient and intended use or effect.

It should be understood, of course, that the foregoing relates to preferred embodiments of the invention and that modifications may be made without departing from the spirit and
10 scope of the invention as set forth in the following claims.

What is claimed is:

1. A compound of formula (I)



wherein

R1 and R2 are independently selected from the group consisting of C₁-C₇ alkyl, C₁-C₇ alkoxy, aminoalkyl having 1 to 20 carbon atoms, phenyl, naphthyl, phenyl-C₁-C₇-alkyl, naphthyl-C₁-C₇-alkyl, substituted C₁-C₇ alkyl, substituted C₁-C₇ alkoxy, substituted C₁-C₂₀ aminoalkyl, substituted phenyl or naphthyl and substituted phenyl-C₁-C₇-alkyl or naphthyl-C₁-C₇-alkyl, wherein the substituent is selected from the group consisting of one or more of halogen, hydroxy, C₁-C₇ alkoxy, amino, alkylamino having 1 to 20 carbon atoms, dialkylamino, each alkyl group having 1 to 20 carbon atoms, cyano and nitro; or

R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a 4 – 8 membered, substituted or unsubstituted, carbocyclic or heterocyclic ring, which heterocyclic ring comprises 1, 2 or 3 atoms selected from nitrogen, oxygen or sulfur, wherein any substituents are independently selected from the group consisting of halogen, hydroxy, C₁-C₇ alkyl, phenyl, naphthyl, phenyl-C₁-C₇-alkyl, naphthyl-C₁-C₇-alkyl, amino, alkylamino having 1 to 20 carbon atoms, dialkylamino, each alkyl group having 1 to 20 carbon atoms, alkylsulfonyl having 1 to 20 carbon atoms, and C₁-C₂₀ alkoxy carbonyl;

R3 is selected from the group consisting of C₁-C₇ alkyl, substituted or unsubstituted phenyl or naphthyl, substituted or unsubstituted phenyl-C₁-C₇-alkyl or naphthyl-C₁-C₇-alkyl, and substituted or unsubstituted C₃-C₁₀ cycloalkyl, wherein any substituents are selected from the group consisting of C₁-C₇ alkyl, C₁-C₇ alkoxy, halogen, cyano, trifluoromethyl, hydroxy, nitro, amino, alkylamino having 1 to 20 carbon atoms,

dialkylamino, each alkyl group having 1 to 20 carbon atoms, carboxy, aminocarbonyl, phenyl, benzyl, phenoxy and benzyloxy; and

R4 is selected from the group consisting of C₁-C₇ alkyl, C₁-C₇ alkoxy, halogen, cyano, trifluoromethyl, hydroxy, nitro, amino, alkylamino having 1 to 20 carbon atoms, dialkylamino, each alkyl group having 1 to 20 carbon atoms, alkylsulfonyl having 1 to 20 carbon atoms, and C₁-C₂₀ alkoxy carbonyl; and

n is 0, 1, 2, 3, 4 or 5;

or a pharmaceutically acceptable salt thereof,

with the proviso that the compound of formula (I) is not

N-(3-cyano-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide,

N-(3-cyano-4,7-dihydro-5,5-dimethyl-5H-thieno[2,3-c]pyran-2-yl)- α -phenyl-benzeneacetamide,

N-(3-cyano-6-ethyl-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide,

N-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thien-2-yl)- α -phenyl-benzeneacetamide, or

N-(3-cyano-5,6,7,8-tetrahydro-4H-cyclohepta[b]thien-2-yl)- α -phenyl-benzeneacetamide.

2. The compound according to claim 1, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a 4 – 8 membered, substituted or unsubstituted, carbocyclic or heterocyclic ring, which heterocyclic ring comprises 1, 2 or 3 atoms selected from nitrogen, oxygen or sulfur, wherein any substituents are selected from the group consisting of halogen, hydroxy, C₁-C₇ alkyl, amino, alkylamino having 1 to 20 carbon atoms, dialkylamino, each alkyl group having 1 to 20 carbon atoms, alkylsulfonyl having 1 to 20 carbon atoms, and C₁-C₂₀ alkoxy carbonyl.

3. The compound according to any one of claims 1 to 2, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a 4-8 membered carbocyclic ring.

4. The compound according to claim 3, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a cyclopentenyl or a cyclohexenyl ring.
5. The compound according to any one of claims 1 to 2, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a substituted or unsubstituted 4-8 membered heterocyclic ring, which heterocyclic ring comprises 1, 2 or 3 atoms selected from nitrogen, oxygen or sulfur.
6. The compound according to claim 5, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a six-membered substituted or unsubstituted heterocyclic ring containing at least one heteroatom selected from nitrogen, oxygen or sulfur.
7. The compound according to claim 6, wherein R1 and R2 are taken together with the carbon atoms to which they are attached and the bond between these carbon atoms to form a six-membered substituted or unsubstituted heterocyclic ring containing one heteroatom selected from nitrogen, oxygen or sulfur.
8. The compound according to claim 7, wherein the heteroatom is a nitrogen atom.
9. The compound according to any one of claims 5 to 8, wherein the heterocyclic ring is substituted with a ring substituent selected from the group consisting of C₁-C₇ alkyl, alkylsulfonyl having 1 to 20 carbon atoms, C₁-C₂₀ alkoxy carbonyl, phenyl, naphthyl, phenyl-C₁-C₇-alkyl and naphthyl-C₁-C₇-alkyl.
10. The compound according to claims 8 and 9, wherein the ring substituent is attached to the nitrogen atom and the ring substituent is C₁-C₇ alkyl, C₁-C₂₀ alkoxy carbonyl or C₁-C₂₀ alkylsulfonyl.
11. The compound according to claim 1, wherein R1 and R2 are independently selected from the group consisting of C₁-C₇ alkyl, C₁-C₇ alkoxy, aminoalkyl having 1 to 20 carbon atoms, phenyl, naphthyl, phenyl-C₁-C₇-alkyl, naphthyl-C₁-C₇-alkyl, substituted C₁-C₇ alkyl, substituted C₁-C₇ alkoxy, substituted C₁-C₇ aminoalkyl,

substituted phenyl or naphthyl and substituted phenyl-C₁-C₇-alkyl or naphthyl-C₁-C₇-alkyl.

12. The compound according to claim 11, wherein R1 and R2 independently are C₁-C₇ alkyl.

13. The compound according to any one of claims 1 to 12, wherein R3 is a substituted or unsubstituted C₃-C₁₀ cycloalkyl.

14. The compound according to claim 13, wherein R3 is cyclopentyl.

15. The compound according to any one of claims 1 to 12, wherein R3 is substituted or unsubstituted phenyl, naphthyl, phenyl-C₁-C₇-alkyl or naphthyl-C₁-C₇-alkyl.

16. The compound according to claim 15, wherein R3 is phenyl or benzyl.

17. The compound according to any one of claims 1 to 12, wherein R3 is C₁-C₇ alkyl.

18. The compound according to claim 17, wherein R3 is selected from the group consisting of methyl, ethyl, propyl, isopropyl and sec-butyl.

19. The compound according to any one of claims 1 to 18, wherein n is 0.

20. The compound according to any one of claims 1 to 19, selected from the group consisting of

N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,2-diphenyl-acetamide,

N-(3-Cyano-4,5-dimethyl-thiophen-2-yl)-2,3-diphenyl-propionamide,

N-(3-Cyano-4-methyl-5-ethyl-thiophen-2-yl)-2,3-diphenyl-propionamide,

N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2,2-diphenyl-acetamide,

3-Cyano-2-diphenylacetylamino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,

3-Cyano-2-(2-phenyl-propionylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,

3-Cyano-2-(3-methyl-2-phenyl-butyrylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,

3-Cyano-2-(3-methyl-2-phenyl-pentanoylamino)-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid tert-butyl ester,

N-(3-Cyano-6-methanesulfonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide,

N-(3-Cyano-6-methyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridin-2-yl)-2,2-diphenyl-acetamide,

3-Cyano-2-diphenylacetyl-amino-4,5,6,7-tetrahydro-5H-thieno[2,3-c]pyridine-6-carboxylic acid methyl ester,

N-(3-Cyano-4,7-dihydro-5H-thieno[2,3-c]pyran-2-yl)-2-phenyl-butylamide,

3-Methyl-2-phenyl-pentanoic acid-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide,


3-Methyl-2-(2-phenyl-propionylamino)-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-amide,

N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-cyclopentyl-2-phenyl-acetamide,


N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2,3-diphenyl-propionamide, and

N-(3-Cyano-5,6-dihydro-4H-cyclopenta[b]thiophen-2-yl)-2-phenyl-propionamide, or a pharmaceutically acceptable salt thereof.

21. A process for the manufacture of the compound defined in any one of claims 1 to 20, which process comprises reacting a compound of formula (II)



with a compound of formula (III)



wherein R1, R2, R3, R4 and n are as defined in any one of claims 1 to 20.

22. The compound according to any one of claims 1 to 20 when manufactured by the process defined in claim 21.
23. A pharmaceutical composition comprising the compound defined in any one of claims 1 to 20 or a compound selected from the group consisting of N-(3-cyano-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-4,7-dihydro-5,5-dimethyl-5H-thieno[2,3-c]pyran-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-6-ethyl-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thien-2-yl)- α -phenyl-benzeneacetamide, or N-(3-cyano-5,6,7,8-tetrahydro-4H-cyclohepta[b]thien-2-yl)- α -phenyl-benzeneacetamide, and a pharmaceutically acceptable carrier and/or adjuvant.
24. The compound according to any one of claims 1 to 20 or a compound selected from the group consisting of N-(3-cyano-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-4,7-dihydro-5,5-dimethyl-5H-thieno[2,3-c]pyran-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-6-ethyl-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thien-2-yl)- α -phenyl-benzeneacetamide, or N-(3-cyano-5,6,7,8-tetrahydro-4H-cyclohepta[b]thien-2-yl)- α -phenyl-benzeneacetamide, for use as therapeutically active substance.
25. The compound according to any one of claims 1 to 20 or a compound selected from the group consisting of N-(3-cyano-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-4,7-dihydro-5,5-dimethyl-5H-thieno[2,3-c]pyran-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-6-ethyl-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thien-2-yl)- α -phenyl-benzeneacetamide, or N-(3-cyano-5,6,7,8-tetrahydro-4H-cyclohepta[b]thien-2-yl)- α -phenyl-benzeneacetamide, for use as a therapeutically active substance for the treatment and/or prophylaxis of diabetes.
26. A use of the compound defined in any one of claims 1 to 20 or a compound selected from the group consisting of N-(3-cyano-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-4,7-dihydro-5,5-dimethyl-5H-

thieno[2,3-c]pyran-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-6-ethyl-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thien-2-yl)- α -phenyl-benzeneacetamide, or N-(3-cyano-5,6,7,8-tetrahydro-4H-cyclohepta[b]thien-2-yl)- α -phenyl-benzeneacetamide, for the preparation of a medicament for the treatment and/or prophylaxis of diabetes.

27. A use of the compound defined in any one of claims 1 to 20 or a compound selected from the group consisting of N-(3-cyano-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-4,7-dihydro-5,5-dimethyl-5H-thieno[2,3-c]pyran-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-6-ethyl-4,5,6,7-tetrahydrobenzo[b]thien-2-yl)- α -phenyl-benzeneacetamide, N-(3-cyano-5,6-dihydro-4H-cyclopenta[b]thien-2-yl)- α -phenyl-benzeneacetamide, or N-(3-cyano-5,6,7,8-tetrahydro-4H-cyclohepta[b]thien-2-yl)- α -phenyl-benzeneacetamide, for the treatment and/or prophylaxis of diabetes.

