


(19)



(11)

**EP 1 123 313 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention  
of the grant of the patent:  
**28.02.2007 Bulletin 2007/09**

(51) Int Cl.:  
**C07K 14/505<sup>(2006.01)</sup> A61P 7/06<sup>(2006.01)</sup>**  
**C12N 5/10<sup>(2006.01)</sup> C12N 15/12<sup>(2006.01)</sup>**

(21) Application number: **99955046.0**

(86) International application number:  
**PCT/US1999/024435**

(22) Date of filing: **18.10.1999**

(87) International publication number:  
**WO 2000/024893 (04.05.2000 Gazette 2000/18)**

(54) **METHODS AND COMPOSITIONS FOR THE PREVENTION AND TREATMENT OF ANEMIA**  
**METHODEN UND ZUSAMMENSETZUNGEN ZUR PRÄVENTION UND BEHANDLUNG DER ANÄMIE**  
**METHODES ET COMPOSITIONS PERMETTANT DE PREVENIR ET DE TRAITER L'ANEMIE**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(56) References cited:  
**EP-A- 0 267 678 WO-A-91/05867**  
**WO-A-95/05465 WO-A-99/11781**  
**US-A- 5 541 458 US-A- 5 716 644**

(30) Priority: **23.10.1998 US 178292**

(43) Date of publication of application:  
**16.08.2001 Bulletin 2001/33**

(73) Proprietor: **AMGEN INC.**  
**Thousand Oaks, CA 91320-1799 (US)**

(72) Inventors:  
• **EGRIE, Joan, C.**  
**Newbury Park, CA 91320 (US)**  
• **ELLIOTT, Steven, G.**  
**Newbury Park, CA 91320 (US)**  
• **BROWN, Jeffrey, K.**  
**Camarillo, CA 93010 (US)**

(74) Representative: **Grünecker, Kinkeldey,**  
**Stockmair & Schwanhäusser**  
**Anwaltssozietät**  
**Maximilianstrasse 58**  
**80538 München (DE)**

- **FIBI M R ET AL: "N- AND O-GLYCOSYLATION MUTEINS OF RECOMBINANT HUMAN ERYTHROPOIETIN SECRETED FROM BHK-21 CELLS" BLOOD, vol. 85, no. 5, 1 March 1995 (1995-03-01), pages 1229-1236, XP002053700**
- **EGRIE J C ET AL.: "The role of carbohydrate on the biological activity of erythropoietin" GLYCOCONJUGATE JOURNAL, 15 - 20 August 1993, page 263 XP000906829 Abstracts of XIIth International Symposium cited in the application**
- **ELLIOTT STEVE ET AL.: "Structural requirements for addition of O-linked carbohydrate to recombinant erythropoietin." BIOCHEMISTRY, vol. 33, no. 37, 17 September 1994 (1994-09-17), pages 11237-11245, XP000906783**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 1 123 313 B1**

**Description**Field of the Invention

5 [0001] The invention relates to increasing hematocrit in a mammal using hyperglycosylated analogs of erythropoietin. More particularly, the invention relates to less frequent dosing of a hyperglycosylated analog compared to recombinant human erythropoietin to raise and maintain hematocrit and treat anemia. The administration of lower amounts of a hyperglycosylated analog compared to recombinant human erythropoietin at an equivalent dosing frequency in order to raise and maintain hematocrit and treat anemia is also described. New hyperglycosylated analogs of erythropoietin are also described.

Background of the Invention

15 [0002] Erythropoietin (Epo) is a glycoprotein hormone necessary for the maturation of erythroid progenitor cells into erythrocytes. It is produced in the kidney and is essential in regulating levels of red blood cells in the circulation. Conditions marked by low levels of tissue oxygen signal increased production of Epo, which in turn stimulates erythropoiesis. A loss of kidney function as is seen in chronic renal failure (CRF), for example, typically results in decreased production of Epo and a concomitant reduction in red blood cells.

20 [0003] Human urinary Epo was purified by Miyake et al. (J. Biol. Chem. 252, 5558 (1977)) from patients with aplastic anemia. However, the amount of purified Epo protein obtained from this source was insufficient for therapeutic applications. The identification and cloning of the gene encoding human Epo and expression of recombinant protein was disclosed in U.S. Patent No. 4,703,008 to Lin, the disclosure of which is incorporated herein by reference. A method for purification of recombinant human erythropoietin from cell medium is disclosed in U.S. Patent No. 4,667,016 to Lai et al., which is incorporated herein by reference. The production of biologically active Epo from mammalian host cells has made available, for the first time, quantities of Epo suitable for therapeutic applications. In addition, knowledge of the gene sequence and the increased availability of purified protein has led to a better understanding of the mode of action of this protein.

30 [0004] Both human urinary derived Epo (Miyake et al. *supra*) and recombinant human Epo expressed in mammalian cells contain three N-linked and one O-linked oligosaccharide chains which together comprise about 40% of the total molecular weight of the glycoprotein. N-linked glycosylation occurs at asparagine residues located at positions 24, 38 and 83 while O-linked glycosylation occurs at a serine residue located at position 126 (Lai et al. J. Biol. Chem. 261, 3116 (1986); Broudy et al. Arch. Biochem. Biophys. 265, 329 (1988)). The oligosaccharide chains have been shown to be modified with terminal sialic acid residues with N-linked chains typically having up to four sialic acids per chain and O-linked chains having up to two sialic acids. An Epo polypeptide may therefore accommodate up to a total of 14 sialic acids.

35 [0005] Various studies have shown that alterations of Epo carbohydrate chains can affect biological activity. In one study, however, the removal of N-linked or O-linked oligosaccharide chains singly or together by mutagenesis of asparagine or serine residues that are glycosylation sites sharply reduces *in vitro* activity of the altered Epo that is produced in mammalian cells (Dube et al. J. Biol. Chem. 263, 17516 (1988)). However, DeLorme et al. (Biochemistry 31, 9871-9876 (1992)) reported that removal of N-linked glycosylation sites in Epo reduced *in vivo* but not *in vitro* biological activity.

40 [0006] The relationship between the sialic acid content of Epo and *in vivo* biological activity was disclosed by determining the *in vivo* activity of isolated Epo isoforms. It was found that a stepwise increase in sialic acid content per Epo molecule gave a corresponding stepwise increase in *in vivo* biological activity as measured by the ability of equimolar concentrations of isolated Epo isoforms to raise the hematocrit of normal mice (Egrie et al. Glycoconjugate J. 10, 263 (1993)). Those Epo isoforms having higher sialic acid content also exhibited a longer serum half-life but decreased affinity for the Epo receptor, suggesting that serum half-life is an important determinant of *in vivo* biological activity.

45 [0007] Introduction of new glycosylation sites in the Epo polypeptide can result in the production of molecules with additional carbohydrate chains. See PCT Publication Nos. WO91/05867 and WO 95/05465. Epo glycosylation analogs having at least one additional N-linked carbohydrate chain and/or having at least one additional O-linked carbohydrate chain are disclosed. A glycosylation analog having one additional N-linked chain was determined to have a longer circulating half-life compared to recombinant human Epo (rHuEpo) (isoforms 9-14) and to a purified isoform of rHuEpo having 14 sialic acids per molecule.

50 [0008] Administration of recombinant human erythropoietin (rHuEpo) is effective in raising red blood cell levels in anemic patients with end stage renal disease (Eschbach et al. New Eng. J. Med. 316, 73-38 (1987)). Subsequent studies have shown that treatment with rHuEpo can correct anemia associated with a variety of other conditions. (Fischl et al. New Eng. J. Med. 322, 1488-1493 (1990); Laupacis, Lancet 341, 1228-1232 (1993). Regulatory approvals have been given for the use of rHuEpo in the treatment of anemia associated with CRF, anemia related to therapy with AZT (zidovudine) in HIV-infected patients, anemia in patients with non-myeloid malignancies receiving chemotherapy, and anemia in patients undergoing surgery to reduce the need of allogenic blood transfusions. Current therapy for all approved

indications (except the surgery indication) involves a starting dose of between 50-150 Units/kg three times per week (TIW) administered either by an intravenous (IV) or subcutaneous (SC) injection to reach a suggested target hematocrit range. For the surgery indication, rHuEpo is administered every day 10 days prior to surgery, on the day of surgery, and four days thereafter (EPOGEN® Package Insert, 12/23/96). In general, the current recommended starting doses for rHuEpo raise hematocrit into the target range in about six to eight weeks. Once the target hematocrit range has been achieved, a maintenance dosing schedule is established which will vary depending upon the patient, but is typically three times per week for anemic patients with CRF. The administration of rHuEpo described above is an effective and well-tolerated regimen for the treatment of anemia.

[0009] It would be desirable to have a therapeutic with greater potency than rHuEpo. An advantage to such a molecule would be that it could be administered less frequently and/or at a lower dose. Current treatments for patients suffering from anemia call for administration of EPOGEN® three times per week and for surgery patients administration once per day. A less frequent dosing schedule would be more convenient to both physicians and patients, especially those patients who do not make regularly scheduled visits to doctor's offices or clinics, or those who self-inject their Epo. Another advantage of a more potent molecule is that less drug is being introduced into patients for a comparable increase in hematocrit.

[0010] It is therefore an object of the invention to identify more potent molecules for the treatment of anemia which will permit a less frequent dosing schedule. It is a further object of the invention to provide molecules which will increase and maintain hematocrit at levels which are at least comparable to that of Epo when administered at a lower dose. It is also an object of the invention that these molecules selected for less frequent dosing is at least as well tolerated as rHuEpo and potentially better tolerated in some patients.

#### Summary of the Invention

[0011] It has been found that a hyperglycosylated Epo analog designated N47 (Asn<sup>30</sup>Thr<sup>32</sup>Val<sup>87</sup>Asn<sup>88</sup>Thr<sup>90</sup> Epo) has a longer serum half-life than recombinant human erythropoietin (rHuEpo) and a greater *in vivo* activity when administered at the same dose and frequency as rHuEpo. Further, the analog has been shown to raise hematocrit in mice at once per week administration that is comparable to hematocrit rise for rHuEpo administered three times per week. The pharmacokinetics of Epo analog N47 administered to mice and to humans were similar.

[0012] The invention provides for a method of raising and maintaining hematocrit in a mammal comprising administering a therapeutically effective amount of an Epo hyperglycosylated analog in a pharmaceutical composition, wherein the analog is administered less frequently than an equivalent molar amount of rHuEpo to obtain a comparable target hematocrit. The dosing frequency of the present invention in order to reach a patient's optimal hematocrit range is less than three times per week. Dosing frequencies may be two times per week, one time per week, or less than one time per week, such as one time every other week, once per month or once every two months. The dosing frequency required to maintain a patient's target hematocrit is less than three times per week. Dosing frequencies may be two times per week, one time per week, or less than one time per week, such as one time every two weeks, once per month or once every two months.

[0013] Also provided for are pharmaceutical compositions comprising Epo hyperglycosylated analogs wherein the compositions are suitable for dosing frequency of less than three times per week. The compositions will include pharmaceutically acceptable adjuvants suitable for use with Epo hyperglycosylated analogs.

[0014] The invention may be employed with any condition resulting in a decrease in red blood cell levels, such as anemia associated with a decline or loss of kidney function, (chronic renal failure) myelosuppressive therapy, cancer, viral infection, chronic disease and excessive loss of blood during surgical procedures. In one embodiment, treatment with once per week dosing, or less frequently, is for anemia resulting from chronic renal failure.

#### Description of the Figures

##### [0015]

Figure 1 shows the amino acid sequence of human erythropoietin.

Figure 2 shows a Western blot analysis of rHuEpo and Epo hyperglycosylated analogs from CHO cell expression in serum free medium. Construction of analogs N53 and N61 are described in Example 1. The number of N-linked carbohydrate chains on each analog is indicated.

Figure 3 compares the activity of rHuEpo, Epo analogs N4, N18, and N50 (containing four N-linked carbohydrate chains), N47 (containing five N-linked carbohydrate chains), and N53 (containing six N-linked carbohydrate chains) in the exhypoxic polycythemic mouse bioassay. Experimental procedures are described in Example 3. Each point

represents the mean response of five animals. Analogs N4, N18 and N47 have been described previously in WO 95/05464.

Figure 4 compares the serum half-life of rHuEpo and Epo analog N47 administered to normal rats by intravenous injection (IV). Experimental procedures are described in Example 4. Results are the mean ( $\pm$ SD) for each group.

Figure 5 compares the serum half-life of rHuEpo and Epo analog N47 administered to Beagle dogs by intravenous injection (IV). Experimental procedures are described in Example 4. Results are the mean ( $\pm$ SD) for each group.

Figure 6 shows the increase in hematocrit in mice in response to varying doses of rHuEpo or Epo analog N47 administered by intraperitoneal injection (IP) three times per week (TIW) for six weeks. Experimental procedures are described in Example 5. Results shown are the group mean ( $\pm$ SD) of the change in hematocrit for each dose group.

Figure 7 compares the relative potency in mice of rHuEpo and Epo analog N47 injected by either the intraperitoneal (IP) or intravenous (IV) routes of administration at a frequency of once weekly (QW) or three times a week (TIW). Experimental procedures are described in Example 5. Each point represents the mean ( $\pm$ SD) of data from separate experiments as follows: N47, IP, TIW (n=5); N47, IV, TIW (n=1); N47, IP, QW (n=2); N47, IV, QW (n=3); rHuEpo, IP, TIW (n=5); rHuEpo, IV, QW (n=2). Each experiment used 7 - 13 mice per dose.

Figure 8 shows the increase in hematocrit in mice in response to varying doses of rHuEpo or Epo analog N47 administered by intravenous (IV) injection one time per week (QW) for approximately six weeks. Experimental procedures are described in Example 5. Results shown are the group mean ( $\pm$ SD) of the change in hematocrit for each dose group.

Figure 9 shows the increase in hematocrit in mice in response to varying doses of Epo analog N47 administered by intravenous (IV) injection one time per week (QW) or once every other week (EOW) for approximately six weeks. Experimental procedures are described in Example 5. Results shown are the group mean ( $\pm$ SD) of the change in hematocrit for each dose group.

#### Detailed Description of the Invention

**[0016]** The invention provides for a method of raising and maintaining hematocrit comprising administering a therapeutically effective amount of a hyperglycosylated analog of erythropoietin in a pharmaceutical composition. The analog is administered less frequently than an equivalent molar amount of rHuEpo to obtain a comparable target hematocrit. The composition may be administered by intravenous, subcutaneous or intraperitoneal routes.

**[0017]** Surprisingly, it has been found that analog N47, a hyperglycosylated Epo analog described in WO 95/05465, could achieve an increase in hematocrit administered once a week that was comparable to that observed for rHuEpo given three times per week. Analog N47 has the following amino acid changes: ala to asn at 30; his to thr at 32; pro to val at 87; trp to asn at 88; and pro to thr at 90 which resulted in the addition of two N-linked carbohydrate chains at asparagine residues 30 and 88. The analog was expressed in Chinese hamster ovary (CHO) cells (as described in Example 1) and purified as described in Example 2 to give isoforms of 17 to 22 sialic acids. Analog N47 showed a greater serum half-life in rats and beagle dogs than rHuEpo when injected intravenously (Figures 4 and 5). When injected intraperitoneally three times per week, N47 induced increases in hematocrit of normal mice comparable to rHuEpo at lower concentrations (Figure 6). The potency of N47 was demonstrated to be about 3 to 4-fold higher than rHuEpo when administered three times per week (Figures 6 and 7). When given once per week, at similar doses, rHuEpo showed little stimulation of hematocrit in normal mice while N47 gave a marked increase (Figure 8). The potency of N47 was about 14-fold higher than rHuEpo for once per week dosing (Figure 7). Significantly, the hematocrit response for analog N47 given once per week is comparable to that for rHuEpo given three times per week. Even when administered once every other week, N47 still produced significant increases in the hematocrit of normal mice (Figure 9). Taken together, the data indicated that Epo hyperglycosylated analogs, and analog N47 in particular, can be used advantageously to raise hematocrit using less frequent dosing than for current treatment with rHuEpo.

**[0018]** It has also been shown that the results described above obtained in mice may be extrapolated to humans. Pharmacokinetic parameters for administration of rHuEpo and analog N47 to 11 Continuous Ambulatory Peritoneal Dialysis (CAPD) patients demonstrate that analog N47 has a three-fold longer serum half-life than rHuEpo (Example 6 and Table 5). These results suggest that Epo hyperglycosylated analogs allow less frequent dosing than rHuEpo in humans.

**[0019]** As used herein, the term "hyperglycosylated Epo analog" refers to Epo comprising at least one additional

glycosylation site with an additional carbohydrate chain added to the site. Glycosylation sites may be for N-linked or O-linked carbohydrate chains. New N-linked glycosylation sites are introduced by alterations in the DNA sequence to encode the consensus site for N-linked carbohydrate addition (the amino acids Asn-X-Ser/Thr) in the polypeptide chain, while new O-linked sites are introduced by alterations in the DNA sequence to encode a serine or a threonine residue.

5 The analogs are constructed by mutagenesis techniques for introducing additions, deletions or substitutions of amino acid residues that increase or alter sites in the Epo polypeptide that are available for glycosylation. DNA encoding an Epo hyperglycosylated analog is transfected into a eucaryotic host cell and the expressed glycoprotein is analyzed for the presence of an additional carbohydrate chain.

10 **[0020]** Epo hyperglycosylated analogs have shown *in vitro* activity which was comparable to or even less than that determined for rHuEpo, suggesting that binding to the Epo receptor is not enhanced, and may in some cases be diminished, by addition of carbohydrate chains. However, hyperglycosylation can typically increase serum half-life and potentially lead to increased *in vivo* biological activity. One Epo analog having an additional N-linked carbohydrate chain at position 88 exhibited decreased affinity for receptor compared to rHuEpo (isoforms 9-14) or to a purified isoform of rHuEpo having 14 sialic acids per molecule, yet demonstrated a longer circulating half-life and enhanced *in vivo* activity

15 compared to either a mixture of Epo isoforms 9-14 or isolated Epo isoform 14.

**[0021]** The Epo hyperglycosylated analogs which may be administered according to the present invention will have at least one additional N-linked or O-linked carbohydrate chain. In one embodiment, the analogs will have two additional N-linked carbohydrate chains. In other embodiments, the analogs will have three, four or more additional N-linked carbohydrate chains. As examples, the analogs of the invention will have at least one additional N-linked chain at one

20 or more of amino acid residues 30, 51, 57, 69, 88, 89, 136 and 138 of the sequence of human Epo. In one embodiment, the analog has additional N-linked carbohydrate chains at residues 30 and 88 of human Epo. The numbering of amino acid residues of human Epo is as shown in Figure 1 and SEQ ID NO:1. Figure 1 shows a predicted mature Epo polypeptide of 166 amino acids whereas recombinant produced Epo has 165 amino acids after removal of the C-terminal arginine residue. It is understood that rHuEpo and hyperglycosylated Epo analogs may have either 165 or 166 amino acids.

25 **[0022]** The analogs of the invention will have at least four N-linked carbohydrate chains. Of the four chains, three may be at the naturally occurring sites at positions 24, 38, and 83. However, it is contemplated that some analogs of the invention may have alterations of one or more of the naturally-occurring glycosylation sites such that one or more of the sites are deleted and substituted with a new site. Such analogs are also provided by the invention. For example, any one of sites at positions 24, 38 and 83 may be deleted and substituted with a site at position 88. Optionally, the analogs

30 may have an O-linked site at position 126.

**[0023]** The invention also provides for new Epo hyperglycosylated analogs having at least one additional carbohydrate chain. It has been found that an additional N-linked carbohydrate chain is added at any of positions 52, 53, 55, 86 and 114 which have been modified to be a glycosylation site. Specific embodiments include analogs N49 through N61 as described in Table 1. The new analogs will have at least one new N-linked glycosylation site at any of positions 52, 53,

35 55, 86 and 114 and may further comprise additional N-linked or O-linked carbohydrate chains at other sites. The analogs may have one, two, three or four additional carbohydrate chains, or more than four additional chains. In one preferred embodiment, the analogs will have three additional N-linked carbohydrate chains (six N-linked chains total). In another preferred embodiment, the analogs will have four additional N-linked chains (seven N-linked chains total). The analogs having three or four, or more than four, additional N-linked carbohydrate chains may have, but are not limited to, an additional chain at any of positions 52, 53, 55, 86 and 114.

**[0024]** Surprisingly, it has been found that a hyperglycosylated analog with three additional N-linked chains at positions 30, 53 and 88 (six N-linked chains total) has a greater *in vivo* activity than analog N47 with two additional chains (five total). The results are shown in Figure 3. It is clear that the *in vivo* activity of the analogs is directly dependent on the number of N-linked carbohydrate chains. These results may be extrapolated to the therapeutic setting wherein the

45 analogs having more N-linked carbohydrate chains than N47 may be dosed even less frequently.

**[0025]** The analogs may be prepared by a variety of mutagenesis techniques available to one skilled in the art, such as site-directed mutagenesis, PCR mutagenesis and cassette mutagenesis (Zoller et al. *Meth. Enz.* **100**, 468-500 (1983); Higuchi, in *PCR Protocols* pp. 177-183 (Academic Press, 1990); Wells et al. *Gene* **34**, 315-323 (1985)). Example 1 describes the use of PCR mutagenesis techniques to construct new Epo hyperglycosylated analogs.

50 **[0026]** An Epo DNA sequence which has undergone mutagenesis is inserted into an expression vector using standard techniques with the vector being suitable for maintenance in a mammalian host cell. The vector will typically contain the following elements: promoter and other "upstream" regulatory elements, origin of replication, ribosome binding site, transcription termination site, polylinker site, and selectable marker that are compatible with use in a mammalian host cell. Vectors may also contain elements that allow propagation and maintenance in procaryotic host cells as well.

55 **[0027]** Suitable cells or cell lines include any from mammalian sources, including human sources. Examples include COS-7 (ATCC accession no. CRL 1651), human 293, baby hamster kidney (BHK, ATCC accession no. CCL 10), Chinese hamster ovary cells (including dihydrofolate reductase (DHFR)-deficient cells, Urlab et al. *Proc. Natl. Acad. Sci. USA* **77**, 4216-4220 (1980)) Other suitable mammalian cell lines include, but are not limited to, HeLa, mouse L-929 and 3T3.

In a preferred embodiment, DHFR-deficient CHO cells are used.

**[0028]** Vectors comprising sequences encoding Epo hyperglycosylation analogs are introduced into host cells by standard transformation or transfection techniques. Culturing, amplifying and screening transformed or transfected host cells are accomplished using publicly available methods (Gething et al. *Nature* 293, 620-625 (1981); Kaufman et al. *Mol Cell. Biol.* 5, 1750-1759 (1985); U.S. Patent No. 4,419,446). Host cells harboring DNA sequences encoding Epo hyperglycosylated analogs are cultured under conditions that permit expression of the analogs. The analogs are recovered from the cell media and purified using procedures essentially as described previously (WO94/09257) and those in Example 2. The purification procedures allow for the isolation of higher sialic acid containing Epo isoforms resulting from adding additional carbohydrate chains.

**[0029]** The Epo hyperglycosylated analogs may include, in addition to new glycosylation sites, additions, deletions or substitutions of amino acid residues which do not create new glycosylation sites and do not substantially alter the biological activity of the hyperglycosylated analog. Those individual sites or regions of Epo which may be altered without affecting biological activity may be determined by examination of the structure of the Epo-Epo receptor complex as described in Syed et. al. *Nature* 395, 511 (1998). Examination of the structure of the Epo-Epo receptor complex reveals those residues which interact with, or are in close proximity to, the receptor binding site of Epo and which should be avoided when making alterations in the Epo amino acid sequence. Alternatively, one may empirically determine those regions which would tolerate amino acid substitutions by alanine scanning mutagenesis (Cunningham et al. *Science* 244, 1081-1085 (1989)). In this method, selected amino acid residues are individually substituted with a neutral amino acid (e.g., alanine) in order to determine the effects on biological activity.

**[0030]** It is generally recognized that conservative amino acid changes are least likely to perturb the structure and/or function of a polypeptide. Accordingly, the invention encompasses one or more conservative amino acid changes within an Epo hyperglycosylated analog. Conservative amino acid changes generally involve substitution of one amino acid with another that is similar in structure and/or function (e.g., amino acids with side chains similar in size, charge and shape). The nature of these changes are well known to one skilled in the art and are summarized in Table 1 below. Such conservative substitutions are shown under the heading of "Preferred substitutions". Also contemplated are more substantial changes ("Exemplary substitutions") which may also be introduced. A skilled artisan will appreciate that initially the sites should be modified by substitution in a relatively conservative manner. If such substitutions result in a retention in biological activity, then more substantial changes (Exemplary Substitutions) may be introduced and/or other additions/deletions may be made and the resulting products screened.

TABLE 1: Amino Acid Substitutions

Original Residue	Preferred Substitutions	Exemplary Substitutions
Ala (A)	Val	Val ; Leu; Ile
Arg (R)	Lys	Lys; Gln ; Asn
Asn (N)	Gln	Gln ; His; Lys; Arg
Asp (D)	Glu	Glu
Cys (C)	Ser	Ser
Gln (Q)	Asn	Asn
Glu (E)	Asp	Asp
Gly (G)	Pro	Pro
His (H)	Arg	Asn; Gln; Lys; Arg
Ile (I)	Leu	Leu; Val; Met; Ala; Phe; norleucine
Leu (L)	Ile	norleucine; Ile; Val; Met; Ala; Phe
Lys (K)	Arg	Arg; Gln; Asn
Met (M)	Leu	Leu; Phe; Ile
Phe (F)	Leu	Leu; Val; Ile; Ala
Pro (P)	Gly	Gly
Ser (S)	Thr	Thr

(continued)

Original Residue	Preferred Substitutions	Exemplary Substitutions
Thr (T)	Ser	Ser
Trp (W)	Tyr	Tyr
Tyr (Y)	Phe	Trp; Phe; Thr; Ser
Val (V)	Leu	Ile; Leu; Met; Phe; Ala; norleucine

**[0031]** Also described are deletions or additions of amino acids in a hyperglycosylated Epo analog which do not substantially affect biological activity. Such additions and deletions may be at the N-terminal or C-terminal of the polypeptide, or may be internal to it. In general, relatively small deletions or additions are less likely to affect structure and/or function of Epo or a hyperglycosylated analog. In one embodiment, deletions or additions can be from 5-10 residues, alternatively from 2-5 amino acid residues, or from 1-2 residues.

**[0032]** The term "molar amount" refers to an amount of a hyperglycosylated analog or rHuEpo which is based upon the molecular weight of the corresponding erythropoietin polypeptide without glycosylation. Equivalent amounts of rHuEpo and analog refer to amounts which are equal when taking into account normal variations in procedure used to determine such amounts. It is necessary to determine equivalent amounts in this manner since the molecular weight of rHuEpo and analogs will vary depending upon the number of carbohydrate chains. For rHuEpo, the molecular weight of erythropoietin polypeptide is calculated based upon amino acid residues 1-165 as shown in Figure 1 and SEQ ID NO: 1. For hyperglycosylated analogs, the molecular weights are adjusted depending upon the amino acid changes in residues 1-165 of Figure 1 and SEQ ID NO: 1.

**[0033]** The dosing frequency for a hyperglycosylated analog will vary depending upon the condition being treated and the target hematocrit, but in general will be less than three times per week. The dosing frequency will be about two times per week, about one time per week. The dosing frequency may also be less than about one time per week, for example about one time every two weeks (about one time per 14 days), one time per month or one time every two months. It is understood that the dosing frequencies actually used may vary somewhat from the frequencies disclosed herein due to variations in responses by different individuals to the Epo analogs; the term "about" is intended to reflect such variations.

**[0034]** As used herein, the term "therapeutically effective amount" refers to an amount of a hyperglycosylated analog which gives an increase in hematocrit to a target hematocrit, or to a target hematocrit range that provides benefit to a patient or, alternatively, maintains a patient at a target hematocrit, or within a target hematocrit range. The amount will vary from one individual to another and will depend upon a number of factors, including the overall physical condition of the patient, severity and the underlying cause of anemia and ultimate target hematocrit for the individual patient. A target hematocrit is typically at least about 30%, or in a range of 30%-38%, preferably above 38% and more preferably 40%-45%. General guidelines relating to target hematocrit ranges for rHuEpo are also found in the EPOGEN® package insert dated 12/23/96 and are 30%-36%, or alternatively 32%-38% as stated therein. It is understood that such targets will vary from one individual to another such that physician discretion may be appropriate in determining an actual target hematocrit for any given patient. Nonetheless, determining a target hematocrit is well within the level of skill in the art.

**[0035]** A therapeutically effective amount of the present compositions may be readily ascertained by one skilled in the art. Example 6 sets forth a clinical protocol which has as one objective to determine a therapeutically effective amount of analog N47 in both once per week and three times per week dosing. A dose range for once per week administration is from about 0.075 to about 4.5 µg erythropoietin peptide per kg per dose. A dose range for three times per week administration is 0.025 to 1.5 µg erythropoietin peptide per kg per dose. This dose range may be employed with other Epo hyperglycosylated analogs, with any adjustments in the dosing range being routine to one skilled in the art.

**[0036]** A significant advantage to the present invention is the ability to correlate the extent of hyperglycosylation either with a dose amount or with a dosing interval that would allow one to "tailor" an Epo analog to a given dose or dosing schedule. Based upon the increasing *in vivo* activities of Epo analogs having one, two or three additional carbohydrate chains as shown in Figure 3, the treating physician can select an analog that is appropriate and convenient for the anemic condition being treated. For example, in patients who are acutely anemic and in need of a large effective dose, or in patients which require a longer-lasting treatment, administration of a hyperglycosylated analog with three or four or even more additional carbohydrate chains may be preferred. For other patients who experience less severe anemia or require treatment for a relatively short time, an analog with one or two additional carbohydrate chains may be preferred. The analogs of the present invention provide the physician with considerable flexibility in preventing and treating anemia that may result from a wide variety of underlying conditions.

**[0037]** The invention also provides for administration of a therapeutically effective amount of iron in order to maintain

increased erythropoiesis during therapy. The amount to be given may be readily determined by one skilled in the art based upon therapy with rHuEpo.

**[0038]** The present invention may be used to stimulate red blood cell production and prevent and treat anemia. Among the conditions treatable by the present invention include anemia associated with a decline or loss of kidney function (chronic renal failure), anemia associated with myelosuppressive therapy, such as chemotherapeutic or anti-viral drugs (such as AZT), anemia associated with the progression of non-myeloid cancers, anemia associated with viral infections (such as HIV), and anemia of chronic disease. Also treatable are conditions which may lead to anemia in an otherwise healthy individual, such as an anticipated loss of blood during surgery. In general, any condition treatable with rHuEpo may also be treated with the Epo hyperglycosylated analogs of the invention.

**[0039]** Also described are pharmaceutical compositions comprising a therapeutically effective amount of an Epo hyperglycosylated analog together with a pharmaceutically acceptable diluent, carrier, solubilizer, emulsifier, preservative and/or adjuvant. The composition will be suitable for a dosing schedule of less than three times per week. The composition may be in a liquid or lyophilized form and comprises a diluent (Tris, citrate, acetate or phosphate buffers) having various pH values and ionic strengths, solubilizer such as Tween or Polysorbate, carriers such as human serum albumin or gelatin, preservatives such as thimerosal, parabens, benzylalconium chloride or benzyl alcohol, antioxidants such as ascorbic acid or sodium metabisulfite, and other components such as lysine or glycine. Selection of a particular composition will depend upon a number of factors, including the condition being treated, the route of administration and the pharmacokinetic parameters desired. A more extensive survey of components suitable for pharmaceutical compositions is found in Remington's Pharmaceutical Sciences, 18th ed. A.R. Gennaro, ed. Mack, Easton, PA (1980). In a preferred embodiment, the Epo glycosylation analogs of the invention are formulated in liquid form in an isotonic sodium chloride/sodium citrate buffered solution containing human albumin, and optionally containing benzyl alcohol as a preservative. The compositions preferably contain analogs having one, two, three, four, or more additional carbohydrate chains.

**[0040]** Compositions of the invention are preferably administered by injection, either subcutaneous or intravenous. The route of administration eventually chosen will depend upon a number of factors and may be ascertained by one skilled in the art.

**[0041]** The following examples are offered to more fully illustrate the invention, but are not to be construed as limiting the scope thereof.

#### EXAMPLE 1

##### Construction of Hyperglycosylated Epo Analogs

##### Construction of cDNAs encoding Hyperglycosylated Epo Analogs

**[0042]** Epo analogs were made by in vitro mutagenesis using several different methods. Analogs N49 and N50 were constructed as described in WO 95/05465. Analogs were also constructed by variations of overlap PCR (polymerase chain reaction) methods. The basic procedure included two successive steps. In the first step, two reactions (PCR1 and PCR2) were performed on Epo or Epo analog template DNA using a total of four oligonucleotides: a 5' (forward) primer, a reverse mutagenic primer, a forward mutagenic primer (usually complementary to the reverse mutagenic primer) and a 3' (reverse) primer. The mutagenic primers contained the desired nucleotide changes as well as 6-14 exact match nucleotides on each side of these changes. PCR1 used the 5' (forward) primer and the reverse mutagenic primer. PCR2 used the 3' (reverse) primer and the forward mutagenic primer. The amplified DNA fragments were separated by agarose gel electrophoresis. Small pieces of agarose containing DNA fragments of the correct size were excised from the gel. The DNA fragments from PCR1 and PCR2 were combined together and a third PCR reaction was performed using only the 5' forward and 3' reverse primers. Thus, a full length DNA segment containing the desired mutations was amplified. In several cases, two or three mutations were combined by introducing a new substitution into DNA already containing a change, using the same PCR process. To construct these multiple glycosylation site analogs, single double or triple site analogs (produced as described above) were used as PCR template, and an additional glycosylation site was introduced by site directed mutagenesis with the appropriate primers.

**[0043]** The Epo analogs N51, N52 and N53 were constructed by the overlap PCR (polymerase chain reaction) method 1. One additional N-glycosylation site was introduced in each case. N56 added a glycosylation site (N114 T116) to native sequence HuEpo by using pDSR $\alpha$ 2 Epo as PCR template, N51 added an O-linked glycosylation site (Thr125) to N47 Epo by using pDSR $\alpha$ 2 Epo N47 template (Asn30, Thr32, Val187, Asn88, Thr90) and analog N59 added a glycosylation site (Asn53) to analog N47 using pDSR $\alpha$ 2 EpoN47 template.

**[0044]** Polymerase chain reactions for method 1 were performed using a protocol adapted from Cheng et. al., (Proc. Natl. Acad. Sci. USA 91, 5695 (1994)). The 3' (reverse) primer contained sequences that introduced a stop codon followed by a Xba I restriction site: ATCTAGAAGTTGCTCTCTGGACAGTTCCT (SEQ ID NO: 2). The 5' forward reaction primer: GAAGCTTGCGCCACCATGGGGTGCACGAATG (SEQ ID NO: 3) had an Hind III restriction site followed by



a Kozak sequence upstream of the Epo initiator codon (ATG). The typical PCR reaction mix contained: 4  $\mu$ l each of forward and reverse primers (5 pmol/ $\mu$ l), 1  $\mu$ l template (25 ng), 10  $\mu$ l of 5X LP buffer (100 mM Tricine pH 8.7/25% glycerol/425 mM KOAc), 10  $\mu$ l dNTP stock (1 mM each of dATP, dTTP, dCTP, dGTP), 0.8  $\mu$ l rTh polymerase (Perkin Elmer; 2.5 U/ $\mu$ l), and 2  $\mu$ l Vent polymerase (NEB; 0.01 U/ $\mu$ l after 1:100 fresh dilution in 1X LP buffer). H<sub>2</sub>O was added to bring the final volume to 50  $\mu$ l. All the components were added together in the order shown and the PCR was started when the temperature during the first cycle was above 60°C by adding 1  $\mu$ l of 50 mM MgOAc. Typical reaction conditions were: 2 cycles of 94°C, 10 sec/50°C, 1 min./ 68°C, 5 min. followed by 25 cycles of 94°C, 10 sec/55°C, 1 min./ 68°C, 5 min. The amplified fragments were separated by agarose gel electrophoresis and the correct sized DNA fragment was purified using a GeneClean™ kit and procedures supplied by the manufacturer (Bio 101, Inc.). The purified DNA was digested with Hind III and Xba I, then it was purified again using the GeneClean™ kit. The fragment was then ligated into Hind III and Xba I cut pDSR $\alpha$ 2 vector. Ligated DNA was precipitated with 2 volumes of ethanol in 0.3M NaOAc pH 5.2 in the presence of carrier tRNA and transformed into *E. coli*. Epo analogs were screened by restriction digest on mini DNA preps. Plasmids from positive clones were then prepared and the insert was sequenced to confirm the presence of the desired mutations and to ensure that no additional amino acid changes were introduced.

**[0045]** Analogs N54 to N61 were constructed using overlap PCR strategy method 2. The 3'(reverse) primer contained sequences that introduced a stop codon followed by a  $\times$ bal restriction site: GATCCTCTAGAGTTGCTCTCTGGACAG (SEQ ID NO: 4). The 5' forward reaction primer: CAACAAGCTTGCGCCGCCATGGGGG (SEQ ID NO: 5) had a HindIII restriction site followed by a Kozak sequence upstream of the Epo initiator codon (ATG). A high fidelity PCR strategy was performed using Perkin Elmer UITma DNA Polymerase and accompanying reagents; 10  $\mu$ l 10X PCR buffer, 3  $\mu$ l 1 mM dNTPS, 5 pmol of each primer, and water in a final volume of 100  $\mu$ l. 0.5 units of UITma polymerase was added after the PCR mixture reached 94°C. PCR reactions were then carried out for 5 cycles at 94°C for 30 seconds, 50°C for 30 seconds, and 72°C for 90 seconds. A subsequent 25 cycles were performed at 94°C for 30 seconds, and 72°C for 90 seconds. Product bands of the correct sizes were excised from an agarose gel following electrophoresis.

**[0046]** The resulting PCR products for each analog were cleaned using the Qiagen gel extraction kit. The purified DNA was digested in a 100  $\mu$ l restriction digest with HindIII and  $\times$ bal restriction enzymes (Boehringer Mannheim) at 37° C for 1 hour. The digests were again gel purified and the digested fragment was then ligated into HindIII and XbaI digested pDSR $\alpha$ 2 vector.

**[0047]** Ligated DNA was precipitated with 2 volumes of ethanol in 0.3M NaOAc pH 5.2 in the presence of carrier tRNA and transformed into *E. coli*. Epo analogs were initially screened by colony PCR to identify clones containing the correctly sized and type of DNA insert. With this procedure, cells containing plasmids were placed into PCR tubes in the presence of Epo forward and reverse primers. The mixture was then subjected to PCR using the reaction conditions described above. Plasmids from positive clones were then prepared and the Epo analog insert was sequenced to confirm the presence of the desired mutations and to ensure that no additional amino acid changes were introduced.

TABLE 1  
ERYTHROPOIETIN ANALOGS HAVING SITES FOR N-LINKED CARBOHYDRATE CHAINS

Analog	Amino Acid Substitution	Sequence Changes
N49	LYS, Met→Asn52, Thr54	AAG, ATG→AAT, ACC
N50	Arg, Glu→Asn53, Thr55	AGG, GAG → AAT, ACG
N51	Ala, His, Pro, Trp, Pro, Ala → N30,T32,V87,N88,T90 T125	GCT,CAC,CCG,TGG,CCC, GCC→AAT, ACG, GTG, AAT, ACC,ACC
N52	Ala, Lys→Asn114, Thr116 →	GCC, AAG→AAC, ACG
N53	Ala, His, Arg, Glu Pro,Trp,Pro → N30,T32,N53,T55,V87,N88,T90	GCT,CAC,AGG,GAG,CCG, TGG, CCC→AAT, ACG, AAT ACG,GTG,AAT,ACC
N54	Glu, Gly → Asn55, Thr57	GAG, GGG → AAT, ACT
N55	Gln, Pro, Trp → Asn86, Val87, Thr88	CAG, CCG, TGG → ACC, GTG, ACG
N56	Pro, Trp, Pro → Ala87, Asn88, Thr90	CCG, TGG, CCC → GCG, AAT, ACC

(continued)

## ERYTHROPOIETIN ANALOGS HAVING SITES FOR N-LINKED CARBOHYDRATE CHAINS

Analog	Amino Acid Substitution	Sequence Changes
N57	Pro, Trp, Pro → Val87, Asn88, Ser90	CCG, TGG, CCC → GTG,AAT,ACG
N58	Pro, Trp, Glu, Pro → Val87, Asn88, Gly89, Thr90	CCG, TGG, GAG, CCC → GTG, AAT, GGG, ACC
N59	Ala, His, Arg, Glu → Asn30, Thr32, Asn53, Asn55	GCT,CAC AGG GAG → AAT,ACG,AAT,ACG
N60	Ala, His, Ala, Lys → Asn30, Thr32, Asn114, Thr116	GCT, CAC, GCC, AAG → AAT,ACG,AAC, ACG
N61	A,H,R,E,P,W,P,A,K → 80, T32, N53, T55, V87, N88, T90, N114, T115	GCT,CAC,ACG,GAG,CCG, TGG, CCC, GCC, AAG → AAT, ACG, AAT, A CG,GTG, AAT,ACC,AAC,ACG

Analysis of carbohydrate addition

**[0048]** The constructs for the hyperglycosylated Epo analogs which were inserted into the expression vector pDSRα2 were transfected into COS cells. Supernatants from the transfected COS cells were analyzed by western blot to determine whether the expressed and secreted Epo analog contained additional carbohydrate. Samples were loaded directly into wells of SDS-PAGE gels then analyzed by immunoblot using the monoclonal antibody, 9G8A (Elliott et al (1996) Blood 87:p2714). Mobilities of analog samples were compared to that of samples containing rHuEpo. Figure 1 shows decreased mobility of analogs N53 and N61 compared to analogs N4 (four carbohydrate chains) and N47 (five carbohydrate chains). The mobility is consistent with the presence of six carbohydrate chains for analog N53 and seven carbohydrate chains for analog N61. Data for all hyperglycosylated analogs are shown in Table 2.

In vitro bioassays

**[0049]** Media conditioned by COS or CHO cells expressing rHuEpo or analogs were assayed for stimulation of 3H-thymidine uptake by UT7-Epo cells (Komatsu et al., Blood 82, 456). UT7-Epo cells are responsive to Epo and express human Epo receptors on their cell surface. UT7-Epo cells were grown in Growth medium (1X Iscove's Modified Dulbecco's Medium with L-glutamine, 25 mM HEPES buffer, and 3024 mg/L sodium bicarbonate, but without either alpha-thioglycerol or betamercaptoethanol (GIBCO)/ 10% v/v Fetal Bovine Serum/ 1% v/v L-glutamine-Penicillin-Streptomycin solution (Irvine Scientific)/ 1 Unit/mL rHuEpo) to approximately 3x10<sup>5</sup> cells/mL. Cells were collected by centrifugation (approx. 500xG) washed twice with phosphate buffered saline and resuspended at 5x10<sup>4</sup> cells/mL in Assay medium (1x RPMI Medium 1640 without L-glutamine (Gibco)/1% L-glutamine/4% fetal bovine serum). Test samples or Epo standard (rHuEpo), 100 uL diluted in assay medium at least 5-fold, were added to wells in a 96 well microtiter plate. 50μL of suspended cells were then added (5000 cells/well) and plates were incubated in a humidified incubator at 37°C and 5% CO<sub>2</sub>. After 72 hours, 50 uL methyl-<sup>3</sup>H-Thymidine (1 mCi/mL; 20 Ci/mMole) diluted 1:100 in assay medium was added. Cells were incubated for an additional 4 hours at 37°C and 5% CO<sub>2</sub>. Labeled cells were harvested onto glass fiber filtermats, washed with deionized water followed by 2-propanol, dried and counted. Activity was determined by comparing the response determined for each analog to that of the rHuEpo standard. The specific biological activity was then determined by dividing in vitro activity by the concentration of each analog as determined by immunoassay (Elliott et al (1996) Blood 87:p2714). The results are shown in Table 2.

TABLE 2

ANALOG	Number of N-linked Carbohydrate Chains	In Vitro Activity**
rHuEpo	3	+++
N49	4	+++
N50	4	+++
N51	5*	+++

(continued)

ANALOG	Number of N-linked Carbohydrate Chains	In Vitro Activity **
N52	3 - 4	+++
N53	6	++
N54	4	NT
N55	4	+++
N56	4	+++
N57	3 - 4	+++
N58	4	+++
N59	5	++
N60	4 - 5	+++
N61	6 - 7	NT

\* contains 1-2 O-linked chains  
 \*\* In vitro activity is relative to rHuEpo activity  
 +++ activity equivalent to rHuEpo  
 ++ activity is 25-75% of rHuEpo  
 NT Not Tested

## EXAMPLE 2

## Preparation of Recombinant Human Erythropoietin and Hyperglycosylated Erythropoietin Analogs

**[0050]** Recombinant human erythropoietin (rHuEpo) used for the experiments described herein was expressed by Chinese hamster ovary (CHO) cells transfected with a recombinant plasmid carrying the human erythropoietin gene. The recombinant product was recovered from the conditioned medium and purified essentially as described by Lai et al. *supra*. The resulting rHuEpo preparation has predominantly isoforms of 9 to 14 sialic acids as determined by isoelectric focusing.

**[0051]** Recombinant hyperglycosylated erythropoietin analogs were expressed in CHO cells transfected with a recombinant plasmid carrying the Epo analog gene as described WO91/05867 and WO94/09257 hereby incorporated by reference. The hyperglycosylated analogs were purified from culture supernatants as described below.

Concentration and diafiltration of conditioned media

**[0052]** Conditioned medium (serum free) from three successive harvests (5-8 days each) of the transfected CHO cell line was collected, filtered through a 0.45  $\mu$ m filter, concentrated about thirty fold, and diafiltered into 10 mM Tris, 20  $\mu$ M CuSO<sub>4</sub>, pH 7.0 using a tangential-flow ultrafiltration system (Millipore) with a 10,000 molecular weight cutoff membrane. The diafiltered media (DFM) was filtered (0.45  $\mu$ m) a second time and stored at -20°C until used for purification.

Purification

**[0053]** All procedures were carried out at 2 to 8°C.

Anion - Exchange Chromatography (10)

**[0054]** The clarified DFM was applied to a Q-Sepharose Fast Flow column (Pharmacia, 6 cm x 18 cm) equilibrated in 10 mM bis Tris propane (BTP), pH 7.0 and washed with two column volumes of 10 mM BTP to elute all non-binding species. The following gradients were run depending upon whether the hyperglycosylated analog had four, five or six N-linked carbohydrate chains. All buffers used at this stage contain 1 mM glycine, 20  $\mu$ M CuSO<sub>4</sub>, 6 M urea, 5  $\mu$ g/mL leupeptin and 1  $\mu$ g/mL pepstatin. For analogs with four N-linked carbohydrate chains, the gradient was 10 mM acetic acid, 0.1 mM NaCl to 500 mM acetic acid, 5 mM NaCl over 49 column volumes with a two column volume hold at high salt conditions. For analogs with five N-linked carbohydrate chains, the gradient was 0.7 M acetic acid, 7 mM NaCl to 1.0 M acetic acid, 12 mM NaCl over 30 column volumes with a two column volume hold at high salt conditions. For analogs with six N-linked carbohydrate chains, the gradient was 1.0 M acetic acid, 10 mM NaCl to 1.5 M acetic acid, 20 mM NaCl over 50 column volumes with a two column volume hold at high salt conditions. Following the gradient, the column was washed with two column volumes of 10 mM BTP, pH 7.0 and the high isoform fraction was eluted with 0.6

M NaCl, 100 mM BTP, pH 7.0.

#### Reversed Phase Chromatography (C4)

**[0055]** The high salt strip from the Q-Sepharose column (1Q) was applied to a Vydac C4 reversed phase column (30  $\mu$  particles, 4 cm x 18 cm) equilibrated in 20% ethanol, 10 mM BTP, pH 7.0 and eluted from the column with a thirty column volume gradient to 94% ethanol buffered in 10 mM BTP, pH 7.0. The pooled product peak, eluting in approximately 60% ethanol, was diluted with four volumes of 10 mM BTP, pH 7.0 to minimize possibility of aggregation in the presence of ethanol.

#### Anion - Exchange Chromatography (20)

**[0056]** The diluted eluate from the reversed phase column was applied to a second Q-Sepharose Fast Flow (Pharmacia, 3 cm x 9 cm) column equilibrated with 10 mM BTP, pH 7.0. The column was washed with equilibration buffer, and the hyperglycosylated Epo analog was eluted with 0.6 M sodium chloride, 20 mM sodium citrate, pH 6.0.

**[0057]** The purified protein was exchanged into 20 mM NaPO<sub>4</sub>, pH 6.0, 140 mM NaCl via centricon (10,000 molecular weight cutoff), followed by passage through a 0.2  $\mu$ m filter and storage at 2-8°C.

#### EXAMPLE 3

##### In Vivo Bioactivity of rHuEpo and rHuEpo Analogs Containing four, five and six N-linked Carbohydrate Chains

**[0058]** The *in vivo* activity of Epo analogs containing four, five and six N-linked carbohydrate chains was compared with that of rHuEpo in the exhypoxic polycythemic mouse bioassay. This assay quantifies the incorporation of <sup>59</sup>Fe into newly synthesized red blood cells as a measure of the increase in erythropoiesis in mice in response to an exogenously-administered test sample. The assay, performed as described below, is a modification of the method of Cotes and Bangham (Nature 191, 1065 (1961)).

**[0059]** In this assay, female BDF<sub>1</sub> mice are first preconditioned by exposure to low oxygen conditions in a hypobaric chamber (0.4 -0.5 atm.) for approximately 18 hours a day for 14 days. To compensate for the low oxygen conditions the mice respond by stimulating erythropoiesis to increase the number of red blood cells and thereby the relative oxygen-carrying capacity. After completion of the final hypobaric exposure, the mice are allowed to remain at ambient pressure for approximately 72 hours prior to the administration of test samples by intraperitoneal injection. At ambient pressure the mice are relatively polycythemic and respond by decreasing endogenous erythropoietin production and the rate of erythropoiesis. Five days after sample administration, 0.2 -0.3  $\mu$ Ci of <sup>59</sup>FeCl<sub>3</sub> in 0.2 mL is injected intravenously into the tail vein. Forty-eight hours later, the animals are sacrificed and the increase in erythropoiesis produced by the test samples is determined by measuring the amount of <sup>59</sup>Fe incorporated in a 0.5 mL sample of whole blood.

**[0060]** An example of the results obtained when rHuEpo and five different Epo analogs, containing four, five or six N-linked carbohydrate chains were tested in this assay is shown in Figure 3. Each sample was assayed at six or seven different dilutions within an appropriate concentration range. All samples were diluted in phosphate-buffered saline containing 0.5% bovine serum albumin, and 0.4 mL of each dilution was administered to five preconditioned mice. Forty-eight hours after the administration of <sup>59</sup>Fe, the amount incorporated in 0.5 mL of blood was measured by gamma counting. The results for each of the samples are plotted as the percent <sup>59</sup>Fe incorporated versus the log of the administered dose.

**[0061]** As shown in Figure 3, all five of the hyperglycosylated Epo analogs tested in this assay were more potent than rHuEpo. In addition, the potency of each analog was directly dependent on the number of N-linked carbohydrate chains, with those analogs having an increased number of carbohydrate chains having the greater activity. Thus, analog N53, which contains six N-linked carbohydrate chains, was the most potent analog. Analog N47, which contains five N-linked carbohydrate chains, was in turn, more potent than those analogs containing four N-linked chains. The potencies of the three analogs containing four N-linked carbohydrate chains (N4, N18 and N50) were approximately equal to each other and greater than that of rHuEpo.

**[0062]** In this experiment, the doses of rHuEpo and analogs containing four, five or six N-linked carbohydrate chains required to produce 40% <sup>59</sup>Fe incorporation were 10,700 ng, 640 ng, 140 ng and 38 ng, respectively. Based on the amount of material required to produce this level of erythropoiesis, Epo analogs containing four, five, or six N-linked carbohydrate chains are 17-fold, 77-fold and 280-fold more potent than rHuEpo.

## EXAMPLE 4

## IV Pharmacokinetics of rHuEpo and Epo analog N47 in Rats and Beagle Dogs

**[0063]** Two separate studies in rats and dogs were performed to compare the pharmacokinetic parameters of Epo N47 analog and rHuEpo.

**[0064]** In the rat studies, 1  $\mu\text{Ci}$  ( $\sim 0.1 \mu\text{g}$  of peptide/kg) of either  $^{125}\text{I}$ -Epo N47 analog, or  $^{125}\text{I}$ -recombinant human erythropoietin (Amersham) was injected intravenously into a surgically implanted carotid cannula in normal male Sprague-Dawley rats weighing between 314-363 g. At various time points after administration, 0.3 mL of blood was collected and serum was prepared by centrifugation. The level of  $^{125}\text{I}$ -rHuEpo or  $^{125}\text{I}$  Epo N47 analog in 0.1 mL of each serum sample was then determined following an overnight  $4^\circ\text{C}$  incubation with 90% ethanol. The ethanol-precipitated protein in each serum sample was collected by centrifugation and the radioactivity was counted in a gamma counter. The resulting serum concentration vs time pharmacokinetic curves are shown in Figure 4. Each point represents a group mean of five rats in the N47 analog group and six rats in the rHuEpo group. The pharmacokinetic parameters were determined for each rat using PCNONLIN 4.0 nonlinear regression analysis (Statistical Consultants, 1992) and the results for each group were averaged. The results are shown in Table 3.

TABLE 3  
Comparison of IV Pharmacokinetic Parameters of N47 and r-HuEpo in Rats

Sample Test	Half-life		$V_d$ (mL/kg)	Serum Clearance (mL/kg-hr)
	$\alpha$ (hours)	$\beta$ (hours)		
N47 (n = 5 rats)	$0.57 \pm 0.49$	$6.9 \pm 0.3$	$33 \pm 5$	$4.8 \pm 1.2$
r-HuEpo (n = 6 rats)	$0.18 \pm 0.03$	$2.5 \pm 0.2$	$36 \pm 6$	$17.7 \pm 3.4$

a The results are presented as the group average  $\pm$  SD for five rats in the N47 group and six rats in the r-HuEpo group.

**[0065]** In the dog studies, normal Beagle dogs weighing between 7.8-9.5 kg received an intravenous bolus injection of  $\sim 29 \mu\text{Ci}$  of either  $^{125}\text{I}$ -rHuEpo or  $^{125}\text{I}$ -N47 ( $\sim 0.1 \mu\text{g}$  peptide/kg) into the cephalic vein. At various time points through 24 hours post-administration, approximately 1 to 2 mL of blood was collected and serum prepared. The concentration of  $^{125}\text{I}$ -rHuEpo and  $^{125}\text{I}$ -N47 in 0.1 mL serum was determined and pharmacokinetic parameters were calculated as described above. The serum concentration vs time pharmacokinetic curves for the dog studies are shown in Figure 5. The time points are the group means of two animals in each group. The pharmacokinetic parameters are summarized in Table 4.

TABLE 4  
Comparison of IV Pharmacokinetic Parameters of N47 and r-HuEpo in Dogs

Sample Test	Half-life		$V_d$ (mL/kg)	Serum Clearance (mL/kg-hr)
	$\alpha$ (hours)	$\beta$ (hours)		
N47	0.34	25.0	55.9	2.4
r-HuEpo	0.40	7.2	60.8	8.4

a The results presented are the average parameters for the two dogs in each group.

**[0066]** In both the rat and dog studies, rHuEpo and Epo N47 analog exhibited a biphasic serum clearance. Clearance in rats was about 3.7-fold faster for rHuEpo than for Epo N47 analog and the  $\beta$ -half-life was about 2.8-fold longer for Epo N47 analog than for rHuEpo. The pharmacokinetic parameters in the dog studies were generally consistent with those observed in rat. In dogs, the clearance of rHuEpo was 3.5-fold faster than for Epo N47 analog and the  $\beta$ -half-life was 3.5-fold longer for Epo N47 analog compared with that for rHuEpo.

## EXAMPLE 5

Dose response of hematocrit after administration of rHuEpo and Epo analog N47.

Hematocrit Dose Response Studies at three times per week (TIW)

**[0067]** The *in vivo* biological effects of rHuEpo and Epo analog N47 in normal mice were compared after administering a range of doses by either intraperitoneal or intravenous injection three times per week for up to six weeks. Hematocrit determinations were performed twice weekly by retro-orbital bleed.

**[0068]** Normal CD1 mice weighing approximately 30g (10-13 mice per group) were injected intraperitoneally three times per week for a total of six weeks with either rHuEpo (over the dose range of 0.625-10 µg peptide/kg/dose), Epo N47 analog (over the dose range of 0.156-1.25 µg peptide/kg/dose) or vehicle control. The vehicle control and diluent for the various rHuEpo and Epo N47 analog dosing preparations was phosphate-buffered saline (PBS), containing 0.025% mouse serum albumin. The hematocrits of all mice were determined at baseline and twice weekly thereafter by retro-orbital bleeds. At the conclusion of the experiment, serum from all animals was collected and assayed for antibodies to the injected product by a solution radioimmunoprecipitation assay. Hematocrit data from animals judged to be negative for neutralizing antibodies were used for subsequent analysis.

**[0069]** As shown in Figure 6 both rHuEpo and Epo N47 analog produce a dose-dependent increase in hematocrit in the six week study, although N47 analog promotes a greater increase in hematocrit compared to rHuEpo at a given concentration. In this experiment the Epo N47 analog is about 3 to 4-fold more potent when dosed three times per week by intraperitoneal injection.

**[0070]** Dose response studies of rHuEpo and analog N47 were carried out by intravenous injection three times per week using procedures similar to those for intraperitoneal injection. The results obtained were similar to those for intraperitoneal administration and, in particular, the studies further confirmed that Epo N47 analog had a greater potency than rHuEpo when administered three times per week.

**[0071]** To better compare and quantify the biological activity of rHuEpo and Epo N47 analog in raising the hematocrit of normal mice, results of experiments were also analyzed by relative potency plots. For each experiment, the activity of rHuEpo or N47 analog at each dose was determined by summing the increase in hematocrit over the first 38 days of the study by trapezoidal summation to obtain the area under the curve (AUC). This was then plotted versus the log dose in µg peptide/kg/week. Potency difference between compounds administered by the same or different routes of administration or dosing frequencies can be determined by measuring the distance between the relevant log-dose response lines. Figure 7 summarizes the relative potency data for all experiments performed comparing the activity of rHuEpo and Epo N47 analog administered by two different routes (intraperitoneal and intravenous) and at two different dosing schedules.

**[0072]** As shown in Figure 7 when administered three times per week, Epo N47 analog has the same potency when injected by either the intravenous or intraperitoneal route and was 3.6-fold more potent than rHuEpo injected intraperitoneally three times weekly.

Hematocrit Dose Response Studies at one time per week (OW)

**[0073]** Comparisons of rHuEpo and Epo analog N47 at increasing hematocrit in normal mice were undertaken with once weekly dosing by either the intraperitoneal or intravenous routes of administration for six weeks.

**[0074]** Normal CD1 mice weighing approximately 30g (8-10 mice per group) were injected intravenously once weekly for a total of six weeks with varying concentrations of either rHuEpo or Epo N47 analog prepared in PBS containing 0.025% mouse serum albumin, or with vehicle control (PBS with 0.025% mouse serum albumin). The analog dose varied from 6.25-25 µg of peptide/kg/dose and the dose of rHuEpo varied from 25-200 µg/kg/dose. The hematocrits of all mice were determined at baseline and twice weekly thereafter by retro-orbital bleeds. At the conclusion of the experiment, serum from all animals was collected and assayed for antibodies to the injected product by a solution radioimmunoprecipitation. Data from animals judged to be negative for neutralizing antibodies were used for subsequent analysis.

**[0075]** As shown in Figure 8, whereas both rHuEpo and analog N47 can increase the hematocrit of normal mice when dosed once weekly, the dose of rHuEpo required to produce a response was significantly greater than that for analog N47. For instance, in this experiment 25 µg peptide/kg/week of N47 increased the hematocrit of mice by 41.2 points in six weeks, whereas the same dose of rHuEpo produced only a 12.5 point hematocrit rise.

**[0076]** Dose response studies of rHuEpo and analog N47 were performed by intraperitoneal injection once weekly using procedures similar to those described above. The results obtained were consistent with the results for intravenous administration and further confirmed the greater potency of analog N47 compared with rHuEpo when administered one time per week.

**[0077]** To quantify the activity difference between rHuEpo and N47 analog when each is dosed once weekly, relative

potency plots were generated from all relevant experiments as described above. As shown in Figure 7, when administered one time per week, analog N47 has the same potency when injected by the intravenous and intraperitoneal route. Analog N47 is approximately 14-fold more potent than rHuEpo when each is administered once weekly.

[0078] In addition, the log-dose response plots in Figure 6 also illustrate the following: (1) A given dose of analog N47 administered once weekly (QW) is approximately as effective as the same total weekly dose of rHuEpo given as three divided doses (TIW); (2) A given dose of rHuEpo administered once weekly (QW) is only approximately 2% as effective as the same total weekly dose of analog N47 given as three divided doses (TIW); (3) Analog N47 is approximately 4-fold more potent in mice when administered TIW compared to QW.

#### Hematocrit Dose Response Studies at Every Other Week (EOW)

[0079] Experiments were also undertaken to assess the ability of analog N47 to increase the hematocrit of mice when injected once every other week. Normal CD-1 mice (10 mice per group) were injected intravenously either once weekly or once every other week for a total of approximately six weeks with varying concentrations of Epo N47 analog prepared in PBS containing 0.025% mouse serum albumin. Analog N47 was administered at either 200, 100 or 25 µg/kg/dose every other week or at 12.5 µg/kg/dose once weekly. The hematocrits of all mice were determined at baseline and twice weekly thereafter by retroorbital bleeds.

[0080] As shown in Figure 9, analog N47 can increase the hematocrit of normal mice in a dose-dependent fashion even when administered bi-monthly. As expected when dosed less frequently, a greater amount of N47 analog is required to increase the hematocrit. A dose of 200 µg/kg of N47 analog administered every other week increased the hematocrit to approximately the same extent in six weeks as did 12.5 µg/kg when dosed weekly.

#### EXAMPLE 6

##### IV Pharmacokinetics of Epo N47 analog and rHuEpo in Continuous Ambulatory Peritoneal Dialysis (CAPD) patients

[0081] In view of the marked increase in serum half-life of Epo N47 analog compared to rHuEpo in rat and beagle dog, it was of interest to determine whether an increase could also be observed in humans.

[0082] A double-blind, randomized cross-over design study of eleven stable CAPD patients (7 males, 4 females, aged 27-75 years) was undertaken. One patient group received 100U/kg of rHuEpo (equivalent to 0.5 µg of peptide/kg) while a second group of patients received 0.5 µg peptide/kg of Epo N47 analog, both administered as a single bolus injection intravenously. Venous blood samples (3mL) were drawn via an indwelling cannula and were taken pre-dose and at 5, 10, 15, 30 minutes and 1, 2, 5, 8, 12, 16, 24, 30, 36, 48, 60, 72 and 96 hours after the intravenous bolus. After a 28 day washout period, the first patient group received a single intravenous dose of Epo analog N47 while the second group received a single intravenous dose of rHuEpo. Blood samples were taken as in the first cycle of treatment. Levels of rHuEpo and Epo N47 analog were determined in serum by ELISA after subtraction of baseline endogenous Epo levels. The pharmacokinetic parameters (mean ± SE) estimated after adjustment for cross-over design effects are shown in Table 5. The serum concentration AUC was calculated using the linear trapezoidal summation.  $t_{1/2z}$  is defined as:  $\log(2)/K_z$ , where  $K_z$  is calculated as the slope of the terminal portion of the  $\ln$  (serum concentration) time curve. The clearance (Cl) is defined as: dose/AUC. The volume of distribution ( $V_d$ ) is defined as:  $Cl/K$ .

TABLE 5

Dose Group	AUC (ng.h/mL)	$t_{1/2z}$ (h)	Cl (mL/h/kg)	Vd (mL/kg)
N47	291.0±7.6	25.3±2.2	1.6±0.3	52.4±2.0
rHuEpo	131.9±8.3	8.5±2.4	4.0±0.3	48.7±2.1

[0083] The mean serum half-life for Epo N47 analog (25.3 hr) was three times longer than for rHuEpo (8.5 hr) and the clearance was 2.5-fold faster for rHuEpo than for analog N47.

#### EXAMPLE 7

##### A Phase II Dose Finding and Dose Scheduling Study of Epo N47 Analog

[0084] Multicenter, randomized, sequential dose-escalation studies are initiated to investigate the optimum dose and dose schedule for analog N47 when administered by subcutaneous or intravenous injection in patients with CRF receiving dialysis.

**[0085]** The dosing schedule is as follows:

Once per week dosing: 0.075, 0.225, 0.45, 0.75, 1.5 and 4.5  $\mu\text{g}$  of peptide/kg/dose.

Three times per week dosing: 0.025, 0.075, 0.15, 0.25, 0.5 and 1.5  $\mu\text{g}$  of peptide/kg/dose.

**[0086]** The studies are carried out in two parts: the first part is a dose-escalation study designed to evaluate the dose of analog N47 given either one or three times per week which increases hemoglobin at an optimum rate over four weeks (greater than or equal to 1 g/dL but less than 3 g/dL). The second part of each study is designed to determine the doses required (when administered once or three times per week by either the intravenous or subcutaneous routes of administration) to maintain the hematocrit at the therapeutic target.

**[0087]** Preliminary results indicate that once weekly dosing with analog N47 can be used to both increase and maintain the hematocrit of anemic CRF patients. Initial results suggest that the preferred doses to initiate therapy on a three times a week dosing schedule are 0.15 and 0.25  $\mu\text{g}$ /peptide/kg/dose, and on a one time per week dosing schedule are 0.45 and 0.75  $\mu\text{g}$ /peptide/kg/dose for both routes of administration.

#### SEQUENCE LISTING

**[0088]**

<110> Egrie C., Joan  
Elliott G., Steven  
Browne K., Jeffrey

<120> Methods and Compositions for the Prevention and Treatment of Anemia

<130> A-460

<140> 09/178,292

<141> 1998-10-23

<160> 5

<170> PatentIn Ver. 2.0

<210> 1

<211> 193

<212> PRT

<213> Human

<220>

<221> SIGNAL

<222> (1)..(27)

<223> signal sequence is residue 1-27

<400> 1



EP 1 123 313 B1

Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu  
1 5 10 15  
5 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu  
20 25 30  
Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu  
35 40 45  
10 Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His Cys Ser Leu Asn Glu  
50 55 60  
Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe Tyr Ala Trp Lys Arg  
65 70 75 80  
15 Met Glu Val Gly Gln Gln Ala Val Glu Val Trp Gln Gly Leu Ala Leu  
85 90 95  
Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu Leu Val Asn Ser Ser  
100 105 110  
20 Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp Lys Ala Val Ser Gly  
115 120 125  
Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu Gly Ala Gln Lys Glu  
130 135 140  
25 Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala Pro Leu Arg Thr Ile  
145 150 155 160  
Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val Tyr Ser Asn Phe Leu  
165 170 175  
30 Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala Cys Arg Thr Gly Asp  
180 185 190  
Arg

35  
<210> 2  
<211> 29  
<212> DNA  
40 <213> Human  
<400> 2  
atctagaagt tgctctctgg acagttcct 29  
45 <210> 3  
<211> 32  
<212> DNA  
<213> Human  
50 <400> 3  
gaagcttgcg ccaccatggg ggtgcacgaa tg 32  
<210> 4  
<211> 27  
55 <212> DNA  
<213> Human  
<400> 4

gacacctag agttgctctc tggacag 27

<210> 5

<211> 25

<212> DNA

<213> Human

<400> 5

caacaagctt gcgccgcat ggggg 25

## Claims

1. Use of a hyperglycosylated analog of erythropoietin for the preparation of a pharmaceutical composition for raising and maintaining hematocrit in a mammal, wherein the analog is to be administered less frequently than an equivalent molar amount of recombinant human erythropoietin to obtain a comparable target hematocrit, and wherein the analog comprises at least one additional glycosylation site compared to human erythropoietin at any one of positions 30 and 88 of the sequence of human erythropoietin such that the analog comprises at least one additional N-linked carbohydrate chain.
2. The use according to claim 1, wherein the hyperglycosylated analog of erythropoietin is to be administered about two times per week.
3. The use according to claim 1, wherein the hyperglycosylated analog of erythropoietin is to be administered about one time per week.
4. The use according to claim 1, wherein the hyperglycosylated analog of erythropoietin is to be administered about one time every other week.
5. The use according to claim 1, wherein the hyperglycosylated analog of erythropoietin is to be administered about one time per month.
6. The use according to claim 3, wherein the hyperglycosylated analog of erythropoietin that is to be administered is at an amount of about 0.075 to 4.5 ug per erythropoietin peptide per kg per dose.
7. The use according to claim 1, wherein the target hematocrit is at least about 30%.
8. The use according to claim 1, wherein the pharmaceutical composition is for the treatment of a mammal suffering from anemia associated with a decline or a loss of kidney function.
9. The use according to claim 9, wherein the pharmaceutical composition is for the treatment of a mammal suffering from anemia associated with myelosuppressive therapy.
10. The use according to claim 9, wherein the myelosuppressive therapy comprises chemotherapeutic or anti-viral drugs.
11. The use according to claim 1, wherein the pharmaceutical composition is for the treatment of a mammal suffering from anemia associated with excessive blood loss.
12. The use according to claim 1, wherein the pharmaceutical composition further comprises a therapeutically effective amount of iron.
13. The use according to claim 1, wherein the analog is Asn30 Thr32 Val87 Asn88 Thr90 Epo.
14. The use according to claim 1, wherein the analog comprises two additional N-linked carbohydrate chains.
15. The use according to claim 1, wherein the analog comprises three additional N-linked carbohydrate chains.
16. The use according to claim 1, wherein the analog comprises four additional N-linked carbohydrate chains.

## Patentansprüche

1. Verwendung eines hyperglykosylierten Analogons von Erythropoietin zur Herstellung einer pharmazeutischen Zusammensetzung zur Erhöhung und Aufrechterhaltung von Hämatokrit in einem Säuger, wobei das Analogon weniger häufig verabreicht werden soll als eine äquivalente molare Menge eines rekombinanten humanen Erythropoietins, um einen vergleichbaren Zielwert von Hämatokrit zu erreichen, und wobei das Analogon an einer der Positionen 30 und 88 der Sequenz des humanen Erythropoietins wenigstens eine zusätzliche Glykosylierungsstelle im Vergleich zu humanem Erythropoietin umfasst, so dass das Analogon wenigstens eine zusätzliche N-verknüpfte Kohlenhydratkette umfasst.
2. Die Verwendung gemäß Anspruch 1, wobei das hyperglykosylierte Analogon von Erythropoietin etwa zweimal pro Woche verabreicht werden soll.
3. Die Verwendung gemäß Anspruch 1, wobei das hyperglykosylierte Analogon von Erythropoietin etwa einmal pro Woche verabreicht werden soll.
4. Die Verwendung gemäß Anspruch 1, wobei das hyperglykosylierte Analogon von Erythropoietin etwa einmal alle zwei Wochen verabreicht werden soll.
5. Die Verwendung gemäß Anspruch 1, wobei das hyperglykosylierte Analogon von Erythropoietin etwa einmal pro Monat verabreicht werden soll.
6. Die Verwendung gemäß Anspruch 3, wobei das hyperglykosylierte Analogon von Erythropoietin, das verabreicht werden soll, in einer Menge von etwa 0,075 bis 4,5 µg pro Erythropoietinpeptid pro kg pro Dosis vorliegt.
7. Die Verwendung gemäß Anspruch 1, wobei der Zielwert von Hämatokrit wenigstens etwa 30% beträgt.
8. Die Verwendung gemäß Anspruch 1, wobei die pharmazeutische Zusammensetzung zur Behandlung eines Säugers bestimmt ist, der an Anämie leidet, die mit einer Verschlechterung oder einem Verlust der Nierenfunktion einhergeht.
9. Die Verwendung gemäß Anspruch 1, wobei die pharmazeutische Zusammensetzung zur Behandlung eines Säugers bestimmt ist, der an Anämie leidet, die mit einer myelosuppressiven Therapie verbunden ist.
10. Die Verwendung gemäß Anspruch 1, wobei die myelosuppressive Therapie chemotherapeutische oder antivirale Arzneimittel umfasst.
11. Die Verwendung gemäß Anspruch 1, wobei die pharmazeutische Zusammensetzung zur Behandlung eines Säugers bestimmt ist, der an Anämie leidet, die mit einem starken Blutverlust einhergeht.
12. Die Verwendung gemäß Anspruch 1, wobei die pharmazeutische Zusammensetzung weiterhin eine therapeutisch wirksame Menge an Eisen umfasst.
13. Die Verwendung gemäß Anspruch 1, wobei das Analogon Asn30 Thr32 Val87 Asn88 Thr90 Epo ist.
14. Die Verwendung gemäß Anspruch 1, wobei das Analogon zwei zusätzliche N-verknüpfte Kohlenhydratketten umfasst.
15. Die Verwendung gemäß Anspruch 1, wobei das Analogon drei zusätzliche N-verknüpfte Kohlenhydratketten umfasst.
16. Die Verwendung gemäß Anspruch 1, wobei das Analogon vier zusätzliche N-verknüpfte Kohlenhydratketten umfasst.

## Revendications

1. Utilisation d'un analogue hyperglycosylé de l'érythropoïétine pour la préparation d'une composition pharmaceutique destinée à augmenter et à maintenir l'hématocrite chez un mammifère, où l'analogue doit être administré moins fréquemment qu'une quantité molaire équivalente d'érythropoïétine humaine recombinante pour obtenir un hématocrite cible comparable, et où l'analogue comprend au moins un site de glycosylation supplémentaire par compa-

## EP 1 123 313 B1

raison à l'érythropoïétine humaine au niveau de l'une quelconque des positions 30 et 88 de la séquence de l'érythropoïétine humaine de telle manière que l'analogue comprend au moins une chaîne glucidique N-liée supplémentaire.

- 5      2. Utilisation selon la revendication 1, où l'analogue hyperglycosylé de l'érythropoïétine doit être administré environ deux fois par semaine.
3. Utilisation selon la revendication 1, où l'analogue hyperglycosylé de l'érythropoïétine doit être administré environ une fois par semaine.
- 10     4. Utilisation selon la revendication 1, où l'analogue hyperglycosylé de l'érythropoïétine doit être administré environ une fois toutes les deux semaines.
5. Utilisation selon la revendication 1, où l'analogue hyperglycosylé de l'érythropoïétine doit être administré environ une fois par mois.
- 15     6. Utilisation selon la revendication 3, où l'analogue hyperglycosylé de l'érythropoïétine qui doit être administré est à une quantité d'environ 0,075 à 4,5 µg par peptide de l'érythropoïétine par kg par dose.
7. Utilisation selon la revendication 1, où l'hématocrite cible est au moins d'environ 30 %.
- 20     8. Utilisation selon la revendication 1, où la composition pharmaceutique est destinée au traitement d'un mammifère souffrant d'une anémie associée à un déclin ou à une perte de la fonction rénale.
9. Utilisation selon la revendication 1, où la composition pharmaceutique est destinée au traitement d'un mammifère souffrant d'une anémie associée à une thérapie myélosuppressive.
- 25     10. Utilisation selon la revendication 9, où la thérapie myélosuppressive comprend des médicaments chimiothérapeutiques ou antiviraux.
- 30     11. Utilisation selon la revendication 1, où la composition pharmaceutique est destinée au traitement d'un mammifère souffrant d'une anémie associée à une perte de sang excessive.
12. Utilisation selon la revendication 1, où la composition pharmaceutique comprend en outre une quantité thérapeutiquement efficace de fer.
- 35     13. Utilisation selon la revendication 1, où l'analogue est l'Asn30 Thr32 Val87 Asn88 Thr90 Epo.
14. Utilisation selon la revendication 1, où l'analogue comprend deux chaînes glucidiques N-liées supplémentaires.
- 40     15. Utilisation selon la revendication 1, où l'analogue comprend trois chaînes glucidiques N-liées supplémentaires.
16. Utilisation selon la revendication 1, où l'analogue comprend quatre chaînes glucidiques N-liées supplémentaires.

## FIG. 1

Met	Gly	Val	His	Glu	Cys	Pro	Ala	Trp	Leu	Trp	Leu	Leu	Leu	Ser	Leu	-25	-20	-15	
Leu	Ser	Leu	Pro	Leu	Gly	Leu	Pro	Val	Leu	Gly	Ala	Pro	Pro	Arg	Leu	-10	-5	1	5
Ile	Cys	Asp	Ser	Arg	Val	Leu	Glu	Arg	Tyr	Leu	Leu	Glu	Ala	Lys	Glu	10	15	20	
Ala	Glu	Asn	Ile	Thr	Thr	Gly	Cys	Ala	Glu	His	Cys	Ser	Leu	Asn	Glu	25	30	35	
Asn	Ile	Thr	Val	Pro	Asp	Thr	Lys	Val	Asn	Phe	Tyr	Ala	Trp	Lys	Arg	40	45	50	
Met	Glu	Val	Gly	Gln	Gln	Ala	Val	Glu	Val	Trp	Gln	Gly	Leu	Ala	Leu	55	60	65	
Leu	Ser	Glu	Ala	Val	Leu	Arg	Gly	Gln	Ala	Leu	Leu	Val	Asn	Ser	Ser	70	75	80	85
Gln	Pro	Trp	Glu	Pro	Leu	Gln	Leu	His	Val	Asp	Lys	Ala	Val	Ser	Gly	90	95	100	
Leu	Arg	Ser	Leu	Thr	Thr	Leu	Leu	Arg	Ala	Leu	Gly	Ala	Gln	Lys	Glu	105	110	115	
Ala	Ile	Ser	Pro	Pro	Asp	Ala	Ala	Ser	Ala	Ala	Pro	Leu	Arg	Thr	Ile	120	125	130	
Thr	Ala	Asp	Thr	Phe	Arg	Lys	Leu	Phe	Arg	Val	Tyr	Ser	Asn	Phe	Leu	135	140	145	
Arg	Gly	Lys	Leu	Lys	Leu	Tyr	Thr	Gly	Glu	Ala	Cys	Arg	Thr	Gly	Asp	150	155	160	165
Arg																			

FIG. 2





FIG. 3



◆ N53      ▲ N4  
× N18      ● rHuEPO  
■ N47  
\* N50

FIG.4






FIG. 5






FIG. 6




- N47 1.25 µg/kg/dose (IP-TIW)
- ▲ N47 0.625 µg/kg/dose (IP-TIW)
- N47 0.313 µg/kg/dose (IP-TIW)
- ▼ N47 0.156 µg/kg/dose (IP-TIW)
- vehicle (IV-TIW)
- rHuEPO 10 µg/kg/dose (IP-TIW)
- △ rHuEPO 2.5 µg/kg/dose (IP-TIW)
- rHuEPO 1.25 µg/kg/dose (IP-TIW)
- ▽ rHuEPO 0.625 µg/kg/dose (IP-TIW)

FIG. 7



- N47 (IV TIW)
- ▲ N47 (IP TIW)
- N47 (IV QW)
- × N47 (IP QW)
- △ rHuEPO (IP TIW)
- rHuEPO (IV QW)


FIG.8



● N47 25 µg/kg/dose (IV-QW)  
 ◆ N47 12.5 µg/kg/dose (IV-QW)  
 ▲ N47 6.25 µg/kg/dose (IV-QW)  
 ■ vehicle (IV-QW)

△ rHuEPO 200 µg/kg/dose (IV-QW)  
 ◇ rHuEPO 100 µg/kg/dose (IV-QW)  
 □ rHuEPO 50 µg/kg/dose (IV-QW)  
 ○ rHuEPO 25 µg/kg/dose (IV-QW)

FIG. 9



- N47 200 µg/kg/dose (IV-EOW)
- ▲ N47 100 µg/kg/dose (IV-EOW)
- N47 25 µg/kg/dose (IV-EOW)
- N47 12.5 µg/kg/dose (IV-QW)
- vehicle