



US00PP15800P2

(12) **United States Plant Patent**  
**Dümmen**

(10) **Patent No.:** **US PP15,800 P2**

(45) **Date of Patent:** **Jun. 14, 2005**

(54) **GERANIUM PLANT NAMED ‘DUEBERTINO’**

(56) **References Cited**

(50) Latin Name: *Pelargonium peltatum*  
Varietal Denomination: **Duebertino**

PUBLICATIONS

(75) Inventor: **Marga Dümmen**, Rheinberg (DE)

UPOV-ROM GTITM Computer Database 2004/04, GTI Jouve Retrieval Software, Citation for ‘Duebertino’.\*

(73) Assignee: **Dümmen Jungpflanzen GbR**,  
Rheinberg (DE)

\* cited by examiner

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

*Primary Examiner*—Anne Marie Grunberg  
*Assistant Examiner*—Annette H Para  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(21) Appl. No.: **10/859,509**

(57) **ABSTRACT**

(22) Filed: **Jun. 1, 2004**

A new and distinct cultivar of Ivy *Geranium* plant named ‘Duebertino’, characterized by its upright and outwardly spreading plant habit; freely basal branching habit; freely and early flowering habit; and red-colored double flowers.

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./332**

(58) **Field of Search** ..... **Plt./332**

**1 Drawing Sheet**

**1**

**2**

Botanical classification/cultivar denomination: *Pelargonium peltatum* cultivar Duebertino.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of Ivy *Geranium* plant, botanically known as *Pelargonium peltatum*, and hereinafter referred to by the name ‘Duebertino’.

The new Ivy *Geranium* is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. The objective of the breeding program was to develop new freely-flowering Ivy *Geraniums* with attractive flower and foliage colors.

The new Ivy *Geranium* originated from a cross-pollination made by the Inventor in May, 1999, of a proprietary selection of *Pelargonium peltatum* identified as code number E-15-14, not patented, as the female, or seed, parent with a proprietary selection of *Pelargonium peltatum* identified as code number S-15-14, not patented, as the male, or pollen, parent. The cultivar Duebertino was discovered and selected by the Inventor as a flowering plant within the progeny from this cross in a controlled environment in Rheinberg, Germany in April, 2002.

Asexual reproduction of the new cultivar by terminal vegetative cuttings at Rheinberg, Germany since June, 2002 has shown that the unique features of this new Ivy *Geranium* are stable and reproduced true to type in successive generations of asexual reproduction.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Duebertino’. These characteristics in combination distinguish ‘Duebertino’ as a new cultivar and distinguish it from other known Ivy *Geranium* cultivars:

1. Upright and outwardly spreading plant habit.
2. Freely basal branching habit.

3. Freely and early flowering habit.

4. Red-colored double flowers.

Compared to plants of the female and male parent selections, plants of the new Ivy *Geranium* differ in flower coloration.

The new Ivy *Geranium* can be compared to plants of the cultivar Narina 98, not patented. In side-by-side comparisons conducted in Rheinberg, Germany, plants of the new Ivy *Geranium* differed from plants of the cultivar Narina 98 in the following characteristics:

1. Plants of the new Ivy *Geranium* were taller than plants of the cultivar Narina 98.
2. Flowers of plants of the new Ivy *Geranium* were more intense in color than flowers of plants of the cultivar Narina 98.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Flower and foliage colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Ivy *Geranium*.

The photograph comprises a side perspective view of typical flowering plants of ‘Duebertino’ grown in a container.

**DETAILED BOTANICAL DESCRIPTION**

The cultivar Duebertino has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment, such as temperature and light intensity, without, however, any variance in genotype.

The aforementioned photograph, following observations and measurements describe plants grown in Rheinberg, Germany during the spring under commercial practice in a

glass-covered greenhouse with day and night temperatures about 18° C. and light levels about 4,500 foot-candles. Plants were grown with three plants per container. Plants were pinched about three weeks after planting. Plants were about eight weeks from unrooted cuttings when the photograph and the detailed botanical description were taken.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Pelargonium peltatum* cultivar Duebertino.

Parentage:

*Female parent*.—Proprietary selection of *Pelargonium peltatum* identified as code number E-15-14, not patented.

*Male parent*.—Proprietary selection of *Pelargonium peltatum* identified as code number S-15-14, not patented.

Propagation:

*Type cutting*.—Terminal vegetative cuttings.

*Time to initiate roots*.—Summer: About 5 days at 20° C.

Winter: About 7 days at 20° C.

*Time to develop roots*.—Summer: About three weeks at 20° C. Winter: About four weeks at 20° C.

*Root description*.—Fine; fibrous; white in color.

*Rooting habit*.—Freely branching.

Plant description:

*General appearance*.—Upright and outwardly spreading plant habit, rounded; densely foliated.

*Growth and branching habit*.—Moderately vigorous. Freely basal branching, about five lateral branches per plant.

*Plant height*.—About 28 cm.

*Plant width*.—About 12 cm.

*Lateral branches*.—Length: About 34.5 cm. Internode length: About 3.2 cm. Texture: Smooth. Color: 144A.

*Foliage description*.—Arrangement: Alternate, single. Length: About 4.3 cm. Width: About 7.5 cm. Shape: Reniform. Apex: Acute. Base: Peltate. Margin: Crenate. Venation pattern: Palmate. Texture, upper and lower surfaces: Smooth. Color: Developing and fully expanded foliage, upper surface: 137A; no distinct zonation pattern. Developing and fully expanded foliage, lower surface: 137C. Venation, upper surface: 144A. Venation, lower surface: 144B. Petiole: Length: About 3.6 cm. Diameter: About 2.1 mm. Color, upper and lower surfaces: 144A.

Flower description:

*Flower arrangement*.—Red-colored double flowers arranged in rounded hemispherical umbels arising from apical leaf axils. Umbels displayed above the foliage on upright peduncles. Flowers rounded in form. Umbels persistent, flowers not persistent. Flowers not fragrant.

*Quantity of flowers*.—Freely flowering; at full flower, plants have about six open umbels with about eight or nine flowers per umbel.

*Flowering season*.—Flowering continuous spring through summer.

*Flower longevity*.—Flowers last about five to seven days on the plant.

*Umbel size*.—Diameter: About 7.3 cm. Height: About 4.9 cm.

*Flower size*.—Diameter: About 3.7 cm. Depth (height): About 1.9 cm.

*Flower buds*.—Length: About 11.3 mm. Diameter: About 5.3 mm. Shape: Ovoid. Color: 144A.

*Petals*.—Quantity per flower: About five or six. Length: About 2.3 cm. Width: About 1.4 cm. Shape: Obovate. Apex: Rounded. Base: Attenuate. Margin: Sinuate. Texture, upper and lower surfaces: Smooth. Color: When opening and fully opened, upper surface: 44A to 44B; towards base, 60B; color becoming closer to 50A with development. When opening and fully opened, lower surface: 44C to 44D; towards base, 61A to 61B. Venation, upper surface: 44A to 44B. Venation, lower surface: 44C to 44D.

*Petaloids*.—Quantity per flower: About six. Length: About 2 cm. Width: About 1 cm. Shape: Obovate. Apex: Rounded. Base: Attenuate. Margin: Sinuate. Texture, upper and lower surfaces: Smooth. Color: When opening and fully opened, upper surface: 44A to 44B; towards base, 60B; color becoming closer to 50A with development. When opening and fully opened, lower surface: 44C to 44D; towards base, 61A to 61B. Venation, upper surface: 44A to 44B. Venation, lower surface: 44C to 44D.

*Sepals*.—Quantity per flower: About five to six, arranged in a single whorl. Length: About 1.3 cm. Width: About 4.1 mm. Shape: Elongated, tapering. Apex: Apiculate. Margin: Entire. Texture, upper and lower surfaces: Smooth. Color, upper and lower surfaces: 144A to 144B.

*Peduncle (umbel stem)*.—Length: About 10.3 cm. Diameter: About 2 cm. Angle: Erect. Strength: Moderately strong. Texture: Smooth. Color: 144A.

*Pedice (individual flower stem)*.—Length: About 2.3 cm. Diameter: About 2 mm. Angle: Erect. Strength: Moderately strong. Texture: Pubescent. Color: 144A overlain with 183C.

*Reproductive organs*.—Androecium: Anther quantity per flower: About eight or nine. Anther length: About 3 mm. Anther shape: Ovate. Anther color: 61A to 61B. Pollen amount: Moderate. Pollen color: 28A. Gynoecium: Pistil quantity per flower: One. Pistil length: About 1.1 cm. Stigma shape: Five-parted, star-shaped. Stigma color: 60B. Style length: About 4.6 mm. Style color: 2D. Ovary color: 144A.

*Seed/fruit*.—Development of seeds and fruit have not been observed.

Disease/pest resistance: Plants of the new Ivy *Geranium* have not been observed to be resistant to pathogens and pests common to Ivy *Geraniums*.

It is claimed:

1. A new and distinct cultivar of Ivy *Geranium* plant named 'Duebertino', as herein illustrated and described.

\* \* \* \* \*

