



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2285383 C 2006/09/26

(11)(21) **2 285 383**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**


(86) Date de dépôt PCT/PCT Filing Date: 1998/03/24
(87) Date publication PCT/PCT Publication Date: 1998/10/01
(45) Date de délivrance/Issue Date: 2006/09/26
(85) Entrée phase nationale/National Entry: 1999/09/24
(86) N° demande PCT/PCT Application No.: EP 1998/001714
(87) N° publication PCT/PCT Publication No.: 1998/042754
(30) Priorité/Priority: 1997/03/24 (ITMI97A000678)

(51) Cl.Int./Int.Cl. *C08B 37/00* (2006.01),
C08B 37/10 (2006.01)
(72) Inventeurs/Inventors:
CASU, BENITO, IT;
NAGGI, ANNAMARIA, IT;
TORRI, GIANGIACOMO, IT
(73) Propriétaire/Owner:
INALCO S.P.A., IT
(74) Agent: RIDOUT & MAYBEE LLP

(54) Titre : GLYCOSAMINOGLYCANS PRESENTANT UNE EFFICACITE ANTI-THROMBOSE ELEVEE
(54) Title: GLYCOSAMINOGLYCANS HAVING HIGH ANTITHROMBOTIC ACTIVITY

(57) **Abrégé/Abstract:**

Glycosaminoglycans having high antithrombotic activity in vitro, obtained by various kinds of glycosaminoglycans supersulfated by the preparation of the salt of an organic base of the starting supersulfated glycosaminoglycan, by partial solvolithic desulfation of said salt and N-resulfation of said partially desulfated product.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C08B 37/00, 37/10, A61K 31/725	A3	(11) International Publication Number: WO 98/42754 (43) International Publication Date: 1 October 1998 (01.10.98)
(21) International Application Number: PCT/EP98/01714 (22) International Filing Date: 24 March 1998 (24.03.98) (30) Priority Data: MI97A000678 24 March 1997 (24.03.97) IT (71) Applicant (for all designated States except US): INALCO S.P.A. [IT/IT]; Via C. Goldoni, 11, I-20129 Milan (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): CASU, Benito [IT/IT]; Via G. Colombo, 81/A, I-20133 Milan (IT). NAGGI, Annamaria [IT/IT]; Via Monte Nevoso, 32, I-20025 Legnano (IT). TORRI, Giangiacomo [IT/IT]; Via G. Colombo, 81/A, I-20133 Milan (IT). (74) Agent: PASSINI, Angelo; Notarbartolo & Gervasi S.p.A., Corso di Porta Vittoria, 9, I-20122 Milan (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 23 December 1998 (23.12.98)
(54) Title: GLYCOSAMINOGLYCANS HAVING HIGH ANTITHROMBOTIC ACTIVITY (57) Abstract Glycosaminoglycans having high antithrombotic activity in vitro, obtained by various kinds of glycosaminoglycans supersulfated by the preparation of the salt of an organic base of the starting supersulfated glycosaminoglycan, by partial solvolithic desulfation of said salt and N-resulfation of said partially desulfated product.		

GLYCOSAMINOGLYCANS HAVING HIGH ANTITHROMBOTIC ACTIVITY


Field of the invention

Object of the present invention are glycosaminoglycans having high antithrombotic activity in vitro, obtained by various kinds of glycosaminoglycans supersulfated by the preparation of the salt of an organic base of the starting supersulfated glycosaminoglycan, by partial solvolytic desulfation of said salt and N-resulfation of said partially desulfated product.

Prior art

Heparin is a polysaccharide extracted by animal organs, employed for more than fifty years as an anticoagulant and antithrombotic agent. Together with heparan sulfate, it belongs to the family of the glycosaminoglycans, consisting of alternate sequences of an uronic acid (iduronic or glucuronic) and glucosamine, variously sulfated depending on the tissue and the animal species from which they have been obtained and, to a certain extent, on the isolation processes too.

The structure of heparin may be represented in statistical terms by formula



where the U units represent the iduronic (IdoA) or glucuronic (GlcA) acid and the A units represent the N-sulfated (GlcNAc) glucosamine; R₂, R₃ and R₆ represent sulfate groups or hydrogen and R' represents SO₃ or Ac.

The sequences mostly represented in the heparins used in clinic are those of the trisulfated disaccharide (IdoA₂SO₃-GlcNSO₃SO₃). On the contrary, a minor

pentasaccharidic sequence contained only in about one third of the chains forming the common heparins and characterized by a glucosaminic unit sulfated in position 3 (GlcNSO₃SO₃) and forming the active site of heparin and heparan sulfate for the antithrombin III is essential for the expression of significant anticoagulant and antithrombotic activities. The 3-O-sulfated glucosamine unit is considered the marker of said active site and the antithrombotic activity, generally expressed by the inhibition capability of the X activated (Xa) and thrombin (IIa) coagulation factors, is correlable with the content per cent of said unit in the heparins.

The anticoagulant and antithrombotic properties of heparin and heparan sulfate are also modulated by the length of the polysaccharidic chains composing them. For example the heparins having low molecular weight (LMWH) have lower anticoagulant activity but antithrombotic activity similar to the traditional heparins one, and they are inclined to substitute the latter ones in several therapeutic applications, especially in order to reduce the haemorrhagic risks and other side effects of heparin, such as the thrombocytopenia. In addition, the LMWH are characterized by a bioavailability better than the traditional heparins one when they are administered by subcutaneous route, as it is common in the prevention of the venous thrombosis. (B. Casu Heparin structure, Hemostasis 20/1, 66-73 (1990)); (D.A. Lane, J. Björk, U. Lindall (Ed. s). Heparin and Related Polysaccharides, Plenum Press, New York, 1992).

Since, as above remembered, the pentasaccharidic sequences of heparin mostly responsible of the antithrombotic activity are contained only in about one third of the natural heparin chains, it is of practical interest to strengthen said activity concentrating the chains containing said sequences, or generating new active sites either in the chains which already contain one, or in those ones lacking active sites. The first goal, attainable by affinity chromatography on antithrombin III or (less effectively) by treatment of heparin with cationic resins, is still considered expensive, also because it does not use a substantial (about two thirds) part of the common heparins.

The goal to generate further active sites for antithrombin is not on the other hand attainable by sulfation of heparin with classical methods, which result in structures

wherein the active site for antithrombin is masked by an excess of sulfate groups. Even if fortuitously they show an anticoagulant activity greater than the starting heparin one (for the most part by action mechanisms different from the ones mediated by antithrombin), said products are generally less active than heparin as
5 antithrombotic agents, and they may generate undesirable side effects due to the aspecific interactions with other plasmatic proteins. [B. Casu, Structure and biological activity of heparin, Advances Carbohydr. Chem. Biochem. 43, 51-134 (1985)].

On the other hand, attempts to "reconstruct" the active sequences of heparin by resulfation of partially or totally desulfated heparins result in products with a reduced,
10 rather than strengthened, antithrombotic activity. This is due, in particular, to the tendency of the iduronic units to sulfate in position 3 rather than in position 2, and/or to insufficient sulfation in position 3 of the glucosaminic units. [R. N. Rej, K. G. Ludwig-Baxter, A. S. Perlin, Carbohydr. Res. 210, 299-310 (1991)].

Summary of the invention

15 Now we have found a process which makes possible to obtain glycosaminoglycans having high antithrombotic activity in vitro, containing at least 20% of iduronic units sulfated in position 2 and not sulfated in position 3 and at least 30% of sulfaminoglucosaminic units sulfated in position 3 and in position 6, and having a sulfates/carboxyls molar ratio ranging from 1.5 to 3.5.

20 Said process employs as starting materials various kinds of supersulfated glycosaminoglycans and it is characterized by the following steps:

- a) preparation of the salt of an organic base of the starting supersulfated glycosaminoglycan;
- b) partial solvolytic desulfation of the salt of the organic base of the step a);
- 25 c) N-resulfation of the partially desulfated product of the step b);
- d) possible 6-O-resulfation of the product of the step c).

The obtained product may be used as an active substance in the preparation of pharmaceutical compositions suitable to the antithrombosis treatment.

Detailed description of the invention

30 The characteristics and the advantages of the glycosaminoglycans having high

antithrombotic activity according to the present invention and the related preparation process will be mostly illustrated during the following detailed description.

In the process according to the present invention as starting materials may be used several kinds of supersulfated glycosaminoglycans and in particular may be employed: low molecular weight supersulfated heparins (ssLMWH), from 1,500 to 8,000 daltons, high molecular weight supersulfated heparins, from 8,000 to 20,000 daltons, low molecular weight supersulfated heparan sulfates, from 1,500 to 8,000 daltons, high molecular weight supersulfated heparan sulfates, from 8,000 to 25,000 daltons, "biotechnological" supersulfated heparan sulfates and heparins, epimerized and not epimerized, obtained from the N-sulfated K5 polysaccharide.

As it is known, the supersulfated glycosaminoglycans are glycosaminoglycans in which all the hydroxylic hydrogens (or the most part of them) have been substituted with SO_3^- groups, and they are prepared according to procedures described by various authors (M. L. Wolfrom et al, J. Am. Chem. Soc. 75, 1519 (1953); Nagasawa et al., Carbohydr. Res. 158, 183-190 (1986); EP 214,879 (1986); US Patent 4,727,063; Naggi et al., Biochem. Pharmacol. 36, 1895-1900 (1987); Ogamo et al., Carbohydr. Res. 193, 165-172 (1989)).

The process is carried out according to the following steps:

- a) Preparation of the salt of an organic base of the starting supersulfated glycosaminoglycan, said organic base being selected among pyridine, tetramethyl ammonium and tetrabutyl ammonium salts.

The starting supersulfated glycosaminoglycan in form of a sodium salt is dissolved in distilled water and passed through a cationic exchange column.

- To the obtained solution the organic base is added at room temperature and in an amount such that pH 9 is reached obtaining the salt which is then lyophilized.

- b) Partial solvolytic desulfation of the salt of the step a). The organic base salt obtained in the step a) is treated with a solution of an aprotic polar solvent containing methanol or H_2O in a percentage ranging from 5 to 10%, preferably DMSO containing 10% of methanol, with a ratio by weight between this solution and said organic base salt from 50:1 to 100:1.

The obtained solution is heated under stirring at a temperature ranging from 50 to

AMENDED SHEET

90 °C for a time ranging from 5 to 480 minutes obtaining the partial desulfation of said salt.

The reaction mixture is then added with water and 0.25 N NaOH to a neutral pH and the product is precipitated by addition of EtOH saturated with sodium acetate. The
5 product is then dialyzed through cut-off 2000 membranes and desalted by gel filtration.

c) N-resulfation of the product obtained from the step b).

The product obtained from the step b) is dissolved in an amount of water from 10/0.1 to 10/1 ml/g and the solution is added with NaHCO₃ to pH 9 and then trimethylamine
10 sulfur trioxide in an amount by weight equal to said product one. It is heated to 50-60 °C for two hours and then an equal amount of trimethylamine sulfur trioxide is added again and the heating is continued at the same temperature for 24 hours.

The product is precipitated by addition of EtOH saturated with sodium acetate, dialyzed through cut-off 2000 membranes in NaCl solutions first and then in distilled
15 water and finally it is desalted by gel filtration.

d) Possible 6-O-resulfation of the product obtained from the step c). The products which, basing on the NMR analysis result to be desulfated for more than 30% in position 6 of the glucosaminic unit, may be 6-O-resulfated according to known processes (i. e. according to K. Nagasawa, H. Vchiyama and N. Wajama, Carbohydr.
20 Res. 158 (1986)183-190).

The obtained product exhibits the following characteristics:

- at the least 20% of iduronic units sulfated in position 2 and not sulfated in position 3;
- at least 30% of sulfaminoglucosaminic units sulfated in position 3 and in position 6, as in the site active for the antithrombin of the heparins;
- 25 - sulfates/carboxyls molar ratio ranging from 1.5 to 3.5;
- anti-Xa activity in vitro ranging from 100 to 150 U/mg.

The content of iduronic units sulfated in position 2 and not sulfated in position 3, and of sulfaminoglucosaminic units sulfated in position 3 has been determined by NMR analysis based on the integration of signals specifically attributable to
30 determined units and determined types of sulfation [E. A. Yates, F. Santini, M.

Guerrini, A. Naggi, C. Torri, B. Casu: ^1H and ^{13}C NMR spectral assignments of the major sequences of twelve systematically modified heparin derivatives, Carbohydr. Res. 294, 15-27 (1996)].

5 The sulfates/carboxyls molar ratio has been determined by a conductometric method [B. Casu and U. Gennaro: A simple conductometric method for determining the sulfate and carboxylate groups of heparin and chondroitin sulfates, Carbohydr. Res. 39, 168-176 (1975)].

10 The anti-Xa activity in vitro has been determined by a chromogenic method (A. N. Teien et al., Thrombosis Res. 8, 413-416, 1976). The present invention refers particularly to glycosaminoglycans belonging to the group consisting of low molecular weight heparins (from 1,500 to 8,000 daltons), high molecular weight heparins (from 8,000 to 18,000 daltons), low molecular weight heparan sulfates (from 1,500 to 8,000 daltons), high molecular weight heparan sulfates (from 8,000 to 25,000 daltons) and heparan sulfates and heparins from the K5 polysaccharide. Thanks to their
15 characteristics, the glycosaminoglycans according to the present invention may be used in mixture with pharmacologically acceptable excipients or diluents, for the preparation of pharmaceutical compositions suitable to the antithrombosis treatment.

The present invention also relates thereby to the therapeutic method for the treatment and the prevention of thrombosis, consisting in the administration of a dose from 1 to
20 1,500 mg/day of said glycosaminoglycans.

DETAILED DESCRIPTION OF THE DRAWINGS

FIG. 1A: ^{13}C -NMR spectrum of supersaturated low molecular weight heparin (ssLMWH) starting material as described in Example 1a).

25 FIG. 1B: ^{13}C -NMR spectrum of totally N-desulfated and partially 6-O-desulfated ssLMWH as described in Example 1b).

FIG. 1C: ^{13}C -NMR spectrum of desulfated ssLMWH as described in Example 1b) and with a desulfation time of 30 minutes.

FIG. 1D: ^{13}C -NMR spectrum of desulfated ssLMWH as described in Example 1b) and with a desulfation time of 45 minutes.

30 With illustrative aim of the invention the following examples are reported:

EXAMPLE 1 - Preparation of low molecular weight heparin having high antithrombotic activity.

a) Preparation of the pyridine salt of a supersulfated low molecular weight heparin (ssLMWH) (G 1074/4)

360 mg of the sodium salt of ssLMWH having a molecular weight equal to 5,000 daltons, obtained from a commercial heparin from swine intestinal mucosa according to the procedure described in the US patent 4,727,063 and in the publication: Naggi et al., Biochem. Pharmacol. 36, 1895-1900 (1987), having anti Xa activity = 87 U/mg, the
5 ^{13}C -NMR spectrum of Fig. 1A, have been dissolved in distilled water (30 ml) and passed through a cationic exchange column (AMBERLITE[®] 120 in the H^+ form). The obtained solution has been taken to pH 9 with pyridine. The so obtained pyridine salt of the supersulfated low molecular weight heparin has been lyophilized.

b) Desulfation of the pyridine salt of the ssLMWH.

10 The above obtained pyridine salt of the supersulfated low molecular weight heparin has been dissolved in a solution of 10% DMSO/MeOH (25 ml). The solution has been heated under stirring in an oil bath at 80 °C for 15 minutes. At the end of the reaction 15 ml of water have been added and the solution has been neutralized with 0.25 N NaOH. The reaction product has been precipitated by the addition of 60 ml of EtOH
15 saturated with sodium acetate, dialyzed through cut-off 2000 membranes for 48 hours in distilled water and desalted by gel filtration on Sephadex G25. The obtained product exhibits the following characteristics: the ^{13}C -NMR spectrum (Figure 1B) shows typical signals from totally N-desulfated ($\text{C}2\text{NH}_2$:56 ppm) and partially 6-O-desulfated ($\text{C}6\text{OH}$:62 ppm) heparin, and the is signal at 65 ppm typical of iduronic units not
20 sulfated in position 3 and sulfated in position 2.

c) N-resuffation of the partially desulfated product of the step b) (G 2291)

100 mg of the product obtained in the step b) have been dissolved in 10 ml of water and taken to pH 9 with NaHCO_3 . Then the same amount by weight of trimethylamine sulfur trioxide (TMA.SO_3) has been added and the solution has been heated under
25 stirring in an oil bath at 55 °C. After 2 hours a further addition equal by weight of TMA.SO_3 has been carried out and it is left to react for 24 hours. At the end of the reaction the mixture has been taken to pH 9.

The sample has been precipitated by addition of 4 volumes of EtOH saturated with sodium acetate, dialyzed through cut-off 2000 membranes for 24 hours in 1 N NaCl,
30 for 24 hours in 0.5 N NaCl and for 24 hours in distilled water, and desalted by gel filtration on Sephadex G25.

The obtained product exhibits the following characteristics:

Sulfates/carboxyls molar ratio: 3.22

Anti-Xa activity: 152 U/mg.

35 The ^1H -NMR spectrum confirms the complete N-sulfation.

EXAMPLE 2 (G 2309)

The Example 1 has been repeated, with a reaction time of desulfation equal to 30 min. The product obtained from the desulfation shows the ^{13}C -NMR spectrum reported in Fig. 1C, with the characteristics described in the point b) of the Example 1.

After the N-resulfation, the obtained product exhibits the following characteristics:
Anti-Xa activity: 110 U/mg.

The ^1H -NMR spectrum confirms the complete N-sulfation.

EXAMPLE 3 (G 2301)

a) The Example 1 has been repeated, with a reaction time of desulfation equal to 45 min. The product obtained from the desulfation shows the ^{13}C -NMR spectrum reported in Fig. 1D, with the characteristics described in the point b) of the Example 1.

After the N-resulfation, the ^1H -NMR spectrum confirms the complete N-sulfation.

b) The N-resulfated product has been 6-O-resulfated according to the procedure cited in the detailed description.

EXAMPLE 4

The Example 1 has been repeated changing times and temperature of the point b), and precisely operating for 1.5 hours at 65 °C. The products obtained from the desulfations exhibit ^{13}C -NMR spectra analogous to those ones of the Figures 1B, 1C, and 1D.

The signal at 65 ppm (typical of the iduronic units desulfated in position 3 and sulfated in position 2) is present in significant amounts, even if lower than those obtained under the conditions of the Example 1 at the point b).

EXAMPLE 5

a) The Example 1 has been repeated with a desulfation temperature equal to 90 °C and a time of 45 min. The product obtained from the desulfation exhibits a ^{13}C -NMR spectrum with characteristics similar to those described in the point b) of the Example 1. However the signal characteristic of the glucosamine sulfated in position 6 is sensibly lower, and the signal characteristic of the glucosamine not sulfated in said position is sensibly greater, than that one of the spectra of the

products obtained at higher temperatures and shorter times.

b) The product has been 6-O-resulfated according to the procedure cited in the detailed description.

EXAMPLE 6

5 The Example 1 has been repeated, starting from high molecular weight heparin (12,000 daltons) supersulfated treating its pyridine salt in DMF at 55 °C with SO₃/pyridine adduct (molar ratio 3:1 with respect to the free hydroxyls) for 5 hours. The obtained product (sulfates/carboxyls molar ratio > 3; ¹³C-NMR signal at 75 ppm characteristic of the iduronic units sulfated in position 2 not sulfated in
10 position 3) has been N-resulfated as described in the Example 1c.

As illustrated in Table 1 for the low molecular weight heparins, the antithrombotic activity of the obtained products (expressed by the inhibition in vitro of the Xa factor) is noteworthy higher either than the starting supersulfated product one or than the low molecular weight heparins one presently available on the market,
15 used in therapy.

TABLE 1 - Anti-Xa ACTIVITY*

	1st set	2nd set
LMW-H	-	100
ssLMWH (G 1074/4)	100	-
20 15' desulfation from G1074/4 (G 2281)	175	162
30' desulfation from G1074/4 (G 2309)	125	125

*) U/mg. Normalized values in the first set with respect to the supersulfated low molecular weight heparin (ssLMWH), and in the second set with respect to a mixture 1:1 by weight of the commercial products "Fraxieparin" and "Lovenox"
25 (LMWH).

EXAMPLES 7- 17

Description of the used glycosaminoglycans.

- Commercial heparin having high molecular weight HMWH (Mw = 12,000 Da, anti-Xa activity 192 U/mg) obtained from swine intestinal mucosa (Examples 7, 8,
30 9, 10)

- Commercial heparan sulfate (HS). The ratio between iduronic acid and glucuronic acid varies as a function of the used heparan:

30% iduronic acid, 70% glucuronic acid (Example 11)

40% iduronic acid, 60% glucuronic acid (Example 12)

- K5 Polysaccharide (K5-PS) (Examples 13, 14)

- Epimerized K5 Polysaccharide (eK5-PS). The degree of epimerization varies as
5 a function of the used eK5-PS:

30% iduronic acid, 70% glucuronic acid (Example 15)

60% iduronic acid, 30% glucuronic acid (Examples 16, 17)

a) Preparation of the supersulfated substrate

1.5 g of the sodium salt of the used substrate have been dissolved in distilled
10 water (30 ml) and passed through a cationic exchange column (AMBERLITE IR
120 in the H⁺ form). The obtained solution has been taken to pH 9 with a solution
of 10% tetrabutyl ammonium hydroxide in ethanol. The so obtained tetrabutyl
ammonium salt has been lyophilized. The tetrabutyl ammonium salt of the
15 substrate has been dissolved in DMF (60 ml) and then pyridine sulfur trioxide
(Py.SO₃) has been added in the amounts of 3 eq. for free hydroxyl and the
solution has been heated under stirring in an oil bath at 55 °C overnight. At the
end of the reaction 5 ml of water have been added and the reaction product has
been precipitated by addition of 200 ml of EtOH saturated with sodium acetate,
20 dialyzed through cut-off 6000-8000 dialysis membranes for 48 hours in distilled
water and desalted by gel filtration on Sephadex G25. The obtained products
exhibit the following characteristics.

The ¹³C-NMR spectra show the signals typical of a completely N-desulfated
(A2NH₂) C2:56 ppm glucosamine and totally 6-O-sulfated (A6S) C6 69 ppm, and
of uronic units sulfated in position 2 and 3 on the same residue (U2,3S) C1 Å
25 101.5 ppm). The sulfation degree obtained on the uronic units changes as a
function of the used glycosaminoglycan. The data obtained for the obtained
supersulfated products are described in Table 2.

b) Preparation of the pyridine salt of the supersulfated substrate (ssHMWH).

1 g of the sodium salt obtained by the process described in the step a) has been
30 dissolved in distilled water (30 ml) and passed through a cationic exchange
column (AMBERLITE IR 120 in the H⁺ form). The obtained solution has been
taken to pH 9 with pyridine. The pyridine salt of the so obtained supersulfated

substrate has been lyophilized.

c) Desulfation of the pyridine salt of the supersulfated substrate

100 mg of the pyridine salt obtained as described above have been dissolved in a solution of DMSO/MeOH 10% (10 ml). The solution has been heated under
5 stirring (in an oil bath) using the temperatures and times described in Table 3. At the end of the reaction 5 ml of water have been added and the reaction product has been precipitated by the addition of 100 ml of EtOH saturated with sodium acetate, dialyzed through cut-off 6000-8000 dialysis membranes for 48 hours in distilled water and desalted by gel filtration on Sephadex G25.

10 The obtained products exhibit the characteristics described in Table 3.

d) N-resulfation of the partially desulfated products obtained from the step c)

100 mg of the product obtained in the step c) have been dissolved in 10 ml of water and taken to pH 9 with NaHCO_3 . Then the same amount by weight of trimethylamine sulfur trioxide ($\text{TMA}:\text{SO}_3$) has been added and the solution has
15 been heated under stirring in an oil bath at 55 °C. After 2 hours a further addition equal by weight of $\text{TMA}:\text{SO}_3$ has been carried out and it has been left to react for 24 hours. At the end of the reaction the 5 ml of water have been added and the reaction product has been precipitated by the addition of 50 ml of EtOH saturated with sodium acetate, dialyzed through cut-off 6000-8000 dialysis membranes for
20 48 hours in distilled water and desalted by gel filtration on Sephadex G25.

The ^{13}C -NMR spectra show the complete N-resulfation of the obtained products. Of the obtained products the sulfation degree expressed as sulfates/carboxyls (DS) ratio, the molecular weight (M_w) and the biological activities (aPTT, anti IIa and anti Xa) have been determined. The obtained data are described in Table 4.

TABLE 2 Description of the supersulfated glycosaminoglycans obtained as described in point a). In the first column of the Table it is pointed out in which Examples such products are used

Example	Substrate	GAG	% A6S	% UOH+U3S	% U2S+U2,3S
7, 8, 9, 10	ss-HMWH	2522	100	0	100
11	ss-LMWH	1079/4	100	5	95
12	ss-HS	2460	100	50	50
13	ss-HS	2648	n. a.	n. a.	n. a.
14- 15	ss-K5-PS	2444	100	0	100
16	ss-eK5- PS	2611	100	0	100
17	ss-eK5- PS	2721	n. a.	n. a.	n. a.

TABLE 3 Description of the used reaction conditions (substrate, temperatures and times) and characterization of the obtained products

Ex.	Substrate	GAG	T °C	t h	% A6S	% UOH+ U3S	% I2S	% G3S
7	HMWHep	2541A	65	4	39	35	54	n. a.
8	HMWHep	2632A	65	8	30	44	54	n. a.
9	HMWHep	2632B	65	16	5	50	54	n. a.
10	HMWHep	2541B	65	24	5	59	42	n. a.
11	LMWHep	2474	65	4	25	54	46	n. a.
12	HS	2556	65	4	50	60	20	25
13	HS	2734	80	30'	n. a.	n. a.	n. a.	n. a.
14	K5-PS	2451A	80	45'	40	48	-	52
15	K5-PS	2451B	80	75'	20	50	-	50
16	eK5-PS	2622	65	4	30	n. a.	n. a.	20
17	eK5-PS	2737	80	30'	n. a.	n. a.	n. a.	n. a.

TABLE 4 Description of the products obtained after the N-resulfation and their characterization

Ex.	GAG	DS	Mw	aPTT	anti IIa	anti Xa
7	2557	1.8	12000	n. a.	n. a.	110
8	2632BN S	n. a.	10300	n. a.	n. a.	99
9	2632AN S	n. a.	7600	n. a.	n. a.	7
10	2565	1	11000	n. a.	n. a.	3
11	2374NS	1.5	7400	10	273	130
12	2556NS	n. a.	n. a.	70	1230	65
13	2734NS	n. a.	n. a.	n. a.	n. a.	n. a.
14	2459A	2.4	n. a.	97	470	10
15	2459B	2.3	n. a.	32	114	5
16	2822NS	n. a.	n. a.	48	500	13
17	2737NS	n. a.	n. a.	n. a.	n. a.	n. a.

CLAIMS

1. Glycosaminoglycans having high antithrombotic activity in vitro, containing at least 20% of iduronic units sulfated in position 2 and not sulfated in position 3 and at least 30% of sulfaminoglucosaminic units sulfated in position 3 and in position 6, and having a sulfates/carboxyls molar ratio ranging from 1.5 to 3.5.
5
2. Glycosaminoglycans as claimed in claim 1, consisting of low molecular weight heparins from 1,500 to 8,000 daltons.
10
3. Glycosaminoglycans as claimed in claim 1, consisting of high molecular weight heparins from 8,000 to 20,000 daltons.
4. Glycosaminoglycans as claimed in claim 1, consisting of low molecular weight heparan sulfate from 1,500 to 8,000 daltons.
15
5. Glycosaminoglycans as claimed in claim 1, consisting of high molecular weight heparan sulfate from 8,000 to 25,000 daltons.
- 20 6. Glycosaminoglycans as claimed in claim 1, consisting of heparan sulfates and heparins.
7. Process for the preparation of the glycosaminoglycans as claimed in claim 1 starting from various kinds of supersulfated glycosaminoglycans, characterized by the following steps:
25
 - a) preparation of a salt of the starting supersulfated glycosaminoglycan by treatment with an organic base;
 - b) partial solvolytic desulfation of the salt of the organic base of the step a) by treatment with an aprotic polar solvent containing methanol or water at a temperature ranging from
30 50° to 80°C for a time ranging from 5 to 480 minutes;
 - c) N-resulfation of the partially desulfated product of the step b);
 - d) optional 6-O-resulfation of the product of the step c).
8. Process as claimed in claim 7, characterized in that said starting supersulfated
35 glycosaminoglycans are selected from the group consisting of low molecular weight

supersulfated' heparins, from 1,500 to 8,000 daltons, high molecular weight
supersulfated heparins, from 8,000 to 20,000 daltons, low molecular weight
supersulfated heparan sulfates, from 1,500 to 8,000 daltons, high molecular weight
supersulfated heparan sulfates, from 8,000 to 25,000 daltons, biotechnological
5 supersulfated heparan sulfates and heparins, epimerized or not epimerized.

9. Process as claimed in claim 7, characterized in that said preparation of the step a)
is obtained by treating with an organic base to pH 9 a solution of sodium salt of the
starting glycosaminoglycan.

10

10. Process as claimed in claim 7, characterized in that said N-resulfation of the step
c) is carried out by treating an aqueous solution of the product obtained in the step b)
with trimethylamine sulfur trioxide in an amount by weight equal to said product one,
at pH 9, heating to 50-60 °C and adding subsequently again an equal amount of
15 trimethylamine sulfur trioxide.

11. Use of a glycosaminglycan as defined in claim 1 for the preparation of an
antithrombotic medicament.

1/1

FIGURE 1 A




FIGURE 1 B




FIGURE 1 C




FIGURE 1 D

