



(86) Date de dépôt PCT/PCT Filing Date: 2006/05/04
(87) Date publication PCT/PCT Publication Date: 2006/11/16
(85) Entrée phase nationale/National Entry: 2007/11/01
(86) N° demande PCT/PCT Application No.: US 2006/017482
(87) N° publication PCT/PCT Publication No.: 2006/121895
(30) Priorité/Priority: 2005/05/05 (US11/123,873)

(51) Cl.Int./Int.Cl. *A61K 39/00* (2006.01),
A61K 38/00 (2006.01)
(71) Demandeur/Applicant:
LOS ANGELES BIOMEDICAL RESEARCH INSTITUTE
AT HARBOR-UCLA MEDICAL CENTER, US
(72) Inventeurs/Inventors:
EDWARDS, JOHN E., JR., US;
FILLER, SCOTT G., US;
SHEPPARD, DONALD C., CA;
IBRAHIM, ASHRAF, US;
FU, YUE, US;
SPELLBERG, BRAD J., US;
YEAMAN, MICHAEL, US
(74) Agent: SMART & BIGGAR

(54) Titre : COMPOSITIONS PHARMACEUTIQUES ET METHODES DE VACCINATION CONTRE LA CANDIDOSE
DISSEMINEE ET D'AUTRES AGENTS INFECTIEUX
(54) Title: PHARMACEUTICAL COMPOSITIONS AND METHODS TO VACCINATE AGAINST DISSEMINATED
CANDIDIASIS AND OTHER INFECTIOUS AGENTS

(57) **Abrégé/Abstract:**

The invention provides a vaccine including an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, with an adjuvant in a pharmaceutically acceptable medium. The invention also provides a method of treating or preventing hematogenously disseminated or mucocutaneous candidiasis. The method includes administering an immunogenic amount of a vaccine an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium. A method of treating or preventing disseminated candidiasis also is provided that includes administering an effective amount of an isolated Als protein family member having cell adhesion activity, or an functional fragment thereof, to inhibit the binding or invasion of *Candida* to a host cell or tissue. The Als protein family member can be derived from a *Candida* strain selected from the group consisting of *Candida albicans*, *Candida krusei*, *Candida tropicalis*, *Candida glabrata* and *Candida parapsilosis* and the Als protein family member includes Als1p, Als3p, Als5p, Als-p, Als7p or Als9p. Also provided is a method of treating or preventing *Staphylococcus aureus* infections. The method includes administering an immunogenic amount of a vaccine an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 November 2006 (16.11.2006)

PCT

(10) International Publication Number
WO 2006/121895 A3

(51) International Patent Classification:
A61K 39/00 (2006.01) A61K 38/00 (2006.01)

(74) Agent: GAY, David A.; McDERMOTT WILL & EMERY
LLP, 4370 La Jolla Village Drive, Suite 700, San Diego, CA
92122-1252 (US).

(21) International Application Number:
PCT/US2006/017482

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 4 May 2006 (04.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
11/123,873 5 May 2005 (05.05.2005) US

(71) Applicant (*for all designated States except US*): LOS ANGELES BIOMEDICAL RESEARCH INSTITUTE AT HARBOR-UCLA MEDICAL CENTER [US/US]; 1124 West Carson Street, Torrance, CA 90502-2064 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): EDWARDS, Jr., John, E. [US/US]; 3708 Via La Selva, Palos Verdes, CA 90274-1118 (US). FILLER, Scott, G. [US/US]; 28890 King Arthur Court, Rancho Palos Verdes, CA 90275-7209 (US). SHEPPARD, Donald, C. [CA/CA]; 5531 Chemin de La Cote St. Antoine, Montreal, QC, H4A 1R3 (CA). IBRAHIM, Ashraf [US/US]; 8192 Andora Drive, La Mirada, CA 90638-6108 (US). FU, Yue [US/US]; 21501 S. Vermont Ave., Apt. 101A, Torrance, CA 90502-1914 (US). SPELLBERG, Brad, J. [US/US]; 30140 Avenida Celestial, Rancho Palos Verdes, CA 90275-5493 (US). YEAMAN, Michael [US/US]; 504 Esplanade Avenue, Suite 508, Redondo Beach, CA 90277-4066 (US).

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report

(88) Date of publication of the international search report:
7 June 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHARMACEUTICAL COMPOSITIONS AND METHODS TO VACCINATE AGAINST DISSEMINATED CANDIDIASIS AND OTHER INFECTIOUS AGENTS

(57) Abstract: The invention provides a vaccine including an isolated AIs protein family member having cell adhesion activity, or an immunogenic fragment thereof, with an adjuvant in a pharmaceutically acceptable medium. The invention also provides a method of treating or preventing hematogenously disseminated or mucocutaneous candidiasis. The method includes administering an immunogenic amount of a vaccine an isolated AIs protein family member having cell adhesion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium. A method of treating or preventing disseminated candidiasis also is provided that includes administering an effective amount of an isolated AIs protein family member having cell adhesion activity, or an functional fragment thereof, to inhibit the binding or invasion of Candida to a host cell or tissue. The AIs protein family member can be derived from a Candida strain selected from the group consisting of Candida albicans, Candida krusei, Candida tropicalis, Candida glabrata and Candida parapsilosis and the AIs protein family member includes AIs Ip, Als3p, Als5p, Als1p, Als7p or Als9p. Also provided is a method of treating or preventing Staphylococcus aureus infections. The method includes administering an immunogenic amount of a vaccine an isolated AIs protein family member having cell adhesion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium.

WO 2006/121895 A3

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 71

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 71

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

**PHARMACEUTICAL COMPOSITIONS AND METHODS TO
VACCINATE AGAINST DISSEMINATED CANDIDIASIS AND
OTHER INFECTIOUS AGENTS**

5

BACKGROUND OF THE INVENTION

This invention relates to *Candida albicans* surface adhesin proteins, to antibodies resulting from an immune response to vaccination with *C. albicans* surface adhesion proteins and to methods for the prevention and/or treatment of candidiasis and other bacterial infections with *C. albicans* surface adhesion proteins.

10

There has been a dramatic increase in the incidence of nosocomial infections caused by *Candida* species in recent years. The incidence of hematogenously disseminated candidal infections increased 11-fold from 1980 to 1989. This increasing incidence has continued into the 1990s. Infections by *Candida* species are now the fourth most common cause of nosocomial septicemia, are equal to that of *Escherichia coli*, and

15

surpass the incidence caused by *Klebsiella* species. Furthermore *Candida* species are the most common cause of deep-seated fungal infections in patients who have extensive burns. Up to 11% of individuals undergoing bone marrow transplantation and 13% of those having an orthotopic liver transplant will develop an invasive candidal infection.

20

Candida albicans, the major pathogen in this genus, can switch between two morphologies: the blastospore (budding yeast) and filamentous (hyphae and pseudohyphae) phases. *Candida* mutants that are defective in genes regulating filamentation are reported to have reduced virulence in animal models. This reduced virulence suggests that the ability to change from a blastospore to a filament is a key virulence factor of *C. albicans*. To date, no essential effectors of these filamentation pathways have been identified in *C. albicans*. See Caesar-TonThat, T.C. and J.E. Cutler, "A monoclonal antibody to *Candida albicans* enhances mouse neutrophil candidacidal activity," Infect. Immun. 65:5354-5357, 1997.

25

Staphylococcus aureus infections also are common and increasingly result in drug resistance to antibiotics. For example, *S. aureus* is a common cause of skin and skin structure infections, endocarditis and bacteremia in the U.S. and throughout the world. Formerly community acquired *S. aureus* (CA-*S. aureus*) infections were nearly
5 uniformly susceptible to penicillinase-resistant beta lactams such as cefazolin, oxacillin, methicillin, penicillin and amoxicillin. However, over the past decade, epidemics of beta-lactam resistant *S. aureus* (MRSA) infection have been seen in multiple locales throughout the world, especially community acquired MRSA (CA-MRSA). In many places MRSA has become the predominant *S. aureus* strain causing CA infections. A
10 recent, prospective, population-based survey in three states in the U.S. estimated that the incidence of CA-MRSA infections is 500 cases per 100,000 population, which translates to approximately 1.5 million cases per year in the U.S. alone. The increasing frequency of drug-resistant *S. aureus* infections highlights the need for new ways to prevent and treat these infections.

15 The identification of effectors in the regulatory pathways of the organism that contribute to virulence offers the opportunity for therapeutic intervention with methods or compositions that are superior to existing antifungal agents. The identification of cell surface proteins that affect a regulatory pathway involved in virulence is particularly promising because characterization of the protein enable
20 immunotherapeutic techniques that are superior to existing antifungal agents when fighting a candidal infection.

The virulence of *Candida albicans* is regulated by several putative virulence factors of which adherence to host constituents and the ability to transform from yeast-to-hyphae are among the most critical in determining pathogenicity. While potent
25 antifungal agents exist that are microbicidal for *Candida*, the attributable mortality of candidemia is approximately 38%, even with treatment with potent anti-fungal agents such as amphotericin B. Also, existing agents such as amphotericin B tend to exhibit undesirable toxicity. Although additional antifungals may be developed that are less toxic than amphotericin B, it is unlikely that agents will be developed that are more
30 potent. Therefore, either passive or active immunotherapy to treat or prevent disseminated candidiasis is a promising alternative to standard antifungal therapy.

Thus, there exists a need for effective immunogens that will provide host immune protection and passive immunoprotection against *Candida*, *S. aureus* and other immunogenically related pathogens. The present invention satisfies this need and provides related advantages as well.

5

SUMMARY OF THE INVENTION

The invention provides a vaccine including an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, with an adjuvant in a pharmaceutically acceptable medium. The invention also provides a method of treating or preventing disseminated candidiasis. The method includes
10 administering an immunogenic amount of a vaccine an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium. A method of treating or preventing disseminated candidiasis also is provided that includes administering an effective amount of an isolated
15 Als protein family member having cell adhesion activity, or an functional fragment thereof, to inhibit the binding or invasion of *Candida* to a host cell or tissue. The Als protein family member can be derived from a *Candida* strain selected from the group consisting of *Candida albicans*, *Candida krusei*, *Candida tropicalis*, *Candida glabrata* and *Candida parapsilosis* and the Als protein family member includes Als1p, Als3p, Als5p, Als6p, Als7p or Als9p. Also provided is a method of treating or preventing
20 *Staphylococcus aureus* infections. The method includes administering an immunogenic amount of a vaccine an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A, 1B show the mediation of Als1p adherence of *C. albicans* to
25 human umbilical vein endothelial cells. Values represent the mean $\pm$ SD of at least three independent experiments, each performed in triplicate. (A) Endothelial cell adherence of *ALS11/als2 als1/als1* and *ALS1*-complemented mutants and wild-type CAII2(30)(B) Endothelial cell adherence of P_{ADHI}-ALS1 mutant that overexpresses *ALS1*, compared to

wild type *C. albicans*. Statistical treatment was obtained by Wilcoxon ran sum test and corrected for multiple comparisons with the Bonferroni correction. * $P < 0.001$ for all comparisons.

Figure 2A-D shows the cell surface localization of Als1p on filaments of *C. albicans* indirect immunofluorescence. Filamentation of *C. albicans* was induced by incubating yeast cells in RPMI 1640 medium with glutamine for 1.5 hours at 37°C. Als1p was detected by incubating organisms first with anti-Als1p mouse mAb followed by FITC-labeled goat anti-mouse IgG. *C. albicans* cell surface was also stained with anti-*C. albicans* polyclonal Ab conjugated with Alexa 594 (Molecular Probes, Eugene, OR). Areas with yellow staining represent Als1p localization. (A) *C. albicans* wild-type. (B) *als1/als1* mutant strain. (C) *als1/als1* complemented with wild type ALS1 (D) P_{ADHI}-ALS1 overexpression mutant.

Figure 3A, 3B show the mediation of Als1p on *C. albicans* filamentation on solid medium. *C. albicans* blastospores were spotted on Lee's agar plates and incubated at 37°C for 4 days (A) or 3 days (B).

Figure 4A, 4B show the control of *ALS1* expression and the mediation of *C. albicans* filamentation by the *EFG1* filamentation regulatory pathway. (A) Northern blot analysis showing expression of *ALS1* in (i) mutants deficient in different filamentation regulatory pathways. (ii) *efg1/efg1* mutant complemented with either *EFG1* or P_{ADHI}-ALS1. Total RNA was extracted from cells grown in RPMI 1640 + glutamine medium at 37°C for 90 minutes to induce filamentation. Blots were probed with *ALS1* and *TEF1*. (B) Photomicrographs of the *efg1/efg1* mutant and *efg1/efg1* mutant complemented with P_{ADHI}-ALS1 grown on Lee's agar plates at 37°C. for 4 days.

Figure 5A, 5B show the reduction of virulence in the mouse model of hematogenously disseminated candidiasis by (A) Male Balb/C mice (n = 30 for each yeast strain) were injected with stationary phase blastospores (10⁶ per mouse in 0.5 ml of PBS). Curves are the compiled results of three replicate experiments (n = 30 mice for each strain). The doubling times of all strains, grown in YPD at 30°C, ranged between 1.29 to 1.52 hours and were not statistically different from each other. Southern blot analysis of total chromosomal DNA was used to match the identity of the genotype of *C.*

albicans strains retrieved from infected organs with those of *C. albicans* strains used to infect the mice. Statistical analysis was obtained by Wilcoxon rank sum test and corrected for multiple comparisons with the Bonferroni correction. * $P < 0.002$ for the *als1/als1* mutant versus each of the other strains. (B) Histological micrographs of
 5 kidneys infected with *C. albicans* wild-type, homozygous *als1 null* mutant, or heterozygous *ALS1* complemented mutant. Kidney samples were retrieved 28 hours (a) or 40 (b) hours post infection, fixed in paraformaldehyde and sections were stained with silver (magnification X400). Arrows denote *C. albicans* cells.

Figure 6 shows the prophylactic effect of anti-ALS antibody against
 10 disseminated candidiasis as a function of surviving animals over a 30-day period for animals infused with anti-Als1p polyserum.

Figure 7 is polypeptide sequence alignment of the N-terminal portion of select ALS polypeptides arranged by adherence phenotype. The top three lines are the sequences from ALS1, 3 and 5 polypeptides (SEQ ID NOS: 1-3, respectively), which
 15 bind endothelial cells. The bottom three are sequences from ALS6, 7 and 9 polypeptides (SEQ ID NOS: 4-6, respectively), which do not bind endothelial cells. The last line represents the ALS polypeptide family consensus sequence (SEQ ID NO:7).

Figure 8 shows Als proteins confer substrate-specific adherence properties when heterologously expressed in *Saccharomyces cerevisiae*. Each panel demonstrates
 20 the percentage adherence of one Als expression strain (filled bars) to a variety of substrates to which *C. albicans* is known to adhere. Adherence of *S. cerevisiae* transformed with the empty vector (empty bars) is included in each panel as a negative control. Gel, gelatin; FN, fibronectin; LN, laminin; FaDU, FaDU epithelial cells; EC, endothelial cells. *, $p < 0.01$ when compared with empty plasmid control by single factor
 25 analysis of variance. Results are the mean $\pm$ S.D. of at least three experiments performed in triplicate.

Figure 9 shows domain swapping demonstrates that substrate-specific adherence is determined by the composition of the N-terminal domain of Als proteins. A representation of the *ALS* gene or construct being tested is depicted as a bar composed of
 30 sequences from *ALS5* (black) or *ALS6* (white). Adherence properties of each mutant are

displayed as a photomicrograph illustrating the adherence of transformed *S. cerevisiae* to fibronectin-coated beads and a graph demonstrating the adherence to gelatin (black bars) and endothelial cells (gray bars) as measured in the 6-well plate assay. Results are mean $\pm$ S.D. of at least three experiments, each performed in triplicate.

5 Figure 10 shows a subset of Als proteins mediate endothelial cell invasion when expressed in *S. cerevisiae*. A, endothelial cell adherence of *S. cerevisiae* strains expressing Als proteins or transformed with the empty plasmid (control). Data represent the total number of endothelial cell-associated organisms and are expressed as cells per high power field. B, degree of endothelial cell invasion of Als_p expressing *S. cerevisiae*
10 strains presented as the number of intracellular organisms per high power field. *, $p < 0.01$ when compared with empty plasmid control by single factor analysis of variance. Results are the mean $\pm$ S.D. of at least three experiments performed in triplicate.

Figure 11 shows an alignment of the N-terminal amino acid sequence of Als proteins of known function demonstrates an alternating pattern of CRs and HVRs. A,
15 percentage of consensus identity among the N-terminal regions of Als proteins of known function. Note that the signal peptide region (amino acids 1–20) is not shown. Open boxes indicate the regions designated as HVRs 1–7. B, schematic alignment of Als proteins (SEQ ID NOS:1-6, respectively) showing the composition of the individual HVRs. The sequences are arranged to compare proteins with an affinity to multiple
20 substrates with those that bind few or no identified substrates. The number of amino acids in each conserved region is indicated in parentheses.

Figure 12 shows CD and FTIR spectra of the Als1 protein N-terminal domain. A, circular dichroism spectrum of 10 μ M Als1p in phosphate-buffered saline. B, FTIR spectrum of Als1p self-film hydrated with D₂O vapor.

25 Figure 13 shows a comparison of predicted physicochemical properties of N-terminal domains among the Als protein family. Hydrophobic, electrostatic, or hydrogen-bonding features are projected onto water-accessible surfaces of each domain. Hydrophobics are shown as follows: brown, most hydrophobic; blue, most hydrophilic. Electrostatics (spectral continuum) is shown as follows: red, most positive charge (+10
30 kcal/mol); blue, most negative charge (-10 kcal/mol). Hydrogen-bonding potential (H-

binding) is shown as follows: red, donor; blue, acceptor. Als proteins are distinguishable into three groups based on the composite of these properties. For example, note the similar hydrophobic, electrostatic, and hydrogen-bonding profiles among Als group A proteins, Als1p, Als3p, and Als5p. In contrast, Als group B members, Als6p and Als7p, display striking differences in hydrophobic and electrostatic features from those of Als group A. In addition to biochemical profiles, note the differences in predicted structure among these domains.

Figure 14. Conceptual model of structural-functional relationships in Als family proteins. Als proteins are composed of three general components: an N-terminal domain, tandem repeats, and a serine/threonine-rich C-terminal domain containing a glycosylphosphatidylinositol anchor that interfaces with the *C. albicans* cell wall. As illustrated, Als proteins contain multiple conserved anti-parallel β -sheet regions (CR1-n) that are interposed by extended spans, characteristic of the immunoglobulin superfamily. Projecting from the β -sheet domains are loop/coil structures containing the HVRs. The three-dimensional physicochemical properties of specific Als protein HVRs probably govern interactions with host substrates that confer adhesive and invasive functions to *Candida*. For illustrative purposes, only three N-terminal β -sheet/coil domains and their respective CR/HVR components are shown. Note that this projection is viewed at right angles to the structural images shown in Fig. 13.

Figure 15. Immunization of mice (retired breeders) with rAls1p-N improves survival during subsequent disseminated candidiasis. Survival of mice immunized with Als1p plus adjuvant. N = 16 mice per group in duplicate experiments on different days; Adj. = adjuvant. * $p < 0.05$ vs adjuvant.

Figure 16. Immunization with rAls1p-N improves the survival of both retired breeder and juvenile mice. Survival of retired breeder (A) and juvenile (B) mice infected with a rapidly fatal, 10^6 inoculum of *C. albicans*. N = 16 mice per group in duplicate experiments on different days; Adj. = adjuvant. * $p < 0.05$ vs adjuvant control.

Figure 17. Anti-rAls1p-N titers do not correlate with survival. Titers of anti-rAls1p-N polyclonal antibodies raised in Balb/c mice immunized with varying doses of rAls1p-N with or without adjuvant. Adj. = adjuvant. * $p \leq 0.005$ for 200 μg vs. all others.

5 Figure 18. Only the protective dose of rAls1p-N induces an increase in *C. albicans*-stimulated Th1 splenocytes. Induction of Th1 ($\text{CD4}^+\text{IFN-}\gamma^+\text{IL-4}^-$) and Th2 ($\text{CD4}^+\text{IFN-}\gamma^-\text{IL-4}^+$) splenocytes by different doses of the rAls1p-N vaccine. Splenocytes from immunized mice ($n = 9$ per group) were stimulated for 48 h with heat-killed pre-germinated *C. albicans* and then analyzed by 3-color flow cytometry. * $p = 0.03$ vs.
10 adjuvant.

Figure 19. Only the protective dose of rAls1p-N induces an increase in rAls1p-N-stimulated delayed type hypersensitivity. Delayed type hypersensitivity, assessed by footpad swelling, in mice ($n = 9-12$ per group) vaccinated with rAls1p-N or CFA alone. Mice were immunized with the indicated amount of rAls1p-N and then
15 injected with 50 μg of rAls1p-N into the footpad. Footpad swelling was assessed 24 h later. * $p < 0.05$ versus adjuvant, 0.2 μg , and 200 μg .

Figure 20. The rAls1p-N vaccine requires T cells, but not B cells, to induce protective immunity. Survival of B cell-deficient, T cell-deficient (nude), and congenic wild-type Balb/c control mice ($n = 7$ or 8 per group) was simultaneously
20 assessed after vaccination with rAls1p-N + adjuvant or adjuvant alone. * $p < 0.04$ versus adjuvant alone, * $p = 0.003$ versus wild-type adjuvant-treated.

Figure 21. SQ vaccination with rAls1p-N induces an *in vivo* DTH response in immunocompetent mice. Footpad swelling was assessed 24 h after injection of 50 μg of rAls1p-N into the footpad in BALB/c mice ($n = 10$ per group). Median
25 values are displayed as black bars. * $p = 0.002$ vs. control by Wilcoxon Rank Sum test.

Figure 22. The rAls1p-N vaccine improves survival of immunocompetent mice with hematogenously disseminated candidiasis and reduces tissue fungal burden. A) Survival of vaccinated or control BALB/c mice ($n = 7$ or 10 per group for 2.5 or 5×10^5 inocula, respectively) mice subsequently infected via the tail-vein with *C. albicans*. Each

experiment was terminated at 30 days post-infection with all remaining mice appearing well. * $p < 0.05$ vs. Control by Log Rank test. B) Kidney fungal burden in BALB/c mice (n = 7 per group) infected via the tail vein with 5×10^5 blastospores of *C. albicans*. The y axis reflects the lower limit of detection of the assay. Median values are displayed as
5 black bars. * $p = 0.01$ vs control by Wilcoxon Rank Sum test.

Figure 23. The rAls1p-N vaccine induces a DTH reaction in neutropenic mice and improves their survival during subsequent hematogenously disseminated candidiasis. A) Footpad swelling was assessed 24 h after injection of 50 μ g of rAls1p-N into the footpad in BALB/c mice (n = 10 for Control, n = 8 for rAls1p-N). * $p = 0.006$ vs
10 Control by Wilcoxon Rank Sum test. B) Survival of neutropenic BALB/c mice (n = 16 per group from 2 experiments) infected with 2.5×10^4 blastospores of *C. albicans*. * $p = 0.007$ vs adjuvant control by Log Rank test.

Figure 24. The rAls1p-N vaccine diminishes the severity of histopathological fungal lesions on the tongues of mice with oropharyngeal candidiasis.
15 N = 4 mice per group. Inflammatory score generated by a blinded observer as described in the text. * $p = 0.03$ by Wilcoxon Rank Sum test.

Figure 25 shows that rAls3p-N but not rAls1p-N vaccine diminishes fungal colonization of vagina of mice inoculated with *C. albicans* (* $p=0.01$ vs mice vaccinated with CFA alone, by Wilcoxon Rank Sum test) N= 11 mice per group.

20 Figure 26 shows an Als1p homology map versus *S. aureus* clumping factor A (c1n67A). Consensus functional sites from *C. albicans* Als1p and *S. aureus* ClfA were mapped onto the Als1p homology model. Numerous residues from the N-termini of Als1p and ClfA map to a consensus cleft motif, which is where binding to substrate is predicted to occur for both adhesins.

25 Figure 27 shows that rAls1p-N and rAls3p-N vaccines improve the survival of staphylococcemic mice. (* $p<0.003$ vs mice vaccinated with CFA alone, by Log Rank test). N = 22 mice per group.

Figure 28 shows that antibody titers do not correlate with degree of protection in individual vaccinated mice, but they do distinguish unvaccinated from vaccinated mice. Titers of anti-rAls1p-N or anti-rAls3p-N polyclonal antibodies raised in Balb/c mice immunized with CFA alone, or CFA + 20 μ g of rAls1p-N or rAls3p-N, respectively. Overall there is a significant correlation between antibody titers and survival ($\rho = 0.474$, $p = 0.0057$), indicating that antibody titers can be used as a surrogate marker for vaccine protection. However, when data from mice receiving CFA alone are excluded, there is no correlation between antibody titers and survival of mice vaccinated with rAls1p-N or rAls3p-N ($\rho = 0.041143$, $p = 0.847$), indicating that antibodies are likely not the predominant mechanism of protection of the vaccine.

Figure 29 shows that the rAls1p-N vaccine protects outbred, CD1 mice from hematogenously disseminated candidiasis. A) CD1 mice ($n = 8$ per group) were vaccinated SQ with rAls1p-N (20 μ g) + CFA, or CFA alone, and infected via the tail-vein with *C. albicans* SC5314 fourteen days after the boost. B) CD1 mice ($n = 8$ per group) were vaccinated SQ with rAls1p-N at various doses with alum, or with alum alone, and infected via the tail-vein with *C. albicans* SC5314 fourteen days after the boost. * $p < 0.05$ vs. adjuvant control by Log Rank test.

Figure 30 shows that the rAls1p-N vaccine improves the survival of Balb/c mice infected with one of several strains of *C. albicans*. Survival of Balb/c mice immunized with rAls1p-N plus CFA versus CFA alone and infected via the tail-vein with *C. albicans* 15563 (7×10^5 blastospores), 16240 (4×10^5 blastospores), or 36082 (4×10^5 blastospores) ($n = 8$ mice per group). * $p < 0.05$ vs adjuvant control by Log Rank test.

Figure 31 shows that the rAls1p-N vaccine reduces tissue fungal burden in Balb/c mice infected with several non-*albicans* species of *Candida*. Balb/c mice ($n = 5$ per group) were vaccinated with CFA or CFA + rAls1p-N (20 μ g) and infected via the tail-vein with *C. glabrata*, *C. krusei*, *C. parapsilosis*, or *C. tropicalis*. Infectious inocula are shown in parentheses below the species names. Kidney fungal burden was determined on day five post-infection. The y axis reflects the lower limit of detection of the assay. * $p < 0.05$ vs. adjuvant control by non-parametric Steel test for multiple comparisons.

Figure 32 shows that rAls3p-N-immunized mice generated antibodies that cross-reacted against rAls1p-N. Titers of individual mice immunized with CFA alone, CFA + rAls1p-N, or CFA + rAls3p-N. N = 7 mice per group for CFA and CFA + rAls3p-N; n = 6 mice for CFA + rAls1p-N. *p < 0.05 vs. CFA alone; **p < 0.002 vs. CFA alone & p < 0.011 vs. CFA + rAls1p-N by Mann Whitney U test. Bars denote medians.

Figure 33 shows that both rAls1p-N and rAls3p-N primed mice for *in vivo* delayed type hypersensitivity responses. Mice (n = 7 per group for CFA and CFA + rAls3p-N; n = 6 for CFA + rAls1p-N) were vaccinated with CFA alone, CFA + rAls1p-N, or CFA + rAls3p-N. Delayed type hypersensitivity *in vivo* was measured by footpad swelling. *p < 0.05 vs. CFA alone by Mann Whitney U test. Bars denote medians.

Figure 34 shows that the rAls1p-N and rAls3p-N vaccines mediated similar efficacy against murine hematogenously disseminated candidiasis. Survival of Balb/c mice (n = 15 per group from 2 experiments for CFA and CFA + rAls3p-N, and n = 14 from 2 experiments for CFA + rAls1p-N) infected via the tail vein with 5×10^5 blastospores of *C. albicans*. The experiment was terminated at day 28 post-infection with all remaining mice appearing well. *p ≤ 0.0001 vs CFA control by Log Rank test.

Figure 35 shows that *in vivo* delayed-type hypersensitivity correlated with survival during disseminated candidiasis. Anti-rAls1p-N or anti-rAls3p-N antibody titers and footpad swelling reactions were measured in mice (n = 7 per group for CFA or CFA + rAls3p-N, n = 6 for CFA + rAls1p-N) two days prior to infection via the tail-vein with *C. albicans*. Correlations determined with the Spearman Rank sum test.

Figure 36 shows that the rAls3p-N vaccine significantly reduced tissue fungal burden during murine oropharyngeal candidiasis. Tongue fungal burden in mice (n = 7 for CFA and 8 for rAls1p-N or rAls3p-N vaccinated groups) with oropharyngeal candidiasis. The y axis reflects the lower limit of detection of the assay. *p = 0.005 vs. CFA by Mann Whitney U test.

Figure 37 shows that rAls3p-N reduced vaginal fungal burden compared to both CFA alone and CFA + rAls1p-N in murine candidal vaginitis. Vaginal fungal burden in mice (n = 11 per group from 2 experiments) vaccinated with CFA, CFA + rAls1p-N, or CFA + rAls3p-N. The y axis reflects the lower limit of detection of the assay. *p ≤ 0.02 vs CFA and CFA + rAls1p-N by Steel test for multiple comparisons.

DETAILED DESCRIPTION OF THE INVENTION

Candida albicans and *Staphylococcus aureus* are common pathogen in humans. For example, *C. albicans*, while normally a harmless commensal, this organism can cause a variety of conditions ranging from superficial mucocutaneous infection such as vaginal and/or oropharyngeal candidiasis, to deep organ involvement in disseminated candidiasis. Prior to causing disease, the fungus colonizes the gastrointestinal tract, and in some cases skin and mucous membranes. Adherence to host mucosal surfaces is a key prerequisite for this initial step. After colonization, *C. albicans* enters the bloodstream via infected intravascular devices or by transmigration through gastrointestinal mucosa compromised by chemotherapy or stress ulcerations. Organisms then disseminate via the bloodstream, bind to and penetrate the vascular endothelium to egress from the vascular tree, and invade deep organs such as liver, spleen, and kidney.

The identification and functional characterizations of a variety of exemplary Als protein family members described herein allow this family of proteins to be effectively utilized in the treatment of candidiasis. Specific binding activity to diverse substrates and other selective cell adhesion functions can be exploited in the production of vaccines for active or passive immunization, in the production of peptide, analogue of mimetic inhibitors of cell adhesion to reduce or prevent initial infection by inhibiting binding, adhesion or invasion of a host cell. Moreover, the differential binding and invasion profiles allow design and use of broad spectra or targeted inhibition of Als protein family member activities. Additionally, functional fragments that confer binding and/or invasive activity allow elimination of unwanted foreign protein sequences, thus, increasing the efficacy of the Als family protein member vaccine or therapeutic inhibitor.

The nature of the pathogenesis of *C. albicans* by adherence to endothelial cells is discussed in USP 5,578,309 which is specifically incorporated herein by reference in its entirety. For a description of the ALS1 gene and characteristics thereof, including the characterization of the gene product as an adhesin see, Fu, Y., G. Rieg, W.A. Forizi, P.H. Belanger, J.E.J. Edwards, and S.G. Filler. 1998. Expression of the *Candida albicans* gene *ALS1* in *Saccharomyces cerevisiae* induces adherence to endothelial and epithelial cells. *Infect. Immun.* 66:1783-1786; Hoyer, L.L. 1997. Fu Y, Ibrahim AS, Sheppard DC, Chen Y-C, French SW, Cutler JE, Filler SG, Edwards, JE, Jr. 2002. *Candida albicans* Als1p: an adhesin that is a downstream effector of the *EFG1* filamentation pathway. *Molecular Microbiology* 44:61-72. Sheppard DC, Yeaman MR, Welch WH, Phan QT, Fu Y, Ibrahim AS, Filler SG, Zhang M, Waring AJ, Edwards, Jr., JE 2004. Functional and Structural Diversity in the Als Protein Family of *Candida albicans*. *Journal Biological Chemistry*. 279: 30480-30489. The ALS gene family of *Candida albicans*. International Society for Human and Animal Mycology Salsimorge, Italy:(Abstract); Hoyer, L.L., S. Scherer, A.R. Shatzman, and G.P. Livi. 1995. *Candida albicans* *ALS1*: domains related to a *Saccharomyces cerevisiae* sexual agglutinin separated by a repeating motif. *Mol. Microbiol.* 15:39-54.

In this regard, the human fungal pathogen *Candida albicans* colonizes and invades a wide range of host tissues. Adherence to host constituents plays an important role in this process. Two members of the *C. albicans* Als protein family (Als1p and Als5p) have been found to mediate adherence and exemplify the binding, adhesion and cell invasion activities of Als protein family members. As described herein, members of the ALS gene family were cloned and expressed in *S. cerevisiae* to characterize their individual functions. Distinct Als proteins conferred distinct adherence profiles to diverse host substrates. Using chimeric Als5p-Als6p constructs, the regions mediating substrate-specific adherence were localized to the N-terminal domains in Als proteins. In particular, a subset of Als proteins also mediated endothelial cell invasion, a previously unknown function of this family. Consistent with these results, homology modeling revealed that Als members contain anti-parallel β -sheet motifs interposed by extended regions, homologous to adhesions or invasins of the immunoglobulin superfamily. This finding was confirmed using circular dichroism and Fourier transform infrared spectrometric analysis of the N-terminal domain of Als1p. Specific regions of amino acid

hypervariability were found among the N-terminal domains of Als proteins, and energy-based models predicted similarities and differences in the N-terminal domains that probably govern the diverse function of Als family members. Collectively, these results indicate that the structural and functional diversity within the Als family provides *C.*
5 *albicans* with an array of cell wall proteins capable of recognizing and interacting with a wide range of host constituents during infection.

The invention provides a vaccine having an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, and an adjuvant in a pharmaceutically acceptable medium. The vaccine can be an Als protein
10 family member derived from a *Candida* species such as *Candida albicans*, *Candida krusei*, *Candida tropicalis*, *Candida glabrata* or *Candida parapsilosis*. The Als protein family member can be, for example, Als1p, Als3p, Als5p, Als6p, Als7p and Als9p, or an immunogenic fragment thereof. All other Als protein family members within an *Candida* species can similarly be employed as a vaccine of the invention.

15 The present invention utilizes the gene product of *C. albicans* agglutinin like sequence protein family member as a vaccine to treat, prevent, or alleviate disseminated candidiasis. The vaccine is effective against different strains of *C. albicans* as well as against different *Candida* species. The Als protein family member can be, for example, Als1p, Als3p, Als5p, Als6p, Als7p and Als9p. The invention exploits the role
20 of the ALS gene products in the adherence of and invasion by *C. albicans* to endothelial and/or epithelial cells and the susceptibility of the Als protein family member-expressed surface protein for use as a vaccine to retard the pathogenesis of the organism.

Pursuant to this invention, an ALS family member gene encodes a surface adhesin that is selected as the target of an immunotherapeutic strategy against *C.*
25 *albicans*. A demonstration that the expression product of the *ALS1* gene, the Als1p protein, has structural characteristics typical of surface proteins and is, in fact, expressed on the cell surface of *C. albicans* is one criterion for proteins that act as adhesins to host tissues. The Als protein family members can be structurally characterized as having a signal peptide at the N-terminus, a glycosylphosphatidylinosine (GPI) anchorage
30 sequence in the C-terminus, and a central region comprising repeats rich in threonine and serine. Also, Als protein family members have N-, and O- glycosylation sites, typical of

proteins that are expressed on the cell surface. Indirect immunofluorescence using a monoclonal antibody directed against the N-terminus of ALs1p, for example, revealed that ALs1p is expressed during the log phase of blastospores. This expression of ALs1p is increased during hyphal formation and is localized to the junction where the hyphal element extends from the blastospores as indicated by the diffused surface staining. Furthermore, this monoclonal antibody blocked the enhanced adherence of *C. albicans* overexpression mutant to endothelial cells, thereby establishing the principle for immunotherapy applications using ALs1p. Functional characteristics as they relate to cell adhesion and invasion of other Als family members are described further below in Example VI.

Thus, according to one aspect, the invention provides an Als family member surface adhesion protein, designated, for example, Als1p, Als3p, Als5p, Als6p, Als7p and Als9p, or a functional fragment, conjugate or analogue thereof, having useful properties when formulated in a pharmaceutical composition and administered as a vaccine with or without an adjuvant. An Als protein family member, combination of two or more Als protein family members or one or more functional fragments, analogues, conjugates or derivatives thereof, can be obtained from, for example, *Candida albicans*. Similar adhesin or invasin molecules or analogues or derivatives thereof can be of candidal origin and can be obtainable, for example, from species belonging to the genera *Candida*, for example *Candida parapsilosis*, *Candida krusei*, *Candida glabrata* and *Candida tropicalis*. A surface adhesin or invasin protein according to the invention can be obtained in isolated or purified form, and thus, according to one embodiment of the invention a substantially pure Als protein family member *Candida* surface adhesin protein, or functional fragment, immunogenic fragment, analogue, conjugate or derivative thereof, is formulated as a vaccine to cause an immune response in a patient to elicit an immune response against *Candida* and/or to block adhesion of the organism to the endothelial cells. Fragments of Als protein family members that exhibit similar binding, adhesion or invasion activity as an intact Als protein family member is referred to herein as a functional fragment. Fragments of Als protein family members that are capable of eliciting an antibody or cellular immune response against a *Candida* species are referred to herein as an immunogenic fragment. Exemplary functional fragments include the N-terminal polypeptide region of the Als protein family member described further below in

Example VI. Exemplarily immogenic fragments include the N-terminal Als polypeptide region, the C-terminal Als polypeptide region as well as any other Als fragment that is sufficient to generate an antibody, cellular or both an antibody and cellular immune response. Such immogenic fragments can be as small as about four amino acids and as
5 large as the intact polypeptide as well as include all polypeptide lengths in between.

An analogue or derivative of the surface adhesion protein according to the invention can be identified and further characterized by the criteria described herein for an ALS family member gene and/or gene product. For example, a null mutant of the analogue or derivative would share markedly reduced adhesion to endothelial cells
10 compared to controls. Similarly, over-expression of the analogue or derivative in an appropriate model would show an increased adherence to endothelial cells compared to controls and would be confirmed as a cell surface adhesin in accord with the criteria described above. Also, antisera to an analogue or derivative can cross-react with anti-Als protein family member antibodies and can exhibit increased survival times when
15 administered in a mouse model of disseminated *candidiasis* as disclosed herein.

The invention also provides a method of treating or preventing disseminated candidiasis. The method includes administering an immunogenic amount of a vaccine an isolated Als protein family member having cell adhesion or invasion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium. The
20 vaccine can be administered with or without an adjuvant. The Als protein family member can be derived from different *Candida* strains as well as from different *Candida* species such as *Candida albicans*, *Candida krusei*, *Candida tropicalis*, *Candida glabrata* and *Candida parapsilosis*. An Als protein family member used in the method of treating or prevention disseminated candidiasis includes Als1p, Als3p, Als5p, Als6p, Als7p and
25 Als9p.

The effectiveness of the vaccines of the invention against different *Candida* strains, different *Candida* species, other bacteria and infectious agents and their wide range of immune activity are described further below and exemplified in the Examples. For example, Example V shows that anti-ALS antibodies are effective against
30 mucosal and hematogenously disseminated *candidal* infections. Example VII shows that vaccination with rAls1p-N improves survival during murine disseminated candidiasis by

enhancing cell-mediated immunity. Example VIII shows that the vaccines of the invention reduce fungal burden and improve survival in both immunocompetent and immunocompromised mice. Example IX shows the effectiveness of the ALS vaccines of the invention against *S. aureus* infections. Example X exemplifies that the vaccines of the invention are effective against different strains of *C. albicans* and against different species such as *C. glabrata*, *C. krusei*, *C. parapsilosis* and *C. tropicalis* as well as effectiveness in different animal models. Example XI also exemplifies the effectiveness of the different vaccines of the invention in different animal models as well as provides a comparison of the different responses elicited and potency of two representative ALS vaccines.

The invention further provided is a method of treating or preventing disseminated candidiasis that includes administering an effective amount of an isolated Als protein family member having cell adhesion activity, or an functional fragment thereof, to inhibit the binding or invasion of *Candida* to a host cell or tissue. The Als protein family member can be derived from *Candida albicans*, *Candida krusei*, *Candida tropicalis*, *Candida glabrata*, and *Candida parapsilosis*. An Als protein family member used in the method of treating or prevention disseminated candidias includes Als1p, Als3p, Als5p, Als6p, Als7p and Als9p. The cell adhesion activity includes binding to gelatin, fibronectin, laminin, epithelial cells or endothelial cells and/or promoting cell invasion.

In addition, the invention also provides a method of treating or preventing *Staphylococcus aureus* infections using the Als protein family members described herein. In particular, the method of treating or preventing *Staphylococcus aureus* infections includes administering an immunogenic amount of a vaccine an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium.

Als1p and Als3p are particularly efficacious because of significant homology to *S. aureus* cell surface proteins. The sequence and structural homology of, for example, Als1p and Als3p, are described further below in Example IX. Given the teachings and guidance provided herein, those skilled in the art will understand that the vaccines and methods of the invention can be applied to the treatment of *Candida* and

Staphylococcus infections alike. Similarly, given the teachings and methods described herein, those skilled in the art also will understand that the vaccines and methods of the invention also can be applied to other pathogens having cell surface polypeptides with similar immunogenicity, sequence and/or structural homology to the Als protein family members described herein, including fungus, bacteria and the like.

Immunotherapeutic and/or Als polypeptide inhibition of cell adhesion or invasion strategies against *Candida* or *Staphylococcus* infection can operate at the level of binding to the vascular endothelial cells as well as through a downstream effector of the filamentation regulatory pathway. An immunotherapeutic strategy or inhibition of binding using a soluble Als protein family member or functional fragment is useful in this context because: (i) the morbidity and mortality associated with hematogenously disseminated candidiasis and other infectious pathogens remains unacceptably high, even with currently available antifungal therapy; (ii) a rising incidence of antifungal and antibiotic resistance is associated with the increasing use of antifungal and antibacterial agents, iii) the population of patients at risk for serious *Candida* and *Staphylococcus* infections is well-defined and very large, and includes post-operative patients, transplant patients, cancer patients and low birth weight infants; and iv) a high percentage of the patients who develop serious *Candida* infections are not neutropenic, and thus may respond to a vaccine or a competitive polypeptide or compound inhibitor. For these reasons, *Candida* and *Staphylococcus* are attractive fungal and bacterial targets for passive immunotherapy, active immunotherapy or a combination of passive or active immunotherapy. Additionally, *Candida* also is attractive for competitive inhibition using an Als protein family member polypeptide, functional fragment thereof and/or a compound or mimetic thereof that binds to one or more Als family members and prevents binding of *Candida* to a host cell receptor.

Given the teachings and guidance provided herein, those skilled in the art will understand that immunotherapeutic methods well known in the art can be employed with the Als protein family members of the invention, immunogenic fragments, analogues, conjugates, and/or derivatives thereof, to use one or more of the molecule as an immunogen in a pharmaceutically acceptable composition administered as a vaccine with or without an adjuvant. For the purposes of this invention, the terms “pharmaceutical” or “pharmaceutically acceptable” refer to compositions formulated by

known techniques to be non-toxic and, when desired, used with carriers or additives that can be safely administered to humans. Administration can be performed using well known routes including, for example, intravenous, intramuscular, intraperitoneal or sub-cutaneous injection. Such vaccines of the inventions also can include buffers, salts or
5 other solvents known to those skilled in the art to preserve the activity of the vaccine in solution. Similarly, any of a wide range of adjuvants well known in the art can be employed with the vaccine of the invention to elicit, promote or enhance a therapeutically effective immune response capable of reducing or blocking binding, invasion and/or infection of *Candida* or *Staphylococcus* to a susceptible host cell.

10 Similarly, given the teachings and guidance provided herein, those skilled in the art also will understand that therapeutic methods well known in the art for administering and selectively blocking the binding of cell surface molecules to their cognate receptors also can be employed with the Als protein family members of the invention, functional fragments, analogues, conjugates and/or derivatives thereof, to use
15 one or more of the Als protein family member as an inhibitor in a pharmaceutically acceptable composition. As with vaccine formulations, inhibitory formulations can similarly be administered using well known method in the art including, for example, intravenous intramuscular, intraperitoneal or sub-cutaneous injection. Such inhibitory compositions that bind Als family member receptors and block an Als protein family
20 member binding also can include buffers, salts or other solvents known to those skilled in the art to preserve the activity of the vaccine in solution. Further, any of a wide range of formulations well known in the art can be employed with the inhibitory compositions of the invention to target and/or enhance delivery or uptake so as to reduce or inhibit binding, invasion and/or infection of *Candida* or *Staphylococcus* to a susceptible host
25 cell.

With respect to the molecule used as a therapeutic immunogen or receptor binding inhibitor pursuant to the present invention, those of skill in the art will recognize that the Als protein family member molecules can be truncated or fragmented without losing the essential qualities as an immunogenic vaccine or cell adhesion or invasion
30 inhibitor. For example, an Als protein family member can be truncated to yield an N-terminal fragment by truncation from the C-terminal end with preservation of the functional properties described above and further below in the Examples. Similarly, C-

terminal fragments can be generated by truncation from the N-terminal end with preservation of their functional properties. Other modifications in accord with the teachings and guidance provided herein can be made pursuant to this invention to create other Als protein family member functional fragments, immunogenic fragments, analogs
5 or derivatives thereof, to achieve the therapeutically useful properties described herein with the native protein.

One aspect of the therapeutic effectiveness of Als protein family members and methods of the invention achieves interference with regulation of filamentation, to block adherence of the organism to host constituents, and to enhance clearance of the
10 organism by immunoeffector cells and other physiological mechanisms. Since endothelial cells cover the majority of the vasculature, strategies to block the adherence, invasion and/or both of the organism to endothelial cells using antibodies, Als family member proteins, polypeptide or peptides or any combination thereof include useful embodiment of the present invention. As described previously, such adherence and/or
15 invasion blocking therapies include active or passive immunotherapy or inhibitory binding directed against the candidal adhesins, invasins, or cognate receptors disclosed herein. Thus, for example, any suitable host can be injected with protein and the serum collected to yield the desired anti-adhesin antibody after appropriate purification and/or concentration. Prior to injection, the adhesin or invasin protein or a combination thereof,
20 can be formulated in a suitable vehicle preferably a known immunostimulant such as a polysaccharide or delivery formulation such as liposomes or time-released compositions. Thus, according to a further aspect, invention provides a pharmaceutical composition comprising a candidal adhesin or invasin protein together with one or more pharmaceutically acceptable excipients in a formulation for use as a vaccine or Als
25 receptor inhibitor.

The method of the invention is ameliorating and/or preventing candidal or *Staphylococcus* infection by blocking the adherence of *C. albicans* to the endothelial or epithelial cells of a host constituent or by, for example, antibody binding to the *Staphylococcus* and allowing immune mechanisms remove the pathogen. Thus,
30 according to one aspect of the invention, a pharmaceutical composition comprising an Als protein family member adhesin or invasin protein, functional or immunogenic fragment, derivative, analogue, or conjugate thereof is formulated as a vaccine or Als receptor

inhibitor in a pharmaceutical composition containing a biocompatible carrier for injection or infusion and is administered to a patient. Also, direct administration of antiserum raised against Als family member protein or isolated or recombinant Als family member protein can be used to block the adherence of *C. albicans* to a mammalian host

5 constituent or effect the removal of a *Staphylococcus* pathogen. Antiserum against adhesin protein can be obtained by known techniques, Kohler and Milstein, Nature 256: 495-499 (1975), and may be humanized to reduce antigenicity, see USP 5,693,762, or produced in transgenic mice leaving an unrearranged human immunoglobulin gene, see USP 5,877,397. Similarly, isolated or recombinant Als protein family member also can

10 be produced using methods well known to those skilled in the art including, for example, the recombinant production described in the Examples below.

A still further use of the invention, for example, is using the Als protein family member adhesin or invasin protein to develop vaccine strategies for the prevention and/or amelioration of candidal or *Staphylococcus* infections. Thus, according to one

15 aspect of the invention, for example, standard immunology techniques can be employed to construct a multi-component vaccine strategy that can enhance and/or elicit immune response from a host constituent to block adherence of *C. albicans* or to effect the elimination of *Staphylococcus* pathogens.

A still further use of the invention, for example, is developing DNA

20 vaccine strategies. Thus, according to one aspect of the invention, for example, the ALS family member polynucleotides encoding Als protein family member adhesin or invasin or a functional fragment thereof is administered according to a protocol designed to yield an immune response to the gene product. See e.g., Felgner USP 5,703,055.

A still further use of the invention, for example, is developing combination

25 vaccine strategies. Thus, according to one aspect of the invention, for example, anti-ALS protein family member antibodies may be used with antibodies in treating and/or preventing candidal or *Staphylococcus* infections. See USP 5,578,309.

The following Examples illustrate the immunotherapeutic utility of the ALS1 adhesin as the basis for preventive measures or treatment of disseminated

30 candidiasis. Example 1 describes the preparation of an ALS1 null mutant and a strain of

C. albicans characterized by overexpression of ALS1 to confirm the mediation of adherence to endothelial cells. Example 2 describes the localization of Als1p and the implication of the efg filamentation regulatory pathway. Example 3 describes the purification of ALS1 adhesin protein. Example 4 describes the preparation of rabbit polyclonal antibodies raised against the ALS1 surface adhesin protein to be used to demonstrate the blocking of the surface adhesin protein. Example 5, describes the blocking of adherence *in vivo*, using polyclonal antibodies raised against the ALS1 surface adhesion protein as described herein according to the invention to protect against disseminated candidiasis in a mouse model. Example VI describes the structural and functional characteristics of Als protein family members.

It is understood that modifications which do not substantially affect the activity of the various embodiments of this invention are also included within the definition of the invention provided herein. Accordingly, the following examples are intended to illustrate but not limit the present invention.

15

EXAMPLE I

Als1 Mediates Adherence of *C. albicans* to Endothelial Cells

The URA blaster technique was used to construct a null mutant of *C. albicans* that lacks express of the Als1p. The als1/als1 mutant was constructed in *C. albicans* strain CAI4 using a modification of the Ura-blaster methodology (Fonzi and Irwin, *Genetics* 134, 717 (1993)) as follows: Two separate als1-hisG-URA3-hisG-als1 constructs were utilized to disrupt the two different alleles of the gene. A 4.9 kb ALS1 coding sequence was generated with high fidelity PCR (Boehringer Mannheim, Indianapolis, IN) using the primers: 5'-CCCTCGAGATGCTTCAACAATTTACATTGTTA-3' (SEQ ID NO:8) and 5'-CCGCTCGAGTCACTAAATGAACAAGGACAATA-3' (SEQ ID NO:9). Next, the PCR fragment was cloned into pGEM-T vector (Promega, Madison, WI), thus obtaining pGEM-T-ALS1. The hisG-URA3-hisG construct was released from pMG-7 by digestion with Kpn1 and Hind3 and used to replace the portion of ALS1 released by Kpn1 and Hind3 digestion of pGEM-T-ALS1. The final als1-hisG-URA3-hisG-als1 construct was released from the plasmid by digestion with Xho1 and used to disrupt the first allele of ALS1 by transformation of strain CAI-4.

A second als1-hisG-URA3-hisG-als1 construct was generated in two steps. First, a Bgl2-Hind3 hisG-URA3-hisG fragment of pMB7 was cloned into the BamH1-Hind3 sites of pUC19, thereby generating pYC2. pYC2 was then digested with Hind3, partially filled in with dATP and dGTP using T4 DNA polymerase, and then digested
5 with Sma1 to produce a new hisGURA3-hisG fragment. Second, to generate ALS1 complementary flanking regions, pGEM-T-ALS1 was digested with Xba1 and then partially filled in with dCTP and dTTP. This fragment was digested with Hpa1 to delete the central portion of ALS1 and then ligated to the hisG-URA3-hisG fragment generating pYC3. This plasmid was then digested by Xho1 to release a construct that was used to
10 disrupt the second allele of the ALS1. Growth curves were done throughout the experiment to ensure that the generated mutations had no effect on growth rates. All integrations were confirmed by Southern blot analysis using a 0.9kb ALS1 specific probe generated by digestion of pYF5 with XbaI and HindIII.

The null mutant was compared to *C. albicans* CAI-12 (a URA + revertant strain) for its ability to adhere *in vitro* to human umbilical vein endothelial cells. For
15 adherence studies, yeast cells from YPD (2% glucose, 2% peptone, and 1 % yeast extract) overnight culture, were grown in RPMI with glutamine at 25°C for 1 hour to induce Als1p expression. 3×10^2 organisms in Hanks balanced salt solution (HBSS) (Irvine Scientific, Irvine, CA) were added to each well of endothelial cells, after which the plate
20 was incubated at 37°C for 30 minutes. The inoculum size was confirmed by quantitative culturing in YPD agar. At the end of incubation period, the nonadherent organisms were aspirated and the endothelial cell monolayers were rinsed twice with HBSS in a standardized manner. The wells were overlaid with YPD agar and the number of adherent organisms were determined by colony counting. Statistical treatment was
25 obtained by Wilcoxon rank sum test and corrected for multiple comparisons with the Bonferroni correction. $P < 0.001$.

Referring to Figure 1, a comparison of the ALS1/ALS1 and als1/als1 strain showed that the ALS1 null mutant was 35% less adherent to endothelial cells than *C. albicans* CAI-12. To reduce background adherence, the adherence of the wild-type strain
30 grown under non-ALS1 expressing conditions was compared with a mutant autonomously expressing Als1p. This mutant was constructed by integrating a third copy of ALS1 under the control of the constitutive ADH1 promoter into the wild-type *C.*

albicans. To achieve constitutive expression of the ALS1 in *C. albicans*, a blunt-ended PCR generated URA3 gene is ligated into a blunt-edged Bgl2 site of pOCUS-2 vector (Novagen, Madison, WI), yielding pOU-2. A 2.4 kb Not1-Stul fragment, which contained *C. albicans* alcohol dehydrogenase gene (ADH1) promoter and terminator
5 (isolated from pLH-ADHpt, and kindly provided by A. Brown, Aberdeen, UK), was cloned into pOU-2 after digestion with Not1 and Stul. The new plasmid, named pOAU-3 had only one Bgl2 site between the ADH1 promoter and terminator. ALS1 coding sequence flanked by BamH1 restriction enzyme sites was generated by high fidelity PCR using pYF-5 as a template and the following primers: 5'-CGGGATCCAGATGCTTCA-
10 ACAATTTACATTG-3' (SEQ ID NO:10) and 5'-CGGGATCCTCACTAATGAACAAGGACAATA-3' (SEQ ID NO:11). This PCR fragment was digested with BamH1 and then cloned into the compatible Bgl2 site of pOAU-3 to generate pAU-1. Finally, pAU-1 was linearized by XbaI prior to transforming *C. albicans* CAI-4. The site-directed integration was confirmed by Southern Blot
15 analysis. Referring to Figure 1B, overexpressing ALS1 in this P_{ADH1}-ALS1 strain resulted in a 76% increase in adherence to endothelial cells compared to the wild-type *C. albicans*. In comparing endothelial cell adherence of the wild-type to that of the overexpressing mutant, yeast cells were grown overnight in YPD at 25°C (non-inducing condition of Als1p). Als1p expression was not induced to reduce the background
20 adherence of the wild-type, thus magnifying the role of Als1p in adherence through P_{ADH1}-ALS1 hybrid gene. The adherence assay was carried out as described above. Statistical treatment was obtained by Wilcoxon rank sum test and corrected for multiple comparisons with the Bonferroni correction. $P < 0.001$.

A monoclonal anti-Als1p murine IgG antibody was raised against a
25 purified and truncated N-terminus of Als1p (amino acid #17 to #432) expressed using Clontech YEXpress (TM) Yeast Expression System (Palo Alto, CA). The adherence blocking capability of these monoclonal anti-Als1p antibodies was assessed by incubating *C. albicans* cells with either anti-Als1 antibodies or mouse IgG (Sigma, St. Louis, MO) at a 1:50 dilution. After which the yeast cells were used in the adherence assay as described
30 above. Statistical treatment was obtained by Wilcoxon rank sum test and corrected for multiple comparisons with the Bonferroni correction. $P < 0.001$. The results revealed that the adherence of the P_{ADH1}-ALS1 strain was reduced from $26.8\% \pm 3.5\%$ to $14.7\% \pm 5.3\%$.

Thus, the effects of ALS1 deletion and overexpression demonstrate that Als1p mediates adherence of *C. albicans* to endothelial cells.

EXAMPLE II

Localization of Als1p

5 For Als1p to function as an adhesin, it must be located on the cell surface. The cell surface localization of Als1p was verified using indirect immunofluorescence with the anti-Als1p monoclonal antibody. Diffuse staining was detected on the surface of blastospores during exponential growth. This staining was undetectable on blastospores in the stationary phase. Referring to Figure 2A, when blastospores were induced to
10 produce filaments, intense staining was observed that localized exclusively to the base of the emerging filament. No immunofluorescence was observed with the *als1/als1* mutant, confirming the specificity of this antibody for Als1p. See Figure 2B. These results establish that Als1p is a cell surface protein.

 The specific localization of Als1p to the blastospore-filament junction
15 implicates Als1p in the filamentation process. To determine the mechanism, the filamentation phenotype of the *C. albicans* ALS1 mutants was analyzed. Referring to Figure 3A, the *als1/als1* mutant failed to form filaments after a 4 day incubation on Lee's solid medium, while both the ALS1/ALS1 AND ALS1/*als1* strains as well as the ALS1-complemented mutant produced abundant filaments at this time point. The *als1/als1*
20 mutant was capable of forming filaments after longer periods of incubation. Furthermore, overexpressing ALS1 augmented filamentation: the P_{ADHI}-ALS1 strain formed profuse filaments after a 3 day incubation, whereas the wild-type strain produced scant filaments at this time point. See Figure 3B. To further confirm the role of Als1p in filamentation, a negative control was provided using mutant similar to the ALS1 overexpression mutant,
25 except the coding sequence of the ALS1 was inserted in the opposite orientation. The filamentation phenotype of the resulting strain was shown to be similar to that of the wild-type strain. The filament-inducing properties of Als1p are specific to cells grown on solid media, because all of the strains described above filamented comparably in liquid media. The data demonstrates that Als1p promotes filamentation and implicates ALS1
30 expression in the regulation of filamentation control pathways. Northern blot analysis of ALS1 expression in mutants with defects in each of these pathways, including *efg1/efg1*,

cph1/cph1, efg1/efg cph1/cph1, tup1/tup1, and cla4/cla4 mutants were performed. Referring to Figure 4A, mutants in which both alleles of EFG1 had been disrupted failed to express ALS1. Introduction of a copy of wild-type EFG1 into the efg1/efg1 mutant restored ALS1 expression, though at a reduced level. See Figure 4B. Also, as seen in
5 Figure 4A, none of the other filamentation regulatory mutations significantly altered ALS1 expression (Fig. 4A). Thus, Efg1p is required for ALS1 expression.

If Efg1p stimulates the expression of ALS1, which in turn induces filamentation, the expression of ALS1 in the efg1/efg1 strain should restore filamentation. A functional allele of ALS1 under the control of the ADH1 promoter was integrated into
10 the efg1/efg1 strain. To investigate the possibility that ALS1 gene product might complement the filamentation defect in efg1 null mutant, an Ura efg1 null mutant was transformed with linearized pAU- 1. Ura⁺ clones were selected and integration of the third copy of ALS1 was confirmed with PCR using the primers: 5'-
CCGTTTATACCATCCAATC-3' (SEQ ID NO:12) and 5'-CTACA
15 TCCTCCAATGATATAAC-3' (SEQ ID NO:13). The resulting strain expressed ALS1 autonomously and regained the ability to filament on Lee's agar. See Figures 4B and C. Therefore, Efg1p induces filamentation through activation of ALS1 expression.

Because filamentation is a critical virulence factor in *C. albicans* delineation of a pathway that regulates filamentation has important implications for
20 pathogenicity. Prior to ALS1, no gene encoding a downstream effector of these regulatory pathways had been identified. Disruption of two other genes encoding cell surface proteins, HWP1 AND INT1, results in mutants with filamentation defects. Although HWP1 expression is also regulated by Efg1p, the autonomous expression of HWP1 in the efg1/efg1 mutant fails to restore filamentation. Therefore Hwp1p alone
25 does not function as an effector of filamentation downstream of EFG1. Also, the regulatory elements controlling INT1 expression are not know. Thus, Als1p is the first cell-surface protein identified that functions as a downstream effector of filamentation, thereby suggesting a pivotal role for this protein in the virulence of *C. albicans*.

The contribution of Als1p to *C. albicans* virulence was tested in a model
30 of hematogenously disseminated candidiasis, A.S. Ibrahim *et al.*, *Infect. Immun.* 63, 1993 (1995). Referring to Figure 5A, mice infected with the als1/als1 null mutant survived

significantly longer than mice infected with the ALS1/ALS1 strain, the ALS1/als1 mutant or the ALS1-complemented mutant. After 28 hours of infection, the kidneys of mice infected with the als1/als1 mutant contained significantly fewer organisms (5.70 ± 0.46 10^6 CFU/g) ($P < 0.0006$ for both comparisons). No difference was detected in colony counts of organisms recovered from spleen, lungs, or liver of mice infected with either of the strains at any of the tested time points. These results indicate that Als1p is important for *C. albicans* growth and persistence in the kidney during the first 28 hours of infection. Referring to Figure 5B, examination of the kidneys of mice after 28 hours of infection revealed that the als1/als1 mutant produced significantly shorter filaments and elicited a weaker inflammatory response than did either the wild-type or ALS1-complemented strains. However, by 40 hours of infection, the length of the filaments and the number of leukocytes surrounding them were similar for all three strains.

The filamentation defect of the als1/als1 mutant seen on histopathology paralleled the *in vitro* filamentation assays on solid media. This mutant showed defective filamentation at early time points both *in vivo* and *in vitro*. This defect eventually resolved with prolonged infection/incubation. These results suggest that a filamentation regulatory pathway that is independent of ALS1 may become operative at later time points. The activation of this alternative filamentation pathway by 40 hours of infection is likely the reason why mice infected with the als1/als1 mutant subsequently succumbed in the ensuing 2-3 days.

Collectively, these data demonstrate that *C. albicans* ALS1 encodes a cell surface protein that mediates both adherence to endothelial cells and filamentation. Als1p is the only identified downstream effector of any known filamentation regulatory pathway in *C. albicans*. Additionally, Als1p contributes to virulence in hematogenous candidal infection. The cell surface location and dual functionality of Als1p make it an attractive target for both drug and immune-based therapies.

EXAMPLE III

Purification of ALS1 Adhesin Protein

The ALS1 protein synthesized by *E. coli* is adequate as an immunogen. However eukaryotic proteins synthesized by *E. coli* may not be functional due to

improper folding or lack of glycosylation. Therefore, to determine if the ALS1 protein can block the adherence of *C. albicans* to endothelial cells, the protein is, preferably, purified from genetically engineered *C. albicans*.

5 PCR was used to amplify a fragment of ALS1, from nucleotides 52 to 1296. This 1246 bp fragment encompassed the N-terminus of the predicted ALS1 protein from the end of the signal peptide to the beginning of the tandem repeats. This region of ALS1 was amplified because it likely encodes the binding site of the adhesin, based on its homology to the binding region of the *S. cerevisiae* Aga1 gene product. In addition, this portion of the predicted ALS1 protein has few glycosylation sites and its size is
10 appropriate for efficient expression in *E. coli*.

 The fragment of ALS1 was ligated into pQE32 to produce pINS5. In this plasmid, the protein is expressed under control of the *lac* promoter and it has a 6-hits tag fused to its N-terminus so that it can be affinity purified. We transformed *E. coli* with pINS5, grew it under inducing conditions (in the presence of IPTG), and then lysed the
15 cells. The cell lysate was passed through a Ni²⁺-agarose column to affinity purify the ALS1-6His fusion protein. This procedure yielded substantial amounts of ALS1-6His. The fusion protein was further purified by SDS-PAGE. The band containing the protein was excised from the gel so that polyclonal rabbit antiserum can be raised against it. It will be appreciated by one skilled in the art that the surface adhesin protein according to
20 the invention may be prepared and purified by a variety of known processes without departing from the spirit of the present invention. The sequence of Als1p is listed in Figure 7.

EXAMPLE IV

Raising Polyclonal Antisera against ALS1 Protein

25 To determine whether antibodies against the ALS1 protein block the adherence of *Candida albicans* to endothelial and epithelial cells, and the selected host constituent *in vitro*, rabbits were inoculated with *S. cerevisiae* transformed with ALS1 protein. The immunization protocol used was the dose and schedule used by Hasenclever and Mitchell for production of antisera that identified the antigenic relationship among
30 various species of *Candida*. Hasenclever, H. F. and W. O. Mitchell. 1960. Antigenic

relationships of *Torulopsis glabrata* and seven species of the genus *Candida*. J. Bacteriol. 79:677-681. Control antisera were also raised against *S. cerevisiae* transformed with the empty plasmid. All yeast cells were grown in galactose to induce expression of the ALS genes. Before being tested in the adherence experiments, the serum was heat-inactivated at 56 C to remove all complement activity.

Sera from immunized rabbits were absorbed with whole cells of *S. cerevisiae* transformed with empty plasmid to remove antibodies that are reactive with components of the yeast other than ALS1 protein. The titer of the antisera was determined by immunofluorescence using *S. cerevisiae* that express the ALS1 gene.

10 FITC-labeled anti-rabbit antibodies were purchased from commercial sources (Southern Biotechnology, Inc). Affinity-purified secondary antibodies were essential because many commercially available sera contain antibodies reactive with yeast glucan and mannan. The secondary antibodies were pretested using *Candida albicans* as well as *S. cerevisiae* transformed with the plasmid and were absorbed as needed to remove any anti-*S.*

15 *cerevisiae* or anti-*Candida* antibodies. Negative controls were 1) preimmune serum 2) *S. cerevisiae* transformed with the empty plasmid, and 3) *S. cerevisiae* transformed with the ALS gene but grown under conditions that suppress expression of the ALS gene (glucose).

In addition to the above experiments, Western blotting was used to provide

20 further confirmation that an antiserum binds specifically to the ALS protein against which it was raised. *S. cerevisiae* transformed with the ALS1 were grown under inducing conditions and their plasma membranes were isolated by standard methods. Panaretou R and P. Piper. 1996. Isolation of yeast plasma membranes. p. 117- 121. In I.H. Evans. (ed.), Yeast Protocols. Methods in Cell and Molecular Biology. Humana Press, Totowa,

25 New Jersey. Plasma membranes were also prepared from *S. cerevisiae* transformed with the empty plasmid and grown under identical conditions. The membrane proteins were separated by SDS-PAGE and then transferred to PVDF membrane by electroblotting. Harlow, E. and D. Lane. 1988. Antibodies: a laboratory manual. Cold Spring Harbor Laboratory Press. After being blocked with nonfat milk, the blot was incubated with the

30 ALS antiseru. The preabsorbed antiserum did not react with proteins extracted from *S. cerevisiae* containing empty plasmid. This antiserum blocked the adherence of *S. cerevisiae* pYF5 (a clone that expresses *Candida albicans* ALS1) to endothelial cells.

EXAMPLE V

Polyclonal Antibodies Against Specific ALS Proteins Prophylactically Protect Mice from Mucosal and Hematogenously Disseminated Candidal Infections

Having identified the antisera that block the adherence of a clone of *S. cerevisiae* transformed with an ALS gene under the above conditions, these antisera were demonstrated to protect mice from intravenous challenge with *Candida albicans*.

The antisera against the ALS proteins were first tested in the murine model of hematogenously disseminated candidiasis. Affinity-purified anti-ALS antibodies are effective in preventing adhesion of yeast cells to various substrates (see EXAMPLE 3). Affinity-purification is useful in this system because antibody doses can be accurately determined. Moreover, the unfractionated antisera will undoubtedly contain large amounts of antibody directed toward antigens on the *S. cerevisiae* carrier cells. Many of these anti-*Saccharomyces* antibodies would likely bind to *C. albicans* and make interpretation of the results impossible. Additionally, it is quite possible that the procedure used to elute antibodies from *S. cerevisiae* that express the ALS protein may also elute small amounts of yeast mannan or glucan that could have adjuvant-like activity. The immunoaffinity-purified antibodies are further purified before use. They may also be preabsorbed with mouse splenocytes.

Antibody doses may be administered to cover the range that brackets the levels of serum antibody that can be expected in most active immunization protocols and to cover the range of antibody doses that are typically used for passive immunization in murine models of candidiasis. See Dromer, F., J. Charreire, A. Contrepois, C. Carbon, and P. Yeni. 1987, Protection, of mice against experimental cryptococcosis by *anti-Cryptococcus neoformans* monoclonal antibody, Infect. Immun. 55:749-752; Han, Y. and J.E. Cutler. 1995, Antibody response that protects against disseminated candidiasis, Infect. Immun. 63:2714-2719; Mukherjee, J., M.D. Scharff, and A. Casadevall. 1992, Protective murine monoclonal antibodies to *Cryptococcus neoformans*, Infect. Immun. 60:4534-4541; Sanford, J.E., D.M. Lupan, A.M. Schlageter, and T.R. Kozel. 1990, Passive immunization against *Cryptococcus neoformans* with an isotype-switch family of monoclonal antibodies reactive with cryptococcal polysaccharide, Infect. Immun. 58:1919-1923. BALB/c Mice (femal, 7 week old, the NCI) were given anti-ALS that had

been absorbed with mouse splenic cells by an intraperitoneal (i.p.) injection. Control mice received prebled serum that had been absorbed with mouse splenic cells, intact anti-ALS serum, or DPBS, respectively. For the pre-absorption, 2 ml of anti-ALS or prebled sera were mixed with 100 μ l of mouse (BALB/c, 7 weeks old female, NCI) splenic cells
5 (app. 9×10^6 cells per ml) at room temperature for 20 minutes. The mixture was washed with warm sterile DPBS by centrifugation (@300 xg) for 3 minutes. This procedure was repeated three times. The volume of i.p. injection was 0.4 ml per mouse. Four hours later, the mice were challenged with *C. albicans* (strain CA-1; 5×10^5 hydrophilic yeast cells per mouse by i.v. injection. Then, their survival times were measured. See Figure 6.

10 Previous studies have shown that antibodies administered via the intraperitoneal route are rapidly (within minutes) and almost completely transferred to the serum (Kozel and Casadevall, unpublished observations). As a control for effects of administering the antibody preparations, a parallel group of mice were treated with antibodies isolated from pre-immune serum that has been absorbed with *S. cerevisia*
15 transformed with the ALS gene. The survival time and numbers of yeast per gram of kidney were measured. Again, referring to Figure 6, mice infected intravenously with 10^6 blastopores of ALS1 null mutant had a longer median survival time when compared to mice infected with *Candida albicans* CAI-12 or *Candida albicans* in which one allele of the ALS1 had been deleted (p=0.003).

20 These results indicate that immunotherapeutic strategies using the ALS1 proteins as a vaccine have a protective prophylactic effect against disseminated candidiasis.

EXAMPLE VI

Functional and Structural Diversity in the Als Protein Family of *Candida albicans*

25 Isolation and characterized of the *C. albicans* *ALS1* gene by heterologous complementation of nonadherent *S. cerevisiae* has been previously described (Fu et al., Infect. Immun. 66:1783-1786 (1998)). *ALS1* encodes a cell surface protein that mediates

adherence to endothelial and epithelial cells. Disruption of both copies of this gene in *C. albicans* is associated with a 35% reduction in adherence to endothelial cells, and overexpression of *ALS1* increases adherence by 125% (Fu et al., Mol. Microbiol. 44:61-72 (2002)).

5 *ALS1* is a member of a large *C. albicans* gene family consisting of at least eight members originally described by Hoyer et al. (Hoyer et al., Trends Microbiol. 9:176-180 (2001), Zhao et al., Microbiology 149:2947-2960 (2003)). These genes encode cell surface proteins that are characterized by three domains. The N-terminal region contains a putative signal peptide and is relatively conserved among Als proteins.
10 This region is predicted to be poorly glycosylated (Zhao et al., Microbiology 149:2947-2960 (2003), Hoyer et al., Genetics 157:1555-1567 (2001)). The central portion of these proteins consists of a variable number of tandem repeats (~36 amino acids in length) and is followed by a serine-threonine-rich C-terminal region that contains a glycosylphosphatidylinositol anchor sequence (*supra*). Whereas the proteins encoded by
15 this gene family are known to be expressed during infection (Hoyer et al., Infect. Immun. 67:4251-4255 (1999), Zhang et al., Genome Res. 13:2005-2017 (2003)), the function of the different Als proteins has not been investigated in detail.

Heterologous expression of Als proteins in nonadherent *S. cerevisiae* was performed to evaluate the function of Als proteins and to avoid the high background
20 adherence mediated by the multiple other adhesins expressed by *C. albicans*. This heterologous expression system has been used extensively for the study of *C. albicans* genes, including the isolation and characterization of the adhesins *ALS1*, *ALS5*, and *EAP1* (Li et al., Eukaryot Cell 2:1266-1273 (2003), Fu et al., Infect. Immun. 66:1783-1786 (1998), Gaur et al., Infect. Immun. 65:5289-5294 (1997)). As described further below,
25 using this model system Als proteins were demonstrated to have diverse adhesive and invasive functions. Consistent with these results, homology modeling indicated that Als proteins are closely related in structure to adhesin and invasin members of the immunoglobulin superfamily of proteins. Structural analyses using CD and Fourier transform infrared (FTIR)1 spectrometry confirmed that the N-terminal domain of Als1p
30 is composed of anti-parallel β sheet, turn, α -helical, and unstructured domains consistent with the structures of other members of the immunoglobulin superfamily. Finally,

comparative energy-based models suggest differences in key physicochemical properties of the N-terminal domains among different Als proteins that may govern their distinct adherence and invasive biological functions.

To clone *ALS* family members and express them in *S. cerevisiae*, *ALS1*, -3, -5, -6, -7, and -9 were successfully amplified and expressed as described below. Briefly, for cloning and other culture steps, *S. cerevisiae* strain S150-2B (leu2 his3 trp1 ura3) was used for heterologous expression as has been described previously (Fu et al., Infect. Immun. 66:2078-2084 (1998)). *C. albicans* strain SC5314 was used for genomic cloning. All strains were grown in minimal defined medium (1x yeast nitrogen base broth (Difco), 2% glucose, and 0.5% ammonium sulfate, supplemented with 100 µg/ml L-leucine, - L tryptophan, L-histidine, and adenine sulfate) solidified with 1.5% bacto-agar (Difco) as needed. Growth of ura-strains was supported by the addition of 80 µg/ml uridine (Sigma). Plasmids pGK103, containing ALS5, pYF5, containing ALS1, and pALS_n, containing ALS9, have been described previously (Fu et al., Infect. Immune. 66:1783-1786 (1998), Gaur et al., Infect. Immune. 65:5289-5297 (1997), Lucinod et al., *Proceedings of the 102nd Annual Meeting of the American Society for Microbiology*, pp. 204, American Society for Microbiology, Salt Lake City, Ut. (2002)). Plasmid pADH1, obtained from A. Brown (Aberdeen, UK) contains the *C. albicans* alcohol dehydrogenase gene (ADH1) promoter and terminator, which are functional in *S. cerevisiae* (Bailey et al., J. Bacteriol. 178:5353-5360 (1996)). This plasmid was used for constitutive expression of ALS genes in *S. cerevisiae*.

Human oral epithelial and vascular endothelial cells were obtained and cultured as follows. The FaDu oral epithelial cell line, isolated from a pharyngeal carcinoma, was purchased from the American Type Culture Collection (ATCC) and maintained as per their recommended protocol. Endothelial cells were isolated from umbilical cord veins and maintained by our previously described modification of the method of Jaffe et al. (Fu et al., Mol. Microbiol. 44:61-72 (2002), Jaffe et al., J. Clin. Invest. 52:2745-2756 (1973)). All cell cultures were maintained at 37 °C in a humidified environment containing 5% CO₂.

For cloning the ALS genes, genomic sequences of members of the ALS family were identified by BLAST searching of the Stanford data base (available on the

World Wide Web at URL: sequence.stanford.edu/group/candida/search.html). PCR primers were generated to specifically amplify each of the open reading frames that incorporated a 5' BglII and a 3' XhoI restriction enzyme site and are shown below in Table I (SEQ ID NOS:14-19 (ALS 1, 3, 5, 6, 7 and 9 sense primers, respectively); SEQ ID NOS:20-25 ((ALS 1, 3, 5, 6, 7 and 9 antisense primers, respectively)). Each gene was cloned by PCR using the Expand[®] High Fidelity PCR system (Roche Applied Science). ALS3, ALS6, and ALS7 were amplified from *C. albicans* SC5314 genomic DNA, whereas ALS1, ALS5, and ALS9 were amplified from plasmids that had been previously retrieved from *C. albicans* genomic libraries (Fu et al., *Infect. Immune.* 66:1783-1786 (1998), Gaur et al., *Infect. Immune.* 65:5289-5297 (1997), Lucinod et al., *Proceedings of the 102nd Annual Meeting of the American Society for Microbiology*, pp. 204, American Society for Microbiology, Salt Lake City, Ut. (2002)). PCR products were ligated into pGEM-T-Easy (Promega) for sequencing. Sequence-verified ALS open reading frames were then released from pGEM-T-Easy by BglII-XhoI co-digestion and ligated into pADH1, such that the ALS gene of interest was under the control of the ADH1 promoter. *S. cerevisiae* strain S150-2B was transformed with each of the ALS overexpression constructs as well as the empty pADH1 construct using the lithium acetate method. Expression of each ALS gene in *S. cerevisiae* was verified by Northern blot analysis before phenotypic analyses were performed.

20

Table I

PCR primers used to amplify the coding regions of ALS gene for heterologous expression in S. cerevisia

<i>ALS gene</i>	Sense (5' -3')	Antisense (5' -3')
<i>ALS1</i>	AGATCTCAGATGCTTCAACAATTT ACATTG	CTCGAGTCACTAAATGAACAAGGACA ATA
<i>ALS3</i>	GAAGATCTATGCTACAACAATATA CATTGTTACTC	CCGCTCGAGTTAAATAAACAAGGATAA TAATGTGATC
<i>ALS5</i>	AGATCTCAACTACCAACTGCTAAC A	CTCGAGACCATATTATTTGGTACAATC
<i>ALS6</i>	AGATCTCATTCACCGACAATGAAG ACA	CTCGAGTTGGTACAATCCCGTTTGA
<i>ALS7</i>	AGATCTTCAACAGTCTAATACCTA TGA	CTCGAGACTTGATTGAATTATACCATA TA
<i>ALS9</i>	AGATCTCGAATGCTACCACAATTC CTA	CTCGAGTCTTAGCACCCTGACGTAGCT

ALS mRNA expression was detected by Northern blot analysis for each construct. Despite the use of three sets of primers, amplification of ALS2 and ALS4 from genomic DNA of *C. albicans* SC5314 was unsuccessful. Given the difficulty of sequencing and assembling across the tandem repeats of ALS genes, it is possible that
5 this outcome reflects errors in the sequence assembly currently available on the published genome data base.

Flow cytometry confirmed that each of the Als proteins was expressed on the surface of their respective *S. cerevisiae* hosts. Briefly, confirmation of cell surface expression for each of the Als constructs was determined using indirect
10 immunofluorescence employing two different polyclonal anti-Als antisera. Antiserum A consisted of anti-Als1p antibodies, generated by immunization of rabbits with a 417-amino acid N-terminal fragment of Als1p. Antiserum B was rabbit anti-*C. albicans* mannan factor 5 that recognizes *C. albicans* cell wall components but does not cross-react with *S. cerevisiae* (Iatron Laboratories).

15 For each strain, 10^7 blastospores were isolated from overnight culture, blocked with 100 μ l of goat serum, and then stained with either polyclonal antiserum A or B at a 1:25 dilution, followed by fluorescein isothiocyanate-labeled goat anti-rabbit IgG at 1:100. A FACSCaliber (Becton Dickinson) instrument equipped with an argon laser emitting at 488 nm was used for flow cytometric analyses. Fluorescence emission was
20 detected with a 515/40-nm bandpass filter. Fluorescence data for 10,000 events were collected, and the distribution of cells with fluorescence above base line (i.e. *S. cerevisiae* transformed with the empty plasmid) was analyzed for each strain using CELLQUEST software (Becton Dickinson).

As shown in Table II, two distinct antisera demonstrated that all of the
25 Als_p-expressing strains exhibited at least a 4-fold increase in fluorescence when compared with *S. cerevisiae* transformed with the empty plasmid. Consistent with the predicted structural diversity among members of the Als family, the antisera displayed differences in recognition of individual Als expression strains.

Table II

Detection of Als proteins on the surface of S. cerevisiae by flow cytometric analysis

5 Blastospores of each strain were stained using indirect immunofluorescence with either polyclonal anti-Als1p antiserum (A) or polyclonal anti-*C. albicans* cell wall antiserum (B) and then analyzed using flow cytometry. Results are expressed as percentage of positive cells above background (*S. cerevisiae* transformed with empty plasmid), with – fold increase in parentheses.

Percentage of cells above background (-fold increase)		
<i>Als construct</i>	Antiserum A	Antiserum B
<i>Empty plasmid</i>	(1)	(1)
<i>Als1p</i>	47.8 (17)	50.1 (19)
<i>Als3p</i>	24.5 (9)	54.0 (20)
<i>Als5p</i>	23.5 (8)	28.2 (11)
<i>Als6p</i>	12.7 (4)	16.2 (6)
<i>Als7p</i>	22.1 (8)	15.7 (6)
<i>Als9p</i>	11.4 (4)	33.9 (13)

10 *S. cerevisiae* clones that expressed the various Als proteins were examined for their ability to adhere to a variety of host substrates. As described below, the results show that Als proteins display different profiles of substrate-specific adherence.

Fungal adherence assays were performed to determine the adherence properties of transformed *S. cerevisiae* strains. Briefly, a modification of previously described adherence assay (8) was employed as follows. Adherence plates were coated by adding 1 ml of a 0.01 mg/ml solution of gelatin (Sigma), laminin (Sigma), or fibronectin (Becton Dickinson) to each well of a 6-well tissue culture plate (Costar) and incubating overnight at 37 °C. For endothelial cells, second passage cells were grown to confluence in 6-well tissue culture plates coated with a 0.2% gelatin matrix, and for epithelial cells, FaDU cells were grown to confluence (3 days) in 6-well tissue culture plates coated with a 0.1% fibronectin matrix. Before adherence testing, wells were washed twice with 1 ml of warm Hanks' balanced salt solution (HBSS). *S. cerevisiae* strains to be tested were grown overnight in minimal defined media at 30 °C and then harvested by centrifugation, washed with HBSS (Irvine Scientific), and enumerated using a hemacytometer. Three hundred organisms were added to each well of a 6-well tissue

culture plate coated with the substrate of interest and incubated for 30 min at 37 °C in CO₂. Nonadherent organisms were removed by washing twice in a standardized manner with 10 ml of HBSS. The wells were overlaid with YPD agar (1% yeast extract (Difco), 2% bacto-peptone (Difco), 2% D-glucose, 1.5% agar), and the inoculum was confirmed
5 by quantitative culture. Plates were incubated for 48 h at 30 °C, and the colonies were counted. Adherence was expressed as a percentage of the initial inoculum. Differences in adherence were compared using a single factor analysis of variance test, with $p < 0.01$ considered significant.

There were striking differences in the adherence profiles of the *S. cerevisiae* transformants to the different substrates (Fig. 8). Whereas Als1p-, Als3p-, and Als5p-expressing strains bound to all substrates tested, Als6p-expressing *S. cerevisiae* adhered only to gelatin, and Als9p-expressing *S. cerevisiae* adhered above background levels only to laminin. Further, there were quantitative differences in adherence to the various substrates. For example, when compared with Als3p, Als1p conferred greater
10 adherence to gelatin but less adherence to epithelial cells ($p < 0.01$, single factor analysis of variance). Only *S. cerevisiae* expressing Als7p adhered to none of the substrates tested. Whereas small differences in levels of Als protein expression cannot be ruled out by the immunofluorescence studies shown in Table II, such differences are unlikely to be responsible for the substrate-specific binding patterns found in this study. Such a global
15 increase or decrease in the amount of Als protein expressed on the cell surface would be expected to produce a commensurate increase or decrease in adherence across all substrates and not result in the substrate-specific differences that were observed.
20

As described below, the substrate binding specificity for Als proteins resides in the N-terminal sequences of Als Proteins. Briefly, Als5p expression in *S. cerevisiae* conferred adherence to multiple substrates, including gelatin and endothelial
25 cells, whereas Als6p expression resulted in adherence to gelatin alone. Despite this marked difference in function, Als5p and Als6p are more than 80% identical at the amino acid level. The tandem repeat and C-terminal portions of these proteins are virtually identical, and the majority of the sequence differences are concentrated in the N termini
30 of these two proteins. These data indicate that N-terminal sequence variability confers substrate specificity.

The above result was supported by the results of studies determining the adherence phenotypes of chimeric ALS5/ALS6 constructs. Briefly, chimeric Als5/Als6 proteins were constructed by exchanging the N termini of each protein. Chimeric ALS5/6 genes were constructed as follows. A BglII-HpaI fragment of ALS5 encompassing the 5' 2117 bp of the gene was isolated. pGEM-T-ALS6 was then digested with BglII and HpaI to release the corresponding 5' 2126 bp of ALS6, and the fragment consisting of pGEM-T-Easy plus the 3' sequences of ALS6 was isolated and ligated to the 5' ALS5 fragment to generate plasmid pGEM-T-5N6C. An identical approach using the corresponding 5' fragment of ALS6 and 3' fragment of ALS5 was used to generate plasmid p-GEM-T-6N5C. After sequence confirmation, each chimeric ALS gene was released by BglII-XhoI digestion and subcloned into pADH1 as above. *S. cerevisiae* S150-2B was then transformed with these constructs, and expression was verified by Northern blot analysis before characterization of their adherence properties.

S. cerevisiae expressing a chimeric fusion of the N terminus of Als5p to the C terminus of Als6p adhered to both gelatin and endothelial cells in a manner similar to Als5p (Fig. 9). Likewise, strains expressing the chimeric fusion of the Als6 N terminus to the C terminus of Als5p adhered only to gelatin, as did *S. cerevisiae* expressing Als6p (Fig. 9). Further, strains expressing Als5p and chimeric Als5N6C protein agglutinated fibronectin-coated beads, whereas those expressing Als6p and chimeric Als6N5C protein had little to no affinity for these beads. Collectively, these data indicate that the adherence profiles of these transformed *S. cerevisiae* strains were governed by the N-terminal portion of the Als protein.

In addition to the differences in substrate specificity demonstrated between the Als protein family members, differences in other biological functions also were observed. For example, a subset of Als proteins was shown to mediate endothelial cell invasion by *S. cerevisiae*. *C. albicans* invades endothelial cells by inducing its own endocytosis (Filler et al., Infect. Immun. 63976-983 (1995), Belanger et al., Cell Microbiol., in press (2002)). This endocytosis occurs after the organism adheres to endothelial cells; however, the *C. albicans* ligands required for this process are unknown. Further, it is unclear if distinct candidal ligands are required for both adherence and endocytosis. In addition to being nonadherent, *S. cerevisiae* does not undergo significant endocytosis by endothelial cells. Therefore, to test whether Als proteins could serve as

invasins as well as adhesins, the ability of *S. cerevisiae* strains expressing Als proteins to invade endothelial cells was determined.

The ability of Als proteins to mediate endothelial cell invasion was determined using a modification of a previously described differential fluorescence assay (Phan et al., Infect. Immun. 68:3485-3490 (2000)). Briefly, endothelial cells were grown to confluence on 12-mm diameter glass coverslips coated with fibronectin and placed in a 24-well tissue culture plate (Corning). Cells were then infected with 10^5 blastospores of each *S. cerevisiae* strain in RPMI 1640 medium (Irvine Scientific). As a positive control, cells were infected with a similar number of *C. albicans* blastospores. After incubation for 90 min, the cells were rinsed twice with 0.5 ml of HBSS in a standardized manner and fixed with 3% paraformaldehyde. Organisms remaining adherent to the surface of the endothelial cells were stained for 1 h with the rabbit anti-*C. albicans* antiserum (Biodesign), which had been conjugated with Alexa 568 (Molecular Probes, Inc., Eugene, OR), which fluoresces red. This antiserum cross-reacts with *S. cerevisiae* at a 2-fold higher dilution. The endothelial cells were then permeabilized in 0.2% Triton X-100 in phosphate-buffered saline for 10 min, after which the cell-associated organisms (the internalized plus adherent organisms) were again stained with the anti-*C. albicans* antiserum conjugated with Alexa 488, which fluoresces green. The coverslips were then observed under epifluorescence. The number of organisms that had been internalized by the endothelial cells was determined by subtracting the number of adherent organisms (fluorescing red) from the number of cell-associated organisms (fluorescing green). At least 100 organisms were counted on each coverslip, and all experiments were performed in triplicate on at least three separate occasions.

Fibronectin bead adherence assays also was performed to further characterize the binding characteristics of certain Als proteins. In this regard, Als5p was originally identified by virtue of the protein's ability to induce agglutination of fibronectin-coated beads when expressed on the surface of *S. cerevisiae* (Gaur et al., Infect. Immune. 65:5289-5297 (1997)). Therefore, *S. cerevisiae* strains transformed with ALS5, ALS6, 5N6C, and 6N5C for fibronectin were tested for bead adherence using this methodology (Gaur et al., Infect. Immune. 65:5289-5297 (1997), Gaur et al., Infect. Immun. 67:6040-6047 (1999)). Briefly, tosylated magnetic beads (DynaL Biotech) were coated with fibronectin following the manufacturer's instructions. Next, 10 μ l of coated

beads (10^6 beads) were mixed with 1×10^8 transformed *S. cerevisiae* in 1 ml of 1x Tris-EDTA (TE) buffer, pH 7.0, and incubated with gentle mixing for 45 min. The tubes were placed in a magnet to separate beads and adherent *S. cerevisiae* from nonadherent organisms. The supernatant containing nonadherent organisms was removed by
5 aspiration, and the remaining beads were washed three times by resuspending in 1 ml of TE buffer, followed by magnetic separation and aspiration of the supernatant. Finally, the washed beads and adherent organisms were resuspended in 100 μ l of TE buffer and examined microscopically for co-agglutination.

The results show that *S. cerevisiae* expressing Als1p, Als3p, and Als5p
10 displayed a significant increase in the percentage of cell-associated organisms, reflecting their ability to adhere to endothelial cells. In addition, organisms expressing Als3p and, to a lesser extent, Als1p and Als5p demonstrated significant endothelial cell invasion (Fig. 10).

In addition to the functional studies described above, Als proteins also
15 were found to be homologous to adhesins and invasins of the immunoglobulin superfamily. As an initial step in the molecular modeling of Als proteins, a knowledge-based search algorithm was used to identify molecules that share significant structural similarity with Als family members. Briefly, homology and energy-based modeling was conducted to compare overall physicochemical features of Als proteins. First, a
20 knowledge-based method (SWISS-MODEL) (Guex et al., Electrophoresis 18:2714-2723 (1997), Schwede et al., Nucleic Acids Res. 31:3381-3385 (2003)) was used to analyze and compare combinatorial extension structural alignments of structures in the Swiss and Brookhaven protein data bases for proteins with homologous conformation (Shindyalov et al., Protein Eng. 11:739-747 (1998)). This approach included the BLASTP2 algorithm
25 (Altschul et al., Mol. Biol. 215:403-410 (1990)) to search for primary sequence similarities in the ExNRL-3D data base. In parallel, the dynamic sequence alignment algorithm SIM (Huang et al., Adv. Appl. Math. 12:337-367 (1991)) was used to select candidate templates with greatest sequence identity. Subsequently, ProModII was used to conduct primary and refined match analyses. Resulting proteins were used as templates
30 for homology modeling of Als protein backbone trajectories.

Robust models of the N-terminal domains of Als proteins (e.g. amino acids 1–480; preceding initial tandem repeats) were generated through complementary approaches. The N-terminal domains of Als proteins were converted to putative solution conformations by sequence homology (Composer (Topham et al. Biochem. Soc. Symp. 57:1-9 (1990)) and threading methods (Matchmaker (Godzik et al., J. Mol. Biol. 227:227-238 (1992)) and Gene-Fold (Jaroszewski et al., Protein Sci. 7:1431-1440 (1998), Godzik et al., Protein Eng. 8:409-416 (1995), Godzik et al., Proc. Natl. Acad. Sci. U.S.A. 89:12098-12102 (1992), Godzik et al., J. Comput. Aided Mol. Des. 7:397-438 (1993)) using SYBYL 6.9.1 software (Tripos Associates) operating on Silicon Graphics workstations (SGI, Inc.). Resulting conformers and amino acid side chains of target Als domains were refined by molecular dynamics, and strain energies were minimized using the AMBER95 force field method (Duan et al., J. Comput. Chem. 24:1999-2012 (2003)) and the Powell minimizer (Powell et al., Math. Program 12:241-254 (1977)).

These approaches optimize side chain interactions where positions of the peptide backbone atoms are fixed. Preferred conformations were determined from extended molecular dynamics in aqueous solvent. Next, the torsion angles of all peptide bonds were adjusted to $180 \pm 15^\circ$, with minimal constraints. In some cases, molecular dynamics were executed, either with no constraints or with α -helical regions constrained by applying a 0.4-kJ penalty to the canonical Ramachandran ϕ and ψ angles. Final global energy minimizations were performed for each model after the removal of all constraints and aggregates. Resulting Als N-terminal domain models were prioritized based on three criteria: (i) most favorable strain energy (molecular mechanics); (ii) empirical positional energy functions; and (iii) preservation of the spatial arrangement of potential disulfide bridging (Godzik et al., J. Mol. Biol. 227:227-238 (1992), Bowie et al., Science 253:164-170 (1991), Eisenberg et al., Methods Enzymol. 277:396-404 (1997), Fischer et al., FASEB J. 10:126-136 (1996), Luthy et al., Nature 356:83-85 (1992)). Als models were assessed for validity in relationship to homology templates using standard measures (e-values (Welch et al., Biochemistry 35:7165-7173 (1996), Welch et al., Biochemistry 33:6074-6085 (1994)). Finally, the physicochemical properties of the Als models were visualized by MOLCAD (Heiden et al., J. Comput. Chem. 14:246-250 (1993)), as

implemented in SYBYL and HINT platforms (Kellog et al., J. Comput. Aided Mol. Des. 5:545-552 (1991)), such that the physical properties were projected onto the water-accessible surface of the Als N-terminal domains.

These models indicate that the N-terminal domains of all Als proteins
5 contain multiple anti-parallel β -sheet domains, consistent with members of the immunoglobulin superfamily. The results are summarized below in Table III. These proteins typically consist of complex seven-stranded anti-parallel β -sheet domains, from which project loop/coil structures. The β -sheet domains are separated from one another by interposing regions. This structure is often referred to as a beads-on-a-string motif.
10 Particularly noted is that virtually all of the Als proteins modeled to known adhesin or invasin homologs (Table III). Different patterns of similarity were observed among the Als proteins analyzed. For example, all Als proteins examined, except Als7p, shared significant homology with collagen-binding protein of *Staphylococcus aureus*. However, the specific primary, secondary, and tertiary homologs varied for most family members
15 (Table III). For example, Als2p and Als9p shared an identical primary, secondary, and tertiary homolog.

Table III

Comparison of homologs among Als proteins

Homologues of each Als protein were identified by the knowledge-based algorithm described and were ranked in descending order of structural correlation from 1 to 3. NS, no significant model identified for homology modeling (correlation coefficient (r^2) $\leq$ 70%. PDB, Protein Data Bank code per the National Center for Biotechnology Information format.

Protein	Homolog 1	Homolog 2	Homolog 3
Als1p	Invasin/integrin-binding protein Yersinia pseudotuberculosis (PDB 1cwv) ^a	Collagen-binding protein <i>Staphylococcus aureus</i> (PDB 1d2p) ^a	Clumping factor <i>S. aureus</i> (PDB 1n67A) ^b
Als2p	Collagen-binding protein <i>S. aureus</i> (PDB 1d2p) ^a	Invasin/integrin-binding protein <i>Y. pseudotuberculosis</i> (PDB 1cwv) ^b	Surface layer protein <i>Methanosarcina mazei</i> (PDB 1LOQA) ^c
Als3p	Collagen-binding protein <i>S. aureus</i> (PDB 1d2p) ^a	Invasin/integrin-binding protein <i>Y. pseudotuberculosis</i> (PDB 1cwv) ^b	Clumping factor <i>S. aureus</i> (PDB 1n67A) ^c
Als4p	Collagen-binding protein <i>S. aureus</i> (PDB 1d2p) ^a	Invasin/integrin-binding protein <i>Y. pseudotuberculosis</i> (PDB 1cwv) ^b	NS
Als5p	Invasin/integrin-binding protein Yersinia pseudotuberculosis (PDB 1cwv) ^b	Surface layer protein <i>M. mazei</i> (PDB 1LOQA) ^b	Collagen-binding protein <i>S. aureus</i> (PDB 1d2p) ^c
Als6p	Collagen-binding protein <i>S. aureus</i> (PDB 1d2p) ^b	Invasin/integrin-binding protein <i>Y. pseudotuberculosis</i> (PDB 1cwv) ^b	Neuraminidase Influenza virus type B (PDB 1nsca) ^c
Als7p	Surface layer protein <i>M. mazei</i> (PDB 1LOQA) ^b	NS	NS
Als9p	Collagen-binding protein <i>S. aureus</i> (PDB 1d2p) ^a	Invasin/integrin-binding protein <i>Y. pseudotuberculosis</i> (PDB 1cwv) ^b	Surface layer protein <i>M. mazei</i> (PDB 1LOQA) ^c

Als proteins were also determined to contain N-terminal hypervariable regions that map to predicted loop/coil structures. In this regard, despite the observed differences in substrate-specific adherence mediated by individual Als proteins, large regions of sequence in the N-terminal domains are conserved across this family. However, seven regions of significant divergence among Als proteins designated hypervariable regions (HVRs) 1–7, were found. These regions (composed of 8 or more amino acids) contained no apparent consensus identity across Als proteins and less than 50% consensus conservation. In contrast, the intervening conserved regions (CRs) 1–7, displayed more than 30% consensus identity and more than 50% consensus conservation across Als proteins. An identity plot and schematic alignment of these amino acid sequences comprising the N-terminal domains (residues 1–420) of Als proteins with known function is presented in Fig. 11, A and B. In particular, homology modeling revealed that the HVRs of different Als proteins, while distinguishable in sequence, are predicted to conform to similar loop/coil structures that project from the β -sheet components of the CRs. Thus, the presence of these conserved HVRs indicate that they are available to interact with host constituents.

In addition to the homology modeling and related determinations described above, empirical determinations additionally confirm the predicted structure of the N-terminal domain of Als1p. To test the hypotheses generated by our homology modeling, the structural features of the N-terminal domain of Als1p was determined using the complementary approaches of CD and FTIR spectrometry. This protein, encompassing amino acids 17–432 of Als1p, was produced in *S. cerevisiae* and has been described previously by Fu, et al., Molecular Microbiology, 44:61-72 (2002).

Briefly, circular dichroic spectra were recorded with an AVIV 62DS spectropolarimeter (Aviv Biomedical Inc.) fitted with a thermoelectric temperature controller. Aqueous solutions of Als1p (10 μ M in phosphate-buffered saline) were scanned using 0.1-mm light path demountable quartz cells at a rate of 10 nm/min from 260 to 185 nm and a sample interval of 0.2 nm. Spectra from buffer lacking peptide were subtracted from sample solutions to minimize light scattering artifacts, and final spectra were an average of 8 scans recorded at 25 °C. The instrument was routinely calibrated

with (+)-10-camphorsulfonic acid (1 mg/ml in a 1-mm path length cell) (Johnson et al., Proteins 7:205-214 (1990)), and ellipticity was expressed as the mean residue ellipticity (1)MRE (degrees-cm² dmol⁻¹). The protein concentration was determined by absorbance at 280 nm based on aromatic amino acid composition of the expressed Als1p domain (Pace et al, Protein Sci 4:2411-2423 (1995)). The CD spectra were deconvoluted into helix, β -sheet, turn, and disordered structures using Selcon (Sreerama et al, Protein Sci. 8:370-380 (1999)) through the internet-based Dichroweb (Lobley et al., Bioinformatics 18:211-212 (2002)) interface (cryst.bbk.ac.uk/cdweb/html/home.html).

Infrared spectra of Als1p self-films were recorded at 25 °C on a Bruker Vector 22 FTIR spectrometer (Bruker Optics) fitted with a deuterated triglycine sulfate detector at a gain of 4, averaged over 256 scans, and at a resolution of 2 cm⁻¹. Fifty micrograms of the protein in 50 μ l of phosphate-buffered saline were spread onto the surface of a 50 x 20 x 2-mm germanium attenuated total reflectance sample crystal (Pike Technologies) and allowed to dry. The dry protein self-film was then hydrated with D₂O for 1 h prior to recording the infrared spectra. Amide I bands of the infrared spectra were analyzed for secondary conformations by area calculations of component peaks with curve-fitting software (GRAMS/32, Version 5; Galactic). The frequency limits for the various conformations were as follows: α -helix (1662–1645 cm⁻¹), β -sheet (1637–1613 and 1710–1682 cm⁻¹), β -turn loops (1682–1662 cm⁻¹), and disordered structures (1645–1637 cm⁻¹) (50–52).

Circular dichroism results of the N-terminal domain of Als1p are shown in Figure 12A and reveal a dichroic minimum at 217 nm and strong positive dichroic maximum near 200 nm. These features are characteristic of a protein having a dominant anti-parallel β sheet component. Deconvolution of the CD spectra indicated that the protein assumed conformations of 50.1% β sheet, whereas other structure class contributions include disordered structures (26.9%), turn structures (19.3%), and α -helix (3.7%).

As shown in Figure 12B, FTIR measurements of a self-film of the hydrated Als1p strongly corroborated that the sample has a dominant β -sheet conformation. These spectra revealed strong low frequency amide I bands with peaks centered at 1634 and 1628 cm⁻¹ and a weak high frequency band centered at 1685 cm⁻¹.

This frequency splitting of the protein amide I infrared spectra into high and low frequency components has been shown to be typical of the effect of transition dipole coupling between intermolecular anti-parallel β -sheets (Halverson et al., J. Am. Chem. Soc. 113:6701-6703 (1991)). Curve fitting of the spectra indicated that the protein
5 construct is ~57.2% antiparallel β -sheet. Other secondary structural conformations from curve fitting of the IR spectra include disordered structures (20.5%), turn components (13.3%), and α -helix (9.0%).

Taken together, the FTIR and CD data further corroborate that the N terminus of Als1p contains predominant domains of anti-parallel β -sheet structure
10 containing minor α -helical and turn components, interposed by less structured regions.

Three-dimensional models further indicate Physicochemical distinctions among Als N-terminal domains. In this regard, molecular models indicated differences in predicted physicochemical attributes of the N-terminal domains of Als proteins that likely influence their interactions with host cells and several substrates. As shown in Figure 13,
15 Als proteins are separable into three distinct groups based on surface distributions of hydrophobicity, charge, and hydrogen bonding potential. Als1p, Als3p, and Als5p each share similar patterns of these properties and thus are considered the Als group A. In contrast, the predicted physicochemical properties of Als6p and Als7p N-terminal domains (Als group B) have striking differences from those of the Als group A (Fig. 13).
20 Whereas the cationic potential in Als group A members is typically segregated from their neutral or anionic facets, positive charge is broadly distributed across the entire surface of the Als group B members Als6p and Als7p. Finally, the N termini of Als2p, Als4p, and Als9p appear to constitute a third group of Als proteins (the Als group C) that differ structurally from either the Als group A or B proteins. The Als group C proteins would
25 appear to be more similar to the Als group A than Als group B proteins in terms of hydrophobic or electrostatic distribution.

Several proteins with adhesive function have been identified in *C. albicans*. Hwplp has been shown to mediate adherence to buccal epithelial cells by acting as a substrate for mammalian transglutaminase (5). *EAP1* was recently identified by
30 heterologous expression in *S. cerevisiae* and mediates adherence to polystyrene and renal epithelial cells *in vitro* (7). Of the eight members of the Als protein family, only Als1p

and Als5p have been studied from a functional perspective. Heterologous expression of Als1p has been shown to mediate binding to human vascular endothelial cells and epithelial cells, a finding that has been confirmed in *C. albicans* through gene disruption studies (Fu et al., Mol. Microbiol. 44:61-72 (2002), Fu et al., Infect. Immune. 66:1783-1786 (1998)). Heterologous expression of *ALS5* in *S. cerevisiae* confers adherence to collagen, fibronectin, bovine serum albumin, and laminin (Gaur et al., Infect. Immune. 65:5289-5297 (1997), Gaur et al., Infect. Immun. 67:6040-6047 (1999), Gaur et al., Cell Commun. Adhes. 9:45-57 (2002)). No large scale comparison of the substrate specificities of *C. albicans* adhesins has been performed. In this study, we compared the adhesive properties of a structurally diverse group of Als protein family members. Our data demonstrate that the Als proteins comprise a diverse family of surface proteins with an overlapping spectrum of specificities for adherence to a variety of human substrates (Fig. 8). Further, results from the present domain exchange experiments indicate that the N-terminal domains of Als proteins confer the specificity of their substrate adherence profiles.

In addition to mediating adherence, our data suggest that Als proteins also can function as invasins. Interestingly, whereas both Als1p and Als3p expressing *S. cerevisiae* demonstrated similar endothelial cell adherence, Als3p-expressing *S. cerevisiae* underwent internalization at a much higher rate. These results indicate that endocytosis is not simply an extension of adherence but rather a distinct process that can be influenced by the ligand-receptor interaction. It is likely that differences in N-terminal sequences in Als proteins mediate these distinct functions, as is the case with adherence.

The physicochemical properties of protein domains as distributed in three-dimensional space are crucial structural features governing receptor-ligand interactions (Eisenberg et al, J. Mol. Biol. 179:125-142 (1984), Waring et al., Protein Peptidew Lett. 3:177-184 (1996), Hancock et al., Lancet 349:418-422 (1997)). The Als proteins share conformational features characteristic of other adhesins and invasins of the immunoglobulin superfamily. However, individual Als proteins differed in their primary homolog, a finding consistent with the experimental data indicating that members of the Als family exhibit different substrate-binding profiles. Collectively, these patterns of Als

homologies indicate that, whereas Als protein members share a global similarity in structure and predicted fold, there exists structural differences among distinct Als proteins that are responsible for their differences in function.

The results described above relating to the Als family member structural determinations corroborate the homology modeling, which indicates that the N-terminal regions of Als1p are composed predominantly of anti-parallel β -sheet domains containing loop/coil structures, with lesser amounts of relatively unstructured regions. These features are indicative motifs of members of the immunoglobulin superfamily. These results show significant predictive correlation with circular dichroism studies of Als5p (Hoyer et al., Yeast 18:49-60 (2001)), indicating that the N-terminal domain of Als5p is characterized by a relative predominance of anti-parallel β -sheet and loop/coil regions. Thus, it is highly likely that all members of the Als protein family exhibit this overall structure. In particular, the structural results above are also consistent with the homology models that indicate that many of the HVRs correspond to the flexible loop/coil structures projecting from β -sheet domains in the N termini of distinct Als proteins. Collectively, these results indicate that these structures are integral to substrate-specific binding by Als proteins (Fig. 14). Consistent with the results above, analogous regions of mannose-binding lectin, α -agglutinin, and other members of the immunoglobulin superfamily appear to confer substrate binding specificity (Zhao et al., Hybrid Hybridomics 21:25-36 (2002), Wojciechowicz et al., Mol. Cell. Biol. 13:2554-2563 (1993)). Furthermore, mutations of these variable loop regions significantly alter substrate binding in these homologous proteins (Renz et al., J. Cell Biol. 125-1395-1406 (1994), Viney et al., J. Immunol. 157:2488-2497 (1996)).

The three-dimensional modeling results further indicate that N-terminal domains of individual Als proteins possess distinctive molecular signatures that relate to their adhesive profiles. These signatures incorporate parameters such as surface area, hydrophobicity, and electrostatic charge, yielding configurations that distinguish structural relationships among Als proteins. For example, Als proteins that bind to multiple substrates, such as the Als group A members (Als1p, Als3p, and Als5p), have similar predicted N-terminal profiles in terms of steric bulk, hydrophobic distribution, and electrostatic potential. Yet, even within this group, specific physicochemical distinctions exist that can govern functional differences within the group (Fig. 13). In contrast, Als

proteins with reduced adhesive capacity have surface features predicted to be distinct from the Als group A proteins in multiple physicochemical properties, including hydrophobicity and electrostatic potential. It is highly likely that the aggregate effects of differences in these structural features confer the specific functional properties of distinct
5 Als proteins.

Extensive genetic variability has been demonstrated within the ALS gene family. Sequence variation in specific ALS genes of different isolates of *C. albicans* has been observed (Zhang et al., Genome Res. 13:2005-2017 (2003), Hoyer et al., Yeast 18:49-60 (2001)), and not all members of the ALS family are present in all isolates. Even
10 significant sequence divergence between two different alleles in a single isolate have been found (Zhao et al., Microbiology 149:2947-2960 (2003), Zhang et al., Genome Res. 13:2005-2017 (2003)). This degree of genetic variability would suggest that these proteins may undergo rearrangement or mutation at a relatively high frequency. Such a mechanism would provide the organism with the ability to generate the high degree of
15 structural and functional diversity demonstrated in this study. Indirect support for this hypothesis is provided by a recent study of allelic variation of ALS7, which suggested both that this gene is both hypermutable and that these mutations are subject to selective pressure (Zhang et al., Genome Res. 13:2005-2017 (2003)).

Collectively, the above results indicate an analogy between antibodies and
20 Als proteins at both the structural and functional level. For example, the homology modeling underscores the similarities in structural configurations of these families, with hypervariability targeted to localized domains within an otherwise stable framework (e.g. HVRs of Als proteins and Fab regions in immunoglobulins). Further, as with antibodies, the genetic variability of the ALS gene family may provide the opportunity for *Candida*
25 to display a diverse array of proteins with a spectrum of specificity in adherence and invasion. The availability of such a group of related proteins is likely to improve the ability of the organism to colonize and invade different anatomical and physiological niches during infection.

Throughout this application various publications have been referenced
30 within parentheses. The disclosures of these publications in their entireties are hereby

incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains.

Although the invention has been described with reference to the disclosed embodiments, those skilled in the art will readily appreciate that the specific examples and studies detailed above are only illustrative of the invention. It should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

EXAMPLE VII

Vaccination with rAls1p-N improves survival during murine disseminated candidiasis by enhancing cell-mediated, not humoral, immunity

This example shows that immunizing BALB/c mice with the recombinant N-terminus of Als1p (rAls1p-N) improved survival during subsequent challenge with a lethal inoculum of *C. albicans*. The protective 20 µg dose of rAls1p-N significantly increased *Candida*-stimulation of Th1 splenocytes and increased *in vivo* delayed type hypersensitivity. In contrast, antibody titers did not correlate with protection. Finally, the vaccine was not protective in T cell-deficient mice but was protective in B cell-deficient mice. These data indicate that the mechanism of action of the rAls1p-N vaccine is stimulation of cell mediated, rather than humoral, immunity against *C. albicans*.

The *C. albicans* used in the study was SC5314, a well-characterized clinical isolate that is highly virulent in animal models (Spellberg et al., Infect Immun. 71:5756-5764 (2003)) was supplied by W. Fonzi (Georgetown University). The organism was serially passaged three times in yeast peptone dextrose broth (Difco) prior to infection.

The mice strains used in the study were female BALB/c mice obtained from the National Cancer Institute (Bethesda, MD). To explore the impact of age on vaccine efficacy, both juvenile mice (8-10 weeks) and retired breeders (≥ 6 months) were utilized. Female B cell-deficient mice bearing a homozygous deletion of the igh loci (C.129B6-IgH-Jhdtm1Dhu), T cell-deficient nude mice (C.Cg/AnBomTac-Foxn1nuN20), and congenic wild-type BALB/c littermates were obtained from Taconic Farms (Germantown, NY). Mice were housed in filtered cages with irradiated food and

autoclaved water ad libitum. For survival experiments, mice were immunized with varying doses of antigen (see below) and subsequently infected via the tail vein with the appropriate inoculum of *C. albicans* SC5314 blastospores, or PBS (Irvine Scientific, Irvine, CA) control. Results of replicate survival studies were combined if the individual
5 datasets demonstrated no statistical heterogeneity (see below). All procedures involving mice were approved by the institutional animal use and care committee, following the National Institutes of Health guidelines for animal housing and care.

The rAls1p-N immunization procedures described below were performed as follows. Briefly, rAls1p-N (amino acids 17 to 432 of Als1p) was produced in *S. cerevisiae* and purified by gel filtration and Ni-NTA matrix affinity purification (Fu et al.,
10 Molec. Microbiol. 44:61-72 (2002)). The amount of protein was quantified by modified Lowry assay. A high degree of purity ($\approx 90\%$) was confirmed by SDS-polyacrylamide gel electrophoresis as well as circular dichroism and FTIR, as described above. Mice were immunized by intraperitoneal (ip) injection of rAls1p-N mixed with complete Freund's
15 adjuvant (CFA, Sigma-Aldrich) at day 0, boosted with another dose of the antigen with incomplete Freund's adjuvant (IFA, Sigma-Aldrich) at day 21, and infected two weeks following the boost.

Resultant Antibody titers were determined by ELISA in 96 well plates. Briefly, wells were coated with 100 μ l per well of 5 μ g/ml rAls1p-N in PBS. Mouse sera
20 were incubated for 1 h at room temperature following a blocking step with tris buffer saline (TBS) (0.01 M TrisHCl, pH 7.4, 0.15 M NaCl) containing 3% bovine serum albumin. The wells were washed 3 times with TBS containing 0.05% Tween 20, followed by another 3 washes with TBS. Goat anti-mouse secondary antibody conjugated with horseradish peroxidase (Sigma) was added at a final dilution of 1:5000
25 and the plate was further incubated for 1 h at room temperature. Wells were washed with TBS and incubated with substrate containing 0.1 M citrate buffer (pH 5.0), 50 mg/ml of o-phenylenediamine (Sigma), and 10 μ l of 30% H_2O_2 . The color was allowed to develop for 30 min after which the reaction was terminated by adding 10% H_2SO_4 and the optical density (OD) was determined at 490 nm in a microtiter plate reader. Negative control
30 wells received only diluent, and background absorbance was subtracted from the test

wells to obtain final OD readings. The ELISA titer was taken as the reciprocal of the last serum dilution that gave a positive OD reading (i.e. > mean OD of negative control samples + 2 standard deviation).

Other methods described below were performed as follows. Briefly, C. albicans-induced cytokine profiles were performed to determine the effect of the rAls1p-N vaccine on cell-mediated immunity and *in vivo* cytokine profiles. Mice were immunized as described above. Two weeks after the final boost, splenocytes were harvested and cultured in complete media at a density of 4×10^6 cells/ml as previously described (Spellberg et al., Infect. Immun. 71:5756-5764 (2003)). To stimulate cytokine production, splenocytes were co-cultured with heat-killed C. albicans SC5314 germ tubes. We used heat-killed *C. albicans* in lieu of rAls1p-N to stimulate the splenocytes to mimic the *in vivo* situation during infection. The *C. albicans* cells were pre-germinated in RPMI-1640 with glutamine (Gibco BRL) for 90 minutes to induce expression of Als1p (Fu et al., Molec. Microbiol. 44:61-72 (2002)). The resulting *C. albicans* germ tubes were heat-killed by incubation for 90 minutes at 60°C (Ibrahim et al., Infect. Immun. 63:4368-74 (1995)). The heat-killed fungi were added to the splenocyte cultures at a density of 2×10^7 pseudohyphae/ml (ratio of five fungi to one leukocyte). After 48 h, splenocytes were profiled for Th1 (CD4+IFN- γ +IL-4-), Th2 (CD4+IFN- γ -IL-4+), or CD4+IL-10+ frequencies by intracellular cytokine detection and flow cytometry, as previously described (Spellberg et al., Infect. Immun. 71:5756-5764 (2003)). Three-color flow cytometry was performed on a Becton-Dickinson FACScan instrument calibrated with CaliBRITE beads (Becton Dickinson, San Jose, CA) using FACSCComp software as per the manufacturer's recommendations. During data acquisition, CD4+ lymphocytes were gated by concatenation of forward and side scatter, and FITC-anti-CD4 antibody fluorescence properties. Data for each sample were acquired until 10,000 CD4+ lymphocytes were analyzed. Results are presented as the median $\pm$ 25th and 75th quartiles of the percentage of all gated lymphocytes that were Th1 or Th2 cells.

Footpad swelling was determined by the method of Oomura et al (41). Briefly, mice were immunized with the appropriate dose of rAls1p-N or CFA alone as described above. Two weeks following the boost, baseline footpad sizes of immunized mice were measured using an electronic digital caliper. Fifty μ g of rAls1p-N in 25 μ l of

PBS was injected into the right footpads, and PBS alone injected into the left footpads of the immunized mice. Twenty-four hours later the footpads were again measured.

Antigen-specific footpad swelling was calculated as: (right footpad thickness at 24 h – right footpad thickness at baseline) – (left footpad thickness at 24 h – left footpad thickness at baseline).

The non-parametric Log Rank test was utilized to determine differences in survival times of the mice. Titers of antibody, frequency of Th1 or Th2 lymphocytes, and footpad swelling were compared by the Steel test for non-parametric multiple comparisons (Rhyne et al., Biometrics 23:539-49 (1967) or the Mann Whitney U test for unpaired comparisons, as appropriate. Correlations were calculated with the Spearman Rank sum test. To determine if heterogeneity existed in replicate survival studies, the Kolmogorov-Smirnov test was utilized. P values < 0.05 were considered significant.

To determine the most effective dose of the rAls1p-N immunogen, a 10^7 -fold dose range was evaluated (20 pg to 200 µg per mouse). Female retired breeder BALB/c mice were immunized with rAls1p-N plus adjuvant (CFA/IFA) or adjuvant alone. Immunized mice were bled 2 weeks after boosting to determine anti-rAls1p-N antibody titers (see below). The mice were subsequently infected with a lethal inoculum of *C. albicans* (2×10^5 blastospores). The survival data from repeat experiments were combined since the individual experiments demonstrated no statistical heterogeneity ($p > 0.05$ by Kolmogorov-Smirnov test). The 20 µg dose of rAls1p-N resulted in long-term survival of 25% of the infected mice, and a significant increase in overall survival compared to adjuvant alone ($p = 0.044$ by Log Rank test, Figure 1). Neither 10-fold higher (Figure 15) nor lower (data not shown) doses significantly increased survival compared to adjuvant alone. These results indicate that an intermediate dose of the rAls1p-N vaccine induces protection against murine disseminated candidiasis.

The above findings established a protective dose for the rAls1p-N vaccine. Next the efficacy of the vaccine was evaluated in a more rapidly lethal model of mice infected with 10^6 blastospores (median survival 3 vs. 11 days for 10^6 vs. 2×10^5 inocula in unvaccinated mice, respectively). Again the data from repeat studies were combined as the results of the individual experiments demonstrated no statistical heterogeneity ($p > 0.05$ by Kolmogorov-Smirnov test). When administered as a 20 µg dose + CFA to Balb/c

mice infected with 10^6 *C. albicans* blastospores, the rAls1p-N vaccine more than doubled the median survival and resulted in a significant increase in overall survival versus unvaccinated controls ($p = 0.001$ by Log Rank test, Fig. 16A). To determine if the age of the mice influenced their response to the rAls1p-N vaccine, we tested it in juvenile mice.

5 A similar survival benefit was found when juvenile mice were vaccinated and infected with the same high inoculum ($p = 0.02$ by Log Rank test, Fig. 16B).

Although the 200 μ g dose of rAls1p-N resulted in inferior protection compared to the 20 μ g dose, only the 200 μ g dose of antigen induced a significant increase in serum anti-Als1p antibody titers ($p \leq 0.005$ for 200 μ g dose vs. all other
10 groups, Figure 17). No significant increases in anti-Als1p antibody titers were detected at the intermediate, protective antigen dose ($p = 0.1$ for 20 μ g vs. adjuvant). When the serum anti-Als1p antibody titers of individual mice were plotted against the survival time of each mouse, no correlation between antibody titer and survival was found ($R^2 = 0.03$, $p > 0.05$ by the Spearman rank sum test). Indeed, mice immunized with the highest dose of
15 antigen (200 μ g) had anti-rAls1p-N antibody titers in excess of 1:100,000, but had survival durations no different from mice immunized with lower doses of antigen whose titers were at the lower limit of detection ($\sim 1:100$). These results indicate that protection induced by the rAls1p-N vaccine does not appear to correlate with antibody titers.

Since humoral immunity did not correlate with rAls1p-N-induced
20 protection, we examined the cell-mediated immune response induced by protective and non-protective doses of rAls1p-N. Mice were immunized with 0.2, 20, or 200 μ g of rAls1p-N, or adjuvant alone, as above. Two weeks after the boost, splenocytes were harvested and cultured in the presence of heat-killed, pre-germinated *C. albicans*, which are known to express Als1p (Fu et al., Molec. Microbiol. 44:61-72 (2002)). Following
25 h of culture, splenocytes were harvested for intracellular cytokine detection by flow cytometry. Only the lymphocytes from mice immunized with the protective 20 μ g dose of antigen developed a significantly increased frequency of Th1 cells compared to mice given adjuvant alone ($p = 0.03$, Fig. 18). No significant differences in Th2 frequency (Fig. 18) or in the frequency of IL-10⁺ T lymphocytes (data not shown) were detected
30 between mice immunized with adjuvant or any of the doses of antigen.

To confirm that Type 1 immunity was stimulated by r-Als1p-N *in vivo*, delayed type hypersensitivity was tested by footpad swelling. Only mice vaccinated with the protective 20 µg dose of rAls1p-N developed a significantly increased delayed type hypersensitivity reaction compared to control, and this response was also significantly
5 greater than that induced by the non-protective 0.2 and 200 µg doses (Fig. 19, $p < 0.05$ for all comparisons versus 20 µg dose, by the non-parametric Steel test). Collectively, these results indicate that a protective dose of the rAls1p-N antigen induced significant Th1 polarization and delayed type hypersensitivity reaction.

To define the role of antibody and T-cells in vaccine-mediated protection,
10 B cell-deficient, T-cell deficient nude, or congenic BALB/c wild-type control mice were immunized with 20 µg of rAls1p-N plus adjuvant or adjuvant alone, and infected with a lethal inoculum (8×10^5 blastospores) of *C. albicans*. B cell-deficient mice trended to being more resistant to infection, whereas T cell-deficient mice were more susceptible, than were wild-type control mice given adjuvant alone ($p = 0.065$ and 0.01 for B cell-
15 deficient and T cell-deficient mice versus wild-type adjuvant-treated, respectively, Fig. 20). Finally, the rAls1p-N vaccine maintained its efficacy in B cell-deficient mice ($p = 0.04$ for rAls1p-N vaccinated versus adjuvant alone, Fig. 6) but was ineffective in T cell-deficient mice ($p = 0.4$ for rAls1p-N vaccinated versus adjuvant alone, Fig. 20). These results indicate that the Als1p vaccine is effective in B cell-deficient mice but not in T-
20 cell deficient nude mice.

Described above are the results showing that immunization with the N-terminus of this protein improved survival of both juvenile and mature BALB/c mice during subsequent hematogenously disseminated candidiasis. In particular, an intermediate dose of rAls1p-N (20 µg) provided superior protection compared to both
25 lower doses and a higher dose (200 µg). Nevertheless, the non-protective 200 µg dose of rAls1p-N was immunogenic, as it induced 100-fold higher titers of antibody than did the protective 20 µg dose.

The inverted U-shaped dose-response efficacy curve, with lower protection at the highest dose of rAls1p-N, is reminiscent of the classical studies of Parish
30 *et al.*, who first described the inverse relationship between the induction of humoral and

cell-mediated immunity by a given dose of antigen. In the context of Parish's seminal data, an inverted U-shaped dose-response efficacy curve could be explained if: 1) vaccine efficacy depended on cell-mediated immunity and, 2) intermediate doses of rAls1p-N stimulated superior cell-mediated immunity compared to the high, antibody-stimulating dose. We therefore hypothesized that the inverted U-shaped dose response efficacy curve seen with the rAls1p-N vaccine was due to superior induction of cell-mediated immunity by the protective, intermediate doses of antigen.

To test this hypothesis, the ability of high, intermediate, and low doses of antigen to stimulate Th1 cells and delayed-type hypersensitivity were determined. To stimulate cytokine-production from splenocytes, we specifically activated the cells by exposure to heat-killed *C. albicans*, instead of rAls1p-N, to mimic the *in vivo* situation during infection. Only the protective 20 µg dose significantly increased the frequency of *C. albicans*-stimulated, splenic Th1 lymphocytes. The frequency of Th1 cells seen in *ex vivo* *C. albicans*-stimulated splenocytes was similar to that detected *in vivo* during disseminated candidiasis in mice (59), underscoring the relevance of this approach.

To determine if the detected *ex vivo* Th1 cells were of functional significance *in vivo*, we compared the delayed type hypersensitivity induced by different doses of rAls1p-N immunization. Concordant with the frequency of Th1 cells, only the protective 20 µg dose of rAls1p-N stimulated a significant *in vivo* delayed type hypersensitivity reaction. These results are consistent with the hypothesis that vaccine-induced protection was due to induction of Type 1, cell mediated immunity. Surprisingly, despite induction of markedly elevated antibody titers by the 200 µg dose of rAls1p-N, we did not find an increase in splenic Th2 lymphocytes in mice vaccinated with this dose. One possible explanation is that Th2 cells were activated in peripheral lymph nodes rather than the spleen. Alternatively, other T cell populations (e.g. NKT cells) may have been responsible for inducing the high antibody titers seen in response to the 200 µg dose of rAls1p-N.

The lack of correlation between antibody titer and protection did not completely exclude a role of antibodies in mediating vaccine-induced protection. For example, ELISA titers are the result of enumeration of antibodies with a variety of specificities and affinities. Therefore, the possibility that small subsets of antibodies were

generated that did participate in vaccine-mediated protection could not be excluded by measuring antibody titer. To confirm the role of cell-mediated and not humoral immunity in rAls1p-N vaccine-mediated protection, we tested the efficacy of the vaccine in B cell- and T cell-deficient mice. B cell-deficient mice trended to being more resistant to disseminated candidiasis than wild-type controls, and the efficacy of the vaccine was not abrogated in B cell-deficient mice. In contrast, T cell-deficient mice were more susceptible to disseminated candidiasis than were wild-type controls, and the efficacy of the vaccine was lost in T cell-deficient mice. Our findings therefore confirm that the efficacy of the rAls1p-N vaccine is dependent of induction of T-cell mediated, and not primarily humoral, immunity. As well, because B cell-deficient mice were not more susceptible to disseminated candidiasis than congenic wild type littermates, antibody is not a dominant effector against disseminated candidiasis in this model.

In sum, we report that the novel rAls1p-N vaccine mediates protection against experimental disseminated candidiasis by inducing cell-mediated rather than humoral immunity. Enhancement of the modest protective effect of the rAls1p-N vaccine may therefore be accomplished with additional priming of cell-mediated immunity using optimized adjuvants and/or cytokines, or an alternate route of immunization. Indeed, in our ongoing studies we have already found a marked increase in efficacy by administering rAls1p-N subcutaneously as compared to intraperitoneally.

20

EXAMPLE VIII

The anti-*Candida albicans* rAls1p-N vaccine reduces fungal burden and improves survival in both immunocompetent and immunocompromised mice

This example describes enhancement of the efficacy of the rAls1p-N vaccine described in example VII when administered by a subcutaneous (SQ) route in both immunocompetent and immunocompromised mice. Initially, the efficacy of the rAls1p-N vaccine in immunocompetent mice. rAls1p-N, encompassing amino acids 19-433 of the full length protein, was produced in *S. cerevisiae* and purified as described above. Control preparation was similarly purified from *S. cerevisiae* transformed with an empty plasmid. BALB/c retired breeder mice (25-30 g) were immunized by SQ injection of rAls1p-N (20 µg) or control preparation mixed with Complete Freund's Adjuvant (CFA) at day 0, followed by a booster dose in Incomplete Freund's Adjuvant (IFA) at day

21. Two weeks following the boost, the immunogenicity of the vaccine was confirmed by evaluating the intensity of the footpad swelling reaction as a marker of delayed type hypersensitivity (DTH), as previously described. Vaccinated mice had marked increases in rAls1p-N specific DTH (Fig. 21).

5 The efficacy of the rAls1p-N vaccine was evaluated by determining the impact of rAls1p-N vaccination on survival in infected BALB/c mice (Fig. 22A). Vaccinated or control mice were infected via the tail-vein with rapidly lethal inocula (2.5×10^5 blastospores) of *C. albicans*. We have previously shown that mice infected such inocula die of overwhelming septic shock (Spellberg et al., *J. Infect. Dis.* In press (2005)).
10 Vaccination markedly prolonged time to death ($p < 0.05$ for both inocula by Log Rank test) and improved 30 day survival (50-57% vs. 0%, $p < 0.05$ for both inocula by Fisher's Exact test).

 The impact of vaccination on tissue fungal burden during hematogenously disseminated candidiasis was then determined. Fourteen days following the boost,
15 vaccinated and control BALB/c mice were infected with via the tail-vein with 5×10^5 blastospores of *C. albicans* SC5314. Six days following infection, prior to onset of the first deaths in the control arm, kidneys were harvested, homogenized, and quantitatively cultured in Sabouraud dextrose agar (Difco) (18). SQ vaccination with rAls1p-N resulted in a median 1.5 log CFU/g decrease in kidney fungal burden compared to control ($p =$
20 0.01 by Wilcoxon Rank Sum test, Fig. 22B).

 The efficacy of the rAls1p-N vaccine also was assessed in immunocompromised mice. Having demonstrated efficacy in immunocompetent mice, the potential for the rAls1p-N vaccine to induce immunity in and protect neutropenic mice from disseminated candidiasis also was evaluated. Vaccinated BALB/c mice were
25 made neutropenic by administration of cyclophosphamide (200 mg/kg ip on day -2, and 100 mg/kg ip on day +9 relative to infection, resulting in approximately 12 days of neutropenia, as described (Sheppard et al., *Antimicrob. Agents. Chemother.* 48:1908-11 (2004)). Footpad swelling reaction was performed 2 days after the first dose of cyclophosphamide. Vaccinated neutropenic mice developed DTH reactions of similar
30 magnitude to immunocompetent mice (Fig. 23A vs. 1, experiments performed in parallel). In neutropenic mice infected via the tail-vein with 2.5×10^4 blastospores of *C.*

albicans, vaccination also resulted in significant improvements in time to death ($p = 0.007$ by Log Rank test vs. Control), median survival time (> 21 vs 12 d, $p = 0.008$ by Wilcoxon Rank Sum Test), and overall survival (88% vs. 38% , $p = 0.005$ by Fisher's Exact test) (Fig. 23B).

5 To determine the efficacy of rAls1p-N vaccination in mucosal infection, the vaccine was tested in a murine oropharyngeal candidiasis (OPC) model (Kamai et al., Infect. Immun. 70:5256-8 (2002) and Kamai et al., Antimicrob. Agents Chemother. 45:3195-97 (2001)). Vaccinated mice were treated with cortisone acetate (225 mg/kg SQ on days -1, 1, and 3 relative to infection) and infected sublingually as described. Tongues
10 were excised on day 5 post-infection. Because colony forming units of homogenized tongues cannot distinguish between invasive infection and surface-adherent colonization, we evaluated extent of invasion by histopathology. A blinded observer (BJS) scored each section by scanning along the entire length of the tongues and quantifying the severity of fungal lesions per 40x high-powered field ($0 =$ no lesion, $1^+ =$ mild mucosal
15 inflammation, $2^+ =$ significant inflammation restricted to the epithelium, $3^+ =$ inflammation extending through the entire epithelial layer, $4^+ =$ inflammation extending into the subepithelium). To avoid sampling bias, two sections of each tongue, separated by at least five intervening tissue sections, were scored. All control mice developed marked fungal invasion of their tongues in numerous locations, while only two vaccinated
20 mouse developed any tongue lesions. In total, the median number (75^{th} , 25^{th} quartile) of lesions per tongue in control mice was 6.5 (8, 5.75) as compared to 1 (2.5, 0) for vaccinated mice ($p = 0.03$ by Wilcoxon Rank Sum test). Semi-quantitative evaluation of the severity of infection demonstrated a significant reduction in vaccinated mice compared to controls (Fig. 24, $p = 0.03$ by Wilcoxon Rank Sum test).

25 To determine the efficacy of rAls1p-N or rAls3-p-N vaccination in mucosal infection, these two vaccines in a murine model of vaginal colonization (Clemons et al., Infect. Immun. 72: 4878-80 (2004); Fidel. Int Rev Immunol. 21: 515-48 (2002) and Wozniak et al., Infect Immun. 70: 5790-9 (2002)). Vaccinated mice were treated with estrogen ($30 \mu\text{g}$, given SQ) on day -3 relative to infection and then
30 challenged in the vagina with $10 \mu\text{l}$ phosphate buffered saline containing 10^6 blastospores of *C. albicans*. Vaginas were excised on day 3 post-inoculation, homogenized and serial

dilutions were plated on YPD plates. Colony forming units (CFU) were enumerated 24-48 h following incubation of plates at 30-35°C. Vaginas collected from mice vaccinated with rAls3p-N but not those collected from mice vaccinated with rAls1p-N had lower CFU than vaginas collected from control mice (i.e mice vaccinated with CFA alone) (Fig 25, p=0.01 by Wilcoxon Rank Sum test).

In light of the increasing incidence of candidemia and its continuing high mortality rate, development of a vaccine against *Candida spp.* is of great importance. The results described above show that SQ vaccination with rAls1p-N resulted in marked improvement in survival and significant reductions in fungal burden during otherwise rapidly fatal hematogenously disseminated candidiasis in both immunocompetent and immunocompromised mice. Of interest are the kidney fungal burden results from individual vaccinated mice, demonstrating that approximately half the mice had kidney fungal burdens under 5 log CFU/g. We have previously found that the threshold of kidney fungal burden indicative of a fatal infection is 5 log CFU/g; mice with kidney fungal burdens above this level typically die from infection, whereas mice with kidney fungal burdens below this burden survive the infection (Spellberg et al., *J. Infect. Dis.* In press (2005) and (Spellberg et al., *Infect. Immun.* 71:5756-5764 (2003)). Therefore, breakthrough deaths in the vaccinated group likely reflect high fungal burden in spite of vaccination. The mouse to mouse variations in tissue fungal burden may reflect the complexities of host-pathogen interactions and/or variable vaccine responsiveness.

In summary, the rAls1p-N vaccine can be used for the treatment, reduction in severity and/or prevention of increasingly common and highly lethal disseminated candidiasis. The vaccine is efficacious in immunocompetent mice, and efficacy is retained even in neutropenic and corticosteroid-treated hosts. Finally, the vaccine can protect against mucocutaneous candidiasis including vaginal and oropharyngeal candidiasis

EXAMPLE IX

Effectiveness of ALS Vaccines Against *S. aureus* Infections

This Example shows that Als proteins from *C. albicans* improves survival of animal models infected with *S. aureus*.

Als adhesins of *C. albicans* were identified to be significantly homologous to adhesins on *S. aureus*. This characteristic was used to design and implement an effective vaccine against *S. aureus* using Als adhesins. Briefly, the *C. albicans* ALS family is comprised of at least 9 genes (Hoyer et al., *Genetics* 157:1555-67 (2001); Hoyer LL., *Trends Microbiol.* 9:176-80 (2001)). As described previously, Als proteins function as adhesins to biologically relevant substrates (Fu et al., *Molec. Microbiol.* 44:61-72 (2002); Gaur and Klotz, *Infect. Immun.* 65:5289-94 (1997); Zhao et al., *Microbiology* 150:2415-28 (2004); Oh et al., *Microbiology* 151:673-81 (2005); Zhao et al. *Microbiology* 151:1619-30 (2005)); Hoyer et al., *Mol. Microbiol.* 15:39-54 (1995)). In particular, the N-termini of Als1p and Als3p are significantly homologous to surface proteins expressed by pathogenic *S. aureus*, including collagen binding protein and clumping factor (Table IV; Sheppard et al. *J. Biol. Chem.* 279:30480-89 (2004)).

Table IV. Homology of Als proteins to various pathogenic adhesins and invasions		
Protein	Homologue 1	Homologue 2
Als1p	Collagen binding protein of <i>S. aureus</i> : $\geq 95\%$ homology	Clumping factor of <i>S. aureus</i> : $\geq 90\%$ homology
Als3p	Collagen binding protein of <i>S. aureus</i> : $\geq 95\%$ homology	Clumping factor of <i>S. aureus</i> : $\geq 80\%$ homology
Als5p	Invasin/integrin-binding protein <i>Y. pseudotuberculosis</i>	Surface layer protein <i>M. mazei</i>

The homology calculation provided above in Table IV takes into account both features of sequence alignment and 3-dimensional surface structure. Homology of Als1p was calculated to be greater than 95% or 90% compared to collagen binding protein or clumping factor of *S. aureus* ($r^2 \geq 90\%$; Sheppard et al., *supra*). Similarly, homology of Als3p was calculated to be greater than 95% or 80% compared to collagen binding protein or clumping factor of *S. aureus* ($r^2 \geq 90\%$).

To corroborate the above findings, homology and threading methods were employed to model structure-function congruence between Als1p and *S. aureus* clumping factor A (ClfA – PDB code: c1n67A). These methods assessed specific homologies in primary structure, 3-D conformation and pattern analyses were conducted to seek analogous functional motifs. For example, BLASTP, PROSITE and JALVIEW methods were employed to analyze similarities and differences in ALS versus ClfA primary

sequences (Yount et al. Antimicrob. Agents Chemother. 48:4395-4404 (2004) and Yount and Yeaman. Proc. Natl. Acad. Sci. USA 101:7363-7368 (2004)). Internet-based applications including 3-D PSSM were then used to prioritize potential ALS homologues for further analysis (Sheppard, et al. J. Biol. Chem. 279:30480-30489 (2004)). Along with resulting data, the PHYRE application (Kelley, L., R. Bennett-Lovsey, A. Herbert, and K. Fleming; website is as follows: <http://www.sbg.bio.ic.ac.uk/~phyre/>) was used to conduct topology mapping and to identify 3-dimensional motifs shared by proteins with greatest structural or functional homology to selected ALS proteins for the purpose of identifying putative shared functional motifs. The above methods are widely available in public domain and used in a variety of proteomic and structural biology applications. Based on the above homology and threading method results a consensus of functional site homologies between Als1p and ClfA was generated and mapped to specific residues of the Als1p model constructed on ClfA. Several particular findings emanated from these modeling analyses as set forth below.

First, significant homology was identified between the N-terminal regions of Als1p and ClfA in secondary structure and amino acid conservation, particularly in the region encompassed by amino acids 30 – 300 (i.e. the N-termini of both proteins).

Second, consensus mapping of homologous functional sites based on established ClfA adhesin determinants converged on a specific topological motif in Als1p. This topological motif is shown in Figure 26 as a cleft formed by the inflection of adjacent facets of two β -sheet domains.

Third, consistent with primary structure homology, the predicted functional cleft motif in Als1p maps to specific residues originating from hypervariable regions in the N-terminal region encompassing amino acid residues 30 – 300.

These results provided a structural basis for congruent biological functions, as well as immunological responses to Als1p and ClfA. These results also further corroborate our overall model of Als1p structure-activity, and further facilitate targeted approaches to mutational analyses and epitope mapping. Finally, these results indicate that Als1p and ClfA are adhesins of analogous structure and function present on diverse microbial pathogens.

A monoclonal antibody against *S. aureus* also was identified that may reduce infections caused by *C. albicans*. As with the above structural findings, this characteristic also was used to design and implement an effective vaccine against *S. aureus* using Als adhesins.

5 Briefly, a humanized anti-staphylococcal monoclonal antibody (Aurexis[®]) that is known to recognize surface adhesins on *S. aureus* is currently in clinical trials. This monoclonal antibody also cross reacts with Als family members. Favorable results of a phase II clinical trial of Aurexis[®] for the treatment of *staphylococcal* bloodstream infections have been reported (Inhibitex Inc., 2005; accessed September 19, 2005, at
10 [http://phx.corporate-ir.net/phoenix.zhtml?c=176944&p=irol-](http://phx.corporate-ir.net/phoenix.zhtml?c=176944&p=irol-newsArticle&ID=707322&highlight=)
[newsArticle&ID=707322&highlight=](http://phx.corporate-ir.net/phoenix.zhtml?c=176944&p=irol-newsArticle&ID=707322&highlight=)). Briefly, in this report, patients with known *S. aureus* in the blood were administered the Aurexis[®] antibody as treatment for active infection (i.e., this is not an active vaccine strategy or a prophylaxis study). Nine patients receiving placebo experienced breakthrough bloodstream infections caused by *Candida*,
15 while only three patients in the Aurexis[®] arm experienced *Candida* bloodstream infections. Recognizing the decrease in *Candida* blood infection for those patients treated with an antibody to *S. aureus* combined with the above homology and structural findings indicate that immunogenic epitopes are shared between *Candida* and *S. aureus* and that these immunogenic epitopes can be targeted for therapeutic benefit using immune
20 responses, antibodies or effector mechanisms raised against one species for treatment of the other species. Therefore, the above data together provide for immune responses to surface adhesins on *S. aureus* to cross react with *Candida* spp.

Following the above strategy, exemplary Als adhesin vaccines were designed and shown to improve survival of mice infected with *S. aureus*. The exemplary
25 Als adhesins used to vaccinate were rAls1p-N or rAls3p-N, which were produced and used as described above. Briefly, to determine if these Als vaccines against *Candida*, rAls1p-N and rAls3p-N, can mediate cross-species protection against *S. aureus*, female Balb/c mice were vaccinated with the previously described regimen (Complete Freund's Adjuvant + 20 µg of rAls1p-N or rAls3p-N on day 0, followed by a booster dose in
30 Incomplete Freund's Adjuvant at 3 weeks, both administered subcutaneously). Two weeks following vaccination, mice were infected via the tail-vein with a lethal dose of *S.*

aureus strain 67-0, which is methicillin-resistant and known to be virulent in animal models. The results showing mice survival are shown in Figure 26. As indicated, both the rAls1p-N and rAls3p-N vaccines mediated improved long-term survival in these infected mice (Figure 27). Additionally, the mechanism of protection likely to be an enhancement of Th1 rather than Th2 since no correlation between Ab titers and survival of mice vaccinated with either rAls1p-N or rAls3p-N was observed (Figure 28).

EXAMPLE X

The Anti-Candida rAls1p-N Vaccine Mediates a Broad Range of Protection Against Disseminated Candidiasis

This Example show that the rAls1p-N vaccine protects outbred mice from disseminated candidiasis, and protects Balb/c mice against other virulent strains of *C. albicans* and non-*albicans* *Candida*.

The current studies were performed to illustrate the breadth of protection induced by rAls1p-N by specifically evaluating its efficacy in outbred mice, in combination with a second adjuvant other than Freund's adjuvant, against other strains of *C. albicans*, and against non-*albicans* species of *Candida*.

Vaccination with rAls1p-N protected outbred mice from disseminated candidiasis. Briefly, outbred CD1 mice were obtained from the National Cancer Institute (Bethesda, MD). All procedures involving mice were approved by the institutional animal use and care committee, following the National Institutes of Health guidelines for animal housing and care. The mice were vaccinated with rAls1p-N + Freund's adjuvant as previously described above and in, for example, Ibrahim et al., *Infect. Immun.* 73:999-1005 (2005); Spellberg et al., *Infect. Immun.* 73:6191-93 (2005). rAls1p-N (amino acids 17 to 432 of Als1p) was produced in *S. cerevisiae* and purified by gel filtration and Ni-NTA matrix affinity purification. A high degree of purity ($\approx 90\%$) was confirmed by SDS-polyacrylamide gel electrophoresis as well as circular dichroism and FTIR, as described above and in, for example, Sheppard et al., *J Biol Chem* 279:30480-89 (2004). Mice were immunized by SQ injection of rAls1p-N (20 μ g) mixed with Complete Freund's Adjuvant (CFA; Sigma-Aldrich, St. Louis, MO) at day 0, followed by a booster dose in Incomplete Freund's Adjuvant (IFA; Sigma-Aldrich) at day 21. Control mice

were immunized with CFA/IFA alone. Fourteen days following the boost, immunized mice were infected via the tail-vein with *C. albicans* SC5314, as we have described previously Ibrahim et al., (2005) *supra*; and Spellberg et al. (2005), *supra*. Similar to our previous findings in Balb/c mice, the rAls1p-N vaccine markedly improved the survival
5 of infected CD1 mice (Figure 29A).

Because Freund's adjuvant is considered to be too toxic for use in humans, we performed a dose response of rAls1p-N vaccine in alum (2% Alhydrogel, Brenntag Biosector, Frederikssund, Denmark), the only vaccine adjuvant currently approved by the US Food and Drug Administration (FDA) for use in humans. Vaccination with alum was
10 performed on an identical schedule as Freund's adjuvant, with immunization on day 1, boost on day 21, and infection 2 weeks later. We found that higher doses of rAls1p-N combined with alum resulted in significant improvements in survival of mice with disseminated candidiasis (Figure 29B). There are also appeared to be a dose response relationship, with trends to improved survival at higher doses of rAls1p-N when
15 combined with alum.

The rAls1p-N vaccine also was shown to improve the survival of Balb/c mice infected with several strains of *C. albicans*. Particularly useful vaccines utilize an immunogen that can prime the immune system to recognize multiple strains of the target pathogen. By DNA sequence analysis, we found that the predicted amino acid sequence
20 of the N-terminal region of Als1p was 99.9% conserved amongst a diverse group of clinical *C. albicans* isolates from bloodstream (5 strains), urine (5 strains) and oropharyngeal (10 strains) infections (data not shown). These results indicated that the rAls1p-N vaccine can be effective against a broad array of *C. albicans* strains. To confirm the breadth of protection of the rAls1p-N vaccine against other strains of *C.*
25 *albicans*, mice were vaccinated with rAls1p-N + Freund's adjuvant as above, and infected with one of several clinical isolates of *C. albicans* (Ibrahim et al., *Infect Immun* 63:1993-98 (1995)). As shown in Figure 30, the rAls1p-N vaccine significantly improved the survival of mice infected with each of these strains.

The rAls1p-N vaccine also was shown to reduce tissue fungal burden in
30 mice infected with several non-*albicans* species of *Candida*. Briefly, the *ALS* gene family is present in other *Candida* species, including *C. dubliniensis* and *C. tropicalis*

(Hoyer et al., *Genetics* 157:1555-67 (2001)). Similarly, an adhesin analogous to Als family members has been described in *C. glabrata* (Cormack et al., *Science* 285:578-82 (1999); Frieman et al., *Mol Microbiol* 46:479-92 (2002)). To confirm the efficacy of the rAls1p-N against non-*albicans* species, Balb/c mice were vaccinated with rAls1p-N + Freund's adjuvant as above, and infected via the tail-vein with *C. glabrata* 31028 (a clinical bloodstream isolate from the microbiology laboratory at Harbor-UCLA Medical Center), *C. krusei* 91-1159, (generously provided by Michael Rinaldi, San Antonio, Texas), *C. parapsilosis* 22019 (clinical bloodstream isolate from Harbor-UCLA Medical Center), or *C. tropicalis*.4243 (clinical bloodstream isolate from Harbor-UCLA Medical Center). As shown in Figure 31, the rAls1p-N vaccine reduced the kidney fungal burden of mice infected with each of these species.

In summary, the rAls1p-N vaccine is able to prevent and/or reduce the severity of an increasingly common and highly lethal disseminated candidiasis. The vaccine is efficacious in both inbred and outbred mice, when mixed with alum as an adjuvant, against multiple strains of *C. albicans*, and against several non-*albicans* species of *Candida*. These results further corroborate that the ALS vaccines of the invention are effective against a wide variety of candidal and other infections.

EXAMPLE XI

The Anti-Candida rAls3p-N Vaccine is Equally Effective as rAls1p-N Against Disseminated and More Efficacious Against Mucosal Candidiasis

This Example compares the efficacy of rAls3p-N to rAls1p-N vaccines in murine models of hematogenously disseminated, oropharyngeal, and vaginal candidiasis.

Of the *ALS* family members, the *ALS1* and *ALS3* genes encode adhesins with the broadest array of substrate affinity. When compared to one another, Als1p mediated greater adherence to endothelial cells and gelatin, but inferior adherence to epithelial cells (Sheppard et al., *J Biol Chem* 279:30480-89 (2004)). Their differences in adherence qualities suggested that rAls3p-N may have different efficacy as a vaccine immunogen compared to rAls1p-N.

The vaccines and vaccinations were performed as described above. Briefly, rAls1p-N and rAls3p-N (amino acids 17 to 432 of Als1p or Als3p) were

produced in *S. cerevisiae* and purified by gel filtration and Ni-NTA matrix affinity purification, as described above and in Ibrahim et al., (2005), *supra*; Spellberg et al., (2005), *supra*). The amount of protein was quantified by modified Lowry assay. A high degree of purity ($\approx 90\%$) was confirmed by SDS-polyacrylamide gel electrophoresis as well as circular dichroism and FTIR, as described above and in Ibrahim et al., (2005), *supra*; Spellberg et al., (2005), *supra*). Mice were immunized by subcutaneous (SQ) injection of 20 μg of rAls1p-N or rAls3p-N mixed with Complete Freund's adjuvant (CFA, Sigma-Aldrich, St. Louis, MO) at day 0, boosted with another dose of the antigen with Incomplete Freund's adjuvant (IFA, Sigma-Aldrich) at day 21, and infected two weeks following the boost.

Statistical analyses were performed as follows. The non-parametric Log Rank test was utilized to determine differences in survival times of the mice. Antibody titers and footpad swelling were compared by the Steel test for non-parametric multiple comparisons Rhyne et al., *Biometrics* 23:539-49 (1967), or the Mann Whitney U test for unpaired comparisons, as appropriate. Correlations were calculated with the Spearman Rank test. To determine if heterogeneity existed in replicate survival studies, the Kolmogorov-Smirnov test was utilized. P values < 0.05 were considered significant.

Vaccination with rAls3p-N was shown to stimulate a broader array of antibody responses in comparison with rAls1p-N. In this regard, the results shown in Figure 32 show mice vaccinated with CFA + rAls1p-N or rAls3p-N developed antibody titers significantly greater than mice receiving CFA alone. Of note, mice vaccinated with rAls3p-N generated anti-rAls1p-N antibodies at equivalent titers to mice vaccinated with rAls1p-N (Fig. 32, top). In contrast, mice vaccinated with rAls1p-N generated smaller titers against rAls3p-N than did mice vaccinated with rAls3p-N (Fig. 32, bottom). However, both rAls1p-N and rAls3p-N resulted in similar delayed type hypersensitivity responses *in vivo* as shown in Figure 33.

The rAls1p-N and rAls3p-N vaccines also were shown to mediate similar efficacy against disseminated candidiasis. Briefly, to further corroborate that the rAls3p-N vaccine was as effective as rAls1p-N against hematogenously disseminated candidiasis, mice were vaccinated with CFA, CFA + rAls1p-N, or CFA + rAls3p-N, and subsequently

infected via the tail-vein with *C. albicans*. The results shown in Figure 34 demonstrate that both the rAls1p-N and rAls3p-N vaccines resulted in significant improvement in survival.

Correlation of anti-Alsp antibody titers and delayed type hypersensitivity reactions with survival in vaccinated mice subsequently infected with *C. albicans* was also determined. Briefly, antibody titers were determined by ELISA in 96 well plates, as we have described previously and in Ibrahim et al., (2005), *supra*; Spellberg et al., (2005), *supra*. Wells were coated with 100 μ l per well of 5 μ g/ml rAls1p-N or rAls3p-N in PBS. Mouse sera were incubated for 1 h at room temperature following a blocking step with tris buffer saline (TBS) (0.01 M TrisHCl, pH 7.4, 0.15 M NaCl) containing 3% bovine serum albumin. The wells were washed 3 times with TBS containing 0.05% Tween 20, followed by another 3 washes with TBS without Tween. Goat anti-mouse IgG secondary antibody conjugated with horseradish peroxidase (Sigma-Aldrich) was added at a final dilution of 1:5000 and the plate was further incubated for 1 h at room temperature. Wells were washed with TBS and incubated with substrate containing 0.1 M citrate buffer (pH 5.0), 50 mg/ml of *o*-phenylenediamine (Sigma), and 10 μ l of 30% H₂O₂. The color was allowed to develop for 30 min after which the reaction was terminated by adding 10% H₂SO₄ and the optical density (OD) was determined at 490 nm in a microtiter plate reader. Negative control wells received irrelevant antibody, and background absorbance was subtracted from the test wells to obtain final OD readings. The ELISA titer was taken as the reciprocal of the last serum dilution that gave a positive OD reading (i.e. > mean OD of negative control samples + (standard deviation * 2)).

Delayed type hypersensitivity reactions were assessed by measuring the footpad swelling tests. Briefly, mice were immunized with rAls1p-N, rAls3p-N, or CFA alone. Two weeks following the boost, baseline footpad sizes of immunized mice were measured using an electronic digital caliper. Fifty μ g of rAls1p-N or rAls3p-N in 25 μ l of PBS were injected into the right footpads, and PBS alone injected into the left footpads of the immunized mice. Twenty-four hours later the footpads were again measured. Antigen-specific footpad swelling was calculated as: (right footpad thickness at 24 h – right footpad thickness at baseline) – (left footpad thickness at 24 h – left footpad thickness at baseline).

Vaccinated mice were bled for titer determinations and underwent footpad swelling tests two days prior to infection. Vaccinated mice that did not survive the infection nevertheless had a broad range of antibody titers as shown in Figure 35. Many such mice had anti-rAls1p-N and anti-rAls3p-N antibody titers of $\geq 1:50,000$ (≥ 4.5 5 $\log_{10}$). As a result, antibody titers did not significantly correlate with survival. In contrast, the intensity of footpad swelling reactions did correlate with survival (Fig. 35, $\rho = 0.6$ & $p = 0.009$ by Spearman Rank correlation test).

The rAls3p-N vaccine also demonstrated more efficacy than rAls1p-N in two models of mucosal candidiasis. Because Als3p mediated superior adhesion to 10 epithelial cells compared to Als1p, this observation indicates that rAls3p-N can exhibit unique efficacy in mucosal models of infection. The efficacy of rAls1p-N compared to rAls3p-N assessed in a steroid-treated, oropharyngeal model of infection and in a model of candidal vaginitis.

Briefly, vaccine studies in the above murine oropharyngeal candidiasis 15 (OPC) model were performed as previously described and as described in Spellberg et al., (2005), *supra*; Kamai et al., *Antimicrob Agents Chemother* 45:3195-57 (2001), and Kamai et al., *Infect Immun* 70:5256-58 (2002). Vaccinated mice were immunocompromised by treatment with cortisone acetate (225 mg/kg SQ on days -1, 1, and 3 relative to infection). On the day of infection, the mice were anesthetized by 20 intraperitoneal injection with 8 mg xylazine and 110 mg ketamine per kg. Calcium alginate urethral swabs were saturated with *C. albicans* by placing them in a suspension of 10^6 organisms per ml in HBSS at 30°C. The saturated swabs were placed sublingually in the oral cavity of the mice for 75 min. After 5 days of infection, the tongue and hypoglossal tissue were excised, weighed, homogenized, and then quantitatively cultured 25 to determine the oral fungal burden.

Effectiveness of the vaccine against murine vaginal candidiasis was performed by vaccinating female Balb/c mice were treated with 30 μ g of subcutaneous estradiol valerate dissolved in peanut oil (both from Sigma-Aldrich) on day -3 relative to infection to induce pseudoestrus. On the day of infection, mice were sedated by ip 30 administration of 100 mg/kg of ketamine. Sedated mice were infected intravaginally with

10^6 blastospores of *C. albicans* in 10 μ l of HBSS. On day 3 post-infection, vaginas and approximately one centimeter of each uterine horn were dissected en block, homogenized, and quantitatively cultured.

As shown in Figure 36, in cortisone-treated mice with oropharyngeal candidiasis, the rAls1p-N vaccine mediated a strong trend towards reduced tongue fungal burden ($p = 0.054$). The overall magnitude of the benefit was < 0.3 log CFU/gram (Fig. 36). In comparison, the rAls3p-N vaccine mediated a > 0.6 log CFU/gram decrease in tongue fungal burden that was statistically significant ($p = 0.005$, Fig. 36). Similarly, in a non-immunocompromised model of candidal vaginitis, the rAls3p-N vaccine mediated a 0.7 log CFU/gram decrease in vaginal fungal burden compared to CFA alone ($p = 0.02$) as shown in Figure 37. In comparison, rAls1p-N mediated no benefit at all in the vaginitis model, and rAls3p-N was significantly more effective than rAls1p-N ($p = 0.01$).

The above results indicate that a vaccine based on rAls3p-N, which is 85% homologous to rAls1p-N at the amino acid level, was equally effective against disseminated candidiasis, but was more effective than rAls1p-N against mucosal infection. The increased effectiveness of rAls3p-N was seen in both a steroid-treated model of oropharyngeal candidiasis and an immunocompetent model of candidal vaginitis. The above results also show achievement of $\geq 50\%$ long-term survival in a murine model of candidal septic shock with no adjunctive anti-fungal therapy is encouraging, and further corroborates the therapeutic benefit all ALS vaccines of the invention.

Antibody titers did not correlate with the protective effect of either vaccine during disseminated candidiasis, but induction of delayed type hypersensitivity *in vivo* did correlate with protection. These data also further corroborate the mechanism of vaccine-induced protection was induction of Type 1, cell-mediated immunity to the fungus. Both rAls1p-N and rAls3p-N induced equivalent titers of antibody against rAls1p-N, but that rAls3p-N induced significantly higher titers of anti-rAls3p-N antibodies than did rAls1p-N. These data indicated that, despite their high degree of amino acid sequence homology (85%), the humoral immune system can distinguish between rAls1p-N and rAls3p-N. The above results further corroborate that, regardless of differences in Als1p and Als3p

epithelial cell adherence characteristics, the rAls1p-N and rAls3p-N vaccines were equally effective in protecting against hematogenously disseminated (i.e. endovascular) candidiasis.

In sum, the anti-candidal rAls3p-N vaccine induced equivalent cell-mediated but broader antibody-based responses than did the rAls1p-N vaccine. The immunogens resulted in an equivalent degree of protection against hematogenously disseminated candidiasis, but rAls3p-N mediated greater protection against both oropharyngeal and vaginal candidiasis.

Throughout this application various publications have been referenced within parentheses. The disclosures of these publications in their entireties are hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains.

Although the invention has been described with reference to the disclosed embodiments, those skilled in the art will readily appreciate that the specific examples and studies detailed above are only illustrative of the invention. It should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 71

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 71

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

What is claimed is:

1. A vaccine comprising an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, with an adjuvant in a pharmaceutically acceptable medium.
- 5 2. The vaccine of claim 1, wherein said Als protein family member comprises an Als protein derived from a *Candida* strain selected from the group consisting of *Candida albicans*, *Candida krusei*, *Candida tropicalis*, *Candida glabrata* and *Candida, parapsilosis*
- 10 3. The vaccine of claim 1, wherein said Als protein family member is selected from the group consisting of Als1p, Als3p, Als5p, Als6p, Als7p and Als9p.
4. The vaccine of claim 1, wherein said immunogenic fragment comprises an N-terminal region fragment or any other fragment of an Als protein family member.
- 15 5. A method of treating or preventing disseminated candidiasis or a *Staphylococcus aureus* infection, comprising administering an immunogenic amount of a vaccine an isolated Als protein family member having cell adhesion activity, or an immunogenic fragment thereof, in a pharmaceutically acceptable medium.
- 20 6. The method of claim 5, wherein said Als protein family member comprises an Als protein derived from a *Candida* strain selected from the group consisting of *Candida albicans*, *Candida krusei*, *Candida tropicalis*, *Candida glabrata* and *Candida, parapsilosis*
7. The method of claim 5, wherein said Als protein family member is selected from the group consisting of Als1p, Als3p, Als5p, Als6p, Als7p and Als9p.
8. The method of claim 5, wherein said immunogenic fragment comprises an N-terminal region fragment of an Als protein family member.
- 25 9. The method of claim 5, wherein said administering comprising active immunization, passive immunization or a combination thereof.

10. A method of treating or preventing disseminated candidiasis or a *Staphylococcus aureus* infection, comprising administering an effective amount of an isolated Als protein family member having cell adhesion activity, or an functional fragment thereof, to inhibit the binding or invasion of *Candida* or *Staphylococcus aureus* to a host cell or tissue.


11. The method of claim 10, wherein said Als protein family member comprises an Als protein derived from a *Candida* strain selected from the group consisting of *Candida albicans*, *Candida krusei*, *Candida tropicalis*, *Candida glabrata* and *Candida parapsilosis*.


12. The method of claim 10, wherein said Als protein family member is selected from the group consisting of Als1p, Als3p, Als5p, Als6p, Als7p and Als9p.

13. The method of claim 12, wherein said cell adhesion activity comprises binding to gelatin, fibronectin, laminin, epithelial cells or endothelial cells.

14. The method of claim 10, wherein said functional fragment comprises an N-terminal region fragment of an Als protein family member.

15. The method of claim 10, wherein said host cell comprises a cell of endothelial or epithelial cell origin.

**Fig. 1A**

**Fig. 1B**




FIG. 2A




FIG. 2B




FIG. 2C






FIG. 2D

Wild-Type



als1/als1



als1/als1::ALS1





FIG. 3A

Wild-Type



P_{ADH1}-ALS1




FIG. 3B




FIG. 4A






FIG. 4B

efg1/efg1*efg1/efg1, P_{ADH1}ALS1***FIG. 4C**







Fig. 5A

Wild-Type

*als1/als1**als1/als1::ALS1***FIG. 5B**

Wild-Type

*als1/als1**als1/als1::ALS1***FIG. 5C**





Fig. 6

ALS1	.MLQOFTL	FLYLSIASAKTIT	V	DSEF	TWSNAANYAFKGP	GYP	WNAVLGWSLDG	S	NP	TLNMP	CVFKY	TSQT	D	DGVKYATCQ	Y;	99								
ALS3	.MLQOYTL	LIYLSVATAK	IT	V	NSFN	TWSNAATYNYKGP	GTP	WNAVLGWSLDG	S	SP	TLNMP	CVFKFTTSQT	D	HGVKYATCQ	Q;	99								
ALS5	.MIQOFTL	FLYLSFATAK	IT	I	NSID	TYSNAGNYAFKGP	GYP	YNAVLGWSLDG	S	NP	ILNMP	CVFKFTASQK	D	DGVKYATCQ	Y;	99								
ALS6	MKTIVILH	FFYCTIAMAK	TI	S	TSEF	TYTNTGNYPYGP	GYP	YTAVLGWSLDG	L	SP	TLVMP	CVFKFTTQT	D	NGVKYATCT	H;	100								
ALS7	MKKLYLLY	LASFYTVISKE	VT	V	NQFN	IWSYTYRAREYE	EISTL	AKAOLEWALDG	I	SP	TLVMP	CVYKFTMYET	Q	NSIAYATCD	D;	100								
ALS9	.MLPQFL	LLYLTVSTAK	IT	V	NSFN	TWANAANYGYQ	IPETP	WTAVLGWSLNS	T	DA	TLIMP	CVFKFTTSQT	D	DGVSYATCD	N;	99								
Consensus	L	K	G	F	SL	T	A	L	W	L	T	A	GDTF	L	MPCV	K	SV	LTA	YATC	F	;			
ALS1	S	EFTT	TLTCTVNDAL	KSSIKAF	T	TLPIAFNVGGT	STDLE	CFTAGTNT	N	DKDI	IDVEFEKSTV	DP	SAYLYAS	VMPSLNK	VTTL	;	199							
ALS3	A	EFMT	TLTCTVNSNTL	TPSIKAL	T	TLPIAFNVGGT	SVDL	CFTAGTNT	N	GKKI	INVDFFERSNV	DP	KGYLTD	VIPSLNK	VSTL	;	199							
ALS5	S	EFTT	TLTCTVNDAL	KSSIKAF	T	TLPIAFNVGGT	STDLE	CFTAGTNT	N	SKKL	IAVNF	FKSTVDRSG	YLTT	FMPSLNK	VIATL	;	199							
ALS6	A	DFTT	SMSCVNVNGL	SSNIRAF	T	RLPIAFNVGGT	SVNIQ	CFTAGTNT	T	DKHI	TTVNF	PKTPQSSSL	LVYFA	VIPSLDK	LSL	;	200							
ALS7	A	DTKS	SLKCTVTD	DELTE	TSVF	S	ILPIAFNVGGS	KSTIT	CFSSGYNT	F	NNQL	TTANFLPR	RELAFGL	VVSQ	LSMSLD	TMTNF	;	200						
ALS9	A	EFTT	SLSCTVNSVS	VSYDKAS	T	KLPIAFNVGGT	SVDLT	CFTAGTNT	T	DTEI	TSVDFQAS	PISSSG	YIASA	VVPSLNK	KASL	;	199							
CON	GE	FS	C	V	G	V	LP	FN	VGG	GS	DSKCF	G	NTVTF	DG	S	F	R	SL						
ALS1	FVAPQ	EN	TSGTGMG	FSSSSNGDVA	NIHIGITKGL	D	NY	S	ESFSYTKTCT	SG	NIQIKYQNV	PAGYRPF	DAYIS	ATDVNQY	TLAYTNDYT	;	297							
ALS3	FVAPQ	AN	TSGTGMGFANTY	GDVQ	NIHVGITKGL	D	NY	S	ESFSYTKTCS	NGIFITYK	NPAGYRPF	FVDAYIS	ATDVNSY	TL	SYANEYT	;	297							
ALS5	YVAPQ	EN	TSGTGMGFST	SYGDVA	NVHIGISKGV	D	NH	T	ESFSYTKSC	SFGISITYQ	NPAGYRPF	FIDAYIS	PSDNQYQ	LS	YKNDYT	;	297							
ALS6	VVASQ	TA	ASGVLGFS	ATKDDVT	TIHVGITNGL	S	NM	S	ESFSYTKTCT	PN	SIITYENVP	PAGYRPF	FIDSVYKKS	AFATNGFN	LN	TYNIYN	;	300						
ALS7	VMSTP	FM	QSGKLGF	TNSDDDFE	SIHVGITNEI	D	SM	S	VPDFHTIR	CT	SALYIEFK	TIPAGYRPF	FVDAIVQ	IPTEP	PFV	KYTN	NEFA	;	298					
ALS9	FVLPO	EN	TSGIMGF	VTISQ	AT	NINIGISKGL	D	NF	S	ESFTYTKT	CS	SGIIVEYEN	VPAGYRPF	FVDAYIS	SENVEQ	YTLTYANEYT	;	296						
CONSENSUS	C	GY	SG	GF	IDCS	GI	NW	PV	S	F	T	C	PAGYRPF	D	Y	N								
ALS1	AGSR	SQSK	FTLRW	TG	YKNSD	GSN	IVIVAT	TRITVD	T	AV	NPSVDKTK	EILQ	TITTSYV	GV	T	SYL	KT	PIGETAT	VI	VDV	;	396		
ALS3	AGGY	WQRA	FTLRW	TG	YRNSD	GSN	IVIVAT	TRITVD	T	AV	DPNRDKTK	EILK	TITTSYV	GV	T	SYL	KT	PIGETAT	VI	VDI	;	396		
ALS5	VDDY	WQHA	FTLK	WTG	YKNSD	GSN	IVIVAT	TRITVD	T	AV	NPSVDKTK	EILQ	TITTSYV	GV	T	SYL	KT	PIGETAT	VI	VDV	;	396		
ALS6	MDGK	KGND	LIYF	WTS	YTN	SD	GSN	AAV	VT	TRITVD	T	AI	DPTVDKTK	EVIE	TITTSYV	GV	T	SYL	KT	TIGTAT	VI	VDV	;	399
ALS7	VNGI	YTSI	FTS	FSQ	PILYDE	LAI	ADL	VRT	TS	TVIG	I	RT	ISRLQKTK	LVLE	TVTTS	HGFD	WYY	KK	TIGTAT	VI	FIDV	;	398	
ALS9	KNGN	TVVD	F	TLTWIG	YKNS	E	DSN	DIIV	VT	TKT	VT	A	NPTVDKTE	EVIO	TITTSYV	GV	T	SYE	FT	TIGTAT	VI	VDI	;	395
CON	C	P	A	G	V	TT	TV	S	T	TL	PF	KT	TI	PI	TTTT	TTS	G	T						

C. albicans ALS Gene Family**Fig. 8**






FIG. 9

C. albicans ALS Gene Family**Fig. 10**



Residue number
FIG. 11A




FIG. 11B




Fig. 12




FIG. 13





Fig. 14

**Fig. 15**









Fig. 16A, 16B

**Fig. 17**

**Fig. 18**

**Fig. 19**

**Fig. 20**

**Fig. 21**




Fig. 22A, 22B





Fig. 23A, 23B

**Fig. 24**





Figure 25

**FIGURE 26**




Figure 27




Figure 28




FIGURE 29




FIGURE 30




FIGURE 31




FIGURE 32




FIGURE 33




FIGURE 34




FIGURE 35




FIGURE 36




FIGURE 37