



US 20070092505A1

(19) **United States**

(12) **Patent Application Publication**
Buelow et al.

(10) **Pub. No.: US 2007/0092505 A1**

(43) **Pub. Date: Apr. 26, 2007**

(54) **PRODUCTION OF HUMANIZED
ANTIBODIES IN TRANSGENIC ANIMALS**

(76) Inventors: **Roland Buelow**, Palo Alto, CA (US);
Josef Platzer, Muenchen (DE); **Wim
van Schooten**, Sunnyvale, CA (US);
Jens-Ulrich Buelow, Karlsruhe (DE)

Correspondence Address:
HELLER EHRMAN LLP
275 MIDDLEFIELD ROAD
MENLO PARK, CA 94025-3506 (US)

(21) Appl. No.: **11/584,187**

(22) Filed: **Oct. 20, 2006**

Related U.S. Application Data

(62) Division of application No. 09/921,819, filed on Aug. 3, 2001, now Pat. No. 7,129,084.

(60) Provisional application No. 60/222,872, filed on Aug. 3, 2000. Provisional application No. 60/276,156, filed on Mar. 15, 2001.

Publication Classification

(51) **Int. Cl.**
A61K 39/395 (2006.01)
C07K 16/44 (2006.01)
(52) **U.S. Cl.** **424/133.1; 530/388.15**

(57) **ABSTRACT**

This invention relates to humanized antibodies and antibody preparations produced from transgenic non-human animals. The non-human animals are genetically engineered to contain one or more humanized immunoglobulin loci which are capable of undergoing gene rearrangement and gene conversion in the transgenic non-human animals to produce diversified humanized immunoglobulins. The present invention further relates to novel sequences, recombination vectors and transgenic vectors useful for making these transgenic animals. The humanized antibodies of the present invention have minimal immunogenicity to humans and are appropriate for use in the therapeutic treatment of human subjects.

```

1
clone11 [REDACTED] CACGA AACTTTACGG
clone3 [REDACTED] CACGA GGCCCTGCAC
clone5 [REDACTED] CACGA GGCCCTGCAC

151
clone11 AATCACTACA AAGAGAAGTC CACCTCGAGG TCTCCGGGTA AATGAGCCTC
clone3 AATCACTACA CGCAGAAGTC CACCTCTAAG TCTGCGGGTA AATGAGCCTC
clone5 AATCACTACA CGCAGAAGTC CACCTCTAAG TCTGCGGGTA AATGAGCCTC

201
clone11 GCGCCGCTGA TCTAGTGGAC GTTCCTCAT CCACCCACCC CTCCCCCCAC
clone3 ACGTCCCTGC ACCAGCAAGC CCTCACCCAG C..... ..CCACCCTC
clone5 ACGTCCCTGC ACCAGCAAGC CCTCACCCAG C..... ..CCACCCTC

251
clone11 CCCGGGCTCC AGGTCCAGCC AGGGCGCCCT AGCCCCTCCC TGTGTGCATT
clone3 CCCGGGCTCC AAGTCCAGCC AGGACGCCCT AGCCCCTCCC TGTGTGCATT
clone5 CCCGGGCTCC AGGTCCAGCC AGGACGCCCT AGCCCCTCCC TGTGTGCATT

301
clone11 CCTCCTGGGC CGCCGTGAAT AAAGCACCCA GGCCGCCCTG GGACCCTGCA
clone3 CCTCCTGGGC CGCCGTGAAT AAAGCACCCA GGCCACCCTG GGACCCTGCA
clone5 CCTCCTGGGC CGCCGTGAAT AAAGCACCCA GGCCGCCCTG GGACCCTGCA

351
clone11 ACGCTGTGCT GGTTCTTTCC GAGGCAGAGC CCTGGTGGCC GCCAGGCCTG
clone3 ACGCTGTGCT GGTTCTTTCC GAGGCAGAGC CCTGGTGGCC GCCAGGCCTG
clone5 ACGCTGTGCT GGTTCTTTCC GAGGCAGAGC CCTGGTGGCC GCCAGGCCTG

401
clone11 CGGGGGTGGG CTGAGCCGAC TCTGGGCCAC TTTGTTTCAGC ATCTGTGGGG
clone3 CAGGGGTGGG CTGAGCCGAC TCTGGGCCAC TTTGTTTCAGC ATCTGTGGGG
clone5 CGGGGGTGGG CTGAGCCGAC TCTGGGCCAC TTTGTTTCAGC ATCTGTGGGG

451
clone11 GAGCTGACCC CACTCCGGGC CAGACACACA GTGAGTGGGT CCAGCAGGCC
clone3 GAGCTGACCC CGTCCGGGC CAGACACACA GTGAGTGGGT CCAGCAGGCC
clone5 GAGCTGACCC CACTCCGGGC CAGACACACA GTGAGTGGGT CCAGCAGGCC

501
clone11 ACCTGGGGGC TGCCCAAGGC CACAGAGGGG CTTGGCCAGA GGCACAGCTC
clone3 ACCTGGGGGC TGCCGAGGC CACGAGGGG CTTGGCCAGA GGCGTACCTC
clone5 ACCTGGGGGC TGCCCAAGGC CACAGAGGGG CTTGGCCAGA GGCACAGCTC

```

Figure 1(a)

	551		600		
clone11	CACGGTCCCC	TCCAGCCACC	ACCTGCTGGG	CCGGCCTCTG	GACAGGAACC
clone3	CACGGCCCCC	TCCAGCCACC	ACCTGCTGGG	CCGGCCTCTG	GACAGGAACC
clone5	CACGGTCCCC	TCCAGCCACC	ACCTGCTGGG	CCGGCCTCTG	GACAGGAACC
	601		650		
clone11	GGGGAAGCCC	CCGAGACCCT	CAGGGATTGA	GGCCCAATGC	TTCCCGCCTC
clone3	GGGGAAGCCC	CCGAGACCCT	CAGGGATTGA	GGCCCAATGC	TTCCCGCCTC
clone5	GGGGAAGCCC	CCGAGACCCT	CAGGGATTGA	GGCCCAATGC	TTCCCGCCTC
	651		700		
clone11	TGCTCCAGCC	CACGCTGTGG	GGCAGGGCCA	CATCCTTGTC	CCCAGGCCCC
clone3	TGCTCCAGCC	CACGCTGTGG	GGCAGGGCCA	CATCCTTGTC	CCCAGGCCCC
clone5	TGCTCCAGCC	CACGCTGTGG	GGCAGGGCCA	CATCCTTGTC	CCCAGGCCCC
	701		750		
clone11	TGTCCTTGGG	TGTCCAGAGT	CCTTGTGTCC	ACTCTGGGCC	TGCCTGGAGC
clone3	TGTCCTTGGG	TGCCCAGAGT	CCTTGTGTCC	ACTCTGGGCC	TGCCTGGAGC
clone5	TGTCCTTGGG	TGTCCAGAGT	CCTTGTGTCC	ACTCTGGGCC	TGCCTGGAGC
	751		800		
clone11	CACGCATGGC	CAGGGGGTGG	CCCTGCTTCA	CCCTCAGGCT	CCCAAGGTCA
clone3	CACGCATGGC	CGGGGGATGG	CCCTGCTTCA	CCCTCAGGCT	CCCAAGGTCA
clone5	CACGCATGGC	CAGGGGGTGG	CCCTGCTTCA	CCCTCAGGCT	CCCAAGGTCA
	801		850		
clone11	GGCCTCGCCC	TCCCTCGGCC	AGGAGGCTCT	GCCCGGCTCT	CCCTGCCCAG
clone3	GGCCTCGCCC	TCCCTCAGCC	AGGAGGCTCT	GCCCGGCTCT	CCCTGCCCAG
clone5	GGCCTCGCCC	TCCCTCGGCC	AGGAGGCTCT	GCCCGGCTCT	CCCTGCCCAG
	851		900		
clone11	GGCCAGGCCT	GTGCGCCCAT	GGGGAGGTCA	TCCCTGTGCC	TGAAAGGGGT
clone3	GGCCAGGCCT	GTGCGCCCAT	GGGGAGGTCA	TCCCTGTGCC	TGAAAGGGCT
clone5	GGCCAGGCCT	GTGCGCCCAT	GGGGAGGTCA	TCCCTGTGCC	TGAAAGGGGT
	901		950		
clone11	CCAGGCCGAG	AGCCCTGAAT	GTCCAGGGCA	GGGACCTAGC	TGCTCCCTGT
clone3	CCAGGCCGGG	AGCCCTGAAT	GTCCAGGGCA	GGGACCTAGC	TGCTCCCTGC
clone5	CCAGGCCGAG	AGCCCTGAAT	GTCCAGGGCA	GGGACCTAGC	TGCTCCCTGT
	951		1000		
clone11	GGACACGGAG	CCCAGAGCCA	CAGACAACAA	GCCCCAGCCC	CGCACGCACA
clone3	AGACACGGAG	CCCAGAGCCA	CAGACAACAA	GCCCCAGCCC	CGCACGCACA
clone5	GGACACGGAG	CCCAGAGCCA	CAGACAACAA	GCCCCAGCCC	CGCACGCACA
	1001		1050		
clone11	CGAGACAGCC	CGCACCCAGC	CTCCTCCACA	CGCACTCAGG	TGTACATGGC
clone3	CAAGACAGCC	CGCACCCAGC	CTCCTCCACA	CGCACTCAGG	TGTGCATCCG
clone5	CGAGACAGCC	CACACCCCGC	CTCCTCCACA	CGCACTCAGG	TGTGCATCCG

Figure 1(b)

	1051				1100
clone11	CACATGAGCA	CACTTCACCC	CGTCACACCC	ACACACCTAC	ACACACTCAG
clone3	CACATGAGCA	CACTTCACCC	CGTCACACCC	ACACGCCTAC	ACACACTCAG
clone5	CACATGAGCA	CACTTCACCC	CATCACACCC	ACACGCCTAC	ACACACTCAG
	1101				1150
clone11	GTCTCGCACT	CGGGGACCCA	TGGGGTGACC	CCACGGGCCC	AGA.CCAGAG
clone3	GTCTCGCACT	CGGGGACCCA	TGGGGTGACC	CCACAGGCCC	AGACCCAGAG
clone5	GTCTCGCACT	CGGGGACCCA	TGGGGTGACC	CCACAGGCCC	AGACCCAGAG
	1151				1200
clone11	CTGGGTCTTG	TGAGCCCTCC	CTGTGGACAC	CAGCTGGGCC	CCACCCTCCA
clone3	CTGGGTCTTG	TGAGCCCTCC	CTGTGGACAC	CAGCTGGTCC	CCACCCTCCA
clone5	CTGGGTCTTG	TGAGCCCTCC	CTGTGGACAC	CAGCTGGTCC	CCACCCTCCA
	1201				1250
clone11	GCGCCCATGG	GCTGCTCAGC	GGCCCTTTCC	CACACTGACC	ACACTGACCA
clone3	GCGCCCATGG	GCTGCTCAGT	GGCCCTTTCC	CACACTGACC	ACACTGACCA
clone5	GCGCCCGTGG	GCTGCTCAGC	GGTCCTTTCC	CACACTGACC	ACACTGACCA
	1251				1300
clone11	GGTCAGACAT	CCGTCCTTG	CCTCCCCTGG	GACACCCAGC	CCCCTCCCTA
clone3	GGTCAGACAT	CCGTCCTTG	CCTCCCCTGG	GGCACCCAGC	CCCCTCCCTA
clone5	GGTCAGACAT	CCGTCCTTG	CCTCCCCTGG	GGCACCCATG	CCCCTCCCTA
	1301				1350
clone11	GCAGGCTGAG	ATCCCCCCTC	AGCCCCCTCGT	CCTGGCAGCC	TCACCCCTCG
clone3	GCAGGCTGAG	ATCCCCCCTC	AGCCCCCTCGT	CCTGGCACCC	TCACCCCTCA
clone5	GCAGGCTGAG	ATCCCCCCTC	AGCCCCCTCGT	CCTGGCACCC	TCACCCCTCA
	1351				1400
clone11	GGCACAGCAC	CCCTCAGGCC	CGGTGCTGTC	AGCCCTCCCT	CCCCGGGGGC
clone3	GGCACAGGGA	CAC...AGCC	CGGCGCTGTC	TGCCCTCCCT	CCCTGGGGGC
clone5	GGCACAGGGA	CAC...AGCC	CGGTGCTGTC	TGCCCTCCCT	CCCTGGGGGC
	1401				1450
clone11	AGGGCCCAGG	AACGTGCGCT	CTGCTGACCC	TCCCAGCTCC	AGGCCTGGCC
clone3	AGGGCCCAGG	CTCACATGCT	CTGCTGACCC	TCCCGGCTCC	AGGCCTGGCC
clone5	AGGGCCCAGG	CTCACATGCT	CTGCTGACCC	TCCCAGCTCC	AGGCCTGGCC
	1451				1500
clone11	CCCAGGGCAG	AGGAGGCCAG	GAAGTGAAGCC	TCTGTCCTGT	GGGGAGGTAG
clone3	CCCAGGGCAG	AGGAGGCCAG	GAAGTGAAGCC	TCTGTCCTGG	GGGGAGGTGG
clone5	CCCAGGGCAG	AGGAGGCCAG	GAAGTGAAGCC	TCTGTCCTGG	GGGGAGGTGG
	1501				1550
clone11	GGTCAGGGTC	CCAGCTCAGG	GCACAGCTCA	GGATGGGAGC	AGGACCCAC
clone3	GGTCAGGGCC	CCAGCTCAGG	GCACAGCTCA	GGATGGGAAC	AGGACACCAC
clone5	GGTCAGGGCC	CCAGCTCAGG	GCACAGCTCA	GGATGGGAGC	AGGACACCAC

Figure 1(c)




	1551				1600
clone11	AGGCCAGGCC	CAGATAGCAG	CCAGGGCTGG	AGGGGTGGG	GCTGGGGCTG
clone3	AGGCCAGGCC	CAGACAGTGG	CCAGGGCTGG	AGGGGTGGG	TCTGGGGCTG
clone5	AGGCCAGGCC	CAGACAGTGG	CCAGGGCTGG	AGGGGTGGG	TCTGGGGCTG
	1601				1650
clone11	GGCCCCAGAG	ACTGACCTCA	GGTGACCCCT	GCCTGGCCCA	TGGGGAGATC
clone3	GGCCCCAGAG	ACTGACCTCA	GGTGATCCCT	GCCCAGCCCA	TGGGGGGATC
clone5	GGCCCCAGAG	AATGACCTCA	GGTGATCCCT	GCCCAGCCCA	TGGGGGGATC
	1651				1700
clone11	ACGCCACCTT	CCCCCACC	AGAGGGAGCC	CTGCCC...T	ACCCAGTGA
clone3	CTGCCACCTT	CCCCCACC	AGAGGGAGCC	CTGCCCCGAG	GCCCTGATGA
clone5	CTGCCACCTT	CCCCCACC	AGAGGGAGCC	CTGCCCCGAG	GCCCTGATGA
	1701				1750
clone11	CCCTGCCAG	CCCTCCGTGG	GCAGACACAG	CACTGACCAC	CCCTCCCTGT
clone3	TGCCACCCAG	CCCCCGTGG	GCAGACACAG	CACTGACCAC	CCCTCCCTGT
clone5	TGCCACCCAG	CCCCCGTGG	GCAGACACAG	CACTGACCAC	CCCTCCCTGT
	1751				1800
clone11	GCAGACTTGC	TGCTGGAGGA	GGAGATCTGT	GCGGACGACC	TGGATGGGGA
clone3	GCAGACTTGC	TGCTGGAGGA	GGAGATCTGT	GCGGACGCC	AGGACGGGGA
clone5	GCAGACTTGC	TGCTGGAGGA	GGAGATCTGT	GCGGACGCC	AGGACGGGGA
	1801				1850
clone11	GCTGGACGGG	CTC			
clone3	GCTGGACGGG	CTC			
clone5	GCTGGACGGG	CTC			

Figure 1(d)

	101				150
clone11			CACGAGGCTC	TACACAACCA	CTACACACAG
clone1			CACGAGGCTC	TGCACAACCA	CTACACACAG
	151				200
clone11	AAGTCGATCT	CTAAGCCTCC	GGGTAAATGA	GCCACATGCC	CCCGCACCAG
clone1	AAGTCGGTCT	CTAAGCCTCC	GGGTAAATGA	GCCACACGCC	CCCGCACCAG
	201				250
clone11	CAAGCCCTCA	CCCAGCCC GC	CCTCCCCGGG	CTCCAGGTCC	AGCCAGGACG
clone1	CAAGCCCTCA	CCCAGCCC GC	CCTCCCCGGG	CTCCAGGTCC	AGCCAGGACG
	251				300
clone11	CCCTAGCCCC	TCCCTGTGTG	CATGCCTCCT	GGGCCGCCAT	GAATAAAGCA
clone1	CCCTAGCCCC	TCCCTGTGTG	CATGCCTCCT	GGGCCGCCAT	GAATAAAGCA
	301				350
clone11	CCCAGGCCGC	CCTGGGACCC	TGCAACGCTG	TGCTTGTTCT	TTCCGAGGCA
clone1	CCCAGGCCGC	CCTGGGACCC	TGCAGCGCTG	TGCTGGTTCT	TTCCGAGGCA
	351				400
clone11	GAGCCCTGGT	GACCGCCAGG	CCTGCGGGGG	GTGGGCTGAG	CCCACTCTGG
clone1	GAGCCCTGGT	GATCGCCAGG	CCTGCGGGGG	GCGGGCTGAG	CCCACTCTGG
	401				450
clone11	GCCGCTTGGT	TCAGCATCTG	TGGGGGCGCT	GACCCCTCTC	CGGGCCAGAC
clone1	GCCGCTTGGT	TCAGCATCTG	TGGGGGCGCT	GACCCCTCTC	CGGGCCAGAC
	451				500
clone11	ACACAGTGAG	TGGGTCCGGC	AGGGCACCTG	GGGGCTGCCC	GAGGCCTCGG
clone1	ACACAGTGAG	TGGGTCCGGC	AGGGCACCTG	GGGGCTGCCC	GAGGCCTCGG
	501				550
clone11	AGGGGCTTGG	CCAGAGGCGC	AGCTTCACGG	CCCCCTCCAG	CCACCACATT
clone1	AGGGGCTTGG	CCAGAGGCGC	AGCTTCACGG	CCCCCTCCAG	CCACCACATT
	551				600
clone11	CTGGGCCAGA	CTCTGGGCAG	GAACGGGGGA	AGCCCCCGAC	ACCTCAGGGA
clone1	CTGGGCCAGA	CTCTGGGCAG	GAACGGGGGA	AGCCCCCGAC	ACCTCAGGGA
	601				650
clone11	TTGAGGCCCA	ACGCTTCCCG	CCTCTGCTCC	AGCCCACGCT	GAGGGGCAGG
clone1	TTGAGGCCCA	ACGCTTCCCG	CCTCTGCTCC	AGCCCACGCT	GAGGGGCAGG

Figure 2(a)

	651				700
clone11	GCCGCGGCCT	TGTCCCCAGG	CCCCTGTTCC	TGGGTGCCCA	GAGTCCGTGT
clone1	GCCGCGGCCT	TGTCCCCAGG	CCCCTGTTCC	TGGGTGCCCA	GAGTCCGTGT
	701				750
clone11	GTCCACTCTG	GGCCTGCCTG	GAGCCAGACT	GGCCCAGGGG	GAGGCCCTGC
clone1	GTCCACTCTG	GGCCTGCCTG	GAGCCAGACT	GGCCCAGGGG	GAGGCCCTGC
	751				800
clone11	TTCACCCTCA	GGCTCCCGAG	GTCAGGCATC	ATCCTCGTCG	GCCAGTAGCT
clone1	TTCACCCTCA	GGCTCCCGAG	GTCAGGCATC	ATCCTCGTCG	GCCAGTAGCT
	801				850
clone11	CTGCCTGGCT	CTCTCTGCCC	GGGGCCAAGC	CTGTGTGCCC	ATGGGGAGGT
clone1	CTGCCTGGCT	CTCTCTGCCC	GGGGCCAAGC	CTGTGTGCCC	ATGGGGAGGT
	851				900
clone11	CGTCCCTGTG	CCTGAAAAGG	GCCCAGGCTG	GGAGCCCTGA	ACGTCCAGGG
clone1	CGTCCCTGTG	CCTGAAAAGG	GCCCAGGCTG	GGAGCCCTGA	ACGTCCAGGG
	901				950
clone11	CAGGGACCTA	GCTGCTCCCT	GGGGACACTG	AGCCCAGAGC	CCCAGACACC
clone1	CAGGGACCTA	GCTGCTCCCT	GGGGACACTG	AGCCCAGAGC	CCCAGACACC
	951				1000
clone11	AAGCCCCAGC	CCCGCACGCA	CACGAGACAG	CCCACACCCA	GCGTCTCCA
clone1	AAGCCCCAGC	CCCGCACGCA	CACGAGACAG	CCCACACCCA	GCGTCTCCA
	1001				1050
clone11	CACGCACTCA	GGCGTCCACC	CGCACACAAG	CATGCTTCAC	CCCCGTCACA
clone1	CACCCACTCA	GGCGTCCACC	CGCACACAAG	CATGCTTCAC	CCCCGTCACA
	1051				1100
clone11	CACCCACATG	CCTGCACACA	CTCAGGTCTC	ACGCTCCGGG	ACCCATGGAG
clone1	CACCCACATG	CCTGCACACA	CTCAGGTCTC	ACGCTCCGGG	ACCCATGGAG
	1101				1150
clone11	TGATCCCACG	GGCCCAGACC	CAGAGCTGGG	TTCATGAGC	CCTCCCTGTG
clone1	TGATCCCACG	GGCCCAGACC	CAGAGCTGGG	TTCATGAGC	CCTCCCTGTG
	1151				1200
clone11	GACACCAGCT	GGTCCCCATT	CTCCAGCGCC	CTTGGGCTGC	TCAGTGGCCC
clone1	GACACCAGCT	GGTCCCCATC	CTCCAGCGCC	CTTGGGCTGC	TCAGTGGCCC
	1201				1250
clone11	TTTCCCACAC	TGACCACACT	GACCAGGTCA	GACATCCTTC	CTCGCCTCCC
clone1	TTTCCCACAC	TGACCACACT	GACCAGGTCA	GACATCCTTC	CTCGCCTCCC

Figure 2(b)

	1251				1300
clone11	CTGGGGCACC	CACGCCCCTC	CCTCGCAGGC	TGAGACCCCC	CCTCAGCCCC
clone1	CTGGGGCACC	CACGCCCCTC	CCTTGCAGGC	TGAGACCCCC	CCTCAGCCCC
	1301				1350
clone11	TCGTCCTGGC	ACCCTCACCC	CTCGGGCACA	GGGACACAGC	CCGGCACTGT
clone1	TCGTCCTGGC	ACCCTCACCC	CTCGGGCACA	GGGACACAGC	CCGGCACTGT
	1351				1400
clone11	CTGCCCTCCC	TCTCGGGGAC	AGAGCCCAGG	CACGTGTGCT	CTGCTGAGCC
clone1	CTGCCCTCCC	TCTCGGGGAC	AGAGCCCAGG	CACGTGTGCT	CTGCTGAGCC
	1401				1450
clone11	TCCCGGCTCC	AGGCCTGGCC	CCCAGGGCAG	AGGAGGCCAG	GAATTGAGCC
clone1	TCCCGGCTCC	AGGCCTGGCC	CCCAGGGCAG	AGGAGGCCAG	GAATTGAGCC
	1451				1500
clone11	TCTGTCTTGC	GGGGAGGTGG	GGTCAGGGCC	CCAGCTCAGG	GCACAGCTCA
clone1	TCTGTCTTGC	GGGGAGGTGG	GGTCAGGGCC	CCAGCTCAGG	GCACAGCTCA
	1501				1550
clone11	GGATGGGAGC	AGGACCCAC	AGGCCAGGCC	CAGACAGTGG	CCAGGGCTGG
clone1	GGATGGGAGC	AGGACCCAC	AGGCCAGGCC	CAGACAGTGG	CCAGGGCTGG
	1551				1600
clone11	GGCTGGGGCT	GGGGCCCAGA	GACTGACCTC	AGGTGACCCC	TGCCCCGCCC
clone1	GGCTGGGGCT	GGGGCCCAGA	GACTGACCTC	AGGTGACCCC	TGCCCCGCCC
	1601				1650
clone11	ATGGGGGATC	ACACCGCCAT	CCCCCCC GCC	GCAGAGGGAG	CCCTGCCCCG
clone1	ATGGGGGATC	ACACCGCCAT	CCCCCCC GCC	GCAGAGGGAG	CCCTGCCCCG
	1651				1700
clone11	AAGCCCCGAT	GGCCCCGCCC	AGCCCCCGT	GGGCAGACAC	AGCACTGACC
clone1	AAGCCCCGAT	GGCCCCGCCC	AGCCCCCGT	GGGCAGACAC	AGCACTGACC
	1701				1750
clone11	CCCCTCCCTG	TGCAGATCTG	CTGCTGGAGG	AGGAGAGCTG	TGCGGACGCC
clone1	CCCCTCCCTG	TGCAGATCTG	CTGCTGGAGG	AGGAGAGCTG	TGCGGACGCC
	1751				1800
clone11	CAGGACGGGG	AGCTGGACGG	GCTCTGGACG	ACTATCTCCA	TCTTCATCAC
clone1	CAGGACGGGG	AGCTGGACGG	GCTCTGGACG	ACTATCTCCA	TCTTCATCAC
	1801				1850
clone11	GCCCTTCCTG	CTCAGCGTCT	GCTACAGCGC	CACCGTGACC	CTCTTCAAGG
clone1	GCTCTTCCTG	CTCAGCGTCT	GCTACAGTGC	CACCGTGACC	CTCTTCAAGG

Figure 2(c)

	1851		1900
clone11	TGGGGGTCCA	CCCTGCTGGG	CCCTCGGGCC
clone1	TGGGGGCCCA	CCCTGCTGGG	CCCTCGGGCC
	1901		1950
clone11	CCCGCAGAGT	CCCTCCCTGC	CCCTCACTGT
clone1	CCCGCAGAGT	CCCTCCCTGC	CCCTCACTGT
	1951		2000
clone11	CCCTCTCTGT	CCCTCTCTGT	CCCTCTCTGT
clone1	CCCTCTCTGT	CCCTCTCTGT	CCCTCTCTGT
	2001		2050
clone11	GTAAGCTTGA	GACAGATTGG	GGTCATTTCA
clone1	GTAAGCTTGA	GACAGATTGG	GGTCATTTCA
	2051		2100
clone11	TGTGCCGCAC	GCCTCCCTTC	ATGTCAGTGG
clone1	TGTGCCGCAC	GCCTCCCTTC	ATGTCAGTGG
	2101		2150
clone11	AGTGCTGGGT	GAGAAATGAG	GCTTGCGGCG
clone1	AGTGCTGGGT	GAGAAATGAG	GCTTGCGGCG
	2151		2200
clone11	ACTGCTGCTC	CCTGAGACCT	GCGCGGACAC
clone1	ACTGCTGCTC	CCTGAGACCT	GCGCGGACAC
	2201		2250
clone11	GCGGGCAAGG	GAAAACGCCC	TCTTGGTCTC
clone1	GCGGGCAAGG	GAAAACGCCC	TCTTGGTCTC
	2251		2300
clone11	TTGGTCATCA	GTCCTCCCC	CAGTGAGGCT
clone1	TTGGTCATCA	GTCCTCCCC	CAGTGAGGCT
	2301		2350
clone11	CTCGATGGTC	AGGTCAGGAC	TGTCATAGAC
clone1	CTCGATGGTC	AGGTCAGGAC	TGTCATAGAC
	2351		2400
clone11	ATGCTTGCTT	TTTGTGTGCA	GAGAGCCTGT
clone1	ATGCTTGCTT	TTTGTGTGC	GAGAGCCTGT
	2401		2450
clone11	CTCACTGAGC	TCGCGGGGCA	GGGGTGGGCT
clone1	CTCACTGAGC	TCGCGGGGCA	GGGGTGGGCT

Figure 2 (d)

	2451				2500
clone11	GGGAGCGCAT	CTCCAGCATG	CTGTGCGACA	GCTTCGTTGC	TAACAAGACC
clone1	GGGAGCGGCA	TCTCCAGCTG	CTGTGCGACA	GCTTCGTTGC	TAACAAGACC
	2501				2550
clone11	GCTTAGTCTC	GTGGTTAGAC	CAACCTGCTT	TCTCGAGTAA	TTGTTAATTT
clone1	GCTTAGTCTC	GTGGTTAGAC	CAACCTGCTT	TCTCGAGTAA	TTGTTAATTT
	2551				2600
clone11	ACAGGAGTTT	CCTGTATTTT	TCAACTTATA	ATCCCCTAGT	CAGATAACTC
clone1	ACAGGAGTTT	CCTGTATTTT	TCAACTTATA	ATCCCCTAGT	CAGATAACTC
	2601				2650
clone11	TTTAATCACC	TATTCTGCCC	CTTCATTTTC	TCCCTATCGA	TCTCAGCAAC
clone1	TTTAATCACC	TATTCTGCCC	CTTCATTTTC	TCCCTATCGA	TCTCAGCAAC
	2651				2700
clone11	CCATCACTGC	CCTCACTGTC	CTTAAACTGT	CCCTTAACTG	ACCAGACTGT
clone1	CCATCACTGC	CCTCACTGTC	CTTAAACTGT	CCCTTAACTG	ACCAGACTGT
	2701				2750
clone11	CCCTCAGTGT	CCCCTCAGAG	TCACCTCCCT	ATCACCTCAC	TGTCCCTCTC
clone1	CCCTCAGTGT	CCCCTCAGAG	TCACCTCCCT	ATCACCTCAC	TGTCCCTCTC
	2751				2800
clone11	TGCCCTCTC	TGCCCTCTC	TGTCCCTCCC	TGCCCTCCC	CGTCCCTCT
clone1	TGCCCTCTC	TGCCCTCTC	TGTCCCTCTC	TGTCCCTCCC	CGTCCCTCT
	2801				2850
clone11	CTGTCCCTCT	CTGCCCTCA	CTGCTCCTCT	CTGCACCTCA	CTGCTCCTCA
clone1	CTGTCCCTCT	CTGCCCTCA	CTGCTCCTCT	CTGCACCTCA	CTGCTCCTCA
	2851				2900
clone11	CTGCCCTGGG	GGAGGCCCGC	ATCGAGGTGT	CTCTGCTCAC	CCCGTCCCC
clone1	CTGCCCTGGG	GGAGGCCCGC	ATCGAGGTGT	CTCTGCTCAC	CCCGTCCCC
	2901				2950
clone11	ACCCGTACC	CCCCGCCAGG			
clone1	ACCCGTCCC	CCCCGCCAGG			

Figure 2 (e)




Figure 7a




Figure 7b

Figure 8

tgacctacct accctgcca ggtcaggggt cctccaaggc
aagggatcac atggcaccac ctctcttgca gcctccacca agggcccatc ggtcttcccc
ctggcaccct cctccaagag cacctctggg ggcacagcgg ccctgggctg cctgggtcaag
gactacttcc ccgaaccggg gacgggtgctg tggaaactcag gcgccctgac cagcggcgtg
cacaccttcc cggctgtcct acagtcctca ggactctact ccctcagcag cgtggtgacc
gtgccctcca gcagcttggg caccagacc tacatctgca acgtgaatca caagcccagc
aacaccaagg tggacaagaa agttggtgag aggccagcac agggagggag ggtgtctgct
ggaagccagg ctcagcgcct ctgcctggac gcaccccggc tatgcagccc cagtccaggg
cagcaaggca ggccccgtct gcctcttcac ccggaggcct ctgcccggcc cactcatgct
cagggagagg gtcttctggc ttttcccca ggctctgggc aggcacagge taggtgcccc
taaccaggc cctgcacaca aaggggagag tgctgggctc agacctgcca agaccatata
ccgggaggac cctgcccctg acctaagccc accccaaagg ccaaactctc cactccctca
gctcggacac cttctctcct ccagattcc agtaactccc aatcttctct ctgcagagcc
caaatcttgt gacaaaactc acacatgccc accgtgcca ggtaagccag cccaggcctc
gccctccagc tcaaggcggg acaggtgccc tagagtagcc tgcacccagg gacaggcccc
agccgggtgc tgacacgtcc acctccatct ctctctcagc acctgaactc ctggggggac
cgtcagtctt cctcttcccc caaaaccca aggacaccct catgatctcc cggaccctg
aggtcacatg cgtggtggtg gacgtgagcc acgaagacc tggaggtcaag ttcaactggt
acgtggacgg cgtggagggtg cataatgcca agacaaagcc gcgggaggag cagtacaaca
gcacgtaccg tgtggtcagc gtcctcaccg tcctgcacca ggactggctg aatggcaagg
agtacaagtg caaggtctcc aacaaagccc tcccagcccc catcgagaaa accatctcca
aagccaaagg tgggaccctg ggggtgcgag ggccacatgg acagaggccg gctcggccca
ccctctgccc tgagagtgc cgctgtacca acctctgtcc ctacagggca gccccgagaa
ccacaggtgt acaccctgcc cccatcccgg gatgagctga ccaagaacca ggtcagcctg
acctgcctgg tcaaaggctt ctatcccagc gacatcgccg tggagtggga gagcaatggg
cagccggaga acaactaaa gaccagcct cccgtgctgg actccgacgg ctctctcttc
ctctacagca agctcaccgt ggacaagagc aggtggcagc aggggaacgt cttctcatgc
tccgtgatgc atgaggctct gcacaaccac tacacgcaga agagcctctc cctgtctccg
ggtaaatgag cgctgtgccg gcgagctgcc cctctccctc cccccacgc cgcagctgt.

Figure 9.

tgagtgcagc tgtcctgacc atgtcgtctg tgtttgagg tgtccagtgt
gaggtgcagc tgttggagtc cgggggaggt ctctgccagc caggggggac cctgagactc
acctgcgcag tctctggatt caccttcagt agctatgcaa tgagctgggt cggccaggct
ccaggggaagg ggctggaatg ggtcggagcc attagtggta gtggtagcac atactacgcg
gacagcgtga aaggccgatt caccatctcc agagacaact ccaagaacac gctgtatctg
caaatgaaca gtctgagagc cgaggacacg gccgcctatt actgtgcgaa agacacagtg
aggggccttc aggctgagcc cagacacaaa cctccctgca

Figure 10.

ggagatgtcc actggtacct aagcctcgcc atcctgttg cttctttcct catccacat ggcaccagg
ggctgcacca tctgtcttca tcttcccgcc atctgatgag cagttgaaat ctggaactgc
ctctgtttgt tgctgtctga ataacttcta tcccagagag gccaaaagtac agtggagggt
ggataacgcc ctccaatcgg gtaactccca ggagagtgtc acagagcagg acagcaagga
cagcacctac agcctcagca gcaccctgac gctgagcaaa gcagactacg agaaacacaa
agtctacgcc tgcgaagtca cccatcaggg cctgagctcg cccgtcacia agagcttcaa
caggggagag tgtttagagcg agacgcctgc cagggcaccg ccagcgacc tgagggccag
cctcgc

Figure 11.

catgcaggag gcagtaccag gcaggaccca gcatggacat gaggtccct gctcccaccc
tgggactcct gctgctctgg ctcccaggta aggagggaaa caacaaaaat tttattcagc
cagtgtagcc actaatgcct ggcacttcag gaaattcttc ttagaacatt actaatcatg
tggatatgtg ttttatggtt cctaatatca gataccagat gttacatcca gatgaccag
tctccatcct ctctgtctgc atctgtggga gacagagtca ccatcacttg ccgagccagt
cagggcatta gcaattactt agcctggat cagcagaaac cagggagggt tcccaagctc
ctgatttatg ctgcatccac tttgcaatct ggggtccat cgcggttcag tggcagtgga
tctgggacag atttactct taccatcagc agcctgcagc ctgaagatgt tgccacctat
tactgtcaaa agtacaacag tgcccctcca cttttcggcg gagggaccaa ggtggagatc
aaacgtaagt gcactttcct aatgttctc accgttctg cctgatttgt ttgcttttc
cattttttcgctat..

Figure 12.

catacacag ccatacatac gcgtgtggcc gctctgcctc tctcttgag gtcagcccaa
ggctgcccc tccgtcactc tgttcccgcc ctctctgag gagcttcaag ccaacaaggc
cacactggg tgtctcataa gtgacttcta cccgggagcc gtgacagtgg cttggaagc
agatagcag cccgtcaagg cgggagtgga gaccaccaca cctccaaac aaagcaaca
caagtacgg gccagcagct atctgagcct gacgcctgag cagtggaagt cccacagaag
ctacagctgc caggtcacgc atgaaggag caccgtggag aagacagtgg cccctacaga
atgttcatag tagtcccact ggggatgcaa tgtgaggaca gtggttcctc accctcctg

Figure 13.




Figure 14.

.ttgccgttt tctccctct ctctctccc tetccaggtt cctgggtgca gtcagtgctg actcagccgc
cctcgggtgc agcagccccg ggacaagaag tcacgatctc ctgctccggg tctagtagca acattggcga
taatttcgtc tottggtacC agcagctgcc tggcactgcc cctaagcttc tgatctatga taacaacAag
agaccctcgg gcatccctga ccgattctcc ggttccaat ccggcacctc agccacatta ggcactcactg
ggtccaac cggcgacgag gctgactatt actgtgggac ttgggacagc agcctttctg ttggtatgtt
tgggggcggg acacgcgtga ccgtcctagg tgagtccgtg acctcgtctc ggtctttctt ccccat...




Figure 15. Humanized chicken light chain locus.

PRODUCTION OF HUMANIZED ANTIBODIES IN TRANSGENIC ANIMALS

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This is a divisional application claiming the benefit of the priority of copending application Ser. No. 09/921,819, filed Aug. 3, 2001 under 35 U.S.C. 120, which is a non-provisional application filed under 37 CFR 1.53(b), claiming priority under 35 USC 119(e) to Provisional Application Ser. No. 60/222,872, filed on Aug. 3, 2000, and Provisional Application Ser. No. 60/276,156, filed on Mar. 15, 2001, the contents of which are incorporated herein by reference.

FIELD OF INVENTION

[0002] This invention relates to humanized antibodies produced from transgenic non-human animals. The non-human animals are genetically engineered to contain one or more humanized immunoglobulin loci which are capable of undergoing gene rearrangement and gene conversion in the transgenic non-human animals to produce diversified humanized immunoglobulins. The present invention further relates to novel sequences, recombination vectors and transgenic vectors useful for making these transgenic animals. The humanized antibodies of the present invention have minimal immunogenicity to humans and are appropriate for use in the therapeutic treatment of human subjects.

BACKGROUND OF THE INVENTION

[0003] The therapy of infectious diseases caused by bacteria, fungi, virus and parasites is largely based on chemotherapy. However, the emergence of drug-resistant organisms requires the continuous development of new antibiotics. Therapies of patients with malignancies and cancer are also based on chemotherapy. However, many of these therapies are ineffective and the mortality of diseased patients is high. For both infectious diseases and cancer, improved and innovative therapies are needed. Therapy of steroid resistant rejection of transplanted organs requires the use of biological reagents (monoclonal or polyclonal antibody preparations) that reverse the ongoing alloimmune response in the transplant recipient. The major problem of antibody preparations obtained from animals is the intrinsic immunogenicity of non-human immunoglobulins in human patients. In order to reduce the immunogenicity of non-human antibodies, genetic engineering of individual antibody genes in animals has been proposed. In particular, it has been shown that by fusing animal variable (V) region exons with human constant (C) region exons, a chimeric antibody gene can be obtained. However, this approach may only eliminate the immunogenicity caused by the non-human Fc region, while the remaining non-human Fab sequences may still be immunogenic. In another approach, human immunoglobulin genes for both, heavy and light chain immunoglobulins have been introduced into the genome of mice. While this genetic engineering approach resulted in the expression of human immunoglobulin polypeptides in genetically engineered mice, the level of human immunoglobulin expression is low. This may be due to species-specific regulatory elements in the immunoglobulin loci that are necessary for efficient expression of immunoglobulins. As demonstrated in transfected cell lines, regulatory elements present in human immunoglobulin genes may not function properly in non-human animals.

[0004] Several regulatory elements in immunoglobulin genes have been described. Of particular importance are enhancers downstream (3') of heavy chain constant regions and intronic enhancers in light chain genes. In addition, other, yet to be identified, control elements may be present in immunoglobulin genes. Studies in mice have shown that the membrane and cytoplasmic tail of the membrane form of immunoglobulin molecules play an important role in expression levels of human-mouse chimeric antibodies in the serum of mice homozygous for the human C γ 1 gene. Therefore, for the expression of heterologous immunoglobulin genes in animals it is desirable to replace sequences that contain enhancer elements and exons encoding transmembrane (M1 exon) and cytoplasmic tail (M2 exon) with sequences that are normally found in the animal in similar positions.

[0005] The introduction of human immunoglobulin genes into the genome of mice resulted in expression of a diversified human antibody repertoire in genetically engineered mice. In both mice and humans, antibody diversity is generated by gene rearrangement. This process results in the generation of many different recombined V(D)J segments encoding a large number of antibody molecules with different antigen binding sites. However, in other animals, like rabbits, pigs, cows and birds, antibody diversity is generated by a substantially different mechanism called gene conversion. For example, it is well established that in rabbit and chicken, VDJ rearrangement is very limited (almost 90% of immunoglobulin is generated with the 3'proximal VH1 element) and antibody diversity is generated by gene conversion and hypermutation. In contrast, mouse and human gene conversion occurs very rarely, if at all. Therefore, it is expected that in animals that diversify antibodies by gene conversion a genetic engineering approach based on gene rearrangement will result in animals with low antibody titers and limited antibody diversity. Thus, the genetic engineering of large animals for the production of non-immunogenic antibody preparations for human therapy requires alternative genetic engineering strategies.

Relevant Literature

[0006] The use of polyclonal antibody preparations for the treatment of transplant rejection was recently reviewed by N. Bonnefoy-Berard et al., *J Heart Lung Transplant* 1996; 15(5): 435-442; C. Colby et al., *Ann Pharmacother* 1996; 30(10):1164-1174; M. J. Dugan et al., *Ann Hematol* 1997; 75(1-2):41-46. The use of polyclonal antibody therapies for autoimmune diseases has been described by W. Cendrowski, *Boll Ist Sieroter Milan* 1997; 58(4):339-343; L. K. Kastrukoff et al., *Can J Neurol Sci* 1978; 5(2):175-178; J. E. Walker et al., *J Neurol Sci* 1976; 29(24):303-309. The depletion of fat cells using antibody preparations has been described by L. De Clercq et al., *J Anim Sci* 1997; 75(7):1791-1797; J. T. Wright et al., *Obes Res* 1995; 3(3):265-272.

[0007] Regulatory elements in immunoglobulin genes have been described by Bradley et al. (1999), *Transcriptional enhancers and the evolution of the IgH locus*; Lauster, R. et al., *Embo J* 12: 4615-23 (1993); Volgina et al., *J Immunol* 165:6400 (2000); Hole et al., *J Immunol* 146:4377 (1991).

[0008] Antibody diversification by gene conversion in chicken and rabbit has been described by Bucchini et al.,

Nature 326: 409-11 (1987); Knight et al., *Advances in Immunology* 56: 179-218 (1994); Langman et al., *Res Immunol* 144: 422-46 (1993). The generation of mice expressing human-mouse chimeric antibodies has been described by Pluschke et al., *Journal of Immunological Methods* 215: 27-37 (1998). The generation of mice expressing human-mouse chimeric antibodies with mouse derived membrane and cytoplasmic tails has been described by Zou et al., *Science* 262: 1271-1274 (1993); Zou et al. *Curr Biol* 4: 1099-1103. The generation of mice expressing human immunoglobulin polypeptides has been described by Bruggemann et al. *Curr Opin Biotechnol* 8(4): 455-8 (1997); Lonberg et al. *Int Rev Immunol* 13(1):65-93 (1995); Neuberger et al., *Nature* 338: 350-2 (1989). Generation of transgenic mice using a BAC clone has been described by Yang et al., *Nat Biotechnol* 15: 859-65 (1997).

[0009] The generation of transgenic rabbits has been described by Fan, J. et al., *Pathol Int* 49: 583-94 (1999); Brem et al., *Mol Reprod Dev* 44: 56-62 (1996). Nuclear transfer cloning of rabbits has been described by Stice et al., *Biology of Reproduction* 39: 657-664 (1988). Rabbits with impaired immunoglobulin expression have been described by McCartney-Francis et al., *Mol Immunol* 24: 357-64 (1987); Allegrucci, et al., *Eur J Immunol* 21: 411-7 (1991).

[0010] The production of transgenic chicken has been described by Etches et al., *Methods in Molecular Biology* 62: 433-450; Pain et al., *Cells Tissues Organs* 1999; 165(3-4): 212-9; Sang, H., "Transgenic chickens—methods and potential applications", *Trends Biotechnol* 12:415 (1994); and in WO 200075300, "Introducing a nucleic acid into an avian genome, useful for transfecting avian blastodermal cells for producing transgenic avian animals with the desired genes, by directly introducing the nucleic acid into the germinal disc of the egg".

[0011] Agammaglobulinemic chicken have been described by Frommel et al., *J Immunol* 105(1): 1-6 (1970); Benedict et al., *Adv Exp Med Biol* 1977; 88(2): 197-205.

[0012] The cloning of animals from cells has been described by T. Wakayama et al., *Nature* 1998; 394:369-374; J. B. Cibelli et al., *Science* 280:1256-1258 (1998); J. B. Cibelli et al., *Nature Biotechnology* 1998; 16:642-646; A. E. Schnieke et al., *Science* 278: 2130-2133 (1997); K. H. Campbell et al., *Nature* 380: 64-66 (1996).

[0013] Production of antibodies from transgenic animals is described in U.S. Pat. Nos. 5,814,318, 5,545,807 and 5,570,429. Homologous recombination for chimeric mammalian hosts is exemplified in U.S. Pat. No. 5,416,260. A method for introducing DNA into an embryo is described in U.S. Pat. No. 5,567,607. Maintenance and expansion of embryonic stem cells is described in U.S. Pat. No. 5,453,357.

[0014] The mechanisms involved in the diversification of the antibody repertoire in pigs, sheep and cows are reviewed in Butler, J. E. (1998), "Immunoglobulin diversity, B-cell and antibody repertoire development in large farm animals", *Rev Sci Tech* 17:43. Antibody diversification in sheep is described in Reynaud, C. A., C. Garcia, W. R. Hein, and J. C. Weill (1995), "Hypermutation generating the sheep immunoglobulin repertoire is an antigen-independent process", *Cell* 80:115; and Dufour, V., S. Malinge, and F. Nau. (1996), "The sheep Ig variable region repertoire consists of a single VH family", *J Immunol* 156:2163.

SUMMARY OF THE INVENTION

[0015] One embodiment of the present invention provides humanized antibodies (humanized immunoglobulins) having at least a portion of a human immunoglobulin polypeptide sequence.

[0016] The humanized antibodies of the present invention are made from transgenic non-human animals genetically engineered to contain one or more humanized Ig loci.

[0017] Preferably, the humanized antibodies of the present invention are prepared from transgenic non-human animals which generate antibody diversity primarily by gene conversion and hypermutation, e.g., rabbit, pigs, chicken, sheep, cow and horse. The antibodies can be made by immunizing transgenic animals with a desired antigen such as an infectious agent (e.g., bacteria or viruses) or parts or fragments thereof.

[0018] Such humanized antibodies have reduced immunogenicity to primates, especially humans, as compared to non-humanized antibodies prepared from non-human animals. Therefore, the humanized antibodies of the present invention are appropriate for use in the therapeutic treatment of human subjects.

[0019] Another embodiment of the present invention provides a preparation of humanized antibodies which can be monoclonal antibodies or polyclonal antibodies. Preferred antibody preparations of the present invention are polyclonal antibody preparations which, according to the present invention, have minimal immunogenicity to primates, especially humans.

[0020] A preferred preparation of polyclonal antibodies is composed of humanized immunoglobulin molecules having at least a heavy chain or light chain constant region polypeptide sequence encoded by a human constant region gene segment. More preferably, the variable domains of the heavy chains or light chains of the immunoglobulins molecules are also encoded by human gene segments.

[0021] In another embodiment, the present invention provides pharmaceutical compositions which include a preparation of humanized antibodies, and a pharmaceutically-acceptable carrier.

[0022] Another embodiment of the present invention provides novel sequences from the 5' and 3' flanking regions of the Ig gene segments of non-human animals, preferably, animals which rely primarily on gene conversion in generating the antibody diversity. In particular, the present invention provides novel nucleotide sequences downstream (3', 3-prime) of the genes coding for C λ in chickens, C γ and C ϵ in rabbits, C γ 1,2,3 in cows and C γ 1,2 in sheep, as well as novel sequences 5' of rabbit C γ .

[0023] In another embodiment, the present invention provides recombination vectors useful for replacing an Ig gene segment of a non-human animal with the corresponding human Ig gene segment. These vectors include a human Ig gene segment which is linked to flanking sequences at the 5' end and the 3' end, wherein the flanking sequences are homologous to the flanking sequences of the target animal Ig gene segment.

[0024] Preferred recombination vectors are those useful for the replacement of the animal's Ig constant region. For

example, recombination vectors useful for replacing the rabbit heavy chain constant region genes are provided. A preferred vector contains from 5' to 3', a nucleotide sequence as set forth in SEQ ID NO: 12 or SEQ ID NO: 13, or a portion of SEQ ID NO: 12 or SEQ ID NO: 13, a human heavy chain constant region gene segment, a nucleotide sequence as set forth in SEQ ID NO: 10 or a portion of or SEQ ID NO: 10. Another preferred vector contains a nucleotide sequence as set forth in SEQ ID NO: 51, which sequence is characterized as having a human C γ 1 gene linked to flanking sequences from the 5' and 3' flanking regions of a rabbit heavy chain constant region gene.

[0025] Recombination vectors are also provided useful for replacing the rabbit light chain constant region genes. A preferred vector contains a nucleotide sequence as set forth in SEQ ID NO: 53, which sequence is characterized as having a human C κ linked to flanking sequences from the 5' and 3' flanking regions of the rabbit light chain C γ 1 gene.

[0026] Other recombination vectors are provided which are useful for replacing the chicken light chain constant region genes. A preferred vector contains a nucleotide sequence as set forth in SEQ ID NO: 57 which is characterized as having a human C λ 2 linked to flanking sequences from the 5' and 3' flanking regions of the chicken light chain C λ gene.

[0027] Other recombination vectors provided include those useful for replacing the animal's Ig V region elements. For example, a recombination vector useful for replacing a rabbit heavy chain V region element is provided and contains SEQ ID NO: 52. A recombination vector useful for replacing a rabbit light chain V region element is provided and contains SEQ ID NO: 54.

[0028] In still another embodiment, the present invention provides transgenic constructs or vectors containing at least one humanized Ig locus, i.e., an Ig locus from a non-human animal or a portion of an Ig locus from a non-human animal wherein the locus or the portion of a locus is genetically modified to contain at least one human Ig gene segment. Such humanized Ig locus has the capacity to undergo gene rearrangement and gene conversion in the non-human animal thereby producing a diversified repertoire of humanized immunoglobulins.

[0029] One humanized Ig locus provided by the invention is a humanized heavy chain locus which includes one or more V gene segments, one or more D gene segments, one or more J gene segments, and one or more constant region gene segments, wherein at least one gene segment is a human heavy chain gene segment. The gene segments in the humanized heavy chain locus are juxtaposed with respect to each other in an unrearranged, or partially or fully rearranged configuration. A preferred humanized heavy chain locus contains a human constant region gene segment, preferably, C α or C γ . A more preferred humanized locus contains multiple V gene segments and at least one human V gene segment, in addition to a human heavy chain constant region segment. The human V gene segment is placed downstream of the non-human V gene segments.

[0030] Another humanized Ig locus is a humanized light chain locus which includes one or more V gene segments, one or more J gene segments, and one or more constant region gene segments, wherein at least one gene segment is

a human light chain gene segment. The gene segments in the humanized light chain locus are juxtaposed with respect to each other in an unrearranged or rearranged configuration. A preferred humanized light chain locus contains a human constant region gene segment, preferably, C λ or C κ . More preferably, the humanized light chain locus further contains multiple V gene segments and at least one human V gene segment. The human V gene segment is placed downstream of the non-human V gene segments. Even more preferably, the humanized light chain locus includes a rearranged human VJ segment, placed downstream of a number of (e.g., 10-100) VL gene segments of either non-human or human origin.

[0031] Another embodiment of the present invention is directed to methods of making a transgenic vector containing a humanized Ig locus by isolating an Ig locus or a portion of an Ig locus from a non-human animal, and integrating the desired human Ig gene segment(s) into the isolated animal Ig locus or the isolated portion of an Ig locus. The human Ig gene segment(s) are integrated into the isolated animal Ig locus or the isolated portion of an Ig locus by ligation or homologous recombination in such a way as to retain the capacity of the locus for undergoing effective gene rearrangement and gene conversion in the non-human animal. Integration of a human Ig gene segment by homologous recombination can be accomplished by using the recombination vectors of the present invention.

[0032] In another embodiment, the present invention provides methods of making transgenic animals capable of producing humanized antibodies. The transgenic animals can be made by introducing a transgenic vector containing a humanized Ig locus, or a recombination vector containing a human Ig gene segment, into a recipient cell or cells of an animal, and deriving an animal from the genetically modified recipient cell or cells.

[0033] Transgenic animals containing one or more humanized Ig loci, and cells derived from such transgenic animals (such as B cells from an immunized transgenic animal) are also provided. The transgenic animals of the present invention are capable of gene rearranging and gene converting the transgenic humanized Ig loci to produce a diversified repertoire of humanized immunoglobulin molecules.

BRIEF DESCRIPTION OF THE DRAWINGS

[0034] FIG. 1. Cow C γ 3' flanking sequences. Primers are shown in shaded boxes. The 5' primer is in CH3, and the 3' primer is in M1. The sequences of clone 11, clone 3, and clone 5 are set forth in SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 5, respectively.

[0035] FIG. 2. Sheep C γ 3' flanking sequences. Primers are shown in shaded boxes. The 5' primer is in CH3, and the 3' primer is in M2. The sequences of clone 11 and clone 1 are set forth in SEQ ID NO: 8 and SEQ ID NO: 9, respectively.

[0036] FIG. 3. A novel 3' flanking sequence (SEQ ID NO: 10) of the rabbit C γ gene.

[0037] FIG. 4. A novel nucleotide sequence (SEQ ID NO: 11) 3' of the rabbit C κ gene.

[0038] FIG. 5. Novel nucleotide sequences (SEQ ID NO: 12 and SEQ ID NO: 13) 5' of the rabbit C γ gene. The

sequences between SEQ ID NO: 12 and SEQ ID NO: 13 (a gap of about 1000 nt) remain to be determined.

[0039] FIG. 6. Comparison of human, mouse, rabbit, sheep, cow and camel sequences for the M1 and M2 regions 3' of the Cgamma gene.

[0040] FIG. 7a. DNA construct for the replacement of rabbit Cκ with human Cκ. A 0.5 kb fragment containing a DNA sequence encoding human Cλ is flanked by sequences from the rabbit Cκ1 gene. The upstream sequence (5'Cκ) is 2.8 kb, the downstream sequence (3'Cκ) is 2.6 kb. The vector also contains a lox-neo cassette for positive selection and a Hsv-Tk cassette for negative selection.

[0041] FIG. 7b. DNA construct for the replacement of rabbit Cγ with human Cγ1. A 1.8 kb fragment containing a DNA sequence encoding human Cγ1 is flanked by sequences from the rabbit Cγ gene. The upstream sequence (5'Cγ) is 1.9 kb, the downstream sequence (3'Cγ) is 3.1 kb. The vector also contains a lox-neo cassette for positive selection and a Hsv-Tk cassette for negative selection. The figure is not up to scale.

[0042] FIG. 8. DNA fragment (SEQ ID NO: 51) containing a human immunoglobulin heavy chain Cγ1 gene segment flanked by 50 nucleotides derived from the flanking regions of rabbit Cγ gene. Flanking sequences derived from the flanking regions of rabbit Cγ gene are underlined.

[0043] FIG. 9. DNA fragment (SEQ ID NO: 52) containing a V gene segment with more than 80% sequence identity with rabbit V elements and encoding a human V element polypeptide sequence. Flanking sequences derived from the flanking regions of rabbit VH1 and J genes are underlined.

[0044] FIG. 10. DNA fragment (SEQ ID NO: 53) containing a human immunoglobulin heavy chain Cκ gene segment flanked by 50 nucleotides derived from the rabbit light chain immunoglobulin Kappal gene. Flanking sequences derived from the flanking regions of rabbit Cκ gene are underlined.

[0045] FIG. 11. DNA fragment (SEQ ID NO: 54) containing a V gene segment with more than 80% sequence identity with rabbit V elements and encoding a human V element polypeptide sequence. Flanking sequences derived from the flanking regions of rabbit immunoglobulin V and J genes are underlined.

[0046] FIG. 12. DNA fragment (SEQ ID NO: 57) containing a gene encoding human immunoglobulin light chain constant region Clambda2 flanked by 50 nucleotides (underlined) derived from the flanking sequences of chicken Clambda gene.

[0047] FIG. 13. Modification of the chicken light chain locus using the ET system. A chicken genomic BAC clone with the full-length light chain locus was modified by homologous recombination. In a first step Cλ was deleted by insertion of a selection cassette which was in a second homologous recombination step exchanged against the human Cλ gene.

[0048] FIG. 14. DNA fragment (SEQ ID NO: 58) containing a VJ gene segment with 80% sequence identity with chicken V gene segments and encoding a human VJ immunoglobulin polypeptide. Flanking sequences derived from the flanking regions of chicken immunoglobulin V and J genes are underlined.

[0049] FIG. 15. Modified chicken light chain locus.

DETAILED DESCRIPTION OF THE INVENTION

[0050] One embodiment of the present invention provides humanized immunoglobulins (antibodies).

[0051] By “a humanized antibody” or “a humanized immunoglobulin” is meant an immunoglobulin molecule having at least a portion of a human immunoglobulin polypeptide sequence (or a polypeptide sequence encoded by a human Ig gene segment). The humanized immunoglobulin molecules of the present invention can be isolated from a transgenic non-human animal engineered to produce humanized immunoglobulin molecules. Such humanized immunoglobulin molecules are less immunogenic to primates, especially humans, relative to non-humanized immunoglobulin molecules prepared from the animal or prepared from cells derived from the animal.

[0052] The term “non-human animals” as used herein includes, but is not limited to, rabbits, pigs, birds (e.g., chickens, turkeys, ducks, geese and the like), sheep, goats, cows and horses. Preferred non-human animals are those animals which rely primarily on gene conversion and/or somatic hypermutation to generate antibody diversity, e.g., rabbit, pigs, birds (e.g., chicken, turkey, duck, goose and the like), sheep, goat, and cow. Particularly preferred non-human animals are rabbit and chicken.

[0053] In animals such as human and mouse, there are multiple copies of V, D and J gene segments on the heavy chain locus, and multiple copies of V and J gene segments on a light chain locus. Antibody diversity in these animals is generated primarily by gene rearrangement, i.e., different combinations of gene segments to form rearranged heavy chain variable region and light chain variable region. In other animals (e.g., rabbit, chicken, sheep, goat, and cow), however, gene rearrangement does not play a significant role in the generation of antibody diversity. For example, in rabbit, only a very limited number of the V gene segments, most often the V gene segments at the 3' end of the V-region, are used in gene rearrangement to form a contiguous VDJ segment. In chicken, only one V gene segment (the one adjacent to the D region, or “the 3' proximal V gene segment”), one D segment and one J segment are used in the heavy chain rearrangement; and only one V gene segment (the 3' proximal V segment) and one J segment are used in the light chain rearrangement. Thus, in these animals, there is little diversity among initially rearranged variable region sequences resulting from junctional diversification.

[0054] Further diversification of the rearranged Ig genes is achieved by gene conversion, a process in which short sequences derived from the upstream V gene segments replace short sequences within the V gene segment in the rearranged Ig gene.

[0055] The term “Ig gene segment” as used herein refers to segments of DNA encoding various portions of an Ig molecule, which are present in the germline of animals and humans, and which are brought together in B cells to form rearranged Ig genes. Thus, Ig gene segments as used herein include V gene segments, D gene segments, J gene segments and C region gene segments.

[0056] The term “human Ig gene segment” as used herein includes both naturally occurring sequences of a human Ig gene segment, degenerate forms of naturally occurring

sequences of a human Ig gene segment, as well as synthetic sequences that encode a polypeptide sequence substantially identical to the polypeptide encoded by a naturally occurring sequence of a human Ig gene segment. By "substantially" is meant that the degree of amino acid sequence identity is at least about 85%-95%.

[0057] A preferred humanized immunoglobulin molecule of the present invention contains at least a portion of a human heavy or light chain constant region polypeptide sequence. A more preferred immunoglobulin molecule contains at least a portion of a human heavy or light chain constant region polypeptide sequence, and at least a portion of a human variable domain polypeptide sequence.

[0058] In another embodiment of the present invention, a preparation of humanized antibodies is provided.

[0059] By "a preparation of humanized antibodies" or "a humanized antibody preparation" is meant an isolated antibody product or a purified antibody product prepared from a transgenic non-human animal (e.g., serum, milk, or egg yolk of the animal) or from cells derived from a transgenic non-human animal (e.g., a B-cell or a hybridoma cell).

[0060] A humanized antibody preparation can be a preparation of polyclonal antibodies, which includes a repertoire of humanized immunoglobulin molecules. A humanized antibody preparation can also be a preparation of a monoclonal antibody.

[0061] Although the immunogenicity to humans of a humanized monoclonal antibody preparation is also reduced as compared to a non-humanized monoclonal antibody preparation, humanized polyclonal antibody preparations are preferred embodiments of the present invention. It has been recognized that humanized monoclonal antibodies still invoke some degree of an immune response (an anti-idiotypic response) in primates (e.g., humans) when administered repeatedly in large quantities because of the unique and novel idiotype of the monoclonal antibody. The present inventors have uniquely recognized that the overall immunogenicity of polyclonal antibodies is less dependent on an anti-idiotypic response. For example, polyclonal antibodies made from non-human animals with only the constant region elements humanized (e.g., polyclonal antibodies having constant regions encoded by human gene segments, and having variable domains encoded by the endogenous genes of the non-human animal), are substantially non-immunogenic to primates.

[0062] Without intending to be bound to any theory, the present inventors have proposed that the reduced immunogenicity of such a humanized polyclonal antibody preparation is due to the fact that the preparation contains a very large number of different antibodies with many different idiotypes which are to a large extent defined by novel amino acid sequences in the complementarity determining regions (CDR) of the heavy and light chain. Therefore, upon administration of such preparation into a primate such as a human, the administered amount of each individual immunoglobulin molecule in the preparation may be too low to solicit immune response against each immunoglobulin molecule. Thus, the humanized polyclonal antibody preparation which has many different idiotypes and variable regions has minimal immunogenicity to a recipient, even if the antibodies in the polyclonal antibody preparation are all directed to the

same antigen. To further reduce any potential residual immunogenicity, a humanized polyclonal antibody preparation may be prepared which is composed of immunoglobulin molecules having both the variable domains and the constant regions encoded by human Ig gene segments.

[0063] In a preferred embodiment, the present invention provides an antibody preparation which includes humanized immunoglobulin molecules having at least a portion of a human heavy or light chain constant region polypeptide sequence. More preferably, the humanized immunoglobulins in the antibody preparation of the present invention further contain at least a portion of a human variable domain polypeptide sequence, in addition to at least a portion of a human constant region polypeptide sequence.

[0064] Preferred humanized antibody preparations of the present invention are composed of humanized antibodies made from transgenic non-human animals whose antibody diversity is generated primarily by gene conversion, such as rabbit, birds (e.g., chicken, turkey, duck, goose and the like), sheep, goat, and cow; preferably, rabbit and chicken.

[0065] Once a transgenic non-human animal capable of producing diversified humanized immunoglobulin molecules is made (as further set forth below), humanized immunoglobulins and humanized antibody preparations against an antigen can be readily obtained by immunizing the animal with the antigen. A variety of antigens can be used to immunize a transgenic host animal. Such antigens include, microorganism, e.g. viruses and unicellular organisms (such as bacteria and fungi), alive, attenuated or dead, fragments of the microorganisms, or antigenic molecules isolated from the microorganisms.

[0066] Preferred bacterial antigens for use in immunizing an animal include purified antigens from *Staphylococcus aureus* such as capsular polysaccharides type 5 and 8, recombinant versions of virulence factors such as alpha-toxin, adhesin binding proteins, collagen binding proteins, and fibronectin binding proteins. Preferred bacterial antigens also include an attenuated version of *S. aureus*, *Pseudomonas aeruginosa*, *enterococcus*, *enterobacter*, and *Klebsiella pneumoniae*, or culture supernatant from these bacteria cells. Other bacterial antigens which can be used in immunization include purified lipopolysaccharide (LPS), capsular antigens, capsular polysaccharides and/or recombinant versions of the outer membrane proteins, fibronectin binding proteins, endotoxin, and exotoxin from *Pseudomonas aeruginosa*, *enterococcus*, *enterobacter*, and *Klebsiella pneumoniae*.

[0067] Preferred antigens for the generation of antibodies against fungi include attenuated version of fungi or outer membrane proteins thereof, which fungi include, but are not limited to, *Candida albicans*, *Candida parapsilosis*, *Candida tropicalis*, and *Cryptococcus neoformans*.

[0068] Preferred antigens for use in immunization in order to generate antibodies against viruses include the envelop proteins and attenuated versions of viruses which include, but are not limited to respiratory syncytial virus (RSV) (particularly the F-Protein), Hepatitis C virus (HCV), Hepatitis B virus (HBV), cytomegalovirus (CMV), EBV, and HSV.

[0069] Therapeutic antibodies can be generated for the treatment of cancer by immunizing transgenic animals with

isolated tumor cells or tumor cell lines; tumor-associated antigens which include, but are not limited to, Her-2-neu antigen (antibodies against which are useful for the treatment of breast cancer); CD20, CD22 and CD53 antigens (antibodies against which are useful for the treatment of B cell lymphomas), (3) prostate specific membrane antigen (PMSA) (antibodies against which are useful for the treatment of prostate cancer), and 17-1A molecule (antibodies against which are useful for the treatment of colon cancer).

[0070] The antigens can be administered to a transgenic host animal in any convenient manner, with or without an adjuvant, and can be administered in accordance with a predetermined schedule.

[0071] After immunization, serum or milk from the immunized transgenic animals can be fractionated for the purification of pharmaceutical grade polyclonal antibodies specific for the antigen. In the case of transgenic birds, antibodies can also be made by fractionating egg yolks. A concentrated, purified immunoglobulin fraction may be obtained by chromatography (affinity, ionic exchange, gel filtration, etc.), selective precipitation with salts such as ammonium sulfate, organic solvents such as ethanol, or polymers such as polyethyleneglycol.

[0072] For making a monoclonal antibody, spleen cells are isolated from the immunized transgenic animal and used either in cell fusion with transformed cell lines for the production of hybridomas, or cDNAs encoding antibodies are cloned by standard molecular-biology techniques and expressed in transfected cells. The procedures for making monoclonal antibodies are well established in the art. See, e.g., European Patent Application 0 583 980 A1 ("Method For Generating Monoclonal Antibodies From Rabbits"), U.S. Pat. No. 4,977,081 ("Stable Rabbit-Mouse Hybridomas And Secretion Products Thereof"), WO 97/16537 ("Stable Chicken B-cell Line And Method of Use Thereof"), and EP 0 491 057 B1 ("Hybridoma Which Produces Avian Specific Immunoglobulin G"), the disclosures of which are incorporated herein by reference. In vitro production of monoclonal antibodies from cloned cDNA molecules has been described by Andris-Widhopf et al., "Methods for the generation of chicken monoclonal antibody fragments by phage display", *J Immunol Methods* 242:159 (2000), and by Burton, D. R., "Phage display", *Immunotechnology* 1:87 (1995), the disclosures of which are incorporated herein by reference.

[0073] In a further embodiment of the present invention, purified monoclonal or polyclonal antibodies are admixed with an appropriate pharmaceutical carrier suitable for administration in primates especially humans, to provide pharmaceutical compositions.

[0074] Pharmaceutically acceptable carriers which can be employed in the present pharmaceutical compositions can be any and all solvents, dispersion media, isotonic agents and the like. Except insofar as any conventional media, agent, diluent or carrier is detrimental to the recipient or to the therapeutic effectiveness of the antibodies contained therein, its use in the pharmaceutical compositions of the present invention is appropriate. The carrier can be liquid, semi-solid, e.g. pastes, or solid carriers. Examples of carriers include oils, water, saline solutions, alcohol, sugar, gel, lipids, liposomes, resins, porous matrices, binders, fillers, coatings, preservatives and the like, or combinations thereof

[0075] The present invention is further directed to novel nucleotide sequences and vectors, as well as the use of the

sequences and vectors in making a transgenic non-human animal which produces humanized immunoglobulins.

[0076] In general, the genetic engineering of a non-human animal involves the integration of one or more human Ig gene segments into the animal's genome to create one or more humanized Ig loci. It should be recognized that, depending upon the approach used in the genetic modification, a human Ig gene segment can be integrated at the endogenous Ig locus of the animal (as a result of targeted insertion, for example), or at a different locus of the animal. In other words, a humanized Ig locus can reside at the chromosomal location where the endogenous Ig locus of the animal ordinarily resides, or at a chromosomal location other than where the endogenous Ig locus of the animal ordinarily resides. Regardless of the chromosomal location, a humanized Ig locus of the present invention has the capacity to undergo gene rearrangement and gene conversion in the non-human animal thereby producing a diversified repertoire of humanized immunoglobulin molecules. An Ig locus having the capacity to undergo gene rearrangement and gene conversion is also referred to herein as a "functional" Ig locus, and the antibodies with a diversity generated by a functional Ig locus are also referred to herein as "functional" antibodies or a "functional" repertoire of antibodies.

[0077] In one embodiment, the present invention provides novel sequences useful for creating a humanized Ig locus and making transgenic animals capable of producing humanized immunoglobulin molecules. In particular, the present invention provides sequences from the 5' and 3' flanking regions of the Ig gene segments of non-human animals, preferably, animals which rely primarily on gene conversion in generating antibody diversity (e.g., rabbit, pigs, sheep, goat, cow, birds such as chicken, turkey, duck, goose, and the like).

[0078] The 5' and 3' flanking regions of the genes coding for the constant region are particularly important as these sequences contain untranslated regulatory elements (e.g., enhancers) critical for high Ig expression in the serum. The 3' flanking region of the genes coding for the constant region of the heavy chain also contain exons coding for the membranous and cytoplasmic tail of the membrane form of immunoglobulin (Volgina et al. *J Immunol* 165:6400, 2000). It has been previously established that the membrane and cytoplasmic tail of the membrane form of antibodies are critical in achieving a high level of expression of the antibodies in mice sera (Zou et al., *Science* 262:1271, 1993). Thus, the identification of the flanking sequences permits the replacement of exons and intervening introns of the C γ gene with the human equivalent, and the maintenance of the endogenous exons encoding the transmembrane and cytoplasmic tail regions as well as the endogenous non-coding enhancer sequences.

[0079] In one embodiment, the present invention provides 3' flanking sequences of heavy chain constant regions of non-human animals. More particularly, nucleotide sequences downstream (3', 3-prime) of the genes coding for rabbit C γ , cow C γ 1,2,3, and sheep C γ 1,2 are provided. Especially preferred nucleotide sequences include SEQ ID NO: 10 (3' of rabbit C γ), SEQ ID NOS: 3-5 (3' of cow C γ 1,2,3), and SEQ ID NOS: 8-9 (3' of sheep C γ 1,2).

[0080] In another embodiment, the present invention provides 3' flanking sequences of light chain constant regions of

non-human animals. More particularly, the present invention provides nucleotide sequences downstream (3', 3-prime) of the genes coding for C κ in rabbits. Especially preferred nucleotide sequences include SEQ ID NO: 11 (3' of rabbit C κ).

[0081] In still another embodiment, the present invention provides 5' flanking sequences of heavy chain constant regions of non-human animals. More particularly, nucleotide sequences upstream (5', 5-prime) of the rabbit C γ gene are provided. Especially preferred sequences include SEQ ID NO: 12 and SEQ ID NO: 13.

[0082] Another embodiment of the present invention provides 5' flanking sequences of light chain constant regions of non-human animals.

[0083] Portions of the above novel flanking sequences are provided by the present invention. By "a portion" is meant a fragment of a flanking nucleotide sequence capable of mediating homologous recombination between the human Ig gene segment and the target animal Ig gene segment. Generally, a portion is at least about 200 base pairs, preferably, at least about 400 base pairs, for recombination in animal cells such as ES cells or fibroblasts, and at least about 40 base pairs, preferably at least about 50 base pairs, for recombination in *E. coli*. Examples of portions of the above novel flanking-sequences include SEQ ID NOS: 59-60, 61-62, 63-64, 65-66, 67-68 and 69-70 (represented by the underlined sequences in FIGS. 8-12 and 14, respectively).

[0084] In a further aspect, the present invention provides vectors useful for the replacement of an Ig gene segment of a non-human animal with the corresponding human Ig gene segment. These vectors, also referred to herein as "recombination vectors", include a human Ig gene segment which is linked to flanking sequences at the 5' end and the 3' end, wherein the flanking sequences have a degree of homology with the flanking sequences of the target animal Ig gene segment sufficient to mediate homologous recombination between the human gene and the animal gene segments. Generally, at least about 200 bases should be identical between the flanking regions in a recombination vector and the flanking regions of the target gene to achieve efficient homologous recombination in animal cells such as ES cells and fibroblasts; and at least about 40 bases should be identical to achieve efficient homologous recombination in *E. coli*.

[0085] Recombination vectors useful for replacing the animal's immunoglobulin heavy chain constant region genes are provided, which contain from 5' to 3', a nucleotide sequence homologous to the 5' flanking region of the target animal heavy chain constant region gene, a human heavy chain constant region gene (e.g., human C γ 1), and a nucleotide sequence homologous to the 3' flanking region of the target animal heavy chain constant region gene.

[0086] Preferred recombination vectors are provided for the replacement of the rabbit heavy chain constant region genes. One such vector contains from 5' to 3', a nucleotide sequence as set forth in SEQ ID NO: 12 or SEQ ID NO: 13 or a portion thereof, a human heavy chain constant region gene segment, a nucleotide sequence as set forth in SEQ ID NO: 10 or a portion of or SEQ ID NO: 10. Another such vector contains SEQ ID NO: 51 (FIG. 8) which is characterized as having a human C γ 1 gene linked to flanking

sequences from the 5' and 3' flanking regions of a rabbit heavy chain constant region gene.

[0087] Recombination vectors are also provided which are useful for replacing the animal's immunoglobulin light chain constant region genes. Such vectors contain from 5' to 3', a nucleotide sequence homologous to the 5' flanking region of the target light chain constant region gene, a human light chain constant region gene (e.g., human C λ), and a nucleotide sequence homologous to the 3' flanking region of the target light chain constant region gene.

[0088] Preferred vectors include those useful for replacing the rabbit light chain constant region genes. A preferred vector contains a nucleotide sequence as set forth in SEQ ID NO: 53, which sequence is characterized as having a human C κ linked to flanking sequences from the 5' and 3' flanking regions of the rabbit light chain C κ 1 gene.

[0089] Other recombination vectors provided include those useful for replacing the animal's Ig V region elements. For example, a recombination vector useful for replacing a rabbit heavy chain V region element is provided and contains SEQ ID NO: 52. A recombination vector useful for replacing a rabbit light chain V region element is provided and contains SEQ ID NO: 54.

[0090] The recombination vectors of the present invention can include additional sequences that facilitate the selection of cells which have undergone a successful recombination event. For example, marker genes coding for resistance to neomycin, bleomycin, puromycin and the like can be included in the recombination vectors to facilitate the selection of cells which have undergone a successful recombination event.

[0091] In a further aspect of the present invention, transgenic constructs or vectors carrying one or more humanized Ig loci are provided.

[0092] In one embodiment, the present invention provides transgenic constructs containing a humanized Ig heavy chain locus which includes one or more V gene segments, one or more D gene segments, one or more J gene segments, and one or more constant region gene segments, wherein at least one gene segment is a human heavy chain gene segment. The gene segments in such humanized heavy chain locus are juxtaposed with respect to each other in an unrearranged configuration (or "the germline configuration"), or in a partially or fully rearranged configuration. The humanized heavy chain locus has the capacity to undergo gene rearrangement (if the gene segments are not fully rearranged) and gene conversion in the non-human animal thereby producing a diversified repertoire of heavy chains having human polypeptide sequences, or "humanized heavy chains".

[0093] In a preferred embodiment, the humanized heavy chain locus contains at least one C-region gene segment that is a human constant region gene segment, preferably, C α or C γ (including any of the C γ subclasses 1, 2, 3 and 4).

[0094] In another more preferred embodiment, the humanized heavy chain locus of the transgene contains a humanized V-region and a humanized C-region, i.e., a V-region having at least one human VH gene segment and a C-region having at least one human C gene segment (e.g., human C α or C γ).

[0095] Preferably, the humanized V-region includes at least about 10-100 heavy chain V (or "VH") gene segments, at least one of which is a human VH gene segment. In accordance with the present invention, the human VH gene segment included in the transgene shares at least about 75% to about 85% homology to the VH gene segments of the host animal, particularly those animal VH gene segments included in the upstream region of the transgene. As described above, a human VH segment encompasses naturally occurring sequences of a human VH gene segment, degenerate forms of naturally occurring sequences of a human VH gene segment, as well as synthetic sequences that encode a polypeptide sequence substantially (i.e., at least about 85%-95%) identical to a human heavy chain V domain polypeptide.

[0096] Preferably, the human VH gene segment(s) is placed downstream of the non-human VH segments in the transgene locus. Preferably, the non-human VH gene segments in the transgene are the VH gene segments from the 3' VH-region in the Ig locus of the host animal, including the 3' proximal VH1.

[0097] In another embodiment, the present invention provides transgenic constructs containing a humanized light chain locus capable of undergoing gene rearrangement and gene conversion in the host animal thereby producing a diversified repertoire of light chains having human polypeptide sequences, or "humanized light chains".

[0098] The humanized light locus includes one or more V gene segments, one or more J gene segments, and one or more constant region gene segments, wherein at least one gene segment is a human light chain gene segment. The gene segments in the humanized light chain locus are juxtaposed in an unrearranged configuration (or "the germline configuration"), or fully rearranged configuration.

[0099] In a preferred embodiment, the humanized light chain locus contains at least one C-region gene segment that is a human constant region gene segment, preferably, C λ or C κ .

[0100] In another preferred embodiment, the humanized light chain locus of the transgene contains a humanized V-region and a humanized C-region, e.g., a V-region having at least one human VL gene and/or at least one rearranged human VJ segment, and a C-region having at least one human C gene segment (e.g., human C λ or C κ).

[0101] Preferably, the humanized V-region includes at least about 10-100 light chain V (or "VL") gene segments, at least one of which is a human VL gene segment. The human VL gene segment included in the transgene shares at least about 75% to about 85% homology to the VL gene segments of the host animal, particularly those animal VL gene segments included in the upstream region of the transgene. Consistently, a human VL segment encompasses naturally occurring sequences of a human VL gene segment, degenerate forms of naturally occurring sequences of a human VL gene segment, as well as synthetic sequences that encode a polypeptide sequence substantially (i.e., at least about 85%-95%) identical to a human light chain V domain polypeptide.

[0102] Preferably, the human VL gene segment(s) is placed downstream of the non-human VL segments in the transgene locus. The non-human VL gene segments in the

transgene construct are selected from the VL gene segments in the 3'VL-region in the light chain locus of the host animal, including the 3' proximal VL1.

[0103] In still another preferred embodiment, the humanized light chain locus includes a rearranged human VJ segment, placed downstream of a number of (e.g., 10-100) VL gene segments of either non-human or human origin.

[0104] Another aspect of the present invention is directed to methods of making a transgenic vector containing a humanized Ig locus. Such methods involve isolating an Ig locus or a portion thereof from a non-human animal, and inserting the desired human Ig gene segment(s) into the isolated animal Ig locus or the isolated portion of an animal Ig locus. The human Ig gene segment(s) are inserted into the isolated animal Ig locus or a portion thereof by ligation or homologous recombination in such a way as to retain the capacity of the locus of undergoing effective gene rearrangement and gene conversion in the non-human animal.

[0105] Preferably, DNA fragments containing an Ig locus to be humanized are isolated from animals which generate antibody diversity by gene conversion, e.g., rabbit and chicken. Such large DNA fragments can be isolated by screening a library of plasmids, cosmids, YACs or BACs, and the like, prepared from the genomic DNA of the non-human animal. An entire animal C-region can be contained in one plasmid or cosmid clone which is subsequently subjected to humanization. YAC clones can carry DNA fragments of up to 2 megabases, thus an entire animal heavy chain locus or a large portion thereof can be isolated in one YAC clone, or reconstructed to be contained in one YAC clone. BAC clones are capable of carrying DNA fragments of smaller sizes (about 150-250 kb). However, multiple BAC clones containing overlapping fragments of an Ig locus can be separately humanized and subsequently injected together into an animal recipient cell, wherein the overlapping fragments recombine in the recipient animal cell to generate a continuous Ig locus.

[0106] Human Ig gene segments can be integrated into the Ig locus on a vector (e.g., a BAC clone) by a variety of methods, including ligation of DNA fragments, or insertion of DNA fragments by homologous recombination. Integration of the human Ig gene segments is done in such a way that the human Ig gene segment is operably linked to the host animal sequence in the transgene to produce a functional humanized Ig locus, i.e., an Ig locus capable of gene rearrangement and gene conversion which lead to the production of a diversified repertoire of humanized antibodies.

[0107] Preferably, human Ig gene segments are integrated into the Ig locus by homologous recombination. Homologous recombination can be performed in bacteria, yeast and other cells with a high frequency of homologous recombination events. For example, a yeast cell is transformed with a YAC containing an animal's Ig locus or a large portion thereof. Subsequently, such yeast cell is further transformed with a recombination vector as described hereinabove, which carries a human Ig gene segment linked to a 5' flanking sequence and a 3' flanking sequence. The 5' and the 3' flanking sequences in the recombination vector are homologous to those flanking sequences of the animal Ig gene segment on the YAC. As a result of a homologous' recombination, the animal Ig gene segment on the YAC is replaced with the human Ig gene segment. Alternatively, a

bacterial cell such as *E. coli* is transformed with a BAC containing an animal's Ig locus or a large portion thereof. Such bacterial cell is further transformed with a recombination vector which carries a human Ig gene segment linked to a 5' flanking sequence and a 3' flanking sequence. The 5' and the 3' flanking sequences in the recombination vector mediate homologous recombination and exchange between the human Ig gene segment on the recombination vector and the animal Ig gene segment on the BAC. Humanized YACs and BACs can be readily isolated from the cells and used in making transgenic animals.

[0108] In a further aspect of the present invention, methods of making transgenic animals capable of producing humanized immunoglobulins are provided.

[0109] According to the present invention, a transgenic animal capable of making humanized immunoglobulins are made by introducing into a recipient cell or cells of an animal one or more of the transgenic vectors described herein above which carry a humanized Ig locus, and deriving an animal from the genetically modified recipient cell or cells.

[0110] Preferably, the recipient cells are from non-human animals which generate antibody diversity by gene conversion and hypermutation, e.g., bird (such as chicken), rabbit, cows and the like. In such animals, the 3'proximal V gene segment is preferentially used for the production of immunoglobulins. Integration of a human V gene segment into the Ig locus on the transgene vector, either by replacing the 3'proximal V gene segment of the animal or by being placed in close proximity of the 3'proximal V gene segment, results in expression of human V region polypeptide sequences in the majority of immunoglobulins. Alternatively, a rearranged human V(D)J segment may be inserted into the J locus of the immunoglobulin locus on the transgene vector.

[0111] The transgenic vectors containing a humanized Ig locus is introduced into the recipient cell or cells and then integrated into the genome of the recipient cell or cells by random integration or by targeted integration.

[0112] For random integration, a transgenic vector containing a humanized Ig locus can be introduced into an animal recipient cell by standard transgenic technology. For example, a transgenic vector can be directly injected into the pronucleus of a fertilized oocyte. A transgenic vector can also be introduced by co-incubation of sperm with the transgenic vector before fertilization of the oocyte. Transgenic animals can be developed from fertilized oocytes. Another way to introduce a transgenic vector is by transfecting embryonic stem cells and subsequently injecting the genetically modified embryonic stem cells into developing embryos. Alternatively, a transgenic vector (naked or in combination with facilitating reagents) can be directly injected into a developing embryo. Ultimately, chimeric transgenic animals are produced from the embryos which contain the humanized Ig transgene integrated in the genome of at least some somatic cells of the transgenic animal.

[0113] In a preferred embodiment, a transgene containing a humanized Ig locus is randomly integrated into the genome of recipient cells (such as fertilized oocyte or developing embryos) derived from animal strains with an impaired expression of endogenous immunoglobulin genes. The use of such animal strains permits preferential expres-

sion of immunoglobulin molecules from the humanized transgenic Ig locus. Examples for such animals include the Alicia and Basilea rabbit strains, as well as Agammaglobinemic chicken strain. Alternatively, transgenic animals with humanized immunoglobulin transgenes or loci can be mated with animal strains with impaired expression of endogenous immunoglobulins. Offspring homozygous for an impaired endogenous Ig locus and a humanized transgenic Ig locus can be obtained.

[0114] For targeted integration, a transgenic vector can be introduced into appropriate animal recipient cells such as embryonic stem cells or already differentiated somatic cells. Afterwards, cells in which the transgene has integrated into the animal genome and has replaced the corresponding endogenous Ig locus by homologous recombination can be selected by standard methods. The selected cells may then be fused with enucleated nuclear transfer unit cells, e.g. oocytes or embryonic stem cells, cells which are totipotent and capable of forming a functional neonate. Fusion is performed in accordance with conventional techniques which are well established. See, for example, Cibelli et al., *Science* (1998) 280:1256. Enucleation of oocytes and nuclear transfer can also be performed by microsurgery using injection pipettes. (See, for example, Wakayama et al., *Nature* (1998) 394:369.) The resulting egg cells are then cultivated in an appropriate medium, and transferred into synchronized recipients for generating transgenic animals. Alternatively, the selected genetically modified cells can be injected into developing embryos which are subsequently developed into chimeric animals.

[0115] Further to the present invention, a transgenic animal capable of producing humanized immunoglobulins can also be made by introducing into a recipient cell or cells, one or more of the recombination vectors described herein above, which carry a human Ig gene segment, linked to 5' and 3' flanking sequences that are homologous to the flanking sequences of the endogenous Ig gene segment, selecting cells in which the endogenous Ig gene segment is replaced by the human Ig gene segment by homologous recombination, and deriving an animal from the selected genetically modified recipient cell or cells.

[0116] Similar to the target insertion of a transgenic vector, cells appropriate for use as recipient cells in this approach include embryonic stem cells or already differentiated somatic cells. A recombination vector carrying a human Ig gene segment can be introduced into such recipient cells by any feasible means, e.g., transfection. Afterwards, cells in which the human Ig gene segment has replaced the corresponding endogenous Ig gene segment by homologous recombination, can be selected by standard methods. These genetically modified cells can serve as nuclei donor cells in a nuclear transfer procedure for cloning a transgenic animal. Alternatively, the selected genetically modified embryonic stem cells can be injected into developing embryos which can be subsequently developed into chimeric animals.

[0117] Transgenic animals produced by any of the foregoing methods form another embodiment of the present invention. The transgenic animals have at least one, i.e., one or more, humanized Ig loci in the genome, from which a functional repertoire of humanized antibodies is produced.

[0118] In a preferred embodiment, the present invention provides transgenic rabbits having one or more humanized

Ig loci in the genome. The transgenic rabbits of the present invention are capable of rearranging and gene converting the humanized Ig loci, and expressing a functional repertoire of humanized antibodies.

[0119] In another preferred embodiment, the present invention provides transgenic chickens having one or more humanized Ig loci in the genome. The transgenic chickens of the present invention are capable of rearranging and gene converting the humanized Ig loci, and expressing a functional repertoire of humanized antibodies.

[0120] Cells derived from the transgenic animals of the present invention, such as B cells or cell lines established from a transgenic animal immunized against an antigen, are also part of the present invention.

[0121] In a further aspect of the present invention, methods are provided for treating a disease in a primate, in particular, a human subject, by administering a purified humanized antibody composition, preferably, a humanized polyclonal antibody composition, desirable for treating such disease.

[0122] The humanized polyclonal antibody compositions used for administration are generally characterized by containing a polyclonal antibody population, having immunoglobulin concentrations from 0.1 to 100 mg/ml, more usually from 1 to 10 mg/ml. The antibody composition may contain immunoglobulins of various isotypes. Alternatively, the antibody composition may contain antibodies of only one isotype, or a number of selected isotypes.

[0123] In most instances the antibody composition consists of unmodified immunoglobulins, i.e., humanized antibodies prepared from the animal without additional modification, e.g., by chemicals or enzymes. Alternatively, the immunoglobulin fraction may be subject to treatment such as enzymatic digestion (e.g. with pepsin, papain, plasmin, glycosidases, nucleases, etc.), heating, etc. and/or further fractionated.

[0124] The antibody compositions generally are administered into the vascular system, conveniently intravenously by injection or infusion via a catheter implanted into an appropriate vein. The antibody composition is administered at an appropriate rate, generally ranging from about 10 minutes to about 24 hours, more commonly from about 30 minutes to about 6 hours, in accordance with the rate at which the liquid can be accepted by the patient. Administration of the effective dosage may occur in a single infusion or in a series of infusions. Repeated infusions may be administered once a day, once a week once a month, or once every three months, depending on the half-life of the antibody preparation and the clinical indication. For applications on epithelial surfaces the antibody compositions are applied to the surface in need of treatment in an amount sufficient to provide the intended end result, and can be repeated as needed.

[0125] The antibody compositions can be used to bind and neutralize antigenic entities in human body tissues that cause disease or that elicit undesired or abnormal immune responses. An "antigenic entity" is herein defined to encompass any soluble or cell-surface bound molecules including proteins, as well as cells or infectious disease-causing organisms or agents that are at least capable of binding to an antibody and preferably are also capable of stimulating an immune response.

[0126] Administration of an antibody composition against an infectious agent as a monotherapy or in combination with chemotherapy results in elimination of infectious particles. A single administration of antibodies decreases the number of infectious particles generally 10 to 100 fold, more commonly more than 1000-fold. Similarly, antibody therapy in patients with a malignant disease employed as a monotherapy or in combination with chemotherapy reduces the number of malignant cells generally 10 to 100 fold, or more than 1000-fold. Therapy may be repeated over an extended amount of time to assure the complete elimination of infectious particles, malignant cells, etc. In some instances, therapy with antibody preparations will be continued for extended periods of time in the absence of detectable amounts of infectious particles or undesirable cells. Similarly, the use of antibody therapy for the modulation of immune responses may consist of single or multiple administrations of therapeutic antibodies. Therapy may be continued for extended periods of time in the absence of any disease symptoms.

[0127] The subject treatment may be employed in conjunction with chemotherapy at dosages sufficient to inhibit infectious disease or malignancies. In autoimmune disease patients or transplant recipients, antibody therapy may be employed in conjunction with immunosuppressive therapy at dosages sufficient to inhibit immune reactions.

[0128] The invention is further illustrated, but by no means limited, by the following examples.

EXAMPLE 1

Novel Sequences 3'Prime of the C γ Gene from Cows, Sheep and Rabbits

[0129] Genomic DNA was isolated from blood of a Simmental cow using the QIAamp DNA Blood Kit (QIAGEN). The genomic region 3' of the cow C γ gene (i.e., the cow C γ gene 3' flanking sequence) was PCR-amplified using the isolated genomic DNA as template and the following primers:

5' primer:
5'cgcaagcttCCTACACGTGTGGTGATG3'; (SEQ ID NO: 1)

3' primer:
5'cgcaagcttAAGATGGWGATGGTSGTCCA3' (SEQ ID NO: 2)

The upper-case portion of the 5' primer was from exon 3 of C γ , and the lower-case portion represented a terminal HindIII restriction site. The upper-case portion of the 3' primer was a degenerate sequence designed according to the published sequences from the human M1 exon and the mouse M1 exon, and the lower-case portion represented a terminal HindIII restriction site. A 1.3 kb PCR fragment was obtained using the EXPAND long template PCR system (Roche). The fragment was gel purified, digested with HindIII, and cloned into a Bluescript cloning vector. The resulting clones fell into three populations, which differ from one another in the pattern of the restriction fragments obtained with BamHI, EcoRI and XhoI. One clone from each population was sequenced, and the sequences are shown in FIG. 1 (SEQ ID NOS: 3-5).

[0130] Genomic DNA was isolated from blood of a Merino sheep using the QIAamp DNA Blood Kit

(QIAGEN). The genomic region 3' of the sheep C γ gene (i.e., the sheep C γ gene 3' flanking sequence) was PCR-amplified using the isolated genomic DNA as template and the following primers:

5' primer:
5'cgcgatcccCTACGCGTGTGGTGTATG3' (SEQ ID NO: 6)

3' primer:
5'cgcgatcccACCGAGGAGAAGATCCACTT3' (SEQ ID NO: 7)

The upper-case portion of the 5' primer was from exon 3 of C γ , and the lower-case portion represented a terminal BamHI restriction site. The upper-case portion of the 3' primer was designed according to the published sequences from the human M2 exon and the mouse M2 exon, and the lower-case portion represented a terminal BamHI restriction site. A 2.9 kb PCR fragment was obtained using the EXPAND long template PCR system (Roche). The fragment was gel purified, digested with BamHI, and cloned into a Bluescript cloning vector. The resulting clones fell into two populations, which differ from each other in the pattern of the restriction fragments obtained with HindIII, EcoRI and XhoI. One clone from each population was sequenced, and the sequences are shown in FIG. 2 (SEQ ID NOS: 8-9).

[0131] A 10 kb EcoRI fragment containing the C γ gene and its flanking sequences from A2 allotype rabbit was subcloned from a genomic cosmid clone (cos 8.3 from Knight et al., *J Immunol* (1985) 1245-50, "Organization and polymorphism of rabbit immunoglobulin heavy chain genes"). The nucleotide sequences 5' and 3' of C γ were determined using standard methods and are set forth in FIG. 3 and 5, SEQ ID NO: 10, 12, 13, respectively.

[0132] Sequences 3' of rabbit Ckappa were determined from an EcoRI/BamHI subclone from VJk2C λ In pSV2neo. The nucleotide sequence is set forth in FIG. 4, SEQ ID NO: 11.

[0133] The amino acid sequences encoded by the M1 and M2 exons from cow, sheep and rabbit were deduced from the above 3' flanking sequence. These amino acid sequences were aligned with the published M1 and M2 sequences from camel, human and mouse, as shown in FIG. 6.

EXAMPLE 2

A Vector for Replacing the Rabbit Endogenous C γ Gene Segment with the Human C γ 1 Segment

[0134] Genomic DNA is isolated from rabbit fetal fibroblasts of an α 2-homozygous rabbit. The DNA sequence upstream of rabbit C γ (i.e., the 5' flanking sequence of rabbit C γ) is amplified by PCR using the following primers:

5' taattatgcgccgctTTCAGCGTGAACCAC (SEQ ID NO: 39)
GCCCTC 3'
with a 5' NotI site and

5' GTCGACGCCCTCGATGCACTCCAGAG (SEQ ID NO: 40)
3'.

[0135] The DNA sequence downstream of rabbit C γ (i.e., the 3' flanking sequence of rabbit C γ) is amplified with the following primers:

5' ggtaccCTCTCCCTCCCCACGCCGAGC (SEQ ID NO: 41)
3'
with a 5' KpnI site and

5' atatctcagaACTGGCTGTCCCTGCTAGT (SEQ ID NO: 42)
ACACGG 3'
with a 5' XhoI site.

[0136] Human genomic DNA is isolated from human peripheral blood lymphocytes. The DNA fragment encoding human C γ 1 is amplified using the following primers:

5' GTCGACACTGGACGCTGAACCTCGCGG 3' (SEQ ID NO: 43)
and

5' GGTACCGGGGGCTTGCCGGCCGTCGCAC (SEQ ID NO: 44)
3'.

[0137] The fragments are digested with restriction enzymes and cloned into a Bluescript vector. Subsequently, a lox neo-cassette is inserted into the Sall site and an Hsv-tk cassette into the XhoI site. A schematic drawing of the final construct is shown in FIG. 7a.

EXAMPLE 3

A Vector for Replacing the Rabbit Endogenous Ck Gene Segment with the Human Ck Segment

[0138] Genomic DNA was isolated from rabbit fetal fibroblasts of a b5-homozygous rabbit. The DNA sequence upstream of rabbit Ck1 (i.e., the 5' flanking sequence of rabbit Ck1) was amplified by PCR using the following primers:

5' gggggcgcTGGCGAGGAGACCAAGCTGGAGA (SEQ ID NO: 45)
TCAAACG 3'
with a 5' NotI site

5' GTCGACGCAGCCCAAAGCTGTTGCAATGGGG (SEQ ID NO: 46)
CAGCG 3'.

[0139] The DNA sequence downstream of rabbit Ck1 (i.e., the 5' flanking sequence of rabbit Ck1) was amplified with the following primers:

5' atatggtaccGCGAGAGCCTGCCAGGGCAC (SEQ ID NO: 47)
CGCC 3'
with a 5' KpnI site

5' GGATCCCGAGCTTTATGGGAGGGTGGGG (SEQ ID NO: 48)
3'.

[0140] Human genomic DNA was isolated from human peripheral blood lymphocytes. The DNA fragment encoding human Ck was amplified using the following primers:

5' ATATGTCGACCTGGGATAAGCATGCTGTTTT (SEQ ID NO: 49)
CTGTCTGTCCC 3'

5' CTAGGTACCAGCAGGTGGGGCACTTCTCCC (SEQ ID NO: 50)
3'.

The fragments were digested with restriction enzymes and cloned into a Bluescript vector. Subsequently, a lox neo-

cassette was inserted into the Sall site and an Hsv-tk cassette into the XhoI site. A schematic drawing of the final construct is shown in FIG. 7b.

EXAMPLE 4

Replacement of the Endogenous C γ and C κ Gene Segments in Rabbit Fetal Fibroblasts with the Corresponding Human Gene Segments

[0141] Rabbit fetal fibroblast cells are prepared by standard methods. After one passage, fibroblasts are transfected with 5 μ g of the NotI-linearized targeting vector as shown in FIG. 5a for C γ or FIG. 51b for C κ , and are seeded in 96-well plates (2×10^3 cells/well). After a positive selection with 600 μ g/ml G418 and a negative selection with 200 nM FIAU, resistant colonies are replica-plated to two 96-well plates for DNA analysis and cryopreservation, respectively. PCR and/or Southern blot analysis is performed to identify cells with the human C γ 1 gene segment integrated in the genome. The cells having the integrated human C γ 1 gene are used in rabbit cloning as described in Example 5.

EXAMPLE 5

Cloning of Rabbits

[0142] Mature Dutch Belton rabbits are superovulated by subcutaneous injection of follicle stimulating hormone (FSH) every 12 hours (0.3 mg \times 2 and 0.4 mg \times 4). Ovulation is induced by intravenous administration of 0.5 mg luteinizing hormone (LH) 12 hours after the last FSH injection. Oocytes are recovered by ovidual flush 17 hours after LH injection. Oocytes are mechanically enucleated 16-19 hours after maturation. Chromosome removal is assessed with bisBENZIMIDE (HOECHST 33342, Sigma, St. Louis, Mo.) dye under ultraviolet light. Enucleated oocytes are fused with actively dividing fibroblasts by using one electrical pulse of 180 V/cm for 15 μ s (Electrocell Manipulator 200, Genetronics, San Diego, Calif.). After 3-5 hours oocytes are chemically activated with calcium ionophore (6 μ M) for 4 min (#407952, Calbiochem, San Diego, Calif.) and 2 mM 6-dimethylaminopurine (DMAP, Sigma) in CR2 medium (Specialty Media, Lavalett, N.J.) with 3 mg/ml bovine serum albumin (fatty acid free, Sigma) for 3 hours. Following the activation, the embryos are washed in hamster embryo culture medium (HECM)-Hepes five times and subsequently, cultivated in CR2 medium containing 3 mg/ml fatty-acid free BSA for 248 hours at 37.8 $^\circ$ C. and 5% CO $_2$ in air. Embryos are then transferred into synchronized recipients. Offsprings are analyzed by PCR for a segment of the transgene.

EXAMPLE 6

Construction of a DNA Fragment Containing a Portion of a Rabbit Heavy Chain Locus with a Human C γ 1 Gene Segment and a VH Gene Segment Encoding a Human VH Domain Polypeptide Sequence

[0143] The upstream and downstream regions (i.e., the 5' and 3' flanking regions) of the rabbit heavy chain C γ gene from an a2-allotype rabbit were sequenced. A DNA fragment (SEQ ID NO: 51) is generated by PCR using overlapping oligonucleotides wherein the DNA fragment contains

from 5' to 3', a sequence derived from the 5' flanking region of the rabbit C γ gene, the human C γ 1 gene, and a sequence derived from the 3' flanking region of the rabbit C γ gene (FIG. 8).

[0144] A genomic BAC library derived from an a2-allotype rabbit is generated by standard procedures and screened with probes specific for rabbit C γ . A BAC clone containing rabbit heavy chain gene segments is identified. The rabbit C γ gene on this BAC clone is replaced with the human C γ 1 gene by homologous recombination in *E. coli* using the DNA fragment of SEQ ID NO: 51 and the pET system. This replacement is accomplished by two consecutive recombination steps: first the rabbit C γ gene segment is replaced with a marker gene; then the marker gene is replaced the human C γ 1 gene segment.

[0145] The modified BAC clone containing rabbit heavy chain genes and the inserted human C γ 1 gene is further modified by replacing the 3'proximal VH1 segment with a synthetic VH gene segment (FIG. 9). This synthetic VH gene segment (SEQ ID NO: 52) is made using overlapping oligonucleotides and includes a 5' flanking sequence, a 3' flanking sequence, and a sequence coding for a polypeptide nearly identical to the human immunoglobulin heavy chain variable domain polypeptide sequence described by Huang and Stollar (*J. Immunol.* 151: 5290-5300, 1993). The coding sequence of the synthetic VH gene segment is designed based on the published sequence of a rabbit VH1 gene (a2, Knight and Becker, *Cell* 60:963-970, 1990) and is more than 80% identical to rabbit VH gene segments. The 5' and the 3' flanking sequences in the synthetic VH segment are derived from the upstream and downstream regions of the a2-allotype rabbit VH1 gene. The synthetic VH gene of SEQ ID NO: 52 is used to replace the rabbit VH1 gene on the BAC clone by homologous recombination using the pET or the red ϵ β γ system. The modified BAC clone is amplified and purified using standard procedures.

EXAMPLE 7

Construction of a DNA Fragment Containing a Portion of a Rabbit Light Chain Locus with a Human C κ Gene Segment and a VJ Gene Segment Encoding a Human VL Domain Polypeptide Sequence

[0146] The upstream and downstream regions (i.e., the 5' and 3' flanking regions) of the rabbit light chain C κ 1 gene from a b5-allotype rabbit were sequenced. A DNA fragment (SEQ ID NO: 53) is generated by PCR using overlapping oligonucleotides wherein the DNA fragment contains from 5' to 3', a sequence derived from the 5' flanking region of the rabbit C κ 1 gene, the human C κ 1 gene, and a sequence derived from the 3' flanking region of the rabbit C κ 1 gene (FIG. 10).

[0147] A genomic BAC library derived from a b5-allotype rabbit is generated by standard procedures and screened with probes specific for rabbit C κ 1. A BAC clone containing rabbit light chain gene segments is identified. The rabbit C κ 1 gene on this BAC clone is replaced with the human C κ 1 gene on the DNA fragment of SEQ ID NO: 53 by homologous recombination in *E. coli* using the pET or the red ϵ β γ system. This replacement is accomplished by two consecutive recombination steps: first the rabbit C κ 1 gene segment

is replaced with a marker gene; then the marker gene is replaced the human Ck1 gene segment.

[0148] The modified BAC clone containing rabbit light chain genes and the inserted human Ck1 gene is further modified by inserting a rearranged VJ DNA fragment into the J region of the rabbit light chain locus. The rearranged VJ DNA fragment encodes a human immunoglobulin variable domain polypeptide described by Pritsch et al. (*Blood* 82(10):3103-3112, 1993) and Lautner-Rieske et al. (*Eur. J. Immunol.* 22 (4), 1023-1029, 1992)) (FIG. 7). The nucleotide sequence of the rearranged VJ fragment is designed to maximize the sequence homology at the nucleotide level to the rabbit Vkappa sequence published by Lieberman et al. (*J. Immunol.* 133 (5), 2753-2756, 1984). This rearranged VJ DNA sequence is more than 80% identical with known rabbit Vk genes. Using overlapping oligonucleotides in PCR, the rearranged VJ DNA fragment is linked to a 5' and a 3' flanking sequence, resulting the DNA fragment of SEQ ID NO: 54 (FIG. 11). The 5'flanking sequence is derived from 5' of a rabbit VK, the 3'flanking sequence is derived from 3' of rabbit J2. The DNA fragment of SEQ ID NO: 54 is subsequently inserted into the rabbit light chain locus by homologous recombination in *E. coli* using the pET or the red β y system. The insertion is performed in such a way that the rabbit light chain region containing the rabbit Vk1 gene segment, the rabbit J1 and J2 segments, and the sequences in between, is replaced with the rearranged VJ DNA fragment. Again, this insertion is accomplished by replacement of the rabbit V to J region with a marker gene, followed by the replacement of the marker gene with the rearranged VJ DNA fragment. The modified BAC clone is amplified and purified using standard procedures.

EXAMPLE 8

Transgenic Rabbits Expressing the Humanized Immunoglobulin Light and/or Heavy Chain Transgene

[0149] Transgenic rabbits are generated as described by Fan et al. (*Pathol Int.* 49: 583-594, 1999). Briefly, female rabbits are superovulated using standard methods and mated with male rabbits. Pronuclear-stage zygotes are collected from oviduct and placed in an appropriate medium such as Dulbecco's phosphate buffered saline supplemented with 20% fetal bovine serum. The exogenous DNA (e.g., the humanized BAC clone from Example 4 and/or 5 which has been linearized prior to injection) is microinjected into the male pronucleus with the aid of a pair of manipulators. Morphological surviving zygotes are transferred to the oviducts of pseudopregnant rabbits. Pseudopregnancy is induced by the injection of human chorionic gonadotrophin (hCG). Between about 0.1-1% of the injected zygotes develop into live transgenic rabbits. Integration of the transgene in the genome is confirmed by Southern blots analysis using a probe specific for the transgene.

[0150] cDNA is prepared using RNA isolated from B cells (in blood, spleen and/or lymph nodes) of a transgenic rabbit. Primers specific for the human transgene (human CH gene segment or the synthetic humanized VH gene segment) are used to generate amplified products from cDNA. The observation of amplified products indicates that the transgene is rearranged in the transgenic animal and the rearranged transgene is transcribed in the animal. Amplified products

are sequenced and the presence of donor sequences from upstream V genes indicates that the transgene introduced into the germline of the animal undergoes gene conversion.

[0151] The presence of antibodies containing human IgG and/or human kappa light chain antigenic determinants in the serum of transgenic founder rabbits is determined using an ELISA assay.

EXAMPLE 9

Production of Humanized Antibodies From Transgenic Rabbits with the Genetic Background of the Alicia and/or Basilea Rabbit Strain

[0152] The Alicia strain lacks the VH1 gene segment and therefore has an impaired Ig heavy chain expression. Transgenic founder rabbits capable of expressing humanized heavy chain molecules in the genetic background of the Alicia rabbit strain are generated, e.g., by using fetal fibroblasts established from Alicia rabbits in Examples 4-5 above, or by using zygotes from female Alicia rabbits mated with male Alicia rabbits in Example 8 above. Transgenic animals are also obtained which are homozygous for the Alicia Ig phenotype and are also homozygous for a humanized heavy chain transgene. Serum is tested in ELISA for the presence of humanized heavy chain (e.g., a human heavy chain constant region). The concentration of antibodies with humanized Ig heavy chains in these homozygous Alicia animals is substantially higher, e.g., about 10 to 100 fold higher, than that produced from a transgene integrated in the genome of wild type (non-Alicia) rabbits.

[0153] The Basilea strain does not express κ 1 light chain and in its place exclusively express the κ 2 and λ light chains. Transgenic founder rabbits capable of expressing humanized light chain molecules in the genetic background of the Basilea rabbit strain are generated, e.g., by using fetal fibroblasts established from Basilea rabbits in Examples 4-5 above, or by using zygotes from female Basilea rabbits mated with male Basilea rabbits in Example 8 above. Transgenic animals are obtained which are homozygous for the Basilea light chain phenotype, and are also homozygous for a humanized light chain transgene. Serum is tested in ELISA for the presence of the humanized light chain. The concentration of the humanized light chain in the homozygous Basilea animals is substantially higher, about 10-100 fold higher, than the concentration of a humanized light chain in a transgenic rabbit with the wild type (non-Basilea) genetic background. Transgenic founder rabbits are mated with each other to generate transgenic rabbits with the following traits: (1) having at least one humanized light chain transgene, (2) having at least one humanized heavy chain transgene, (3) homozygous for the Alicia heavy chain locus, and (4) homozygous for the Basilea light chain locus.

EXAMPLE 10

Construction of a DNA Fragment Containing a Modified Chicken Light Chain Locus Having a Human Clambda2 Gene Segment and a VJ Gene Segment Encoding a Human VL Domain

[0154] A genomic BAC library derived from a jungle fowl chicken was screened with radiolabeled probes specific for chicken light chain Clambda and chicken Vpsi25 (the V gene segment at the very 5' end of the light chain locus). A

BAC clone containing the entire lambda light chain locus was identified. The chicken C λ gene on this BAC clone is replaced with the human C λ 2 gene by homologous recombination in *E. coli* using the pET system (Zhang et al., *Nat. Biotechnol.* 18(12):1314-7, 2000) as follows.

[0155] A first DNA fragment containing a kanamycin selection cassette was generated by PCR using primers specific for Tn5 gene. The 5' primer (5'catcacagccatacat-acgcgtgtggccctctcctctctct-tgcaggTATGGACAGCAAGCGAACCG 3', SEQ ID NO: 55) was designed to include 50 bp at the 5' end (lower case), derived from the 5' flanking region of the chicken light chain C λ gene. The 3' primer (5'atcagggtgaccctacgttacactct-gtcaccaaggagtgaggaggacTCAGAAGAACTCGTCAAGAAG3', SEQ ID NO: 56) was designed to include about 50 bp at the end (lower case), derived from the 3' flanking region of the chicken light chain C λ gene.

[0156] A second DNA fragment (SEQ ID NO: 57) was synthesized using overlapping oligonucleotides wherein the DNA fragment contains from 5' to 3', a sequence derived from the 5' flanking region of the chicken light chain Clambda gene, the human Clambda2 gene, and a sequence derived from the 3' flanking region of the chicken Clambda gene (FIG. 12).

[0157] *E. coli* cells of the chicken light chain BAC clone were transformed with a recombination plasmid expressing the recE and recT functions under an inducible promoter. Cells transformed with the recombination plasmid were then transformed with the first DNA fragment above and selected afterwards in media containing kanamycin. Clones resistant to kanamycin were identified, and the replacement of the chicken C λ segment by the kanamycin selection cassette via homologous recombination was confirmed by restriction enzyme digest.

[0158] In the second homologous recombination step, cells positive for the kanamycin selection cassette were transformed with the second DNA fragment above. Transformed cells were screened for the loss of kanamycin resistance as indicative of the replacement of the kanamycin selection cassette by the human C λ 2 gene. The exchange was confirmed by restriction enzyme digest and/or sequence analysis.

[0159] The ET cloning procedure is summarized in FIG. 13.

[0160] The BAC clone containing the chicken light chain locus and the inserted human Clambda2 gene segment was further modified by inserting a rearranged VJ DNA fragment. The rearranged VJ DNA fragment encodes a human immunoglobulin variable domain polypeptide described by Kametani et al. (*J. Biochem.* 93 (2), 421-429, 1983) as IG LAMBDA CHAIN V-I REGION NIG-64 (P01702) (FIG. 14). The nucleotide sequence of the rearranged VJ fragment was so designed as to maximize the sequence homology at the nucleotide level to the chicken Vlambd1 sequence published by McCormack et al. (*Cell* 56, 785-791, 1989). This rearranged VJ DNA sequence is more than 80% identical with known chicken light chain V genes. The rearranged VJ DNA fragment was linked to a 5' flanking sequence and a 3' flanking sequence, resulting in the DNA fragment of SEQ ID NO: 58 (FIG. 14). The 5' flanking sequence was derived from 5' of chicken Vlambd1, and the

3'flanking sequence was derived from 3' of chicken J. The DNA fragment of SEQ ID NO: 58 was subsequently inserted into the chicken light chain locus in *E. coli* using the pET system as shown in FIG. 15. The insertion was performed in such a way that the region on the chicken light chain locus from the 5' end of the chicken Vlambd1 gene segment to the 3' end of the chicken J region was replaced with the rearranged, synthetic VJ DNA fragment. Again, this insertion was accomplished by the replacement of the chicken V-J region with a marker gene, followed by the replacement of the marker gene with the rearranged VJ DNA fragment. The modified region of the chicken light chain locus is shown in FIG. 15. The modified BAC clone was amplified and purified using standard procedures.

EXAMPLE 11

Construction of a DNA Fragment Containing a Portion of a Chicken Heavy Chain Locus With a Human C γ 1 Gene Segment and a VH Gene Segment Encoding a Human VH Domain Polypeptide Sequence

[0161] A jungle fowl chicken genomic BAC library was generated by standard procedures and screened with probes specific for chicken C γ . A BAC clone containing chicken heavy chain gene segments is identified. The upstream and downstream regions (i.e., the 5' and 3' flanking regions) of the heavy chain C γ gene are sequenced. The chicken C γ gene on this BAC clone is replaced with the human C γ 1 gene by homologous recombination in *E. coli* using the pET system as follows.

[0162] A first DNA fragment containing a kanamycin selection cassette is generated by PCR using primers specific for Tn5 gene. The 5' and 3' primers are designed to include about 50 bp at the end, derived from the 5' and 3' flanking regions of the chicken heavy chain C γ gene.

[0163] A second DNA fragment is generated by PCR using overlapping oligonucleotides wherein this second DNA fragment contains from 5' to 3', a sequence of about 50 bp derived from the 5' flanking region of the chicken C γ gene, the human C γ 1 gene, and a sequence of about 50 bp derived from the 3' flanking region of the chicken C γ gene.

[0164] *E. coli* cells of the chicken CY BAC clone are transformed with a recombination plasmid expressing the recE and recT functions under an inducible promoter. Cells transformed with the recombination plasmid are further transformed with the first DNA fragment and selected in media containing kanamycin. Clones resistant to kanamycin are identified, and the replacement of the chicken CY segment by the kanamycin selection cassette via homologous recombination is confirmed by restriction enzyme digest.

[0165] In the second homologous recombination step, cells positive for the kanamycin selection cassette are now transformed with the second DNA fragment described above. Transformed cells are screened for loss of kanamycin resistance as indicative of the replacement of the kanamycin selection cassette by the human C γ 1 gene. The exchange is confirmed by restriction enzyme digest and/or sequence analysis.

[0166] The BAC clone containing the inserted human C γ 1 gene is further modified by replacing the 3'proximal VH1

segment (i.e., the 3'proximal VH1 gene in the V region) with a synthetic VH gene segment. This synthetic VH gene segment is designed based on the published sequence of a chicken VH1 gene (Arakawa et al., *EMBO J* 15(10): 2540-2546, 1996). The synthetic gene segment is more than 80% identical to chicken VH gene segments and encodes an amino acid sequence that is identical to the amino acid sequence of a human immunoglobulin heavy chain variable domain polypeptide described by Matthyssens and Rabbitts (in Steinberg CM and Lefkovits I, (eds). *The Immune System*: 132-138, S. Karger, NY 1981). This synthetic VH segment including 5' and 3' flanking sequences is synthesized by PCR using overlapping oligonucleotides. The 5' and the 3' flanking sequences are derived from the upstream and downstream regions of chicken VH1 gene. This synthetic VH segment is used to replace the chicken VH1 gene on the BAC clone by homologous recombination using the pET system. The modified BAC clone is amplified and purified using standard procedures.

EXAMPLE 12

Transgenic Chicken Expressing the Humanized Immunoglobulin Light and/or Heavy Chain Transgenes

[0167] The production of transgenic chicken is carried out using techniques as described by Etches et al., *Methods in Molecular Biology* 62: 433-450; Pain et al., *Cells Tissues Organs* 1999; 165(34): 212-9; Sang, H., "Transgenic chickens—methods and potential applications", *Trends Biotechnol* 12:415 (1994); and in WO 200075300, "Introducing a nucleic acid into an avian genome, useful for transfecting avian blastodermal cells for producing transgenic avian animals with the desired genes, by directly introducing the nucleic acid into the germinal disc of the egg".

[0168] Briefly, the modified BAC clones are linearized and mixed with a transfection reagent to promote uptake of DNA into cells. The formulations are injected into a multi-cell stage chicken embryo in close proximity to the germinal disc. The window in the egg shell is closed and the eggs are incubated. After hatching chimeric chickens are identified by PCR and Southern blot analysis using transgene specific sequences. Integration of the transgene in the genome is confirmed by Southern blots analysis using a probe specific for the transgene. Heavy and light chain transgenic animals are bred with each other to generate transgenic chickens expressing antibodies having humanized heavy and light chains.

[0169] cDNA is prepared using RNA isolated from B cells (in blood, spleen and/or lymph nodes) from transgenic chickens. Primers specific for the human transgene (e.g., human CH gene segments and/or the synthetic humanized VH gene segments) are used to generate amplified products from cDNA. The observation of amplified products indicates that the transgene is rearranged in the transgenic animal and the rearranged transgene is transcribed in the animal. Amplified products are sequenced and the presence of donor sequences from upstream V genes indicates that the transgene introduced into the germline of the animal undergoes gene conversion.

[0170] The presence of antibodies containing human IgG and/or human kappa light chain antigenic determinants in the serum of transgenic chickens is determined using an ELISA assay.

EXAMPLE 13

Production of Functional Humanized Antibodies in Transgenic Chicken with the Agammaglobulinemic Phenotype

[0171] Transgenic chickens with the following traits are produced: (1) having at least one humanized light chain transgene, (2) having at least one humanized heavy chain transgene, and (3) homozygous for the agammaglobulinemic phenotype. These animals produce antibodies into the blood and eggs, and antibodies can be purified from either source. In general, antibody concentrations in the eggs are about 5% to 50% of antibodies concentration in the blood. Animals that contain humanized antibodies at high levels in eggs can be selected and bred to produce offspring. Alternatively, transgenic animals can be generated that specifically secrete humanized antibodies into their eggs.

EXAMPLE 14

Generation of Transgenic Chickens Expressing Humanized Immunoglobulin

[0172] Chicken embryonic stem cells are isolated and cultured as described by Pain et al. (*Development* 122, 2339-2348; 1996). Chicken embryos are obtained from eggs immediately after they are laid. The entire blastoderm is removed by gentle aspiration, embryos are slowly dissociated mechanically and cells are seeded in ESA complete medium on inactivated STO feeder cells. ESA medium is composed of MEM medium containing 10% FCS, 2% chicken serum, 1% bovine serum albumin, 10 ng/ml ovalbumin, 1 mM sodium pyruvate, 1% non-essential amino acids, 1 μ M of each nucleotide adenosine, guanosine, cytidine, uridine, thymidine, 0.16 mM β -mercaptoethanol, ESA complete medium is supplemented with 10 ng/ml bFGF, 20 ng/ml h-IGF-1, 1% vol/vol avian-SCF and 1% vol/vol h-LIF, 1% vol/vol h-IL-11. Cell cultures are incubated wt 37° C. in 7.5 CO₂ and 90% humidity. After 48 hours fresh blastodermal cells are added to the culture in half of the original volume of ESA complete medium. After an additional incubation for three days, the culture medium is partially (50%) replaced with fresh ESA complete medium, and totally every day thereafter. For cell harvesting, cultures are washed with PBS and incubated in a pronase solution (0.025% w/v). Dissociated cells are transfected with various linearized transgenic constructs containing a humanized Ig locus. Transfected cells are incubated with STO feeder cells (as described above) in the presence of selective antibiotics. Cells are transferred onto fresh feeder cells twice per week. Antibiotic resistant cells are isolated and the integration of a humanized Ig gene fragments at a random site or at the corresponding chicken immunoglobulin gene loci is confirmed by PCR.

[0173] Subsequently, genetically modified cells are injected into recipient embryos. As recipient embryos, freshly laid eggs are irradiated (6Gy—Cobalt source). Between 100 to 200 genetically modified cells are injected into the subgerminal cavity using a micropipet. The window in the egg shell is closed and the eggs are incubated. Somatic chimerism of hatched chickens is evaluated by PCR. Germ-line chimerism is assessed by mating of somatic chimeras.

EXAMPLE 15

Immunization of Transgenic Animals

[0174] Genetically engineered chickens are immunized intramuscularly with purified Hepatitis B surface antigen (HBsAg) (5 μ g in incomplete Freund's adjuvant) on day 0, 14 and day 28. On day 35 animals are bled and serum is prepared. ELISA plates (NUNC, Denmark) are coated with 1 μ g/ml HBsAg in PBS for 1 hour at room temperature. Subsequently, available binding sites are blocked by incubation with 1% non-fat dry milk (NFM) in PBS (300 μ l/well). Chicken serum is diluted in PBS/1% NFM and added to the coated wells. After an incubation of 1 hour, the plates are washed 3 times with PBS/0.05% Tween 20 and bound Ig is detected using goat anti-human Ig conjugated with horseradish peroxidase. Conjugated goat antibody is detected using o-phenylenediamine dihydrochloride (Sigma) at 1 mg/ml. The colorimetric reaction is stopped by addition of 1 M HCl solution and the absorbance is measured at 490 nm. As a control, serum from non-immunized chicken is used. Serum from non-immunized chickens does not react with HBsAg. At a dilution of 1:250 the optical density measured in uncoated and HBsAg coated wells is below 0.2. In contrast, serum from immunized chickens contains humanized antibodies reactive with HBsAg. At a serum dilution of 1:250 the measured optical density is 2.3. Upon further dilution of the serum the measured optical density declines to 0.1 (at a dilution of 25600). No antibodies reactive with a goat anti-chicken IgG-HRP conjugate can be detected. This demonstrates that the genetically engineered chickens produce humanized anti-HBsAg antibodies following immunization.

[0175] Genetically engineered rabbits are immunized intramuscularly with purified Hepatitis B surface antigen (HBsAg) (10 μ g in incomplete Freund's adjuvant) on day 0 and day 14. On day 28 animals are bled from the ear and serum is prepared. ELISA plates (NUNC, Denmark) are coated with 1 μ g/ml HBsAg in PBS for 1 hour at room temperature. Subsequently, available binding sites are blocked by incubation with 1% non-fat dry milk (NFM) in PBS (300 μ l/well). Rabbit serum is diluted in PBS/1% NFM and added to the coated wells. After an incubation of 1 hour, the plates are washed 3 times with PBS/0.05% Tween 20 and bound Ig is detected using goat anti-human Ig conjugated with horse-radish peroxidase. Conjugated goat antibody is detected using o-phenylenediamine dihydrochloride (Sigma) at 1 mg/ml. The calorimetric reaction is stopped by addition of 1 M HCl solution and the absorbance is measured at 490 nm. As a control serum from non-immunized rabbits is used. Serum from non-immunized rabbits does not react with HBsAg. At a dilution of 1:100 the optical density measured in uncoated and HBsAg coated wells is below 0.4. In contrast, serum from immunized rabbits contains partially human antibodies reactive with HBsAg. At a serum dilution of 1:100 the measured optical density is 2.8. Upon further dilution of the serum the measured optical density declines to 0.2 (at a dilution of 25600). No antibodies reactive with a goat anti-rabbit IgG-HRP conjugate can be detected. This demonstrates that the genetically engineered rabbits produce humanized anti-HBsAg antibodies following immunization.

EXAMPLE 16

Complement Mediated Cytotoxicity of Virus Infected Cell Line Using Humanized Antibodies

[0176] A human liver carcinoma cell line expressing HBsAg is labeled with 0.1 mCi 51 Cr in 100 μ l PBS for 1 hr at 37° C. Two thousand 51 Cr-labeled cells are incubated with serum from genetically engineered rabbits or chickens expressing anti-HbsAg humanized immunoglobulins. After two hours at 37° C. the release of 51 Cr into the supernatant is determined by measuring radioactivity using a scintillation counter. For the determination of maximum release, 1% Triton X100 is added. The degree of cell lysis is calculated as follows:

$$\% \text{ Lysis} = \frac{\text{CPM}_{\text{experimental}} \pm \text{CPM}_{\text{spontaneous}}}{\text{CPM}_{\text{total}} \pm \text{CPM}_{\text{spontaneous}}} \times 100$$

Incubation of labeled cells with serum (diluted 1:30) from non-immunized rabbits does not result in cell lysis (<10%). However, incubation of cells with serum from immunized rabbits causes 80% cell lysis. Inactivation of complement in the serum by heat treatment (56° C. for 30 minutes) renders the serum from immunized rabbits inactive. These results demonstrate that humanized antibodies produced by genetically engineered rabbits bind to HBsAg-positive cells and cause complement dependent lysis.

EXAMPLE 17

Immunization of Transgenic Animals against *Staphylococcus aureus*

[0177] Genetically engineered chickens are immunized intramuscularly with a recombinant fragment of the *Staphylococcus aureus* collagen adhesin protein (100 μ g in incomplete Freund's adjuvant) on day 0, 14 and day 28. On day 35 animals are bled and serum is prepared. ELISA plates (NUNC, Denmark) are coated with 2 μ g/ml collagen adhesin protein in PBS for 1 hour at room temperature. Subsequently, available binding sites are blocked by incubation with 1% non-fat dry milk (NFM) in PBS (300 μ l/well). Chicken serum is diluted in PBS/1% NFM and added to the coated wells. After an incubation of 1 hour, the plates are washed 3 times with PBS/0.05% Tween 20 and bound Ig is detected using goat anti-human Ig conjugated with horseradish peroxidase. Conjugated goat antibody is detected using o-phenylenediamine dihydrochloride (Sigma) at 1 mg/ml. The calorimetric reaction is stopped by addition of 1 M HCl solution and the absorbance is measured at 490 nm. As a control, serum from non-immunized chicken is used. Serum from non-immunized chickens does not react with collagen adhesin protein. At a dilution of 1:250 the optical density measured in uncoated and collagen adhesin protein coated wells is below 0.2. In contrast, serum from immunized chickens contains humanized antibodies reactive with collagen adhesin. At a serum dilution of 1:250 the measured optical density is 2.3. Upon further dilution of the serum the measured optical density declines to 0.1 (at a dilution of 25600). No antibodies reactive with a goat anti-chicken IgG-HRP conjugate can be detected. This demonstrates that the genetically engineered chickens produce humanized anti-Staph. aureus collagen adhesin antibodies following immunization.

[0178] Genetically engineered rabbits are immunized intramuscularly with recombinant fragment of the *Staphylococcus aureus* collagen adhesin protein (100 μ g in incom-

plete Freund's adjuvant) on day 0 and day 14. On day 35 animals are bled and serum is prepared. ELISA plates (NUNC, Denmark) are coated with 2 µg/ml collagen adhesin protein in PBS for 1 hour at room temperature. Subsequently, available binding sites are blocked by incubation with 1% non-fat dry milk (NFM) in PBS (300 µl/well). Rabbit serum is diluted in PBS/1% NFM and added to the coated wells. After an incubation of 1 hour, the plates are washed 3 times with PBS/0.05% Tween 20 and bound Ig is detected using goat anti-human Ig conjugated with horseradish peroxidase. Conjugated goat antibody is detected using o-phenylenediamine dihydrochloride (Sigma) at 1 mg/ml. The colorimetric reaction is stopped by addition of 1 M HCl solution and the absorbance is measured at 490 nm. As a control, serum from non-immunized rabbit is used. Serum from non-immunized rabbits does not react with collagen adhesin protein. At a dilution of 1:250 the optical density measured in uncoated and collagen adhesin protein coated wells is below 0.2. In contrast, serum from immunized rabbits contains humanized antibodies reactive with collagen adhesin. At a serum dilution of 1:250 the measured optical density is 2.3. Upon further dilution of the serum the measured optical density declines to 0.1 (at a dilution of 25600). No antibodies reactive with a goat anti-rabbit IgG-HRP conjugate can be detected. This demonstrates that the genetically engineered rabbits produce humanized anti-*Staph. aureus* collagen adhesin antibodies following immunization.

EXAMPLE 18

Protection Against *Staphylococcus Aureus* Infection in a Mouse Model

[0179] Naive mice are passively immunized i.p. on day -1 with 16 mg of the immunoglobulin fraction containing antibodies specific for the *S. aureus* collagen adhesin protein (from Example 17) or with the immunoglobulin fraction from non-immunized animals. On day 0, the mice are challenged i.v. with 4×10^7 CFU *S. aureus* per mouse and mortality is monitored over the next 7 days. Mortality rate in the control groups is 80% and 10% in the group treated with the immunoglobulin fraction containing antibodies specific for the *S. aureus* collagen adhesin protein. The data indicate that anticollagen adhesin antibodies can protect mice against lethal *S. aureus* challenge.

EXAMPLE 19

Antigen-Specific Hybridomas Made from Transgenic Animals

[0180] Transgenic animals are immunized with an antigen (e.g., KLH, human red blood cells or sheep red blood cells). Spleen cells are removed at various times after immuniza-

tion and fused with myeloma cell lines derived from rabbit and chicken, respectively. After fusion cells are plated into 96 well plates and supernatants are tested for the presence of humanized antibodies. To demonstrate that the antibodies contain human immunoglobulin sequences, hybridomas are stained with fluorescent-labeled antibodies reactive with human heavy and light chain immunoglobulins. Limiting dilution is conducted to purify hybridomas to monoclonality.

EXAMPLE 20

Evaluation of Immunogenicity

[0181] Serum samples are collected from five cynomolgous monkeys on day 0. Subsequently, a purified partially human polyclonal antibody preparation (5 mg/kg) is administered into five cynomolgous monkeys by intravenous administration. The administration is repeated six times in bi-weekly intervals. Monkeys are monitored closely for any side-effects (e.g., anaphylactic shock, reflected by an elevated body temperature). After seven months serum is collected from blood samples. Affinity resins containing purified human IgG or partially human IgG are produced by standard procedure using CNBr-activated Sepharose. Monkey serum samples (3 ml) are added to the IgG-affinity resin (4 ml) containing 10 mg human or partially human IgG. Subsequently, the columns are washed with PBS. Bound monkey immunoglobulin is eluted from the column with 0.1M glycyl/HCl pH2.5 and dialyzed 2 times against PBS. The protein content of the eluted fractions is determined using the BCA assay using human IgG as a standard. The total amounts of protein in these fractions demonstrate that therapy with partially human IgG does not lead to a significant antibody response in the treated animals.

EXAMPLE 21

Treating Animals Using Humanized Antibodies

[0182] Humanized polyclonal immunoglobulins are purified from the serum of genetically engineered rabbits, or from egg yolk of genetically engineered chickens, by ammonium sulfate precipitation and ion exchange chromatography. SCID-mice are injected with one million human liver carcinoma cells expressing HBsAg. Subsequently, 25 µg immunoglobulin is injected peritoneally once per day. Animals treated with antibodies isolated from non-immunized rabbit serum die after about 60 days. This is similar to untreated recipients of liver carcinoma cells. In contrast, mice treated with antibodies isolated from immunized rabbit serum survive for more than 150 days. This demonstrates that human antibodies produced in genetically engineered rabbits are capable of eliminating human carcinoma cells from SCID-mice.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 58

<210> SEQ ID NO 1

<211> LENGTH: 29

<212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 1

cgcaagcttc ctacacgtgt gtggtgatg                29

<210> SEQ ID NO 2
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 2

cgcaagctta agatggwgat gtsgtcca                29

<210> SEQ ID NO 3
<211> LENGTH: 1714
<212> TYPE: DNA
<213> ORGANISM: cow

<400> SEQUENCE: 3

cctacacgtg tgtggtgatg cacgaaactt tacggaatca ctacaaagag aagtccacct    60
cgaggtctcc gggtaaatga gcctcgcgcc gctgatctag tggacgttcc ctcatccacc    120
caccctccc cccaccccg gctccaggtc cagccagggc gccctagccc ctccctgtgt    180
gcattctctc tgggcccgcg tgaataaagc acccaggccg ccctgggacc ctgcaacgct    240
gtgtgtggtc tttccgaggc agagcccttg tggccgccag gcctgcgggg gtgggctgag    300
ccgactctgg gccactttgt tcagcatctg tgggggagct gaccccactc cgggccagac    360
acacagttag tgggtccagc aggccacctg ggggctgccc aaggccacag aggggcttgg    420
ccagaggcac agctccacg tccccccag ccaccacctg ctggggccgc ctctggacag    480
gaaccgggga agcccccgag accctcaggg attgaggccc aatgcttccc gcctctgctc    540
cagcccacgc tgtggggcag ggccacatcc ttgtccccag gccctgttcc ttgggtgtcc    600
agagtcttgg tgtcactct gggctgcct ggagccacgc atggccaggg ggtggccctg    660
cttcaccctc aggtcccaa ggtcaggcct cgccctccct cggccaggag gctctgcccg    720
gtctccctg cccagggccg gccctgtgag cccatgggga ggtcatcct gtgctgaaa    780
gggttccag cagagagccc tgaatgtcca gggcagggac ctactgtctc cctgtggaca    840
cggagcccag agccacagac aacaagcccc agccccgcac gcacacgaga cagcccgcac    900
ccagcctcct ccacacgcac tcaggtgtac atgcgcacat gagcacactt caccocgtca    960
caccacaca cctacacaca ctcaggtctc gcaactcggg acccatgggg tgaccccacg    1020
ggcccagacc agagctgggt cttgtgagcc ctccctgtgg acaccagctg ggccccacc    1080
tccagcgcct atgggctgct cagcggccct tccccacact gaccacactg accaggtcag    1140
acatccgttc cttgcctccc ctgggacacc cacgcccctc cctagcaggc tgagatcccc    1200
cctcagcccc tcgtcctgac agcctcacc ctcgggcaca gcaccctca ggcccgtgc    1260
tgtcagccct ccctccccgg gggcagggcc caggaactg cgctctgctg acctccccag    1320
ctccaggcct ggcccccag gcagaggagg ccaggaactg agcctctgct ctgtggggag    1380
gtagggtcag ggtcccagct cagggcacag ctccaggtgg gagcaggacc ccacaggcca    1440

```

-continued

ggcccagata gcagccaggg ctggaggggt tggggctggg gctgggcccc agagactgac	1500
ctcaggtgac ccctgcctgg cccatgggga gatcacgcca ccttcccccc acccagaggg	1560
agccctgccc taccceagtg accctgccc gccctccgtg ggcagacaca gcaactgacca	1620
ccccccctg tgcagacttg ctgctggagg aggagatctg tgcggacgac ctggatgggg	1680
agctggacgg gctctggacc accatctcca tctt	1714

<210> SEQ ID NO 4
 <211> LENGTH: 1704
 <212> TYPE: DNA
 <213> ORGANISM: cow

<400> SEQUENCE: 4

cctacacgtg tgtggtgatg cacgaggccc tgcacaatca ctacacgag aagtccacct	60
ctaagtctgc gggtaaata gcctcacgtc cctgcaccag caagccctca cccagcccac	120
cctccccggg ctccaagtcc agccaggacg ccctagcccc tccctgtgtg cattcctcct	180
ggggcccggt gaataaagca cccagggcac cctgggaccc tgcaacgctg tgctggttct	240
ttccgaggca gagccctggt ggccgccagg cctgcagggg tgggctgagc cgactctggg	300
ccactttggt cagcatctgt gggggagctg accccgctcc gggccagaca cacagtgagt	360
gggtccagca ggccacctgg gggtgcccg agggccacgga ggggcttggc cagaggcgta	420
cctccacggc ccctccagc caccacctgc tgggcccggc tctggacagg aaccggggaa	480
gccccgaga ccctcaggga ttgaggccca atgcttccc cctctgctcc agcccagct	540
gtggggcagg gccacatcct tgtccccagg cccctgtccc tgggtgcccc gagtccctgt	600
gtccactctg ggctgcctg gagccacgag tggccggggg atggccctgc tccacctca	660
ggctcccaag gtcaggcctc gccctcccctc agccaggagg ctctgcccgg ctctccctgc	720
ccagggccag gcctgtgcgc ccattgggag gtcattccctg tgcctgaaag ggtccaggc	780
cgggagccct gaatgtccag ggcagggacc tagctgctcc ctgcagacac ggagcccaga	840
gccacagaca acaagcccc gcccgcacg cacacaagac agcccgcacc cagcctcctc	900
cacacgcact caggtgtgca tccgcacatg agcacacttc accccgtcac acccacagc	960
ctacacacac tcaggtctcg cactcgggga cccatggggg gaccccacag gccagacct	1020
agagctgggt cttgtgagcc ctcccctgtg acaccagctg gtcccccccc tccagcgccc	1080
atgggctgct cagtggccct tccccacct gaccacctg accaggtcag acatccgttc	1140
cttgcctccc ctggggcacc cacgcccctc cctagcaggc tgagatcccc cctcagcccc	1200
tgtcctggc accctcacc ctccaggcaca gggacacagc ccggcgctgt ctgccctccc	1260
tcccggggg cagggcccag gctcacatgc tctgtgacc ctcccggctc caggcctggc	1320
ccccagggca gaggaggcca ggaactgagc ctctgtcctg gggggagggt gggtcagggc	1380
cccagctcag ggcacagctc aggatgggaa caggacacca caggccaggc ccagacagt	1440
gccagggctg gagggggtgg gtctggggct ggggcccaga gactgacctc aggtgatccc	1500
tgcccagccc atggggggat cctgccacct tccccccacc cagagggagc cctgccccga	1560
ggccctgatg atgccacca gccccctgtg ggcagacaca gcaactgacca cccctccctg	1620
tgcagacctg ctgctggagg aggagatctg tgcggacgac caggacgggg agctggacgg	1680
gctctggacg accatcacca tctt	1704

-continued

<210> SEQ ID NO 5
<211> LENGTH: 1704
<212> TYPE: DNA
<213> ORGANISM: cow

<400> SEQUENCE: 5

```
cctacacgtg tgtggtgatg cacgaggccc tgcacaatca ctacacgcag aagtccacct    60
ctaagtctgc gggtaaatga gcctcacgtc cctgcaccag caagccctca cccagcccac    120
ctccccggg ctccaggctc agccaggacg ccctagcccc tccctgtgtg cattcctcct    180
gggcccggct gaataaagca cccagggccg cctgggacct tgcaacgctg tctggttct    240
ttccgaggca gagccctggt ggccgccagg cctgcggggg tgggctgagc cgactctggg    300
ccactttggt cagcatctgt gggggagctg accccactcc gggccagaca cacagtgagt    360
gggtccagca ggccacctgg gggctgccc aaggccacaga ggggcttggc cagaggcaca    420
gtccacaggt cccctccagc caccacctgc tgggcccggc tctggacagg aaccggggaa    480
gcccccgaga ccctcaggga ttgaggccca atgcttccc cctctgctcc agcccacgct    540
gtggggcagg gccacatcct tgtccccagg cccctgtcct tgggtgtcca gagtcttgt    600
gtccactctg ggctgcctg gagccacgca tggccagggg gtggccctgc ttcacctca    660
ggctcccaag gtcaggcctc gccctccctc ggccaggagg ctctgcccgg ctctccctgc    720
ccagggccag gcctgtgctc ccatggggag gtcacccctg tgctgaaag gggccaggc    780
cgagagccct gaatgtccag ggcagggacc tagctgctcc ctgtggacac ggagcccaga    840
gccacagaca acaagcccca gccccgcagc cacacgagac agcccacacc ccgctcctc    900
cacacgcact cagggtgtgca tccgcacatg agcacacttc accccatcac acccacagc    960
ctacacacac tcaggctctg cactcgggga cccatggggg gaccccacag gccagacct    1020
agagctgggt cttgtgagcc ctccctgtgg acaccagctg gtccccaccc tccagcgccc    1080
gtgggctgct cagcggctct tcccacact gaccacactg accaggctcag acatccgttc    1140
cttgctccc ctggggcacc catgcccctc cctagcaggc tgagatcccc cctcagcccc    1200
tcgtcctggc accctaccc ctcaggcaca gggacacagc ccggtgctgt ctgccctccc    1260
tccctggggg cagggcccag gctcacatgc tctgtgacc ctcccagctc caggcctggc    1320
ccccagggca gaggaggcca ggaactgagc ctctgtcctg gggggagggt gggtcagggc    1380
cccagctcag gccacagctc aggatgggag caggacacca caggccaggc ccagacagtg    1440
gccagggctg gaggggtggg gtctggggct gggcccaga gaatgacctc aggtgatccc    1500
tgccagccc atggggggat cctgccacct tccccccacc cagagggagc cctgcccga    1560
ggcctgatg atgccacca gcccccgctg ggcagacaca gactgaoca cccctcctg    1620
tgcagacctg ctgctggagg aggagatctg tgcggagccc caggacgggg agctggacgg    1680
gctctggacc accateacca tctt    1704
```

<210> SEQ ID NO 6
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 6

-continued

cgcggatccc ctacgcgtgt gtggtgatg 29

<210> SEQ ID NO 7
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:primer
 <400> SEQUENCE: 7

cgcggatcca ccgaggagaa gatccactt 29

<210> SEQ ID NO 8
 <211> LENGTH: 2842
 <212> TYPE: DNA
 <213> ORGANISM: sheep
 <400> SEQUENCE: 8

cctacgcgtg tgtggtgatg caccaggctc tacacaacca ctacacacag aagtcgatct 60
 ctaagcctcc gggtaaatga gccacatgcc cccgcaccag caagccctca cccagcccgc 120
 cctccccggg ctccaggctc agccaggacg ccctagcccc tccctgtgtg catgcctcct 180
 gggccgcat gaataaagca cccagggcgc cctgggaccc tgcaacgctg tgcttgttct 240
 ttccaggca gaggcctggt gaccgccagg cctgcggggg gtgggctgag cccactctgg 300
 gccgcttgtt tcagcatctg tgggggcgct gaccctctc cgggccagac acacagtgag 360
 tgggtccggc agggcacctg ggggtgccc gaggcctcgg aggggcttgg ccagagggcg 420
 agcttcacgg cccctccag ccaccacatt ctgggcccaga ctctgggag gaacggggga 480
 agccccgac acctcaggga ttgaggccca acgcttccc cctctgctcc agccccgct 540
 gaggggcagg gccgcggcct gtccccagg cccctgttcc tgggtgccc gagtccgtgt 600
 gtccactctg ggctgctg gagccagact ggcccagggg gaggccctgc ttcaccctca 660
 ggctcccag gtcaggcatc atcctcgtcg gccagtagct ctgctggct ctctctgccc 720
 ggggccaagc ctgtgtgccc atggggaggt cgtccctgtg cctgaaaagg gcccaggctg 780
 ggagccctga acgtccagg cagggaccta gctgctccct ggggacactg agcccagagc 840
 cccagacacc aagccccagc cccgcacgca caccagacag cccacacca gcgtcctcca 900
 caccgactca ggcgtccacc cgcacacaag catgcttcc cccgctcaca cccccatg 960
 cctgcacaca ctcaggctct acgctccggg acccatggag tgatcccacg ggcccagacc 1020
 cagagctggg tctcatgagc cctccctgtg gacaccagct ggtcccatt ctccagcgcc 1080
 cttgggctgc tcagtggccc ttccccacac tgaccacact gaccaggctc gacatccttc 1140
 ctgcctccc ctggggcacc caccgcccct cctgcaggc tgagacccc cctcagcccc 1200
 tcgtcctggc acctcaccct ctcgggcaca gggacacagc ccggcactgt ctgccctccc 1260
 tctcggggac agagcccagg cacgtgtgct ctgctgagcc tcccggctcc aggcctggcc 1320
 cccagggcag aggagggcag gaattgagcc tctgtcctgc ggggaggtgg ggtcagggcc 1380
 ccagctcagg gcacagctca gcatgggagc agggccccc agggcaggcc cagacagtgg 1440
 ccagggtggt ggctggggct ggggcccaga gactgacctc aggtgacccc tgcccggccc 1500
 atgggggatc acaccgcat ccccccgccc gcagagggag ccctgcccgc aagccccgat 1560
 ggccccgccc agcccccggt gggcagacac agcactgacc cccctccctg tgcagatctg 1620

-continued

```

ctgctggagg aggagagctg tgcggacgcc caggacgggg agctggacgg gctctggacg 1680
actatctcca tttcatcac gcccttcctg ctacagctct gctacagcgc caccgtgacc 1740
ctcttcaagg tgggggtcca cctctctggg ccctcgggcc ccctctctgt ccccagggtc 1800
ccccagagct ccctccctgc ccctcactgt ccctccctgt ccctctctgt ccctctctgt 1860
ccctctctgt ccctctctgt ccgttcattt tcccttcacc gtaagcttga gacagattgg 1920
ggtcatttca gagggcgtct gaagagtctc tgtgccgcac gcctcccttc atgtcagtgg 1980
ggagaattca gcaaggggtg agtgctgggt gagaaatgag gcttgcggcg ctcacgagca 2040
gtgatggggc actgctgctc cctgagacct gcgcggacac cgttttccat cgcaggagaa 2100
gcgggcaagg gaaaacgccc tcttggctct tcttgagtaa atgtcgcgtt ttggtcatca 2160
gtccctcccc cagtgaggct agaggagttt acttctccct ctcatgggc aggtcaggac 2220
tgtcatagac tccgatcac cttcctgtaa atgcttgctt tttgtgtgca gagagcctgt 2280
tttagctcgg gggctcctag ctcaactgac tcgcggggca ggggtgggct cgggctggcg 2340
ccgctgttc gggagcgc atccagcatg ctgtcgcaca gcttcgttgc taacaagacc 2400
gcttagtctc gtggttagac caacctgctt tctcgagtaa ttgttaattt acaggagttt 2460
cctgtatttt tcaacttata atcccctagt cagataactc tttaatcacc tattctgccc 2520
cttcattttc tccctatcga tctcagcaac ccataactgc cctcactgtc cttaaactgt 2580
cccttaactg accagactgt ccctcagtgt cccctcagag tcacctcctc atcacctcac 2640
tgtccctctc tgcccctctc tgcccctctc tgtccctccc tgcccctccc cgtcccctct 2700
ctgtccctct ctgcccctca ctgctcctct ctgcacctca ctgctcctca ctgcccctgg 2760
ggaggcccg c atcgagggt ctctgctcac cccgtccccc acccctgacc ccccgccagg 2820
tgaagtggat cttctcctcg gt 2842

```

```

<210> SEQ ID NO 9
<211> LENGTH: 2840
<212> TYPE: DNA
<213> ORGANISM: sheep

```

```

<400> SEQUENCE: 9

```

```

cctacgcgtg tgtggtgatg cacgaggctc tgcacaacca ctacacacag aagtcggtct 60
ctaagcctcc gggtaaatga gccacacgcc cccgcaccag caagccctca cccagcccgc 120
ctccccggg ctccaggtcc agccaggacg ccctagcccc tccctgtgtg catgctcct 180
gggcccgat gaataaagca cccagggccc cctgggaccc tgcagcgtg tgcgtgttct 240
ttccagggca gagccctggt gatcgccagg cctgcggggg gcgggctgag cccactctgg 300
gccgcttggc tcagcatctg tggggggcgt gaccctctc cgggccagac acacagtgg 360
tgggtccggc agggcacctg ggggtgccc gaggcctcgg aggggcttgg ccagaggcgc 420
agetccacgg cccccctcag ccaccacatt ctggggcaga ctctgggag gaacggggga 480
agccccgac acctcagggg ttgagggcca acgcttccc cctctgctcc agcccacgt 540
gaggggcagg gccgcggcct tgcctccagg cccctgttcc tgggtgcccc gagtccgtgt 600
gtccactctg ggctcctg gagccagact ggcccagggg gagccctgc ttcacctca 660
ggctcccag gtcaggcatc atcctctcgc gccagtagct ctgcttggct ctctctgccc 720
ggggccaagc ctgtgtgccc atggggaggt cgtccctgtg cctgaaaagg gccagggctg 780

```

-continued

ggagccctga acgtccaggg cagggacctg gctgctccct ggggacctg agcccagagc	840
cccagacacc aagccccagc cccgcacgca cacgagacag cccacaccca gcgtccctcca	900
caccactca ggcgccacc cgcacacaag catgcttcac ccccgtcaca ccccacatg	960
ctcgcacaca ctcaagtctc acgctccggg acccatggag tgatcccacg ggcccagacc	1020
cagagctggg tctcatgagc cctccctgtg gacaccagct ggtccccatc ctccagcgcc	1080
cttgggctgc tcagtggccc tttcccacac tgaccacact gaccaggcca gacatccttc	1140
ctcgcctccc ctggggcacc cacgcccctc ccttgcaagg tgagaccccc cctcagcccc	1200
tcgtcctggc accctcacc ctcgggcaca gggacacagc ccggcactgt ctgccctccc	1260
tctcggggac agagcccagg cacgtgtgct ctgctgagcc tcccggctcc aggcctggcc	1320
cccagggcag aggagggcag gaattgagcc tctgtcctgc ggggaggtgg ggtcagggcc	1380
ccagctcagg gcacagctca ggatgggagc aggacccac aggccaggcc cagacagtgg	1440
ccagggctgg ggctggggct ggggcccaga gactgacctc aggtgacccc tgcccggccc	1500
atgggggatc acaccgccat cccccccgcc gcagagggag ccctgccccg aagccccgat	1560
ggccccgcc agccccctg gggcagacac agcactgacc cccctccctg tgcagatctg	1620
ctgctggagg aggagagctg tgcggacgcc caggacgggg agctggacgg gctctggacg	1680
actatctcca tcttcatcac gctcttctg ctcagcgtct gctacagtgc caccgtgacc	1740
ctcttaagg tggggcccac ccctgctggg ccctcgggcc ccctctctgt ccccagggtc	1800
cccgcagagt ccctccctgc ccctcactgt ccctccctgt ccctctctgt ccctctctgt	1860
ccctctctgt ccctctctgt ccgctcattt tcccttcacc gtaagcttga gacagattgg	1920
ggtcatttca gagggcgtct gaagagtctc tgtgccgcac gcctcccttc atgtcagtgg	1980
ggagaattca gcaagggctg agtgtgggt gagaaatgag gcttgcggcg ctccagagca	2040
gtgatggggc actgctgctc cctgagacct gcgcggacac cgttttccat cgcaggagaa	2100
gcgggcaagg gaaaacgccc tcttggctctc tcttgagtaa atgtcgcggt ttggtcatca	2160
gtccctcccc cagtgggct agaggagttt acttctccct ctcatgggtc aggtcaggac	2220
tgtoatagac tccgatcac cttcctgtaa atgcttgctt tttgtgtgca gagcctgttt	2280
tagctcgggg gtctcagct cactgagctc gcggggcagg ggtgggctcg ggtggcgcc	2340
gcctgttcgg gagcggcatc tccagctgct gtcgcacagc ttcggtgcta acaagaccgc	2400
ttagtctcgt ggttagacca acctgcttcc togagtaatt gtaatttac aggagtttcc	2460
tgtatttttc aacttataat cccctagtc gataactctt taatcaccta ttctgcccct	2520
tcattttctc cctatcgatc tcagcaaccc atcactgccc tcaactgctc taaactgtcc	2580
cttaactgac cagactgtcc ctcagtgtcc cctcagatc acctccctat cacctcactg	2640
tcccctctctg cccctctctg cccctctctg tcccctctctg tcccctcccg tcccctctct	2700
gtccctctctg gcccctcact gctcctctct gcacctcact gctcctcact gcccctgggg	2760
aggcccgcac cagaggtgct ctgctcacc cgtcccccac cccgtccccc ccgccaggtg	2820
aagtggatct tctcctcggg	2840

<210> SEQ ID NO 10
 <211> LENGTH: 3120
 <212> TYPE: DNA
 <213> ORGANISM: rabbit

-continued

<400> SEQUENCE: 10

```

ctcccccca cgccgcagct gtgcaccccg cacacaaata aagcacccag ctctgcacctg    60
agaggctgtc ctgattcctt ccaaggcaga ggcttccact cgggccggac agggttgggc    120
ggggcccgctg ggctctgctg tggccagcag ccagaacggt caacagtggg acagggggcag    180
accacacaga caggggcctg ccaagaactg ggctcagccg gagtgtgtg gcaggtcccc    240
ccttgacgct agcacgtgtg tgctgggcag gcagaggccc ccaggggagg agcacacagc    300
taccacctct gcaagagcct ggctggcgc ccaggtccca gtccacaggg tgtgtagtac    360
acagagcctc atcttaccac agatgtaggg acagaccac cacgccctg caccaccacc    420
agcctcgccc ctgtggggac cagggctacc actccactcc cccgccaga gcagcagaag    480
caggtggcat cctcagcaga gggacagtct caccctcca cggcactgag ccctgacca    540
tcaaacaagc ccctcctgct gcacagcacc tgtgtgcaca tcacacacac acacacacac    600
acactgaggc ctgaccccat caaacaagcc cctcctgctg cacagcacct gtgtgcacat    660
cacacacaca cacacacaca cactgaggcc tgaccccatc ctgccctcct gctgcatggc    720
acctgtgtgc acatcacaca cacatgcaca cacacactca cacacactga gccctgacct    780
catcctgccc tcctgtgtga tggcacctgt gtgcacatca cacacacaca cacacacaca    840
cacacactga gccctgacct catcctgccc tcctgtgtga tggcacctgt gcacacatca    900
caaacacgcc tgccatac actggcactc agaaggggcc cctgtacacg catacacatg    960
cacacacctt gacacatggg cccctacac acgcatcaca cacactcatg cacactcctc   1020
acacatggcc ctctgcaca tacattgcac acacatgtgc acagacctca caatggggccc   1080
ctgcacacac attgtacaca cgcattgtga cacacttcac acatggggccc cctgacatg   1140
cattgcacac acagacacac acatgtgcat tcctcacaca ttggggcctt gcaagggatg   1200
ccctgcacac acattgcaca tgctcacatg tgacacacacc ccacactgga gccttgcata   1260
gggcccctg tacacacacc atgcatacac acacacctca cacaaggggc cccctacata   1320
cgaaaaacac acacacacac atgcacacac ctcatacacg ggccccctac acacatcaca   1380
cacacacaca cacacacgta tgcatgcctc acacacagac cttgcaaggg gcccccctgca   1440
catgcatcaa acacatatgc acatgtttca cacacacggt cccctacaca cactgcacac   1500
gcacacatgt gtacatgctt cacacactgg ggccttgcac ggggtcccct gcatagcata   1560
gcacccagag ccacgccagg tgccctggca catggacact ggtgcacaca cagcacccaa   1620
gcccagctct cccatccaag gggcaccagc acccccact cacgagcacc ctgaattcct   1680
gtccccaca agcgaacgtg caccctacct ctccagacgt ccctttcctg tggccactcc   1740
cataggtatt ggcgagaccc tcccttgacc cttgggcctg gtcaccacag ggacaggaga   1800
gggccaagtt gggccacagt accactgccc agcaggggtg aggcaagcag aggggtgggtc   1860
tgtgaggcgt ctggccagcc gtgctggggc ccaggtgggg agcagctggg tggctgaggt   1920
ggcttccttg caggtggttg ggggagctg gcccacaag tgccactgcc cagcactgtc   1980
cagtgcttcc ccctgaacct cccggccacc catccccagc tgacgccgca gagggagtgc   2040
ccctcggcct cctcggaag acgcacgctg actgcccctc cccatccaga gctgcagctg   2100
gacgagagct gtgccgagcc ccaggacggg gagctggacg ggtgtggac caccatcacc   2160
atcttcatct ccctcttctt gctcagcgtg tgctacagcg ccacagtcac cctcttcaag   2220

```

-continued

```

gtgggtgctg caccggcac ggggtggctg ggggccagg gcgggggccg ggggccaggc 2280
cctcctcacc ccgcccgcgc gctgctgcag gtgaagtgga tcttctcgtc cgtgggtggag 2340
ctgaaacaca ccatcgctcc cgactacagg aacatgatcg ggcagggggc ctaggccctt 2400
cgttctcaca gcctgcctcc ctggccagca ggagcccccg cctccgcctc ggaccccatg 2460
gctctctgct ctggccgctc cggacccctc cgcctcggga gaagcgcgca gctgatgcct 2520
gccggccctt ccacgcagca gtgcggacag cacgcatctg tcgtccaccg ggcaggaccc 2580
caccagggc cagccctgac cgcagcctc ctggactcag ggctcctctg agaaaaggcc 2640
cacttgttgg tcccctcagc ccacaccag gcagcctccg gtgggtgctt ccctggaccc 2700
cagcctgagg cctatgcttg ttctcctgtg gctcttactc agaggccctg gctggactcc 2760
caccacagc gacagtgcct tgctccaacc ctactgcac tgggggtcat ggggccacct 2820
tctgtgcagg ggtcctggct ccaaggagaa cactcgaagg gcctgcttgg ccacctggca 2880
ccacgggagc cccgctgggt agcttggcag ggacccctga gtagaggtgg gtgcaccag 2940
ccagaaagcc tgctggatgg acaggagcct ggcgtccggg cccagggcag gcagacagg 3000
cttcatggac aggagaggcc aaggaacatc agcaaagaga gacagctggg ccgggcgttc 3060
cagccagacc catcctgcag cccagcatcg gcccgtgta ctacagcagg gacagccagt 3120

```

```

<210> SEQ ID NO 11
<211> LENGTH: 2587
<212> TYPE: DNA
<213> ORGANISM: rabbit

```

```

<400> SEQUENCE: 11

```

```

gcgagacgcc tgccagggca ccgcccagtg ccctgaggcc cagcctcgcc gctccctccc 60
ctcagtggac ccattcccac cacagtcctc cagcccctcc cctcccggcc ctcacccctt 120
ccttggtctt aaccttgcga atgttgggtg gatggatgaa taaagtgaat ctttgcactt 180
gtgacttctc tctgcttctt catttaattg ttattactca tggtttccca gttgccctaa 240
agtcaccgcc atttcatcct ccatcccacc ctgcccctgt gtcctccggg agacaccact 300
ccctgaaacc cacagggccc tgtcttcaca ccgcccagcc cgaccacacg tgaggggctt 360
gcttcgtgtc tactccctc catcgagccc cagagtcctc ctttagtggt cttacagtca 420
catacagtta tacagtttga gtcaatccaa cctgccctgc caatttccca aaacaaagat 480
tttcagaata aaacagctat gaagaaagtc atttatggaa gcatgatata caacaacaaa 540
acaatgcaaa caacctaact gaataagcag agggaaatgt tcagacacac tatggggctt 600
gggcttcagt gagtattaca ccttcattac atttttaaac ttgtattaag gagctcctat 660
attacaagga ttatactaga gcactttcca tgacctaat aattctcatt acactgtgag 720
gttaaaagca ttagttaaaa tattgggcag gctccctata gccaacagtt gttcatattc 780
cataacccaa ccatcattta ggtgactcag ggtccttgtc caccaagaac tttggcaaga 840
atgttcagag caacttcctt tataaaagtc aaaaattgga agtaactcaa atgtctacca 900
acagtagaat gggctgttaa ttggcatatg tttacatatt agaatgctgt ttaataaaga 960
gaattaacaa actacaacta tccctaataa cataggtgac tcataaacat gatgttaagc 1020
acaagaaccc aaacacaaaa gacacactgt gtatgttttc atccatagga agttcaaac 1080
tagttaaaaa ttgaattaga aattgagatg aagtttactc ttggctgggg gtgtggagtg 1140

```

-continued

```

aggcgggtgcc tgggtggggga cagaaagtgg ctgctgggggt cttgggtgatg ttctagtcct 1200
cactgtgggtg tgtgctactc tgaaaatgta ttgagtacac aattaggttt tgtgctttca 1260
ttatactcca aagtaagttc tcataaacat tgccttacac ggggtctaca gataagagag 1320
actaagagga atgagtaaca gatcaaggcc acacagctgg taggcatggg cctgggatca 1380
aacctgtct gcccaattct gctctcttga gccctacact attctttcca gcaactggaat 1440
gccatgcaga acagggagta ggacatgcta cctccctagg gtctcctcct ttaccacact 1500
aacccaggagc acccatacat agaaacagga tggaaaagac catcagcaat ggaacaaggg 1560
agagattaac cttgttcagt attgtgatcc catgtaggaa agattgtggg aggaggctg 1620
cacacagagc accgtcccc ttctatgtgc ccaccgtct gtgccccctt atctgctcac 1680
ccgccagcg tgcattcact cagcaccctt ttgcctgcc ctctgaaaga ggtgcagaag 1740
taactaaacc agcttcctc cttcagtgc ttggaatcca gtttccctc tctatttccc 1800
cctccttttc agtgcaggag cctggagaaa tgtgatttgt gttattataa atttcccaca 1860
tcattttgtg taaggaaaa tatactcaac agtcataact ggtaaaactg ctgtgaaaac 1920
taagagaagt aattcatgcg aaggttgagc accagccttg tatatactaa gagatccaga 1980
agtgttagtc accgttagaa ataagaagga gtactcaat ttgactagtt cctggttcac 2040
tccttgaaca tgttcttcag ttatcatctt tcagtccaa atgattgaac ttggaattaa 2100
ctcacatgga ttctagacct gtgccgagaa tggctgccac tcgtgctcta gagctctggg 2160
gatgaggctg tccctactgt ggtgtgctac aggtctaaca acacaccagg ttttgaagac 2220
ttagcactat gaatatata atatatata ttccaataaa ttaacatac tttctacttt 2280
cattgcatgt tgagatagta atctactttg gatatatttg gttaaaccaa actattctca 2340
agacaaattt cataggttta tggttttttt acaatttaat caaaatataa acatagtcca 2400
aacaattaat ccatttaaag tggagaatgg cccaagtgtt tgggccctg ctaccattt 2460
ttaagaccag atgttctct tggcttctgg cttttgcttg gctcagcct ggccattgca 2520
gccatctgag gagtaaacag tggatggaag acatctcccc ccaccctgcc cataaagctc 2580
gggatcc 2587

```

```

<210> SEQ ID NO 12
<211> LENGTH: 1205
<212> TYPE: DNA
<213> ORGANISM: rabbit
<220> FEATURE:
<221> NAME/KEY: Unsure
<222> LOCATION: (997)
<223> OTHER INFORMATION: n at position 997 is uncertain
<220> FEATURE:
<221> NAME/KEY: Unsure
<222> LOCATION: (1127)
<223> OTHER INFORMATION: n at position 1127 is uncertain

```

```

<400> SEQUENCE: 12

```

```

cttcagcgtg aaccacgcc tcccgtgag tctcacgcac ccccggtgca ggcagtccg 60
cctcacctgg aaggtgcact gactgaagac actgcaaggg gtgagagcat ttctcaggaa 120
agagccctga gtttagaagg ccagagagca gagggctgag ggctgccttg cgctgcaacc 180
catgaaaca caggcttagc agatgttcaa gctccgggag tccacactgg gtgagggcag 240
gcgtccagcc tgacatggcc cccacagact cgcccacagg tgacgccaga tgaggacggt 300

```

-continued

```

caaggatcgg gggatcctac atgccacagg gcaccaagac agccaggaga gcaccagagg 360
ccacaagaga ggctcgggac agtctccctg ctgacatcca gagcccaggc cccacttggc 420
agagctggct gagaacacgt ctctcgggtg gaagctgccc cgtcctgggt gttgctcggc 480
gggctaagcc gactgacgcg gggcgggcca ggccatcggc cccacggcct gcagcttcct 540
ccccagccca ggccacgtgg gctcctggct gaactggccg ctgctgagc tctcaccccc 600
ccaccagca gcaggccggg cggtgctgcc atgagctcca tccccaccac acaagcgaca 660
ggccgggag cggccaggc ccacggggcg tttgctgtgc ggctcgcact cgtgctcag 720
ggccagcgca gggtgacgca gggactcacc aaccgcccc gactcggctg gcacgtttac 780
tggaggcctc tgagcctgac cgtggcagt gggcccagc aggctccagg ctgcccctg 840
cacctgggca ttgcccctcc gggaccctg gtgggacact tcccagatgt gctcccaccg 900
tgctccttg gggctctggg ctcatagcgg tcaactctcg ccttctctcc tcccagcct 960
ttctgcctc cctatggccc catctagctc tgccctntct agagcctcta cctggaagga 1020
atctgctgtt ggaccaagac accacccgca gcacaggtg gcgccttga ctgtgctagg 1080
ccctccccgc acagaaaagg gccctaggct ctggaggctg ctgctgnctc tggggctggc 1140
atcgggcgca cctgcaccc tgcaccctga ggaaactcag gcctgcccgc tccaggcctg 1200
tccct 1205

```

```

<210> SEQ ID NO 13
<211> LENGTH: 668
<212> TYPE: DNA
<213> ORGANISM: rabbit

```

```

<400> SEQUENCE: 13

```

```

gaagctttac ttgttggggg cgggcaggtc taagggaact gccagggtg ggggctgggc 60
ttgactcagc aggagccttc tagaaggaaa gctctggaga aggtgggggc agagggcggg 120
aaagccctgt gaggaggcgg gtggtgggca gggccactgg gaaggaggg ctgggggtga 180
cactcagggt ggcaactggg aggacctgag gaggcaggtg ccaggcacag agctgaacct 240
gggcagggca gggcagggta acaagaagga ttctccttg agcctggtcc aggggtgtcc 300
agggcgtcc agggcctggg gtttgcaagc tgggctgtga caggccctct ctcccaggg 360
gcaagcagca aagcctgggc acagagccca aagccccac acagagaagc tcccagggc 420
agggcctgca gggcttgggg gaccttctg gagcaggcag aggacagagg catgagatca 480
gcctcccaga ggctggaatg ataggtcag caggaggggc ccacatggc tctggttagc 540
aggagaaaac agccccagg tcccctggc caccacgac cgactgctg tgaagctttg 600
ggtggcagc gagagccaca tggcagctgc tctgtcact ctctgggagt gcatcgagg 660
gcgtcgac 668

```

```

<210> SEQ ID NO 14
<211> LENGTH: 45
<212> TYPE: PRT
<213> ORGANISM: camel

```

```

<400> SEQUENCE: 14

```

```

Glu Pro Leu Leu Glu Glu Glu Ser Cys Ala Glu Ala Gln Ser Gly Glu
  1           5           10           15
Leu Asp Gly Leu Trp Thr Thr Ile Ser Ile Phe Ile Thr Leu Phe Leu

```

-continued

```

                20           25           30
Leu Ser Val Cys Tyr Ser Ala Thr Val Thr Leu Phe Lys
                35           40           45

```

```

<210> SEQ ID NO 15
<211> LENGTH: 44
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 15

```

```

Glu Leu Gln Leu Glu Glu Ser Cys Ala Glu Ala Gln Asp Gly Glu Leu
 1           5           10           15
Asp Gly Leu Trp Thr Thr Ile Thr Ile Phe Ile Thr Leu Phe Leu Leu
                20           25           30
Ser Val Cys Tyr Ser Ala Thr Val Thr Phe Phe Lys
                35           40

```

```

<210> SEQ ID NO 16
<211> LENGTH: 44
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 16

```

```

Glu Leu Gln Leu Glu Glu Ser Cys Ala Glu Ala Gln Asp Gly Glu Leu
 1           5           10           15
Asp Gly Leu Trp Thr Thr Ile Thr Ile Leu Ile Thr Leu Phe Leu Leu
                20           25           30
Ser Val Cys Tyr Ser Ala Thr Val Thr Phe Phe Lys
                35           40

```

```

<210> SEQ ID NO 17
<211> LENGTH: 44
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 17

```

```

Glu Leu Gln Leu Glu Glu Ser Cys Ala Glu Ala Gln Asp Gly Glu Leu
 1           5           10           15
Asp Gly Leu Trp Thr Thr Ile Thr Ile Phe Ile Thr Leu Phe Leu Leu
                20           25           30
Ser Val Cys Tyr Ser Ala Thr Val Thr Phe Phe Lys
                35           40

```

```

<210> SEQ ID NO 18
<211> LENGTH: 44
<212> TYPE: PRT
<213> ORGANISM: mouse

```

```

<400> SEQUENCE: 18

```

```

Gly Leu Gln Leu Asp Glu Thr Cys Ala Glu Ala Gln Asp Gly Glu Leu
 1           5           10           15
Asp Gly Leu Trp Thr Thr Ile Thr Ile Phe Ile Ser Leu Phe Leu Leu
                20           25           30
Ser Val Cys Tyr Ser Ala Ala Val Thr Leu Phe Lys
                35           40

```

```

<210> SEQ ID NO 19
<211> LENGTH: 44

```

-continued

<212> TYPE: PRT
 <213> ORGANISM: mouse
 <400> SEQUENCE: 19
 Gly Leu Asp Leu Asp Asp Val Cys Ala Glu Ala Gln Asp Gly Glu Leu
 1 5 10 15
 Asp Gly Leu Trp Thr Thr Ile Thr Ile Phe Ile Ser Leu Phe Leu Leu
 20 25 30
 Ser Val Cys Tyr Ser Ala Ser Val Thr Leu Phe Lys
 35 40

<210> SEQ ID NO 20
 <211> LENGTH: 45
 <212> TYPE: PRT
 <213> ORGANISM: mouse
 <400> SEQUENCE: 20
 Pro Gly Leu Gln Leu Asp Glu Thr Cys Ala Glu Ala Gln Asp Gly Glu
 1 5 10 15
 Leu Asp Gly Leu Trp Thr Thr Ile Thr Ile Phe Ile Ser Leu Phe Leu
 20 25 30
 Leu Ser Val Cys Tyr Ser Ala Ala Val Thr Leu Phe Lys
 35 40 45

<210> SEQ ID NO 21
 <211> LENGTH: 44
 <212> TYPE: PRT
 <213> ORGANISM: mouse
 <400> SEQUENCE: 21
 Glu Leu Glu Leu Asn Glu Thr Cys Ala Glu Ala Gln Asp Gly Glu Leu
 1 5 10 15
 Asp Gly Leu Trp Thr Thr Ile Thr Ile Phe Ile Ser Leu Phe Leu Leu
 20 25 30
 Ser Val Cys Tyr Ser Ala Ser Val Thr Leu Phe Lys
 35 40

<210> SEQ ID NO 22
 <211> LENGTH: 44
 <212> TYPE: PRT
 <213> ORGANISM: mouse
 <400> SEQUENCE: 22
 Glu Leu Glu Leu Asn Gly Thr Cys Ala Glu Ala Gln Asp Gly Glu Leu
 1 5 10 15
 Asp Gly Leu Trp Thr Thr Ile Thr Ile Phe Ile Ser Leu Phe Leu Leu
 20 25 30
 Ser Val Cys Tyr Ser Ala Ser Val Thr Leu Phe Lys
 35 40

<210> SEQ ID NO 23
 <211> LENGTH: 44
 <212> TYPE: PRT
 <213> ORGANISM: sheep
 <400> SEQUENCE: 23
 Leu Leu Leu Glu Glu Glu Ser Cys Ala Asp Ala Gln Asp Gly Glu Leu
 1 5 10 15

-continued

Pro Asp Tyr Arg Asn Met Ile Gly Gln Gly Ser
 20 25

<210> SEQ ID NO 29
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

Val Lys Trp Ile Phe Ser Ser Val Val Asp Leu Lys Gln Thr Ile Ile
 1 5 10 15

Pro Asp Tyr Arg Asn Met Ile Gly Gln Gly Ala
 20 25

<210> SEQ ID NO 30
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

Val Lys Trp Ile Phe Ser Ser Val Val Asp Leu Lys Gln Thr Ile Ile
 1 5 10 15

Pro Asp Tyr Arg Asn Met Ile Gly Gln Gly Ala
 20 25

<210> SEQ ID NO 31
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

Val Lys Trp Ile Phe Ser Ser Val Val Asp Leu Lys Gln Thr Ile Ile
 1 5 10 15

Pro Asp Tyr Arg Asn Met Ile Gly Gln Gly Ala
 20 25

<210> SEQ ID NO 32
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: mouse

<400> SEQUENCE: 32

Val Lys Trp Ile Phe Ser Ser Val Val Glu Leu Lys Gln Thr Leu Val
 1 5 10 15

Pro Glu Tyr Lys Asn Met Ile Gly Gln Ala Pro
 20 25

<210> SEQ ID NO 33
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: mouse

<400> SEQUENCE: 33

Val Lys Trp Ile Phe Ser Ser Val Val Glu Leu Lys Gln Thr Ile Ser
 1 5 10 15

Pro Asp Tyr Arg Asn Met Ile Gly Gln Gly Ala
 20 25

<210> SEQ ID NO 34

-continued

<211> LENGTH: 27
 <212> TYPE: PRT
 <213> ORGANISM: mouse

<400> SEQUENCE: 34

Val Lys Trp Ile Phe Ser Ser Val Val Glu Leu Lys Gln Thr Leu Val
 1 5 10 15

Pro Glu Tyr Lys Asn Met Ile Gly Gln Ala Pro
 20 25

<210> SEQ ID NO 35
 <211> LENGTH: 27
 <212> TYPE: PRT
 <213> ORGANISM: mouse

<400> SEQUENCE: 35

Val Lys Trp Ile Phe Ser Ser Val Val Gln Val Lys Gln Thr Ala Ile
 1 5 10 15

Pro Asp Tyr Arg Asn Met Ile Gly Gln Gly Ala
 20 25

<210> SEQ ID NO 36
 <211> LENGTH: 27
 <212> TYPE: PRT
 <213> ORGANISM: mouse

<400> SEQUENCE: 36

Val Lys Trp Ile Phe Ser Ser Val Val Gln Val Lys Gln Thr Ala Ile
 1 5 10 15

Pro Asp Tyr Arg Asn Met Ile Gly Gln Gly Ala
 20 25

<210> SEQ ID NO 37
 <211> LENGTH: 27
 <212> TYPE: PRT
 <213> ORGANISM: rabbit

<400> SEQUENCE: 37

Val Lys Trp Ile Phe Ser Ser Val Val Glu Leu Lys His Thr Ile Ala
 1 5 10 15

Pro Asp Tyr Arg Asn Met Met Gly Gln Gly Ala
 20 25

<210> SEQ ID NO 38
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: sheep

<400> SEQUENCE: 38

Val Lys Trp Ile Phe Ser Ser Val
 1 5

<210> SEQ ID NO 39
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 39

cgcaagcttc ctacacgtgt gtggtgatg

-continued

<210> SEQ ID NO 40
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 40

gtcgcgccc ctcgatgcac tcccagag 28

<210> SEQ ID NO 41
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 41

ggtaccctct cctccccca cgccgcagc 29

<210> SEQ ID NO 42
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 42

atatctcaga actggctgtc cctgctgtag tacacgg 37

<210> SEQ ID NO 43
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 43

gtcgcactg gacgctgaac ctcgcgg 27

<210> SEQ ID NO 44
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 44

ggtaccgggg gcttgccggc cgtcgcac 28

<210> SEQ ID NO 45
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 45

gcggccgctg gcgaggagac caagctggag atcaaacy 38

<210> SEQ ID NO 46
<211> LENGTH: 36

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 46
gtcgacgcag cccaaagctg ttgcaatggg gcagcg          36

<210> SEQ ID NO 47
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 47
atatggtacc gcgagacgcc tgccagggca ccgcc          35

<210> SEQ ID NO 48
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 48
ggatcccgag ctttatgggc aggggtgggg          30

<210> SEQ ID NO 49
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 49
atatgtcgac ctgggataag catgctgttt tctgtctgtc cc          42

<210> SEQ ID NO 50
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer

<400> SEQUENCE: 50
ctaggtacca gcaggtgggg gcacttctcc c          31

<210> SEQ ID NO 51
<211> LENGTH: 1719
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:human
immunoglobulin heavy chain C?1 gene segment
flanked by nucleotides derived from the rabbit
heavy chain.

<400> SEQUENCE: 51
tgacctacct accctgcaa ggtcaggggt cctccaaggc aagggatcac atggcaccac          60
ctctcttgca gcctccacca agggcccatc ggtcttcccc ctggcacccct cctccaagag          120
cacctctggg ggcacagcgg ccctgggctg cctgggtcaag gactacttcc ccgaaccggt          180

```

-continued

gacggtgctg tggaaactcag gcgcctgac cagcggcgtg cacaccttcc cggctgtcct	240
acagtccctca ggactctact ccctcagcag cgtggtgacc gtgccctcca gcagcttggg	300
caccagacc tacatctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa	360
agttggtgag aggccagcac agggaggag ggtgtctgct ggaagccagg ctacagcctc	420
ctgcctggac goatcccggc tatgcagccc cagtccaggg cagcaaggca ggccccgtct	480
gcctcttcac ccggaggcct ctgcccggcc cactcatgct cagggagagg gtcttctggc	540
tttttcccca ggctctgggc aggcacagcc taggtgcccc taaccaggc cctgcacaca	600
aaggggcagg tgctgggctc agacctgcca agagccatat ccgggaggac cctgcccctg	660
acctaagccc accccaaagg ccaaactctc cactcccctca gctcggacac cttctctcct	720
cccagattcc agtaactccc aatcttctct ctgcagagcc caaatcttgt gacaaaactc	780
acacatgccc accgtgccc a gtaagccag cccaggcctc gccctccagc tcaaggcggg	840
acaggtgccc tagagtagcc tgcattccagg gacaggcccc agccgggtgc tgacacgtcc	900
acctccatct cttcctcagc acctgaactc ctggggggac cgtcagctct cctcttcccc	960
ccaaaaccca aggacaccct catgatctcc cggaccctg aggtcacatg cgtggtggtg	1020
gacgtgagcc acgaagacc tgaggtaag ttcaactggt acgtggacgg cgtggagggtg	1080
cataatgcca agacaaagcc gcgggaggag cagtacaaca gcacgtaccg tgtggtcagc	1140
gtcctcaccg tcctgcacca ggactggctg aatggcaagg agtacaagtg caaggctctc	1200
aacaaagccc tcccagcccc catcgagaaa accatctcca aagccaaagg tgggaccctg	1260
ggggtgctgag ggccacatgg acagaggccg gctcggccca ccctctgccc tgagagtgac	1320
cgctgtacca acctctgtcc ctacagggca gccccgagaa ccacaggtgt acaccctgcc	1380
cccatcccgg gatgagctga ccaagaacca ggtcagcctg acctgcctgg tcaaaggctt	1440
ctatcccagc gacatcgccc tggagtggga gagcaatggg cagccggaga acaactaaa	1500
gaccacgcct cccgtgctgg actccgacgg ctccttcttc ctctacagca agctcaccgt	1560
ggacaagagc aggtgacagc aggggaacgt cttctcatgc tccgtgatgc atgaggctct	1620
gcacaaccac tacacgcaga agagcctctc cctgtctccg ggtaaagtag cgctgtgccc	1680
gcgagctgcc cctctccctc cccccacgc cgcagctgt	1719

<210> SEQ ID NO 52

<211> LENGTH: 390

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: a VH gene segment encoding a human VH element polypeptide sequence, with flanking sequences derived from rabbit immunoglobulin DNA sequences.

<400> SEQUENCE: 52

tgagtgcagc tgcctgacc atgtcgtctg tgtttgcagg tgtccagtgt gaggtgcagc	60
tggtggagtc cggggagggt ctcgtccagc caggggggac cctgagactc acctgcgcag	120
ctcttgatt caccttcagt agctatgcaa tgagctgggt ccgccaggct ccaggggaagg	180
ggctggaatg ggtcggagcc attagtggta gtggtagcac atactacgcg gacagcgtga	240
aagggcgatt caccatctcc agagacaact ccaagaacac gctgtatctg caaatgaaca	300
gtctgagagc cgaggacagc gccgcctatt actgtgcgaa agacacagtg aggggcctc	360

-continued

aggctgagcc cagacacaaa cctccctgca 390

<210> SEQ ID NO 53
 <211> LENGTH: 445
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:containing a human immunoglobulin light chain C??gene segment flanked by 50 nucleotides derived from the rabbit light chain immunoglobulin Kappal gene.

<400> SEQUENCE: 53
 catccacatg gcaccaggg gagatgtcca ctggtaccta agcctcgcca tcctgtttgc 60
 ttctttcctc aggaactgtg gctgcacat ctgtcttcat cttcccgcca tctgatgagc 120
 agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taacttctat cccagagagg 180
 ccaaagtaca gtggaagggtg gataacgccc tccaatcggg taactcccag gagagtgtca 240
 cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg ctgagcaaag 300
 cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc ctgagctcgc 360
 ccgtcacaaa gagcttcaac aggggagagt gttagagcga gacgcctgcc agggcaccgc 420
 cagcgaccct gagggcccgc ctgcg 445

<210> SEQ ID NO 54
 <211> LENGTH: 632
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:a V??gene segment encoding a human V??element polypeptide sequence, flanking by sequences derived from rabbit immunoglobulin DNA.

<400> SEQUENCE: 54
 ggcaggctgc tcccacccca tgcaggaggc agtaccaggc aggaccagc atggacatga 60
 ggggtccctgc tcagctcctg ggactcctgc tgctctggct cccaggtaag gagggaaaca 120
 acaaaaattt tattcagcca gtgtagccac taatgcctgg cacttcagga aattcttctt 180
 agaacattac taatcatgtg gatatgtgtt tttatgttcc taatatacaga taccagatgt 240
 tacatccaga tgaccagtc tccatcctct ctgtctgcat ctgtgggaga cagagtcacc 300
 atcacttgcc gagccagtca gggcattagc aattacttag cctggatca gcagaaacca 360
 ggaaggttc ccaagctcct gatttatgct goatccactt tgcaatctgg ggtccatcg 420
 cggttcagtg gcagtggtgc tgggacagat ttcactctta ccatcagcag cctgcagcct 480
 gaagatgttg ccacctatta ctgtcaaaag tacaacagtg cccctccact tttcggcgga 540
 gggaccaagg tggagatcaa acgtaagtgc actttcctaa tgttcctcac cgtttctgcc 600
 tgatttgttt gctttttcca ttttttcgct at 632

<210> SEQ ID NO 55
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:primer
 <400> SEQUENCE: 55

-continued

```

catcacacagc catacatacgc cgtgtggccg ctctgcctct ctcttgacag tatggacagc    60
aagcgaaccg                                                                    70

```

```

<210> SEQ ID NO 56
<211> LENGTH: 71
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:primer
<400> SEQUENCE: 56

```

```

atcaggggtga cccctacggtt acactcctgt caccaaggag tgggagggac tcagaagaac    60
tcgtcaagaa g                                                                    71

```

```

<210> SEQ ID NO 57
<211> LENGTH: 419
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:a gene
encoding human immunoglobulin light chain constant region
Clambda2 flanked by nucleotides derived from the chicken light
chain gene.

```

```

<400> SEQUENCE: 57

```

```

catcacacagc catacatacgc cgtgtggccg ctctgcctct ctcttgacag tcagcccaag    60
gtgccccct cgtcactct gttcccgccc tcctctgagg agcttcaagc caacaaggcc    120
acactggtgt gtctcataag tgactttctac cggggagccg tgacagtggc ttggaaagca    180
gatagcagcc ccgtcaaggc gggagtggag accaccacac cctccaaaca aagcaacaac    240
aagtacgagg ccagcagcta tctgagcctg acgcctgagc agtggaaagtc ccacagaagc    300
tacagctgcc aggtcacgca tgaagggagc accgtggaga agacagtggc ccctacagaa    360
tgttcatagt agtcccactg gggatgcaat gtgaggacag tggttcctca cctccctg    419

```

```

<210> SEQ ID NO 58
<211> LENGTH: 416
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence:a VJ gene
segment encoding a human VJ immunoglobulin
polypeptide, with flanking sequences derived from
chicken immunoglobulin DNA.

```

```

<400> SEQUENCE: 58

```

```

ttgcccgtttt ctcccctctc tcctctccct ctccaggttc cctgggtcag tcagtgtgta    60
ctcagccgcc ctccggtgtca gcagccccgg gacaagaagt cacgatctcc tgctccgggt    120
ctagtagcaa cattggcgtat aatttctgtct cttggtacca gcagctgctt ggcactgccc    180
ctaagcttct gatctatgat aacaacaaga gaccctcggg catccctgac cgattctccg    240
gttccaaatc cggcacctca gccacattag gcatcactgg gctccaaacc ggcgacgagg    300
ctgactatta ctgtgggact tgggacagca gcctttctgt tggatggttt gggggcggga    360
cacgcgtgac cgtcctaggt gagtcgtgta cctcgtctcg gtctttcttc ccccat    416

```

What is claimed:

1. A humanized immunoglobulin molecule comprising at least a portion of a human immunoglobulin heavy chain constant (C) region, and variable region amino acids sequences encoded by more than one immunoglobulin variable (V) gene segment.

2. The humanized immunoglobulin molecule of claim 1 which is derived from a nonhuman transgenic animal relying primarily on gene conversion and/or somatic hypermutation to generate antibody diversity.

3. The humanized immunoglobulin molecule of claim 2 wherein said transgenic animal is a rabbit, chicken, sheep or cow.

4. The humanized immunoglobulin molecule of claim 1 wherein said human immunoglobulin heavy chain C region sequence is a C_γ, C_κ, or C_λ sequence.

5. The humanized immunoglobulin molecule of any one of claims 1-4, which is specific for an antigen.

6. The humanized immunoglobulin molecule of claim 5, wherein said antigen is a microorganism selected from bacterium, fungus, or virus; an antigenic portion of said organism; an antigenic molecule derived from said microorganism; or a tumor-associated antigen.

7. The humanized immunoglobulin molecule of claim 6, wherein said bacterium is selected from *S. aureus*, *Pseudomonas aeruginosa*, *enterococcus*, *enterobacter*, and *Klebsiella pneumoniae*.

8. The humanized immunoglobulin molecule of claim 6, wherein said fungus is selected from *Candida albicans*, *Candida parapsilosis*, *Candida tropicalis*, and *Cryptococcus neoformans*.

9. The humanized immunoglobulin molecule of claim 6, wherein said virus is selected from respiratory syncytial virus

(RSV), Hepatitis C virus (HCV), Hepatitis virus (HBV), cytomegalovirus (CMV), EBV, and HSV.

10. The humanized immunoglobulin molecule of claim 6, wherein said antigen is selected from Her-2-neu antigen, CD20, CD22, CD53, prostate specific membrane antigen (PMSA), and 17-1A molecule.

11. A monoclonal antibody preparation comprising a humanized immunoglobulin molecule of claim 5.

12. A monoclonal antibody preparation comprising a humanized immunoglobulin molecule of any one of claims 6-10.

13. The monoclonal antibody preparation of claim 11 which is substantially non-immunogenic to human.

14. The monoclonal antibody preparation of claim 12 which is substantially non-immunogenic to human.

15. A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and the monoclonal antibody preparation of claim 11.

16. A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and the monoclonal antibody preparation of claim 12.

17. A method of treating a disease in a human subject comprising administering to said subject a therapeutically effective amount of a pharmaceutical composition of claim 15.

18. The method of claim 17 wherein said disease is caused by bacterial, fungal or viral infection, or said disease is a cancer.

* * * * *