



US 20050186205A1

(19) **United States**

(12) **Patent Application Publication**

**Anderson et al.**

(10) **Pub. No.: US 2005/0186205 A1**

(43) **Pub. Date: Aug. 25, 2005**

(54) **THERAPEUTIC APPLICATION OF CHIMERIC AND RADIOLABELED ANTIBODIES TO HUMAN B LYMPHOCYTE RESTRICTED DIFFERENTIATION ANTIGEN FOR TREATMENT OF B CELL LYMPHOMA**

(76) Inventors: **Darrell R. Anderson**, Escondido, CA (US); **Nabil Hanna**, Olivenhain, CA (US); **John E. Leonard**, Encinitas, CA (US); **Roland A. Newman**, San Diego, CA (US); **Mitchell E. Reff**, San Diego, CA (US); **William H. Rastetter**, Rancho Santa Fe, CA (US)

Correspondence Address:

**SIDLEY AUSTIN BROWN & WOOD LLP**  
**1501 K STREET, NW**  
**WASHINGTON, DC 20009 (US)**

(21) Appl. No.: **10/956,039**

(22) Filed: **Oct. 4, 2004**

**Related U.S. Application Data**

(60) Continuation of application No. 09/905,928, filed on Jul. 17, 2001, now abandoned, which is a continua-

tion of application No. 08/475,813, filed on Jun. 7, 1995, now Pat. No. 6,682,734, which is a division of application No. 08/149,099, filed on Nov. 3, 1993, now Pat. No. 5,736,137, which is a continuation-in-part of application No. 07/978,891, filed on Nov. 13, 1992, now abandoned.

**Publication Classification**

(51) **Int. Cl.<sup>7</sup>** ..... **A61K 39/395**; A61K 31/704; A61K 31/573; A61K 31/4745; A61K 31/66

(52) **U.S. Cl.** ..... **424/144.1**; 514/34; 514/171; 514/283; 514/109

(57) **ABSTRACT**

Disclosed herein are therapeutic treatment protocols designed for the treatment of B cell lymphoma. These protocols are based upon therapeutic strategies which include the use of administration of immunologically active mouse/human chim eric anti-CD20 antibodies, radiolabeled anti-CD20 antibodies, and cooperative strategies comprising the use of chimeric anti-CD20 antibodies and radiolabeled anti-CD20 antibodies.




FIG. 1

LINKER #1 15bp | SV40 ORIGIN=332bp  
 GACGTCGCGG CCGCTCTAGG CCTCCAAAAA AGCCTCCTCA CTACTTCTGG AATAGCTCAG 60  
 AGGCCGAGGC GGCCTCGGCC TCTGCATAAA TAAAAAAAAT TAGTCAGCCA TGCATGGGGC 120  
 GGAGAATGGG CGGAACCTGGG CGGAGTTAGG GGCGGGATGG GCGGAGTTAG GGGCGGGACT 180  
 ATGGTTGCTG ACTAATTGAG ATGCATGCTT TGCATACTTC TGCTGCTGG GGAGCCTGGG 240  
 GACTTTCCAC ACCTGGTTGC TGACTAATTG AGATGCATGC TTTGCATACT TCTGCCTGCT 300  
 GGGGAGCCTG GGGACTTTCC ACACCCCTAAC TGACACACAT TCCACAGAAAT TAATTCCCT LINKER #2=13bp 360 1  
 347 8 360  
 AGTTATTAAAT AGTAATCAAT TACGGGGTCA TTAGTTCAT A GCCCATATAT GGAGTTCCGC 420  
 GTTACATAAC TTACGGTAAA TGGCCCGCCT GGCTGACCGC CCAACGACCC CCGCCCATTG 480  
 CMV PROMOTER-ENHANCER=567bp  
 ACGTCAATAA TGACGTATGT TCCCATAGTA ACGCCAAATAG GGACTTCCA TTGACGTCAA 540  
 TGGGTGGACT ATTTACGGTA AACTGCCAC TTGGCAGTAC ATCAAGTGT A TCATATGCCA 600  
 AGTACGCCCT CTATTGACGT CAATGACGGT AAATGGCCG CCTGGCATT TGCCCAGTAC 660  
 ATGACCTTAT GGGACTTTCC TACTTGGCAG TACATCTACG TATTAGTCAT CGCTATTACC 720  
 ATGGTGTGATGC GGTTTGGCA GTACATCAAT GGGCGTGGAT AGCGGTTGA CTCACGGGG 780  
 TTTCCAAGTC TCCACCCCAT TGACGTCAAT GGGAGTTGT TTTGGCACCA AAATCAACGG 840  
 GACTTTCCAA AATGTCGTA CAACTCCGCC CCATTGACGC AAATGGCCG TAGGCGTGT 900  
 CCGTGGGAGG TCTATATAAG CAGAGCTGGG TACGTGAACC GTCAAGATGC CTGGAGACGC 960  
 727 8  
 Bgl II | LEADER=60bp  
 CATCACAGAT CTCTCACCAT GAGGGTCCCC GCTCAGCTCC TGGGCTCCT GCTGCTCTGG 1020  
 978 9  
 +1 101 102 107 108  
 CTCCCAGGTG CACGATGTGA TGGTACCAAG GTGGAAATCA AAAGTACCGT GGCTGCACCA 1080  
 1038 9 1062 3 Bsi WI  
 TCTGTCTTCA TCTTCCCGCC ATCTGATGAG CAGTTGAAAT CTGGAACTGC CTCTGTTGTG 1140  
 TGCCTGCTGA ATAACCTCTA TCCCAGAGAG GCCAAAGTAC AGTGGAAAGGT GGATAACGCC 1200  
 HUMAN KAPPA CONSTANT 324bp 107 AMINO ACID & STOP CODON  
 CTCCAATCGG GTAATCTCCA GGAGAGTGTGTC ACAGAGCAGG ACAGCAAGGA CAGCACCTAC 1260  
 AGCCTCAGCA GCACCCCTGAC GCTGAGCAAA GCAGACTACG AGAAACACAA AGTCTACGCC 1320  
 TGCAGTCAGTCAGGG CCTGAGCTCG CCCGTACAA AGAGCTTCAA CAGGGGAGAG 1380  
 STOP  
 LIGHT  
 CHAIN | Eco RI LINKER #4=85bp  
 TGTGAATTC AGATCCGTTA ACGGTTACCA ACTACCTAGA CTGGATTCTG GACAACATGC 1440  
 1386 17  
 GGCCTGTATA TCTACGTATG ATCAGCCTCG ACTGTGCCCT CTAGTTGCCA GCCATCTGTT 1500  
 1471 2

FIG. 2A

GTTTGCCCT CCCCCGTGCC TTCCTGACC CTGGAAAGGTG CCACTCCCAC TGTCCCTTCC 1560  
 TAATAAAATG AGGAAATTGC ATCGCATTGT CTGAGTAGGT GTCATTCTAT TCTGGGGGAT 1620  
 GGGGTGGGC AGGACAGCAA GGGGGAGGAT TGGGAAGACA ATAGCAGGCA TGCTGGGAT 1680  
 GCGGTGGGCT CTATGGAACC AGCTGGGCT CGACAGCTAT GCCAAGTACG CCCCCTATT: 1740  
 ACGTCAATGA CGGTAAATGG CCCGCCTGGC ATTATGCCA GTACATGACC TTATGGGACT 1800  
 TTCCTACTTG GCAGTACATC TACGTATTAG TCATGCTAT TACCATGGTG ATGCGGTTT 1860  
 CMV PROMOTER-ENHANCER=334bp  
 GGCAGTACAT CAATGGCGT GGATAGCGGT TTGACTCACG GGGATTTCCA AGTCTCCACC 1920  
 CCATTGACGT CAATGGGAGT TTGTTTGGC ACCAAAATCA ACGGGACTTT CCAAAATGTC 1980  
 GTAACAACTC CGCCCCATTG ACGCAAATGG GCGGTAGGCG TGTACGGTGG GAGGTCTATA 2040  
 TAAGCAGAGC TGGGTACGTC CTCACATTCA GTGATCAGCA CTGAACACAG ACCCGTCGAC 2100  
 ATGGTTGGA GCCTCATCTT GCTCTTCCTT GTCGCTGTTG CTACGCGTGT CGCTAGCACC 2160  
 START HEAVY CHAIN -5 -4 -3 114 115  
 AAGGGCCCAT CGGTCTTCCC CCTGGCACCC TCCCTCAAGA GCACCTCTGG GGGCACACG 2220  
 GCCCTGGGCT GCCTGGTCAA GGACTACTTC CCCGAACCGG TGACGGTGTC GTGGAACTCA 2280  
 GGCGCCCTGA CCAGGGCGT GCACACCTTC CCGGCTGTCC TACAGTCCTC AGGACTCTAC 2340  
 HUMAN GAMMA 1 CONSTANT  
 TCCCTCAGCA GCGTGGTGAC CGTCCCTCC AGCAGCTTG GCACCCAGAC CTACATCTGC 2400  
 993bp=330 AMINO ACID & STOP CODON  
 AACGTGAATC ACAAGCCCAG CAACACCAAG GTGGACAAGA AAGCAGAGCC CAAATCTTGT 2460  
 GACAAAACTC ACACATGCC ACCGTGCCCA GCACCTGAAC TCCCTGGGGGG ACCGTAGTC 2520  
 TTCCTCTTCC CCCCCAAAACC CAAGGACACC CTCATGATCT CCCGGACCCC TGAGGTCACA 2580  
 TGGGTGGTGG TGGACGTGAG CCACGAAGAC CCTGAGGTCA AGTTCAACTG GTACGTGGAC 2640  
 GGCCTGGAGG TGCATAATGC CAAGACAAAG CCGCGGGAGG AGCAGTACAA CAGCACGTAC 2700  
 CGTGTGGTCA GCGTCCTCAC CGTCCCTGCAC CAGGACTGGC TGAATGGCAA GGACTACAAG 2760  
 TGCAAGGTCT CCAACAAAGC CCTCCCAGCC CCCATCGAGA AAACCATCTC CAAAGCCAAA 2820  
 GGGCAGCCCC GAGAACACCA GGTGTACACC CTGCCCCAT CCCGGGATGA GCTGACCAGG 2880  
 AACCAGGTCA GCCTGACCTG CCTGGTCAAA GGCTTCTATC CCAGCGACAT CGCCGTGGAG 2940  
 TGGGAGAGCA ATGGGAGGCC GGAGAACAC TACAAGACCA CGCCTCCCGT GCTGGACTCC 3000

FIG. 2B

GACGGCTCCT TCTTCCTCTA CAGCAAGCTC ACCGTGGACA AGAGCAGGTG GCAGCAGGGG 3060  
 AACGTCTTCT CATGCTCCGT GATGCATGAG GCTCTGCACA ACCACTACAC GCAGAAGAGC 3120  
 STOP HEAVY CHAIN Bam HI LINKER #7=81bp  
 CTCTCCCTGT CTCCGGGTAA ATGAGGATCC GTTAACGGTT ACCAACTACCC TAGACTGGAT 3180  
 3144 5  
 TCGTGACAAC ATGCGGCCGT GATATCTACG TATGATCAGC CTCGACTGTG CCTTCTAGTT 3240  
 3225 6  
 GCCAGCCATC TGTTGTTGC CCCTCCCCCG TGCCCTCCTT GACCCCTGGAA GGTGCCACTC 3300  
 BOVINE GROWTH HORMONE POLYADENYLATION REGION=231bp  
 CCACTGTCTT TTCTAATAA AATGAGGAAA TTGCATCGCA TTGTCTGAGT AGGTGTCATT 3360  
 CTATTCTGGG GGGTGGGGTG GGGCAGGACA GCAAGGGGGA GGATTGGGAA GACAATACCA 3420  
 LINKER #8=34bp  
 GGCATGCTGG GGATGCGGTG GGCTCTATGG AACCAGCTGG GGCTCGACAG CGCTGGATCT 3480  
 3456 7  
 CCCGATCCCC AGCTTGCTT CTCATTCT TATTCATA ATGAGAAAAA AAGGAAAATT 3540  
 3490 1  
 AATTTTAACA CCAATTCACT AGTTGATTGA GCAAATGCGT TGCCAAAAAG GATGCTTTAG 3600  
 MOUSE BETA GLOBIN MAJOR PROMOTER=366bp  
 AGACAGTGTT CTCTGCACAG ATAAGGACAA ACATTATTCA GAGGGAGTAC CCAGAGCTGA 3660  
 GACTCTAAG CCAGTGAGTG GCACAGCATT CTAGGGAGAA ATATGCTTGT CATCACCGAA 3720  
 GCCTGATTCC GTAGAGCCAC ACCTTGGTAA GGGCCAATCT GCTCACACAG GATAGAGAGG 3780  
 GCAGGAGCCA GGGCAGAGCA TATAAGGTGA GGTAGGATCA GTTGCTCCTC ACATTTGCTT 3840  
 LINKER #9=19bp 5' UNTRANSLATED DHFR=82bp  
 CTGACATAGT TGTGTTGGGA GCTTGGATAG CTTGGACAGC TCAGGGCTGC GATTCGCGC 3900  
 3856 7 3875 6  
 START DHFR  
 CAAACTTGAC GGCAATCTA GCGTGAAGGC TGGTAGGATT TTATCCCCGC TGCCATCA 3960  
 3957 8  
 GTTCGACCAT TGAACTGCAT CGTCGCCGTG TCCCCAAAATA TGGGGATTGG CAAGAA~~CGGA~~ 4020  
 GACCTACCCCT GGCTCCGCT CAGGAACGAG TTCAAGTACT TCCAAAGAAT GACCACAAACC 4080  
 TCTTCAGTGG AAGGTAACAA GAATCTGGTG ATTATGGTA GGAAAACCTG GTTCTCCATT 4140  
 MOUSE DHFR=564bp=187 AMINO ACID & STOP CODON  
 CCTGAGAAGA ATCGACCTT AAAGGACAGA ATTAATATAG TTCTCAGTAG AGAACTCAAA 4200  
 GAACCACAC GAGGAGCTCA TTTTCTTGCC AAAAGTTGG ATGATGCCTT AAGACTTATT 4260  
 GAACAAACCGG AATTGGCAAG TAAAGTAGAC ATGGTTGGA TAGTCGGAGG CAGTTCTGTT 4320  
 TACCAAGGAAG CCATGAATCA ACCAGGCCAC CTTAGACTCT TTGTGACAAG GATCATGCAG 4380  
 GAATTTGAAA GTGACACGTT TTTCCCAGAA ATTGATTTGG GGAAATATAA ACTTCTCCCA 4440  
 GAATACCCAG GCGTCCTCTC TGAGGTCCAG GAGGAAAAAG GCATCAAGTA TAAGTTGAA 4500

FIG. 2C

STOP DHFR  
 GTCTACGAGA AGAAAGAC~~TA~~ ACAGGAAGAT GCTTTCAAGT TCTCTGCTCC CCTCCTAAAG 4560  
 4521 2

3' UNTRANSLATED DHFR=82bp LINKER #10=10bp  
 TCATGCATT TTATAAGACC ATGGGACTTT TGCTGGCTTT AGATCAGCCT CGACTGTGCC 4620  
 4603 4 4613 4

TTCTAGTTGC CAGCCATCTG TTGTTTGCCTC CTCCCCGTG CCTTCCTTGA CCCTGGAAAGG 4680  
 BOVINE GROWTH HORMONE POLYADENYLATION REGION=231bp  
 TGCCACTCCC ACTGTCCTT CCTAATAAAA TGAGGAAATT GCATCGCATT GTCTGAGTAG 4740

GTGTCATTCT ATTCTGGGGG GTGGGGTGGG GCAGGACAGC AAGGGGGAGG ATTGGGAAGA 4800  
 CAATAGCAGG CATGCTGGGG ATGCGGTGGG CTCTATGGAA CCAGCTGGGG CTCGAGCTAC 4860  
 4844 5

TAGCTTGCT TCTCAATTTC TTATTTGCAT AATGAGAAAA AAAGGAAAAT TAATTTAAC 4920

ACCAATTCA TAGTTGATTG AGCAAATGCG TTGCCAAAAA GGATGCTTTA GAGACAGTGT 4980  
 MOUSE BETA GLOBIN MAJOR PROMOTER=366bp  
 TCTCTGCACA GATAAGGACA AACATTATTC AGAGGGAGTA CCCAGAGCTG AGACTCCTAA 5040

GCCAGTGAGT GGCACACGCAT TCTAGGGAGA AATATGCTTG TCATCACCGA AGCCTGATTG 5100

CGTAGAGCCA CACCTTGGTA AGGGCCAATC TGCTCACACA GGATAGAGAG GGCAGGAGCC 5160

AGGGCAGAGC ATATAAGGTG AGGTAGGATC AGTTGCTCCT CACATTGCT TCTGACATAG 5220  
 LINKER #12=21bp START NEO  
 TTGTGTTGGG AGCTTGGATC GATCCTCT~~TA~~ ATGGTGAACAA GATGGATTGC ACGCAGGTTG 5280  
 5227 8 5248 9

TCCGGCCGCT TGGGTGGAGA GGCTATTGGG CTATGACTGG GCACAAACAGA CAATGGCTG 5340

CTCTGATGCC GCCGTGTTCC GGCTGTCAGC GCAGGGGCGC CCGGTTCTTT TTGTCAAGAC 5400  
 NEOMYCIN PHOSPHOTRANSFERASE  
 CGACCTGTCC GGTGCCCTGA ATGAACATGCA GGACGAGGCA GCGGGGCTAT CGTGGCTGGC 5460

795bp=264 AMINO ACIDS & STOP CODON  
 CACGACGGGC GTTCCTGCG CAGCTGTGCT CGACGTTGTC ACTGAAGCGG GAAGGGACTG 5520

GCTGCTATTG GGCGAAGTGC CGGGGCAGGA TCTCCTGTCA TCTCACCTG CTCCCTGCCG 5580

GAAAGTATCC ATCATGGCTG ATGCAATGCG GCGGCTGCAT ACGCTTGATC CGGCTACCTG 5640

CCCATTCGAC CACCAAGCGA AACATCGCAT CGAGCGAGCA CGTACTCGGA TGGAAAGCCGG 5700

TCTTGTGAT CAGGATGATC TGGACGAAGA GCATCAGGGG CTCGCGCCAG CCGAACTGTT 5760

CGCCAGGCTC AAGGCCGCA TGCCCGACGG CGAGGATCTC GTCTGACCC ATGGCGATGC 5820

CTGCTTGGCG AATATCATGG TGAAAATGG CCGCTTTCT GGATTCAATCG ACTGTGGCCG 5880

GCTGGCTGTG GCGGACCGCT ATCAGGACAT AGCGTTGGCT ACCCGTGTATA TTGCTGAAGA 5940

GCTTGGCCGC GAATGGGCTG ACCGCTTCT CGTGCTTTAC GGTATGCCG CT~~T~~CCCGATTG 6000

FIG. 2D

STOP NEO  
 GCAGCGCATC GCCTTCTATC GCCTTCTTGA CGAGTTCTTC TGAGCGGGAC TCTGGGGTTC 6060  
 6043 4

GAAATGACCG ACCAACCGAC GCCCAACCTG CCATCACGAG ATTCGATTC CACCGCCGCC 6120  
 3' UNTRANSLATED NEO=173bp  
 TTCTATGAAA GGTTGGGCTT CGGAATCGTT TTCCGGGACG CGGGCTGGAT GATCCTCCAG 6180  
 CGCGGGGATC TCATGCTGGA GTTCTTCGCC CACCCCAACT TGTTTATTCC AGCTTATAAT 6240  
 6216 7

GGTTACAAAT AAAGCAATAG CATCACAAAT TTCACAAATA AAGCATTTC TTCACTGCAT 6300  
 SV40 POLY A EARLY=133bp LINKER #13=19bp  
 TCTAGTTGTG GTTTGTCCAA ACTCATCAAT CTATCTTATC ATGTCGGAT CGCGGCCGCG 6360  
 6349 50

ATCCCGTGA GAGCTTGGCG TAATCATGGT CATAGCTGTT TCCGTGTGA AATTGTTATC 6420  
 6368 9

CGCTCACAAAT TCCACACAAAC ATACGAGCCG GAAGCATAAA GTGAAAGCC TGGGGTGCCT 6480  
 AATGAGTGAG CTAACTCACA TTAATTGCGT TCGCCTCACT GCCCCCTTC CAGTCGGGAA 6540  
 ACCTGTCGTG CCAGCTGCAT TAATGAATCG GCCAACGCGC GGGGAGAGGC GGTTTGCCTA 6600  
 PVC 19  
 TTGGGCGCTC TTCCGCTTCC TCGCTCACTG ACTCGCTGCG CTCGGTCGTT CGGCTGCGGC 6660  
 GAGCGGTATC AGCTCACTCA AAGGCGGTAAC TACGGTTATC CACAGAATCA GGGGATAACG 6720  
 CAGGAAAGAA CATGTGAGCA AAAGGCCAGC AAAAGGCCAG GAACCGTAAA AAGGCCGCGT 6780  
 6792=BACTERIAL ORIGIN OF REPLICATION  
 TGCTGGCGTT ~~T~~TCCATAGG CTCCGCCCGG CTGACGAGCA TCACAAAAAT CGACGCTCAA 6840  
 GTCAGAGGTG GCGAAACCCG ACAGGACTAT AAAGATAACCA GGCCTTCCC CCTGGAAAGCT 6900  
 CCCTCGTGCCTC CTCTCCTGTT CGAACCTGC CGCTTACCGG ATACCTGTCC GCCTTCTCC 6960  
 CTTCGGGAAG CGTGGCGCTT TCTCAATGCT CACGCTGTAG GTATCTCAGT TCGGTGTAGG 7020  
 TCGTTCGCTC CAAGCTGGGC TGTGTGCACG AACCCCCCGT TCAGCCCCGAC CGCTGCGCCT 7080  
 TATCCGGTAA CTATCGTCTT GAGTCCAACC CGGTAAGACA CGACTTATCG CCACTGGCAG 7140  
 CAGCCACTGG TAACAGGATT AGCAGAGCGA GGTATGTAGG CGGTCTACAGA GAGTTCTTGA 7200  
 AGTGGTGGCC TAACTACGGC TACACTAGAA GGACAGTATT TGGTATCTGC GCTCTGCTGA 7260  
 AGCCAGTTAC CTTGGAAAAA AGAGTTGGTA GCTCTTGATC CGGAAACAA ACCACCGCTG 7320  
 GTAGCGGTGG TTTTTTGTT TGCAAGCAGC AGATTACGGG CAGAAAAAAA CGATCTCAAG 7380  
 AAGATCCTTT GATCTTTCT ACGGGGTCTG ACGCTCAGTG GAAAGAAAAC TCACGTTAAC 7440  
 GGATTTGGT CATGAGATTA TCAAAAAGGA TCTTCACCTA GATCCTTTA AATTAAAAAT 7500

FIG. 2E

STOP BETA LACTAMASE  
GAAGTTTAA ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT TACCAATGCT 7560  
7550  
TAATCAGTGA GGCACCTATC TCAGCGATCT GTCTATTTCG TTCAATCCATA GTTGCCTGAC 7620  
TCCCCGTCGT GTAGATAACT ACGATAACGGG AGGGCTTACC ATCTGGCCCC AGTGCTGCAA 7680  
TGATACCGCG AGACCCACGC TCACCGGCTC CAGATTATC AGCAATAAAC CAGCCAGCCG 7740  
BETA LACTAMASE=861bp  
GAAGGGCCGA GCGCAGAAGT GGTCTGCAA CTTTATCCGC CTCCATCCAG TCTATTAATT 7800  
286 AMINO ACID & STOP CODON  
GTTGCCGGGA AGCTAGAGTA AGTAGTTCGC CAGTTAATAG TTTGCGCAAC GTTGTTCCTA 7860  
TTGCTACAGG CATCGTGGTG TCACGCTCGT CGTTTGGTAT GGCTTCATTG AGCTCCGGTT 7920  
CCCAACGATC AAGGCCAGTT ACATGATCCC CCATGTTGTG CAAAAAAAGCG GTTAGCTCCT 7980  
TCGGTCCTCC GATCGTTGTC AGAAGTAAGT TGGCCGCAGT GTTATCACTC ATGGTTATGG 8040  
CAGCACTGCA TAATTCTCTT ACTGTCATGC CATCCGTAAG ATGCTTTCT GTGACTGGTG 8100  
AGTACTCAAC CAAGTCATTC TGAGAATAGT GTATGCCGG ACCGAGTTGC TCTTGCCCGG 8160  
CGTCAATAACG GGATAATACC GCGCCACATA GCAGAACTTT AAAAGTGCTC ATCATTGGAA 8220  
AACGTTCTTC GGGGGCAAAA CTCTCAAGGA TCTTACCGCT GTTGAGATCC AGTTCGATGT 8280  
AACCCACTCG TGCAACCAAC TGATCTTCAG CATCTTTAC TTTCACCAGC GTTTCTGGGT 8340  
GAGCAAAAAC AGGAAGGCAA AATGCCGCAA AAAAGGGAAT AACGGCGACA CGGAAATGTT 8400  
START BETA LACTAMASE  
GAATACTCAT ACTCTTCCTT TTCAATATT ATTGAAGCAT TTATCAGGGT TATTGTCTCA 8460  
8410  
TGAGCGGATA CATATTGAA TGTATTTAGA AAAATAAAC AATAGGGGTT CCGCCGACAT 8520  
TTCCCCGAAA AGTGCCACCT

FIG. 2F

**LINKER #1=15bp**  
 GACGTCGCGG CCGCTCTAGG CCTCCAAAAA AGCCTCCTCA CTACTTCTGG AATAGCTAG 60  
 15 6

AGGCCGAGGC GGCGCTCGGCC TCTGCATAAA TAAAAAAAAT TAGTCAGCCA TGCATGGGC 120  
 SV40 ORIGIN=332bp  
 GGAGAAATGGG CGGAACTGGG CGGAGTTAGG GGCGGGATGG GCGGAGTTAG GGGCGGGACT 180

ATGGTTGCTG ACTAATTGAG ATGCATGCTT TGCACTACTC TGCTGCTGG GGAGCCTGGG 240

GACTTTCCAC ACCTGGTTGC TGACTAATTG AGATGCATGC TTTGCATACT TCTGCCTGCT 300

GGGGAGCCTG GGGACTTTCC ACACCCCTAAC TGACACACAT TCCACAGAAAT TAATTCCCCCT 360  
 347 8

**LINKER #2=13bp**  
 AGTTATTAAT AGTAATCAAT TACGGGGTCA TTAGTTCATATA CCCATATAT GGAGTTCCGC 420

GTTACATAAC TTACGGTAAA TGGCCCGCCT GGCTGACCGC CCAACGACCC CCGCCCCATTG 480

ACGTCAATAA TGACGTATGT TCCCAGTAGTA ACGCCAATAG GGACTTTCCA TTGACGTCAA 540

**CVM PROMOTER-ENHANCER=567bp**  
 TGGGTGGACT ATTTACGGTA AACTGCCAC TTGGCAGTAC ATCAAGTGTA TCATATGCCA 600

AGTACGCCCC CTATTGACGT CAATGACGGT AAATGGCCCC CCTGGCATT TGCCCAGTAC 660

ATGACCTTAT GGGACTTTCC TACTTGGCAG TACATCTACG TATTAGTCAT CGCTATTACC 720

ATGGTGTGATGC GGTTTGGCA GTACATCAAT GGGCGTGGAT AGCGGTTGA CTCACGGGG 780

TTTCCAAGTC TCCACCCCAT TGACGTCAAT GGGAGTTGT TTTGGCACCA AAATCAACGG 840

GACTTTCCAA AATGTGTAA CAACTCCGCC CCATTGACGC AAATGGGGGG TAGGGGTGTA 900

**LINKER #3=7bp**  
 CGGTGGGAGG TCTATATAAG CAGAGCTGGG TACGTGAACC GTCAGATCGC CTGGAGACGC 960  
 927 8 934 5

**Bgl 2** **START LIGHT CHAIN** **NATURAL LEADER=66bp**  
 CATCACAGAT CTCTCACTAT GGATTTCAAG GTGCAGATTAA TCAGCTTCCT GCTAACTAGT 1020  
 978 9

GCTTCAGTCA TAATGTCCAG AGGACAAATT GTTCTCTCCC AGTCTCCAGC AATCCTGTCT 1080  
 1044 5<sup>+1</sup>

GCATCTCCAG GGGAGAAGGT CACAATGACT TGCAGGGCCA GCTGAAGTGT AAGTTACATC 1140

CACTGGTTCC AGCAGAAGCC AGGATCCTCC CCCAAACCCCT GGATTTATGC CACATCCAAC 1200

**LIGHT CHAIN VARIABLE REGION 318bp 106 AMINO ACID**  
 CTGGCTTCTG GAGTCCCTGT TCGCTTCAGT GGCAGTGGGT CTGGGACTTC TTACTCTCTC 1260

ACCATCAGCA GAGTGGAGGC TGAAGATGCT GCCACTTATT ACTGCCAGCA GTGGACTAGT 1320

**BsiWI**  
 AACCCACCCA CGTTGGAGG GGGGACCAAG CTGGAAATCA AACGTACGGT GGCTGCACCA 1380  
 1362 3

TCTGTCTTCA TCTTCCCGCC ATCTGATGAG CAGTTGAAAT CTGGAACTGC CTCTGTTGTG 1440

TGCCTGCTGA ATAACCTCTA TCCCAGAGAG GCCAAAGTAC AGTGGAAAGGT GGATAACGCC 1500

*FIG. 3A*

HUMAN KAPPA CONSTANT=324bp=107 AMINO ACID & STOP CODON  
 CTCCAATCGG GTAACTCCCA GGAGAGTGT ACAGAGCAGG ACAGCAAGGA CAGCACCTAC 1560  
 AGCCTCAGCA GCACCCCTGAC GCTGAGCAAA GCAGACTACG AGAAACACAA AGTCTACGCC 1620  
 TGCAGTCA CCCATCAGGG CCTGAGCTCG CCCGTCACAA AGAGCTTCAA CAGGGGAGAG 1680  
 STOP  
 LIGHT  
 CHAIN Eco RI LINKER #4=81bp  
 TGT ~~GAATT~~ AGATCCGTTA ACGGTTACCA ACTACCTAGA CTGGATTCTGT GACAACATGC 1740  
 1646 7  
 GGCGTGATA TCTACGTATG ATCAGCCTCG ~~ACTGTGCCTT~~ CTAGTTGCCA GCCATCTGTT 1800  
 1771 2  
 GTTGGCCCT CCCCCGTGCC TTCTTGACC CTGGAAAGGTG CCACTCCCAC TGTCTTTCC 1860  
 TAATAAAATG AGGAAATTGC ATCGCATTGT CTGAGTAGGT GTCATTCTAT TCTGGGGGGT 1920  
 BOVINE GROWTH HORMONE POLYADENYLATION REGION=231bp  
 GGGGTGGGGC AGGACACCAA GGGGGAGGAT TGGGAAGACA ATAGCAGGCA TGCTGGGGAT 1980  
 LINKER #5=15bp  
 GCGGTGGGCT CTATGGAACC AGCTGGGCT CGACAGCTAT GCCAAGTACG CCCCCTATTG 2040  
 2002 3 2017 8  
 ACGTCAATGA CGGTAAATGG CCCGCCTGGC ATTATGCCCA GTACATGACC TTATGGACT 2100  
 TTCCTACTTG GCAGTACATC TACGTATTAG TCATCGCTAT TACCATGGTG ATGCGTTTT 2160  
 CMV PROMOTER-ENHANCER=334bp  
 GGCAGTACAT CAATGGCGT GGATAGCGGT TTGACTCACG GGGATTCCA AGTCTCCACC 2220  
 CCATTGACGT CAATGGAGT TTGTTTGGC ACCAAAATCA ACGGGACTTT CCAAAATGTC 2280  
 GTAACAACTC CGCCCCATTG ACCCAAATGG GCGGTAGGGG TGTACGGTGG GAGGTCTATA 2340  
 LINKER #6=7bp Sal I  
 TAAGCAGAGC ~~TGGGTACGTC~~ CTCACATTCA GTGATCAGCA CTGAACACAG ACCCGTCGAC 2400  
 START 2351 2 2358 9  
 HEAVY CHAIN SYNTHETIC & NATURAL LEADER Mlu I 2457 8  
 ATGGGTTGGA GCCTCATCTT GCTCTCCCTT GTCGCTGTTG ~~CTACGGGTGT CCTGTCAG~~ 2460  
 2401 -5 -4 -3 -2 -1 +1  
 GTACAACCTGC AGCAGCCTGG GGCTGAGCTG GTGAAGCCTG GGGCCTCAGT GAAGATGTCC 2520  
 TGCAAGGCTT CTGGCTACAC ATTACCACT TACAATATGC ACTGGTAAA ACAGACACCT 2580  
 HEAVY CHAIN VARIABLE=363bp=121 AMINO ACID  
 GGTGGGGGCC TGGAAATGGAT TGGAGCTATT TATCCCCGAA ATGGTGTACAC TTCTACAAT 2640  
 CAGAAGTTCA AAGGCAAGGC CACATTGACT GCAGACAAAT CCTCCAGCAC AGCCTACATG 2700  
 CAGCTCAGCA CCCTGACATC TGAGGACTCT GCGGTCTATT ACTGTGCAAG ATCGACTTAC 2760  
 TACGGCGGTG ACTGGTACTT CAATGTCTGG GGCGCAGGGA CCACGGTCAC CGTCTCTGCA 2820  
Nhe I  
 GCTACACCA AGGGCCCATC GGTCTTCCCC CTGGCACCCCT CCTCCAAGAG CACCTCTGG 2880  
 GGCACAGCGG CCCTGGGCTG CCTGGTCAAG GACTACTTCC CGGAACCGGT GACGGTGTCC 2940  
 HUMAN GAMMA 1 CONSTANT=993bp  
 TGGAACCTCAG GCGCCCTGAC CAGCGGGCGTG CACACCTTCC CGGCTGTCTT ACAGTCTCA 3000

FIG. 3B

330 AMINO ACID & STOP CODON

GGACTCTACT CCCTCAGCAAG CGTGGTGACC GTGCCCTCCA GCAGCTTGGG CACCCAGACC 3060  
 TACATCTGCA ACGTGAATCA CAAGCCCAGC AACACCAAGG TGGACAAGAA AGCAGAGCCC 3120  
 AAATCTTGTG ACAAAACTCA CACATGCCCA CCGTGCCAG CACCTGAAC TCTGGGGGA 3180  
 CCGTCAGTCT TCCTCTTCCC CCCAAAACCC AAGGACACCC TCATGATCTC CCGGACCCCT 3240  
 GAGGTCACAT GCGTGGTGGT GGACGTGAGC CACGAAGACC CTGAGGTCAA GTTCAACTGG 3300  
 TACGTGGACG GCGTGGAGGT GCATAATGCC AAGACAAAGC CGCGGGAGGA GCAGTACAAC 3360  
 AGCACGTACC GTGTGGTCAG CGTCCTCACC GTCCCTGCACC AGGACTGGCT GAATGGCAAG 3420  
 GAGTACAAGT GCAAGGTCTC CAACAAAGCC CTCCCAGGCC CCATCGAGAA AACCATCTCC 3480  
 AAAGCCAAAG GGCAGCCCCG AGAACACACAG GTGTACACCC TGCCCCCATC CCGGGATGAG 3540  
 CTGACCAAGA ACCAGGTCAAG CCTGACCTGC CTGGTCAAAG GCTTCTATCC CAGCGACATC 3600  
 GCGGTGGAGT GGGAGAGCAA TGGGCAGCCG GAGAACAACT ACAAGACCAC GCCTCCCGTG 3660  
 CTGGACTCCG ACGGCTCCTT CTTCTCTAC AGCAAGCTCA CCGTGGACAA GAGCAGGTGG 3720  
 CAGCAGGGGA ACGTCTTCTC ATGCTCCGTG ATGCATGAGG CTCTGCACAA CCACTACACG 3780  

STOP HEAVY CHAIN Bam HI LINKER #7=81bp

CAGAAGAGCC TCTCCCTGTC TCCGGGTAAA TGAGGATCCG TTAACGGTTA CCAACTACCT 3840  
 3813 4

AGACTGGATT CGTGACAACA TGGGGCCGTG ATATCTACGT ATGATCAGCC TCGACTGTGC 3900  
 3894 5

CTTCTAGTTG CCAGCCATCT GTTGTGGCC CCTCCCCCGT GCCTTCCTTG ACCCTGGAA 3960

GTGCCACTCC CACTGTCCTT TCCTAATAAA ATGAGGAAAT TGATCGCAT TGTCTGAGTA 4020

BOVINE GROWTH HORMONE POLYADENYLATION REGION=231bp

GGTGTCAATT TATTCTGGGG GGTGGGGTGG GGCAGGACAG CAAGGGGGAG GATTGGGAAG 4080

LINKER #8=34bp

ACAATAGCAG GCATGCTGGG GATGCGGTGG GCTCTATGGA ACCAGCTGGG GCTCGACAGC 4140  
 4125 6

GCTGGATCTC CCGATCCCCA GCTTTGCTTC TCAATTCTT ATTTGCATAA TGAGAAAAAA 4200

AGGAAAATTA ATTTAACAC CAATTCACTA GTTGATTGAG CAAATGCGTT GCCAAAAAGG 4260

MOUSE BETA GLOBIN MAJOR PROMOTER=366bp

ATGCTTTAGA GACAGTGGTC TCTGCACAGA TAAGGACAAA CATTATTCAAG AGGGAGTACC 4320

CAGAGCTGAG ACTCCTAACGC CAGTGAGTGG CACAGCATTG TAGGGAGAAA TATGCTTGTC 4380

ATCACCGAAG CCTGATTCCG TAGAGCCACA CCTTGGTAAG GGCACATCTG CTCACACAGG 4440

ATAGAGAGGG CAGGAGCCAG GGCAGAGCAT ATAAGGTGAG GTAGGATCAAG TTGCTCCTCA 4500

*FIG. 3C*

CATTTGCTTC TGACATAGTT **LINKER #9=19bp** 15' UNTRANSLATED DHFR=82bp  
 4525 6 4544 5 4560  
 ATTTGGCGCC AAACTTGACG GCAATCCTAG CGTGAAGGCT GGTAGGATT TATCCCCGCT 4620  
**START DHFR**  
 GCCATC**ATG** TTGACCATT GAACTGCATC GTCGCCGTGT CCCAAATAT GGGGATTGGC 4680  
 4626 7  
 AAGAACGGAG ACCTACCCCTG GCCTCCGCTC AGGAACGAGT TCAAGTACTT CCAAAGAATG 4740  
 ACCACAAACCT CTTCAAGTGGA AGGTAAACAG AATCTGGTGA TTATGGTAG GAAAACCTGG 4800  
**DHFR=564bp=187 AMINO ACID & STOP CODON**  
 TTCTCCATTG CTGAGAAGAAA TCGACCTTTA AAGGACAGAA TTAATATAGT TCTCAAGTAGA 4860  
 GAACTCAAAG AACCACCACG AGGAGCTCAT TTTCTTGCCA AAAGTTGGA TGATGCCCTA 4920  
 AGACTTATTG AACAACCGGA ATTGGCAAGT AAAGTAGACA TGGTTTGGAT AGTCGGAGGC 4980  
 AGTTCTGTTT ACCAGGAAGC CATGAATCAA CCAGGCCACC TTAGACTCTT TGTGACAAGG 5040  
 ATCATGCAGG AATTGAAAG TGACACGTTT TTCCCAGAAA TTGATTTGGG GAAATATAAA 5100  
 CTTCTCCCAG AATAACCCAGG CGTCCTCTCT GAGGTCCAGG AGGAAAAAGG CATCAAGTAT 5160  
**STOP DHFR** 3' UNTRANSLATED DHFR=82bp  
 AAGTTGAAG TCTACGAGAA GAAAGAC**TAA** CAGGAAGATG CTTCAAGTT CTCTGCTCCC 5220  
 5140 1 **LINKER #10**  
 CTCCTAAAGC TATGCATTT TATAAGACCA TGGGACTTTT GCTGGCTTTA GATCAGCCCTC 5280  
 5272 3  
**=10bp**  
 GACTGTGCCT TCTAGTTGCC AGCCATCTGT TGTTTGCCCC TCCCCCGTGC CTTCCATTGAC 5340  
**BOVINE GROWTH HORMONE POLYADENYLATION=231bp**  
 CCTGGAAAGGT GCCACTCCCCA CTGTCCTTTA CTAATAAAAT GAGGAAATTG CATCGCATTG 5400  
 TCTGAGTAGG TGTCAATTCTA TTCTGGGGGG TGGGGTGGGG CAGGACAGCA AGGGGGAGGA 5460  
**LINKER #11**  
 TTGGGAAGAC AATAGCAGGC ATGCTGGGA TGCCTGGGC TCTATGGAAC CAGCTGGGGC 5520  
 5513 4  
**=17bp**  
 TCGAGCTACT AGCTTTGCTT CTCAATTCT TATTTGCATA ATGAGAAAAA AAGGAAAATT 5580  
 5530 1  
 AATTTAACCA CCAATTCACT AGTTGATTGA GCAAATGCGT TGCCAAAAAG GATGCTTTAG 5640  
**MOUSE BETA GLOBIN MAJOR PROMOTER=366bp**  
 AGACAGTGTG CTCTGCACAG ATAAGGACAA CTAGGGAGAA ATATGCTTGT CATCACCGAA 5700  
 GACTCCTAAG CCAGTGAGTG GCACAGCATT CTAGGGAGAA ATATGCTTGT CATCACCGAA 5760  
 GCCTGATTCC GTAGAGCCAC ACCTTGGTAA GGGCCAATCT GCTCACACAG GATAGAGAGG 5820  
 GCAGGAGCCA GGGCAGAGCA TATAAGGTGA GGTAGGATCA GTTGCCTCTC ACATTTGCTT 5880  
**LINKER #12=21bp** **START NEO**  
 CTGACATAGT TGTGTTGGGA GCTTGGATCG ATCCTCT**ATG** GTTGAACAAG ATGGATTGCA 5940  
 5896 7 5917 8  
 CGCAGGTTCT CGGGCCCGCTT GGGTGGAGAG GCTATTCCGC TATGACTGGG CACAACACAC 6000

FIG. 3D

AATCGGCTGC TCTGATGCCG CGGTGTTCCG GCTGTCAGCG CAGGGGGCGC CGGTTCTTT 6060  
 NEOMYCIN PHOSPHOTRANSFERASE=795bp=264 AMINO ACID & STOP CODON  
 TGTCAGGACC GACCTGTCCG GTGCCCTGAA TGAACTGCAG GACGAGGCAG CGCGGCTATC 6120  
 GTGGCTGGCC ACGACGGGCG TTCCTTGCGC AGCTGTGCTC GACGTTGTCA CTGAAGCGGG 6180  
 AAGGGACTGG CTGCTATTGG GCGAAGTGCC GGGGCAGGAT CTCTGTCTAT CTCACCTTGC 6240  
 TCCCTGCCAG AAAGTATCCA TCATGGCTGA TGCAATGCCG CGGCTGCATA CGCTTGATCC 6300  
 GGCTACCTGC CCATTGACCC ACCAAGCGAA ACATCGCATC GAGCGAGCAC GTACTCGGAT 6360  
 GGAAGCCGGT CTTGTCGATC AGGATGATCT GGACGAAGAG CATCAGGGGC TCGCGCCAGC 6420  
 CGAACTGTTG GCCAGGCTCA AGGCGCGCAT GCCCGACGGC GAGGATCTG TCGTGAACCA 6480  
 TGGCGATGCC TGCTTGCCTGA ATATCATGGT GGAAAATGGC CGCTTTCTG GATTATCGA 6540  
 CTGTGGCCGG CTGGGTGTGG CGGACCGCTA TCAGGACATA GCGTTGGCTA CCCGTGATTAT 6600  
 TGCTGAAGAG CTTGGCGCCG AATGGGCTGA CCGCTTCCTC GTGCTTTAAG GTATCGCCG 6660  
 STOP NEO  
 TCCCGATTCTG CAGCGCATCG CCTTCTATCG CCTTCTTGAC GAGTTCTTC ~~GAGCGGGACT~~ 6720  
 6712 3  
 CTGGGGTTCG AAATGACCGA CCAAGCGACG CCCAACCTGC CATCACGAGA TTTCGATTCC 6780  
 3' UNTRANSLATED NEO=173bp  
 ACCGGCCGCT TCTATGAAAG GTTGGGCTTC GGAATCGTTT TCCGGGACGC CGGCTGGATG 6840  
 ATCCTCCAGC GCGGGGATCT CATGCTGGAG TTCTTCCGCAC ACCCCAAACTT GTTTATTGCA 6885 6  
 GCTTATAATG GTTACAAATA AAGCAATAGC ATCACAAATT TCACAAATAA AGCATTTC 6960  
 SV40 EARLY POLYADENYLATION REGION=133bp  
 TCACTGCATT CTAGTTGTGG TTTGTCCAAA CTCATCAATC TATCTTATCA TGTCTGGATC 7020  
 7018 9  
 LINKER #13=19bp  
 GCGGCCGCGA TCCCGTCGAG AGCTTGGCGT AATCATGGTC ATAGCTGTTT CCTGTGTGAA 7080  
 7037 8  
 PUC 19  
 ATTGTTATCC GCTCACAAATT CCACACAAACA TACGAGCCGG AAGCATAAAG TGTAAAGCCT 7140  
 GGGGTGCCTA ATGAGTGAGC TAACTCACAT TAATTGCGTT GCGCTCACTG CCCGCTTTCC 7200  
 AGTCGGGAAA CCTGTCGTGC CAGCTGCATT AATGAATCG CCAACGGCGG GGGAGAGGCG 7260  
 GTTTGCGTAT TGGGCCCTCT TCCGCTTCCT CGCTCACTGA CTCGCTGCAG TCGGTGTTTC 7320  
 GGCTGCCGGCG AGCGGTATCA GCTCACTCAA AGGCGGTAAT ACGGTTAAGC ACAGAATCAG 7380  
 GGGATAACGC AGGAAAGAAC ATGTGAGCAA AAGGCCAGCA AAAGGCCAAG AACCGTAAAAA 7440  
 7461=BACTERIAL ORIGIN OF REPLICATION  
 AGGGCCGCGTT TCCATAGGC TCCGCCCGCC TGACGAGCAT CACAAAATC 7560

FIG. 3E

GACGCTCAAG TCAGAGGTGG CGAAACCCGA CAGGACTATA AAGATACCAAG GCGTTTCCCC 7560  
 CTGGAAAGCTC CCTCGTGCAGC TCTCCTGTT CGACCCCTGCC GCTTACCGGA TACCTGTCCG 7620  
 CCTTTCTCCC TTGGGGAAAGC GTGGCGCTTT CTCATGCTC ACGCTGTAGG TATCTCAGTT 7680  
 CGGTGTAGGT CGTTCGCTCC AAGCTGGGCT GTGTGCACGA ACCCCCCGTT CAGCCCCGACC 7740  
 GCTGCGCCCTT ATCCGGTAAC TATCGTCTTG AGTCCAACCC GGTAAGACAC GACTTATCGC 7800  
 CACTGGCAGC AGCCACTGGT AACAGGATTA GCAGAGCGAG GTATGTAGGC GGTGCTACAG 7860  
 AGTTCTTGAA GTGGTGGCCT AACTACGGCT ACACTAGAAG GACAGTATTG GGTATCTGCG 7920  
 CTCTGCTGAA GCCAGTTACC TTGGGAAAAA GAGTTGGTAG CTCTTGATCC GGCAAAACAAA 7980  
 CCACCGCTGG TAGCGGTGGT TTTTTGTTT GCAAGCAGCA GATTACGCGC AGAAAAAAAG 8040  
 GATCTCAAGA AGATCCTTG ATCTTTCTA CGGGGTCTGA CGCTCACTGG AACGAAAAGT 8100  
 CACGTTAAGG GATTTGGTC ATGAGATTAT CAAAAAGGAT CTTCACCTAG ATCCTTTAA 8160  
 STOP  
 ATTAAAAATG AAGTTTTAAA TCAATCTAAA GTATATATGA GTAAACTTGG TCTGACAGT 8220  
**BETA LACTAMASE**  
ACCAATGCTT AATCAGTGAG GCACCTATCT CAGCGATCTG TCTATTTCGT TCATCCATAG 8280  
 TTGCCTGACT CCCCGTCTG TAGATAACTA CGATAACGGGA GGGCTTACCA TCTGGCCCCA 8340  
 GTGCTGCAAT GATACCGCGA GACCCACGCT CACCGGCTCC AGATTTATCA GCAATAAAGC 8400  
 BETA LACTAMASE=861bp=286 AMINO ACID & STOP CODON  
 AGCCAGCCGG AAGGGCCGAG CGCAGAAGTG GTCCTGCAAC TTTATCCGCC TCCATCCAGT 8460  
 CTATTAATTG TTGCCGGGAA GCTAGAGTAA GTAGTTGCC AGTTAATAGT TTGCGCAACG 8520  
 TTGTTGCCAT TGCTACAGGC ATCGTGGTGT CACGCTCGTC GTTTGGTATG GCTTCATTCA 8580  
 GCTCCGGTTC CCAACGATCA AGGGCAGTTA CATGATCCCC CATGTTGTGC AAAAAAGCGG 8640  
 TTAGCTCCTT CGGTCTCTCG ATCGTTGTCA GAAGTAAGTT GGCGCGAGTG TTATCACTCA 8700  
 TGGTTATGGC AGCACTGCAT AATTCTCTTA CTGTCATGCC ATCCGTAAGA TGCTTTCTG 8760  
 TGACTGGTGA GTACTCAACC AAGTCATTCT GAGAATAGTG TATGCGCCGA CCGAGTTGCT 8820  
 CTTGCCCGGC GTCAATAACCG GATAATAACCG CGCCACATAG CAGAACTTTA AAAGTGCTCA 8880  
 TCATTGGAAA ACGTTCTTCG GGGCGAAAAC TCTCAAGGAT CTTACCGCTG TTGAGATCCA 8940  
 GGTGATGTA ACCCACTCGT GCACCCAACT GATCTTCAGC ATCTTTACT TTCACCGCG 9000  
 TTTCTGGGTG AGCAAAAACA GGAAGGCAAA ATGCCGCAAA AAAGGGAAATA AGGGCGACAC 9060  
 START BETA LACTAMASE  
 GGAAATGTTG AATACTCATA CTCTTCTTT TTCAATATTAA TTGAAGCATT TATCAGGGTT 9120  
 ATTGTCTCAT GAGCGGATAC ATATTGAAT GTATTTAGAA AAATAAACAA ATAGGGGTTC 9180  
 CGCGCACATT TCCCCGAAAAA GTGCCACCT

FIG. 3F

**LEADER**

FIG. 4

## LEADER

-19	-15	-10	-5
FRAME 1 Met Gly Trp Ser Leu Ile Leu Leu Phe Leu Val Ala Val Ala Thr Arg Val			
ATG GGT TGG AGC CTC ATC TTG CTC TTC CTT GTC GCT GTT GCT ACG CGT GTC			
2409	2418	2427	2436
2445			
15			
Leu Ser Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys <i>Ala</i> Gly Ala Ser			
CTG TCC CAG GTA CAA CTG CAG CAG CCT GGG GCT GAG CTG GTG AAG CCT GGG GCC TCA			
2460	2469	2478	2487
2505			
2505			
20	25	30	31
Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr Asn Met His Trp			
GTG AAG ATG TCC TGC AAG GCT TCT GGC TAC ACA TTT ACC AGT TAC AAT ATG CAC TGG			
2517	2526	2536	2544
2562			
40 FR2	45	49	50
Val Lys Gln Thr Pro Gly Arg Gly Leu Glu Trp Ile Gly Ala Ile Tyr Pro Gly Asn			
GTA AAA CAG ACA CCT GGT CGG GGC CTG GAA TGG ATT GGA GCT ATT TAT CCC CGA AAT			
2574	2583	2592	2601
2619			
55 CDR2	60	65	66 FR3
Gly Asp Thr Ser Tyr Asn Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys			
GGT GAT ACT TCC TAC AAT CAG AAG TTC AAA GGC AAG GCC ACA TTG ACT GCA GAC AAA			
2631	2640	2649	2658
2676			
75	80	82 82A 82B 82C 83	85
Ser Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val			
TCC TCC AGC ACA GCC TAC ATG CAG CTC AGC ACC CTC ACA TCT GAG GAC TCT GCG GTC			
2688	2697	2706	2715
2733			
90	94 95	CDR3	100 100A 100B 100C 100D 101
Tyr Tyr Cys Ala Arg Ser Thr Tyr Tyr Gly Gly Asp Trp Tyr Phe Asn Val Trp Gly			
TAT TAC TGT GCA AGA TCG ACT TAC TAC GGC GGT GAC TGG TAC TTC AAT GTC TGG GGC			
2745	2754	2763	2772
2790			
105 FR4	110	113	
Ala Gly Thr Thr Val Thr Val Ser Ala			
GCA GGG ACC ACG GTC ACC GTC TCT GCA			
2802	2811	2820	

FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14A




FIG. 14B

**THERAPEUTIC APPLICATION OF CHIMERIC  
AND RADIOLABELED ANTIBODIES TO HUMAN  
B LYMPHOCYTE RESTRICTED  
DIFFERENTIATION ANTIGEN FOR TREATMENT  
OF B CELL LYMPHOMA**

[0001] This application is a continuation of U.S. application Ser. No. 09/905,928, filed Jul. 17, 2001; which is a continuation of U.S. application Ser. No. 08/475,813, filed Jun. 7, 1995, now U.S. Pat. No. 6,672,813; which is a divisional of U.S. application Ser. No. 08/149,099, filed Nov. 3, 1993, now U.S. Pat. No. 5,736,137; which is a continuation-in-part of U.S. application Ser. No. 07/978,891, filed Nov. 13, 1992, now abandoned. Each of these prior applications is incorporated herein by reference in its entirety.

**TABLE OF CONTENTS**

- [0002] A. FIELD OF THE INVENTION
- [0003] B. BACKGROUND OF THE INVENTION
- [0004] C. SUMMARY OF THE INVENTION
- [0005] D. BRIEF DESCRIPTION OF THE DRAWINGS
- [0006] E. DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS
- [0007] F. EXAMPLES
- [0008] I. RADIOLABELED ANTI-CD20 ANTI-BODY 2B8
  - [0009] A. Anti-CD20 Monoclonal Antibody (Murine) Production (“2B8”)
  - [0010] B. Preparation of 2B8-MX-DTPA Conjugate
    - [0011] i. MX-DTPA
    - [0012] ii. Preparation of 2B8
    - [0013] iii. Conjugation of 2B8 with MX-DTPA
    - [0014] iv. Determination of MX-DTPA Incorporation
    - [0015] v. Immunoreactivity of 2B8-MX-DTPA
    - [0016] vi. Preparation of Indium-[111]-labeled 2B8-MX-DTPA (“I2B8”)
    - [0017] vii. Preparation of Yttrium-[90]-labeled 2B8-MX-DTPA (“Y2B8”)
  - [0018] C. Non-Human Animal Studies
    - [0019] i. Distribution of Radiolabeled 2B8-MX-DTPA
    - [0020] ii. Tumor Localization of I2B8
    - [0021] iii. Biodistribution and Tumor Localization Studies with Radiolabeled 2B8-MX-DTPA
  - [0022] D. Human Studies
    - [0023] i. 2B8 and 2B8-MX-DTPA: Immunohistochemistry Studies with Human Tissues
    - [0024] ii. Clinical Analysis of I2B8 (Imaging and Y2B8 (Therapy)
- [0025] a. Phase I/II Clinical Trial: Single Dose Therapy Study
- [0026] b. Phase I/II Clinical Trial: Multiple Dose Therapy Study
- [0027] II. CHIMERIC ANTI-CD20 PRODUCTION
- [0028] A. Construction of Chimeric Anti-CD20 Immunoglobulin DNA Expression Vectors
- [0029] B. Creation of Chimeric Anti-CD20 Producing CHO and SP2/0 Transfectomas
- [0030] C. Determination of Immunological Activity of Chimeric Anti-CD20 Antibodies
  - [0031] i. Human C1q Analysis
  - [0032] ii. Complement Dependent Cell Lyses
  - [0033] iii. Antibody Dependent Cellular Cytotoxicity Effector Assay
- [0034] III. DEPLETION OF B CELLS IN VIVO USING CHIMERIC ANTI-CD20
- [0035] A. Non-Human Primate Study
- [0036] B. Clinical Analysis of C2B8
  - [0037] i. Phase I/II Clinical Trial of C2B8: Single Dose Therapy Study
  - [0038] ii. Phase I/II Clinical Trial of C2B8: Multiple Dose Therapy Study
- [0039] IV. COMBINATION THERAPY: C2B8 AND Y2B8
  - [0040] A. Preparation of Y2B8
  - [0041] B. Preparation of C2B8
  - [0042] C. Results
- [0043] V. ALTERNATIVE THERAPY STRATEGIES
- [0044] VI. DEPOSIT INFORMATION
- [0045] G. SEQUENCE LISTING
- [0046] H. CLAIMS

**A. FIELD OF THE INVENTION**

[0047] The references to be discussed throughout this document are set forth merely for the information described therein prior to the filing date of this document, and nothing herein is to be construed as an admission, either express or implied, that the references are “prior art” or that the inventors are not entitled to antedate such descriptions by virtue of prior inventions or priority based on earlier filed applications.

[0048] The present invention is directed to the treatment of B cell lymphoma using chimeric and radiolabeled antibodies to the B cell surface antigen Bp35 (“CD20”).

**B. BACKGROUND OF THE INVENTION**

[0049] The immune system of vertebrates (for example, primates, which include humans, apes, monkeys, etc.) consists of a number of organs and cell types which have evolved to: accurately and specifically recognize foreign

microorganisms (“antigen”) which invade the vertebrate-host; specifically bind to such foreign microorganisms; and, eliminate/destroy such foreign microorganisms.

[0050] Lymphocytes, amongst others, are critical to the immune system. Lymphocytes are produced in the thymus, spleen and bone marrow (adult) and represent about 30% of the total white blood cells present in the circulatory system of humans (adult). There are two major sub-populations of lymphocytes: T cells and B cells. T cells are responsible for cell mediated immunity, while B cells are responsible for antibody production (humoral immunity). However, T cells and B cells can be considered as interdependent—in a typical immune response, T cells are activated when the T cell receptor binds to fragments of an antigen that are bound to major histocompatibility complex (“MHC”) glycoproteins on the surface of an antigen presenting cell; such activation causes release of biological mediators (“interleukins”) which, in essence, stimulate B cells to differentiate and produce antibody (“immunoglobulins”) against the antigen.

[0051] Each B cell within the host expresses a different antibody on its surface—thus, one B cell will express antibody specific for one antigen, while another B cell will express antibody specific for a different antigen. Accordingly, B cells are quite diverse, and this diversity is critical to the immune system. In humans, each B cell can produce an enormous number of antibody molecules (ie about  $10^7$  to  $10^8$ ). Such antibody production most typically ceases (or substantially decreases) when the foreign antigen has been neutralized. Occasionally, however, proliferation of a particular B cell will continue unabated; such proliferation can result in a cancer referred to as “B cell lymphoma.”

[0052] T cells and B cells both comprise cell surface proteins which can be utilized as “markers” for differentiation and identification. One such human B cell marker is the human B lymphocyte-restricted differentiation antigen Bp35, referred to as “CD20.” CD20 is expressed during early pre-B cell development and remains until plasma cell differentiation. Specifically, the CD20 molecule may regulate a step in the activation process which is required for cell cycle initiation and differentiation and is usually expressed at very high levels on neoplastic (“tumor”) B cells. CD20, by definition, is present on both “normal” B cells as well as “malignant” B cells, ie those B cells whose unabated proliferation can lead to B cell lymphoma. Thus, the CD20 surface antigen has the potential of serving as a candidate for “targeting” of B cell lymphomas.

[0053] In essence, such targeting can be generalized as follows: antibodies specific to the CD20 surface antigen of B cells are, eg injected into a patient. These anti-CD20 antibodies specifically bind to the CD20 cell surface antigen of (ostensibly) both normal and malignant B cells; the anti-CD20 antibody bound to the CD20 surface antigen may lead to the destruction and depletion of neoplastic B cells. Additionally, chemical agents or radioactive labels having the potential to destroy the tumor can be conjugated to the anti-CD20 antibody such that the agent is specifically “delivered” to, eg, the neoplastic B cells. Irrespective of the approach, a primary goal is to destroy the tumor: the specific approach can be determined by the particular anti-CD20 antibody which is utilized and, thus, the available approaches to targeting the CD20 antigen can vary considerably.

[0054] For example, attempts at such targeting of CD20 surface antigen have been reported. Murine (mouse) monoclonal antibody 1F5 (an anti-CD20 antibody) was reportedly administered by continuous intravenous infusion to B cell lymphoma patients. Extremely high levels (>2 grams) of 1F5 were reportedly required to deplete circulating tumor cells, and the results were described as being “transient.” Press et al., “Monoclonal Antibody 1F5 (Anti-CD20) Serotherapy of Human B-Cell Lymphomas.” *Blood* 69/2: 584-591 (1987). A potential problem with this approach is that non-human monoclonal antibodies (eg, murine monoclonal antibodies) typically lack human effector functionality, ie they are unable to, *inter alia*, mediate complement dependent lysis or lyse human target cells through antibody dependent cellular toxicity or Fc-receptor mediated phagocytosis. Furthermore, non-human monoclonal antibodies can be recognized by the human host as a foreign protein; therefore, repeated injections of such foreign antibodies can lead to the induction of immune responses leading to harmful hypersensitivity reactions. For murine-based monoclonal antibodies, this is often referred to as a Human Anti-Mouse Antibody response, or “HAMA” response. Additionally, these “foreign” antibodies can be attacked by the immune system of the host such that they are, in effect, neutralized before they reach their target site.

[0055] Lymphocytes and lymphoma cells are inherently sensitive to radiotherapy for several reasons: the local emission of ionizing radiation of radiolabeled antibodies may kill cells with or without the target antigen (eg CD20) in close proximity to antibody bound to the antigen; penetrating radiation may obviate the problem of limited access to the antibody in bulky or poorly vascularized tumors; and, the total amount of antibody required may be reduced. The radionuclide emits radioactive particles which can damage cellular DNA to the point where the cellular repair mechanisms are unable to allow the cell to continue living; therefore, if the target cells are tumors, the radioactive label beneficially kills the tumor cells. Radiolabeled antibodies, by definition, include the use of a radioactive substance which may require the need for precautions for both the patient (ie possible bone marrow transplantation) as well as the health care provider (ie the need to exercise a high degree of caution when working with the radioactivity).

[0056] Therefore, an approach at improving the ability of murine monoclonal antibodies to be effective in the treatment of B-cell disorders has been to conjugate a radioactive label or toxin to the antibody such that the label or toxin is localized at the tumor site. For example, the above-referenced 1F5 antibody has been “labeled” with iodine-131 (“ $^{131}\text{I}$ ”) and was reportedly evaluated for biodistribution in two patients. See Eary, J. F. et al., “Imaging and Treatment of B-Cell Lymphoma” *J. Nuc. Med.* 31/8: 1257-1268 (1990); see also, Press, O. W. et al., “Treatment of Refractory Non-Hodgkin’s Lymphoma with Radiolabeled MB-1 (Anti-CD37) Antibody” *J. Clin. Onc.* 7/8: 1027-1038 (1989) (indication that one patient treated with  $^{131}\text{I}$ -labeled 1F5 achieved a “partial response”); Goldenberg, D. M. et al., “Targeting, Dosimetry and Radioimmunotherapy of B-Cell Lymphomas with Iodine-131-Labeled LL2 Monoclonal Antibody” *J. Clin. Onc.* 9/4: 548-564 (1991) (three of eight patients receiving multiple injections reported to have developed a HAMA response); Appelbaum, F. R. “Radiolabeled Monoclonal Antibodies in the Treatment of Non-Hodgkin’s Lymphoma” *Hem./One. Clinics of N.A.* 5/5: 1013-1025

(1991) (review article); Press, O. W. et al “Radiolabeled Antibody Therapy of B-Cell Lymphoma with Autologous Bone Marrow Support.” *New England Journal of Medicine* 329/17: 1219-1223 (1993) (iodine-131 labeled anti-CD20 antibody IF5 and B1); and Kaminski, M. G. et al “Radioimmunotherapy of B-Cell Lymphoma with [<sup>131</sup>I] Anti-B1 (Anti-CD20) Antibody”. *NEJM* 329/7 (1993) (iodine-131 labeled anti-CD20 antibody B1; hereinafter “Kaminski”).

[0057] Toxins (ie chemotherapeutic agents such as doxorubicin or mitomycin C) have also been conjugated to antibodies. See, for example, PCT published application WO 92/07466 (published May 14, 1992).

[0058] “Chimeric” antibodies, ie antibodies which comprise portions from two or more different species (eg, mouse and human) have been developed as an alternative to “conjugated” antibodies. For example, Liu, A. Y. et al., “Production of a Mouse-Human Chimeric Monoclonal Antibody to CD20 with Potent Fc-Dependent Biologic Activity” *J. Immun.* 139/10: 3521-3526 (1987), describes a mouse/human chimeric antibody directed against the CD20 antigen. See also, PCT Publication No. WO 88/04936. However, no information is provided as to the ability, efficacy or practicability of using such chimeric antibodies for the treatment of B cell disorders in the reference. It is noted that in vitro functional assays (eg complement dependent lysis (“CDC”); antibody dependent cellular cytotoxicity (“ADCC”), etc.) cannot inherently predict the in vivo capability of a chimeric antibody to destroy or deplete target cells expressing the specific antigen. See, for example, Robinson, R. D. et al., “Chimeric mouse-human anti-carcinoma antibodies that mediate different anti-tumor cell biological activities,” *Hum. Antibod. Hybridomas* 2: 84-93 (1991) (chimeric mouse-human antibody having undetectable ADCC activity). Therefore, the potential therapeutic efficacy of chimeric antibody can only truly be assessed by in vivo experimentation.

[0059] What is needed, and what would be a great advance in the art, are therapeutic approaches targeting the CD20 antigen for the treatment of B cell lymphomas in primates, including, but not limited to, humans.

#### C. SUMMARY OF THE INVENTION

[0060] Disclosed herein are therapeutic methods designed for the treatment of B cell disorders, and in particular, B cell lymphomas. These protocols are based upon the administration of immunologically active chimeric anti-CD20 antibodies for the depletion of peripheral blood B cells, including B cells associated with lymphoma; administration of radiolabeled anti-CD20 antibodies for targeting localized and peripheral B cell associated tumors; and administration of chimeric anti-CD20 antibodies and radiolabeled anti-CD20 antibodies in a cooperative therapeutic strategy.

#### D. BRIEF DESCRIPTION OF THE DRAWINGS

[0061] **FIG. 1** is a diagrammatic representation of a tandem chimeric antibody expression vector useful in the production of immunologically active chimeric anti-CD20 antibodies (“TCAE 8”);

[0062] **FIGS. 2A through 2E** are the nucleic acid sequence of the vector of **FIG. 1**;

[0063] **FIGS. 3A through 3F** are the nucleic acid sequence of the vector of **FIG. 1** further comprising murine light and heavy chain variable regions (“anti-CD20 in TCAE 8”);

[0064] **FIG. 4** is the nucleic acid and amino acid sequences (including CDR and framework regions) of murine variable region light chain derived from murine anti-CD20 monoclonal antibody 2B8;

[0065] **FIG. 5** is the nucleic acid and amino acid sequences (including CDR and framework regions) of murine variable region heavy chain derived from murine anti-CD20 monoclonal antibody 2B8;

[0066] **FIG. 6** are flow cytometry results evidencing binding of fluorescent-labeled human C1q to chimeric anti-CD20 antibody, including, as controls labeled C1q; labeled C1q and murine anti-CD20 monoclonal antibody 2B8; and labeled C1q and human IgG1,k;

[0067] **FIG. 7** represents the results of complement related lysis comparing chimeric anti-CD20 antibody and murine anti-CD20 monoclonal antibody 2B8;

[0068] **FIG. 8** represents the results of antibody mediated cellular cytotoxicity with in vivo human effector cells comparing chimeric anti-CD20 antibody and 2B8;

[0069] **FIGS. 9A, 9B and 9C** provide the results of non-human primate peripheral blood B lymphocyte depletion after infusion of 0.4 mg/kg (A); 1.6 mg/kg (B); and 6.4 mg/kg (C) of immunologically active chimeric anti-CD20 antibody;

[0070] **FIG. 10** provides the results of, inter alia, non-human primate peripheral blood B lymphocyte depletion after infusion of 0.01 mg/kg of immunologically active chimeric anti-CD20 antibody;

[0071] **FIG. 11** provides results of the tumoricidal impact of Y2B8 in a mouse xenographic model utilizing a B cell lymphoblastic tumor;

[0072] **FIG. 12** provides results of the tumoricidal impact of C2B8 in a mouse xenographic model utilizing a B cell lymphoblastic tumor;

[0073] **FIG. 13** provides results of the tumoricidal impact of a combination of Y2B8 and C2B8 in a mouse xenographic model utilizing a B cell lymphoblastic tumor; and

[0074] **FIGS. 14A and 14B** provide results from a Phase I/II clinical analysis of C2B8 evidencing B-cell population depletion over time for patients evidencing a partial remission of the disease (14A) and a minor remission of the disease (14B).

#### E. DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0075] Generally, antibodies are composed of two light chains and two heavy chain molecules; these chains form a general “Y” shape, with both light and heavy chains forming the arms of the Y and the heavy chains forming the base of the Y. Light and heavy chains are divided into domains of structural and functional homology. The variable domains of both the light (“V<sub>L</sub>”) and the heavy (“V<sub>H</sub>”) chains determine recognition and specificity. The constant region domains of light (“C<sub>L</sub>”) and heavy (“C<sub>H</sub>”) chains confer important

biological properties, eg antibody chain association, secretion, transplacental mobility, Fc receptor binding complement binding, etc. The series of events leading to immunoglobulin gene expression in the antibody producing cells are complex. The variable domain region gene sequences are located in separate germ line gene segments referred to as " $V_H$ ," " $D$ ," and " $J_H$ ," or " $V_L$ " and " $J_L$ ." These gene segments are joined by DNA rearrangements to form the complete V regions expressed in heavy and light chains, respectively. The rearranged, joined V segments ( $V_L$ - $J_L$  and  $V_H$ - $D$ - $J_H$ ) then encode the complete variable regions or antigen binding domains of light and heavy chains, respectively.

[0076] Serotherapy of human B cell lymphomas using an anti-CD20 murine monoclonal antibody (1F5) has been described by Press et al., (69 *Blood* 584, 1987, *supra*); the reported therapeutic responses, unfortunately, were transient. Additionally, 25% of the tested patients reportedly developed a human anti-mouse antibody (HAMA) response to the serotherapy. Press et al., suggest that these antibodies, conjugated to toxins or radioisotopes, might afford a more lasting clinical benefit than the unconjugated antibody.

[0077] Owing to the debilitating effects of B cell lymphoma and the very real need to provide viable treatment approaches to this disease, we have embarked upon different approaches having a particular antibody, 2B8, as the common link between the approaches. One such approach advantageously exploits the ability of mammalian systems to readily and efficiently recover peripheral blood B cells; using this approach, we seek to, in essence, purge or deplete B cells in peripheral blood and lymphatic tissue as a means of also removing B cell lymphomas. We accomplish this by utilization of, *inter alia*, immunologically active, chimeric anti-CD20 antibodies. In another approach, we seek to target tumor cells for destruction with radioactive labels.

[0078] As used herein, the term "anti-CD20 antibody" is an antibody which specifically recognizes a cell surface non-glycosylated phosphoprotein of 35,000 Daltons, typically designated as the human B lymphocyte restricted differentiation antigen Bp35, commonly referred to as CD20. As used herein, the term "chimeric" when used in reference to anti-CD20 antibodies, encompasses antibodies which are most preferably derived using recombinant deoxyribonucleic acid techniques and which comprise both human (including immunologically "related" species, eg, chimpanzee) and non-human components: the constant region of the chimeric antibody is most preferably substantially identical to the constant region of a natural human antibody; the variable region of the chimeric antibody is most preferably derived from a non-human source and has the desired antigenic and specificity to the CD20 cell surface antigen. The non-human source can be any vertebrate source which can be used to generate antibodies to a human CD20 cell surface antigen or material comprising a human CD20 cell surface antigen. Such non-human source includes, but is not limited to, rodents (eg, rabbit, rat, mouse, etc.) and non-human primates (eg, Old World Monkey, Ape, etc.). Most preferably, the non-human component (variable region) is derived from a murine source. As used herein, the phrase "immunologically active" when used in reference to chimeric anti-CD20 antibodies, means a chimeric antibody which binds human C1q, mediates complement dependent lysis ("CDC") of human B lymphoid cell lines, and lyses human target cells through antibody dependent cellular

cytotoxicity ("ADCC"). As used herein, the phrases "indirect labeling" and "indirect labeling approach" both mean that a chelating agent is covalently attached to an antibody and at least one radionuclide is inserted into the chelating agent. Preferred chelating agents and radionuclides are set forth in Srivagta, S. C. and Mease, R. C., "Progress in Research on Ligands, Nuclides and Techniques for Labeling Monoclonal Antibodies," *Nucl. Med. Bio.* 18/6: 589-603 (1991) ("Srivagta") which is incorporated herein by reference. A particularly preferred chelating agent is 1-isothiocyanato-benzyl-3-methylthiophene triaminepent acetic acid ("MX-DTPA"); particularly preferred radionuclides for indirect labeling include indium [111] and yttrium [90]. As used herein, the phrases "direct labeling" and "direct labeling approach" both mean that a radionuclide is covalently attached directly to an antibody (typically via an amino acid residue). Preferred radionuclides are provided in Srivagta; a particularly preferred radionuclide for direct labeling is iodine [131] covalently attached via tyrosine residues. The indirect labeling approach is particularly preferred.

[0079] The therapeutic approaches disclosed herein are based upon the ability of the immune system of primates to rapidly recover, or rejuvenate, peripheral blood B cells. Additionally, because the principal immune response of primates is occasioned by T cells, when the immune system has a peripheral blood B cell deficiency, the need for "extraordinary" precautions (ie patient isolation, etc.) is not necessary. As a result of these and other nuances of the immune systems of primates, our therapeutic approach to B cell disorders allows for the purging of peripheral blood B cells using immunologically active chimeric anti-CD20 antibodies.

[0080] Because peripheral blood B cell disorders, by definition, can indicate a necessity for access to the blood for treatment, the route of administration of the immunologically active chimeric anti-CD20 antibodies and radioalabeled anti-CD20 antibodies is preferably parenteral; as used herein, the term "parenteral" includes intravenous, intramuscular, subcutaneous, rectal, vaginal or intraperitoneal administration. Of these, intravenous administration is most preferred.

[0081] The immunologically active chimeric anti-CD20 antibodies and radiolabeled anti-CD20 antibodies will typically be provided by standard technique within a pharmaceutically acceptable buffer, for example, sterile saline, sterile buffered water, propylene glycol, combinations of the foregoing, etc. Methods for preparing parenteraly administerable agents are described in *Pharmaceutical Carriers & Formulations*, Martin, Remington's Pharmaceutical Sciences, 15th Ed. (Mack Pub. Co., Easton, Pa. 1975), which is incorporated herein by reference.

[0082] The specific, therapeutically effective amount of immunologically active chimeric anti-CD20 antibodies useful to produce a unique therapeutic effect in any given patient can be determined by standard techniques well known to those of ordinary skill in the art.

[0083] Effective dosages (ie therapeutically effective amounts) of the immunologically active chimeric anti-CD20 antibodies range from about 0.001 to about 30 mg/kg body weight, more preferably from about 0.01 to about 25 mg/kg body weight, and most preferably from about 0.4 to about 20.0 mg/kg body weight. Other dosages are viable; factors

influencing dosage include, but are not limited to, the severity of the disease; previous treatment approaches; overall health of the patient; other diseases present, etc. The skilled artisan is readily credited with assessing a particular patient and determining a suitable dosage that falls within the ranges, or if necessary, outside of the ranges.

[0084] Introduction of the immunologically active chimeric anti-CD20 antibodies in these dose ranges can be carried out as a single treatment or over a series of treatments. With respect to chimeric antibodies, it is preferred that such introduction be carried out over a series of treatments; this preferred approach is predicated upon the treatment methodology associated with this disease. While not wishing to be bound by any particular theory, because the immunologically active chimeric anti-CD20 antibodies are both immunologically active and bind to CD20, upon initial introduction of the immunologically active chimeric anti-CD-20 antibodies to the individual, peripheral blood B cell depletion will begin; we have observed a nearly complete depletion within about 24 hours post treatment infusion. Because of this, subsequent introduction(s) of the immunologically active chimeric anti-CD20 antibodies (or radiolabeled anti-CD20 antibodies) to the patient is presumed to: a) clear remaining peripheral blood B cells; b) begin B cell depletion from lymph nodes; c) begin B cell depletion from other tissue sources, eg, bone marrow, tumor, etc. Stated again, by using repeated introductions of the immunologically active chimeric anti-CD20 antibodies, a series of events take place, each event being viewed by us as important to effective treatment of the disease. The first "event" then, can be viewed as principally directed to substantially depleting the patient's peripheral blood B cells; the subsequent "events" can be viewed as either principally directed to simultaneously or serially clearing remaining B cells from the system clearing lymph node B cells, or clearing other tissue B cells.

[0085] In effect, while a single dosage provides benefits and can be effectively utilized for disease treatment/management, a preferred treatment course can occur over several stages; most preferably, between about 0.4 and about 20 mg/kg body weight of the immunologically active chimeric anti-CD20 antibodies is introduced to the patient once a week for between about 2 to 10 weeks, most preferably for about 4 weeks.

[0086] With reference to the use of radiolabeled anti-CD20 antibodies, a preference is that the antibody is non-chimeric; this preference is predicated upon the significantly longer circulating half-life of chimeric antibodies vis-a-vis murine antibodies (ie with a longer circulating half-life, the radionuclide is present in the patient for extended periods). However, radiolabeled chimeric antibodies can be beneficially utilized with lower milli-Curies ("mCi") dosages used in conjunction with the chimeric antibody relative to the murine antibody. This scenario allows for a decrease in bone marrow toxicity to an acceptable level, while maintaining therapeutic utility.

[0087] A variety of radionuclides are applicable to the present invention and those skilled in the art are credited with the ability to readily determine which radionuclide is most appropriate under a variety of circumstances. For example, iodine [131] is a well known radionuclide used for targeted immunotherapy. However, the clinical usefulness of

iodine [131] can be limited by several factors including: eight-day physical half-life; dehalogenation of iodinated antibody both in the blood and at tumor sites; and emission characteristics (eg large gamma component) which can be suboptimal for localized dose deposition in tumor. With the advent of superior chelating agents, the opportunity for attaching metal chelating groups to proteins has increased the opportunities to utilize other radionuclides such as indium [131] and yttrium [90]. Yttrium [90] provides several benefits for utilization in radioimmunotherapeutic applications: the 64 hour half-life of yttrium [90] is long enough to allow antibody accumulation by tumor and, unlike eg iodine [131], yttrium [90] is a pure beta emitter of high energy with no accompanying gamma irradiation in its decay, with a range in tissue of 100 to 1000 cell diameters. Furthermore, the minimal amount of penetrating radiation allows for outpatient administration of yttrium [90]-labeled antibodies. Furthermore, internalization of labeled antibody is not required for cell killing, and the local emission of ionizing radiation should be lethal for adjacent tumor cells lacking the target antigen.

[0088] One non-therapeutic limitation to yttrium [90] is based upon the absence of significant gamma radiation making imaging therewith difficult. To avoid this problem, a diagnostic "imaging" radionuclide, such as indium [111], can be utilized for determining the location and relative size of a tumor prior to the administration of therapeutic doses of yttrium [90]-labeled anti-CD20. Indium [111] is particularly preferred as the diagnostic radionuclide because: between about 1 to about 10 mCi can be safely administered without detectable toxicity; and the imaging data is generally predictive of subsequent yttrium [90]-labeled antibody distribution. Most imaging studies utilize 5 mCi indium [111]-labeled antibody because this dose is both safe and has increased imaging efficiency compared with lower doses, with optimal imaging occurring at three to six days after antibody administration. See, for example, Murray J. L., 26 *J. Nuc. Med.* 3328 (1985) and Carraguillo, J. A. et al, 26 *J. Nuc. Med.* 67 (1985).

[0089] Effective single treatment dosages (ie therapeutically effective amounts) of yttrium [90] labeled anti-CD20 antibodies range from between about 5 and about 75 mCi, more preferably between about 10 and about 40 mCi. Effective single treatment non-marrow ablative dosages of iodine [131] labeled anti-CD20 antibodies range from between about 5 and about 70 mCi, more preferably between about 5 and about 40 mCi. Effective single treatment ablative dosages (ie may require autologous bone marrow transplantation) of iodine [131] labeled anti-CD20 antibodies range from between about 30 and about 600 mCi, more preferably between about 50 and less than about 500 mCi. In conjunction with a chimeric anti-CD20 antibody, owing to the longer circulating half life vis-a-vis murine antibodies, an effective single treatment non-marrow ablative dosages of iodine [131] labeled chimeric anti-CD20 antibodies range from between about 5 and about 40 mCi, more preferably less than about 30 mCi. Imaging criteria for, eg the indium [111] label, are typically less than about 5 mCi.

[0090] With respect to radiolabeled anti-CD20 antibodies, therapy therewith can also occur using a single therapy treatment or using multiple treatments. Because of the radionuclide component, it is preferred that prior to treatment, peripheral stem cells ("PSC") or bone marrow ("BM")

be "harvested" for patients experiencing potentially fatal bone marrow toxicity resulting from radiation. BM and/or PSC are harvested using standard techniques, and then purged and frozen for possible reinfusion. Additionally, it is most preferred that prior to treatment a diagnostic dosimetry study using a diagnostic labeled antibody (eg using indium [111]) be conducted on the patient, a purpose of which is to ensure that the therapeutically labeled antibody (eg using yttrium [90]) will not become unnecessarily "concentrated" in any normal organ or tissue.

[0091] Chimeric mouse/human antibodies have been described. See, for example, Morrison, S. L. et al., *PNAS* 11: 6851-6854 (November 1984); European Patent Publication No. 173494; Boulianne, G. L. et al., *Nature* 312: 643 (December 1984); Neubeiger, M. S. et al., *Nature* 314: 268 (March 1985); European Patent Publication No. 125023; Tan et al., *J. Immunol.* 135: 8564 (November 1985); Sun, L. K. et al., *Hybridoma* 5/1: 517 (1986); Sahagan et al., *J. Immunol.* 137: 1066-1074 (1986). See generally, Muron, *Nature* 312: 597 (December 1984); Dickson, *Genetic Engineering News* 5/3 (March 1985); Marx, *Science* 229 455 (August 1985); and Morrison *Science* 229: 1202-1207 (September 1985). Robinson et al., in PCT Publication Number WO 88/04936 describe a chimeric antibody with human constant region and murine variable region, having specificity to an epitope of CD20; the murine portion of the chimeric antibody of the Robinson references is derived from the 2H7 mouse monoclonal antibody (gamma 2b, kappa). While the reference notes that the described chimeric antibody is a "prime candidate" for the treatment of B cell disorders, this statement can be viewed as no more than a suggestion to those in the art to determine whether or not this suggestion is accurate for this particular antibody, particularly because the reference lacks any data to support an assertion of therapeutic effectiveness, and importantly, data using higher order mammals such as primates or humans.

[0092] Methodologies for generating chimeric antibodies are available to those in the art. For example, the light and heavy chains can be expressed separately, using, for example, immunoglobulin light chain and immunoglobulin heavy chains in separate plasmids. These can then be purified and assembled in vitro into complete antibodies; methodologies for accomplishing such assembly have been described. See, for example, Scharff, M., *Harvey Lectures* 69: 125 (1974). In vitro reaction parameters for the formation of IgG antibodies from reduced isolated light and heavy chains have also been described. See, for example, Beychok, S., *Cells of Immunoglobulin Synthesis*, Academic Press, New York, p. 69, 1979. Co-expression of light and heavy chains in the same cells to achieve intracellular association and linkage of heavy and light chains into complete H<sub>2</sub>L<sub>2</sub> IgG antibodies is also possible. Such co-expression can be accomplished using either the same or different plasmids in the same host cell.

[0093] Another approach, and one which is our most preferred approach for developing a chimeric non-human/human anti-CD20 antibody, is based upon utilization of an expression vector which includes, ab initio, DNA encoding heavy and light chain constant regions from a human source. Such a vector allows for inserting DNA encoding non-human variable region such that a variety of non-human anti-CD20 antibodies can be generated, screened and ana-

lyzed for various characteristics (eg type of binding specificity, epitope binding regions, etc.); thereafter, cDNA encoding the light and heavy chain variable regions from a preferred or desired anti-CD20 antibody can be incorporated into the vector. We refer to these types of vectors as Tandem Chimeric Antibody Expression ("TCAE") vectors. A most preferred TCAE vector which was used to generate immunologically active chimeric anti-CD20 antibodies for therapeutic treatment of lymphomas is TCAE 8. TCAE 8 is a derivative of a vector owned by the assignee of this patent document, referred to as TCAE 5.2 the difference being that in TCAE 5.2, the translation initiation start site of the dominant selectable marker (neomycin phosphotransferase, "NEO") is a consensus Kozak sequence, while for TCAE 8, this region is a partially impaired consensus Kozak sequence. Details regarding the impact of the initiation start site of the dominant selectable marker of the TCAE vectors (also referred to as "ANEX vector") vis-a-vis protein expression are disclosed in detail in the co-pending application filed herewith.

[0094] TCAE 8 comprises four (4) transcriptional cassettes, and these are in tandem order, ie a human immunoglobulin light chain absent a variable region; a human immunoglobulin heavy chain absent a variable region; DHFR; and NEO. Each transcriptional cassette contains its own eukaryotic promoter and polyadenylation region (reference is made to **FIG. 1** which is a diagrammatic representation of the TCAE 8 vector). Specifically:

[0095] 1) the CMV promoter/enhancer in front of the immunoglobulin heavy chain is a truncated version of the promoter/enhancer in front of the light chain, from the Nhe I site at -350 to the Sst I site at -16 (see, 41 *Cell* 521, 1985).

[0096] 2) a human immunoglobulin light chain constant region was derived via amplification of cDNA by a PCR reaction. In TCAE 8, this was the human immunoglobulin light chain kappa constant region (Kabat numbering, amino acids 108-214, allotype Km 3, (see, Kabat, E. A "Sequences of proteins of immunological interest," NIH Publication, Fifth Ed. No. 91-3242, 1991)), and the human immunoglobulin heavy chain gamma 1 constant region (Kabat numbering amino acids 114-478, allotype Gmla, Gmlz). The light chain was isolated from normal human blood (IDEC Pharmaceuticals Corporation, La Jolla, Calif.); RNA therefrom was used to synthesize cDNA which was then amplified using PCR techniques (primers were derived vis-a-vis the consensus from Kabat). The heavy chain was isolated (using PCR techniques) from cDNA prepared from RNA which was in turn derived from cells transfected with a human IgG1 vector (see, 3 *Prot. Eng.* 531, 1990; vector pN<sub>γ1</sub>62). Two amino acids were changed in the isolated human IgG1 to match the consensus amino acid sequence from Kabat, to wit: amino acid 225 was changed from valine to alanine (GTT to GCA), and amino acid 287 was changed from methionine to lysine (ATG to AAG);

[0097] 3) The human immunoglobulin light and heavy chain cassettes contain synthetic signal sequences for secretion of the immunoglobulin chains;

[0098] 4) The human immunoglobulin light and heavy chain cassettes contain specific DNA restriction sites

which allow for insertion of light and heavy immunoglobulin variable regions which maintain the transitional reading frame and do not alter the amino acids normally found in immunoglobulin chains;

[0099] 5) The DHFR cassette contained its own eukaryotic promoter (mouse beta globin major promoter, "BETA") and polyadenylation region (bovine growth hormone polyadenylation, "BGH"); and

[0100] 6) The NEO cassette contained its own eukaryotic promoter (BETA) and polyadenylation region (SV40 early polyadenylation, "SV").

[0101] With respect to the TCAE 8 vector and the NEO cassette, the Kozak region was a partially impaired consensus Kozak sequence (which included an upstream Cla I site):

ClaI      -3      +1  
GGGAGCTTGG ATCGAT ccTct ATG Gtt

[0102] (In the TCAE 5.2 vector, the change is between the ClaI and ATG regions, to wit: ccAcc.)

[0103] The complete sequence listing of TCAE 8 (including the specific components of the four transcriptional cassettes) is set forth in **FIG. 2** (SEQ. ID. NO. 1).

[0104] As will be appreciated by those in the art, the TCAE vectors beneficially allow for substantially reducing the time in generating the immunologically active chimeric anti-CD20 antibodies. Generation and isolation of non-human light and heavy chain variable regions, followed by incorporation thereof within the human light chain constant transcriptional cassette and human heavy chain constant transcriptional cassette, allows for production of immunologically active chimeric anti-CD20 antibodies.

[0105] We have derived a most preferred non-human variable region with specificity to the CD20 antigen using a murine source and hybridoma technology. Using polymerase chain reaction ("PCR") techniques, the murine light and heavy variable regions were cloned directly into the TCAE 8 vector—this is the most preferred route for incorporation of the non-human variable region into the TCAE vector. This preference is principally predicated upon the efficiency of the PCR reaction and the accuracy of insertion. However, other equivalent procedures for accomplishing this task are available. For example, using TCAE 8 (or an equivalent vector), the sequence of the variable region of a non-human anti-CD20 antibody can be obtained, followed by oligonucleotide synthesis of portions of the sequence or, if appropriate, the entire sequence; thereafter, the portions or the entire synthetic sequence can be inserted into the appropriate locations within the vector. Those skilled in the art are credited with the ability to accomplish this task.

[0106] Our most preferred immunologically active chimeric anti-CD20 antibodies were derived from utilization of TCAE 8 vector which included murine variable regions derived from monoclonal antibody to CD20; this antibody (to be discussed in detail, infra), is referred to as "2B8." The complete sequence of the variable regions obtained from 2B8 in TCAE 8 ("anti-CD20 in TCAE 8") is set forth in **FIG. 3** (SEQ. ID. NO. 2).

[0107] The host cell line utilized for protein expression is most preferably of mammalian origin; those skilled in the art

are credited with ability to preferentially determine particular host cell lines which are best suited for the desired gene product to be expressed therein. Exemplary host cell lines include, but are not limited to, DG44 and DUXBII (Chinese Hamster Ovary lines, DHFR minus), HEGLA (human cervical carcinoma), CVI (monkey kidney line), COS (a derivative of CVI with SV40 T antigen), R1610 (Chinese hamster fibroblast) BALBC/3T3 (mouse fibroblast), HAK (hamster kidney line), SP2/O (mouse myeloma), P3x63-Ag3.653 (mouse myeloma), BFA-IclBPT (bovine endothelial cells), RAJI (human lymphocyte) and 293 (human kidney). Host cell lines are typically available from commercial services, the American Tissue Culture Collection or from published literature.

[0108] Preferably the host cell line is either DG44 ("CHO") or SP2/O. See Urland, G. et al., "Effect of gamma rays and the dihydrofolate reductase locus: deletions and inversions." *Som. Cell & Mol. Gen.* 12/6: 555-566 (1986), and Shulman, M. et al., "A better cell line for making hybridomas secreting specific antibodies." *Nature* 276: 269 (1978), respectively. Most preferably, the host cell line is DG44. Transfection of the plasmid into the host cell can be accomplished by any technique available to those in the art. These include, but are not limited to, transfection (including electrophoresis and electroporation), cell fusion with enveloped DNA, microinjection, and infection with intact virus. See, Ridgway, A. A. G. "Mammalian Expression Vectors." Chapter 24.2, pp. 470-472 Vectors, Rodriguez and Denhardt, Eds. (Butterworths, Boston, Mass. 1988). Most preferably, plasmid introduction into the host is via electroporation.

#### F. EXAMPLES

[0109] The following examples are not intended, nor are they to be construed, as limiting the invention. The examples are intended to evidence: dose-imaging using a radiolabeled anti-CD20 antibody ("12B8"); radiolabeled anti-CD20 antibody ("Y2B8"); and immunologically active, chimeric anti-CD20 antibody ("C2B8") derived utilizing a specific vector ("TCAE 8") and variable regions derived from murine anti-CD20 monoclonal antibody ("2B8").

[0110] I. Radiolabeled ANTI-CD20 Antibody 2B8

[0111] A. Anti-CD20 Monoclonal Antibody (Murine) Production ("2B8")

[0112] BALB/C mice were repeatedly immunized with the human lymphoblastoid cell line SB (see, Adams, R. A. et al., "Direct implantation and serial transplantation of human acute lymphoblastic leukemia in hamsters, SB-2." *Can Res* 28: 1121-1125 (1968); this cell line is available from the American Tissue Culture Collection, Rockville, Md., under ATCC accession number ATCC CCL 120), with weekly injections over a period of 3-4 months. Mice evidencing high serum titers of anti-CD20 antibodies, as determined by inhibition of known CD20-specific antibodies (anti-CD20 antibodies utilized were Leu 16, Beckton Dickinson, San Jose, Calif., Cat. No. 7670; and B1, Coulter Corp., Hialeah, Fla., Cat. No. 6602201) were identified; the spleens of such mice were then removed. Spleen cells were fused with the mouse myeloma SP2/0 in accordance with the protocol described in Einfeld, D. A. et al., (1988) *EMBO* 7: 711 (SP2/0 has ATCC accession no. ATCC CRL 8006).

[0113] Assays for CD20 specificity were accomplished by radioimmunoassay. Briefly, purified anti-CD20 B1 was

radiolabeled with  $I^{125}$  by the iodobead method as described in Valentine, M. A. et al., (1989) *J. Biol. Chem.* 264: 11282. ( $I^{125}$  Sodium Iodide, ICN, Irvine, Calif., Cat. No. 28665H). Hybridomas were screened by co-incubation of 0.05 ml of media from each of the fusion wells together with 0.05 ml of  $I^{125}$  labeled anti-CD20 B1 (10 ng) in 1% BSA, PBS (pH 7.4), and 0.5 ml of the same buffer containing 100,000 SB cells. After incubation for 1 hr at room temperature, the cells were harvested by transferring to 96 well titer plates (V&P Scientific, San Diego, Calif.), and washed thoroughly. Duplicate wells containing unlabeled anti-CD20 B1 and wells containing no inhibiting antibody were used as positive and negative controls, respectively. Wells containing greater than 50% inhibition were expanded and cloned. The antibody demonstrating the highest inhibition was derived from the cloned cell line designated herein as "2B8."

[0114] B. Preparation of 2B8-MX-DTPA Conjugate

[0115] i. MX-DTPA

[0116] Carbon-14-labeled 1-isothiocyanatobenzyl-3-methylidihethylene triaminepentaacetic acid ("carbon-14 labeled MX-DTPA") was used as a chelating agent for conjugation of radiolabel to 2B8. Manipulations of MX-DTPA were conducted to maintain metal-free conditions, ie metal-free reagents were utilized and, when possible, polypropylene plastic containers (flasks, beakers, graduated cylinders, pipette tips) washed with Alconox and rinsed with Milli-Q water, were similarly utilized. MX-DTPA was obtained as a dry solid from Dr. Otto Gansow (National Institute of Health, Bethesda, Md.) and stored desiccated at 4° C. (protected from light), with stock solutions being prepared in Milli-Q water at a concentration of 2-5 mM, with storage at -70° C. MX-DTPA was also obtained from Coulter Immunology (Hialeah, Fla.) as the disodium salt in water and stored at -70° C.

[0117] ii. Preparation of 2B8

[0118] Purified 2B8 was prepared for conjugation with MX-DTPA by transferring the antibody into metal-free 50 mM bicine-NaOff, pH 8.6, containing 150 mM NaCl, using repetitive buffer exchange with CENTRICON 30™ spin filters (30,000D, MWCO; Amicon). Generally, 50-200  $\mu$ L of protein (10 mg/ml) was added to the filter unit, followed by 2 mL of bicine buffer. The filter was centrifuged at 4° C. in a Sorval SS-34 rotor (6,000 rpm, 45 min.). Retentate volume was approximately 50-100  $\mu$ L; this process was repeated twice using the same filter. Retentate was transferred to a polypropylene 1.5 mL screw cap tube, assayed for protein, diluted to 10.0 mg/mL and stored at 4° C. until utilized; protein was similarly transferred into 50 mM sodium citrate, pH 5.5, containing 150 mM NaCl and 0.05% sodium azide, using the foregoing protocol.

[0119] iii. Conjugation of 2B8 with MX-DTPA

[0120] Conjugation of 2B8 with MX-DTPA was performed in polypropylene tubes at ambient temperature. Frozen MX-DTPA stock solutions were thawed immediately prior to use. 50-200 mL of protein at 10 mg/mL were reacted with MX-DTPA at a molar ratio of MX-DTPA-to-2B8 of 4:1. Reactions were initiated by adding the MX-DTPA stock solution and gently mixing; the conjugation was allowed to proceed overnight (14 to 20 hr), at ambient temperature. Unreacted MX-DTPA was removed from the conjugate by dialysis or repetitive ultrafiltration, as described above in

Example I.B.ii, into metal-free normal saline (0.9% w/v) containing 0.05% sodium azide. The protein concentration was adjusted to 10 mg/mL and stored at 4° C. in a polypropylene tube until radiolabeled.

[0121] iv. Determination of MX-DTPA Incorporation

[0122] MX-DTPA incorporation was determined by scintillation counting and comparing the value obtained with the purified conjugate to the specific activity of the carbon-[14]-labeled MX-DTPA. For certain studies, in which non-radioactive MX-DTPA (Coulter Immunology) was utilized, MX-DTPA incorporation was assessed by incubating the conjugate with an excess of a radioactive carrier solution of yttrium-[90] of known concentration and specific activity.

[0123] A stock solution of yttrium chloride of known concentration was prepared in metal-free 0.05 N HCl to which carrier-free yttrium-[90] (chloride salt) was added. An aliquot of this solution was analyzed by liquid scintillation counting to determine an accurate specific activity for this reagent. A volume of the yttrium chloride reagent equal to 3-times the number of mols of chelate expected to be attached to the antibody, (typically 2 mol/mol antibody), was added to a polypropylene tube, and the pH adjusted to 4.0-4.5 with 2 M sodium acetate. Conjugated antibody was subsequently added and the mixture incubated 15-30 min. at ambient temperature. The reaction was quenched by adding 20 mM EDTA to a final concentration of 1 mM and the pH of the solution adjusted to approximately pH 6 with 2M sodium acetate.

[0124] After a 5 min. incubation, the entire volume was purified by high-performance, size-exclusion chromatography (described infra). The eluted protein-containing fractions were combined, the protein concentration determined, and an aliquot assayed for radioactivity. The chelate incorporation was calculated using the specific activity of the yttrium-[90] chloride preparation and the protein concentration.

[0125] v. Immunoreactivity of 2B8-MX-DTPA

[0126] The immunoreactivity of conjugated 2B8 was assessed using whole-cell ELISA. Mid-log phase SB cells were harvested from culture by centrifugation and washed two times with 1xHBSS. Cells were diluted to 1-2 $\times$ 10<sup>6</sup> cells/mL in HBSS and aliquoted into 96-well polystyrene microtiter plates at 50,000-100,000 cells/well. The plates were dried under vacuum for 2 h. at 40-45° C. to fix the cells to the plastic; plates were stored dry at -20° C. until utilized. For assay, the plates were warmed to ambient temperature immediately before use, then blocked with 1xPBS, pH 7.2-7.4 containing 1% BSA (2 h). Samples for assay were diluted in 1xPBS/1% BSA, applied to plates and serially diluted (1:2) into the same buffer. After incubating plates for 1 h. at ambient temperature, the plates were washed three times with 1xPBS. Secondary antibody (goat anti-mouse IgG1-specific HRP conjugate 50  $\mu$ L) was added to wells (1:1500 dilution in 1xPBS/1% BSA) and incubated 1 h. at ambient temperature. Plates were washed four times with 1xPBS followed by the addition of ABTS substrate solution (50 mM sodium citrate, pH 4.5 containing 0.01% ATBS and 0.001% H<sub>2</sub>O<sub>2</sub>). Plates were read at 405 nm after 15-30 min. incubation. Antigen-negative HSB cells were included in assays to monitor non-specific binding. Immunoreactivity of the conjugate was calculated by plotting the absorbance

values vs. the respective dilution factor and comparing these to values obtained using native antibody (representing 100% immunoreactivity) tested on the same plate; several values on the linear portion of the titration profile were compared and a mean value determined (data not shown).

[0127] vi. Preparation of Indium-[111]-Labeled 2B8-MX-DTPA ("I2B8")

[0128] Conjugates were radiolabeled with carrier-free indium-[111]. An aliquot of isotope (0.1-2 mCi/mg antibody) in 0.05 M HCl was transferred to a polypropylene tube and approximately one-tenth volume of metal-free 2 M HCl added. After incubation for 5 min., metal-free 2 M sodium acetate was added to adjust the solution to pH 4.0-4.4. Approximately 0.5 mg of 2B8-MX-DTPA was added from a stock solution of 10.0 mg/mL DTPA in normal saline, or 50 mM sodium citrate/150 mM NaCl containing 0.05% sodium azide, and the solution gently mixed immediately. The pH solution was checked with pH paper to verify a value of 4.0-4.5 and the mixture incubated at ambient temperature for 15-30 min. Subsequently, the reaction was quenched by adding 20 mM EDTA to a final concentration of 1 mM and the reaction mixture was adjusted to approximately pH 6.0 using 2 M sodium acetate.

[0129] After a 5-10 min. incubation, uncomplexed radioisotope was removed by size-exclusion chromatography. The HPLC unit consisted of Waters Model 6000 or Toso-Haas Model TSK-6110 solvent delivery system fitted, respectively, with a Waters U6K or Rheodyne 700 injection valve. Chromatographic separations were performed using a gel permeation column (BioRad SEC-250; 7.5×300 mm or comparable TosoHaas column) and a SEC-250 guard column (7.5×100 mm). The system was equipped with a fraction collector (Pharmacia Frac200) and a UV monitor fitted with a 280 nm filter (Pharmacia model UV-1). Samples were applied and eluted isocratically using 1×PBS, pH 7.4, at 1.0 mL/min flow rate. One-half milliliter fractions were collected in glass tubes and aliquots of these counted in a gamma counter. The lower and upper windows were set to 100 and 500 KeV respectively.

[0130] The radioincorporation was calculated by summing the radioactivity associated with the eluted protein peak and dividing this number by the total radioactivity eluted from the column; this value was then expressed as a percentage (data not shown). In some cases, the radioincorporation was determined using instant thin-layer chromatography ("ITLC"). Radiolabeled conjugate was diluted 1:10 or 1:20 in 1×PBS containing or 1×PBS/1 mM DTPA, then 1  $\mu$ L was spotted 1.5 cm from one end of a 1×5 cm strip of ITLC SG paper. The paper was developed by ascending chromatography using 10% ammonium acetate in methanol:water (1:1;v/v). The strip was dried, cut in half crosswise, and the radioactivity associated with each section determined by gamma counting. The radioactivity associated with the bottom half of the strip (protein-associated radioactivity) was expressed as a percentage of the total radioactivity, determined by summing the values for both top and bottom halves (data not shown).

[0131] Specific activities were determined by measuring the radioactivity of an appropriate aliquot of the radiolabeled conjugate. This value was corrected for the counter efficiency (typically 75%) and related to the protein concentra-

tion of the conjugate, previously determined by absorbance at 280 nm, and the resulting value expressed as mCi/mg protein.

[0132] For some experiments, 2B8-MX-DTPA was radiolabeled with indium [111] following a protocol similar to the one described above but without purification by HPLC; this was referred to as the "mix-and-shoot" protocol.

[0133] vii. Preparation of Yttrium-[90]-Labeled 2B8-MX-DTPA ("Y2B8")

[0134] The same protocol described for the preparation of I2B8 was followed for the preparation of the yttrium-[90]-labeled 2B8-MX-DTPA ("Y2B8") conjugate except that 2 ng HCl was not utilized; all preparations of yttrium-labeled conjugates were purified by size-exclusion chromatography as described above.

[0135] C. Non-Human Animal Studies.

[0136] i. Biodistribution of Radiolabeled 2B8-MX-DTPA

[0137] I2B8 was evaluated for tissue biodistribution in six-to-eight week old BALB/c mice. The radiolabeled conjugate was prepared using clinical-grade 2B8-MX-DTPA following the "mix and shoot" protocol described above. The specific activity of the conjugate was 2.3 mCi/mg and the conjugate was formulated in PBS, pH 7.4 containing 50 mg/mL HSA. Mice were injected intravenously with 100  $\mu$ L of I2B8 (approximately 21  $\mu$ Ci) and groups of three mice were sacrificed by cervical dislocation at 0, 24, 48, and 72 hours. After sacrifice, the tail, heart, lungs, liver, kidney, spleen, muscle, and femur were removed, washed and weighed; a sample of blood was also removed for analysis. Radioactivity associated with each specimen was determined by gamma counting and the percent injected dose per gram tissue subsequently determined. No attempt was made to discount the activity contribution represented by the blood associated with individual organs.

[0138] In a separate protocol, aliquots of 2B8-MX-DTPA incubated at 4° C. and 30° C. for 10 weeks were radiolabeled with indium-[111] to a specific activity of 2.1 mCi/mg for both preparations. These conjugates were then used in biodistribution studies in mice as described above.

[0139] For dosimetry determinations, 2B8-MX-DTPA was radiolabeled with indium-[111] to a specific activity of 2.3 mCi/mg and approximately 1.1  $\mu$ Ci was injected into each of 20 BALB/c mice. Subsequently, groups of five mice each were sacrificed at 1, 24, 48 and 72 hours and their organs removed and prepared for analysis. In addition, portions of the skin, muscle and bone were removed and processed for analysis; the urine and feces were also collected and analyzed for the 24-72 hour time points.

[0140] Using a similar approach, 2B8-MX-DTPA was also radiolabeled with yttrium-[90] and its biological distribution evaluated in BALB/c mice over a 72-hour time period. Following purification by HPLC size exclusion chromatography, four groups of five mice each were injected intravenously with approximately 1  $\mu$ Ci of clinically-formulated conjugate (specific activity: 12.2 mCi/mg); groups were subsequently sacrificed at 1, 24, 48 and 72 hours and their organs and tissues analyzed as described above. Radioactivity associated with each tissue specimen was determined by measuring bremsstrahlung energy with a gamma scintillation counter. Activity values were subsequently expressed

as percent injected dose per gram tissue or percent injected dose per organ. While organs and other tissues were rinsed repeatedly to remove superficial blood, the organs were not perfused. Thus, organ activity values were not discounted for the activity contribution represented by internally associated blood.

[0141] ii. Tumor Localization of I2B8

[0142] The localization of radiolabeled 2B8-MX-DTPA was determined in athymic mice bearing Ramos B cell tumors. Six-to-eight week old athymic mice were injected subcutaneously (left-rear flank) with 0.1 mL of RPMI-1640 containing  $1.2 \times 10^7$  Ramos tumor cells which had been previously adapted for growth in athymic mice. Tumors arose within two weeks and ranged in weight from 0.07 to 1.1 grams. Mice were injected intravenously with 100  $\mu$ L of indium-[111]-labeled 2B8-MX-DTPA (16.7  $\mu$ Ci) and groups of three mice were sacrificed by cervical dislocation at 0, 24, 48, and 72 hours. After sacrifice the tail, heart, lungs, liver, kidney, spleen, muscle, femur, and tumor were removed, washed, weighed; a sample of blood was also removed for analysis. Radioactivity associated with each specimen was determined by gamma counting and the percent injected dose per gram tissue determined.

[0143] iii. Biodistribution and Tumor Localization Studies with Radiolabeled 2B8-MX-DTPA

[0144] Following the preliminary biodistribution experiment described above (Example I.B.viii.a.), conjugated 2B8 was radiolabeled with indium-[111] to a specific activity of 2.3 mCi/mg and roughly 1.1  $\mu$ Ci was injected into each of twenty BALB/c mice to determine biodistribution of the radiolabeled material. Subsequently, groups of five mice each were sacrificed at 1, 24, 48 and 72 hours and their organs and a portion of the skin, muscle and bone were removed and processed for analysis. In addition, the urine and feces were collected and analyzed for the 24-72 hour time-points. The level of radioactivity in the blood dropped from 40.3% of the injected dose per gram at 1 hour to 18.9% at 72 hours (data not shown). Values for the heart, kidney, muscle and spleen remained in the range of 0.7-9.8% throughout the experiment. Levels of radioactivity found in the lungs decreased from 14.2% at 1 hour to 7.6% at 72 hours; similarly the respective liver injected-dose per gram values were 10.3% and 9.9%. These data were used in determining radiation absorbed dose estimates I2B8 described below.

[0145] The biodistribution of yttrium-[90]-labeled conjugate, having a specific activity of 12.2 mCi/mg antibody, was evaluated in BALB/c mice. Radioincorporations of >90% were obtained and the radiolabeled antibody was purified by HPLC. Tissue deposition of radioactivity was evaluated in the major organs, and the skin, muscle, bone, and urine and feces over 72 hours and expressed as percent injected dose/g tissue. Results (not shown) evidenced that while the levels of radioactivity associated with the blood dropped from approximately 39.2% injected dose per gram at 1 hour to roughly 15.4% after 72 hours the levels of radioactivity associated with tail, heart, kidney, muscle and spleen remained fairly constant at 10.2% or less throughout the course of the experiment. Importantly, the radioactivity associated with the bone ranged from 4.4% of the injected dose per gram bone at 1 hour to 3.2% at 72 hours. Taken together, these results suggest that little free yttrium was

associated with the conjugate and that little free radiometal was released during the course of the study. These data were used in determining radiation absorbed dose estimates for Y2B8 described below.

[0146] For tumor localization studies, 2B8-MX-DTPA was prepared and radiolabeled with <sup>111</sup>Indium to a specific activity of 2.7 mCi/mg. One hundred microliters of labeled conjugate (approximately 24  $\mu$ Ci) were subsequently injected into each of 12 athymic mice bearing Ramos B cell tumors. Tumors ranged in weight from 0.1 to 1.0 grams. At time points of 0, 24, 48, and 72 hours following injection, 50  $\mu$ L of blood was removed by retro-orbital puncture, the mice sacrificed by cervical dislocation, and the tail, heart, lungs, liver, kidney, spleen, muscle, femur, and tumor removed. After processing and weighing the tissues, the radioactivity associated with each tissue specimen was determined using a gamma counter and the values expressed as percent injected dose per gram.

[0147] The results (not shown) evidenced that the tumor concentrations of the <sup>111</sup>In-2B8-MX-DTPA increased steadily throughout the course of the experiment. Thirteen percent of the injected dose was accumulated in the tumor after 72 hours. The blood levels, by contrast, dropped during the experiment from over 30% at time zero to 13% at 72 hours. All other tissues (except muscle) contained between 1.3 and 6.0% of the injected dose per gram tissue by the end of the experiment; muscle tissue contained approximately 13% of the injected dose per gram.

[0148] D. Human Studies

[0149] i. 2B8 and 2B8-MX-DTPA: Immunohistology Studies with Human Tissues

[0150] The tissue reactivity of murine monoclonal antibody 2B8 was evaluated using a panel of 32 different human tissues fixed with acetone. Antibody 2B8 reacts with the anti-CD20 antigen which had a very restricted pattern of tissue distribution, being observed only in a subset of cells in lymphoid tissues including those of hematopoietic origin.

[0151] In the lymph node, immunoreactivity was observed in a population of mature cortical B-lymphocytes as well as proliferating cells in the germinal centers. Positive reactivity was also observed in the peripheral blood, B-cell areas of the tonsils, white pulp of the spleen, and with 40-70% of the medullary lymphocytes found in the thymus. Positive reactivity was also seen in the follicles of the lamina propria (Peyer's Patches) of the large intestines. Finally, aggregates or scattered lymphoid cells in the stroma of various organs, including the bladder, breast, cervix, esophagus, lung, parotid, prostate, small intestine, and stomach, were also positive with antibody 2B8 (data not shown).

[0152] All simple epithelial cells, as well as the stratified epithelia and epithelia of different organs, were found to be unreactive. Similarly, no reactivity was seen with neuroectodermal cells, including those in the brain, spinal cord and peripheral nerves. Mesenchymal elements, such as skeletal and smooth muscle cells, fibroblasts, endothelial cells, and polymorphonuclear inflammatory cells were also found to be negative (data not shown).

[0153] The tissue reactivity of the 2B8-MX-DTPA conjugate was evaluated using a panel of sixteen human tissues which had been fixed with acetone. As previously demon-

strated with the native antibody (data not shown), the 2B8-MX-DTPA conjugate recognized the CD20 antigen which exhibited a highly restricted pattern of distribution, being found only on a subset of cells of lymphoid origin. In the lymph node, immunoreactivity was observed in the B cell population. Strong reactivity was seen in the white pulp of the spleen and in the medullary lymphocytes of the thymus. Immunoreactivity was also observed in scattered lymphocytes in the bladder, heart, large intestines, liver, lung, and uterus, and was attributed to the presence of inflammatory cells present in these tissues. As with the native antibody, no reactivity was observed with neuroectodermal cells or with mesenchymal elements (data not shown).

[0154] ii. Clinical Analysis of I2B8 (Imaging) and Y2B8 (Therapy)

[0155] a. Phase I/II Clinical Trial Single Dose Therapy Study

[0156] A Phase I/II clinical analysis of I2B8 (imaging) followed by treatment with a single therapeutic dose of Y2B8 is currently being conducted. For the single-dose study, the following schema is being followed:

[0157] 1. Peripheral Stem Cell (PSC) or Bone Marrow (BM) Harvest with Purging;

[0158] 2. I2B8 Imaging;

[0159] 3. Y2B8 Therapy (three Dose Levels); and

[0160] 4. PSC or Autologous BM Transplantation (if necessary based upon absolute neutrophil count below 500/mm<sup>3</sup> for three consecutive days or platelets below 20,000/mm<sup>3</sup> with no evidence of marrow recovery on bone marrow examination).

[0161] The Dose Levels of Y2B8 are as follows:

Dose Level	Dose (mCi)
1.	20
2.	30
3.	40

[0162] Three patients are to be treated at each of the dose levels for determination of a Maximum Tolerated Dose ("MTD").

[0163] Imaging (Dosimetry) Studies are conducted as follows: each patient is involved in two in vivo biodistribution studies using I2B8. In the first study, 2 mg of I2B8 (5 mCi), is administered as an intravenous (i.v.) infusion over one hour; one week later 2B8 (ie unconjugated antibody) is administered by i.v. at a rate not to exceed 250 mg/hr followed immediately by 2 mg of I2B8 (5 mCi) administered by i.v. over one hour. In both studies, immediately following the I2B8 infusion, each patient is imaged and imaging is repeated at time t=14-18 hr (if indicated), t=24 hr; t=72 hr; and t=96 hr (if indicated). Whole body average retention times for the indium [111] label are determined; such determinations are also made for recognizable organs or tumor lesions ("regions of interest").

[0164] The regions of interest are compared to the whole body concentrations of the label; based upon this compari-

son, an estimate of the localization and concentration of Y2B8 can be determined using standard protocols. If the estimated cumulative dose of Y2B8 is greater than eight (8) times the estimated whole body dose, or if the estimated cumulative dose for the liver exceeds 1500 cGy, no treatment with Y2B8 should occur.

[0165] If the imaging studies are acceptable, either 0.0 or 1.0 mg/kg patient body weight of 2B8 is administered by i.v. infusion at a rate not to exceed 250 mg/h. This is followed by administration of Y2B8 (10, 20 or 40 mCi) at an i.v. infusion rate of 20 mCi/hr.

[0166] b. Phase I/II Clinical Trial: Multiple Dose Therapy Study

[0167] A Phase I/II clinical analysis of Y2B8 is currently being conducted. For the multiple-dose study, the following schema is being followed:

[0168] 1. PSC or BM Harvest;

[0169] 2. I2B8 Imaging,

[0170] 3. Y2B8 Therapy (three Dose Levels) for four doses or a total cumulative dose of 80 mCi; and

[0171] 4. PSC or Autologous BM Transplantation (based upon decision of medical practitioner).

[0172] The Dose Levels of Y2B8 are as follows:

Dose Level	Dose (mCi)
1.	10
2.	15
3.	20

[0173] Three patients are to be treated at each of the dose levels for determination of an MTD.

[0174] Imaging (Dosimetry) Studies are conducted as follows: A preferred imaging dose for the unlabeled antibody (ie 2B8) will be determined with the first two patients. The first two patients will receive 100 mg of unlabeled 2B8 in 250 cc of normal saline over 4 hrs followed by 0.5 mCi of I2B8—blood will be sampled for biodistribution data at times t=0, t=10 min., t=120 min., t=24 hr, and t=48 hr. Patients will be scanned with multiple regional gamma camera images at times t=2 hr, t=24 hr and t=48 hr. After scanning at t=48 hr, the patients will receive 250 mg of 2B8 as described, followed by 4.5 mCi of I2B8—blood and scanning will then follow as described. If 100 mg of 2B8 produces superior imaging, then the next two patients will receive 50 mg of 2B8 as described, followed by 0.5 mCi of I2B8 followed 48 hrs later by 100 mg 2B8 and then with 4.5 mCi of I2B8. If 250 mg of 2B8 produces superior imaging, then the next two patients will receive 250 mg of 2B8 as described, followed by 0.5 mCi of I2B8 followed 48 hrs later with 500 mg 2B8 and then with 4.5 mCi of I2B8. Subsequent patients will be treated with the lowest amount of 2B8 that provides optimal imaging. Optimal imaging will be defined by: (1) best effective imaging with the slowest disappearance of antibody; (2) best distribution minimizing compartmentalization in a single organ; and (3) best subjective resolution of the lesion (tumor/background comparison).

**[0175]** For the first four patients, the first therapeutic dose of Y2B8 will begin 14 days after the last dose of I2B8; for subsequent patients, the first therapeutic dose of Y2B8 will begin between two to seven days after the I2B8.

**[0176]** Prior to treatment with Y2B8, for the patients other than the first four, 2B8 will be administered as described, followed by i.v. infusion of Y2B8 over 5-10 min. Blood will be sampled for biodistribution at times t=0, t=10 min., t=120 min., t=24 hr and t=48 hr. Patients will receive repetitive doses of Y2B8 (the same dose administered as with the first dose) approximately every six to eight weeks for a maximum of four doses, or total cumulative dose of 80 mCi. It is most preferred that patients not receive a subsequent dose of Y2B8 until the patients' WBC is greater than/equal to 3,000 and AGC is greater than/equal to 100,000.

**[0177]** Following completion of the three-dose level study, an MTD will be defined. Additional patients will then be enrolled in the study and these will receive the MTD.

**[0178]** II. Chimeric ANTI-CD20 Antibody Production ("C2B8")

**[0179]** A. Construction of Chimeric Anti-CD20 Immuno-globulin DNA Expression Vector

**[0180]** RNA was isolated from the 2B8 mouse hybridoma cell (as described in Chomczynki, P. et al., "Single step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction." *Anal. Biochem.* 162: 156-159 (1987)), and cDNA was prepared therefrom. The mouse immunoglobulin light chain variable region DNA was isolated from the cDNA by polymerase chain reaction using a set of DNA primers with homology to mouse light chain signal sequences at the 5' end and mouse light chain J region at the 3' end. Primer sequences were as follows:

1.  $V_L$  Sense (SEQ. ID. NO. 3)

5' ATC AC AGATCT CTC ACC ATG GAT TTT CAG GTG  
CAG ATT ATC AGC TTC 3'

(The underlined portion is a Bgl II site; the above-lined portion is the start codon.)

2.  $V_L$  Antisense (SEQ. ID. NO. 4)

5' TGC AGC ATC CGTAGC TTT GAT TTC CAG CTT 3'  
(The underlined portion is a Bsi WI site.)

**[0181]** See, **FIGS. 1 and 2** for the corresponding Bgl II and Bsi WI sites in TCAE 8, and **FIG. 3** for the corresponding sites in anti-CD20 in TCAE 8.

**[0182]** These resulting DNA fragment was cloned directly into the TCAE 8 vector in front of the human kappa light chain constant domain and sequenced. The determined DNA sequence for the murine variable region light chain is set forth in **FIG. 4** (SEQ. ID. NO. 5); see also **FIG. 3**, nucleotides 978 through 1362. **FIG. 4** further provides the amino acid sequence from this murine variable region, and the CDR and framework regions. The mouse light chain variable region from 2B8 is in the mouse kappa VI family. See, Kabat, *supra*.

**[0183]** The mouse heavy chain variable region was similarly isolated and cloned in front of the human IgG1 constant domains. Primers were as follows:

1.  $V_H$  Sense (SEQ. ID. NO. 6)

5' GCG GCT CCC ACGGGT GTC CTG TCC CAG 3'

(The underlined portion is an Mlu I site.)

2.  $V_H$  Antisense (SEQ. ID. NO. 7)

5' GG(G/C) TGT TGT GCTAGC TG(A/C) (A/G)GA GAC  
(G/A)GT GA 3'

(The underlined portion is an Nhe I site.)

**[0184]** See, **FIGS. 1 and 2** for corresponding Mlu I and Nhe I sites in TCAE 8, and **FIG. 3** for corresponding sites in anti-CD20 in TCAE 8.

**[0185]** The sequence for this mouse heavy chain is set forth in **FIG. 5** (SEQ. ID. NO. 8); see also **FIG. 3**, nucleotide 2401 through 2820. **FIG. 5** also provides the amino acid sequence from this murine variable region, and the CDR and framework regions. The mouse heavy chain variable region from 2B8 is in the mouse VH 2B family. See, Kabat, *supra*.

**[0186]** B. Creation of Chimeric Anti-CD20 Producing CHO and SP2/0 Transfectomas

**[0187]** Chinese hamster ovary ("CHO") cells DG44 were grown in SSFM II minus hypoxanthine and thymidine media (Gibco, Grand Island, N.Y., Form No. 91-0456PK); SP2/0 mouse myeloma cells were grown in Dulbecco's Modified Eagles Medium media ("DMEM") (Irvine Scientific, Santa Ana, Calif., Cat. No. 9024) with 5% fetal bovine serum and 20 ml/L glutamine added. Four million cells were electroporated with either 25  $\mu$ g CHO or 50  $\mu$ g SP2/0 plasmid DNA that had been restricted with Not I using a BTX 600 electroporation system (BTX, San Diego, Calif.) in 0.4 ml disposable cuvettes. Conditions were either 210 volts for CHO or 180 volts for SP2/0, 400 microfaradays, 13 ohms. Each electroporation was plated into six 96 well dishes (about 7,000 cells/well). Dishes were fed with media containing G418 (GENETICIN, Gibco, Cat. No. 860-1811) at 400  $\mu$ g/ml active compound for CHO (media further included 50  $\mu$ M hypoxanthine and 8  $\mu$ M thymidine) or 800  $\mu$ g/ml for SP2/0, two days following electroporation and thereafter 2 or 3 days until colonies arose. Supernatant from colonies was assayed for the presence of chimeric immunoglobulin via an ELISA specific for human antibody. Colonies producing the highest amount of immunoglobulin were expanded and plated into 96 well plates containing media plus methotrexate (25 nM for SP2/0 and 5 nM for CHO) and fed every two or three days. Supernatants were assayed as above and colonies producing the highest amount of immunoglobulin were examined. Chimeric anti-CD20 antibody was purified from supernatant using protein A affinity chromatography.

**[0188]** Purified chimeric anti-CD20 was analyzed by electrophoresis in polyacrylamide gels and estimated to be greater than about 95% pure. Affinity and specificity of the chimeric antibody was determined based upon 2B8. Chimeric anti-CD20 antibody tested in direct and competitive

binding assays, when compared to murine anti-CD20 monoclonal antibody 2B8, evidenced comparable affinity and specificity on a number of CD20 positive B cells lines (data not presented). The apparent affinity constant ("K<sub>a</sub>") of the chimeric antibody was determined by direct binding of <sup>125</sup>I radiolabeled chimeric anti-CD20 and compared to radiolabeled 2B8 by Scatchard plot; estimated K<sub>a</sub> for CHO produced chimeric anti-CD20 was  $5.2 \times 10^{-9}$  M and for SP2/0 produced antibody,  $7.4 \times 10^{-9}$  M. The estimated K<sub>a</sub> for 2B8 was  $3.5 \times 10^{-9}$  M. Direct competition by radioimmunoassay was utilized to confirm both the specificity and retention of immunoreactivity of the chimeric antibody by comparing its ability to effectively compete with 2B8. Substantially equivalent amounts of chimeric anti-CD20 and 2B8 antibodies were required to produce 50% inhibition of binding to CD20 antigens on B cells (data not presented), ie there was a minimal loss of inhibiting activity of the anti-CD20 antibodies, presumably due to chimerization.

[0189] The results of Example II.B indicate, inter alia, that chimeric anti-CD20 antibodies were generated from CHO and SP2/0 transfectomas using the TCAE 8 vectors, and these chimeric antibodies had substantially the same specificity and binding capability as murine anti-CD20 monoclonal antibody 2B8.

[0190] C. Determination of Immunological Activity of Chimeric Anti-CD20 Antibodies

[0191] i. Human C1q Analysis

[0192] Chimeric anti-CD20 antibodies produced by both CHO and SP2/0 cell lines were evaluated for human C1q binding in a flow cytometry assay using fluorescein labeled C1q (C1q was obtained from Quidel, Mira Mesa, Calif., Prod. No. A400 and FITC label from Sigma, St. Louis Mo., Prod. No. F-7250; FITC. Labeling of C1q was accomplished in accordance with the protocol described in *Selected Methods In Cellular Immunology*, Michell & Shiigi, Ed. (W.H. Freeman & Co., San Francisco, Calif., 1980, p. 292). Analytical results were derived using a Becton Dickinson FACScan™ flow cytometer (fluorescein measured over a range of 515-545 nm). Equivalent amounts of chimeric anti-CD20 antibody, human IgG1,K myeloma protein (Binding Site, San Diego, Calif., Prod. No. BP078), and 2B8 were incubated with an equivalent number of CD20-positive SB cells, followed by a wash step with FACS buffer (0.2% BSA in PBS, pH 7.4, 0.02% sodium azide) to remove unattached antibody, followed by incubation with FITC labeled C1q. Following a 30-60 min. incubation, cells were again washed. The three conditions, including FITC-labeled C1q as a control, were analyzed on the FACScan™ following manufacturing instructions. Results are presented in FIG. 6.

[0193] As the results of FIG. 6 evidence, a significant increase in fluorescence was observed only for the chimeric anti-CD20 antibody condition; ie only SB cells with adherent chimeric anti-CD20 antibody were C1q positive, while the other conditions produced the same pattern as the control.

[0194] ii. Complement Dependent Cell Lyses

[0195] Chimeric anti-CD20 antibodies were analyzed for their ability to lyse lymphoma cell lines in the presence of human serum (complement source). CD20 positive SB cells were labeled with <sup>51</sup>Cr by admixing 1001  $\mu$ Ci of <sup>51</sup>Cr with  $1 \times 10^6$  SB cells for 1 hr at 37° C.; labeled SB cells were then

incubated in the presence of equivalent amounts of human complement and equivalent amounts (0-50  $\mu$ g/ml) of either chimeric anti-CD20 antibodies or 2B8 for 4 hrs at 37° C. (see, Brunner, K. T. et al., "Quantitative assay of the lytic action of immune lymphoid cells on <sup>51</sup>Cr-labeled allogeneic target cells in vitro." *Immunology* 14: 181-189 (1968). Results are presented in FIG. 7.

[0196] The results of FIG. 7 indicate, inter alia, that chimeric anti-CD20 antibodies produced significant lysis (49%) under these conditions.

[0197] iii. Antibody Dependent Cellular Cytotoxicity Effector Assay

[0198] For this study, CD20 positive cells (SB) and CD20 negative cells (T cell leukemia line HSB; see, Adams, Richard, "Formal Discussion," *Can. Res.* 27: 2479-2482 (1967); ATCC deposit no. ATCC CCL 120.1) were utilized; both were labeled with <sup>51</sup>Cr. Analysis was conducted following the protocol described in Brunner, K. T. et al., "Quantitative assay of the lytic action of immune lymphoid cells on <sup>51</sup>Cr-labeled allogeneic target cells in vitro; inhibition by isoantibody and drugs." *Immunology* 14: 181-189 (1968); a substantial chimeric anti-CD20 antibody dependent cell mediated lysis of CD20 positive SB target cells (<sup>51</sup>Cr-labeled) at the end of a 4 hr, 37° C. incubation, was observed and this effect was observed for both CHO and SP2/0 produced antibody (effector cells were human peripheral lymphocytes; ratio of effector cells:target was 100:1). Efficient lysis of target cells was obtained at 3.9  $\mu$ g/ml. In contrast, under the same conditions, the murine anti-CD20 monoclonal antibody 2B8 had a statistically insignificant effect, and CD20 negative HSB cells were not lysed. Results are presented in FIG. 8.

[0199] The results of Example II indicate, inter alia, that the chimeric anti-CD20 antibodies of Example I were immunologically active.

[0200] III. Depletion of B Cells In Vivo using Chimeric ANTI-CD20

[0201] A. Non-Human Primate Study

[0202] Three separate non-human primate studies were conducted. For convenience, these are referred to herein as "Chimeric Anti-CD20: CHO & SP2/0;" "Chimeric Anti-CD20: CHO;" and "High Dosage Chimeric Anti-CD20." Conditions were as follows:

[0203] Chimeric Anti-CD20: CHO & SP2/0

[0204] Six cynomolgus monkeys ranging in weight from 4.5 to 7 kilograms (White Sands Research Center, Alamogordo, N. Mex.) were divided into three groups of two monkeys each. Both animals of each group received the same dose of immunologically active chimeric anti-CD20 antibody. One animal in each group received purified antibody produced by the CHO transfectoma; the other received antibody produced by the SP2/0 transfectoma. The three groups received antibody dosages corresponding to 0.1 mg/kg, 0.4 mg/kg, and 1.6 mg/kg each day for four (4) consecutive days. The chimeric immunologically active anti-CD20 antibody, which was admixed with sterile saline, was administered by intravenous infusion; blood samples were drawn prior to each infusion.

[0205] Additional blood samples were drawn beginning 24 hrs after the last injection (T=0) and thereafter on days

1, 3, 7, 14 and 28; blood samples were also taken thereafter at biweekly intervals until completion of the study at day 90.

[0206] Approximately 5 ml of whole blood from each animal was centrifuged at 2000 RPM for 5 min. Plasma was removed for assay of soluble chimeric anti-CD20 antibody levels. The pellet (containing peripheral blood leukocytes and red blood cells) was resuspended in fetal calf serum for fluorescent-labeled antibody analysis (see, "Fluorescent Antibody Labeling of Lymphoid Cell Population," infra.).

[0207] Chimeric Anti-CD20: CHO

[0208] Six cynomolgus monkeys ranging in weight from 4 to 6 kilograms (White Sands) were divided into three groups of two monkeys each. All animals were injected with immunologically active chimeric anti-CD20 antibodies produced from the CHO transfecoma (in sterile saline). The three groups were separated as follows: subgroup 1 received daily intravenous injections of 0.01 mg/kg of the antibody over a four (4) day period; subgroup 2 received daily intravenous injections of 0.4 mg/kg of the antibody over a four (4) day period; subgroup 3 received a single intravenous injection of 6.4 mg/kg of the antibody. For all three subgroups, a blood sample was obtained prior to initiation of treatment; additionally, blood samples were also drawn at T=0, 1, 3, 7, 14 and 28 days following the last injection, as described above, and these samples were processed for fluorescent labeled antibody analysis (see, "Fluorescent Antibody Labeling," infra.). In addition to peripheral blood B cell quantitation, lymph node biopsies were taken at days 7, 14 and 28 following the last injection, and a single cell preparation stained for quantitation of lymphocyte populations by flow cytometry.

[0209] High Dosage Chimeric Anti-CD20

[0210] Two cynomolgus monkeys (White Sands) were infused with 16.8 mg/kg of the immunologically active chimeric anti-CD20 antibodies from the CHO transfecomas (in sterile saline) weekly over a period of four consecutive weeks. At the conclusion of the treatment, both animals were anesthetized for removal of bone marrow; lymph node biopsies were also taken. Both sets of tissue were stained for the presence of B lymphocytes using Leu 16 by flow cytometry following the protocol described in Ling, N. R. et al., "B-cell and plasma cell antigens." *Leucocyte Typing III: White Cell Differentiations Antigens*, A. J. McMichael, Ed. (Oxford University Press, Oxford UK, 1987), p. 302.

[0211] Fluorescent Antibody Labeling of Lymphoid Cell Population

[0212] After removal of plasma, leukocytes were washed twice with Hanks Balanced Salt Solution ("HBSS") and resuspended in a plasma equivalent volume of fetal bovine serum (heat inactivated at 56° C. for 30 min.). A 0.1 ml volume of the cell preparation was distributed to each of six (6), 15 ml conical centrifuge tubes. Fluorescein labeled monoclonal antibodies with specificity for the human lymphocyte surface markers CD2 (AMAC, Westbrook, Me.), CD20 (Becton Dickinson) and human IgM (Binding Site, San Diego, Calif.) were added to 3 of the tubes for identifying T and B lymphocyte populations. All reagents had previously tested positive to the corresponding monkey lymphocyte antigens. Chimeric anti-CD20 antibody bound to monkey B cell surface CD20 was measured in the fourth tube using polyclonal goat anti-human IgG coupled with

phycoerythrin (AMAC). This reagent was pre-adsorbed on a monkey Ig-sepharose column to prevent cross-reactivity to monkey Ig, thus allowing specific detection and quantitation of chimeric anti-CD20 antibody bound to cells. A fifth tube included both anti-IgM and anti-human IgG reagents for double stained B cell population. A sixth sample was included with no reagents for determination of autofluorescence. Cells were incubated with fluorescent antibodies for 30 min., washed and fixed with 0.5 ml of fixation buffer (0.15 M NaCl, 1% paraformaldehyde, pH7.4) and analyzed on a Becton Dickinson FACScan™ instrument. Lymphocyte populations were initially identified by forward versus right angle light scatter in a dot-plot bitmap with unlabeled leucocytes. The total lymphocyte population was then isolated by gating out all other events. Subsequent fluorescence measurements reflected only gated lymphocyte specific events.

[0213] Depletion of Peripheral Blood B Lymphocytes

[0214] No observable difference could be ascertained between the efficacy of CHO and SP2/0 produced antibodies in depleting B cells in vivo, although a slight increase in B cell recovery beginning after day 7 for monkeys injected with chimeric anti-CD20 antibodies derived from CHO transfecomas at dosage levels 1.6 mg/kg and 6.4 mg/kg was observed and for the monkey injected with SP2/0 producing antibody at the 0.4 mg/kg dose level. FIGS. 9A, B and C provide the results derived from the chimeric anti-CD20:CHO & SP2/0 study, with FIG. 9A directed to the 0.4 mg/kg dose level; FIG. 9B directed to the 1.6 mg/kg dose level; and FIG. 9C directed to the 6.4 mg/kg dose level.

[0215] As is evident from FIG. 9, there was a dramatic decrease (>95%) in peripheral B cell levels after the therapeutic treatment across all tested dose ranges, and these levels were maintained up to seven (7) days post infusion; after this period, B cell recovery began, and, the time of recovery initiation was independent of dosage levels.

[0216] In the Chimeric Anti-CD20:CHO study, a 10-fold lower antibody dosage concentration (0.01 mg/kg) over a period of four daily injections (0.04 mg/kg total) was utilized. FIG. 10 provides the results of this study. This dosage depleted the peripheral blood B cell population to approximately 50% of normal levels estimated with either the anti-surface IgM or the Leu 16 antibody. The results also indicate that saturation of the CD20 antigen on the B lymphocyte population was not achieved with immunologically active chimeric anti-CD20 antibody at this dose concentration over this period of time for non-human primates; B lymphocytes coated with the antibody were detected in the blood samples during the initial three days following therapeutic treatment. However, by day 7, antibody coated cells were undetectable.

[0217] Table I summarizes the results of single and multiple doses of immunologically active chimeric anti-CD20 antibody on the peripheral blood populations; single dose condition was 6.4 mg/kg; multiple dose condition was 0.4 mg/kg over four (4) consecutive days (these results were derived from the monkeys described above).

TABLE I

PERIPHERAL BLOOD POPULATION FROM C2B8 PRIMATE STUDY				
Monkey	Dose	Day	CD2	Anti-Hu IgG
A	0.4 mg/kg (4 doses)	Prebleed	81.5	—
		0	86.5	0.2
		7	85.5	0.0
		21	93.3	—
B	0.4 mg/kg (4 doses)	Prebleed	81.7	—
		0	94.6	0.1
		7	92.2	0.1
		21	84.9	—
C	6.4 mg/kg (1 dose)	Prebleed	77.7	0.0
		7	85.7	0.1
		21	86.7	—
		28	76.7	—
D	6.4 mg/kg (1 dose)	Prebleed	85.7	0.1
		7	94.7	0.1
		21	85.2	—
		28	85.9	—
Monkey	Anti-Hu IgG + Anti-Hu IgM*	Leu-16	% B Cell Depletion	
A	— 0.3 0.1 — —	9.4 0.0 1.2 2.1	0 97 99 78	
		4.1	66	
		14.8	0	
		0.2 0.1 — — —	0.1 0.1 6.9 8.7	99 99 53 41
B	— — 0.2 0.1 — — —	17.0 0.0 14.7 8.1	0 99 15 62	
		17.0	0	
		0.0	99	
		14.7	15	
C	0.2 0.1 — — — —	14.4 0.0 8.7 14.4 0.0 9.2	0 99 41 0 99 46	
		14.4	0	
		0.0	99	
		8.7	41	
D	0.1 0.2 — — — —	14.4 0.0 9.2 6.7	0 99 46 53	

\*Double staining population which indicates extent of chimeric anti-CD20 coated B cells.

[0218] The data summarized in Table I indicates that depletion of B cells in peripheral blood under conditions of antibody excess occurred rapidly and effectively, regardless of single or multiple dosage levels [text missing or illegible when filed] observed for at least seven (7) days following the last injection, with partial B cell recovery observed by day 21.

[0219] Table II summarizes the effect of immunologically active, chimeric anti-CD20 antibodies on cell populations of lymph nodes using the treatment regimen of Table I (4 daily doses of 0.4 mg/kg, 1 dose of 6.4 mg/kg); comparative values for normal lymph nodes (control monkey, axillary and inguinal) and normal bone marrow (two monkeys) are also provided.

TABLE II

CELL POPULATIONS OF LYMPH NODES				
Monkey	Dose	Day	CD2	Anti-Hu IgM
A	0.4 mg/kg (4 doses)	7	66.9	—
		14	76.9	19.6
		28	61.6	19.7

TABLE II-continued

CELL POPULATIONS OF LYMPH NODES				
Monkey	Anti-Hu IgG + Anti-Hu IgM	Leu-16	% B Lymphocyte Depletion	
A	0.4 mg/kg (4 doses)	7.4 0.8 — 29.9	40.1 22.6 26.0 52.2	1 44 36 0
		0.7 — — 0.7	14.5 14.6 13 64	64 64 13 47
		22.3 1.1 — 12.5	35.2 23.9 21.4 19.7	41 47 51 51
		0.2 — — 0.2	8.7 12.9	78 68
CD2	Anti-Hu IgG + Anti-Hu IgM	Leu-16	% B Lymphocyte Depletion	
Normal Lymph Nodes				
Control 1				
Axillary	55.4	25.0	41.4	NA
Inguinal	52.1	31.2	39.5	NA
Normal Bone Marrow				
Control 2	65.3	19.0	11.4	NA
Control 3	29.8	28.0	16.6	NA

[0220] The results of Table II evidence effective depletion of B lymphocytes for both treatment regimens. Table II further indicates that for the non-human primates, complete saturation of the B cells in the lymphatic tissue with immunologically active, chimeric anti-CD20 antibody was not achieved; additionally, antibody coated cells were observed seven (7) days after treatment, followed by a marked depletion of lymph node B cells, observed on day 14.

[0221] Based upon this data, the single High Dosage Chimeric Anti-CD20 study referenced above was conducted, principally with an eye toward pharmacology/toxicology determination. Ie this study was conducted to evaluate any toxicity associated with the administration of the chimeric antibody, as well as the efficacy of B cell depletion from peripheral blood lymph nodes and bone marrow. Additionally, because the data of Table II indicates that for that study, the majority of lymph node B cells were depleted between 7 and 14 days following treatment, a weekly dosing regimen might evidence more efficacious results. Table III summarizes the results of the High Dosage Chimeric Anti-CD20 study.

TABLE III

Monkey	CELL POPULATIONS OF LYMPH NODES AND BONE MARROW				
	CD2	CD20 <sup>a</sup>	mIgM + anti-C2B8 <sup>b</sup>	C2B8 <sup>c</sup>	Day <sup>d</sup>
<u>Lymphocyte Populations (%)</u>					
<u>Inguinal Lymph Node</u>					
E	90.0	5.3	4.8	6.5	22
F	91.0	6.3	5.6	6.3	22
G	89.9	5.0	3.7	5.8	36
H	85.4	12.3	1.7	1.8	36
<u>Bone Marrow</u>					
E	46.7	4.3	2.6	2.8	22
F	41.8	3.0	2.1	2.2	22
G	35.3	0.8	1.4	1.4	36
H	25.6	4.4	4.3	4.4	36

<sup>a</sup>Indicates population stained with Leu 16.<sup>b</sup>Indicates double staining population, positive for surface IgM cells and chimeric antibody coated cells.<sup>c</sup>Indicates total population staining for chimeric antibody including double staining surface IgM positive cells and single staining (surface IgM negative) cells.<sup>d</sup>Days after injection of final 16.8 mg/kg dose.

**[0222]** Both animals evaluated at 22 days post treatment cessation contained less than 5% B-cells, as compared to 40% in control lymph nodes (see, Table II, supra). Similarly, in the bone marrow of animals treated with chimeric anti-CD20 antibody, the levels of CD20 positive cells were less than 3% as compared to 11-15% in the normal animals (see, Table II, supra). In the animals evaluated at 36 days post treatment cessation, one of the animals (H) had approximately 12% B cells in the lymph node and 4.4% B cells in bone marrow, while the other (G) had approximately 5% B cells in the lymph node and 0.8% in the bone marrow—the data is indicative of significant B cell depletion.

**[0223]** The results of Example III.A indicate, inter alia, that low doses of immunologically active, chimeric anti-CD20 leads to long-term peripheral blood B cell depletion in primates. The data also indicates that significant depletion of B cell populations was achieved in peripheral lymph nodes and bone marrow when repetitive high doses of the antibody were administered. Continued follow-up on the test animals has indicated that even with such severe depletion of peripheral B lymphocytes during the first week of treatment, no adverse health effects have been observed. Furthermore, as recovery of B cell population was observed, a conclusion to be drawn is that the pluripotent stem cells of these primates were not adversely affected by the treatment.

#### **[0224] B. Clinical Analysis of C2B8**

##### **[0225] i. Phase I/II Clinical Trial of C2B8: Single Dose Therapy Study**

**[0226]** Fifteen patients having histologically documented relapsed B cell lymphoma have been treated with C2B8 in a Phase I/II Clinical Trial. Each patient received a single dose of C2B8 in a dose-escalating study; there were three patients per dose: 10 mg/m<sup>2</sup>; 50 mg/m<sup>2</sup>; 100 mg/m<sup>2</sup>; 250 mg/m<sup>2</sup> and 500 mg/m<sup>2</sup>. Treatment was by i.v. infusion through an 0.22 micron in-line filter with C2B8 being diluted in a final volume of 250 cc or a maximal concen-

tration of 1 mg/ml of normal saline. Initial rate was 50 cc/hr for the first hour; if no toxicity was seen, dose rate was able to be escalated to a maximum of 200 cc/hr.

**[0227]** Toxicity (as indicated by the clinician) ranged from “none”, to “fever” to “moderate” (two patients) to “severe” (one patient); all patients completed the therapy treatment. Peripheral Blood Lymphocytes were analyzed to determine, inter alia, the impact of C2B8 on T-cells and B-cells. Consistently for all patients, Peripheral Blood B Lymphocytes were depleted after infusion with C2B8 and such depletion was maintained for in excess of two weeks.

**[0228]** One patient (receiving 100 mg<sup>2</sup> of C2B8) evidenced a Partial Response to the C2B8 treatment (reduction of greater than 50% in the sum of the products of the perpendicular diameters of all measurable indicator lesions lasting greater than four weeks, during which no new lesions may appear and no existing lesions may enlarge); at least one other patient (receiving 500 mg/m<sup>2</sup>) evidenced a Minor Response to the C2B8 treatment (reduction of less than 50% but at least 25% in the sum of the products of the two longest perpendicular diameters of all measurable indicator lesions). For presentational efficiency, results of the PBLs are set forth in FIG. 14; data for the patient evidencing a PR is set forth in FIG. 14A; for the patient evidencing an MR, data is set forth in FIG. 14B. In FIG. 14, the following are applicable: ■=Lymphocytes; □=CD3+ cells (T cells); ▨=CD20+cells; ●=CD19+cells; ⊖=Kappa; ▲=lambda; and ▽=C2B8. As evidenced, the B cell markers CD20 and CD19, Kappa and Lambda, were depleted for a period in excess of two weeks; while there was a slight, initial reduction in T-cell counts, these returned to an approximate base-line level in a relatively rapid time-frame.

##### **[0229] U. Phase I/II Clinical Trial of C2B8: Multiple Dose Therapy Study**

**[0230]** Patients having histologically confirmed B cell lymphoma with measurable progressive disease are eligible for this study which is separated into two parts: in Phase I, consisting of a dose escalation to characterize dose limiting toxicities and determination of biologically active tolerated dose level, groups of three patients will receive weekly i.v. infusions of C2B8 for a total of four (4) separate infusions. Cumulative dose at each of the three levels will be as follows: 500 mg/m<sup>2</sup> (125 mg/m<sup>2</sup>/infusion); 1000 mg/m<sup>2</sup> (250 mg/m<sup>2</sup>/infusion); 1500 mg/m<sup>2</sup> (375 mg/m<sup>2</sup>/infusion). A biologically active tolerated dose is defined, and will be determined, as the lowest dose with both tolerable toxicity and adequate activity); in Phase II, additional patients will receive the biologically active tolerated dose with an emphasis on determining the activity of the four doses of C2B8.

#### **[0231] IV. Combination Therapy: C2B8 and Y2B8**

**[0232]** A combination therapeutic approach using C2B8 and Y2B8 was investigated in a mouse xenographic model (nu/nu mice, female, approximately 10 weeks old) utilizing a B cell lymphoblastic tumor (Ramos tumor cells). For comparative purposes, additional mice were also treated with C2B8 and Y2B8.

**[0233]** Ramos tumor cells (ATCC, CRL 1596) were maintained in culture using RPMI-1640 supplemented with 10% fetal calf serum and glutamine at 37° C. and 5% CO<sub>2</sub>. Tumors were initiated in nine female nude mice approxi-

mately 7-10 weeks old by subcutaneous injection of  $1.7 \times 10^6$  Ramos cells in a volume of 0.10 ml (HBSS) using a 1 cc syringe fitted with 25 g needle. All animals were manipulated in a laminar flow hood and all cages, bedding, food and water were autoclaved. Tumor cells were passaged by excising tumors and passing these through a 40 mesh screen; cells were washed twice with 1×HBSS (50 ml) by centrifugation (1300 RPM), resuspended in 1×HBSS to  $10 \times 10^6$  cells/ml, and frozen at  $-70^\circ\text{C}$ . until used.

[0234] For the experimental conditions, cells from several frozen lots were thawed, pelleted by centrifugation (1300 RPM) and washed twice with 1×HBSS. Cells were then resuspended to approximately  $2.0 \times 10^6$  cells/ml. Approximately 9 to 12 mice were injected with 0.10 ml of the cell suspension (s.c.) using a 1 cc syringe fitted with a 25 g needle; injections were made on the animal's left side, approximately mid-region. Tumors developed in approximately two weeks. Tumors were excised and processed as described above. Study mice were injected as described above with  $1.67 \times 10^6$  cells in 0.10 ml HBSS.

[0235] Based on preliminary dosing experiments, it was determined that 200 mg of C2B8 and 100  $\mu\text{Ci}$  of Y2B8 would be utilized for the study. Ninety female nu/nu mice (approximately 10 weeks old) were injected with the tumor cells.

[0236] Approximately ten days later, 24 mice were assigned to four study groups (six mice/group) while attempting to maintain a comparable tumor size distribution in each group (average tumor size, expressed as a product of length×width of the tumor, was approximately  $80 \text{ mm}^2$ ). The following groups were treated as indicated via tail-vein injections using a 100  $\mu\text{l}$  Hamilton syringe fitted with a 25 g needle:

- [0237] A. Normal Saline
- [0238] B. Y2B8 (100  $\mu\text{Ci}$ )
- [0239] C. C2B8 (200  $\mu\text{g}$ ); and
- [0240] D. Y2B8 (100  $\mu\text{Ci}$ )+C2B8 (200  $\mu\text{g}$ ).

[0241] Groups tested with C2B8 were given a second C2B8 injection (200  $\mu\text{g}/\text{mouse}$ ) seven days after the initial injection. Tumor measurements were made every two or three days using a caliper.

[0242] Preparation of treatment materials were in accordance with the following protocols:

[0243] A. Preparation of Y2B8

[0244] Yttrium-[90] chloride (6 mCi) was transformed to a polypropylene tube and adjusted to pH 4.1-4.4 using metal free 2M sodium acetate. 2B8-MX-DTPA (0.3 mg in normal saline; see above for preparation of 2B8-MX-DTPA) was added and gently mixed by vortexing. After 15 min. incubation, the reaction was quenched by adding 0.05× volume 20 mM EDTA and 0.05× volume 2M sodium acetate. Radioactivity concentration was determined by diluting 5.0  $\mu\text{l}$  of the reaction mixture in 2.5 ml 1×PBS containing 76 m/ml HSA and 1 nM DTPA ("formulation buffer"); counting was accomplished by adding 10.0  $\mu\text{l}$  to 20 ml of Ecolume™ scintillation cocktail. The remainder of the reactive mixture was added to 3.0 ml formulation buffer, sterile filtered and stored at  $2\text{--}8^\circ\text{C}$ . until used. Specific activity (14 mCi/mg at time of injection) was calculated using the radioactivity

concentration and the calculated protein concentration based upon the amount of antibody added to the reaction mixture. Protein-associated radioactivity was determined using instant thin-layer chromatography. Radioincorporation was 95%. Y2B8 was diluted in formulation buffer immediately before use and sterile-filtered (final radioactivity concentration was 1.0 mCi/ml).

[0245] B. Preparation of C2B8

[0246] C2B8 was prepared as described above. C2B8 was provided as a sterile reagent in normal saline at 5.0 mg/ml. Prior to injection, the C2B8 was diluted in normal saline to 2.0 mg/ml and sterile filtered.

[0247] C. Results

[0248] Following treatment, tumor size was expressed as a product of length and width, and measurements were taken on the days indicated in **FIG. 11** (Y2B8 vs. Saline); **FIG. 12** (C2B8 vs. Saline); and **FIG. 13** (Y2B8+C2B8 vs. Saline). Standard error was also determined.

[0249] As indicated in **FIG. 13**, the combination of Y2B8 and C2B8 exhibited tumoricidal effects comparable to the effects evidenced by either Y2B8 or C2B8.

[0250] V. Alternative Therapy Strategies

[0251] Alternative therapeutic strategies recognized in view of the foregoing examples are evident. One such strategy employs the use of a therapeutic dose of C2B8 followed within about one week with a combination of either 2B8 and radioabeled 2B8 (eg Y2B8); or 2B8, C2B8 and, eg Y2B8; or C2B8 and, eg Y2B8. An additional strategy is utilization of radiolabeled C2B8—such a strategy allows for utilization of the benefits of the immunologically active portion of C2B8 plus those benefits associated with a radiolabel. Preferred radiolabels include yttrium-90 given the larger circulating half-life of C2B8 versus the murine antibody 2B8. Because of the ability of C2B8 to deplete B-cells, and the benefits to be derived from the use of a radiolabel, a preferred alternative strategy is to treat the patient with C2B8 (either with a single dose or multiple doses) such that most, if not all, peripheral B cells have been depleted. This would then be followed with the use of radiolabeled 2B8; because of the depletion of peripheral B cells, the radiolabeled 2B8 stands an increased chance of targeting tumor cells. Iodine [131] labeled 2B8 is preferably utilized, given the types of results reported in the literature with this label (see Kaminski). An alternative preference involves the use of a radiolabeled 2B8 (or C2B8) first in an effort to increase the permeability of a tumor, followed by single or multiple treatments with C2B8; the intent of this strategy is to increase the chances of the C2B8 in getting both outside and inside the tumor mass. A further strategy involved the use of chemotherapeutic agent in combination with C2B8. These strategies include so-called "staggered" treatments, ie, treatment with chemotherapeutic agent, followed by treatment with C2B8, followed by a repetition of this protocol. Alternatively, initial treatment with a single or multiple doses of C2B8, thereafter followed with chemotherapeutic treatment, is viable. Preferred chemotherapeutic agents include, but are not limited to: cyclophosphamide; doxorubicin; vincristine; and prednisone. See Armitage, J. O. et al., *Cancer* 50: 1695 (1982), incorporated herein by reference.

[0252] The foregoing alternative therapy strategies are not intended to be limiting, but rather are presented as being representative.

[0253] VI. Deposit Information

[0254] Anti-CD20 in TCAE 8 (transformed in *E. coli* for purposes of deposit) was deposited with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Md., 20852, under the provisions of the Budapest Treaty for the International Recognition of the Deposit of

Microorganisms for the Purpose of Patent Procedure ("Budapest Treaty"). The microorganism was tested by the ATCC on Nov. 9, 1992, and determined to be viable on that date. The ATCC has assigned this microorganism for the following ATCC deposit number: ATCC 69119 (anti-CD20 in TCAE 8). Hybridoma 2B8 was deposited with the ATCC on Jun. 22, 1993 under the provisions of the Budapest Treaty. The viability of the culture was determined on Jun. 25, 1993 and the ATCC has assigned this hybridoma the following ATCC deposit number: HB 11388.

---

SEQUENCE LISTING

```

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1
<211> LENGTH: 8540
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: vector

<400> SEQUENCE: 1

gacgtcgccg ccgctctagg cctccaaaaa agcctcctca ctacttctgg aataagctcag      60
aggccgaggg ggcctcgcc tctgcataaa taaaaaaaaat tagtcagcca tgcatggggc      120
ggagaatggg cggaaactggg cggaggttagg ggcgggatgg gccggatgg gggccggact      180
atggttgctg actaatttagat atgcattgtt tgcataacttc tgcctgtgg ggagcctggg      240
gactttccac acctgggttc tgactaatgg agatgcattgc tttgcataact tctgcctgt      300
ggggagcctg gggactttcc acaccctaaac tgacacacat tccacagaat taattccct      360
atttatataat agtaatcaat tacgggtca ttagttcata gcccataatgg gtagttccgc      420
gttacataac ttacggtaaa tggcccgct ggctgaccgc ccaacgaccc ccggccattg      480
acgtcaataa tgacgtatgt tcccatatgtt acgcaatag ggactttcca ttgacgtcaa      540
tgggtggact atttacggta aactgcccac ttggcagtttac atcaagtgtt tcatatgcca      600
agtacgcccc ctatttgcgtt caatgcgggaaatggcccg cctggcattt tgcccgatgt      660
atgacccat gggactttcc tacttggcgtt tacatctacg tattatgttcat cgctattacc      720
atgggtgttc ggttttggca gtacatcaat gggcgtggat agcgttttgc ctcacgggg      780
tttccaagtc tccacccat tgacgtcaat gggagttgtt tttggcatttca aaatcaacgg      840
gactttccaa aatgtcgtaa caactccggcc ccattgacgc aaatggccgg taggcgtgt      900
cggtgggagg tctatataag cagagctggg tacgtgaaacc gtcagatcgc ctggagacgc      960
catcacatgtt ctctcaccat gagggtcccc gctcagatcc tggggctctt gctgtctgg      1020
ctccccaggtt cacatgttca tggtaaccat gttggaaatca aacgtacggt ggctgcacca      1080
tctgtcttca tcttcccgcc atctgtatgtt cagttgaaat ctggactgc ctctgttgc      1140
tgcctgtca ataacttca tcccagatgtt gccaatgtt acgtggaaatgg ggataacgcc      1200
ctccaaatcggtt gtaactccca ggagagtgcc acagagcagg acagaacggg cagcacatcc      1260
agcctcagca gcaccctgac gctgagggaa gcaatgtt acgtggaaatgg ggataacgcc      1320
tgcgtgttca cccatcggtt cctgagatcc tgggttccaa agagttcaaa caggggagag      1380
tggtaatttca agatccgttca acgggttacca actacccatgtt ctggattcgtt gacaacatgc      1440

```

**-continued**


---

ggccgtgata	tctacgtatg	atcagcctcg	actgtgcctt	ctagttgcca	gccatctgtt	1500	
gtttgc	ccccctgtcc	ttcccttgc	ctggaaagg	tgccactccac	tgtccttcc	1560	
taataaaatg	aggaaatgc	atcgcat	ctgagtaggt	gtcatttat	tctgggggt	1620	
ggggtggggc	aggacagca	gggggaggat	tggaaagaca	atagcaggca	tgctgggat	1680	
gcggtgggct	ctatggaacc	agctgggct	cgacagctat	gccaagtacg	ccccctattt	1740	
acgtcaatga	cggtaatgg	cccgccctggc	attatgccca	gtacatgacc	ttatggact	1800	
ttcctacttg	gcagta	atc	tacgtattag	tcatcgat	taccatgg	atgcgg	1860
ggcagta	catatggcgt	ggatagcgg	ttgactc	acg	gggat	ttcca	1920
ccattgacgt	caatgggagt	ttgtttggc	acaaaatca	acgggactt	ccaaaatgtc	1980	
gtacaactc	cggcc	catttgc	acgcaatgg	g	gggtt	ggcg	2040
taagcagagc	tgggtacgtc	ctcacattca	gtgatcagca	ctgaacacag	acc	cg	2100
atgggttgg	gcctcat	ttt	gtcgctgt	ctac	cg	ctc	2160
aaggccc	cgttcc	cctggc	accc	tc	cc	atgg	2220
gcctgg	gc	ctgg	tca	gg	actt	tc	2280
ggcgc	cc	cc	ctgt	cc	actt	cc	2340
tcc	c	cc	atc	cc	act	cc	2400
aacgt	gatc	aca	ggcc	ca	atc	ttgt	2460
gacaaaactc	acacatgccc	accgtgccc	gcac	ctga	ac	tc	2520
ttc	cc	cc	aaaacc	ca	agg	acacc	2580
tgc	gtt	gg	tg	ac	gt	ttcaca	2640
ggc	gtt	gg	tg	ca	at	ctgt	2700
cgt	gtt	ca	gg	tc	act	cc	2760
tgcaagg	tct	cc	aa	cc	at	cc	2820
ggc	ca	cc	cc	cc	at	cc	2880
aaccagg	tct	cc	cc	cc	at	cc	2940
tgg	gag	ca	gg	ac	ac	cc	3000
gac	gg	cc	cc	cc	cc	cc	3060
aa	cg	cc	cc	cc	cc	cc	3120
ct	cc	cc	cc	cc	cc	cc	3180
tc	cc	cc	cc	cc	cc	cc	3240
cc	cc	cc	cc	cc	cc	cc	3300
cc	cc	cc	cc	cc	cc	cc	3360
ct	at	at	at	at	at	at	3420
gg	at	at	at	at	at	at	3480
gg	at	at	at	at	at	at	3540
aa	tt	tt	tt	tt	tt	tt	3600
aa	ttt	ttt	ttt	ttt	ttt	ttt	3660
gact	cct	taa	gg	gg	gg	gg	3720

---

## -continued

---

gcctgattcc	gtagagccac	accttggtaa	ggccaaatct	gctcacacag	gatagagagg	3780
gcaggagcca	gggcagagca	tataaggta	ggtaggatca	gttgctcctc	acatttgctt	3840
ctgacatagt	tgtgttggga	gcttggatag	cttggacagc	tcagggctgc	gatttcgcgc	3900
caaacttgac	ggcaatccta	gctgtgaaggc	tggttaggatt	ttatccccgc	tgccatcatg	3960
gttcgaccat	tgaactgcat	cgtcgccgtg	tcccaaata	tggggatgg	caagaacgga	4020
gacctaccct	ggcctccgct	caggaacgag	ttcaagtact	tccaaagaat	gaccacaacc	4080
tcttcagtgg	aaggtaaaca	gaatctggtg	attatggta	ggaaaacctg	gttctccatt	4140
cctgagaaca	atcgacccctt	aaaggacaga	attaatatag	ttctcagttag	agaactcaaa	4200
gaaccaccac	gaggagctca	ttttcttgcc	aaaagtttg	atgatgcctt	aagacttatt	4260
gaacaaccgg	aattggcaag	taaagtagac	atggtttgg	tagtcggagg	cagttctgtt	4320
taccaggaag	ccatgaatca	accaggccac	cttagactct	ttgtgacaag	gatcatgcag	4380
gaatttgaaa	gtgacacgtt	tttcccagaa	attgattttgg	ggaaatataa	acttctccca	4440
gaataccccag	gcgtcctctc	tgaggtccag	gaggaaaaag	gcatcaagta	taagttgaa	4500
gtctacgaga	agaaaagacta	acaggaagat	gcttcaagt	tctctgcctc	cctcctaaag	4560
tcatgcattt	ttataagacc	atgggacttt	tgctggctt	agatcgcct	cgactgtgcc	4620
ttcttagttgc	cagccatctg	ttgtttgcc	ctccccctg	ccttccttga	ccctggaaagg	4680
tgccactccc	actgtccctt	cctaataaaa	tgagggaaatt	gcategcatt	gtctgagtag	4740
gtgtcattct	attctggggg	gtggggtggg	gcaggacagc	aagggggagg	attgggaaga	4800
caatagcagg	catgctgggg	atgcgggtgg	ctctatggaa	ccagctgggg	ctcgagctac	4860
tagcttgct	tctcaatttc	ttatttgcatt	aatgagaaaa	aaaggaaaat	taattttaac	4920
accaattcag	tagttgattt	agcaaatgcg	ttgccaaaaaa	ggatgctta	gagacagtg	4980
tctctgcaca	gataaggaca	aacattattc	agagggagta	cccagagctg	agactcctaa	5040
gccagtggat	ggcacagcat	tctagggaga	aatatgctt	tcatcaccga	agcctgattc	5100
cgttagagcca	caccttggta	agggccaatc	tgctcacaca	ggatagagag	ggcaggagcc	5160
agggcagagc	atataaggtt	agtttaggatc	agttgctcct	cacatttgct	tctgacatag	5220
ttgtgttggg	agcttggatc	gatcccttat	ggttgaacaa	gatggattgc	acgcaggttc	5280
tccggccgct	tgggtggaga	ggctattcgg	ctatgactgg	gcacaacaga	caatcggtct	5340
ctctgatgcc	gccgtgttcc	ggctgtcagc	gcagggggcgc	ccggttcttt	ttgtcaagac	5400
cgacctgtcc	ggtgccctga	atgaactgca	ggacgaggca	gcgcggctat	cgtggctggc	5460
cacgacgggc	gttccttgcg	cagctgtct	cgacgttgc	actgaagcg	gaagggactg	5520
gctgctatttgc	ggcgaagtgc	cggggcagga	tctcctgtca	tctcaccccttgc	ctcctgcccga	5580
gaaagtatcc	atcatggctg	atgcaatgcg	cgggctgcat	acgcttgcatt	cggtctacctg	5640
cccatcgac	caccaagcga	aacatcgcat	cgagcgagca	cgtactcgg	tggaaaggccgg	5700
tcttgcgtat	caggatgatc	tggacgaaga	gcatcagggg	ctcgcccaag	ccgaactgtt	5760
cggcaggctc	aaggcgcgc	tgcccgacgg	cgaggatctc	gtcgtgaccc	atggcgatgc	5820
ctgcttgccg	aatatcatgg	tggaaaatgg	ccgcttttct	ggattcatcg	actgtggccg	5880
gctgggtgttgc	gcggaccgct	atcaggacat	agcggttgc	acccggtata	ttgtctgaaga	5940
gcttggccggc	gaatgggctg	accgcttccct	cgtgcttac	ggtatcgccg	ctcccgattc	6000

**-continued**


---

gcagcgcatac	gccttctatac	gccttcttga	cgagttcttc	tgagcgggac	tctggggttc	6060
gaaatgaccg	accaagcgcac	gccccaaacct	ccatcacagag	atttcgattc	caccggccgc	6120
ttctatgaaa	ggttgggctt	cggaatcggt	ttccgggacg	ccggctggat	gatcccccag	6180
cgcggggatc	tcatgctgga	gttcttcgc	caccccaact	tgtttattgc	agcttataat	6240
gtttacaaat	aaagcaatag	catcacaaat	ttcacaaata	aagcattttt	ttcactgcat	6300
tctagttgt	gtttgtccaa	actcatcaat	ctatcttac	atgtctggat	cgcggccgcg	6360
atcccgtcga	gagcttggcg	taatcatgg	catagctgtt	tcctgtgtga	aattgttata	6420
cgctcacaat	tccacacaac	atacgagccg	gaagcataaa	gtgttaagcc	tggggtgcc	6480
aatgagtgag	ctaactcaca	ttaattgcgt	tgcgctact	gcccgttac	cagtcggaa	6540
acctgtcgt	ccagctgcat	taatgaatcg	gccaacgcgc	ggggagaggc	gttttgcgt	6600
tttggcgctc	ttccgcttcc	tcgctactg	actcgctgcg	ctcggctgtt	cggctgcggc	6660
gagcggatc	agtcactca	aaggcgttac	tacggttatc	cacagaatca	ggggataacg	6720
cagggaaagaa	catgtgagca	aaaggccagc	aaaaggccag	gaaccgtaaa	aggccgcgt	6780
tgctggcggtt	tttccatagg	ctccggcccc	ctgacgagca	tcacaaaaat	cgacgctcaa	6840
gtcagaggtg	gcgaaacccg	acaggactat	aaagatacca	ggcggttccc	cctggaaagct	6900
ccctcgtcgc	ctctccctgtt	ccgaccctgc	cgcttacccg	atacctgtcc	gcctttctcc	6960
cttcgggaag	cgtggcgctt	tctcaatgct	cacgctgttag	gtatctcagt	tcgggttagg	7020
tcgttcgctc	caagctggc	tgtgtgcacg	aaccccccgt	tcagccgcac	cgctgcgcct	7080
tatccggtaa	ctatcgctt	gagtccaaacc	cggttaagaca	cgacttacg	ccactggcag	7140
cagccactgg	taacaggatt	agcagagcga	ggtatgttag	cggtgctaca	gagtttga	7200
agtgggtggc	taactacggc	tacactagaa	ggacagtatt	tggtatctgc	gctctgctga	7260
agccagttac	cttcggaaaa	agagttggta	gctcttgc	cggcaaacaa	accaccgctg	7320
gtagcgggtt	ttttttgtt	tgcaaggcgc	agattacgcg	cagaaaaaaa	ggatctcaag	7380
aaagatcctt	gatctttct	acggggctc	acgctcagtg	gaacgaaaac	tcacgttaag	7440
ggattttgtt	catgagatta	tcaaaaagg	tcttcacca	gatcctttta	aattaaaaat	7500
gaagttttaa	atcaatctaa	agtatataat	agtaaacttg	gtctgacagt	taccaatgct	7560
taatcagtga	ggcacctatc	tcagcgatct	gtcttattcg	ttcatccata	gttgcctgac	7620
tccccgtcgt	gtagataact	acgatacggg	agggcttacc	atctggcccc	agtgcgtcaa	7680
tgataaccgcg	agacccacgc	tcaccggctc	cagatttatc	agcaataaac	cagccagccg	7740
gaaggggccga	gcgcagaagt	ggtcctgca	ctttatccgc	ctccatccag	tctattaatt	7800
gttgcgggaa	agctagagta	agtagttcgc	cagttaatag	tttgcgaac	gttggcc	7860
ttgtctacagg	catcgtggtg	tcacgctcgt	cgtttggat	ggcttcattc	agctccggtt	7920
cccaacgcgc	aaggcgagtt	acatgatccc	ccatgttg	caaaaaagcg	gttagctcct	7980
tcggtcctcc	gatcgttg	agaagtaatg	tggccgcagt	gttatcactc	atggttatgg	8040
cagcaactgca	taattctt	actgtcatgc	catccgtaag	atgcattttct	gtgactgg	8100
agtactcaac	caagtcatc	tgagaatagt	gtatgcggcg	accgagttgc	tcttgc	8160
cgtcaatacg	ggataatacc	gcccacata	gcagaactt	aaaagtgc	atcatggaa	8220
aacgttcttc	ggggcgaaaa	ctctcaagga	tcttaccgct	gtttagatcc	agttcgatgt	8280

**-continued**


---

aaccttactcg	tgcacccaaac	tgtatcttcag	catctttac	tttcaccagc	gtttctgggt	8340
gagcaaaaac	aggaaggcaa	aatgcccaca	aaaaggaaat	aagggcgaca	cggaatgtt	8400
gaatactcat	actcttcctt	tttcaatattt	attgaaggat	ttatcagggt	tattgtctca	8460
tgagcggata	cataatttcaa	tgtatattga	aaaataaaca	aatagggtt	ccgcgcacat	8520
ttccccgaaa	agtgccacct					8540

<210> SEQ\_ID NO 2  
 <211> LENGTH: 9209  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: vector with chimeric antibody sequence

<400> SEQUENCE: 2						
gacgtcgcgg	ccgctctagg	cctccaaaaa	agcctctca	ctacttcgg	aatagctcag	60
aggccgaggc	ggectcggcc	tctgcataaa	taaaaaaaat	tagtcagcca	tgcatagggc	120
ggagaatggg	cggaaactggg	cggagttagg	ggcgggatgg	gcggagttag	ggcgggact	180
atggttctg	actaattttag	atgcatacttc	tgcctctgg	ggagcctgg		240
gactttccac	acctgggttc	tgactaattt	agatgcatac	tttgcatact	tctgcctgt	300
ggggagctg	gggactttcc	acaccctaa	tgacacacat	tccacagaat	taattccct	360
agttattaaat	agtaatcaat	tacggggtca	ttagttcata	gcccatatata	ggagttccgc	420
gttacataac	ttacggtaaa	tggccgcct	ggctgaccgc	ccaacgaccc	ccgcccattt	480
acgtcaataa	tgacgtatgt	tccatagta	acgccaatag	ggactttcca	ttgacgtcaa	540
tgggtggact	atttacggta	aactgcccac	ttggcagttac	atcaagtgt	tcatatgc	600
agtacgcccc	ctatttgcgt	aatggccgt	aaatggcccg	cctggcatta	tgcccaagtac	660
atgaccttat	gggactttcc	tacttggcag	tacatctac	tatttgtcat	cgctattacc	720
atgggtatgc	ggttttggca	gtacatcaat	gggcgtggat	accgggttga	ctcacgcgga	780
tttccaagtc	tccacccat	tgacgtcaat	gggagtttgt	tttggcacc	aaatcaacgg	840
gactttccaa	aatgtcgtaa	caactccgccc	ccattgacgc	aaatggccgg	taggcgtgt	900
cggtgggagg	tctatataag	cagagctggg	tacgtgaacc	gtcagatcgc	ctggagacgc	960
catcacagat	ctctcaactat	ggattttcag	gtgcagat	ttagtcat	cgctattacc	1020
gcttcagtca	taatgtccag	aggacaaattt	gttctctccc	agtctccac	aatcctgtct	1080
gcatttccag	gggagaaggt	cacaatgact	tgcaggccca	gctcaagtgt	aagtacatc	1140
cactggttcc	agcagaagcc	aggatccccc	cccaaaccct	ggattttatgc	cacatccaa	1200
ctggcttctg	gagtccctgt	tgccttca	ggcagttgggt	ctgggacttc	ttactctctc	1260
acaatcagca	gagtggaggc	tgaagatgt	gccacttatt	actgcccac	gtggactagt	1320
aacccaccca	cgttcggagg	ggggaccaag	ctggaaatca	aacgtacgg	ggctgcacca	1380
tctgtcttca	tcttccgc	atctgtatgag	cagttgaaat	ctggaaactgc	ctctgttgt	1440
tgcctgctga	ataacttcta	tcccagagag	gccaaagtac	agtggaaagg	ggataacgccc	1500
ctccaatcg	gttaactccca	ggagagtg	acagagcagg	acagcaagga	cagcacat	1560
agcctcagca	gcacccctgac	gtcgacaa	gcagactac	agaaacacaa	agtctacgccc	1620
tgcgaagtca	cccatcagg	cctgagctcg	cccgatcaca	agagcttcaa	cagggagag	1680

-continued

tggtgaattt agatccgtta acgggtacca actacccatg ctggattcgt gacaacatgc 1740  
ggccgtata tctacgtatg atcagccctcg actgtgcctt ctatgtccca gccatctgtt 1800  
gtttggccctt ccccccgtgcc ttcccttgacc ctggaaagggtt ccactccac tgcctttcc 1860  
taataaaatg aggaaattgc atcgcattgt ctgagtaggtt gtcattctat tctgggggtt 1920  
gggggtgggc aggacagcaa gggggaggat tggaaagaca atagcaggca tgctgggat 1980  
gccccgtggct ctatggacc agctgggggtt cgacagctat gccaagtagc ccccttattt 2040  
acgtcaatga cggtaaatgg cccgcctggc attatgccc gatcatgacc ttatggactt 2100  
ttccctacttg gcaagtacate tacgtattag tcatcgctat taccatgggtt atgcgggtttt 2160  
ggcagttacat caatgggcgtt ggatagcgggtt ttgactcaccg gggatttcca agtctccacc 2220  
ccattgtacgtt caatgggagt ttgttttggc accaaaatca acgggactttt cccaaatgtc 2280  
gtaaacaactc cgccccatgtt acgcaaatggt gcggttagggc tgcgtacgtt gaggctata 2340  
taaggcagacg tgggtacgtc ctcacattca gtatcagca ctgaacacag acccgctcgcac 2400  
atgggttggat gcctcatctt gctttccctt gtcgctgtt gtcgtcgat cctgtcccag 2460  
gtacaaactgc agcagccctgg ggctgagatgtt gtgaaggctt gggcctcagt gaagatgtcc 2520  
tgcaaggctt ctggctacac atttaccatgt tacaatatgc actgggtaaa acagacacctt 2580  
ggtcggggcc tggaaatggat tggagctattt tattccggaa atgggtatac ttcctacaat 2640  
cagaagttca aaggcaaggc cacattgactt gcaagacaat cctccagcac agcctacatgt 2700  
cagctcagca gcctgacatc tgaggactctt gcggtctattt actgtgtcaag atcgcattac 2760  
tacggcgggtt actggtaactt caatgtctgg ggcgcaggga ccacggtcac cgtctctgca 2820  
gtctgacccatc agggccccc ggtttccccc ctggcacccctt cctccaaagag cacctctggg 2880  
ggcacacagcg ccctgggctt cctggtaag gactacttcc ccgaaccgggtt gacgggtcg 2940  
tggaaactca ggcgcctgac cagcggcgtt cacacccctt cggctgtcctt acagtctca 3000  
ggactctactt ccctcagcag cgtggtaacc gttccctcca gcaacttggg caccctgacc 3060  
tacatctgca acgtgaatca caagccccagc aacaccaagg tggacaagaa agcagagcccc 3120  
aaatcttggat acaaaaactca cacatgccc ccgtgcccag cacctgaactt cctgggggg 3180  
ccgtcagtctt tccttccccc cccaaaaccc aaggacaccc tcatgtatcc cccggacccctt 3240  
gagggtcacat gctgtgggtt ggacgtgagc cacgaagacc ctgaggtaa gtcaacttgg 3300  
tacgtggaccc gctgtggaggtt gcaataatgcc aagacaaaggc cgcggggggg gcaactacaac 3360  
agcacgtacc gttgtggtaacc cgtcctcacc gtcctgcacc aggactggctt gaatggcaag 3420  
gagttacaatgtt gcaaggcttc caacaaaggcc ctcccaagccccc ccacgtggaa aaccatctcc 3480  
aaagccaaag ggcagccccg agaaccacag gtgtacaccc tggcccccattt ccgggatgag 3540  
ctgaccaaga accagggtcag cctgacccctgc ctggtaaaatg gtttctatcc cagcgcacatc 3600  
gccccgtggat gggagagcaatggggcagccg gagaacaactt acaagaccac gctccctgtt 3660  
ctggactcccg acggctccctt cttccctctac agcaaggctt ccgtggacaa gagcgggtttt 3720  
cagcagggggacgttcccttc atgtccctgtt atgcattggg ctctgcacaa ccactacac 3780  
cagaagaccc tctccctgttccgggtttttt tgaggatccg ttaacggtaa ccaactacat 3840  
agactggattt cgtgacaaca tggccggccgtt atatctacgtt atgatcagcc tcgactgtcc 3900  
cttcttagttt ccagccatctt gttttttttcc cttcccccgtt gctttccctt accccctggaaat 3960

## -continued

---

gtgccactcc cactgtcctt tcctaataaa atgagggaaat tgcacatcgcat tgtctgagta	4020
ggtgtcattc tattctgggg ggtgggggtgg ggcaggacag caagggggag gattggaaag	4080
acaatagcag gcacatgcgtgg gatgcgggtgg gctctatgga accagctggg gctcgacacgc	4140
gctggatctc ccgatccccca gctttgcctt tcaatttctt atttgcataa tgagaaaaaa	4200
aggaaaatta attttaaacac caattcagta gttgatttag caaatgcgtt gccaaaaagg	4260
atgcctttaga gacagtgttc tctgcacaga taaggacaaa cattattcg agggagtagcc	4320
cagagctgag actcctaagc cagttagtgg cacagcattc tagggagaaa tatgcttgc	4380
atcacccgaag cctgattccg tagagccaca ccttggtaag ggcacatctg ctcacacagg	4440
atagagaggg caggagccag ggcagagcat ataaggtag ctaggatcatg ttgctctca	4500
catttgcttc tgacatagtt gtgttgggg ctggatagc ttggacagct cagggtgc	4560
atttcgcgcc aaacttgacg gcaatccatcg cgtgaaggct ggttaggattt tatccccgt	4620
gccatcatgg ttcgaccatt gaactgcac gtcgcgggtgt cccaaaatat ggggattggc	4680
aagaacggag acctaccctg gcctccgctc aggaacgagt tcaagtactt ccaaagaatg	4740
accacaacccctt cttcagtgga aggttaaacag aatctggta ttatggtag gaaaacctgg	4800
ttctccattc ctgagaagaa tcgaccttta aaggacagaa ttaatatagt tctcagtaga	4860
gaactcaaag aaccaccacg aggagctcat tttcttgcca aaagtttggta tgatgcctta	4920
agacttattg aacaaccgga attggcaagt aaagtagaca tggtttggat agtcggaggc	4980
agttctgttt accaggaagc catgaatcaa ccaggccacc ttagactctt tgtgacaagg	5040
atcatgcagg aatttggaaag tgacacgttt ttcccagaaa ttgattttgg gaaatataaa	5100
cttctcccgaa aatacccgagg cgtccctctc gaggtccagg agggaaaagg catcaagtt	5160
aagtttgaag tctacgagaa gaaagactaa caggaagatg ctttcaagtt ctctgcctcc	5220
ctcctaaagc tatgcatttt tataagacca tggactttt gctggcttta gatcaggctc	5280
gactgtgcct tctagttgcc agccatctgt tggggccccc tccccgtgc cttccttgac	5340
cctggaaagg gccactccca ctgcctttc ctaataaaat gaggaaatttgcatttgc	5400
tctgagtagg tgcattcta ttctgggggg tgggggtgggg caggacagca agggggagga	5460
ttgggaagac aatagcaggc atgctggggta tgcgggtggc tctatggaaac cagctggggc	5520
tcgagctact agctttgtttt ctcattttt tatttgacata atgagaaaaa aaggaaaattt	5580
aattttaca ccaatttgcgtt agttgattga gcaaatgcgt tgccaaaaag gatgcttttag	5640
agacagtgtt ctctgcacag ataaggacaa acattttca gaggagttac ccagagctga	5700
gactcctaag ccagttagtg gcacagcattt ctagggagaa atatgcgttgc catcaccgaa	5760
gcctgattcc gttagagccac accttggtaa gggccaaatct gctcacacag gatagagagg	5820
gcaggagcca gggcagagca tataaggtaa ggttaggatca gttgctctc acatttgc	5880
ctgacatagt tgggttggga gcttggatcg atccctatg gttgaacaag atggattgca	5940
cgcaggttcc cccggccgtt gggggggag gctattcggc tatgactggg cacaacagac	6000
aatcggttgc tctgatgcgg ccgtgttccg gctgtcagcg cagggggcc cgggtttttt	6060
tgtcaagacc gacctgtccg gtgcctgaa tgaactgcag gacgaggcag cgcggctatc	6120
gtggctggcc acgacggcg ttccttgcgc agctgtgc tgcgttgca ctgaagcggg	6180
aagggactgg ctgttattgg gcgaaatgcgc gggggcaggat ctctgtcat ctcacccgtc	6240

## -continued

---

tcctgccgag	aaagtatcca	tcatggctga	tgcaatgcgg	cggtgcata	cgttgatcc	6300
ggctacctgc	ccattcgacc	accaagcgaa	acatcgcatc	gagcgagcac	gtactcgat	6360
ggaagccggt	cttgtcgatc	aggatgtatc	ggacgaagag	catcaggggc	tcgcgcacgc	6420
cgaactgttc	gccaggctca	aggcgccatc	gcccgcacggc	gaggatctcg	tcgtgaccca	6480
tggcgatgcc	tgcttgccga	atatcatggt	ggaaaatggc	cgctttctg	gattcatcga	6540
ctgtggccgg	ctgggtgtgg	cggaccgcta	tcagggacata	gcgttggcta	cccggtat	6600
tgctgaagag	cttggcggcg	aatgggctga	ccgcttcctc	gtgcttacg	gtatcgccgc	6660
tcccgatccg	cagcgcatcg	ccttctatcg	ccttcttgac	gagttcttct	gagcgggact	6720
ctggggatcg	aaatgaccga	ccaagcgacg	cccaacctgc	catcacgaga	tttcgattcc	6780
accggccgcct	tctatgaaag	gttgggatcc	ggaatcgtt	tccgggacgc	cggctggatg	6840
atccctccagc	gcggggatct	catgctggag	ttcttcgccc	accccaactt	gtttatcgca	6900
gcttataatg	gttacaaata	aagcaatagc	atcacaattt	tcacaaataa	agcattttt	6960
tcactgcatt	ctagttgtgg	tttgcataaa	ctcatcaatc	tatcttatca	tgtctggatc	7020
gcggccgcga	tcccgtcgag	agcttggcg	aatcatggc	atagctttt	cctgtgtgaa	7080
attgttatcc	gctcacaattt	ccacacaaca	tacgagccgg	aagcataaaag	tgtaaagcct	7140
gggggtgccta	atagtgagc	taactcacat	taattgcgtt	gcgctactg	cccgcttcc	7200
agtcgggaaa	cctgtcgatc	cagctgcatt	aatgaatcgg	ccaaacgcgc	gggagaggcg	7260
gtttgcgtat	tggcgctct	tccgcttcct	cgctcactga	ctcgctgcgc	tcggcgatcc	7320
ggctgcggcg	agcggatata	gctcactcaa	aggcggtat	acgggtatcc	acagaatcag	7380
gggataacgc	aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	7440
aggccgcgtt	gctggcggtt	ttccataggc	tccggccccc	tgacgagcat	cacaaaaatc	7500
gacgctcaag	tcagagggtgg	cggaaacccga	caggactata	aagataccag	gcgtttcccc	7560
ctggaaagctc	cctcgtgcgc	tctcctgttc	cgaccctgccc	gcttaccgg	tacctgtccg	7620
ccttctccc	tccgggaagc	gtggcgctt	ctcaatgctc	acgctgttag	tatctcagtt	7680
cggtgttagt	cgttcgctcc	aagctgggt	gtgtgcacga	accccccgtt	cagcccgacc	7740
gctgcgcctt	atccggtaac	tatcgtcttg	agtccaaaccc	ggtaagacac	gacttatcgc	7800
cactggcagc	agccactggt	aacaggatta	gcagagcgag	gtatgttaggc	ggtgctacag	7860
agttcttcaa	gtgggtggct	aactacggct	acactagaag	gacagtattt	ggtatctcg	7920
ctctgctgaa	gccagttacc	tccggaaaaaa	gagttggtag	ctcttgatcc	ggcaaaacaaa	7980
ccaccgctgg	tagcggtgg	ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	8040
gatctcaaga	agatcccttg	atctttctca	cggggtctga	cgctcgtgg	aacgaaaaact	8100
cacgttaagg	gattttggtc	atgagattat	caaaaaggat	cttcacatcg	atccttttaa	8160
ataaaaatg	aagttttaaa	tcaatctaaa	gtatatatga	gtaaacttgg	tctgacagtt	8220
accaatgctt	aatcagttag	gcacctatct	cagcgatctg	tctatttcgt	tcatccatag	8280
ttgcctgact	ccccgtcgat	tagataacta	cgatacggga	gggcttacca	tctggcccca	8340
gtgctgcaat	gataccgcga	gacccacgct	caccggctcc	agatttatca	gcaataaacc	8400
agccagccgg	aaggcccgag	cgcagaatgt	gtcctgcac	tttatccgccc	tccatccagt	8460
ctattaattt	ttgcccggaa	gctagagtaa	gtatgtcgcc	agttatagt	ttgcgcacg	8520

**-continued**


---

ttgttgccat tgctacaggc atcgtggtgt cacgctcgac	8580
gctccgggttc ccaacgatca aggcgagttt catgatcccc	8640
ttagctccctt cggtccctcg atcgttgtca gaagtaagttt	8700
tggttatggc agcactgcat aattcttta ctgtcatgcc	8760
tgactggtaa gtactcaacc aagtcatattt gagaatagt	8820
cttgcccgcc gtaatacgg gataataccg cgccacatag	8880
tcattggaaa acgttctcg gggcgaaaac tctcaaggat	8940
gttcgatgtt acccactcgat gcacccaaactt gatcttc	9000
tttctgggtg agcaaaaaca ggaaggcaaa atgcccggaaa	9060
gaaatgttg aataactcata ctcttcctt ttcaatattt	9120
attgtctcat gagcggatac atatttgaat gtattttagaa	9180
cgcgcacatt tccccgaaaa gtgccacatt	9209

&lt;210&gt; SEQ ID NO 3

&lt;211&gt; LENGTH: 384

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Mus musculus

&lt;400&gt; SEQUENCE: 3

atggattttc aggtgcagat tatcagtttc ctgctaatca	60
gtgcttcaatca gtgtttttttt cataatgtcc	120
agagggcaaa ttgttcttc ccagtctcca gcaatccctgt	180
ctgcacaaatga cttgcaggcc cagcctgtct gcatctccag	240
ggggagaag cttttttttt tttttttttt tttttttttt	300
tttcgtttca gtggcagtgg gtctggact tttttttttt tcacaatccat	360
ttttttttttt tttttttttt tttttttttt tttttttttt	384

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 128

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Mus musculus

&lt;400&gt; SEQUENCE: 4

Met Asp Phe Gln Val Gln Ile Ile Ser Phe Leu Leu Ile Ser Ala Ser	
1 5 10 15	
Val Ile Met Ser Arg Gly Gln Ile Val Leu Ser Gln Ser Pro Ala Ile	
20 25 30	
Leu Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser	
35 40 45	
Ser Ser Val Ser Tyr Ile His Trp Phe Gln Gln Lys Pro Gly Ser Ser	
50 55 60	
Pro Lys Pro Trp Ile Tyr Ala Thr Ser Asn Leu Ala Ser Gly Val Pro	
65 70 75 80	
Val Arg Phe Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile	
85 90 95	
Ser Arg Val Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp	
100 105 110	
Thr Ser Asn Pro Pro Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys	

---

-continued

---

115

120

125

```

<210> SEQ_ID NO 5
<211> LENGTH: 420
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 5

atgggttgg a gcctcattt gctttcctt gtcgctgtt ctacgcgtt cctgtccca 60
gtacaactgc a gcagcctgg ggctgagctg gtgaagcctg gggcctcagt gaagatgtcc 120
tgcaaggctt ctggctacac atttaccagt tacaatatgc actggtaaa acagacacct 180
ggtcggggcc tggatggat tggagctatt tatccggaa atggtgatac ttccataat 240
cagaagttca aaggcaaggc cacattgact gcagacaaat cctccagcac agcctacatg 300
cagctcagca gcctgacatc tgaggactct gcggctatt actgtcaag atcgacttac 360
tacggcggtg actggactt caatgtctgg ggccgaggga ccacggtcac cgtctctgca 420

```

```

<210> SEQ_ID NO 6
<211> LENGTH: 140
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 6

Met Gly Trp Ser Leu Ile Leu Leu Phe Leu Val Ala Val Ala Thr Arg
1 5 10 15

Val Leu Ser Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys
20 25 30

Ala Gly Ala Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe
35 40 45

Thr Ser Tyr Asn Met His Trp Val Lys Gln Thr Pro Gly Arg Gly Leu
50 55 60

Glu Trp Ile Gly Ala Ile Tyr Pro Gly Asn Gly Asp Thr Ser Tyr Asn
65 70 75 80

Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser
85 90 95

Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val
100 105 110

Tyr Tyr Cys Ala Arg Ser Thr Tyr Tyr Gly Gly Asp Trp Tyr Phe Asn
115 120 125

Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ala
130 135 140

```

```

<210> SEQ_ID NO 7
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: impaired Kozak sequence and restriction enzyme
site

<400> SEQUENCE: 7

gggagcttgg atcgatcctc tatggtt

```

```

<210> SEQ_ID NO 8
<211> LENGTH: 47
<212> TYPE: DNA

```

---

-continued

---

<213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: PCR Primer  
 <400> SEQUENCE: 8

atcacagatc tctcaccatg gatttcagg tgcagattat cagcttc

47

<210> SEQ ID NO 9  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: PCR Primer  
 <400> SEQUENCE: 9

tgcagcatcc gtacgtttga tttccagctt

30

<210> SEQ ID NO 10  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: PCR Primer  
 <400> SEQUENCE: 10

gcggctccca cgcgtgtcct gtcccaag

27

<210> SEQ ID NO 11  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: PCR Primer  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(29)  
 <223> OTHER INFORMATION: s is g or c  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(29)  
 <223> OTHER INFORMATION: m is a or c  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(29)  
 <223> OTHER INFORMATION: r is g or a

<400> SEQUENCE: 11

ggstgttgtg ctagctgmrg agacrgtgta

29

---

What is claimed is:

1. A method of lymphoma therapy in a patient in need of such therapy which comprises:

(a) performing a first treatment wherein an unlabelled anti-CD20 antibody that binds to the CD20 antigen present on lymphoma cells is administered to the patient; and

(b) performing a second treatment which comprises administration of at least one chemotherapeutic agent; wherein said first and second treatments are effected in any order or concurrently.

2. The method of claim 1 wherein said second treatment comprises administration of more than one chemotherapeutic agent.

3. The method of claim 1 wherein said first and second steps are effected concurrently.

4. The method of claim 1 wherein said anti-CD20 antibody is administered prior to said administration of chemotherapeutic agent.

5. The method of claim 1 wherein said anti-CD20 antibody is administered after administration of said chemotherapeutic agent.

6. A method for lymphoma therapy which comprises:

(a) performing a first treatment wherein an unlabelled anti-CD20 antibody that binds to the CD20 antigen present on lymphoma cells is administered to the patient; and

(b) performing a second treatment which comprises administration of at least one chemotherapeutic agent

selected from the group consisting of cyclophosphamide, doxorubicin, vincristine and prednisone; wherein said first and second treatments are effected in any order or concurrently.

7. The method of claim 4 wherein said chemotherapeutic agent is selected from the group consisting of cyclophosphamide, vincristine, prednisone and doxorubicin.

8. The method of claim 6 wherein said chemotherapeutic agent is doxorubicin.

9. The method of claim 6 wherein said chemotherapeutic agent is vincristine.

10. The method of claim 6 wherein said chemotherapeutic agent is cyclophosphamide.

11. The method of claim 6 wherein said chemotherapeutic agent is prednisone.

12. The method of claim 1 wherein said unlabelled anti-CD20 antibody is a chimeric anti-CD20 antibody.

13. The method of claim 2 wherein said unlabelled anti-CD20 antibody is a chimeric anti-CD20 antibody.

14. The method of claim 3 wherein said unlabelled anti-CD20 antibody is a chimeric anti-CD20 antibody.

15. The method of claim 4 wherein said anti-CD20 antibody is a chimeric anti-CD20 antibody.

\* \* \* \* \*