



(86) Date de dépôt PCT/PCT Filing Date: 1996/05/14

(87) Date publication PCT/PCT Publication Date: 1996/11/21

(45) Date de délivrance/Issue Date: 2009/08/04

(85) Entrée phase nationale/National Entry: 1997/11/14

(86) N° demande PCT/PCT Application No.: US 1996/006849

(87) N° publication PCT/PCT Publication No.: 1996/036599

(30) Priorité/Priority: 1995/05/18 (US08/443,992)

(51) Cl.Int./Int.Cl. *C07D 333/28* (2006.01),  
*A61K 31/557* (2006.01), *A61K 9/08* (2006.01),  
*C07D 263/30* (2006.01), *C07D 277/30* (2006.01),  
*C07D 277/32* (2006.01), *C07D 307/54* (2006.01),  
*C07D 307/56* (2006.01), *C07D 307/79* (2006.01),  
*C07D 333/24* (2006.01), *C07D 333/32* (2006.01),  
*C07D 333/60* (2006.01)


(72) Inventeur/Inventor:  
BURK, ROBERT M., US

(73) Propriétaire/Owner:  
ALLERGAN, INC., US

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : UTILISATION DE DERIVES D'ACIDE CYCLOPENTANE HEPTAN(ENE)OIQUE, 2-HETEROARYLALKENYLE  
EN TANT QU'AGENTS THERAPEUTIQUES POUR LE TRAITEMENT DE L'HYPERTENSION OCULAIRE

(54) Title: CYCLOPENTANE HEPTAN(ENE)OIC ACID, 2-HETEROARYLALKENYL DERIVATIVES AS THERAPEUTIC  
AGENTS FOR THE TREATMENT OF OCULAR HYPERTENSION



(57) Abrégé/Abstract:

The invention relates to the use of derivatives of F-type prostaglandins as ocular hypotensives. The PGF derivatives used in accordance with the invention are represented by formula (I), wherein wavy line attachments indicate either the alpha  $\alpha$  or beta  $\beta$

**(57) Abrégé(suite)/Abstract(continued):**

configuration; hatched segments indicate  $\alpha$  configuration; solid triangles are used to indicate  $\beta$  configuration, dashed bonds represent a double bond, or a single bond, R is a substituted heteroaryl radical,  $R^1$  is hydrogen or a lower alkyl radical having up to six carbon atoms, X is selected from the group consisting of  $-OR^1$  and  $-N(R^1)_2$ , Y is = O or represents 2 hydrogen radicals. Certain of the compounds represented by formula (I) comprise another aspect of the present invention.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification<sup>6</sup>:

C07C 405/00, A61K 31/557

A1

(11) International Publication Number:

WO 96/36599

(43) International Publication Date: 21 November 1996 (21.11.96)

(21) International Application Number: PCT/US96/06849

(22) International Filing Date: 14 May 1996 (14.05.96)

(30) Priority Data:  
08/443,992 18 May 1995 (18.05.95) US

(71) Applicant: ALLERGAN [US/US]; 8301 Mars Drive, Waco, TX 76712 (US).

(72) Inventor: BURK, Robert, M.; 1337 Cerritos Drive, Laguna Beach, CA 92651 (US).

(74) Agents: BARAN, Robert, J. et al.; Allergan, Inc., P.O. Box 19534, 2525 Dupont Drive, T-2, 2-E, Irvine, CA 92713-9534 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).



Published

With international search report.

(54) Title: CYCLOPENTANE HEPTAN(ENE)OIC ACID, 2-HETEROARYLALKENYL DERIVATIVES AS THERAPEUTIC AGENTS FOR THE TREATMENT OF OCULAR HYPERTENSION

## (57) Abstract

The invention relates to the use of derivatives of F-type prostaglandins as ocular hypotensives. The PGF derivatives used in accordance with the invention are represented by formula (I), wherein wavy line attachments indicate either the alpha  $\alpha$  or beta  $\beta$  configuration; hatched segments indicate  $\alpha$  configuration; solid triangles are used to indicate  $\beta$  configuration, dashed bonds represent a double bond, or a single bond, R is a substituted heteroaryl radical, R<sup>1</sup> is hydrogen or a lower alkyl radical having up to six carbon atoms, X is selected from the group consisting of -OR<sup>1</sup> and -N(R<sup>1</sup>)<sub>2</sub>, Y is = O or represents 2 hydrogen radicals. Certain of the compounds represented by formula (I) comprise another aspect of the present invention.



CYCLOPENTANE HEPTAN(ENE)OIC ACID, 2-HETEROARYLALKENYL DERIVATIVES  
AS THERAPEUTIC AGENTS FOR THE TREATMENT OF OCULAR HYPERTENSION

5

1. Field of the Invention

1 0 The present invention relates to cyclopentane heptanoic acid, 2  
heteroarylalkenyl derivatives which may be substituted in the 1-  
position with hydroxyl, alkyloxy, amino and amido groups, e.g. 1-OH  
cyclopentane heptanoic acid, 2 heteroarylalkenyl derivatives. These  
compounds are potent ocular hypotensive and are particularly suited  
for the management of glaucoma.

1 5

Background of the Invention

2. Description of Related Art

2 0 Ocular hypotensive agents are useful in the treatment of a  
number of various ocular hypertensive conditions, such as post-  
surgical and post-laser trabeculectomy ocular hypertensive episodes,  
glaucoma, and as presurgical adjuncts.

2 5 Glaucoma is a disease of the eye characterized by increased  
intraocular pressure. On the basis of its etiology, glaucoma has been  
classified as primary or secondary. For example, primary glaucoma in  
adults (congenital glaucoma) may be either open-angle or acute or  
chronic angle-closure. Secondary glaucoma results from pre-existing  
ocular diseases such as uveitis, intraocular tumor or an enlarged  
cataract.

3 0

The underlying causes of primary glaucoma are not yet known.  
The increased intraocular tension is due to the obstruction of aqueous  
humor outflow. In chronic open-angle glaucoma, the anterior  
chamber and its anatomic structures appear normal, but drainage of  
the aqueous humor is impeded. In acute or chronic angle-closure  
3 5 angle-closure glaucoma, the anterior chamber is shallow, the filtration  
angle is narrowed, and the iris may obstruct the trabecular meshwork

2


at the entrance of the canal of Schlemm. Dilation of the pupil may push the root of the iris forward against the angle, and may produce pupillary block and thus precipitate an acute attack. Eyes with narrow anterior chamber angles are predisposed to acute angle-closure glaucoma attacks of various degrees of severity.

Secondary glaucoma is caused by any interference with the flow of aqueous humor from the posterior chamber into the anterior chamber and subsequently, into the canal of Schlemm. Inflammatory disease of the anterior segment may prevent aqueous escape by causing complete posterior synechia in iris bombe, and may plug the drainage channel with exudates. Other common causes are intraocular tumors, enlarged cataracts, central retinal vein occlusion, trauma to the eye, operative procedures and intraocular hemorrhage.

Considering all types together, glaucoma occurs in about 2% of all persons over the age of 40 and may be asymptotic for years before progressing to rapid loss of vision. In cases where surgery is not indicated, topical  $\beta$ -adrenoreceptor antagonists have traditionally been the drugs of choice for treating glaucoma.

Certain eicosanoids and their derivatives have been reported to possess ocular hypotensive activity, and have been recommended for use in glaucoma management. Eicosanoids and derivatives include numerous biologically important compounds such as prostaglandins and their derivatives. Prostaglandins can be described as derivatives of prostanoic acid which have the following structural formula:

25



Various types of prostaglandins are known, depending on the structure and substituents carried on the alicyclic ring of the prostanoic

3

acid skeleton. Further classification is based on the number of unsaturated bonds in the side chain indicated by numerical subscripts after the generic type of prostaglandin [e.g. prostaglandin E<sub>1</sub> (PGE<sub>1</sub>), prostaglandin E<sub>2</sub> (PGE<sub>2</sub>)], and on the configuration of the substituents on the alicyclic ring indicated by  $\alpha$  or  $\beta$  [[e.g. prostaglandin F<sub>2</sub> $\alpha$  (PGF<sub>2</sub> $\alpha$ )].

Prostaglandins were earlier regarded as potent ocular hypertensives, however, evidence accumulated in the last decade shows that some prostaglandins are highly effective ocular hypotensive agents, and are ideally suited for the long-term medical management of glaucoma (see, for example, Bito, L.Z. Biological Protection with Prostaglandins, Cohen, M.M., ed., Boca Raton, Fla, CRC Press Inc., 1985, pp. 231-252; and Bito, L.Z., Applied Pharmacology in the Medical Treatment of Glaucomas Drance, S.M. and Neufeld, A.H. eds., New York, Grune & Stratton, 1984, pp. 477-505. Such prostaglandins include PGF<sub>2</sub> $\alpha$ , PGF<sub>1</sub> $\alpha$ , PGE<sub>2</sub>, and certain lipid-soluble esters, such as C<sub>1</sub> to C<sub>2</sub> alkyl esters, e.g. 1-isopropyl ester, of such compounds.

Although the precise mechanism is not yet known experimental results indicate that the prostaglandin-induced reduction in intraocular pressure results from increased uveoscleral outflow [Nilsson et.al., Invest. Ophthalmol. Vis. Sci. (suppl), 284 (1987)].

The isopropyl ester of PGF<sub>2</sub> $\alpha$  has been shown to have significantly greater hypotensive potency than the parent compound, presumably as a result of its more effective penetration through the cornea. In 1987, this compound was described as "the most potent ocular hypotensive agent ever reported" [see, for example, Bito, L.Z., Arch. Ophthalmol. 105, 1036 (1987), and Siebold et.al., Prodrug 5 3 (1989)].

Whereas prostaglandins appear to be devoid of significant intraocular side effects, ocular surface (conjunctival) hyperemia and foreign-body sensation have been consistently associated with the topical ocular use of such compounds, in particular PGF<sub>2</sub> $\alpha$  and its prodrugs, e.g., its 1-isopropyl ester, in humans. The clinical potentials

4


of prostaglandins in the management of conditions associated with increased ocular pressure, e.g. glaucoma are greatly limited by these side effects.

In a series of co-pending United States patent applications assigned to Allergan, Inc. prostaglandin esters with increased ocular hypotensive activity accompanied with no or substantially reduced side-effects are disclosed. The co-pending U.S. Patent 5,446,041 (filed 10 October 1990), relates to certain 11-acyl-prostaglandins, such as 11-pivaloyl, 11-acetyl, 11-isobutyryl, 11-valeryl, and 11-isovaleryl PGF<sub>2</sub>α.

Similarly, 11,15- 9,15 and 9,11-diesters of prostaglandins, for example 11,15-dipivaloyl PGF<sub>2</sub>α are known to have ocular hypotensive activity. See the co-pending patent applications USSN Nos. 385,645 (filed 07 July 1989, now U.S. Patent 4,994,274), 584,370 (filed 18 September 1990, now U.S. Patent 5,028,624) and 585,284 (filed 18 September 1990, now U.S. Patent 5,034,413).

### Summary of the Invention

The present invention concerns a method of treating ocular hypertension which comprises administering to a mammal having ocular hypertension a therapeutically effective amount of a compound of formula I



wherein the hatched segments represent α bonds, the solid triangle represents a β bond, the wavy segment represents α or β bond, dashed

5

lines represent a double bond or a single bond, R is a substituted heteroaryl radical, R<sup>1</sup> is hydrogen or a lower alkyl radical having up to six carbon atoms, X is selected from the group consisting of -OR<sup>1</sup> and -N(R<sup>1</sup>)<sub>2</sub>, Y is = O or represents 2 hydrogen radicals. In a further aspect, the present invention relates to an ophthalmic solution comprising a therapeutically effective amount of a compound of formula (I), wherein the symbols have the above meanings, or a pharmaceutically acceptable salt thereof, in admixture with a non-toxic, ophthalmically acceptable liquid vehicle, packaged in a container suitable for metered application. In particular, the substituents on the heteroaryl radical may be selected from the group consisting of lower alkyl, e.g. C<sub>1</sub> to C<sub>6</sub> alkyl; halogen, e.g. fluoro, chloro and bromo; trifluoromethyl (CF<sub>3</sub>); COR<sup>1</sup>, e.g. COCH<sub>3</sub>; COCF<sub>3</sub>; SO<sub>2</sub>NR<sup>1</sup>, e.g. SO<sub>2</sub>NH<sub>2</sub>; NO<sub>2</sub>; CN; etc.

In a still further aspect, the present invention relates to a pharmaceutical product, comprising

a container adapted to dispense its contents in a metered form; and

an ophthalmic solution therein, as hereinabove defined.

Finally, certain of the compounds represented by the above formula, disclosed below and utilized in the method of the present invention are novel and unobvious.

#### Brief Description of the Drawing Figures

Figure 1 is a schematic of the chemical synthesis of certain 1-carboxylic acid compounds of the invention specifically disclosed as Example 4(a)-(v) below.

Figure 2 is a schematic of the chemical synthesis of certain  $\delta$ -substituted thienyl 1-carboxylic acid compound of the invention specifically disclosed as Examples 6 and 6(a), below.


Figure 3 is a schematic of the chemical synthesis of certain 1-amido compounds of the invention specifically disclosed as Examples 8(p)-(q), below.

Figure 4 is a schematic of the chemical synthesis of certain -substituted thienyl 1-carboxylic acid compounds.

Figure 5 is a schematic of the chemical synthesis of  $\delta$ -substituted furanyl-1-carboxylic acid compounds specifically disclosed as Example 15.

### Detailed Description of the Invention


The present invention relates to the use of nonacidic cyclopentane heptan(ene)oic acid, 2-heteroaryl alkenyl derivatives as therapeutic agents, e.g. as ocular hypotensives. The compounds used in accordance with the present invention are encompassed by the following structural formula I:




wherein the substituents and symbols are as hereinabove defined. The dotted lines on bonds between carbons 5 and 6 (C-5) and carbons 13 and 14 (C-13) indicate a single or double bond. If two solid lines are used at C-5, or C-13, it indicates a specific configuration for that double bond. Hatched lines used at position C-8, C-9 and C-11 indicate the  $\alpha$  configuration. A triangle at position C-12 represents  $\beta$  orientation

A preferred group of the compounds of the present invention includes compounds that have the following structural formula II:

7




wherein Z is selected from the group consisting of O and S, A is selected from the group consisting of N, -CH, and C,  $R^2$  is selected from the group consisting of hydrogen, halogen, and lower alkyl having from 1 to 6 carbon atoms,  $R^3$  and  $R^4$  are selected from the group consisting of hydrogen, halogen, lower alkyl having from 1 to 6

carbon atoms, or, together with ,  $R^3$  and  $R^4$  forms a condensed aryl ring. Preferably, when X is  $-N(R^1)_2$ , Y is = O.

More preferably, at least one of  $R^2$ ,  $R^3$  or  $R^4$  are independently selected from the group consisting of chloro, bromo and lower alkyl. In one aspect of the invention, at least one of  $R^2$ ,  $R^3$  or  $R^4$  is chloro or bromo, and more preferably at least one of  $R^2$ ,  $R^3$  or  $R^4$  is bromo or at least two of  $R^2$ ,  $R^3$  or  $R^4$  are chloro. In another aspect of this invention, at least one of  $R^2$ ,  $R^3$  or  $R^4$  is ethyl, propyl, or butyl.

Another preferred group includes compounds having the formula III:



In the above formulae, the substituents and symbols are as hereinabove defined and R<sup>5</sup> is hydrogen or methyl.

5 The above compounds of the present invention may be prepared by methods that are known in the art or according to the working examples below. The compounds, below, are especially preferred representative of the compounds of the present invention.

1 0 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2-methyl)-thienyl-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid

7-[3 $\alpha$ ,5 $\alpha$ , -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(4-(2-methyl)-thienyl-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid

1 5 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(5-methyl)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

2 0 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2-chloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(4-bromo)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

2 5 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(5-bromo)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2,5-dichloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

3 0 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(3-chloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

3 5 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-benzothienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid .

9

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-benzofuranyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid

5

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(3-furanyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-furanyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

1 0

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-thiazolyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

1 5

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\beta$ -hydroxy-5-(2-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

2 0

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2,5-dichloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenamide.

2 5

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\beta$ -hydroxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-thienyl)-1E-pentyl)cyclopentyl]-5Z-heptenoic acid.

3 0

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -methoxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

3 5

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenamide.

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\beta$ -hydroxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenamide.

5           A pharmaceutically acceptable salt is any salt which retains the activity of the parent compound and does not impart any deleterious or undesirable effect on the subject to whom it is administered and in the context in which it is administered. Of particular interest are salts formed with inorganic ions, such as sodium, potassium, calcium,  
10           magnesium and zinc.

          Pharmaceutical compositions may be prepared by combining a therapeutically effective amount of at least one compound according to the present invention, or a pharmaceutically acceptable acid addition salt thereof, as an active ingredient, with conventional  
15           ophthalmically acceptable pharmaceutical excipients, and by preparation of unit dosage forms suitable for topical ocular use. The therapeutically efficient amount typically is between about 0.0001 and about 5% (w/v), preferably about 0.001 to about 1.0% (w/v) in liquid formulations.

20           For ophthalmic application, preferably solutions are prepared using a physiological saline solution as a major vehicle. The pH of such ophthalmic solutions should preferably be maintained between 6.5 and 7.2 with an appropriate buffer system. The formulations may also contain conventional, pharmaceutically acceptable preservatives,  
25           stabilizers and surfactants.

          Preferred preservatives that may be used in the pharmaceutical compositions of the present invention include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate and phenylmercuric nitrate. A preferred surfactant is, for  
30           example, Tween 80. Likewise, various preferred vehicles may be used in the ophthalmic preparations of the present invention. These vehicles include, but are not limited to, polyvinyl alcohol, povidone, hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose and purified water.

11

Tonicity adjustors may be added as needed or convenient. They include, but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

5 Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include acetate buffers, citrate buffers, phosphate buffers and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

10 In a similar vein, an ophthalmically acceptable antioxidant for use in the present invention includes, but is not limited to, sodium metabisulfite, sodium thiosulfate, acetylcysteine, butylated hydroxyanisole and butylated hydroxytoluene.

15 Other excipient components which may be included in the ophthalmic preparations are chelating agents. The preferred chelating agent is edentate disodium, although other chelating agents may also be used in place or in conjunction with it.

The ingredients are usually used in the following amounts:

|    | <u>Ingredient</u> | <u>Amount (% w/v)</u>  |
|----|-------------------|------------------------|
| 20 | active ingredient | about 0.001-5          |
|    | preservative      | 0-0.10                 |
|    | vehicle           | 0-40                   |
|    | tonicity adjustor | 1-10                   |
| 25 | buffer            | 0.01-10                |
|    | pH adjustor       | q.s. pH 4.5-7.5        |
|    | antioxidant       | as needed              |
|    | surfactant        | as needed              |
| 30 | purified water    | as needed to make 100% |

The actual dose of the active compounds of the present invention depends on the specific compound, and on the condition to be treated; the selection of the appropriate dose is well within the knowledge of the skilled artisan.

35 The ophthalmic formulations of the present invention are conveniently packaged in forms suitable for metered application, such as in containers equipped with a dropper, to facilitate the application to the eye. Containers suitable for dropwise application are usually made

1 2

of suitable inert, non-toxic plastic material, and generally contain between about 0.5 and about 15 ml solution.

5 The invention is further illustrated by the following non-limiting Examples, which are summarized in the reaction schemes of Figures 1 through 4, wherein the compounds are identified by the same designator in both the Examples and the Figures.

#### Compound 4a

10 7-[3 $\alpha$ ,5 $\alpha$ , -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2-methyl)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid

#### Step 1: Preparation of Enone 2a

15 To a suspension of sodium hydride (36 mg, 1.50 mmol) in THF (4.5 mL) cooled to 0°C was added dimethyl 4-(3-(2-methyl) thienyl)-2-oxo-butylphosphonate (414 mg, 1.50 mmol) in THF (3.0 mL). After 0.25h a solution of the aldehyde 1 (438 mg, 1.00 mmol) in THF (3.0 mL) was added and the reaction was allowed to slowly warm to 23° C over a period of 8 h. The reaction solution was quenched with saturated aqueous NH<sub>4</sub>Cl and extracted w/ EtOAc. The aqueous phase was  
20 made slightly acidic before extraction with EtOAc. The combined organics were washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*. Flash column chromatography (silica gel, 2:1 hex/EtOAc) gave 544 mg (93%) of the enone 2a.

#### 25 Step 2: Preparation of alcohol 3a

Sodium tetrahydriodoborate (35 mg, 0.93 mmol) was added to a solution of the enone 2a (544 mg, 0.93 mmol) in MeOH (4.5 mL) at 0°C. After 2h the solvent was removed *in vacuo* and the residue was stirred with 1N NaOH/EtOAc. The organic portion was separated,  
30 dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*. The 3a -alcohol was separated by flash column chromatography or HPLC (silica gel, 3:1 Hex/EtOAc).

35 Step 3: Preparation of 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2-methyl)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

13

5 A solution of alcohol 3a and pyridinium p-toluene sulfonate (116 mg, 0.462 mmol) in MeOH(4.5mL) was heated at 40°C for 4h. The solvent was removed *in vacuo* and the residue was diluted with EtOAc followed by washing with 1N HCl, saturated aqueous NaHCO<sub>3</sub> and brine. The organic portion was dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*.

10 The residue was diluted with THF (0.78 mL) and lithium hydroxide (0.39mL of a 0.5N solution in H<sub>2</sub>O, 0.194 mmol) was added. After 16h the reaction was acidified with 1N HCl and extracted with EtOAc. The organic portion was washed with brine, dried (MgSO<sub>4</sub>), and concentrated *in vacuo* to give 37.6 mg of the free acid 4a.

The title compound was identified by the following NMR spectrum.

15 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 12.1(brs, 1H), 6.98 (d,J=5.1Hz), 6.81(d, J=5.1Hz, 1H), 5.30-5.64(m, 4H), 4.92(brs, 3H), 4.07-4.17 (m, 2H), 3.89-3.93(m, 1H), 2.55-2.59 (m,2H), 2.35(δ,3H), 2.07-2.33(m,10H), 1.42-1.86(m,4H).

20 By methods described for compound 4a, steps 1 through 3, the following compounds were prepared. (The compounds below are also identified by their NMR spectra.)

Compound 4b

25 7-[3α,5α-Dihydroxy-2-(3α-hydroxy-5-(2-(5-methyl)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid .

According to the procedures described above for 4a, the use of dimethyl 4-(2-(5-methyl)thienyl)-2-oxo-butylphosphonate afforded 26 mg of free acid 4b.

30 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 12.1(brs, 1H), 6.56 (d,J=3.4Hz, 1H), 6.54(d, J=3.4Hz, 1H), 5.34-5.64(m, 4H), 4.70(brs, 3H), 4.13-4.20 (m, 2H), 3.91-3.93(m, 1H), 2.82 (t, J= 7.7 Hz, 2H), 2.42(δ,3H), 2.05-2.38(m,11H), 1.44-1.96(m,5H).

35

14

## Compound 4c

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(5-methyl)thienyl)-1E-pentynyl)cyclopentyl]-5Z-heptenoic acid .

5 According to the procedures described above for 4a, the use of dimethyl 4-(3-(2-methyl)thienyl)-2-oxo-butyl phosphonate afforded 25 mg of free acid 4c.

10 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  12.0(brs, 1H), 6.67 (s, 1H), 6.60(s, 1H), 5.34-5.62(m, 4H), 4.42(brs, 3H), 4.10-4.17 (m, 2H), 3.89-3.93(m, 1H), 2.57-2.60(m, 2H), 2.44( $\delta$ ,3H), 2.09-2.36(m, 8H), 1.44-1.87(m,6H).

## Compound 4d

15 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2-chloro)thienyl)-1E-pentynyl)cyclopentyl]-5Z-heptenoic acid.

According to the procedures described above for 4a, the use of dimethyl 4-(3-(2-chloro)thienyl)-2-oxo-butylphosphonate afforded 25 mg of free acid 4d.

20 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  12.0(brs, 1H), 6.99 (d, J=5.7Hz), 6.78(d, J=5.7Hz,1H), 5.29-5.60(m, 4H), 4.02-4.11 (m, 2H), 3.84-3.87(m, 1H), 3.37(brs, 3H), 2.56-2.63(m, 2H), 2.01-2.32(m, 8H), 1.38-1.83(m,7H).

## 25 Compound 4e

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(4-bromo)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

30 According to the procedures described above for 4a, the use of dimethyl 4-(2-(4-bromo)thienyl)-2-oxo-butylphosphonate afforded 10 mg of free acid 4e.

35 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  12.01(brs, 1H), 6.95 (s, 1H), 6.65(s, 1H), 5.24-5.53(m, 4H), 3.99-4.08 (m, 2H), 3.76-3.80(m, 1H), 2.70-2.79(m, 2H), 2.44(s,3H), 2.09-2.36(m, 8H), 1.44-1.87(m,6H).

## 15

## Compound 4f

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(5-bromo)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

5

According to the procedures described above for 4a, the use of dimethyl 4-(2-(5-bromo)thienyl)-2-oxo-butylphosphonate afforded 50 mg of free acid 4f.

10

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  12.0(brs, 1H), 6.80 (d,  $J=3.6$  Hz, 1H), 6.51(d,  $J=3.9$  Hz, 1H), 5.33-5.55(m, 4H), 4.05-4.12(m, 2H), 3.82-3.88(m, 1H), 2.75-2.81(m, 2H), 1.38-2.28 (m,14H).

## Compound 4g

15

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2,5-dichloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

According to the procedures described above for 4a, the use of dimethyl 4-(3-(2,5-dichloro)thienyl)-2-oxo-butylphosphonate afforded 18 mg of free acid 4g.

20

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  12.01(brs, 1H), 6.64 (s, 1H), 5.27-5.56(m, 4H), 4.05-4.15(m, 2H), 3.85-3.92(m, 1H),1.42-2.31(m,18H).

25

## Compound 4h

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(3-chloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

According to the procedures described above for 4a, the use of dimethyl 4-(2-(3-chloro)thienyl)-2-oxo-butylphosphonate afforded 10 mg of free acid 4h.

30

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  12.0(brs, 1H), 7.13 (d,  $J=5.4$  Hz, 1H), 6.75(d,  $J=5.4$  Hz, 1H), 5.19-5.46(m, 4H), 3.96-3.98(m, 2H), 3.69-3.76(m, 1H), 2.72-2.75(m, 2H), 1.35-2.28(m, 17H).

35

## Compound 4i

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-benzothienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid .

5

According to the procedures described above for 4a, the use of dimethyl 4-(2-benzothienyl)-2-oxo-butylphosphonate afforded 22 mg of free acid 4i.

1 0  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  11.8(brs, 1H), 7.73(d,  $J=7.7$  Hz, 1H), 7.63(d,  $J=6.9$  Hz, 1H), 7.23-7.31(m, 2H), 7.00(s, 1H), 5.31-5.65(m, 4H), 4.86(brs, 3H), 4.16-4.22 (m, 2H), 3.89-3.93(m, 1H), 2.96 (t,  $J=7.6$  Hz, 2H) 1.86-2.35(m, 10H), 1.44-1.78(m, 4H).

1 5

## Compound 4j

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-benzofuranyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

2 0 According to the procedures described above for 4a, the use of dimethyl 4-(2-benzofuranyl)-2-oxo-butylphosphonate afforded 30.5 mg of free acid 4j.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  12.1(brs, 1H), 7.37-7.47 (m, 2H), 7.15-7.19(m, 2H), 6.38(s, 1H), 5.30-5.66 (m, 4H), 5.04 (brs, 3H), 4.10-4.20(m, 2H), 3.86-3.94(m, 1H), 2.84(t,  $J=7.6$  Hz, 2H) 1.90-2.33(m, 10H), 1.38-1.78(m, 4H).

2 5

## Compound 4k

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-furanyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

3 0

According to the procedures described above for 4a, the use of dimethyl 4-(3-furanyl)-2-oxo-butylphosphonate afforded 10.1 mg of free acid 4k.

17

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  12.0(brs, 1H), 7.35 (d,  $J=1.7$  Hz, 1H), 7.23(s, 1H), 6.28 (d,  $J=1.7$  Hz, 1H), 5.34-5.64(m, 4H), 4.15-4.20(m, 2H), 3.90-3.94 (m, 1H), 3.70 (brs, 3H), 2.09-2.52 (m, 12H), 1.40-1.88 (m, 4H).

## 5 Compound 4l

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-furanyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

10 According to the procedures described above for 4a, the use of dimethyl 4-(2-furanyl)-2-oxo-butylphosphonate afforded 33.3 mg of free acid 4l.

15  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  11.8(brs, 1H), 7.29(s, 1H), (d,  $J=3.0$  Hz 1H), 6.26(dd,  $J=3.0$ , 1.8 Hz, 1H), 5.99 (d,  $J=1.8$  Hz, 1H) 5.34-5.64 (m, 4H), 4.82 (brs, 3H), 4.11-4.17 (m, 2H), 3.89-3.93 (m, 1H), 2.69 (t,  $J=8.4$  Hz, 2H), 2.05-2.34 (m, 10H), 1.40-1.94(m, 4H).

## Compound 4m

20 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-thiazolyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

According to the procedures described above for 4a, the use of dimethyl 4-(2-thiazolyl)-2-oxo-butylphosphonate afforded 32.2 mg of free acid 4m.

25

$^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.47 (d,  $J=3.3$  Hz, 1H), 7.23(d,  $J=3.3$  Hz, 1H), 3.86-3.93 (m, 2H), 3.60-3.65 (m, 1H), 2.88-2.95 (m, 2H), 1.75-2.20 (m, 10H), 1.22-1.44(m, 4H).

30

## Compound 4n

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

18

According to the procedures described above for 4a, the use of dimethyl 4-(2-thienyl)-2-oxo-butylphosphonate afforded 15.0 mg of free acid 4n.

5  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  11.9(brs, 1H), 7.11, (d,  $J=5.1$  Hz, 1H), 6.92(dd,  $J=5.1, 3.3$  Hz, 1H), 6.00 (d,  $J=3.3$  Hz, 1H) 5.32-5.64 (m, 4H), 4.15-4.19(m, 2H), 3.93-3.97 (m, 1H), 3.61 (brs, 3H), 2.89-2.95 (m, 2H), 2.09-2.35 (m, 8H), 1.46-1.98(m, 6H).

10

Compound 4o

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

15

The 3 $\beta$ -isomer of 3n was isolated from the reaction mixture obtained in step 2 during preparation of 4n and subjected to step 3 to afford 14.3 mg of free acid 4o.

20  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  11.5(brs, 1H), 7.11, (d,  $J=5.1$  Hz 1H), 6.92(dd,  $J=5.1, 3.3$  Hz, 1H), 6.81 (d,  $J=3.3$  Hz, 1H) 5.36-5.64 (m, 4H), 4.62 (brs, 3H), 4.17-4.21 (m, 2H), 3.95-3.98 (m, 1H), 2.90-2.96 (m, 2H), 2.08-2.34 (m, 8H), 1.40-1.98(m, 6H).

Compound 4p

25 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

30

According to the procedures described above for 4a, the use of dimethyl 4-(3-thienyl)-2-oxo-butylphosphonate afforded 9.6 mg of free acid 4p. While this compound is not a substituted heteroaryl derivative within the scope of general Formula I, above, it represents another aspect of this invention in view of its excellent ability to lower intraocular pressure as shown below.

19

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 12.0(brs, 1H), 7.23-7.27(m, 1H), 6.94-6.95(m, 2H), 5.36-5.65(m, 4H), 4.10-4.17(m, 2H), 3.94(brs, 3H), 3.90-3.94(m, 1H), 2.68-2.74(m, 2H), 2.00-2.35(m, 8H), 1.44-1.96(m, 6H).

5 Compound 4q  
7-[3α,5α-Dihydroxy-2-(3β-hydroxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

10 The 3β-isomer of 3p was isolated from the reaction mixture obtained in step 2 during preparation of 4p and subjected to step 3 to afford 36.2 mg of free acid 4q.

15 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 11.9(brs, 1H), 7.23-7.27(m, 1H), 6.94-6.95(m, 2H), 5.32-5.65(m, 4H), 4.72(brs, 3H), 4.12-4.19(m, 2H), 3.93-3.97(m, 1H), 2.69-2.76(m, 2H), 2.07-2.33(m, 8H), 1.39-1.90(m, 6H).

Compound 4r

20 7-[3α,5α-Dihydroxy-2-(3α-hydroxy-5-(2-(5-ethyl)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

According to the procedures described above for 4a, the use of dimethyl 4-(2(5-ethyl)thienyl)2-oxo-butylphosphonate will result in the free acid 4r.

25

Compound 4s

30 7-[3α,5α-Dihydroxy-2-(3α-hydroxy-5-(2-(5-butyl)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

35

According to the procedures described above for 4a, the use of dimethyl 4-(2-(5-butyl)thienyl)2-oxo-butylphosphonate will result in the free acid 4s.

Compound 4t

20

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(5-propyl)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

5           According to the procedures described above for 4a, the use of dimethyl 4-(2-(5-propyl)thienyl)2-oxo-butylphosphonate will result in the free acid 4t.

Compound 4u

10

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(2-(5-methoxy)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

15           According to the procedures described above for 4a, the use of dimethyl 4-(2-(5-methoxy)thienyl)2-oxo-butylphosphonate will result in the free acid 4u.

Compound 4v

20           7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\beta$ -hydroxy-5-(2-thiazolyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

25           The 3 $\beta$  isomer of 3m was isolated from the reaction mixture obtained in Step 2 during preparation of 4m and subjected to Step 3 to afford the free acid 4v.

Compound 4w

30           7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3-hydroxy-5-(2-(3-chloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

Compound 6

35           7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -methoxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

21

Alcohol (400 mg, 0.694 mmol) obtained in step 2 of preparation of 4p was treated with silver triflate (803 mg, 3.12 mmol), 2,6-di-t-butylpyridine (0.89 mL, 3.98 mmol) and iodomethane (0.21 mL, 3.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (11 mL) at 0°C. After 12h the reaction mixture was filtered through celite, concentrated *in vacuo* and purified by flash column chromatography to give the 3 $\alpha$ -methoxy product 5. Further subjection of 5 to the procedures described above in step 3 of preparation of 4a provided 41.2 mg of free acid 6.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  11.6(brs, 1H), 7.24-7.28(m, 2H), 6.93 (d, J=3.3 Hz, 1H), 5.34-5.60 (m, 4H), 4.90 (brs, 3H), 4.20-4.23 (m, 1H), 3.99-4.02(m, 1H), 3.54-3.64 (m, 1H), 3.30 (s, 3H), 2.69 (t, J=7.3 Hz, 2H), 2.07-2.42 (m, 9H), 1.50-2.01 (m, 5H).

Compound 6a

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3-hydroxy-5-(2-(3-chloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenoic acid.

The racemate of the alcohol prepared according to step 2 of preparation 4h is treated according to the procedures described above for 6 and results in the free acid 6a.

Compound 8p

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenamide.

The 3 $\alpha$ -alcohol 3p, isolated from step 2 during preparation of 4p, was deprotected with pyridinium p-toluenesulfonate in MeOH at 45°C for 4h and after the usual work-up gave triol 7p.

A mixture of 7p and ammonium chloride in liquid ammonia was heated to 55°C for 48h in a sealed tube. The tube was recooled to -70°C, vented, and then allowed to warm to room temperature on its own accord. The residue was dissolved in 1:1 EtOAc/H<sub>2</sub>O. The organic

WO 96/36599

PCT/US96/06849

2 2

portion was separated, dried ( $\text{MgSO}_4$ ), filtered and concentrated *in vacuo*. Flash column chromatography (silica gel, 9:1  $\text{CH}_2\text{Cl}_2/\text{MeOH}$ ) gave 10.9 mg of the title compound 8p.

5  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.24-7.27(m, 1H), 6.95-6.96 (m, 2H), 5.76 (brs, 1H), 5.34-5.63 (m, 4H), 4.08-4.19 (m, 2H), 3.94-3.98(m, 1H), 2.95 (brs, 3H), 2.69-2.76 (m, 2H), 2.05-2.39 (m, 8H), 1.48-1.96 (m, 6H).

1 0 Compound 8q  
7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\beta$ -hydroxy-5-(3-thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenamide.

According to the procedures described for preparation of 8p the 3 $\beta$ -alcohol 3g was converted to the title compound 8q.

1 5  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.24-7.27 (m, 1H) 6.95-6.97 (m, 2H), 5.72 (brs, 2H), 5.34-5.66 (m, 4H), 4.08--4.19 (m, 2H), 3.95-3.99 (m, 1H), 3.04 (brs, 1H), 2.70-2.84 (m, 4H), 2.08-2.36 (m, 9H), 1.42-1.89 (m, 5H).

2 0 Compound 8r  
7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2,5-dichloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenamide.

2 5 According to the procedures described for preparation of 8p the alcohol 3r was converted to the title compound 8r.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  6.64 (s, 1H), 5.26-5.68 (m, 6H), 4.07-4.10 (m, 1H), 3.97-4.03 (m, 1H), 3.83-3.86 (m, 1H), 2.50-2.56 (m, 2H), 1.96-2.30(m, 11H), 1.39-1.80(m, 6H).

3 0 Compound 13

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-thienyl)pentyl)cyclopentyl]-5Z-heptenoic acid.

3 5

## 23

## Step 1: Preparation of Alcohol

To a suspension of sodium hydride 271 mg (11.30 mmol) in THF (21 mL) cooled to 0°C was added a solution of dimethyl 4-(3-thienyl)-2-oxo-butylphosphonate (2.96g, 11.30 mmol) in THF (10 mL). After stirring for 0.5h a solution of aldehyde 9 (2.80 g, 10.28 mmol) in THF (10 mL) was added dropwise. The reaction was allowed to warm to room temperature and stirred for a total of 12 h before quenching with saturated aqueous NH<sub>4</sub>Cl. The mixture was extracted with EtOAc and the organic portion was washed with saturated aqueous NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*. The residue was purified by flash column chromatography (silica gel, 1:1 hex/EtOAc) to afford 3.98, (95%) of enone.

Immediately, a solution of the enone (3.98 g, 9.75 mmol) in MeOH (22mL) was cooled to 0°C and sodium tetrahydridoborate (369 mg, 9.75 mmol) was added. After 2h the reaction was quenched with saturated aqueous NH<sub>4</sub>Cl and extracted with EtOAc. The organic portion was washed with brine, dried (MgSO<sub>4</sub>) filtered and concentrated *in vacuo*. Purification by HPLC (Waters Partisil-10, 1:1 hex/EtOAc) afforded 1.30 g (33%) of the  $\alpha$ -alcohol 10.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.97 (d, J=7.2 Hz, 2H), 7.21-7.57, (m, 4H), 6.88 (d, J=4.1 Hz, 2H), 5.54-5.70 (m, 2H), 5.23 (q, J=6.1 Hz, 1H), 5.04 (t, J=6.5 Hz, 1H), 4.10 (q, J=7.1 Hz 1H), 2.45-2.89 (m, 7H), 2.18-2.26 (m, 2H), 1.76-1.84 (m, 2H).

## Step 2: Preparation of Bis-Silyl ether 11

Potassium carbonate (523 mg, 3.78 mmol) was added to a solution of benzoate 10 (1.3g, 3.15 mmol) in MeOH(6.5 mL). After 16h the reaction solvent was removed *in vacuo* and the residue was dissolved in EtOAc/saturated aqueous NH<sub>4</sub>Cl. The organic portion was washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*.

The residue was dissolved in THF (6.5 mL) and triphenylphosphine rhodium chloride (400 mg) was added. The

24

solution was degassed and purged under an atmosphere of hydrogen gas at 40-45 psi. After 16h the reaction was concentrated *in vacuo*, and the residue purified by flash column chromatography (silica gel, 3:1 hex/EtOAc) to afford the apparent saturated diol after evaporation of the solvents.

The apparent diol was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (6.5 mL) and 2,6-lutidine (2.0 mL, 16.5 mmol) was added followed by t-butyltrimethylsilyl triflate (3.0 mL, 13.2 mmol). The reaction was quenched with saturated aqueous NaHCO<sub>3</sub> and washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*. Flash column chromatography (9:1 hex/EtOAc) gave 1.29g (94%) of the bis-TBDMS ether 11.

#### Compound 11

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.22-7.25 (m, 1H), 7.22-7.23 (m, 2H), 4.92-4.98 (m, 1H), 3.90-3.94 (m, 1H), 3.64-3.70 (m, 1H), 2.45-2.82 (m, 5H), 1.95-2.16 (m, 2H), 1.71-1.77 (m, 3H), 1.05-1.51 (m, 4H), 0.88 (s, 9H), 0.85 (s, 9H), 0.03 (s, 9H), 0.02 (s, 3H).

#### Step 3: Preparation of ester 12

Lactone 11 (170 mg, 0.315 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1.0 mL) and cooled to -70°C. Dibal-H (0.47 mL of a 1.0 M solution in CH<sub>2</sub>Cl<sub>2</sub>, 0.47 mmol) was added. After 2h the reaction was quenched with MeOH, allowed to warm to room temperature, and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic portion was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated *in vacuo* to give the lactol as a clear, viscous oil.

To a suspension of (4-carboxybutyl)triphenylphosphonium bromide (558 mg, 1.26 mmol) in THF (2.5mL) was added potassium bis(trimethylsilyl)amide (503 mg, 2.52 mmol) at 0°C. After 0.5 h the solution was cooled to -70°C and a solution of the lactol in THF (2.5

25

mL) was added. The reaction was allowed to warm to room temperature on its own accord, quenched with saturated aqueous  $\text{NH}_4\text{Cl}$ , and extracted with EtOAc. The organic portion was washed with saturated aqueous  $\text{NaHCO}_3$ , brine, dried ( $\text{MgSO}_4$ ), filtered and concentrated *in vacuo*. The residue was diluted with  $\text{Et}_2\text{O}$  and excess diazomethane in  $\text{Et}_2\text{O}$  was added until the reaction solution persisted yellow. Evaporation of the solvent gave 140 mg (70%) of ester 12.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.22-7.24 (m, 1H), 6.90-6.92 (m, 2H), 5.28-5.52 (m, 2H), 4.02-4.06 (m, 1H), 3.95-3.96 (m, 1H), 3.63-3.67 (m, 1H), 3.63 (s, 3H), 2.52-2.70 (m, 2H), 2.00-2.32 (m, 5H), 1.20-1.84 (m, 14H), 0.88 (s, 9H), 0.86 (s, 9H), 0.108-0.055 (m, 12H), (d,  $J=7.2$  Hz, 2H), 7.21-7.57, (m, 4H), 6.88 (d,  $J=4.1$  Hz, 2H), 5.54-5.70 (m, 2H), 5.23 (q,  $J=6.1$  Hz, 1H), 5.04 (+,  $J=6.5$  Hz, 1H), 4.10 (q,  $J=7.1$  Hz, 1H), 2.45-2.89 (m, 7H), 2.18-2.26 (m, 2H), 1.76-1.84 (m, 2H).

#### Step 4: Preparation carboxylic acid 13

To a solution of bis-TBDMS ether 12 (25 mg, 0.040 mmol) in THF (0.24 mL) was added  $\text{Bu}_4\text{NF}$  (0.12 mL of a 1.0M solution in THF, 0.12 mmol). After 16 h the reaction was concentrated *in vacuo* and purified by flash column chromatography (silica gel, 3:1 hex/EtOAc) to yield 13.0 mg (79%) of the triol.

Lithium hydroxide (0.15 mL of a 0.5N solution in  $\text{H}_2\text{O}$ , 0.073 mmol) was added to a solution of the ester (13.0 mg, 0.0316 mmol) in THF (0.3 mL). After 16h the reaction was acidified with 1N HCl and extracted with EtOAc. The organic portion was dried ( $\text{MgSO}_4$ ), filtered and concentrated *in vacuo* to give 7.0 mg (56%) of free acid 13.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  12.0 (brs, 1H), 7.19-7.22 (m, 1H), 6.90-6.92 (m, 2H), 5.31-5.48 (m, 2H), 4.10 (+,  $J=3.9$  Hz, 1H), 3.86-3.88 (m, 1H), 3.59-3.65 (m, 1H), 2.65-2.82 (m, 2H), 1.20-2.30 (m, 21H).

Compound 15

7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3-hydroxy-5-(3-furanyl)pentyl) cyclopentyl]-5Z-heptenoic acid 15.

5 Step 1: Preparation of ketone 14.

1 0 A mixture of the enone (137 mg, 0.245 mmol) obtained in step 2 of preparation of 4k above, Aliquat 336 (34  $\mu$ L, 0.074 mmol), sodium dithionite (384.7 mg, 2.21 mmol) and sodium bicarbonate (371.3 mg, 4.42 mmol) in benzene: H<sub>2</sub>O (1:1, 6.0 mL) was heated to 75°C for 1.5 h. The reaction mixture was allowed to cool to room temperature, was diluted with EtOAc, and was washed with H<sub>2</sub>O and brine. The organic portion was dried (MgSO<sub>4</sub>), filtered and the filtrate was concentrated *in vacuo*. Purification by flash column chromatography (silica gel, 4:1 hex/EtOAc) gave 113.3 mg (83%) of the ketone 14.

1 5

Step 2: Preparation of 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3-hydroxy-5-(3-furanyl)pentyl) cyclopentyl]-5Z-heptenoic acid 15.

2 0 Sodium tetrahydridoborate (14.2 mg, 0.375 mmol) was added to a solution of the ketone (210 mg, 0.375 mmol) in MeOH (3.0 mL) cooled to 0°C. After 30 minutes the reaction was quenched with saturated aqueous ammonium chloride and allowed to warm to room temperature. The mixture was extracted with Et<sub>2</sub>O and the organic portion was dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*.

2 5

The residue was diluted with MeOH (3.0 mL) and pyridinium p-toluene sulfonate (141 mg, 0.562 mmol) was added. After heating to 45°C for 16 h the reaction was concentrated *in vacuo*, diluted with EtOAc and washed with 1 N HCl, saturated aqueous sodium bicarbonate, brine, dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo*. Flash column chromatography (silica gel, 2:1 hex/EtOAc) followed by 100% EtOAc) gave 123 mg (83%) of a mixture of alcohols which were homogenous by TLC.

3 0

3 5 The mixture of alcohols (52.3 mg, 0.132 mmol) was diluted with THF (1.0 mL) and lithium hydroxide (0.53 mL of a 0.5 N solution in

27

H<sub>2</sub>O, 0.265 mmol) was added. After 16 h the reaction was acidified with 1 N HCl and extracted with EtOAc. The organic portion was washed with brine, dried (MgSO<sub>4</sub>) filtered and concentrated *in vacuo* to afford 44.6 mg (89%) of free acid 15.

5

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.34 (d, J= 1.8 Hz, 1H), 7.23 (s, 1H), 6.2 (d, J=1.8 Hz, 1H), 5.30-5.52 (m, 2H), 4.81 (brs, 3H) 4.16 (brs, 1H), 3.95 (brs, 1H), 3.61-3.72 (m, 1H), 2.10-2.64 (m, 7H), 2.17 (s, 3H), 1.34-1.91 (m, 10H).

10

Certain of the above compounds were tested for activity in the various *in vitro* assays described below.

15

Activity at different prostanoid receptors was measured *in vitro* in isolated smooth muscle preparations. FP-activity was measured as contraction of the isolated feline iris sphincter. EP<sub>1</sub>-activity was measured as contraction of the longitudinal smooth muscle of the isolated guinea pig ileum. EP<sub>3</sub>-activity was measured as inhibition of the twitch response induced by electrical field stimulation in the isolated guinea pig vas deferens and as contraction of the longitudinal smooth muscle of the isolated chick ileum. Activity was also measured as relaxation of smooth muscle of isolated rabbit jugular vein a preparation which appears to contain a unique PGF<sub>2</sub>α-sensitive receptor provisionally termed FPVASC. TP-vasoconstrictor activity was measured as contraction of rings of the isolated rat thoracic aorta. Effects on platelets from healthy human donors were measured by incubating platelet-rich plasma with the compounds described herein. Inhibition of aggregation was determined by the ability of the compounds described herein to inhibit platelet aggregation in platelet-rich plasma induced by 20 μM ADP.

20

25

30

35

In addition to stimulating the FP receptor associated with the cat iris, several examples also stimulated the EP<sub>3</sub> receptor. Compounds with agonist activity at EP<sub>3</sub> receptors may also be used for treating gastric or duodenal ulcer by virtue of their cytoprotective and anti-secretory properties. They may also be used as adjunctive therapy in combination with aspirin-like drugs and steroids to limit

gastrointestinal side effects. EP<sub>3</sub> agonists stimulate uterine smooth muscle and may be used to terminate pregnancy in human females. EP<sub>3</sub> agonists are also useful in the cervical ripening process and could be used for inducing labor.

5            Other potential therapeutic applications are in osteoporosis, constipation, renal disorders, sexual dysfunction, baldness, diabetes, cancer and in disorder of immune regulation.

Many examples also have pronounced activity at the FP receptor, provisionally termed FPVASC associated with the vascular endothelium in the rabbit jugular vein preparation. Since such agents would be vasodilators they have potential in hypertension and any disease where tissue blood perfusion is compromised. Such indications include, but are not limited to, systemic hypertension, angina, stroke, retinal vascular diseases, claudication, Raynauds disease, diabetes, and pulmonary hypertension.

The compounds were prepared at the said concentrations in a vehicle comprising 0.1% polysorbate 80 and 10 mM TRIS base. Dogs were treated by administering 25  $\mu$ l to the ocular surface, the contralateral eye received vehicle as a control. Intraocular pressure was measured by applanation pneumatonometry. Dog intraocular pressure was measured immediately before drug administration and at 6 hours thereafter.

25 Compound 4g was examined and showed a pronounced ocular hypotensive effect in dogs.

5 The compounds of the invention may also be useful in the treatment of various pathophysiological diseases including acute myocardial infarction, vascular thrombosis, hypertension, pulmonary hypertension, ischemic heart disease, congestive heart failure, and angina pectoris, in which case the compounds may be administered by any means that effect vasodilation and thereby relieve the symptoms of the disease. For example, administration may be by oral, transdermal, parenteral, subcutaneous, intravenous, 10 intramuscular, intraperitoneal, transdermal, or buccal routes.

The compounds of the invention may be used alone, or in combination with other of the known vasodilator drugs.

15 The compounds of the invention may be formulated into an ointment containing about 0.10 to 10% of the active ingredient in a suitable base of, for example, white petrolatum, mineral oil and petroatum and lanolin alcohol. Other suitable bases will be readily apparent to those skilled in the art.

20 The pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, for example, by means of conventional dissolving or suspending the compounds, which are all either water soluble or suspendable. For administration in the treatment of the other mentioned pathophysiological disorders. The pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed 25 capsules make of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules can contain the active compounds in liquid form that may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are 30 preferably dissolved or suspended in suitable liquids, such as in buffered salt solution. In addition, stabilizers may be added.

35 In addition to being provided in a liquid form, for example in gelatin capsule or other suitable vehicle, the pharmaceutical preparations may contain suitable excipients to facilitate the processing of the active compounds into preparations that can be

5 used pharmaceutically. Thus, pharmaceutical preparations for oral use can be obtained by adhering the solution of the active compounds to a solid support, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

10 Suitable excipients are, in particular, fillers such as sugars, for example lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, as well as inders such as starch, paste using for example, maize starch, wheat starch, rich starchy, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, 15 and/or polyvinyl pyrrolidone. If desired, disintegrating agents may be added such as the above-mentioned starches and also carboxymethyl-starch, crosslinked polyvinyl pyrrolidone, agar, or algenic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, 20 talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings which if desired, are resistant to gastric juices. For this purpose, concentrated sugar solutions may be used, which may optionally containing gum arabic, talc, polyvinyl 25 pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations such as acetylcellulose phthalate or hydroxypropylmethyl-cellulose phthalate, are used. Dye stuffs or 30 pigments may be added to the tables or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

Suitable formulations for intravenous or parenteral administration include aqueous solutions of the active compounds.


SUBSTITUTE PAGE

31

In addition, suspensions of the active compounds as oily injection suspensions may be administered. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, soribitol, and/or dextran. Optionally, the suspension may also contain stabilizers.


## CLAIMS

1. Use of a compound represented by formula I for treating ocular hypertension:




wherein the hatched segments represent  $\alpha$  bonds, the solid triangle represents a  $\beta$  bond, wavy line attachments indicate either the alpha ( $\alpha$ ) or beta ( $\beta$ ) configuration; dashed bonds represent a double bond or a single bond, R is a substituted hetero aryl radical, wherein the substituent is selected from the group consisting of  $C_1$  to  $C_6$  alkyl, trifluoromethyl ( $CF_3$ ),  $COCF_3$ , halogen  $SO_2NR_1$ ,  $SO_2NH_2$ ,  $NO_2$ , and  $CN$ , X is  $N(R^1)_2$ ,  $R^1$  is hydrogen or a lower alkyl radical having from 1 to 6 carbon atoms, Y is  $=O$  or represents 2 hydrogen radicals.


2. The use of Claim 1 wherein said compound is represented by formula II:



wherein the hatched segments represent  $\alpha$  bonds, the solid triangle represents a  $\beta$  bond, wavy line attachments indicate either the alpha ( $\alpha$ ) or beta ( $\beta$ ) configuration; dashed bonds represent a double bond or a single bond,  $R^1$ , X and Y are as defined in claim 1, Z is selected from the group consisting of O and S, A is selected from the group consisting of N, -CH, and C,  $R^2$  is selected from the group consisting of hydrogen, halogen and lower alkyl having from 1 to 6 carbon atoms,  $R^3$  and  $R^4$  are selected from the group consisting of hydrogen, halogen, and lower alkyl having from 1 to 6

carbon atoms, or, together with ,  $R^3$  and  $R^4$  forms a condensed aryl ring.


3. The use of claim 2 wherein said compound is represented by formula III:



wherein the hatched segments represent  $\alpha$  bonds, the solid triangle represents a  $\beta$  bond, wavy line attachments indicate either the alpha ( $\alpha$ ) or beta ( $\beta$ ) configuration; dashed bonds represent a double bond or a single bond, X, Y, A, Z,  $R^2$ ,  $R^3$  and  $R^4$  are as defined in claim 2 and  $R^5$  is hydrogen or methyl.


4. The use of claim 3 wherein X is  $\text{-NH}_2$ .
5. The use of claim 3 wherein Y is  $= \text{O}$ .
6. The use of claim 3 wherein Y is  $= \text{O}$  and X is  $\text{-NH}_2$ .
7. The use of claim 3 wherein Z is S.
8. The use of claim 7 wherein at least one of  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  is selected from the group consisting of halogen and lower alkyl having from 1 to 4 carbon atoms.
9. The use of claim 7 wherein at least one of  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  is selected from the group consisting of chloro and bromo.
10. The use of claim 7 wherein at least one of  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  is chloro.
11. The use of claim 10 wherein at least two of  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  are chloro.
12. The use of claim 3 wherein Y is  $= \text{O}$ , X is  $\text{-NH}_2$  and Z is S.
13. The use of claim 12 wherein at least one of  $\text{R}^2$ ,  $\text{R}^3$  and  $\text{R}^4$  is selected from the group consisting of chloro and bromo.
14. The use of claim 12 wherein at least one of  $\text{R}^2$ ,  $\text{R}^3$  or  $\text{R}^4$  is bromo or at least two of  $\text{R}^2$ ,  $\text{R}^3$  or  $\text{R}^4$  are chloro.

15. The use of claim 14 wherein said compound is 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2,5-dichloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenamide.
16. The use of claim 12 wherein at least one of R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup> is a lower alkyl radical having from 1 to 4 carbon atoms.
17. The use of claim 16 wherein at least one of R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup> is ethyl, propyl or butyl.
18. An ophthalmic solution comprising a compound of Formula I:




or a pharmaceutically acceptable salt thereof, in admixture with a non-toxic, ophthalmically acceptable liquid vehicle, packaged in a container suitable for metered application, wherein the hatched segments represent  $\alpha$  bonds, the solid triangle represents a  $\beta$  bond, wavy line attachments indicate either the alpha ( $\alpha$ ) or beta ( $\beta$ ) configuration; dashed bonds represent a double bond or a single bond, R is a substituted hetero aryl radical, wherein the substituent is selected from the group consisting of C<sub>1</sub> to C<sub>6</sub> alkyl, trifluoromethyl (CF<sub>3</sub>), COCF<sub>3</sub>, halogen SO<sub>2</sub>NR<sub>1</sub>, SO<sub>2</sub>NH<sub>2</sub>, NO<sub>2</sub>, and CN, X is N(R<sup>1</sup>)<sub>2</sub>, R<sup>1</sup> is hydrogen or a lower alkyl radical having from 1 to 6 carbon atoms, Y is =O or represents 2 hydrogen radicals.


19. The ophthalmic solution of claim 18 wherein said compound is a compound of Formula III:



wherein the hatched segments represent  $\alpha$  bonds, the solid triangle represents a  $\beta$  bond, wavy line attachments indicate either the alpha ( $\alpha$ ) or beta ( $\beta$ ) configuration; dashed bonds represent a double bond or a single bond, X, Y and R are as defined in claim 18, Z is selected from the group consisting of O and S, A is selected from the group consisting of N, -CH, and C,  $R^2$  is selected from the group consisting of hydrogen, halogen and lower alkyl having from 1 to 6 carbon atoms,  $R^3$  and  $R^4$  are selected from the group consisting of hydrogen, halogen, and lower alkyl having from 1 to 6

carbon atoms, or, together with ,  $R^3$  and  $R^4$  forms a condensed aryl ring and  $R^5$  is hydrogen or methyl.


20. A pharmaceutical product, comprising a container adapted to dispense the contents of said container in metered form; and an ophthalmic solution in said container comprising a compound of Formula I:




or a pharmaceutically acceptable salt thereof, in admixture with a non-toxic, ophthalmically acceptable liquid vehicle,

wherein the hatched segments represent  $\alpha$  bonds, the solid triangle represents a  $\beta$  bond, wavy line attachments indicate either the alpha ( $\alpha$ ) or beta ( $\beta$ ) configuration; dashed bonds represent a double bond or a single bond, R is a substituted hetero aryl radical, wherein the substituent is selected from the group consisting of  $C_1$  to  $C_6$  alkyl, trifluoromethyl ( $CF_3$ ),  $COCF_3$ , halogen,  $SO_2NR_1$ ,  $SO_2NH_2$ ,  $NO$ , and  $CN$ , X is  $N(R^1)_2$ ,  $R^1$  is hydrogen or a lower alkyl radical having from 1 to 6 carbon atoms, Y is  $=O$  or represents 2 hydrogen radicals.


21. The product of claim 20 wherein said compound is a compound of Formula III:



wherein the hatched segments represent  $\alpha$  bonds, the solid triangle represents a  $\beta$  bond, wavy line attachments indicate either the alpha ( $\alpha$ ) or beta ( $\beta$ ) configuration; dashed bonds represent a double bond or a single bond, X, Y and R are as defined in claim 18, Z is selected from the group consisting of O and S, A is selected from the group consisting of N,  $-\text{CH}$ , and C,  $\text{R}^2$  is selected from the group consisting of hydrogen, halogen and lower alkyl having from 1 to 6 carbon atoms,  $\text{R}^3$  and  $\text{R}^4$  are selected from the group consisting of hydrogen, halogen, and lower alkyl having from 1 to 6


carbon atoms, or, together with ,  $\text{R}^3$  and  $\text{R}^4$  forms a condensed aryl ring and  $\text{R}^5$  is hydrogen or methyl.

22. The compound represented by Formula III:



wherein the hatched segments represent  $\alpha$  bonds, the solid triangle represents a  $\beta$  bond, X and Y are defined in claim 21, Z is selected from the group consisting of O and S, A is selected from the group consisting of N,  $-\text{CH}$  and C,  $\text{R}^2$  is selected from the

group consisting of hydrogen, halogen and lower alkyl having from one to six carbon atoms,  $R^3$  and  $R^4$  are selected from the group consisting of hydrogen, halogen and lower alkyl having from 1 to 6 carbon atoms, or together



23. A compound according to claim 22 wherein Z is S.
24. The compound of claim 23 wherein at least one of  $R^2$ ,  $R^3$  and  $R^4$  is selected from the group consisting of halogen and lower alkyl having from 1 to 4 carbon atoms.
25. The compound of claim 23 wherein at least one of  $R^2$ ,  $R^3$  and  $R^4$  is selected from the group consisting of chloro and bromo.
26. The compound of claim 23 wherein at least one of  $R^2$ ,  $R^3$  and  $R^4$  is chloro.
27. The compound of claim 26 wherein at least two of  $R^2$ ,  $R^3$  and  $R^4$  are chloro.
28. The compound of claim 22 wherein Y is = O, X is  $-NH_2$  and Z is S.
29. The compound of claim 28 wherein at least one of  $R^2$ ,  $R^3$  and  $R^4$  is selected from the group consisting of chloro and bromo.
30. The compound of claim 28 wherein at least one of  $R^2$ ,  $R^3$  or  $R^4$  is bromo or at least two of  $R^2$ ,  $R^3$  or  $R^4$  are chloro.

31. The compound of claim 30 wherein said compound is 7-[3 $\alpha$ ,5 $\alpha$ -Dihydroxy-2-(3 $\alpha$ -hydroxy-5-(3-(2,5-dichloro)thienyl)-1E-pentenyl)cyclopentyl]-5Z-heptenamide.
32. The compound of claim 28 wherein at least one of R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup> is a lower alkyl radical having from 1 to 4 carbon atoms.
33. The compound of claim 32 wherein at least one of R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup> is ethyl, propyl or butyl.
34. The use of claim 1 wherein R<sup>1</sup> is hydrogen.
35. The solution of claim 18, wherein R<sup>1</sup> is hydrogen.
36. The product of claim 20, wherein R<sup>1</sup> is hydrogen.
37. The compound of claim 22, wherein R<sup>5</sup> is hydrogen.

FIGURE

1




FIGURE 2







FIGURE 3

3p  $\alpha$ -OH3q  $\beta$ -OH3r  $\alpha$ -OH

PPTs, MeOH, 40 °C

7p  $\alpha$ -OH7q  $\beta$ -OH7r  $\alpha$ -OHNH<sub>4</sub>Cl  
NH<sub>3</sub>  
55-60 °C8p  $\alpha$ -OH8q  $\beta$ -OH8r  $\alpha$ -OH

FIGURE

4







Figure 5





1


$NaH$ , THF,  $0^\circ C$



2

1.  $NaBH_4$ , MeOH,  $0^\circ C$


2. separation




3

1. PPTs, MeOH,  $40^\circ C$

2. 0.5 N LiOH in THF



4a-q



(I)