

**(12) PATENT
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 200067824 B2
(10) Patent No. 780191**

(54) Title
Heteropolycyclic compounds and their use as metabotropic glutamate receptor antagonists

(51)⁷ International Patent Classification(s)
C07D 413/04 A61K 031/497
A61K 031/4439 A61P 025/00
A61K 031/4725 C07D 401/04

(21) Application No: **200067824** (22) Application Date: **2000.08.18**

(87) WIPO No: **WO01/12627**

(30) Priority Data

(31) Number **60/149464** (32) Date **1999.08.19** (33) Country **US**

(43) Publication Date : **2001.03.13**

(43) Publication Journal Date : **2001.05.17**

(44) Accepted Journal Date : **2005.03.03**

(71) Applicant(s)
NPS Pharmaceuticals, Inc.

(72) Inventor(s)
Bradford C Van Wagenen; Thomas M Stormann; Scott T Moe; Susan M. Sheehan; Donald A McLeod; Daryl L. Smith; Methvin Benjamin Isaac; Abdelmalik Slassi

(74) Agent/Attorney
Griffith Hack,GPO Box 1285K,MELBOURNE VIC 3001

(56) Related Art
US 3647809
US 4022901
WO 1998/017652

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12627 A1(51) International Patent Classification⁷: C07D 413/04,
401/04, A61K 31/4439, 31/4725, 31/497, A61P 25/00SMITH, Daryl, L. [US/US]; 1592 East Parkridge Drive,
Salt Lake City, UT 84121 (US). ISAAC, Methvin, Ben-
jamin [KN/CA]; 1507-10 Markbrook Lane, Etobicoke,
Ontario M9V SE3 (CA). SLASSI, Abdelmalik [CA/CA];
3237 Cairnburne Crescent, Mississauga, Ontario L5N
5G4 (CA).

(21) International Application Number: PCT/US00/22618

(74) Agents: BENT, Stephen, A. et al.; Foley & Lardner, 3000
K. Street, NW, Washington, DC 20007-5109 (US).

(22) International Filing Date: 18 August 2000 (18.08.2000)

(81) Designated States (national): AF, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,
VN, YU, ZA, ZW.

(25) Filing Language: English

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(26) Publication Language: English

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

(30) Priority Data:
60/149,464 19 August 1999 (19.08.1999) US(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:US 60/149,464 (CIP)
Filed on 19 August 1999 (19.08.1999)(71) Applicant (for all designated States except US): NPS
PHARMACEUTICALS, INC. [US/US]; 420 Chipeta
Way, Suite 240, Salt Lake City, UT 84108 (US).(72) Inventors; and
(75) Inventors/Applicants (for US only): VAN WAGENEN,
Bradford, C. [US/US]; 3969 South 3250 East, Salt
Lake City, UT 84127 (US). STORMANN, Thomas, M.
[US/US]; 1327 East Harrison, Salt Lake City, UT 84105
(US). MOE, Scott, T. [US/US]; 6152 South Vinefield
Lane, Salt Lake City, UT 84121 (US). SHEEHAN, Susan,
M. [US/US]; 1803 East Redondo Avenue, Salt Lake City,
UT 84108 (US). MCLEOD, Donald, A. [US/US]; 7740
South Newport Way, Salt Lake City, UT 84121 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/12627 A1

(54) Title: HETEROPOLYCYCLIC COMPOUNDS AND THEIR USE AS METABOTROPIC GLUTAMATE RECEPTOR ANTAGONISTS



metabotropic glutamate receptors, and are useful for treating neurological diseases and disorders.

(57) Abstract: Compound of formula (II) wherein X, Y, and Z are independently selected from the group consisting of N, O, S, C, and CO wherein at least one of X, Y, and Z is a heteroatom; Ar¹ and Ar² are independently selected from the group consisting of a heterocyclic or fused heterocyclic moiety and an aromatic moiety; act as antagonist at

HETEROPOLYCYCLIC COMPOUNDS AND THEIR USE AS MATABOTROPIC GLUTAMATE RECEPTOR ANTAGONISTS

FIELD OF THE INVENTION

The invention provides compounds active at metabotropic glutamate receptors and that are useful for treating neurological and psychiatric diseases and disorders.

BACKGROUND OF THE INVENTION

5 Recent advances in the elucidation of the neurophysiological roles of metabotropic glutamate receptors have established these receptors as promising drug targets in the therapy of acute and chronic neurological and psychiatric disorders and diseases. However, the major challenge to the realization of this promise has been the development of metabotropic glutamate receptor subtype-selective compounds.

10 Glutamate is the major excitatory neurotransmitter in the mammalian central nervous system (CNS). Glutamate produces its effects on central neurons by binding to and thereby activating cell surface receptors. These receptors have been divided into two major classes, the ionotropic and metabotropic glutamate receptors, based on the structural features of the receptor proteins, the means by which the receptors transduce signals into 15 the cell, and pharmacological profiles.

16 The metabotropic glutamate receptors (mGluRs) are G protein-coupled receptors that activate a variety of intracellular second messenger systems following the binding of glutamate. Activation of mGluRs in intact mammalian neurons elicits one or more of the following responses: activation of phospholipase C; increases in phosphoinositide (PI) 20 hydrolysis; intracellular calcium release; activation of phospholipase D; activation or inhibition of adenyl cyclase; increases or decreases in the formation of cyclic adenosine monophosphate (cAMP); activation of guanylyl cyclase; increases in the formation of cyclic guanosine monophosphate (cGMP); activation of phospholipase A₂; increases in

arachidonic acid release; and increases or decreases in the activity of voltage- and ligand-gated ion channels. Schoepp *et al.*, *Trends Pharmacol. Sci.* 14:13 (1993); Schoepp, *Neurochem. Int.* 24:439 (1994); Pin *et al.*, *Neuropharmacology* 34:1 (1995).

Eight distinct mGluR subtypes, termed mGluR1 through mGluR8, have been identified by molecular cloning. *See, for example*, Nakanishi, *Neuron* 13:1031 (1994); Pin *et al.*, *Neuropharmacology* 34:1 (1995); Knopfel *et al.*, *J. Med. Chem.* 38:1417 (1995). Further receptor diversity occurs via expression of alternatively spliced forms of certain mGluR subtypes. Pin *et al.*, *PNAS* 89:10331 (1992); Minakami *et al.*, *BBRC* 199:1136 (1994); Joly *et al.*, *J. Neurosci.* 15:3970 (1995).

Metabotropic glutamate receptor subtypes may be subdivided into three groups, Group I, Group II, and Group III mGluRs, based on amino acid sequence homology, the second messenger systems utilized by the receptors, and by their pharmacological characteristics. Nakanishi, *Neuron* 13:1031 (1994); Pin *et al.*, *Neuropharmacology* 34:1 (1995); Knopfel *et al.*, *J. Med. Chem.* 38:1417 (1995).

Group I mGluRs comprise mGluR1, mGluR5, and their alternatively spliced variants. The binding of agonists to these receptors results in the activation of phospholipase C and the subsequent mobilization of intracellular calcium. Electrophysiological measurements have been used to demonstrate these effects, for example, in *Xenopus* oocytes that express recombinant mGluR1 receptors. *See, for example*, Masu *et al.*, *Nature* 349:760 (1991); Pin *et al.*, *PNAS* 89:10331 (1992). Similar results have been achieved with oocytes expressing recombinant mGluR5 receptors. Abe *et al.*, *J. Biol. Chem.* 267:13361 (1992); Minakami *et al.*, *BBRC* 199:1136 (1994); Joly *et al.*, *J. Neurosci.* 15:3970 (1995). Alternatively, agonist activation of recombinant mGluR1 receptors expressed in Chinese hamster ovary (CHO) cells stimulates PI hydrolysis, cAMP formation, and arachidonic acid release as measured by standard biochemical assays. Aramori *et al.*, *Neuron* 8:757 (1992).

By comparison, the activation of mGluR5 receptors, expressed in CHO cells, stimulates PI hydrolysis and subsequent intracellular calcium transients, but no stimulation of cAMP formation or arachidonic acid release is observed. Abe *et al.*, *J. Biol. Chem.* 267:13361 (1992). However, activation of mGluR5 receptors expressed in LLC-PK1 cells results in PI hydrolysis and increased cAMP formation. Joly *et al.*, *J. Neurosci.* 15:3970 (1995). The agonist potency profile for Group I mGluRs is quisqualate > glutamate = ibotenate > (2S,1'S,2'S)-2-carboxycyclopropyl)glycine (L-CCG-I) > (1S,3R)-1-

aminocyclopentane-1,3-dicarboxylic acid (ACPD). Quisqualate is relatively selective for Group I receptors, as compared to Group II and Group III mGluRs, but it also is a potent activator of ionotropic AMPA receptors. Pin *et al.*, *Neuropharmacology* 34:1, Knopfel *et al.*, *J. Med. Chem.* 38:1417 (1995).

5 The lack of subtype-specific mGluR agonists and antagonists has impeded elucidation of the physiological roles of particular mGluRs, and the mGluR-associated pathophysiological processes that affect the CNS have yet to be defined. However, work with the available non-specific agonists and antagonists has yielded some general insights about the Group I mGluRs as compared to the Group II and Group III mGluRs.

10 Attempts at elucidating the physiological roles of Group I mGluRs suggest that activation of these receptors elicits neuronal excitation. Various studies have demonstrated that ACPD can produce postsynaptic excitation upon application to neurons in the hippocampus, cerebral cortex, cerebellum, and thalamus, as well as other brain regions. Evidence indicates that this excitation is due to direct activation of postsynaptic mGluRs, 15 but it also has been suggested that activation of presynaptic mGluRs occurs, resulting in increased neurotransmitter release. Baskys, *Trends Pharmacol. Sci.* 15:92 (1992); Schoepp, *Neurochem. Int.* 24:439 (1994); Pin *et al.*, *Neuropharmacology* 34:1(1995).

20 Pharmacological experiments implicate Group I mGluRs as the mediators of this excitatory mechanism. The effects of ACPD can be reproduced by low concentrations of quisqualate in the presence of iGluR antagonists. Hu *et al.*, *Brain Res.* 568:339 (1991); Greene *et al.*, *Eur. J. Pharmacol.* 226:279 (1992). Two phenylglycine compounds known to activate mGluR1, namely (S)-3-hydroxyphenylglycine ((S)-3HPG) and (S)-3,5-dihydroxyphenylglycine ((S)-DHPG), also produce excitation. Watkins *et al.*, *Trends Pharmacol. Sci.* 15:33 (1994). In addition, the excitation can be blocked by (S)-4-carboxyphenylglycine ((S)-4CPG), (S)-4-carboxy-3-hydroxyphenylglycine ((S)-4C3HPG), 25 and (+)-alpha-methyl-4-carboxyphenylglycine ((+)-MCPG), compounds known to be mGluR1 antagonists. Eaton *et al.*, *Eur. J. Pharmacol.* 244:195 (1993); Watkins *et al.*, *Trends Pharmacol. Sci.* 15:333 (1994).

30 Metabotropic glutamate receptors have been implicated in a number of normal processes in the mammalian CNS. Activation of mGluRs has been shown to be required for induction of hippocampal long-term potentiation and cerebellar long-term depression. Bashir *et al.*, *Nature* 363:347 (1993); Bortolotto *et al.*, *Nature* 368:740 (1994); Aiba *et al.*, *Cell* 79:365 (1994); Aiba *et al.*, *Cell* 79:377 (1994). A role for mGluR activation in

nociception and analgesia also has been demonstrated. Meller *et al.*, *Neuroreport* 4: 879 (1993). In addition, mGluR activation has been suggested to play a modulatory role in a variety of other normal processes including synaptic transmission, neuronal development, apoptotic neuronal death, synaptic plasticity, spatial learning, olfactory memory, central 5 control of cardiac activity, waking, motor control, and control of the vestibulo-ocular reflex. Generally, see Nakanishi, *Neuron* 13: 1031 (1994); Pin *et al.*, *Neuropharmacology* 34:1; Knopfel *et al.*, *J. Med. Chem.* 38:1417 (1995).

Metabotropic glutamate receptors also have been suggested to play roles in a variety of pathophysiological processes and disease states affecting the CNS. These include stroke, 10 head trauma, anoxic and ischemic injuries, hypoglycemia, epilepsy, and neurodegenerative diseases such as Alzheimer's disease. Schoepp *et al.*, *Trends Pharmacol. Sci.* 14:13 (1993); Cunningham *et al.*, *Life Sci.* 54:135 (1994); Hollman *et al.*, *Ann. Rev. Neurosci.* 17:31 (1994); Pin *et al.*, *Neuropharmacology* 34:1 (1995); Knopfel *et al.*, *J. Med. Chem.* 38:1417 (1995). Much of the pathology in these conditions is thought to be due to 15 excessive glutamate-induced excitation of CNS neurons. Because Group I mGluRs appear to increase glutamate-mediated neuronal excitation via postsynaptic mechanisms and enhanced presynaptic glutamate release, their activation probably contributes to the pathology. Accordingly, selective antagonists of Group I mGluR receptors could be therapeutically beneficial, specifically as neuroprotective agents, analgesics, or 20 anticonvulsants.

Preliminary studies assessing therapeutic potentials with the available mGluR agonists and antagonists have yielded seemingly contradictory results. For example, it has been reported that application of ACPD onto hippocampal neurons leads to seizures and neuronal damage (Sacaan *et al.*, *Neurosci. Lett.* 139:77 (1992); Lipparti *et al.*, *Life Sci.* 52:85 (1993). Other studies indicate, however, that ACPD inhibits epileptiform activity, 25 and also can exhibit neuroprotective properties. Taschenberger *et al.*, *Neuroreport* 3:629 (1992); Sheardown, *Neuroreport* 3:916 (1992); Koh *et al.*, *Proc. Natl. Acad. Sci. USA* 88:9431 (1991); Chiamulera *et al.*, *Eur. J. Pharmacol.* 216:335 (1992); Siliprandi *et al.*, *Eur. J. Pharmacol.* 219:173 (1992); Pizzi *et al.*, *J. Neurochem.* 61:683 (1993).

30 It is likely that these conflicting results are due to the lack of selectivity of ACPD, which causes activation of several different mGluR subtypes. In the studies finding neuronal damage it appears that Group I mGluRs were activated, thereby enhancing undesirable excitatory neurotransmission. In the studies showing neuroprotective effects it

appears that activation of Group II and/or Group III mGluRs occurred, inhibiting presynaptic glutamate release, and diminishing excitatory neurotransmission.

This interpretation is consistent with the observation that (S)-4C3HPG, a Group I mGluR antagonist and Group II mGluR agonist, protects against audiogenic seizures in DBA/2 mice, while the Group II mGluR selective agonists DCG-IV and L-CCG-I protect neurons from NMDA- and KA-induced toxicity. Thomsen *et al.*, *J. Neurochem.* 62:2492 (1994); Bruno *et al.*, *Eur. J. Pharmacol.* 256:109 (1994); Pizzi *et al.*, *J. Neurochem.* 61:683 (1993).

Based on the foregoing, it is clear that currently available mGluR agonists and antagonists have limited value, due to their lack of potency and selectivity. In addition, most currently available compounds are amino acids or amino acid derivatives that have limited bioavailabilities, thereby hampering *in vivo* studies to assess mGluR physiology, pharmacology and their therapeutic potential. Compounds that selectively inhibit activation of metabotropic glutamate receptor Group I subtypes should be useful for treatment of neurological disorders and diseases such as senile dementia, Parkinson's disease, Alzheimer's disease, Huntington's Chorea, pain, migraine headaches, epilepsy, head trauma, anoxic and ischemic injuries, psychiatric disorders such as schizophrenia, depression, and anxiety, ophthalmological disorders such as various retinopathies, for example, diabetic retinopathies, glaucoma, and neurological disorders of a auditory nature such as tinnitus, and neuropathic pain disorders, including neuropathic diseases states such as diabetic neuropathies, chemotherapy induced neuropathies, post-herpetic neuralgia, and trigeminal neuralgia.

Accordingly, a need exists for potent mGluR agonists and antagonists that display a high selectivity for a mGluR subtype, particularly a Group I receptor subtype.

25 SUMMARY OF THE INVENTION

It is an object of the present invention, therefore, to identify metabotropic glutamate receptor-active compounds which exhibit a high degree of potency and selectivity for individual metabotropic glutamate receptor subtypes, and to provide methods of making these compounds.

30 It is a further object of this invention to provide pharmaceutical compositions containing compounds which exhibit a high degree of potency and selectivity for individual

metabotropic glutamate receptor subtypes, and to provide methods of making these pharmaceutical compositions.

It is yet another object of this invention to provide methods of inhibiting activation of an mGluR Group I receptor, and of inhibiting neuronal damage caused by excitatory activation of an mGluR Group I receptor, specifically mGluR5.

It is still another object of the invention to provide methods of treating a disease associated with excitatory activation of an mGluR Group I receptor, specifically mGluR5.

To accomplish these and other objectives, the present invention provides potent antagonists of Group I mGluRs, specifically, mGluR5. These antagonists fall within the general class of compounds represented by the formula I below, although this application now relates to a more confined group of these compounds, as set out in the appended claims. The general class of antagonists are represented by the formular I,

15 $\text{Ar}^1\text{-G-}\text{Ar}^2$


wherein Ar^1 is an optionally substituted heteroaromatic moiety and Ar^2 is an optionally substituted benzene ring. The G moiety is a group that not only covalently binds to the Ar^1 and Ar^2 moieties, and facilitates adoption of the correct spatial orientation of Ar^1 and Ar^2 , but also itself may interact with the protein, to effect receptor binding.

In general sense, G could be selected from the group consisting of – NH-, -S-, -O-, -CO-, -CONH-, -CONHCH₂-, -CH₂CONH-, -CNHNH-, -CNHNHCH₂-, -C=NO-CH₂-, -CH₂NHCH₂-, -CH₂CH₂NH-, -NHCH₂CO-, -NHCH₂CHOH-, -NHCNHNH-, -NHCONH-, cyclopentane, cyclopentadiene, furan, thifuran, pyrrolidine, 25 pyrrole, 2-imidazoline, 3-imidazoline, 4-imidazoline, imidazole, pyrazoline, pyrazolidine, imidazolidine, oxazole, 2-oxazole, thiazole, isoxazole, isothiazole, 1*H*-1,2,4-triazole, 1*H*-1,2,3-triazole, 1,2,4-oxathiazole, 1,3,4-oxathiazole, 1,4,2-dioxazole, 1,4,2-oxathiazole, 1,2,4-oxadiazole, 1,2,4-thiadiazole, 1,2,5-oxadiazole, 1,2,5-thiadiazole, 1,3,4-oxadiazole, 30 1,3,4-thiadiazole, 1*H*-tetrazole, cyclohexane, piperidine, tetrahydropyridine, 1,4-dihydropyridine, pyridine, benzene, tetrahydropyran, 3,4-dihydro-2*H*-pyran, 2*H*-pyran, 4*H*-pyran, tetrahydrothiopyran, 3,4-dihydro-2*H*-thiopyran, 2*H*-thiin, 4*H*-thiopyran, morpholine, thiomorpholine, piperazine, pyridazine, pyrimidine, pyrazine, 1,2,4-triazine, 1,2,3-triazine, 1,3,5-triazine, and 1,2,4,5-tetrazine.

35 This application is now limited to specific groups within this set.

In a general sense, Ar¹ could be selected from the group consisting of phenyl, benzyl, naphthyl, fluorenyl, anthrenyl, indenyl, phenanthrenyl, and benzonaphthetyl, and Ar² could be selected from the group consisting of thiazoyl, furyl, pyranyl, 2H-pyrrolyl, thienyl, pyrroyl, imidazoyl, pyrazoyl, pyridyl, pyrazinyl, pyrimidinyl, 5 pyridazinyl, benzothiazole, benzimidazole, 3H-indolyl, indolyl, indazoyl, purinyl, quinolizinyl, isoquinolyl, quinolyl, phthalizinyl, naphthyridinyl, quinazolinyl, cinnolinyl, isothiazolyl, quinoxalinyl indolizinyl, isoindolyl, benzothienyl, benzofuranyl, isobenzofuranyl, and chromenyl. This application now relates to specific groups within this set.

10 In a general sense compounds of the present invention are within the class represented by formula II:



wherein X, Y, and Z are independently selected from the group consisting of N, O, S, C, and CO wherein at least one of X, Y, and Z is a heteroatom;

20 Ar¹ and Ar² are independently selected from the group consisting of a heterocyclic or fused heterocyclic moiety containing 1 to 4 heteroatoms selected from the group consisting of N, O, and S and an aromatic moiety selected from the group consisting of phenyl, benzyl, 1-naphthyl, 2-naphthyl, fluorenyl, anthrenyl, indenyl, phenanthrenyl, and benzonaphthetyl, wherein the Ar¹ and Ar² moieties are optionally substituted with one or more moieties selected from the group consisting of -F, -Cl, -Br, -I, -OR, -SR, -SOR, -SO₂R, -SO₂NRR', -OCOR, -OCONRR', -NRCOR', -NRCO₂R', -CN, -NO₂, -CO₂R, -CONRR', -C(O)R, -CH(O)R', -CH₂(OR), -R, and -A-(CH₂)_n-NRR'; wherein R or R' is selected from the group consisting of H, CF₃, C₁-C₁₀ alkyl, cycloalkyl, alkyl-aryl, alkyl-heteroaryl, heterocycloalkyl, aryl and where R and R' may combine to form a ring, and A is defined as CH₂, O, NH, S, SO, SO₂ and n is 1, 2, 3, or 4. The heterocyclic or fused heterocyclic moiety preferably is selected from the group consisting of quinolyl, quinazolyl, quinoxalyl, 2-pyrimidyl, 4-pyrimidyl, 5-pyrimidyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, and pyrazyl.

30 The present application is now limited to a more limited class within this general class, as outlined in the appended claims.

35

In a preferred embodiment of the invention, the compound is selected from the group consisting of 3-(2-pyridyl)-5-(3,5-dichlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-methoxyphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-methylphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(1-naphthyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,3-difluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,5-difluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3,5-difluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(2-

5

10

pyridyl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazol, 3-(2-pyridyl)-5-(2,3-dichlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chloro-5-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-fluoro-5-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chloro-5-fluorophenyl)-1,2,4-oxadiazole, 3-(5-chloropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(5-fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(5-fluoropyrid-2-yl)-5-(3-cyano-5-fluorophenyl)-1,2,4-oxadiazole, 3-(3-fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(5-fluoropyrid-2-yl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole, 3-(5-methoxypyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(2-quinolinyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(3-chloro-5-trifluoromethylpyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(5-chloro-2-methoxyphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2-chloro-5-methylthiophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2-bromo-5-methoxyphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,5,6-trifluorophenyl)-1,2,4-oxadiazole, 2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole and 3-(2-pyridyl)-5-(2,5,6-trifluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-nitrophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-bromophenyl)-1,2,4-oxadiazole and pharmaceutically acceptable salts thereof.

In another embodiment of the invention, the compound is selected from the group consisting of 2-(3,5-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-methylphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(1-naphthyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-trifluoromethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2,3-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3,5-dimethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2,3-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-chloro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-chloro-5-fluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(5-chloropyrid-2-yl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(3-fluoropyrid-2-yl)-1,3-oxazole, 2-(3,5-dimethoxyphenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(5-methoxypyrid-2-yl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(2-quinolinyl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(3-chloro-5-trifluoromethylpyrid-2-yl)-1,3-oxazole, 2-(5-chloro-2-methoxyphenyl)-4-(2-pyridyl)-1,3-

oxazole, 2-(2-chloro-5-methylthiophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2-bromo-5-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole and 2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-nitrophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-bromophenyl)-4-(2-pyridyl)-1,3-oxazole and pharmaceutically acceptable salts thereof.

In accordance with another embodiment of the invention, there has been provided a pharmaceutical composition comprising a compound as set forth above in Formula I and formula II, together with a pharmaceutically acceptable diluent or excipient.

In accordance with still another embodiment of the invention, there has been provided a method of making a compound as set forth above. Specifically, compounds of the invention generally can be prepared by formation of the G moiety between two precursor compounds containing suitable Ar¹ and Ar² moieties. When the linker contains an 1,2,4-oxadiazole, the heterocycle may be formed using well known techniques, such as reaction between an amidoxime and an acid chloride, or by the reaction of an amidoxime and an acylimidazole. An illustration of such a transformation is provided in Examples 3 through 6, below.

Amidoximes can be prepared using well known techniques by the reaction of an Ar¹ substituted nitrile with hydroxylamine. An illustration of such a transformation is provided below in Example 1.

In most cases, the precursor Ar² carbonyl chlorides are readily available, or may be prepared using straightforward techniques of organic chemistry. For example, carboxylic acids may be converted into the corresponding acid chlorides by reaction with, for example, thionyl chloride or oxalyl chloride.

In the case where the linker contains a 1,3-oxazole, compounds were prepared from the procedure similar to that given by Kelly *et al.*, *J. Org. Chem.* 61, 4623-4633 (1996). 3,5-Disubstituted-1,3-Oxazoles were prepared by reacting a haloketone with carboxamide in refluxing toluene for 3 days. The resulting mixture was allowed to cool to room temperature, the solvent was removed and the residue was purified.

In accordance with a still further embodiment of the invention, there has been provided a method of inhibiting activation of an mGluR Group I receptor, specifically mGluR5, comprising treating a cell containing said mGluR Group I receptor with an effective amount of a compound as set forth above.

In yet another embodiment of the invention, there has been provided a method of inhibiting neuronal damage caused by excitatory activation of an mGluR Group I receptor, comprising treating neurons with an effective amount of a compound as set forth above.

5 In accordance with a further embodiment of the invention, there has been provided a method of treating a disease or disorder associated with glutamate-induced neuronal damage, or a method of treating a disease or disorder associated with Group I mGluR activation or amenable to therapeutic intervention with a mGluR Group I antagonist, comprising administering to a patient suffering from said disease or disorder an effective amount of a composition as set forth above, wherein said disease or disorder is selected 10 from the group consisting of as senile dementia, Parkinson's disease, Alzheimer's disease, Huntington's Chorea, pain, migraine headaches, epilepsy, head trauma, anoxic and ischemic injuries, psychiatric disorders such as schizophrenia, depression, anxiety, diabetic retinopathies, glaucoma, tinnitus, diabetic neuropathies, chemotherapy induced neuropathies, post-herpetic neuralgia, and trigeminal neuralgia.

15

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments 20 of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Figure 1 shows illustrative compounds of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides compounds that are potent and selective 30 antagonists of mGluR5. The compounds contemplated by the invention are within the class represented by the general formula I:



35 where Ar^1 is an optionally substituted heterocyclic moiety and Ar^2 is an optionally substituted carbocyclic moiety. The G moiety is a group that not only covalently binds to

the Ar¹ and Ar² moieties and facilitates adoption of the correct spatial orientation of Ar¹ and Ar², but may itself interact with the protein to allow receptor binding.

Structure of the Ar¹ and Ar² moieties

The Ar¹ moiety is generally defined as a heterocyclic moiety, and the Ar² moiety is generally defined as a carbocyclic moiety. Ar¹ and Ar² can be monocyclic or fused bicyclic groups. Ar² is preferably defined as an aryl or alkaryl moiety. Ar¹ is preferably defined as a heterocyclic, heteroaryl or heteroarylalkyl moiety. The ring systems encompassed by Ar¹ can contain up to four heteroatoms, independently selected from the group consisting of N, S, and O. When Ar¹ is a heteroaryl ring or ring system, it preferably contains one or two heteroatoms. At least one of the heteroatoms preferably is nitrogen (N). The heterocyclic or fused heterocyclic moiety preferably is selected from the group consisting of quinolyl, quinazolyl, quinoxalyl, 2-pyrimidyl, 4-pyrimidyl, 5-pyrimidyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, and pyrazyl.

Monocyclic Ar¹ groups include, but are not limited to: thiazoyl, furyl, pyranyl, 2H-pyrrolyl, thienyl, pyrrolyl, imidazoyl, pyrazoyl, pyridyl, pyrazinyl, pyrimidinyl, and pyridazinyl moieties. Monocyclic Ar² group include but are not limited to phenyl and benzyl. Fused bicyclic Ar² include, but are not limited to, naphthyl, fluorenyl, anthrenyl, indenyl, phenanthrenyl, and benzonaphthyl. Fused bicyclic Ar¹ groups include, but are not limited to: benzothiazole, benzimidazole, 3H-indolyl, indolyl, indazoyl, purinyl, quinolizinyl, isoquinolyl, quinolyl, phthalizinyl, naphthyridinyl, quinazolinyl, cinnolinyl, isothiazoyl, quinoxalinyl indolizinyl, isoindolyl, benzothienyl, benzofuranyl, isobenzofuranyl, and chromenyl moieties. Ar¹ preferably is a 2-pyridyl moiety. Ar² preferably is a substituted phenyl moiety.

The Ar¹ and Ar² moieties optionally may independently be substituted with one or more moieties selected from the group consisting of halogen, C₁-C₃ alkyl, C₁-C₃ O-alkyl, -OH, -OCF₃, -COOR, -COR, -SOR, -SO₂NRR', -NRR', -CN, -CF₃, -CO-NRR', -A-(CH₂)_n-NRR', wherein A is C, O, N, SO, SO₂, and R and R' are independently selected from the group consisting of C₁-C₃ alkyl, H, cycloalkyl, heterocycloalkyl, aryl, and n is 1, 2, 3, or 4.

Structure of the G moiety

The G moiety is generally made up of 1-14 atoms. G can be independently selected from the group of atoms: C, H, N, O, and S.

The G moiety can thus be made of a non-cyclic moiety. Several examples of these 5 are -NH- (amine), -S- (thioether), -O- (ether), -CO- (ketone), -CONH- (amide), - CONHCH₂-, -CH₂CONH-, -CNHNH- (amidine), -CNHNHCH₂-, -C=NO-CH₂- (methoxime), -CH₂NHCH₂-, -CH₂CH₂NH-, -NHCH₂CO-, -NHCH₂CHOH-, - NHCNHNH- (guanidine), and -NHCONH- (urea), for example.


The atomic arrangement in the G moiety can also be made to form a five-membered 10 ring. Several examples of these are cyclopentane, cyclopentadiene, furan, thifuran, pyrrolidine, pyrrole, 2-imidazoline, 3-imidazoline, 4-imidazoline, imidazole, pyrazoline, pyrazolidine, imidazolidine, oxazole, 2-oxazole, thiazole, isoxazole, isothiazole, 1*H*-1,2,4-triazole, 1*H*-1,2,3-triazole, 1,2,4-oxathiazole, 1,3,4-oxathiazole, 1,4,2-dioxazole, 1,4,2-oxathiazole, 1,2,4-oxadiazole, 1,2,4-thiadiazole, 1,2,5-oxadiazole, 1,2,5-thiadiazole, 15 1,3,4-oxadiazole, 1,3,4-thiadiazole, and 1*H*-tetrazole, for example. The 1,2,4-oxadiazole is most preferred.

The atomic arrangement in the G moiety can also be made to form a six-membered ring. Several examples of these are cyclohexane, piperidine, tetrahydropyridine, 1,4-dihydropyridine, pyridine, benzene, tetrahydropyran, 3,4-dihydro-2*H*-pyran, 2*H*-pyran, 20 4*H*-pyran, tetrahydrothiopyran, 3,4-dihydro-2*H*-thiopyran, 2*H*-thiin, 4*H*-thiopyran, morpholine, thiomorpholine, piperazine, pyridazine, pyrimidine, pyrazine, 1,2,4-triazine, 1,2,3-triazine, 1,3,5-triazine, and 1,2,4,5-tetrazine, for example.

The atomic arrangement in the G moiety can also be made to form a five- or six-membered ring containing one or more carbonyl groups. Several examples of these are 25 2-azetidinone, 1,2-diazetidin-3-one, cyclopentanone, 2-cyclopentenone, 2-pyrrolidinone, 3-pyrrolin-2-one, succinimide, maleimide, 3-pyrazolidinone, 2-imidazolidone, 4-imidazolin-2-one, 2*H*-imidazol-2-one, 4-imidazolinone, 3-pyrazolin-5-one, hydantoin, 1*H*-imidazole-2,5-dione, 2-oxazoline-4-one, 2-oxazolidinone, 3-oxazolin-5-one, 3(2*H*)-isoxazolone, 2,4-oxazolidinedione, 1,2,4-triazoline-3,5-dione, 2,4-dihydro-3*H*-1,2,4-triazol-3-one, 2*H*-30 pyran-2-one, 2(1*H*)-pyridone, 2(1*H*)-pyrazinone, 4(3*H*)-pyrimidone, 3,4-dihydropyrimidin-4-one, glutarimide, 4,6-(1*H*,5*H*)-pyrimidinedione, 1,3,5-triazin-2(1*H*)-one, and cyanuric acid, for example.

In a preferred embodiment, G comprises a heterocyclic 5-membered ring system. Preferably, G is an oxazole or an 1,2,4-oxadiazole ring. The G moiety may have either one of two possible orientations with respect to the Ar¹ and Ar² groups. Thus, for example, the invention prefers compounds having the configuration 4-(Ar¹)-2-(Ar²)-oxazole or 3-(Ar¹)-5-(Ar²)-1,2,4-oxadiazole.

In yet another embodiment, compounds of the present invention are within the class represented by formula II:



wherein X, Y, and Z are independently selected from the group consisting of N, O, S, C, and CO wherein at least one of X, Y, and Z is a heteroatom;

15 Ar¹ and Ar² are independently selected from the group consisting of a heterocyclic or fused heterocyclic moiety containing 1 to 4 heteroatoms selected from the group consisting of N, O, and S and an aromatic moiety selected from the group consisting of phenyl, benzyl, 1-naphthyl, 2-naphthyl, fluorenyl, anthrenyl, indenyl, phenanthrenyl, and benzonaphthyl, wherein the Ar¹ and Ar² moieties are optionally substituted with one or more moieties selected from the group consisting of -F, -Cl, -Br, -I, -OR, -SR, -SOR, -SO₂R, -SO₂NRR', -OCOR, -OCONRR', -NRCOR', -NRCO₂R', -CN, -NO₂, -CO₂R, -CONRR', -C(O)R, -CH(O)R', -CH₂(OR), -R, and -A-(CH₂)_n-NRR'; wherein R or R' is selected from the group consisting of H, CF₃, C₁-C₁₀ alkyl, cycloalkyl, alkyl-aryl, alkyl-heteroaryl, heterocycloalkyl, aryl and where R and R' may combine to form a ring, and A is defined as CH₂, O, NH, S, SO, SO₂ and n is 1, 2, 3, or 4.

20

25

30

35

In a preferred embodiment of the invention, the compound is selected from the group consisting of 3-(2-pyridyl)-5-(3,5-dichlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-methoxyphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-methylphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(1-naphthyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,3-difluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,5-difluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3,5-difluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(2-

pyridyl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazol, 3-(2-pyridyl)-5-(2,3-dichlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chloro-5-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-fluoro-5-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chloro-5-fluorophenyl)-1,2,4-oxadiazole, 3-(5-chloropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(5-fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(5-fluoropyrid-2-yl)-5-(3-cyano-5-fluorophenyl)-1,2,4-oxadiazole, 3-(3-fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(5-fluoropyrid-2-yl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole, 3-(5-methoxypyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(2-quinolinyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(3-chloro-5-trifluoromethylpyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(5-chloro-2-methoxyphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2-chloro-5-methylthiophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2-bromo-5-methoxyphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,5,6-trifluorophenyl)-1,2,4-oxadiazole, 2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole and 3-(2-pyridyl)-5-(2,5,6-trifluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-nitrophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-bromophenyl)-1,2,4-oxadiazole and pharmaceutically acceptable salts thereof.

In another embodiment of the invention, the compound is selected from the group consisting of 2-(3,5-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-methylphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(1-naphthyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-trifluoromethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2,3-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3,5-dimethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2,3-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-chloro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-fluoro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-chloro-5-fluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(5-chloropyrid-2-yl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole, 2-(3-cyano-5-fluorophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(3-fluoropyrid-2-yl)-1,3-oxazole, 2-(3,5-dimethoxyphenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(5-methoxypyrid-2-yl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(2-quinolinyl)-1,3-oxazole, 2-(3-cyanophenyl)-4-(3-chloro-5-trifluoromethylpyrid-2-yl)-1,3-oxazole, 2-(5-chloro-2-methoxyphenyl)-4-(2-pyridyl)-1,3-

oxazole, 2-(2-chloro-5-methylthiophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2-bromo-5-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole and 2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-nitrophenyl)-4-(2-pyridyl)-1,3-oxazole, 2-(3-bromophenyl)-4-(2-pyridyl)-1,3-oxazole and pharmaceutically acceptable salts thereof.

Preparation of mGluR Group I antagonists

Many starting materials for preparing the compounds of the present invention are available from commercial sources, such as Aldrich Chemical Company (Milwaukee, WI).

Moreover, compounds of the invention are readily prepared, from available precursors, using straightforward transformations which are well known in the art. The skilled artisan will recognize that mGluR Group I antagonists, according to the invention, can be prepared via methodology that is well known, using widely recognized techniques of organic chemistry. Suitable reactions are described in standard textbooks of organic chemistry.

For example, see March, *ADVANCED ORGANIC CHEMISTRY*, 2d ed., McGraw Hill (1977).

More specifically, compounds of the invention generally can be prepared by formation of the G moiety between two precursor compounds containing suitable Ar¹ and Ar² moieties. When the linker contains a 1,2,4-oxadiazole, the heterocycle may be formed using well known techniques, such as reaction between an amidoxime and an acid chloride, or by the reaction of an amidoxime and an acylimidazole. An illustration of such a transformation is provided in Examples 3 through 6, below.

Amidoximes can be prepared using well known techniques by the reaction of an Ar¹ substituted nitrile with hydroxylamine. An illustration of such a transformation is provided below in Example 1.

In most cases, the precursor Ar² acid chlorides are readily available, or may be prepared using straightforward techniques of organic chemistry. For example, carboxylic acids may be converted into the corresponding acid chlorides by reaction with, for example, thionyl chloride or oxalyl chloride.

In the case where the linker contains a 1,3-oxazole, compounds were prepared using a procedure similar to that given by Kelly *et al.*, *J. Org. Chem.* 61, 4623-4633 (1996). Thus, 3,5-Disubstituted-1,3-Oxazoles were prepared by mixing a haloketone with

carboxamide in refluxing toluene for 3 days. The resulting mixture was allowed to cool to room temperature, the solvent was removed and the residue was purified.

Testing of compounds for mGluR Group I antagonist activity

The pharmacological properties of the compounds of the invention can be analyzed 5 using standard assays for functional activity. Examples of glutamate receptor assays are well known in the art, for example, see Aramori *et al.*, *Neuron* 8:757 (1992); Tanabe *et* al., *Neuron* 8:169 (1992); Miller *et al.*, *J. Neuroscience* 15: 6103 (1995); Balazs, *et al.*, *J. Neurochemistry* 69:151 (1997). The methodology described in those publications is incorporated herein by reference.

10 Conveniently, the compounds of the invention can be studied by means of an assay that measures the mobilization intracellular calcium, $[Ca^{2+}]_i$ in cells expressing mGluR5 that can bind the compounds. A well-known cell line which is suitable for this purpose is described in Miller *et al.*, *J. Neuroscience* 15: 6103 (1995), the contents of which are hereby incorporated by reference. It has been shown that exposure of rat astrocytes to the 15 growth factors basic fibroblast growth factor, EGF, or transforming growth factor- α markedly increased the protein expression and functional activity of endogenous mGluR5 (Miller *et al.*, *J. Neuroscience*, 15(9): 6103-6109, 1995).

In brief, primary astrocyte cultures were prepared from 3-5 day old Sprague-Dawley rat pups using a modification of Miller *et al.* and were plated on poly-L lysine 20 coated flasks in Dulbecco's modified Eagle's medium (DMEM) containing fetal calf serum (FCS). For cuvette analysis, cultures were up-regulated with growth factors in flasks for 3-5 days, then harvested and prepared for measurement of $[Ca^{2+}]_i$ mobilization as previously described (Nemeth *et al.*, 1998).

For fluorescent imaging plate reader (FLIPR) analysis, cells were seeded on poly-D 25 lysine coated clear bottom 96-well plates with black sides and analysis of $[Ca^{2+}]_i$ mobilization was performed 3 days following the growth factor up-regulation.

FLIPR experiments were carried out using a laser setting of 0.800 W and a 0.4 second CCD camera shutter speed. Each FLIPR experiment was initiated with 180 μ L of buffer present in each well of the cell plate. After each addition of compound, the 30 fluorescence signal was sampled 50 times at 1 second intervals followed by 3 samples at 5 second intervals. Responses were measured as the peak height of the response within the sample period.

EC₅₀ and IC₅₀ determinations were made from data obtained from 8 point concentration response curves (CRC) performed in duplicate. Agonist CRC were generated by scaling all responses to the maximal response observed for the plate. Antagonist block of the agonist challenge was normalized to the average response of the 5 agonist challenge in 14 control wells on the same plate. A detailed protocol for testing the compounds of the invention is provided below at Example 4.

Preparation of pharmaceutical compositions containing mGluR antagonists, and their use in treating neurological disorders

The compounds of the present invention are useful for treating neurological 10 disorders or diseases. While these compounds typically will be used in therapy for human patients, they also can be used in veterinary medicine, to treat similar or identical diseases.

In therapeutic and/or diagnostic applications, the compounds of the invention can be formulated for a variety of modes of administration, including systemic and topical or localized administration. Techniques and formulations generally may be found in 15 REMINGTON'S PHARMACEUTICAL SCIENCES (18th ed.), Mack Publishing Co. (1990).

The compounds according to the invention are effective over a wide dosage range. For example, in the treatment of adult humans, dosages from about 0.01 to about 1000 mg per 60-70 kg adult, preferably from about 0.5 to about 100 mg per 60-70 kg adult, per 20 [day] dose may be used. A more preferable dosage is about 2 mg to about 70 mg per 60-70 kg adult per [day] dose. The exact dosage will depend upon the route of administration, the form in which the compound is administered, the subject to be treated, the body weight of the subject to be treated, and the preference and experience of the attending physician.

Pharmaceutically acceptable salts are generally well known to those of ordinary skill in the art, and may include, by way of example but not limitation, acetate, 25 benzenesulfonate, besylate, benzoate, bicarbonate, bitartrate, bromide, calcium edetate, camsylate, carbonate, citrate, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isethionate, lactate, lactobionate, malate, maleate, mandelate, mesylate, mucate, napsylate, nitrate, pamoate (embonate), 30 pantothenate, phosphate/disphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, sulfate, tannate, tartrate, or teoclate. Other pharmaceutically acceptable salts

may be found, for example, in REMINGTON'S PHARMACEUTICAL SCIENCES (18th ed.), *supra*.

Preferred pharmaceutically acceptable salts include, for example, acetate, benzoate, bromide, carbonate, citrate, gluconate, hydrobromide, hydrochloride, maleate, mesylate, 5 napsylate, pamoate (embonate), phosphate, salicylate, succinate, sulfate, or tartrate.

Depending on the specific conditions being treated, such agents may be formulated into liquid or solid dosage forms and administered systemically or locally. The agents may be delivered, for example, in a timed- or sustained-release form as is known to those skilled in the art. Techniques for formulation and administration may be found in 10 REMINGTON'S PHARMACEUTICAL SCIENCES; (18th ed.), *supra*. Suitable routes may include oral, buccal, sublingual, rectal, transdermal, vaginal, transmucosal, nasal or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections, *inter alia*.

15 For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants generally are known in the art.

20 Use of pharmaceutically acceptable carriers to formulate the compounds herein disclosed for the practice of the invention into dosages suitable for systemic administration is within the scope of the invention. With proper choice of carrier and suitable manufacturing practice, the compositions of the present invention, in particular, those formulated as solutions, may be administered parenterally, such as by intravenous 25 injection. The compounds can be formulated readily using pharmaceutically acceptable carriers well known in the art into dosages suitable for oral administration. Such carriers enable the compounds of the invention to be formulated as tablets, pills, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

30 Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. The preparations formulated for oral administration may be in the form 5 of tablets, dragees, capsules, or solutions.

Pharmaceutical preparations for oral use can be obtained by combining the active compounds with solid excipients, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including 10 lactose, sucrose, mannitol, or sorbitol; cellulose preparations, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethyl-cellulose (CMC), and/or polyvinylpyrrolidone (PVP: povidone). If desired, disintegrating agents may be added, such as the cross-linked polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as 15 sodium alginate.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol (PEG), and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dye-stuffs or 20 pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin, and a plasticizer, such as 25 glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols (PEGs). In addition, stabilizers may be added.


The present invention, thus generally described, will be understood more readily by 30 reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the present invention.

EXAMPLES**General Experimental Methods**


Capillary gas chromatographic and mass spectral data were obtained using a Hewlett-Packard (HP) 5890 Series II Gas Chromatograph coupled to an HP 5971 Series

5 Mass Selective Detector [Ultra-2 Ultra Performance Capillary Column (crosslinked 5% PhMe silicone); column length, 25 m; column i.d., 0.20 mm; helium flow rate, 60 mL/min; injector temp., 250 °C; temperature program, 20 °C/min from 125 to 325 °C for 10 min, then held constant at 325 °C for 6 min]. Thin-layer chromatography was performed using Analtech Uniplate 250-μm silica gel HF TLC plates. UV light sometimes 10 in conjunction with ninhydrin and Dragendorff's spray reagents (Sigma Chemical Co.) were used for detecting compounds on the TLC plates. Most reagents used in reactions were purchased from the Aldrich Chemical Co. (Milwaukee, WI), Sigma Chemical Co. (Saint Louis, MO), Fluka Chemical Corp. (Milwaukee, WI), Fisher Scientific (Pittsburgh, PA), TCI America (Portland, OR), or Lancaster Synthesis (Windham, NH).


15 **Example 1: Synthesis of amidoxime intermediates**

Pyrid-2-ylamidoxime

Using the procedure of Shine et al., *J. Heterocyclic Chem.* (1989) 26:125-128, hydroxylamine hydrochloride (7.65 g, 110 mmol) in ethanol (100 mL) was treated with a 20 solution of sodium hydroxide (11 mL of 10 N, 110 mmol). A precipitate quickly formed and the reaction mixture was stirred at ambient temperature for 30 min. The inorganic precipitate was filtered and rinsed with ethanol (100 mL). The filtrate and ethanol washings were combined and treated with 2-cyanopyridine (10.4 g, 100 mmol). The reaction mixture was heated at reflux for 20 hours. The volatiles were then removed *in* 25 *vacuo* to afford 13.3 g (97%) of pyrid-2-ylamidoxime.


3-Methoxybenzamidoxime

Using the general procedure for the synthesis of amidoximes, hydroxylamine hydrochloride (7.65 g, 110 mmol), sodium hydroxide (11 mL of 10 *N*, 110 mmol), and 3-methoxybenzyl nitrile (12.2 mL, 100 mmol) afforded 9.9 g (60%) of 3-methoxybenzamidoxime.

5-Chloropyrid-2-ylamidoxime

10 A mixture of 2,5-dichloropyridine (1.48 g, 10 mmol), zinc cyanide (705 mg, 6 mmol), zinc (dust, 29 mg, 0.45 mmol), [1,1'-bis(diphenylphosphino)ferrocene] dichloropalladium(II) complex with dichloromethane (1:1) (0.18 g, 0.22 mmol) in *N,N*-dimethylformamide (10 mL) was heated at reflux for 5 hours. After cooling, the reaction was diluted with ethyl acetate and extracted with water and brine. Silica gel chromatography afforded 735 mg (53%) of 2-cyano-5-chloropyridine.

15 Using the general procedure for the synthesis of amidoximes, 2-cyano-5-chloropyridine (735 mg, 5.3 mmol), a solution of hydroxylamine hydrochloride (1.2 mL of 5 M, 6 mmol) in ethanol (7 mL), and sodium hydroxide (0.61 mL of 10 *N*, 6.1 mmol), were heated at reflux for 24 hours. Standard work up afforded 707 mg (77%) of 5-chloropyrid-2-ylamidoxime.


5-Fluoropyrid-2-ylamidoxime

20 A mixture of 2-cyano-5-chloropyridine (1 g, 7.22 mmol) and potassium fluoride (1.26 g, 21.68 mmol) in 1-methyl-2-pyrrolidinone (25 mL) was heated at reflux 18 hours. After cooling, the reaction was diluted with ethyl acetate and extracted with water and

brine. The organic solvents were then removed *in vacuo*. Silica gel chromatography of the residue afforded 425 mg (48%) of 2-cyano-5-fluoropyridine.

Using the general procedure for the synthesis of amidoximes, 2-cyano-5-fluoropyridine (425 mg, 3.48 mmol), a solution of hydroxylamine hydrochloride (0.79 ml of 5 M, 3.95 mmol) in ethanol (5 mL), and sodium hydroxide (0.398 mL of 10 N, 3.98 mmol) were heated at reflux for 24 hours. Standard work up afforded 330 mg (61%) of 5-fluoropyrid-2-ylamidoxime.

5-Methoxypyrid-2-ylamidoxime



10

A solution of 2-cyano-5-fluoropyridine (0.65 g, 5.3 mmol) in sodium methoxide (1.83 mL of 25% wt. solution in methanol, 7.95 mmol) was stirred at 0 °C for 1.5 hours and 2 hours at ambient temperature. The reaction was then diluted with ethyl acetate and washed with water and brine. Removal of the solvent *in vacuo* afforded 304 mg (43%) of 15 2-cyano-5-methoxypyridine.

Using the general procedure for the synthesis of amidoximes, 2-cyano-5-methoxypyridine (270 mg, 2.01 mmol), a solution of hydroxylamine hydrochloride (0.457 ml of 5 M, 2.28 mmol) in ethanol (4 mL), and sodium hydroxide (0.230 mL of 10 N, 2.30 mmol) were heated at reflux for 24 hours. Standard work up afforded 79 mg (24%) of 20 5-methoxypyrid-2-ylamidoxime.

3-Fluoropyrid-2-ylamidoxime




A mixture of 2,3-dichloropyridine (1.48 g, 10 mmol), zinc cyanide (705 mg, 6 mmol,), zinc (dust, 29 mg, 0.45 mmol), [1,1'-bis(diphenylphosphino)ferrocene] 25 dichloropalladium(II), complex with dichloromethane (1:1) (0.18 g, 0.22 mol) in *N,N*-dimethylformamide (10 mL) was heated at reflux for 5 hours. After cooling, the reaction was diluted with ethyl acetate and extracted with water and brine. Removal of the solvent and silica gel chromatography afforded 1.05 g (76%) of 2-cyano-3-chloropyridine.

A solution of 2-cyano-3-chloropyridine (1 g, 7.22 mmol) in 1-methyl-2-pyrrolidinone (25 mL) was treated with potassium fluoride (1.26 g, 21.68 mmol) and heated at reflux for 18 hours. After cooling, the reaction was diluted with ethyl acetate and extracted with water and brine. Silica gel chromatography afforded 442 mg (50%) of 5 2-cyano-3-fluoropyridine.

Using the general procedure for the synthesis of amidoximes, 2-cyano-3-fluoropyridine (442 mg, 3.62 mmol), a solution of hydroxylamine hydrochloride (0.82 mL of 5 M, 4.1 mmol) in ethanol (5 mL), and sodium hydroxide (0.415 ml of 10 N, 4.15 mmol) were heated at reflux for 24 hours. Standard work up afforded 368 mg (66%) of 10 3-fluoropyrid-2-ylamidoxime.


Quinol-2-ylamidoxime



Using the general procedure for the synthesis of amidoximes, 2-quinolincarbonitrile (1.02 g, 6.6 mmol), a solution of hydroxylamine hydrochloride (1.44 mL of 5 N solution, 7.2 mmol) in ethanol (10 mL), and sodium hydroxide (0.72 mL of 10 N solution, 7.2 mmol) were heated at reflux for 18 hours. Standard work up afforded 990 mg (80%) of quinol-2-ylamidoxime.

20 Example 2: Synthesis of carboxylic acid intermediates


3-Chloro-5-cyanobenzoic acid



A mixture of methyl 3,5-dichlorobenzoate (14.66 g, 71.5 mmol), zinc cyanide (5.04 g, 42.9 mmol) zinc (dust, 0.21 g, 3.21 mmol), [1,1'Bis(diphenylphosphino)ferrocene] dichloropalladium(II), complex with dichloromethane (1:1) (1.3 g, 1.57 mmol) in N,N-dimethylformamide (70 mL) was heated at reflux for 5 hours. After cooling the reaction was diluted with ethyl acetate and extracted with water and brine. Silica gel chromatography afforded 2.34g (17%) methyl 2-chloro-5-cyanobenzoate.

The intermediate ester was treated with a solution of sodium hydroxide (7.5 mL of 4 N solution, 30 mmol) in methanol (50 mL) and stirred at ambient temperature for 18 hours. The solvent was removed *in vacuo* and the residue dissolved in ethyl acetate. The organic solution was washed with 5% HCl and brine. Removal of the solvent afforded 1.8 g (83%) of 3-chloro-5-cyanobenzoic acid.

3-Chloro-5-fluorobenzoic acid




A mixture of 1-bromo-3-chloro-5-fluorobenzene (25.0 g, 120 mmol), zinc cyanide (8.45 g, 72 mmol) zinc (dust, 235 mg, 3.6 mmol), [1,1'Bis(diphenylphosphino)ferrocene] dichloropalladium(II), complex with dichloromethane (1:1) (1.5 g, 1.8 mmol) in *N,N*-dimethylformamide (70 ml) was heated at reflux for 1 hour. After cooling the reaction was diluted with ethyl acetate and extracted with water and brine. Silica gel chromatography afforded 15.9g (85%) 3-chloro-5-fluorobenzonitrile.

The intermediate nitrile was treated with a solution of sodium hydroxide (100 mL of 10 N solution, 1 mol) in 100 mL water and heated at reflux for 2 hours. After this time the solution was cooled and acidified with concentrated hydrochloric acid. Extraction with dichloromethane and evaporation of the solvent, afforded 15.14g (85%) of 3-chloro-5-fluorobenzoic acid.

20

3-Fluoro-5-cyanobenzoic acid



3-Chloro-5-fluorobenzoic acid (13.74g, 78.7 mmol) was treated with 50 ml thionyl chloride and heated at reflux for 2 hours. The excess thionyl chloride was removed *in vacuo* and the residue treated with 100 ml dry methanol to afford 13.6g (92%) of methyl 3-chloro-5-fluorobenzoate.


A mixture of the methyl 3-chloro-5-fluorobenzoate, zinc cyanide (8.46g, 72.3 mmol) zinc (dust, 235 mg, 3.6mmol), [1,1'bis(diphenylphosphino)ferrocene] dichloropalladium(II), complex with dichloromethane (1:1) (1.5 g, 1.8 mmol) in *N,N*-dimethylformamide (70 ml) was heated at reflux for 1 hour. The reaction was cooled to 5 ambient temperature and diluted with ethyl acetate. The organic solution was extracted with water and brine and concentrated *in vacuo*, to afford crude methyl 3-chloro-5-cyanobenzoate.

The crude methyl 3-chloro-5-cyanobenzoate was treated with a solution of sodium hydroxide (45 ml of 4 *N* solution, 180 mmol) in methanol (350 mL) at ambient temperature 10 for 4 hours. The solvent was removed *in vacuo* and the residue dissolved in ethyl acetate. The organic solution was washed with 5% aqueous HCl and brine. Silica gel chromatography afforded 7.0 g (54%) of 3-fluoro-5-cyanobenzoic acid.

Example 3: Synthesis of 3,5-disubstituted-1,2,4-oxadiazoles from acid chlorides

15 In general, modifications were made from the procedures given Shine et al., *J. Heterocyclic Chem.* (1989) 26:125-128. 3,5-Disubstituted-1,2,4-oxadiazoles were typically made by adding an acylchloride to a solution of an amidoxime in pyridine after which the reaction mixture was either heated to reflux or placed in a sealed tube and heated. Typically, the oxadiazoles were isolated by precipitating with cold water and 20 filtering or by extraction with an organic solvent. If necessary, the oxadiazoles were purified by chromatography or recrystallization.

3-(2-Pyridyl)-5-(3,5-dichlorophenyl)-1,2,4-oxadiazole (NPS 64982) (404) B2




25 A mixture of 3,5-dichlorobenzoyl chloride (2.1 g, 10 mmol) and pyrid-2-ylamidoxime (1.37 g, 10 mmol) in pyridine (5 mL) was heated in sealed tube at 190 °C for 2 hours. After this time, the reaction mixture was added to ice cold water to precipitate the oxadiazole. The solid was collected by filtration, washed with water and then recrystallized from ethanol to yield 2.1 g (72%) of 3-(2-pyridyl)-5-(3,5-dichlorophenyl)-

1,2,4-oxadiazole: mp 162-166 °C; GC/EI-MS gave *m/z* (rel. int.) 291 (M⁺, 38), 293 (25), 261 (1), 173 (6), 145 (13), 120 (100), 90 (20), 78 (28), 51 (15).


3-(2-Pyridyl)-5-(3-chlorophenyl)-1,2,4-oxadiazole (NPS 64983) (405) B3

5




Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-chlorobenzoyl chloride (127 μ L, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at reflux for 4 hours. Standard work up afforded 156 mg (61%) of 3-(2-pyridyl)-5-(3-chlorophenyl)-1,2,4-oxadiazole: mp 136-140 °C; GC/EI-MS gave *m/z* (rel. int.) 257 (M⁺, 64), 259 (21), 227 (3), 120 (100), 111 (22), 90 (24), 78 (32), 75 (26), 51 (20).

3-(2-Pyridyl)-5-(3-methoxyphenyl)-1,2,4-oxadiazole (B1)



15 Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-anisoyl chloride (151 μ L, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at reflux for 4 hours. Standard work up afforded 200 mg (79%) of 3-(2-pyridyl)-5-(3-methoxyphenyl)-1,2,4-oxadiazole: mp 96-99 °C; GC/EI-MS gave *m/z* (rel. int.) 253 (M⁺, 100), 223 (3), 179 (3), 135 (74), 133 (90), 92 (27), 78 (29), 77 (32), 64 (23), 63 (23).

3-(2-Pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole (B5)




25 Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 2-chlorobenzoyl chloride (127 μ L, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at reflux for 4 hours. Standard work up afforded 157 mg (61%) of 3-(2-pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole: mp 93-94 °C; GC/EI-MS gave *m/z* (rel.

int.) 257 (M⁺, 76), 259 (26), 227 (4), 139 (11), 120 (100), 111 (21), 90 (27), 78 (35), 75 (29), 51 (21).

3-(2-Pyridyl)-5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazole (B6)


5



Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-(trifluoromethyl)benzoyl chloride (151 μ L, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at reflux for 16 hours. Standard work up afforded 10 233 mg (80%) of 3-(2-pyridyl)-5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazole: mp 116-118 °C; GC/EI-MS gave *m/z* (rel. int.) 291 (M⁺, 81), 272 (7), 173 (6), 145 (25), 120 (100), 90 (20), 78 (23), 51 (11).


3-(2-Pyridyl)-5-(3-fluorophenyl)-1,2,4-oxadiazole (B7)

15



Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-fluorobenzoyl chloride (122 μ L, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at reflux for 16 hours. Standard work up afforded 176 mg (73%) of 3-(2-pyridyl)-5-(3-fluorophenyl)-1,2,4-oxadiazole: mp 88-98 °C; GC/EI-MS gave *m/z* (rel. int.) 241 (M⁺, 95), 211 (5), 120 (100), 107 (13), 95 (30), 90 (21), 78 (27), 75 (19), 51 (15).


3-(2-Pyridyl)-5-(3-methylphenyl)-1,2,4-oxadiazole (B9)



25 Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-toluoyl chloride (264 μ L, 2 mmol) and pyrid-2-ylamidoxime (274 mg, 2 mmol) in pyridine (1 mL) were heated in a sealed tube at 200 °C for 2 hours. Standard work up afforded 387 mg (82%) of 3-(2-pyridyl)-5-(3-toluoyl)-1,2,4-oxadiazole: mp 127-128 °C; GC/EI-MS gave

m/z (rel. int.) 237 (M^+ , 100), 222 (2), 207 (8), 120 (68), 117 (24), 91 (29), 90 (29), 78 (32), 65 (26), 51 (23).


3-(2-Pyridyl)-5-(1-naphthyl)-1,2,4-oxadiazole (B10)



5


Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 1-naphthoyl chloride (150 μ L, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated in a sealed tube at 200 $^{\circ}$ C for 3 hours. Standard work up afforded 50 mg (18%) of 3-(2-pyridyl)-5-(1-naphthyl)-1,2,4-oxadiazole: mp 132-136 $^{\circ}$ C; GC/EI-MS gave *m/z* (rel. int.) 273 (M^+ , 75), 195 (5), 169 (88), 153 (100), 139 (12), 127 (66), 126 (29), 105 (23), 78 (14), 51 (14).

3-(2-Pyridyl)-5-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazole (B11)



15 Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-(trifluoromethoxy)benzoyl chloride (220 mg, 1 mmol), and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated in a sealed tube at 200 $^{\circ}$ C for 3 hours. Standard work up afforded 175 mg (57%) of 3-(2-pyridyl)-5-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazole: mp 86-88 $^{\circ}$ C; GC/EI-MS gave *m/z* (rel. int.) 307 (M^+ , 73), 277 (3), 222 (3), 189 (6), 161 (5), 120 (100), 78 (21), 69 (17), 51 (10).

3-(2-Pyridyl)-5-(2,3-difluorophenyl)-1,2,4-oxadiazole (B16)




25 Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 2,3-difluorobenzoyl chloride (124 μ L, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at 100 $^{\circ}$ C for 16 hours. Standard work up afforded 158 mg

(61%) of 3-(2-pyridyl)-5-(2,3-difluorophenyl)-1,2,4-oxadiazole: mp 120-121 °C; GC/EI-MS gave *m/z* (rel. int) 259 (M⁺, 97), 229 (5), 228 (4), 141 (11), 120 (100), 113 (26), 90 (27), 78 (34), 51 (17).


5

3-(2-Pyridyl)-5-(2,5-difluorophenyl)-1,2,4-oxadiazole (B17)



Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 2,5-difluorobenzoyl chloride (124 μ L, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at 100 °C for 16 hours. Standard work up afforded 3-(2-pyridyl)-5-(2,5-difluorophenyl)-1,2,4-oxadiazole: mp 120-126 °C; GC/EI-MS gave *m/z* (rel. int) 259 (M⁺, 91), 229 (5), 228 (4), 141 (13), 120 (100), 113 (25), 90 (23), 78 (27), 51 (14).


3-(2-Pyridyl)-5-(3,5-difluorophenyl)-1,2,4-oxadiazole (B18)



15

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3,5-difluorobenzoyl chloride (1.25 mL, 10 mmol) and pyrid-2-ylamidoxime (1.37 g, 10 mmol) in pyridine (5 mL) were heated in a sealed tube at 200 °C for 4 hours. Standard work up afforded 1.2 g (46%) of 3-(2-pyridyl)-5-(3,5-difluorophenyl)-1,2,4-oxadiazole: mp 115-119 °C; GC/EI-MS gave *m/z* (rel. int) 259 (M⁺, 100), 229 (4), 228 (5), 141 (9), 125 (13), 113 (30), 90 (19), 78 (27), 63 (23), 51 (15).

3-(2-Pyridyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole (B21)




25

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-cyanobenzoyl chloride (165 mg, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at 100 °C for 72 hours. Standard work up afforded 158 mg (64%) of 3-(2-

pyridyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole: mp 148-149 °C; GC/EI-MS gave *m/z* (rel. int.) 248 (M⁺, 85), 218 (5), 130 (6), 120 (100), 114 (9), 102 (28), 90 (26), 78 (37), 75 (19), 51 (30).

5


3-(2-Pyridyl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole (B23)



Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3,5-dimethoxybenzoyl chloride (200 mg, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at 100 °C for 72 hours. Standard work up afforded 210 mg (74%) of 3-(2-pyridyl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole: mp 145-148 °C; GC/EI-MS gave *m/z* (rel. int.) 283 (M⁺, 100), 253 (3), 165 (69), 163 (19), 137 (36), 122 (33), 107 (17), 90 (10), 78 (25), 63 (19), 51 (19).


15

3-(2-Pyridyl)-5-(2,3-dichlorophenyl)-1,2,4-oxadiazole (B25)



Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 2,3-dichlorobenzoyl chloride (209 mg, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at 100 °C for 48 hours. Standard work up afforded 236 mg (81%) of 3-(2-pyridyl)-5-(2,3-dichlorophenyl)-1,2,4-oxadiazole: mp 128-133 °C; GC/EI-MS gave *m/z* (rel. int.) 291 (M⁺, 66), 293 (43), 256 (6), 173 (10), 145 (11), 120 (100), 90 (19), 78 (27), 51 (14).

3-(2-Pyridyl)-5-(3-chloro-5-cyanophenyl)-1,2,4-oxadiazole (B26)




25 3-Chloro-5-cyanobenzoic acid (0.82 g, 4.97 mmol) was treated with a solution of oxalyl chloride (10 mL of 2.5 M in dichloromethane, 25 mmol) and a catalytic amount of

N,N-dimethylformamide. The reaction was stirred at ambient temperature for 2.5 hours. The excess oxalyl chloride was removed *in vacuo* to afford 3-chloro-5-cyanobenzoyl chloride.

5 Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 3-chloro-5-cyanobenzoyl chloride and pyrid-2-ylamidoxime (682 mg, 5 mmol, 1 equivalent) in pyridine (5 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up and recrystallization from 2-propanol afforded 250 mg (19%) of 3-(2-pyridyl)-5-(3-chloro-5-cyanophenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 282 (M⁺, 100), 283 (18),
10 284 (34), 251 (4), 136 (10), 120 (53), 100 (10), 78 (15), 51 (6).


3-(2-Pyridyl)-5-(3-fluoro-5-cyanophenyl)-1,2,4-oxadiazole (B27)



3-Fluoro-5-cyanobenzoic acid (2.5 g, 15.14 mmol) was treated with a solution of 15 oxalyl chloride (30 mL of 2.5 M in dichloromethane, 75 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred at ambient temperature for 2.5 hours. The excess oxalyl chloride was removed *in vacuo* to afford 3-fluoro-5-cyanobenzoyl chloride.

20 Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 3-fluoro-5-cyanobenzoyl chloride and pyrid-2-ylamidoxime (2.076 g, 15.15 mmol, 1 equivalent) in pyridine (5 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up and recrystallization from 2-propanol afforded 1.5 g (37%) of 3-(2-pyridyl)-5-(3-fluoro-5-cyanophenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 266 (M⁺, 81), 267 (13), 235 (5), 132 (12), 120 (100), 100 (18), 90 (18), 78 (35), 51 (20).


3-(2-Pyridyl)-5-(3-chloro-5-fluorophenyl)-1,2,4-oxadiazole (B28)



3-Chloro-5-fluorobenzoic acid (400 mg, 2.3 mmol) was treated with a solution of oxalyl chloride (4.6 mL of 2.5 M in dichloromethane, 11.5 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred at ambient temperature for 2.5 hours. The excess oxalyl chloride was removed *in vacuo* to afford 3-chloro-5-fluorobenzoyl chloride.


Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 3-chloro-5-fluorobenzoyl chloride and pyrid-2-ylamidoxime (314 mg, 2.3 mmol, 1 equivalent) in pyridine (5 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up and recrystallization from 2-propanol afforded 250 mg (39%) of 3-(2-pyridyl)-5-(3-chloro-5-fluorophenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 275(M⁺, 89), 276 (14), 277 (29), 129 (26), 120 (100), 109 (7), 90 (20), 78 (31), 51 (14).

15 **3-(5-Chloropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole (B29)**



Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-cyanobenzoyl chloride (675 mg, 4 mmol) and 5-chloropyrid-2-ylamidoxime (686 mg, 4 mmol) in pyridine (5 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up and recrystallization from 2-propanol afforded 357 mg (32%) of 3-(5-chloropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 282 (M⁺, 85), 283 (14), 284 (27), 156 (31), 154 (100), 112 (19), 102 (30), 76 (28), 64 (13).

3-(5-Fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole (B30)




25 Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-cyanobenzoyl chloride (0.534 g, 3.2 mmol) and 5-fluoropyrid-2-ylamidoxime (0.5 g, 3.2 mmol) in

pyridine (5 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up and recrystallization from 2-propanol afforded 370 mg (43%) of 3-(5-fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 266 (M⁺, 100), 267 (10), 138 (80), 114 (8), 102 (19), 96 (22), 76 (17), 57 (8).

5


3-(5-Fluoropyrid-2-yl)-5-(3-cyano-5-fluorophenyl)-1,2,4-oxadiazole (B31)



3-Fluoro-5-cyanobenzoic acid (1.0 g, 6 mmol) was treated with a solution of oxalyl chloride (12 mL of 2.5 M in dichloromethane, 30 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred at ambient temperature for 2.5 hours. The excess oxalyl chloride was removed *in vacuo* to afford 3-fluoro-5-cyanobenzoyl chloride.


Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 3-fluoro-5-cyanobenzoyl chloride (1.1 g, 6 mmol) and 5-fluoropyrid-2-ylamidoxime (0.93 g, 6 mmol) in pyridine (5 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up and recrystallization from 2-propanol afforded 0.41 g (24%) of 3-(5-fluoropyrid-2-yl)-5-(3-cyano-5-fluorophenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 284 (M⁺, 100), 285 (16), 253 (2), 138 (99), 120 (23), 108 (16), 96 (25), 82 (15), 57 (11).

3-(3-Fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole (B32)




20

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-cyanobenzoyl chloride (107 mg, 0.64 mmol) and 3-fluoropyrid-2-ylamidoxime (0.1 g, 0.64 mmol) in pyridine (5 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up, silica gel chromatography, and recrystallization from 2-propanol, afforded 32 mg (19%) of 3-(3-fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 266 (M⁺, 75), 267 (12), 138 (100), 114 (11), 102 (19), 96 (17), 76 (16), 57 (5), 51 (5).

3-(5-Fluoropyrid-2-yl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole (B33)


Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3,5-dimethoxybenzoyl chloride (0.10 g, 0.5 mmol) and 5-fluoropyrid-2-ylamidoxime (78 mg, 0.5 mmol) in pyridine (3 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up, silica gel chromatography, and recrystallization from 2-propanol afforded 94 mg (62%) of 3-(5-fluoropyrid-2-yl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 301 (M⁺, 100), 302 (17), 165 (41), 137 (23), 122 (27), 96 (15), 77 (11), 63 (12).

10

3-(5-Methoxypyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole (B34)

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-cyanobenzoyl chloride (79 mg, 0.47 mmol) and 5-methoxypyrid-2-ylamidoxime (79 mg, 0.47 mmol) in pyridine (2.5 mL) were heated in a sealed tube at 175 °C for 4 hours. Standard work up, silica gel chromatography, and recrystallization from 2-propanol afforded 59 mg (45%) of 3-(5-methoxypyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole: GC/EI-MS gave *m/z* (rel. int.) 278 (M⁺, 100), 279 (16), 150 (56), 128 (7), 107 (21), 102 (17), 80 (12), 64 (5).


20

3-(2-Quinolinyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole (B35)

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-cyanobenzoyl chloride (68 mg, 0.41 mmol) and quinol-2-ylamidoxime (75.9 mg, 0.405 mmol) in pyridine (0.5 mL) were heated in a sealed tube at 165 °C for 22 hours. Standard work up, recrystallization from ethanol, and solid phase extraction (SPE) afforded 23.7 mg (20%) of 3-(2-quinolinyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole. ¹H-NMR (CDCl₃), δ (ppm): 8.62 (s,


1H), 8.54 (d, 1H), 8.36 (d, 2H), 8.28 (d, 1H), 7.90 (d, 2H), 7.80 (t, 1H), 7.72 (t, 1H), 7.64 (t, 1H).

3-(3-chloro-5-trifluoromethylpyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole (B36)




Using the general procedure for the synthesis of 1,2,4-oxadiazoles, 3-cyanobenzoyl chloride (66 mg, 0.40 mmol) and 3-chloro-5-trifluoromethylpyrid-2-ylamidoxime (96.5 mg, 0.403 mmol) in pyridine (0.5 mL) were heated in a sealed tube at 165 °C for 22 hours. Standard work up and solid phase extraction (SPE) afforded 45.9 mg (33%) of 3-(3-chloro-5-trifluoromethylpyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole. ¹H-NMR (CDCl₃), δ (ppm): 8.99 (s, 1H), 8.57 (s, 1H), 8.49 (d, 1H), 8.19 (s, 1H), 7.92 (d, 1H), 7.72 (t, 1H).

15 3-(2-pyridyl)-5-(5-chloro-2-methoxyphenyl)-1,2,4-oxadiazole (B37)




5-Chloro-*O*-anisic acid (187 mg, 1 mmol) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred at ambient temperature for 2 hours. The excess oxalyl chloride was removed *in vacuo* to afford 5-chloro-2-methoxybenzoyl chloride.

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 5-chloro-2-methoxybenzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at 115 °C for 17 hours. Standard work up, and silica gel chromatography afforded 49 mg (17%) of 3-(2-pyridyl)-5-(5-chloro-2-methoxyphenyl)-1,2,4-oxadiazole. ¹H-NMR (CDCl₃), δ (ppm): 4.00 (s, 3H), 7.03 (d, J = 8.9 Hz, 1H), 7.42-7.47 (m, 1H), 7.50 (dd, J = 8.9 Hz, 2.8 Hz, 1H), 7.87 (ddd, J = 1.4 Hz, 7.4 Hz, 8.2 Hz, 1H), 8.22 (d, J = 8.2 Hz, 1H), 8.28 (d, J = 2.4 Hz, 1H), 8.84 (m, 1H).

3-(2-pyridyl)-5-(2,3-dimethoxyphenyl)-1,2,4-oxadiazole (B38)

2,3-Dimethoxybenzoic acid (182 mg, 1 mmol) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred at ambient temperature for 2 hours. The excess oxalyl chloride was removed *in vacuo* to afford 2,3-dimethoxybenzoyl chloride.


Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 2,3-dimethoxybenzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at 115 °C for 17 hours. Standard work up and silica gel chromatography afforded 120 mg (42%) of 3-(2-pyridyl)-5-(2,3-dimethoxyphenyl)-1,2,4-oxadiazole.

3-(2-pyridyl)-5-(2-chloro-5-methylthiophenyl)-1,2,4-oxadiazole (B39)

2-Chloro-5-methylthiobenzoic acid (182 mg, 1 mmol) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred at ambient temperature for 2 hours. The excess oxalyl chloride was removed *in vacuo* to afford 2-chloro-5-methylthiobenzoyl chloride.


Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 2-chloro-5-methylthiobenzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated at 115 °C for 17 hours. Standard work up and silica gel chromatography afforded 250 mg (82%) of 3-(2-pyridyl)-5-(2-chloro-5-methylthiophenyl)-1,2,4-oxadiazole.

¹H-NMR (CDCl₃), δ (ppm): 7.37 (dd, J = 2.4 Hz, 8.2 Hz, 1H), 7.40-7.50 (m, 2H), 7.89 (ddd, J = 1.4 Hz, 7.4 Hz, 8.2 Hz, 1H), 8.05 (d, J = 2.4 Hz, 1H), 8.23 (dd, J = 2.2 Hz, 25 8.0 Hz, 1 x H), 8.85 (m, 1H).

3-(2-Pyridyl)-5-(3-phenoxyphenyl)-1,2,4-oxadiazole (B40)


3-Phenoxybenzoic acid (214 mg, 1.0 mmol) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred overnight at ambient temperature. The excess oxalyl chloride was removed *in vacuo* to afford 3-phenoxybenzoyl chloride.

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 3-phenoxybenzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated in a sealed vial overnight at 110 °C. Standard work up afforded 118 mg (37%) of 3-(2-pyridyl)-5-(3-phenoxyphenyl)-1,2,4-oxadiazole as a white solid.

3-(2-Pyridyl)-5-(3-benzoylphenyl)-1,2,4-oxadiazole (B41)


3-Benzoylbenzoic acid (226 mg, 1.0 mmol) in dichloromethane (1.5 mL) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred overnight at ambient temperature. The excess oxalyl chloride was removed *in vacuo* to afford 3-benzoylbenzoyl chloride.

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 3-benzoylbenzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated in a sealed vial overnight at 110 °C. Standard work up and filtration through silica gel (with dichloromethane) afforded 200 mg (61%) of 3-(2-pyridyl)-5-(3-benzoylphenyl)-1,2,4-oxadiazole as a white solid. ¹H NMR (CDCl₃), δ (ppm): 8.85 (d, 1H), 8.68 (m, 1H), 8.53 (dd, 1H), 8.23 (d, 1H), 8.07 (m, 1H), 7.88 (m, 3H), 7.70 (m, 2H), 7.49 (m, 3H).

3-(2-Pyridyl)-5-(2-bromo-5-methoxyphenyl)-1,2,4-oxadiazole (B42)


2-Bromo-5-methoxybenzoic acid (231 mg, 1.0 mmol) in dichloromethane (1.5 mL) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred overnight at ambient temperature. The excess oxalyl chloride was removed *in vacuo* to afford 2-bromo-5-methoxybenzoyl chloride.

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 2-bromo-5-methoxybenzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated in a sealed vial overnight at 110 °C. Standard work up and filtration through silica gel (with dichloromethane) afforded 147 mg (44%) of 3-(2-pyridyl)-5-(2-bromo-5-methoxyphenyl)-1,2,4-oxadiazole. ¹H NMR (CDCl₃), δ (ppm): 8.85 (d, 1H), 8.24 (d, 1H), 7.89 (m, 1H), 7.65 (m, 2H), 7.47 (m, 1H), 6.99 (m, 1H), 3.89 (s, 3H).

15 3-(2-Pyridyl)-5-(2-chloro-5-(trifluoromethyl)phenyl)-1,2,4-oxadiazole (B43)


2-Chloro-5-(trifluoromethyl)benzoic acid (224 mg, 1.0 mmol) in dichloromethane (1.5 mL) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred overnight at ambient temperature. The excess oxalyl chloride was removed *in vacuo* to afford 2-chloro-5-(trifluoromethyl)benzoyl chloride.

Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 2-chloro-5-(trifluoromethyl)benzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated in a sealed vial overnight at 110 °C. Standard work up and filtration through silica gel (with dichloromethane) afforded 136 mg (42%) of 3-(2-pyridyl)-5-(2-chloro-5-(trifluoromethyl)phenyl)-1,2,4-oxadiazole as a beige solid. ¹H NMR (CDCl₃), δ (ppm): 8.87 (d, 1H), 8.56 (s, 1H), 8.25 (d, 1H), 7.89 (m, 1H), 7.78 (m, 2H), 7.50 (m, 1H).

3-(2-Pyridyl)-5-(3,4,5-trifluorophenyl)-1,2,4-oxadiazole (B44)

3,4,5-Trifluorobenzoic acid (0.176 g, 1.0 mmol) in dichloromethane (1.5 mL) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred overnight at ambient temperature. The excess oxalyl chloride was removed *in vacuo* to afford 3,4,5-trifluorobenzoyl chloride.


Using the general procedure for the synthesis of 1,2,4-oxadiazoles, the 3,4,5-trifluorobenzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) were heated in a sealed vial overnight at 110 °C. Standard work up and silica gel chromatography (with 10-30% ethyl acetate in hexane) afforded 15 mg (5%) of 3-(2-pyridyl)-5-(3,4,5-trifluorophenyl)-1,2,4-oxadiazole as a white solid.

3-(2-pyridyl)-5-(2,5,6-trifluorophenyl)-1,2,4-oxadiazole (B45)

2,5,6-Trifluorobenzoic acid (176 mg, 1 mmol) was treated with a solution of oxalyl chloride (1.5 mL of 2 M in dichloromethane, 3 mmol) and a catalytic amount of *N,N*-dimethylformamide. The reaction was stirred at ambient temperature for 16 hours. The excess oxalyl chloride was removed *in vacuo* to afford 2,5,6-trifluorobenzoyl chloride.


A solution of the intermediate 2,5,6-trifluorobenzoyl chloride and pyrid-2-ylamidoxime (137 mg, 1 mmol) in dichloromethane was stirred at ambient temperature for 0.5 hours. Silica gel chromatography afforded 151 mg (51%) of *N*-(2,5,6-trifluorobenzoyl)oxy]pyridine-2-carboximidamide.

A solution of *N*-(2,5,6-trifluorobenzoyl)oxy]pyridine-2-carboximidamide (50 mg, 0.169 mmol) in pyridine (0.3 mL) was heated at 115 °C for 17 hours. Standard work up, and silica gel chromatography, afforded 9.5 mg (20%) of 3-(2-pyridyl)-5-(2,5,6-trifluorophenyl)-1,2,4-oxadiazole.


Example 4: Synthesis of 3,5-disubstituted-1,2,4-oxadiazoles from acylimidazoles**3-(3-Methoxyphenyl)-5-(2-pyridyl)-1,2,4-oxadiazole (B8)**

5 Using modifications of the method of Shine et al., *J. Heterocyclic Chem.* (1989) 26:125-128, a solution of picolinic acid (123 mg, 1 mmol) in pyridine (1 mL) was treated with 1,1'-carbonyldiimidazole (162 mg, 1 mmol) and the reaction stirred at ambient temperature until the evolution of carbon dioxide ceased (30 min). The intermediate acylimidazole was then treated with 3-methoxybenzamidoxime (166 mg, 1 mmol) and the reaction heated at reflux for 1 hour. Ice cold water was added to the reaction mixture to precipitate the oxadiazole. The solid was collected by filtration, washed with water and dried to afford 80 mg (32%) of 3-(3-methoxyphenyl)-5-(2-pyridyl)-1,2,4-oxadiazole: mp 90-94 °C; GC/EI-MS gave *m/z* (rel. int.) 253 (M⁺, 100), 254 (17), 179 (2), 175 (2), 149 (77), 133 (33), 119 (4), 106 (29), 78 (45), 51 (18).


15

Example 5: Synthesis of 3,5-disubstituted-1,2,4-oxadiazoles from esters**3-(Pyrid-2-yl)-5-(2-hydroxyphenyl)-1,2,4-oxadiazole (B46)**

Using the method of Korbonits et al., *J. Chem. Soc. Perkin Trans. I* (1982) 759-20 766, a mixture of ethyl salicylate (200 mg, 1.2 mmol), pyrid-2-ylamidoxime (82.5 mg, 0.6 mmol), 21% sodium ethoxide (19.4 mL, 6 mmol) in ethanol (12mL) was heated at reflux for 16 hours. After cooling, the reaction mixture was diluted with dichloromethane (50 mL) and washed with water and saturated sodium hydrocarbonate. The organic layer was dried with sodium sulfate and concentrated *in vacuo*. Recrystallization from diethyl ether afforded 15 mg (5%) of 3-(Pyrid-2-yl)-5-(2-hydroxyphenyl)-1,2,4-oxadiazole.


3-(2-pyridyl)-5-(5-chloro-2-hydroxyphenyl)-1,2,4-oxadiazole (B47)

In a similar fashion, methyl 5-chloro-2-hydroxybenzoate (372 mg, 2 mmol), pyrid-2-ylamidoxime (137 mg, 1 mmol), 21 % sodium ethoxide (32.4 mL, 10 mmol) in ethanol (20 mL) were heated at reflux for 16 hours. Standard work up and recrystallization from diethyl ether afforded 14.2 mg (5%) of 3-(2-pyridyl)-5-(5-chloro-2-hydroxyphenyl)-1,2,4-oxadiazole.


Example 6: Synthesis of 3,5-disubstituted-1,2,4-oxadiazoles from isatoic anhydrides10 **3-(2-pyridyl)-5-(2-aminophenyl)-1,2,4-oxadiazole (B48)**

Using modifications from the procedure of Nagahara et al., *Chem. Pharm. Bull.*, (1975) 23:3178-3183, a mixture of isatoic anhydride (163 mg, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) was heated at 115 °C for 17 hours. 15 After cooling the reaction, the mixture was diluted with 50 mL of dichloromethane and washed with water and saturated sodium hydrocarbonate. The organic layer was dried over sodium sulfate, filtered through silica gel and concentrated *in vacuo*. Recrystallization from diethyl ether afforded 45.6 mg (19%) of 3-(2-pyridyl)-5-(2-aminophenyl)-1,2,4-oxadiazole.


20

3-(2-pyridyl)-5-(2-aminophenyl)-1,2,4-oxadiazole (B49)


In a similar fashion, 5-chloroisatoic anhydride (197 mg, 1 mmol) and pyrid-2-ylamidoxime (137 mg, 1 mmol) in pyridine (1 mL) was heated at 115 °C for 17 hours. 25 Work up afforded 138 mg (51%) of 3-(2-pyridyl)-5-(2-aminophenyl)-1,2,4-oxadiazole.

Example 7: Synthesis of 2,4-disubstituted-1,3-oxazoles**2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole (B50)**

Using the procedures of Kelly et al., *J. Org. Chem.*, (1996) 61:4623-4633, a
 5 solution of 2-bromoacetylpyridine (120 mg, 0.6 mmol) in toluene (5mL) was treated with
 3-chlorobenzamide (300 mg, 1.9 mmol) and the mixture heated in a sealed vial at reflux
 for 60 hours. The mixture was then cooled and the solvent was removed *in vacuo*. Silica
 gel chromatography using a gradient of hexane to ethyl acetate afforded 38 mg (9%) of 2-
 [3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole as a pale yellow solid. ¹H-NMR (CDCl₃), δ
 10 (ppm): 8.62 (d, 1H), 8.35 (s, 1H), 8.15 (m, 1H), 8.00 (m, 2H), 7.80 (td, 1H), 7.42 (m,
 2H), 7.23 (m, 1H).

2-[3-Bromophenyl]-4-[pyridin-2-yl]-1,3-oxazole (B51)

15 In a similar fashion 2-bromoacetylpyridine (500 mg, 2.5 mmol) and 3-
 chlorobenzamide (1.2g, 6 mmol) in toluene (10 mL) was heated in a sealed vial at reflux
 for 60 hours. Work up and silica gel chromatography using a gradient of hexane to ethyl
 acetate afforded 50 mg (7%) of 2-[3-bromophenyl]-4-[pyridin-2-yl]-1,3-oxazole as a white
 solid. ¹H-NMR (CDCl₃), δ (ppm): 8.60 (d, 1H), 8.34 (s, 1H), 8.30 (t, 1H), 8.00 (m,
 20 2H), 7.80 (td, 1H), 7.60 (dd, 1H), 7.35 (t, 1H), 7.23 (m, 1H).

2-[3-cyanophenyl]-4-[pyridin-2-yl]-1,3-oxazole (B52)


A mixture of 2-[3-bromophenyl]-4-[pyridin-2-yl]-1,3-oxazole (23 mg, 0.076 mmol)
 25 and zinc cyanide (112 mg, 0.96 mmol) in *N,N*-dimethylformamide (2 mL) was treated with
 Pd(PPh₃)₄ (74 mg, 0.064 mmol) and heated overnight at 80°C. Standard work up and
 chromatography afforded 6 mg (32 %) of 2-[3-cyanophenyl]-4-[pyridin-2-yl]-1,3-oxazole

as a white solid. $^1\text{H-NMR}$ (CDCl_3), δ (ppm): 8.61 (d, 1H), 8.45 (s, 1H), 8.38 (s, 1H), 8.36 (m, 1H), 8.00 (d, 1H), 7.80 (m, 2H), 7.61 (t, 1H), 7.23 (m, 1H).

Example 8: Synthesis of 3,5-disubstituted-1,2-oxazoles

5


5-[3-hydroxyphenyl]-3-[pyridin-2-yl]-1,2-oxazole (B53)



A stirred solution of pyridine-2-carbohydroximoyl chloride (300 mg, 1.9 mmol) and 3-hydroxyphenylacetylene (760 mg, 6.4 mmol) in a 1:1 mixture of $\text{THF}/\text{CH}_2\text{Cl}_2$ (10 mL) at 0 °C was treated with triethylamine (2 mL, 1.45 g, 15 mmol). The mixture was allowed to warm to room temperature overnight. The solvent was removed *in vacuo*. The residue was dissolved in dichloromethane, washed with brine and dried over anhydrous sodium sulphate. Removal of the solvent *in vacuo* followed by trituration with 10% ethylacetate in hexane afforded 200 mg (44%) of 5-[3-hydroxyphenyl]-3-[pyridin-2-yl]-1,2-oxazole as a beige solid.

10


5-[3-cyanophenyl]-3-[pyridin-2-yl]-1,2-oxazole (B54)



A mixture of 5-[3-trifluoromethanesulfonylphenyl]-3-[pyridin-2-yl]-1,2-oxazole (98 mg, 0.26 mmol), KCN (230 mg, 4 mmol), $\text{NiBr}_2(\text{PPh}_3)_2$ (52.4 mg, 0.07 mmol), and PPh_3 (42 mg, 0.16 mmol) in acetonitrile (1 mL) was treated with zinc powder (20 mg, 0.3 mmol) and the mixture was heated overnight at 60 °C. Silica gel chromatography of the resulting mixture using a gradient of hexane to ethyl acetate afforded 15 mg (23 %) of 5-[3-cyanophenyl]-3-[pyridin-2-yl]-1,2-oxazole as a white solid.


Example 9: Synthesis of 3,5-disubstituted-1,2,4-triazoles

3-Chlorobenzhydrazide




5 A mixture of 3-chlorobenzoic acid (0.5 g, 3.19 mmol), 1,3-dicyclohexylcarbodiimide (0.72 g, 3.51 mmol), 4-dimethylaminopyridine (0.04 g, 0.32 mmol) in ethanol was stirred at ambient temperature for 1.5 hour. The white solid was filtered off and the filtrate diluted with dichloromethane (100 mL). The organic solution was washed with 1 N sodium hydrogen sulfate (100 mL), saturated sodium bicarbonate (100 mL), water (100 mL) and brine (100 mL). The organic phase was dried over anhydrous magnesium sulfate and filtered. The filtrate was concentrated *in vacuo*. The crude residue was dissolved in ethanol (15 mL) and treated with hydrazine monohydrate (0.46 mL, 9.58 mmol). The resulting clear solution was stirred overnight at ambient temperature. The reaction mixture was then concentrated to dryness *in vacuo*. Silica gel chromatography of the residue, using 3% methanol in dichloromethane, afforded 0.29 g (53%) of 3-chlorobenzhydrazide as a white solid.

3-(2-Pyridyl)-5-(3-chlorophenyl)-1,2,4-triazole (B55)



20 Using the procedures of Browne et al., *Aust. J. Chem.*, (1975) 28:2543-2546, a solution of 2-cyanopyridine (0.1 mL, 1.00 mmol) in methanol (5 mL) was treated with sodium metal (6.9 mg, 0.30 mmol) and stirred for at ambient temperature for 1 hour. After this time, a solution of 3-chlorobenzhydrazide (0.17 g, 1.0 mmol) in methanol (5 mL) was added and the resulting solution heated at reflux for 3 hours. The reaction mixture 25 was concentrated *in vacuo*, and the resulting yellow solid (100 mg) dissolved in toluene (2 mL). The mixture was heated at 175 °C for 3 hours and then stirred overnight at ambient temperature. Evaporation of the solvent *in vacuo* and silica gel chromatography using 1% methanol in dichloromethane afforded 29 mg (11%) of 3-(2-pyridyl)-5-(3-chlorophenyl)-1,2,4-triazole as an off-white solid.

3-(2-Pyridyl)-5-(3-iodophenyl)-1,2,4-triazole (B56)

In a similar fashion, 2-cyanopyridine (0.15 mL, 1.53 mmol), sodium metal (10.5 mg, 0.46 mmol) and 3-iodobenzhydrazide (0.40 g, 1.53 mmol) afforded, after work up 5 and chromatography, 210 mg (40%) of 3-(2-pyridyl)-5-(3-iodophenyl)-1,2,4-triazole as a white solid.

Example 10: Assays of Group I receptor antagonist activity**Astrocyte Screening Assay**

Primary astrocyte cultures were prepared from 3-5 day old Sprague-Dawley rat 10 pups using a modification of Miller (Miller *et al*, *J. Neuroscience*, 15(9): 6103-6109, 1995). In brief, primary cultures were plated on poly-L lysine coated flasks in Dulbecco's modified Eagle's medium (DMEM) containing fetal calf serum (FCS). After 6 days, cell cultures were shaken over night at 280 rpm, then transferred to astrocyte-defined media 15 (ADM) containing growth factors that up-regulate the expression of mGluR5 (Miller *et al.*, 1995). For cuvette analysis, cultures were up-regulated with growth factors in flasks for 3-5 days, then harvested and prepared for measurement of $[Ca^{2+}]_i$ mobilization as previously described (Nemeth *et al.*, 1998).

For FLIPR analysis, cells were seeded on poly-D lysine coated clear bottom 96-well plates with black sides and analysis of $[Ca^{2+}]_i$ mobilization was performed 20 3 days following the growth factor up-regulation. Cell cultures in the 96-well plates were loaded with a 4 μ M solution of acetoxyethyl ester form of the fluorescent calcium indicator fluo-3 (Molecular Probes, Eugene, Oregon) in 0.01 % pluronic. All assays were performed in a buffer containing 127 mM NaCl, 5 mM KCl, 2 mM MgCl₂, 0.7 mM NaH₂PO₄, 2 mM CaCl₂, 0.422 mg/ml NaHCO₃, 2.4 mg/ml HEPES, 1.8 mg/ml glucose and 1 mg/ml BSA 25 Fraction IV (pH 7.4).

FLIPR experiments were done using a laser setting of 0.800 W and a 0.4 second CCD camera shutter speed. Each FLIPR experiment was initiated with 180 μ L of buffer present in each well of the cell plate. A 20 μ L addition from the antagonist plate was

followed by a 50 μ L addition from the agonist plate. After each addition the fluorescence signal was sampled 50 times at 1 second intervals followed by 3 samples at 5 second intervals. Responses were measured as the peak height of the response within the sample period.

5 EC₅₀/IC₅₀ determinations were made from data obtained from 8 point concentration response curves (CRC) performed in duplicate. Agonist CRC were generated by scaling all responses to the maximal response observed for the plate. Antagonist block of the agonist challenge was normalized to the average response of the agonist challenge in 14 control wells on the same plate.

10 **CaR/mGluR5d Screening Assay**

15 HEK 293 cells expressing the chimeric CaR/mGluR5d receptor (clonal cell line hCaR/hmGluR5d_hek6) are plated 24 hours prior to assay at a density of 100,000 cells per well in Collagen I-coated 96-well black, clear bottom plates (Becton Dickenson) in DMEM supplemented with 10% FBS (Hyclone).

20 On the day of the assay, tissue culture medium is aspirated from the wells of a plate and 80 μ L of Assay Buffer (Assay Buffer is: 20 mM HEPES, 146 mM NaCl, 5 mM KCl, 1 mM MgCl₂, 1 mM CaCl₂, 1 mg/ml BSA, 1 mg/ml glucose, pH 7.4) supplemented with 6 μ M of the Ca²⁺-sensitive dye, Fluo-3 AM (Molecular Probes) and 0.025% Pluronic 25 (Molecular Probes) is added to each well. The plate is then incubated in the dark for 1 hour at room temperature to efficiently load the cells with Fluo-3. At the end of the incubation, extracellular Fluo-3 is removed by washing the plate with Assay buffer. Assay Buffer is added back to each well (final volume = 160 μ L) prior to beginning the assay.

30 The plate is loaded into a FLIPR robotic device (Molecular Devices) with the laser setting at 0.8 Watts. At a time of 10 seconds after initiation of the assay, 40 μ L of Assay Buffer containing 62.5 μ M test substance and 2% DMSO is added to the 160 μ L of Assay Buffer in each well of the plate to yield a final concentration of 12 μ M test substance and 0.4% DMSO. At a time of 75 seconds after initiation of the assay, 50 μ L of Assay Buffer containing 6 mM CaCl₂ is added to the 200 μ L present in each well to yield a final Ca²⁺ concentration of 2.0 mM, and a final concentration of test substance of 10 nM. Relative fluorescence intensity (excitation λ = 488 nm / emission λ = 510 nM) is monitored at

relevant time intervals throughout the assay period to measure receptor activation and/or inhibition.


By way of illustration, a 1,2,4-oxadiazole disclosed above, designated "B21" (see Example 3), had an IC₅₀ value of 43 nM in relation to CaR/mGluR_{5a} and an IC₅₀ value of 5 121 nM on the native receptor, mGluR_{5a}. A corresponding 1,3 oxazole, designated "B52" (see Example 7), was found to be equipotent on the CaR/mGluR_{5a} chimera, with an IC₅₀ value of 45 nM, but displayed increased potency on the native mGluR_{5a} receptor, with an IC₅₀ value of 74 nM.

The invention thus has been disclosed broadly and illustrated in reference to 10 representative embodiments described above. Those skilled in the art will recognize that various modifications can be made to the present invention without departing from the spirit and scope thereof.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A compound selected from

a) a compound of Formula II:



wherein X and Y are N,

Z is O;

Ar¹ is 2-pyridyl;

Ar² is phenyl;

10 wherein at least one of Ar¹ and Ar² is substituted with one or more moieties selected from the group consisting of -F, -Cl, -Br, -I, -SR, -SOR, -SO₂R, -SO₂NRR', -OCOR, -OCONRR', -NRCOR', -NRCO₂R', -CN, -CO₂R, -CONRR', -C(O)R, -CH(OR)R', -CH₂(OR), CF₃, C₁-C₁₀ alkyl, cycloalkyl, aryl-alkyl, heterocycloalkyl and aryl;

15 wherein R or R' is selected from the group consisting of H, CF₃, C₁-C₁₀ alkyl, cycloalkyl, alkyl-aryl, heterocycloalkyl, aryl and where R and R' may combine to form a ring, and

with the proviso that the compound is not 3-(2-Pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole, or 3-(2-Pyridyl)-5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazole; or

20 b) 3-(2-pyridyl)-5-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazole,

3-(2-pyridyl)-5-(2,3-difluorophenyl)-1,2,4-oxadiazole,

3-(2-pyridyl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole,

3-(5-fluoropyrid-2-yl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole,

3-(5-methoxypyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole,

25 3-(2-pyridyl)-5-(3-nitrophenyl)-1,2,4-oxadiazole,

2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole,

2-(3,5-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole,

2-(3-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole,

2-(3-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,

30 2-(2-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole,


2-(3-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-methylphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-trifluoromethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
5 2-(2,3-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3,5-dimethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
10 2-(2,3-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-chloro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-fluoro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-chloro-5-fluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(5-chloropyrid-2-yl)-1,3-oxazole,
15 2-(3-cyanophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyano-5-fluorophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(3-fluoropyrid-2-yl)-1,3-oxazole,
2-(3,5-dimethoxyphenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(5-methoxypyrid-2-yl)-1,3-oxazole,
20 2-(3-cyanophenyl)-4-(3-chloro-5-trifluoromethylpyrid-2-yl)-1,3-oxazole,
2-(5-chloro-2-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2-chloro-5-methylthiophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2-bromo-5-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
25 2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole,
2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-nitrophenyl)-4-(2-pyridyl)-1,3-oxazole, or
2-(3-bromophenyl)-4-(2-pyridyl)-1,3-oxazole; or
a pharmaceutically acceptable salt thereof.
30 2. The compound of claim 1, wherein Ar² is substituted with one or more moieties selected from the group consisting of CF₃, Cl, F, Br, CH₃, SCH₃, and CN.

3. The compound of claim 1 or claim 2, wherein Ar^1 is substituted with one or more moieties selected from the group consisting of CF_3 , F, and Cl.

5 4. A compound of claim 1, said compound being selected from the group consisting of 3-(2-pyridyl)-5-(3,5-dichlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-methylphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,5-difluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3,5-difluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,3-dichlorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chloro-5-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-fluoro-5-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-chloro-5-fluorophenyl)-1,2,4-oxadiazole, 3-(5-chloropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(5-fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(5-fluoropyrid-2-yl)-5-(3-cyano-5-fluorophenyl)-1,2,4-oxadiazole, 3-(3-fluoropyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(3-chloro-5-trifluoromethylpyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(5-chloro-2-methoxyphenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2-chloro-5-methylthiophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,5,6-trifluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(2,5,6-trifluorophenyl)-1,2,4-oxadiazole, 3-(2-pyridyl)-5-(3-bromophenyl)-1,2,4-oxadiazole and pharmaceutically acceptable salts thereof.

5. A pharmaceutical composition comprising a compound selected from:

a) a compound of Formula II:


(II)

25 wherein X and Y are N,
 Z is O;
 Ar^1 is 2-pyridyl;
 Ar^2 is phenyl;

wherein at least one of Ar^1 and Ar^2 is substituted with one or more moieties

30 selected from the group consisting of -F, -Cl, -Br, -I, -SR, -SOR, - SO_2R , - $\text{SO}_2\text{NRR}'$, -

OCOR, -OCONRR', -NRCOR', -NRCO₂R', -CN, -CO₂R, -CONRR', -C(O)R, -CH(OR)R', -CH₂(OR), CF₃, C₁-C₁₀ alkyl, cycloalkyl, aryl-alkyl, heterocycloalkyl and aryl;

5 wherein R or R' is selected from the group consisting of H, CF₃, C₁-C₁₀ alkyl, cycloalkyl, alkyl-aryl, heterocycloalkyl, aryl and where R and R' may combine to form a ring,

with the proviso that the compound is not 3-(2-Pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole, or 3-(2-Pyridyl)-5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazole; or

b) 3-(2-pyridyl)-5-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazole,

10 3-(2-pyridyl)-5-(2,3-difluorophenyl)-1,2,4-oxadiazole,
 3-(2-pyridyl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole,
 3-(5-fluoropyrid-2-yl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole,
 3-(5-methoxypyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole,
 3-(2-pyridyl)-5-(3-nitrophenyl)-1,2,4-oxadiazole,
 15 2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole,
 2-(3,5-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(2-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 20 2-(3-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-methylphenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-trifluoromethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(2,3-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(2,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 25 2-(3,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3,5-dimethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(2,3-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-chloro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
 30 2-(3-fluoro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-chloro-5-fluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-cyanophenyl)-4-(5-chloropyrid-2-yl)-1,3-oxazole,
 2-(3-cyanophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,

2-(3-cyano-5-fluorophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(3-fluoropyrid-2-yl)-1,3-oxazole,
2-(3,5-dimethoxyphenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(5-methoxypyrid-2-yl)-1,3-oxazole,
5 2-(3-cyanophenyl)-4-(3-chloro-5-trifluoromethylpyrid-2-yl)-1,3-oxazole,
2-(5-chloro-2-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2-chloro-5-methylthiophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2-bromo-5-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
10 2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole,
2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-nitrophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-bromophenyl)-4-(2-pyridyl)-1,3-oxazole, or
a pharmaceutically acceptable salt thereof;
15 and a pharmaceutically acceptable excipient.

6. The composition of claim 5, wherein Ar² is substituted with one or more
moieties selected from the group consisting of CF₃, Cl, F, Br, CH₃, CN, and SCH₃.

20 7. The composition of claim 5 or claim 6, wherein Ar¹ is substituted with one or
more moieties selected from the group consisting of CF₃, F, and Cl.

8. Use of a pharmaceutically acceptable amount of a compound selected from:
a) a compound of Formula II:

25

(II)

wherein X and Y are N,
Z is O;
Ar¹ is 2-pyridyl;
Ar² is phenyl;

wherein at least one of Ar¹ and Ar² is substituted with one or more moieties selected from the group consisting of -F, -Cl, -Br, -I, -SR, -SOR, -SO₂R, -SO₂NRR', -OCOR, -OCONRR', -NRCOR', -NRCO₂R', -CN, -CO₂R, -CONRR', -C(O)R, -CH(OR)R', -CH₂(OR), and -R;

5 wherein R or R' is selected from the group consisting of H, CF₃, C₁-C₁₀ alkyl, cycloalkyl, alkyl-aryl, heterocycloalkyl, aryl and where R and R' may combine to form a ring,

with the proviso that the compound is not 3-(2-Pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole, or 3-(2-Pyridyl)-5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazole; or

10 b) 3-(2-pyridyl)-5-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazole,
 3-(2-pyridyl)-5-(2,3-difluorophenyl)-1,2,4-oxadiazole,
 3-(2-pyridyl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole,
 3-(5-fluoropyrid-2-yl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole,
 3-(5-methoxypyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole,
15 3-(2-pyridyl)-5-(3-nitrophenyl)-1,2,4-oxadiazole,
 2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole,
 2-(3,5-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
20 2-(2-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-methylphenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-trifluoromethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(2,3-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
25 2-(2,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3,5-dimethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(2,3-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
30 2-(3-chloro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-fluoro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
 2-(3-chloro-5-fluorophenyl)-4-(2-pyridyl)-1,3-oxazole,

2-(3-cyanophenyl)-4-(5-chloropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyano-5-fluorophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(3-fluoropyrid-2-yl)-1,3-oxazole,
5 2-(3,5-dimethoxyphenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(5-methoxypyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(3-chloro-5-trifluoromethylpyrid-2-yl)-1,3-oxazole,
2-(5-chloro-2-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2-chloro-5-methylthiophenyl)-4-(2-pyridyl)-1,3-oxazole,
10 2-(2-bromo-5-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole,
2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-nitrophenyl)-4-(2-pyridyl)-1,3-oxazole, or
15 2-(3-bromophenyl)-4-(2-pyridyl)-1,3-oxazole;


with the proviso that the compound is not 3-(2-Pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole, for the preparation of a medicament for the treatment of diseases associated with metabotropic glutamate receptors.

20 9. The use according to claim 8, wherein Ar² is substituted with one or more moieties selected from the group consisting of CF₃, Cl, F, Br, CH₃, CN, and SCH₃.

10. The use according to claim 8 or claim 9, wherein Ar¹ is substituted with one or more moieties selected from the group consisting of CF₃, F, and Cl.

25 11. A method of treating diseases associated with metabotropic glutamate receptors comprising administering to a patient in need thereof a pharmaceutically acceptable amount of a compound selected from:

a) a compound of Formula II:



wherein X and Y are N,

5 Z is O;

Ar¹ is 2-pyridyl;

Ar² is phenyl;

wherein at least one of Ar¹ and Ar² is substituted with one or more moieties selected from the group consisting of -F, -Cl, -Br, -I, -SR, -SOR, -SO₂R, -SO₂NRR', -OCOR, -OCONRR', -NRCOR', -NRCO₂R', -CN, -CO₂R, -CONRR', -C(O)R, -CH(OR)R', -CH₂(OR), and -R;

wherein R or R' is selected from the group consisting of H, CF₃, C₁-C₁₀ alkyl, cycloalkyl, alkyl-aryl, heterocycloalkyl, aryl and where R and R' may combine to form a ring,

15 with the proviso that the compound is not 3-(2-Pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole, or 3-(2-Pyridyl)-5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazole; or

b) 3-(2-pyridyl)-5-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazole,

3-(2-pyridyl)-5-(2,3-difluorophenyl)-1,2,4-oxadiazole,

3-(2-pyridyl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole,

20 3-(5-fluoropyrid-2-yl)-5-(3,5-dimethoxyphenyl)-1,2,4-oxadiazole,

3-(5-methoxypyrid-2-yl)-5-(3-cyanophenyl)-1,2,4-oxadiazole,

3-(2-pyridyl)-5-(3-nitrophenyl)-1,2,4-oxadiazole,

2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole,

2-(3,5-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole,

25 2-(3-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole,

2-(3-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,

2-(2-chlorophenyl)-4-(2-pyridyl)-1,3-oxazole,

2-(3-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,

2-(3-methylphenyl)-4-(2-pyridyl)-1,3-oxazole,

30 2-(3-trifluoromethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,

2-(2,3-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3,5-difluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
5 2-(3,5-dimethoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2,3-dichlorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-chloro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-fluoro-5-cyanophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-chloro-5-fluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
10 2-(3-cyanophenyl)-4-(5-chloropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyano-5-fluorophenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(3-fluoropyrid-2-yl)-1,3-oxazole,
2-(3,5-dimethoxyphenyl)-4-(5-fluoropyrid-2-yl)-1,3-oxazole,
15 2-(3-cyanophenyl)-4-(5-methoxypyrid-2-yl)-1,3-oxazole,
2-(3-cyanophenyl)-4-(3-chloro-5-trifluoromethylpyrid-2-yl)-1,3-oxazole,
2-(5-chloro-2-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2-chloro-5-methylthiophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(2-bromo-5-methoxyphenyl)-4-(2-pyridyl)-1,3-oxazole,
20 2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-[3-chlorophenyl]-4-[pyridin-2-yl]-1,3-oxazole,
2-(2,5,6-trifluorophenyl)-4-(2-pyridyl)-1,3-oxazole,
2-(3-nitrophenyl)-4-(2-pyridyl)-1,3-oxazole, or
2-(3-bromophenyl)-4-(2-pyridyl)-1,3-oxazole;
25 with the proviso that the compound is not 3-(2-Pyridyl)-5-(2-chlorophenyl)-1,2,4-oxadiazole.

12. The method of claim 11, wherein Ar² is substituted with one or more moieties selected from the group consisting of CF₃, Cl, F, Br, CH₃, CN, and SCH₃.

30 13. The method of claim 11 or claim 12 wherein Ar¹ is substituted with one or more moieties selected from the group consisting of CF₃, F, and Cl.

14. The method of any one of claims 11 to 13, wherein the disease associated with metabotropic glutamate receptors is a neurological disease or disorder.

5 15. The method of any one of claims 11 to 13, wherein the disease associated with metabotropic glutamate receptors is a psychiatric disease.

10 16. The method of any one of claims 11 to 13, wherein the disease or disorder is selected from the group consisting of stroke, head trauma, anoxic injury, ischemic injury, hypoglycemia, epilepsy, pain, migraine headaches, Parkinson's disease, senile dementia, Huntington's Chorea and Alzheimer's disease.

15 17. The method of any one of claims 11 to 13, wherein the disease or disorder is selected from the group consisting of schizophrenia and depression.

18. Compound of claim 1, pharmaceutical composition of claim 5, use of claim 8 or method of claim 11, substantially as herein described with reference to the accompanying Examples and/or Figures.

Dated this 29th day of September 2004
20 NPS PHARMACEUTICALS, INC.
By their Patent Attorneys
GRIFFITH HACK
Fellows Institute of Patent and
Trade Mark Attorneys of Australia

FIGURE 1

















Compound #	Structure	Compound #	Structure
B1		B9	
B2		B10	
B3		B11	
B4		B12	
B5		B13	
B6		B15	
B7		B16	
B8		B17	

FIGURE 1

Compound #	Structure	Compound #	Structure
B18		B26	
B19		B27	
B20		B28	
B21		B29	
B22		B30	
B23		B31	
B24		B32	
B25		B33	

FIGURE 1
























Compound #	Structure	Compound #	Structure
B34		B42	
B35		B43	
B36		B44	
B37		B45	
B38		B46	
B39		B47	
B40		B48	
B41		B49	

FIGURE 1

Compound #	Structure	Compound #	Structure
B50			
B51			
B52			
B53			
B54			
B55			
B56			

4/4