




(86) Date de dépôt PCT/PCT Filing Date: 2016/03/04
(87) Date publication PCT/PCT Publication Date: 2016/09/15
(45) Date de délivrance/Issue Date: 2021/01/05
(85) Entrée phase nationale/National Entry: 2017/08/23
(86) N° demande PCT/PCT Application No.: US 2016/020947
(87) N° publication PCT/PCT Publication No.: 2016/144778
(30) Priorité/Priority: 2015/03/06 (US62/129,074)

(51) Cl.Int./Int.Cl. *A23K 10/30* (2016.01)
(72) Inventeur/Inventor:
SMALLWOOD, NORMAN J., US
(73) Propriétaire/Owner:
SMALLWOOD, NORMAN J., US
(74) Agent: TORYS LLP

(54) Titre : ADDITION DE MATIERE GRASSE COMESTIBLE (LIPIDES) A DU FOIN POUR AMELIORER SA VALEUR NUTRITIONNELLE ET DIGESTIBILITE ET REDUIRE LE RISQUE DE TOXICITE
(54) Title: ADDITION OF EDIBLE FAT (LIPIDS) TO HAY FOR ENHANCING THE NUTRITIONAL VALUE AND DIGESTIBILITY AND REDUCING THE TOXICITY RISK

Proposed Method for Spent Edible Oil Bleaching Earth Utilization



(57) **Abrégé/Abstract:**

A method for improving the nutritional value of hay used for livestock feed is disclosed. The methods includes the application of one or more lipids to the hay prior to feeding to animals. Additional components may also be added such as vitamins, minerals, nitrogen, amino acids and the like.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2016/144778 A3

(43) International Publication Date
15 September 2016 (15.09.2016)

WIPO | PCT

(51) International Patent Classification:

A23K 1/14 (2006.01) A23K 1/16 (2006.01)
A23K 1/18 (2006.01)

(21) International Application Number:

PCT/US2016/020947

(22) International Filing Date:

4 March 2016 (04.03.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/129,074 6 March 2015 (06.03.2015) US

(72) Inventor; and

(71) Applicant : SMALLWOOD, Norman J. [US/US]; 2601 Marsh Lane, Unit 303, Plano, Texas 75093 (US).

(74) Agents: NEBEL, Heidi S. et al.; McKee, Voorhees & Sease, P.L.C., 801 Grand Avenue, Suite 3200, Des Moines, Iowa 50309-2721 (US).

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(88) Date of publication of the international search report:

3 November 2016

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

(54) Title: ADDITION OF EDIBLE FAT (LIPIDS) TO HAY FOR ENHANCING THE NUTRITIONAL VALUE AND DIGESTIBILITY

Proposed Method for Spent Edible Oil Bleaching Earth Utilization




FIG. 2

(57) Abstract: A method for improving the nutritional value of hay used for livestock feed is disclosed. The methods includes the application of one or more lipids to the hay prior to feeding to animals. Additional components may also be added such as vitamins, minerals, nitrogen, amino acids and the like.

WO 2016/144778 A3

**TITLE: ADDITION OF EDIBLE FAT (LIPIDS) TO HAY FOR ENHANCING
THE NUTRITIONAL VALUE AND DIGESTIBILITY
AND REDUCING THE TOXICITY RISK**

5

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority under 35 U.S.C. § 119 to provisional application
Serial No. 62/129,074 filed March 6, 2015.

10 **FIELD OF THE INVENTION**

The present invention relates to nutritional value and digestibility improvement
and toxicity reduction of hay for livestock feed by the addition of edible lipids.

BACKGROUND OF THE INVENTION

15 Hay is the term covering grass, legumes, or other herbaceous plants that have
been cut, dried, baled or loose stacked, and stored for use as livestock feed. Hay is feed to
livestock when growing plants for grazing are not available. The nutritional value,
digestibility and health risk of hay varies by type and handling conditions.

Nutritionally, hay is low in metabolizable energy and essential fatty acids. Protein
20 level is low with the exception of legume hay like alfalfa. All kinds of hay have a high
fiber content.

Hay-fed livestock are exposed to several health risks. Most kinds of hay are
inadequate in essential fatty acid content. Hay exposed to moisture in the handling steps
after drying results in the growth of mold. Respiratory impairment from breathing mold
25 dust in the hay is often a problem. With the fiber content including rough stems, the
digestibility of hay is troublesome. Digestive track inflammation is problematic, especially
for young animals.

It is common to feed hay to livestock hay when pasture grass is not available. Hay
alone is nutritionally deficient due to the inadequate content of metabolizable energy from
30 lipids. The lipid deficiency results in insufficient essential fatty acid intake. In most cases,
this deficiency is not addressed by providing supplemental feed containing the optimum
quantity of lipids. It is well-known that animal growth and fertility is limited by the level of
nutrients in the feed available for consumption.

SUMMARY OF THE INVENTION

The present invention provides a novel method for treating hay with lipids to improve animal nutrition, provide essential fatty acids for health, reduce animal digestive system irritation and inflammation, and ameliorate the mold-dust health problem inherent with hay. Application of the invention offers an attractive financial benefit from animal weight gain and health-problem avoidance.

Lipid treatment of hay not only improves the nutritional value, but ameliorates the rough texture and mold problems. The lubrication and fiber softening effect of lipids adhering to the hay-fiber surface area and penetrating into the stems and leaves significantly improves digestibility. Irritation and inflammation of the digestive system are reduced.

Any of a number of methods may be used to incorporate lipids into hay. A liquid state is required for the best application of lipids. In the case of phospholipids, achieving a fluid state is usually impractical due to the high viscosity. For hay treatment with phospholipids, there are fewer options for application.

Edible lipids for livestock feed comprise both animal and vegetable sources. It includes all states of lipids from crude to fully refined that are classified as edible for animals. Lipid degradation products like free fatty acids, monoglycerides and diglycerides classified as edible are used in animal feed. Complex lipids like phospholipids are included.

Lipids applied to hay soften and lubricate the rough hay fiber, including stems that reduce the irritation and inflammation in the animal digestive system. With the exception of those classified as emulsifiers (monoglycerides, diglycerides and phospholipids), lipids applied to properly dried hay during the baling or stacking operations provide moisture protection against bio-gradation and mold growth. Hay treatment by any kind of lipid before feeding to livestock minimizes mold dust that causes respiratory problems.

The quantity and type of lipid added to the hay can be determined by considering the type of hay, quality of hay including the moisture content, storage conditions provided for the hay before feeding, the weight gain objective for the animals being fed, and the financial benefit to be gained. For treatment of hay immediately before feeding, the

quantity of lipid added can best determined by what is needed to control the mold dust. The minimum amount of lipid added should be not less than 2% by weight.

Lipid selection decisions are dictated by economics, availability, and cultural and/or religious restrictions. Lipid melting point is another consideration. High melting
5 point lipids require heating to liquefy before application to the hay.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a flow chart illustrating the current method for feeding hay to livestock.

10 Figure 2 is a flow chart illustrating the general method of treating hay with lipids for feeding livestock.

Figure 3 is a flow chart illustrating specific options for treating hay with lipids.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

15 The embodiments of this invention are not limited to particular hay (forage) types and composition and lipid compositions and methods of use thereof, which can vary and are understood by skilled artisans. It is further to be understood that all terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting in any manner or scope. For example, as used in this specification and the
20 appended claims, the singular forms "a," "an" and "the" can include plural referents unless the content clearly indicates otherwise. Further, all units, prefixes, and symbols may be denoted in their SI accepted form. Numeric ranges recited within the specification are inclusive of the numbers defining the range and include each integer within the defined range.

25 So that the present invention may be more readily understood, certain terms are first defined. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which embodiments of the invention pertain. Many methods and materials similar, modified, or equivalent to those described herein can be used in the practice of the embodiments of the
30 present invention without undue experimentation, the preferred materials and methods are described herein. In describing and claiming the embodiments of the present invention, the following terminology will be used in accordance with the definitions set out below. The

term "about," as used herein, refers to variation in the numerical quantity that can occur, for example, through typical measuring and liquid handling procedures used for making concentrates or use solutions in the real world; through inadvertent error in these procedures; through differences in the manufacture, source, or purity of the ingredients
5 used to make the compositions or carry out the methods; and the like. The term "about" also encompasses amounts that differ due to different equilibrium conditions for a composition resulting from a particular initial mixture. Whether or not modified by the term "about", the claims include equivalents to the quantities.

The terms "weight percent," "wt-%," "wt %" "percent by weight," "% by weight,"
10 and variations thereof, as used herein, refer to the concentration of a substance as the weight of that substance divided by the total weight of the composition and multiplied by 100. It is understood that, as used here, "percent," "%," and the like are intended to be synonymous with "weight percent," "wt-%," etc.

Ingredients

15 Hay

Commonly used plants for hay include mixtures of grasses such as ryegrass (*Lolium* species), timothy, brome, fescue, Bermuda, orchard grass, and other species, depending on region. Hay may also include legumes, such as alfalfa (lucerne) and clovers (red, white and subterranean). Legumes in hay are ideally cut pre-bloom. Other pasture
20 forbs are also sometimes a part of the mix, though these plants are not necessarily desired as certain forbs are toxic to some animals.

Oat, barley, and wheat plant materials are occasionally cut green and made into hay for animal fodder; however they are more usually used in the form of straw, a harvest byproduct where the stems and dead leaves are baled after the grain has been harvested and
25 threshed. Straw is used mainly for animal bedding. Although straw is also used as fodder, particularly as a source of dietary fibers, it has lower nutritional value than hay.

Lipids

Suitable lipid sources for the composition of the present disclosure may be any known or used in the art, including but not limited to, animal sources, e.g., milk fat, butter,
30 butter fat, egg yolk lipid; marine sources, such as fish oils, marine oils, single cell oils; vegetable and plant oils, such as corn oil, canola oil, sunflower oil, soybean oil, palm olein

oil, coconut oil, high oleic sunflower oil, evening primrose oil, rapeseed oil, olive oil, flaxseed (linseed) oil, cottonseed oil, high oleic safflower oil, palm stearin, palm kernel oil, wheat germ oil; medium chain triglyceride oils and emulsions and esters of fatty acids; and any combinations thereof. Examples include, but are not limited to, fatty acids (e.g.,

5 stearic, palmitic, oleic, linoleic, and lauric acid), complex lipids (e.g., phospholipids), and monoglycerides and diglycerides. Sources of edible fats may include, but are not limited to, coconut oil, corn oil, cottonseed oil, fish oil, olive oil, palm oil, sesame oil, soybean oil, canola oil, sunflower seed oil, tallow, greases, beef fat, restaurant fats, and mixtures thereof.

10

Optional Ingredients

Lipids are excellent carriers for some feed supplements including vitamins. Examples of such vitamins include, but are not limited to, vitamins A, E, K, and the B group vitamins.

15

Optionally, dietary nitrogen may be included in the lipids. Optional dietary nitrogen sources include, but are not limited to, ammonia, ammonium polyphosphate, animal protein products, oilseed meals, synthetic amino acids, and urea.

Optionally, various vitamins may be added to the mixture.

Optionally, various trace minerals and elements may be added to the mixture.

20

Examples of such trace minerals and elements include, but are not limited to, cobalt sulfate, copper sulfate, ferrous sulfate, ferrous oxide, iodines, manganese sulfate, potassium iodate, selenium and its compounds, sulphur, zinc oxide, and zinc sulfate, etc.

Optionally, various drugs, medicaments, insecticides, enzymes, antimicrobials, probiotics and the like may be added to the mixture.

25

Treatment of Hay with Lipids

The preferred and most beneficial method for treating hay with lipids is by use of a spray system installed on each baler used. Spray nozzles are installed at the optimum location to uniformly distribute the lipid to the hay on entry to the baler. The nozzle type,

30 number and location is determined on the basis of the specific bailer design. Lipid is fed to the nozzles by means of piping and hose from a pump driven by a variable speed motor. The motor speed is adjusted to deliver the specified quantity of lipid to the hay. A lipid

supply tank attached to the baler or the tractor pulling the baler is piped to the feed pump suction. This method is not applicable to lipids classified as emulsifiers like phosphatides, monoglycerides and diglycerides.

For loose hay that is stacked, lipid is sprayed on top of each layer of hay added
5 (about 18-inch-thick layer). The spray system may be a handheld spray nozzle or series of nozzles mounted on a suspended rack that can be posited above the haystack as layers of hay are added. A supply tank, piping, hose, pump with variable-speed drive are provided at the stacking site to feed lipid to the spraying nozzle(s). This method is not applicable to lipids classified as emulsifiers like phosphatides, monoglycerides and diglycerides.

10 The advantage of treating hay with non-emulsifying lipids immediately after cutting and drying is to waterproof the hay during the time it is exposed to rain or snow. This minimizes bio-degradation and mold growth. The additional advantage is long exposure of the lipids to the hay for the softening effect to improve digestibility.

Baled or loose hay can be treated with lipids just prior to feeding. Lipid can be
15 applied to baled or loose hay by standard spray apparatus. To avoid lipid waste, each bale or lot of loose hay should be placed in a large pan during the spraying operation. Lipid draining of the hay collects in the pan for reuse. This method is not applicable to the phosphatide class of lipids because of the high viscosity.

Lipid can be injected into round bales by inserting a sparge pipe into the center
20 axis. Using a high pressure pump, lipid is distributed in the hay uniformly through the sparge pipe nozzles. The inserted end of the sparge pipe is closed with a conical cap to facilitate penetration. This method is not applicable to the phosphatide class of lipids because of the high viscosity.

For round hay bales, lipid can be applied, by rotating the bale in a pan containing
25 the specified quantity of lipid. After the specified quantity of lipid has been absorbed by the rotating bale, the bale is lifted up above the pan and placed on a drainage rack. Lipid draining from the bale flows back to the pan for use in the next application. This method is applicable to the phosphatide class of lipids that have a high viscosity.

Square (rectangular) bales can be treated with lipid by emersion in a vat of lipid
30 for sufficient time to absorb the specified quantity. The treated bale is lifted out of the vat and placed on a rack to enable free lipid to drain back into the vat for further use. This method is applicable to the phosphatide class of lipids that have a high viscosity.

For high-melting-point lipids, heat exchange capability must be included in the treatment systems to liquefy the lipid before use.

Livestock animals

5 Any grazing animal that is fed hay can be fed the animal feed according to the invention, this includes animals such as cattle, horses, goats, and sheep. This may also include smaller animals such as rabbits and guinea pigs. The hay may be fed at any time during the animal's life and in any amount sufficient for traditional grazing applications.

10 **EXAMPLE 1**

To validate the invention, feeding tests were conducted using 10 six-month old weaned calves over a period of 60 days. Five calves were fed hay without lipid treatment. Five calves were fed with hay from the same source with treatment of 3% by weight lipid (PBSY cottonseed oil). All of the calves were given with the same salt and mineral supply.

15 The results of the test is outlined as follows:

UNTREATED-HAY FEEDING RESULTS

Average Daily Gain (ADG): 2.1 pounds

Body Condition Score (BCS): Moderate (5)

20

LIPID-TREATED-HAY FEEDING RESULTS

Average Daily Gain (ADG): 2.6 pounds

Body Condition Score (BCS): Very Good (7)

Lipid-Feed Conversion Ratio: 1 to 0.93

25

Claims


1. A method for improving the nutritional value and reducing the health risk of hay for livestock feed comprising:
treating a hay component with a treatment composition comprising a
lipid component prior to feeding to the livestock;
wherein the lipid component is present in a minimum amount of about 2% by weight;
wherein the amount of the hay component is at a maximum of about 98% by weight; and
wherein the hay component is in the form of loose hay, a hay stack, and/or a hay bale.
2. The method of claim 1, wherein the hay component includes one or more of the following: grass, legumes, pasture forbs, oats, barley, wheat, and straw.
3. The method of claim 2, wherein the grass includes one or more of: ryegrass, timothy, brome, fescue, Bermuda, and orchard grass.
4. The method of claim 2, wherein the legumes include one or more of the following: alfalfa, and clover.
5. The method of claim 4, wherein the legumes are cut pre-bloom.
6. The method or claim 2, wherein the oat, barley, and wheat are cut green and then dried.
7. The method of claim 2, wherein the straw has been harvested.
8. The method of any one of claims 1-7, wherein the lipid component includes one or more of the following: fatty acids, complex lipids, monoglycerides, diglycerides, and edible fats.
9. The method of claim 8, wherein the lipid component is an edible fat.
10. The method of claim 9, wherein the edible fat includes one or more of the following: coconut oil, com oil, cottonseed oil, fish oil, olive oil, palm oil, sesame oil, soybean oil, canola oil, sunflower seed oil, tallow, greases, beef fat, and mixtures thereof.


11. The method of claim 8, wherein the lipid component is a fatty acid.
12. The method of claim 11, wherein the fatty acid includes one or more of: stearic acid, palmitic acid, oleic acid, linoleic acid, and lauric acid.
13. The method of claim 8, where the lipid component is one or more complex lipids.
14. The method of claim 13, wherein the one or more complex lipids are a phospholipid.
15. A composition made by the method of claim 1.
16. A feed composition comprising:
lipid treated hay, wherein the feed composition comprises about 2% by weight of a
lipid component and about 98% by weight of a hay component; and
wherein the hay component is in the form of loose hay, a hay stack, and/or a hay bale.
17. The feed composition of claim 16, further comprising a vitamin.
18. The feed composition of claim 17, wherein the vitamin is one or more of vitamin A, vitamin E, vitamin K, and the B group vitamins.
19. The feed composition of any one of claims 16-18, further comprising nitrogen.
20. The feed composition of claim 19, wherein nitrogen is from a source of one or more of ammonia, ammonium polyphosphate, animal protein products, oilseed meals, synthetic amino acids, and urea.
21. The feed composition of any one of claims 16-20, further comprising minerals, elements, or a combination thereof.
22. The feed composition of claim 21, where the minerals and/or elements include one or more of: cobalt sulfate, copper sulfate, ferrous sulfate, ferrous oxide, iodines, manganese sulfate, potassium iodate, selenium and its compounds, sulphur, zinc oxide, and zinc

sulfate.

23. The feed composition of any one of claims 16-22, further comprising one or more of the following: drugs, medicaments, insecticides, enzymes, antimicrobials, probiotics.
24. A method for producing a hay feed product with improved nutritional value and reduced health risks comprising:
treating hay with a treatment composition comprising a lipid component,
wherein the hay feed product comprises about 2% by weight of the lipid component and
about 98% by weight of the hay; and
wherein the hay component is in the form of loose hay, a hay stack, and/or a hay bale.
25. The method of claim 24, wherein the treating comprises spraying loose hay with the treatment composition by using a spraying system.
26. The method of any one of claims 24-25, wherein the treating comprises spraying the hay component with the treatment composition during hay stacking.
27. The method of any one of claims 24-26, wherein the treating comprises spraying the hay component with the treatment composition during baling of hay.
28. The method of any one of claims 24-27, wherein the treating comprises applying the treatment composition to hay bales, wherein the treatment composition is applied to the hay bales individually.
29. The method of claim 28, wherein the treating comprises injecting the hay bales with the treatment composition by using a sparge pipe inserted in a hay bale.
30. The method of claim 28, wherein the treating comprises rotating the hay bales, partially submerged, in an open container containing the treatment composition.
31. The method of claim 28, wherein the treating comprises immersing the hay bales in a vat filled with the treatment composition.

32. Use of a lipid treated animal feed for improvement of livestock health, wherein the lipid treated animal feed comprises about 2% by weight of a lipid component and about 98% by weight hay;
wherein the hay is in the form of loose hay, a hay stack, and/or a hay bale.
33. The use of claim 32, wherein the livestock is an animal selected from the group comprising: cattle, horses, goats, and sheep.

*1/4*Current Method for Feeding Hay to Livestock*FIG. 1*


*2/4*Proposed Method for Spent Edible Oil Bleaching Earth Utilization*FIG. 2*

3/4


Flow Chart Illustrating Specific Options for Treating Hay with Lipids

Treatment at During Hay Harvest

1, Spray lipid on hay as baled:




2, Spray lipid on hay as stacked



Treatment of Hay Before Feeding

1. Spray lipid on bales or loose hay



2. Inject Lipid into baled hay by inserting a sparge pipe





FIG. 3A

4/4


Flow Chart Illustrating Specific Options for Treating Hay with Lipids – Page 2

3. Apply lipid by rotating hay bale in open-top container (pan) containing product:

Round Day Bale supported by spindles on both ends
and rotated partially immersed in an open-top pan containing lipid.



4. Apply lipid by emersion of hay bale into a vat containing the product



Hay bale fully immersed in lipid. Immersion time is provided to achieve the proper quantity of lipid absorption. The hay bale is supported by a perforated metal basket connected to the ho

FIG. 3B

Proposed Method for Spent Edible Oil Bleaching Earth Utilization

