




(86) Date de dépôt PCT/PCT Filing Date: 2014/02/28
(87) Date publication PCT/PCT Publication Date: 2014/09/04
(45) Date de délivrance/Issue Date: 2021/02/09
(85) Entrée phase nationale/National Entry: 2015/08/31
(86) N° demande PCT/PCT Application No.: US 2014/019622
(87) N° publication PCT/PCT Publication No.: 2014/134554
(30) Priorités/Priorities: 2013/03/01 (US61/771,534);
2013/03/01 (US61/771,642); 2013/06/26 (US61/839,753)

(51) Cl.Int./Int.Cl. A61K 38/07 (2006.01),
A61P 3/00 (2006.01)
(72) Inventeurs/Inventors:
WILSON, D. TRAVIS, US;
BAMBERGER, MARK, US
(73) Propriétaire/Owner:
STEALTH BIOTHERAPEUTICS CORP, MC
(74) Agent: OSLER, HOSKIN & HARCOURT LLP

(54) Titre : METHODES ET COMPOSITIONS DE PREVENTION OU DE TRAITEMENT DU SYNDROME DE BARTH
(54) Title: METHODS AND COMPOSITIONS FOR THE PREVENTION OR TREATMENT OF BARTH SYNDROME



(57) Abrégé/Abstract:

The disclosure provides methods of preventing or treating Barth Syndrome in a mammalian subject, reducing risk factors associated with Barth Syndrome, and/or reducing the likelihood or severity of Barth Syndrome. The methods comprise administering to the subject an effective amount of an aromatic-cationic peptide to increase expression of TAZ1 in subjects in need thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2014/134554 A1

(43) International Publication Date
4 September 2014 (04.09.2014)(51) International Patent Classification:
A61K 38/06 (2006.01)

(74) Agents: VAVRA, Stephanie H. et al.; Foley & Lardner LLP, 3000 K Street N.W., Suite 600, Washington, District of Columbia 20007-5109 (US).

(21) International Application Number:

PCT/US2014/019622

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:

28 February 2014 (28.02.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/771,642	1 March 2013 (01.03.2013)	US
61/771,534	1 March 2013 (01.03.2013)	US
61/839,753	26 June 2013 (26.06.2013)	US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW).

(71) Applicant: STEALTH PEPTIDES INTERNATIONAL, INC. [MC/MC]; 2nd Floor, Le Prince de Galles, 3-5 Avenue des Citronniers, MC-98000 Mc (MC).


(72) Inventor; and

(71) Applicant : WILSON, D. Travis [US/US]; 22 Brewster Road, Newton, Massachusetts 02461 (US).

(72) Inventor: BAMBERGER, Mark; 257 Old Farms Road, South Glastonbury, Connecticut 06073 (US).

[Continued on next page]

(54) Title: METHODS AND COMPOSITIONS FOR THE PREVENTION OR TREATMENT OF BARTH SYNDROME

FIG. 1**Cardiolipin 18:2-18:2-18:2-18:2
nmol/mg non-collagen protein**

(57) Abstract: The disclosure provides methods of preventing or treating Barth Syndrome in a mammalian subject, reducing risk factors associated with Barth Syndrome, and/or reducing the likelihood or severity of Barth Syndrome. The methods comprise administering to the subject an effective amount of an aromatic-cationic peptide to increase expression of TAZ1 in subjects in need thereof.

WO 2014/134554 A1



TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, — *as to the applicant's entitlement to claim the priority of KM, ML, MR, NE, SN, TD, TG).*

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))* — **Published:** *with international search report (Art. 21(3))*

METHODS AND COMPOSITIONS FOR THE PREVENTION OR TREATMENT OF BARTH SYNDROME

TECHNICAL FIELD

[0002] The present technology relates generally to compositions and methods for preventing or treating Barth Syndrome, reducing risk factors associated with Barth Syndrome, and/or reducing the severity of Barth Syndrome. In particular, the present technology relates to administering an effective amount of an aromatic-cationic peptide to a subject in need thereof to normalize expression levels of TAZ1.

BACKGROUND

[0003] The following description is provided to assist the understanding of the reader. None of the information provided or references cited is admitted to be prior art to the present invention.

[0004] Barth Syndrome is a heritable disorder of phospholipid metabolism characterized by dilated cardiomyopathy (DCM), skeletal myopathy, neutropenia, growth delay and organic aciduria. The prevalence of Barth Syndrome is estimated at 1/454,000 live births, with an estimated incidence ranging from 1/400,000 to 1/140,000 depending on geographic location. Barth Syndrome is an X-linked disorder, and so disproportionately affects male patients.

[0005] Barth Syndrome is caused by mutations in the TAZ gene (tafazzin; Xq28), which encodes TAZ1, an acyltransferase involved in the metabolism of cardiolipin, a phospholipid localized to the inner mitochondrial membrane. Defective TAZ1 function results in abnormal remodeling of cardiolipin and compromises mitochondrial structure and respiratory chain function.

SUMMARY

[0006] In one aspect, the present disclosure provides a method for treating or preventing Barth Syndrome in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of the peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ or a pharmaceutically acceptable salt thereof.

[0007] In some embodiments, the subject displays reduced levels of TAZ1 expression compared to a normal control subject. In some embodiments, the peptide is administered daily for 6 weeks or more. In some embodiments, the peptide is administered daily for 12 weeks or more.

[0008] In some embodiments, the subject has been diagnosed as having Barth Syndrome. In some embodiments, the Barth Syndrome comprises one or more of cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and frequent bacterial infections.

[0009] In some embodiments, the subject is human. In some embodiments, the peptide is administered orally, topically, systemically, intravenously, subcutaneously, intraperitoneally, or intramuscularly.

[0010] In some embodiments, the method further comprises separately, sequentially or simultaneously administering a cardiovascular agent to the subject. In some embodiments, the cardiovascular agent is selected from the group consisting of: an anti-arrhythmia agent, a vasodilator, an anti-anginal agent, a corticosteroid, a cardiotrope, a diuretic, a sedative, an angiotensin converting enzyme (ACE) inhibitor, an angiotensin II antagonist, a thrombolytic agent, a calcium channel blocker, a thromboxane receptor antagonist, a radical scavenger, an anti-platelet drug, a β -adrenaline receptor blocking drug, α -receptor blocking drug, a sympathetic nerve inhibitor, a digitalis formulation, an inotrope, and an antihyperlipidemic drug.

[0011] In some embodiments, the pharmaceutically acceptable salt comprises acetate or trifluoroacetate salt.

[0012] In one aspect, the present disclosure provides a method for increasing the expression of TAZ1 in a mammalian subject in need thereof, the method comprising:

administering to the subject a therapeutically effective amount of the peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ or a pharmaceutically acceptable salt thereof.

[0013] In some embodiments, the expression of TAZ1 in the subject is about 2-5 fold less than the level of TAZ1 expression in a normal control subject. In some embodiments, the peptide is administered daily for 6 weeks or more. In some embodiments, the peptide is administered daily for 12 weeks or more.

[0014] In some embodiments, the subject has been diagnosed has having, is suspected of having, or is at risk of having Barth Syndrome. In some embodiments, the Barth Syndrome comprises one or more of cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and frequent bacterial infections.

[0015] In some embodiments, the subject is human. In some embodiments, the peptide is administered orally, topically, systemically, intravenously, subcutaneously, intraperitoneally, or intramuscularly

[0016] In some embodiments, the method further comprises separately, sequentially or simultaneously administering a cardiovascular agent to the subject. In some embodiments, the cardiovascular agent is selected from the group consisting of: an anti-arrhythmia agent, a vasodilator, an anti-anginal agent, a corticosteroid, a cardiotrope, a diuretic, a sedative, an angiotensin converting enzyme (ACE) inhibitor, an angiotensin II antagonist, a thrombolytic agent, a calcium channel blocker, a thromboxane receptor antagonist, a radical scavenger, an anti-platelet drug, a β -adrenaline receptor blocking drug, α -receptor blocking drug, a sympathetic nerve inhibitor, a digitalis formulation, an inotrope, and an antihyperlipidemic drug.

[0017] In some embodiments, the pharmaceutically acceptable salt comprises acetate or trifluoroacetate salt.

[0018] In one aspect, the present disclosure provides a method for reducing the risk of Barth Syndrome in a mammalian subject having decreased expression of TAZ1 compared to a normal control subject, the method comprising: administering to the subject a therapeutically effective amount of the peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ or a pharmaceutically acceptable salt thereof.

[0019] In one aspect, the present disclosure provides a method for stabilizing cardiolipin remodeling in a mammalian subject having or suspected of having Barth Syndrome. In some embodiments, the mammalian subject has decreased expression of TAZ1 compared to a normal control subject. In some embodiments, the cardiolipin is 18:2 species of cardiolipin.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] FIG. 1 is a chart showing the effects of D-Arg-2'6'-Dmt-Lys-Phe-NH₂ on levels of cardiolipin species 18:2-18:2-18:2-18:2 in a dog heart failure model.

[0021] FIG. 2 is a chart showing the effects of D-Arg-2'6'-Dmt-Lys-Phe-NH₂ on levels of TAZ1 expression in a dog heart failure model.

[0022] FIG. 3 is an electron microscopy image of mitochondria in a Barth Syndrome patient.

[0023] FIG. 4A is an electron microscopy image of the ultrastructure of mitochondria in cardiac disease.

[0024] FIG. 4B is an electron microscopy image of the ultrastructure of mitochondria in cardiac disease treated with D-Arg-2'6'-Dmt-Lys-Phe-NH₂.

[0025] FIG. 5A is an electron microscopy image of the organization of mitochondria in cardiac disease.

[0026] FIG. 5B is an electron microscopy image of the organization of mitochondria in cardiac disease treated with D-Arg-2'6'-Dmt-Lys-Phe-NH₂.

DETAILED DESCRIPTION

[0027] It is to be appreciated that certain aspects, modes, embodiments, variations and features of the invention are described below in various levels of detail in order to provide a substantial understanding of the present invention. The definitions of certain terms as used in this specification are provided below. Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0028] As used in this specification and the appended claims, the singular forms “a”, “an” and “the” include plural referents unless the content clearly dictates otherwise. For example, reference to “a cell” includes a combination of two or more cells, and the like.

[0029] As used herein, the “administration” of an agent, drug, or peptide to a subject includes any route of introducing or delivering to a subject a compound to perform its intended function. Administration can be carried out by any suitable route, including orally, intranasally, parenterally (intravenously, intramuscularly, intraperitoneally, or subcutaneously), or topically. Administration includes self-administration and the administration by another.

[0030] As used herein, the term “amino acid” includes naturally-occurring amino acids and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally-occurring amino acids. Naturally-occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, *e.g.*, hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally-occurring amino acid, *i.e.*, an α -carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, *e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (*e.g.*, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally-occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally-occurring amino acid. Amino acids can be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission.

[0031] As used herein, the term “effective amount” refers to a quantity sufficient to achieve a desired therapeutic and/or prophylactic effect, *e.g.*, an amount which results in an increase in (*e.g.*, normalization of) the expression level of *e.g.*, TAZ1 in a subject in need thereof. In the context of therapeutic or prophylactic applications, in some embodiments, the amount of a composition administered to the subject will depend on the type and severity of the disease and on the characteristics of the individual, such as general health, age, sex, body weight and tolerance to drugs. In some embodiments, it will also

depend on the degree, severity and type of disease. The skilled artisan will be able to determine appropriate dosages depending on these and other factors. The compositions can also be administered in combination with one or more additional therapeutic compounds. In the methods described herein, aromatic-cationic peptides, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, may be administered to a subject having one or more signs, symptoms, or risk factors of Barth Syndrome, such as, *e.g.*, cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and/or frequent bacterial infections, such as pneumonia. For example, a “therapeutically effective amount” of the aromatic-cationic peptides includes levels at which a subject’s levels of TAZ1 expression are increased after administration, and/or at which the presence, frequency, or severity of one or more signs, symptoms, or risk factors of Barth Syndrome are reduced or eliminated. In some embodiments, a therapeutically effective amount reduces or ameliorates the physiological effects of a Barth Syndrome, and/or the risk factors of Barth Syndrome, and/or the likelihood of developing Barth Syndrome.

[0032] As used herein, the term “Barth Syndrome” refers to a heritable disorder of phospholipid metabolism caused by deficiencies of the TAZ1 acyltransferase. Signs and Symptoms of Barth Syndrome include, but are not limited to, cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and/or frequent bacterial infections, such as pneumonia.

[0033] As used herein, the term “TAZ1” or “tafazzin” refers to the human X chromosome acyltransferase encoded by the *TAZ* gene. Illustrative sequences of TAZ1 isoforms are given by, for example, GenBank Accession Numbers NM_000116.3, NM_181311.2, NM_181312.2, and NM_181313.2.

[0034] As used herein, “isolated” or “purified” polypeptide or peptide refers to a polypeptide or peptide that is substantially free of cellular material or other contaminating polypeptides from the cell or tissue source from which the agent is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. For example, an isolated aromatic-cationic peptide would be free of materials that would interfere with diagnostic or therapeutic uses of the agent. Such

interfering materials may include enzymes, hormones and other proteinaceous and nonproteinaceous solutes.

[0035] As used herein, “normalizing” a subject’s levels of TAZ1 expression refers to altering the subject’s levels of TAZ1 expression in the direction of “normal” or wild-type expression levels. For example, normalizing TAZ1 expression levels in a subject with reduced TAZ1 expression compared to a normal subject refers to increasing the levels of TAZ1 expression. In some embodiments, normalizing TAZ1 expression in a subject refers to attenuating or reducing the degree of reduced TAZ1 expression compared to *e.g.*, an untreated control subject.

[0036] As used herein “increasing” a subject’s TAZ1 expression level means increasing the level of TAZ1 in the subject (*e.g.*, a subject’s TAZ1 expression level such as RNA and/or protein level) in an organ or tissue.. In some embodiments, increasing TAZ1 expression level is an increase by about 1%, about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, or more. Alternatively, or additionally, in some embodiments, increasing TAZ1 expression level is measured as an attenuation or reduction in the extent to which TAZ1 expression is decreased in a subject. In some embodiments, the TAZ1 reduction is decreased about 0.25 fold to about 0.5 fold, about 0.5 fold to about 0.75 fold, about 0.75 fold to about 1.0 fold, or about 1.0 fold to about 1.5 fold.

[0037] As used herein, the terms “polypeptide,” “peptide,” and “protein” are used interchangeably herein to mean a polymer comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, *i.e.*, peptide isosteres. Polypeptide refers to both short chains, commonly referred to as peptides, glycopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. Polypeptides include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques that are well known in the art.

[0038] As used herein, the term “simultaneous” therapeutic use refers to the administration of at least two active ingredients by the same route and at the same time or at substantially the same time.

[0039] As used herein, the term “separate” therapeutic use refers to an administration of at least two active ingredients at the same time or at substantially the same time by different routes.

[0040] As used herein, the term “sequential” therapeutic use refers to administration of at least two active ingredients at different times, the administration route being identical or different. More particularly, sequential use refers to the whole administration of one of the active ingredients before administration of the other or others commences. It is thus possible to administer one of the active ingredients over several minutes, hours, or days before administering the other active ingredient or ingredients. There is no simultaneous treatment in this case.

[0041] As used herein, the terms “treating” or “treatment” or “alleviation” refers to therapeutic treatment, wherein the object is to prevent, reduce, alleviate or slow down (lessen) the targeted pathologic condition or disorder. A subject is successfully “treated” for Barth Syndrome if, after receiving a therapeutic amount of the aromatic-cationic peptides, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, according to the methods described herein, the subject shows observable and/or measurable reduction in or absence of one or more signs and symptoms of Barth Syndrome, such as, *e.g.*, cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and/or frequent bacterial infections, such as pneumonia. It is also to be appreciated that the various modes of treatment or prevention of medical conditions as described are intended to mean “substantial,” which includes total but also less than total treatment or prevention, and wherein some biologically or medically relevant result is achieved. Treating Barth Syndrome, as used herein, also refers to treating reduced TAZ1 expression levels characteristic of the Syndrome, thereby causing an increase in TAZ1 expression compared to the subject’s level of TAZ1 expression prior to treatment.

[0042] As used herein, “prevention” or “preventing” of a disorder or condition refers to a compound that, in a statistical sample, reduces the occurrence of symptoms of a disorder or condition in the treated sample relative to an untreated control sample, or delays the onset or reduces the severity of one or more symptoms of the disorder or condition relative to the untreated control sample. As used herein, preventing Barth Syndrome includes

preventing or delaying the initiation of, preventing, delaying, or slowing the progression or advancement of, and/or reversing the progression of Barth Syndrome. As used herein, prevention of Barth Syndrome also includes preventing a recurrence of one or more signs or symptoms of Barth Syndrome.

Aromatic-Cationic Peptides

[0043] The present technology relates to methods and compositions for preventing or treating Barth Syndrome in a subject in need thereof. In some embodiments, the methods and compositions prevent one or more signs or symptoms of Barth Syndrome in a subject. In some embodiments, the methods and compositions increase the level of TAZ1 expression in a subject. In some embodiments, the methods and compositions reduce the likelihood that a subject with risk factors for Barth Syndrome will develop one or more signs or symptoms of Barth Syndrome.

[0044] The aromatic-cationic peptides are water-soluble and highly polar. Despite these properties, the peptides can readily penetrate cell membranes. The aromatic-cationic peptides typically include a minimum of three amino acids or a minimum of four amino acids, covalently joined by peptide bonds. The maximum number of amino acids present in the aromatic-cationic peptides is about twenty amino acids covalently joined by peptide bonds. Suitably, the maximum number of amino acids is about twelve, more preferably about nine, and most preferably about six.

[0045] The amino acids of the aromatic-cationic peptides can be any amino acid. As used herein, the term “amino acid” is used to refer to any organic molecule that contains at least one amino group and at least one carboxyl group. Typically, at least one amino group is at the α position relative to a carboxyl group. The amino acids may be naturally occurring. Naturally occurring amino acids include, for example, the twenty most common levorotatory (L) amino acids normally found in mammalian proteins, *i.e.*, alanine (Ala), arginine (Arg), asparagine (Asn), aspartic acid (Asp), cysteine (Cys), glutamine (Gln), glutamic acid (Glu), glycine (Gly), histidine (His), isoleucine (Ile), leucine (Leu), lysine (Lys), methionine (Met), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr), tryptophan, (Trp), tyrosine (Tyr), and valine (Val). Other naturally occurring amino acids include, for example, amino acids that are synthesized in metabolic processes not associated with protein synthesis. For example, the amino acids ornithine and citrulline

are synthesized in mammalian metabolism during the production of urea. Another example of a naturally occurring amino acid includes hydroxyproline (Hyp).

[0046] The peptides optionally contain one or more non-naturally occurring amino acids. Optimally, the peptide has no amino acids that are naturally occurring. The non-naturally occurring amino acids may be levorotatory (L-), dextrorotatory (D-), or mixtures thereof. Non-naturally occurring amino acids are those amino acids that typically are not synthesized in normal metabolic processes in living organisms, and do not naturally occur in proteins. In addition, the non-naturally occurring amino acids suitably are also not recognized by common proteases. The non-naturally occurring amino acid can be present at any position in the peptide. For example, the non-naturally occurring amino acid can be at the N-terminus, the C-terminus, or at any position between the N-terminus and the C-terminus.

[0047] The non-natural amino acids may, for example, comprise alkyl, aryl, or alkylaryl groups not found in natural amino acids. Some examples of non-natural alkyl amino acids include α -aminobutyric acid, β -aminobutyric acid, γ -aminobutyric acid, δ -aminovaleric acid, and ε -aminocaproic acid. Some examples of non-natural aryl amino acids include ortho-, meta, and para-aminobenzoic acid. Some examples of non-natural alkylaryl amino acids include ortho-, meta-, and para-aminophenylacetic acid, and γ -phenyl- β -aminobutyric acid. Non-naturally occurring amino acids include derivatives of naturally occurring amino acids. The derivatives of naturally occurring amino acids may, for example, include the addition of one or more chemical groups to the naturally occurring amino acid.

[0048] For example, one or more chemical groups can be added to one or more of the 2', 3', 4', 5', or 6' position of the aromatic ring of a phenylalanine or tyrosine residue, or the 4', 5', 6', or 7' position of the benzo ring of a tryptophan residue. The group can be any chemical group that can be added to an aromatic ring. Some examples of such groups include branched or unbranched C₁-C₄ alkyl, such as methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, or t-butyl, C₁-C₄ alkyloxy (*i.e.*, alkoxy), amino, C₁-C₄ alkylamino and C₁-C₄ dialkylamino (*e.g.*, methylamino, dimethylamino), nitro, hydroxyl, halo (*i.e.*, fluoro, chloro, bromo, or iodo). Some specific examples of non-naturally occurring derivatives of naturally occurring amino acids include norvaline (Nva) and norleucine (Nle).

[0049] Another example of a modification of an amino acid in a peptide is the derivatization of a carboxyl group of an aspartic acid or a glutamic acid residue of the peptide. One example of derivatization is amidation with ammonia or with a primary or secondary amine, *e.g.* methylamine, ethylamine, dimethylamine or diethylamine. Another example of derivatization includes esterification with, for example, methyl or ethyl alcohol. Another such modification includes derivatization of an amino group of a lysine, arginine, or histidine residue. For example, such amino groups can be acylated. Some suitable acyl groups include, for example, a benzoyl group or an alkanoyl group comprising any of the C₁-C₄ alkyl groups mentioned above, such as an acetyl or propionyl group.

[0050] The non-naturally occurring amino acids are suitably resistant or insensitive to common proteases. Examples of non-naturally occurring amino acids that are resistant or insensitive to proteases include the dextrorotatory (D-) form of any of the above-mentioned naturally occurring L-amino acids, as well as L- and/or D- non-naturally occurring amino acids. The D-amino acids do not normally occur in proteins, although they are found in certain peptide antibiotics that are synthesized by means other than the normal ribosomal protein synthetic machinery of the cell. As used herein, the D-amino acids are considered to be non-naturally occurring amino acids.

[0051] In order to minimize protease sensitivity, the peptides should have less than five, preferably less than four, more preferably less than three, and most preferably, less than two contiguous L-amino acids recognized by common proteases, irrespective of whether the amino acids are naturally or non-naturally occurring. Optimally, the peptide has only D-amino acids, and no L-amino acids. If the peptide contains protease sensitive sequences of amino acids, at least one of the amino acids is preferably a non-naturally-occurring D-amino acid, thereby conferring protease resistance. An example of a protease sensitive sequence includes two or more contiguous basic amino acids that are readily cleaved by common proteases, such as endopeptidases and trypsin. Examples of basic amino acids include arginine, lysine and histidine.

[0052] The aromatic-cationic peptides should have a minimum number of net positive charges at physiological pH in comparison to the total number of amino acid residues in the peptide. The minimum number of net positive charges at physiological pH will be referred to below as (p_m). The total number of amino acid residues in the peptide will be

referred to below as (r). The minimum number of net positive charges discussed below are all at physiological pH. The term “physiological pH” as used herein refers to the normal pH in the cells of the tissues and organs of the mammalian body. For instance, the physiological pH of a human is normally approximately 7.4, but normal physiological pH in mammals may be any pH from about 7.0 to about 7.8.

[0053] “Net charge” as used herein refers to the balance of the number of positive charges and the number of negative charges carried by the amino acids present in the peptide. In this specification, it is understood that net charges are measured at physiological pH. The naturally occurring amino acids that are positively charged at physiological pH include L-lysine, L-arginine, and L-histidine. The naturally occurring amino acids that are negatively charged at physiological pH include L-aspartic acid and L-glutamic acid.

[0054] Typically, a peptide has a positively charged N-terminal amino group and a negatively charged C-terminal carboxyl group. The charges cancel each other out at physiological pH. As an example of calculating net charge, the peptide Tyr-Arg-Phe-Lys-Glu-His-Trp-D-Arg has one negatively charged amino acid (*i.e.*, Glu) and four positively charged amino acids (*i.e.*, two Arg residues, one Lys, and one His). Therefore, the above peptide has a net positive charge of three.

[0055] In one embodiment, the aromatic-cationic peptides have a relationship between the minimum number of net positive charges at physiological pH (p_m) and the total number of amino acid residues (r) wherein $3p_m$ is the largest number that is less than or equal to $r + 1$. In this embodiment, the relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) is as follows:

TABLE 1. Amino acid number and net positive charges ($3p_m \leq p+1$)

(r)	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
(p_m)	1	1	2	2	2	3	3	3	4	4	4	5	5	5	6	6	6	7

[0056] In another embodiment, the aromatic-cationic peptides have a relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) wherein $2p_m$ is the largest number that is less than or equal to $r + 1$. In

this embodiment, the relationship between the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) is as follows:

TABLE 2. Amino acid number and net positive charges ($2p_m \leq p+1$)

(r)	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
(p_m)	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9	9	10	10

[0057] In one embodiment, the minimum number of net positive charges (p_m) and the total number of amino acid residues (r) are equal. In another embodiment, the peptides have three or four amino acid residues and a minimum of one net positive charge, suitably, a minimum of two net positive charges and more preferably a minimum of three net positive charges.

[0058] It is also important that the aromatic-cationic peptides have a minimum number of aromatic groups in comparison to the total number of net positive charges (p_t). The minimum number of aromatic groups will be referred to below as (a). Naturally occurring amino acids that have an aromatic group include the amino acids histidine, tryptophan, tyrosine, and phenylalanine. For example, the hexapeptide Lys-Gln-Tyr-D-Arg-Phe-Trp has a net positive charge of two (contributed by the lysine and arginine residues) and three aromatic groups (contributed by tyrosine, phenylalanine and tryptophan residues).

[0059] The aromatic-cationic peptides should also have a relationship between the minimum number of aromatic groups (a) and the total number of net positive charges at physiological pH (p_t) wherein 3a is the largest number that is less than or equal to $p_t + 1$, except that when p_t is 1, a may also be 1. In this embodiment, the relationship between the minimum number of aromatic groups (a) and the total number of net positive charges (p_t) is as follows:

TABLE 3. Aromatic groups and net positive charges ($3a \leq p_t+1$ or $a = p_t = 1$)

(p_t)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
(a)	1	1	1	1	2	2	2	3	3	3	4	4	4	5	5	5	6	6	6	7

[0060] In another embodiment, the aromatic-cationic peptides have a relationship between the minimum number of aromatic groups (a) and the total number of net positive charges (p_t) wherein 2a is the largest number that is less than or equal to $p_t + 1$. In this

embodiment, the relationship between the minimum number of aromatic amino acid residues (a) and the total number of net positive charges (p_t) is as follows:

TABLE 4. Aromatic groups and net positive charges (2a ≤ p_t+1 or a= p_t=1)

(p _t)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
(a)	1	1	2	2	3	3	4	4	5	5	6	6	7	7	8	8	9	9	10	10

[0061] In another embodiment, the number of aromatic groups (a) and the total number of net positive charges (p_t) are equal.

[0062] Carboxyl groups, especially the terminal carboxyl group of a C-terminal amino acid, are suitably amidated with, for example, ammonia to form the C-terminal amide. Alternatively, the terminal carboxyl group of the C-terminal amino acid may be amidated with any primary or secondary amine. The primary or secondary amine may, for example, be an alkyl, especially a branched or unbranched C₁-C₄ alkyl, or an aryl amine. Accordingly, the amino acid at the C-terminus of the peptide may be converted to an amido, N-methylamido, N-ethylamido, N, N-dimethylamido, N, N-diethylamido, N-methyl-N-ethylamido, N-phenylamido or N-phenyl-N-ethylamido group. The free carboxylate groups of the asparagine, glutamine, aspartic acid, and glutamic acid residues not occurring at the C-terminus of the aromatic-cationic peptides may also be amidated wherever they occur within the peptide. The amidation at these internal positions may be with ammonia or any of the primary or secondary amines described above.

[0063] In one embodiment, the aromatic-cationic peptide is a tripeptide having two net positive charges and at least one aromatic amino acid. In a particular embodiment, the aromatic-cationic peptide is a tripeptide having two net positive charges and two aromatic amino acids.

[0064] Aromatic-cationic peptides include, but are not limited to, the following peptide examples:

TABLE 5: EXEMPLARY PEPTIDES

2',6'-Dmp-D-Arg-2',6'-Dmt-Lys-NH ₂
2',6'-Dmp-D-Arg-Phe-Lys-NH ₂
2',6'-Dmt-D-Arg-PheOrn-NH ₂
2',6'-Dmt-D-Arg-Phe-Ahp(2-aminoheptanoicacid)-NH ₂
2',6'-Dmt-D-Arg-Phe-Lys-NH ₂

2',6'-Dmt-D-Cit-PheLys-NH ₂
Ala-D-Phe-D-Arg-Tyr-Lys-D-Trp-His-D-Tyr-Gly-Phe
Arg-D-Leu-D-Tyr-Phe-Lys-Glu-D-Lys-Arg-D-Trp-Lys-D-Phe-Tyr-D-Arg-Gly
Asp-Arg-D-Phe-Cys-Phe-D-Arg-D-Lys-Tyr-Arg-D-Tyr-Trp-D-His-Tyr-D-Phe-Lys-Phe
Asp-D-Trp-Lys-Tyr-D-His-Phe-Arg-D-Gly-Lys-NH ₂
D-Arg-2',6'-Dmt-Lys-Phe-NH ₂
D-Glu-Asp-Lys-D-Arg-D-His-Phe-Phe-D-Val-Tyr-Arg-Tyr-D-Tyr-Arg-His-Phe-NH ₂
D-His-Glu-Lys-Tyr-D-Phe-Arg
D-His-Lys-Tyr-D-Phe-Glu-D-Asp-D-Asp-D-His-D-Lys-Arg-Trp-NH ₂
D-Tyr-Trp-Lys-NH ₂
Glu-Arg-D-Lys-Tyr-D-Val-Phe-D-His-Trp-Arg-D-Gly-Tyr-Arg-D-Met-NH ₂
Gly-Ala-Lys-Phe-D-Lys-Glu-Arg-Tyr-His-D-Arg-D-Arg-Asp-Tyr-Trp-D-His-Trp-His-D-Lys-Asp.
Gly-D-Phe-Lys-His-D-Arg-Tyr-NH ₂
His-Tyr-D-Arg-Trp-Lys-Phe-D-Asp-Ala-Arg-Cys-D-Tyr-His-Phe-D-Lys-Tyr-His-Ser-NH ₂
Lys-D-Arg-Tyr-NH ₂
Lys-D-Gln-Tyr-Arg-D-Phe-Trp-NH ₂
Lys-Trp-D-Tyr-Arg-Asn-Phe-Tyr-D-His-NH ₂
Met-Tyr-D-Arg-Phe-Arg-NH ₂
Met-Tyr-D-Lys-Phe-Arg
Phe-Arg-D-His-Asp
Phe-D-Arg-2',6'-Dmt-Lys-NH ₂
Phe-D-Arg-His
Phe-D-Arg-Lys-Trp-Tyr-D-Arg-His
Phe-D-Arg-Phe-Lys-NH ₂
Phe-Phe-D-Tyr-Arg-Glu-Asp-D-Lys-Arg-D-Arg-His-Phe-NH ₂
Phe-Tyr-Lys-D-Arg-Trp-His-D-Lys-D-Lys-Glu-Arg-D-Tyr-Thr
Thr-Gly-Tyr-Arg-D-His-Phe-Trp-D-His-Lys
Thr-Tyr-Arg-D-Lys-Trp-Tyr-Glu-Asp-D-Lys-D-Arg-His-Phe-D-Tyr-Gly-Val-Ile-D-His-Arg-Tyr-Lys-NH ₂
Trp-D-Lys-Tyr-Arg-NH ₂
Trp-Lys-Phe-D-Asp-Arg-Tyr-D-His-Lys
Tyr-Asp-D-Lys-Tyr-Phe-D-Lys-D-Arg-Phe-Pro-D-Tyr-His-Lys
Tyr-D-Arg-Phe-Lys-Glu-NH ₂
Tyr-D-Arg-Phe-Lys-NH ₂
Tyr-D-His-Phe-D-Arg-Asp-Lys-D-Arg-His-Trp-D-His-Phe
Tyr-His-D-Gly-Met
Val-D-Lys-His-Tyr-D-Phe-Ser-Tyr-Arg-NH ₂

[0065] In one embodiment, the peptides have mu-opioid receptor agonist activity (*i.e.*, they activate the mu-opioid receptor). Peptides, which have mu-opioid receptor agonist activity, are typically those peptides that have a tyrosine residue or a tyrosine derivative at the N-terminus (*i.e.*, the first amino acid position). Suitable derivatives of tyrosine include 2'-methyltyrosine (Mmt); 2', 6'-dimethyltyrosine (2'6'-Dmt); 3', 5'-dimethyltyrosine (3'5'Dmt); N, 2', 6'-trimethyltyrosine (Tmt); and 2'-hydroxy-6'-methyltryosine (Hmt).

[0066] In one embodiment, a peptide that has mu-opioid receptor agonist activity has the formula Tyr-D-Arg-Phe-Lys-NH₂. Tyr-D-Arg-Phe-Lys-NH₂ has a net positive charge of three, contributed by the amino acids tyrosine, arginine, and lysine and has two aromatic groups contributed by the amino acids phenylalanine and tyrosine. The tyrosine of Tyr-D-Arg-Phe-Lys-NH₂ can be a modified derivative of tyrosine such as in 2', 6'-dimethyltyrosine to produce the compound having the formula 2', 6'-Dmt-D-Arg-Phe-Lys-NH₂. 2', 6'-Dmt-D-Arg-Phe-Lys-NH₂ has a molecular weight of 640 and carries a net three positive charge at physiological pH. 2', 6'-Dmt-D-Arg-Phe-Lys-NH₂ readily penetrates the plasma membrane of several mammalian cell types in an energy-independent manner (Zhao, *et al.*, *J. Pharmacol Exp Ther.*, 304:425-432, 2003).

[0067] Alternatively, in other instances, the aromatic-cationic peptide does not have mu-opioid receptor agonist activity. For example, during long-term treatment, such as in a chronic disease state or condition, the use of an aromatic-cationic peptide that activates the mu-opioid receptor may be contraindicated. In these instances, the potentially adverse or addictive effects of the aromatic-cationic peptide may preclude the use of an aromatic-cationic peptide that activates the mu-opioid receptor in the treatment regimen of a human patient or other mammal. Potential adverse effects may include sedation, constipation and respiratory depression. In such instances an aromatic-cationic peptide that does not activate the mu-opioid receptor may be an appropriate treatment. Peptides that do not have mu-opioid receptor agonist activity generally do not have a tyrosine residue or a derivative of tyrosine at the N-terminus (*i.e.*, amino acid position 1). The amino acid at the N-terminus can be any naturally occurring or non-naturally occurring amino acid other than tyrosine. In one embodiment, the amino acid at the N-terminus is phenylalanine or its derivative. Exemplary derivatives of phenylalanine include 2'-methylphenylalanine (Mmp), 2', 6'-dimethylphenylalanine (2', 6'-Dmp), N, 2', 6'-trimethylphenylalanine (Tmp), and 2'-hydroxy-6'-methylphenylalanine (Hmp).

[0068] An example of an aromatic-cationic peptide that does not have mu-opioid receptor agonist activity has the formula Phe-D-Arg-Phe-Lys-NH₂. Alternatively, the N-terminal phenylalanine can be a derivative of phenylalanine such as 2', 6'-dimethylphenylalanine (2'6'-Dmp). Tyr-D-Arg-Phe-Lys-NH₂ containing 2', 6'-dimethylphenylalanine at amino acid position 1 has the formula 2', 6'-Dmp-D-Arg-Phe-Lys-NH₂. In one embodiment, the amino acid sequence of 2', 6'-Dmt-D-Arg-Phe-Lys-

NH₂ is rearranged such that Dmt is not at the N-terminus. An example of such an aromatic-cationic peptide that does not have mu-opioid receptor agonist activity has the formula D-Arg-2'6'-Dmt-Lys-Phe-NH₂.

[0069] Suitable substitution variants of the peptides listed herein include conservative amino acid substitutions. Amino acids may be grouped according to their physicochemical characteristics as follows:

- (a) Non-polar amino acids: Ala(A) Ser(S) Thr(T) Pro(P) Gly(G) Cys (C);
- (b) Acidic amino acids: Asn(N) Asp(D) Glu(E) Gln(Q);
- (c) Basic amino acids: His(H) Arg(R) Lys(K);
- (d) Hydrophobic amino acids: Met(M) Leu(L) Ile(I) Val(V); and
- (e) Aromatic amino acids: Phe (F) Tyr(Y) Trp (W) His (H).

[0070] Substitutions of an amino acid in a peptide by another amino acid in the same group is referred to as a conservative substitution and may preserve the physicochemical characteristics of the original peptide. In contrast, substitutions of an amino acid in a peptide by another amino acid in a different group are generally more likely to alter the characteristics of the original peptide.

[0071] Examples of peptides that activate mu-opioid receptors include, but are not limited to, the aromatic-cationic peptides shown in Table 6.

TABLE 6. Peptide Analogs with Mu-Opioid Activity

Amino Acid Position 1	Amino Acid Position 2	Amino Acid Position 3	Amino Acid Position 4	C-Terminal Modification
Tyr	D-Arg	Phe	Lys	NH ₂
Tyr	D-Arg	Phe	Orn	NH ₂
Tyr	D-Arg	Phe	Dab	NH ₂
Tyr	D-Arg	Phe	Dap	NH ₂
2'6'Dmt	D-Arg	Phe	Lys	NH ₂
2'6'Dmt	D-Arg	Phe	Lys-NH(CH ₂) ₂ -NH-dns	NH ₂
2'6'Dmt	D-Arg	Phe	Lys-NH(CH ₂) ₂ -NH-atn	NH ₂
2'6'Dmt	D-Arg	Phe	dnsLys	NH ₂
2'6'Dmt	D-Cit	Phe	Lys	NH ₂
2'6'Dmt	D-Cit	Phe	Ahp	NH ₂
2'6'Dmt	D-Arg	Phe	Orn	NH ₂
2'6'Dmt	D-Arg	Phe	Dab	NH ₂
2'6'Dmt	D-Arg	Phe	Dap	NH ₂

TABLE 6. Peptide Analogs with Mu-Opioid Activity

Amino Acid Position 1	Amino Acid Position 2	Amino Acid Position 3	Amino Acid Position 4	C-Terminal Modification
2'6'Dmt	D-Arg	Phe	Ahp(2-aminoheptanoic acid)	NH ₂
Bio-2'6'Dmt	D-Arg	Phe	Lys	NH ₂
3'5'Dmt	D-Arg	Phe	Lys	NH ₂
3'5'Dmt	D-Arg	Phe	Orn	NH ₂
3'5'Dmt	D-Arg	Phe	Dab	NH ₂
3'5'Dmt	D-Arg	Phe	Dap	NH ₂
Tyr	D-Arg	Tyr	Lys	NH ₂
Tyr	D-Arg	Tyr	Orn	NH ₂
Tyr	D-Arg	Tyr	Dab	NH ₂
Tyr	D-Arg	Tyr	Dap	NH ₂
2'6'Dmt	D-Arg	Tyr	Lys	NH ₂
2'6'Dmt	D-Arg	Tyr	Orn	NH ₂
2'6'Dmt	D-Arg	Tyr	Dab	NH ₂
2'6'Dmt	D-Arg	Tyr	Dap	NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Lys	NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Orn	NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Dab	NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Dap	NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Arg	NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Lys	NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Orn	NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Dab	NH ₂
Tyr	D-Lys	Phe	Dap	NH ₂
Tyr	D-Lys	Phe	Arg	NH ₂
Tyr	D-Lys	Phe	Lys	NH ₂
Tyr	D-Lys	Phe	Orn	NH ₂
2'6'Dmt	D-Lys	Phe	Dab	NH ₂
2'6'Dmt	D-Lys	Phe	Dap	NH ₂
2'6'Dmt	D-Lys	Phe	Arg	NH ₂
2'6'Dmt	D-Lys	Phe	Lys	NH ₂
3'5'Dmt	D-Lys	Phe	Orn	NH ₂
3'5'Dmt	D-Lys	Phe	Dab	NH ₂
3'5'Dmt	D-Lys	Phe	Dap	NH ₂
3'5'Dmt	D-Lys	Phe	Arg	NH ₂
Tyr	D-Lys	Tyr	Lys	NH ₂
Tyr	D-Lys	Tyr	Orn	NH ₂
Tyr	D-Lys	Tyr	Dab	NH ₂
Tyr	D-Lys	Tyr	Dap	NH ₂
2'6'Dmt	D-Lys	Tyr	Lys	NH ₂
2'6'Dmt	D-Lys	Tyr	Orn	NH ₂
2'6'Dmt	D-Lys	Tyr	Dab	NH ₂

TABLE 6. Peptide Analogs with Mu-Opioid Activity

Amino Acid Position 1	Amino Acid Position 2	Amino Acid Position 3	Amino Acid Position 4	C-Terminal Modification
2'6'Dmt	D-Lys	Tyr	Dap	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Lys	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Orn	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Dab	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Dap	NH ₂
2'6'Dmt	D-Arg	Phe	dnsDap	NH ₂
2'6'Dmt	D-Arg	Phe	atnDap	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Lys	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Orn	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Dab	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Dap	NH ₂
Tyr	D-Lys	Phe	Arg	NH ₂
Tyr	D-Orn	Phe	Arg	NH ₂
Tyr	D-Dab	Phe	Arg	NH ₂
Tyr	D-Dap	Phe	Arg	NH ₂
2'6'Dmt	D-Arg	Phe	Arg	NH ₂
2'6'Dmt	D-Lys	Phe	Arg	NH ₂
2'6'Dmt	D-Orn	Phe	Arg	NH ₂
2'6'Dmt	D-Dab	Phe	Arg	NH ₂
3'5'Dmt	D-Dap	Phe	Arg	NH ₂
3'5'Dmt	D-Arg	Phe	Arg	NH ₂
3'5'Dmt	D-Lys	Phe	Arg	NH ₂
3'5'Dmt	D-Orn	Phe	Arg	NH ₂
Tyr	D-Lys	Tyr	Arg	NH ₂
Tyr	D-Orn	Tyr	Arg	NH ₂
Tyr	D-Dab	Tyr	Arg	NH ₂
Tyr	D-Dap	Tyr	Arg	NH ₂
2'6'Dmt	D-Arg	2'6'Dmt	Arg	NH ₂
2'6'Dmt	D-Lys	2'6'Dmt	Arg	NH ₂
2'6'Dmt	D-Orn	2'6'Dmt	Arg	NH ₂
2'6'Dmt	D-Dab	2'6'Dmt	Arg	NH ₂
3'5'Dmt	D-Dap	3'5'Dmt	Arg	NH ₂
3'5'Dmt	D-Arg	3'5'Dmt	Arg	NH ₂
3'5'Dmt	D-Lys	3'5'Dmt	Arg	NH ₂
3'5'Dmt	D-Orn	3'5'Dmt	Arg	NH ₂
Mmt	D-Arg	Phe	Lys	NH ₂
Mmt	D-Arg	Phe	Orn	NH ₂
Mmt	D-Arg	Phe	Dab	NH ₂
Mmt	D-Arg	Phe	Dap	NH ₂
Tmt	D-Arg	Phe	Lys	NH ₂
Tmt	D-Arg	Phe	Orn	NH ₂
Tmt	D-Arg	Phe	Dab	NH ₂
Tmt	D-Arg	Phe	Dap	NH ₂

TABLE 6. Peptide Analogs with Mu-Opioid Activity

Amino Acid Position 1	Amino Acid Position 2	Amino Acid Position 3	Amino Acid Position 4	C-Terminal Modification
Hmt	D-Arg	Phe	Lys	NH ₂
Hmt	D-Arg	Phe	Orn	NH ₂
Hmt	D-Arg	Phe	Dab	NH ₂
Hmt	D-Arg	Phe	Dap	NH ₂
Mmt	D-Lys	Phe	Lys	NH ₂
Mmt	D-Lys	Phe	Orn	NH ₂
Mmt	D-Lys	Phe	Dab	NH ₂
Mmt	D-Lys	Phe	Dap	NH ₂
Mmt	D-Lys	Phe	Arg	NH ₂
Tmt	D-Lys	Phe	Lys	NH ₂
Tmt	D-Lys	Phe	Orn	NH ₂
Tmt	D-Lys	Phe	Dab	NH ₂
Tmt	D-Lys	Phe	Dap	NH ₂
Tmt	D-Lys	Phe	Arg	NH ₂
Hmt	D-Lys	Phe	Lys	NH ₂
Hmt	D-Lys	Phe	Orn	NH ₂
Hmt	D-Lys	Phe	Dab	NH ₂
Hmt	D-Lys	Phe	Dap	NH ₂
Hmt	D-Lys	Phe	Arg	NH ₂
Mmt	D-Lys	Phe	Arg	NH ₂
Mmt	D-Orn	Phe	Arg	NH ₂
Mmt	D-Dab	Phe	Arg	NH ₂
Mmt	D-Dap	Phe	Arg	NH ₂
Mmt	D-Arg	Phe	Arg	NH ₂
Tmt	D-Lys	Phe	Arg	NH ₂
Tmt	D-Orn	Phe	Arg	NH ₂
Tmt	D-Dab	Phe	Arg	NH ₂
Tmt	D-Dap	Phe	Arg	NH ₂
Tmt	D-Arg	Phe	Arg	NH ₂
Hmt	D-Lys	Phe	Arg	NH ₂
Hmt	D-Orn	Phe	Arg	NH ₂
Hmt	D-Dab	Phe	Arg	NH ₂
Hmt	D-Dap	Phe	Arg	NH ₂
Hmt	D-Arg	Phe	Arg	NH ₂

Dab = diaminobutyric

Dap = diaminopropionic acid

Dmt = dimethyltyrosine

Mmt = 2'-methyltyrosine

Tmt = N, 2',6'-trimethyltyrosine

Hmt = 2'-hydroxy,6'-methyltyrosine

dnsDap = β -dansyl-L- α , β -diaminopropionic acidatnDap = β -anthraniloyl-L- α , β -diaminopropionic acid

Bio = biotin

[0072] Examples of peptides that do not activate mu-opioid receptors include, but are not limited to, the aromatic-cationic peptides shown in Table 7.

TABLE 7. Peptide Analogs Lacking Mu-Opioid Activity

Amino Acid Position 1	Amino Acid Position 2	Amino Acid Position 3	Amino Acid Position 4	C-Terminal Modification
D-Arg	Dmt	Lys	Phe	NH ₂
D-Arg	Dmt	Phe	Lys	NH ₂
D-Arg	Phe	Lys	Dmt	NH ₂
D-Arg	Phe	Dmt	Lys	NH ₂
D-Arg	Lys	Dmt	Phe	NH ₂
D-Arg	Lys	Phe	Dmt	NH ₂
Phe	Lys	Dmt	D-Arg	NH ₂
Phe	Lys	D-Arg	Dmt	NH ₂
Phe	D-Arg	Phe	Lys	NH ₂
Phe	D-Arg	Dmt	Lys	NH ₂
Phe	D-Arg	Lys	Dmt	NH ₂
Phe	Dmt	D-Arg	Lys	NH ₂
Phe	Dmt	Lys	D-Arg	NH ₂
Lys	Phe	D-Arg	Dmt	NH ₂
Lys	Phe	Dmt	D-Arg	NH ₂
Lys	Dmt	D-Arg	Phe	NH ₂
Lys	Dmt	Phe	D-Arg	NH ₂
Lys	D-Arg	Phe	Dmt	NH ₂
Lys	D-Arg	Dmt	Phe	NH ₂
D-Arg	Dmt	D-Arg	Phe	NH ₂
D-Arg	Dmt	D-Arg	Dmt	NH ₂
D-Arg	Dmt	D-Arg	Tyr	NH ₂
D-Arg	Dmt	D-Arg	Trp	NH ₂
Trp	D-Arg	Phe	Lys	NH ₂
Trp	D-Arg	Tyr	Lys	NH ₂
Trp	D-Arg	Trp	Lys	NH ₂
Trp	D-Arg	Dmt	Lys	NH ₂
D-Arg	Trp	Lys	Phe	NH ₂
D-Arg	Trp	Phe	Lys	NH ₂
D-Arg	Trp	Lys	Dmt	NH ₂
D-Arg	Trp	Dmt	Lys	NH ₂
D-Arg	Lys	Trp	Phe	NH ₂
D-Arg	Lys	Trp	Dmt	NH ₂
Cha	D-Arg	Phe	Lys	NH ₂
Ala	D-Arg	Phe	Lys	NH ₂

Cha = cyclohexyl alanine

[0073] The amino acids of the peptides shown in Tables 5-7 may be in either the L- or the D- configuration.

[0074] The peptides may be synthesized by any of the methods well known in the art. Suitable methods for chemically synthesizing the protein include, for example, those described by Stuart and Young in *Solid Phase Peptide Synthesis*, Second Edition, Pierce Chemical Company (1984), and in *Methods Enzymol.*, 289, Academic Press, Inc., New York (1997).

Cardiolipin Remodeling

[0075] Cardiolipin (cardiolipin) is an important component of the inner mitochondrial membrane, where it constitutes about 20% of the total lipid composition. In mammalian cells, cardiolipin is found almost exclusively in the inner mitochondrial membrane where it is essential for the optimal function of enzymes involved in mitochondrial metabolism.

[0076] Cardiolipin is a species of diphosphatidylglycerol lipid comprising two phosphatidylglycerols connected with a glycerol backbone to form a dimeric structure. It has four alkyl groups and potentially carries two negative charges. As there are four distinct alkyl chains in cardiolipin, the molecule has the potential for great complexity. However, in most animal tissues, cardiolipin contains 18-carbon fatty alkyl chains with 2 unsaturated bonds on each of them (18:2). It has been proposed that the 18:2 configuration is an important structural requirement for the high affinity of cardiolipin to inner membrane proteins in mammalian mitochondria. However, studies with isolated enzyme preparations indicate that its importance may vary depending on the protein examined.

[0077] Each of the two phosphates in cardiolipin can capture one proton. Although it has a symmetric structure, ionization of one phosphate happens at different levels of acidity than ionizing both, with $pK1 = 3$ and $pK2 > 7.5$. Hence, under normal physiological conditions (a pH of approximately 7.0), the molecule may carry only one negative charge. Hydroxyl groups ($-OH$ and $-O-$) on the phosphate form stable intramolecular hydrogen bonds, forming a bicyclic resonance structure. This structure traps one proton, which is conducive to oxidative phosphorylation.

[0078] During the oxidative phosphorylation process catalyzed by Complex IV, large quantities of protons are transferred from one side of the membrane to another side

causing a large pH change. Without wishing to be bound by theory, it has been suggested that cardiolipin functions as a proton trap within the mitochondrial membranes, strictly localizing the proton pool and minimizing pH in the mitochondrial intermembrane space. This function is thought to be due to the unique structure of cardiolipin, which, as described above, can trap a proton within the bicyclic structure while carrying a negative charge. Thus, cardiolipin can serve as an electron buffer pool to release or absorb protons to maintain the pH near the mitochondrial membranes.

[0079] In addition, cardiolipin has been shown to play a role in apoptosis. An early event in the apoptosis cascade involves cardiolipin. As discussed in more detail below, a cardiolipin-specific oxygenase produces cardiolipin-hydroperoxides which causes the lipid to undergo a conformational change. The oxidized cardiolipin then translocates from the inner mitochondrial membrane to the outer mitochondrial membrane where it is thought to form a pore through which cytochrome c is released into the cytosol. Cytochrome c can bind to the IP3 receptor stimulating calcium release, which further promotes the release of cytochrome c. When the cytoplasmic calcium concentration reaches a toxic level, the cell dies. In addition, extra-mitochondrial cytochrome c interacts with apoptotic activating factors, causing the formation of apoptosomal complexes and activation of the proteolytic caspase cascade.

[0080] Other roles proposed for cardiolipin are: 1) participation in stabilization of the physical properties of the membrane (Schlame, *et al.*, 2000; Koshkin and Greenberg, 2002; Ma, *et al.*, 2004), for example, membrane fluidity and osmotic stability and 2) participation in protein function via direct interaction with membrane proteins (Schlame, *et al.*, 2000; Palsdottir and Hunte, 2004). Cardiolipin has been found in tight association with inner membrane protein complexes such as the cytochrome *bc1* complex (complex III). As well, it has been localized to the contact sites of dimeric cytochrome c oxidase, and cardiolipin binding sites have also been found in the ADP/ATP carrier (AAC; for review see Palsdottir and Hunte, 2004). Recent work also suggests a role of cardiolipin in formation of respiratory chain supercomplexes (respirasomes).

[0081] The major tetra-acyl molecular species are 18:2 in each of the four fatty acyl positions of the cardiolipin molecule (referred to as the 18:2-18:2-18:2-18:2 cardiolipin species). Remodeling of cardiolipin is essential to obtain this enrichment of cardiolipin with linoleate because cardiolipin synthase has no molecular species substrate specificity for cytidine-5'-diphosphate-1,2-diacyl-sn-glycerol. In addition, the species pattern of

cardiolipin precursors is similar enough to imply that the enzymes of the cardiolipin synthetic pathway are not molecular species-selective. Alterations in the molecular composition of cardiolipin are associated with various disease states.

[0082] Remodeling of cardiolipin occurs via at least three enzymes. Mitochondrial cardiolipin is remodeled by a deacylation-reacylation cycle in which newly synthesized cardiolipin was rapidly deacylated to monolysocardiolipin (MLCL) and then reacylated back to cardiolipin. MLCL AT1 is responsible for the deacylation and ALCAT1 is responsible for the reacylation. In addition to these mitochondrial and microsomal acyltransferase activities, mitochondrial cardiolipin may be remodeled by a mitochondrial cardiolipin transacylase. Tafazzin (TAZ1) is a cardiolipin transacylase that specifically remodels mitochondrial cardiolipin with linoleic acid.

Barth Syndrome

[0083] Barth Syndrome is a heritable disorder of phospholipid metabolism characterized by dilated cardiomyopathy (DCM), skeletal myopathy, neutropenia, growth delay and organic aciduria. The prevalence of Barth Syndrome is estimated at 1/454,000 live births, with an estimated incidence ranging from 1/400,000 to 1/140,000 depending on geographic location. Barth Syndrome is an X-linked disorder, and so disproportionately affects male patients.

[0084] Barth Syndrome is caused by mutations in the TAZ gene (tafazzin; Xq28), which encodes TAZ1, an acyltransferase involved in the metabolism of cardiolipin, a phospholipid localized to the inner mitochondrial membrane. Defective TAZ1 function results in abnormal remodeling of cardiolipin and compromises mitochondrial structure and respiratory chain function. TAZ1 is expressed at high levels in cardiac and skeletal muscle and is involved in the maintenance of the inner membrane of mitochondria. TAZ1 is involved in maintaining levels of cardiolipin, which is essential for energy production in the mitochondria.

[0085] Clinical presentation of Barth Syndrome is highly variable. Most subjects develop DCM during the first decade of life, and typically during the first year of life, which may be accompanied by endocardial fibroelastosis (EFE) and/or left ventricular noncompaction (LVNC). The manifestations of Barth Syndrome may begin in utero, causing cardiac failure, fetal hydrops and miscarriage or stillbirth during the 2nd/3rd

trimester of pregnancy. Ventricular arrhythmia, especially during adolescence, can lead to sudden cardiac death. There is a significant risk of stroke. Skeletal (mostly proximal) myopathy causes delayed motor milestones, hypotonia, severe lethargy or exercise intolerance. There is a tendency to hypoglycemia during the neonatal period. Ninety percent of patients show mild to severe intermittent or persistent neutropenia with a risk of septicemia, severe bacterial sepsis, mouth ulcers and painful gums. Lactic acidosis and mild anemia may occur. Affected boys usually show delayed puberty and growth delay that is observed until the late teens or early twenties, when a substantial growth spurt often occurs. Patients may also present severe difficulties with adequate food intake. Episodic diarrhea is common. Many patients have a similar facial appearance with chubby cheeks, deep-set eyes and prominent ears.

[0086] In some embodiments, treatment with an aromatic-cationic peptide, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, increases the expression of TAZ1 in a tissue or an organ in mammalian subjects that have suffered or are at risk of suffering Barth Syndrome. By way of example, but not by way of limitation, in some embodiments, the level of TAZ1 expression is increased in the myocardium of a subject in need thereof.

[0087] In some embodiments, increasing TAZ1 expression level is measured as a attenuation or reduction in the extent to which TAZ1 expression is decreased in a subject. In some embodiments, the TAZ1 reduction is decreased about 0.25 fold to about 0.5 fold, about 0.5 fold to about 0.75 fold, about 0.75 fold to about 1.0 fold, or about 1.0 fold to about 1.5 fold.

Therapeutic Methods

[0088] The following discussion is presented by way of example only, and is not intended to be limiting.

[0089] It is to be understood that increasing the expression level of TAZ1 in a subject in need thereof (e.g., RNA and/or protein level) will reduce the risk, severity, presentation/onset of any number of negative physical effects. One aspect of the present technology includes methods of treating reduced TAZ1 expression in a subject diagnosed as having, suspected as having, or at risk of having reduced TAZ1 expression levels. One aspect of the present technology includes methods of treating Barth Syndrome in a subject

diagnosed as having, suspected as having, or at risk of having Barth Syndrome. In therapeutic applications, compositions or medicaments are administered to a subject suspected of, or already suffering from such a disease, such as, *e.g.*, decreased TAZ1 expression levels or Barth Syndrome, in an amount sufficient to cure, or at least partially arrest, the symptoms of the disease, including its complications and intermediate pathological phenotypes in development of the disease.

[0090] Subjects suffering from decreased TAZ1 expression levels or Barth Syndrome can be identified by any or a combination of diagnostic or prognostic assays known in the art. For example, typical symptoms of Barth Syndrome include symptoms such as, *e.g.*, cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and/or frequent bacterial infections, such as pneumonia. In some embodiments, the subject may exhibit reduced levels of TAZ1 expression compared to a normal subject, which is measureable using techniques known in the art. In some embodiments, the subject may exhibit one or more mutations in the *TAZ* gene associated with Barth Syndrome, which are detectable using techniques known in the art.

Prophylactic Methods

[0091] In one aspect, the present technology provides a method for preventing or delaying the onset of Barth Syndrome or symptoms of Barth Syndrome in a subject at risk of having reduced levels of TAZ1 expression compared to a normal subject. In some embodiments, the subject may exhibit one or more mutations in the *TAZ* gene associated with Barth Syndrome, which are detectable using techniques known in the art. Subjects at risk for reduced TAZ1 expression levels or Barth Syndrome can be identified by, *e.g.*, any or a combination of diagnostic or prognostic assays known in the art. In prophylactic applications, pharmaceutical compositions or medicaments of aromatic-cationic peptides, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, are administered to a subject susceptible to, or otherwise at risk of a disease or condition such as *e.g.*, Barth Syndrome, in an amount sufficient to eliminate or reduce the risk, lessen the severity, or delay the onset of the disease, including biochemical, histologic and/or behavioral symptoms of the disease, its complications and intermediate pathological phenotypes presenting during development of the disease. Administration of a prophylactic aromatic-cationic can occur prior to the

manifestation of symptoms characteristic of the disease or disorder, such that symptoms of the disease or disorder is prevented or, alternatively, delayed in its progression.

[0092] Subjects or at risk for reduced TAZ1 expression levels or Barth Syndrome may exhibit one or more of the following non-limiting risk factors: cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and/or frequent bacterial infections, such as pneumonia.

Determination of the Biological Effect of the Aromatic-Cationic Peptide-Based Therapeutic

[0093] In various embodiments, suitable *in vitro* or *in vivo* assays are performed to determine the effect of a specific aromatic-cationic peptide-based therapeutic and whether its administration is indicated for treatment. In various embodiments, *in vitro* assays can be performed with representative animal models, to determine if a given aromatic-cationic peptide-based therapeutic exerts the desired effect increasing TAZ1 expression, and preventing or treating Barth Syndrome. Compounds for use in therapy can be tested in suitable animal model systems including, but not limited to rats, mice, chicken, cows, monkeys, rabbits, and the like, prior to testing in human subjects. Similarly, for *in vivo* testing, any of the animal model system known in the art can be used prior to administration to human subjects. In some embodiments, *in vitro* or *in vivo* testing is directed to the biological function of D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt.

[0094] Heart failure has been induced in different species with volume overload, pressure overload, fast pacing, myocardial ischemia, cardiotoxic drugs, or genetically modified models. Hypertension is associated with an increased risk for the development of heart failure. In one mouse model, angiotensin II (Ang II) increases blood pressure and induces cardiomyocyte hypertrophy, increased cardiac fibrosis, and impaired cardiomyocyte relaxation. Infusion of angiotensin to mice by mini osmotic pump increases systolic and diastolic blood pressure, increases heart weight and left ventricular thickness (LVMI), and impaired myocardial performance index (MPI). TAZ1 expression levels are monitored at various time points before, during and after heart failure induction.

[0095] In a second illustrative mouse model, sustained high level expression of G α q can lead to marked myocyte apoptosis, resulting in cardiac hypertrophy and Heart failure by 16 weeks of age (D'Angelo, *et al.*, 1998). The β -adrenergic receptors (β ARs) are primarily coupled to the heterotrimeric G protein, Gs, to stimulate adenylyl cyclase activity. This association generates intracellular cAMP and protein kinase A activation, which regulate cardiac contractility and heart rate. Overexpression of G α q leads to decreased responsiveness to β -adrenergic agonists and results in heart failure. TAZ1 expression levels are monitored at various time points before, during and after heart failure induction.

[0096] Experimental constriction of the aorta by surgical ligation is also widely used as a model of heart failure. Transaortic constriction (TAC) results in pressure overload induced heart failure, with increase in left ventricular (LV) mass. TAC is performed as described by Tamavski O, *et al.* (2004) using a 7-0 silk double-knot suture to constrict the ascending aorta. After TAC, mice develop heart failure within a period of 4 weeks. TAZ1 expression levels are monitored at various time points before, during and after heart failure induction.

Modes of Administration and Effective Dosages

[0097] Any method known to those in the art for contacting a cell, organ or tissue with an aromatic-cationic peptide of the present technology, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, may be employed. Suitable methods include *in vitro*, *ex vivo*, or *in vivo* methods. *In vivo* methods typically include the administration of an aromatic-cationic peptide, such as those described above, to a mammal, suitably a human. When used *in vivo* for therapy, the aromatic-cationic peptides, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, are administered to the subject in effective amounts (*i.e.*, amounts that have desired therapeutic effect). The dose and dosage regimen will depend upon the degree of the infection in the subject, the characteristics of the particular aromatic-cationic peptide used, *e.g.*, its therapeutic index, the subject, and the subject's history.

[0098] The effective amount may be determined during pre-clinical trials and clinical trials by methods familiar to physicians and clinicians. An effective amount of a peptide useful in the methods may be administered to a mammal in need thereof by any of a

number of well-known methods for administering pharmaceutical compounds. The peptide may be administered systemically or locally.

[0099] The peptide may be formulated as a pharmaceutically acceptable salt. The term "pharmaceutically acceptable salt" means a salt prepared from a base or an acid which is acceptable for administration to a patient, such as a mammal (e.g., salts having acceptable mammalian safety for a given dosage regime). However, it is understood that the salts are not required to be pharmaceutically acceptable salts, such as salts of intermediate compounds that are not intended for administration to a patient. Pharmaceutically acceptable salts can be derived from pharmaceutically acceptable inorganic or organic bases and from pharmaceutically acceptable inorganic or organic acids. In addition, when a peptide contains both a basic moiety, such as an amine, pyridine or imidazole, and an acidic moiety such as a carboxylic acid or tetrazole, zwitterions may be formed and are included within the term "salt" as used herein. Salts derived from pharmaceutically acceptable inorganic bases include ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium, and zinc salts, and the like. Salts derived from pharmaceutically acceptable organic bases include salts of primary, secondary and tertiary amines, including substituted amines, cyclic amines, naturally-occurring amines and the like, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperadine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like. Salts derived from pharmaceutically acceptable inorganic acids include salts of boric, carbonic, hydrohalic (hydrobromic, hydrochloric, hydrofluoric or hydroiodic), nitric, phosphoric, sulfamic and sulfuric acids. Salts derived from pharmaceutically acceptable organic acids include salts of aliphatic hydroxyl acids (e.g., citric, gluconic, glycolic, lactic, lactobionic, malic, and tartaric acids), aliphatic monocarboxylic acids (e.g., acetic, butyric, formic, propionic and trifluoroacetic acids), amino acids (e.g., aspartic and glutamic acids), aromatic carboxylic acids (e.g., benzoic, p-chlorobenzoic, diphenylacetic, gentisic, hippuric, and triphenylacetic acids), aromatic hydroxyl acids (e.g., o-hydroxybenzoic, p-hydroxybenzoic, 1-hydroxynaphthalene-2-carboxylic and 3-hydroxynaphthalene-2-carboxylic acids), ascorbic, dicarboxylic acids (e.g., fumaric, maleic, oxalic and succinic

acids), glucuronic, mandelic, mucic, nicotinic, orotic, pamoic, pantothenic, sulfonic acids (e.g., benzenesulfonic, camphosulfonic, edisyllic, ethanesulfonic, isethionic, methanesulfonic, naphthalenesulfonic, naphthalene-1,5-disulfonic, naphthalene-2,6-disulfonic and p-toluenesulfonic acids), xinafoic acid, and the like. In some embodiments, the salt is an acetate or trifluoroacetate salt.

[0100] The aromatic-cationic peptides described herein, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, can be incorporated into pharmaceutical compositions for administration, singly or in combination, to a subject for the treatment or prevention of a disorder described herein. Such compositions typically include the active agent and a pharmaceutically acceptable carrier. As used herein the term “pharmaceutically acceptable carrier” includes saline, solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Supplementary active compounds can also be incorporated into the compositions.

[0101] Pharmaceutical compositions are typically formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral (e.g., intravenous, intradermal, intraperitoneal or subcutaneous), oral, inhalation, transdermal (topical), intraocular, iontophoretic, and transmucosal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. For convenience of the patient or treating physician, the dosing formulation can be provided in a kit containing all necessary equipment (e.g., vials of drug, vials of diluent, syringes and needles) for a treatment course (e.g., 7 days of treatment).

[0102] Pharmaceutical compositions suitable for injectable use can include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the

extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, a composition for parenteral administration must be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi.

[0103] The aromatic-cationic peptide compositions can include a carrier, which can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thiomerasol, and the like. Glutathione and other antioxidants can be included to prevent oxidation. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, aluminum monostearate or gelatin.

[0104] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle, which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, typical methods of preparation include vacuum drying and freeze drying, which can yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0105] Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules, *e.g.*, gelatin capsules. Oral

compositions can also be prepared using a fluid carrier for use as a mouthwash. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[0106] For administration by inhalation, the compounds can be delivered in the form of an aerosol spray from a pressurized container or dispenser, which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer. Such methods include those described in U.S. Pat. No. 6,468,798.

[0107] Systemic administration of a therapeutic compound as described herein can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art. In one embodiment, transdermal administration may be performed my iontophoresis.

[0108] A therapeutic protein or peptide can be formulated in a carrier system. The carrier can be a colloidal system. The colloidal system can be a liposome, a phospholipid bilayer vehicle. In one embodiment, the therapeutic peptide is encapsulated in a liposome while maintaining peptide integrity. One skilled in the art would appreciate, there are a variety of methods to prepare liposomes. (*See* Lichtenberg, *et al.*, *Methods Biochem. Anal.*, 33:337-462 (1988); Anselem, *et al.*, *Liposome Technology*, CRC Press (1993)). Liposomal formulations can delay clearance and increase cellular uptake (*See* Reddy, *Ann. Pharmacother.*, 34(7-8):915-923 (2000)). An active agent can also be loaded into a particle prepared from pharmaceutically acceptable ingredients including, but not limited to, soluble, insoluble, permeable, impermeable, biodegradable or gastroretentive polymers

or liposomes. Such particles include, but are not limited to, nanoparticles, biodegradable nanoparticles, microparticles, biodegradable microparticles, nanospheres, biodegradable nanospheres, microspheres, biodegradable microspheres, capsules, emulsions, liposomes, micelles and viral vector systems.

[0109] The carrier can also be a polymer, *e.g.*, a biodegradable, biocompatible polymer matrix. In one embodiment, the therapeutic peptide can be embedded in the polymer matrix, while maintaining protein integrity. The polymer may be natural, such as polypeptides, proteins or polysaccharides, or synthetic, such as poly α -hydroxy acids. Examples include carriers made of, *e.g.*, collagen, fibronectin, elastin, cellulose acetate, cellulose nitrate, polysaccharide, fibrin, gelatin, and combinations thereof. In one embodiment, the polymer is poly-lactic acid (PLA) or copoly lactic/glycolic acid (PGLA). The polymeric matrices can be prepared and isolated in a variety of forms and sizes, including microspheres and nanospheres. Polymer formulations can lead to prolonged duration of therapeutic effect. (*See* Reddy, *Ann. Pharmacother.*, 34(7-8):915-923 (2000)). A polymer formulation for human growth hormone (hGH) has been used in clinical trials. (*See* Kozarich and Rich, *Chemical Biology*, 2:548-552 (1998)).

[0110] Examples of polymer microsphere sustained release formulations are described in PCT publication WO 99/15154 (Tracy, *et al.*), U.S. Pat. Nos. 5,674,534 and 5,716,644 (both to Zale, *et al.*), PCT publication WO 96/40073 (Zale, *et al.*), and PCT publication WO 00/38651 (Shah, *et al.*). U.S. Pat. Nos. 5,674,534 and 5,716,644 and PCT publication WO 96/40073 describe a polymeric matrix containing particles of erythropoietin that are stabilized against aggregation with a salt.

[0111] In some embodiments, the therapeutic compounds are prepared with carriers that will protect the therapeutic compounds against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Such formulations can be prepared using known techniques. The materials can also be obtained commercially, *e.g.*, from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to specific cells with monoclonal antibodies to cell-specific antigens) can also be used as pharmaceutically acceptable

carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

[0112] The therapeutic compounds can also be formulated to enhance intracellular delivery. For example, liposomal delivery systems are known in the art, see, e.g., Chonn and Cullis, "Recent Advances in Liposome Drug Delivery Systems," *Current Opinion in Biotechnology* 6:698-708 (1995); Weiner, "Liposomes for Protein Delivery: Selecting Manufacture and Development Processes," *Immunomethods*, 4(3):201-9 (1994); and Gregoriadis, "Engineering Liposomes for Drug Delivery: Progress and Problems," *Trends Biotechnol.*, 13(12):527-37 (1995). Mizguchi, *et al.*, *Cancer Lett.*, 100:63-69 (1996), describes the use of fusogenic liposomes to deliver a protein to cells both *in vivo* and *in vitro*.

[0113] Dosage, toxicity and therapeutic efficacy of the therapeutic agents can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds that exhibit high therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[0114] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the methods, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to determine useful doses in humans accurately. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0115] Typically, an effective amount of the aromatic-cationic peptides, sufficient for achieving a therapeutic or prophylactic effect, range from about 0.000001 mg per kilogram body weight per day to about 10,000 mg per kilogram body weight per day. Suitably, the dosage ranges are from about 0.0001 mg per kilogram body weight per day to about 100 mg per kilogram body weight per day. For example dosages can be 1 mg/kg body weight or 10 mg/kg body weight every day, every two days or every three days or within the range of 1-10 mg/kg every week, every two weeks or every three weeks. In one embodiment, a single dosage of peptide ranges from 0.001-10,000 micrograms per kg body weight. In one embodiment, aromatic-cationic peptide concentrations in a carrier range from 0.2 to 2000 micrograms per delivered milliliter. An exemplary treatment regime entails administration once per day or once a week. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the subject shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

[0116] In some embodiments, a therapeutically effective amount of an aromatic-cationic peptide may be defined as a concentration of peptide at the target tissue of 10^{-12} to 10^{-6} molar, *e.g.*, approximately 10^{-7} molar. This concentration may be delivered by systemic doses of 0.001 to 100 mg/kg or equivalent dose by body surface area. The schedule of doses would be optimized to maintain the therapeutic concentration at the target tissue, most preferably by single daily or weekly administration, but also including continuous administration (*e.g.*, parenteral infusion or transdermal application).

[0117] The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to, the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of the therapeutic compositions described herein can include a single treatment or a series of treatments.

[0118] The subject treated in accordance with present methods can be any mammal or animal, including, for example, farm animals, such as sheep, pigs, cows, and horses; pet animals, such as dogs and cats; laboratory animals, such as rats, mice and rabbits. In a preferred embodiment, the mammal is a human.

Combination Therapy with an Aromatic-Cationic Peptide and Other Therapeutic Agents

[0119] In some embodiments, the aromatic-cationic peptides, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, may be combined with one or more additional agents for the prevention or treatment of reduced TAZ1 expression levels or Barth Syndrome. Drug treatment for reduced TAZ1 expression levels or Barth Syndrome typically involves antibiotics, granulocyte colony stimulating factor (GCSF), and agents for the control of cardiac conditions, including but not limited to, for example, diuretics, ACE inhibitors, digoxin (digitalis), calcium channel blockers, and beta-blockers. In mild cases, thiazide diuretics, such as hydrochlorothiazide at 25-50 mg/day or chlorothiazide at 250-500 mg/day, are useful. However, supplemental potassium chloride may be needed, since chronic diuresis causes hypokalemis alkalosis. Moreover, thiazide diuretics usually are not effective in patients with advanced symptoms of Barth Syndrome. Typical doses of ACE inhibitors include captopril at 25-50 mg/day and quinapril at 10 mg/day.

[0120] In one embodiment, the aromatic-cationic peptide, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, is combined with an adrenergic beta-2 agonist. An “adrenergic beta-2 agonist” refers to adrenergic beta-2 agonists and analogues and derivatives thereof, including, for example, natural or synthetic functional variants, which have adrenergic beta-2 agonist biological activity, as well as fragments of an adrenergic beta-2 agonist having adrenergic beta-2 agonist biological activity. The term “adrenergic beta-2 agonist biological activity” refers to activity that mimics the effects of adrenaline and noradrenaline in a subject and which improves myocardial contractility in a patient having Barth Syndrome. Commonly known adrenergic beta-2 agonists include, but are not limited to, clenbuterol, albuterol, formeoterol, levalbuterol, metaproterenol, pirbuterol, salmeterol, and terbutaline.

[0121] In one embodiment, the aromatic-cationic peptide, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, is combined with an adrenergic beta-1 antagonist. Adrenergic beta-1 antagonists and adrenergic beta-1 blockers refer to adrenergic beta-1 antagonists and analogues and derivatives thereof, including, for example, natural or synthetic functional variants which have adrenergic beta-1 antagonist biological activity, as well as fragments of an adrenergic beta-1 antagonist having adrenergic beta-1 antagonist biological activity. Adrenergic beta-

1 antagonist biological activity refers to activity that blocks the effects of adrenaline on beta receptors. Commonly known adrenergic beta-1 antagonists include, but are not limited to, acebutolol, atenolol, betaxolol, bisoprolol, esmolol, and metoprolol.

[0122] Clenbuterol, for example, is available under numerous brand names including Spiropent® (Boehringer Ingelheim), Broncodil® (Von Boch I), Broncoterol® (Quimedical PT), Cesbron® (Fidelis PT), and Clenbuter® (Biomedica Foscama). Similarly, methods of preparing adrenergic beta-1 antagonists such as metoprolol and their analogues and derivatives are well-known in the art. Metoprolol, in particular, is commercially available under the brand names Lopressor® (metoprolol tartate) manufactured by Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, N.J. 07936-1080. Generic versions of Lopressor® are also available from Mylan Laboratories Inc., 1500 Corporate Drive, Suite 400, Canonsburg, Pa. 15317; and Watson Pharmaceuticals, Inc., 360 Mt. Kemble Ave. Morristown, N.J. 07962. Metoprolol is also commercially available under the brand name Toprol XL®, manufactured by Astra Zeneca, LP.

[0123] In one embodiment, an additional therapeutic agent is administered to a subject in combination with an aromatic cationic peptide, such that a synergistic therapeutic effect is produced. Therefore, lower doses of one or both of the therapeutic agents may be used in treating Barth Syndrome, resulting in increased therapeutic efficacy and decreased side-effects.

[0124] In any case, the multiple therapeutic agents may be administered in any order or even simultaneously. If simultaneously, the multiple therapeutic agents may be provided in a single, unified form, or in multiple forms (by way of example only, either as a single pill or as two separate pills). One of the therapeutic agents may be given in multiple doses, or both may be given as multiple doses. If not simultaneous, the timing between the multiple doses may vary from more than zero weeks to less than four weeks. In addition, the combination methods, compositions and formulations are not to be limited to the use of only two agents.

EXAMPLES

[0125] The present invention is further illustrated by the following examples, which should not be construed as limiting in any way.

Example 1 – Effects of Aromatic-Cationic Peptides on Heart Mitochondrial Cardiolipin in a Dog Model of Heart Failure

[0126] This example demonstrates the effect of the aromatic-cationic peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ on levels of heart mitochondrial cardiolipin in dogs with coronary microembolization-induced heart failure. In particular, the effects of D-Arg-2'6'-Dmt-Lys-Phe-NH₂ on levels of the 18:2-18:2-18:2-18:2 cardiolipin species are evaluated.

Methods

[0127] Heart failure was induced in dogs via multiple sequential intracoronary microembolizations as described in Sabbah, *et al.*, *Am J Physiol.* (1991) 260:H1379-84. Half the dogs were subsequently treated with the mitochondrial peptide; the other half were treated with drug vehicle and served as controls. Peptide treatment was started upon induction of heart failure (HF), defined as left ventricular ejection fraction of approximately 30%. The daily dose of the peptide was 0.5 mg/kg/day administered intravenously. At the end of the treatment phase (12 weeks) dogs in both the vehicle and treatment groups were sacrificed and a sample of heart muscle from the left ventricle was removed, washed with saline, and immediately frozen and stored at -80° C. For cardiolipin analysis, lipids were extracted from the heart tissue sample with a chloroform/methanol solution (Bligh Dyer extraction). Individual lipid extracts were reconstituted with chloroform:methanol (1:1), flushed with N₂, and then stored at -20° C before analysis via electrospray ionization mass spectroscopy using a triple-quadrupole mass spectrometer equipped with an automated nanospray apparatus. Enhanced multidimensional mass spectrometry-based shotgun lipidomics for cardiolipin was performed as described by Han, *et al.*, “*Shotgun lipidomics of cardiolipin molecular species in lipid extracts of biological samples*,” *J Lipid Res* 47(4):864-879 (2006).

Results

[0128] The 18:2 cardiolipin species was significantly reduced in untreated heart failure dogs (Heart Failure, Control) ($p < 0.05$) as compared to cardiac tissue from normal subjects (Normal). FIG. 1. However, heart failure dogs treated with D-Arg-2'6'-Dmt-Lys-Phe-NH₂ (Heart Failure, Peptide) had levels of 18:2 cardiolipin that were similar to normal subjects, and greater than the heart failure control subjects ($p < 0.05$). FIG. 1.

Conclusions

[0129] The 18:2 cardiolipin species is reduced in heart failure subjects. The reduction of 18:2 cardiolipin leads to poor oxidative phosphorylation and subsequent LV dysfunction. Chronic treatment with D-Arg-2'6'-Dmt-Lys-Phe-NH₂ normalized 18:2 cardiolipin, which leads to improved LV function and rate of mitochondrial ATP synthesis.

[0130] These results show that aromatic-cationic peptides of the present invention, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, are useful in the prevention and treatment of diseases and conditions associated with aberrant cardiolipin levels. In particular, these results show that aromatic-cationic peptides of the present invention, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, are useful in methods comprising administration of the peptide to subjects in need of normalization of cardiolipin levels and remodeling.

Example 2 – Effects of Aromatic-Cationic Peptides on TAZ1 Expression in a Dog Model of Heart Failure

[0131] This example demonstrates the effect of the aromatic-cationic peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ on levels of TAZ1 expression in dogs with coronary microembolization-induced heart failure. In particular, the effects of D-Arg-2'6'-Dmt-Lys-Phe-NH₂ on levels of TAZ1 mRNA are evaluated.

Methods

[0132] Twelve dogs were subject to coronary microembolization-induced heart failure (LV ejection fraction ~30%) as described above in Example 1. Subjects were randomized into D-Arg-2'6'-Dmt-Lys-Phe-NH₂-treated and control groups for a three-month trial. Subjects received subcutaneous injections of D-Arg-2'6'-Dmt-Lys-Phe-NH₂ (0.5 mg/kg once daily, n=6) or saline (Untreated-HF Control, n=6). RNA was prepared from LV tissue of all subjects at the end of the treatment phase and from the LV of six normal subject controls. Levels of TAZ1 mRNA were determined by real-time PCR. Changes in mRNA levels were expressed as fold reduction using the CT Method, with normalization to a glyceraldehyde 1,3 diphosphate dehydrogenase (GAPDH) internal control.

Results

[0133] Levels of TAZ1 mRNA were reduced 2.25-fold in heart failure subjects receiving saline control as compared to normal subjects. FIG. 2. Treatment with D-Arg-2'6'-Dmt-Lys-Phe-NH₂ attenuated the decrease in TAZ1 to only 1.23-fold, relative to normal subjects. FIG. 2.

Conclusions

[0134] Heart failure is associated with dysregulation of cardiolipin remodeling enzymes that can lead to pathologic remodeling of cardiolipin and to structural and functional mitochondrial abnormalities. Chronic therapy with D-Arg-2'6'-Dmt-Lys-Phe-NH₂ partially reverses these maladaptations thus allowing for resumption of physiologic post-biosynthesis remodeling of cardiolipin.

[0135] These results show that aromatic-cationic peptides of the present invention, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, are useful in the prevention and treatment of diseases and conditions associated with reduced TAZ1 expression levels. In particular, these results show that aromatic-cationic peptides of the present invention, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, are useful in methods comprising administration of the peptide to subjects in need of normalization of TAZ1 expression levels, such as, for example, subjects having Barth Syndrome.

Example 3 – Effects of Aromatic Cationic Peptides on Mitochondrial Ultrastructure and Organization

[0136] This example demonstrates that aromatic-cationic peptides, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, are useful in the treatment of Barth Syndrome.

[0137] A tissue sample from a subject with Barth Syndrome and tissue samples from a subject with cardiac disease was prepared for electron microscopy imaging of the mitochondria using standards known in the art. The tissue sample from the Barth Syndrome subject was stained and showed abnormal features or structures within the

mitochondria's ultrastructure and organization (see arrows in FIG. 3), some of which are highlighted in boxes b-d. FIG. 3.

[0138] Similar derangement of mitochondrial ultrastructure was seen in the tissue sample from a cardiac disease subject. FIG. 4A. Treating a cardiac disease subject with an effective amount of D-Arg-2'6'-Dmt-Lys-Phe-NH₂ ameliorated the abnormal features of the ultrastructure of the mitochondria. FIG. 4B.

[0139] Furthermore, the amelioration of the pathological effects of cardiac disease in mitochondria was further shown in the improved organization of the mitochondria in tissue from a cardiac disease subject treated with D-Arg-2'6'-Dmt-Lys-Phe-NH₂ as compared to mitochondria in tissue from a subject not treated with D-Arg-2'6'-Dmt-Lys-Phe-NH₂. FIGs. 5A-5B.

[0140] The results show that aromatic-cationic peptides such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂ are useful for reducing the number of mitochondria with abnormal mitochondrial ultrastructure, and/or ameliorating abnormal mitochondrial ultrastructure, and maintaining mitochondrial organization in cardiac disease. It is anticipated that aromatic-cationic peptides such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂ will have a similar effect on abnormal mitochondria ultrastructure in subjects with Barth Syndrome. As such, aromatic-cationic peptides of the present disclosure are useful in methods for the treatment of Barth Syndrome.

Example 4 – Use of Aromatic-Cationic Peptides in the Treatment of Barth Syndrome

[0141] This example will demonstrate the use of aromatic-cationic peptides, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt, in the treatment of Barth Syndrome.

Methods

[0142] Barth Syndrome patients will receive daily administrations of a therapeutically effective amount of aromatic-cationic peptide, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt. Peptides may be administered orally, topically, systemically, intravenously, subcutaneously, intraperitoneally, or intramuscularly according to methods known in the art. Subjects will be evaluated weekly for the presence and/or severity of signs and symptoms associated

with Barth Syndrome, including, but not limited to, *e.g.*, cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and frequent bacterial infections. Treatments will be maintained until such a time as Symptoms of Barth Syndrome are ameliorated or eliminated.

Results

[0143] It is predicted that Barth Syndrome subjects receiving therapeutically effective amounts of aromatic-cationic peptide, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt will display reduced severity or elimination of symptoms associated with Barth Syndrome.

[0144] These results will show that aromatic-cationic peptides, such as D-Arg-2'6'-Dmt-Lys-Phe-NH₂, or a pharmaceutically acceptable salt thereof, such as acetate or trifluoroacetate salt are useful in the treatment of Barth Syndrome. Accordingly, the peptides are useful in methods comprising administering aromatic-cationic peptides to a subject in need thereof for the treatment of Barth Syndrome.

EQUIVALENTS

[0145] The present invention is not to be limited in terms of the particular embodiments described in this application, which are intended as single illustrations of individual aspects of the invention. Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. Functionally equivalent methods and apparatuses within the scope of the invention, in addition to those enumerated herein, will be apparent to those skilled in the art from the foregoing descriptions. Such modifications and variations are intended to fall within the scope of the appended claims. The present invention is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled. It is to be understood that this invention is not limited to particular methods, reagents, compounds compositions or biological systems, which can, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

[0146] In addition, where features or aspects of the disclosure are described in terms of Markush groups, those skilled in the art will recognize that the disclosure is also thereby

described in terms of any individual member or subgroup of members of the Markush group.

[0147] As will be understood by one skilled in the art, for any and all purposes, particularly in terms of providing a written description, all ranges disclosed herein also encompass any and all possible subranges and combinations of subranges thereof. Any listed range can be easily recognized as sufficiently describing and enabling the same range being broken down into at least equal halves, thirds, quarters, fifths, tenths, *etc.* As a non-limiting example, each range discussed herein can be readily broken down into a lower third, middle third and upper third, *etc.* As will also be understood by one skilled in the art all language such as “up to,” “at least,” “greater than,” “less than,” and the like, include the number recited and refer to ranges which can be subsequently broken down into subranges as discussed above. Finally, as will be understood by one skilled in the art, a range includes each individual member. Thus, for example, a group having 1-3 cells refers to groups having 1, 2, or 3 cells. Similarly, a group having 1-5 cells refers to groups having 1, 2, 3, 4, or 5 cells, and so forth.

[0148] Other embodiments are set forth within the following claims.

The embodiments of the present invention for which an exclusive property or privilege is claimed are defined as follows:

1. Use of the peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ or a pharmaceutically acceptable salt thereof for treatment or prevention of Barth Syndrome in a subject.
2. The use of claim 1, wherein the subject displays reduced levels of TAZ1 expression compared to a normal control subject.
3. The use of claim 1 or 2, wherein the peptide is for use daily for 6 weeks or more.
4. The use of any one of claims 1-3, wherein the peptide is for use daily for 12 weeks or more.
5. The use of any one of claims 1-4, wherein the subject has been diagnosed as having Barth Syndrome.
6. The use of claim 5, wherein the Barth Syndrome comprises one or more of cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and frequent bacterial infections.
7. The use of any one of claims 1-6, wherein the subject is human.
8. The use of any one of claims 1-7, wherein the peptide is in a form for administration orally, topically, systemically, intravenously, subcutaneously, intraperitoneally, or intramuscularly.
9. The use of any one of claims 1-8, wherein the peptide is for use separately, sequentially or simultaneously with use of a cardiovascular agent.


10. The use of claim 9, wherein the cardiovascular agent is selected from the group consisting of: an anti-arrhythmia agent, a vasodilator, an anti-anginal agent, a corticosteroid, a cardiotropin, a diuretic, a sedative, an angiotensin converting enzyme (ACE) inhibitor, an angiotensin II antagonist, a thrombolytic agent, a calcium channel blocker, a thromboxane receptor antagonist, a radical scavenger, an anti-platelet drug, a β -adrenaline receptor blocking drug, α -receptor blocking drug, a sympathetic nerve inhibitor, a digitalis formulation, an inotrope, and an antihyperlipidemic drug.
11. The use of any one of claims 1-10, wherein the pharmaceutically acceptable salt comprises acetate or trifluoroacetate salt.
12. Use of the peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ or a pharmaceutically acceptable salt thereof for increasing expression of TAZ1 in a mammalian subject.
13. The use of claim 12, wherein the expression of TAZ1 in the subject is about 2-5 fold less than the level of TAZ1 expression in a normal control subject.
14. The use of claim 12 or 13, wherein the peptide is for use daily for 6 weeks or more.
15. The use of any one of claims 12-14, wherein the peptide is for use daily for 12 weeks or more.
16. The use of any one of claims 12-15, wherein the subject has been diagnosed as having, is suspected of having, or is at risk of having Barth Syndrome.
17. The use of claim 16, wherein the Barth Syndrome comprises one or more of cardiomyopathy, skeletal muscle abnormalities, neutropenia, slow development, weak muscle tone, increased levels of organic acids in the urine and blood, and frequent bacterial infections.
18. The use of any one of claims 12-17, wherein the subject is human.
19. The use of any one of claims 12-18, wherein the peptide is in a form for administration orally, topically, systemically, intravenously, subcutaneously, intraperitoneally, or intramuscularly.

20. The use of any one of claims 12-19, wherein the peptide is for use separately, sequentially or simultaneously with a cardiovascular agent.
21. The use of claim 20, wherein the cardiovascular agent is selected from the group consisting of: an anti-arrhythmia agent, a vasodilator, an anti-anginal agent, a corticosteroid, a cardiotropine, a diuretic, a sedative, an angiotensin converting enzyme (ACE) inhibitor, an angiotensin II antagonist, a thrombolytic agent, a calcium channel blocker, a thromboxane receptor antagonist, a radical scavenger, an anti-platelet drug, a β -adrenaline receptor blocking drug, α -receptor blocking drug, a sympathetic nerve inhibitor, a digitalis formulation, an inotrope, and an antihyperlipidemic drug.
22. The use of any one of claims 12-21, wherein the pharmaceutically acceptable salt comprises acetate or trifluoroacetate salt.
23. Use of the peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ or a pharmaceutically acceptable salt thereof for reducing the risk of Barth Syndrome in a mammalian subject having decreased expression of TAZ1 compared to a normal control subject.
24. Use of the peptide D-Arg-2'6'-Dmt-Lys-Phe-NH₂ or a pharmaceutically acceptable salt thereof for stabilizing cardiolipin remodeling in a mammalian subject having or suspected of having Barth Syndrome.
25. The use of claim 24, wherein the mammalian subject has decreased expression of TAZ1 compared to a normal control subject.
26. The use of claim 24, wherein the cardiolipin is 18:2 species of cardiolipin.


1/5

FIG. 1


Cardiolipin 18:2-18:2-18:2-18:2
nmol/mg non-collagen protein



2/5

FIG. 2

3/5

FIG. 3

4/5

FIG. 4A





FIG. 4B



5/5

FIG. 5A





FIG. 5B



Cardiolipin 18:2-18:2-18:2-18:2
nmol/mg non-collagen protein

