

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
5 July 2012 (05.07.2012)



(10) International Publication Number
WO 2012/091571 A3

(51) International Patent Classification:

A61K 31/14 (2006.01) A61K 31/7072 (2006.01)
A61K 31/202 (2006.01) A61K 31/714 (2006.01)
A61K 31/355 (2006.01) A61K 33/04 (2006.01)
A61K 31/375 (2006.01) A61K 45/06 (2006.01)
A61K 31/4415 (2006.01) A61P 3/02 (2006.01)
A61K 31/519 (2006.01) A61P 21/00 (2006.01)
A61K 31/683 (2006.01) A61P 25/28 (2006.01)
A61K 31/685 (2006.01)

(21) International Application Number:

PCT/NL2011/050908

(22) International Filing Date:

28 December 2011 (28.12.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PCT/NL2010/050892
28 December 2010 (28.12.2010) NL
PCT/NL2011/050555
11 August 2011 (11.08.2011) NL
PCT/NL2011/050556
11 August 2011 (11.08.2011) NL

(71) Applicant (for all designated States except US): **N.V. NUTRICIA** [NL/NL]; Eerste Stationsstraat 186, NL-2712 HM Zoetermeer (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HAGEMAN, Robert Johan Joseph** [NL/NL]; Hamsterlaan 12, NL-6705 CT Wageningen (NL). **DE WILDE, Mattheus Cornelis** [NL/NL]; Zonnegloren 14, NL-6661 RK Elst (NL). **GROENENDIJK, Martine** [NL/NL]; Mazurkastraat 50, NL-2992 GP Barendrecht (NL). **KAMPHUIS, Patrick**

Joseph Gerardus Hendrikus [NL/NL]; Adriaen van Ostadelaan 118, NL-3583 AM Utrecht (NL).

(74) Agents: **JANSEN, C.M.** et al.; Vereenigde, Johan de Wittlaan 7, NL-2517 JR Den Haag (NL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SI, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

7 September 2012

(54) Title: COMBINATION OF COMPONENTS FOR THE PREVENTION AND TREATMENT OF FRAILTY

(57) Abstract: Composition comprising at least two components, more preferably at least three components, more preferably at least four components, selected from the group of (i) a nucleoside equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, (iv) a phospholipid, (v) an antioxidant and (vi) a choline, with the proviso that at least component (i) or (iii) is present, for use in the prevention or treatment of frailty in a mammal, wherein frailty is determined by compliance with at least 2 criteria, preferably 3 criteria, selected from the group of muscle weakness, excessive feelings of exhaustion or fatigue, abnormally low physical activity, slow or unsteady gait, weight loss, and neurological dysfunction.



WO 2012/091571 A3

COMBINATION OF COMPONENTS FOR THE PREVENTION AND TREATMENT OF FRAILTY

5 FIELD OF THE INVENTION

The invention relates to a combination of specific components for use in the prevention or treatment of frailty in a mammal, in particular for use in increasing one or more of body weight, body mass index, lean body weight, muscle mass, muscle strength, or muscle function, especially in an elderly human, more in particular in a human prone to or suffering from cognitive decline or dementia, in particular senile dementia, more in particular Alzheimer's disease.

The invention also relates to the specific clinically relevant benefits that can be achieved in such persons, like improvement of stamina, an increased degree of activity during daytime and associated change in lifestyle. The invention also relates to a composition comprising said specific combination of components, either as a nutritional or pharmaceutical composition, which are suitable for achieving said effects in a frail mammal, especially in an elderly human, most in particular in a human prone to or suffering from dementia, in particular senile dementia, more in particular Alzheimer's disease.

20

BACKGROUND OF THE INVENTION

Body weight of humans is determined by the weight of the different parts in the body, like that of bones, muscle, organs, vessels, adipose tissue, et cetera. During lifetime the contributions of each body part to the total body weight changes. After maturation of the human body has stopped, typically muscle mass will gradually and steadily decrease with time. This decrease results in a decrease of lean body mass (LBM, which is the body mass minus the mass of adipose tissue), despite the fact that total body mass (or body weight) may increase, e.g. due to an increase of the mass of adipose tissue and changed masses of other parts of the body. Ferruci et al. reported about the progression of loss of muscle strength during aging (Ferruci et al. (1996) J Gerontol Med Sci, 51AM123-M130). The strength of a muscle is considered to be dependent on its mass. Its mass depends on the number of muscle fibers, which decreases only after 55 years of age to about 50 % at the age of about 80 years, their length, which depends on their trophic condition, and

the cross-sectional area, which depends on training (Paukner et al. (2007) Proc Au P S, 38; 69-75). Muscle strength (strength of handgrip or quadriceps) in normal persons appeared to decrease about 0.5 % per year after the 30th year. Typically, this rate of decline of muscle strength increases with age. After the 65th year of age, this decrease in muscle strength has reached a magnitude of 1.5 % per year for arm muscles and about 3.5 % for leg muscle (Skelton et al (1994) Age and Aging, 23, 371-377). Similar age-associated changes in body composition and muscle strength have also been described by Evans, Cyr-Campbell (1997), J American Dietetic Association, 97(6), 632, and by Campion (1998) N Engl J med, 338(15), 1064-66.

10 It is important to note that, apart from these chronic age-associated decreases in muscle mass and lean body weight, also temporal and mostly reversible changes therein can occur, which in most cases depend on the applied exercise efforts. For example, Muller observed that during long term bed rest, muscle strength decreased at a rate of about 1 % per day (Muller (1970) Arch Phys Med Rehabil, 51, 449-462).

15 These losses in body weight, lean body weight, muscle mass and muscle strength during aging are considered to be normal and physiological, though undesirable. However, in part of the human population these losses have occurred at a greater speed or have occurred for a longer period of time to reach a critical level. This abnormally large weight loss is in most cases associated with several health problems which occur at the same time. It is thought that a complex general and non-specific malfunction of the human body causes a low capability of the human body to adapt to the prevalent circumstances to which the individual is exposed. This general condition is recognized by physicians as an independent health problem for which the word "frailty" was proposed. In the past, different tools have been used to arrive at the diagnosis of frailty (mild or moderate) or prefrailty.

25 Frailty is a large problem to the individual which experiences it, to the environment and to society. It has a large impact on the patient's life and creates huge costs for medical care. For this reason, the problem is recognized in the prior art as a geriatric syndrome that is distinct from disability and comorbidity. In addition, a relatively low lean body mass and body weight in elderly and especially in persons experiencing neurological problems, is common and a large problem, which has not yet been solved in the prior art.

30 Food intake, lifestyle and metabolic properties, including energy expenditure, of an individual change with increasing age, which may lead to what has been called a

“physiologic anorexia of aging”. Dietary habits, nutrient intake, life style and the aging process are interrelated. For example, with decreasing activity during the applied life style, and the age-associated decline in basal metabolic rate, neuro-endocrine function, immune function and taste and smell perception, older people tend to consume less food, and consequently fewer nutrients, which may lead to a nutritional status, which does not the specific requirements of the elderly of the frail individuals. This complex combination of events which is specific to aging individuals, can result in a further decrease in body weight, lean body mass or body mass index (BMI). This may even result in frailty, as defined above.

10 In one embodiment of the invention, it is therefore an objective to meet the specific nutritional requirements associated with aging above 65 years of age, preferably of aging above 75 years and to include the active components in a food for this special medical purpose, and to include them in such amounts to compensate for the deficiencies, which would eventually arise when conventional food patterns would be applied, as already applied in the prior art. In this way, a nutritional composition according the invention preferably comply with the regulations for Food of Special Medical Purposes (FSMP), as currently valid in the European Union. In addition, the invention aims to increase lean body mass or significantly decrease the rate of loss of lean body mass, muscle mass or body weight.

20 Involuntary weight loss during aging above 65 years is strongly associated with decreased immuno-competence, impaired mood and low stamina and an increased prevalence of health complications in the same patient. Involuntary weight loss is therefore typically a great concern to the clinical practitioner and has been included in the list of symptoms for diagnosed frailty by Fried and others (Fried et al. (2001) J Gerontol, 56, M146-156).

Weight loss may already be present in the early stages of dementia and increases with the severity and progression of AD.

Many studies have been carried out to compare the annual weight variation in Alzheimer's disease patients and healthy control subjects. These studies reported that weight loss frequently occurs in the first stages of the disease, although patients usually have adequate energy intakes. In addition, there is no compelling evidence that energy expenditure is abnormally elevated relative to body size in Alzheimer's disease patients.

Typically, the prognosis of developing more diseases, or developing more severe grades of disease, or experiencing a more rapid progression of diseases increases with

lower BMI, in particular when it starts to become lower than 23.5 kg/m². This is especially true for patients suffering from a neurological problem, such as an impairment of brain function or the function of peripheral nerves. An example of such neurological dysfunction may be an impaired cognitive function, like a form of cognitive impairment or dementia.

Though the prior art is consistent about the progression of loss of lean body mass during Alzheimer's disease or dementia, and the increased risk of co-morbidity when lean body mass deteriorates, it is inconsistent about the risk of developing more or a more rapid cognitive decline when BMI increases. For example, the review of Gorospe and Dave (2007) Age and Aging, 36, 23-29) concluded that increased BMI is independently associated with increased risk of dementia. In one embodiment of the invention, the specific combination of components according to the invention was found not only to increase BMI in frail elderly suffering from dementia, with a subsequent improvement of the activities of daily living (ADL), but in addition to improve the cognitive function. This helps to obtain a superior effect of the composition according the invention on the rate and extent of decrease of frailty, compared to prior art compositions, because at least 2 symptoms which contribute to the diagnosis of frailty are simultaneously improved. This applies also to the effect of the composition according to the invention on other criterions that define frailty, like the effect on exhaustion or fatigue, as will be explained below. In one embodiment of the invention, the efficacy of the composition according to the invention can be represented as done in Figure 1.

In frail elderly, the combination of low stamina, low drive to perform normal activities to keep independence and low abilities of skeletal muscles to allow the activities of daily living may result in a low degree of activity during daily life, including the ability to purchase, prepare or consume adequate food quality and quantity.

Function, mass and strength of a skeletal muscle also depends on its use. Fiatarone et al (1994) N Engl J Med, 330 (25), 1769-750 found that high intensity training indeed decreased muscle weakness in frail elderly people, but administration of a soy protein based vitaminized energy drink (360 kcal) did not. However, high intensity exercise programmes and protocols become more difficult to apply, when the patients become older or experience more severe cognitive dysfunction or when the patients suffer from low stamina or depression and become little motivated. Application of the protocols demands special equipment and skilled personnel to make them more successful. Despite the availability of these programs, the problem of frailty and too low body mass

index persists in the elderly population and impairs the amount of activities that are applied during daytime, especially in disabled or institutionalized aged persons and persons having moderate to severe dementia. This illustrates the desirability for having available a composition, preferably a nutritional or pharmaceutical composition, which supports the treatment of frailty and prefrailty and that may have a positive effect on the amount of activities done in daily living and to the degree frail individuals can live independently.

Though frailty is associated with an increased risk for experiencing e.g. a stroke, Alzheimer's disease and other dementias, a delirium, Parkinson's Disease, hip fracture, incontinence, pneumonia and dehydration, pre-death, and several end stage diseases, frailty has to be considered as an independent state of the body. It is known in the prior art that the "nutritional status", in terms of concentrations of nutrients in blood (plasma), in frail elderly does not comply with the values that are typically measured in the blood of healthy elderly or healthy middle-aged adults. It is also known that bad nutritional status, in terms of blood concentrations of specific nutrients is associated with specific diseases, though in most cases causal relationships are difficult to recognize, if present at all. Nevertheless attempts have been made in the past to normalize nutrient concentrations in blood in frail persons and especially frail elderly, and subsequently improve BMI and the amount of activities of daily living, but this has appeared to be very difficult. For example Fiatarone, et al, have reported in 1994 a failure to achieve this, when a protein energy supplement with 30 % of the recommended intake of vitamins per serving is administered.

Borsheim et al. reported an improvement of lean body mass, muscle mass and muscle strength, but this study was not placebo-controlled and in 12 glucose-intolerant elderly subjects having a relatively high body weight of 74 kg. The positive effects were observed when for 16 weeks every day 2 times 11 g free amino acids (8 essential amino acids and arginine, of which about 1/3 was leucine) were administered; these are so unpalatable, that typically compliance to a therapeutic protocol using the effective amount of amino acids is low and thus efficacy in clinical practice is not obtained (Borsheim et al. (2008) Clin Nutr. 27, 189-195).

In a small study (5 healthy active non-frail elderly) Volpi E, et al found that oral administration of 40 g crystalline amino acids together with glucose in small portions every 10 minutes during 3 hours, resulted in a small decrease of leg muscle catabolic rate (Volpi et al. (2007) J Clin Endocrinol Metab 85 (12), 4481-90). De Jong and co-

workers have compared the effect of administrating mineral and vitamin-enriched food products (fruit and dairy products) and non-enriched foods and found no difference in lean body mass (de Jong et al. (2000) Am J Public health, 90 (6), 947-954) in frail elderly. Promoting and stimulating eating more volumes or more healthy and regular food products does also not work to improve body weight, BMI or frailty characteristics.

Therefore, there is a need to effectively improve the nutritional status in frail patients, in elderly having a low BMI and in frail elderly in particular, but in such a way that health is improved, as can be recognized by increasing lean body mass, healthy body weight, muscle capacity, and decreasing frailty score. Preferably, these effects are achieved in frail or prefrail persons, frail or prefrail elderly or elderly having a BMI below 26 and more specifically below 23.5 kg/m². More preferably, these improvements in health result in a higher amount of activities during daytime, especially during wake time and in general a better functioning in life and quality of life.

This specific improvement of the nutritional status of a frail or prefrail person or elderly person having a relatively low body weight is defined to be the nutritional management of the frail or prefrail consumer. Disease-specific requirements, in particular frailty-specific requirements are met, by administering the composition according to the invention. In one embodiment of the invention, the nutritional composition according to the invention achieves the nutritional management of frailty and prefrailty. This is done so not by providing nutrients in general which have been attempted in the prior art, but failed or had severe disadvantages.

In the nutritional management of frail persons, and especially frail elderly it is also important to recognize the problem of xerostomia or a dry mouth in general or during eating, and the problem of hypochlorhydria, i.e. the reduced secretion of hydrochloric acid by the stomach, in general or after food intake or after smelling or seeing the food. In one embodiment of the invention it is an objective to provide a nutritional or pharmaceutical composition which, when consumed, is well tolerated and even appreciated by persons who suffer from xerostomia or hypochlorhydria.

It is known in the prior art that supplementation of protein and energy to elderly who are at risk from malnutrition produces a small but consistent weight gain (Milne et al. (2009) Cochrane review). Healthy persons like athletes can increase their BMI, when very high amounts of specific proteins, peptides or amino acids are consumed, in particular when this is combined with an exercise protocol. The known approach to supplement additional protein and energy demands consumption of food on top of daily

meals, and a proper organ function, for dealing with the large amount of dietetic protein (i.e. nitrogen). As explained above, the elderly, and especially the elderly with impaired cognitive function and the frail elderly generally experience difficulties with consuming large quantities or volumes of food, may suffer from impairments in body and organ function or from early satiety and low appetite, have practical problems with cooking and with consuming the food products, and are not keen on or capable of applying exercise programs (Holmes (2008) Nursing standard, 22 (26), 47-57).

Accordingly, there is a need for an alternative way of nutritional management of elderly which are in need of an increased body weight, body mass index, lean body weight, muscle mass or muscle strength, to perform appropriately, especially frail elderly, in particular the frail elderly which experiences cognitive decline or a dementia. Hence, this forms a way of treatment of frailty, when body weight increases to such extent that this criterion of frailty is no longer fulfilled.

Preferably, the composition according the invention achieves all benefits as described above at about the same time. This is observed in elderly in general (above 65 years of age), but also in a subgroup thereof, the oldest one (persons above 75 years of age). In a preferred embodiment, the nutritional or pharmaceutical composition according to the invention is to be used for treating at least two symptoms of frailty, i.e. for increasing low body weight or BMI and for improving activities of daily living (ADL), especially in persons older than 50 years of age, more in particular those older than 65 years of age (elderly) and in particular older than 75 years of age (the oldest). The composition according to the invention may also be used to combat other symptoms of frailty, such as exhaustion or fatigue by its effect on muscle power, and thrive or stamina and the effect on neurological performance, in particular cognition, as disclosed before. Thus, it constitutes a composition that not only solves these symptoms, but also provides superior effectivity against frailty in comparison to prior art compositions, and without adverse effect.

BACKGROUND PRIOR ART

In WO2009/002146 (NV Nutricia), a composition is disclosed which comprises DIIA or EPA, in combination with uridine or its equivalent and optionally a range of other components to support activities of daily living. The composition can be administered to elderly and locomotor function appeared improved by administering these components. Also, protein could be included in the composition of the invention, in order to improve

muscle strength, when administered to frail elderly. In order to achieve this, in particular 1 to 5 g protein per 100 ml of a liquid composition was included, wherein the protein comprises more than 80 weight% milk-derived proteins.

WO2010/002257 (NV Nutricia) discloses the use of a nutritional preparation
5 comprising more than 18 energy percent protein (preferably 22 to 32 en%), whey protein, at least 12 g leucine per 100 g proteinaceous matter and a lipid fraction which comprises at least one of EPA, (n-3)DPA, DHA, and (n-3) ETA for improving muscle function in a mammal. The improvement of the function of muscle was in terms of maximum force, maximum contraction velocity and maximum relaxation velocity, all corrected for muscle
10 mass. This improvement was claimed to occur when the patient suffered from specific diseases, in particular cancer or during "aging" and was claimed to result in improving daily activity, physical performance and quality of life. Frail individuals as such were not mentioned and neither were elderly with a low BMI or elderly being frail. The composition as disclosed comprises preferably the components as mentioned and in
15 addition specific indigestible oligosaccharides, glutamine, cysteine, oligosaccharides, carnitine and taurine. Though soy protein and wheat protein were mentioned the inclusion of casein instead of these vegetable proteins was preferred. Nucleotides and uridine sources were not mentioned.

WO 2005/060952 (NV Nutricia) relates to a composition comprising in a daily
20 dosage form 14 to 1000 mg panthothenic acid (vitamin B5) for stimulating appetite, whereby body weight and muscle mass is increased in specific groups of diseased humans. This was surprising, since panthothenic acid had been reported before as a hunger suppressant. The composition may further comprise folic acid, vitamin C (as antioxidant), and vitamin B6 and B12 as part of a common vitamin premix.

25 WO2007/073178 (NV Nutricia) discloses a drink liquid for Alzheimer patients (Example 3) comprising a nucleoside equivalent (UMP), fish oil comprising DHA and EPA, vitamin B6, folic acid and vitamin B12, phospholipids, vitamin C (as antioxidant), and choline. The claimed effect is not disclosed.

WO2004/026294 (Nestec S.A.) discloses a nutritional composition which comprises
30 leucine and at least one of isoleucine, lysine, methionine, phenylalanine, threonine, tryptophan, valine or histidine in free or salt form, wherein free leucine is present in an amount of 10 to 35 weight% of the total amount of amino acids. Such composition was claimed to be useful for controlling tumour-induced weight loss, for stimulating protein synthesis, ameliorating loss of muscle in a human or for the dietary management of

malnutrition.

SUMMARY DESCRIPTION OF THE INVENTION

It has now been found that lean body mass (LBW) or the body mass index (BMI)
5 can be increased in frail mammals, or frailty can effectively be treated therapeutically or prophylactically by providing a nutritional or pharmaceutical composition comprising a specific combination of active components. Accordingly, the present invention relates to a composition comprising at least two, preferably three, and more preferably four components (as active ingredients) selected from the group of (i) a nucleoside equivalent,
10 (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, (iv) a phospholipid, (v) an antioxidant and (vi) a choline, with the proviso that at least component (i) or (iii) is present, for use in the prevention or treatment of (pathological) frailty in a mammal, in particular for use in increasing one or more of body weight, body mass index, and lean body weight, especially in a human,
15 especially in an elderly human.

Preferably, the vitamin b is selected from the group consisting of vitamin B6, vitamin B9 and vitamin B12. Further preferably, the anti-oxidant is selected from the group of vitamin C, vitamin E and selenium (including selenium compounds)

Further, it has been found that LBW or BMI can be increased. Accordingly, the
20 invention in particular relates to a composition comprising at least two, preferably three, and more preferably four components (as active ingredients) selected from the group of (i) a nucleoside equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of docosahexaenoic acid (DHA), docosapentaenoic acid (DPA) and eicosapentaenoic acid (EPA), (iii) a vitamin B, preferably selected from the group of vitamin B6, vitamin B9
25 and vitamin B12, (iv) a phospholipid, (v) an antioxidant, preferably selected from the group of vitamin C, vitamin E and selenium (including selenium compounds) and (vi) a choline, with the proviso that at least component (i) or (iii) is present, for use in increasing one or more of the following: body weight, body mass index, and lean body weight, especially in a human, especially in an elderly human.

30 A use according to the invention can be accomplished by administering the components used according to the invention, wherein the components are administered to the mammal in a diet without essentially increasing the daily caloric intake of the mammal respectively patient. The combination can be used for increasing body weight

for a medical reason, in particular can be administered to a subject that is underweight and as a result thereof has a medical problem, or has an increased risk of acquiring a health problem. For example the subject may be anorexic, or a frail or prefrail person, having a body weight that is undesirably low.

5 In particular, elderly with an age above 65 years of age, preferably with an age may benefit from a use according to the invention.

In particular the increase in or maintenance of in body weight comprises one or more of: (i) increasing or maintaining lean body weight and (ii) increasing or maintaining muscle mass. In case the subject is a human, the use preferably comprises
10 increasing or maintaining the body mass index.

In a specific embodiment of the invention at least two components selected from the group of: (i) nucleoside equivalents, (ii) n-3 polyunsaturated fatty acids selected from the group of DHA, DPA and EPA, (iii) vitamins B, (iv) phospholipids, (v) antioxidants and (vi) cholines - with the proviso that at least one (i) nucleoside or at least one (iii)
15 vitamin B is present – are used for improving the ability to perform an activity of daily living of a mammal, for maintaining the ability to perform an activity of daily living (ADL) of a mammal, or for reducing a deterioration in the ability to perform an activity of daily living of a mammal.

The ADL-status of a subject or a change in ADL of a subject may be determined
20 with a score-test such as the Katz score or the PASE score.

The Katz Index of Independence in Activities of Daily Living, commonly referred to as the Katz ADL, is the most appropriate instrument to assess functional status as a measurement of the client's ability to perform activities of daily living independently. Clinicians typically use the tool to detect problems in performing activities of daily living
25 and to plan care accordingly. The Index ranks adequacy of performance in the six functions of bathing, dressing, toileting, transferring, continence, and feeding. Clients are scored yes/no for independence in each of the six functions. A score of 6 indicates full function, 4 indicates moderate impairment, and 2 or less indicates severe functional impairment, see 'M. Wallace *et al.* 'Try This: Best Practices in Nursing Care to Older
30 Adults, Issue Number 2, revised 2007, from the Hartford Institute for Geriatric Nursing, New York University, College of Nursing'.

(http://consultgericn.org/uploads/File/trythis/try_this_2.pdf)

In general, a medical use regarding ADL involves treatment of a subject that has a

severe functional impairment. Common causes of severe functional impairment are motor dysfunctional causes, *e.g.* because of joint problems (*e.g.* due to arthrosis or rheumathoid arthritis), severe vascular problems (*e.g.* diabetic feet) or severe obesity (Quetelet Index > 40). In particular, the subject of a medical use for improving the ability to perform an activity of daily living of a mammal, for maintaining the ability to perform an activity of daily living of a mammal, or for reducing a deterioration in the ability to perform an activity of daily living of a mammal according to the invention may have a score of 2 or less on the Katz scale, or a PASE score of less than 30, in particular of less than 27.5 (for women) or less than 30 (for men).

Further, the invention relates to a nutritional composition comprising (i) at least two components selected from the group of uridine and iuridine monophosphate, (ii) DHA and EPA, (iii) a vitamin B, (iv) a phospholipid, v) an antioxidant, and (vi) a choline.

The term 'active ingredient' or 'active component' is used herein in particular for (i) nucleoside equivalents, (ii) n-3 polyunsaturated fatty acids selected from the group of DHA, DPA and EPA, (iii) vitamins B, (iv) phospholipids, (v) antioxidants and (vi) cholines, which are effective in one or more of the claimed uses.

If a specific compound falls in more than one of these groups (i)-(vii) of active ingredients it typically provides an active ingredient of both groups. The actual dosage or concentration provided for each group can be determined based on how the compound is used/metabolised after administration. For example, phosphatidyl choline (PC) is both a phospholipid and a choline. As a phospholipid it can be active in the distribution/emulsification of lipids, and thereafter it can be metabolised to release the choline, which is an essential component for mammals, such as humans. Thus, a composition comprising PC comprises both an active ingredient of the group of phospholipids and of the group of cholines. Another example of a compound providing more than one type of active ingredients is citicoline (cytidine diphosphate-choline), which provides both choline and a nucleoside equivalent.

In an advantageous embodiment, a nutritional or pharmaceutical composition for use according to the invention (hereafter also called "composition according to the invention") not only has an effect on LBW or the BMI in (frail) elderly, or may be used to treat frailty, especially in an elderly mammal, but that the administration of a nutritional or pharmaceutical composition of the invention wherein a uridine-source and ω -3 type polyunsaturated fatty acids (ω -3 PUFAs) are present, results in a further

improvement in the amount of activities of daily living that can be performed and an increased muscle mass, muscle strength or muscle function, when also a specific protein amount is included in the composition according to the invention. In particular, the composition according the invention may be used to improve stamina, to increase degree
5 and frequency of feeling more energetic, to decrease duration, severity or frequency of feeling tired or exhausted or feeling fatigue, or to increase eagerness to demonstrate initiatives and become more active. In addition the composition according the invention can improve gait.

It has also been found that mammals physically benefit from treatment with a
10 specific combination of active ingredients such that one or more of body weight, body mass index, or lean body weight are better maintained or even increased without the need to increase the daily caloric intake of said mammals. It has also been found that the specific combination of components is simultaneously capable of decreasing other symptoms of frailty, like an improvement of neurological dysfunction or an improvement
15 of longer term muscle use leading to an increase in the amount of activities that are applied during wake time. This is mandatory to the effective treatment of frailty. For this reason, the specific combination of active ingredients may be used for the treatment of frailty or prefrailty, thus resulting in achieving a lower risk of developing frailty in the course of time.

20 The components as described above can be combined with a specific protein composition to obtain a stronger effect on LBM or BMI.

In a very specific embodiment, the invention relates to a composition comprising at least (i) a uridine equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, (iv) a phospholipid, (v) an antioxidant,
25 preferably selected from the group consisting of vitamin C, vitamin E and selenium, and (vi) a choline, for use in the prevention or treatment of (pathological) frailty in a mammal, wherein the mammal is a patient suffering from dementia, in particular from Alzheimer's disease, the patient having an BMI below 26 and more preferably below 23.5.

30 In a very specific embodiment, the invention also relates to a composition comprising at least (i) a uridine equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, (iv) a phospholipid, (v) an antioxidant, preferably selected from the group consisting of vitamin C, vitamin E and selenium, and (vi) a choline, for improving the (pathological) frailty in a mammal,

and/or the quality of life, and/or the daily activities (ADL), wherein the mammal is a patient suffering from dementia, , in particular from Alzheimer's disease, the patient having an BMI below 26 and more preferably below 23.5.

5 The term "or", as used herein, means also "and", unless specified otherwise or the context dictates otherwise . Hence, "option A or B" means any of the options A, B and A and B.

The term "a" or "an", as used herein, means "at least one" unless specified otherwise.

10 When referring to a noun (*e.g.* a compound, an additive *etc.*) in singular, the plural is meant to be included, unless specified otherwise.

When referring hereinafter to a 'vitamin B', in general a vitamin B selected from the group of vitamin B6(pyridoxins), vitamin B9 (folates) and vitamin B12 (Cobalamines) is meant, unless specifically stated otherwise. When referred hereinafter to an 'antioxidant', in general an antioxidant selected from the group of vitamin C (ascorbates), vitamin E (tocopherols) and selenium (including selenium compounds) (v) is
15 meant, unless specifically stated otherwise.

When referring herein to an acid, *e.g.* a fatty acid or folic acid, this term is meant to include the conjugated bases of said acid (*e.g.* folate), salts of the acids and derivatives of the acid of which the body is capable of converting it into the acid (*e.g.* fatty acid esters, such as triglycerides), unless specified otherwise.
20

When referring herein to dosages, these are in particular intended for adult humans. The skilled person will be able to determine a suitable dosage for other mammals based on common general knowledge, the information disclosed herein and optionally a limited amount of routine testing.

25 When referring to a mammal, preferably the mammal is a human mammal, more preferably an elderly human mammal.

DETAILED DESCRIPTION OF THE INVENTION

Definition of frailty

30 For the purpose of this document, the inventors apply the following definition for frailty, which includes the practical approaches of many scientists (like Fried, Ory, and Chin A Paw). A person is deemed to be frail when the individual's condition in a recent

period of time complies with at least 3 of the following list of (6) classes of symptoms (frailty criteria):

1. Muscle weakness,
2. Excessive feelings of exhaustion or fatigue,
- 5 3. Abnormally low physical activity,
4. Slow or unsteady gait,
5. Weight loss, and
6. Neurological dysfunction

The degree of malfunction of each of the mentioned classes of symptoms can be measured by applying recognized methods that are known in the prior art. For the purpose of this invention, the application of the following methods is preferred :

Criterion 1 (muscle weakness) - For determining muscle weakness one of the following list: a) for muscles of the arm: hand grip strength or quadriceps strength, b) for the leg the method as described by Børshiem et al. (2008) Clin Nutr. 27, 189-195 ; or as described in Pijnappels et al. (2008) Eur J Appl Physiol. 102(5): 585–592 can be applied. Belonging to the lower 20% of the group of age and gender matched persons resulted in fulfilment of this frailty criterion.

Criterion 2 (feeling of exhaustion or fatigue) - For determining the degree of exhaustion or fatigue, one can assess one or more of a) mental fatigue, b) the perception of exerting a standardized exercise protocol, muscle capacity or muscle power, or c) lung capacity. Preferably, method a) is done by applying a questionnaire as known in the prior art, e.g. the CES-D scale as described by Roberts (1980) Psychiatr Res, 2, 125-134, so scoring on statements like “I cannot get going” or “I feel that everything I do is an effort”. This measure is strongly related to the degree of stamina of the individual. Method b) is a measure of physical exhaustion and can for example be done by applying the Borg scale for measuring perceived exertion (Borg (1998) Human kinetics, Champaign, Ill ; Nybo (2003) Med Sci Sports Exerc, 35, 589-594). Method c) is a measure of the capacity of a human being to provide sufficient oxygen (air) to the body. The preferred method for applying it, is measuring forced expiratory volume (F'EV1) in one second, which should be below 30 % of the normal value, for individuals of similar age and gender, in order to meet the criterion.

Measurement of the degree of physical exhaustion or fatigue, as recommended in method b) can be done in several ways, which all represent ways that reflect the capacity of the neuro-muscular system to exert labour and be active. For example, the

amount of physical activities can be measured, by attachment of devices to the person to be tested, and measuring the amount of physical activity which is voluntarily applied normally during wake time. This amount can also be related to the amount of sleep or rest that is needed to recover from it, and be put in a score. Secondly, the capacity of the muscle to maintain a certain force for a certain time is a useful measure. Preferably, this force that needs to be maintained is at 15 to 80 % of the maximal force the muscle can provide, in order to reflect better the activities as applied in normal life. Thirdly, the speed with which physical exhaustion is developed can be determined. For example, the amount of power a muscle can exert in a certain period of time can be determined; e.g. by measuring the force multiplied with the distance (in a movement) that the force was exerted, e.g. by swinging a leg at a certain velocity. Equipment for measuring these advanced parameters is commercially available. The speed with which a muscle can become tired, or in other words, the time that a certain force can be maintained, is a valuable measure of the capability of an organism to apply daily activities. When the measured values for these parameters in the tested person belong to the lower 20% of such values for a group of persons of similar age and gender, the person is deemed to comply with this criterion of frailty.

It is important to recognise that larger muscle strength does not in all circumstances necessarily lead to a higher muscle capacity or power or a better endurance. The larger force which is provided by the muscle may have exhausted endogenous energy supplies more, which deteriorates the time or distance that the muscle can exert the force, and therefore does not increase muscle capacity or power. A larger force may also increase negative feedback signals from peripheral joints, muscles and tendons. Muscle capacity or muscle power also better reflects the muscle effort that needs to be done to apply an activity of daily living (Theou et al. (2008) Appl Physiol, Katsiaris, et al, 2005). Therefore, muscle capacity or muscle power, and not muscle strength, provides useful information about the state of exhaustion and fatigue, and especially physical exhaustion and fatigue of a patient. The force that a skeletal muscle can apply also influences the maximal velocity with which a certain movement can be made. A high movement velocity does not automatically result in a large endurance or a good recovery after exercise, which are required for performing a large amount of activities or showing a high mobility during wake time over the day.

Criterion 3 (physical activity) - For determining the magnitude of physical activity of a human, one or more methods of the following list can be applied: a) determination of the amount of basic activities as applied during daily life (as described

above), and b) determination of the amount of instrumental activities, as applied during daily life. Especially the measurement of instrumental activities is a valuable measure. A reliable method to do this is assessment of the "Physical Activity Scale for the Elderly" (PASE), as described by Washburn et al. (1999) J Clin Epidemiol 52, 643-651; "The
5 physical activity scale for the elderly; evidence for validity". For men, a score of less than 30 and belonging to lowest 20 % part fulfilled this criterion for frailty. For women, the threshold value was < 27.5 (Graham et al. (2009) Gerontol, 55, 644-651).

Criterion 4 (gait) - For determining the gait of a human, tests can be applied, like those described in the "Modified Physical Performance Test" (PTT), as disclosed by
10 Binder et al. (1999) J Gerontol, 54, M428-432, like the preferred a) measure of "balance", as e.g. done by determining the way how a frail person is capable of taking a penny from the floor, or b) time and way to walk 5 to 15 meters, or c) time and way to raise from a chair. For example, for method b) the criterion for frailty was fulfilled when the person belonged to 20 % of persons with regard to the time needed. The threshold values per
15 gender as mentioned in Graham et al. (2009) Gerontol, 55, 644-651 can suitably be applied.

Criterion 5 (Weight loss) - For determining the presence of undesired weight loss, it is important to avoid confusion with acute weight loss as caused by a (serious) disease or trauma or an underlying pre-death syndrome. In one embodiment of the
20 invention, frailty as the result of cachexia is excluded from the group of persons wherein the composition according to the invention is effective. In one embodiment of the invention, this exclusion of persons who have become frail because of cachexia is preferred. In this way, frail individuals which comply with the definition as given above differ from the individuals which have developed a frailty that results from these
25 diagnosed conditions, related to acute phases during diseases and briefly before death.

With regard to determining the value of BMI, which should give concerns for becoming "frail", it is important to be aware of the fact that optimal BMI values change with age. While at young and middle age a BMI value between 20 to 25 kg per square meter is considered to be most healthy, at older age, e.g. above 65 years of age and
30 especially above 75 years of age, a BMI below 26 and more specifically below 23.5 is considered to be undesirable (Chin A Paw et al. (2003) J Nutr Health Aging 7(1), 55-60, or Chin (1999) J Clin Epidemiol 52 (11), 1015-1021). The preferred list of diagnostic tools for establishing a too low body weight, lean body mass or muscle mass aims to be suitable for the aged population.

For the purpose of the invention, the inventors feel that it is desirable to look carefully at the metabolic condition of the individual who suffers from a low BMI, body weight or muscle capacity. When such low body weight is the result of an acute process, the metabolic condition of the patient differs on essential points from that of a person who experiences a more generalized metabolic problem. This difference has also been made by others, for example for Alzheimer patients (Guerin et al. (2005) Am J Clin Nutr, 82 (2), 435-441). Part of the Alzheimer patients appear to experience sudden and severe weight loss as a result of intercurrent events, like institutionalization, a major change in life style, a trauma or the presence of a major disease, while a different part suffered from a more chronic weight loss. Therefore it is preferred that the BMI value, as proposed as a limit for meeting one criterion of frailty, is assessed in the absence of an acute phase condition, i.e. the absence of a diagnosed disease selected from a cancer, AIDS, a COPD, an infection, pre-death, pre-death associated with anorexia, surgery, an accident and similar major trauma, or in an alternative embodiment the acute phases during these diseases or conditions. The presence of the acute phase during these diseases can be determined by measuring a biomarker as known in the prior art, e.g. measuring plasma concentrations of acute phase proteins, like transthyretin, alpha1-acid glycoprotein, C-reactive protein, or measuring concentrations of cytokines which are representative of acute infections or severely progressed disease, like IL-1, IL-6 or TNF- α . It is important to note that a disease, like cancer, chronic obstructive pulmonary diseases, AIDS and other diseases or pre-death, at some stage can cause a frailty condition that is acute and severe. The inventors find that the metabolic condition of such individual is completely different from that of an individual suffering from a frailty that originates from a generalized metabolic inability to cope with the daily stresses from the outside world. Therefore, in the context of this invention, frailty, which is treated and solved by administering the nutritional or pharmaceutical composition according to the invention, is not this acute -, and disease- or trauma-related type of advanced weight loss or frailty. Instead, it is the frailty as defined below and caused by a general and complex deterioration of the body to adapt the external stresses, as can have been acquired during life e.g., by applying bad life style including bad dietary habits, exposure to toxicants or as a result of time, as occurs during aging, in particular by aging after 65 years of age and more in particular the frailty as caused by a chronic deterioration of adaptive responses.

Further, a method for determining involuntary weight loss can be used for assessing whether criterion 5 applies, is applying the criterion of 4.5 kg undesired

weight loss in the previous year or 5-10% of body weight within 6-12 months or less. Involuntary weight loss of a subject is weight loss occurring despite the subject's aim to maintain or to increase its body weight (on the subject's own behalf or after advice of carer or health care professional). Such weight loss can be determined routinely by
5 comparing present weight with the weight 1-2 years earlier and asking the subject whether he/she aimed to maintain or increase the body weight.

Further, a method can be used applying the criterion of 6 kg undesired or unexplainable weight loss during previous 2 years. Such weight loss can be determined routinely by comparing present weight with the weight in a period 2 years earlier and
10 asking the subject whether he/she aimed to maintain or increase the body weight. A pre-death condition can be diagnosed by a physician. Measurement of transthyretin and α 1-acid glycoprotein in blood plasma of both genders and the determination of low blood albumin and high C-reactive protein in blood of males, as applied by Carriere (POLA Group) et al. (1998) Arch Ophthalmology, 116, 1031) is recommended for establishing
15 predeath risk.

Criterion 6 (Neurological dysfunction) - For determining the degree of neurological dysfunction application of the following list is preferred: a) establishment of a tremor or a locomotor dysfunction different from the balance and performance tests applied in measuring gait performance ; b) measurement of cognitive function
20 (impairment) ; c) determination of verbal fluency ; d) measurement of speed of conductance of electrical signals over nerves ; e) analyses of sensory functioning, to establish a problem related to hearing, vision, tasting, smelling and touch) ; f) measurement of a emotional or psychological condition, like establishing the presence of major depression, an affect disorder or an anxiety disorder ; g) determination of
25 incontinence or the daily occurrence of significant involuntary urinary loss, and h) the presence of a major sleep disorder, like chronic insomnia or sleep apnea. Methods to assess neurological dysfunction have been described in the prior art and include the DSM IV methods, and the measurements as described in the corresponding domains in common frailty assessment methods, like the Groningen frailty indicator (Schuurmans
30 et al. (2004) J Gerontol, Biol Sci, Med Sci, 59, M962-5), the comprehensive geriatric assessment (FI-SGA: Goggins et al. (2005) 60, 1046 ; Jones et al. (2004) J Am Geriatr Soc, 52, 1929-33) or the clinical global impression of change in physical frailty (CGIC-PF: Studenski et al. (2004) J AM Geriatr Soc, 52, 1560-6). Involuntary urine loss is deemed to be significant when it exceeds 10 ml a day.

A patient is considered to meet this neurological dysfunction criterion of frailty when b) is met (cognitive impairment) or, when a combination of at least two of the other criteria has been fulfilled. Cognitive impairment can be assessed by methods known in the prior art, for example by applying measures of the domains related to the measurement of verbal memory, visuospatial memory and attention-executive abilities. Preferred methods include application of the ADAS-cog assessment, the MMSE assessment, the Montreal cognition assessment or the CERAD methods.

The analyses of sensory function can be applied by using methods known in the prior art to determine threshold values for tastes, odours and sounds, or to determine the ability to differentiate between different odours and tastes.

A person is considered to be mildly frail when it complies with three of the six criteria, and moderately frail when it complies with four criteria or when the score of the three symptoms is so bad, that it seriously impairs the patient's condition, as diagnosed by the clinician or physician.

A person is considered to be in a prefrail condition or prodromal frail condition, when his or her condition complies with only two of the six frailty criteria. When the composition according to the invention treats prefrailty, it is defined to act in a preventive manner on the development of frailty. So, prefrailty and prodromal frailty is defined to be synonymous.

A person is defined to be a frail elderly when it complies with the above-mentioned criteria for frailty and in addition is older than 65, preferably 75 years of age.

An individual is defined to be a prefrail elderly when the person's condition complies with the criteria of prefrailty and the person is older than 65, preferably 75 years of age.

Within the context of this invention, the term "frailty" or "frail" refers to pathological conditions, and is used to comprise conditions indicated above as prefrail, prodromal frail, mildly frail or moderately frail.

Though some scientists recognize various forms of frailty, including medical, functional, social, psychological and physical frailty (Faber et al. (2006) Arch Phys Med Rehabil. 87, 885-96), the composition according the invention aims to have its efficacy only in those patients which comply exactly with the criteria as set above.

Elderly or the aged population is a group that is defined in different ways in the prior art. For the purpose of defining the invention, the inventors have applied the

following definition. Elderly or the aged population is defined to be all persons being older than 65 years of age. The “oldest old” are those persons being older than 75 years. The claimed combination for use in accordance with the invention is in particular suitable for treatment of a mammal. In a preferred embodiment, said combination is to be used for the treatment of a human, in particular an elderly person. In the context of this application, an elderly person is a person of the age of 50 or more, in particular of the age of 55 or more, more in particular of the age of 60 or more, more in particular of the age of 65 or more. This rather broad definition takes into account the fact that the average age varies between different populations, on different continents, *etc.* Most developed world countries have accepted the chronological age of 65 years as a definition of 'elderly' or older person (associated with the age at which one may begin to receive pension benefits), but like many westernized concepts, this does not adapt well to *e.g.* the situation in Africa. At the moment, there is no United Nations (UN) standard numerical criterion, but the UN agreed cut-off is 60+ years to refer to the older population in Western world. The more traditional African definitions of an elder or 'elderly' person correlate with the chronological ages of 50 to 65 years, depending on the setting, the region and the country.

With the “nutritional management of frail individuals, prefrail individuals and persons experiencing a too low LBM or BMI” according the invention is meant the administration of nutritional components to frail individuals, prefrail individuals or persons having a too low BMI or LBM, in such a way that not only the endogenous concentrations of nutritional components are influenced, but that also the health-beneficial effect is obtained in persons suffering from frailty, frailty symptoms, prefrail persons or individuals suffering from a too low LBM or BMI as claimed in this application. This means that the composition according the invention may induce a concentration in blood of that nutrient, which is outside the normal range as typically observed in the same tissue of healthy individuals, for example of healthy individuals which consume regular food or a normal diet. In the context of this invention, an alternative way of saying the same is that the composition according to the invention also aims to therapeutically nourish frail or prefrail persons or persons having too low LBM or BMI.

It is important to notice that many elderly have become malnourished and have developed deficiencies in one or more nutrients like protein, energy or micro-ingredients or combinations thereof, like PEM (protein-energy malnourishment). Such “malnourishment” is currently assessed by using tools like the “MUST” or “MNA”, but it

is envisaged by the inventors that alternative tools will be developed, like one that is specifically focussing on nourishment status and one other tool which specifically focusses on general and quick screening of the condition of an individual in order to assess whether a condition is present wherein additional nutrition or other therapy can be useful. Therefore, in one embodiment of the invention the composition according to the invention is used in combination with an assessment of malnourishment of a mammal, preferably using MUST or MNA. In this embodiment, the invention comprises at least the following steps:

- 1- Assessment of the nutritional status of a mammal, and
- 2- Intervention with the nutritional or pharmaceutical composition according the invention.

Preferably, the assessment of nutritional status is applied at least twice in combination and relation to the nutritional intervention. More preferably, the assessment is done at least one time before and at least one time after the intervention to measure any changes in nutritional status. The initial nutritional assessment may reveal deficiencies which can be resolved by adapting the intervention composition according to the invention. In order to allow such nutritional management according the invention, separate tools have been defined which is a specific nutritional assessment tool and a module of food components which allows convenient adaptation and fortification of the nutritional intervention composition according to the invention.

A preferred tool for assessing nutritional status comprises several parts, including a questionnaire, which asks the right questions to assess nutritional status, a database which is filled in by answering the questions, and an algorithm which compares the answers with predefined normal values. Optionally conclusions are drawn to arrive at an advice. These functionalities can be incorporated into an electronic device like a computer or minicomputer, by loading and running an appropriate software program. The questions to ask are at least covering biomarkers for the determination of nutrient status using methods as applied in the state of the prior art clinical chemistry laboratories.

Apart from the “nutritional management”, the inventors also identify the “therapeutic management” of frailty and prefrailty. In this definition of “therapeutic management”, the management of the individual to be treated comprises at least the following steps:

a) diagnosis for frailty, using an assessment of the six criteria defined above for assessing the presence of frailty,

b) an intervention using the composition according to the invention, and

c) measurement of progress made by assessing at least again the same symptom
5 classes as done in step a).

Preferably, the assessment of symptoms is done by using the tool, in particular a form or document or an electronic device loaded with appropriate software, as described below.

Optionally, in the therapeutic management of frailty and prefrailty, the
10 intervention with the nutritional composition according to the invention can be combined with one or more of 1) an exercise program, 2) a program to improve social interactions, 3) a program to become more exposed to sunlight and fresh air, 4) an adaptation of the diet or of the general food intake practices of the patient, and 5) an intervention with a drug or medicine. It is preferred that the therapeutic management comprises at least a
15 program to be exposed to sunlight and fresh air or an exercise program as a fourth and fifth step in the therapeutic management of the condition of the patient.

In principle, the mammal to be treated may suffer from any of the frailty symptoms as defined above. In particular, the mammal is a human which suffer from at least a neurological or brain problem and one selected from the group of weight loss and an
20 abnormally low physical activity. Preferably, the patient experiences all three types of symptoms. When the patient experiences a neurological or brain problem, this is preferably a form of dementia, in particular senile dementia. For the purpose of the invention, persons that suffer from "senile dementia" are defined as suffering from one or more dementias. Senile dementia or dementia is considered to comprise Alzheimer's
25 disease (AD).

The dementia may be in any stage. In particular, the mammal may be in a prodromal stage of dementia. A "prodromal dementia patient" is a person who does not suffer from a senile dementia as defined above, but has an increased likelihood to develop senile dementia. Likewise, a "prodromal Alzheimer patient" is a person who does
30 not suffer from AD, but has an increased likelihood to develop AD. In principle, any diagnostic tool for determining prodromal dementia patient may be used. Several diagnostic tools that can be used to classify the patients as prodromal dementia patients are described below and include an accurate diagnosis of brain lesions and biochemical

problems and careful setting of criteria. Hence, the invention is independently directed at prodromal dementia patients or prodromal Alzheimer's patients.

In particular, persons that score positively on at least one, preferably at least two, more preferably at least three of the following criteria, are considered to be
5 prodromal dementia patients as defined herein:

- a level of more than 350 ng Total-tau per litre cerebrospinal fluid (CSF);
- a weight ratio of abeta-42/ Phospho-tau-181 of less than 6.5 in CSF;
- presence of medial temporal lobe (MTL) atrophy, existing of volume loss of hippocampus, entorhinal cortex, or amygdala evidenced on Magnetic Resonance
10 Imaging (MRI) with either qualitative ratings using visual scoring (referenced to well characterised population with age norms) or quantitative volumetry of regions of interest (referenced to well characterized population with age norms)
- presence of fronto-temporal lobe (FTL) atrophy evidenced on MRI with qualitative ratings or quantitative volumetry;
- 15 - a level of more than 25 pg F2-iso-prostane (F2-IsoP, isoprostane 8,12-iso-iPF2alpha-VI) per mL CSF.

Further explanations of the significance of concentrations of T-tau, P-tau181, Abeta42 and F2-Isoprostane in CSF for future development of Alzheimer's disease can be found in Hansson et al. (2006) Lancet Neurol 5:228-234 ; and in Pratico et al. (2002) Arch
20 Neurol 59:972-976.

In the context of this document, patients who are in a prodromal state of Alzheimer's Disease (AD) are defined to be in a predementia stage of AD. The values of the biomarkers indicate a condition of the body, in particular the central nervous system, wherein the risk of developing Alzheimer's disease is significantly increased, no matter
25 whether the final form of Alzheimer's disease will be typical Alzheimer's disease, atypical Alzheimer's disease or mixed Alzheimer's disease. This predementia state of the body can be without significant clinical symptoms, the so called preclinical state of Alzheimer's disease, wherein a patient is asymptomatic at risk for Alzheimer's disease or is experiencing presymptomatic Alzheimer's disease. It is also important to note that a
30 patient may experience mild cognitive impairment without having an increased risk for developing Alzheimer's disease as determined by measuring the above-mentioned parameters (Dubois et al. (2010) Lancet Neurol, 9, 1118-1127).

The composition according to the invention has its effects on body weight and frailty, no matter the patient is a real Alzheimer's disease patient, is a prodromal Alzheimer's disease patient, is a normal elderly which experiences age-associated memory impairment or mild cognitive decline without having increased risk of developing AD, or is in the preclinical state of being at increased risk of developing Alzheimer's disease or a different dementia, like a vascular dementia.

In a preferred embodiment, a prodromal Alzheimer patient can be identified as such because he meets at least the first two criteria (total tau and ratio abeta-42/P-tau-181). More preferably, one of the three other criteria (MTL atrophy, FTL atrophy, F2-IsoP) also applies.

In addition to or instead of one or more of the above criteria, the following can be advantageously used:

- reduced glucose metabolism in bilateral temporal parietal areas of the brain, as is detectable by Positron Emission Tomography (PET);
- reduced glucose metabolism in the posterior cingulate cortex, as is detectable by PET;
- impaired blood flow in the brain as measurable by applying Single-Photon Emission Computed Tomography (SPECT), for example by applying the radioisotope ^{99m}Tc -HMPAO);
- impaired glucose metabolism in the brain as measurable by applying SPECT;
- abnormalities in the histology of the medial or inferior temporal lobes as can be determined by MRI or in the rate of glucose utilisation;
- abnormalities in histology or glucose utilization in the temporal parietal cortex or posterior cingulate cortex.

Abnormalities in the condition of the brain or parts thereof can be established by either taking the person's own condition under healthy circumstances as a reference, or, when this is not available, by taking the average condition of a representative group (so matched for e.g. age) as a reference. The latter will occur most frequently. By comparison of the patient's condition with the reference situation and the average situation when the pathological condition would have been developed to its full extent, the clinician is capable of recognizing a prodromal phase. In particular an intermediate situation wherein the patient demonstrates a deviation of x% from the value of a healthy

individual in the direction of the pathological conditions is for the purpose of this invention considered to be a prodromal patient. The value of x for the determination of blood flow and glucose metabolism is 20% when determined under standardised conditions in terms of feeding and exercise.

5 It is important to notice that the effect of the composition according to the invention on body weight occurs in the presence of relatively little energy and protein in the composition according to the invention and that this effect on body weight occurs simultaneously with an effect on other symptoms of frailty, like neurological or brain function, or the physical activity, in particular instrumental activities, which makes it
10 suitable for the treatment of frailty and prefrailty.

Composition according to the invention

The composition according the invention is a nutritional or a pharmaceutical composition. When referring to pharmaceutical composition, it is understood that such
15 composition has only a medical effect or benefit, i.e. the composition comprises essentially no ingredients which provide a substantial source of energy, other than the active components. When referring to nutritional composition, it is understood that such composition has both a nutritional and a medical effect or benefit, i.e. the composition comprises macronutrients which provide a substantial source of energy, other than the
20 active components, in particular proteinaceous matter, fat, other than component (ii), and digestible carbohydrates. It preferably comprises food grade components, which make it suitable for safe oral intake or enteral administration. The components can also be dissolved in a matrix which makes it suitable for parenteral administration.

The composition according to the invention can have any form or physical
25 condition. Preferably, it is a sterile composition or a composition which comprises a defined micro-organism population, like a dairy product, for example which is fermented under controlled conditions or a dry product to which probiotics have been added. An example of such fermented product is a yoghurt. The composition according the invention can be solid, semi-solid or a drink. Such forms have been widely disclosed in
30 the prior art. Preferably it is a drink, though for patients which experience dysphagia, a high-viscosity product or a semi-solid form is preferred.

The composition according to the invention may also be a kit-of-parts comprising the components according to the invention, packed for simultaneous or sequential

administration to a person in need thereof. Hence, each component may be packed separately, or some may be packed together, for example in a sachet, bottle, etc.

The composition according the invention is preferably a composition which complies with the criterions as set for the ruling regulations for food for special medical purposes or of a medical food. These regulations are distributed by the Food and Drug Administration or as directives from the European Union or by recognized authorities in other jurisdictions. In particular the composition according to the invention provides those nutrients and those amounts as required by the patient because of the specific disease state of the patient. These amounts of the active components cannot be consumed by adapting the normal diet.

In an embodiment, the composition according the invention is meant to therapeutically improve one or more of body weight, body mass index, lean body weight, muscle mass, muscle strength or muscle function. This improvement can occur in elderly, in particular frail elderly. The composition according to the invention can suitably be used in the nutritional management of individuals suffering from frailty symptoms or the therapeutic management of frailty (or prefrailty), by providing a certain amount of the active components per day, and preferably per period of about 4 hours.

A daily amount as described herein means in particular an amount in a daily dosage unit provided by the combination of the invention. Such a daily dosage unit may be a single dosage, but it may also be divided over two or three, or even more daily servings. If the combination, according to a preferred embodiment, is intended for administration as a single unit, the daily amounts as described herein are preferably the amounts present in the (preferably packaged) combination unit.

It has been found that the administration of at least two components selected from the group of (i) a nucleoside equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, preferably selected from the group of vitamin B6, B11 and B12, (iv) a phospholipid, (v) an antioxidant, preferably selected from the group of vitamin C and vitamin E and (vi) a choline, with the proviso that at least component (i) or (ii) is present, is suitable for use in the prevention or treatment of frailty in a mammal, in one embodiment together with or as part of a diet providing energy sources (carbohydrate, protein, fat), compared to the same isocaloric diet without said combination. Thus, it is concluded that the composition according to the invention is effective, without needing an increase in caloric intake. Accordingly, in a specific embodiment, the effect of said combination on increasing one or more of body

weight, body mass index, or lean body weight is not attributed to an increase in caloric intake.

(i) Nucleoside equivalent

The composition according to the invention may comprise a nucleoside equivalent.
5 As used herein, nucleosides include nucleosides as such deoxynucleosides as such, and equivalents of nucleosides as such or deoxynucleosides as such. Thus, when referring to a nucleoside, this term is meant to include the corresponding deoxynucleoside.

Equivalents in particular are compounds comprising a nucleobase, such as mononucleotides (mono-, di- or triphosphates of nucleosides), oligonucleotides,
10 polynucleotides, nucleobases and physiologically acceptable derivatives thereof that may be converted into the nucleoside as such or a nucleotide as such *in vivo*. Examples of such derivatives include various esters. WO 2002/088159 (Trommsdorff GmbH) relates to uridine esters, which may be used in accordance with the present invention. The contents of this publication regarding (deoxy)uridine esters is incorporated by reference.
15 Such equivalents are capable of increasing endogenous levels of the active forms of nucleosides in body tissues such as blood, liver and brain. Also synthetic compounds can be suitably included as nucleoside source, e.g. acylated derivatives of the nucleosides, for example triacetyl-uridine.

The composition according to the invention preferably comprises a pyrimidine
20 nucleoside or equivalent thereof, such as cytidine or equivalent thereof or a uridine or equivalent equivalent. More preferably, the composition according to the invention comprises a uridine or an equivalent thereof, preferably at least one uridine or an equivalent thereof selected from the group consisting of uridine (i.e. ribosyl uracil), deoxyuridine (deoxyribosyl uracil), uridine phosphates (UMP, dUMP, UDP, UTP),
25 nucleobase uracil and acylated uridine derivatives. Preferably, the composition according to the invention comprises an uridine phosphate selected from uridine monophosphate (UMP), uridine diphosphate (UDP) and uridine triphosphate (UTP). Most preferably, the composition according to the invention comprises UMP, as UMP is most efficiently being taken up by the body after oral administration. Hence, inclusion of UMP in the
30 composition according to the invention enables a high efficacy at the lowest dosage, the administration of a low volume to the subject or both. Uridine derivatives like UDP, which is readily formed from dietetic UMP, also appear to be important, in particular for transport of glycoproteins and glycolipids within the cell and availability thereof in the cytosol and plasma membrane.

Preferably, at least 20 weight% of the uridine or an equivalent thereof in the composition according to the invention is provided by UMP, more preferably at least 50 weight%, most preferably at least 90 weight%.

Preferably, the present use comprises the daily administration of uridine or an
5 equivalent thereof in a daily dosage of 0.08 to 3 g per day, preferably 0.1 to 2 g per day, more preferably 0.12 to 1 g per day.

Preferably, the present use comprises the daily administration of UMP in a daily dosage of 1.3 to 37.5 mg UMP per kilogram body weight of the subject to be treated. The required dosages of the equivalents of uridine on a weight basis can be calculated from
10 the daily dosage for UMP by taking equimolar amounts using the molecular weight of the equivalent and of UMP, the latter being 324 Dalton. The daily dosage of equivalents is preferably 3 to 115 μ mol per kg body weight per day, preferably 5 to 35 μ mol per kg body weight per day, or 0.25 to 9 mmol, preferably 0.3 to 6, most preferably 0.45 to 2.8 mmol per day.

Preferably, the present use comprises the daily administration of a combination
15 comprising uridine or an equivalent thereof in an amount of 0.06 to 2.4 g UMP per 100 ml liquid composition, preferably 0.08 to 1.6 g UMP per 100 ml liquid composition, more preferably 0.12 to 0.8 g per 100 ml liquid composition. Alternatively, the optimal dose for uridine monophosphate per 100 g dry matter is 0.18 to 7.2 g, preferably 0.24 to 5.4 g and
20 more preferably 0.36 to 2.4 g.

As a suitable cytidine equivalent cytidine can be used, for example as free base or as a salt, as an ester, as a phosphate derivative, like CMP, CDP or CTP, as cytosine, and as choline derivative, e.g. as citicoline. However, when both an uridine equivalent and a
25 cytidine equivalent are included simultaneously in the composition according to the invention it is preferred that the weight ratio of the sum of uridine and equivalents thereof to the sum of cytidine and equivalents thereof is larger than 1.0, more preferably at least 2.0, most preferably more than 5.0. Although cytidine is a precursor of uridine, it is more efficient and effective to include uridine in the composition according to the invention, because it passes more easily the blood brain barrier.

30 In some embodiments of the invention, useful sources of nucleosides include extracts of plant, animal, bacterial, algae or yeast material, e.g. in a composition according to the invention for individuals which don't suffer from a kidney disease. Examples of such extracts include heat-treated aqueous extracts from baker's yeast or brewer's yeast. In a further preferred embodiment, the composition according to the

invention preferably does not contain high amounts of other nucleotides. Hence, preferably the weight ratio sum of uridine and equivalents thereof to adenosine or its equivalents in the composition according to the invention is below 0.1, more preferably below 0.01, most preferably 0. Preferably, the weight ratio of the sum of the amount of uridine and equivalents thereof to the amount of guanosine or its equivalents in the composition according to the invention is below 0.1, more preferably below 0.01, most preferably 0. Preferably, the weight ratio of sum of uridine and equivalents thereof to inosine in the composition according to the invention is below 0.1, more preferably below 0.01, most preferably 0.

10

(ii) ω -3 polyunsaturated fatty acids (ω -3 PUFA's)

The composition according to the invention may comprise an ω -3 polyunsaturated fatty acid (ω -3 PUFA), in particular an ω -3 long chain polyunsaturated fatty acid (LCPUFA), more in particular selected from the group of docosahexaenoic acid (22:6 ω -3; DHA), docosapentaenoic acid (22:5 ω -3; DPA) and eicosapentaenoic acid (20:5 ω -3; EPA). Useful sources include fish oil, algae oil, eggs lipids and genetically modified organisms.

Preferably, the composition according to the invention comprises at least DHA, preferably DHA and EPA. More preferably, the combination comprises DHA and at least one precursor of DHA selected from EPA and DPA. More preferably, the composition according to the invention comprises DHA and EPA. The inventors recognized that only a part of the DHA incorporated in the brain originates from orally ingested DHA. An important part of the DHA incorporated in the brain is derived from conversion of DPA to DHA in the brain. In a further aspect, the composition according to the invention preferably contains a significant amount of EPA. EPA is converted to DPA (ω -3), increasing subsequent conversion of DPA (ω -3) to DHA in the brain. Hence, the composition according to the invention preferably also contains a significant amount of EPA, so to further stimulate in-vivo DHA formation.

The ω -3 PUFA's, in particular the LCPUFA's, more in particular DHA, DPA and EPA, may be provided in any form such as, but not limited to, triglycerides, diglycerides, monoglycerides, free fatty acids or their salts or esters, phospholipids, lysophospholipids, glycerol ethers, lipoproteins, ceramides, glycolipids or combinations thereof. Preferably, the composition according to the invention comprises at least DHA in triglyceride form.

Preferably, the present method comprises the daily administration of 200 to 5000

mg, more preferably 400 to 3000 mg, most preferably 800 to 2500 mg of the sum of DHA and EPA. The proportion of (DHA+EPA) relative to the total amount of fatty acids in the combination is preferably 5 to 50 weight%, more preferably 10 to 45 weight%, most preferably 15 to 40 weight%. Preferably, the present method comprises the daily
5 administration of 100 to 4000 mg, more preferably 120 to 1800 mg of DHA.

Preferably, the composition according to the invention comprises 1 to 40 weight% DHA based on total amount of fatty acids, preferably 3 to 36 weight% DHA based on total amount of fatty acids, more preferably 10 to 30 weight% DHA based on total amount of fatty acids in the composition according to the invention. The composition
10 according to the invention preferably comprises 0.5 to 20 weight% EPA based on total amount of fatty acids, preferably 2 to 10 weight% EPA based on total amount of fatty acids, more preferably 5 to 10 weight% EPA based on total fatty acids. The weight ratio of DHA to the sum of EPA and DPA is preferably larger than 1.0, more preferably 1.2 to 10, more preferably 2 to 8. The above-mentioned ratios and amounts take into account
15 and optimise several aspects, including taste (too high LC-PUFA levels reduce taste, resulting in a reduced compliance), balance between DHA and precursors thereof to ensure optimal effectiveness in relation to maximum dosage and possibility of product formulations such as liquid form, bar or capsule.

In one embodiment, the composition according to the invention contains a low
20 amount of arachidonic acid (AA; 20:4 ω -6). Arachidonic acid is believed to counteract the effects of the composition according to the invention. The present subjects normally ingest sufficient AA, or precursors thereof, and an excess daily dosage may stimulate inflammatory responses, inhibiting daily activities. Preferably, the weight ratio DHA/AA in the composition according to the invention is at least 5, preferably at least 10, more
25 preferably at least 15, up to e.g. 100. Preferably, the weight ratio EPA/AA is at least 2. The present method preferably comprises the administration of a composition comprising less than 5 weight% AA based on total amount of fatty acids, more preferably below 2.5 weight%. The ω -6/ ω -3 weight ratio of long-chain polyunsaturated fatty acids with at least 20 carbon atoms in the composition according to the invention is
30 advantageously below 0.5, preferably below 0.2. If the long-chain polyunsaturated fatty acids with 18 carbon atoms are also included in the ratio, the preferred ω -6/ ω -3 weight ratio is 0.05 to 1, more preferably 0.1 to 0.6, most preferably 0.15 to 0.4.

(iii) Vitamin B

The composition according to the invention comprises at least one vitamin B. The vitamin B is selected from the group of vitamin B1 (thiamine), vitamin B2 (riboflavin), vitamin B3 (niacin or niacinamide), vitamin B5 (pantothenic acid), vitamin B6 (pyridoxine, pyridoxal, or pyridoxamine, or pyridoxine hydrochloride), vitamin B7 (biotin), vitamin B9 (folic acid or folate), and vitamin B12 (various cobalamins). Preferably, at least one vitamin B is selected from the group of vitamin B6, vitamin B12 and vitamin B9. In particular, good results have been achieved with a combination comprising vitamin B6, vitamin B12 and vitamin B9. Advantageously, vitamin B12 and vitamin B9 are included because low plasma B12 or vitamin B9 levels are a risk factor for the development of Alzheimer's disease.

The vitamin B is to be administered in an effective dose, which dose depends on the type of vitamin B used. As a rule of thumb, a suitable minimum or a maximum dose may be chosen based on known dietary recommendations, for instance as recommended by Institute of Medicine (IOM) of the U.S. National Academy of Sciences or by Scientific Committee on Food (a scientific committee of the EU), the information disclosed herein and optionally a limited amount of routine testing. A minimum dose may be based on the estimated average requirement (EAR), although a lower dose may already be effective. A maximum dose usually does not exceed the tolerable upper intake levels (UL), as recommended by IOM.

If present in the nutritional or pharmaceutical composition according to the invention, the vitamin B6 is usually present in an amount to provide a daily dosage in the range of 0.5 to 100 mg, in particular in the range of 0.75 to 25 mg, more in particular in the range of 0.9 to 5 mg. If present in the nutritional or pharmaceutical composition according to the invention, the vitamin B12 is usually present in an amount to provide a daily dosage in the range of 0.5 to 1000 µg, in particular in the range of 1 to 100 µg, more in particular in the range of 1.5 to 10. If present in the nutritional or pharmaceutical composition according to the invention, the vitamin B9 is usually present in an amount to provide a daily dosage in the range of 50 to 5000 µg, in particular in the range of 150 to 1000 µg, more in particular in the range of 200 to 1000 µg.

In a preferred embodiment of the invention, the active components are included in a drink, preferably having a volume of about 125 ml, or in an alternative preferred embodiment in a product having a dry mass content of about 30 g, per packaging each of them being for consumption once a day. This means that the amounts per daily dose as mentioned above can be recalculated to a concentration per millilitre, by dividing the

above-mentioned value with 125, or to a concentration per g dry mass of the product by dividing by 30.

This way of calculation also applies to the other active components (i) to (vi) in the composition according to the invention when a desired dose per daily dose is disclosed.

5

(iv) Phospholipids

The composition according to the invention may comprise a phospholipid. As used herein, the term phospholipid includes lyso-phospholipids, de-acylated phospholipids and glycerophospholipids. It is preferred to include a phospholipid which is capable of increasing chylomicrons formation in elderly after administration of triglyceride lipids and can provide useful fatty acids. In particular, the phospholipid is selected from the group consisting of phosphatidylcholine(PC), phosphatidylethanolamine (PE), phosphatidylserine (PS), phosphatidic acid (phosphatidate), phosphoinositides (such as phosphatidylinositol (PI), phosphatidylinositol phosphate, phosphatidylinositol bisphosphate, phosphatidylinositol triphosphate) and sphingomyelin. In particular, the combination according to the invention comprises at least two different phospholipids selected from the group consisting of phosphatidylserine, phosphatidylinositol, phosphatidylcholine and phosphatidylethanolamine. Preferably the combination according to the invention comprises phosphatidylcholine, phosphatidylethanolamine and phosphatidylserine. Good results have been achieved with a combination of phosphatidylcholine (PC) and phosphatidylserine (PS), preferably in a weight ratio of 3:1.

For instance, lecithin may be used as a source for the phospholipids. Optionally the phospholipids are fortified with one or more phospholipids, such as a ceramide, a sphingolipid or a specific phospholipid, such as a phosphatidylcholine.

The phospholipid is to be administered in an effective dose. Usually, the total phospholipid daily dosage is in the range of 50 to 5000 mg, in particular in the range of 100 to 2000 mg, more in particular in the range of 150 to 1200 mg

Inclusion of phospholipids further beneficially improves membrane function, thereby enabling an improved functioning of the different parts of the brain that may be affected in prodromal subjects. Furthermore, the phospholipids improve stability and shelf life of the composition according to the invention. Phospholipids further enable the manufacturing of palatable compositions. Also, phospholipids are a source for choline

and prevent the decline in plasma choline levels after exercise. Choline is necessary for the formation of acetylcholine, a neurotransmitter involved in learning and memory and in the activation of muscles. These advantages are already achieved at relatively low phospholipid levels.

5

(v) Anti-oxidant

The composition according to the invention may comprise an antioxidant, in particular one or more antioxidants selected from the group of vitamin C, vitamin E and selenium. The vitamin C may be present as free acid (ascorbic acid) or as a salt, *e.g.* sodium ascorbate or potassium ascorbate. Suitable sources of vitamin E include (alpha-)tocopherol and tocotrienol. Suitable sources of selenium include selenate and selenite.

10

The anti-oxidant is to be administered in an effective dose. As a rule of thumb, a suitable minimum or a maximum dose may be chosen based on known dietary recommendations, for instance as recommended by the Institute of Medicine (IOM) of the U.S. National Academy of Sciences or by the Scientific Committee on Food (a scientific committee of the EU), the information disclosed herein and optionally a limited amount of routine testing. A minimum dose may be based on the estimated average requirement (EAR), although a lower dose may already be effective. A maximum dose usually does not exceed the tolerable upper intake levels (UL), as recommended by IOM. If present in the combination, vitamin C is usually present in an amount to provide a daily dosage in the range of 20 to 1200 mg, in particular in the range of 30 to 400 mg, more in particular in the range of 35 to 120 mg. If present in the nutritional or pharmaceutical composition according to the invention, the vitamin E is usually present in an amount to provide a daily dosage in the range of 8 to 200 mg, in particular in the range of 20 to 140 mg, more in particular in the range of 35 to 100 mg.

15

20

25

30

If present in the nutritional or pharmaceutical composition according to the invention, the selenium is usually present in an amount to provide a daily dosage in the range of 40 to 400 µg, in particular in the range of 50 to 200 µg, more in particular in the range of 55 to 80 µg.

Optionally one or more antioxidants may be present other than (v) the antioxidant selected from the group of vitamin C, vitamin E and selenium.

(vi) Choline

The nutritional or pharmaceutical composition according to the invention may comprise a choline. Choline refers to the various quaternary ammonium salts containing the N,N,N-trimethylethanolammonium cation. More specifically, choline is selected from the group of the choline cation, choline salts or esters, such as choline chloride, choline bitartrate, choline stearate or the like, or compounds that dissociate to choline, such as choline alfoscerate, sphingomyelin, cytidine-diphospho-choline or citicoline or CDP-choline, acylglycerophosphocholines, e.g., lecithin, lysolecithin, glycerophosphatidylcholine, and any mixture thereof. It is preferred to include a choline salt or choline alfoscerate into the composition according to the invention.

In particular, choline is to be administered in an amount to provide a daily dosage of 100 to 4000 mg, more in particular of 200 to 2000 mg.

Furthermore, the nutritional or pharmaceutical composition according to the invention may comprise one or more further micronutrients, for instance one or more micronutrients selected from the group of vitamins, minerals, and trace elements, taurine and inositol.

Protein and energy content

The composition, preferably the nutritional composition according to the invention allows improvement of BMI, or LBM, or frailty, without demanding to consume large amounts of protein or additional energy. Actually, the amount of energy in the composition according to the invention can remain limited to a value of less than 400 kcal (1680 kJ), preferably less than 280 kcal (1178 kJ) or more preferably less than 210 kcal (882 kJ), all expressed per daily dose for an adult human. The amount of energy is not needed to create the effect (as can be concluded from the experiment, which has an isocaloric control), but is simply the result of incorporating the active components into a small sized food product, which is thus used as a vehicle. In one embodiment, the components (i) to (v) are included in a pharmaceutical composition according to the invention having an energy content per serving unit of less than 130 kcal (546 kJ) or preferably less than 80 kcal (336 kJ).

In one embodiment, the amount of active components (defined as components (i) to (v)), which is needed to achieve the effect on BMI, etc, delivers more than 50 %, preferably more than 58 %, most preferably 62 to 88 % of the total amount of calories of

the composition as claimed. For the calculation of energy, the inventors use 9 kcal (37.8 kJ) per g of lipids or phospholipids, zero kiloJoule per g of nucleosides, vitamins and a choline equivalent, 4 kcal (16.8 kJ) per g of protein and digestible carbohydrate and 2 kcal (8.4 kJ) per g of fibre.

5 The protein amount in the composition according to the invention, if present, can remain relatively low, which can have important advantages to the largest part of individuals which suffer from weight loss or symptoms of frailty. In particular, it is preferred that composition according to the invention should not impair normal eating patterns, allow manufacture of a palatable product and not induce heartburn or
10 gastrointestinal discomfort after consumption thereof. The composition according to the invention appears effective, also when the concentration of protein in the composition according to the invention remains below 11 g per 100 ml and preferably below 9, more preferably 3 to 8.4 g, and most preferably 5.2 to 8.2 g per 100 ml of the composition according to the invention.

15 For concentrates, semi-dry and dry products, it is more convenient to express the concentration of protein on gram dry matter basis. The amount of protein is in these cases usually less than 400 mg, preferably less than 360 mg, more preferably 100-340, most preferably 150 to 330 mg per g dry matter. Simple proteins can be used in these amounts like milk protein. These technical features of the composition according to the
20 invention result in good compliance with the feeding protocol with such composition according to the invention and in very little adaptation of the normal diet after the composition according to the invention have been consumed according their recommended use.

 In order to avoid an undesired satiating effect, especially in elderly, the amount of
25 caseinates preferably is 4.5 weight% or vol% or less for liquid formula, and more preferably 0.5 to 4.0 weight% or vol% or most preferably 0.8 to 3.3 weight% or vol%. Therefore, but also for efficacy reasons, it is preferred to include a non-caseinate in the composition according to the invention, which is elaborated below.

 The effect of the active components (specific combinations of nucleosides, LC-
30 PUFAs, phospholipids, vitamins and a choline equivalent) allows efficacy by administering a nutritional composition according to the invention with the minimum amount of food volume, e.g. less than 150 ml per serving unit, which is also important because elderly, and especially frail elderly experience much earlier satiety when consuming food. In one embodiment of the invention the actives are provided to the

consumer, in a ready to use serving unit which provides 15 to 160 g of the effective composition according to the invention.

It has been suggested before, that additional protein of specific quality could improve LBM or increase body weight in elderly. Focus in the most relevant papers in this field is on the amount of protein (which should preferably be more than a few dozen g additional protein per daily dose), inclusion of L-leucine or including a large portion of essential amino acids in the protein amount, e.g. more than 15 g of the pure crystalline amino acids. Complying to such diet can be very demanding to many frail elderly, also because the taste of free amino acids is often considered as offensive or because consumption of such large amounts of protein has a significant impact on the consumption of regular diet.

The invention therefore provides a solution for the problem that an undesired amount of additional protein has to be administered for anabolic purposes, i.e. the improvement in BMI of elderly and it increases the improvement of the brain function as observed in frail elderly by consuming the combination of one or more of the active components (i) to (v) according to the invention.

The protein amount preferably comprises a non-caseinate protein for supporting an effect of the composition according the invention on BMI in frail elderly. In particular, specific whey protein relatively low in phosphorous, fish proteins, in particular cod protein, or a protein derived from eggs and proteins derived from vegetables, like potato, soy, pea, beans, lupin, quinoa and amaranth appeared suitable. The proteins can be intact, either heat-treated or non-denatured, or be partially hydrolyzed. Hydrolyses of the intact protein aims to improve its solubility, but degree of hydrolyses should be kept to the minimum, preferably to a degree of hydrolyses between 2 and 12, in order to maintain good organoleptic properties of the ready to use composition according to the invention. In a preferred embodiment, the amount of non-dairy protein is more than 21 weight%, more preferably more than 25 weight%, most preferably more than 42 weight%, in particular more than 51 weight% of the protein amount.

Suitable whey proteins include those which have a phosphorous content less than 400 mg/l when 100 g of the protein is dissolved in one litre of water. Preferably, this phosphorous content is 70 to 340 mg/l. The amount of protein can be calculated from the label of the product, or, when this is impossible or not justified by measuring Kjeldahl nitrogen by a method accepted as preferred in the prior art for the specific matrix, and multiplying this by 6.25 for mixtures of proteins and peptides.

If dairy proteins are included, it is preferred to include a whey protein. Such protein is preferably enriched in serum albumin or alpha lactalbumin. Such dairy amount also preferably comprises a lactoferrin. The concentration of alpha-lactalbumin as fraction of all whey proteins is preferably more than 25 weight%, that of serum albumin 5 to 12 weight%. The concentration of lactoferrin is preferably in the range 0.25 to 3 weight%, preferably 0.29 to 1.4 weight%, more preferably 0.34 to 1.2 weight% of the protein amount.

In cases where the fatty acids comprise that much omega 3 fatty acids that the ratio of the weight amounts of omega 3 to omega 6 exceed 5, the amount of whey protein is preferably less than 50 weight% of the protein amount, in order to keep the satiating character of the composition according to the invention as low as possible, while maintaining efficacy of the composition according to the invention.

It is also useful to include non-essential amino acids or their salts or esters. Examples of suitable amino acids include serine and aspartic acid. These amino acids can be administered as L-isomer or as a racemic mixture of L and D isomers of the particular amino acid. The amount aspartate plus asparagine in the composition according to the invention is preferably more than 8.4, more preferably 9.0 to 16, most preferably 9.5 to 15 g per 100 g amino acids in the composition according to the invention. The amount of serine in the ready to use composition according to the invention is preferably more than 4.1 g per 100 g amino acids. The presence of these two amino acids in the formula is thought to be at least partially responsible for the anabolic character of the formula and the observed efficacy in treating symptoms of frailty. Increasing the concentrations to the indicated amounts in the preferred embodiment increases their effect. In this respect, the effect on neurological symptoms in the low BMI elderly includes the effect on depressed mood, stamina and the activities of daily living. The inventors believe that an effect on metabolic pathways is responsible for this, in particular an effect on glucose metabolism and gene expression and not the amount of calories or amount of protein that are provided by the composition according to the invention.

The selected types and amounts of protein appeared to have the unexpected advantage in that long term administration to persons experiencing frailty or low body weight, especially in the aged population achieved better results in increasing BMI, lean body mass or body weight and muscle function, muscle strength and muscle mass, than most high protein dairy protein based formula, fortified with essential amino acids, and

demonstrated less adverse effects in sensitive persons.

It is of importance that in a preferred embodiment, the protein amount contributes to a better functioning of metabolism in order to support the maintenance of BMI, and lean body mass, and other symptoms of frailty, like exhaustion and fatigue and neurological function. Therefore, the protein preferably comprises a whey protein and more preferably a whey protein and a vegetable protein.

In a preferred embodiment, the osmotic value of the composition according to the invention is as low as possible, in order to facilitate easy stomach emptying. In a preferred embodiment, the composition according to the invention demonstrates an osmolality below 450 mEq/l.

In addition, the buffer strength of the composition according to the invention preferably low, in order to achieve rapid digestion and gut transfer of the composition according to the invention after consumption. This is done by using the amounts and types of proteins as indicated above and by preventing the use of high salt loads, in particular of citrates and phosphates. The amount of phosphorous in the composition according to the invention is preferably less than 150 mg, more preferably 20 to 110 mg, most preferably 50 to 72 mg per 100 ml. In one embodiment, the nucleotides in the composition according to the invention, like a uridine phosphate, or cytidine phosphate are replaced by their base. In a preferred embodiment, the composition according to the invention comprises a nucleobase and a nucleoside or nucleotide. In a preferred embodiment, the ratio of the weight amount of nucleobase to the sum of the corresponding nucleosides and nucleotides is more than 0.06, preferably 0.2 to 0.9.

The amount of organic acids, like citrates, is preferably less than 2, more preferably less than 1.2 weight%, most preferably 0.1 to 0.9 weight% of the digestible carbohydrate amount. These features are in particular relevant when the elderly suffer from diagnosed anchlorhydria. This is a major problem in elderly and in particular in institutionalized elderly or elderly having a BMI under 23.5 kg/m² or frail elderly.

In a preferred embodiment, the composition according the invention is a liquid formula having a viscosity of less than 60, more preferably 2 to 30 mPa.s, measured at 20 degrees °C. As used herein, the viscosity is the viscosity as measurable using a Anton Paar Physica MCR301 rheometer with aCP50-1/PC cone (diameter 50 mm, 1° difference between middle and outside) at 20 °C at 100s⁻¹.

Dosage form

Further, one or more additional ingredients may be present that are commonly used in the prior art, dependent on the form - nutritional or pharmaceutical composition - in which the combination is provided.

5 If the dosage form is a pharmaceutical composition, the pharmaceutical composition may comprise one or more excipients known in the prior art to provide a pharmaceutical composition in a dosage form of choice. The pharmaceutical composition is preferably for enteral application (orally or via tube-feeding). Examples of solid formulations are tablets, capsules (*e.g.* hard or soft shell gelatine capsules), pills,
10 sachets, powders, granules and the like which contain the active ingredients together with a conventional carrier. Any conventional carrier material can be utilized. The carrier material can be organic or inorganic inert carrier material suitable for oral administration. Suitable carriers include water, gelatine, gum Arabic, lactose, starch, magnesium stearate, talc, vegetable oils, and the like. Additionally, additives such as
15 flavouring agents, preservatives, stabilizers, emulsifying agents, pH-buffers and the like may be added in accordance with accepted practices of pharmaceutical compounding. While the individual active ingredients are suitably administered in a single composition they may also be administered in individual dosage units.

 If the dosage form is a nutritional composition, the composition generally
20 comprises at least one macronutrient for providing (additional) energetic value to the nutritional composition. The macronutrient may suitably be selected from the group of proteinaceous matter (proteins, peptides, amino acids), fat, other than component (ii), and digestible carbohydrates.

 Suitable proteinaceous matter, lipids and carbohydrates, and suitable
25 concentrations of the macronutrients may be based on known dietary guidelines for food products, in particular for food products for the elderly. Suitable formulations may *e.g.* be based on known commercially available clinical foods, or foods advertised for feeding elderly people or for feeding people suffering from dementia.

 Regarding the lipid, preferably one or more triglycerides are present. These may be
30 selected from vegetable oils and fats and animal oils and fats.

 Regarding the digestible carbohydrates, these may in particular be selected from digestible pentoses, digestible hexoses digestible oligosaccharides, *e.g.* digestible disaccharides and digestible trisaccharides, and digestible polysaccharides (*e.g.* starch).

More specifically one or more digestible carbohydrates may be chosen selected from the group of galactose, mannose, ribose sucrose, trehalose, palatinose, lactose, maltodextrose, maltose, glucose, fructose, including oligomers and polymers thereof.

Optionally, a nutritional composition according to the invention comprises one or
5 more non-digestible carbohydrates (dietary fibres) such as oligosaccharides. As used herein, the term oligosaccharides in particular refers to saccharides comprising 3 to 25 monosaccharide units per molecule. The oligosaccharide(s) may in particular be selected from the group of fructo-oligosaccharides (FOS), galacto-oligosaccharides (GOS), trans-
10 galacto-oligosaccharides (TOS), xylo-oligosaccharides (XOS), soy oligosaccharides, and the like. Optionally, also higher molecular weight compounds such as inulin, resistant starch and the like may be incorporated in the composition according to the invention.

Further, the nutritional composition may comprise a probiotic.

Further, the nutritional composition may comprise one or more additives commonly used in food technology, such as one or more additives selected from the
15 group of flavourings, stabilisers, preservatives, colourants, emulsifiers, pH-buffers *etc.*

The nutritional composition for use in a accordance with the invention may be a solid composition, a semi-solid composition (such as a paste or a gel) or a liquid composition, such as a beverage of a drinkable food product.

The nutritional composition according to the invention may in particular be
20 intended for enteral administration (orally or by tube feeding). Alternative forms of administration may be applied, in particular parenteral administration. The skilled person will be able to formulate a suitable product for parenteral administration, in particular by preventing inclusion of non-endogenous proteinaceous material which may induce an allergic reaction or other adverse effects. The administration may be carried
25 out based on a manner known *per se* for a specific type of nutritional composition.

In particular, the nutritional composition may be selected from the group of spreads; yoghurts, custards, ice-creams, butter, and other dairy products; dairy-substitute products; drinks, such as fruit drinks; candy bars; cookies, cakes and other bakery products; and drinkable foods.

30 The total energetic value of the composition may be chosen within wide limits and may range, *e.g.*, from 0.2 to 4 kcal/g. In particular, the energetic value may be at least 0.4 kcal/g, more in particular at least 0.8 kcal/g. In particular, the energetic value may be 5 kcal/g or less, more in particular 3 kcal/g or less.

In case the nutritional composition is a fluid, it usually has a nutritional value of at least 20 kcal/100 ml, preferably of at least 50 kcal/100 ml, in particular of at least 75 kcal/100 ml or at least 100 kcal/100 ml. For a fluid composition the nutritional value is usually 300 kcal/100 ml or less, in particular 200 kcal/100 ml or less, more in particular
5 150 kcal/100 ml or less.

Suitable dosage forms, active ingredients, further components that may be co-administered and ways of administration are as described for the nutritional or pharmaceutical composition as described herein above, the claims, or the examples herein below.

10 The vitamin B, the phospholipid, the antioxidant and – if present – further active ingredients, may be administered under the supervision of a medical specialist or be self-administered.

Specific embodiment: nutritional composition

As mentioned above, the invention also relates to a nutritional composition
15 comprising (i) at least one component selected from the group of uridine and uridine monophosphate, (ii) DHA and EPA (which may be present in bound form for e.g. as a triglyceride ester) (iii) a vitamin B, (iv) a phospholipid, v) an antioxidant, and (vi) a choline. Preferred sources, preferred specific compounds for each group, concentrations, dosages and other product properties may generally be as described herein above. The
20 nutritional composition according to the invention usually also comprises, protein. Further, digestible carbohydrate and fibre, are usually present

Preferably, the nutritional composition comprises i-a) the nucleoside uridine and i-b) the nucleotide uridine monophosphate. i-a) Uridine and i-b) uridine monophosphate are preferably present in a weight to weight ratio in the range of 0.2:1
25 to 0.7:1. The presence of both i-a) uridine and i-b) uridine monophosphate, especially in said ratio, is in particular advantageous because it provides a more uniform bioavailability after oral administration, which increases the number of responders to the composition (used) according to the invention (subjects reacting positively on treatment with a composition in accordance with the invention), without any
30 unacceptable adverse effects. In addition, the nucleoside uridine may decrease activation of P2Y receptors in the gut.

In a preferred embodiment, the total content of uridine plus uridine monophosphate is in the range of 5-30 mg per gram dry weight, more preferably in the

range of 8-20 mg per gram dry weight, in particular in the range of 10-18 mg per gram dry weight

The total content of uridine plus UMP as a weight percentage of total nucleoside equivalents preferably is more than 28 wt. %, more preferably 40-100 wt. %, most preferably 60-100 wt. %, in order to obtain a selective effect of the nucleotide fraction in the product of the invention, such as in order to avoid triggering an undesired effect in the enterocytes or liver or the enteral nervous system, due to activation of receptors, for example the P2X or P2Y receptors, by adenosine-based or guanosine-based nucleotides.

The weight to weight ratio of UMP to GMP (guanosinemonophosphat) preferably is more than 10. The weight to weight ratio of UMP to UMP (inosine monophosphate) preferably is more than 10.

In a specific embodiment, the nutritional composition according to the invention, comprises 3-14 mg EPA per gram dry matter, preferably 5-10 mg EPA per gram dry matter and 12-56 mg DHA per gram dry matter, preferably 2.5-20 mg DHA per gram dry matter. Preferably, the sum of DHA and EPA is 5-50 wt. % based on total fatty acids.

A composition according to the invention preferably has a low arachidonic acid (AA), content, if present at all, especially in an embodiment wherein on or more of component (ii) ω -3 PUFAs selected from EPA, DHA and DPA are present. AA is thought to have a disadvantageous effect on the effectiveness of these components in a use according to the invention. In view thereof, the weight to weight ratio of the sum of DHA+DPA+EPA to AA preferably is higher than 5, in particular 6 or higher, more in particular 12 or higher. In a particularly preferred embodiment, the weight to weight ratio of DHA to AA is higher than 5. The ratio long chain ω -6 PUFAs to long chain ω -3 PUFAs preferably is 0.05 to 1. The term 'long-chain' is used herein for PUFA's having a carbon chain of at least 20 carbon atoms.. In a particularly preferred embodiment, the weight to weight ratio of DHA to AA is higher than 5. The ratio ω -6 PUFAs to ω -3 PUFAs preferably is 0.05 to 1.

The ratio [uridine monophosphate + uridine]/[phospholipids] of a nutritional composition according to the invention is usually less than 5.9, preferably 0.15-4, in particular 0.20-2.4, more in particular 0.25-0.71. A relatively low ratio of [uridine monophosphate + uridine]/[phospholipids] is in particular preferred in a composition comprising a protein. In particular in such embodiment, it contributes to a better efficacy with respect to a use of the invention, in particular with respect to providing a palatable effective product a decrease of uridine concentration and an increase of phospholipid

concentration allows better tasting while maintaining efficacy. A relatively low ratio is also is advantageous for improving the palatability, especially during shelf-life.

The nutritional composition of the invention usually has an energy density of less than 13 kcal per gram dry matter, preferably of 3-9.3 kcal per gram dry matter, more preferably of 4.0-7.0 kcal per gram dry matter.

The nutritional composition according to the invention preferably has a phospholipid content in the range of 3.3-67 mg/gram dry matter, more preferably in the range of 5-33 3.3-67 mg/gram dry matter, in particular in the range of 7-14 mg/gram dry matter.

In a specific embodiment, the nutritional composition according to the invention, comprises phosphatidylcholine (PC). In such embodiment the weight to weight ratio phosphatidylcholine to choline is usually more than 0.1, preferably more than 0.26, in particular 0.30-6, more preferably 0.36-3. Herein, the amount of choline in grams is to be calculated as the molar contribution of choline as provided by all choline sources (when orally digested and assuming 100% bioavailability, including PC), times the molecular weight of choline (104 g/mol). Herein., the molecular weight of PC 810 gram/mol. So for example, including 400 mg choline chloride and 200 mg phosphatidylcholine and 200 mg PL's other than PC would then result in a weight ratio of PC to choline of $200 / [(104/139.6) \times 400 + (104/810) \times 200] = 200 / [298 + 25.7] = 0.62$

The presence of PC is in particular preferred as a source for choline, because less PC is required than equimolar choline ingredient (*i.e.* *N,N,N*-trimethylethanolammonium cation) for obtaining the same choline concentrations in blood. Herewith choline (salt) content can be reduced, whilst still providing a choline source. Advantageous thereof include avoiding a fishy odour of product and/or subject treated with the product; avoiding irritation of the mucous membranes in case of xerostomia.

The nutritional product usually comprises a protein. The protein content is preferably less than 400 mg per gram dry matter, more, preferably 100-340 mg protein per gram dry matter. A relatively low protein content is preferred to decrease consequences on diet intake. In particular, a relatively low protein content is desired in view of a satiating effect a high protein content has, which may cause the subject to consume an insufficient amount of the nutritional product. For improved palatability and/or for improving brain function. In a liquid product, the protein content preferably is less than 11 g/100 ml, more preferably 5.2-8.2 g/l. In a specific embodiment, the protein

content is more than 7 g/100 ml.

In an advantageous embodiment, the nutritional composition comprises at least a whey protein. In a liquid product, the dairy protein content (whey protein, casein, caseinate) preferably is less than 4 wt./vol %.

5 In an advantageous embodiment, the nutritional composition according to the invention comprises a protein selected from the group of fish proteins (in particular cod protein), egg-protein and vegetable proteins. If present, the total content of non-dairy protein is usually more than 21 wt. % preferably 22-80 wt. %, in particular 25-40 wt. %, based on total protein content. Non-dairy proteins, such as vegetable, fish or egg protein
10 is an advantageous protein source, amongst others because they are less satiating than dairy proteins. Another reason to include such non-dairy protein is an improved endocrine response action, which results in good postprandial glucose response, compared to a dairy protein, in particular compared to casein/caseinate.

A vegetable protein may in particular be selected from the group of potato protein ,
15 soy protein, pea protein, beans protein, lupin protein, quinoa protein and amaranth protein.

A composition comprising whey protein and amaranth or pea protein is in particular suitable for increasing BMI. The weight ratio whey protein to the sum of amaranth and pea protein may in particular be 50:50 to 90:10, more in particular 60:40
20 to 80:20.

A composition comprising whey protein, soy protein and pea protein is in particular suitable for increasing ADL in subjects suffering from early physical exhaustion after exercise. When present in combination, the whey protein content preferably is 30-70wt. % of the protein fraction, the pea protein fraction preferably is 15-
25 35 wt. % of the protein fraction, and the soy protein fraction preferably is 15-35 wt. % of the protein fraction. One or more other proteins from other sources may be present in a product comprising proteins from these three protein sources, e.g. casein, usually in a total concentration of up to 40 wt. % of the protein fraction.

The dairy protein or the non-dairy protein may be non-hydrolysed or may be
30 partially hydrolysed. The degree of hydrolysis may in particular be 2-12.

In a specific embodiment, a composition (for use) according to the invention comprises at least one amino source providing L-serine and/or L-aspartic acid. The amino acid source can be selected from free amino acids, including salts thereof, peptides

(oligopeptides, polypeptides, proteins), comprising and L-serine unit and/or L-aspartic acid units. If present, the L-serine content preferably is higher than 4.1 g per 100 gram amino acids, including amino acids in peptides and other compounds providing an amino acid when digested. If present, the L-aspartic acid content is preferably 8.4-15 g per 100 gram amino acids including amino acids in peptides and other compounds providing an amino acid when digested.

Preferably, the nutritional composition comprises 0.15-0.5 g digestible carbohydrate per gram dry weight, in particular 0.20-0.40 g digestible carbohydrate per gram dry weight. Preferably, the digestible carbohydrate fraction provides glucose, fructose and galactose (present as monosaccharide or in oligo/polysaccharide form).

Preferably 0.01-0.1 g fibre per g dry weight is present in a composition according to the invention. In particular, the fibre content may be in the range of 0.02-0.08 g per g dry weight.

The lipid content in a nutritional composition according to the invention preferably is 0.15-0.3 g lipids per gram dry weight.

Further, the composition may comprises additional vitamins and/or minerals.

In particular good results have been achieved with a liquid composition according to the invention.

The liquid composition usually has a dry matter content of 15-30 g per 100 ml. The dry matter content preferably is 24 g per 100 ml or less, in order to aid water homeostasis, in particular 16-24 g per 100 ml, more in particular 17-22 g per 100 ml.

The liquid composition preferably has an osmolarity of less than 450 mEq/l, in particular of 120 to 150 mEq/l.

The viscosity of a liquid nutritional composition according to the invention is usually less than 200 mPa.s

The liquid composition may in particular be packaged as a unit dose packaging, which may in particular have a liquid composition content in the range of 50-250 ml, more in particular in the range of 100-150 ml.

In a specific embodiment, the composition is a powder, in particular a powder that is reconstitutable with water to provide a liquid product according to the invention

The invention will now be illustrated by the following examples and the experimental part, without being bound or restricted thereto.

EXPERIMENTAL

Example 1

APP/PS1 mice, a mouse model for AD, were provided with a diet enriched with DHA and UMP (intake per day: DHA=22.8 mg; UMP=23.1 mg) or with a control diet for 3 months. The two diets were isocaloric and differed only with respect to DHA and UMP content. The amount of fat, carbohydrates and protein was the same between diets. During the 3 months of diet intervention body weight was monitored as was food intake.

Results. Mice fed the enriched diet showed – on average – a 17.5% increase in body weight after 3 months compared to 14% for the control mice. Food intake was slightly lower in the group which were fed the enriched diet (on average 2.88 gram per day) compared to control mice (on average 3.08 gram per day).

Conclusion. Alzheimer's disease mice showed an increase in body weight when fed with the composition according to the invention (the ω -3 polyunsaturated fatty acid DHA and the nucleoside UMP), not attributed to a caloric increase in body weight.

Example 2

APP/PS1 mice, a mouse model for AD, were provided with a diet enriched with B-vitamins (B6, B12 and folic acid; intake per day: B6=0.1 mg ; B12=0.11 μ g ; folic acid=18.1 μ g), phospholipids (lecithin; intake per day: 12.4 mg) and antioxidants (vitamin C, vitamin E, selenium; intake per day: vitamin C=4.8 mg; vitamin E=4.5 mg; selenium=3.2 μ g) or with a control diet for 3 months. The two diets were isocaloric and differed only with respect to B-vitamins, phospholipids and antioxidants content. The amount of fat, carbohydrates and protein was the same between diets. During the 3 months of diet intervention body weight was monitored as was food intake.

Results. Mice fed the B-vitamins+phospholipids+antioxidants enriched diet showed a 18 % increase in body weight after 3 months compared to 14 % for the control mice. Food intake was slightly lower in the B-vitamins+phospholipids+antioxidants group (on average 2.86 gram per day) compared to control mice (on average 3.08 gram per day).

Conclusion. Alzheimer's disease mice showed an increase in body weight when fed with the composition according to the invention (B vitamins, a phospholipid, and antioxidants), not attributed to a caloric increase in body weight.

Example 3

APP/PS1 mice, a mouse model for AD, were provided with a diet enriched with DHA+UMP (intake per day: DHA=22.8 mg; UMP=23.1 mg) (as in Example 1), B-
5 vitamins (B6, B12 and folic acid; intake per day: B6=0.1 mg; B12=0.11 µg; folic acid=18.1 µg), and phospholipids (lecithin; intake per day: 12.4 mg) or with a control diet for 3 months. The two diets were isocaloric and differed only with respect to DHA, UMP, B-
vitamins and phospholipids content. The amount of fat, carbohydrates and protein was the same between diets. During the 3 months of diet intervention, body weight was
10 monitored as was food intake.

Results. Mice fed the enriched diet showed a 20 % increase in body weight after 3 months compared to 14 % for the control mice. Food intake was the same in the ω-3 PUFA+nucleoside+B-vitamins+phospholipids group (on average 3.02 gram per day) compared to control mice (on average 3.08 gram per day).

15 *Conclusion.* Alzheimer's disease mice showed an increase in body weight when fed with the composition according to the invention (ω-3 polyunsaturated fatty acids, a nucleoside, B vitamins, and phospholipids), not attributed to a caloric increase in body weight.

20 Example 4

APP/PS1 mice, a mouse model for AD, were provided with a nutritional composition comprising uridine-5'-monophosphate (intake per day: UMP=23.1 mg); omega-3 fatty acids (including DHA) (intake per day: DHA=22.8 mg); choline (intake per day: 12 mg); phospholipids (intake per day: 12.4 mg); B vitamins (intake per day: B6=0.1 mg;
25 B12=0.11µg; folic acid=18.1 ug), and antioxidants (intake per day: vitamin C=4.8 mg; vitamin E=4.5 mg; selenium=3.2 ug) or with a control diet for 3 months. The two diets were isocaloric. The amount of fat, carbohydrates and protein was the same between diets. During the 3 months of diet intervention, body weight was monitored as was food intake.

30

Results. Mice fed the nutritional composition showed a 25 % increase in body weight after 3 months compared to 14 % for the control mice. Food intake was the same in the

group fed the nutritional composition (on average 3.10 gram per day) compared to control mice (on average 3.08 gram per day).

Conclusion. Alzheimer's disease mice showed an increase in body weight when fed with the composition according to the invention (nucleoside : UMP; ω -3 polyunsaturated fatty acids (including DHA); choline; phospholipids; B vitamins, and antioxidants), not attributed to a caloric increase in body weight.

Example 5

Human patients older than 65 years of age and suffering from a cognitive impairment were provided for 3 months with a once a day drink (125 ml) enriched with EPA (300 mg) and DHA (1200), phospholipids (160 mg lecithin), choline (400 mg), UMP (625 mg), antioxidants (40 mg vitamin E, 80 mg vitamin C, 60 μ g selenium) B-vitamins (3 μ g B12, 1 mg B6, and 400 μ g folic acid), and or with a control drink (125 ml). The two drinks were isocaloric and differed only with respect to DHA, UMP, B-vitamins, phospholipids and antioxidants content. The amount of fat, carbohydrates and protein was the same for both diets.

The compositions of the drinks were as given in Table 1.

Results Patients provided with the enriched drink showed an increase in BMI compared to Alzheimer's disease patients provided with the control drink (Table 2). Furthermore, the enriched drink was effective on ADL (ADCS score) in elderly of BMI < 26 (Figure 2).

Table 1 : Isocaloric drinks used in Example 5.

Component	Invention	Control
Amount	125 ml	125 ml
Macronutrients		
Energy, kcal	125	125
Protein (caseinate/WPI), g	3.8	3.8
Carbohydrate, g	16.5	16.5
Fat, g	4.9	4.9
Components according to invention		
EPA, mg	300	0
DHA, mg	1200	0
Phospholipids, mg	106	0
Choline, mg	400	0
UMP (uridine monophosphate), mg	625	0
Vitamin E (alpha-TE), mg	40	0
Vitamin C, mg	80	0
Selenium, µg	60	0
Vitamin B12, µg	3	0
Vitamin B6, mg	1	0
Folic acid, µg	400	0
Minerals		
Sodium, mg	125	125
Potassium, mg	187.5	187.5
Chloride, mg	156.3	156.3
Calcium, mg	100	100
Phosphorus, mg	87.5	87.5
Magnesium, mg	25.0	25.0
Other trace elements		
Iron, mg	2	2
Zinc, mg	1.5	1.5
Iodine, µg	16.3	16.3
Manganese, mg	0.41	0.41
Copper, µg	225	225
Molybdenum, µg	12.5	12.5
Chromium, µg	8.4	8.4
Other vitamins		
Vitamin A, µg	200	200
Thiamin (B1), mg	0.19	0.19
Riboflavin (B2), mg	0.20	0.20
Niacin (B3), mg NE	2.25	2.25
Pantothenic acid (B5), mg	0.66	0.66
Vitamin D, µg	0.88	0.88
Biotin, µg	5.0	5.0
Vitamin K, µg	6.6	6.6

Abbreviations : EPA, eicosapentaenoic acid; DHA, docosahexaenoic acid; TE tocopherol equivalents, NE niacin equivalent.

5 **Table 2 : Results of BMI increase (Example 5)**

	BMI at Baseline	BMI after 3 months	BMI after 6 months
Invention	26.198	26.469	26.58*
Control	26.175	26.353	26.348

* $p = 0.088$

Example 6: Example compositions according to the invention.

The following compositions according to the invention may be used for the healthy
 5 improvement of body weight in an elderly person experiencing one or more symptoms of
 frailty, preferably in elderly having a BMI below 26 and more specifically below 23.5
 kg/m² :

Table 3 : Composition A (per 100 g of ready-to-use composition)

Pyrimidine nucleoside	0.3 g of a mixture of uridine and uridine monophosphate
Vitamin B	1 mg B6, 2 µg B12 and 300 µg folic acid or folate
and optionally :	
A phospholipid	0.2 g lecithin
A choline	1 g choline as choline chloride
Lipids	2 g marine or algae oil, comprising 30-60 g of DHA, DPA or EPA per 100 g fatty acids

10

Table 4 : Composition B (sip feed, amount per 100 ml)

Ingredient	Amount
Energy	≥150 kcal (≥ 630kJ)
Protein	≥ 3.5 g
Digestible carbohydrates	16 g
Fat	5.2 g
DHA	300 mg
EPA	75 mg
Phospholipids	50 mg
UMP	160 mg

Ingredient	Amount
Choline	100 mg
Vitamin B6	0.3 mg
Vitamin B12	0.8 µg
Folic acid	100 µg
Vitamin C	20 mg
Vitamin E	10 mg
Selenium	15 µg
Vitamin D	1.2 µg
Calcium	200 mg

Table 5 : Composition C to F

Components	Composition C	Composition D	Composition E	Composition F
Macronutrients				
Energy (kcal)	140	100	150	140
Protein (g)	8 (milk protein + blend soy prot/ α -lac)	3 (milk protein)	10 (ultra filtrated milk protein + free leucine)	8 (pea protein, casein, α -lac)
Lipids (EPA, DHA, phospholipids) (g)	5.8	3.9	5.1	5.1
Digestible carbohydrates (g)	14	13.2	16	14
Fiber (g)	0.2	0	0.2	0
EPA (mg)	120	240	240	200
DHA (mg)	480	960	960	800
Phospholipids (mg)	210	128	128	160
Choline (mg)	200-300	320	320	260
UMP (mg)	250	500	500	400
Vitamin E (mg α -TE)	16 32	32 64	32 64	25 50

Components	Composition C	Composition D	Composition E	Composition F
Vitamin C (mg)	67	48	48	50
Selenium (µg)	2.4	2.4	2.4	2
Vitamin B12 (µg)	0.8	0.8	0.8	1
Vitamin B6 (mg)	320	320	320	280
Folates (µg)				
Sodium (mg)	100	100	100	100
Potassium (mg)	100-200	150	100-200	100-200
Chloride (mg)	100-150	125	100-150	100-150
Calcium (mg)	100-300	80	100-300	100-300
Phosphorus (mg)	100-300	70	100-300	100-300
Magnesium (mg)	32	23.2	23.2	23.2
Iron (mg)	1.6	1.6	1.6	1.4
Zinc (mg)	2.4	1.2	1.2	2
Iodine (µg)	26	13	13	20
Manganese (mg)	0.88	0.33	0.33	0.4
Copper (µg)	180	180	180	150
Molybdenum (µg)	26	10	10	20
Chromium (µg)	6.7	6.7	6.7	8
Vitamin A (µg)	160	160	160	140
Thiamine (mg)	0.21	0.15	0.15	0.16
Riboflavin (mg)	0.23	0.16	0.16	0.18
Niacin (mg)	2.5	1.8	1.8	2
Pantothenic acid (mg)	1.4	0.53	0.53	0.6
Vitamin D (µg)	4-7	0.7	0.7	2
Biotin (µg)	10.5	4.0	4	6
Vitamin K (µg)	14	5.3	5.3	6

Composition G : Product comprising per 1000 litres (about)

Ingredient	Amount (kg)
Ultrafiltrated dairy protein	700
Superrefined Tuna Fish Oil	18

Lecithin Powder	0.9
Sugar	42
Maltodextrin	118
Municipal Water	164
Rapeseed Lear-Sunflower-High Oleic Blend	32
Choline chloride	2
UMP	3.7
Potassium hydroxide	0.27
Citric Acid Monohydraat	1.3
Magnesium hydroxide	0.6
Tri-potassium citrate.1aq	0.9
Mineral Premix	0.2
Vitamin Premix	1.0
Vanilla Flavour	1.2

Example 7: Tool for assessing the degree of frailty of a patient

Consists of answering a fixed set of questions, each related to one or more of the aspects
5 of frailty, and scoring the replies by comparison with normal values.

In order to correctly answer the questions, specific measurements are recommended, as described in the text.

The questions can be asked by an expert and orally replied to; alternatively they can also be put on a paper or form and the replies written down; however, preferably the
10 questions can be raised on a screen or monitor of an electronic device, like a small computer, tablet or peripheral station of a central computer system.

The electronic device can have the normal values stored and can be equipped with software which allows automatic calculation of scores per parameter and of the frailty index.

15 The algorithm to calculate the final score can be subject to individual medical expertise. However, it is preferred to operate with a common algorithm to allow comparison of diagnosis. In the text of the document, it is described how this can occur. It is preferred to score the value of each individual parameter by comparison with age-matched controls; when the patient scores positive on two of the parameters as given below the
20 patient is defined to be prefrail and can be helped by the composition according to the invention according the invention. When the patient scores positive on at least three parameters, the individual is diagnosed to be frail and to be receptive for improvement by use of the composition according to the invention.

Table 6 : Example of a questionnaire for scoring frailty.

Parameter	Measurements	Normal value/ Score
Muscle weakness	Muscle: method: Value	
Excessive feelings of exhaustion or fatigue	Mental fatigue Perceived fatigue after exercise: Activities of daily living (ADL)/ sleep needed Muscle capacity or power Speed of developing fatigue Lung capacity	
Abnormally low physical activity	Voluntary normal activities Capabilities to apply instrumental activities	
Slow or unsteady gait	One or more or combinations of balance, effort (walking time) and coordination	
Weight loss	Average involuntary weight loss (BMI, LBM) Absence of acute weight loss or an acute phase response	
Neurological dysfunction	Cognitive impairment Memory disorder Sensory impairment Motoric impairment Chronic pain Depression Sleep disorder Anxiety disorder	
TOTAL SCORE	Prefrail: 2 Mildly frail: 3 Moderately frail: 4	

	Severely frail: 5-6	
--	---------------------	--

Example: 8 Further formulation examples

5 **Table 7: Ready to feed liquid formulations**

Component	Composition H ¹	Composition I ²	Composition J ³
Volume per packaging	125 ml	125 ml	125 m;
Energy density (kcal/ml)	1.0	1.6	1.6
Protein (g/100 ml) Ingredients/ components	7.5 (i) Whey and (ii) amaranth - or pea protein (70/30)	10 Whey protein	8 Whey, -soy and wheat protein and casein (35:20:20:25)
Lipids (g/100 ml) Ingredients/ components	5.1 Phospholipids = 220 mg Fish oil/ rape seed/ lecithin 10:7:1 by wt, providing 600 mg DHA and 150 mg EPA	8 PL's 400 mg Marine oil/ corn oil/ palm oil/lecithin to result in 600 mg and 150 mg EPA	8 Lecithin 420 mg Marine oil/ soy/ canola/ lecithin to get 600 mg DHA and 150 mg EPA
Digestible Carbohydrates (g/100 ml)	6 after hydrolyses of ingredients by weight: 10 % galactose, 60% glucose, 10% fructose, and 10% other mono- saccharides	9.5	11.5
Fiber (g/100 ml)	1.5 Inuline hydrolysate + galacto-OS and manno OS (60: 20: 20 by wt)	1.5 Inuline hydrolysate + galacto-OS and manno OS (60: 20: 20 by wt)	1.5 Inuline hydrolysate + galacto-OS and manno OS (60: 20: 20 by wt)
Choline (mg/100 ml)	200 Choline alfosclerate	200 Choline alfosclerate	200 Choline alfosclerate
Uridine source (mg/100 ml)	310 Uridine + UMP 4:6 by wt	320 Uridine + UMP 2 : 8 by weight	310 Uridine + UMP 3: 7 by weight
Vitamins (per 100 ml)	Folates: 200 ug B12: 6 ug B6: 1 mg Vitamin E: 20 mg Vitamin C: 40 mg Vitamin premix providing all vitamins except the	Folates: 200 ug B12: 6 ug B6: 1 mg Vitamin E: 20 mg Vitamin C: 40 mg Vitamin premix providing all vitamins except the	Folates: 200 ug B12: 6 ug B6: 1 mg Vitamin E: 20 mg Vitamin C: 40 mg Vitamin premix providing all vitamins except the above in

	above in 0.2xRDA of FDA 1988	above in 0.2xRDA of FDA 1988	0.2xRDA of FDA 1988
Minerals (per 100 ml)	Sc: 30 ug Fe: 2 mg Zn: 1.5 mg Mg: 25 mg Ca: 120 mg Na : 120 mg K: 180 mg Cl: 150 mg P: 90 mg I: 16 ug Mn: 0.5 mg Cu: 0.23 mg Mo: 13 ug Cr: 8 ug	Sc: 30 ug Fe: 2 mg Zn: 1.5 mg Mg: 25 mg Ca: 120 mg Na : 120 mg K: 180 mg Cl: 150 mg P: 90 mg I: 16 ug Mn: 0.5 mg Cu: 0.23 mg Mo: 13 ug Cr: 8 ug	Sc: 30 ug Fe: 2 mg Zn: 1.5 mg Mg: 25 mg Ca: 120 mg Na : 120 mg K: 180 mg Cl: 150 mg P: 90 mg I: 16 ug Mn: 0.5 mg Cu: 0.23 mg Mo: 13 ug Cr: 8 ug

¹Product in particular suitable for increasing body mass index

²Product in particular suitable for increasing lean body mass (LBM) in non-frail elderly

³ Product in particular suitable for increasing ADL in persons suffering from early
5 physical exhaustion after exercise

Example 9: improving BMI and/or ADL in patients with mild AD

Methods

- 10 The composition according to the invention shown in Example 5 (Table 1) was used for
was used for a 12-week, double-blind, randomized, controlled, multicenter study with a
similarly designed, exploratory optional 12-week extension period. The detailed
methodology and primary results of the study have been described previously (Scheltens
P, Kamphuis PJ, Verhey FR, Olde Rikkert MG, Wurtman R-J, Wilkinson D, et al.
15 Efficacy of a medical food in mild Alzheimer's disease: A randomized, controlled trial.
Alzheimers Dement. 2010 Jan;6(1):1-10 e1). Briefly, patients with mild AD (MMSE score
of 20–26) were recruited and randomized 1:1 to receive either the composition of the
invention or an iso-caloric control product as a 125 ml drink to be taken every day. In
line with current guidelines for clinical trials in AD((CIIMP) CfMPfIU. Guideline on
20 Medicinal Products for the Treatment of Alzheimer's Disease and Other Dementias
(Document CPMP/EWP/553/95 Rev 1). London, UK: European Medicines Agency; 2009.),
the study included measures of cognition (modified ADAS-cog, primary outcome) and

functional abilities (23-item ADCS-ADL, secondary outcome) (Mohs RC, Knopman D, Petersen RC, Ferris SH, Ernesto C, Grundman M, et al. Development of cognitive instruments for use in clinical trials of antidementia drugs: additions to the Alzheimer's Disease Assessment Scale that broaden its scope. The Alzheimer's Disease Cooperative Study. Alzheimer Dis Assoc Disord. 1997;11 Suppl 2:S13-21). This ADCS-ADL scale is rated by the caregiver and ranges from 0 (need for extensive help) to 78 (independent performance). Baseline height and weight measurements were used to calculate BMI (kg/m^2), which was included as an exploratory outcome measure. Both ADCS-ADL and body weight were assessed at baseline, Week 6, 12 and 24 (height was measured at baseline only).

The study was conducted in accordance with the Declaration of Helsinki and the International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use / WHO Good Clinical Practice (ICH-GCP) guidelines, as appropriate to the legislation of the involved countries. The clinical trial registration number is ISRCTN72254645.

Modeling analysis

The primary analysis population was the intent-to-treat (ITT) efficacy population from the proof-of-concept study, defined as all patients who received at least one dose of study product and one post-baseline assessment. The per protocol (PP) efficacy population consisted of all patients who completed the study without major protocol deviations. Patients who did not receive any study product for > 25% of days, and/or patients who received < 70% of the prescribed dosage overall, were excluded from the PP group.

Data were analyzed using a repeated-measures mixed model (RMMM), in which time was treated as a categorical variable and represented by dummy codes. This model allows a comparison between the active group and the control group at each individual time point. Additional comparisons (active *versus* control group) were performed using contrast statements. For example, comparisons of the difference between baseline value and all post baseline values, or comparisons of the difference between baseline and Week 6 ADCS-ADL (combined) and Week 12 and Week 24 ADCS-ADL (combined). The Heterogeneous Compound Symmetry variance-covariance structure was used as it provided the best fit to the data.

Prior to the modeling analyses, the ADCS-ADL data were transformed using a quadratic transformation to adjust for the skewed nature of the data. Similarly, a square-root transformation was applied to BMI data.

5 Results

In the initial proof-of-concept study, 225 patients were randomized to treatment, of which 112 received Souvenaid and 113 received control product. Overall, 161 patients completed the 24-week study. Of the 212 patients in the ITT efficacy population, 199 were included in the PP analysis population at Week 12 and 161 at Week 24. No statistically significant differences between active or control groups were reported for any baseline characteristic (Table 8).

Table 8. Baseline characteristics for the ITT population.

Characteristics	Randomized treatment group	
	Control product	Active product
Total patient population	(n = 106)	(n = 106)
Men, n (%)	54 (51)	52 (49)
Age $\pm$ SD, yr	74.1 $\pm$ 7.2	73.3 $\pm$ 7.8
MMSE, mean $\pm$ SD	23.8 $\pm$ 2.7	24.0 $\pm$ 2.5
Total ADCS-ADL score	61.1 $\pm$ 10.5	61.9 $\pm$ 10.9
BMI	26.2 $\pm$ 4.8	26.2 $\pm$ 3.5
'low' baseline BMI group	(n = 60)	(n = 52)
Men, n (%)	31 (52)	28 (56)
Age $\pm$ SD, yr	74.9 $\pm$ 7.0	72.2 $\pm$ 8.4
MMSE, mean $\pm$ SD	23.9 $\pm$ 2.5	24.4 $\pm$ 2.4
Total ADCS-ADL, mean $\pm$ SD	60.7 $\pm$ 10.4	62.3 $\pm$ 11.5
BMI	23.2 $\pm$ 2.6	23.4 $\pm$ 1.7
'high' baseline BMI group	(n = 40)	(n = 51)
Men, n (%)	20 (50)	23 (45)
Age $\pm$ SD, yr	73.0 $\pm$ 7.5	74.3 $\pm$ 7.1
MMSE, mean $\pm$ SD	23.6 $\pm$ 2.5	23.5 $\pm$ 2.6
Total ADCS-ADL, mean $\pm$ SD	61.8 $\pm$ 11.2	61.9 $\pm$ 10.5

BMI	30.8 ± 3.5	29.0 ± 2.4
-----	------------	------------

Intention-to-treat efficacy population

ADCS-ADL = Alzheimer's Disease Cooperative Study – Activities of Daily Living (0–78; higher scores indicate greater functioning); MMSE = Mini-Mental State Examination (0–30; lower scores indicate greater cognitive dysfunction)

5

Treatment effect on BMI

In the entire study population, data suggested an increased BMI in the active study group *versus* the control group at Week 24. (ITT population: $p = 0.07$; PP population: $p = 0.03$), but not at Week 12 (ITT population: $p = 0.39$; PP population: $p = 0.42$, Table 9).

10

Additional analysis indicated that this treatment effect was primarily driven by changes observed in the 'low' BMI subgroup (for definition, see below), indicated by a trend towards an increase in BMI in the PP subgroup population (ITT population: $p = 0.23$; PP population: $p = 0.07$)

15

Table 9. Estimated marginal mean (95% confidence interval) BMI scores over 24 weeks for patients receiving active or control product (back transformed data; transformed [square-root] data were squared).

	Estimated Marginal Mean BMI (overall ITT population)			
	n	Control	n	Active
Baseline	103*	26.1 (25.3 – 26.9)	100	26.0 (25.2 – 26.8)
Week 6	97	26.2 (25.4 – 27.0)	96	26.2 (25.4 – 27.0)
Week 12	97	26.2 (25.4 – 27.0)	98	26.3 (25.5 – 27.1)
Week 24	76	26.2 (25.4 – 27.0)	76	26.4 (25.6 – 27.2)

BMI = Body mass index; ITT = Intention to treat

20

* BMI data were available for only part of the ITT population

Treatment effect on ADL

Overall, no treatment effect on ADCS-ADL was observed. However, baseline BMI was found to be a significant treatment effect modifier (ITT population: $p = 0.0393$) indicating that BMI significantly affected the ADCS-ADL response of the composition according to the invention.

5

In order to study whether this effect was predominantly driven by low or high BMI, patients were divided into two subgroups: those with a baseline BMI less than or equal to the mean (26.2 kg/m^2) and those above the mean. No statistically significant differences between active or control groups were reported for any baseline characteristic in any of the BMI subgroups (Table 8). The treatment effect of being in the 'high' or 'low' baseline subgroup was analysed by including a treatment*subgroup indicator interaction in the model.

Subgroup analysis of patients with 'high' or 'low' baseline BMI

15 There was no observed trend towards an intervention effect in the subgroup of patients with 'high' baseline BMI (ITT population: $p = 0.1675$; PP population: $p = 0.1701$). Within the 'low' baseline BMI subgroup, the RMMM model indicated a treatment effect at Week 12 (ITT population: $p = 0.0173$; PP population: $p = 0.0184$). However, this difference was not found to be significant at Week 24 (ITT population: $p = 0.2049$; PP population: $p =$
20 0.1042; Table 10).

Table 10. Estimated marginal mean (95% confidence interval) ADCS-ADL scores over 24 weeks for patients receiving active or control product with a 'low' (< mean) baseline BMI score at baseline (back transformed data; transformed [squared] data).

Estimated Marginal Mean ADCS-ADL in 'low' BMI subgroup (ITT population)				
	n	Control	n	Active
Baseline	52	63.4 (60.6 – 66.0)	60	61.7 (59.0 – 64.2)
Week 6	50	64.5 (61.8 – 67.1)	57	63.5 (60.9 – 66.0)
Week 12	49	62.8 (60.0 – 65.5)	56	64.0 (61.5 – 66.5)
Week 24	37	62.5 (59.5 – 65.3)	45	62.5 (59.8 – 65.2)

5 ADCS-ADL = Alzheimer's Disease Cooperative Study–Activities of Daily Living (0–78; higher scores indicate greater functioning); BMI = Body mass index; ITT = Intention to treat

10 Exploratory analyses using contrast statements were conducted to further investigate the effect of treatment on ADL response in patients with 'low' baseline BMI. When baseline ADCS-ADL was contrasted against all post-baseline assessments, no significant treatment effect was observed (ITT population: $p = 0.0759$, PP population: $p = 0.1289$). However, a significant treatment effect was observed when baseline and Week 6 ADSC-ADL (combined) were contrasted against Week 12 and Week 24 ADCS-ADL (combined)

15 (ITT population: $p = 0.0326$, PP population: $p = 0.0071$). These exploratory analyses suggest that the effect of the composition on ADL performance within the 'low' BMI subgroup is observable after at least 6 weeks of treatment.

20 Sensitivity analyses showed that the results were independent of transformation used and type of covariance structure chosen.

Discussion

This example study presents the positive effects of a multi-nutrient food product according to the invention on BMI. In addition, the modeling analyses presented here indicated that the composition significantly improved ADCS-ADL performance in a subgroup of mild AD patients with a 'low' baseline BMI.

This is the first study to indicate that improved (increased) BMI can be achieved with a specific combination of nutrients, independent of the energy content of the product.

- 10 The mean BMI for the 'low' baseline BMI subgroup was approximately 23 kg/m², which can be regarded as suboptimal (Chin APMJ, Dekker JM, Feskens EJ, Schouten EG, Kromhout D. How to select a frail elderly population? A comparison of three working definitions. J Clin Epidemiol. 1999 Nov;52(11):1015-21.), as previous research has shown a BMI < 23 kg/m² to be associated with poorer 7-year survival in patients with
- 15 AD(Faxen-Irving G, Basun H, Cederholm T. Nutritional and cognitive relationships and long-term mortality in patients with various dementia disorders. Age Ageing. 2005 Mar;34(2):136-41; Irving GF, Freund-Levi Y, Eriksdotter-Jonhagen M, Basun H, Brismar K, Hjorth E, et al. Omega-3 fatty acid supplementation effects on weight and appetite in patients with Alzheimer's disease: the omega-3 Alzheimer's disease study. J
- 20 Am Geriatr Soc. 2009 Jan;57(1):11-7.)

Example 10

Tablet comprising 500 mg uridinemonophosphate, uridine or citicoline (cytidine diphosphate-choline), combined with 300 microgram sodium folate monoglutamate and 5

25 microgram vitamin B12 and conventional tableting aids, like binders, inert filling aids, colorants, etc., to provide a tablet of about 1 g.

The tablet is in particular suitable for increasing BMI in an elderly person. The skilled person will be able to provide a different dosage form providing the same or similar

30 active ingredients in the same or other suitable dosage, based on the information disclosed herein and common general knowledge.

Example 11

Tablet comprising 400 mg of a nucleoside equivalent, like uridine, 6 mg vitamin B6, 200 microgram folate, 5 microgram vitaminB12 and 400 mg of a choline equivalent included in conventional tableting aids to provide a tablet of about 1 g.

5

The tablet is in particular suitable for increasing BMI in an elderly person. The skilled person will be able to provide a different dosage form providing the same or similar active ingredients in the same or other suitable dosage, based on the information disclosed herein and common general knowledge.

10

Example 12

Capsule weighing about 1.9 g, comprising 700 mg marine oil, 300 mg soy lecithin, 100 mg of uridine, 100 mg of UMP, 100 micogram of folate, 5 mg vitamin B6, 20 microgram vitamin B12, 10 mg vitamin E (as alpha-tocopherol) and 20 mg vitamin C, the remainder being capsulating material, *e.g.* gelatin,

15

The capsule is in particular suitable for increasing BMI in an elderly person. The skilled person will be able to provide a different dosage form providing the same or similar active ingredients in the same or other suitable dosage, based on the information disclosed herein and common general knowledge.

20

25

CLAIMS

1. Composition comprising at least two components selected from the group of (i) a nucleoside equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, (iv) a phospholipid, (v) an antioxidant and (vi) a choline, with the proviso that at least component (i) or (iii) is present, for use in the prevention or treatment of frailty in a mammal.
2. Composition comprising at least three components selected from the group of (i) a nucleoside equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, (iv) a phospholipid, (v) an antioxidant and (vi) a choline, with the proviso that at least component (i) or (iii) is present, for use in the prevention or treatment of frailty in a mammal.
3. Composition comprising at least four components selected from the group of (i) a nucleoside equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, (iv) a phospholipid, (v) an antioxidant and (vi) a choline, with the proviso that at least component (i) or (iii) is present, for use in the prevention or treatment of frailty in a mammal.
4. Composition according to any one of claims 1 to 3, wherein frailty is determined by compliance with at least 2 criteria, preferably 3 criteria, selected from the group of muscle weakness, excessive feelings of exhaustion or fatigue, abnormally low physical activity, slow or unsteady gait, weight loss, and neurological dysfunction.
5. Composition according to any one of claims 1 to 4, for use in increasing one or more of body weight, body mass index, lean body weight, muscle mass, muscle strength or muscle function, especially in an elderly human.
6. Composition according to any one of claims 1 to 4, for use in increasing one or more of body weight, body mass index, lean body weight, muscle mass, muscle strength or muscle function and for improving activities of daily living.
7. Composition according to any one of claims 1 to 6, wherein the composition is to be administered to a human prone to or suffering from cognitive decline or dementia, in particular senile dementia, more in particular Alzheimer's disease.
8. Composition according to any one of the preceding claims, wherein the mammal is

suffering from a prodromal form of dementia.

9. Composition according to any one of the preceding claims, wherein the mammal has a BMI below 26 and preferably below 23.5 kg/m².
10. Composition according to any one of the preceding claims, wherein the composition is a nutritional or a pharmaceutical composition.
11. Composition according to any one of the preceding claims, wherein the composition comprises UMP.
12. Composition according to any one of the preceding claims, wherein the composition comprises DHA and EPA.
- 10 13. Composition according to any one of the preceding claims, wherein the composition comprises vitamin B6, vitamin B12 and vitamin B9.
14. Composition according to any one of the preceding claims, wherein the composition comprises at least two different phospholipids selected from the group consisting of phosphatidylserine, phosphatidylinositol, phosphatidylcholine and phosphatidylethanolamine.
- 15 15. Composition according to any of the preceding claims, wherein the composition comprises vitamin C, vitamin E and selenium.
16. Composition according to any of the preceding claims, wherein the composition comprises choline chloride.
- 20 17. Composition according to any of the preceding claims, wherein the composition comprises (i) UMP, (ii) DHA and EPA, (iii) vitamin B6, vitamin B12 and vitamin B9, (iv) at least two different phospholipids selected from the group consisting of phosphatidylserine, phosphatidylinositol, phosphatidylcholine and phosphatidylethanolamine, (v) vitamin C, vitamin E and selenium and (vi) choline chloride.
- 25 18. Composition comprising at least (i) a uridine equivalent, (ii) an ω -3 polyunsaturated fatty acid selected from the group of DHA, DPA and EPA, (iii) a vitamin B, (iv) a phospholipid, (v) an antioxidant, preferably selected from the group consisting of vitamin C, vitamin E and selenium, and (vi) a choline, for use in the prevention or treatment of pathological frailty in a mammal, wherein the mammal is a patient suffering from Alzheimer's disease or dementia, the patient
- 30


having an BMI below 26 and more preferably below 23.5.

- 5 **19.** Nutritional composition comprising a composition according to any of the preceding claims, wherein the nutritional composition comprises at least a whey protein, most preferably a whey protein enriched in serum albumin and optionally a vegetable protein.

- 10 **20.** Nutritional composition according to claim 19, wherein the nutritional composition has a nutritional value of at least at least 50 kcal per 100 ml, in particular 100 to 300 kcal per 100 ml of composition.

1/2

Figure 1: Uses of the composition according to the invention in frail and pre-frail



2/2

Figure 2 : Composition according to the invention effective on ADL (ADCS score) in elderly of BMI < 26 and not in elderly having BMI > 26.

