



US 20090203142A1

(19) **United States**

(12) **Patent Application Publication**
Booth et al.

(10) **Pub. No.: US 2009/0203142 A1**

(43) **Pub. Date: Aug. 13, 2009**

(54) **NUCLEOTIDE SEQUENCES OF A NEW CLASS OF DIVERGED DELTA-9 STEAROYL-ACP DESATURASE GENES**

(75) Inventors: **John R. Booth**, Boothwyn, PA (US); **Rebecca E. Cahoon**, Lincoln, NE (US); **William D. Hitz**, Wilmington, DE (US); **Anthony J. Kinney**, Wilmington, DE (US); **Narendra S. Yadav**, Wilmington, DE (US)

Correspondence Address:
E I DU PONT DE NEMOURS AND COMPANY
LEGAL PATENT RECORDS CENTER
BARLEY MILL PLAZA 25/1122B, 4417 LANCASTER PIKE
WILMINGTON, DE 19805 (US)

(73) Assignee: **E.I. DU PONT DE NEMOURS AND COMPANY**

(21) Appl. No.: **12/357,474**

(22) Filed: **Jan. 22, 2009**

Related U.S. Application Data

(60) Division of application No. 10/981,293, filed on Nov. 4, 2004, now Pat. No. 7,498,427, which is a continuation of application No. 09/934,900, filed on Aug. 22, 2001, now abandoned.

(60) Provisional application No. 60/226,996, filed on Aug. 22, 2000.

Publication Classification

(51) **Int. Cl.**
C12N 15/82 (2006.01)
C07H 21/04 (2006.01)
C12N 15/63 (2006.01)
C40B 30/04 (2006.01)

(52) **U.S. Cl.** **435/468**; 536/23.2; 435/320.1; 506/9

(57) **ABSTRACT**

An isolated nucleic acid fragment encoding a diverged delta-9 fatty acid desaturase is disclosed. Also the construction of a chimeric gene encoding all or a portion of the diverged delta-9 fatty acid desaturase is disclosed, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the diverged delta-9 fatty acid desaturase in a transformed host cell.

344

345

SID2 (soy) PVLARLCGTIAADEKCRHENAYSRIVEKLLLEVPDTGAMVAIGNMMEKKTITMPAHLMYDGGDDPRLFEHYSAVAQRIQVYVTANDYADILEFLV
SID10 (corn) HVLARVCGAIMADEKCRHETAYTRIVAKLFEVDPDAAVRALGYMMRHRITMPAALMTDGRDAHLYAHYAAAAQQTGVYFASDYRSILEHLI
SID12 (corn) DVLARACGTIAADEKCRHETAYGRIVEQLLQLDPEGAVLAVADMRRKRI TMPAHLMHGDRMDLFEHFHFAVAQRLGVYVTARDYADIVEFLV
SID14 (rice) AALARACGTVAADEKCRHEAAAYTRIVSRLLLEADPDAGVRAVARMRLREGVAMPTSPISDGRRDDLYACVVSLAEQAGIYTVSDYCSIVEHLV
SID16 (rice) DVLARTCGTIAADEKCRHETAYGRIVEQLLRLDPPDGAMLAIDMMHKRI TMPAHLMHGDRMNLFDHFAVAQRLNVYVTARDYADIVEFLV
gi4704824 PVLARICGTIAADEKCRHENAYSRIVEKLLLELDPDTGAMVAIGDMMQKKTITMPAHLMYDGEDPKLFDHFSAVAQRMVYVTANDYADILEFLI
gi267036 FQLAQVCGIIAADEKCRHETAYTKIVEKLFEDIIDPGAVLALADMRRKVSMPAHLMYDGKDDNLFENYSAVAQQIGVYTAADYADILEHLV
gi6957724 TTLAKICGTIAADEKCRHETAYTRIVEKLFEDIIDPGTVQALASMMRKRITMPAHLMHGDRDDDLFDHYAAVAQRIQVYVTATDYAGILEFLI
gi3355632 MKLAQICGLIAADEKCRHETAYTKIVEKLFEDIIDPGTIVLALADMRRKKTISMPAHLMYDGEDDNLFDNYSVAQRIQVYTAADYADILEFLV
SID23 (soy) IKLAQICGMIASDEKRIUETAYTKIVEKLFVEVDPDGVMAFADMRRKKTAMPAILMYDGRDDNLFDNYSAVAQRIQVYTAADYADILEFLV

345

405

SID2 (soy) ERWRLEKLE-GLMAEGKRAQDFVCGLAPRIRRLQERADERARKMKKHHG-VKFSWIFNKE---LLL (405 aa)
SID10 (corn) RQWRVEELAAGLSGEGRRARDYVCGLPKHIRRMEKKAHDRAAQTKKPTSVPFWSWIFDRSVNVVIP (424 aa)
SID12 (corn) KRWKLETLESGLSGEGRRARDYVCGLAPRMRRAEAEADRACK-DEPRM-VKFSWIFDRE---AVV (380 aa)
SID14 (rice) REWRVEELAAGLSGEGRRARDYVCELQKIRRMKEKAHERAVKAQKPIIPINWIFDRHVSVMLP (219 aa)
SID16 (rice) KRWKLETLETGLSGEGRRARDYVCGLAKRMRRAEAEADRACK-DEQRK-VKFSWIYDRE---VIV (381 aa)
gi4704824 GRWRLEKVQ-DLKDEGKKAQDFVCGLAPRIRRLQERADERARKMK-PHA-VKFSWIFNKE---IIL (384 aa)
gi267036 NRWKVENL-MGLSGEGHKAQDFVCGLAPRIRKLGERAQSLK?VS-L---VPPFSWIFNKE---LKV (398 aa)
gi6957724 RRWEVEKLGMLSGEGRRARDYVCGLAPRIRRLQERANDRVKLSKSPSVSFSWIYGRE---VEL (411 aa)
gi3355632 GRWKVDAF-TGLSGEGNKAQDFVCGLAPRIRKLEERAAGRAKQTSKS---VPPFSWIFSRE---LVL (396 aa)
SID23 (soy) GRWKVEQL-TGLSGEGRKAQDFVCGLPPRIRRLLEERAQARGKESST---LKFSWIHDRE---VLL (391 aa)

FIG. 1B

**NUCLEOTIDE SEQUENCES OF A NEW
CLASS OF DIVERGED DELTA-9
STEAROYL-ACP DESATURASE GENES**

[0001] This application claims the benefit of U.S. Provisional Application No. 60/226,996, filed Aug. 22, 2000, the entire contents of which are herein incorporated by reference.

FIELD OF THE INVENTION

[0002] This invention relates to the field of plant molecular biology and, in particular, to nucleic acid fragments encoding a diverged delta-9 fatty acid desaturase in plants and seeds.

BACKGROUND OF THE INVENTION

[0003] Soybean oil accounts for about 70% of the 14 billion pounds of edible oil consumed in the United States and is a major edible oil worldwide. It is used in baking, frying, salad dressing, margarine, and a multitude of processed foods. In 1987/88 60 million acres of soybean were planted in the U.S. Soybean is the lowest-cost producer of vegetable oil, which is a by-product of soybean meal. Soybean is agronomically well-adapted to many parts of the U.S. Machinery and facilities for harvesting, storing, and crushing are widely available across the U.S. Soybean products are also a major element of foreign trade since 30 million metric tons of soybeans, 25 million metric tons of soybean meal, and 1 billion pounds of soybean oil were exported in 1987/88. Nevertheless, increased foreign competition has led to recent declines in soybean acreage and production. The low cost and ready availability of soybean oil provides an excellent opportunity to upgrade this commodity oil into higher value specialty oils to both add value to soybean crop for the U.S. farmer and enhance U.S. trade.

[0004] Soybean oil derived from commercial varieties is composed primarily of 11% palmitic (16:0), 4% stearic (18:0), 24% oleic (18:1), 54% linoleic (18:2) and 7% linolenic (18:3) acids. Palmitic and stearic acids are, respectively, 16- and 18-carbon-long saturated fatty acids. Oleic, linoleic and linolenic are 18-carbon-long unsaturated fatty acids containing one, two and three double bonds, respectively. Oleic acid is also referred to as a monounsaturated fatty acid, while linoleic and linolenic acids are also referred to as polyunsaturated fatty acids. The specific performance and health attributes of edible oils is determined largely by their fatty acid composition.

[0005] Soybean oil is high in saturated fatty acids when compared to other sources of vegetable oil and contains a low proportion of oleic acid, relative to the total fatty acid content of the soybean seed. These characteristics do not meet important health needs as defined by the American Heart Association.

[0006] More recent research efforts have examined the role that monounsaturated fatty acid plays in reducing the risk of coronary heart disease. In the past, it was believed that monounsaturates, in contrast to saturates and polyunsaturates, had no effect on serum cholesterol and coronary heart disease risk. Several recent human clinical studies suggest that diets high in monounsaturated fat may reduce the "bad" (low-density lipoprotein) cholesterol while maintaining the "good" (high-density lipoprotein) cholesterol. [See Mattson et al. (1985) *Journal of Lipid Research* 26:194-202, Grundy

(1986) *New England Journal of Medicine* 314:745-748, and Mensink et al. (1987) *The Lancet* 1:122-125, all collectively herein incorporated by reference.] These results corroborate previous epidemiological studies of people living in Mediterranean countries where a relatively high intake of monounsaturated fat and low consumption of saturated fat correspond with low coronary heart disease mortality. [Keys, A., *Seven Countries: A Multivariate Analysis of Death and Coronary Heart Disease*, Cambridge: Harvard University Press, 1980, herein incorporated by reference.] The significance of monounsaturated fat in the diet was further confirmed by international researchers from seven countries at the Second Colloquium on Monounsaturated Fats held Feb. 26, 1987, in Bethesda, Md., and sponsored by the National Heart, Lung and Blood Institutes [Report, *Monounsaturates Use Said to Lower Several Major Risk Factors*, Food Chemical News, Mar. 2, 1987, p. 44, herein incorporated by reference].

[0007] Soybean oil is also relatively high in polyunsaturated fatty acids—at levels in far excess of our essential dietary requirement. These fatty acids oxidize readily to give off-flavors and result in reduced performance associated with unprocessed soybean oil. The stability and flavor of soybean oil is improved by hydrogenation, which chemically reduces the double bonds. However, the need for this processing reduces the economic attractiveness of soybean oil.

[0008] A soybean oil low in total saturates and polyunsaturates and high in monounsaturate would provide significant health benefits to the United States population, as well as, economic benefit to oil processors. Soybean varieties which produce seeds containing the improved oil will also produce valuable meal as animal feed.

[0009] Another type of differentiated soybean oil is an edible fat for confectionary uses. More than 2 billion pounds of cocoa butter, the most expensive edible oil, are produced worldwide. The U.S. imports several hundred million dollars worth of cocoa butter annually. The high and volatile prices and uncertain supply of cocoa butter have encouraged the development of cocoa butter substitutes. The fatty acid composition of cocoa butter is 26% palmitic, 34% stearic, 35% oleic and 3% linoleic acids. About 72% of cocoa butter's triglycerides have the structure in which saturated fatty acids occupy positions 1 and 3 and oleic acid occupies position 2. Cocoa butter's unique fatty acid composition and distribution on the triglyceride molecule confer on it properties eminently suitable for confectionary end-uses: it is brittle below 27° C. and depending on its crystalline state, melts sharply at 25-30° C. or 35-36° C. Consequently, it is hard and non-greasy at ordinary temperatures and melts very sharply in the mouth. It is also extremely resistant to rancidity. For these reasons, producing soybean oil with increased levels of stearic acid, especially in soybean lines containing higher-than-normal levels of palmitic acid, and reduced levels of unsaturated fatty acids is expected to produce a cocoa butter substitute in soybean. This will add value to oil and food processors as well as reduce the foreign import of certain tropical oils.

[0010] Only recently have serious efforts been made to improve the quality of soybean oil through plant breeding, especially mutagenesis, and a wide range of fatty acid composition has been discovered in experimental lines of soybean (Table 1). These findings (as well as those with other oilcrops) suggest that the fatty acid composition of soybean oil can be significantly modified without affecting the agronomic performance of a soybean plant. However, there is no soybean

mutant line with levels of saturates less than those present in commercial canola, the major competitor to soybean oil as a “healthy” oil.

TABLE 1

Range of Fatty Acid Percentages Produced by Soybean Mutants	
Fatty Acids	Range of Percentages
Palmitic Acid	6-28
Stearic Acid	3-30
Oleic Acid	17-50
Linoleic Acid	35-60
Linolenic Acid	3-12

[0011] There are serious limitations to using mutagenesis to alter fatty acid composition. It is unlikely to discover mutations a) that result in a dominant (“gain-of-function”) phenotype, b) in genes that are essential for plant growth, and c) in an enzyme that is not rate-limiting and that is encoded by more than one gene. Even when some of the desired mutations are available in soybean mutant lines their introgression into elite lines by traditional breeding techniques will be slow and expensive, since the desired oil compositions in soybean are most likely to involve several recessive genes.

[0012] Recent molecular and cellular biology techniques offer the potential for overcoming some of the limitations of the mutagenesis approach, including the need for extensive breeding. Particularly useful technologies are: a) seed-specific expression of foreign genes in transgenic plants [see Goldberg et al. (1989) *Cell* 56:149-160], b) use of antisense RNA to inhibit plant target genes in a dominant and tissue-specific manner [see van der Krol et al. (1988) *Gene* 72:45-50], c) transfer of foreign genes into elite commercial varieties of commercial oilcrops, such as soybean [Chee et al. (1989) *Plant Physiol.* 91:1212-1218; Christou et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:7500-7504; Hinchey et al. (1988) *Bio/Technology* 6:915-922; EPO publication 0 301 749 A2], rapeseed [De Block et al. (1989) *Plant Physiol.* 91:694-701], and sunflower [Everett et al. (1987) *Bio/Technology* 5:1201-1204], and d) use of genes as restriction fragment length polymorphism (RFLP) markers in a breeding program, which makes introgression of recessive traits into elite lines rapid and less expensive [Tanksley et al. (1989) *Bio/Technology* 7:257-264]. However, application of each of these technologies requires identification and isolation of commercially-important genes.

[0013] Oil biosynthesis in plants has been fairly well-studied [see Harwood (1989) in *Critical Reviews in Plant Sciences*, Vol. 8(1):1-43]. The biosynthesis of palmitic, stearic and oleic acids occur in the plastids by the interplay of three key enzymes of the “ACP track”: palmitoyl-ACP elongase, stearyl-ACP desaturase and acyl-ACP thioesterase. Stearyl-ACP desaturase introduces the first double bond on stearyl-ACP to form oleoyl-ACP. It is pivotal in determining the degree of unsaturation in vegetable oils. Because of its key position in fatty acid biosynthesis it is expected to be an important regulatory step. While the enzyme’s natural substrate is stearyl-ACP, it has been shown that it can, like its counterpart in yeast and mammalian cells, desaturate stearyl-CoA, albeit poorly [McKeon et al. (1982) *J. Biol. Chem.* 257:12141-12147]. The fatty acids synthesized in the plastid are exported as acyl-CoA to the cytoplasm. At least three different glycerol acylating enzymes (glycerol-3-P

acyltransferase, 1-acyl-glycerol-3-P acyltransferase and diacylglycerol acyltransferase) incorporate the acyl moieties from the cytoplasm into triglycerides during oil biosynthesis. These acyltransferases show a strong, but not absolute, preference for incorporating saturated fatty acids at positions 1 and 3 and monounsaturated fatty acid at position 2 of the triglyceride. Thus, altering the fatty acid composition of the acyl pool will drive by mass action a corresponding change in the fatty acid composition of the oil. Furthermore, there is experimental evidence that, because of this specificity, given the correct composition of fatty acids, plants can produce cocoa butter substitutes [Bafar et al. (1990) *JAOCs* 67:217-225].

[0014] Based on the above discussion, one approach to altering the levels of stearic and oleic acids in vegetable oils is by altering their levels in the cytoplasmic acyl-CoA pool used for oil biosynthesis. There are two ways of doing this genetically: a) altering the biosynthesis of stearic and oleic acids in the plastid by modulating the levels of stearyl-ACP desaturase in seeds through either overexpression or antisense inhibition of its gene, and b) converting stearyl-CoA to oleoyl-CoA in the cytoplasm through the expression of the stearyl-ACP desaturase in the cytoplasm.

[0015] In order to use antisense inhibition of stearyl-ACP desaturase in the seed, it is essential to isolate the gene(s) or cDNA(s) encoding the target enzyme(s) in the seed, since antisense inhibition requires a high-degree of complementarity between the antisense RNA and the target gene that is expected to be absent in stearyl-ACP desaturase genes from other species.

[0016] The purification and nucleotide sequences of mammalian microsomal stearyl-CoA desaturases have been published [Thiede et al. (1986) *J. Biol. Chem.* 262:13230-13235; Ntambi et al. (1988) *J. Biol. Chem.* 263:17291-17300; Kaestner et al. (1989) *J. Biol. Chem.* 264:14755-14761]. However, the plant enzyme differs from them in being soluble, in utilizing a different electron donor, and in its substrate-specificities. The purification and the nucleotide sequences for animal enzymes do not teach how to purify the plant enzyme or isolate a plant gene. The purification of stearyl-ACP desaturase was reported from safflower seeds [McKeon et al. (1982) *J. Biol. Chem.* 257:12141-12147]. However, this purification scheme was not useful for soybean, either because the desaturases are different or because of the presence of other proteins such as the soybean seed storage proteins in seed extracts.

[0017] The rat liver stearyl-CoA desaturase protein has been expressed in *E. coli* [Strittmatter et al. (1988) *J. Biol. Chem.* 263:2532-2535] but, as mentioned above, its substrate specificity and electron donors are quite distinct from that of the plant.

[0018] Plant stearyl-ACP desaturase cDNAs have been cloned from soybean [U.S. Pat. No. 5,760,206, the disclosure of which is hereby incorporated by reference], safflower [Thompson et al. (1991) *Proc. Natl. Acad. Sci.* 88:2578], castor [Shanklin and Somerville (1991) *Proc. Natl. Acad. Sci.* 88:2510-2514], and cucumber [Shanklin et al. (1991) *Plant Physiol.* 97:467-468]. Kutzon et al. [(1992) *Proc. Natl. Acad. Sci.* 89:2624-2648] have reported that rapeseed stearyl-ACP desaturase when expressed in *Brassica rapa* and *B. napus* in an antisense orientation can result in increase in 18:0 level in transgenic seeds. All of the reported genes have 59-80% identity to each other at the nucleotide and polypeptide level.

[0019] U.S. Pat. No. 5,723,595, issued to Thompson et al. on Mar. 3, 1998, describes stearyl-ACP desaturases from castor and safflower.

[0020] U.S. Pat. No. 5,443,974, issued to Hitz et al., on Aug. 22, 1995, describes soybean stearyl-ACP desaturase.

[0021] U.S. Pat. No. 5,760,206, issued to Hitz et al, on Jun. 2, 1998, describes soybean stearyl-ACP desaturase.

SUMMARY OF THE INVENTION

[0022] The present invention concerns an isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding a polypeptide having delta-9 fatty acid desaturase activity that has at least 80%, 85%, 90%, or 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, or 16, or (b) the complement of the nucleotide sequence.

[0023] In a second embodiment, it is preferred that the isolated polynucleotide of the claimed invention comprises a nucleotide sequence which comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, or that codes for a polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, or 16.

[0024] In a third embodiment, this invention relates to a chimeric gene comprising an isolated polynucleotide of the present invention operably linked to at least one suitable regulatory sequence.

[0025] In a fourth embodiment, the present invention concerns an isolated host cell comprising a chimeric construct of the present invention or an isolated polynucleotide of the present invention. The host cell may be eukaryotic, such as a yeast or a plant cell, or prokaryotic, such as a bacterial cell.

[0026] In a fifth embodiment, the invention also relates to a process for producing an isolated host cell comprising a chimeric construct of the present invention or an isolated polynucleotide of the present invention, the process comprising either transforming or transfecting an isolated compatible host cell with a chimeric gene or isolated polynucleotide of the present invention.

[0027] In a sixth embodiment, the invention concerns a diverged delta-9 stearyl desaturase polypeptide of at least 400 amino acids comprising at least 80% identity based on the Clustal method of alignment compared to a polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, or 16.

[0028] In a seventh embodiment, the invention relates to a method of selecting an isolated polynucleotide that affects the level of expression of a diverged delta-9 stearyl desaturase polypeptide or enzyme activity in a host cell, preferably a plant cell, the method comprising the steps of: (a) constructing an isolated polynucleotide of the present invention or an isolated chimeric construct of the present invention; (b) introducing the isolated polynucleotide or the isolated chimeric construct into a host cell; (c) measuring the level of the diverged delta-9 stearyl desaturase polypeptide or enzyme activity in the host cell containing the isolated polynucleotide; and (d) comparing the level of the diverged delta-9 stearyl desaturase polypeptide or enzyme activity in the host cell containing the isolated polynucleotide with the level of the diverged delta-9 stearyl desaturase polypeptide or enzyme activity in the host cell that does not contain the isolated polynucleotide.

[0029] In an eighth embodiment, the invention concerns a method of obtaining a nucleic acid fragment encoding a substantial portion of a diverged delta-9 stearyl desaturase polypeptide, preferably a plant diverged delta-9 stearyl desaturase polypeptide, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 30 (preferably at least one of 40, most preferably at least one of 60) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, or 15, and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a substantial portion of a diverged delta-9 stearyl desaturase amino acid sequence.

[0030] In a ninth embodiment, this invention relates to a method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a diverged delta-9 stearyl desaturase polypeptide comprising the steps of: probing a cDNA or genomic library with an isolated polynucleotide of the present invention; identifying a DNA clone that hybridizes with an isolated polynucleotide of the present invention; isolating the identified DNA clone; and sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

[0031] In a tenth embodiment, this invention relates to a method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a diverged delta-9 stearyl desaturase polypeptide comprising the steps of: probing a cDNA or genomic library with an isolated polynucleotide of the present invention; identifying a DNA clone that hybridizes with an isolated polynucleotide of the present invention; isolating the identified DNA clone; introducing said clone into a construct for expression in a bacteria or yeast; and assaying for delta-9 desaturase activity in the bacteria or yeast.

[0032] In an eleventh embodiment, this invention relates to a method of identifying an isolated polynucleotide that encodes a delta-9 fatty acid desaturase comprising the steps of: determining an amino acid sequence of the polypeptide encoded by the isolated DNA; determining if the amino acid sequence comprises at least two amino acid sequences selected from the group consisting of HSMPPPEK corresponding to amino acids 67-73 of SEQ ID NO:2, LPLLKPVE corresponding to amino acids 89-96 of SEQ ID NO:2, EYFV-VLVGDM corresponding to amino acids 132-141 of SEQ ID NO:2, EKTIV corresponding to amino acids 205-208 of SEQ ID NO:2, GMDPGT corresponding to amino acids 215-220 of SEQ ID NO:2, NNPYLGFVYTSFQERAT corresponding to amino acids 222-238 of SEQ ID NO:2, VLAR corresponding to amino acids 256-259 of SEQ ID NO:2, RIVE corresponding to amino acids 277-280 of SEQ ID NO:2, ITMPAHL corresponding to amino acids 302-308 of SEQ ID NO:2, or DFVCGLA corresponding to amino acids 364-370 of SEQ ID NO:2.

[0033] In a twelfth embodiment, this invention relates to a method of identifying an isolated polynucleotide that encodes a delta-9 fatty acid desaturase comprising the steps of: determining the polypeptide sequence by one of the aforementioned methods; determining that the amino acid sequence of the polypeptide does not contain at least one of the following amino acid sequences KEIPDDYFWLVGDMITEEALP-

TYQTMLNT corresponding to positions 116-145 of SEQ ID NO:23; or DYADILEFLVGRWK corresponding to positions 324-337 of SEQ ID NO:23.

[0034] In an thirteenth embodiment, this invention relates to a method of altering the level of expression of a diverged delta-9 fatty acid desaturase in a host cell comprising: (a) transforming a host cell with a chimeric construct of the present invention; and (b) growing the transformed host cell under conditions that are suitable for expression of the chimeric construct wherein expression of the chimeric construct results in production of altered levels of the a diverged delta-9 fatty acid desaturase in the transformed host cell.

BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCE LISTINGS

[0035] The invention can be more fully understood from the following detailed description and the accompanying drawings and Sequence Listing which form a part of this application.

[0036] FIG. 1 shows a comparison of the amino acid stearoyl-ACP desaturase sequences of the soybean enzyme [SEQ ID NO:2], corn [SEQ ID NOs:10 and 12], rice [SEQ ID NOs:14 and 16], to the lupine [gi 4704824, SEQ ID NO:17], jojoba [gi 267036, SEQ ID NO:20], *Arabidopsis* [gi 6957724, SEQ ID NO:21], flax [gi 3355632, SEQ ID NO:22], and to the soybean stearoyl-ACP desaturase [SEQ ID NO:23] found in U.S. Pat. No. 5,760,206.

[0037] Table 2 lists the polypeptides that are described herein, the designation of the cDNA clones that comprise the nucleic acid fragments encoding polypeptides representing all or a substantial portion of these polypeptides, and the corresponding identifier (SEQ ID NO:) as used in the attached Sequence Listing. The sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

TABLE 2

Diverged Delta-9 Fatty Acid Desaturase			
Protein	Clone Designation	SEQ ID NO:	
		(Nucleotide)	(Amino Acid)
Soybean [<i>Glycine max</i>]	se6.pk0026.a8	1	2
Corn [<i>Zea mays</i>]	cbn10.pk0061.a3	3	4
Corn [<i>Zea mays</i>]	contig of: cen7f.pk001.k12 cpd1c.pk012.n9 cpd1c.pk014.l18 p0103.ciaad81r p0106.cjlp88r	5	6
Rice [<i>Oryza sativa</i>]	rds1c.pk007.g19	7	8
Corn [<i>Zea mays</i>]	cbn10.pk0061.a3:fis	9	10
Corn [<i>Zea mays</i>]	cpd1c.pk014.l18:fis	11	12
Rice [<i>Oryza sativa</i>]	rds1c.pk007.g19:fis	13	14
Rice [<i>Oryza sativa</i>]	rsl1n.pk008.j18:fis	15	16

[0038] The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Res.* 13:3021-3030 (1985) and in the *Biochemical J.* 219 (No. 2):345-373 (1984) which are herein incorporated by reference. The sym-

bols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

DETAILED DESCRIPTION OF THE INVENTION

[0039] A new diverged class of delta-9 steroyl desaturases are disclosed herein. These desaturases were obtained from soybean, corn, and rice and are less than 60% identical to the previously characterized class. This new diverged class of delta-9 steroyl desaturases still performs the substantially identical biochemical function in plants as the previously characterized class. In addition, evidence is presented to show that the new class of desaturases may play a more important role in regulating fatty acid synthesis than the previous class.

[0040] The terms “diverged delta-9 fatty acid desaturase”, “diverged delta-9 steroyl desaturase”, or “diverged delta-9 desaturase” are used interchangeably herein and include, but are not limited to, all plant delta-9 steroyl desaturases that are less than 60% identical to the previously characterized delta-9 steroyl desaturases (PCT Publication Nos. WO 91/13972 and WO 91/18985). This new diverged class of delta-9 steroyl desaturases still performs the substantially identical biochemical function in plants as the previously characterized class, namely the introduction of a double bond between carbon atoms 9 and 10 of stearoyl-ACP to form oleoyl-ACP.

[0041] In the context of this disclosure, a number of terms shall be utilized. The terms “polynucleotide”, “polynucleotide sequence”, “nucleic acid sequence”, and “nucleic acid fragment”/“isolated nucleic acid fragment” are used interchangeably herein. These terms encompass nucleotide sequences and the like. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally contains synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA, synthetic DNA, or mixtures thereof. An isolated polynucleotide of the present invention may include at least one of 30 contiguous nucleotides, preferably at least one of 40 contiguous nucleotides, most preferably one of at least 60 contiguous nucleotides derived from SEQ ID NO:1, 3, 5, 7, 9, 11, 13, or 15, or the complement of such sequences.

[0042] The term “isolated” polynucleotide refers to a polynucleotide that is substantially free from other nucleic acid sequences, such as and not limited to other chromosomal and extrachromosomal DNA and RNA. Isolated polynucleotides may be purified from a host cell in which they naturally occur. Conventional nucleic acid purification methods known to skilled artisans may be used to obtain isolated polynucleotides. The term also embraces recombinant polynucleotides and chemically synthesized polynucleotides. Nucleotides (usually found in their 5'-monophosphate form) are referred to by their single letter designation as follows: “A” for adenylate or deoxyadenylate (for RNA or DNA, respectively), “C” for cytidylate or deoxycytidylate, “G” for guanylate or deoxyguanylate, “U” for uridylate, “T” for deoxythymidylate, “R” for purines (A or G), “Y” for pyrimidines (C or T), “K” for G or T, “H” for A or C or T, “I” for inosine, and “N” for any nucleotide.

[0043] The term “host” refers to any organism, or cell thereof, whether human or non-human into which a recombinant construct can be stably or transiently introduced in order to alter gene expression in the host.

[0044] The term “recombinant” means, for example, that a nucleic acid sequence is made by an artificial combination of

two otherwise separated segments of sequence, e.g., by chemical synthesis or by the manipulation of isolated nucleic acids by genetic engineering techniques.

[0045] As used herein, "contig" refers to a nucleotide sequence that is assembled from two or more constituent nucleotide sequences that share common or overlapping regions of sequence homology. For example, the nucleotide sequences of two or more nucleic acid fragments can be compared and aligned in order to identify common or overlapping sequences. Where common or overlapping sequences exist between two or more nucleic acid fragments, the sequences (and thus their corresponding nucleic acid fragments) can be assembled into a single contiguous nucleotide sequence.

[0046] As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the polypeptide encoded by the nucleotide sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by gene silencing through for example antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate gene silencing or alteration of the functional properties of the resulting protein molecule. It is therefore understood that the invention encompasses more than the specific exemplary nucleotide or amino acid sequences and includes functional equivalents thereof. The terms "substantially similar" and "corresponding substantially" are used interchangeably herein.

[0047] Substantially similar nucleic acid fragments may be selected by screening nucleic acid fragments representing subfragments or modifications of the nucleic acid fragments of the instant invention, wherein one or more nucleotides are substituted, deleted and/or inserted, for their ability to affect the level of the polypeptide encoded by the unmodified nucleic acid fragment in a plant or plant cell. For example, a substantially similar nucleic acid fragment representing at least one of 30 contiguous nucleotides derived from the instant nucleic acid fragment can be constructed and introduced into a plant or plant cell. The level of the polypeptide encoded by the unmodified nucleic acid fragment present in a plant or plant cell exposed to the substantially similar nucleic acid fragment can then be compared to the level of the polypeptide in a plant or plant cell that is not exposed to the substantially similar nucleic acid fragment.

[0048] For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by using nucleic acid fragments that do not share 100% sequence identity with the gene to be suppressed. In a preferred embodiment, it has been found that suitable nucleic sequences and their reverse complement can be used to alter the expression of any homologous, endogenous RNA which is in proximity to the suitable nucleic acid and its reverse complement. This is described in greater detail in Applicant's Assignee's co-pending provisional application having Appli-

cation No. 60/213,961 filed Jun. 23, 2000, the disclosure of which is hereby incorporated by reference.

[0049] In addition, alterations in a nucleic acid fragment which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded polypeptide, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the polypeptide molecule would also not be expected to alter the activity of the polypeptide. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Consequently, an isolated polynucleotide comprising a nucleotide sequence of at least one of 30 (preferably at least one of 40, most preferably at least one of 60) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO: 1, 3, 5, 7, 9, 11, 13, or 15, and the complement of such nucleotide sequences may be used in methods of selecting an isolated polynucleotide that affects the expression of a diverged delta-9 stearoyl desaturase polypeptide in a host cell. A method of selecting an isolated polynucleotide that affects the level of expression of a polypeptide in a virus or in a host cell (eukaryotic, such as plant or yeast, prokaryotic such as bacterial) may comprise the steps of: constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; introducing the isolated polynucleotide or the isolated chimeric gene into a host cell; measuring the level of a polypeptide or enzyme activity in the host cell containing the isolated polynucleotide; and comparing the level of a polypeptide or enzyme activity in the host cell containing the isolated polynucleotide with the level of a polypeptide or enzyme activity in a host cell that does not contain the isolated polynucleotide.

[0050] Moreover, substantially similar nucleic acid fragments may also be characterized by their ability to hybridize. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.). Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. Post-hybridization washes determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6×SSC, 0.5% SDS at room temperature for 15 min, then repeated with 2×SSC, 0.5% SDS at 45° C. for 30 min, and then repeated twice with 0.2×SSC, 0.5% SDS at 50° C. for 30 min. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2×SSC, 0.5% SDS was increased to 60° C. Another preferred set of highly stringent conditions uses two final washes in 0.1×SSC, 0.1% SDS at 65° C.

[0051] Substantially similar nucleic acid fragments of the instant invention may also be characterized by the percent identity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Suitable nucleic acid fragments (isolated polynucleotides of the present invention) encode polypeptides that are at least about 70% identical, preferably at least about 80% identical to the amino acid sequences reported herein. Preferred nucleic acid fragments encode amino acid sequences that are about 85% identical to the amino acid sequences reported herein. More preferred nucleic acid fragments encode amino acid sequences that are at least about 90% identical to the amino acid sequences reported herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are at least about 95% identical to the amino acid sequences reported herein. Suitable nucleic acid fragments not only have the above identities but typically encode a polypeptide having at least 50 amino acids, preferably at least 100 amino acids, more preferably at least 150 amino acids, still more preferably at least 200 amino acids, and most preferably at least 250 amino acids. Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, Wis.). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

[0052] It should be appreciated by one skilled in the art that genes encoding delta-9 desaturases can be identified in a number of ways. Conserved sequence motifs such as HSMPPEK corresponding to amino acids 67-73 of SEQ ID NO:2, LPLKPVVE corresponding to amino acids 89-96 of SEQ ID NO:2, EYFVVLVGDM corresponding to amino acids 132-141 of SEQ ID NO:2, EKTIV corresponding to amino acids 205-208 of SEQ ID NO:2, GMDPGT corresponding to amino acids 215-220 of SEQ ID NO:2, NNPYLGFVYTSFQERAT corresponding to amino acids 222-238 of SEQ ID NO:2, VLAR corresponding to amino acids 256-259 of SEQ ID NO:2, RIVE corresponding to amino acids 277-280 of SEQ ID NO:2, ITMPAHL corresponding to amino acids 302-308 of SEQ ID NO:2, or DFVCGLA corresponding to amino acids 364-370 of SEQ ID NO:2, can be used once several members of a diverged class are identified (as is the case in the present invention). In addition one can use hybridization, sequencing, and electronic alignment to aid the identification of gene candidates. These approaches can be coupled to assay of the polypeptide activity in bacterial, yeast, or plant host cells. Stable transgenic plants would provide a preferred method of determining the identity of a nucleic acid sequence encoding a delta-9 desaturase.

[0053] A “substantial portion” of an amino acid or nucleotide sequence comprises an amino acid or a nucleotide sequence that is sufficient to afford putative identification of the protein or gene that the amino acid or nucleotide sequence comprises. Amino acid and nucleotide sequences can be evaluated either manually by one skilled in the art, or by using computer-based sequence comparison and identification tools that employ algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In

general, a sequence of ten or more contiguous amino acids or thirty or more contiguous nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene-specific oligonucleotide probes comprising 30 or more contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., in situ hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12 or more nucleotides may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a “substantial portion” of a nucleotide sequence comprises a nucleotide sequence that will afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches amino acid and nucleotide sequences encoding polypeptides that comprise one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

[0054] “Codon degeneracy” refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment comprising a nucleotide sequence that encodes all or a substantial portion of the amino acid sequences set forth herein. The skilled artisan is well aware of the “codon-bias” exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a nucleic acid fragment for improved expression in a host cell, it is desirable to design the nucleic acid fragment such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

[0055] “Synthetic nucleic acid fragments” can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form larger nucleic acid fragments which may then be enzymatically assembled to construct the entire desired nucleic acid fragment. “Chemically synthesized”, as related to a nucleic acid fragment, means that the component nucleotides were assembled in vitro. Manual chemical synthesis of nucleic acid fragments may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the nucleic acid fragments can be tailored for optimal gene expression based on optimization of the nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

[0056] “Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory

sequences. “Chimeric gene” refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. “Endogenous gene” refers to a native gene in its natural location in the genome of an organism. A “foreign-gene” refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

[0057] “Coding sequence” refers to a nucleotide sequence that codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

[0058] “Promoter” refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a nucleotide sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or may be composed of different elements derived from different promoters found in nature, or may even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamoto and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

[0059] “Translation leader sequence” refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

[0060] “3' non-coding sequences” refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation

signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

[0061] “RNA transcript” refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from post-transcriptional processing of the primary transcript and is referred to as the mature RNA. “Messenger RNA (mRNA)” refers to the RNA that is without introns and that can be translated into polypeptides by the cell. “cDNA” refers to DNA that is complementary to and derived from an mRNA template. The cDNA can be single-stranded or converted to double stranded form using, for example, the Klenow fragment of DNA polymerase I. “Sense-RNA” refers to an RNA transcript that includes the mRNA and so can be translated into a polypeptide by the cell. “Antisense RNA” refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (see U.S. Pat. No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific nucleotide sequence, i.e., at the 5' non-coding sequence, 3' non coding sequence, introns, or the coding sequence. “Functional RNA” refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

[0062] The term “operably linked” refers to the association of two or more nucleic acid fragments on a single polynucleotide so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

[0063] An “intron” is an intervening sequence in a gene that does not encode a portion of the protein sequence. Thus, such sequences are transcribed into RNA but are then excised and are not translated. The term is also used for the excised RNA sequences. An “exon” is a portion of the sequence of a gene that is transcribed and is found in the mature messenger RNA derived from the gene, but is not necessarily a part of the sequence that encodes the final gene product.

[0064] The term “expression”, as used herein, refers to the production of a functional end-product. Expression of a gene involves transcription of the gene and translation of the mRNA into a precursor or mature protein. “Antisense inhibition” refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. “Overexpression” refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. “Co-suppression” refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Pat. No. 5,231,020, the disclosure of which is hereby incorporated by reference).

[0065] A “protein” or “polypeptide” is a chain of amino acids arranged in a specific order determined by the coding sequence in a polynucleotide encoding the polypeptide. Each protein or polypeptide has a unique function.

[0066] “Altered levels” or “altered expression” refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

[0067] “Null mutant” as used herein refers to a host cell which either does not express a certain polypeptide or expresses a polypeptide which is inactive or does not have any detectable expected enzymatic function.

[0068] “Mature protein” or the term “mature” when used in describing a protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. “Precursor protein” or the term “precursor” when used in describing a protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

[0069] A “chloroplast transit peptide” is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made. “Chloroplast transit sequence” refers to a nucleotide sequence that encodes a chloroplast transit peptide. A “signal peptide” is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (supra) can further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (supra) may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

[0070] The present invention describes a nucleic acid fragment that encodes a diverged delta-9 fatty acid desaturase. This enzyme catalyzes the introduction of a double bond between carbon atoms 9 and 10 of stearoyl-ACP to form oleoyl-ACP. It can also convert stearoyl-CoA into oleoyl-CoA, albeit with reduced efficiency. Transfer of the nucleic acid fragment of the invention, or a part thereof that encodes a functional enzyme, with suitable regulatory sequences into a living cell will result in the production or over-production of stearoyl-ACP desaturase, which in the presence of an appropriate electron donor, such as ferredoxin, may result in an increased level of unsaturation in cellular lipids, including oil, in tissues when the enzyme is absent or rate-limiting.

[0071] “Transformation” refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or “gene gun” transformation technology (Klein et al. (1987) *Nature* (London) 327:70-73; U.S. Pat. No. 4,945,050, incorporated herein by reference). Thus, isolated polynucleotides of the present invention can be incorporated into recombinant constructs, typically DNA constructs, capable of introduction into and replication in a host cell. Such a construct can be a vector that includes a replication system and sequences that are capable of transcription and translation of a polypeptide-encoding sequence in a given host cell. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants have been

described in, e.g., Pouwels et al., *Cloning Vectors: A Laboratory Manual*, 1985, supp. 1987; Weissbach and Weissbach, *Methods for Plant Molecular Biology*, Academic Press, 1989; and Flevin et al., *Plant Molecular Biology Manual*, Kluwer Academic Publishers, 1990. Typically, plant expression vectors include, for example, one or more cloned plant genes under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant expression vectors also can contain a promoter regulatory region (e.g., a regulatory region controlling inducible or constitutive, environmentally- or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

[0072] Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook et al. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter “Maniatis”).

[0073] “PCR” or “polymerase chain reaction” is well known by those skilled in the art as a technique used for the amplification of specific DNA segments (U.S. Pat. Nos. 4,683,195 and 4,800,159).

[0074] The present invention concerns an isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of: (a) first nucleotide sequence encoding a polypeptide of at least 400 amino acids having at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, or 16, or (b) a second nucleotide sequence comprising the complement of the first nucleotide sequence.

[0075] Preferably, the first nucleotide sequence comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, or 15, that codes for the polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, or 16.

[0076] Nucleic acid fragments encoding at least a portion of several diverged delta-9 fatty acid desaturases have been isolated and identified by comparison of random plant cDNA sequences to public databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization, and methods of DNA and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction, ligase chain reaction).

[0077] For example, genes encoding other diverged delta-9 stearoyl desaturases, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis). Moreover, an entire sequence can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primer DNA

labeling, nick translation, end-labeling techniques, or RNA probes using available in vitro transcription systems. In addition, specific primers can be designed and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

[0078] In addition, two short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) to generate cDNAs by using PCR to amplify copies of the region between a single point in the transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5673-5677; Loh et al. (1989) *Science* 243:217-220). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman and Martin (1989) *Techniques* 1:165). Consequently, a polynucleotide comprising a nucleotide sequence of at least one of 30 (preferably one of at least 40, most preferably one of at least 60) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, or 15 and the complement of such nucleotide sequences may be used in such methods to obtain a nucleic acid fragment encoding a substantial portion of an amino acid sequence of a polypeptide.

[0079] The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a diverged delta-9 stearoyl desaturase polypeptide, preferably a substantial portion of a plant diverged delta-9 stearoyl desaturase polypeptide, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 30 (preferably at least one of 40, most preferably at least one of 60) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, or 15, and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a diverged delta-9 stearoyl desaturase polypeptide.

[0080] Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expres-

sion libraries to isolate full-length cDNA clones of interest (Lerner (1984) *Adv. Immunol.* 36:1-34; Maniatis).

[0081] In another embodiment, this invention concerns viruses and host cells comprising either the chimeric constructs of the invention as described herein or an isolated polynucleotide of the invention as described herein. Examples of host cells which can be used to practice the invention include, but are not limited to, yeast, bacteria, and plants.

[0082] As was noted above, the nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed polypeptides are present at higher or lower levels than normal or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the level of mono-, poly- and unsaturated fatty acids in those cells.

[0083] Overexpression of the proteins of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. The chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

[0084] The terms "chimeric construct", "recombinant construct", "expression construct" and "recombinant expression construct" are used interchangeably herein. Such construct may be itself or may be used in conjunction with a vector. If a vector is used then the choice of vector is dependent upon the method that will be used to transform host plants as is well known to those skilled in the art. For example, a plasmid vector can be used. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al. (1985) *EMBO J.* 4:2411-2418; De Almeida et al. (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

[0085] For some applications it may be useful to direct the instant polypeptides to different cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by directing the coding sequence to encode the instant polypeptides with appropriate intracellular targeting sequences such as transit sequences (Keegstra (1989) *Cell* 56:247-253), signal sequences or sequences encoding endoplasmic reticulum localization (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel (1992) *Plant Phys.* 100:1627-1632) with or without removing targeting sequences that are already present. While the references cited give examples of each of these, the list is not exhaustive and more targeting signals of use may be discovered in the future.

[0086] It may also be desirable to reduce or eliminate expression of genes encoding the instant polypeptides in

plants for some applications. In order to accomplish this, a chimeric construct designed for co-suppression of the instant polypeptide can be constructed by linking a gene or gene fragment encoding that polypeptide to plant promoter sequences. Alternatively, a chimeric construct designed to express antisense RNA for all or part of the instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either the co-suppression or antisense chimeric constructs could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated. In a preferred embodiment, it has been found that suitable nucleic sequences and their reverse complement can be used to alter the expression of any homologous, endogenous RNA which is in proximity to the suitable nucleic acid and its reverse complement. This is described in greater detail in Applicant's Assignee's co-pending provisional application having Application No. 60/213,961 filed Jun. 23, 2000, the disclosure of which is hereby incorporated by reference.

[0087] Molecular genetic solutions to the generation of plants with altered gene expression have a decided advantage over more traditional plant breeding approaches. Changes in plant phenotypes can be produced by specifically inhibiting expression of one or more genes by antisense inhibition or cosuppression (U.S. Pat. Nos. 5,190,931, 5,107,065 and 5,283,323). An antisense or cosuppression construct would act as a dominant negative regulator of gene activity. While conventional mutations can yield negative regulation of gene activity these effects are most likely recessive. The dominant negative regulation available with a transgenic approach may be advantageous from a breeding perspective. In addition, the ability to restrict the expression of a specific phenotype to the reproductive tissues of the plant by the use of tissue specific promoters may confer agronomic advantages relative to conventional mutations which may have an effect in all tissues in which a mutant gene is ordinarily expressed.

[0088] The person skilled in the art will know that special considerations are associated with the use of antisense or cosuppression technologies in order to reduce expression of particular genes. For example, the proper level of expression of sense or antisense genes may require the use of different chimeric genes utilizing different regulatory elements known to the skilled artisan. Once transgenic plants are obtained by one of the methods described above, it will be necessary to screen individual transgenics for those that most effectively display the desired phenotype. Accordingly, the skilled artisan will develop methods for screening large numbers of transformants. The nature of these screens will generally be chosen on practical grounds. For example, one can screen by looking for changes in gene expression by using antibodies specific for the protein encoded by the gene being suppressed, or one could establish assays that specifically measure enzyme activity. A preferred method will be one which allows large numbers of samples to be processed rapidly, since it will be expected that a large number of transformants will be negative for the desired phenotype.

[0089] In another embodiment, the present invention concerns a polypeptide of at least 400 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, or 16.

[0090] The instant polypeptides (or portions thereof) may be produced in heterologous host cells, particularly in the

cells of microbial hosts, and can be used to prepare antibodies to these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting the polypeptides of the instant invention in situ in cells or in vitro in cell extracts. Preferred heterologous host cells for production of the instant polypeptides are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to make a chimeric construct for production of the instant polypeptides. This chimeric construct could then be introduced into appropriate microorganisms via transformation to provide high level expression of the encoded diverged delta-9 fatty acid desaturase. An example of a vector for high level expression of the instant polypeptides in a bacterial host is provided (Example 6).

[0091] All or a substantial portion of the polynucleotides of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part of, and used as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al. (1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein et al. (1980) *Am. J. Hum. Genet.* 32:314-331).

[0092] The production and use of plant gene-derived probes for use in genetic mapping is described in Bernatzky and Tanksley (1986) *Plant Mol. Biol. Reporter* 4:37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology outlined above or variations thereof. For example, F2 intercross populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

[0093] Nucleic acid probes derived from the instant nucleic acid sequences may also be used for physical mapping (i.e., placement of sequences on physical maps; see Hoheisel et al. in: *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

[0094] In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence in situ hybridization (FISH) mapping (Trask (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; see Laan et al. (1995) *Genome Res.* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

[0095] A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using

the instant nucleic acid sequences. Examples include allele-specific amplification (Kazazian (1989) *J. Lab. Clin. Med.* 11:95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter et al. (1997) *Nat. Genet.* 7:22-28) and Happy Mapping (Dear and Cook (1989) *Nucleic Acid Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

[0096] Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these genes contained in a maize population carrying mutations in all possible genes (Ballinger and Benzer (1989) *Proc. Natl. Acad. Sci. USA* 86:9402-9406; Koes et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:8149-8153; Bensen et al. (1995) *Plant Cell* 7:75-84). The latter approach may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, supra). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the instant polypeptides. Alternatively, the instant nucleic acid fragment may be used as a hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding the instant polypeptides can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the instant polypeptides disclosed herein.

[0097] Methods for assaying delta-9 fatty acid desaturase activities in *E. coli* have been previously described (U.S. Pat. Nos. 5,443,974 and 5,760,206). Fatty acid analysis of oil samples is performed by gas chromatography. Briefly, fatty acid (FA) determination was done from a total of 300-400 mg of tissue lyophilized for 24 hours. The tissue was then ground using a FastPrep mill (Bio101) at 4.5 speed and 20 seconds in the presence of 0.5 ml of 2.5% Sulfuric Acid+97.5% Methanol and Heptadecanoic acid (17:0, stock 10 mg/ml in Toluene) as an external standard. Thereafter, another 0.5 ml 2.5% Sulfuric Acid+97.5% Methanol was used to wash each tube and incubate in 95° C. for 1 hour for transesterification. The tubes were removed from the water bath and allowed to cool down to room temperature. FAs were extracted in one volume of heptane:H₂O (1:1) and cleared by centrifugation. The supernatant (50 ul) containing the fatty acid methyl esters were loaded into a Hewlett Packard 6890 gas chromatograph

fitted with a 30 mx0.32 mm Omegawax column and the separated peaks were analyzed and characterized.

EXAMPLES

[0098] The present invention is further defined in the following Examples, in which parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

[0099] The disclosure of each reference set forth herein is incorporated herein by reference in its entirety.

Example 1

Composition of cDNA Libraries; Isolation and Sequencing of cDNA Clones

[0100] cDNA libraries representing mRNAs from various soybean, corn, and rice tissues were prepared. The characteristics of the libraries are described below.

TABLE 3

cDNA Libraries from Soybean, Corn, and Rice		
Library	Description	Clone
se6	Soybean Embryo, 26 Days After Flowering	se6.pk0026.a8
cbn10	Corn Developing Kernel (Embryo and Endosperm); 10 Days After Pollination	cbn10.pk0061.a3
cpd1c	Corn (<i>Zea mays</i> L.) pooled BMS treated with chemicals related to protein kinases	cpd1c.pk014.i18
rds1c	Rice (<i>Oryza sativa</i> , YM) developing seeds 1	rds1c.pk007.g19
rsl1n	Rice (<i>Oryza sativa</i> , YM) 15 day old seedling normalized	rsl1n.pk008.j18:fs

[0101] cDNA libraries may be prepared by any one of many methods available. For example, the cDNAs may be introduced into plasmid vectors by first preparing the cDNA libraries in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene Cloning Systems, La Jolla, Calif.). The Uni-ZAP™ XR libraries are converted into plasmid libraries according to the protocol provided by Stratagene. Upon conversion, cDNA inserts will be contained in the plasmid vector pBluescript. In addition, the cDNAs may be introduced directly into precut Bluescript II SK(+) vectors (Stratagene) using T4 DNA ligase (New England Biolabs), followed by transfection into DH10B cells according to the manufacturer's protocol (GIBCO BRL Products). Once the cDNA inserts are in plasmid vectors, plasmid DNAs are prepared from randomly picked bacterial colonies containing recombinant pBluescript plasmids, or the insert cDNA sequences are amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted cDNA sequences. Amplified insert DNAs or plasmid DNAs are

sequenced in dye-primer sequencing reactions to generate partial cDNA sequences (expressed sequence tags or "ESTs"; see Adams et al., (1991) *Science* 252:1651-1656). The resulting ESTs are analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

[0102] Full-insert sequence (FIS) data is generated utilizing a modified transposition protocol. Clones identified for FIS are recovered from archived glycerol stocks as single colonies, and plasmid DNAs are isolated via alkaline lysis. Isolated DNA templates are reacted with vector primed M13 forward and reverse oligonucleotides in a PCR-based sequencing reaction and loaded onto automated sequencers. Confirmation of clone identification is performed by sequence alignment to the original EST sequence from which the FIS request is made.

[0103] Confirmed templates are transposed via the Primer Island transposition kit (PE Applied Biosystems, Foster City, Calif.) which is based upon the *Saccharomyces cerevisiae* Ty1 transposable element (Devine and Boeke (1994) *Nucleic Acids Res.* 22:3765-3772). The in vitro transposition system places unique binding sites randomly throughout a population of large DNA molecules. The transposed DNA is then used to transform DH10B electro-competent cells (Gibco BRL/Life Technologies, Rockville, Md.) via electroporation. The transposable element contains an additional selectable marker (named DHFR; Fling and Richards (1983) *Nucleic Acids Res.* 11:5147-5158), allowing for dual selection on agar plates of only those subclones containing the integrated transposon. Multiple subclones are randomly selected from each transposition reaction, plasmid DNAs are prepared via alkaline lysis, and templates are sequenced (ABI Prism dye-terminator ReadyReaction mix) outward from the transposition event site, utilizing unique primers specific to the binding sites within the transposon.

[0104] Sequence data is collected (ABI Prism Collections) and assembled using Phred/Phrap (P. Green, University of Washington, Seattle). Phred/Phrap is a public domain software program which re-reads the ABI sequence data, re-calls the bases, assigns quality values, and writes the base calls and quality values into editable output files. The Phrap sequence assembly program uses these quality values to increase the accuracy of the assembled sequence contigs. Assemblies are viewed by the Consed sequence editor (D. Gordon, University of Washington, Seattle).

Example 2

Identification of cDNA Clones

[0105] cDNA clones encoding a diverged delta-9 fatty acid desaturase were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTn algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database

using the BLASTX algorithm (Gish and States (1993) *Nat. Genet.* 3:266-272) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as "pLog" values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST "hit" represent homologous proteins.

[0106] ESTs submitted for analysis are compared to the genbank database as described above. ESTs that contain sequences more 5- or 3-prime can be found by using the BLASTn algorithm (Altschul et al (1997) *Nucleic Acids Res.* 25:3389-3402.) against the Du Pont proprietary database comparing nucleotide sequences that share common or overlapping regions of sequence homology. Where common or overlapping sequences exist between two or more nucleic acid fragments, the sequences can be assembled into a single contiguous nucleotide sequence, thus extending the original fragment in either the 5 or 3 prime direction. Once the most 5-prime EST is identified, its complete sequence can be determined by Full Insert Sequencing as described in Example 1. Homologous genes belonging to different species can be found by comparing the amino acid sequence of a known gene (from either a proprietary source or a public database) against an EST database using the tBLASTn algorithm. The tBLASTn algorithm searches an amino acid query against a nucleotide database that is translated in all 6 reading frames. This search allows for differences in nucleotide codon usage between different species, and for codon degeneracy.

Example 3

Characterization of cDNA Clones Encoding a Diverged Delta-9, or Stearoyl-ACP, Desaturase

[0107] The BLASTX search using the EST sequences from clones listed in Table 3 revealed similarity of the polypeptides encoded by the cDNAs to a diverged delta-9, or stearoyl-ACP, desaturase from lupine (*Lupinus luteus*), cucumber (*Cucumis sativus*), *Arabidopsis* (*Arabidopsis thaliana*), jojoba (*Simmondsia chinensis*), *Arabidopsis* (*Arabidopsis thaliana*), and flax (*Linum usitatissimum*) (NCBI General Identifier Nos. gi 4704824, gi 417820, gi 7523660, gi 267036, gi 6957724, and gi 3355632 respectively). Shown in Table 4 are the BLAST results for individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), the sequences of contigs assembled from two or more ESTs ("Contig"), sequences of contigs assembled from an FIS and one or more ESTs ("Contig*"), or sequences encoding an entire protein derived from an FIS, a contig, or an FIS and PCR ("CGS"):

TABLE 4

BLAST Results for Sequences Encoding Polypeptides Homologous to a Diverged Delta-9, or Stearoyl-ACP, Desaturase			
Clone	Status	BLAST pLog	gi #
se6.pk0026.a8	CGS	254.00	4704824
cbn10.pk0061.a3	EST	2.52	4704824
cpd1c.pk014.118	contig	107.00	417820
rds1c.pk007.g19	EST	33.04	7523660
cbn10.pk0061.a3:fis	CGS	113.00	267036
cpd1c.pk014.118:fis	CGS	149.00	4704824

TABLE 4-continued

BLAST Results for Sequences Encoding Polypeptides Homologous to a Diverged Delta-9, or Stearoyl-ACP, Desaturase			
Clone	Status	BLAST pLog	gi #
rds1c.pk007.g19:fis	FIS	60.52	6957724
rsl1n.pk008.j18:fis	CGS	147.00	3355632

[0108] FIG. 1 presents an alignment of the amino acid sequences set forth in SEQ ID NO:2, 10, 12, 14, and 16, and the lupine, jojoba, *Arabidopsis*, and flax sequences (SEQ ID NO:17, 20, 21, and 22) and the original soybean delta-9 desaturase presented in U.S. Pat. No. 5,760,206 (SEQ ID NO:23). The data in Table 5 represents a calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, and 16, and the lupine, cucumber, *Arabidopsis*, jojoba, *Arabidopsis*, and flax sequences (SEQ ID NOs:17, 18, 19, 20, 21, and 22; NCBI General Identifier Nos. gi 4704824, gi 417820, gi 7523660, gi 267036, gi 6957724, and gi 3355632 respectively).

TABLE 5

Percent Identity of Polypeptides Homologous to a Diverged Delta-9, or Stearoyl-ACP, Desaturase		
SEQ ID NO.	Percent Identity	gi #
2	77.6%	4704824
4	16.4%	4704824
6	62.3%	417820
8	59.5%	7523660
10	49.2%	267036
12	64.2%	4704824
14	50.2%	6957724
16	64.0%	3355632

[0109] Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, Wis.). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode a substantial portion of a diverged delta-9, or stearyl-ACP, desaturase. Confirmation of the biochemical identity of each clone is accomplished according to methods well known to those skilled in the art (U.S. Pat. No. 5,760,206).

Example 4

Expression of Chimeric Constructs in Monocot Cells

[0110] A chimeric construct comprising a cDNA encoding the instant polypeptides in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide

primers. Cloning sites (NcoI or SmaI) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below. Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes NcoI and SmaI and fractionated on an agarose gel. The appropriate band can be isolated from the gel and combined with a 4.9 kb NcoI-SmaI fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas, Va. 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb Sall-NcoI promoter fragment of the maize 27 kD zein gene and a 0.96 kb SmaI-Sall fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15° C. overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli* XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U.S. Biochemical). The resulting plasmid construct would comprise a chimeric construct encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment encoding the instant polypeptides, and the 10 kD zein 3' region.

[0111] The chimeric construct described above can then be introduced into corn cells by the following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated 10 to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al. (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27° C. Friable embryogenic callus consisting of undifferentiated masses of cells with somatic proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

[0112] The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable marker. This plasmid contains the Pat gene (see European Patent Publication 0 242 236) which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The pat gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

[0113] The particle bombardment method (Klein et al. (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1 μm in diameter) are coated with DNA using the following technique. Ten μg of plasmid DNAs are added to 50 μL of a suspension of gold particles (60 mg per mL). Calcium chloride (50 μL of a 2.5 M solution) and spermidine free base (20 μL of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 minutes, the tubes are briefly centrifuged (5 sec at

15,000 rpm) and the supernatant removed. The particles are resuspended in 200 μ L of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30 μ L of ethanol. An aliquot (5 μ L) of the DNA-coated gold particles can be placed in the center of a Kapton™ flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic™ PDS-1000/He (Bio-Rad Instruments, Hercules Calif.), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

[0114] For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

[0115] Seven days after bombardment the tissue can be transferred to N6 medium that contains glufosinate (2 mg per liter) and lacks casein or proline. The tissue continues to grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing glufosinate. After 6 weeks, areas of about 1 cm in diameter of actively growing callus can be identified on some of the plates containing the glufosinate-supplemented medium. These calli may continue to grow when sub-cultured on the selective medium.

[0116] Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the tissue can be transferred to regeneration medium (Fromm et al. (1990) *Bio/Technology* 8:833-839).

Example 5

Expression of Chimeric Constructs in Dicot Cells

[0117] A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the β subunit of the seed storage protein phaseolin from the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used for expression of the instant polypeptides in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the 5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), Sma I, Kpn I and Xba I. The entire cassette is flanked by Hind III sites.

[0118] The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

[0119] Soybean embryos may then be transformed with the expression vector comprising sequences encoding the instant polypeptides. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature

seeds of the soybean cultivar A2872, can be cultured in the light or dark at 26° C. on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

[0120] Soybean embryogenic suspension cultures can be maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26° C. with florescent lights on a 16:8 hour day/night schedule. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

[0121] Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein et al. (1987) *Nature* (London) 327:70-73, U.S. Pat. No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used for these transformations.

[0122] A selectable marker gene which can be used to facilitate soybean transformation is a chimeric construct composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al. (1983) *Gene* 25:179-188) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression cassette comprising the phaseolin 5' region, the fragment encoding the instant polypeptides and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

[0123] To 50 μ L of a 60 mg/mL 1 μ m gold particle suspension is added (in order): 5 μ L DNA (1 μ g/ μ L), 20 μ L spermidine (0.1 M), and 50 μ L CaCl₂ (2.5 M). The particle preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 μ L 70% ethanol and resuspended in 40 μ L of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five μ L of the DNA-coated gold particles are then loaded on each macro carrier disk.

[0124] Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

[0125] Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcul-

tured and maintained as clusters of immature embryos or regenerated into whole plants by maturation and germination of individual somatic embryos.

Example 6

Expression of Chimeric Constructs in Microbial Cells

[0126] The cDNAs encoding the instant polypeptides can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter system. Plasmid pBT430 was constructed by first destroying the EcoR I and Hind III sites in pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using oligonucleotide-directed mutagenesis. The DNA sequence of pET-3aM in this region, 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

[0127] Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% low melting agarose gel. Buffer and agarose contain 10 µg/ml ethidium bromide for visualization of the DNA fragment. The fragment can then be purified from the agarose gel by digestion with GELase™ (Epicentre Technologies, Madison, Wis.) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20 µL of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase (New England Biolabs (NEB), Beverly, Mass.). The fragment containing the ligated adapters can be purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16° C. for 15 hours followed by transformation into DH5 electrocompetent cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and 100 µg/mL ampicillin. Transformants containing the gene encoding the instant polypeptides are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

[0128] For high level expression, a plasmid clone with the cDNA insert in the correct orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21 (DE3) (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25° C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant determined. One µg of protein from the soluble fraction of the culture can

be separated by SDS-polyacrylamide gel electrophoresis. Gels can be observed for protein bands migrating at the expected molecular weight.

Example 7

Transformation of Somatic Soybean Embryo Cultures

[0129] Soybean embryogenic suspension cultures were maintained in 35 ml liquid media (SB55 or SBP6) on a rotary shaker, 150 rpm, at 28° C. with mixed fluorescent and incandescent lights on a 16:8 h day/night schedule. Cultures were subcultured every four weeks by inoculating approximately 35 mg of tissue into 35 ml of liquid medium.

TABLE 6

Stock Solutions (g/L):	
<u>MS Sulfate 100X Stock</u>	
MgSO ₄ 7H ₂ O	37.0
MnSO ₄ H ₂ O	1.69
ZnSO ₄ 7H ₂ O	0.86
CuSO ₄ 5H ₂ O	0.0025
<u>MS Halides 100X Stock</u>	
CaCl ₂ 2H ₂ O	44.0
KI	0.083
CoCl ₂ 6H ₂ O	0.00125
KH ₂ PO ₄	17.0
H ₃ BO ₃	0.62
Na ₂ MoO ₄ 2H ₂ O	0.025
<u>MS FeEDTA 100X Stock</u>	
Na ₂ EDTA	3.724
FeSO ₄ 7H ₂ O	2.784
<u>B5 Vitamin Stock</u>	
10 g m-inositol	
100 mg nicotinic acid	
100 mg pyridoxine HCl	
1 g thiamine	
SB55 (per Liter, pH 5.7)	
10 mL each MS stocks	
1 mL B5 Vitamin stock	
0.8 g NH ₄ NO ₃	
3.033 g KNO ₃	
1 mL 2,4-D (10 mg/mL stock)	
60 g sucrose	
0.667 g asparagine	
SBP6	
same as SB55 except 0.5 mL 2,4-D	
SB103 (per Liter, pH 5.7)	
1X MS Salts	
6% maltose	
750 mg MgCl ₂	
0.2% Gelrite	
SB71-1 (per Liter, pH 5.7)	
1X B5 salts	
1 ml B5 vitamin stock	
3% sucrose	
750 mg MgCl ₂	
0.2% Gelrite	

[0130] Soybean embryogenic suspension cultures were transformed with pTC3 by the method of particle gun bombardment (Kline et al. (1987) *Nature* 327:70). A DuPont Biolistic PDS1000/HE instrument (helium retrofit) was used for these transformations.

[0131] To 50 ml of a 60 mg/ml 1 mm gold particle suspension was added (in order); 5 μ l DNA (1 μ g/ μ l), 20 μ l spermidine (0.1 M), and 50 μ l CaCl₂ (2.5 M). The particle preparation was agitated for 3 min, spun in a microfuge for 10 sec and the supernatant removed. The DNA-coated particles were then washed once in 400 μ l 70% ethanol and are suspended in 40 μ l of anhydrous ethanol. The DNA/particle suspension was sonicated three times for 1 sec each. Five μ l of the DNA-coated gold particles were then loaded on each macro carrier disk.

[0132] Approximately 300-400 mg of a four week old suspension culture was placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue were normally bombarded. Membrane rupture pressure was set at 1000 psi and the chamber was evacuated to a vacuum of 28 inches of mercury. The tissue was placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue was placed back into liquid and cultured as described above.

[0133] Eleven days post bombardment, the liquid media was exchanged with fresh SB55 containing 50 mg/ml hygromycin. The selective media was refreshed weekly. Seven weeks post bombardment, green, transformed tissue was observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue was removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Thus each new line was treated as independent transformation event. These suspensions can then be maintained as suspensions of embryos clustered in an immature developmental stage through subculture or regenerated into whole plants by maturation and germination of individual somatic embryos.

[0134] Independent lines of transformed embryogenic clusters are removed from liquid culture and placed on a solid agar media (SB103) containing no hormones or antibiotics. Embryos are cultured for four weeks at 26° C. with mixed fluorescent and incandescent lights on a 16:8 h day/night schedule. During this period, individual embryos are removed from the clusters and screened for alterations in their fatty acid compositions (Example 8). Co-suppression of Fad2 results in a reduction of polyunsaturated fatty acids and an increase in oleic acid content. Co-suppression of the delta-9 desaturases of the instant invention result in an increase in the accumulation of stearic acid (18:0 fatty acid).

Example 8

The Phenotype of Transgenic Soybean Somatic Embryos is Predictive of Seed Phenotypes from Resultant Regenerated Plants

[0135] Mature somatic soybean embryos are a good model for zygotic embryos. While in the globular embryo state in liquid culture, somatic soybean embryos contain very low amounts of triacylglycerol or storage proteins typical of maturing, zygotic soybean embryos. At this developmental stage, the ratio of total triacylglyceride to total polar lipid (phospholipids and glycolipid) is about 1:4, as is typical of zygotic soybean embryos at the developmental stage from which the somatic embryo culture was initiated. At the globular stage as well, the mRNAs for the prominent seed proteins, α' subunit of β -conglycinin, kunitz trypsin inhibitor 3, and seed lectin are essentially absent. Upon transfer to hormone-

free media to allow differentiation to the maturing somatic embryo state, triacylglycerol becomes the most abundant lipid class. As well, mRNAs for α' -subunit of β -conglycinin, kunitz trypsin inhibitor 3 and seed lectin become very abundant messages in the total mRNA population. On this basis somatic soybean embryo system behaves very similarly to maturing zygotic soybean embryos in vivo, and is therefore a good and rapid model system for analyzing the phenotypic effects of modifying the expression of genes in the fatty acid biosynthesis pathway.

[0136] Most importantly, the model system is also predictive of the fatty acid composition of seeds from plants derived from transgenic embryos. This is illustrated with two different antisense constructs in two different types of experiment that were constructed following the protocols set forth in the PCT Publication Nos. WO 93/11245 and WO 94/11516. Liquid culture globular embryos were transformed with a chimeric gene comprising a soybean microsomal Δ^{15} desaturase as described in PCT Publication No. WO 93/11245 which was published on Jun. 10, 1993, the disclosure of which is hereby incorporated by reference (experiment 1,) or a soybean microsomal Δ^{12} desaturase as described in PCT Publication No. WO 94/11516 which was published on May 26, 1994, the disclosure of which is hereby incorporated by reference (experiment 2). Both gene constructs were introduced in antisense orientation under the control of a seed-specific promoter (β -conglycinin promoter) and gave rise to mature embryos. The fatty acid content of mature somatic embryos from lines transformed with vector only (control) and the vector containing the antisense chimeric genes as well as of seeds of plants regenerated from them was determined.

[0137] One set of embryos from each line was analyzed for fatty acid content and another set of embryos from that same line was regenerated into plants. Fatty acid analysis of single embryos was determined either by direct trans-esterification of individual seeds in 0.5 mL of methanolic H₂SO₄ (2.5%) or by hexane extraction of bulk seed samples followed by trans-esterification of an aliquot in 0.8 mL of 1% sodium methoxide in methanol. Fatty acid methyl esters were extracted from the methanolic solutions into hexane after the addition of an equal volume of water. In all cases, if there was a reduced 18:3 content in a transgenic embryo line when compared to an untransformed control, then a corresponding reduction in 18:3 content was also observed in the segregating seeds of the plant derived from that transformed line (Table 7).

TABLE 7

Percent 18:3 Content Of Embryos and Seeds of Control and Δ^{15} Antisense Construct Transgenic Soybean Lines		
Transformant Line	Embryo Average (SD, n = 10)	Seed Average (SD, n = 10)
Control	12.1 (2.6)	8.9 (0.8)
Δ^{15} antisense, line 1	5.6 (1.2)	4.3 (1.6)
Δ^{15} antisense, line 2	8.9 (2.2)	2.5 (1.8)
Δ^{15} antisense, line 3	7.3 (1.1)	4.9 (1.9)
Δ^{15} antisense, line 4	7.0 (1.9)	2.4 (1.7)
Δ^{15} antisense, line 5	8.5 (1.9)	4.5 (2.2)
Δ^{15} antisense, line 6	7.6 (1.6)	4.6 (1.6)

*[Seeds which were segregating with wild-type phenotype and without a copy of the transgene are not included in these averages]

[0138] In addition, different lines containing the same antisense construct, were used for fatty acid analysis in somatic embryos and for regeneration into plants. About 55% of the

transformed embryo lines showed an increased 18:1 content when compared with control lines (Table 8). Soybean seeds, of plants regenerated from different somatic embryo lines containing the same antisense construct, had a similar frequency (53%) of high oleate transformants as the somatic embryos (Table 8). On occasion, an embryo line may be chimeric. That is, 10-70% of the embryos in a line may not contain the transgene. The remaining embryos that do contain the transgene, have been found in all cases to be clonal. In such a case, plants with both wild type and transgenic phenotypes may be regenerated from a single, transgenic line, even if most of the embryos analyzed from that line had a transgenic phenotype. An example of this is shown in Table 9, in which, of 5 plants regenerated from a single embryo line, 3 have a high oleic phenotype and two were wild type. In most cases, all the plants regenerated from a single transgenic line will have seeds containing the transgene. Thus, it was concluded that an altered fatty acid phenotype observed in a transgenic, mature somatic embryo line is predictive of an altered fatty acid composition of seeds of plants derived from that line.

TABLE 8

Oleate Levels in Somatic Embryos and Seeds of Regenerated Soybeans Transformed With, or Without, Δ^{12} Desaturase Antisense Construct			
	# of Vector Lines	# of Lines with High 18:1	Average* %18:1
Somatic embryos:			
Control	19	0	12.0
Δ^{12} antisense	20	11	35.3
Seeds of regenerated plants:			
Control	6	0	18.2
Δ^{12} antisense	17	9	44.4

*average 18:1 of transgenics is the average of all embryos or seeds transformed with the Δ^{12} antisense construct in which at least one embryo or seed from that line had an 18:1 content greater than 2 standard deviations from the control value (12.0 in embryos, 18.2 in seeds). The control average is the average of embryos or seeds which do not contain any transgenic DNA but have been treated in an identical manner to the transgenics.

TABLE 9

Analysis of Seeds From Five Independent Plants Segregating From Plant Line 4		
Plant #	Average seed 18:1%	Highest seed 18:1%
1	18.0	26.3
2	33.6	72.1
7	13.6	21.2
9	32.9	57.3
11	24.5	41.7

[0139] Mean of 15-20 seeds from 5 different plants regenerated from a single embryo line. Only plants # 2, 9 and 11 have seeds with a high 18:1 phenotype.

Example 9

Suppression in Soybean of Fad2 by ELVISLIVES Complementary Region

[0140] Cosuppression of plant genes is covered in a U.S. provisional patent application 60/213,961 filed on Jun. 23, 2000, and nationally filed in the USPTO as application Ser. No. 09/887,194 on Jun. 22, 2001, the contents of which are hereby incorporated by reference. Constructs have now been made which have "synthetic complementary regions" (SCR). Since complementary regions from Fad 2 can successfully suppress a thioesterase target, and a Cer3 complementary region can suppress Fad2, it was deduced that it may be possible to use any complementary sequence to reduce the expression of a target. In this example the target sequence is placed between complementary sequences that are not known to be part of any biologically derived gene or genome (i.e. sequences that are "synthetic" or conjured up from the mind of the inventor). The target DNA would therefore be in the sense or antisense orientation and the complementary RNA would be unrelated to any known nucleic acid sequence. It is possible to design a standard "suppression vector" into which pieces of any target gene for suppression could be dropped. The plasmids pKS106, pKS124, and pKS133 exemplify this. One skilled in the art will appreciate that all of the plasmid vectors contain antibiotic selection genes such as, but not limited to, hygromycin phosphotransferase with promoters such as the T7 inducible promoter.

[0141] pKS106 uses the beta-conglycinin promoter while the pKS124 and 133 plasmids use the Kti promoter, both of these promoters exhibit strong tissue specific expression in the seeds of soybean. pKS106 uses a 3' termination region from the phaseolin gene, and pKS124 and 133 use a Kti 3' termination region. pKS106 and 124 have single copies of the 36 nucleotide EagI-ELVISLIVES sequence surrounding a NotI site (the amino acids given in parentheses are back-translated from the complementary strand): SEQ ID NO:24.

EagI E L V I S L I V E S NotI
CGGCCG GAG CTG GTC ATC TCG CTC ATC GTC GAG TCG GCGGCCG

(S) (E) (V) (I) (L) (S) (I) (V) (L) (E) EagI
CGA CTC GAC GAT GAG CGA GAT GAC CAG CTC CCGCCG

pKS 133 has 2X copies of ELVISLIVES surrounding the NotI site:
SEQ ID NO:25

EagI E L V I S L I V E S EagI E L V I S
cgccggagctggtcctctcgctcatcgctcgagtcg gcgccg gagctggctctctg

L I V E S NotI (S) (E) (V) (I) (L) (S) (I) (V) (L) (E) EagI
ctcatcgctcgagtcg gcgccgc cgactcgacgatgagcgagatgaccagctc cgcccg

(S) (E) (V) (I) (L) (S) (I) (V) (L) (E) EagI
cgactcgacgatgagcgagatgaccagctc cgcccg

[0142] The idea is that the single EL linker (SCR) can be duplicated to increase stem lengths in increments of approximately 40 nucleotides. A series of vectors will cover the SCR lengths between 40 bp and the 300 bp. Various target gene lengths are also under evaluation. It is believed that certain combinations of target lengths and complementary region lengths will give optimum suppression of the target, although preliminary results would indicate that the suppression phenomenon works well over a wide range of sizes and sequences. It is also believed that the lengths and ratios providing optimum suppression may vary somewhat given different target sequences and/or complementary regions.

[0143] The efficiency of Fad2 suppression using 1XEL (pKS132) was compared to Fad2 suppression using the 2XEL (pKS136) construct. Hygromycin resistant lines of soybean embryos were isolated from independent transformation experiments with pKS132 and pKS136. Out of 98 lines containing pKS132, 69% displayed the high oleic phenotype. Out of 54 lines containing pKS136, 70% displayed the high oleic acid phenotype. Thus, both 1X and 2XEL constructs efficiently suppressed the Fad2 target gene.

Example 10

Suppression the Diverged Delta-9 Desaturase Results in High Stearate Phenotypes

[0144] The two soybean delta-9 desaturase genes previously identified, designated pDS 1 and 2 (U.S. Pat. Nos. 5,443,974 and 5,760,206) share a high degree of homology to other known delta-9 desaturase genes such as castor and safflower (U.S. Pat. No. 5,723,595). The genes of the present invention have less than 65% amino acid sequence identity to these previously described plant delta-9 desaturase polypeptides. All of the soybean delta-9 desaturase genes were placed into *E. coli* and shown to have delta-9 desaturase activity. To test if the three genes had comparable activities in vivo, transgenic plants were constructed.

[0145] Delta-9 desaturase enzymes introduce a double bond into stearic acid to form oleic acid. Inhibition of this activity should result in an increase in stearic acid content and a correlative reduction in unsaturated fatty acids in the oil. An antisense construct of pDS1 (pKS6) was made using the entire coding region in reverse orientation inserted into the SmaI/XbaI site of pCST2 (PCT Publication No. WO 94/11516, published May 26, 1994) behind a beta-conglycinin promoter. A cosuppression construct was made (pRB1) where the HindIII fragment containing the beta-conglycinin promoter and the phaseolin 3' terminator from pAW35 (U.S. Pat. No. 5,952,544) was inserted into the HindIII site of

pML18 (PCT Publication No. WO 94/11516, published May 26, 1994) to form pBS19. The coding region of pDS1 was inserted into the Not I site of pBS19 to form pRB1. Finally, a cosuppression construct was made using pDS3 (pBS68, SEQ ID NO:26) by placing approximately 950 basepairs of pDS3 in the sense orientation between 2XEL complementary regions as described in Example 9 (the pDS region of pBS68 is from positions 6054-6611 linked to 1-411 of SEQ ID NO:26). The construct has a Kti3 promoter (position 3260-5348 of SEQ ID NO:26), a Kti3 terminator (position 523-725 of SEQ ID NO:26), and hygromycin selection (position 1920-880 of SEQ ID NO:26). Soybean transformations were done as previously described (Example 7), and soybean embryo tissue was assayed. As outlined in Example 8, soybean embryo tissue is representative of seed tissues when seed specific promoters such as beta-conglycinin or Kti3 are used.

[0146] The results shown in Table 10 demonstrate that pDS3 is as good, or better, than pDS1 at increasing stearic acid content in oils when cosuppressed in plant tissue. On average there is a 7.4-fold increase in 18:0 content with pDS3 (pBS68) versus 4.5 for the cosuppressed pDS1. Antisense or cosuppression gave similar results. The transformants that showed the highest levels of stearic acid are shown in the "best" columns.

TABLE 10

	18:0 Content of Wild Type and Transgenic High 18:0 Soybean Somatic Embryos					
	wild type (s.d.)	high 18:0 (s.d.)	fold increase	best 18:0	best fold increase	# of high 18:0 events
pKS6	3.6 (0.8)	16.7 (5.6)	4.6x	34.5	9.6x	40
pRB1	3.4 (0.5)	15.4 (3.7)	4.5x	20.6	6.1x	10
pBS68	2.5 (0.8)	18.6 (6.9)	7.4x	29.1	11.6x	10

[0147] These results confirm that the diverged delta-9 desaturase sequences do encode functional enzymes. Furthermore, pDS3 may be the dominant activity found in soybeans. The conserved sequence elements KEIPDDYF-WLVGDMITTEEALPTYQTMLNT corresponding to positions 116-145 of SEQ ID NO:23; and DYADILE-FLVGRWK corresponding to positions 324-337 of SEQ ID NO:23 from the Thompson patent (U.S. Pat. No. 5,723,595) that are claimed to be indicative of delta-9 desaturases are not conserved in the diverged sequences of the instant invention. Therefore, the sequences of the instant invention define a new functional class of plant delta-9 desaturase genes.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 26

<210> SEQ ID NO 1

<211> LENGTH: 1560

<212> TYPE: DNA

<213> ORGANISM: Glycine max

<400> SEQUENCE: 1

gaggcggttg atctggcact cgttttgctg tggctgctct ctgaaactga aagcgaagca 60

-continued

```

gcagccactg aaaagcagaa aacaaagggg aagaacaagc ttagccatgc tgagtattat 120
attcaaggaa ttcgtcaagt acaatagaca cgtaatcaa accatgcaga tacgaacctg 180
ccactccatc accacccaaa cccttcacac acttccgtgt tcttctagaa aagcccacca 240
ccgccacctt cttccgccgt taaacgctgc ggtttccgcy ggcgccgtca aagcccggaa 300
ggcccactca atgcctccag aaaagaaaga aattttcaag tccttgaggg gatgggcctc 360
ggagtggggtc ctaccgctgc tgaagcccgt ggagcaatgc tggcagccac aaaacttctc 420
ccctgacccc tcccttccgc atgaagagtt cagccatcag gtgaaggagc ttcgcgaaag 480
cactaaagag ttacctgatg agtactttgt ggtgctggtg ggtgatatgg tcaccgagga 540
cgcgcttccc acttaccaga ccatgatcaa caaccttgat ggagtgaag atgacagcgg 600
cacgagcccc agcccgtggg ccgtgtggac cggggcctgg accccgagg aaaacagaca 660
cggggatctg ctcaaacctt atttgtatct ctctgggagg gttgacatgg ctaaggtcga 720
aaagaccgta cattacctca tttcagctgg catggacct gggacagaca acaaccata 780
tttggggtt gtgtacagct cattccaaga gcgagcaaca tttgtggcgc acgggaacac 840
ggctcggctc gcgaaggagg gcggggatcc agtgctggcg cgcctatgcy ggaccatcgc 900
agcggacgag aagcggcacg agaacgcgta ctcaagaatc gtggagaagc ttctggaagt 960
ggaccccacc ggggcaatgg tggccatagg gaacatgatg gagaagaaga tcaccgatgc 1020
ggcgcacctt atgtacgatg gggatgaccc caggctattc gagcactact ccgctgtggc 1080
gcagcgcata ggcggtgata ccgccaacga ctacgcagac atcttgaggt ttctcgttga 1140
acggtgagga ttggagaagc ttgaaggatt gatggctgag gggaaagcggg cgcaggattt 1200
cgtgtgtggg ttggcgccga ggattaggag gttgcaagaa cgcgctgatg agcagagcgcg 1260
taagatgaag aagcatcatg gcgttaagtt cagttggatt ttcaataaag aattgctttt 1320
gtgaaatttc agttaagact taagagataa gagatagagg tcaacgtgag tcaacagggt 1380
tttgctttg tgactatttt gagtttttgt ttgtaggtgg catttttagt acgaataatg 1440
aacaatttaa catggattgc gtgtaatgga cattgttggg tccatggttg ttgttctggt 1500
ggatacacia ccagtaggac tttttgttg taacgttttg cttgcatatt agcttagctt 1560

```

```

<210> SEQ ID NO 2
<211> LENGTH: 405
<212> TYPE: PRT
<213> ORGANISM: Glycine max

```

```

<400> SEQUENCE: 2

```

```

Met Leu Ser Ile Ile Phe Lys Glu Phe Val Lys Tyr Asn Arg His Val
1           5           10           15
Ile Lys Thr Met Gln Ile Arg Thr Cys His Ser Ile Thr Thr Gln Thr
20           25           30
Leu Pro Gln Leu Pro Cys Ser Ser Arg Lys Ala His His Arg His Leu
35           40           45
Leu Pro Pro Leu Asn Ala Ala Val Ser Ala Ala Pro Phe Lys Ala Arg
50           55           60
Lys Ala His Ser Met Pro Pro Glu Lys Lys Glu Ile Phe Lys Ser Leu
65           70           75           80
Glu Gly Trp Ala Ser Glu Trp Val Leu Pro Leu Leu Lys Pro Val Glu
85           90           95

```

-continued

Gln Cys Trp Gln Pro Gln Asn Phe Leu Pro Asp Pro Ser Leu Pro His
 100 105 110
 Glu Glu Phe Ser His Gln Val Lys Glu Leu Arg Glu Arg Thr Lys Glu
 115 120 125
 Leu Pro Asp Glu Tyr Phe Val Val Leu Val Gly Asp Met Val Thr Glu
 130 135 140
 Asp Ala Leu Pro Thr Tyr Gln Thr Met Ile Asn Asn Leu Asp Gly Val
 145 150 155 160
 Lys Asp Asp Ser Gly Thr Ser Pro Ser Pro Trp Ala Val Trp Thr Arg
 165 170 175
 Ala Trp Thr Ala Glu Glu Asn Arg His Gly Asp Leu Leu Arg Thr Tyr
 180 185 190
 Leu Tyr Leu Ser Gly Arg Val Asp Met Ala Lys Val Glu Lys Thr Val
 195 200 205
 His Tyr Leu Ile Ser Ala Gly Met Asp Pro Gly Thr Asp Asn Asn Pro
 210 215 220
 Tyr Leu Gly Phe Val Tyr Thr Ser Phe Gln Glu Arg Ala Thr Phe Val
 225 230 235 240
 Ala His Gly Asn Thr Ala Arg Leu Ala Lys Glu Gly Gly Asp Pro Val
 245 250 255
 Leu Ala Arg Leu Cys Gly Thr Ile Ala Ala Asp Glu Lys Arg His Glu
 260 265 270
 Asn Ala Tyr Ser Arg Ile Val Glu Lys Leu Leu Glu Val Asp Pro Thr
 275 280 285
 Gly Ala Met Val Ala Ile Gly Asn Met Met Glu Lys Lys Ile Thr Met
 290 295 300
 Pro Ala His Leu Met Tyr Asp Gly Asp Asp Pro Arg Leu Phe Glu His
 305 310 315 320
 Tyr Ser Ala Val Ala Gln Arg Ile Gly Val Tyr Thr Ala Asn Asp Tyr
 325 330 335
 Ala Asp Ile Leu Glu Phe Leu Val Glu Arg Trp Arg Leu Glu Lys Leu
 340 345 350
 Glu Gly Leu Met Ala Glu Gly Lys Arg Ala Gln Asp Phe Val Cys Gly
 355 360 365
 Leu Ala Pro Arg Ile Arg Arg Leu Gln Glu Arg Ala Asp Glu Arg Ala
 370 375 380
 Arg Lys Met Lys Lys His His Gly Val Lys Phe Ser Trp Ile Phe Asn
 385 390 395 400
 Lys Glu Leu Leu Leu
 405

<210> SEQ ID NO 3
 <211> LENGTH: 563
 <212> TYPE: DNA
 <213> ORGANISM: Zea mays
 <220> FEATURE:
 <221> NAME/KEY: unsure
 <222> LOCATION: (308)
 <223> OTHER INFORMATION: n = a, c, g or t
 <220> FEATURE:
 <221> NAME/KEY: unsure
 <222> LOCATION: (458)
 <223> OTHER INFORMATION: n = a, c, g or t
 <220> FEATURE:
 <221> NAME/KEY: unsure
 <222> LOCATION: (483)

-continued

```

<223> OTHER INFORMATION: n = a, c, g or t
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (494)
<223> OTHER INFORMATION: n = a, c, g or t
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (519)
<223> OTHER INFORMATION: n = a, c, g or t
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (521)
<223> OTHER INFORMATION: n = a, c, g or t
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (545)
<223> OTHER INFORMATION: n = a, c, g or t
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (550)
<223> OTHER INFORMATION: n = a, c, g or t
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: (557)
<223> OTHER INFORMATION: n = a, c, g or t

<400> SEQUENCE: 3

agcgacaaa cccgggcacc tcgtctagct cgccttccat ttcgtccctt cctattcata    60
ctaccttcta cgagtttgag cagccatggc ggcaacaaca ccaactgcttg ctgtggctgg    120
acatggagta tctacaaaac cagcaaatgc taaagacagc tactactgct tcaaatttgc    180
atcatcggca agaacaagag tcaccctccc acagatcatc cactggagggt gcaggagcag    240
tcatagcagc acggggacca cgaccatggc cgtccctgtc ctcaagcggc gggagaagca    300
ggacgaanag caggaatgga tgggggtacct ggccccggag aagctggagg tgctagcaca    360
cctggagccg tgggcccagg cgcacgtgct gccgctgctg aagcccgcgg aggagggtgg    420
aacgcgcgga catctccgga ccggcgcgct ggcgacangg ctcacaccgt gccgcaactc    480
gcnccggggg caantgccga cccactgggt gctggtggna natatacgag gaggctgcca    540
gtcanagcgn ccaacgntca ggg                                           563

<210> SEQ ID NO 4
<211> LENGTH: 110
<212> TYPE: PRT
<213> ORGANISM: Zea mays
<220> FEATURE:
<221> NAME/KEY: UNSURE
<222> LOCATION: (75)
<223> OTHER INFORMATION: Xaa = any amino acid

<400> SEQUENCE: 4

Met Ala Ala Thr Thr Pro Leu Leu Ala Val Ala Gly His Gly Val Ser
 1             5             10             15
Tyr Lys Pro Ala Asn Ala Lys Asp Ser Tyr Tyr Cys Phe Lys Phe Ala
 20            25            30
Ser Ser Ala Arg Thr Arg Val Thr Leu Pro Gln Ile Ile His Trp Arg
 35            40            45
Cys Arg Ser Ser His Ser Ser Thr Gly Thr Thr Thr Met Ala Val Pro
 50            55            60
Val Leu Lys Arg Arg Glu Lys Gln Asp Glu Xaa Gln Glu Trp Met Gly
 65            70            75            80
Tyr Leu Ala Pro Glu Lys Leu Glu Val Leu Ala His Leu Glu Pro Trp

```


-continued

85	90	95
Ala Glu Ala His Val Leu Pro Leu Leu Lys Pro Ala Glu Glu		
100	105	110

<210> SEQ ID NO 5
 <211> LENGTH: 880
 <212> TYPE: DNA
 <213> ORGANISM: Zea mays

<400> SEQUENCE: 5

```

cgtcggcaeg agcggcaega gctcgtgccg cgtccactcc acagtcaccc accgcccct      60
cctccagcgt ccggccccta cgccgcgcag ccaacccagc gggcacgatg caggcccacg      120
gcatcgccat ccgcgcccgc gggccgggtg cggcgacgca ggccccgcg cgccgacggc      180
aatgcccgt gtctgcccgc gcggtcggcg cgcccgcgc gcgcccgcg gtgacgcact      240
cgatgccgcc ggagaaggcg gaggtgttcc gctcgtgga gggctgggcg gcgcggtcgc      300
tgctgcccgt gctcaagccc gtggaggagt gctggcagcc ggcggacttc ctcccggact      360
cctcgtccga gatgttcggg cacgaggtcc gcgagctgcg cgcccgcgcc gggggctcc      420
ccgacgagta cttcgtcgtg ctcgtgggcy acatggtcac ggaagaggcg ctgcccacgt      480
accagaccat gatcaacacg ctcgacggcg tccgcgacga gaccggcgcc agcaactgcc      540
cctgggcccgt ctggacgcgc gcctggaccg ccgaggagaa ccgccacggc gacatcctcg      600
gcaagtacat gtacctatcc ggcccgcgcg acatgcgcac ggtcggagaag accgtccagt      660
acctcatcgg ctccggcatg gatcccggaa cggagaacaa cccgtacctg ggcttcgtgt      720
acacgagcct ccaggagcgc gcgacggcgc tctcgcacgg caacaccgcg cggctcccca      780
ggcgcacagg ggacgacttc ttggcgcgcg cctgcgggac caaccgcccg caacaagaaa      840
cgaaacaaaa cgggttaagg ggcacccctc aagaagttgg      880
    
```

<210> SEQ ID NO 6
 <211> LENGTH: 257
 <212> TYPE: PRT
 <213> ORGANISM: Zea mays

<400> SEQUENCE: 6

Met	Gln	Ala	His	Gly	Ile	Ala	Ile	Arg	Ala	Arg	Gly	Pro	Val	Ala	Ala
1			5					10						15	
Thr	Gln	Ala	Pro	Ala	Arg	Arg	Arg	Gln	Cys	Arg	Val	Ser	Ala	Ala	Ala
		20						25					30		
Val	Gly	Ala	Pro	Ala	Ala	Arg	Ala	Arg	Val	Thr	His	Ser	Met	Pro	Pro
		35					40					45			
Glu	Lys	Ala	Glu	Val	Phe	Arg	Ser	Leu	Glu	Gly	Trp	Ala	Ala	Arg	Ser
	50					55					60				
Leu	Leu	Pro	Leu	Leu	Lys	Pro	Val	Glu	Glu	Cys	Trp	Gln	Pro	Ala	Asp
65					70					75					80
Phe	Leu	Pro	Asp	Ser	Ser	Ser	Glu	Met	Phe	Gly	His	Glu	Val	Arg	Glu
			85						90					95	
Leu	Arg	Ala	Arg	Ala	Ala	Gly	Leu	Pro	Asp	Glu	Tyr	Phe	Val	Val	Leu
		100						105						110	
Val	Gly	Asp	Met	Val	Thr	Glu	Glu	Ala	Leu	Pro	Thr	Tyr	Gln	Thr	Met
		115						120					125		
Ile	Asn	Thr	Leu	Asp	Gly	Val	Arg	Asp	Glu	Thr	Gly	Ala	Ser	Asn	Cys

-continued

```

cgaggccgccc tacaccgca tcgtctccag gctcctcgag gccgaccgag acgcccggcgt 300
gcgcgcccgtg gcgcgcgatgc tacggcgagg ggtcccccaa tgccgaactn ggcccatnct 360
ccgacggccg ccgcgacgac ctctaaccgc tgcgtcggtg tcccctccgc cgaagcaggg 420
ccgggacgta nagnngggtc ggantaactg gntcaatccg tcn 463

```

```

<210> SEQ ID NO 8
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Oryza sativa

```

```

<400> SEQUENCE: 8

```

```

Tyr Leu Ser Gly Arg Phe Asp Met Ala Glu Val Glu Arg Ala Val His
1           5           10           15
Arg Leu Ile Arg Ser Gly Met Ala Val Asp Pro Pro Cys Ser Pro Tyr
          20           25           30
His Ala Phe Val Tyr Thr Ala Phe Gln Glu Arg Ala Thr Ala Val Ala
          35           40           45
His Gly Asn Thr Ala Arg Leu Val Gly Ala Arg Gly His Gly Asp Ala
          50           55           60
Ala Leu Ala Arg Val Cys Gly Thr Val Ala Ala Asp Glu Lys Arg His
65           70           75           80
Glu Ala Ala Tyr Thr Arg Ile Val Ser Arg Leu Leu Glu Ala Asp Pro
          85           90           95
Asp Ala Gly Val Arg Ala Val Ala Arg Met Leu Arg Arg Gly Val
          100          105          110

```

```

<210> SEQ ID NO 9
<211> LENGTH: 1483
<212> TYPE: DNA
<213> ORGANISM: Zea mays

```

```

<400> SEQUENCE: 9

```

```

gcacgagagc gaccaaacc gggcacctcg tctagctcgc cttccatttc gtccttcct 60
attcatacta ctttctacga gtttgagcag ccatggcggc aacaacacca ctgcttgctg 120
tggttgga ca tggagtatcc tacaaccag caaatgctaa agacagctac tactgcttca 180
aatttgcatc atcggcaaga acaagagtca cctcccaca gatcatccac tggaggtgca 240
ggagcagtca tagcagcacg gggaccacga ccatggccgt cctgtcctc aagcggcggg 300
agaagcagga cgaagagcag gaatggatgg ggtacctggc cccggagaag ctggagggtc 360
tagcacacct ggagccgtgg gcggaggcgc acgtgctgcc gctgctgaag cccgaggagg 420
aggcgtggca gccgtcggac atgctcccgg acccggcggc gctgggcgac gagggttcc 480
acgacgcgtg ccgcgagctc cgcgcgcccgg cggccagcgt gcccgacgcc cacctggtgt 540
gctggtggg caacatgatc actgaggagg cctgcccac gtaccagagc gtgcctaacc 600
gcttcgaggc cgtgcgcgac ctcacggcg cagactccac gcctggggcg cgtggatcc 660
gcggctggtc cgcgaggag aaccgccacg gcgacgcct cagccaactac atgtacctct 720
cgggcccgtg cgacatgcgc caggtogacc gcaccgtgca ccgctcctc gctcccggca 780
tggccatgaa cgcgccagg agcccctacc acggcttcat ctacgtcgtc ttccaggagc 840
gcgccaccgc catctcgcac ggcaacatgg cgcggcacgt cggcgcgcac ggcgaccacg 900

```

-continued

```

tgctcgcccc cgtatgcggc gccatcatgg ccgacgagaa gcgccacgag accgcataca    960
cccgcacatcgt cgccaagctc ttcgaggctc acccggacgc ggcctgctgc gcgctcggct    1020
acatgatgctc ccaccggatc accatgcccg cagcgtctcat gaccgacggc cgcgacgccc    1080
acctctacgc ccactacgcc gccgcggcgc agcagaccgg cgtgtacact gcgtctgact    1140
accgaagcat cctggagcac ctcatacggc agtggcgcgt ggaggagctc gcggcggggc    1200
tctccggcga ggggaggcgc gcgcgggact acgtgtgcgg gctgccgcac aagatccgga    1260
ggatggagga gaaggcccat gacagggcgg cccagaccca gaaaaagccc acgtctgtcc    1320
cgtttagctg gatcttcgat agatccgtca atgtcgtgat tccgtaattt tcctcaaaaa    1380
aaattgagaa tcaggttatg cttagaggtg cattcactgt tgtgtggatt atccttgcaa    1440
taaaaaaaca acgccttgcg ggtgaaaaaa aaaaaaaaaa aaa                        1483

```

<210> SEQ ID NO 10

<211> LENGTH: 424

<212> TYPE: PRT

<213> ORGANISM: Zea mays

<400> SEQUENCE: 10

```

Met Ala Ala Thr Thr Pro Leu Leu Ala Val Ala Gly His Gly Val Ser
1          5          10          15
Tyr Lys Pro Ala Asn Ala Lys Asp Ser Tyr Tyr Cys Phe Lys Phe Ala
20        25        30
Ser Ser Ala Arg Thr Arg Val Thr Leu Pro Gln Ile Ile His Trp Arg
35        40        45
Cys Arg Ser Ser His Ser Ser Thr Gly Thr Thr Thr Met Ala Val Pro
50        55        60
Val Leu Lys Arg Arg Glu Lys Gln Asp Glu Glu Gln Glu Trp Met Gly
65        70        75        80
Tyr Leu Ala Pro Glu Lys Leu Glu Val Leu Ala His Leu Glu Pro Trp
85        90        95
Ala Glu Ala His Val Leu Pro Leu Leu Lys Pro Ala Glu Glu Ala Trp
100       105       110
Gln Pro Ser Asp Met Leu Pro Asp Pro Ala Ala Leu Gly Asp Glu Gly
115       120       125
Phe His Asp Ala Cys Arg Glu Leu Arg Ala Arg Ala Ala Ser Val Pro
130       135       140
Asp Ala His Leu Val Cys Leu Val Gly Asn Met Ile Thr Glu Glu Ala
145       150       155       160
Leu Pro Thr Tyr Gln Ser Val Pro Asn Arg Phe Glu Ala Val Arg Asp
165       170       175
Leu Thr Gly Ala Asp Ser Thr Ala Trp Ala Arg Trp Ile Arg Gly Trp
180       185       190
Ser Ala Glu Glu Asn Arg His Gly Asp Ala Leu Ser His Tyr Met Tyr
195       200       205
Leu Ser Gly Arg Val Asp Met Arg Gln Val Asp Arg Thr Val His Arg
210       215       220
Leu Ile Ala Ser Gly Met Ala Met Asn Ala Ala Arg Ser Pro Tyr His
225       230       235       240
Gly Phe Ile Tyr Val Ala Phe Gln Glu Arg Ala Thr Ala Ile Ser His
245       250       255

```


-continued

```

ctggagacac tggagagcgg gctctccggc gagggccgca gggccagggg cttcgtctgc 1140
gggctcgcgc cgaggatgcg cggggccgcg gagcgcgccc aggacagggc caagaaggac 1200
gagcccagga tggtaagatt cagctggatc tttgataggg aagccgttgt ttaggcactt 1260
gttgtaact gtgatatgtg ctcagatcat gtcgctagct gtcagtgtct ttgtcacatt 1320
gtgtttatgt gtttgaatg ccgtaagagt gtttttttcc tgctattatc acaaaattct 1380
gcagaaatat atgttctaaa aaaaaaaaaa aaaaa 1415

```

<210> SEQ ID NO 12

<211> LENGTH: 380

<212> TYPE: PRT

<213> ORGANISM: Zea mays

<400> SEQUENCE: 12

```

Met Gln Ala His Gly Ile Ala Ile Arg Ala Arg Gly Pro Val Ala Ala
 1           5           10          15
Thr Gln Ala Pro Ala Arg Arg Arg Gln Cys Arg Val Ser Ala Ala Ala
          20          25          30
Val Gly Ala Pro Ala Ala Arg Ala Arg Val Thr His Ser Met Pro Pro
          35          40          45
Glu Lys Ala Glu Val Phe Arg Ser Leu Glu Gly Trp Ala Ala Arg Ser
          50          55          60
Leu Leu Pro Leu Leu Lys Pro Val Glu Glu Cys Trp Gln Pro Ala Asp
65          70          75          80
Phe Leu Pro Asp Ser Ser Ser Glu Met Phe Gly His Glu Val Arg Glu
          85          90          95
Leu Arg Ala Arg Ala Ala Gly Leu Pro Asp Glu Tyr Phe Val Val Leu
          100         105         110
Val Gly Asp Met Val Thr Glu Glu Ala Leu Pro Thr Tyr Gln Thr Met
          115         120         125
Ile Asn Thr Leu Asp Gly Val Arg Asp Glu Thr Gly Ala Ser Asn Cys
          130         135         140
Pro Trp Ala Val Trp Thr Arg Ala Trp Thr Ala Glu Glu Asn Arg His
145         150         155         160
Gly Asp Ile Leu Gly Lys Tyr Met Tyr Leu Ser Gly Arg Val Asp Met
          165         170         175
Arg Met Val Glu Lys Thr Val Gln Tyr Leu Ile Gly Ser Gly Met Asp
          180         185         190
Pro Gly Thr Glu Asn Asn Pro Tyr Leu Gly Phe Val Tyr Thr Ser Phe
          195         200         205
Gln Glu Arg Ala Thr Ala Val Ser His Gly Asn Thr Ala Arg Leu Ala
          210         215         220
Arg Ala His Gly Asp Asp Val Leu Ala Arg Ala Cys Gly Thr Ile Ala
225         230         235         240
Ala Asp Glu Lys Arg His Glu Thr Ala Tyr Gly Arg Ile Val Glu Gln
          245         250         255
Leu Leu Gln Leu Asp Pro Glu Gly Ala Val Leu Ala Val Ala Asp Met
          260         265         270
Met Arg Lys Arg Ile Thr Met Pro Ala His Leu Met His Asp Gly Arg
          275         280         285
Asp Met Asp Leu Phe Glu His Phe Ala Ala Val Ala Gln Arg Leu Gly

```


-continued

Asp Ala Gly Val Arg Ala Val Ala Arg Met Leu Arg Arg Gly Val Ala
 100 105 110

Met Pro Thr Ser Pro Ile Ser Asp Gly Arg Arg Asp Asp Leu Tyr Ala
 115 120 125

Cys Val Val Ser Leu Ala Glu Gln Ala Gly Thr Tyr Thr Val Ser Asp
 130 135 140

Tyr Cys Ser Ile Val Glu His Leu Val Arg Glu Trp Arg Val Glu Glu
 145 150 155 160

Leu Ala Ala Gly Leu Ser Gly Glu Gly Arg Ala Arg Asp Tyr Val
 165 170 175

Cys Glu Leu Pro Gln Lys Ile Arg Arg Met Lys Glu Lys Ala His Glu
 180 185 190

Arg Ala Val Lys Ala Gln Lys Lys Pro Ile Ser Ile Pro Ile Asn Trp
 195 200 205

Ile Phe Asp Arg His Val Ser Val Met Leu Pro
 210 215

<210> SEQ ID NO 15
 <211> LENGTH: 1318
 <212> TYPE: DNA
 <213> ORGANISM: *Oryza sativa*

<400> SEQUENCE: 15

```

gcacgagaac tagctactgt agttgactga cagtgatagt ggcagtcacg caggtcgtgg      60
gaaccgtgcg tgtcagtggc tgcgcgcgcg tggtagcgcc ctccgcccgg cagtgcgcgc      120
tgtccgcgcg ggtgctgacg gcccgggaga cggcgacggc gacgcggcgc cgcgtgacgc      180
actcgatgcc gccggagaag gcggaggtgt tccggtcgtc ggaaggggtg gcgaggtcgt      240
cgctgctgcc gctgctcaag cccgtggagg agtgctggca gccgacggac ttctgcccgg      300
actcgtcgtc ggagatgttc gagcaccagg tccacgagct ccgcccgcgc gccgccccgc      360
tccccgacga gtacttctgc gtgctggtcg gggacatgat taccgaggag gcgctgccga      420
cgtaccagac catgatcaac acgctcgacg gcgtccgca cgagaccggc gccagccct      480
gcccctgggc cgtctggacg cgcacctgga ccgcccagga gaaccgccac ggcgacatcc      540
tcggcaagta catgtacctc tccggcccgc tcgacatgcg catggtcgag aagaccgtcc      600
agtacctcat cggctccgca atggatccgg ggacggagaa caaccctac ctggggttcg      660
tgtacaccag cttccaggag cgcgcgacgg ccgtgtcgca cgggaacacg gcgcccctcg      720
ccagggcgca cggggacgac gtccctggcg gcacctgccc caccatcgcc gccgacgaga      780
agcggcacga gacggcgtae gggcgcatcg tggagcagct gctgcccgtc gacccggacg      840
gcgccatgct cgcctcgccc gacatgatgc acaagcggat caccatgccc gcgcacctca      900
tgcacgacgg ccgcgacatg aacctgttcg accacttcgc ccgctggggc cagcgcctca      960
acgtctacac cgcgcccgac tacgcccaca tcgtcgagtt cctcgtcaag cggtaggaagc     1020
tgagaccctt ggagactggg ctctccggcg agggccggag ggcccgggac ttctgtgctg     1080
ggctcgcgaa gaggatgccc cgggcccggc agcgggctga ggacagggct aagaaggatg     1140
agcagaggaa ggtcaagttc agctggatct atgataggga agtgattgct tagtttaact     1200
tgtcttggtt gaattctgaa ttcccagtc tagatgatca tgccatttcg ttatcatctc     1260
tgttcttggt ttctctttgc aatgcagtaa attgtaata aaaaaaaaaa aaaaaaaaaa     1318
    
```


-continued

```

<210> SEQ ID NO 16
<211> LENGTH: 381
<212> TYPE: PRT
<213> ORGANISM: Oryza sativa

<400> SEQUENCE: 16

Met  Gln  Val  Val  Gly  Thr  Val  Arg  Val  Ser  Gly  Cys  Gly  Ala  Val  Val
 1      5      10     15

Ala  Pro  Ser  Arg  Arg  Gln  Cys  Arg  Val  Ser  Ala  Ala  Val  Leu  Thr  Ala
 20     25     30

Ala  Glu  Thr  Ala  Thr  Ala  Thr  Arg  Arg  Arg  Val  Thr  His  Ser  Met  Pro
 35     40     45

Pro  Glu  Lys  Ala  Glu  Val  Phe  Arg  Ser  Leu  Glu  Gly  Trp  Ala  Arg  Ser
 50     55     60

Ser  Leu  Leu  Pro  Leu  Leu  Lys  Pro  Val  Glu  Glu  Cys  Trp  Gln  Pro  Thr
 65     70     75     80

Asp  Phe  Leu  Pro  Asp  Ser  Ser  Ser  Glu  Met  Phe  Glu  His  Gln  Val  His
 85     90     95

Glu  Leu  Arg  Ala  Arg  Ala  Ala  Gly  Leu  Pro  Asp  Glu  Tyr  Phe  Val  Val
 100    105    110

Leu  Val  Gly  Asp  Met  Ile  Thr  Glu  Glu  Ala  Leu  Pro  Thr  Tyr  Gln  Thr
 115    120    125

Met  Ile  Asn  Thr  Leu  Asp  Gly  Val  Arg  Asp  Glu  Thr  Gly  Ala  Ser  Ala
 130    135    140

Cys  Pro  Trp  Ala  Val  Trp  Thr  Arg  Thr  Trp  Thr  Ala  Glu  Glu  Asn  Arg
 145    150    155    160

His  Gly  Asp  Ile  Leu  Gly  Lys  Tyr  Met  Tyr  Leu  Ser  Gly  Arg  Val  Asp
 165    170    175

Met  Arg  Met  Val  Glu  Lys  Thr  Val  Gln  Tyr  Leu  Ile  Gly  Ser  Gly  Met
 180    185    190

Asp  Pro  Gly  Thr  Glu  Asn  Asn  Pro  Tyr  Leu  Gly  Phe  Val  Tyr  Thr  Ser
 195    200    205

Phe  Gln  Glu  Arg  Ala  Thr  Ala  Val  Ser  His  Gly  Asn  Thr  Ala  Arg  Leu
 210    215    220

Ala  Arg  Ala  His  Gly  Asp  Asp  Val  Leu  Ala  Arg  Thr  Cys  Gly  Thr  Ile
 225    230    235    240

Ala  Ala  Asp  Glu  Lys  Arg  His  Glu  Thr  Ala  Tyr  Gly  Arg  Ile  Val  Glu
 245    250    255

Gln  Leu  Leu  Arg  Leu  Asp  Pro  Asp  Gly  Ala  Met  Leu  Ala  Ile  Ala  Asp
 260    265    270

Met  Met  His  Lys  Arg  Ile  Thr  Met  Pro  Ala  His  Leu  Met  His  Asp  Gly
 275    280    285

Arg  Asp  Met  Asn  Leu  Phe  Asp  His  Phe  Ala  Ala  Val  Ala  Gln  Arg  Leu
 290    295    300

Asn  Val  Tyr  Thr  Ala  Arg  Asp  Tyr  Ala  Asp  Ile  Val  Glu  Phe  Leu  Val
 305    310    315    320

Lys  Arg  Trp  Lys  Leu  Glu  Thr  Leu  Glu  Thr  Gly  Leu  Ser  Gly  Glu  Gly
 325    330    335

Arg  Arg  Ala  Arg  Asp  Phe  Val  Cys  Gly  Leu  Ala  Lys  Arg  Met  Arg  Arg
 340    345    350

Ala  Ala  Glu  Arg  Ala  Glu  Asp  Arg  Ala  Lys  Lys  Asp  Glu  Gln  Arg  Lys

```

-continued

355	360	365
Val Lys Phe Ser Trp Ile Tyr Asp Arg Glu Val Ile Val		
370	375	380

<210> SEQ ID NO 17
 <211> LENGTH: 384
 <212> TYPE: PRT
 <213> ORGANISM: *Lupinus luteus*

<400> SEQUENCE: 17

Met Gln Ile Gln Thr Cys Tyr Ser Ile Arg Ile Gln Ile Leu Pro Leu		
1	5	10
Pro Trp Ala Arg Arg Thr Gly Arg His Lys Met Leu Pro Pro Ile Ala		
20	25	30
Ala Ile Ser Ala Thr Pro Pro Ser Leu Lys Ser Pro Lys Thr His Ser		
35	40	45
Met Pro Pro Glu Lys Ile Glu Ile Phe Lys Ser Leu Glu Ser Trp Ala		
50	55	60
Ser Gln Ser Val Leu Pro Leu Leu Lys Pro Val Glu Gln Cys Trp Gln		
65	70	75
Pro Gln Glu Phe Val Pro Asp Ser Ser Leu Pro Phe Gly Asp Phe Thr		
85	90	95
Asp Gln Val Lys Ala Leu Arg Asp Arg Thr Ala Glu Leu Pro Glu Glu		
100	105	110
Tyr Phe Val Val Leu Val Gly Asp Met Ile Thr Glu Asp Ala Leu Pro		
115	120	125
Thr Tyr Gln Ser Met Ile Asn Asn Leu Asp Gly Val Arg Asp Glu Thr		
130	135	140
Gly Ser Ser Pro Ser Pro Trp Ala Leu Trp Thr Arg Ala Trp Thr Ala		
145	150	155
Glu Glu Lys Arg His Gly Asp Leu Leu Arg Thr Tyr Leu Tyr Leu Ser		
165	170	175
Gly Arg Val Asp Met Lys Lys Ile Glu Lys Thr Val Gln Tyr Leu Ile		
180	185	190
Gly Ser Gly Met Asp Pro Gly Thr Glu Asn Asn Pro Tyr Leu Gly Phe		
195	200	205
Val Tyr Thr Ser Phe Gln Glu Arg Ala Thr Phe Val Ser His Gly Asn		
210	215	220
Thr Ala Arg Leu Ala Lys Glu Gly Gly Asp Pro Val Leu Ala Arg Ile		
225	230	235
Cys Gly Thr Ile Ala Ala Asp Glu Lys Arg His Glu Asn Ala Tyr Ser		
245	250	255
Arg Ile Val Glu Lys Leu Leu Glu Leu Asp Pro Thr Gly Ala Met Val		
260	265	270
Ala Ile Gly Asp Met Met Gln Lys Lys Ile Thr Met Pro Ala His Leu		
275	280	285
Met Tyr Asp Gly Glu Asp Pro Lys Leu Phe Asp His Phe Ser Ala Val		
290	295	300
Ala Gln Arg Met Gly Val Tyr Thr Ala Asn Asp Tyr Ala Asp Ile Leu		
305	310	315
Glu Phe Leu Ile Gly Arg Trp Arg Leu Glu Lys Val Gln Asp Leu Lys		
325	330	335

-continued

Asp Glu Gly Lys Lys Ala Gln Asp Phe Val Cys Gly Leu Ala Pro Arg
 340 345 350

Ile Arg Arg Leu Gln Glu Arg Ala Asp Glu Arg Ala Arg Lys Met Lys
 355 360 365

Pro His Ala Val Lys Phe Ser Trp Ile Phe Asn Lys Glu Ile Ile Leu
 370 375 380

<210> SEQ ID NO 18
 <211> LENGTH: 396
 <212> TYPE: PRT
 <213> ORGANISM: Cucumis sativus

<400> SEQUENCE: 18

Met Ala Leu Lys Phe His Pro Leu Thr Ser Gln Ser Pro Lys Leu Pro
 1 5 10 15

Ser Phe Arg Met Pro Gln Leu Ala Ser Leu Arg Ser Pro Lys Phe Val
 20 25 30

Met Ala Ser Thr Leu Arg Ser Thr Ser Arg Glu Val Glu Thr Leu Lys
 35 40 45

Lys Pro Phe Met Pro Pro Arg Glu Val His Leu Gln Val Thr His Ser
 50 55 60

Met Pro Pro Gln Lys Met Glu Ile Phe Lys Ser Leu Glu Asp Trp Ala
 65 70 75 80

Glu Glu Asn Leu Leu Val His Leu Lys Pro Val Glu Arg Cys Trp Gln
 85 90 95

Pro Gln Asp Phe Leu Pro Asp Ser Ala Phe Glu Gly Phe His Glu Gln
 100 105 110

Val Arg Glu Leu Arg Glu Arg Ala Lys Glu Leu Pro Asp Glu Tyr Phe
 115 120 125

Val Val Leu Val Gly Asp Met Ile Thr Glu Glu Ala Leu Pro Thr Tyr
 130 135 140

Gln Thr Met Leu Asn Thr Leu Asp Gly Val Arg Asp Glu Thr Gly Ala
 145 150 155 160

Ser Pro Thr Pro Trp Ala Ile Trp Thr Arg Ala Trp Thr Ala Glu Glu
 165 170 175

Asn Arg His Gly Asp Leu Leu Asn Lys Tyr Leu Tyr Leu Ser Gly Arg
 180 185 190

Val Asp Met Arg Gln Val Glu Lys Thr Ile Gln Tyr Leu Ile Gly Ser
 195 200 205

Gly Met Asp Pro Arg Thr Glu Asn Asn Pro Tyr Leu Gly Phe Ile Tyr
 210 215 220

Thr Ser Phe Gln Glu Arg Ala Thr Phe Ile Ser His Gly Asn Thr Ala
 225 230 235 240

Arg Leu Ala Lys Glu His Gly Asp Ile Lys Leu Ala Gln Ile Cys Gly
 245 250 255

Thr Ile Thr Ala Asp Glu Lys Arg His Glu Thr Ala Tyr Thr Lys Ile
 260 265 270

Val Glu Lys Leu Phe Glu Ile Asp Pro Glu Gly Thr Val Ile Ala Phe
 275 280 285

Glu Glu Met Met Arg Lys Lys Val Ser Met Pro Ala His Leu Met Tyr
 290 295 300

Asp Gly Arg Asp Asp Asn Leu Phe His His Phe Ser Ala Val Ala Gln
 305 310 315 320

-continued

Arg Leu Gly Val Tyr Thr Ala Lys Asp Tyr Ala Asp Ile Leu Glu Phe
 325 330 335

Leu Val Gly Arg Trp Lys Val Glu Ser Leu Thr Gly Leu Ser Gly Glu
 340 345 350

Gly Gln Lys Ala Gln Asp Tyr Val Cys Ala Leu Pro Ala Arg Ile Arg
 355 360 365

Lys Leu Glu Glu Arg Ala Gln Gly Arg Ala Lys Glu Gly Pro Thr Ile
 370 375 380

Pro Phe Ser Trp Ile Phe Asp Arg Gln Val Lys Leu
 385 390 395

<210> SEQ ID NO 19
 <211> LENGTH: 374
 <212> TYPE: PRT
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 19

Met Pro Ser Pro Ser Thr Phe Leu Ala Ser Arg Pro Arg Gly Pro Ala
 1 5 10 15

Lys Ile Ser Ala Val Ala Ala Pro Val Arg Pro Ala Leu Lys His Gln
 20 25 30

Asn Lys Ile His Thr Met Pro Pro Glu Lys Met Glu Ile Phe Lys Ser
 35 40 45

Leu Asp Gly Trp Ala Lys Asp Gln Ile Leu Pro Leu Leu Lys Pro Val
 50 55 60

Asp Gln Cys Trp Gln Pro Ala Ser Phe Leu Pro Asp Pro Ala Leu Pro
 65 70 75 80

Phe Ser Glu Phe Thr Asp Gln Val Arg Glu Leu Arg Glu Arg Thr Ala
 85 90 95

Ser Leu Pro Asp Glu Tyr Phe Val Val Leu Val Gly Asp Met Ile Thr
 100 105 110

Glu Asp Ala Leu Pro Thr Tyr Gln Thr Met Ile Asn Thr Leu Asp Gly
 115 120 125

Val Arg Asp Glu Thr Gly Ala Ser Glu Ser Ala Trp Ala Ser Trp Thr
 130 135 140

Arg Ala Trp Thr Ala Glu Glu Asn Arg His Gly Asp Leu Leu Arg Thr
 145 150 155 160

Tyr Leu Tyr Leu Ser Gly Arg Val Asp Met Leu Met Val Glu Arg Thr
 165 170 175

Val Gln His Leu Ile Gly Ser Gly Met Asp Pro Gly Thr Glu Asn Asn
 180 185 190

Pro Tyr Leu Gly Phe Val Tyr Thr Ser Phe Gln Glu Arg Ala Thr Phe
 195 200 205

Val Ser His Gly Asn Thr Ala Arg Leu Ala Lys Ser Ala Gly Asp Pro
 210 215 220

Val Leu Ala Arg Ile Cys Gly Thr Ile Ala Ala Asp Glu Lys Arg His
 225 230 235 240

Glu Asn Ala Tyr Val Arg Ile Val Glu Lys Leu Leu Glu Ile Asp Pro
 245 250 255

Asn Gly Ala Val Ser Ala Val Ala Asp Met Met Arg Lys Lys Ile Thr
 260 265 270

Met Pro Ala His Leu Met Thr Asp Gly Arg Asp Pro Met Leu Phe Glu

-continued

Cys Gly Ile Ile Ala Ala Asp Glu Lys Arg His Glu Thr Ala Tyr Thr
 260 265 270
 Lys Ile Val Glu Lys Leu Phe Glu Ile Asp Pro Asp Gly Ala Val Leu
 275 280 285
 Ala Leu Ala Asp Met Met Arg Lys Lys Val Ser Met Pro Ala His Leu
 290 295 300
 Met Tyr Asp Gly Lys Asp Asp Asn Leu Phe Glu Asn Tyr Ser Ala Val
 305 310 315 320
 Ala Gln Gln Ile Gly Val Tyr Thr Ala Lys Asp Tyr Ala Asp Ile Leu
 325 330 335
 Glu His Leu Val Asn Arg Trp Lys Val Glu Asn Leu Met Gly Leu Ser
 340 345 350
 Gly Glu Gly His Lys Ala Gln Asp Phe Val Cys Gly Leu Ala Pro Arg
 355 360 365
 Ile Arg Lys Leu Gly Glu Arg Ala Gln Ser Leu Ser Lys Pro Val Ser
 370 375 380
 Leu Val Pro Phe Ser Trp Ile Phe Asn Lys Glu Leu Lys Val
 385 390 395

<210> SEQ ID NO 21
 <211> LENGTH: 411
 <212> TYPE: PRT
 <213> ORGANISM: Arabidopsis thaliana
 <400> SEQUENCE: 21

Met Ala Leu Leu Leu Asn Ser Thr Ile Thr Val Ala Met Lys Gln Asn
 1 5 10 15
 Pro Leu Val Ala Val Ser Phe Pro Arg Thr Thr Cys Leu Gly Ser Ser
 20 25 30
 Phe Ser Pro Pro Arg Leu Leu Arg Val Ser Cys Val Ala Thr Asn Pro
 35 40 45
 Ser Lys Thr Ser Glu Glu Thr Asp Lys Lys Lys Phe Arg Pro Ile Lys
 50 55 60
 Glu Val Pro Asn Gln Val Thr His Thr Ile Thr Gln Glu Lys Leu Glu
 65 70 75 80
 Ile Phe Lys Ser Met Glu Asn Trp Ala Gln Glu Asn Leu Leu Ser Tyr
 85 90 95
 Leu Lys Pro Val Glu Ala Ser Trp Gln Pro Gln Asp Phe Leu Pro Glu
 100 105 110
 Thr Asn Asp Glu Asp Arg Phe Tyr Glu Gln Val Lys Glu Leu Arg Asp
 115 120 125
 Arg Thr Lys Glu Ile Pro Asp Asp Tyr Phe Val Val Leu Val Gly Asp
 130 135 140
 Met Ile Thr Glu Glu Ala Leu Pro Thr Tyr Gln Thr Thr Leu Asn Thr
 145 150 155 160
 Leu Asp Gly Val Lys Asp Glu Thr Gly Gly Ser Leu Thr Pro Trp Ala
 165 170 175
 Val Trp Val Arg Ala Trp Thr Ala Glu Glu Asn Arg His Gly Asp Leu
 180 185 190
 Leu Asn Lys Tyr Leu Tyr Leu Ser Gly Arg Val Asp Met Arg His Val
 195 200 205
 Glu Lys Thr Ile Gln Tyr Leu Ile Gly Ser Gly Met Asp Ser Lys Phe
 210 215 220

-continued

Glu Asn Asn Pro Tyr Asn Gly Phe Ile Tyr Thr Ser Phe Gln Glu Arg
 225 230 235 240
 Ala Thr Phe Ile Ser His Gly Asn Thr Ala Lys Leu Ala Thr Thr Tyr
 245 250 255
 Gly Asp Thr Thr Leu Ala Lys Ile Cys Gly Thr Ile Ala Ala Asp Glu
 260 265 270
 Lys Arg His Glu Thr Ala Tyr Thr Arg Ile Val Glu Lys Leu Phe Glu
 275 280 285
 Ile Asp Pro Asp Gly Thr Val Gln Ala Leu Ala Ser Met Met Arg Lys
 290 295 300
 Arg Ile Thr Met Pro Ala His Leu Met His Asp Gly Arg Asp Asp Asp
 305 310 315 320
 Leu Phe Asp His Tyr Ala Ala Val Ala Gln Arg Ile Gly Val Tyr Thr
 325 330 335
 Ala Thr Asp Tyr Ala Gly Ile Leu Glu Phe Leu Leu Arg Arg Trp Glu
 340 345 350
 Val Glu Lys Leu Gly Met Gly Leu Ser Gly Glu Gly Arg Arg Ala Gln
 355 360 365
 Asp Tyr Leu Cys Thr Leu Pro Gln Arg Ile Arg Arg Leu Glu Glu Arg
 370 375 380
 Ala Asn Asp Arg Val Lys Leu Ala Ser Lys Ser Lys Pro Ser Val Ser
 385 390 395 400
 Phe Ser Trp Ile Tyr Gly Arg Glu Val Glu Leu
 405 410

<210> SEQ ID NO 22

<211> LENGTH: 396

<212> TYPE: PRT

<213> ORGANISM: *Linum usitatissimum*

<400> SEQUENCE: 22

Met Ala Leu Lys Leu Asn Pro Val Thr Thr Phe Pro Ser Thr Arg Ser
 1 5 10 15
 Leu Asn Asn Phe Ser Ser Arg Ser Pro Arg Thr Phe Leu Met Ala Ala
 20 25 30
 Ser Thr Phe Asn Ser Thr Ser Thr Lys Glu Ala Glu Lys Leu Lys Lys
 35 40 45
 Ser His Gly Pro Pro Lys Glu Val His Met Gln Val Thr His Ser Met
 50 55 60
 Pro Pro Gln Lys Leu Glu Ile Phe Lys Ser Leu Glu Gly Trp Ala Glu
 65 70 75 80
 Asp Val Leu Leu Pro His Leu Lys Pro Val Glu Lys Cys Trp Gln Pro
 85 90 95
 Gln Asp Phe Leu Pro Glu Pro Glu Ser Asp Gly Phe Glu Glu Gln Val
 100 105 110
 Lys Glu Leu Arg Ala Arg Ala Lys Glu Leu Pro Asp Asp Tyr Phe Val
 115 120 125
 Val Leu Val Gly Asp Met Ile Thr Glu Glu Ala Leu Pro Thr Tyr Gln
 130 135 140
 Thr Met Leu Asn Thr Leu Asp Gly Val Arg Asp Glu Thr Gly Ala Ser
 145 150 155 160
 Leu Thr Pro Trp Ala Ile Trp Thr Arg Ala Trp Thr Ala Glu Glu Asn

-continued

Asp Met Ile Thr Glu Glu Ala Leu Pro Thr Tyr Gln Thr Met Leu Asn
 130 135 140
 Thr Leu Asp Gly Val Arg Asp Glu Thr Gly Ala Ser Leu Thr Ser Trp
 145 150 155 160
 Ala Ile Trp Thr Arg Ala Trp Thr Ala Glu Glu Asn Arg His Gly Asp
 165 170 175
 Leu Leu Asn Lys Tyr Leu Tyr Leu Ser Gly Arg Val Asp Met Lys Gln
 180 185 190
 Ile Glu Lys Thr Ile Gln Tyr Leu Ile Gly Ser Gly Met Asp Pro Arg
 195 200 205
 Thr Glu Asn Ser Pro Tyr Leu Gly Phe Ile Tyr Thr Ser Phe Gln Glu
 210 215 220
 Arg Ala Thr Phe Ile Ser His Gly Asn Thr Ala Arg Leu Ala Lys Glu
 225 230 235 240
 His Gly Asp Ile Lys Leu Ala Gln Ile Cys Gly Met Ile Ala Ser Asp
 245 250 255
 Glu Lys Arg His Glu Thr Ala Tyr Thr Lys Ile Val Glu Lys Leu Phe
 260 265 270
 Glu Val Asp Pro Asp Gly Thr Val Met Ala Phe Ala Asp Met Met Arg
 275 280 285
 Lys Lys Ile Ala Met Pro Ala His Leu Met Tyr Asp Gly Arg Asp Asp
 290 295 300
 Asn Leu Phe Asp Asn Tyr Ser Ala Val Ala Gln Arg Ile Gly Val Tyr
 305 310 315 320
 Thr Ala Lys Asp Tyr Ala Asp Ile Leu Glu Phe Leu Val Gly Arg Trp
 325 330 335
 Lys Val Glu Gln Leu Thr Gly Leu Ser Gly Glu Gly Arg Lys Ala Gln
 340 345 350
 Glu Tyr Val Cys Gly Leu Pro Pro Arg Ile Arg Arg Leu Glu Glu Arg
 355 360 365
 Ala Gln Ala Arg Gly Lys Glu Ser Ser Thr Leu Lys Phe Ser Trp Ile
 370 375 380
 His Asp Arg Glu Val Leu Leu
 385 390

<210> SEQ ID NO 24

<211> LENGTH: 80

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: ELVISLIVES
complementary region of pKS106 and pKS124

<400> SEQUENCE: 24

cggccgggagc tggatcatctc gctcatcgtc gaggcggcgg ccgccgactc gacgatgagc 60

gagatgacca gctccggcgg 80

<210> SEQ ID NO 25

<211> LENGTH: 154

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: ELVISLIVES
complementary region of pKS133

<400> SEQUENCE: 25

-continued

```

cggccgggagc tggatcatctc gctcatcgtc gagtcggcgg ccggagctgg tcatctcgct    60
catcgctcgag tcggcggccg ccgactcgac gatgagcgag atgaccagct ccggccgccc    120
actcgacgat gagcgagatg accagctccg gccg                                     154

```

```

<210> SEQ ID NO 26
<211> LENGTH: 6611
<212> TYPE: DNA
<213> ORGANISM: Plasmid pBS68
<220> FEATURE:
<221> NAME/KEY: Unsure
<222> LOCATION: (4436)..(4436)
<223> OTHER INFORMATION: n = A, C, G, or T

```

```

<400> SEQUENCE: 26

```

```

cgcgccatag cgggaccatc gcagcggacg agaagcggca cgagaacgcg tactcaagaa    60
tcgtggagaa gcttctgtaa gtggacccca ccggggcaat ggtggccata gggaaacatga    120
tggagaagaa gatcacgatg ccggcgcacc ttatgtacga tggggatgac cccaggctat    180
tcgagcacta ctccgctgtg gcgcagcgca taggcgtgta caccgccaac gactacgcag    240
acatcttgga tttctcgttg acggtgaaga ttggagaagc ttgaaggatt gatgcctgag    300
gggaagcggg ccccaggatt tccgtgtgtg gggtgcccc gaggattagg aggttccaag    360
aacgcgctga tgagcgcgag cgtaagatga agaagcatca tgccgttaag ttcagttgga    420
ttttcaataa agaattgctt ttgtgagcgg ccgccgactc gacgatgagc gagatgacca    480
gctccggccc ccgactcgac gatgagcgag atgaccagct ccggccgcca cacaagtgtg    540
agagtactaa ataaatgctt tggttgtacg aaatcattac actaaataaa ataatacaag    600
cttatatatg ccttccgcta aggccgaatg caaagaaatt ggttctttct cgttatcttt    660
tgccactttt actagtagct attaattact acttaatcat ctttgtttac ggctcattat    720
atccgctgac ggcgccccc atcatccgga tatagttect cctttcagca aaaaaccct    780
caagaccctg ttagaggccc caaggggta tgctagtatt tgctcagcgg tggcagcagc    840
caactcagct tcctttcggg ctttgttagc agccggatcg atccaagctg tacctcacta    900
ttcctttgcc ctccgacgag tgctggggcg tcggtttcca ctatccgga gtacttctac    960
acagccatcg gtccagacgg ccgcgcttct gcggggcatt tgtgtacgcc cgacagtccc    1020
ggctccggat cggacgattg cgtcgcacgc accctgcgcc caagctgcat catcgaaatt    1080
gccgtcaacc aagctctgat agagttggtc aagaccaatg cggagcatat acgcccggag    1140
ccggcggcat cctgcaagct ccggatgcct ccgctcgaag tagcgcgtct gctgctccat    1200
acaagccaac cacggcctcc agaagaagat gttggcgacc tcgtattggg aatccccgaa    1260
catcgctcgc ctccagctca tgaccgctgt tatgcccga ttgtccgtea ggacattggt    1320
ggagccgaaa tccgcgtgca cgaggtgccg gacttcgggg cagtccctcg cccaaagcat    1380
cagctcatcg agagcctcgc cgacggacgc actgacggtg tcgtocatca cagtttgcca    1440
gtgatacaca tggggatcag caatcgcgca tatgaaatca cgccatgtag tgtattgacc    1500
gattccttgc ggtccgaatg ggcccgaacc gctcgtcttg ctaagatcgg ccgcagcagat    1560
cgcatccata gcctccgcca ccggctgcag aacagcgggc agttcggttt caggcaggtc    1620
ttgcaacgty acaccctgtg cacggcggga gatgcaatag gtcaggctct cgetgaattc    1680
cccaatgtca agcaattccg gaatcgggag cggggccgat gcaaagtgcc gataaacata    1740

```

-continued

acgatctttg tagaaacat cggcgcagct atttaccgc aggacatatc cagccctcc	1800
tacatcgaag ctgaaagcac gagattcttc gccctccgag agctgcatca ggtcggagac	1860
gctgtcgaac ttttcgatca gaaacttctc gacagacgtc gcggtgagtt caggcttttc	1920
catgggtata tctccttctt aaagttaaac aaaattattt ctagagggaa accgttgtgg	1980
tctccctata gtgagtcgta ttaatttcgc gggatcgaga tctgatcaac ctgcattaat	2040
gaatcggcca acgcgcgggg agagcggtt tgcgtattgg gcgctcttcc gcttcctcgc	2100
tcactgactc gctgcctcg gtcgttcggc tgcggcagc ggatcagct cactcaaagg	2160
cggtaatacg gttatccaca gaatcagggg ataacgcagg aaagaacatg tgagcaaaag	2220
gccagcaaaa gcccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc cataggctcc	2280
gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga aaccgcacag	2340
gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct cctgttccga	2400
ccctgcgct taccgatac ctgtccgct ttctccctc ggaagcgtg gcgctttctc	2460
aatgtcaecg ctgtaggtat ctccagttcgg tgtaggtcgt tcgctccaag ctgggctgtg	2520
tgcacgaacc ccccgctcag ccgcaccgct gcgccttacc cggtaactat cgtcttgagt	2580
ccaaccgggt aagacacgac ttatcgccac tggcagcagc cactggtaac aggattagca	2640
gagcgaggta ttagcgggt gctacagagt tcttgaagtg gtggcctaac tacggctaca	2700
ctagaaggac agtatttggt atctgcgctc tgctgaagcc agttaccctc ggaaaaagag	2760
ttggtagctc ttgatccgce aaacaaacca ccgctggtag cggtggtttt tttgttgca	2820
agcagcagat tacgcgcaga aaaaaggat ctcaagaaga tcctttgatc ttttctacgg	2880
ggtctgacgc tcagtggaac gaaaactcac gttaagggtat tttggtcatg acattaacct	2940
ataaaaatag gcgtatcacg aggcccttcc gtctcgcgcg ttcgggtgat gacggtgaaa	3000
acctctgaca catgcagctc ccggagacgg tcacagcttg tctgtaagcg gatgccggga	3060
gcagacaagc ccgtcagggc gcgtcagcgg gtgttggcgg gtgtcggggc tggcttaact	3120
atcggcctac agagcagatt gtactgagag tgcaccatat ggacatattg tcgttagaac	3180
gcggctacaa ttaatacata acctatgta tcatacacat acgatttagg tgacactata	3240
gaacggcgcg ccaagcttgg atcctcgaag agaagggtta ataacacatt ttttaacatt	3300
tttaacacaa attttagtta tttaaaaatt tattaaaaaa tttaaaataa gaagaggaac	3360
tctttaaata aatctaactt acaaaattta tgatttttaa taagttttca ccaataaaaa	3420
atgtcataaa aatattgtaa aaagtatatt atcaatattc tctttatgat aataaaaaag	3480
aaaaaaaaaa taaaagttaa gtgaaaatga gattgaagtg actttaggtg tgtataaata	3540
tatcaacccc gccacaatt tatttaatcc aaatatattg aagtatatta ttccatagcc	3600
tttatttatt tatataatata ttatataaaa gctttatttg ttctaggttg ttcatgaaat	3660
atttttttgg ttttatctcc gttgtaagaa aatcatgtgc tttgtgtcgc cactcactat	3720
tgcagctttt tcatgcatg gtcagattga cggttgattg tatttttgggt ttttatgggt	3780
ttgtgttatg acttaagtct tcatctcttt atctcttcat caggtttgat ggttacctaa	3840
tatgggccat gggatcatgc atgggttaaat taggtggcca actttgttgt gaacgataga	3900
atttttttta tattaagtaa actattttta tattatgaaa taataataaa aaaaatattt	3960
tatcattatt aacaaaatca tattagttaa tttgttaact ctataataaa agaataactg	4020

-continued

taacattcac attacatggt aacatcttcc caccctttca tttgtttttt gtttgatgac 4080
tttttttctt gtttaaatth atttcccttc ttttaaatth ggaatacatt atcatcatat 4140
ataaactaaa atactaaaa caggattaca caaatgataa ataataacac aaatatttat 4200
aaatctagct gcaatatatt taaactagct atatcgatat tgtaaaataa aactagctgc 4260
attgatactg ataaaaaaat atcatgtgct ttctggactg atgatgcagt atacttttga 4320
cattgccttt attttatttt tcagaaaagc tttcttagtt ctgggttctt cattatttgt 4380
ttcccatctc cattgtgaat tgaatcattt gcttcgtgtc acaaatataa tttagntagg 4440
tacatgcatt ggtcagattc acggtttatt atgtcatgac ttaagttcat ggtagtacat 4500
tacctgccac gcatgcatta tattggttag atttgatagg caaatgtgtt tgtaacaat 4560
ataaatataa ataattgttt tatattacga aataacagtg atcaaaataa acagttttat 4620
ctttattaac aagattttgt tttgttttga tgacgttttt taatgtttac gctttccccc 4680
ttcttttgaa tttagaacac tttatcatca taaaatcaaa tactaaaaaa attacatatt 4740
tcataaataa taacacaaat atttttaaaa aatctgaaat aataatgaac aatattacat 4800
attatcacga aaattcatta ataaaaatat tatataaata aaatgtaata gtagttatat 4860
gtaggaaaaa agtactgcac gcataatata tacaaaaaga ttaaatgaa ctattataaa 4920
taataacact aaattaatgg tgaatcatat caaaataatg aaaaagtaa taaaatttgt 4980
aattaacttc tatatgtatt acacacacaa ataataaata atagtaaaaa aaattatgat 5040
aaatatttac catctcataa gatatttaaa ataatgataa aaatagat tattttttat 5100
gcaactagct agccaaaaag agaacacggg tatatataaa aagagtacct ttaatttcta 5160
ctgtacttcc tttattctg acgtttttat atcaagtgga catactgaa gatttttaatt 5220
atcagtctaa atatttcatt agcacttaat acttttctgt tttattccta tcctataagt 5280
agtcccatt ctcccaaat tgcttattca cacaactaac taagaaagtc ttccatagcc 5340
ccccaaagg ccggagctgg tcatctgct catcgtcgag tcggcggccg gagctggta 5400
tctcgtcat cgtcagctg gcggcogctg agtgattgct cagcagtggt gtcaccatgc 5460
cttcagcaag taccatggg ttgatgatgt tgtgggtttg acccttcaact caacactttt 5520
agtcccttat ttctcatgga aaataagcca tcgccgccat cactccaaca caggttcctt 5580
tgaccgtgat gaagtgtttg tcccaaaacc aaaatccaaa gttgcatggt tttccaagta 5640
cttaaacac cctctaggaa gggctgtttc tctctcgtc aactcaca taggggtggc 5700
tatgtattta gccttcaatg tctctggtag accctatgat agttttgcaa gccactacca 5760
cccttatgct ccatatatt ctaacgtga gagcttctg atctatgtct ctgatgttgc 5820
tttgtttct gtgacttact ctctctacg tgttgcaacc ctgaaagggt tggtttggt 5880
gctatgtgtt tatgggtgct ctttctcat tgtgaacggt tttcttga ctatcacata 5940
tttgagcac acacactttg ccttgctca ttacgattca tcagaatggg actggctgaa 6000
gggagctttg gcaactatgg acagagatta agcggccgca tgcctccaga aaagaaagaa 6060
atthtcaagt ccttgagggt atgggctcgt gagtgggtcc taccgtgct gaagccctg 6120
gagcaatgct ggcagccaca aaacttctc cctgacctt cccttccgca tgaaggttc 6180
agccatcagg tgaaggagct tcgcaacgc actaaagagt tacctgatga gtactttgtg 6240
gtgtgtgtg gtgataggt caccgaggac gccttccca cttaccagac catgatcaac 6300

-continued

aaccttgatg gagtgaaga tgacagcggc acgagcccga gcccgtaggc cgtgtggacc	6360
cgggcctgga ccgcccagga aaacagacac ggggatctgc tcagaactta tttgatctc	6420
tctgggaggg ttgacatggc taaggtcgaa aagaccgtac attacctcat ttcagctggc	6480
atggaccctg ggacagacaa caaccatat ttggggtttg tgtacacgtc attccaagag	6540
cgagcaacat ttgtggcgca cgggaacacg gctcggctcg cgaaggaggg cggggatcca	6600
gtgctggcgc g	6611

- 1.** An isolated polynucleotide comprising:
- a nucleotide sequence encoding a polypeptide having delta-9 fatty acid desaturase activity that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NO:2, 4, 6, 8, 10, 12, 14, or 16; or
 - the complement of (a).
- 2-5.** (canceled)
- 6.** A chimeric construct comprising the isolated polynucleotide of claim **1** operably linked to at least one suitable regulatory sequence.
- 7-9.** (canceled)
- 10.** A method of obtaining a nucleic acid fragment encoding a delta-9 fatty acid desaturase polypeptide comprising the steps of:
- probing a cDNA or genomic library with an isolated polynucleotide comprising at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1 and/or a complement of the nucleotide sequence;
 - identifying a DNA clone that hybridizes with the isolated polynucleotide;
 - isolating the identified DNA clone; and
 - sequencing a cDNA or genomic fragment that comprises the isolated DNA clone.
- 11.** A method of obtaining a nucleic acid fragment encoding a delta-9 fatty acid desaturase polypeptide comprising the steps of:
- probing a cDNA or genomic library with an isolated polynucleotide comprising at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NO:1, 3, 5, 7, 9, 11, 13, or 15 and/or a complement of such nucleotide sequences;
 - identifying a DNA clone that hybridizes with the isolated polynucleotide;
 - isolating the identified DNA clone;
 - inserting the DNA clone into a plasmid suitable for expression in a bacterial or yeast cell; and
 - assaying for delta-9 desaturase activity or alterations in fatty acid composition of the host cell.
- 12.** A method of identifying an isolated polynucleotide that encodes a delta-9 fatty acid desaturase comprising the steps of:
- determining an amino acid sequence of the polypeptide encoded by the isolated DNA;
 - determining if the amino acid sequence comprises at least two amino acid sequences selected from the group consisting of HSMPEK corresponding to amino acids 67-73 of SEQ ID NO:2, LPLKPE corresponding to amino acids 89-96 of SEQ ID NO:2, EYFVVLVGD corresponding to amino acids 132-141 of SEQ ID NO:2, EKT corresponding to amino acids 205-208 of SEQ ID NO:2, GMDPGT corresponding to amino acids 215-220 of SEQ ID NO:2, NNPYLGFVYTSFQERAT corresponding to amino acids 222-238 of SEQ ID NO:2, VLAR corresponding to amino acids 256-259 of SEQ ID NO:2, RIVE corresponding to amino acids 277-280 of SEQ ID NO:2, ITMPAHL corresponding to amino acids 302-308 of SEQ ID NO:2, or DFVCGLA corresponding to amino acids 364-370 of SEQ ID NO:2.
- 13.** A method of identifying an isolated polynucleotide that encodes a delta-9 fatty acid desaturase comprising the steps of:
- determining the polypeptide sequence of claim **10**, **11**, or **12**;
 - determining that the amino acid sequence of the polypeptide does not contain at least one of the following amino acid sequences KEIPDDYFWLVGDMITEE-ALPTYQTMINT corresponding to positions 116-145 of SEQ ID NO:23; or DYADILEFLVGRWK corresponding to positions 324-337 of SEQ ID NO:23.
- 14.** A method of altering the level of expression of a delta-9 fatty acid desaturase in a host cell comprising:
- transforming a host cell with the chimeric gene of claim **6**; and
 - growing the transformed host cell produced in step (a) under conditions that are suitable for expression of the chimeric construct wherein expression of the chimeric construct results in production of altered levels of a delta-9 fatty acid desaturase in the transformed host cell.
- 15.** A method of altering the level of expression of a delta-9 fatty acid desaturase in a host cell comprising:
- transforming a host cell with the chimeric construct of claim **6**; and
 - growing the transformed host cell produced in step (a) under conditions that are suitable for expression of the chimeric construct wherein expression of the chimeric gene results in production of altered levels of a delta-9 fatty acid desaturase in the transformed host cell.

* * * * *