

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 March 2008 (06.03.2008)

PCT

(10) International Publication Number
WO 2008/025784 A1

(51) International Patent Classification:

A23J 1/00 (2006.01) A23J 1/02 (2006.01)
A23J 1/20 (2006.01) A61K 9/127 (2006.01)
A23J 1/06 (2006.01) A23L 1/00 (2006.01)

(21) International Application Number:

PCT/EP2007/058964

(22) International Filing Date: 29 August 2007 (29.08.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

06018271.4 31 August 2006 (31.08.2006) EP

(71) Applicant (for all designated States except US): NESTEC S.A. [CH/CH]; Avenue Nestlé 55, CH-1800 Vevey (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **POUZOT, Matthieu** [FR/CH]; Avenue de l'Esplanade 4Bis, -, CH-1012 Lausanne (CH). **SCHMITT, Christoph** [FR/CH]; Chemin Aux Channey 17, -, CH-1077 Servion (CH). **MEZZENGA, Raffaele** [IT/CH]; Ch. des Uttins 2, -, CH-1028 Préverenges (CH).

(74) Agent: **MARQUARDT, Ulf**; Avenue Nestlé 55, -, CH-1800 Vevey (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

[Continued on next page]

(54) Title: FOOD PROTEIN AND CHARGED EMULSIFIER INTERACTION



(57) Abstract: The present invention relates to structures obtained from protein and emulsifier interaction, more particularly to structures comprising a protein supramolecular core coated with at least a lipidic layer. The invention also encompasses methods for obtaining these structures and food compositions comprising them.

WO 2008/025784 A1



— *of inventorship (Rule 4.17(iv))*

Published:

— *with international search report*

Food protein and charged emulsifier interaction**5 Field of the invention**

The present invention relates to structures obtained from protein and emulsifier interaction, more particularly to structures comprising a protein supramolecular core coated with at least a lipidic layer. The invention also
10 encompasses methods for obtaining these structures and food compositions comprising them.

Background of the invention

15 Proteins are complex structures which, in solution, can be easily disrupted by a number of factors (heat, pH, salt concentration etc.)

Disruption can be controlled so as to form supramolecular
20 assemblies of protein which are biologically useful structures.

Supramolecular assemblies have been used for example, in the form of protein aggregates, in food applications and
25 are increasingly being used as an emulsifier and as a partial substitute for fat.

US 6767575 B1 discloses a preparation of an aggregate whey protein product, whereby whey protein is denatured by
30 acidification and heating. The protein aggregates thus obtained are used in food application.

GB 1079604 describes improvements in the manufacture of cheese, whereby whey proteins undergo heat treatment at an
35 optimum pH value, in order to obtain insoluble whey proteins which are then added to raw milk.

WO 93/07761 is concerned with the provision of a dry microparticulated protein product which can be used as a fat substitute.

- 5 US 5750183 discloses a process for producing proteinaceous microparticles which are useful as fat substitute containing no fat.

A proteinaceous fat substitute is also disclosed in WO
10 91/17665 whereby the proteins are in the form of a water-dispersible microparticulated denatured whey protein.

A whey derived fat substitute product for use in foods is disclosed in WO 92/18239. It is manufactured by encasing
15 particles in a liposome membrane to give a good mouth-feel.

Apart from the food applications, proteins are also present in many pharmaceutical and cosmetic compositions.

20

Problems encountered with these structures however may include, amongst others, the fact that they are sensitive to their environment, that their taste or texture is not always desirable and that their solubility is limited to
25 certain pH values and media (generally hydrophilic solvents).

Therefore there still remains a need to overcome these disadvantages.

30

Object of the invention

Thus, the object of the present invention is to provide protein supramolecular structures which can be used in a broader range of applications.

35

Summary of the invention

Accordingly, the present invention proposes, in a first aspect, a coated denatured supramolecular protein core structure, wherein the coating comprises at least a first lipid monolayer essentially electrostatically bound to the protein core.

In a second aspect, the invention relates to a liposome-like structure comprising a denatured supramolecular protein core coated with a lipidic bilayer shell.

A supramolecular protein rod structure coated with lipids falls under a further aspect of the invention.

The present invention further encompasses a method of forming a coated denatured supramolecular protein core comprising the steps of:

- a. Preparing a solution of denatured supramolecular protein structures
- b. Adjusting the pH of the solution such that the protein structures are oppositely charged to the lipids used in step c and
- c. Electrostatically binding lipids to the supramolecular structures in order to form a lipid monolayer around a supramolecular protein core.

In a further aspect is provided a method of solubilising a protein supramolecular structure in a solution having a pH equivalent to the isoelectric pH of the protein comprising the step of:

- a. Coating the protein supramolecular structure with a coating comprising a lipidic bilayer such that the lipidic bilayer is essentially electrostatically bound to the protein supramolecular structure.

Similarly, a method of solubilising a protein supramolecular structure in a hydrophobic medium is provided, said method comprising the step of

- 5 a. Coating the protein supramolecular structure with a coating comprising at least a first lipid monolayer such that the lipid monolayer is essentially electrostatically bound to the protein supramolecular structure.

10 The use of a structure according to any of claims 1 to 21 in food compositions, in cosmetic compositions and their use as a vehicle for bioactive substances also form part of the invention.

15 Finally, a food composition and a cosmetic composition comprising a structure according to any of claims 1 to 21 fall under other aspects of the invention.

Figures

20 The present invention is further described hereinafter with reference to some embodiments shown in the accompanying figures in which:

- 25 - Fig. 1 shows a positively charged supramolecular core being electrostatically coated with a charged lipid,
- Fig. 2 shows a second layer coating step which yields a liposome-like structure,
- 30 - Fig. 3 shows the steps in forming a protein rod having a lipid monolayer,
- Fig. 4 compares Differential Interference Contrast (DIC) images of a supramolecular whey protein core without (top images) and

with (bottom images) a lipidic layer of sulfated butyl oleate at pH 4.3,

- 5
- Fig. 5 depicts the behaviour of whey protein aggregates and negatively charged lipids at a pH greater than the isoelectric pH of the protein, at a pH below the isoelectric pH of the protein and at a pH close to the isoelectric pH of the protein,
- 10
- Fig. 6 is a graph of mobility vs lipid concentration,
 - Fig. 7 is a graph of the diameter of the structures of the invention during formation vs the lipid concentration,
- 15
- Fig. 8 shows transmission electron microscopy images of β -lactoglobulin rods and DIC and polarised light images of the resulting complexes obtained with sulfated butyl oleate,
- 20
- Fig. 9 shows DIC images of β -lactoglobulin rod-sodium stearyl lactylate complexes, and
 - Fig. 10 shows images of β -lactoglobulin rod-DATEM (diacetyl tartaric acid esters of monoglycerides) complexes.
- 25

Detailed description of the invention

30 The present invention relates to a supramolecular protein core which is coated with lipids. By "supramolecular protein core" is meant any type of structure comprising at least more than one protein molecule and wherein the protein is in a denatured state. Such protein may be denatured either thermally, physically or chemically.

Referring to fig. 1 and fig. 3, the protein core is charged and coated with at least one layer of charged lipids.

- 5 The present invention provides a method of forming a coated denatured supramolecular protein core comprising the steps of firstly preparing a solution of denatured supramolecular protein structures, secondly adjusting the pH of the solution such that the protein structures are
- 10 oppositely charged to the lipids used in the subsequent step and finally, electrostatically binding lipids to the supramolecular structures in order to form a lipid monolayer around a supramolecular protein core.
- 15 The first step in the method consists of preparing a solution of denatured supramolecular protein structures. The supramolecular core therefore consists of an assembly of denatured proteins. The core may adopt the form of a micelle, an aggregate (fibrillar such as a rod or
- 20 spherical shape), or a gel.

- Methods for generating these supramolecular structures are well known in the art. They usually involve heat denaturation of a native protein under certain pH, certain
- 25 protein and salt concentration conditions in order to induce aggregation or gelation of the protein aqueous solution. The core may therefore be a protein micelle, a protein aggregate, a protein rod or a protein gel.

- 30 In order to form the supramolecular protein core of the invention, any protein selected from vegetal or animal sources may be used. It may include soy protein, milk protein (whey protein, β -lactoglobulin, casein, bovine serum albumin etc.), ovalbumin, meat protein etc.

Preferably however, the supramolecular core is not casein-based.

In a second step, the pH of the solution comprising the
5 supramolecular protein core is adjusted such that the
protein structures are oppositely charged to the lipids
used to coat them. The particles of aggregated denatured
proteins may bear an overall positive charge, or an
overall negative charge. Preferably, the particles are
10 positively charged at a pH below the isoelectric pH of the
native protein from which they are obtained.

This pH value may be different to the pH value needed to
form the supramolecular core. Preferably, the pH will be
15 adjusted to less than 5, even less than 4, preferably to
pH 3, depending on the lipids used for the coating in the
subsequent step. At these pH values, the supramolecular
structures are preferably positively charged, such that
they can be electrostatically bound to a negatively
20 charged lipid in a subsequent step.

The ionic complexation step consists then in providing the
negatively charged lipids to the solution of
supramolecular protein structures.

25

Thus, the resulting structures comprise a charged protein
core with at least a lipid monolayer coating.

The size of the protein core may vary from 100nm to 100µm,
30 preferably between 100nm and 10µm and can be controlled by
the method used for the formation of the protein core. The
person of skill in the art would know which method to use
in order to obtain the desired core size. The advantage of
the wide size variability is that, depending on the
35 desired application, the size of the core may be tailored

accordingly. The core may be spherical in shape or may be rod-like.

According to an embodiment of the present invention, the structure of the invention comprises a supramolecular protein rod coated with lipids. In order to produce rod protein supramolecular cores, protein such as β -lactoglobulin, bovine serum albumin or ovalbumin may be used. Preferably, β -lactoglobulin is used as the protein.

10

A method for obtaining such structures includes heating an aqueous solution (pH 2) comprising the native protein in a concentration of 25 g/L and sodium chloride (0.01M) at 80°C for 10 hours. Under these conditions, the denatured proteins assemble so as to form a supramolecular protein rod. The size of the rod may be monitored by the forming conditions and may range from 2 μ m to 7 μ m. According to the invention, the rod is coated with a lipid coating (as shown in Fig. 3). Preferably, the lipid coating is essentially electrostatically bound to the protein rod.

20

This process is further illustrated in Fig. 8 according to which a solution of rods is adjusted to pH 3 after formation and complexed with sulfated butyl oleate. Polarised light imaging and Differential Interference Contrast (DIC) imaging in Fig. 8 show the precipitation of rod/sulfated butyl oleate (SBO) complexes at pH 3. Fig. 9 and 10 further show the precipitation at pH 4.2 of β -lactoglobulin rods with sodium stearoyl lactylate (SSL) and β -lactoglobulin rods with diacetyl tartaric acid esters of monoglycerides (DATEM) respectively.

25

30

Referring to Fig. 1 and Fig. 3, the charged supramolecular assemblies are thus coated with at least a first lipid

monolayer essentially electrostatically bound to the protein core.

In order to have an essentially electrostatic binding, the
5 lipid is selected such that it is oppositely charged to the protein core. In a preferred embodiment, the lipids are negatively charged. Negatively charged lipids may be selected from sulfated butyl oleate, diacetyl tartaric
10 acid esters of monoglycerides, citric acid esters of monoglycerides, sodium stearyl-2 lactylate, lactic acid esters of monoglycerides, calcium stearyl lactylate, sodium lauryl sulphate etc.

The resulting interaction between the core and lipids of
15 opposite charge is essentially electrostatic. Indeed, in Fig. 6 showing a graph of mobility versus charged lipid concentration, it can be seen that, upon increasing the lipid concentration, the mobility is decreased. This observation confirms that the binding between the lipid
20 layer and the protein core is essentially electrostatic. Moreover, measurements of charge and size have shown that no detectable interactions occur between lipid and protein core at pH above isoelectric pH (tested at pH7 in the case of whey protein micelles and sulfated butyl oleate).

25 According to an embodiment of the invention, the supramolecular core may further encapsulate food-grade substances. The food-grade substance which may be entrapped in the particulate protein assemblies may be
30 flavours, for example, or may be selected from any bioactives such as, bacteria, metal ions, enzymes etc. Preferably, the substance is hydrophilic.

Thus the structures of the invention may serve as a
35 vehicle for these bioactives. They may therefore find

cosmetic, pharmaceutical and/or nutritional applications, whereby delivery of a sensitive active agent is needed.

The coating of the protein core may further comprise a
5 second lipid monolayer. This second layer is typically hydrophobically bound to the first lipid monolayer. A bilayer is thus formed which may, in a preferred embodiment, consist of intercalated monolayers. This bilayer forms a lipidic shell around the protein core (cf.
10 Fig. 2) and confers to the structure a liposome-like function, such that these structures may be used for transporting proteins through membranes in biological systems, for colloidal stability, for slow-release of entrapped particles etc.

15 The lipids used for the second monolayer may be charged or neutral. They may be the same as those used for the first monolayer or they may be different. Neutral lipids (including zwitterionic lipids) may be selected from
20 phospholipids.

Referring to Fig. 7 representing an embodiment where the lipids used for the first monolayer are the same as those used for the second monolayer, it can be seen that in
25 order to form the lipidic bilayer, the concentration of lipid has to be increased. The formation of the lipidic bilayer may be monitored by measuring the diameter size of the structures obtained or it may be monitored by monitoring the charge of the supramolecular protein core -
30 lipid complex. At a certain concentration of lipid, the structures consisting of a protein core coated with one lipid monolayer tend to attract each other thus forming larger structures. Above a certain lipid concentration threshold, the bilayer is formed and the size decreases.
35 This hydrophobically driven formation of the second layer

of lipids results in the charged heads of the lipid being exposed towards the aqueous phase.

Thus, according to the invention, when two lipid
5 monolayers are used for coating the protein core, a liposome-like structure is obtained (as shown in Fig. 2).

If charged lipids are used for the second monolayer, the liposome-like structure will have an overall charged
10 surface. Alternatively, if neutral lipids are used for the second monolayer, the surface of the liposome-like structure will be neutral.

The second layer and more precisely the hydrophilic head
15 borne by the lipid used for the second layer provides the essential properties of the liposome-like structure with respect to colloidal stability in solution or feasibility of transvection of the protein core through biological membranes for instance. Thus, the charge, steric hindrance
20 of the lipid used for the second lipid layer is an important feature which may be tuned for dedicated specific purposes.

With the liposome-like structure of the invention, many
25 improvements in the field of protein solubilisation, dairy powder protection etc. can be achieved due to the fact that the structures are purely self-assembled generated food-grade structures.

30 For instance, as shown in Fig. 4, the charged liposome-like structures may allow solubilisation of proteins at a pH close to the isoelectric pH of the protein. For whey protein, this value is between 3.5 and 4.6. Indeed, without a coating, the protein supramolecular assemblies
35 (e.g. micelles) tend to agglomerate due to the

neutralisation of charges at their surface at isoelectric pH, resulting in aggregation through dominating hydrophobic interactions. With a coating according to the invention, the structures will not flocculate at a pH
5 close to the isoelectric pH of the protein due to their surfaces being only positively or only negatively charged, such that the structures repel each other (cf. Fig. 5).

Thus the invention provides a method of solubilising a
10 protein supramolecular structure in a solution having a pH equivalent to the isoelectric pH of the protein comprising the step of coating the protein supramolecular structure with a coating comprising a lipidic bilayer which is essentially electrostatically bound to the protein
15 supramolecular structure.

This can find applications in sports drinks for example, which can have a low pH (about 4) and still have a high protein content, without loss of stability.

20

An advantage of the present invention is that the lipidic shell may be used as a protective barrier for the protein core against humidity, oxygen, protease etc. The liposome-like structure of the invention may also provide
25 protection against agglomeration of protein powders during the drying process.

An increase in the amount of protein content of fat matrices is possible with the structures of the invention
30 due to the solubilisation of proteins in hydrophobic media (oil, fatty matrices etc.). Thus, the present invention also provides a method of solubilising a protein supramolecular structure in a hydrophobic medium comprising the step of coating the protein supramolecular
35 structure with a coating comprising at least a first lipid

monolayer such that the lipid monolayer is essentially electrostatically bound to the protein supramolecular structure.

- 5 According to the invention, the surface properties of proteins may thus be changed such that a wider scope of applications for proteins may be contemplated.

Another advantage of the invention is that oils may be
10 solidified using the rods of the present invention. Thus, it represents an alternative to hydrogenation of lipids for the manufacture of products such as margarine etc. The resulting products have therefore not only a reduced amount of hydrogenated fats but also contain a
15 considerable amount of protein.

Due to the lipidic bilayer surrounding the protein core, a reduction of the astringency of protein supramolecular structures (in particular micelles) may be achieved. The
20 invention thus allows the sensory attributes of proteins to be improved.

As a summary, the structures of the invention may be used in food compositions.

25

Food compositions which comprise the structures of the invention may include beverage, yogurt, ice cream, sorbet, pet food, biscuits, dried food, milk powder, oil, fat, solidified oil, butter, margarine, food supplement, water-
30 in-oil emulsion etc.

The food compositions of the present invention may be used in a wide range of nutritional, pharmaceutical, and/or cosmetic applications.

35

These structures may also serve as nanovehicles for encapsulation and delivery of hydrophilic compounds.

5 The use of these structures in cosmetic compositions, and cosmetic compositions comprising these structures are also part of the invention. Typical cosmetic compositions may be selected from creams, lotions, gels, shampoos, soaps etc.

10 The present invention is further illustrated by means of the following non-limiting examples.

Examples

15

Liposome-like structure formation

A whey protein aggregates solution was prepared by subjecting a solution of native whey protein to a
20 temperature of 85°C for 15 minutes at pH 5.8. The aggregates are then isolated and used in the preparation of an aqueous solution comprising a concentration in protein of 1.511g/L and a concentration of sulfated butyl oleate greater than 0.4 g/L. The pH of the solution is
25 adjusted to pH3 and a temperature of 25°C. Under these conditions, immediate formation of a liposome-like structure comprising the whey protein aggregate core and a lipidic bilayer (Sulfated butyl oleate) is observed, due to the electrostatic self-assembly between the whey
30 protein core and the sulfated butyl oleate.

Mobility and size measurements

A mixed sample comprising a supramolecular protein
35 assembly (e.g. micelles) and lipids (e.g. sulfated butyl

oleate) was subjected to *in situ* measurements using a Zetasizer Nano-ZS (Malvern, UK).

5 The mobility (the sign of which is equivalent to the charge of the complexes) was determined by the electrophoretic mobility module (determination of the displacement of the particle under an imposed electric field). The results are shown in Fig. 6.

10 The size of complexes were measured by the light scattering module of the apparatus (fit of the autocorrelation fonction $g_2(t)$ with determination of the diffusion coefficient then related to the size by the Stokes- Einstein relation for spherical particles). The
15 results are shown in Fig. 7.

Claims

1. Coated denatured supramolecular protein core
structure, wherein the coating comprises at least a
5 first lipid monolayer essentially electrostatically
bound to the protein core.
2. Coated denatured supramolecular protein core
structure according to claim 1, wherein the coating
10 comprises a second lipid monolayer hydrophobically
bound to the first lipid monolayer.
3. Coated denatured supramolecular protein core
structure according to any of claims 1 or 2, wherein
15 the supramolecular core is a protein micelle, a
protein rod, a protein aggregate or a protein gel.
4. Coated denatured supramolecular protein core
according to any of the preceding claims, wherein a
20 food-grade substance is entrapped in the
supramolecular core.
5. Coated denatured supramolecular protein core
structure according to claim 4, wherein the food-
25 grade substance is selected from bacteria, metal
ions, bioactives etc.
6. Coated denatured supramolecular protein core
structure according to any of the preceding claims,
30 wherein the protein core is not casein-based.
7. Coated denatured supramolecular protein core
structure according to any of the preceding claims,
wherein the first lipid monolayer comprises charged
35 lipids selected from sulfated butyl oleate, diacetyl

- 5 tartaric acid esters of monoglycerides, citric acid
esters of monoglycerides, sodium stearyl-2
lactylate, lactic acid esters of monoglycerides,
calcium stearyl lactylate, sodium lauryl sulphate
etc.
8. Coated denatured supramolecular protein core
structure according to any of claims 2 to 7, wherein
the second lipid monolayer comprises charged or
10 neutral lipids.
9. Liposome-like structure comprising a denatured
supramolecular protein core coated with a lipidic
bilayer shell.
- 15 10. Liposome-like structure according to claim 9,
wherein at least the lipids used for the first
monolayer of the shell are charged lipids such that
the interaction between the core and the first
20 monolayer is essentially electrostatic and
wherein the lipids used for the second monolayer are
selected such that they hydrophobically interact
with the first monolayer.
- 25 11. Liposome-like structure according to any of claims 9
or 10, wherein the lipids used for the first monolayer
are selected from sulfated butyl oleate, diacetyl
tartaric acid esters of monoglycerides, citric acid
esters of monoglycerides, sodium stearyl-2 lactylate,
30 lactic acid esters of monoglycerides, calcium stearyl
lactylate etc.
12. Liposome-like structure according to any of claims 9
to 11, wherein the lipids used for the first monolayer
35 are the same as those used for the second monolayer.

13. Liposome-like structure according to any of claims 9 to 11, wherein the lipids used for the first monolayer are different to those used for the second monolayer.
- 5
14. Liposome-like structure according to any of claims 9 to 12, wherein the supramolecular core is a protein micelle, a protein rod, a protein aggregate or a protein gel.
- 10
15. Liposome-like structure according to any of claims 9 to 14, wherein a food-grade substance is entrapped in the supramolecular core.
- 15
16. Liposome-like structure according to claim 15, wherein the food-grade substance is selected from bacteria, metal ions, bioactives etc.
17. Liposome-like structure according any of claims 9 to 16, wherein the surface of the liposome is charged or neutral.
- 20
18. Supramolecular protein rod structure coated with lipids.
- 25
19. Supramolecular protein rod structure of claim 18, wherein the coating comprises at least one lipid monolayer electrostatically bound to the protein rod.
- 30
20. Supramolecular protein rod structure according to any of claims 18 or 19, wherein the protein is β -lactoglobulin, bovine serum albumin or ovalbumin.
21. Supramolecular protein rod structure according to any of claims 18 to 20, wherein the protein is denatured.
- 35

22. Method of forming a coated denatured supramolecular protein core comprising the steps of:
- a. Preparing a solution of denatured supramolecular protein structures
 - b. Adjusting the pH of the solution such that the protein structures are oppositely charged to the lipids used in step c
 - c. Electrostatically binding lipids to the supramolecular structures in order to form a lipid monolayer around a supramolecular protein core.
23. Method of claim 22, wherein the method comprises a further step of hydrophobically binding further lipids to the lipid monolayer such as to form a lipid-bilayer around the protein core.
24. Method of solubilising a protein supramolecular structure in a solution having a pH equivalent to the isoelectric pH of the protein comprising the step of:
- a. Coating the protein supramolecular structure with a coating comprising a lipidic bilayer such that the lipidic bilayer is essentially electrostatically bound to the protein supramolecular structure.
25. Method of solubilising a protein supramolecular structure in a hydrophobic medium comprising the step of:
- a. Coating the protein supramolecular structure with a coating comprising at least a first lipid monolayer such that the lipid monolayer is essentially electrostatically bound to the protein supramolecular structure.

26. Method of claim 25, wherein the coating comprises a
5 second lipid monolayer hydrophobically bound to the
first lipid monolayer.
27. Use of a structure according to any of claims 1 to 21
in food compositions.
- 10 28. Use of a structure according to any of claims 1 to 21
in cosmetic compositions.
29. Use of a structure according to any of claims 1 to 21
15 as a vehicle for bioactive substances.
30. Food composition comprising a structure according to
any of claims 1 to 21.
- 20 31. Food composition according to claim 30, wherein the
food composition is a beverage, yogurt, ice cream,
sorbet, pet food, biscuits, dried food, milk powder,
oil, fat, solidified oil, butter, margarine, food
supplement, water-in-oil emulsion etc.
- 25 32. Food composition according to any of claims 29 or 30,
wherein the food composition is used in nutritional,
pharmaceutical and/or cosmetic applications.
- 30 33. Cosmetic composition comprising a structure according
to any of claims 1 to 21.

Figures




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7






Figure 8



Complexes $R_{\beta LG}$ / SSL
pH 4.2 (Precipitates)

Figure 9



Complexes $R_{\beta LG}$ / DATEM
pH 4.2 (Precipitates)

Figure 10

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2007/058964

A. CLASSIFICATION OF SUBJECT MATTER

INV. A23J1/00 A23J1/20 A23J1/06 A23J1/02 A61K9/127
A23L1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A23J A61K A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 413 804 A (RHODES KENNETH H [US]) 9 May 1995 (1995-05-09) column 7, line 35 - column 8, line 25 column 137, lines 20-32; claim 1 -----	1-23, 27, 29-33
X	US 7 060 291 B1 (MEERS PAUL R [US] ET AL) 13 June 2006 (2006-06-13) claims 1,3,7; examples 1-16 -----	1-33
A	WO 2006/033985 A2 (SECRETARY OF AGRICULTURE THE U [US]; HARDEN JEROME W [US]; ONWULATA CH) 30 March 2006 (2006-03-30) the whole document ----- -/--	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

11 October 2007

Date of mailing of the international search report

25/10/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Adechy, Miriam

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2007/058964

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 6 767 575 B1 (HUSS MANFRED [DE] ET AL) 27 July 2004 (2004-07-27) cited in the application the whole document -----	
A	US 5 589 189 A (MOYNIHAN KAREN L [US]) 31 December 1996 (1996-12-31) the whole document -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2007/058964

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5413804	A	09-05-1995	NONE
US 7060291	B1	13-06-2006	US 2006172003 A1 03-08-2006
WO 2006033985	A2	30-03-2006	EP 1814401 A2 08-08-2007 US 2007190213 A1 16-08-2007
US 6767575	B1	27-07-2004	AT 239381 T 15-05-2003 AU 3145600 A 04-09-2000 WO 0048473 A1 24-08-2000 DE 19906379 A1 17-08-2000 DE 19964370 B4 11-05-2006 EP 1154700 A1 21-11-2001
US 5589189	A	31-12-1996	NONE