



(86) Date de dépôt PCT/PCT Filing Date: 2015/04/10
 (87) Date publication PCT/PCT Publication Date: 2015/10/15
 (45) Date de délivrance/Issue Date: 2023/04/18
 (85) Entrée phase nationale/National Entry: 2016/10/07
 (86) N° demande PCT/PCT Application No.: US 2015/025391
 (87) N° publication PCT/PCT Publication No.: 2015/157683
 (30) Priorité/Priority: 2014/04/11 (US14/251,220)

(51) Cl.Int./Int.Cl. *A61K 8/21* (2006.01),
A61K 8/19 (2006.01), *A61K 8/72* (2006.01),
A61K 8/81 (2006.01), *A61K 8/86* (2006.01),
A61K 8/89 (2006.01), *A61P 1/02* (2006.01),
A61P 19/08 (2006.01), *A61Q 11/00* (2006.01)
 (72) Inventeurs/Inventors:
 MCHALE, WILLIAM A., US;
 BROWN, DALE G., US
 (73) Propriétaire/Owner:
 PREMIER DENTAL PRODUCTS COMPANY, US
 (74) Agent: MOFFAT & CO.

(54) Titre : PATE DENTIFRICE POUR LA PROTECTION ET LA REPARATION DE L'EMAIL
 (54) Title: ENAMEL PROTECTANT AND REPAIR TOOTHPASTE

(57) **Abrégé/Abstract:**

Substantially aqueous-free, enamel protectant and enamel repair toothpastes containing: stannous fluoride, calcium and a substantivity agent comprising: an emulsion of polydimethylsiloxane in a nonionic surfactant, wherein: (a) substantivity of said stannous fluoride and calcium into biofilm present on enamel is enhanced through calcium binding shifting from bidentate to monodentate in the presence of stannous fluoride; and (b) said toothpastes indicate substantially improved, enamel protectant factor (EPF) and enamel repair factor (ERF) values compared to fluoride brushing treatments with comparable or higher fluoride levels.

Abstract

Substantially aqueous-free, enamel protectant and enamel repair toothpastes containing: stannous fluoride, calcium and a substantivity agent comprising: an emulsion of polydimethylsiloxane in a nonionic surfactant, wherein: (a) substantivity of said stannous fluoride and calcium into biofilm present on enamel is enhanced through calcium binding shifting from bidentate to monodentate in the presence of stannous fluoride; and (b) said toothpastes indicate substantially improved, enamel protectant factor (EPF) and enamel repair factor (ERF) values compared to fluoride brushing treatments with comparable or higher fluoride levels.

Enamel protectant and Repair Toothpaste

BACKGROUND OF THE INVENTION

1. Field of the Invention

5 The present invention is directed to advances in enamel protectant and enamel repair, aqueous-free, stannous fluoride, toothpastes; whereby the protectant and repair ingredients are substantive to enamel surfaces, thereby extending the protecting and repairing processes with improved stannous fluoride effectiveness. Key protectant and repair combinations for the toothpastes of the present invention comprise: stannous
10 fluoride and calcium in aqueous-free, substantivity agents.

2. Description of the Related Art

The use of fluoride in anticaries drug products, marketed in the U.S., is carried out under the guidance of the FDA's Fluoride Monograph, 21 CFR 355.10 (revised April 1,
15 2012).

Table 1

Concentration and Dosage of Stannous Fluoride in Dentifrice/Rinse/Gel products according to the Federal Register 21 CFR 355.10

20

Dentifrices	Dentifrices containing 850 to 1,150 ppm theoretical total fluorine in a gel or paste dosage form. Stannous fluoride 0.351 to 0.474% with an available fluoride ion concentration = 700 ppm for products containing abrasives other than calcium pyrophosphate. Stannous fluoride 0.351 to 0.474% with an available fluoride ion concentration = 290 ppm for products containing the abrasive calcium pyrophosphate.
Preventive treatment gel	Stannous fluoride 0.4% in an anhydrous glycerin gel, made from anhydrous glycerin and the addition of suitable thickening agents to adjust viscosity.
Treatment rinse	Stannous fluoride concentrate marketed in a stable form and containing adequate directions for mixing with water immediately before using to result in a 0.1% aqueous solution.

Dentifrices

Fluoride dentifrices have been shown in numerous clinical trials to be effective anticaries agents [Stookey, J. Dent. Res. 1990, 69(Special Issue): 805-812] and have been recognized as a major cause of the remarkable decline in caries prevalence in many developed countries. Dentifrices have been widely adopted around the world as the principle means of delivering topical fluoride and obtaining caries preventive benefits.

“Washout” of various enamel protectant and enamel repair ingredients from enamel surfaces by saliva flow, eventually controls the effective residence time of various commercial fluoride, enamel protectant and enamel repair, brushing compositions. To improve enamel protectant and enamel repair effectiveness, commercial, professionally prescribed, fluoride, brushing compositions resort to high levels of fluoride, i.e. 5000 ppm for Rx toothpastes, gels and rinses and to approximately 22,000 ppm fluoride for “in-chair”, professionally applied varnishes. In addition, standard OTC, fluoride toothpastes can contain up to 1500 ppm fluoride under the FDA’s Fluoride Monograph.

The current market for fluoride brushing products includes: professional and consumer, oral care, fluoride treatments, both OTC and Rx brushing products; including: toothpastes, gels, pastes and varnishes. As noted above, Rx fluoride toothpastes and Rx fluoride toothpastes are well outside fluoride Monograph levels containing up to 5000 ppm fluoride. Professional oral care, in-chair, fluoride varnishes contain up to about 22,000 ppm fluoride, while OTC fluoride toothpastes can contain up to 1500 ppm fluoride, the maximum level provided for the Monograph.

The: American Dental Association (ADA); the Food & Drug Administration (FDA) and oral care professionals including: general practitioners, periodontists, orthodontists, pediatric dentists, etc. as a group; are generally concerned over the trend of increasing fluoride levels. These organizations and oral care professionals generally favor using lower levels of fluoride in various in-chair treatments and various OTC and Rx, oral care, home treatments for patients, provided... enamel protection and repair, achieved with lower fluoride levels, are comparable to the results reported for brushing products with higher levels of fluoride. This preference for lower fluoride-brushing products is driven by the concern over toxicity, fluorosis in children, etc., associated with exposure to high fluoride levels, long term.

It is generally accepted, approximately 90% of the fluoride used in OTC and Rx, fluoride, brushing treatments is expectorated after use. Thus, the window for fluoride treatment of enamel is essentially limited to the time fluoride is being brushed onto the enamel. In contrast, fluoride varnishes containing 22,000 ppm fluoride, applied to the enamel by an oral care professional, are designed to maintain substantive fluoride levels on the enamel after patient expectoration.

Fluoride varnishes are generally applied professionally, at a frequency of about once every six months with the target audience comprising primarily children.

Dietary fluoride levels have gradually increased due to fluoridated drinking water and the fluoride in water used in food preparation, etc. In addition, most consumers use fluoride: toothpastes, rinses, gels, etc. Extensive literature citations indicate topical fluoride treatments are more effective in protecting and repairing enamel than treatment with systemic fluorides.

See: Ripa, Public Health Dent., 1991; 51:23-41.

Yet, with all this fluoride available, caries continues to pose a challenge: in children, as well as adults including coronal caries in the elderly, caries in dry mouth patients, caries in immunocompromised patients, caries in patients undergoing medical or dental treatment, etc.

There is a need to improve enamel protectant and enamel repair effectiveness for professional oral care, fluoride treatments, as well as for OTC fluoride products for patient use, while reducing the risk associated with exposure to high fluoride levels.

Additionally, there is a need to improve the efficacy of fluoride products in the area of enamel protection and enamel repair, where the efficacy of various fluoride treatments is assessed as a function of the fluoride level used to effect treatment of various conditions of the enamel.

OBJECTS OF THE PRESENT INVENTION

To provide stannous fluoride toothpastes with improved enamel protectant factor (EPF) values and improved enamel repair factor (ERF) values.

5 To provide stannous fluoride toothpastes with improved EPF and ERF values, where improvements in EPF and ERF values are accompanied by reduced fluoride content in the toothpastes.

10 To reduce fluoride levels in toothpastes, while attaining comparable or improved EPF and ERF values, compared to current, commercial, fluoride brushing products, as well as fluoride brushing products described in the prior art.

To improve EPF and ERF values for the enamel of "at-risk" patients.

15 To improve enamel conditions of "at-risk" patients including immunocompromised patients; cancer therapy, cardio treatment, diabetes, COP patients; etc.

DEFINITION OF TERMS

The following terms used throughout this specification and claims to describe features of the toothpastes of the present invention are described below:

20

"Aqueous-free" is defined as: substantially free from water.

"Enamel Protectant Factor (EPF)" is defined as: the percent reduction in enamel solubility divided by the fluoride level in parts per million using FDA method #40.

25

"Enamel Repair Factor (ERF)" is defined as: the average increase in enamel fluoride concentration divided by the fluoride level of the fluoride brushing product tested using FDA method #33.

30 "Toothpaste" is defined as: an abrasive-based paste suitable for administration of stannous fluoride to enamel surfaces by brushing with a toothbrush.

35 "Mucoadhesive" is defined as: a substance that is retained for a period of time onto surfaces in the mouth that is not easily removed by the mechanical action of the tongue nor by flow of saliva.

"Stable stannous fluoride" is defined as: compositions that, when chemically assayed, substantially retains the level of stannous and/or fluoride in an unreacted state.

5 "Biofilm" is defined as: a surface adherent film comprised of bacteria , exuded polysaccharides, etc., that is not easily removed by mechanical means or saliva flow.

"Substantivity agent" is defined as: a composition that improves the mucosal retention of the desired agents.

10

"Cation bridging" is defined as: electrical attraction between two films or membranes initiated by cation moieties.

15 "Shift in calcium binding from bidentate to monodentate" is defined as: the loss of "chelate"-like binding by calcium cations with a corresponding increase in single ligand binding by calcium cations.

"Nonionic surfactant" is defined as: a composition that indicates surface active properties with the absence of charged species.

20

"CaF⁺ moiety" is defined as: a monodentate calcium fluoride ion.

25 "Linear, polymeric, polycarboxylates, substantivity enhancer" is defined as: a linear polymer with carboxylate substituents that increases retention of compositions onto charged surfaces.

"Emulsion discontinuous phase" is defined as: the minor component in an emulsion that is surrounded by a continuous phase.

30 "Emulsion continuous phase" is defined as: an emulsion component that surrounds discontinuous phase component.

SUMMARY OF THE INVENTION

The present invention is directed to aqueous-free, toothpaste compositions and associated methods for treatment and methods of manufacture, wherein the

toothpastes contain stannous fluoride from between 850 and about 1500 ppm fluoride, in an aqueous-free, substantivity agent. The toothpastes of the present invention protect and repair enamel more effectively than toothpastes and toothpaste compositions containing comparable or substantially higher levels of fluoride, as indicated by comparative EPF and ERF values reported herein.

In at least one embodiment, the present invention provides substantially aqueous-free, enamel protectant and enamel repair, toothpastes containing: stannous fluoride and calcium in a substantivity agent, wherein:

- 10 (a) Substantivity of said stannous fluoride and calcium into biofilm present on enamel is enhanced through a shift in calcium binding from bidentate to monodentate in the presence of stannous fluoride; and
- (b) EPF and ERF values of at least about 2.5 and 200, respectively, are achieved with periodic administration of said toothpastes onto enamel surfaces with biofilm present.

In at least one embodiment, said substantivity agent is selected from emulsions of polymethylsiloxanes in nonionic poloxamer surfactants.

20 In at least one embodiment, some of said stannous fluoride and calcium is present as CaF^+ .

In at least one embodiment, said substantivity agent contains linear, polymeric polycarboxylate, substantivity enhancers.

25 In at least one embodiment, the substantially aqueous-free, enamel protectant and enamel repair, toothpastes contain stannous fluoride and calcium in a substantivity agent comprising an emulsion of polydimethylsiloxane polymer at viscosities from between about 10,000 cs and about 2.5 million cs as the discontinuous phase in a nonionic surfactant, continuous phase.

In at least one embodiment, the nonionic poloxamer surfactants, suitable for said toothpastes, are represented by the structural formula:



35 wherein the sum of x, y and z is from between 125 and 175.

In at least one embodiment, the nonionic surfactants, suitable for said toothpastes, are represented by the structural formula:



5 wherein $x = 76$, $y = 25$ and $z = 56$.

In at least one embodiment, the toothpastes contain unreacted, calcium and phosphate components.

10 In at least one embodiment, the present invention provides substantially aqueous-free, enamel protectant and enamel repair, toothpastes containing stannous fluoride and calcium in a substantivity agent comprising an emulsion of polydimethylsiloxane polymer at viscosities from between about 10,000 cs and about 2.5 million cs as the discontinuous phase in a nonionic poloxamer surfactant, continuous phase containing a
15 linear polymeric polycarboxylates substantivity enhancer.

In at least one embodiment, the nonionic poloxamer surfactants, suitable for said toothpastes, are represented by the structural formula:



20 wherein the sum of x , y and z is from between 120 and 150.

In at least one embodiment, the nonionic poloxamer surfactants, suitable for said toothpastes, are represented by the structural formula:



25 wherein $x = 76$, $y = 0$ and $z = 56$.

In at least one embodiment, the toothpastes contain unreacted, calcium and phosphate components.

30 In at least one embodiment, said nonionic surfactant is a surfactant selected from the group consisting of surfactants having the following, general, structural formula:



wherein the sum of x , y and z is from between 125 and 175.

In at least one embodiment, the present invention provides substantially, aqueous-free, enamel protectant and enamel repair, toothpastes, featuring monodentate-bidentate bonding of calcium in the presence of stannous fluoride; containing: stannous fluoride, calcium and phosphate components, and a substantivity agent emulsion containing
5 nonionic poloxamer surfactant as the continuous phase and polydimethylsiloxane polymer as the discontinuous phase; with an EPF of at least about 2.5 and ERF of at least about 200.

In at least one embodiment, said polydimethylsiloxane polymer discontinuous phase
10 comprises up to 20% of said emulsion.

In at least one embodiment, the present invention provides substantially aqueous-free, enamel protectant and enamel repair, toothpastes containing stable stannous fluoride and calcium on a substantivity agent substantive to biofilm coated enamel, wherein:

- 15 (a) substantivity of said stannous fluoride and calcium into said biofilm is enhanced by a substantivity enhancer and by a shift from bidentate binding of calcium to monodentate in the presence of stable stannous fluoride; and
- (b) said stannous fluoride, upon release from said substantivity agent, converts to the moiety CaF^+ which effects EPF and ERF values of at least about 2.5
20 and about 200, respectively.

In at least one embodiment, said stannous fluoride, upon release from said substantivity agent, includes a CaF^+ moiety in the presence of said calcium monodentate binding to said biofilm present on enamel.

25

In at least one embodiment, the pH of said toothpastes, upon administration to enamel with biofilm, is at least about 3.

In at least one embodiment, the level of said substantivity agent and said substantivity
30 enhancers is between about 0.5 and about 5% by wt. and about 0.1 and about 3% by wt., respectively.


In at least one embodiment, said EPF and ERF values range from between about 380 and about 1070.

35

In at least one embodiment, said calcium is selected from the group of calcium compositions comprising: calcium fumarate, calcium sulfate, calcium gluconate, mixed sodium and calcium salts of methyl/vinyl/ether/maleic copolymers and combinations thereof.

5

In at least one embodiment, said substantivity enhancer has the structural formula:



where m is an integer that provides weight between about 60,000 and about 1,000,000.

10 In at least one embodiment, the total calcium level is between about 0.5 and about 5.0.

In at least one embodiment, the concentration of polydimethylsiloxane is up to about 20% by wt. of said emulsion.

15 In at least one embodiment, the ratio of unreacted calcium and phosphate components is from between about 2 and about 1, and the level of calcium and phosphate mixture in said toothpastes ranges from between about 0.2% and about 3%.

20 In at least one embodiment, said emulsion substantivity agent comprises from between about 0.5 and about 3.0 wt. percent of said toothpastes and said copolymer substantivity agent comprises from between about 0.5 and about 2.8 wt. percent of said toothpastes.

25 In at least one embodiment, the pH of said toothpaste, upon exposure to saliva, is from between about 4 and about 8.

The unexpected enamel protectant and enamel repair features of the stannous fluoride toothpastes of the present invention, as detailed below in the Examples, Tables and Drawings; are attributed to unique, aqueous-free, toothpaste compositions, which feature:

- 5 stannous fluoride at fluoride levels from between about 850 and about 1500 ppm fluoride and calcium;
a substantivity agent;
cation bridging associated with microbial fluoride binding; and
a pH of at east about 4.0, when the toothpastes are administered to enamel in the
10 presence of saliva.

The enamel protectant and enamel repair, toothpaste compositions of the present invention, when brushed onto enamel, form substantive, mucoadhesive gels on enamel in the presence of saliva; which mucoadhesive gels gradually release the stannous
15 fluoride onto enamel. This slow release continues until the mucoadhesive gel is eventually totally solubilized by saliva. This gradual release minimizes the “wash-out” effect traditionally experienced with fluoride brushing products. The resultant enamel protectant and enamel repair increases in EPF and ERF values, resulting from the extended enamel residence time of stannous fluoride, calcium and cation bridging
20 associated with microbial fluoride binding to biofilm. This improved stannous fluoride efficiency reduces risks associated with elevated fluoride levels.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 of the Drawings summarizes comparative, in vitro, enamel protectant factor
25 (EPF) values for enamel treated with a toothpaste of the present invention with stannous fluoride at 1150 ppm fluoride; compared with: (a) an Rx 5000 ppm, sodium fluoride toothpaste, and (b) an OTC, 900 ppm, stannous fluoride toothpaste.

Figures 2 through 5 of the Drawings illustrate enamel repair factor (ERF) values for toothpastes of the present invention with compositions, as described in Examples 1 through 5, with stannous fluoride at: 1100, 1148 and 1150 ppm stannous fluoride compared with toothpastes having stannous fluoride at 1100 ppm and sodium fluoride at 900 ppm or sodium fluoride at 5000 ppm, respectively.

Figures 2 through 5 present ERF values for toothpastes of the present invention comprising compositions, as described in Examples 1 through 5. These toothpastes of the present invention contain stannous fluoride at levels of: 1100, 1148 and 1150 ppm stannous fluoride, compared with: (a) a 900 ppm sodium fluoride toothpaste, and (b) an 1100 ppm stannous fluoride toothpaste, (c) a 5000 ppm sodium fluoride toothpaste or (d) a 900 ppm fluoride toothpaste.

Figures 2 through 5 of the Drawings illustrate enamel repair factor (ERF) for toothpastes of the present invention with compositions as described in Examples 1 through 5. These toothpastes of the present invention contain stannous fluoride at levels of: 1100, 1148 and 1150 ppm stannous fluoride compared with 1000 ppm stannous fluoride toothpaste, 900 ppm sodium fluoride toothpaste and 5000 ppm sodium fluoride toothpaste.

DETAILED DESCRIPTION OF THE PRESENTLY PREFERRED EMBODIMENTS

Aqueous-free, stannous fluoride, toothpastes of the present invention comprise substantivity agents that contain various enamel protectant and enamel repair ingredients. These substantivity agents function as carriers for various enamel protectant and repair ingredients. These substantivity agents are characterized by their ability, in the presence of saliva, to form mucoadhesive gels which are substantive to enamel with biofilm. These substantive, mucoadhesive gels are further characterized by their ability to: (a) gradually dissolve when exposed to saliva flow, and (b) gradually release various enamel protectant and enamel repair ingredients onto enamel surfaces with biofilm as they dissolve. This gradual dissolution feature of these mucoadhesive gels minimizes saliva "wash-out" of enamel protectant and enamel repair ingredients by gradually releasing these ingredients onto enamel surfaces with biofilm. The substantivity agents of the present invention extend the duration of enamel protectant

and enamel repair treatments and support cation bridging associated with microbial fluoride binding, thereby enhancing the EPF and ERF values of various toothpastes of the present invention, while simultaneously reducing the level of fluoride required to achieve the unexpected increases in EPF and ERF values.

5

In as preferred embodiment of the invention, calcium and phosphate components are included in the toothpastes. These are described by Ming Tung in U.S. Patents: 5,037,639; 5,268,167; 5,427,768; 5,437,857; 5,460,803; 5,562,895; by Tung in the American Dental Association Foundation publication, "ACP Technology,"; by Schemehorn, et. al., in The Journal of Clinical Dentistry Vol. XXII: No 2. 51-54, 2011; by 10 the 19 references cited by Schemehorn, et. al.; and by the description of various Gantrez® resins containing calcium, including Gantrez® M955 available from International Specialty Products, Wayne, NJ, USA.

15 The aqueous-free, substantivity agents of the present invention hold the various enamel protectant and enamel repair ingredients, including stannous fluoride, calcium and phosphate components, in a condition where these ingredients remain stable and unreacted. When this aqueous-free, substantivity agent is exposed to saliva, it forms a mucoadhesive gel that is substantive to enamel with biofilm. This mucoadhesive gel 20 continues to hold the enamel protectant and enamel repair ingredients onto enamel surfaces with biofilm without the ingredients reacting. These ingredients eventually react upon being released onto saliva and biofilm coated, enamel surfaces.

Eventually, this mucoadhesive, substantivity agent is totally dissolved by saliva, 25 releasing the balance of unreacted enamel protectant and enamel repair ingredients onto saliva and biofilm coated, enamel surfaces.

Aqueous-free, toothpastes of the present invention contain enamel protectant and enamel repair ingredients, suitable for protecting and repairing dental enamel; wherein: 30 said aqueous-free, toothpastes inhibit premature reaction of the enamel protectant and enamel repair ingredients;

the enamel protectant and enamel repair ingredients are introduced onto enamel with biofilm via saliva that solubilizes substantivity agents that are substantive to: enamel, dentin, biofilm and pellicle;

the enamel protectant and enamel repair ingredients contained in the substantivity agents are gradually released onto the enamel in an unreacted state as the saliva soluble, substantivity agent undergoes saliva dissolution at rates, which are controlled by saliva flow and the composition of the substantivity agent; and

5 bidentate binding of calcium shifts to monodentate binding of calcium in the presence of stannous fluoride.

For purposes of the present invention, saliva soluble, aqueous-free emulsions used as substantivity agents include those emulsions that are comprised of

10 polydimethylsiloxane polymers in nonionic surfactants, as described in U.S. Patents: 5,032,387; 5,098,711; 5,538,667; 5,651,959; having the structural formula:

$$\text{HO}[(\text{C}_2\text{H}_4\text{O})_x / (\text{C}_3\text{H}_6\text{O})_y] - [\text{C}_3\text{H}_6\text{O}]_z - [(\text{C}_2\text{H}_4\text{O})_x / (\text{C}_3\text{H}_6\text{O})_y]\text{H}$$

wherein the sum of x, y and z is between 120 and 150. In a preferred embodiment, x = 76, y = 0 and z = 56.

15 Preferred aqueous-free, saliva soluble emulsions for use as substantivity agents of the present invention include aqueous-free emulsions comprising a nonionic surfactant continuous phase and a discontinuous phase of polydimethylsiloxane (PDMS) at viscosities ranging from between about 1500 cs and about 2.5 million cs. Particularly

20 preferred, aqueous-free emulsions include a nonionic surfactant continuous phase and a discontinuous phase PDMS at viscosities between 10,000 cs and 2.5 million cs.

Preferred polydimethylsiloxanes are selected from the group consisting of polydimethylsiloxane: at 1500 cs, at 10,000 cs, at 100,000 cs, at 250,000 cs, at 500,000

25 cs, at 750,000 cs, at 1.5 million cs, at 2.2 million cs, at 2.5 million cs and combinations thereof.

Foam modulators are useful in the present invention. These include, without limitation: materials operable to control amount, thickness or stability of foam generated by the

30 composition (e.g., dentifrice composition) upon agitation. Any orally acceptable, foam modulator can be used, including polyethylene glycols (PEGs), also known as polyoxyethylenes. High molecular weight PEGs are suitable, including those having an average molecular weight of about 200,000 to about 7,000,000, for example about 500,000 to about 5,000,000 or about 1,000,000 to about 2,500,000. One or more PEGs

are optionally present in a total amount of about 0.1% to about 10%, for example about 0.2% to about 5% or about 0.25% to about 2%.

5 Humectants useful for the present invention include, without limitation: polyhydric alcohols such as glycerin, sorbitol, xylitol or low molecular weight PEGs. In various embodiments, humectants can prevent hardening of the toothpastes upon exposure to air. In various embodiments, humectants also function as sweeteners.

10 Any other desired components may be added to the compositions, including, for example, additional: mouth-feel agents, pH modifying agents, flavorants, sweeteners, additional anticalculus and Antiplaque agents, abrasives, polishing agents, antimicrobial (e.g. antibacterial) agents such as those described in U.S. Pat. No. 5,776,435, saliva
15 stimulants, anti-inflammatory agents, H₂ antagonists, nutrients, vitamins, proteins, antioxidants, colorants, or additional active materials useful for the prevention or treatment of a condition or disorder of hard or soft tissue of the oral cavity, the prevention or treatment of a physiological disorder or condition, or to provide a cosmetic benefit

20 Tables 2 through 8 summarize:

EPF and ERF data for toothpastes of the invention at varying compositions, as described in Examples 1 through 5, compared with commercial, fluoride, brushing products;

Actual Examples of toothpastes of the invention;

25 Illustrative Examples of toothpastes of the invention; and
Illustrative Treatments with toothpastes of the invention.

In a preferred embodiment of the invention, the toothpastes of the present invention contain ingredients that substantially effect enamel protection factor (EPF) and enamel
30 repair factor (ERF) values. These include:

Stannous fluoride and calcium,

Phosphate components, and

Substantivity agents and substantivity enhancers, including: mixed sodium and calcium salt copolymers of methyl/vinyl/ether/maleic acid;

35 where the stannous fluoride, calcium and phosphate components remain unreacted

and the pH of the toothpaste when administered to saliva coated enamel is at least about 4.

The foregoing ingredients are described in detail below.

5

Stannous Fluoride Concentration

The amount of stannous fluoride in the toothpastes of the present invention, applied to the toothbrush (dose) is not as important as the concentration of available stannous fluoride in the toothpaste. Heretofore, reducing fluoride concentration in brushing products has been reported not to be as effective as regular concentration fluoride products.

10

Petersson, et. al., Swed. Dent. 1982, 6:233-238

Metropoulos, et. al., Community Dent. Health, 2002, 1:193-200.

15

The extraordinary EPF and ERF values of the toothpastes of the present invention allow for reducing stannous fluoride concentrations while effecting acceptable fluoride protection and uptake results.

The fluoride dose is important in regard to enamel fluorosis in children under six years of age, due to fluoride toothpaste ingestion. For this reason, reducing the amount of stannous fluoride applied with the toothpastes of the present invention is a preferred strategy over lowering the dose of stannous fluoride toothpastes of the invention intended for use by children under six years of age.

20

While fluoride brushing products have a long history of safety, there is a continuing concern associated with dental fluorosis due to fluoride ingestion in children under age six. See: Dendrys, J. Am. Dent. Assoc. 2000, 131(6): 746-755.

Studies have shown that for children 1-3 years of age, 30 to 75% of the fluoride brushing product is ingested; and for children 4-6 years of age, 14 to 48% is ingested. Warren and Levy, Pediatr. Dent., 199, 21:265-271.

25

30

Stannous fluoride toothpastes of the present invention, with their improved efficacy can be used at reduced stannous fluoride levels, and thereby substantially lower the risk of overdosing and the onset of fluorosis, while delivering effective EPF and ERF results.

- 5 See also Zero, BMC Oral Health, 2006, 6(Suppl 1): 59; 1-13.

Monitoring the fluoride-mineral phase formed on enamel as a function of the concentration of fluoride ions [F⁻] in the demineralizing medium

See: Mohammed, et. al., Caries Res., 2013; 47:421-428.

- 10 ▪ At below 45 ppm [F⁻] in the solution ¹⁹F MAS-NMR showed fluoride-substituted apatite formation, 1B ¹⁹F magic angle, spinning nuclear magnetic resonance was used to characterize the solid phase precipitated on enamel as a function of fluoride concentration during exposure of the enamel to an *in vitro* demineralization system. The cariostatic effect of fluoride is due to the formation of F_sa HAP and CaF₂ depending on the [F⁻] level in the solution.
- 15 ▪ Above 45 ppm, calcium fluoride (CaF₂) formed in increasing proportions
- Further increases in [F⁻] caused no further reduction in demineralization, but increased the proportion of CaF₂ formed.
- As to the mechanism of fluoride anticaries efficacy.... fluoroapatite formation in enamel is investigated.
- 20

Advantages of ¹⁹F MAS-NMR:

- (1) selectively probes the local environment of only fluorine atoms in the sample, permitting direct identification of the possible structural forms in which [F⁻] may exist within the enamel.
- 25 (2) detects all fluorine present, whether:
crystalline,
amorphous, or
adsorbed.
- (3) measures very low concentrations of fluoride on the order of 0.1%.

30 ¹⁹F MAS-NMR established the effects of varying fluoride concentrations on fluoride-enamel interactions under acidic conditions using bulk enamel blocks rather than powder.

For the samples demineralized in the presence of [F⁻]:

- (1) chemical shifts were identified;
- 35 (2) formulation of fluoride-substituted apatite: ^(a)(Ca₁₀CPO₄)₆F_{2-x}.

(F_s-HAP); and

(3) formation of CaF₂;

- were observed in the aged portion at 45 ppm [F⁻] solutions,
- At [F⁻] above 45 ppm less F_s-HAP forms and an increased signal for CaF₂,
5 and
- For [F⁻] above 136 ppm mostly CaF₂ was identified.

The present study demonstrates that the addition of fluoride produces F_s-HAP as a major chemical species only at low concentrations of fluoride.

There is overwhelming evidence that low fluoride levels found in:

- 10 (a) saliva can significantly reduce enamel demineralization, and
- (b) plaque has the potential to remineralize even at pH values typically regarded as demineralizing.

Calcium, CaF₂, CaF⁺ and Phosphate

15 Christofferson, et. al., in ACTA ODONTOL. SCAND. 1988, 46:325-336, reports:

“It is suggested that the calcium fluoride-like material formed on dental enamel during treatment of enamel with acidified solutions of high fluoride content is a phosphate-containing calcium fluoride.”

20

“The aims of the present work are to determine the rates of growth and dissolution of pure calcium fluoride in aqueous suspensions and possible mechanisms controlling these processes, and to study the properties of the calcium fluoride-like material formed by adding fluoride to systems containing hydroxyapatite crystals and/or dissolved
25 calcium and phosphate, simulating the type of calcium fluoride-like material formed on dental enamel as a result of topical treatment with acidified solutions of high fluoride content.”

30 “From our results of dissolution of pure CaF₂ in systems containing phosphate it can be seen that 1 μm phosphate has a dramatic effect on the rate of dissolution of CaF₂.”

“The calcium fluoride-like materials containing phosphate appear to be more likely candidates to serve as slow fluoride release agents.”

B. Øgaard's "CaF₂ Formation: Cariostatic Properties and Factors of Enhancing the Effect," *Caries Res.*, 2001; 35 (Suppl) 11:40-41, teaches:

5 "CaF₂ or a CaF₂-like material/phosphate-contaminated CaF₂ is a major reaction product during topical treatment of dental hard tissues. Recently, evidence has suggested that CaF₂ is formed not only on surfaces but also to some extent in the enamel. The minimum concentration of fluoride required for CaF₂ formation is not well known and may depend on whether calcium is available from plaque fluid or only through dissolution of the dental hard tissue. Furthermore surface adsorption of fluoride
10 to crystals may cause local concentrations necessary for CaF₂ formation. It has been suggested that CaF₂ acts as a pH-controlled reservoir of fluoride. The rate-controlling factor appears to be phosphate, which controls the dissolution rate of CaF₂ at high pH. Increasing fluoride concentration, prolonging the exposure time or using a fluoride solution with low pH can increase CaF₂ formation. CaF₂ formed at low pH contains
15 less internal phosphate which has been shown to be less soluble. This may be of clinical significance for fluoride applied topically a few times per year."

"The interaction between the fluoride ion and dental hard tissues has been investigated extensively since modern fluoride research started in the 1940s. The chemistry of this
20 process is complicated due to many impurities in hydroxyapatite-like carbonate and magnesium and to a large variety of fluoride concentrations, pH and composition of the agents used in caries prevention. During pH cycling in plaque, fluoride may exchange with hydroxyl in the apatite and a series of solids with intermediate composition and crystallographic properties are formed known as fluorhydroxyapatite."

25 "CaF₂ is the major or probably the only reaction product on dental hard tissues from short treatments with relatively concentrated fluoride agents (Cruz et. al., *Scand. J. Dent. Res.*, 1992; 100:154-158). Without doubt, this pH-controlled depot of CaF₂ plays a major role in the cariostatic effect of topical fluoride. CaF₂ has been detected on
30 dental hard tissues weeks and months after a single topical fluoride treatment (*Caries Res.*, 1991, 25:21-26) and is the only logical way to explain that such treatments have a cariostatic effect. By treating enamel samples subjected to topical fluoride treatment with KOH, the cariostatic effect is lost (Øgaard, et al., *J. Dent. Res.*, 1990, 69:1505-1507)."

35

J.M. ten Cate's "Review on Fluoride, with special emphasis on calcium fluoride mechanisms in caries prevention", *Eur. J. Oral Sci.*, 1997, 105:461-465. teaches:

5 "For treatments to be effective over periods longer than the brushing and the following salivary clearance, fluoride needs to be deposited and slowly released. Calcium fluoride (or like) deposits act in such a way, owing to a surface covering of phosphate and/or proteins, which makes the CaF_2 less soluble under in vivo conditions than in a pure form in inorganic solutions. Moreover, due to the phosphate groups on the surface of the calcium fluoride globules, fluoride is assumed to be released with decreasing pH
10 when the phosphate groups are protonated in the dental plaque."

"In the presence of low concentrations of fluoride in solution (such as saliva or plaque fluid), hydroxyapatite might be dissolved below the critical pH (for hydroxyapatite), but the released mineral ions could be reprecipitated as fluoroapatite or a mixed fluor-
15 hydroxyapatite. This mechanism prevents the loss of mineral ions, while providing additional protection to mineral crystallites by laying fluoride-rich other layers onto the apatite crystallites."

"These observations point to the presence of slowly releasing fluoride reservoirs, either
20 on the dentition or the mucosal surfaces. Recent work has shown that in particular the oral mucosa, both by its chemical and morphological nature and the large surface area, is an underestimated retention site of fluoride."

"Research has shown that small amounts of fluoride in plaque and saliva are sufficient
25 to shift the de, remineralization balance favorably. Such levels should then be available throughout the day, in particular during periods of carbohydrate fermentation in the plaque. A fluoride-releasing reservoir system, effective at low pH, such as shown for calcium fluoride, would be a preferred system."

30 Vogel, et al., in "No Calcium-Fluoride-Like Deposits Detected in Plaque shortly after a Sodium Fluoride Mouthrinse", *Caries Res.*, 2010; 44:108-115, reported:

"Plaque 'calcium-fluoride-like' (CaF_2 -like) and fluoride deposits held by biological/bacterial calcium fluoride (Ca-F) bonds appear to be the source of cariostatic

concentration of fluoride in plaque fluid. The aim of this study was to quantify the amounts of plaque fluoride held in these reservoirs after a sodium fluoride rinse."

5 "The results suggest that either CaF₂-like deposits were not formed in plaque or, if these deposits had been formed, they were rapidly lost. The inability to form persistent amounts of CaF₂-like deposits in plaque may account for the relatively rapid loss of plaque fluid fluoride after the use of conventional fluoride dentifrices or rinses."

10 "Based on laboratory [Margolis and Moreno, J. Dent. Res., 1990, 69 (Spec. Issue) 606-613; J. Am. Dent. Assoc., 2000, 13:887-889; and clinical observations (reviewed by Featherstone, J. American Dent. Assoc., 2000, 13:887-889; the current models for increasing the anticaries effects of fluoride (F) agents emphasize the importance of maintaining a cariostatic concentration of F in oral fluids."

15 "This inability to form potentially more persistent calcium fluoride deposits, which appears to be due to the low concentration of oral fluid Ca, may account for the relatively rapid loss of F in plaque after the use of current over-the-counter topical F agents. It should be noted in this regard that (1) studies in which a Ca preapplication was used to ameliorate this situation have produced very large and persistent increases
20 in both plaque fluid and that salivary fluoride (Vogel, et. al., Caries Res., 2006, 40:449-454; Vogel, et. al., Caries Res. 2008 (a) 421;401-404; and Vogel, et. al., J. Dent. Res. 2008(b) 87:466-469; and that (2) preliminary studies (unpubl.) using modifications of the techniques described here confirm that the use of a Ca prerinse prior to a F rinse indeed forms large amounts of CaF₂-like deposits."

25

Substantivity Agents

For purposes of the present invention, substantivity agent refers to a composition or combination of compositions that, when administered to oral cavity surfaces with biofilm, enhance the retention of stannous fluoride and calcium to said biofilm
30 containing oral cavity surfaces.

The unexpected enamel protectant and enamel repair features of the aqueous-free, stannous fluoride toothpastes of the present invention are attributed to the unique substantivity properties indicated by the toothpastes of the invention.

35

For purposes of the present invention, preferred substantivity agents for the stannous fluoride toothpastes of the present invention include various aqueous-free emulsions of polydimethylsiloxane/polymers in nonionic surfactants at viscosities of at least about 10,000 cs.

5

These substantivity agents form mucoadhesive gels in the presence of saliva, which are substantive to biofilm-coated enamel and gradually dissolve under saliva flow, releasing stannous fluoride onto the biofilm on the enamel at a pH of at least about 3; thereby effecting EPF and ERF values of at least about 2.5 and about 200, respectively.

10

For purposes of the present invention, substantivity agents include saliva soluble, aqueous-free emulsions comprised of:

polydimethylsiloxane polymers in solid, nonionic surfactants, as described in U.S.

15 Patents: 5,032,387; 5,098,711; 5,538,667; 5,645,841; 5,651,959; having the following structural formula:



wherein $y = 0$ and the sum of x and z is between 120 and 150. In a preferred embodiment, $x = 76$, $y = 0$ and $z = 56$.

20

Preferred aqueous-free, saliva soluble emulsions for use in the substantivity agents of the present invention include aqueous-free emulsions comprising a nonionic surfactant, continuous phase and a discontinuous phase of polydimethylsiloxane (PDMS) at viscosities ranging from between about 1500 cs and about 2.5 million cs. Particularly preferred are aqueous-free emulsions with a nonionic surfactant continuous phase and a discontinuous PDMS phase at viscosities between 10,000 cs and 2.5 million cs.

25

Preferred polydimethylsiloxanes are selected from the group consisting of polydimethylsiloxane: at 1500 cs, at 10,000 cs, at 100,000 cs, at 250,000 cs, at 500,000 cs, at 750,000 cs, at 1.5 million cs, at 2.2 million cs, at 2.5 million cs and combinations thereof.


30

In a preferred embodiment of the invention, copolymers described below are useful as substantivity enhancers; when combined with the aqueous-free, substantivity agents of

35

the present invention. These substantivity enhancers include various linear polymeric, polycarboxylates, such as: copolymers of sodium and calcium salts of methyl/vinyl/ether/maleic acid including those copolymers available commercially as Gantrez® MS-955 polymer, a mixed sodium and calcium salt of methyl/vinyl/ether/maleic acid copolymer; where the cations form salt bridges which cross-link the polymer chains.

The chemical structure of this copolymer is represented by the following chemical structure:



where m is an integer that provides molecular weight for the polymer between about 60,000 and about 500,000.

Preferred are 1:4 to 4:1 copolymers of maleic anhydride or acid with another polymerizable ethylenically unsaturated monomer, preferably methyl vinyl ether (maleic anhydride) having a molecular weight of about 30,000 to about 1,000,000.

Sodium and calcium salts of carboxymethyl cellulose ether polymers can also be used including sodium and calcium salts of carboxymethyl cellulose ether, hydroxyethyl cellulose ether, sodium cellulose ether, etc.

The contribution of Ca^{++} from the copolymer substantivity enhancer to ERF values is illustrated in Figures 5 and 6 of the Drawings and Tables 6 and 7.

Examples 1 through 5

The following stannous fluoride toothpaste samples were prepared, as described below, and subsequently tested for EPF and/or ERF values, as described in Tables 1 through 8 and Figures 1 through 5 below:

Example 1

A 5 liter Ross/Olsa vacuum mixer kettle with internal homogenizer was heated to 60 degrees C while the vessel was charged with 592.225 gm of anhydrous glycerin.

5 Crodasinic L, 53.5 gm and TEGO betaine, 53.5, were added and homogenized at 1500 RPM for 10 minutes under vacuum. Then 942.1 gm of PEG 400 was added along with ULTRAMULSION® [poloxamer 407 (80%)] and 2.5 million cs polydimethylsiloxane (20%), 64.5 gm, with homogenizer speed at 1500 rpm for 10 minutes. Additional glycerin, 592.225 gm, was added along with acesulfame K, 21.0 gm, titanium dioxide,

10 53.5 gm, calcium sulfate, 97.0 gm, micronized (20 micron) sodium phosphate monobasic, 17.4 gm, and Gantrez MS-955, 107.5 gm. The homogenizer at 1500 rpm was begun and continued for 15 minutes under vacuum. The contents were cooled to 25 degrees C and 1.64% stannous fluoride in glycerin, 1443.5 gm, was added to the kettle. The homogenizer was set at 1000 rpm under vacuum for 10 minutes and

15 stopped. To the kettle was added Sident 22S, 650 gm, and Zeodent 113, 268 gm, with the anchor stirrer at medium speed. Sucralose, 4.5 gm, was added and the homogenizer set at 1000 rpm for 5 minutes under vacuum. Vanillamint, 33 gm, and Multisensate flavor, 4.55 gm, were added and homogenized at 1000 rpm for 5 minutes. The homogenizer stopped and the anchor stirrer continued for 10 minutes with cooling

20 to 25 degrees C. The contents was dispersed into tubes for use. Upon dispensing, the toothpaste indicated a pleasant taste with no stannous fluoride aftertaste.

Example 2

25 In a stainless steel 500 mL mixing vessel, using an overhead stirrer, 34.26 gm of PEG 400 were added along with 49.406 gm of glycerin, and 57.74 gm of 1.64% stannous fluoride in glycerin. Stirring was begun at a low speed while heating to 80°C. Then 2.58 gm of poloxamer 407/2.5 million cs (10%) ULTRAMULSION® substantivity agent was added with stirring for 15 minutes. The overhead stirrer speed was increased to

30 medium and 4.3 gm of Gantrez MS-955 substantivity enhancer was added with stirring for 5 minutes. Then Crodasinic L, 2.14 gm; Sucralose, 0.344 gm; and acesulfame K, 1.074 gm were added with continued stirring for 10 minutes. TEGO betaine CKD, 2.14 gm, was added and stirred for 5 minutes. Calcium fumarate, anhydrous, micronized, 2.84 gm, was added with stirring for 5 minutes followed by addition of sodium

dihydrogen phosphate, anhydrous, micronized, 0.696 gm, was added with continued stirring for 5 minutes.

In multiple increments, Sident 22S, 26.0 gm was added with stirring 2 minutes between
5 each addition. When the Sident 22S was all added, then Zeodent 113, 10.72 gm added
with stirring continuing for 15 minutes after which vanillamint, 2.14 gm, spearmint, 1.074
gm, TiO₂, 2.14 gm, ICE1500, 0.16 gm, and Multisensate flavor, 0.250 gm, were added
with continued stirring for 5 minutes. The stirrer was removed and the toothpaste was
10 added to dispensing tubes. When used as a toothpaste, a pleasant, refreshing
mouthfeel and very little metallic taste is perceived.

Example 3

In a stainless steel 500 mL mixing vessel, using an overhead stirrer, 32.26 gm of PEG
15 400 were added along with 45.446 gm of glycerin, and 57.74 gm of 1.64% stannous
fluoride in glycerin. Stirring was begun at a low speed while heating to 80°C. Then 2.58
gm of poloxamer 407/2.5 million cs (10%) ULTRAMULSION® substantivity agent was
added with stirring for 15 minutes. The overhead stirrer speed was increased to
medium and 4.3 gm of Gantrez MS-955 substantivity enhancer was added with stirring
20 for 5 minutes. Then Crodasinic L, 2.14 gm; Sucralose, 0.344 gm; and acesulfame K,
1.074 gm were added with continued stirring for 10 minutes. TEGO betaine CKD, 2.14
gm, was added and stirred for 5 minutes. Calcium lactate gluconate, anhydrous,
micronized, 8.8 gm, was added with stirring for 5 minutes followed by addition of
25 sodium dihydrogen phosphate, anhydrous, micronized, 0.696 gm, was added with
continued stirring for 5 minutes.

In multiple increments, Sident 22S, 26.0 gm was added with stirring 2 minutes between
each addition. When the Sident 22S was all added, then Zeodent 113, 10.72 gm added
with stirring continuing for 15 minutes after which vanillamint, 2.14 gm, spearmint, 1.074
30 gm, TiO₂, 2.14 gm, ICE1500, 0.16 gm, and Multisensate flavor, 0.250 gm, were added
with continued stirring for 5 minutes. The stirrer was removed and the toothpaste was
added to dispensing tubes. When used as a toothpaste, a pleasant, refreshing
mouthfeel and very little metallic taste is perceived.

Example 4

In a stainless steel 500 mL mixing vessel, using an overhead stirrer, 35.18 gm of PEG 400 were added along with 47.446 gm of glycerin, and 57.74 gm of 1.64% stannous fluoride in glycerin. Stirring was begun at a low speed while heating to 80°C. Then 2.58 gm of poloxamer 407/2.5 million cs (10%) ULTRAMULSION® substantivity agent was added with stirring for 15 minutes. The overhead stirrer speed was increased to medium and 4.3 gm of Gantrez MS-955 substantivity enhancer was added with stirring for 5 minutes. Then Crodasinic L, 2.14 gm; Sucralose, 0.344 gm; and acesulfame K, 1.074 gm were added with continued stirring for 10 minutes. TEGO betaine CKD, 2.14 gm, was added and stirred for 5 minutes. Calcium sulfate, anhydrous, 3.88 gm, was added with stirring for 5 minutes followed by addition of sodium dihydrogen phosphate, anhydrous, micronized, 0.696 gm, was added with continued stirring for 5 minutes.

In multiple increments, Sident 22S, 26.0 gm was added with stirring 2 minutes between each addition. When the Sident 22S was all added, then Zeodent 113, 10.72 gm added with stirring continuing for 15 minutes after which vanillamint, 2.14 gm, spearmint, 1.07 gm, TiO₂, 2.14 gm, ICE1500, 0.16 gm, and Multisensate flavor, 0.250 gm, were added with continued stirring for 5 minutes. The stirrer was removed and the toothpaste was added to dispensing tubes. When used as a toothpaste, a pleasant, refreshing mouthfeel and very little metallic taste is perceived.

Example 5

A 5 liter Ross/Olsa vacuum mixer with internal homogenizer was heated to 80 degrees C while the vessel was charged with 942.1 gm of PEG 400, 1184.45 gm of anhydrous glycerin and 1443.5 gm of 1.64% stannous fluoride/glycerin. Anchor stirring at slow speed was begun and continued for 7 minutes. ULTRAMULSION® [poloxamer 407 (80%)] and 2.5 million cs polydimethylsiloxane (20%), 64.5gm, was added with homogenizer speed adjusted to 2500 rpm for 15 minutes. The anchor stirrer was increased to medium speed and Gantrez MS-955, 107.5 gm, was added with stirring and homogenizing for 5 minutes. Crodasinic L, 53.5 gm, was added with continued stirring for 5 minutes. TEGO Betaine CKD, 53.5 gm, was added and stirring continued for 5 minutes. Micronized (20 micron D50) calcium sulfate, anhydrous, 97 gm, was added with stirring for 5 minutes. Micronized (20 micron) sodium phosphate monobasic,

anhydrous, 17.4 gm, was added with stirring for 5 minutes. Sident 22S, 650 gm, was added in increments at 2 minutes between additions until all was added. Stirring was continued for 15 minutes.

- 5 Vanillamint flavor, 33 gm, and spilanthus extract, 4.55 gm, were added with continued stirring for 5 minutes. The vessel was cooled to ambient temperature over 15 minutes. The contents were dispensed into tubes for use. Upon dispensing, the toothpaste indicated a pleasant taste with no stannous fluoride aftertaste. Stannous fluoride stability testing was performed on the product which exhibited a stable characteristic

10

In Vitro Testing

In vitro determination of EPF values attributed to administration of various fluoride containing: toothpastes and a test, stannous fluoride toothpastes onto human enamel subjected to acid challenge.

15

The following study was carried out according to the FDA Monograph on Anticaries Drug Products for over-the-counter, human use. The study was performed following FDA good laboratory practices.

- 20 Purpose of the following in vitro study: to determine the effect of acid challenge to human enamel treated with various fluoride containing brushing products. The effect of the acid challenge was established by measuring the resistance of enamel specimens treated with various fluoride brushing products to an acid challenge; before and after treatment with various fluoride brushing products.

25

Tooth Preparation:

- Three sound human molars were placed in a disc of red boxing wax so that only the enamel surfaces were exposed. Twelve set of three teeth each were prepared for the study. All specimens were cleaned and polished with a flour of pumice slurry and a rag wheel to remove any deposits or stains.

30

Preparation of Buffered Lactic Acid Challenge Solution:

Two moles (203.58 g of 88.5% pure lactic acid were diluted with approximately 500 ml of distilled water. To this was added a solution of 84 g NaOH dissolved in about 600 ml

of distilled water. The total volume was then adjusted to 2000 ml. This was the buffered 1.0 M lactic acid challenge solution.

Another lactic acid solution was prepared by diluting two moles lactic acid to 2000 ml with distilled water. The solution of lactic acid and sodium hydroxide was placed in a 4000 ml beaker, and pH electrodes placed in the solution. The 1.0 M lactic acid solution was used to adjust the pH of the buffered solution to 4.5. To obtain a 0.1 working concentration (for all decalcifications) the 1.0 M buffer was diluted by a factor of 10 with distilled water. Another lactic acid solution was prepared by diluting two moles lactic acid to 2000 ml with distilled water. The solution of lactic acid and sodium hydroxide was placed in a 4000 ml beaker, and pH electrodes placed in the solution. The 1.0 M lactic acid solution was used to adjust the pH of the buffered solution to 4.5. To obtain a 0.1 working concentration (for all decalcifications) the 1.0 M buffer was diluted by a factor of 10 with distilled water.

15

Deprotection:

Before every use, any residual anti-solubility protection afforded by the previous treatment was eliminated. Deprotection of these specimens was accomplished by etching the teeth in the above prepared 0.1 M lactate buffer solution for a two-hour period. Each disc of three specimens was agitated (450 rpm) in about 50 ml of lactate buffer at room temperature during the deprotection period. The teeth were rinsed well with distilled water immediately following deprotection.

Pre-Treatment Etch:

The test was performed using preheated (37°C) tooth sets and lactate buffer. The deprotected tooth sets were mounted on 1/4 inch diameter acrylic rods with molten red boxing wax. Multiplaced stirrers were used for treatments and the etches. All slurries and solutions were pre-heated to 37°C. The actual treatments and etches were carried out on the bench top with the preheated solutions. Plastic specimen containers (120 ml) were used for the etching procedure. A 1/4 inch hole was drilled in each container lid to accommodate the plastic rod to which the tooth sets were mounted. A 40 ml portion of 0.1 M lactic acid buffer was placed in each container along with a one-inch magnetic stirring bar. The rod of the first tooth set was pushed through the hole in the lid, placed in the first container and adjusted so that all enamel surfaces were immersed into the buffer solution. The container was then placed on the first magnetic stirrer and stirring

35

was begun. The timer was started at this time. At 30-second intervals the other tooth sets were started in the same manner. After 15 minutes of exposure to the buffered lactate solution, the first set was stopped and the lid and tooth set immediately removed from the container and placed in a tray of distilled water to terminate etching. The other sets were similarly removed at 30 second intervals in the same order that they were initiated and the lactate buffer solutions was retained for phosphorus analysis. The tooth sets were placed back in the 37°C water bath in preparation for the fluoride treatment step.

10 **Treatment:**

The treatments were performed using slurries of the fluoride brushing products. The slurries consisted of 1 part fluoride brushing product and 3 parts preheated (37°C) distilled water (9g:27ml). Each slurry was mixed for exactly one minute after adding the water. The slurries were NOT prepared ahead of time. They were NOT centrifuged. All tooth sets were treated at the same time (one for each fluoride brushing product). The treatment procedure was similar to the etching procedure with the exception of the slurry in place of the acid. A 30 ml portion of the preheated fluoride brushing slurry was added to the first tooth set, the teeth were immersed in the slurry and the container placed on the first stirrer. The stirrer and timer were started. At 90-second intervals (to allow time for stirring), the other tooth sets were started in the same manner. At the end of the five minutes of treatment, the first set was stopped, the tooth set removed and rinsed well with distilled water. The other sets were removed at 90-second intervals and rinsed well. The treatment fluoride brushing slurries were discarded.

25 **Post-Treatment:**

A second acid exposure was then performed by the same method as the pre-treatment etch and the lactate buffer solutions were again retained for phosphorus analysis. The pre and post-treatment solutions were analyzed using a Klett-Summerson Photoelectric Colorimeter.

30

Repeat Analyses:

The tooth sets were deprotected and the procedure repeated additional times so that each fluoride brushing product was treated and assayed on each tooth set. The treatment design was a Latin Square design so that no treatment followed another treatment consistently.

35

Calculation of Enamel Solubility Reduction:

The percent of enamel solubility reduction was computed as the difference between the amount of phosphorus in the pre and post acidic solutions, divided by the amount of phosphorus in the pre solution and multiplied by 100.

Treatment Groups:

- A. Placebo (deionized water)
- B. Positive control, MI Paste Plus with fluoride @ 900 ppm fluoride
- 10 C. Positive control, ClinPro® 5000 Toothpaste with sodium fluoride at 5000 ppm fluoride
- D. Test, stannous fluoride toothpaste at 1150 ppm fluoride composition as described hereinafter in Example 1.

Statistical Analyses:

Statistical analyses of the individual means were performed with a one-way analysis of variance model using Sigma Stat (3.1) Software. Since the ANOVA indicated significant differences, the individual means were analyzed by the Student Newman-Keuls (SNK) test.

20

Results and Discussion:

The deionized water negative control was not effective in reducing enamel solubility. The positive fluoride containing controls and the test toothpaste were significantly more effective than the deionized water negative control. The Clinpro 5000® toothpaste was slightly, significantly more effective than the negative control. The test toothpaste was dramatically superior to the two positive controls (by a factor of 10).

25

The results are shown in Table 2 below:

30

Table 2

ENAMEL SOLUBILITY REDUCTION**Summary of Results**

Group	Treatment	Pre-Etch μP	Post-Etch μP	Delta μP	Percent Reduction	EPF **
A	Deionized Water	680 \pm 16*	780 \pm 19	-100 \pm 15	-15.03 \pm 2.35	--
B	MI Paste Plus (@ 900 ppm fluoride)	677 \pm 28	697 \pm 19	-20 \pm 20	3.92 \pm 2.83	0.4
C	Clinpro 5000® (sodium fluoride @ 5000 ppm)	702 \pm 20	582 \pm 14	120 \pm 9	16.96 \pm 0.83	0.34
D	Test Toothpaste (stannous fluoride @ 1150 ppm fluoride)	717 \pm 18	359 \pm 11	358 \pm 18	49.71 \pm 1.62	4.3
	See Example 1					

5

* Mean \pm SEM (N=12)

** To establish the enamel protection factor (EPF) values for each fluoride brushing product tested, the percent reduction in enamel solubility was divided by the fluoride level in parts per million of the brushing product tested. The resultant number was multiplied by 100.

See Fig. 1 of the Drawings.

***In vitro* determination of ERF values attributed to administration of various fluoride containing toothpastes and a test toothpaste onto incipient enamel lesions in bovine enamel.**

The following study was carried out according to the FDA Monograph on Anticaries Drug Products for over-the-counter, human use, following FDA good laboratory practices.

Purpose of the following *in vitro* study: to determine the fluoride uptake into incipient enamel lesions in bovine incisors, treated with various fluoride containing, brushing products.

25

The test procedure was identical to the procedure identified as Procedure 40 in the FDA anticaries Monograph, except the lesions were formed using a solution comprising 0.1 M lactic acid and 0.2% Carbopol 907, wherein the solution was saturated with HAP(hydroxyapatite) at a pH of 5.0.

5

The fluoride uptake was established by analyzing fluoride and calcium levels of enamel pre-treatment and enamel post-treatment to determine the change in enamel fluoride attributed to treatment with fluoride containing brushing products.

10 **Procedure:**

Sound, upper, central, bovine incisors were selected and cleaned of all adhering soft tissue. A core of enamel 3mm in diameter was prepared from each tooth by cutting perpendicularly to the labial surface with a hollow-core diamond drill bit. This was performed under water to prevent overheating of the specimens. Each specimen was
15 embedded in the end of a plexiglass rod (1/4" diameter x 2" long) using methylmethacrylate. The excess acrylic was cut away exposing the enamel surface. The enamel specimens were polished with 600 grit wet/dry paper and then micro-fine Gamma Alumina. The resulting specimens were a 3mm disk of enamel with all but the exposed surface covered with acrylic. Twelve specimens per group were prepared.

20

Each enamel specimen was then etched by immersion into 0.5 ml of 1M HClO₄ for 15 seconds. Throughout the etch period, the etch solutions were continuously agitated. A sample of each solution was then buffered with TISAB (fluoride ion probe buffer) to a pH of 5.2 (0.25 ml sample, 0.5 ml TISAB and 0.25 ml 1N NaOH) and the fluoride
25 content of the solution determined by comparison to a similarly prepared standard curve (1 ml std + 1 ml TISAB). For use in depth of each calculation, the Ca content of the etch solution was determined by taking 50 µl and analyzing for Ca by atomic absorption (0.05 ml qs to 5 ml). These data was the indigenous fluoride level of each specimen prior to treatment.

30

The specimens were once again ground and polished as described above. An incipient lesion was formed in each enamel specimen by immersion into a 0.1M lactic acid/0.2% Carbopol 907 solution for 24 hours at room temperature. These specimens were then rinsed well with distilled water and stored in a humid environment until used.

The treatments were performed using slurries of the various fluoride containing brushing products. The slurries consisted of 1 part fluoride containing brushing product and 3 parts distilled water (9g:27ml). Each slurry was mixed for exactly one minute after adding the water. The slurries were NOT prepared ahead of time. They were NOT centrifuged. The 12 specimens of each group were then immersed into 25 ml of their assigned slurry with constant stirring (350 rpm) for 30 minutes. Following treatment, the specimens were rinsed with distilled water. One layer of enamel was then removed from each specimen and analyzed for fluoride and calcium as outlined above (i.e. 15 second etch). The pretreatment fluoride (indigenous) level of each specimen was then subtracted from post treatment, fluoride value to determine the change in enamel fluoride due to the last treatment.

Statistical Analyses:

The results are shown on attached tables. All raw data (individual specimen Enamel Fluoride Uptake (EFU) values are reported. In addition, the mean, S.D.(standard deviation) and SEM(scanning electron micrograph) for each group was calculated. Statistical analysis were performed by a one-way analysis of variance model using Sigma Stat Software (3.1). Since significant differences are indicated, the individual means were analyzed by the Student Newman Keuls (SNK) test.

Test Products:

The test fluoride containing toothpastes were coded as follows:

1. The test fluoride containing toothpastes were coded as follows:
2. Placebo (deionized water)
3. Positive Control MI Paste Plus with 900 ppm fluoride
4. Positive Control 2 Clinpro 5000® with sodium fluoride at 5000 ppm fluoride
5. Test Toothpaste with stannous fluoride at 1150 ppm fluoride composition as described in Example 1.

Results:

The results are shown in Table 3 below:

Table 3

Change in Incipient Lesion Enamel Fluoride Content

Fluoride Containing Brushing Product	Enamel Fluoride Concentration (ppm)				ERF **
	Pre Treatment	Post Treatment	Increase	Etch Depth	
Placebo, deionized water	39 ± 3*	47 ± 4	8 ± 3	16.19 ± 0.45	--
MI Paste Plus @ 900 ppm fluoride	38 ± 3	323 ± 9	285 ± 8	15.84 ± 0.42	31
OTC SnF ₂ Paste @ 5000 ppm sodium fluoride	37 ± 3	1716 ± 22	1679 ± 23	11.30 ± 0.10	34
Clinpro 5000® @ 5000 ppm sodium fluoride	37 ± 3	2616 ± 54	2579 ± 54	14.55 ± 0.36	51
Test Toothpaste @ 1150 ppm stannous fluoride	39 ± 2	6459 ± 451	6420 ± 451	13.23 ± 0.31	540
See Example 1					

5 * Mean ± SEM (N=12)

** To establish the enamel repair factor (ERF) values for each fluoride brushing product tested, the increase in enamel fluoride concentration was divided by the fluoride level of the fluoride brushing product. The resultant number was multiplied by 100.

10

See Fig. 2 of the Drawings.

15

In vitro determination of ERF values attributed to administration of various fluoride containing: toothpastes and a test toothpaste onto incipient enamel lesions in human enamel was carried out following the Enamel Fluoride protocol described above for the results reported in Table 2. Some of the fluoride containing brushing products tested in this fluoride uptake study are different than those reported on in Table 3 above.

The results are shown in Table 4 below:

20

Table 4

Change in Incipient Lesion Enamel Fluoride Content

Fluoride containing Brushing Product		Pre-Treatment		Post-Treatment			ERF*
		Enamel Fluoride Concentration	Depth of Etch (microns)	Enamel Fluoride Concentration	Depth of Etch (microns)	Change in Depth of Etch	
Crest® PRO- HEALTH® Toothpaste (Stannous Fluoride, 1100 ppm fluoride)	Tooth 1	31	45	692	30	33	
	Tooth 2	25	76	1111	35	55	
	Tooth 3	134	31	1320	14	54	
	Average			1008		47	92
	Std Dev			314		12	
Toothpaste with Sodium Fluoride @ 900 ppm fluoride	Tooth 1	95	80	679	21	74	
	Tooth 2	124	72	654	23	68	
	Tooth 3	207	41	1140	13	68	
	Average			824		70	91
	Std Dev			273		3	
Test Toothpaste (Stannous Fluoride, 1100 ppm fluoride)	Tooth 1	165	44	438	23	47	
	Tooth 2	93	61	388	26	57	
	Tooth 3	104	56	723	28	50	
	Average			516		51	50
	Std Dev			181		5	
See Example 2							

- 5 *To establish the enamel repair factor (ERF values for each fluoride brushing product tested, the average increase in Enamel Fluoride Concentration (post treatment) was divided by the fluoride level of the fluoride brushing product tested. The resultant number was multiplied by 100.
- 10 See Fig. 3 of the Drawings.

Table 5

Change in Incipient Lesion Enamel Fluoride Content

Fluoride Containing Brushing Product	Enamel Fluoride Concentration (ppm)				
	Pre Treatment	Post Treatment	Increase	Etch Depth	ERF **
Placebo, deionized water	41 ± 4*	89 ± 5	47 ± 5	17.88 ± 0.70	--
Crest® PRO-HEALTH® Toothpaste (stannous fluoride, 1100 ppm fluoride)	44 ± 5	1530 ± 69	1486 ± 66	10.06 ± 0.25	140
Test Toothpaste (stannous fluoride at 1148 ppm fluoride)	47 ± 5	4857 ± 338	4815 ± 338	13.36 ± 0.26	420
See Example 4					

* Mean ± SEM (N=12)

- 5 ** To establish the enamel repair factor (ERF) values for each fluoride brushing product tested, the average increase in enamel fluoride concentration (post treatment was divided by the fluoride level of the fluoride brushing product tested. The resultant number was multiplied by 100).
- 10 See Fig. 4 of the Drawings

Table 6

Change in Incipient Lesion Enamel Fluoride Content

Fluoride Containing Brushing Product	Enamel Fluoride Concentration (ppm)				
	Pre Treatment	Post Treatment	Increase	Etch Depth	ERF **
Placebo, deionized water	41 ± 4*	89 ± 5	47 ± 5	17.88 ± 0.70	--
Crest® PRO-HEALTH® Toothpaste (stannous fluoride, 1100 ppm fluoride)	44 ± 5	1530 ± 69	1486 ± 66	10.06 ± 0.25	140
Test Toothpaste (stannous fluoride at 1148 ppm fluoride)	52±6	1948±68	1902±70	15.28±0.56	166
See Example 3					

15

* Mean ± SEM (N=12)

** To establish the enamel repair factor (ERF) values for each fluoride brushing product tested, the average increase in enamel fluoride concentration (post treatment was divided by the fluoride level of the fluoride brushing product tested. The resultant number was multiplied by 100).

5

See Fig. 5 of the Drawings

Table 7

10

Illustrative Examples of Toothpastes

Example No.	Stannous Fluoride level in ppm fluoride	Calcium and phosphate components, % by wt. of Toothpaste	Substantivity Agent, Composition and % by wt. of Toothpaste	pH of GEL in saliva
7	970	Calcium Sulfate, 2.0% NaH ₂ PO ₄ , 0.35%	F-127/2.5mmCS PDMS(30%) 1.75%	5.8
8	900	Calcium Sulfate, 2.2% NaH ₂ PO ₄ , 0.35%	F-108/12,500CS PDMS(10%) 2.0%	6.2
9	1000	Calcium Sulfate, 2.5% NaH ₂ PO ₄ , 0.25%	F-127/600KCS PDMS(30%) 2.2%	5.5
10	1100	Calcium Sulfate, 2.8% NaH ₂ PO ₄ , 0.38%	F-127/2.5mmCS PDMS(10%) 1.65%	4.8
11	970	Calcium fumarate, 3.0% NaH ₂ PO ₄ , 0.33%	F-108/600KCS PDMS(20%) 1.4%	5.7
12	900	Calcium fumarate, 2.8% NaH ₂ PO ₄ , 0.30%	F-108/2.5mmCS PDMS(20%) 2.3%	6.3
13	1000	Calcium fumarate, 1.5% NaH ₂ PO ₄ , 0.15%	F-127/12,500CS PDMS(30%) 1.3%	5.6
14	1148	Calcium sulfate, 1.94% NaH ₂ PO ₄ , 0.348%	F-127/2.5mmCS PDMS(20%) 1.29%	4.5
15	970	Calcium gluconate, 2.2% NaH ₂ PO ₄ , 0.25%	F-108/2.5mmCS PDMS(26%) 2.0%	5.9
16	900	Calcium fumarate, 2.4% NaH ₂ PO ₄ , 0.25%	F-127/2.5mmCS PDMS(15%) 2.2%	6.3

* Mean \pm SEM (N=12)

15 ** To establish the enamel repair factor (ERF) values for each fluoride brushing product tested, the average increase in enamel fluoride concentration (post treatment

was divided by the fluoride level of the fluoride brushing product tested. The resultant number was multiplied by 100).

5

Table 8
Change in Incipient Lesion Enamel Fluoride Content

"At-risk" Patient	Specific Toothpaste Recommended	Frequency of Treatment	Duration of Treatment
Caries	Stannous Fluoride: 970 ppm fluoride PDMS/Surfactant Emulsion: PDMS 800,000 cs Calcium and Phosphate component: 3.2% by wt. EPF: at least about 1.8 ERF: at least about 140 pH= 4.9	Once daily for 2 minutes followed by expectorating	To be determined by oral care professionals
Undergoing medical and/or professional oral care treatments with prescribed medications	Stannous Fluoride: 1150 ppm fluoride PDMS/Surfactant Emulsion: PDMS 800,000 cs Calcium and Phosphate component: 3.8% by wt. EPF: at least about 2.2 ERF: at least about 180 pH= 3.8	At least twice daily	Duration of medical/professional oral care treatment
Immunocompromised with chronic conditions	Stannous Fluoride: 900 ppm fluoride PDMS/Surfactant Emulsion: PDMS 600,000 cs Calcium and Phosphate component: 3.4% by wt. EPF: at least about 2.4 ERF: at least about 165 pH= 4.7	At least three times daily	Duration of medical/professional oral care treatment

Table 8 (cont'd)

Diabetes, heart disease, etc.	Stannous Fluoride: 1100 ppm fluoride PDMS/Surfactant Emulsion: PDMS 2.5mm CS Calcium and Phosphate component: 3.0% by wt. EPF: at least about 2.1 ERF: at least about 230 pH= 4.6	At least twice daily	To be determined by oral care professionals
Cancer treatment	Stannous Fluoride: 1150 ppm fluoride PDMS/Surfactant Emulsion: PDMS 2.5mm CS Calcium and Phosphate component: 3.2% by wt. EPF: at least about 2.3 ERF: at least about 200 pH= 4.3	At least twice daily	Duration of medical/professional oral care treatment

* Mean \pm SEM (N=12)

- 5 ** To establish the enamel repair factor (ERF) values for each fluoride brushing product tested, the average increase in enamel fluoride concentration (post treatment was divided by the fluoride level of the fluoride brushing product tested. The resultant number was multiplied by 100).

10 **Discussion of EPF and ERF values established by *in vitro* testing**

Preferred, stannous fluoride toothpastes of the present invention show substantial improvement in EPF and ERF values compared to commercial toothpastes at various fluoride loads as reported in Tables 2 and 8 and Figs. 1 and 5 of the Drawings.

- 15 For example, the ERF values, reported in Table 2 and Fig. 2 for MI Paste Plus, OTC Stannous Fluoride paste and Clinpro 5000® Toothpaste are 31, 34 and 50, respectively, compared to an ERF for the stannous fluoride toothpaste of the present invention described in Example 1 of 540. This 10X plus improvement in ERF value over commercial toothpastes represents a major advance in fluoride uptake and enamel

hardening effectiveness. Such an advantage in enamel hardening efficiency is particularly critical to patients experiencing: rampant caries, coronal caries, cancer therapy treatments, mucositis treatments, immune deficiency treatments, bone marrow transplants, etc.

5

Proposed Mechanism of Action of the compositions of the present invention

The enamel protectant and/or enamel repair (EPF and/or ERF) data reported for formulations described in Examples 1 through 5, as detailed in Tables 1 through 8 and in Figs. 2 through 5 of the Drawings, suggest the substantivity of stannous fluoride and calcium to the biofilm present on enamel surfaces is enhanced by a shift from bidentate to monodentate calcium binding in the presence of stannous fluoride. This shift in calcium binding in the presence of stannous fluoride results in a most effective binding site configuration. See:

15 The enamel protectant and/or enamel repair (EPF and/or ERF) data reported for formulations described in Examples 1 through 5, as detailed in Tables 1 through 8 and in Figs. 2 through 5 of the Drawings, suggest the substantivity of stannous fluoride and calcium to the biofilm present on enamel surfaces is enhanced by a shift from bidentate to monodentate calcium binding in the presence of stannous fluoride. This shift in calcium binding in the presence of stannous fluoride results in a most effective binding site configuration. See:

- Dudey and Lim, J. Phys. Chem B, 2004, 108:4546.
Vogel, et. al., Caries Res., 2010, 94:108-115.
25 Rolla and Bowen, Scand. J. Dent. Res., 1977; 85:149-151.
Rose, et. al., J. Dent. Res., 1993; 72:78-84.

See also:

- Turner, et. al., Ceramics, Silikaty 57(1):1-6 (2013)
30 Mohammed, et. al., Caries Res., 47:421-428 (2013)

Calcium-bridge, fluoride binding

Calcium binding to biofilm shifts from a bidentate chelation to a monodentate chelation in the presence of fluoride, freeing up Calcium+ to bind with fluoride, CaF+ pairs, thereby doubling the calcium binding capacity.

5

Stable fluoride in the toothpastes of the present invention produces marked reduction in calcium binding affinity and approximately doubles calcium binding capacity. In the absence of fluoride, calcium binding to biofilm is bidentate. Stable fluoride in the toothpastes of the present invention compete with biofilm causing calcium binding to biofilm to become monodentate. This allows the binding of about double the quantity of calcium and of CaF+ bound to biofilm. Release of fluoride bound by calcium bridging into biofilm fluid as a result of fluoride clearance into saliva will be accompanied by a corresponding release of calcium which, in turn, potentiates the cariostatic effect of fluoride as indicated in the in vitro testing described in Tables 2 through 8 and Figs. 1 through 5 of the Drawings.

At least some of the stable fluoride present in the toothpastes of the present invention is bound to calcium ions (sourced from various calcium salts in the present invention and/or calcium present in the copolymer substantivity enhancer, such as Gantrez® MS-955). These calcium ions, in turn, are bound to biofilm associated with enamel.

A drop in pH follows exposure of plaque to sucrose which removes some anionic groups by neutralization, thereby liberating calcium and fluoride (as CaF+) at the very sites where these moieties can do the most good.

25

The effectiveness of the toothpastes of the present invention depends on three factors:

substantivity of the formulation to biofilm,
stannous fluoride as the source of CaF+, and
retention of fluoride in a form on biofilm which allows release of CaF+ ions into hydroxyapatite.

Stannous fluoride produces a marked reduction in calcium binding affinity accompanied by an approximate doubling of the calcium binding capacity. In the absence of fluoride, divalent cation binding to plaque is bidentate. Fluoride competes with macromolecular

35

anionic groups, causing binding to become monodentate. Release of fluoride formed by calcium bridging, is accompanied by release of calcium, which potentiates the cariostatic effect of fluoride.

- 5 The presence CaF^+ is required to deliver the enamel protection and repair (EPF and ERF) results required for the toothpaste formulations of the present invention.

Summary as to the role of cation bridging in microbial fluoride binding

- 10 Fluoride binding produces a marked reduction in calcium binding affinity, along with a doubling of calcium binding capacity. This indicates that calcium binding changes from bidentate to monodentate. This shift from bidentate to monodentate is a consequence of fluoride replacing an anionic group as one of the calcium ligands.

- 15 The anionic groups to which calcium is no longer bound are then free to bind a CaF^+ ion pair, resulting in a doubling of the calcium binding capacity. Release of fluoride, bound by calcium bridging into plaque fluid, may be accompanied by a release of calcium which will potentiate the cariostatic effect of fluoride.

- 20 CaF^+ is taken up by hydroxyapatite and is responsible for the EPF and ERF in vitro data reported for the compositions of the present invention. The ERF values reported in Tables 2 through 8 and Figures 2 through 5 of the Drawings suggest that the CaF^+ moiety is incorporated into the hydroxyapatite lattice during remineralization.

- 25 The toothpastes of the present invention set a new, oral care STANDARD for Enamel Protection and Enamel Repair, while dramatically reducing exposure to elevated levels in various fluoride varnishes, gels and toothpastes.

CLAIMS

1. Aqueous-free, enamel protectant and enamel repair toothpastes containing: stannous fluoride and calcium in a substantivity agent selected from emulsions of polydimethylsiloxanes in nonionic poloxamer surfactants, wherein:
- (a) substantivity of said stannous fluoride and calcium into biofilm present on enamel is enhanced through a shift in calcium binding from bidentate to monodentate in the presence of stannous fluoride; and
 - (b) an enamel protectant factor (EPF) value of at least 2.5 and an enamel repair factor (ERF) value of at least 200 are achieved with periodic administration of said toothpastes onto enamel surfaces with biofilm present;
 - (c) wherein the stannous fluoride is present in an amount of between about 850 ppm and about 1500 ppm;
 - (d) wherein the toothpastes have a pH of at least 3;
 - (e) wherein the calcium content is 0.5% to 5.0% by weight; and
 - (f) wherein the calcium is selected from the group consisting of calcium fumarate, calcium sulfate, calcium gluconate, mixed sodium and calcium salts of methyl/vinyl/ether/maleic copolymers and combinations thereof.
2. The toothpastes according to Claim 1, wherein some of said stannous fluoride and calcium is present as CaF^+ .
3. The toothpastes according to Claim 1, wherein said substantivity agent is combined with linear, polymeric polycarboxylate substantivity enhancers.
4. The toothpastes according to Claim 3, wherein the substantivity agent comprises an emulsion of a discontinuous phase of polydimethylsiloxane polymer at viscosities between about 10,000 cs and about 2.5 million cs in a continuous phase of the nonionic poloxamer surfactants.
5. The toothpastes, according to Claim 4, wherein the nonionic poloxamer surfactants, suitable for said toothpastes, are represented by the structural formula:
- $$\text{HO}[(\text{C}_2\text{H}_4\text{O})_x / (\text{C}_3\text{H}_6\text{O})_y]_z - [(\text{C}_2\text{H}_4\text{O})_x / (\text{C}_3\text{H}_6\text{O})_y]_n\text{H}$$
- wherein the sum of x, y and z is between 125 and 175.

6. The toothpastes according to Claim 4, wherein the nonionic poloxamer surfactants, suitable for said toothpastes, are represented by the structural formula:



wherein $x = 76$, $y = 25$ and $z = 56$.

5

7. The toothpastes, according to Claim 1, containing unreacted calcium and phosphate components.

8. Aqueous-free, enamel protectant and enamel repair toothpastes containing stannous fluoride and calcium in a substantivity agent comprising an emulsion of polydimethylsiloxane polymer at viscosities between about 10,000 cs and about 2.5 million cs as a discontinuous phase in a continuous phase of a nonionic poloxamer surfactant containing a linear polymeric polycarboxylate substantivity enhancer, wherein the nonionic poloxamer surfactant is selected from the group consisting of surfactants having the following general structural formula:.

15



wherein the sum of x , y and z is between 125 and 175; with an enamel protectant factor (EPF) value of at least 2.5 and an enamel repair factor (ERF) value of at least 200, wherein:

20

- (a) substantivity of said stannous fluoride and calcium into biofilm present on enamel is enhanced through a shift in calcium binding from bidentate to monodentate in the presence of stannous fluoride; and
- (b) wherein the stannous fluoride is present in an amount of between about 850 ppm and about 1500 ppm;
- (c) wherein the toothpastes have a pH of at least 3;
- (d) wherein the calcium content is 0.5% to 5.0% by weight; and
- (e) wherein the calcium is selected from the group consisting of calcium fumarate, calcium sulfate, calcium gluconate, mixed sodium and calcium salts of methyl/vinyl/ether/maleic copolymers and combinations thereof.

30

9. The toothpastes according to Claim 8, wherein the nonionic poloxamer surfactants, suitable for said toothpastes, are represented by the structural formula:



wherein the sum of x , y and z is between 125 and 150.

35

10. The toothpastes according to Claim 8, wherein the nonionic poloxamer surfactants, suitable for said toothpastes, are represented by the structural formula:



wherein $x = 76$, $y = 0$ and $z = 56$.

5

11. The toothpastes according to Claim 8 containing unreacted calcium and phosphate components.

12. Aqueous-free, enamel protectant and enamel repair toothpastes, featuring monodentate-bidentate bonding of calcium in the presence of stannous fluoride; containing: stannous fluoride, calcium and phosphate components, and a substantivity agent emulsion containing nonionic poloxamer surfactant as a continuous phase and polydimethylsiloxane polymer as a discontinuous phase; with an enamel protectant factor (EPF) of at least 2.5 and an enamel repair factor (ERF) of at least 200, wherein

- 15 (a) substantivity of said stannous fluoride and calcium into biofilm present on enamel is enhanced through a shift in calcium binding from bidentate to monodentate in the presence of stannous fluoride;
- (b) wherein the stannous fluoride is present in an amount of between about 850 ppm and about 1500 ppm;
- 20 (c) wherein the toothpastes have a pH of at least 3;
- (d) wherein the calcium content is 0.5% to 5.0% by weight; and
- (e) wherein the calcium is selected from the group consisting of calcium fumarate, calcium sulfate, calcium gluconate, mixed sodium and calcium salts of methyl/vinyl/ether/maleic copolymers and combinations thereof.

25

13. The toothpastes, according to Claim 5, wherein said polydimethylsiloxane polymer discontinuous phase comprises up to 20% by weight of said emulsion.

14. Aqueous-free, enamel protectant and enamel repair toothpastes containing stable stannous fluoride and calcium on a substantivity agent substantive to biofilm coated enamel, wherein the substantivity agent comprises nonionic poloxamer surfactants and polydimethylsiloxane polymers, wherein:

30


- (a) substantivity of said stannous fluoride and calcium into said biofilm is enhanced by a substantivity enhancer and by a shift from bidentate binding of calcium to monodentate in the presence of stable stannous fluoride; and
- (b) said stannous fluoride, upon release from said substantivity agent, converts to the moiety CaF^+ which effects an enamel protectant factor (EPF) value of at least 2.5 and an enamel repair factor (ERF) value of at least 200;
- (c) wherein the stannous fluoride is present in an amount of between about 850 ppm and about 1500 ppm;
- (d) wherein the toothpastes have a pH of at least 3;
- (e) wherein the calcium content is 0.5% to 5.0% by weight; and
- (f) wherein the calcium is selected from the group consisting of calcium fumarate, calcium sulfate, calcium gluconate, mixed sodium and calcium salts of methyl/vinyl/ether/maleic copolymers and combinations thereof.

15. The toothpastes according to Claim 1, wherein said stannous fluoride, upon release from said substantivity agent, includes a CaF^+ moiety in the presence of said calcium monodentate binding to said biofilm present on enamel.

16. The toothpastes according to Claim 1, wherein the pH of said toothpastes, upon administration to enamel with biofilm, is about 3.

17. The toothpastes according to Claim 3, wherein the level of said substantivity agent and said substantivity enhancers is between about 0.5 and about 5% by wt. and about 0.1 and about 3% by wt., respectively.

18. The toothpastes, according to Claim 3, wherein said substantivity enhancer has the structural formula:



where n and M are integers that provide a molecular weight between about 60,000 and about 1,000,000.

19. The toothpastes according to Claim 14, wherein the substantivity agent comprises an emulsion of polydimethylsiloxane as a discontinuous phase in a continuous phase of the nonionic poloxamer surfactant and wherein the polydimethylsiloxane is at a concentration of up to about 20% by wt. of said emulsion.
- 5
20. The toothpastes according to Claim 14 containing unreacted calcium and phosphate components, wherein: the ratio of unreacted calcium and phosphate components is between about 2 and about 1, and the level of calcium and phosphate mixture in said toothpastes ranges between about 0.2% and about 3% by weight.
- 10
21. The toothpastes according to Claim 19, wherein the substantivity agent is combined with a linear polymeric polycarboxylate substantivity enhancer, wherein said emulsion comprises between about 0.5 and about 3.0 wt. percent of said toothpastes and said substantivity enhancer comprises between about 0.5 and about 2.8 wt. percent of said toothpastes.
- 15
22. The toothpastes according to Claim 14, wherein the pH of said toothpaste, upon exposure to saliva, is between about 4 and about 8.
- 20
23. Use of the toothpastes according to any one of claims 1 to 22 for treating enamel of "at-risk" patients wherein said toothpastes are adapted for administration at a frequency sufficient to effect an EPF value of at least 2.5 and an ERF value of at least 200.
- 25
24. The use according to claim 23 wherein the "at-risk" patients are selected from the group consisting of immunocompromised, cancer therapy, diabetes, COP, mucositis and cardiovascular patients.
- 30
25. The use according to claim 23 or 24 wherein the toothpastes are adapted for administration for up to two minutes, followed by expectoration.
26. The use according to claim 23 or 24 wherein the toothpastes are adapted for administration at least twice daily.

Figure 1
Enamel Protection Factor (EPF)
for three fluoride brushing products at fluoride levels
ranging from 800 ppm stannous fluoride to 5000 ppm sodium fluoride




Figure 2
Enamel Repair Factor (ERF)
for three fluoride brushing products at fluoride levels
ranging from 1150 ppm stannous fluoride to 5000 ppm sodium fluoride




Figure 3
Enamel Repair Factor (ERF)
for three fluoride brushing products at fluoride levels
ranging from 900 ppm sodium fluoride to 1148 ppm stannous fluoride




Figure 4
Enamel Repair Factor (ERF)
for two fluoride brushing products at fluoride levels
ranging from 1100 and 1148 ppm stannous fluoride




Figure 5
Enamel Repair Factor (ERF)
for two fluoride brushing products at fluoride levels
ranging from 1100 and 1148 ppm stannous fluoride

