



(51) International Patent Classification:

C07C 67/08 (2006.01) C07C 201/08 (2006.01)
C07C 69/76 (2006.01) C07C 63/04 (2006.01)

Published:

— with international search report (Art. 21(3))

(21) International Application Number:

PCT/EP2018/069192

(22) International Filing Date:

16 July 2018 (16.07.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

17181783.6 18 July 2017 (18.07.2017) EP
62/533,712 18 July 2017 (18.07.2017) US

(71) Applicant: LONZA LTD [CH/CH]; Lonzastrasse, 3930 Visp (CH).

(72) Inventors: BERSIER, Michael; Haus Wildi, 3938 Ausserberg (CH). KULESZA, Anna; Eggstrasse, 3938 Ausserberg (CH). HANSELMANN, Paul; Holzachra 25, 3902 Brig-Glis (CH). AELLIG, Christof; Bubenbergrasse 15, 3700 Spiez (CH). FILLIGER, Sarah; Landstrasse 47, 3904 Naters (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

(54) Title: METHOD FOR PREPARATION OF 5-FLUORO-2-METHYL-3-NITROBENZOIC ACID AND ITS METHYL ESTER

(57) Abstract: The invention discloses a method for preparation of 5-fluoro-2-methyl-3-nitrobenzoic acid and its methyl ester by conversion of 5-fluoro-2-methylbenzoic acid with fuming nitric acid and oleum and subsequent conversion with methanol.




METHOD FOR PREPARATION OF 5-FLUORO-2-METHYL-3-NITROBENZOIC ACID AND ITS METHYL ESTER

The invention discloses a method for preparation of 5-fluoro-2-methyl-3-nitrobenzoic acid and its methyl ester by conversion of 5-fluoro-2-methylbenzoic acid with fuming nitric acid and oleum and subsequent conversion with methanol.

BACKGROUND OF THE INVENTION

Rucaparib, also known as CO-338, is an inhibitor of poly ADP-ribose polymerase (PARP) inhibitor.



Rucaparib is being developed for the treatment of patients with cancers pre-disposed to PARP inhibitor sensitivity.

Gillmore et al., Org. Process Res. Dev., 2012, 16, 1897–1904, discloses the use of methyl-5-fluoro-2-methyl-3-nitro benzoate for the preparation of Rucaparib. 52% yield of a brown solid is reported for the two steps conversion of 5-fluoro-2-methylbenzoic acid to 5-fluoro-2-methyl-3-nitrobenzoic acid methyl ester by nitration with concentrated sulfuric acid and concentrated nitric acid followed by subsequent esterification.

There was a need for a process that has higher yield and higher purity.

It was found that the use of oleum and fuming nitric acid increases the yield. In addition a colorless product is obtained. Also the content of the undesired dinitro derivative is low.

The following abbreviations are used, if not otherwise stated:

eq equivalent


TFA trifluoro acetic acid

wt% percent by weight

5


SUMMARY OF THE INVENTION

Subject of the invention is a method for the preparation of compound of formula (2)



10

comprising a step STEP1, STEP1 comprises a reaction REAC1, in REAC1 compound of formula (1)



15

is converted to compound of formula (2) by the action of a mixture MIX of concentrated sulphuric acid, oleum and fuming nitric acid.

DETAILED DESCRIPTION OF THE INVENTION

20 Preferably, the concentrated sulfuric acid is conventional concentrated sulfuric acid; more preferably the concentrated sulfuric acid has a content of from 94 to 100 wt% of H₂SO₄; even more preferably of from 95 to 100 wt% of H₂SO₄; especially of from 96 to 100 wt% of H₂SO₄; the wt% based on the weight of the concentrated sulfuric acid.

25

Preferably, the oleum is conventional oleum; more preferably the oleum contains 50 to 70 wt% of SO₃;

even more preferably the oleum contains 60 to 70 wt% of SO_3 .

Preferably, the fuming nitric acid is conventional fuming nitric acid;

preferably the fuming nitric acid has a content of from 95 to 100 wt% of HNO_3 ;

5 more preferably of from 98 to 100 wt% of HNO_3 .

Preferably, MIX contains

concentrated sulphuric acid in an amount of 4 to 12 times of molar equivalents of

H_2SO_4 relative to compound of formula (1),

10 oleum in an amount of 1 to 4 molar times of equivalents of H_2SO_4 relative to compound of formula (1),

fuming nitric acid in an amount of 1 to 2 times of molar equivalents of HNO_3 relative to compound of formula (1);

more preferably, MIX contains

15 concentrated sulphuric acid in an amount of 4 to 6 times of molar equivalents of H_2SO_4 relative to compound of formula (1),

oleum in an amount of 2 to 4 molar times of equivalents of H_2SO_4 relative to compound of formula (1),

20 fuming nitric acid in an amount of 1.5 to 2 times of molar equivalents of HNO_3 relative to compound of formula (1);

preferably the amounts of concentrated sulfuric acid, oleum and fuming nitric acid in MIX add up to 95 to 100 wt%, more preferably to 98 to 100 wt%, the wt% being based on the total weight of MIX; even more preferably MIX consists of concentrated sulfuric acid, oleum and fuming nitric acid.

25

REAC1 can be done batch wise or in a continuous way.

Preferably, the reaction temperature TEMP1 of REAC1 is from -10 to 30°C , more preferably from -5 to 25°C .

30 Preferably, the reaction time TIME1 of REAC1 is from 1 sec to 2 h, more preferably from 10 min to 1.5 h.

When REAC1 is done batch wise, then

preferably, the reaction temperature TEMP1 of REAC1 is from -10 to 20°C, more preferably from -5 to 5°C.

When REAC1 is done in a continuous way, then

preferably, the reaction temperature TEMP1 of REAC1 is from 10 to 30°C, more preferably from 15 to 25°C.

When REAC1 is done batch wise, then

preferably, the reaction time TIME1 of REAC1 is from 30 min to 2 h, more preferably from 45 min to 1.5 h.

When REAC1 is done in a continuous way, then

preferably, the reaction time TIME1 of REAC1 is from 1 sec to 30 min, more preferably from 10 sec to 25 min, even more preferably from 10 sec to 10 min, especially from 10 sec to 5 min, more especially from 10 sec to 1 min.

When REAC1 is done batch wise, then preferably compound of formula (1) is dissolved in a

mixture of concentrated sulfuric acid and oleum to form a solution, and this solution is then mixed with the fuming nitric acid in order to provide for REAC1; more preferably the fuming nitric acid is added to said solution in order to provide for REAC1.

When REAC1 is done in a continuous way, then preferably compound of formula (1) is

dissolved in a mixture of concentrated sulfuric acid and oleum forming a solution SOL1; this solution is then mixed continuously with a mixture MIX1 of concentrated sulfuric acid, oleum and fuming nitric acid in order to provide for REAC1.

Preferably, SOL1 contains

concentrated sulphuric acid in an amount of 2 to 10 times, more preferably of 2 to 7.5 times, even more preferably of 3.5 to 5 times, of molar equivalents of H₂SO₄ relative to compound of formula (1),

oleum in an amount of 1 to 10 times, more preferably of 1.2 to 6 times, even more preferably of 1.8 to 3.6 times, of molar equivalents of H₂SO₄ relative to compound of formula (1).

Preferably, MIX1 contains

concentrated sulphuric acid in an amount of 0.5 to 10 times, more preferably of 0.5 to 5 times, even more preferably of 0.5 to 2.5 times, especially of 0.5 to 1 times, of molar equivalents of H₂SO₄ relative to compound of formula (1),

oleum in an amount of 0.1 to 10 times, more preferably of 0.1 to 5 times, even more preferably of 0.1 to 2 times, especially of 0.1 to 1 times, more especially of 0.2 to 0.4 times, of molar equivalents of H_2SO_4 relative to compound of formula (1), fuming nitric acid in an amount of 1.0 to 5 times, more preferably of 1.2 to 3.5 times, even more preferably of 1.5 to 2 times, of molar equivalents of HNO_3 relative to compound of formula (1).

When REAC1 is done in a continuous way, then preferably REAC1 is done in a mixing device MIXDEV, where a feed FEED1 containing compound of formula (1) is mixed with a feed FEED2 containing the fuming nitric acid, the mixing results is a reaction mixture. Preferably, FEED1 is SOL1 and FEED2 is MIX1.

MIXDEV can be any suitable installation which can be used for mixing two fluids and which is known in the state of the art, such as a common branch connection, e.g. a T or Y piece, a static mixing device or a micro reactor, preferably it is a static mixing device or a micro reactor.

Static mixing devices, e.g. static mixers, are well established and widespread in all fields of chemical process technology. It is characteristic for static mixing devices, that, in contrast to dynamic mixing devices, only the media to be mixed are in motion. The liquids or gases are mixed by pump energy only, while the geometrically strong defined mixing elements in the static mixing devices remain in position. Companies such as Fluotec, Seuzachstrasse, 8413 Neftenbach, Switzerland, or Sulzer Ltd, Neuwiesenstrasse 15, 8401 Winterthur, Switzerland, are well known suppliers among others of such static mixing devices.


Micro reactors, also called micro structured reactors, are devices in which chemical reactions take place in a confinement with typical lateral dimensions below 1 mm; the most typical form of such confinement are micro channels. A micro reactor is a continuous flow reactor. They have been successfully applied in lab, pilot and production scale. E.g. the Fraunhofer Institute for Chemical Technology ICT, Joseph-von-Fraunhofer Strasse 7, 76327 Pfinztal, Germany, develops and offers such micro reactors.

Preferably, the static mixing device has the form of a tube or a plate containing means that present obstacles for the flow of the reaction mixture and thereby effecting the mixing of the components.

Preferably, the micro reactor contains micro channels which are arranged in such a way as to effect the mixing.

After REAC1, compound of formula (2) can be isolated and purified by conventional methods, which are known to those skilled in the art. These conventional methods include quenching the reaction mixture from REAC1 with water, extraction, distillation, preferably
5 fractional distillation, which can be done under reduced pressure, crystallization, chromatography, filtration, washing or any combination of these methods.

Further subject of the invention is a method for the preparation of compound of formula (6)



comprising the STEP1 and a second step STEP2;

STEP2 comprises a reaction REAC2, in REAC2 compound of formula (2), which is obtained by REAC1 in STEP1, is reacted with methanol to provide compound of formula (6);

15 with STEP1 as defined above, also with all its embodiments.

Preferably, the molar amount of methanol in REAC2 is from 1 to 50 times, more preferably from 5 to 20 times, even more preferably from 7.5 to 15 times, of the molar amount of compound of formula (2).

Preferably, REAC2 is done in the presence of an acid ACID2, ACID2 is preferably H₂SO₄.

Preferably, the molar amount of ACID2 is from 1 to 50 times, preferably 1 to 40 times, more preferably 1 to 30 times, more preferably 5 to 20 times, of the molar amount of compound of
25 formula (2).

REAC2 can be done batch wise or in a continuous way.

Preferably, the reaction temperature TEMP2 of REAC2 is from 80 to 120°C, more preferably
30 from 90 to 110°C.

The reaction time TIME2 of REAC2 is preferably from 10 sec to 24 h, more preferably from 1 min to 12 h.

When REAC2 is done batch wise, then

preferably, the reaction time TIME2 of REAC2 is from 30 min to 24 h, more preferably from 1 h min to 12 h.

5

When REAC2 is done in a continuous way, then

preferably, the reaction time TIME2 of REAC2 is from 10 sec to 30 min, more preferably from 1 min to 15 min.

10 Preferably, both REAC1 and REAC2 are done in a continuous way and are preferably done consecutively without isolation of compound of formula (2), preferably the reaction mixture from REAC1 is used as the substrate feed for REAC2, preferably without interruption of the flow of the feeds.

15 After REAC2, compound of formula (6) can be isolated and purified by conventional methods, which are known to those skilled in the art. These conventional methods include quenching the reaction mixture from REAC2 with water, extraction, distillation, preferably fractional distillation, which can be done under reduced pressure, crystallization, chromatography, filtration, washing or any combination of these methods.

20

Examples

HPLC procedure

Column: Phenomenex Kinetex C-18 100x4.6mm, 2.6 micrometer

Temperature: 25°C


5 Solvent A: Acetonitrile

Solvent B: 0.05% (v/v) aqueous TFA

Flow: 1.3 mL/min

Gradient (v/v):

- within 30 min: Solvent A : Solvent B from 5 % : 95 % to 95 % : 5%
- 10 • within 1 min: Solvent A : Solvent B from 95 % : 5 % to 5 % : 95%
- for 4 min: Solvent A : Solvent B 5 % : 95%



dinitro derivative, the compound of formula (4)

15

Comparative Example 1 – with concentrated sulphuric acid and with aqueous nitric acid of 65 wt%

10.0 g (0.065 mol, 1 eq) of compound of formula (1) were dissolved in 67.4 g of concentrated sulphuric acid (0.688 mol, 10.6 eq). The solution was cooled to 0°C and 7.5 g of nitric acid (65 wt% aqueous solution, 0.078 mol, 1.2 eq) were added. After stirring for 1 h at 0°C, an HPLC chromatogram showed 80% conversion and 45% of the dinitro derivative, the compound of formula (4).

Comparative Example 2 – with concentrated sulphuric acid and with fuming nitric acid

25 10.0 g (0.065 mol, 1 eq) of compound of formula (1) were dissolved in 67.4 g of concentrated sulphuric acid (0.688 mol, 10.6 eq). The solution was cooled to 0°C and 5.0 g of fuming nitric acid (99 wt%, 0.078 mol, 1.2 eq) were added. After stirring for 1 h at 0°C, an HPLC chromatogram showed 80% conversion and 42% of the dinitro derivative, the compound of formula (4).

30

Example 1 – with oleum and with fuming nitric acid

10.0 g (0.065 mol, 1 eq) of compound of formula (1) were dissolved in 67.4 g of concentrated sulphuric acid (0.688 mol, 10.6 eq, 96 to 100 wt%) and in 9.6 g of oleum (65% of SO₃, 0.078 mol, 1.2 eq). The solution was cooled to 0°C and 7.5 g of fuming nitric acid (99 wt% , 0.078 mol, 1.2 eq) were added. After stirring for 1 h at 0°C, an HPLC chromatogram showed full conversion and less than 0.5% of the dinitro derivative, the compound of formula (4).

Example 2 – continuous flow nitration with oleum and with fuming nitric acid followed by esterification**Step Nitration**

Two feeds were prepared for the continuous flow nitration, a Feed 1 and a Feed 2:

Feed 1: 125 g of fuming nitric acid (99 wt%, 1.98 mol), 94.54 g of concentrated sulphuric acid (0.90 mol, 96 to 100 wt%) and 31.46 g of oleum (0.32 mol)

Feed 2: 177.55 g of compound of formula (1) (1.12 mol), 522.45 g of concentrated sulphuric acid (5.06 mol, 96 to 100 wt%) and 300 g of oleum (3.05 mol)

Feed 1 and Feed 2 were pumped using a separate pump for each feed. Feed 1 was pumped with 9.139 g/min, Feed 2 was pumped with 55.861 g/min.

The two feeds were initially pre-cooled to 20°C using two plates, a plate with an internal volume of 19.5 mL for Feed 1 and a plate with an internal volume of 25.77 mL for Feed 2.

After said cooling to 20°C, the feeds were mixed in the reactor, which was a FlowPlate® A5 mikroreactor, process plate LL, Ehrfeld Mikrotechnik BTS GmbH, D-55234 Wendelsheim, Germany, with a volume of 10.28 mL, at 20°C and with a residence time of 17 sec. The crude product solution of compound of formula (2) exiting the reactor constituted the third feed, the Feed 3. An HPLC chromatogram of a sample of Feed 3 showed full conversion and less than 0.5% of the dinitro derivative, the compound of formula (4). After 60 min, the system was purged with dichloromethane. Feed 3 was used directly in the next step, the esterification.

Step Esterification

Feed 3 was pumped with 9.834 g/min and a fourth feed, the Feed 4, which was methanol, was pumped with a flow rate of 8.792 g/min (11 eq).


Initially, the two feeds were pre-heated to 100°C using two plates, a plate with an internal volume of 5.13 mL for Feed 3 and a plate with an internal volume of 11.97 mL for Feed 4.

After said heating to 100°C, the feeds were mixed in the reactor, which was a FlowPlate® A5 mikroreactor, process plate LL, Ehrfeld Mikrotechnik BTS GmbH, D-55234 Wendelsheim,


Germany, with a volume of 2.05 mL, at 100°C, the residence time was 6 min. The reaction solution exiting the reactor was then passed through a coil heated at 100°C providing for a residence time of 6 min. An HPLC chromatogram of a sample of the crude product solution of compound of formula (6) exiting the coil showed full conversion and less than 0.5% of the starting material, the compound of formula (2). After 20 min, the system was purged with methanol. The crude product solution was cooled to 20°C whereby compound of formula (6) precipitated, the solid was isolated by filtration and dried under vacuum (40°C, 20 mbar, 10 h). 221 g of compound of formula (6) were obtained as a colorless solid (yield: 90%, purity: 99.2%).

Claims

1. Method for the preparation of compound of formula (2)



comprising a step STEP1, STEP1 comprises a reaction REAC1, in REAC1 compound of formula (1)



is converted to compound of formula (2) by the action of a mixture MIX of concentrated sulphuric acid, oleum and fuming nitric acid.

2. Method according to claim 1, wherein
the concentrated sulfuric acid has a content of from 94 to 100 wt% of H₂SO₄;
the wt% based on the weight of the concentrated sulfuric acid.

3. Method according to claim 1 or 2, wherein
the oleum contains 50 to 70 wt% of SO₃.

4. Method according to one or more of claims 1 to 3, wherein
the fuming nitric acid has a content of from 95 to 100 wt% of HNO₃.

5. Method according to one or more of claims 1 to 4, wherein
MIX contains

concentrated sulphuric acid in an amount of 4 to 12 times of molar equivalents of H_2SO_4 relative to compound of formula (1),

oleum in an amount of 1 to 4 times of molar equivalents of H_2SO_4 relative to compound of formula (1),

5 fuming nitric acid in an amount of 1 to 2 times of molar equivalents of HNO_3 relative to compound of formula (1).


6. Method according to one or more of claims 1 to 5, wherein the amounts of concentrated sulfuric acid, oleum and fuming nitric acid in MIX add up to 95
10 to 100 wt%, more preferably to 98 to 100 wt%, the wt% being based on the total weight of MIX; even more preferably MIX consists of concentrated sulfuric acid, oleum and fuming nitric acid.

7. Method according to one or more of claims 1 to 6, wherein
15 the reaction temperature TEMP1 of REAC1 is from -10 to 30°C .

8. Method according to one or more of claims 1 to 7, wherein the reaction time TIME1 of REAC1 is from 1 sec to 2 h.

20 9. Method according to one or more of claims 1 to 8, wherein REAC1 is done in a continuous way.

10. Method for the preparation of compound of formula (6)



25

comprising the STEP1 and a second step STEP2;

STEP2 comprises a reaction REAC2, in REAC2 compound of formula (2), which is obtained by REAC1 in STEP1, is reacted with methanol to provide compound of formula (6);

30 with STEP1 as defined in claim 1.

11. Method according to claim 10, wherein

REAC2 is done in the presence of an acid ACID2, ACID2 is H₂SO₄.

12. Method according to claim 10 or 11, wherein

5 both REAC1 and REAC2 are done in a continuous way and are done consecutively without isolation of compound of formula (2).

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/069192

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07C67/08 C07C69/76 C07C201/08 C07C63/04
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2009/112832 A1 (ANGELETTI P IST RICHERCHE BIO [IT]; INGENITO RAFFAELE [IT]; JONES PHIL) 17 September 2009 (2009-09-17) page 60, line 27 - line 35 -----	1-12
A	US 2005/272823 A1 (REHWINKEL HARTMUT [DE] ET AL) 8 December 2005 (2005-12-08) page 23, paragraph 495 -----	1-12
A	WO 2015/180613 A1 (GLAXOSMITHKLINE IP DEV LTD [GB]; GLAXOSMITHKLINE CHINA R & D COMPANY L) 3 December 2015 (2015-12-03) page 22, line 21 - line 31 ----- -/-	1-12



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

22 August 2018

Date of mailing of the international search report

04/09/2018

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Stroeter, Thomas

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/069192

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ADAM T. GILLMORE ET AL: "Multikilogram Scale-Up of a Reductive Alkylation Route to a Novel PARP Inhibitor", ORGANIC PROCESS RESEARCH AND DEVELOPMENT, vol. 16, no. 12, 21 December 2012 (2012-12-21), pages 1897-1904, XP055250298, US ISSN: 1083-6160, DOI: 10.1021/op200238p cited in the application page 1901, right-hand column, paragraph 2 -----	1-12
A	US 2 934 571 A (BONETTI GIOVANNI A) 26 April 1960 (1960-04-26) claims -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2018/069192

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2009112832	A1	17-09-2009	
		AU 2009224004 A1	17-09-2009
		CA 2716918 A1	17-09-2009
		EP 2265611 A1	29-12-2010
		JP 2011513475 A	28-04-2011
		US 2011053911 A1	03-03-2011
		WO 2009112832 A1	17-09-2009

US 2005272823	A1	08-12-2005	
		AR 046062 A1	23-11-2005
		AT 348609 T	15-01-2007
		AU 2004280088 A1	21-04-2005
		BR PI0415209 A	05-12-2006
		CA 2539587 A1	21-04-2005
		CR 8357 A	07-12-2006
		CY 1107596 T1	13-03-2013
		DK 1670458 T3	23-04-2007
		EA 200600621 A1	27-10-2006
		EC SP066539 A	17-10-2006
		EP 1670458 A1	21-06-2006
		ES 2279454 T3	16-08-2007
		HK 1097734 A1	28-05-2010
		HN 2004000446 A	17-09-2008
		IL 174405 A	31-01-2011
		JP 4638438 B2	23-02-2011
		JP 2007508268 A	05-04-2007
		KR 20070000395 A	02-01-2007
		ME P15908 A	10-06-2010
		MY 141853 A	16-07-2010
		NZ 546473 A	25-09-2009
		PE 06042006 A1	04-08-2006
		PT 1670458 E	30-03-2007
		RS 20060250 A	07-08-2008
		SI 1670458 T1	30-06-2007
		TW I306455 B	21-02-2009
		US 2005272823 A1	08-12-2005
		WO 2005034939 A1	21-04-2005

WO 2015180613	A1	03-12-2015	
		AU 2015267910 A1	01-12-2016
		CA 2950220 A1	03-12-2015
		CN 106660999 A	10-05-2017
		EP 3148985 A1	05-04-2017
		JP 2017516800 A	22-06-2017
		KR 20170012388 A	02-02-2017
		US 2017121313 A1	04-05-2017
		WO 2015180613 A1	03-12-2015

US 2934571	A	26-04-1960	NONE
