



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2533579 C 2012/09/11

(11)(21) **2 533 579**

(12) **BREVET CANADIEN  
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2004/07/26  
(87) Date publication PCT/PCT Publication Date: 2005/02/03  
(45) Date de délivrance/Issue Date: 2012/09/11  
(85) Entrée phase nationale/National Entry: 2006/01/23  
(86) N° demande PCT/PCT Application No.: GB 2004/003232  
(87) N° publication PCT/PCT Publication No.: 2005/010210  
(30) Priorité/Priority: 2003/07/24 (GB0317343.2)

(51) Cl.Int./Int.Cl. *C12M 1/34* (2006.01),  
*C12Q 1/68* (2006.01), *C40B 30/04* (2006.01),  
*C40B 60/00* (2006.01)  
(72) Inventeur/Inventor:  
DENSHAM, DANIEL HENRY, GB  
(73) Propriétaire/Owner:  
GEN-PROBE INCORPORATED, US  
(74) Agent: TORYS LLP

(54) Titre : DETERMINATION DE LA SEQUENCE D'UN POLYNUCLEOTIDE  
(54) Title: DETERMINING THE SEQUENCE OF A POLYNUCLEOTIDE

(57) **Abrégé/Abstract:**

The sequence of a target polynucleotide can be determined by: (i) contacting the target polynucleotide with a polymerase enzyme and one of the nucleotides A, T(U), G and C under conditions suitable for the polymerase reaction to proceed; (ii) measuring the time taken for the polymerase to bind to and subsequently dissociate from the target polynucleotide, to thereby determine whether the polymerase has incorporated the nucleotide onto the target polynucleotide; (iii) optionally repeating steps (i) and (ii) with additional nucleotides, to thereby identify the sequence of the target polynucleotide.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
3 February 2005 (03.02.2005)

PCT

(10) International Publication Number  
**WO 2005/010210 A3**

(51) International Patent Classification<sup>7</sup>: **C12Q 1/68**

(21) International Application Number:  
PCT/GB2004/003232

(22) International Filing Date: 26 July 2004 (26.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0317343.2 24 July 2003 (24.07.2003) GB

(71) Applicant (for all designated States except US): **MEDICAL BIOSYSTEMS LTD.** [GB/GB]; The Old Mill, Beaston Cross, Broadhempston, Nr. Totnes, Devon TQ9 6BX (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **DENSHAM, Daniel, Henry** [GB/GB]; Mobious Genomics Ltd, Innovation Centre, University of Exeter, Rennes Drive, Exeter EX4 4RN (GB).

(74) Agent: **GILL JENNINGS & EVERY**; Boradgate Hosue, 7 Eldon Street, London EC2M 7LH (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**


- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:  
24 March 2005

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DETERMINING THE SEQUENCE OF A POLYNUCLEOTIDE

(57) Abstract: The sequence of a target polynucleotide can be determined by: (i) contacting the target polynucleotide with a polymerase enzyme and one of the nucleotides A, T(U), G and C under conditions suitable for the polymerase reaction to proceed; (ii) measuring the time taken for the polymerase to bind to and subsequently dissociate from the target polynucleotide, to thereby determine whether the polymerase has incorporated the nucleotide onto the target polynucleotide; (iii) optionally repeating steps (i) and (ii) with additional nucleotides, to thereby identify the sequence of the target polynucleotide.



WO 2005/010210 A3

## **DETERMINING THE SEQUENCE OF A POLYNUCLEOTIDE**

### **Field of the Invention**

The present invention relates to methods for determining the sequence  
5 of a polynucleotide or detecting variations between polynucleotide sequences.

### **Background to the Invention**

The ability to determine the sequence of a polynucleotide is of great  
scientific importance, as demonstrated by the Human Genome Project, which  
has now determined the entire sequence of the three billion bases of the human  
10 genome. However, this sequence information represents an average human and  
there is a considerable need to understand the differences between individuals  
at a genetic level.

The principal method in general use for large-scale DNA sequencing is  
the chain termination method. This method was first developed by Sanger and  
15 Coulson (Sanger et al. Proc. Natl. Acad. Sci. USA 1977; 74: 5463-5467), and  
relies on the use of dideoxy derivatives of the four nucleoside triphosphates  
which are incorporated into the nascent polynucleotide chain in a polymerase  
chain reaction. Upon incorporation, the dideoxy derivatives terminate the  
polymerase reaction and the products are then separated by gel electrophoresis  
20 and analysed to reveal the position at which the particular dideoxy derivative  
was incorporated into the chain.

Although this method is used widely and produces reliable results, it is  
recognised that it is slow, labour-intensive and expensive. Furthermore, it is not  
an effective method for detecting the differences between two sequences, which  
25 may often consist of a single base change (known as a Single Nucleotide  
Polymorphism, or SNP).

Nucleic acid arrays have recently become a preferred method of  
determining polynucleotide sequences and SNPs, usually in the context of  
hybridisation events (Mirzabekov, Trends in Biotechnology (1994) 12:27-32). A  
30 large number of array-based sequencing procedures utilise labelled nucleotides  
in order to obtain the identity of the added (hybridised) bases. These arrays  
rely on the stepwise identification of suitably labelled bases, referred to in US-A-



5634413 as "single base" sequencing methods. Such "single base" procedures utilise two types of label; the radiolabel and the fluorescent label. The radiolabelling of nucleotides has the advantages of high sensitivity and low background. However, radiolabelling suffers from poor resolution.

5 Fluorescently-labelled nucleotides are now used widely in many techniques. Such nucleotides can be incorporated into the nascent polynucleotide chain in a stepwise manner by the polymerase chain reaction. Each of the different nucleotides (A, T, G and C) incorporates a unique fluorophore at the 3' position which can be detected using a sensitive fluorescent  
10 detector, e.g. a charge-coupled detector (CCD). The fluorophore often also acts as a "blocking group", which removes the ability of the incorporated nucleotide to serve as a substrate for further nucleotide addition and therefore prevents uncontrolled polymerisation. Often, a "removable blocking group" is used, which can be removed by a specific treatment that results in cleavage of the covalent  
15 bond between a nucleotide and the blocking group, allowing the sequencing reaction to continue.

Removable blocking groups rely on a number of possible removing treatment strategies, for example, a photochemical, chemical or enzymatic treatment. However, these have been shown to be difficult to control and apply.  
20 Differences in local environments, for example within an array, can result in the removal of an entire nucleotide, or even several nucleotides, instead of just the intended label. Such occurrences have serious consequences for the fidelity of the sequencing method, as uncontrolled removal of nucleotides results in sequencing data becoming out of phase and sequence data becoming corrupted  
25 or unusable.

A further disadvantage of both labelling methods is that repeat sequences can lead to ambiguity of results. This problem is recognised in Automation Technologies for Genome Characterisation, Wiley-Interscience (1997), ed. T.J.Beugelsdijk, Chapter 10:205-225.

30 There is therefore a need for an improved method for identifying the sequence of a polynucleotide, in particular for detecting variations within a polynucleotide sequence, eg for detecting SNPs, which combines the high

sensitivity and low background of radiolabelled nucleotides with the high resolution of fluorescently-labelled labels. Further, the method should be capable of being carried out by high-throughput, automated processes, reducing the cost associated with existing methods.

## 5 **Summary of the Invention**

The present invention is based on the realisation that the identification of a polynucleotide sequence can be carried out by measuring the time during which a polymerase binds to a target polynucleotide during a polymerase reaction. In general, a polymerase will spend less time bound to the target  
10 polynucleotide when there is no nucleotide available for incorporation. Therefore, if the only nucleotide available is non-complementary, the polymerase will bind to the polynucleotide for a shortened period, and this can be determined to reveal the identity of the complementary sequence of the target polynucleotide.

15 According to a first aspect of the present invention, a method of identifying the sequence of a polynucleotide comprises:

- (i) contacting the target polynucleotide with a polymerase enzyme and one of the nucleotides A, T(U), G and C under conditions suitable for the polymerase reaction to proceed;
- 20 (ii) measuring the time taken for the polymerase to bind to and subsequently dissociate from the target polynucleotide, to thereby determine whether the polymerase has incorporated the nucleotide onto the target polynucleotide; and
- (iii) optionally repeating steps (i) and (ii) with additional nucleotides,  
25 to thereby identify the sequence of the target polynucleotide.

According to a second aspect of the present invention, a method for the identification of a mutation in a target polynucleotide, comprises the steps of:

- (i) contacting the target polynucleotide with a polymerase enzyme and one of the nucleotides A, T (U), G and C under conditions suitable for the  
30 polymerase reaction to proceed; and
- (ii) measuring the time taken for the polymerase to bind to and subsequently dissociate from the target polynucleotide, to thereby identify



whether the polymerase has incorporated the nucleotide onto the target polynucleotide, and with reference to the native sequence of the target, determine whether a mutation exists.

### **Description of the Invention**

5           The term "polynucleotide" or "target polynucleotide" as used herein is to be interpreted broadly, and includes DNA and RNA, including modified DNA and RNA, as well as other hybridising nucleic acid-like molecules e.g. peptide nucleic acid (PNA). The target polynucleotide may be double or single-stranded. Preferably, the target polynucleotide is at least partially single-stranded and  
10 more preferably has a primer sequence bound to it, such that the polymerase enzyme can attach to the target polynucleotide, as will be appreciated by one skilled in the art.

          The enzyme is a polymerase enzyme, which interacts with the target polynucleotide in the process of extending a complementary strand, and may be  
15 of any known type. For example, the polymerase may be any DNA-dependent DNA polymerase. If the target polynucleotide is a RNA molecule, then the polymerase may be an RNA-dependent DNA polymerase, i.e. reverse transcriptase, or a RNA-dependent RNA polymerase, i.e. RNA replicase. Primase enzymes are also included within the definition.

20           The target polynucleotide is preferably localised at a specific site on a support material. Preferably, the polynucleotide is localised via immobilisation on a solid support. Supports suitable for use in immobilising the polynucleotide will be apparent to the skilled person, for example silicon, glass or ceramic materials may all be used. Immobilisation may be carried out by covalent or non-  
25 covalent means. For example, covalent linker molecules may be used. In a preferred embodiment, a primer is immobilised onto the support material and the target polynucleotide is hybridised to it. Alternatively, hybridisation of the target polynucleotide and primer may take place in solution and either the primer or target polynucleotide is subsequently, or simultaneously, attached to a support  
30 material.

          There may be one or more target polynucleotides immobilised to the solid support material. In a preferred embodiment, a plurality of target polynucleotides

are attached to the support. These "arrays" of target polynucleotides may be of any known density, including multi-molecule, high-density arrays as well as "single-molecule" arrays in which individual polynucleotide locations may be resolved, i.e. it is possible to monitor the polymerase reaction occurring on a  
5 single polynucleotide.

As a first step in the sequencing method, the target polynucleotide may be brought into contact with an appropriate primer in hybridising/polymerisation buffer. Typically, the buffer will be at a temperature sufficiently high to disrupt (or melt) any secondary structures that exist on the target polynucleotide. On  
10 cooling, the primer will anneal to its complement on the target. This is then brought into contact with the polymerase, to form the target polynucleotide/polymerase complex.

In one embodiment of the invention, the addition of the nucleotides is controlled so that the different nucleotides are added sequentially to the  
15 polymerase/target complex. For example, dGTP may be added and allowed to flow over the polymerase/polynucleotide complex; the time during which the polymerase is complexed to the polynucleotide is then determined. Unbound dGTP flows out of the reaction site and a further nucleotide is introduced. In this manner, the time during which the complex is formed can be correlated to the  
20 particular nucleotide present during that time and the polynucleotide sequence, can therefore be determined.

The reaction may also be started by the addition of the polymerase to a reaction mixture which comprises the target polynucleotide and the nucleotide. After monitoring the time taken by the polymerase to associate with and  
25 dissociate from the target polynucleotide, the unbound nucleotides can be removed and the next nucleotide introduced to carry out further sequencing steps.

The incorporation of a nucleotide onto the target requires the polymerase to associate with the target for a period longer than that taken when no  
30 incorporation is possible. Measurement of the association between the polymerase and the target therefore reveals whether an incorporation event has occurred for regions on the target having sequential nucleotides of the same



type, the time of association will increase proportionally. Therefore the method can be used to identify consecutive nucleotides of the same type.

The detection of the interaction between the polymerase enzyme and the target polynucleotide can be carried out by measuring changes in applied  
5 radiation. Measuring the changes in radiation that occurs on interaction between the polymerase and target polynucleotide may be carried out using conventional apparatus.

In one embodiment, the interaction of the polymerase with the target polynucleotide is measured using a non-linear imaging system.

10 Non-linear imaging systems are known in the art. In general, the non-linear polarisation for a material can be expressed as:

$$P = X^{(2)}E^1 + X^{(2)}E^2 + X^{(3)}E^3 + \dots$$

15 where P is the induced polarisation,  $X^{(n)}$  is the nth-order non-linear susceptibility, and E is the electric field vector. The first term describes normal absorption and reflection of light; the second describes second harmonic generation (SHG), sum and difference frequency generation; and the third describes light scattering, stimulated Raman processes, third harmonic generation (TGH), and both two-  
20 and three-photon absorption.

A preferred imaging system of the present invention relies on the detection of the signal arising from second or third harmonic generation.

Single-molecule resolution using second or third harmonic generation (hereinafter referred to as SHG) is known in the art (Peleg et al., Proc. Natl.  
25 Acad, Sci. USA, 1999;95:6700-6704 and Peleg et al., Bioimaging, 1996; 4: 215-224). Suitable systems are described in WO-A- 02/095070.

In one embodiment, detection is carried out in solution phase (i.e. the polymerase and target polynucleotide are not immobilised), using raman scattering and/or LSPR techniques.

30 When light is directed onto a molecule, the vast majority of the incident photons are elastically scattered without a change in frequency. This is termed Rayleigh scattering. However, the energy of some of the incident photons



(approximately 1 in every  $10^7$  photons) is coupled into distinct vibrational modes of the molecule's bonds. Such coupling causes some of the incident light to be inelastically scattered by the molecule with a range of frequencies that differ from the range of the incident light. This is termed the Raman effect. By plotting the frequency of such inelastically scattered light against intensity, the unique Raman spectrum of the molecule under investigation is obtained. Analysis of the Raman spectrum of an unknown sample can yield information about the samples molecular composition.

The Raman effect can be enhanced significantly by bringing the Raman active molecule(s) close ( $\leq 50\text{\AA}$ ) to a structured metal surface, this field decays exponentially away from the surface. Bringing molecules in close proximity to metal surfaces is typically achieved through adsorption of the Raman-active molecule onto suitably roughened gold, silver copper or other free-electron metals. Surface enhancement of the Raman activity is observed with metal colloidal particles, metal films on dielectric substrates, and with metal particle arrays. The mechanism by which this surface-enhanced Raman scattering takes place is not well understood, but it thought to result from a combination of (i) surface plasmon resonances in the metal that enhance the local intensity of the light, and ; (ii) formation and subsequent transitions of charge-transfer complexes between the metal surface and the Raman-active molecule.

The Raman effect can be utilised to measure the time of association between the polymerase and the target polynucleotide. It is preferred if the target is immobilised on a metal surface, to improve the Raman signal.

In a preferred embodiment of the present invention, Surface Enhanced Raman Scattering (SERS) is employed via the use of a metal colloidal nanoparticle bound to the polymerase eg. a gold or silver nanoparticle. A Raman enhancing metal nanoparticle that has associated or bound to it a Raman-active molecule(s) can have utility as an optical tag. This general concept is outlined in US Patent No. 6,514,767.

In one embodiment, the target polynucleotide and a polymerase can be detected by searching the Raman active molecule's unique raman spectrum. Because a single Raman spectrum (from  $100\text{--}3500\text{ cm}^{-1}$ ) can detect many

different Raman-active molecules, SERS-active nanoparticles bound to or associated with the polymerase may be used in the context of the present invention within multiplexed assay formats.

In a separate embodiment, changes in radiation are monitored by utilising  
5 surface electromagnetic wave technology.

Biosensors incorporating surface electromagnetic wave technology (and, in particular, surface plasmon resonance - SPR - sensors) are based on the sensitivity of surface electromagnetic waves (SEW) to the refractive index of the thin layer adjacent to the surface where the SEW propagates. In the biosensor,  
10 the polymerase and nucleotide(s) are allowed to flow across the surface containing the immobilised target polynucleotide. As binding occurs, the accumulation or redistribution of mass on the surface changes the local refractive index that can be monitored in real time by the sensor.

Several methods utilising SPR technology have been proposed and  
15 realised in biosensors. The most popular methods are based on the Kretschmann-Raether configuration where the intensity of the light reflected from the sensor is monitored. This technique, considered to be one of the most sensitive, is described in J. Homola et al, Sensors and Actuators B 54, p.3-15 (1999) and has a detection limit of  $5 \times 10^{-7}$  refractive index units. Measuring SPR  
20 phase changes can further increase the sensitivity of the sensor by one or two orders of magnitude. This is described in Nelson et al, Sensors and Actuators B 35-36, p. 187 (1996) and in Kabashkin et al, Optics Communications 150, p.5 (1998). Prior art interferometric devices such as a Mach Zehnder device have been configured to measure variations in the refractive index at the sensor  
25 surface via phase shifts. This is disclosed in International Patent Publication WO-A-0120295. The configuration requires four independent components and is sensitive to sub-wavelength relative replacements of these components and hence very small mechanical and environment perturbations. A mechanically more robust monolithic interferometric design is outlined in WO-A-03014715.

30 In a preferred embodiment, a surface electromagnetic wave (SEW) sensor system is used which can compensate for changes in the bulk refractive index of a buffer or which allows the contribution of the bulk refractive index to an



interference pattern to be separated from the contribution of an analyte absorbed on the sensor surface. The biosensor therefore comprises:

a coherent radiation source for producing an incident wave;

a carrier surface for supporting the immobilised polynucleotide, the carrier  
5 surface mounted on a substrate and capable of supporting surface  
electromagnetic waves (SEW);

means for splitting the incident wave into an SEW and a first scattered  
wave, wherein the SEW propagates along the carrier surface and interacts with  
the immobilised polynucleotide;

10 means for generating a second scattered wave from the SEW; and

a detector for monitoring the interference between the first scattered wave  
and the second scattered wave.

In this embodiment, a coherent optical beam generated by a  
monochromatic laser is focussed using a lens, onto the edge of a metallic film  
15 able to support surface electromagnetic waves (SEWs). The optical beam  
passes through a glass prism on which the metallic film is mounted. A near-  
infrared laser is used as the illumination source. Using a near-infrared source  
has the advantage of long propagation length for surface plasmons in gold and  
silver while conventional optics can be still used for imaging and illumination.  
20 However, other monochromatic sources are suitable and may be used.

This system is described in WO-A-04/020985.

In a preferred embodiment, the polymerase is labelled.

The term "label" as used herein may be interpreted broadly. Suitable  
25 labels will be apparent to the skilled person. In a preferred embodiment, the  
label is a fluorophore. In a particular embodiment, the polymerase is prepared  
as a recombinant fusion with GFP (Green Fluorescent Protein). This may either  
be wild type GFP or spectrally shifted mutants thereof (Tsien, Ann. Rev.  
Biochem., 1998; 67:509, US5,777,079 and US 5,625,048). The GFP can be  
30 located at the N- or C- terminus of the enzyme (the C- terminus may be desirable  
if a polymerase is to be used in conjunction with a 'sliding clamp'). Alternatively,  
the GFP molecule may be located anywhere within the enzyme, provided that

enzymatic activity is retained. Alternative labels may be used. A number of strategies for labelling molecules have been reported, such as microspheres (Anal. Chem. (2000) 72, 15:3678-3681), gold nanoparticles (J.Am.Chem. Soc. (2000) 122, 15:3795-3796), silver colloid particles (PNAS, (2000) 97, 3:996-1001) and quantum dots. Any labelling technique that allows unambiguous resolution of the interaction of the polymerase with the target polynucleotide and nucleotide can be utilised within the context of the present invention. Preferably, a label that allows the temporal resolution of enzyme binding and/or unbinding from the target polynucleotide:nucleotide complex is used.

10 Preferably, an array of immobilised target polynucleotides is created and the polymerase enzyme is flowed over the array in conditions sufficient for enzyme activity, together with at least one of the four nucleotides. More preferably, this takes place within a flow cell. Each "array location" within the array, containing a single immobilised target polynucleotide or plurality of  
15 identical target polynucleotides, is then imaged at a time resolution that allows resolution of the binding of the enzyme to the target polynucleotide/nucleotide complex, preferably, from 1 to at least several thousand hertz.

In another preferred embodiment, the polymerase enzyme is labelled with a Fluorescence Resonance Energy Transfer (FRET) pair. This system is  
20 described in WO-A-01/25480. As will be appreciated by one skilled in the art, a FRET pair requires a fluorophore and a second molecule that is capable of interacting with the fluorophore, acting as an energy donor or acceptor. The interaction between the pair alters the emission spectrum of the fluorophore(s), which is detected. The second molecule may be another fluorophore, or any  
25 other suitable molecule, such as a metallic particle. Preferably a classical FRET pair is used, where both labels are fluorophores, chosen so that the emission wavelength of one fluorophore corresponds to the excitation wavelength of the other. The immobilisation locations on the enzyme are chosen such that the fluorophores move relative to one another when the enzyme binds to the  
30 polynucleotide forming the ternary complex. This relative movement between the FRET pair will result in a spectral shift of the emission from the pair, depending upon the way in which the FRET pair has been designed. More or less energy



is then coupled into the acceptor from the donor, leading to excitation or attenuation of the donor fluorescence. This is monitored with high temporal resolution, to obtain the desired information. Preferably the device used to image the extent of fluorescence is a charge-coupled device (CCD), with rapid  
5 read-out time and appropriate optics for high spatial resolution.

The resolution system required to monitor the interaction between the polymerase and the target can be modified to take into account the need to "scan" the sample under study. Such scanning is especially likely in high resolution systems, such as single molecule systems. At the high levels of  
10 spatial resolution required for single molecule experiments, the effective observation area is correspondingly reduced. Since "kinetic" analysis normally requires continuous "real-time" monitoring at a single defined area, scanning introduces the problem of discontinuous monitoring. Such discontinuous monitoring is incompatible with kinetic analysis as interaction events can be  
15 missed. In a preferred aspect of the present invention, therefore, nucleotides labelled at the 5 prime end are employed; the label acting as a removable blocking group. Thus in situations where high spatial resolution is employed (such as single molecule arrays in which the observation region is typically only 100µm by 100µm and the chip is generally at least 100 times larger), the  
20 observation region is "moved" to a new location and that region illuminated with radiation (e.g. UV) suitable to remove the blocking group. This strategy ensures that association/kinetic events only occur in an "observed" region. The "observed" region can then be moved around the chip and by only exposing the region under observation to radiation sufficient to unblock the nucleotide,  
25 selective interactions only take place in this region.

In a further separate preferred embodiment imaging areas are confined to individual flow cells, i.e. in the event of the observation area being limited to 100µm by 100µm, the flow cell is restricted approximately to these relative dimensions. Thus, in this embodiment, nucleotides are only injected into flow  
30 cells that are being actively imaged. The support material onto which the polynucleotides are arrayed, may therefore consist of an "array" of individually addressable flow cells. As the imaging area is moved around the chip, so the

nucleotides are injected into the flow cell(s) in the imaging area only.

In a further embodiment, the binding of the polymerase is detected via a resonance energy transfer between an intercalating molecule and a dye/quencher on the polymerase. The intercalating dye (e.g. CYBRGreen, molecular probes) will only fluoresce when binding to the double stranded primer-template complex. This signal can then be used to identify the location of the target polynucleotide. The polymerase is modified with a dye that is an "acceptor" molecule capable of absorbing and re-emitting energy from the intercalating dye. Alternatively, the "label" may be a quencher. Thus the binding/interaction event may be monitored by observing the change in signal due to energy transfer.

The method of the invention may be used to identify the complete target polynucleotide sequence, or may be used to identify the sequence of a part of the polynucleotide. The method is suitable for determining the presence of mutations within the target, for example determining whether a substitution, deletion or addition has occurred, compared to a control or reference sequence.

In a preferred embodiment, the method can be used to identify a single nucleotide polymorphism in a genetic sample. The term "polymorphism" as used herein refers to the occurrence of two or more alternative genomic sequences or alleles between or among different genomes or subjects. A single nucleotide polymorphism (SNP) is a single base pair change. Typically, a SNP is the replacement of one nucleotide by another nucleotide at the polymorphic site. Deletion of a single nucleotide, or insertion of a single nucleotide, also gives rise to single nucleotide polymorphisms.

The method is used to determine the identity of the nucleotide(s) at the putative site of mutation and the information may then be compared to a reference sequence to reveal whether a mutation is present.

The following Example illustrates the invention.

#### Example

In this experiment, a fusion protein of Green Fluorescent Protein (GFP) and a polymerase was created via recombinant techniques well known in the art.



Quartz chips (14mm in diameter, 0.3mm thick) were spin-coated with a layer of planar dextran modified with immobilised streptavidin (Xantec, Germany). An upright microscope (135TV, Zeiss) equipped with total internal reflection (TIR) illumination served as a platform for the experiment. A laser beam of peak 488 nm (Melles Griot) with a power of 100mW was circularly polarised by quarter-wave plates to undergo TIR in a custom made prism. The prism was coupled optically to the chip via index matching gel. A flow cell with an O-ring seal and transparent optical axis was then assembled over the prism and quartz chip. An objective collected the fluorescence signal through the transparent top surface of the flow cell. Images travelled via a band-pass filter into a back-thinned charge-coupled device (Andor technologies).

Two oligonucleotides were synthesized using standard phosphoramidite chemistry. The oligonucleotide defined as SEQ ID NO.1 was used as the target polynucleotide, and the oligonucleotide defined as SEQ ID NO.2 was biotinated and used as a primer. The biotinated primers (SEQ ID No.2) were deposited onto the streptavidin coated chip by injection into the flow cell 10 pM of SEQ ID NO.2 in Tris buffer containing 100 mM MgCl<sub>2</sub> and allowed to incubate for 10 minutes.

20 SEQ ID NO.1

CAAGGAGAGGACGCTGCTTGTCGAAGGTAAGGAACGGACGAGAGAAGG  
GAGAG

25 SEQ ID NO.2

biotin-CTCTCCCTTCTCTCGTC

The flow buffer was flushed into the flow cell at a rate of 500µl/min. The complementary sequence, SEQ ID NO.1 was then injected into the flow cell at a concentration of 1µM, at a flow rate of 5µl/min. During this injection the cell was heated to 70°C for several minutes via a platinum heating electrode within

the flow cell and allowed to cool to 25°C. The injection was then continued for a further ten minutes. Once the injection was complete, flow buffer was flushed continuously through the system at a flow rate of 500µl/min. After 10 minutes the sequencing reaction was initiated by injection of 0.4mM dATP (8µl) into the buffer  
5 at a flow rate of 500µl/min. One second after commencing the injection a series of CCD images was recorded at maximum frame rate (30 frames per second) and the image containing the highest signal intensity stored. The complementary nucleotide dCTP was then injected at 0.4mM at the further CCD images stored one second after injection.

10 The complementary nucleotide gave a strong signal (Fig 1a) in comparison to the non-complementary nucleotide (Fig 1b) when imaged in a "time-resolved" manner.



## SEQUENCE LISTING

<110> Medical Biosystems Ltd.  
 <120> Determining the sequence of a polynucleotide  
 <130> 31719-2110  
 <140> CA 2,533,579  
 <141> 2004-07-26  
 <150> GB 0317343.2  
 <151> 2003-07-24  
 <160> 2  
 <170> PatentIn version 3.1  
 <210> 1  
 <211> 53  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Synthetic target polynucleotide  
 <400> 1

caaggagagg acgctgcttg tcgaaggtaa ggaacggacg agagaaggga gag

53

<210> 2  
 <211> 17  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Primer  
 <400> 2

ctctcccttc tctcgtc

17

CLAIMS

1. An apparatus for sequencing a polynucleotide, comprising:
  - a. a flow cell comprising a fluid flow path;
  - b. a solid support adapted to immobilize a target polynucleotide thereon,wherein the support is positioned within the fluid flow path of the flow cell such that, upon immobilization of a target polynucleotide, the target polynucleotide is positioned in a reaction site within at least a portion of the fluid flow path;
  - c. a radiation source; and
  - d. an imaging apparatus,wherein the radiation source is configured to emit a radiation signal directed at the position of the solid support adapted to immobilize the target polynucleotide, and wherein the imaging apparatus is configured to focus on the position of the solid support adapted to immobilize the target polynucleotide such that the association and disassociation of a free polymerase enzyme with the immobilized target polynucleotide is monitored over time to determine if a polymerization occurs, thereby identifying nucleotide incorporation events and determining the sequence of the target polynucleotide.
2. The apparatus of claim 1, further comprising one or more target polynucleotides immobilized on the solid support.
3. The apparatus of claim 2, comprising an array of immobilized target polynucleotides.
4. The apparatus of claim 3, wherein the array comprises a high density array.
5. The apparatus of claim 1, wherein the imaging apparatus is a total internal reflection microscope.
6. The apparatus of claim 1, wherein the radiation source comprises one or more laser light emitting devices.
7. The apparatus of claim 1, wherein the imaging apparatus further comprises a charge coupled device (CCD) camera.



8. The apparatus of claim 1, further comprising one or more immobilized primer sequences that are complementary to the target polynucleotide.
9. The apparatus of claim 1, wherein the apparatus is configured to monitor the polymerization reaction by monitoring changes in fluorescent signal intensity over time.
10. The apparatus of claim 9, wherein the wherein the imaging apparatus further comprises a CCD camera that records the fluorescent signal intensity.
11. An apparatus for sequencing a polynucleotide, comprising:
  - a. a flow cell comprising a fluid flow path;
  - b. a solid support adapted to immobilize a target polynucleotide thereon, wherein the support is positioned within the fluid flow path of the flow cell such that, upon immobilization of a target polynucleotide, the target polynucleotide is positioned in a reaction site within at least a portion of the fluid flow path;
  - c. a monochromatic laser; and
  - d. a total internal reflection microscope,wherein the laser is configured to emit an optical beam directed at the reaction site, and wherein a total internal reflection microscope is focused on the reaction such that the association and disassociation of a free polymerase enzyme with the immobilized target polynucleotide is monitored over time to determine if a polymerization occurs, thereby identifying nucleotide incorporation events and determining the sequence of the target polynucleotide.
12. The apparatus of claim 11, further comprising one or more target polynucleotides immobilized on the solid support.
13. The apparatus of claim 11, further comprising one or more immobilized primer sequences that are complementary to the target polynucleotide.
14. The apparatus of claim 11, wherein the apparatus is configured to monitor the polymerization reaction by monitoring changes in fluorescent signal intensity over time.

15. The apparatus of claim 14, wherein the wherein the imaging apparatus further comprises a CCD camera that records the fluorescent signal intensity.
16. A method for sequencing a target polynucleotide, comprising:
  - a. immobilizing a target polynucleotide on the support of the apparatus of claim 1, by hybridizing the target polynucleotide to a complimentary primer sequence immobilized in the reaction site;
  - b. contacting the target polynucleotide with a polymerase enzyme and a nucleotide selected from the group consisting of A, T (U), G, and C in the flow cell under conditions suitable for a polymerase reaction to proceed;
  - c. observing the interaction of the polymerase enzyme, primer sequence, and the nucleotide with the target polynucleotide by irradiating the reaction site using the radiation source and detecting the interaction using the imaging apparatus;
  - d. determining whether the polymerase has incorporated the nucleotide onto the primer sequence by monitoring the duration of time that the polymerase enzyme is associated with the target polynucleotide by detecting a signal produced during the interaction using the imaging apparatus; and
  - e. repeating steps (b)-(d) one or more times, thereby identifying the sequence of the target polynucleotide.
17. The method of claim 16, wherein unbound nucleotides are removed from the reaction site prior to step (c).
18. The method of claim 16, wherein the nucleotide comprises a removable blocking group.
19. A method for sequencing a target polynucleotide, comprising:
  - a. immobilizing the target polynucleotide on the support of the apparatus of claim 11, by hybridizing the target polynucleotide to a complimentary primer sequence immobilized in the reaction site;



- b. contacting the target polynucleotide with a polymerase enzyme and a nucleotide selected from the group consisting of A, T (U), G, and C in the flow cell under conditions suitable for a polymerase reaction to proceed;
- c. observing the interaction of the polymerase enzyme, primer sequence, and the nucleotide with the target polynucleotide by irradiating the reaction site using the monochromatic laser and detecting the interaction using the total internal reflection microscope;
- d. determining whether the polymerase has incorporated the nucleotide onto the primer sequence by monitoring the duration of time that the polymerase enzyme is associated with the target polynucleotide by detecting a signal produced during the interaction using the total internal reflection microscope; and
- e. repeating steps (b)-(d) one or more times, thereby identifying the sequence of the target polynucleotide.

20. The method of claim 19, wherein unbound nucleotides are removed from the reaction site prior to step (c).

21. The method of claim 19, wherein the nucleotide comprises a removable blocking group.

1/2




Fig. 1a



2/2



Fig. 1b