

REPUBLIC OF SOUTH AFRICA

PATENTS ACT, 1978

PUBLICATION PARTICULARS AND ABSTRACT

(Section 32(3)(a) – Regulations 22(i)(g) and 31)

OFFICIAL APPLICATION NO.		LODGING DATE		ACCEPTANCE DATE	
21	01 2006/0376 2	23	11 May 2006	43	03-07-07

INTERNATIONAL CLASSIFICATION		NOT FOR PUBLICATION	
51	A01N	CLASSIFIED BY :	
FULL NAME(S) OF APPLICANT(S)			
71	BAYER CROSCIENCE AG		

FULL NAME(S) OF INVENTOR(S)	
72	FUNKE, Christian; BRETSCHNEIDER, Thomas; FISCHER, Reiner; FISCHER, Rüdiger; HUNGENBERG, Heike; ANDERSCH, Wolfram; THIELERT, Wolfgang; KRAUS, Anton

EARLIEST PRIORITY CLAIMED	COUNTRY	NUMBER	DATE
NOTE : the country must be indicated by its International Abbreviation – see Schedule 4 of the Regulations	33 DE	31 103 53 281.1	32 14 November 2003

TITLE OF INVENTION	
54	ACTIVE AGENT COMBINATIONS WITH INSECTICIDAL AND ACARICIDAL PROPERTIES

57	ABSTRACT (NOT MORE THAN 150 WORDS)	NUMBER OF PAGES	72
----	------------------------------------	-----------------	----

FOR ABSTRACT SEE THE NEXT SHEET

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

- 2006/03762

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
2. Juni 2005 (02.06.2005)

PCT

(10) Internationale Veröffentlichungsnummer
WO 2005/048712 A1

(51) Internationale Patentklassifikation¹: A01N 43/56 //
(A01N 43/56, 55:04, 53:00, 47:38, 47:34, 47:30, 47:24,
47:04, 43:90, 43:76, 43:713, 43:58, 43:56, 43:54, 43:36,
43:24, 43:22, 43:12, 43:10, 41:10, 41:04)

Anton [DE/DE]; Bremersheide 15, 42799 Leichlingen
(DE).

(21) Internationales Aktenzeichen: PCT/EP2004/012329

(74) Gemeinsamer Vertreter: BAYER CROPSCEENCE
AKTIENGESELLSCHAFT; Law and Patents, Patents
and Licensing, 51368 Leverkusen (DE).

(22) Internationales Anmeldedatum:
10. Oktober 2004 (10.10.2004)

(81) Bestimmungsstaaten (soweit nicht anders angegeben, für
jede verfügbare nationale Schutzrechtsart): AE, AG, AL,
AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
103 53 281.1 14. November 2003 (14.11.2003) DE

(84) Bestimmungsstaaten (soweit nicht anders angegeben, für
jede verfügbare regionale Schutzrechtsart): ARIPO (BW,
GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,
ZM, ZW), eurasisches (AM, AZ, BY, KG, KZ, MD, RU,
TJ, TM), europäisches (AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme
von US): BAYER CROPSCEENCE AKTIENGE-
SELLSCHAFT [DE/DE]; Alfred-Nobel-Str. 50, 40789
Monheim (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): FUNKE, Christian
[DE/DE]; Rothenberg 75a, 42799 Leichlingen (DE).
BRETSCHNEIDER, Thomas [DE/DE]; Talstr. 29b,
53797 Lohmar (DE). FISCHER, Kainer [DE/DE];
Nelly-Sachs-Str. 23, 40789 Monheim am Rhein (DE).
FISCHER, Rüdiger [DE/DE]; Zu den Fussfälen 23,
50259 Pulheim (DE). HUNGENBERG, Heike [DE/DE];
Louveciennesstr. 2A, 40764 Langenfeld (DE). ANDER-
SCH, Wolfram [DE/DE]; Schlodderdicher Weg 77,
51468 Bergisch Gladbach (DE). THIELERT, Wolfgang
[DE/DE]; Buschweg 69, 51519 Odenthal (DE). KRAUS,

Veröffentlicht:

- mit internationalem Recherchenbericht
- vor Ablauf der für Änderungen der Ansprüche geltenden
Frist; Veröffentlichung wird wiederholt, falls Änderungen
eintreffen

Zur Erklärung der Zweibuchstaben-Codes und der anderen Ab-
kürzungen wird auf die Erklärungen ("Guidance Notes on Co-
des and Abbreviations") am Anfang jeder regulären Ausgabe der
PCT-Gazette verwiesen.

BCS033093



WO 2005/048712 A1

(54) Title: ACTIVE AGENT COMBINATIONS WITH INSECTICIDAL AND ACARICIDAL PROPERTIES

(54) Bezeichnung: WIRKSTOFFKOMBINATIONEN MIT INSEKTIZIDEN UND AKARIZIDEN EIGENSCHAFTEN

(57) Abstract: The invention relates to novel insecticidal agent combinations, comprising cyclic ketoenols or other acaricidal agents and further insecticidal agents, from the group of anthranilamides, which are particularly suitable for the prevention of animal pests such as insects and undesirable mites.

(57) Zusammenfassung: Die Erfindung betrifft neue insektizide Wirkstoffkombinationen, die aus cyclischen Ketoenolen oder anderen akariziden Wirkstoffen einerseits und weiteren insektiziden Wirkstoffen aus der Gruppe der Anthranilsäureamiden andererseits bestehen und sehr gut zur Bekämpfung von tierischen Schädlingen wie Insekten und unerwünschten Akariden geeignet sind.

Active compound combinations having insecticidal and acaricidal properties

The invention relates to novel active compound combinations comprising, firstly, known cyclic ketoenols and, secondly, further known insecticidally active compounds, which combinations are highly suitable for controlling animal pests, such as insects and unwanted acarids.


It is already known that certain cyclic ketoenols have fungicidal, insecticidal and acaricidal properties (EP-A 0 528 156, EP-A 0 647 637, WO 95/26 345, WO 96/20196, WO 96/25395, WO 96/35664, WO 97/01535, WO 97/02243, WO 97/36868, WO 98/05638, WO 98/25928, WO 99/16748, WO 99/43649, WO 99/48869, WO 99/55673, WO 01/23354 and WO 01/74770). The activity of these compounds is good; however, at low application rates it is sometimes unsatisfactory.

10

It is also known that mixtures of phthalamides and other bioactive compounds have insecticidal and/or acaricidal action (WO 02/087334). The action of these mixtures is not always optimal.

Furthermore, it is already known that numerous heterocycles, organotin compounds, benzoylureas and pyrethroids have insecticidal and acaricidal properties (cf. WO 93/22297, WO 93/10083, DE-A 26 41 343, EP-A 0 347 488, EP-A 0 210 487, US 3,264,177 and EP-A 0 234 045). However, the action of these compounds is not always satisfactory.

It has now been found that active compound combinations comprising compounds of the formula (I) (group 1)



(I)

in which

- X represents C₁-C₆-alkyl, bromine, C₁-C₆-alkoxy or C₁-C₃-haloalkyl,
 Y represents hydrogen, C₁-C₆-alkyl, halogen, C₁-C₆-alkoxy, C₁-C₃-haloalkyl,
 Z represents C₁-C₆-alkyl, halogen, C₁-C₆-alkoxy,
 m represents a number 0-3,
 A³ represents hydrogen or in each case optionally halogen-substituted straight-chain or branched C₁-C₁₂-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, C₁-C₁₀-alkoxy-C₁-C₈-alkyl, C₁-C₈-

polyalkoxy-C₂-C₈-alkyl, C₁-C₁₀-alkylthio-C₂-C₈-alkyl, cycloalkyl having 3-8 ring atoms which may be interrupted by oxygen and/or sulfur or in each case optionally halogen-, C₁-C₆-alkyl-, C₁-C₆-haloalkyl-, C₁-C₆-alkoxy-, C₁-C₆-haloalkoxy, nitro-substituted phenyl or phenyl-C₁-C₆-alkyl,


5 A⁴ represents hydrogen, C₁-C₆-alkyl or C₁-C₆-alkoxy-C₁-C₄-alkyl

or in which

A³ and A⁴ together with the carbon atom to which they are attached form a saturated or unsaturated 3- to 8-membered ring which is optionally interrupted by oxygen and/or sulfur and optionally substituted by halogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio or optionally substituted phenyl or is optionally benzo-fused,

10

G¹ represents hydrogen (a) or represents the groups



in which

R²⁰ represents in each case optionally halogen-substituted C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₈-alkoxy-C₁-C₈-alkyl, C₁-C₈-alkylthio-C₁-C₈-alkyl, C₁-C₈-polyalkoxy-C₂-C₈-alkyl or cycloalkyl having 3-8 ring atoms which may be interrupted by oxygen and/or sulfur atoms,

15

represents optionally halogen-, nitro-, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, C₁-C₆-haloalkyl-, C₁-C₆-haloalkoxy-substituted phenyl;

20

represents optionally halogen-, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, C₁-C₆-haloalkyl-, C₁-C₆-haloalkoxy-substituted phenyl-C₁-C₆-alkyl,

represents in each case optionally halogen- and/or C₁-C₆-alkyl-substituted pyridyl, pyrimidyl, thiazolyl or pyrazolyl,

25

represents optionally halogen- and/or C₁-C₆-alkyl-substituted phenoxy-C₁-C₆-alkyl,

R²¹ represents in each case optionally halogen-substituted C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₈-alkoxy-C₂-C₈-alkyl or C₁-C₈-polyalkoxy-C₂-C₈-alkyl,

represents in each case optionally halogen-, nitro-, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, C₁-C₆-haloalkyl-substituted phenyl or benzyl,

30


R²² represents optionally halogen-substituted C₁-C₈-alkyl, represents in each case optionally C₁-C₄-alkyl-, halogen-, C₁-C₄-haloalkyl-, C₁-C₄-alkoxy-, C₁-C₄-haloalkoxy-, nitro- or cyano-substituted phenyl or benzyl,

R²³ and R²⁴ independently of one another represent in each case optionally halogen-substituted C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylamino, di(C₁-C₈)alkylamino, C₁-C₈-alkylthio, C₂-C₅-alkenylthio, C₂-C₅-alkynylthio, C₃-C₇-cycloalkylthio, represent in each case optionally halogen-, nitro-, cyano-, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy-, C₁-C₄-alkylthio-, C₁-C₄-haloalkylthio-, C₁-C₄-alkyl-, C₁-C₄-haloalkyl-substituted phenyl, phenoxy or phenylthio,

R²⁵ and R²⁶ independently of one another represent in each case optionally halogen-substituted C₁-C₁₀-alkyl, C₁-C₁₀-alkoxy, C₃-C₈-alkenyl, C₁-C₈-alkoxy-C₁-C₈-alkyl, represent optionally halogen-, C₁-C₆-haloalkyl-, C₁-C₆-alkyl- or C₁-C₆-alkoxy-substituted phenyl, represent optionally halogen-, C₁-C₆-alkyl-, C₁-C₆-haloalkyl- or C₁-C₆-alkoxy-substituted benzyl or together represent a 5- to 6-membered ring which is optionally interrupted by oxygen or sulfur and which may optionally be substituted by C₁-C₆-alkyl,

15 or an acaricidally active compound (group 2), preferably

(2-1) the phenylhyrazine derivative of the formula (known from WO 93/10083)




and/or

(2-2) the macrolide with the common name abamectin (known from DE-A 27 17 040)


20 and/or

(2-3) the naphthalenedione derivative of the formula (known from DE-A 26 41 343)



and/or


(2-4) the pyrrole derivative of the formula (known from EP-A 0 347 488)



25


and/or

(2-5) the thiourea derivative of the formula (known from EP-A 0 210 487)



and/or


(2-6) the oxazoline derivative of the formula (known from WO 93/22297)




5

and/or

(2-7) an organotin derivative of the formula



in which

R represents  (2-7-a = azocyclotin),

10


known from The Pesticide Manual, 9th edition, p. 48,

or

R represents -OH (2-7-b = cyhexatin), known from US 3,264,177


and/or

15 (2-8) the pyrazole derivative of the formula (known from EP-A 0 289 879)




and/or

(2-9) the pyrazole derivative of the formula (known from EP-A 0 234 045)



and/or


(2-10) the pyridazinone derivative of the formula (known from EP-A 0 134 439)



(pyridaben)

and/or


5 (2-11) the benzoylurea derivative of the formula (known from EP-A 0 161 019)



(flufenoxuron)

and/or


(2-12) the pyrethroid of the formula (known from EP-A 0 049 977)



(bifenthrin)

10 and/or


(2-13) the tetrazine derivative of the formula (known from EP-A 0 005 912)



(clofentezine)

and/or

(2-14) the organotin derivative of the formula (known from DE-A 21 15 666)




(fenbutatin oxide)

15

and/or


(2-15) the sulfenamide of the formula (known from The Pesticide Manual, 11th edition, 1997, page 1208)



(tolylfluamid)

20 and/or

(2-16) the pyrimidyl phenol ethers of the formula (known from WO 94/02470, EP-A 0 833 991)



in which


R represents fluorine (2-16-a = 4-[(4-chloro- α,α,α -trifluoro-3-tolyl)oxy]-6-[(α,α,α -4-tetrafluoro-3-tolyl)oxy]pyrimidine)

R represents nitro (2-16-b = 4-[(4-chloro- α,α,α -trifluoro-3-tolyl)oxy]-6-[(α,α,α -trifluoro-4-nitro-3-tolyl)oxy]pyrimidine)

R represents bromine (2-16- = 4-[(4-chloro- α,α,α -trifluoro-3-tolyl)oxy]-6-[(α,α,α -trifluoro-4-bromo-3-tolyl)oxy]pyrimidine)

10 and/or

(2-17) the macrolide of the formula (known from EP-A 0 375 316)



(spinosad)

a mixture comprising, preferably,

85% spinosyn A (R = H)

15 15% spinosyn B (R = CH₃)

and/or


(2-18) ivermectin (known from EP-A 0 001 689)

and/or

(2-19) milbemectin (known from The Pesticide Manual, 11th edition, 1997, p. 846)


20 and/or

(2-20) endosulfan (known from DE-A 10 15 797)




and/or

(2-21) fenazaquin (known from EP-A 0 326 329)




and/or

(2-22) pyrimidifen (known from EP-A 0 196 524)




5 and/or

(2-23) triarathen (known from DE-A 27 24 494)



and/or


(2-24) tetradifon (known from US 2,812,281)



10


and/or

(2-25) propargite (known from US 3,272,854)




and/or

15 (2-26) hexythiazox (known from DE-A 30 37 105)




and/or

(2-27) bromopropylate (known from US 3,784,696)



and/or


(2-28) dicofol (known from US 2,812,280)



5

and/or

(2-29) chinomethionat (known from DE-A 11 00 372)



10 and at least one active compound from the group of the anthranilamides of the formula (II) are synergistically active and have very good insecticidal and acaricidal properties.

Surprisingly, the insecticidal and/or acaricidal action of the active compound combinations according to the invention is higher than the sum of the actions of the individual active
15 compounds. Thus, an unforeseeable true synergistic effect is present, and not just an addition of actions.

In addition to at least one active compound of the formula (I) or an active compound of group 2 (compounds (2-1) to (2-29)), the active compound combinations according to the invention
20 comprise at least one active compound of the formula (II).

Preference is given to active compound combinations comprising compounds of the formula (I) in which

- X represents C₁-C₄-alkyl, bromine, C₁-C₄-alkoxy or C₁-C₃-haloalkyl,
25 Y represents hydrogen, C₁-C₄-alkyl, fluorine, chlorine, bromine, C₁-C₄-alkoxy, C₁-C₃-haloalkyl,
Z represents C₁-C₄-alkyl, chlorine, bromine, C₁-C₄-alkoxy,
m represents a number 0-2,
A³ represents hydrogen or in each case optionally mono- to trifluoro-substituted straight-chain
30 or branched C₁-C₆-alkyl, C₂-C₆-alkenyl, C₁-C₄-alkoxy-C₁-C₂-alkyl, cycloalkyl having 3-8


ring atoms which may optionally be interrupted by oxygen and/or sulfur or represents benzyl or phenyl which is optionally mono- to disubstituted by fluorine, chlorine, bromine, C₁-C₂-alkyl, C₁-C₂-haloalkyl, C₁-C₂-alkoxy, C₁-C₂-haloalkoxy, nitro,

A⁴ represents hydrogen, C₁-C₂-alkyl or C₁-C₂-alkoxy-C₁-C₂-alkyl

5 or in which

A³ and A⁴ together with the carbon atom to which they are attached form a saturated or unsaturated 3- to 7-membered ring which is optionally interrupted by oxygen and/or sulfur and optionally mono- to disubstituted by fluorine, chlorine, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₂-haloalkyl, C₁-C₂-haloalkoxy or C₁-C₂-alkylthio,

10 G¹ represents hydrogen (a) or represents groups



in which

R²⁰ represents in each case optionally mono- to pentafluoro- or -chloro-substituted C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkylthio-C₁-C₄-alkyl or cycloalkyl having 3-6 ring atoms which may be interrupted by oxygen and/or sulfur atoms,

15

represents phenyl which is optionally mono- to disubstituted by fluorine, chlorine, bromine, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy,

20

represents benzyl which is optionally mono to disubstituted by fluorine, chlorine, bromine, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy,

represents pyridyl, pyrimidyl, thiazolyl or pyrazolyl, each of which is optionally mono- to disubstituted by chlorine, bromine and/or C₁-C₄-alkyl,

25

R²¹ represents C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₆-alkoxy-C₂-C₆-alkyl, C₁-C₆-polyalkoxy-C₂-C₆-alkyl, each of which is optionally mono- to pentasubstituted by fluorine or chlorine,

represents phenyl or benzyl, each of which is optionally mono- to disubstituted by fluorine, chlorine, bromine, nitro, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-haloalkyl,

30

R²² represents C₁-C₄-alkyl which is optionally mono- to pentasubstituted by fluorine or chlorine, represents phenyl or benzyl, each of which is optionally mono- to disubstituted by C₁-C₄-alkyl, fluorine, chlorine, bromine, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, nitro or cyano,

R²³ and R²⁴ independently of one another represent C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-

alkylamino, di(C₁-C₄)alkylamino, C₁-C₄-alkylthio, C₂-C₄-alkenylthio, C₃-C₆-cycloalkylthio, each of which is optionally mono- to trisubstituted by fluorine or chlorine, represent phenyl, phenoxy or phenylthio, each of which is optionally mono- to disubstituted by fluorine, chlorine, bromine, nitro, cyano, C₁-C₂-alkoxy, C₁-C₂-haloalkoxy, C₁-C₂-alkylthio, C₁-C₂-haloalkylthio, C₁-C₂-alkyl, C₁-C₂-haloalkyl,

R²⁵ and R²⁶ independently of one another represent C₁-C₆-alkyl, C₁-C₆-alkoxy, C₃-C₆-alkenyl, C₁-C₄-alkoxy-C₁-C₂-alkyl, each of which is optionally mono- to trisubstituted by fluorine or chlorine, represent benzyl which is optionally mono- to disubstituted by fluorine, chlorine, bromine, C₁-C₂-haloalkyl, C₁-C₄-alkyl or C₁-C₄-alkoxy or together represent a 5- to 6-membered ring which is optionally interrupted by oxygen or sulfur and which may optionally be substituted by C₁-C₂-alkyl,

and at least one active compound of the formula (II).

In the radicals in the preferred ranges named halogen, halogen preferably represents chlorine and fluorine.

Particular preference is given to active compound combinations comprising compounds of the formula (I) in which

X represents C₁-C₄-alkyl, C₁-C₄-alkoxy or trifluoromethyl,


Y represents hydrogen, C₁-C₄-alkyl, chlorine, bromine, C₁-C₄-alkoxy, C₁-C₂-haloalkyl,

Z represents C₁-C₄-alkyl, chlorine, bromine, C₁-C₄-alkoxy,

m represents 0 or 1,

A³ and A⁴ together with the carbon atom to which they are attached represent a saturated 5- to 6-membered ring which is optionally monosubstituted by C₁-C₄-alkyl or C₁-C₄-alkoxy,

G¹ represents hydrogen (a) or represents the groups



R²⁰ represents in each case optionally mono- to trifluoro- or -chloro-substituted C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₁-C₄-alkoxy-C₁-C₂-alkyl, or cycloalkyl having 3-6 ring atoms which may be interrupted by 1 to 2 oxygen atoms, represents phenyl which is optionally monosubstituted by fluorine, chlorine,

bromine, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, trifluoromethyl or trifluoromethoxy;
 R²¹ represents C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₁-C₄-alkoxy-C₂-C₄-alkyl,
 represents phenyl or benzyl, each of which is optionally monosubstituted by
 fluorine, chlorine, bromine, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy or trifluoromethyl,

5

and at least one active compound of the formula (II).

Very particular preference is given to active compound combinations comprising compounds of
 the formula (I) in which

10 X represents methyl, ethyl, methoxy, ethoxy or trifluoromethyl,

Y represents hydrogen, methyl, ethyl, chlorine, bromine, methoxy or trifluoromethyl,

Z represents methyl, ethyl, chlorine, bromine or methoxy,

m represents 0 or 1,

A³ and A⁴ together with the carbon atom to which they are attached form a saturated 5- to 6-
 15 membered ring which is optionally monosubstituted by methyl, ethyl, propyl, methoxy,
 ethoxy, propoxy, butoxy or isobutoxy,

G¹ represents hydrogen (a) or represents the groups



, in which

(b)

(c)


R²⁰ represents in each case mono- to trifluoro- or -chloro-substituted C₁-C₈-alkyl,
 20 C₂-C₈-alkenyl, C₁-C₃-alkoxy-C₁-C₂-alkyl, or cycloalkyl having 3-6 ring atoms
 which may be interrupted by 1 to 2 oxygen atoms,

represents phenyl which is optionally monosubstituted by fluorine, chlorine,
 bromine, methyl, methoxy, trifluoromethyl or trifluoromethoxy;

R²¹ represents C₁-C₈-alkyl, C₂-C₈-alkenyl, C₁-C₄-alkoxy-C₂-C₃-alkyl,
 25 represents phenyl or benzyl, each of which is optionally monosubstituted by
 fluorine, chlorine, bromine, nitro, methyl, methoxy or trifluoromethyl,


and at least one active compound of the formula (II).

30 Especially preferred are active compound combinations comprising the compound of the formula
 (I-1)



and at least one active compound of the formula (II).

Especially preferred are active compound combinations comprising the compound of the formula
5 (I-2)



and at least one active compound of the formula (II).

Depending inter alia on the nature of the substituents, the compounds of the formula (I) may be
10 present as geometrical and/or optical isomers or isomer mixtures of varying composition which, if appropriate, may be separated in a customary manner. The present invention provides both the pure isomers and the isomer mixtures, their preparation and use and also compositions comprising them. However, hereinbelow, for the sake of simplicity, only compounds of the formula (I) are referred to, although what is meant are both the pure compounds and, if appropriate, also mixtures
15 having varying proportions of isomeric compounds.

Especially preferred are active compound combinations comprising a compound selected from the
group of compounds (2-1) to (2-29) of group 2 and at least one active compound of the formula
(II).

20

Emphasis is given to active compound combinations comprising one of the compounds (2-2) abamectin, (2-5) diafenthiuron, (2-17) spinosad and (2-20) endosulfan and at least one active compound of the formula (II).


25 The anthranilamides of the formula (II) are likewise known compounds which are known from the following publications or embraced by them:

WO 01/70671, WO 03/015518, WO 03/015519, WO 03/016284, WO 03/016282, WO 03/016283,

WO 03/024222, WO 03/062226.

The generic formulae and definitions described in these publications and the individual compounds described therein are expressly incorporated into the present application by way of reference.

- 5 The anthranilamides can be summarized under the formula (II):



in which

A^1 and A^2 independently of one another represent oxygen or sulfur,

X^1 represents N or CR^{10} ,

- 10 R^1 represents hydrogen or represents in each case optionally mono- or polysubstituted C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl or C_3 - C_6 -cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R^6 , halogen, cyano, nitro, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylsulfinyl, C_1 - C_4 -alkylsulfonyl, C_2 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylamino, C_2 - C_8 -dialkylamino, C_3 - C_6 -cycloalkylamino, $(C_1$ - C_4 -alkyl) C_3 - C_6 -cycloalkylamino and R^{11} ,

- 15 R^2 represents hydrogen, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylamino, C_2 - C_8 -dialkylamino, C_3 - C_6 -cycloalkylamino, C_2 - C_6 -alkoxycarbonyl or C_2 - C_6 -alkylcarbonyl,

- 20 R^3 represents hydrogen, R^{11} or represents in each case optionally mono- or polysubstituted C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R^6 , halogen, cyano, nitro, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylsulfinyl, C_1 - C_4 -alkylsulfonyl, C_2 - C_6 -alkoxycarbonyl, C_2 - C_6 -alkylcarbonyl, C_3 - C_6 -trialkylsilyl, R^{11} , phenyl, phenoxy and a 5- or 6-membered heteroaromatic ring, where each phenyl, phenoxy and 5- or 6-membered heteroaromatic ring may optionally be substituted and where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R^{12} , or

- 25 R^2 and R^3 may be attached to one another and form the ring M,

R^4 represents hydrogen, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, C_1 - C_6 -

haloalkyl, C₂-C₆-haloalkenyl, C₂-C₆-haloalkynyl, C₃-C₆-halocycloalkyl, halogen, cyano, nitro, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl, C₁-C₄-haloalkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino, C₃-C₆-trialkylsilyl or
 5 represents in each case optionally mono- or polysubstituted phenyl, benzyl or phenoxy, where the substituents independently of one another may be selected from the group consisting of C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl, C₂-C₄-haloalkenyl, C₂-C₄-haloalkynyl, C₃-C₆-halocycloalkyl, halogen, cyano, nitro, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino, C₃-C₆-(alkyl)cycloalkylamino, C₂-C₄-alkylcarbonyl, C₂-C₆-alkoxycarbonyl, C₂-C₆-alkylaminocarbonyl, C₃-C₈-dialkylaminocarbonyl and C₃-C₆-trialkylsilyl,

R⁵ and R⁸ in each case independently of one another represent hydrogen, halogen or represent in each case optionally substituted C₁-C₄-alkyl, C₁-C₄-haloalkyl, R¹², G, J, -OJ, -OG, -S(O)_p-J, -S(O)_p-G, -S(O)_p-phenyl, where the substituents independently of one another may be
 15 selected from one to three radicals W or from the group consisting of R¹², C₁-C₁₀-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₄-alkoxy and C₁-C₄-alkylthio, where each substituent may be substituted by one or more substituents independently of one another selected from the group consisting of G, J, R⁶, halogen, cyano, nitro, amino, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl, C₁-C₄-haloalkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-trialkylsilyl, phenyl and phenoxy, where each phenyl or phenoxy ring may optionally be substituted and where the substituents independently of one another may be
 20 selected from one to three radicals W or one or more radicals R¹²,

G in each case independently of one another represents a 5- or 6-membered nonaromatic carbocyclic or heterocyclic ring which optionally contains one or two ring members from the group consisting of C(=O), SO and S(=O)₂ and which may optionally be substituted by one to four substituents independently of one another selected from the group consisting of C₁-C₂-alkyl, halogen, cyano, nitro and C₁-C₂-alkoxy, or independently of one another
 30 represents C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₃-C₇-cycloalkyl, (cyano)C₃-C₇-cycloalkyl, (C₁-C₄-alkyl)C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)C₁-C₄-alkyl, where each cycloalkyl, (alkyl)cycloalkyl and (cycloalkyl)alkyl may optionally be substituted by one or more halogen atoms,

J in each case independently of one another represents an optionally substituted 5- or 6-membered heteroaromatic ring, where the substituents independently of one another may
 35 be selected from one to three radicals W or one or more radicals R¹²,

- R⁶ independently of one another represent $-C(=E^1)R^{19}$, $-LC(=E^1)R^{19}$, $-C(=E^1)LR^{19}$, $-LC(=E^1)LR^{19}$, $-OP(=Q)(OR^{19})_2$, $-SO_2LR^{18}$ or $-LSO_2LR^{19}$, where each E¹ independently of the others represents O, S, N-R¹⁵, N-OR¹⁵, N-N(R¹⁵)₂, N-S=O, N-CN or N-NO₂,
- R⁷ represents hydrogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, halogen, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl, C₁-C₄-haloalkylsulfonyl,
- R⁹ represents C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylsulfinyl or halogen,
- R¹⁰ represents hydrogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, halogen, cyano or C₁-C₄-haloalkoxy,
- R¹¹ in each case independently of one another represents in each case optionally mono- to trisubstituted C₁-C₆-alkylthio, C₁-C₆-alkylsulfenyl, C₁-C₆-haloalkylthio, C₁-C₆-haloalkylsulfenyl, phenylthio or phenylsulfenyl, where the substituents independently of one another may be selected from the list consisting of W, $-S(O)_nN(R^{16})_2$, $-C(=O)R^{13}$, $-L(C=O)R^{14}$, $-S(C=O)LR^{14}$, $-C(=O)LR^{13}$, $-S(O)_nNR^{13}C(=O)R^{13}$, $-S(O)_nNR^{13}C(=O)LR^{14}$ and $-S(O)_nNR^{13}S(O)_2LR^{14}$,
- L in each case independently of one another represents O, NR¹⁸ or S,
- R¹² in each case independently of one another represents $-B(OR^{17})_2$, amino, SH, thiocyanato, C₃-C₈-trialkylsilyloxy, C₁-C₄-alkyl disulfide, $-SF_5$, $-C(=E^1)R^{19}$, $-LC(=E^1)R^{19}$, $-C(=E^1)LR^{19}$, $-LC(=E^1)LR^{19}$, $-OP(=Q)(OR^{19})_2$, $-SO_2LR^{19}$ or $-LSO_2LR^{19}$,
- Q represents O or S,
- R¹³ in each case independently of one another represents hydrogen or represents in each case optionally mono- or polysubstituted C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl or C₃-C₆-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R⁶, halogen, cyano, nitro, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino and (C₁-C₄-alkyl)C₃-C₆-cycloalkylamino,
- R¹⁴ in each case independently of one another represents in each case optionally mono- or polysubstituted C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₂-C₂₀-alkynyl or C₃-C₆-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R⁶, halogen, cyano, nitro, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino and (C₁-C₄-alkyl)C₃-C₆-cycloalkylamino or represents optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R¹²,
- R¹⁵ in each case independently of one another represents hydrogen or represents in each case mono- or polysubstituted C₁-C₆-haloalkyl or C₁-C₆-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, nitro,

- hydroxyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl, C₁-C₄-haloalkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₂-C₆-alkoxycarbonyl, C₂-C₆-alkylcarbonyl, C₃-C₆-trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R¹², or N(R¹⁵)₂ represents a cycle which forms the ring M,
- 5 R¹⁶ represents C₁-C₁₂-alkyl or C₁-C₁₂-haloalkyl, or N(R¹⁶)₂ represents a cycle which forms the ring M,
- R¹⁷ in each case independently of one another represents hydrogen or C₁-C₄-alkyl, or B(OR¹⁷)₂ represents a ring, where the two oxygen atoms are attached via a chain to two or three carbon atoms which are optionally substituted by one or two substituents independently of one another selected from the group consisting of methyl and C₂-C₆-alkoxycarbonyl,
- 10 R¹⁸ in each case independently of one another represents hydrogen, C₁-C₆-alkyl or C₁-C₆-haloalkyl, or N(R¹³)(R¹⁸) represents a cycle which forms the ring M,
- 15 R¹⁹ in each case independently of one another represents hydrogen or represents in each case optionally mono- or polysubstituted C₁-C₆-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, nitro, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl, C₁-C₄-haloalkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, CO₂H, C₂-C₆-alkoxycarbonyl, C₂-C₆-alkylcarbonyl, C₃-C₆-trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W, C₁-C₆-haloalkyl, C₃-C₆-cycloalkyl or phenyl or pyridyl, each of which is optionally mono- to trisubstituted by W,
- 20 M in each case represents an optionally mono- to tetrasubstituted ring which, in addition to the nitrogen atom which is attached to the substituent pair R¹³ and R¹⁸, (R¹⁵)₂ or (R¹⁶)₂, contains two to six carbon atoms and optionally additionally a further nitrogen, sulfur or oxygen atom, and where the substituents independently of one another may be selected from the group consisting of C₁-C₂-alkyl, halogen, cyano, nitro and C₁-C₂-alkoxy,
- 25 W in each case independently of one another represent C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl, C₂-C₄-haloalkenyl, C₂-C₄-haloalkynyl, C₃-C₆-halocycloalkyl, halogen, cyano, nitro, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino, (C₁-C₄-alkyl)C₃-C₆-cycloalkylamino, C₂-C₄-alkylcarbonyl, C₂-C₆-alkoxycarbonyl, CO₂H, C₂-C₆-alkylaminocarbonyl, C₃-C₈-dialkylaminocarbonyl or C₃-C₆-trialkylsilyl,
- 30 n in each case independently of one another represents 0 or 1,
- 35

p in each case independently of one another represents 0, 1 or 2.

If (a) R^5 represents hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_2 - C_6 -haloalkenyl, C_2 - C_6 -haloalkynyl, C_1 - C_4 -haloalkoxy, C_1 - C_4 -haloalkylthio or halogen and (b) R^8 represents hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_2 - C_6 -haloalkenyl, C_2 - C_6 -haloalkynyl, C_1 - C_4 -haloalkoxy, C_1 - C_4 -haloalkylthio, halogen, C_2 - C_4 -alkylcarbonyl, C_2 - C_6 -alkoxycarbonyl, C_2 - C_6 -alkylaminocarbonyl or C_3 - C_8 dialkylaminocarbonyl, (c) at least one substituent selected from the group consisting of R^6 , R^{11} and R^{12} is present and (d), if R^{12} is not present, at least one R^6 or R^{11} is different from C_2 - C_6 -alkylcarbonyl, C_2 - C_6 alkoxycarbonyl, C_2 - C_6 -alkylaminocarbonyl and C_3 - C_8 -dialkylaminocarbonyl.


10

The compounds of the general formula (II) include N-oxides and salts.

Depending *inter alia* on the nature of the substituents, the compounds of the formula (II) may be present as geometrical and/or optical isomers or isomer mixtures of varying composition which, if appropriate, may be separated in a customary manner. The present invention provides both the pure isomers and the isomer mixtures, their preparation and use and also compositions comprising them. However, hereinbelow, for the sake of simplicity, only compounds of the formula (II) are referred to, although what is meant are both the pure compounds and, if appropriate, also mixtures having varying proportions of isomeric compounds.

20

Preference is given to active compound combinations comprising compounds of the formula (II-1)



in which

- 25 R^2 represents hydrogen or C_1 - C_6 -alkyl,
 R^3 represents C_1 - C_6 -alkyl which is optionally substituted by a radical R^6 ,
 R^4 represents C_1 - C_4 -alkyl, C_1 - C_2 -haloalkyl, C_1 - C_2 -haloalkoxy or halogen,
 R^5 represents hydrogen, C_1 - C_4 -alkyl, C_1 - C_2 -haloalkyl, C_1 - C_2 -haloalkoxy or halogen,
 R^6 represents $-C(=E^2)R^{19}$, $-LC(=E^2)R^{19}$, $-C(=E^2)LR^{19}$ or $-LC(=E^2)LR^{19}$, where each E^2
30 independently of the others represents O, S, $N-R^{15}$, $N-OR^{15}$, $N-N(R^{15})_2$, and each L independently of the others represents O or NR^{18} ,
 R^7 represents C_1 - C_4 -haloalkyl or halogen,

- R⁹ represents C₁-C₂-haloalkyl, C₁-C₂-haloalkoxy, S(O)_pC₁-C₂-haloalkyl or halogen,
 R¹⁵ in each case independently of one another represents hydrogen or represents in each case optionally substituted C₁-C₆-haloalkyl or C₁-C₆-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl or C₁-C₄-haloalkylsulfonyl,
 R¹⁸ in each case represents hydrogen or C₁-C₄-alkyl,
 R¹⁹ in each case independently of one another represents hydrogen or C₁-C₆-alkyl,
 p independently of one another represents 0, 1, 2.

10

In the radical definitions mentioned as being preferred, halogen represents fluorine, chlorine, bromine and iodine, in particular fluorine, chlorine and bromine.

Particular preference is given to active compound combinations comprising compounds of the formula (II-1), in which


15

- R² represents hydrogen or methyl,
 R³ represents C₁-C₄-alkyl (in particular methyl, ethyl, n-, isopropyl, n-, iso-, sec-, tert-butyl),
 R⁴ represents methyl, trifluoromethyl, trifluoromethoxy, fluorine, chlorine, bromine or iodine,
 R⁵ represents hydrogen, fluorine, chlorine, bromine, iodine, trifluoromethyl or trifluoromethoxy,
 R⁷ represents chlorine or bromine,
 R⁹ represents trifluoromethyl, chlorine, bromine, difluoromethoxy or trifluoroethoxy.

20

Very particular preference is given to active compound combinations comprising the following compounds of the formula (II-1):

25




Example No.	R ²	R ³	R ⁴	R ⁵	R ⁷	R ⁹	m.p. (°C)
II-1-1	H	Me	Me	Cl	Cl	CF ₃	185-186
II-1-2	H	Me	Me	Cl	Cl	OCH ₂ CF ₃	207-208
II-1-3	H	Me	Me	Cl	Cl	Cl	225-226
II-1-4	H	Me	Me	Cl	Cl	Br	162-164
II-1-5	H	Me	Cl	Cl	Cl	CF ₃	155-157
II-1-6	H	Me	Cl	Cl	Cl	OCH ₂ CF ₃	192-195


Example No.	R ²	R ³	R ⁴	R ⁵	R ⁷	R ⁹	m.p. (°C)
II-1-7	H	Me	Cl	Cl	Cl	Cl	205-206
II-1-8	H	Me	Cl	Cl	Cl	Br	245-246
II-1-9	H	i-Pr	Me	Cl	Cl	CF ₃	195-196
II-1-10	H	i-Pr	Me	Cl	Cl	OCH ₂ CF ₃	217-218
II-1-11	H	i-Pr	Me	Cl	Cl	Cl	173-175
II-1-12	H	i-Pr	Me	Cl	Cl	Br	159-161
II-1-13	H	i-Pr	Cl	Cl	Cl	CF ₃	200-201
II-1-14	H	i-Pr	Cl	Cl	Cl	OCH ₂ CF ₃	232-235
II-1-15	H	i-Pr	Cl	Cl	Cl	Cl	197-199
II-1-16	H	i-Pr	Cl	Cl	Cl	Br	188-190
II-1-17	H	Et	Me	Cl	Cl	CF ₃	163-164
II-1-18	H	Et	Me	Cl	Cl	OCH ₂ CF ₃	205-207
II-1-19	H	Et	Me	Cl	Cl	Cl	199-200
II-1-20	H	Et	Me	Cl	Cl	Br	194-195
II-1-21	H	Et	Cl	Cl	Cl	CF ₃	201-202
II-1-22	H	Et	Cl	Cl	Cl	Cl	206-208
II-1-23	H	Et	Cl	Cl	Cl	Br	214-215
II-1-24	H	t-Bu	Me	Cl	Cl	CF ₃	223-225
II-1-25	H	t-Bu	Me	Cl	Cl	Cl	163-165
II-1-26	H	t-Bu	Me	Cl	Cl	Br	159-161
II-1-27	H	t-Bu	Cl	Cl	Cl	CF ₃	170-172
II-1-28	H	t-Bu	Cl	Cl	Cl	Cl	172-173
II-1-29	H	t-Bu	Cl	Cl	Cl	Br	179-180
II-1-30	H	Me	Me	Br	Cl	CF ₃	222-223
II-1-31	H	Et	Me	Br	Cl	CF ₃	192-193
II-1-32	H	i-Pr	Me	Br	Cl	CF ₃	197-198
II-1-33	H	t-Bu	Me	Br	Cl	CF ₃	247-248
II-1-34	H	Me	Me	Br	Cl	Cl	140-141
II-1-35	H	Et	Me	Br	Cl	Cl	192-194
II-1-36	H	i-Pr	Me	Br	Cl	Cl	152-153
II-1-37	H	t-Bu	Me	Br	Cl	Cl	224-225
II-1-38	H	Me	Me	Br	Cl	Br	147-149
II-1-39	H	Et	Me	Br	Cl	Br	194-196
II-1-40	H	i-Pr	Me	Br	Cl	Br	185-187
II-1-41	H	t-Bu	Me	Br	Cl	Br	215-221
II-1-42	H	Me	Me	I	Cl	CF ₃	199-200
II-1-43	H	Et	Me	I	Cl	CF ₃	199-200
II-1-44	H	i-Pr	Me	I	Cl	CF ₃	188-189
II-1-45	H	t-Bu	Me	I	Cl	CF ₃	242-243
II-1-46	H	Me	Me	I	Cl	Cl	233-234
II-1-47	H	Et	Me	I	Cl	Cl	196-197
II-1-48	H	i-Pr	Me	I	Cl	Cl	189-190
II-1-49	H	t-Bu	Me	I	Cl	Cl	228-229
II-1-50	H	Me	Me	I	Cl	Br	229-230
II-1-51	H	iPr	Me	I	Cl	Br	191-192
II-1-52	H	Me	Br	Br	Cl	CF ₃	162-163

Example No.	R ²	R ³	R ⁴	R ⁵	R ⁷	R ⁹	m.p. (°C)
II-1-53	H	Et	Br	Br	Cl	CF ₃	188-189
II-1-54	H	i-Pr	Br	Br	Cl	CF ₃	192-193
II-1-55	H	t-Bu	Br	Br	Cl	CF ₃	246-247
II-1-56	H	Me	Br	Br	Cl	Cl	188-190
II-1-57	H	Et	Br	Br	Cl	Cl	192-194
II-1-58	H	i-Pr	Br	Br	Cl	Cl	197-199
II-1-59	H	t-Bu	Br	Br	Cl	Cl	210-212
II-1-60	H	Me	Br	Br	Cl	Br	166-168
II-1-61	H	Et	Br	Br	Cl	Br	196-197
II-1-62	H	i-Pr	Br	Br	Cl	Br	162-163
II-1-63	H	t-Bu	Br	Br	Cl	Br	194-196
II-1-64	H	t-Bu	Cl	Br	Cl	CF ₃	143-145
II-1-65	Me	Me	Br	Br	Cl	Cl	153-155
II-1-66	Me	Me	Me	Br	Cl	CF ₃	207-208
II-1-67	Me	Me	Cl	Cl	Cl	Cl	231-232
II-1-68	Me	Me	Br	Br	Cl	Br	189-190
II-1-69	Me	Me	Cl	Cl	Cl	Br	216-218
II-1-70	Me	Me	Cl	Cl	Cl	CF ₃	225-227
II-1-71	Me	Me	Br	Br	Cl	CF ₃	228-229
II-1-72	H	i-Pr	Me	H	Cl	CF ₃	237-239


Especially preferred are active compound combinations comprising a compound of the formulae below



(II-1-1)




(II-1-2)




(II-1-3)


5




(II-1-4)




(II-1-5)




(II-1-6)




(II-1-7)




(II-1-8)




(II-1-9)




(II-1-11)




(II-1-12)




(II-1-13)




(II-1-15)




(II-1-16)




(II-1-19)












(II-1-21)






(II-1-22)




(II-1-23)




5







(II-1-43)




(II-1-44)




(II-1-50)




(II-1-51)




(II-1-52)




(II-1-53)




(II-1-54)




(II-1-55)




(II-1-56)




(II-1-57)




(II-1-58)




(II-1-60)




(II-1-61)




(II-1-62)




(II-1-64)




(II-1-65)




(II-1-66)




(II-1-67)



(II-1-68)




(II-1-69)




(II-1-70)

5



(II-1-71)



(II-1-72)

Emphasis is given to the following specifically mentioned active compound combinations (2-compound mixtures) comprising a compound of the formula (I) (group 1) or an acaricidally active compound of group 2 and a compound of the formula (II-1):

No.	Active compound combination comprising	No.	Active compound combination comprising
1a)	(I-1) and (II-1-1)	28a)	(I-1) and (II-1-39)
1b)	(I-2) and (II-1-1)	28b)	(I-2) and (II-1-39)
1c)	(2-2) abamectin and (II-1-1)	28c)	(2-2) abamectin and (II-1-39)
1d)	(2-5) diafenthiuron and (II-1-1)	28d)	(2-5) diafenthiuron and (II-1-39)
1e)	(2-17) spinosad and (II-1-1)	28e)	(2-17) spinosad and (II-1-39)
1f)	(2-20) endosulfan and (II-1-1)	28f)	(2-20) endosulfan and (II-1-39)
2a)	(I-1) and (II-1-2)	29a)	(I-1) and (II-1-40)
2b)	(I-2) and (II-1-2)	29b)	(I-2) and (II-1-40)
2c)	(2-2) abamectin and (II-1-2)	29c)	(2-2) abamectin and (II-1-40)
2d)	(2-5) diafenthiuron and (II-1-2)	29d)	(2-5) diafenthiuron and (II-1-40)
2e)	(2-17) spinosad and (II-1-2)	29e)	(2-17) spinosad and (II-1-40)
2f)	(2-20) endosulfan and (II-1-2)	29f)	(2-20) endosulfan and (II-1-40)
3a)	(I-1) and (II-1-3)	30a)	(I-1) and (II-1-42)
3b)	(I-2) and (II-1-3)	30b)	(I-2) and (II-1-42)
3c)	(2-2) abamectin and (II-1-3)	30c)	(2-2) abamectin and (II-1-42)
3d)	(2-5) diafenthiuron and (II-1-3)	30d)	(2-5) diafenthiuron and (II-1-42)
3e)	(2-17) spinosad and (II-1-3)	30e)	(2-17) spinosad and (II-1-42)
3f)	(2-20) endosulfan and (II-1-3)	30f)	(2-20) endosulfan and (II-1-42)
4a)	(I-1) and (II-1-4)	31a)	(I-1) and (II-1-43)
4b)	(I-2) and (II-1-4)	31b)	(I-2) and (II-1-43)
4c)	(2-2) abamectin and (II-1-4)	31c)	(2-2) abamectin and (II-1-43)
4d)	(2-5) diafenthiuron and (II-1-4)	31d)	(2-5) diafenthiuron and (II-1-43)
4e)	(2-17) spinosad and (II-1-4)	31e)	(2-17) spinosad and (II-1-43)
4f)	(2-20) endosulfan and (II-1-4)	31f)	(2-20) endosulfan and (II-1-43)
5a)	(I-1) and (II-1-5)	32a)	(I-1) and (II-1-44)
5b)	(I-2) and (II-1-5)	32b)	(I-2) and (II-1-44)
5c)	(2-2) abamectin and (II-1-5)	32c)	(2-2) abamectin and (II-1-44)
5d)	(2-5) diafenthiuron and (II-1-5)	32d)	(2-5) diafenthiuron and (II-1-44)
5e)	(2-17) spinosad and (II-1-5)	32e)	(2-17) spinosad and (II-1-44)
5f)	(2-20) endosulfan and (II-1-5)	32f)	(2-20) endosulfan and (II-1-44)
6a)	(I-1) and (II-1-6)	33a)	(I-1) and (II-1-50)
6b)	(I-2) and (II-1-6)	33b)	(I-2) and (II-1-50)
6c)	(2-2) abamectin and (II-1-6)	33c)	(2-2) abamectin and (II-1-50)
6d)	(2-5) diafenthiuron and (II-1-6)	33d)	(2-5) diafenthiuron and (II-1-50)
6e)	(2-17) spinosad and (II-1-6)	33e)	(2-17) spinosad and (II-1-50)
6f)	(2-20) endosulfan and (II-1-6)	33f)	(2-20) endosulfan and (II-1-50)
7a)	(I-1) and (II-1-7)	34a)	(I-1) and (II-1-51)
7b)	(I-2) and (II-1-7)	34b)	(I-2) and (II-1-51)
7c)	(2-2) abamectin and (II-1-7)	34c)	(2-2) abamectin and (II-1-51)

No.	Active compound combination comprising	No.	Active compound combination comprising
7d)	(2-5) diafenthiuron and (II-1-7)	34d)	(2-5) diafenthiuron and (II-1-51)
7e)	(2-17) spinosad and (II-1-7)	34e)	(2-17) spinosad and (II-1-51)
7f)	(2-20) endosulfan and (II-1-7)	34f)	(2-20) endosulfan and (II-1-51)
8a)	(I-1) and (II-1-8)	35a)	(I-1) and (II-1-52)
8b)	(I-2) and (II-1-8)	35b)	(I-2) and (II-1-52)
8c)	(2-2) abamectin and (II-1-8)	35c)	(2-2) abamectin and (II-1-52)
8d)	(2-5) diafenthiuron and (II-1-8)	35d)	(2-5) diafenthiuron and (II-1-52)
8e)	(2-17) spinosad and (II-1-8)	35e)	(2-17) spinosad and (II-1-52)
8f)	(2-20) endosulfan and (II-1-8)	35f)	(2-20) endosulfan and (II-1-52)
9a)	(I-1) and (II-1-9)	36a)	(I-1) and (II-1-53)
9b)	(I-2) and (II-1-9)	36b)	(I-2) and (II-1-53)
9c)	(2-2) abamectin and (II-1-9)	36c)	(2-2) abamectin and (II-1-53)
9d)	(2-5) diafenthiuron and (II-1-9)	36d)	(2-5) diafenthiuron and (II-1-53)
9e)	(2-17) spinosad and (II-1-9)	36e)	(2-17) spinosad and (II-1-53)
9f)	(2-20) endosulfan and (II-1-9)	36f)	(2-20) endosulfan and (II-1-53)
10a)	(I-1) and (II-1-11)	37a)	(I-1) and (II-1-54)
10b)	(I-2) and (II-1-11)	37b)	(I-2) and (II-1-54)
10c)	(2-2) abamectin and (II-1-11)	37c)	(2-2) abamectin and (II-1-54)
10d)	(2-5) diafenthiuron and (II-1-11)	37d)	(2-5) diafenthiuron and (II-1-54)
10e)	(2-17) spinosad and (II-1-11)	37e)	(2-17) spinosad and (II-1-54)
10f)	(2-20) endosulfan and (II-1-11)	37f)	(2-20) endosulfan and (II-1-54)
11a)	(I-1) and (II-1-12)	38a)	(I-1) and (II-1-55)
11b)	(I-2) and (II-1-12)	38b)	(I-2) and (II-1-55)
11c)	(2-2) abamectin and (II-1-12)	38c)	(2-2) abamectin and (II-1-55)
11d)	(2-5) diafenthiuron and (II-1-12)	38d)	(2-5) diafenthiuron and (II-1-55)
11e)	(2-17) spinosad and (II-1-12)	38e)	(2-17) spinosad and (II-1-55)
11f)	(2-20) endosulfan and (II-1-12)	38f)	(2-20) endosulfan and (II-1-55)
12a)	(I-1) and (II-1-13)	39a)	(I-1) and (II-1-56)
12b)	(I-2) and (II-1-13)	39b)	(I-2) and (II-1-56)
12c)	(2-2) abamectin and (II-1-13)	39c)	(2-2) abamectin and (II-1-56)
12d)	(2-5) diafenthiuron and (II-1-13)	39d)	(2-5) diafenthiuron and (II-1-56)
12e)	(2-17) spinosad and (II-1-13)	39e)	(2-17) spinosad and (II-1-56)
12f)	(2-20) endosulfan and (II-1-13)	39f)	(2-20) endosulfan and (II-1-56)
13a)	(I-1) and (II-1-15)	40a)	(I-1) and (II-1-57)
13b)	(I-2) and (II-1-15)	40b)	(I-2) and (II-1-57)
13c)	(2-2) abamectin and (II-1-15)	40c)	(2-2) abamectin and (II-1-57)
13d)	(2-5) diafenthiuron and (II-1-15)	40d)	(2-5) diafenthiuron and (II-1-57)
13e)	(2-17) spinosad and (II-1-15)	40e)	(2-17) spinosad and (II-1-57)
13f)	(2-20) endosulfan and (II-1-15)	40f)	(2-20) endosulfan and (II-1-57)
14a)	(I-1) and (II-1-16)	41a)	(I-1) and (II-1-58)
14b)	(I-2) and (II-1-16)	41b)	(I-2) and (II-1-58)
14c)	(2-2) abamectin and (II-1-16)	41c)	(2-2) abamectin and (II-1-58)
14d)	(2-5) diafenthiuron and (II-1-16)	41d)	(2-5) diafenthiuron and (II-1-58)
14e)	(2-17) spinosad and (II-1-16)	41e)	(2-17) spinosad and (II-1-58)

No.	Active compound combination comprising	No.	Active compound combination comprising
14f)	(2-20) endosulfan and (II-1-16)	41f)	(2-20) endosulfan and (II-1-58)
15a)	(I-1) and (II-1-19)	42a)	(I-1) and (II-1-60)
15b)	(I-2) and (II-1-19)	42b)	(I-2) and (II-1-60)
15c)	(2-2) abamectin and (II-1-19)	42c)	(2-2) abamectin and (II-1-60)
15d)	(2-5) diafenthiuron and (II-1-19)	42d)	(2-5) diafenthiuron and (II-1-60)
15e)	(2-17) spinosad and (II-1-19)	42e)	(2-17) spinosad and (II-1-60)
15f)	(2-20) endosulfan and (II-1-19)	42f)	(2-20) endosulfan and (II-1-60)
16a)	(I-1) and (II-1-21)	43a)	(I-1) and (II-1-61)
16b)	(I-2) and (II-1-21)	43b)	(I-2) and (II-1-61)
16c)	(2-2) abamectin and (II-1-21)	43c)	(2-2) abamectin and (II-1-61)
16d)	(2-5) diafenthiuron and (II-1-21)	43d)	(2-5) diafenthiuron and (II-1-61)
16e)	(2-17) spinosad and (II-1-21)	43e)	(2-17) spinosad and (II-1-61)
16f)	(2-20) endosulfan and (II-1-21)	43f)	(2-20) endosulfan and (II-1-61)
17a)	(I-1) and (II-1-22)	44a)	(I-1) and (II-1-62)
17b)	(I-2) and (II-1-22)	44b)	(I-2) and (II-1-62)
17c)	(2-2) abamectin and (II-1-22)	44c)	(2-2) abamectin and (II-1-62)
17d)	(2-5) diafenthiuron and (II-1-22)	44d)	(2-5) diafenthiuron and (II-1-62)
17e)	(2-17) spinosad and (II-1-22)	44e)	(2-17) spinosad and (II-1-62)
17f)	(2-20) endosulfan and (II-1-22)	44f)	(2-20) endosulfan and (II-1-62)
18a)	(I-1) and (II-1-23)	45a)	(I-1) and (II-1-64)
18b)	(I-2) and (II-1-23)	45b)	(I-2) and (II-1-64)
18c)	(2-2) abamectin and (II-1-23)	45c)	(2-2) abamectin and (II-1-64)
18d)	(2-5) diafenthiuron and (II-1-23)	45d)	(2-5) diafenthiuron and (II-1-64)
18e)	(2-17) spinosad and (II-1-23)	45e)	(2-17) spinosad and (II-1-64)
18f)	(2-20) endosulfan and (II-1-23)	45f)	(2-20) endosulfan and (II-1-64)
19a)	(I-1) and (II-1-24)	46a)	(I-1) and (II-1-65)
19b)	(I-2) and (II-1-24)	46b)	(I-2) and (II-1-65)
19c)	(2-2) abamectin and (II-1-24)	46c)	(2-2) abamectin and (II-1-65)
19d)	(2-5) diafenthiuron and (II-1-24)	46d)	(2-5) diafenthiuron and (II-1-65)
19e)	(2-17) spinosad and (II-1-24)	46e)	(2-17) spinosad and (II-1-65)
19f)	(2-20) endosulfan and (II-1-24)	46f)	(2-20) endosulfan and (II-1-65)
20a)	(I-1) and (II-1-26)	47a)	(I-1) and (II-1-66)
20b)	(I-2) and (II-1-26)	47b)	(I-2) and (II-1-66)
20c)	(2-2) abamectin and (II-1-26)	47c)	(2-2) abamectin and (II-1-66)
20d)	(2-5) diafenthiuron and (II-1-26)	47d)	(2-5) diafenthiuron and (II-1-66)
20e)	(2-17) spinosad and (II-1-26)	47e)	(2-17) spinosad and (II-1-66)
20f)	(2-20) endosulfan and (II-1-26)	47f)	(2-20) endosulfan and (II-1-66)
21a)	(I-1) and (II-1-27)	48a)	(I-1) and (II-1-67)
21b)	(I-2) and (II-1-27)	48b)	(I-2) and (II-1-67)
21c)	(2-2) abamectin and (II-1-27)	48c)	(2-2) abamectin and (II-1-67)
21d)	(2-5) diafenthiuron and (II-1-27)	48d)	(2-5) diafenthiuron and (II-1-67)
21e)	(2-17) spinosad and (II-1-27)	48e)	(2-17) spinosad and (II-1-67)
21f)	(2-20) endosulfan and (II-1-27)	48f)	(2-20) endosulfan and (II-1-67)
22a)	(I-1) and (II-1-29)	49a)	(I-1) and (II-1-68)

No.	Active compound combination comprising	No.	Active compound combination comprising
22b)	(I-2) and (II-1-29)	49b)	(I-2) and (II-1-68)
22c)	(2-2) abamectin and (II-1-29)	49c)	(2-2) abamectin and (II-1-68)
22d)	(2-5) diafenthiuron and (II-1-29)	49d)	(2-5) diafenthiuron and (II-1-68)
22e)	(2-17) spinosad and (II-1-29)	49e)	(2-17) spinosad and (II-1-68)
22f)	(2-20) endosulfan and (II-1-29)	49f)	(2-20) endosulfan and (II-1-68)
23a)	(I-1) and (II-1-30)	50a)	(I-1) and (II-1-69)
23b)	(I-2) and (II-1-30)	50b)	(I-2) and (II-1-69)
23c)	(2-2) abamectin and (II-1-30)	50c)	(2-2) abamectin and (II-1-69)
23d)	(2-5) diafenthiuron and (II-1-30)	50d)	(2-5) diafenthiuron and (II-1-69)
23e)	(2-17) spinosad and (II-1-30)	50e)	(2-17) spinosad and (II-1-69)
23f)	(2-20) endosulfan and (II-1-30)	50f)	(2-20) endosulfan and (II-1-69)
24a)	(I-1) and (II-1-31)	51a)	(I-1) and (II-1-70)
24b)	(I-2) and (II-1-31)	51b)	(I-2) and (II-1-70)
24c)	(2-2) abamectin and (II-1-31)	51c)	(2-2) abamectin and (II-1-70)
24d)	(2-5) diafenthiuron and (II-1-31)	51d)	(2-5) diafenthiuron and (II-1-70)
24e)	(2-17) spinosad and (II-1-31)	51e)	(2-17) spinosad and (II-1-70)
24f)	(2-20) endosulfan and (II-1-31)	51f)	(2-20) endosulfan and (II-1-70)
25a)	(I-1) and (II-1-32)	52a)	(I-1) and (II-1-71)
25b)	(I-2) and (II-1-32)	52b)	(I-2) and (II-1-71)
25c)	(2-2) abamectin and (II-1-32)	52c)	(2-2) abamectin and (II-1-71)
25d)	(2-5) diafenthiuron and (II-1-32)	52d)	(2-5) diafenthiuron and (II-1-71)
25e)	(2-17) spinosad and (II-1-32)	52e)	(2-17) spinosad and (II-1-71)
25f)	(2-20) endosulfan and (II-1-32)	52f)	(2-20) endosulfan and (II-1-71)
26a)	(I-1) and (II-1-33)	53a)	(I-1) and (II-1-72)
26b)	(I-2) and (II-1-33)	53b)	(I-2) and (II-1-72)
26c)	(2-2) abamectin and (II-1-33)	53c)	(2-2) abamectin and (II-1-72)
26d)	(2-5) diafenthiuron and (II-1-33)	53d)	(2-5) diafenthiuron and (II-1-72)
26e)	(2-17) spinosad and (II-1-33)	53e)	(2-17) spinosad and (II-1-72)
26f)	(2-20) endosulfan and (II-1-33)	53f)	(2-20) endosulfan and (II-1-72)
27a)	(I-1) and (II-1-38)		
27b)	(I-2) and (II-1-38)		
27c)	(2-2) abamectin and (II-1-38)		
27d)	(2-5) diafenthiuron and (II-1-38)		
27e)	(2-17) spinosad and (II-1-38)		
27f)	(2-20) endosulfan and (II-1-38)		

However, the general or preferred radical definitions or illustrations listed above can also be combined with one another as desired, i.e. between their respective ranges and preferred ranges. The definitions apply to the end products and, correspondingly, to precursors and intermediates.

5

Preference according to the invention is given to active compound combinations comprising compounds of the formula (I) or a compound of group 2 and at least one compound of the formula

(II), where the individual radicals are a combination of the meanings listed above as being preferred (preferable).

5 Particular preference according to the invention is given to active compound combinations comprising compounds of the formula (I) or a compound of group 2 and at least one compound of the formula (II), where the individual radicals are a combination of the meanings listed above as being particularly preferred.

10 Very particular preference according to the invention is given to active compound combinations comprising compounds of the formula (I) or a compound of group 2 and at least one compound of the formula (II), where the individual radicals are a combination of the meanings listed above as being very particularly preferred.

15 Saturated or unsaturated hydrocarbon radicals, such as alkyl or alkenyl, can in each case be straight-chain or branched as far as this is possible, including in combination with heteroatoms, such as, for example, in alkoxy.

20 Optionally substituted radicals can be mono- or polysubstituted, where in the case of polysubstitution the substituents can be identical or different.

In addition, the active compound combinations may also comprise further fungicidally, acaricidally or insecticidally active additives.

25 If the active compounds in the active compound combinations according to the invention are present in certain weight ratios, the synergistic effect is particularly pronounced. However, the weight ratios of the active compounds in the active compound combinations can be varied within a relatively wide range. In general, the combinations according to the invention comprise active compounds of the formula (I) or a compound of group 2 and the mixing partner of the formula (II) in the preferred and particularly preferred mixing ratios given:

30 The preferred mixing ratio is from 500:1 to 1:50.

The particularly preferred mixing ratio is from 25:1 to 1:10.

35 The mixing ratios are based on weight ratios. The ratio is to be understood as meaning active compound of the formula (I):mixing partner of the formula (II) or as ratio of a compound of group 2:mixing partner of the formula (II).

- The active compound combinations according to the invention are suitable for controlling animal pests, preferably arthropods and nematodes, in particular insects and arachnids, found in agriculture, in animal health, in forests, in the protection of stored products and materials and in the hygiene sector. They are active against normally sensitive and resistant species, and against all or individual developmental stages. The abovementioned pests include:
- 5 From the order of the Isopoda, for example, *Oniscus asellus*, *Armadillidium vulgare*, *Porcellio scaber*.
- From the order of the Diplopoda, for example, *Blaniulus guttulatus*.
- From the order of the Chilopoda, for example, *Geophilus carpophagus*, *Scutigera* spp.
- 10 From the order of the Symphyla, for example, *Scutigera* spp.
- From the order of the Thysanura, for example, *Lepisma saccharina*.
- From the order of the Collembola, for example, *Onychiurus armatus*.
- From the order of the Orthoptera, for example, *Acheta domesticus*, *Gryllotalpa* spp., *Locusta migratoria migratorioides*, *Melanoplus* spp., *Schistocerca gregaria*.
- 15 From the order of the Blattaria, for example, *Blatta orientalis*, *Periplaneta americana*, *Leucophaea maderae*, *Blattella germanica*.
- From the order of the Dermaptera, for example, *Forficula auricularia*.
- From the order of the Isoptera, for example, *Reticulitermes* spp.
- From the order of the Phthiraptera, for example, *Pediculus humanus corporis*, *Haematopinus* spp.,
- 20 *Linognathus* spp., *Trichodectes* spp., *Damalinia* spp.
- From the order of the Thysanoptera, for example, *Hercinothrips femoralis*, *Thrips tabaci*, *Thrips palmi*, *Frankliniella occidentalis*.
- From the order of the Heteroptera, for example, *Eurygaster* spp., *Dysdercus intermedius*, *Piesma quadrata*, *Cimex lectularius*, *Rhodnius prolixus*, *Triatoma* spp.
- 25 From the order of the Homoptera, for example, *Aleurodes brassicae*, *Bemisia tabaci*, *Trialeurodes vaporariorum*, *Aphis gossypii*, *Brevicoryne brassicae*, *Cryptomyzus ribis*, *Aphis fabae*, *Aphis pomi*, *Eriosoma lanigerum*, *Hyalopterus arundinis*, *Phylloxera vastatrix*, *Pemphigus* spp., *Macrosiphum avenae*, *Myzus* spp., *Phorodon humuli*, *Rhopalosiphum padi*, *Empoasca* spp., *Euscelis bilobatus*, *Nephotettix cincticeps*, *Lecanium corni*, *Saissetia oleae*, *Laodelphax striatellus*,
- 30 *Nilaparvata lugens*, *Aonidiella aurantii*, *Aspidiotus hederae*, *Pseudococcus* spp., *Psylla* spp.
- From the order of the Lepidoptera, for example, *Pectinophora gossypiella*, *Bupalus piniarius*, *Cheimatobia brumata*, *Lithocolletis blancardella*, *Hyponomeuta padella*, *Plutella xylostella*, *Malacosoma neustria*, *Euproctis chrysothoea*, *Lymantria* spp., *Bucculatrix thurberiella*, *Phyllocnistis citrella*, *Agrotis* spp., *Euxoa* spp., *Feltia* spp., *Earias insulana*, *Heliiothis* spp.,
- 35 *Mamestra brassicae*, *Panolis flammea*, *Spodoptera* spp., *Trichoplusia ni*, *Carpocapsa pomonella*, *Pieris* spp., *Chilo* spp., *Pyrausta nubilalis*, *Ephestia kuehniella*, *Galleria mellonella*, *Tineola*

Usselliella, *Tinea pellionella*, *Hofmannophila pseudospretella*, *Cacoecia podana*, *Capua reticulana*, *Choristoneura fumiferana*, *Clysia ambiguella*, *Homona magnanima*, *Tortrix viridana*, *Cnaphalocerus* spp., *Oulema oryzae*.

From the order of the Coleoptera, for example, *Anobium punctatum*, *Rhizopertha dominica*,
 5 *Bruchidius obtectus*, *Acanthoscelides obtectus*, *Hylotrupes bajulus*, *Agelastica alni*, *Leptinotarsa decemlineata*, *Phaedon cochleariae*, *Diabrotica* spp., *Psylliodes chrysocephala*, *Epilachna varivestis*, *Atomaria* spp., *Oryzaephilus surinamensis*, *Anthonomus* spp., *Sitophilus* spp., *Otiorrhynchus sulcatus*, *Cosmopolites sordidus*, *Ceuthorrhynchus assimilis*, *Hypera postica*, *Dermestes* spp., *Trogoderma* spp., *Anthrenus* spp., *Attagenus* spp., *Lyctus* spp., *Meligethes aeneus*,
 10 *Ptinus* spp., *Niptus hololeucus*, *Gibbium psylloides*, *Tribolium* spp., *Tenebrio molitor*, *Agriotes* spp., *Conoderus* spp., *Melolontha melolontha*, *Amphimallon solstitialis*, *Costelytra zealandica*, *Lissorhoptrus oryzophilus*.

From the order of the Hymenoptera, for example, *Diprion* spp., *Hoplocampa* spp., *Lasius* spp., *Monomorium pharaonis*, *Vespa* spp.

15 From the order of the Diptera, for example, *Aedes* spp., *Anopheles* spp., *Culex* spp., *Drosophila melanogaster*, *Musca* spp., *Fannia* spp., *Calliphora erythrocephala*, *Lucilia* spp., *Chrysomyia* spp., *Cuterebra* spp., *Gastrophilus* spp., *Hyppobosca* spp., *Stomoxys* spp., *Oestrus* spp., *Hypoderma* spp., *Tabanus* spp., *Tannia* spp., *Bibio hortulanus*, *Oscinella frit*, *Phorbia* spp., *Pegomyia hyoscyami*, *Ceratitis capitata*, *Dacus oleae*, *Tipula paludosa*, *Hylemyia* spp., *Liriomyza* spp.

20 From the order of the Siphonaptera, for example, *Xenopsylla cheopis*, *Ceratophyllus* spp.

From the class of the Arachnida, for example, *Scorpio maurus*, *Latrodectus mactans*, *Acarus siro*, *Argas* spp., *Ornithodoros* spp., *Dermanyssus gallinae*, *Eriophyes ribis*, *Phyllocoptruta oleivora*, *Boophilus* spp., *Rhipicephalus* spp., *Amblyomma* spp., *Hyalomma* spp., *Ixodes* spp., *Psoroptes* spp., *Chorioptes* spp., *Sarcoptes* spp., *Tarsonemus* spp., *Bryobia praetiosa*, *Panonychus* spp.,

25 *Tetranychus* spp., *Hemitarsonemus* spp., *Brevipalpus* spp.

The plant-parasitic nematodes include, for example, *Pratylenchus* spp., *Radopholus similis*, *Ditylenchus dipsaci*, *Tylenchulus semipenetrans*, *Heterodera* spp., *Globodera* spp., *Meloidogyne* spp., *Aphelenchoides* spp., *Longidorus* spp., *Xiphinema* spp., *Trichodorus* spp., *Bursaphelenchus*
 30 spp.

The active compound combinations can be converted into the customary formulations such as solutions, emulsions, wettable powders, suspensions, powders, dusts, pastes, soluble powders, granules, suspension-emulsion concentrates, natural and synthetic materials impregnated with
 35 active compound, and microencapsulations in polymeric materials.

These formulations are produced in a known manner, for example by mixing the active compounds with extenders, that is, liquid solvents and/or solid carriers, optionally with the use of surfactants, that is, emulsifiers and/or dispersants, and/or foam formers.

- 5 If the extender used is water, it is also possible, for example, to use organic solvents as cosolvents. The following are essentially suitable as liquid solvents: aromatics such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example mineral oil fractions, mineral and vegetable oils, alcohols
10 such as butanol or glycol and their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethylformamide and dimethyl sulfoxide, or else water.

Suitable solid carriers are:

- 15 for example ammonium salts and ground natural minerals such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground synthetic materials such as highly disperse silica, alumina and silicates; suitable solid carriers for granules are: for example crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, or else synthetic granules of inorganic and organic meals, and granules of organic material such as
20 sawdust, coconut shells, corn cobs and tobacco stalks; suitable emulsifiers and/or foam formers are: for example nonionic and anionic emulsifiers such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulfonates, alkyl sulfates, arylsulfonates, or else protein hydrolysates; suitable dispersants are: for example
lignosulfite waste liquors and methylcellulose.

- 25 Tackifiers such as carboxymethylcellulose and natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, or else natural phospholipids such as cephalins and lecithins and synthetic phospholipids can be used in the formulations. Other possible additives are mineral and vegetable oils.

- 30 It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic colorants such as alizarin colorants, azo colorants and metal phthalocyanine colorants, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

- 35 The formulations generally comprise between 0.1 and 95% by weight of active compound,

preferably between 0.5 and 90%.

The active compound combinations according to the invention can be present in their commercially available formulations and in the use forms, prepared from these formulations, as a
5 mixture with other active compounds, such as insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth-regulating substances or herbicides. The insecticides include, for example, phosphates, carbamates, carboxylates, chlorinated hydrocarbons, phenylureas and substances produced by microorganisms, inter alia.

10 Mixtures with other known active compounds such as herbicides or with fertilizers and growth regulators are also possible.

When used as insecticides, the active compound combinations according to the invention can furthermore be present in their commercially available formulations and in the use forms, prepared
15 from these formulations, as a mixture with synergists. Synergists are compounds which increase the action of the active compounds, without it being necessary for the synergist added to be active itself.

The active compound content of the use forms prepared from the commercially available
20 formulations can vary within wide limits. The active compound concentration of the use forms can be from 0.0000001 to 95% by weight of active compound, preferably between 0.0001 and 1% by weight.

The compounds are employed in a customary manner appropriate for the use forms.
25

When used against hygiene pests and stored-product pests, the active compound combinations are distinguished by an excellent residual action on wood and clay as well as good stability to alkali on limed substrates.

30 The active compound combinations according to the invention are not only active against plant pests, hygiene pests and stored-product pests, but also, in the veterinary medicine sector, against animal parasites (ectoparasites) such as hard ticks, soft ticks, mange mites, harvest mites, flies (stinging and licking), parasitizing fly larvae, lice, head lice, bird lice and fleas. These parasites include:

35 From the order of the Anoplurida, for example, Haematopinus spp., Linognathus spp., Pediculus spp., Phtirus spp., Solenopotes spp.

From the order of the Mallophagida and the suborders Amblycerina and Ischnocerina, for example, Trimenopon spp., Menopon spp., Trinoton spp., Bovicola spp., Werneckiella spp., Lepikentron spp., Damalina spp., Trichodectes spp., Felicola spp.

From the order Diptera and the suborders Nematocerina and Brachyicerina, for example, Aedes
5 spp., Anopheles spp., Culex spp., Simulium spp., Eusimulium spp., Phlebotomus spp., Lutzomyia
spp., Culicoides spp., Chrysops spp., Hybomitra spp., Atylotus spp., Tabanus spp., Haematopota
spp., Philipomyia spp., Braula spp., Musca spp., Hydrotaea spp., Stomoxys spp., Haematobia spp.,
Morellia spp., Fannia spp., Glossina spp., Calliphora spp., Lucilia spp., Chrysomyia spp.,
Wohlfahrtia spp., Sarcophaga spp., Oestrus spp., Hypoderma spp., Gasterophilus spp., Hippobosca
10 spp., Lipoptena spp., Melophagus spp.

From the order of the Siphonapterida, for example, Pulex spp., Ctenocephalides spp., Xenopsylla
spp., Ceratophyllus spp.

From the order of the Heteropterida, for example, Cimex spp., Triatoma spp., Rhodnius spp.,
Panstrongylus spp.

15 From the order of the Blattarida, for example, Blatta orientalis, Periplaneta americana, Blattella
germanica, Supella spp.

From the subclass of the Acaria (Acarida) and the orders of the Meta- and Mesostigmata, for
example, Argas spp., Ornithodoros spp., Otobius spp., Ixodes spp., Amblyomma spp., Boophilus
spp., Dermacentor spp., Haemophysalis spp., Hyalomma spp., Rhipicephalus spp., Dermanyssus
20 spp., Raillietia spp., Pneumonyssus spp., Sternostoma spp., Varroa spp.

From the order of the Actinedida (Prostigmata) and Acaridida (Astigmata), for example, Acarapis
spp., Cheyletiella spp., Ornithocheyletia spp., Myobia spp., Psorergates spp., Demodex spp.,
Trombicula spp., Listrophorus spp., Acarus spp., Tyrophagus spp., Caloglyphus spp., Hypodectes
spp., Pterolichus spp., Psoroptes spp., Chorioptes spp., Otodectes spp., Sarcoptes spp., Notoedres
25 spp., Knemidocoptes spp., Cytodites spp., Laminosioptes spp.

The active compound combinations according to the invention are also suitable for controlling
arthropods which attack agricultural livestock such as, for example, cattle, sheep, goats, horses,
pigs, donkeys, camels, buffaloes, rabbits, chickens, turkeys, ducks, geese, honey-bees, other
30 domestic animals such as, for example, dogs, cats, caged birds, aquarium fish and so-called
experimental animals such as, for example, hamsters, guinea pigs, rats and mice. By controlling
these arthropods, cases of death and reductions in productivity (for meat, milk, wool, hides, eggs,
honey and the like) should be diminished, so that more economical and simpler animal husbandry
is possible by the use of the active compound combinations according to the invention.

35

The active compound combinations according to the invention are used in the veterinary sector in a

known manner by enteral administration in the form of, for example, tablets, capsules, potions, drenches, granules, pastes, boluses, the feed-through method, suppositories, by parenteral administration such as, for example, by injections (intramuscularly, subcutaneously, intravenously, intraperitoneally and the like), implants, by nasal administration, by dermal administration in the form of, for example, immersing or dipping, spraying, pouring-on, spotting-on, washing, dusting, and with the aid of active-compound-comprising moulded articles such as collars, ear tags, tail tags, limb bands, halters, marking devices and the like.

When used for cattle, poultry, domestic animals and the like, the active compound combinations can be applied as formulations (for example powders, emulsions, flowables) comprising the active compounds in an amount of 1 to 80% by weight, either directly or after 100- to 10 000-fold dilution, or they may be used as a chemical dip.

Moreover, it has been found that the active compound combinations according to the invention show a potent insecticidal action against insects which destroy industrial materials.

The following insects may be mentioned by way of example and with preference, but not by way of limitation:

Beetles such as *Hylotrupes bajulus*, *Chlorophorus pilosis*, *Anobium punctatum*, *Xestobium rufovillosum*, *Ptilinus pecticornis*, *Dendrobium pertinex*, *Ernobius mollis*, *Priobium carpini*, *Lyctus brunneus*, *Lyctus africanus*, *Lyctus planicollis*, *Lyctus linearis*, *Lyctus pubescens*, *Trogoxylon aequale*, *Minthes rugicollis*, *Xyleborus spec.*, *Tryptodendron spec.*, *Apate monachus*, *Bostrychus capucins*, *Heterobostrychus brunneus*, *Sinoxylon spec.*, *Dinoderus minutus*.

Dermapterans such as *Sirex juvencus*, *Urocerus gigas*, *Urocerus gigas taignus*, *Urocerus augur*.

Termites such as *Kaloterms flavicollis*, *Cryptoterms brevis*, *Heteroterms indicola*, *Reticuliterms flavipes*, *Reticuliterms santonensis*, *Reticuliterms lucifugus*, *Mastoterms darwiniensis*, *Zootermopsis nevadensis*, *Coptoterms formosanus*.

Bristle-tails such as *Lepisma saccharina*.

Industrial materials in the present context are understood as meaning non-living materials such as, preferably, polymers, adhesives, glues, paper and board, leather, wood, timber products and paints.

The material which is to be protected from insect attack is very particularly preferably wood and timber products.

35

Wood and timber products which can be protected by the composition according to the invention,

or mixtures comprising it, are to be understood as meaning, for example:

Construction timber, wooden beams, railway sleepers, bridge components, jetties, vehicles made of wood, boxes, pallets, containers, telephone poles, wood lagging, windows and doors made of wood, plywood, chipboard, joinery, or timber products which quite generally are used in house
5 construction or building joinery.

The active compound combinations can be used as such, in the form of concentrates or generally customary formulations such as powders, granules, solutions, suspensions, emulsions or pastes.

10 The abovementioned formulations can be prepared in a manner known per se, for example by mixing the active compounds with at least one solvent or diluent, emulsifier, dispersant and/or binder or fixative, water repellent, if desired desiccants and UV stabilizers, and if desired colorants and pigments and other processing auxiliaries.

15 The insecticidal compositions or concentrates used for protecting wood and timber products comprise the active compound according to the invention in a concentration of 0.0001 to 95% by weight, in particular 0.001 to 60% by weight.

The amount of composition or concentrate employed depends on the species and the abundance of
20 the insects and on the medium. The optimal quantity to be employed can be determined in each case by test series upon application. In general, however, it will suffice to employ 0.0001 to 20% by weight, preferably 0.001 to 10% by weight, of the active compound, based on the material to be protected.

25 A suitable solvent and/or diluent is an organochemical solvent or solvent mixture and/or an oily or oil-type organochemical solvent or solvent mixture of low volatility and/or a polar organochemical solvent or solvent mixture and/or water and, if appropriate, an emulsifier and/or wetter.

30 Organochemical solvents which are preferably employed are oily or oil-type solvents with an evaporation number of above 35 and a flash point of above 30°C, preferably above 45°C. Such oily and oil-type solvents which are insoluble in water and of low volatility and which are used are suitable mineral oils or their aromatic fractions or mineral-oil-containing solvent mixtures, preferably white spirit, petroleum and/or alkylbenzene.

35 Mineral oils with a boiling range of 170 to 220°C, white spirit with a boiling range of 170 to 220°C, spindle oil with a boiling range of 250 to 350°C, petroleum and aromatics with a boiling

range of 160 to 280°C, oil of turpentine, and the like are advantageously used.

In a preferred embodiment, liquid aliphatic hydrocarbons with a boiling range of 180 to 210°C or high-boiling mixtures of aromatic and aliphatic hydrocarbons with a boiling range of 180 to 220°C
5 and/or spindle oil and/or monochloronaphthalene, preferably α -monochloronaphthalene, are used.

The organic oily or oil-type solvents of low volatility and with an evaporation number of above 35 and a flash point of above 30°C, preferably above 45°C, can be replaced in part by organochemical solvents of high or medium volatility, with the proviso that the solvent mixture also has an
10 evaporation number of above 35 and a flash point of above 30°C, preferably above 45°C, and that the mixture is soluble or emulsifiable in this solvent mixture.

In a preferred embodiment, some of the organochemical solvent or solvent mixture or an aliphatic polar organochemical solvent or solvent mixture is replaced. Aliphatic organochemical solvents
15 which contain hydroxyl and/or ester and/or ether groups are preferably used, such as, for example, glycol ethers, esters or the like.

Organochemical binders used for the purposes of the present invention are the synthetic resins and/or binding drying oils which are known per se and which can be diluted in water and/or
20 dissolved or dispersed or emulsified in the organochemical solvents employed, in particular binders composed of, or comprising, an acrylate resin, a vinyl resin, for example polyvinyl acetate, polyester resin, polycondensation or polyaddition resin, polyurethane resin, alkyd resin or modified alkyd resin, phenol resin, hydrocarbon resin such as indene/coumarone resin, silicone resin, drying vegetable and/or drying oils and/or physically drying binders based on a natural
25 and/or synthetic resin.

The synthetic resin employed as binder can be employed in the form of an emulsion, dispersion or solution. Bitumen or bituminous substances may also be used as binders, in amounts of up to 10% by weight. In addition, colorants, pigments, water repellants, odor-masking agents, and inhibitors
30 or anticorrosive agents and the like, all of which are known per se, can be employed.

In accordance with the invention, the composition or the concentrate preferably comprises, as organochemical binders, at least one alkyd resin or modified alkyd resin and/or a drying vegetable oil. Alkyd resins which are preferably used in accordance with the invention are those with an oil
35 content of over 45% by weight, preferably 50 to 68% by weight.

Some or all of the abovementioned binder can be replaced by a fixative (mixture) or plasticizer (mixture). These additives are intended to prevent volatilization of the active compounds, and also crystallization or precipitation. They preferably replace 0.01 to 30% of the binder (based on 100% of binder employed).

5

The plasticizers are from the chemical classes of the phthalic esters, such as dibutyl phthalate, dioctyl phthalate or benzyl butyl phthalate, phosphoric esters such as tributyl phosphate, adipic esters such as di(2-ethylhexyl) adipate, stearates such as butyl stearate or amyl stearate, oleates such as butyl oleate, glycerol ethers or higher-molecular-weight glycol ethers, glycerol esters and

10

p-toluenesulfonic esters.

Fixatives are based chemically on polyvinyl alkyl ethers such as, for example, polyvinyl methyl ether, or ketones such as benzophenone and ethylenebenzophenone.

15 Other suitable solvents or diluents are, in particular, water, if appropriate as a mixture with one or more of the abovementioned organochemical solvents or diluents, emulsifiers and dispersants.

Particularly effective timber protection is achieved by industrial-scale impregnating processes, for example the vacuum, double-vacuum or pressure processes.

20

The active compound combinations according to the invention can equally be employed for protecting objects which come into contact with saltwater or brackish water, in particular hulls, screens, nets, buildings, quaysides and signaling systems, against fouling.

25 Fouling by sessile Oligochaeta, such as Serpulidae, and by shells and species from the Ledamorpha group (goose barnacles), such as various Lepas and Scalpellum species, or by species from the Balanomorpha group (acorn barnacles), such as Balanus or Pollicipes species, increases the frictional drag of ships and, as a consequence, leads to a marked increase in operation costs owing to higher energy consumption and additionally frequent stops in the dry dock.

30

Apart from fouling by algae, for example Ectocarpus sp. and Ceramium sp., in particular fouling by sessile Entomostraka groups, which come under the generic term Cirripedia (cirriped crustaceans), is of particular importance.

35 Surprisingly, it has now been found that the active compound combinations according to the invention have an outstanding antifouling action.

Use of the active compound combinations according to the invention allows the use of heavy metals such as, for example, in bis(trialkyltin) sulfides, tri-*n*-butyltin laurate, tri-*n*-butyltin chloride, copper(I) oxide, triethyltin chloride, tri-*n*-butyl(2-phenyl-4-chlorophenoxy)tin, tributyltin
5 oxide, molybdenum disulfide, antimony oxide, polymeric butyl titanate, phenyl (bispyridine)bismuth chloride, tri-*n*-butyltin fluoride, manganese ethylenebisthiocarbamate, zinc dimethyldithiocarbamate, zinc ethylenebisthiocarbamate, zinc salts and copper salts of 2-pyridinethiol 1-oxide, bisdimethyldithiocarbamoylzinc ethylenebisthiocarbamate, zinc oxide, copper(I) ethylenebisdithiocarbamate, copper thiocyanate, copper naphthenate and tributyltin
10 halides to be dispensed with, or the concentration of these compounds to be substantially reduced.

If appropriate, the ready-to-use antifouling paints can additionally comprise other active compounds, preferably algicides, fungicides, herbicides, molluscicides, or other antifouling active compounds.

15

Preferable suitable components in combinations for the antifouling compositions according to the invention are:

algicides such as 2-*tert*-butylamino-4-cyclopropylamino-6-methylthio-1,3,5-triazine, dichlorophen, diuron, endothal, fentin acetate, isoproturon, methabenzthiazuron, oxyfluorfen, quinoclamine and
20 terbutryn;

fungicides such as benzo[*b*]thiophenecarboxylic acid cyclohexylamide S,S-dioxide, dichlofluanid, fluorfolpet, 3-iodo-2-propynyl butylcarbamate, tolylfluanid and azoles such as azaconazole, cyproconazole, epoxyconazole, hexaconazole, metconazole, propiconazole and tebuconazole;

molluscicides such as fentin acetate, metaldehyde, methiocarb, niclosamid, thiodicarb and
25 trimethacarb;

or conventional antifouling active compounds such as 4,5-dichloro-2-octyl-4-isothiazolin-3-one, diiodomethylparatryl sulfone, 2-(*N,N*-dimethylthiocarbamoylthio)-5-nitrothiazyl, potassium salts, copper salts, sodium salts and zinc salts of 2-pyridinethiol 1-oxide, pyridine-triphenylborane, tetrabutyl-distannoxane, 2,3,5,6-tetrachloro-4-(methylsulfonyl)pyridine, 2,4,5,6-tetrachloro-
30 isophthalonitrile, tetramethylthiuram disulfide and 2,4,6-trichlorophenylmaleimide.

The antifouling compositions used comprise the active compound combinations according to the invention in a concentration of 0.001 to 50% by weight, in particular 0.01 to 20% by weight.

35 Moreover, the antifouling compositions according to the invention comprise the customary components such as, for example, those described in Ungerer, *Chem. Ind.* **1985**, 37, 730-732 and

Williams, Antifouling Marine Coatings, Noyes, Park Ridge, 1973.

Besides the algicidal, fungicidal, molluscicidal active compounds and insecticidal active compounds according to the invention, antifouling paints comprise, in particular, binders.

5

Examples of recognized binders are polyvinyl chloride in a solvent system, chlorinated rubber in a solvent system, acrylic resins in a solvent system, in particular in an aqueous system, vinyl chloride/vinyl acetate copolymer systems in the form of aqueous dispersions or in the form of organic solvent systems, butadiene/styrene/acrylonitrile rubbers, drying oils such as linseed oil, resin esters or modified hardened resins in combination with tar or bitumen, asphalt and epoxy compounds, small amounts of chlorine rubber, chlorinated polypropylene and vinyl resins.

10

If appropriate, paints also comprise inorganic pigments, organic pigments or colorants which are preferably insoluble in seawater. Paints may furthermore comprise materials such as colophonium to allow controlled release of the active compounds. Furthermore, the paints may comprise plasticizers, modifiers which affect the rheological properties and other conventional constituents. The compounds according to the invention or the abovementioned mixtures may also be incorporated into self-polishing antifouling systems.

15

The active compound combinations are also suitable for controlling animal pests, in particular insects, arachnids and mites, which are found in enclosed spaces such as, for example, dwellings, factory halls, offices, vehicle cabins and the like. They can be employed in domestic insecticide products for controlling these pests. They are active against sensitive and resistant species and against all developmental stages. These pests include:

20

25 From the order of the Scorpionidea, for example, *Buthus occitanus*.

From the order of the Acarina, for example, *Argas persicus*, *Argas reflexus*, *Bryobia* ssp., *Dermanyssus gallinae*, *Glyciphagus domesticus*, *Ornithodoros moubat*, *Rhipicephalus sanguineus*, *Trombicula alfreddugesi*, *Neutrombicula autumnalis*, *Dermatophagoides pteronissimus*, *Dermatophagoides forinae*.

30

From the order of the Araneae, for example, Aviculariidae, Araneidae.

From the order of the Opiliones, for example, *Pseudoscorpiones chelifer*, *Pseudoscorpiones cheiridium*, *Opiliones phalangium*.

From the order of the Isopoda, for example, *Oniscus asellus*, *Porcellio scaber*.

From the order of the Diplopoda, for example, *Bianiulus guttulatus*, *Polydesmus* spp.

35

From the order of the Chilopoda, for example, *Geophilus* spp.

From the order of the Zygentoma, for example, *Ctenolepisma* spp., *Lepisma saccharina*,

Lepismodes inquilinus.

From the order of the Blattaria, for example, *Blatta orientalis*, *Blattella germanica*, *Blattella asahinai*, *Leucophaea maderae*, *Panchlora* spp., *Parcoblatta* spp., *Periplaneta australasiae*, *Periplaneta americana*, *Periplaneta brunnea*, *Periplaneta fuliginosa*, *Supella longipalpa*.

5 From the order of the Saltatoria, for example, *Acheta domesticus*.

From the order of the Dermaptera, for example, *Forficula auricularia*.

From the order of the Isoptera, for example, *Kaloterms* spp., *Reticulitermes* spp.

From the order of the Psocoptera, for example, *Lepinatus* spp., *Liposcelis* spp.

From the order of the Coleoptera, for example, *Anthrenus* spp., *Attagenus* spp., *Dermestes* spp.,

10 *Latheticus oryzae*, *Necrobia* spp., *Ptinus* spp., *Rhizopertha dominica*, *Sitophilus granarius*, *Sitophilus oryzae*, *Sitophilus zeamais*, *Stegobium paniceum*.

From the order of the Diptera, for example, *Aedes aegypti*, *Aedes albopictus*, *Aedes taeniorhynchus*, *Anopheles* spp., *Calliphora erythrocephala*, *Chrysozona pluvialis*, *Culex quinquefasciatus*, *Culex pipiens*, *Culex tarsalis*, *Drosophila* spp., *Fannia canicularis*, *Musca domestica*, *Phlebotomus* spp., *Sarcophaga carnaria*, *Simulium* spp., *Stomoxys calcitrans*, *Tipula paludosa*.

15

From the order of the Lepidoptera, for example, *Achroia grisella*, *Galleria mellonella*, *Plodia interpunctella*, *Tinea cloacella*, *Tinea pellionella*, *Tineola bisselliella*.

From the order of the Siphonaptera, for example, *Ctenocephalides canis*, *Ctenocephalides felis*,

20 *Pulex irritans*, *Tunga penetrans*, *Xenopsylla cheopis*.

From the order of the Hymenoptera, for example, *Camponotus herculeanus*, *Lasius fuliginosus*, *Lasius niger*, *Lasius umbratus*, *Monomorium pharaonis*, *Paravespula* spp., *Tetramorium caespitum*.

From the order of the Anoplura, for example, *Pediculus humanus capitis*, *Pediculus humanus corporis*, *Phthirus pubis*.

25 From the order of the Heteroptera, for example, *Cimex hemipterus*, *Cimex lectularius*, *Rhodnius prolixus*, *Triatoma infestans*.

They are used as aerosols, pressureless spray products, for example pump and atomizer sprays, automatic fogging systems, foggers, foams, gels, evaporator products with evaporator tablets made
30 of cellulose or polymer, liquid evaporators, gel and membrane evaporators, propeller-driven evaporators, energy-free, or passive, evaporation systems, moth papers, moth bags and moth gels, as granules or dusts, in baits for spreading or in bait stations.

According to the invention, it is possible to treat all plants and parts of plants. Plants are to be
35 understood here as meaning all plants and plant populations such as desired and undesired wild plants or crop plants (including naturally occurring crop plants). Crop plants can be plants which

can be obtained by conventional breeding and optimization methods or by biotechnological and genetic engineering methods or combinations of these methods, including the transgenic plants and including the plant cultivars which can or cannot be protected by plant breeders' certificates. Parts of plants are to be understood as meaning all above-ground and below-ground parts and organs of plants, such as shoot, leaf, flower and root, examples which may be mentioned being leaves, needles, stems, trunks, flowers, fruit-bodies, fruits and seeds and also roots, tubers and rhizomes. Parts of plants also include harvested plants and vegetative and generative propagation material, for example seedlings, tubers, rhizomes, cuttings and seeds.

10 The treatment according to the invention of the plants and parts of plants with the active compounds is carried out directly or by action on their environment, habitat or storage area according to customary treatment methods, for example by dipping, spraying, evaporating, atomizing, broadcasting, brushing-on and, in the case of propagation material, in particular in the case of seeds, furthermore by one- or multi-layer coating.

15 As already mentioned above, it is possible to treat all plants and their parts according to the invention. In a preferred embodiment, wild plant species and plant cultivars, or those obtained by conventional biological breeding methods, such as crossing or protoplast fusion, and parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant cultivars obtained by genetic engineering methods, if appropriate in combination with conventional methods (Genetic Modified Organisms), and parts thereof are treated. The terms "parts", "parts of plants" and "plant parts" have been explained above.

20 Particularly preferably, plants of the plant cultivars which are in each case commercially available or in use are treated according to the invention.

25 Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the substances and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products are possible which exceed the effects which were actually to be expected.

The transgenic plants or plant cultivars (i.e. those obtained by genetic engineering) which are preferred and to be treated according to the invention include all plants which, in the genetic modification, received genetic material which imparts particularly advantageous useful traits to these plants. Examples of such traits are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products. Further and particularly emphasized examples of such properties are a better defense of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or viruses, and also increased tolerance of the plants to certain herbicidally active compounds. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), corn, soya beans, potatoes, cotton, tobacco, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and grapes), and particular emphasis is given to corn, soya beans, potatoes, cotton, tobacco and oilseed rape. Traits that are particularly emphasized are the increased defense of the plants against insects, arachnids, nematodes and slugs and snails by toxins formed in the plants, in particular those formed by the genetic material from *Bacillus Thuringiensis* (for example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c Cry2Ab, Cry3Bb and CryIF and also combinations thereof) (hereinbelow referred to as "Bt plants"). Traits that are also particularly emphasized are the increased defense of plants to fungi, bacteria and viruses by Systemic Acquired Resistance (SAR), systemin, phytoalexins, elicitors, as well as resistance genes and correspondingly expressed proteins and toxins. Traits that are furthermore particularly emphasized are the increased tolerance of the plants to certain herbicidally active compounds, for example imidazolinones, sulfonylureas, glyphosate or phosphinotricin (for example the "PAT" gene). The genes in question which impart the desired traits can also be present in combination with one another in the transgenic plants. Examples of "Bt plants" which may be mentioned are corn varieties, cotton varieties, soya bean varieties and potato varieties which are sold under the trade names YIELD GARD® (for example corn, cotton, soya beans), KnockOut® (for example corn), StarLink® (for example corn), Bollgard® (cotton), NucoIn® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are corn varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosate, for example corn, cotton, soya bean), Liberty Link® (tolerance to phosphinotricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulfonylureas, for example corn). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned include the varieties sold under the name Clearfield® (for

example corn). Of course, these statements also apply to plant cultivars having these or still-to-be-developed genetic traits, which plants will be developed and/or marketed in the future.

5 The plants listed can be treated according to the invention in a particularly advantageous manner with the active compound mixtures according to the invention. The preferred ranges stated above for the mixtures also apply to the treatment of these plants. Particular emphasis is given to the treatment of plants with the mixtures specifically mentioned in the present text.

10 The good insecticidal and acaricidal action of the active compound combinations according to the invention can be seen from the examples which follow. While the individual active compounds show weaknesses in their action, the combinations show an action which exceeds a simple sum of actions.

15 A synergistic effect in insecticides and acaricides is always present when the action of the active compound combinations exceeds the total of the actions of the active compounds when applied individually.

The expected action for a given combination of two active compounds can be calculated as follows, according to S.R. Colby, Weeds 15 (1967), 20-22:

20

If

X is the kill rate, expressed as a percentage of the untreated control, when employing active compound A at an application rate of \underline{m} g/ha or in a concentration of \underline{m} ppm,

25 Y is the kill rate, expressed as a percentage of the untreated control, when employing active compound B at an application rate of \underline{n} g/ha or in a concentration of \underline{n} ppm and

E is the kill rate, expressed as a percentage of the untreated control, when employing active compounds A and B at application rates of \underline{m} and \underline{n} g/ha or in a concentration of \underline{m} and \underline{n} ppm,

30 then

$$E = X + Y - \frac{X \cdot Y}{100}$$

If the actual insecticidal kill rate exceeds the calculated value, the action of the combination is superadditive, i.e. a synergistic effect is present. In this case, the actually observed kill rate must
35 exceed the value calculated using the above formula for the expected kill rate (E).

After the desired period of time, the kill in % is determined. 100% means that all animals have been killed; 0% means that none of the animals have been killed.

Use examplesExample A**Aphis gossypii test**

5 Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.



10 Cotton leaves (*Gossypium hirsutum*) which are heavily infested by the cotton aphid (*Aphis gossypii*) are treated by being dipped into the preparation of active compound of the desired concentration.

After the desired period of time, the kill in % is determined. 100% means that all aphids have been killed; 0% means that none of the aphids have been killed. The kill rates determined are calculated

15 using Colby's formula (see page 44).

In this test, the following active compound combination in accordance with the present application showed a synergistically enhanced activity compared to the active compounds applied on their own:

Table A
Plant-damaging insects
Aphis gossypii test

Active compounds	Concentration of active compound in ppm	Kill rate in % after 1 ^d	
		found*	calc.**
 (II-1-9)	4	10	
 (2-5) diafenthion	20	15	
(II-1-9) + (2-5) diafenthion (1:5)	4 + 20	55	23.5

* found = activity found

** calc. = activity calculated using Colby's formula

Example B**Heliothis armigera test**

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycol ether


- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of active compound of the desired concentration and are populated with caterpillars of the cotton boll worm
10 (*Heliothis armigera*) while the leaves are still moist.

After the desired period of time, the kill in % is determined. 100% means that all caterpillars have been killed; 0% means that none of the caterpillars have been killed. The kill rates determined are calculated using Colby's formula (see page 44).

In this test, the following active compound combination in accordance with the present application
15 showed a synergistically enhanced activity compared to the active compounds applied on their own:



Table B 1
Plant-damaging insects
Heliothis armigera test

Active compounds	Concentration of active compound in ppm	Kill rate in % after 6 ^d	
		found*	calc.**
 <p>(II-1-9)</p>	0.0064	30	
(2-2) abamectin	0.16	50	
(II-1-9) + (2-2) abamectin (1:25)	0.0064 + 0.16	90	65

* found = activity found

** calc. = activity calculated using Colby's formula

Table B 2
Plant-damaging insects
Heliothis armigera test

Active compounds	Concentration of active compound in ppm	Kill rate in % after 3 ^d	
		found*	calc.**
 <p>(II-1-9)</p>	0.0064	10	
 <p>(2-17) spinosad</p>	0.032	0	
(II-1-9) + (2-17) spinosad (1:5)	0.0064 + 0.032	35	10

* found = activity found

** calc. = activity calculated using Colby's formula

Example C**Myzus persicae test**

Solvent: 7 parts by weight of dimethylformamide

5 Emulsifier: 2 parts by weight of alkylaryl polyglycol ether



To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

10 Cabbage leaves (*Brassica oleracea*) which are heavily infested by the green peach aphid (*Myzus persicae*) are treated by being dipped into the preparation of active compound of the desired concentration.

After the desired period of time, the kill in % is determined. 100% means that all aphids have been killed; 0% means that none of the aphids have been killed. The kill rates determined are calculated using Colby's formula (see page 44).

15 In this test, for example, the following active compound combination in accordance with the present application shows a synergistically enhanced activity compared to the active compounds applied on their own:

Table C
Plant-damaging insects
Myzus persicae test

Active compounds	Concentration of active compound in ppm	Kill rate in % after 6 ^d	
		found*	calc.**
 <p>(II-1-9)</p>	4	25	
 <p>(2-20) endosulfan</p>	20	15	
(II-1-9) + (2-20) endosulfan (1:5)	4 + 20	70	36.25

* found = activity found

** calc. = activity calculated using Colby's formula

Example D**Phaedon cochleariae larvae test**

Solvent: 7 parts by weight of dimethylformamide

5 Emulsifier: 2 parts by weight of alkylaryl polyglycol ether



To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

10 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of active compound of the desired concentration and are populated with larvae of the mustard beetle (*Phaedon cochleariae*) while the leaves are still moist.

After the desired period of time, the kill in % is determined. 100% means that all beetle larvae have been killed; 0% means that none of the beetle larvae have been killed. The kill rates determined are calculated using Colby's formula (see page 44).

15 In this test, the following active compound combination in accordance with the present application showed a synergistically enhanced activity compared to the active compounds applied on their own:



Table D 1
Plant-damaging insects
Phaedon cochleariae larvae test

Active compounds	Concentration of active compound in ppm	Kill rate in % after 4 ^d	
		found*	calc.**
 <p>(II-1-9)</p>	0.16	0	
 <p>(I-2)</p>	100	0	
(II-1-9) + (I-2) (1:625)	0.16 + 100	30	0

* found = activity found

** calc. = activity calculated using Colby's formula

Table D 2
Plant-damaging insects
Phaedon cochleariae larvae test

Active compounds	Concentration of active compound in ppm	Kill rate in % after 4 ^d	
		found*	calc.**
 <p>(II-1-9)</p>	0.16	5	
 <p>(I-1)</p>	100	45	
(II-1-9) + (I-1) (1:625)	0.16 + 100	85	47.75

* found = activity found

** calc. = activity calculated using Colby's formula

Example E

E/P 2006/03762

Tetranychus urticae test (OP-resistant; dip application)

Solvent: 7 parts by weight of dimethylformamide

5 Emulsifier: 2 parts by weight of alkylaryl polyglycol ether



To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

10 Bush beans (*Phaseolus vulgaris*) which are heavily infested by the greenhouse red spider mite (*Tetranychus urticae*) are dipped into a preparation of active compound of the desired concentration.

After the desired period of time, the kill in % is determined. 100% means that all spider mites have been killed; 0% means that none of the spider mites have been killed. The kill rates determined are calculated using Colby's formula (see page 44).

15 In this test, the following active compound combination in accordance with the present application showed a synergistically enhanced activity compared to the active compounds applied on their own:



Table E 1
Plant-damaging mites
Tetranychus urticae test (OP-resistant, dip application)

Active compounds	Concentration of active compound in ppm	Kill rate in % after 7 ^d	
		found*	calc.**
 (II-1-9)	100	0	
 (I-2)	0.8	0	
(II-1-9) + (I-2) (125:1)	100 + 0.8	20	0

* found = activity found

** calc. = activity calculated using Colby's formula

Table E 2
 Plant-damaging mites
 Tetranychus urticae test (OP-resistant, dip application)


Active compounds	Concentration of active compound in ppm	Kill rate in % after 7 ^d	
		found*	calc.**
 <p>(II-1-9)</p>	100	0	
 <p>(I-1)</p>	0.8	65	
(II-1-9) + (I-1) (1:625)	0.16 + 100	95	65

* found = activity found

** calc. = activity calculated using Colby's formula

Patent claims

1. A composition comprising a synergistically effective active compound combination of compounds of the formula (I) (group 1)



5

in which

- X represents C₁-C₆-alkyl, bromine, C₁-C₆-alkoxy or C₁-C₃-haloalkyl,
 Y represents hydrogen, C₁-C₆-alkyl, halogen, C₁-C₆-alkoxy, C₁-C₃-haloalkyl,
 Z represents C₁-C₆-alkyl, halogen, C₁-C₆-alkoxy,
 m represents a number 0-3,

10

- A³ represents hydrogen or in each case optionally halogen-substituted straight-chain or branched C₁-C₁₂-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, C₁-C₁₀-alkoxy-C₁-C₈-alkyl, C₁-C₈-polyalkoxy-C₂-C₈-alkyl, C₁-C₁₀-alkylthio-C₂-C₈-alkyl, cycloalkyl having 3-8 ring atoms which may be interrupted by oxygen and/or sulfur or in each case optionally halogen-, C₁-C₆-alkyl-, C₁-C₆-haloalkyl-, C₁-C₆-alkoxy-, C₁-C₆-haloalkoxy, nitro-substituted phenyl or phenyl-C₁-C₆-alkyl,

15

- A⁴ represents hydrogen, C₁-C₆-alkyl or C₁-C₆-alkoxy-C₁-C₄-alkyl


or in which

- A³ and A⁴ together with the carbon atom to which they are attached form a saturated or unsaturated 3- to 8-membered ring which is optionally interrupted by oxygen and/or sulfur and optionally substituted by halogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio or optionally substituted phenyl or is optionally benzo-fused,

20

- G¹ represents hydrogen (a) or represents the groups

25



in which

- R²⁰ represents in each case optionally halogen-substituted C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₈-alkoxy-C₁-C₈-alkyl, C₁-C₈-alkylthio-C₁-C₈-alkyl,


- C₁-C₈-polyalkoxy-C₂-C₈-alkyl or cycloalkyl having 3-8 ring atoms which may be interrupted by oxygen and/or sulfur atoms,
represents optionally halogen-, nitro-, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, C₁-C₆-haloalkyl-, C₁-C₆-haloalkoxy-substituted phenyl;
- 5 represents optionally halogen-, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, C₁-C₆-haloalkyl-, C₁-C₆-haloalkoxy-substituted phenyl-C₁-C₆-alkyl,
represents in each case optionally halogen- and/or C₁-C₆-alkyl-substituted pyridyl, pyrimidyl, thiazolyl or pyrazolyl,
represents optionally halogen- and/or C₁-C₆-alkyl-substituted phenoxy-
- 10 C₁-C₆-alkyl,
R²¹ represents in each case optionally halogen-substituted C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₈-alkoxy-C₂-C₈-alkyl or C₁-C₈-polyalkoxy-C₂-C₈-alkyl,
represents in each case optionally halogen-, nitro-, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkyl-substituted phenyl or benzyl,
- 15 R²² represents optionally halogen-substituted C₁-C₈-alkyl, represents in each case optionally C₁-C₄-alkyl-, halogen-, C₁-C₄-haloalkyl-, C₁-C₄-alkoxy-, C₁-C₄-haloalkoxy-, nitro- or cyano-substituted phenyl or benzyl,
R²³ and R²⁴ independently of one another represent in each case optionally
- 20 halogen-substituted C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylamino, di(C₁-C₈)alkylamino, C₁-C₈-alkylthio, C₂-C₅-alkenylthio, C₂-C₅-alkynylthio, C₃-C₇-cycloalkylthio, represent in each case optionally halogen-, nitro-, cyano-, C₁-C₄-alkoxy-, C₁-C₄-haloalkoxy-, C₁-C₄-alkylthio-, C₁-C₄-haloalkylthio-, C₁-C₄-alkyl-, C₁-C₄-haloalkyl-substituted
- 25 phenyl, phenoxy or phenylthio,
R²⁵ and R²⁶ independently of one another represent in each case optionally halogen-substituted C₁-C₁₀-alkyl, C₁-C₁₀-alkoxy, C₃-C₈-alkenyl, C₁-C₈-alkoxy-C₁-C₈-alkyl, represent optionally halogen-, C₁-C₆-haloalkyl-, C₁-C₆-alkyl- or C₁-C₆-alkoxy-substituted phenyl, represent optionally
- 30 halogen-, C₁-C₆-alkyl-, C₁-C₆-haloalkyl- or C₁-C₆-alkoxy-substituted benzyl or together represent a 5- to 6-membered ring which is optionally interrupted by oxygen or sulfur and which may optionally be substituted by C₁-C₆-alkyl,

- 56 -

or an acaricidally active compound (group 2).

2. The composition according to claim 1 in which the acaricidally active compound (group 2) is selected from the group consisting of

(2-1) the phenylhyrazine derivative of the formula




and/or

(2-2) the macrolide with the common name abamectin


and/or

(2-3) the naphthalenedione derivative of the formula




and/or

(2-4) the pyrrole derivative of the formula




and/or

(2-5) the thiourea derivative of the formula




and/or

(2-6) the oxazoline derivative of the formula



and/or

(2-7) an organotin derivative of the formula



in which


R represents (2-7-a = azocyclotin),

or

5 R represents -OH (2-7-b = cyhexatin),


and/or

(2-8) the pyrazole derivative of the formula




and/or

10 (2-9) the pyrazole derivative of the formula



and/or


(2-10) the pyridazinone derivative of the formula



15


and/or

(2-11) the benzoylurea derivative of the formula




and/or

(2-12) the pyrethroid of the formula



and/or


(2-13) the tetrazine derivative of the formula



5


and/or

(2-14) the organotin derivative of the formula



and/or


(2-15) the sulfenamide of the formula



10

und/or

(2-16) the pyrimidyl phenol ethers of the formula



in which

15

R represents fluorine (2-16-a = 4-[(4-chloro- α,α,α -trifluoro-3-tolyl)oxy]-6-[(α,α,α -4-tetrafluoro-3-tolyl)oxy]pyrimidine)


R represents nitro (2-16-b = 4-[(4-chloro- α,α,α -trifluoro-3-tolyl)oxy]-6-[(α,α,α -trifluoro-4-nitro-3-tolyl)oxy]pyrimidine)

R represents bromine (2-16- = 4-[(4-chloro- α,α,α -trifluoro-3-tolyl)oxy]-6-[(α,α,α -trifluoro-4-bromo-3-tolyl)oxy]pyrimidine)

20

and/or

(2-17) the macrolide of the formula



(spinosad)

a mixture comprising, preferably,

85% spinosyn A (R = H)

15% spinosyn B (R = CH₃)

5

and/or

(2-18) ivermectin


and/or

(2-19) milbemectin

10


and/or

(2-20) endosulfan



and/or


(2-21) fenazaquin



15

and/or


(2-22) pyrimidifen



and/or


20

(2-23) triarathen



and/or


(2-24) tetradifon



5


and/or

(2-25) propargite



and/or


(2-26) hexythiazox



10

and/or


(2-27) bromopropylate



and/or


15

(2-28) dicofol



and/or

(2-29) chinomethionat




and at least one active compound from the group of the anthranilamides of the formula (II).

3. The composition as claimed in claim 1 or claim 2, comprising at least one compound of the formula (I) in which

- X represents C₁-C₄-alkyl, bromine, C₁-C₄-alkoxy or C₁-C₃-haloalkyl,
 Y represents hydrogen, C₁-C₄-alkyl, fluorine, chlorine, bromine, C₁-C₄-alkoxy, C₁-C₃-haloalkyl,
 Z represents C₁-C₄-alkyl, chlorine, bromine, C₁-C₄-alkoxy,
 m represents a number 0-2,
 A³ represents hydrogen or in each case optionally mono- to trifluoro-substituted straight-chain or branched C₁-C₆-alkyl, C₂-C₆-alkenyl, C₁-C₄-alkoxy-C₁-C₂-alkyl, cycloalkyl having 3-8 ring atoms which may optionally be interrupted by oxygen and/or sulfur or represents benzyl or phenyl which is optionally mono- to disubstituted by fluorine, chlorine, bromine, C₁-C₂-alkyl, C₁-C₂-haloalkyl, C₁-C₂-alkoxy, C₁-C₂-haloalkoxy, nitro,
 A⁴ represents hydrogen, C₁-C₂-alkyl or C₁-C₂-alkoxy-C₁-C₂-alkyl
 or in which

A³ and A⁴ together with the carbon atom to which they are attached form a saturated or unsaturated 3- to 7-membered ring which is optionally interrupted by oxygen and/or sulfur and optionally mono- to disubstituted by fluorine, chlorine, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₂-haloalkyl, C₁-C₂-haloalkoxy or C₁-C₂-alkylthio,

G¹ represents hydrogen (a) or represents groups



in which

R²⁰ represents in each case optionally mono- to pentafluoro- or -chloro-substituted C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-alkylthio-C₁-C₄-alkyl or cycloalkyl having 3-6 ring atoms which may be interrupted by oxygen and/or sulfur atoms,
 represents phenyl which is optionally mono- to disubstituted by fluorine,

chlorine, bromine, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy,
 represents benzyl which is optionally mono to disubstituted by fluorine, chlorine, bromine, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy,
 represents pyridyl, pyrimidyl, thiazolyl or pyrazolyl, each of which is optionally mono- to disubstituted by chlorine, bromine and/or C₁-C₄-alkyl,
 R²¹ represents C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₆-alkoxy-C₂-C₆-alkyl, C₁-C₆-polyalkoxy-C₂-C₆-alkyl, each of which is optionally mono- to pentasubstituted by fluorine or chlorine,
 represents phenyl or benzyl, each of which is optionally mono- to disubstituted by fluorine, chlorine, bromine, nitro, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-haloalkyl,
 R²² represents C₁-C₄-alkyl which is optionally mono- to pentasubstituted by fluorine or chlorine, represents phenyl or benzyl, each of which is optionally mono- to disubstituted by C₁-C₄-alkyl, fluorine, chlorine, bromine, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, nitro or cyano,
 R²³ and R²⁴ independently of one another represent C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylamino, di(C₁-C₄)alkylamino, C₁-C₄-alkylthio, C₂-C₄-alkenylthio, C₃-C₆-cycloalkylthio, each of which is optionally mono- to trisubstituted by fluorine or chlorine, represent phenyl, phenoxy or phenylthio, each of which is optionally mono- to disubstituted by fluorine, chlorine, bromine, nitro, cyano, C₁-C₂-alkoxy, C₁-C₂-haloalkoxy, C₁-C₂-alkylthio, C₁-C₂-haloalkylthio, C₁-C₂-alkyl, C₁-C₂-haloalkyl,
 R²⁵ and R²⁶ independently of one another represent C₁-C₆-alkyl, C₁-C₆-alkoxy, C₃-C₆-alkenyl, C₁-C₄-alkoxy-C₁-C₂-alkyl, each of which is optionally mono- to trisubstituted by fluorine or chlorine, represent benzyl which is optionally mono- to disubstituted by fluorine, chlorine, bromine, C₁-C₂-haloalkyl, C₁-C₄-alkyl or C₁-C₄-alkoxy or together represent a 5- to 6-membered ring which is optionally interrupted by oxygen or sulfur and which may optionally be substituted by C₁-C₂-alkyl,

and at least one anthranilamide of the formula (II).

4. The composition as claimed in claims 1, 2 or 3 comprising at least one compound of the formula (I) in which

- X represents C₁-C₄-alkyl, C₁-C₄-alkoxy or trifluoromethyl,
 Y represents hydrogen, C₁-C₄-alkyl, chlorine, bromine, C₁-C₄-alkoxy, C₁-C₂-haloalkyl,
 Z represents C₁-C₄-alkyl, chlorine, bromine, C₁-C₄-alkoxy,
 m represents 0 or 1,
 A³ and A⁴ together with the carbon atom to which they are attached represent a saturated 5- to 6-membered ring which is optionally monosubstituted by C₁-C₄-alkyl or C₁-C₄-alkoxy,

G¹ represents hydrogen (a) or represents the groups



, in which

(b) (c)

R²⁰ represents in each case optionally mono- to trifluoro- or -chloro-substituted C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₁-C₄-alkoxy-C₁-C₂-alkyl, or cycloalkyl having 3-6 ring atoms which may be interrupted by 1 to 2 oxygen atoms,

represents phenyl which is optionally monosubstituted by fluorine, chlorine, bromine, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, trifluoromethyl or trifluoromethoxy;

R²¹ represents C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, C₁-C₄-alkoxy-C₂-C₄-alkyl, represents phenyl or benzyl, each of which is optionally monosubstituted by fluorine, chlorine, bromine, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy or trifluoromethyl,


and at least one anthranilamide of the formula (II).

5. The composition as claimed in claims 1, 2, 3 or 4, comprising at least one compound of the formula (I) in which

- X represents methyl, ethyl, methoxy, ethoxy or trifluoromethyl,
 Y represents hydrogen, methyl, ethyl, chlorine, bromine, methoxy or trifluoromethyl,
 Z represents methyl, ethyl, chlorine, bromine or methoxy,
 m represents 0 or 1,

A³ and A⁴ together with the carbon atom to which they are attached form a saturated 5- to 6-membered ring which is optionally monosubstituted by methyl, ethyl, propyl, methoxy, ethoxy, propoxy, butoxy or isobutoxy,

G¹ represents hydrogen (a) or represents the groups




R^{20} represents in each case mono- to trifluoro- or -chloro-substituted C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl, C_1 - C_3 -alkoxy- C_1 - C_2 -alkyl, or cycloalkyl having 3-6 ring atoms which may be interrupted by 1 to 2 oxygen atoms,


represents phenyl which is optionally monosubstituted by fluorine, chlorine, bromine, methyl, methoxy, trifluoromethyl or trifluoromethoxy;

R^{21} represents C_1 - C_8 -alkyl, C_2 - C_8 -alkenyl, C_1 - C_4 -alkoxy- C_2 - C_3 -alkyl, represents phenyl or benzyl, each of which is optionally monosubstituted by fluorine, chlorine, bromine, nitro, methyl, methoxy or trifluoromethyl, and at least one anthranilamide of the formula (II).

6. The composition as claimed in claim 1, 2, 3, 4 or 5, comprising the compound of the formula (I-1)




and/or the compound of the formula (I-2)



and at least one anthranilamide of the formula (II).

7. The composition as claimed in claim 1, 2, 3, 4, 5 or 6, comprising at least one anthranilamide of the formula (II)



(II)

in which

A^1 and A^2 independently of one another represent oxygen or sulfur,

X^1 represents N or CR^{10} ,

5 R^1 represents hydrogen or represents in each case optionally mono- or polysubstituted C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl or C_3 - C_6 -cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R^6 , halogen, cyano, nitro, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylsulfinyl, C_1 - C_4 -alkylsulfonyl, C_2 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylamino, C_2 - C_8 -dialkylamino, C_3 - C_6 -cycloalkylamino, $(C_1$ - C_4 -alkyl) C_3 - C_6 -cycloalkylamino and R^{11} ,

R^2 represents hydrogen, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylamino, C_2 - C_8 -dialkylamino, C_3 - C_6 -cycloalkylamino, C_2 - C_6 -alkoxycarbonyl or C_2 - C_6 -alkylcarbonyl,

15 R^3 represents hydrogen, R^{11} or represents in each case optionally mono- or polysubstituted C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R^6 , halogen, cyano, nitro, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylsulfinyl, C_1 - C_4 -alkylsulfonyl, C_2 - C_6 -alkoxycarbonyl, C_2 - C_6 -alkylcarbonyl, C_3 - C_6 -trialkylsilyl, R^{11} , phenyl, phenoxy and a 5- or 6-membered heteroaromatic ring, where each phenyl, phenoxy and 5- or 6-membered heteroaromatic ring may optionally be substituted and where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R^{12} , or

25 R^2 and R^3 may be attached to one another and form the ring M,

R^4 represents hydrogen, C_1 - C_6 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_3 - C_6 -cycloalkyl, C_1 - C_6 -haloalkyl, C_2 - C_6 -haloalkenyl, C_2 - C_6 -haloalkynyl, C_3 - C_6 -halocycloalkyl, halogen, cyano, nitro, hydroxyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylsulfinyl, C_1 - C_4 -alkylsulfonyl, C_1 - C_4 -haloalkylthio, C_1 - C_4 -haloalkyl-

sulfinyl, C₁-C₄-haloalkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino, C₃-C₆-trialkylsilyl or represents in each case optionally mono- or polysubstituted phenyl, benzyl or phenoxy, where the substituents independently of one another may be selected from the group consisting of C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl, C₂-C₄-haloalkenyl, C₂-C₄-haloalkynyl, C₃-C₆-halocycloalkyl, halogen, cyano, nitro, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino, C₃-C₆-(alkyl)-cycloalkylamino, C₂-C₄-alkylcarbonyl, C₂-C₆-alkoxycarbonyl, C₂-C₆-alkylaminocarbonyl, C₃-C₈-dialkylaminocarbonyl and C₃-C₆-trialkylsilyl,

R⁵ and R⁸ in each case independently of one another represent hydrogen, halogen or represent in each case optionally substituted C₁-C₄-alkyl, C₁-C₄-haloalkyl, R¹², G, J, -OJ, -OG, -S(O)_p-J, -S(O)_p-G, -S(O)_p-phenyl, where the substituents independently of one another may be selected from one to three radicals W or from the group consisting of R¹², C₁-C₁₀-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₄-alkoxy and C₁-C₄-alkylthio, where each substituent may be substituted by one or more substituents independently of one another selected from the group consisting of G, J, R⁶, halogen, cyano, nitro, amino, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl, C₁-C₄-haloalkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-trialkylsilyl, phenyl and phenoxy, where each phenyl or phenoxy ring may optionally be substituted and where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R¹²,

G in each case independently of one another represents a 5- or 6-membered nonaromatic carbocyclic or heterocyclic ring which optionally contains one or two ring members from the group consisting of C(=O), SO and S(=O)₂ and which may optionally be substituted by one to four substituents independently of one another selected from the group consisting of C₁-C₂-alkyl, halogen, cyano, nitro and C₁-C₂-alkoxy, or independently of one another represents C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₃-C₇-cycloalkyl, (cyano)C₃-C₇-cycloalkyl, (C₁-C₄-alkyl)C₃-C₆-cycloalkyl, (C₃-C₆-cycloalkyl)C₁-C₄-alkyl, where each cycloalkyl, (alkyl)cycloalkyl and (cycloalkyl)-alkyl may optionally be substituted by one or more halogen atoms,

J in each case independently of one another represents an optionally substituted 5- or 6-membered heteroaromatic ring, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R¹²,

- R⁶ independently of one another represent $-C(=E^1)R^{19}$, $-LC(=E^1)R^{19}$, $-C(=E^1)LR^{19}$, $-LC(=E^1)LR^{19}$, $-OP(=Q)(OR^{19})_2$, $-SO_2LR^{18}$ or $-LSO_2LR^{19}$, where each E¹ independently of the others represents O, S, N-R¹⁵, N-OR¹⁵, N-N(R¹⁵)₂, N-S=O, N-CN or N-NO₂,
- 5 R⁷ represents hydrogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, halogen, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl, C₁-C₄-haloalkylsulfonyl,
- R⁹ represents C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylsulfinyl or halogen,
- R¹⁰ represents hydrogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, halogen, cyano or C₁-C₄-haloalkoxy,
- 10 R¹¹ in each case independently of one another represents in each case optionally mono- to trisubstituted C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, C₁-C₆-haloalkylthio, C₁-C₆-haloalkylsulfinyl, phenylthio or phenylsulfinyl, where the substituents independently of one another may be selected from the list consisting of W,
- 15 $-S(O)_nN(R^{16})_2$, $-C(=O)R^{13}$, $-L(C=O)R^{14}$, $-S(C=O)LR^{14}$, $-C(=O)LR^{13}$, $-S(O)_nNR^{13}C(=O)R^{13}$, $-S(O)_nNR^{13}C(=O)LR^{14}$ and $-S(O)_nNR^{13}S(O)_2LR^{14}$,
- L in each case independently of one another represents O, NR¹⁸ or S,
- R¹² in each case independently of one another represents $-B(OR^{17})_2$, amino, SH, thiocyanato, C₃-C₈-trialkylsilyloxy, C₁-C₄-alkyl disulfide, $-SF_5$, $-C(=E^1)R^{19}$, $-LC(=E^1)R^{19}$, $-C(=E^1)LR^{19}$, $-LC(=E^1)LR^{19}$, $-OP(=Q)(OR^{19})_2$, $-SO_2LR^{19}$ or $-LSO_2LR^{19}$,
- 20 Q represents O or S,
- R¹³ in each case independently of one another represents hydrogen or represents in each case optionally mono- or polysubstituted C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl or C₃-C₆-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R⁶, halogen, cyano, nitro, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino and (C₁-C₄-alkyl)C₃-C₆-cycloalkylamino,
- 25 R¹⁴ in each case independently of one another represents in each case optionally mono- or polysubstituted C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₂-C₂₀-alkynyl or C₃-C₆-cycloalkyl, where the substituents independently of one another may be selected from the group consisting of R⁶, halogen, cyano, nitro, hydroxyl, C₁-C₄-alkoxy, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino and (C₁-C₄-alkyl)C₃-C₆-cycloalkylamino or represents optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R¹²,
- 30
- 35

- 5 R^{15} in each case independently of one another represent hydrogen or represent in each case mono- or polysubstituted C_1-C_6 -haloalkyl or C_1-C_6 -alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, nitro, hydroxyl, C_1-C_4 -alkoxy, C_1-C_4 -haloalkoxy, C_1-C_4 -alkylthio, C_1-C_4 -alkylsulfinyl, C_1-C_4 -alkylsulfonyl, C_1-C_4 -haloalkylthio, C_1-C_4 -haloalkylsulfinyl, C_1-C_4 -haloalkylsulfonyl, C_1-C_4 -alkylamino, C_2-C_8 -dialkylamino, C_2-C_6 -alkoxycarbonyl, C_2-C_6 -alkylcarbonyl, C_3-C_6 -trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W or one or more radicals R^{12} , or $N(R^{15})_2$ represents a cycle which forms the ring M,
- 10 R^{16} represents C_1-C_{12} -alkyl or C_1-C_{12} -haloalkyl, or $N(R^{16})_2$ represents a cycle which forms the ring M,
- 15 R^{17} in each case independently of one another represents hydrogen or C_1-C_4 -alkyl, or $B(OR^{17})_2$ represents a ring, where the two oxygen atoms are attached via a chain to two or three carbon atoms which are optionally substituted by one or two substituents independently of one another selected from the group consisting of methyl and C_2-C_6 -alkoxycarbonyl,
- 20 R^{18} in each case independently of one another represents hydrogen, C_1-C_6 -alkyl or C_1-C_6 -haloalkyl, or $N(R^{13})(R^{18})$ represents a cycle which forms the ring M,
- 25 R^{19} in each case independently of one another represents hydrogen or represents in each case optionally mono- or polysubstituted C_1-C_6 -alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, nitro, hydroxyl, C_1-C_4 -alkoxy, C_1-C_4 -haloalkoxy, C_1-C_4 -alkylthio, C_1-C_4 -alkylsulfinyl, C_1-C_4 -alkylsulfonyl, C_1-C_4 -haloalkylthio, C_1-C_4 -haloalkylsulfinyl, C_1-C_4 -haloalkylsulfonyl, C_1-C_4 -alkylamino, C_2-C_8 -dialkylamino, CO_2H , C_2-C_6 -alkoxycarbonyl, C_2-C_6 -alkylcarbonyl, C_3-C_6 -trialkylsilyl and optionally substituted phenyl, where the substituents independently of one another may be selected from one to three radicals W, C_1-C_6 -haloalkyl, C_3-C_6 -cycloalkyl or phenyl or pyridyl, each of which is optionally mono- to trisubstituted by W,
- 30 M in each case represents an optionally mono- to tetrasubstituted ring which, in addition to the nitrogen atom which is attached to the substituent pair R^{13} and R^{18} , $(R^{15})_2$ or $(R^{16})_2$, contains two to six carbon atoms and optionally additionally a further nitrogen, sulfur or oxygen atom, and where the substituents independently of one another may be selected from the group consisting of C_1-C_2 -alkyl, halogen, cyano, nitro and C_1-C_2 -alkoxy,
- 35 W in each case independently of one another represents C_1-C_4 -alkyl, C_2-C_4 -alkenyl,


C₂-C₃-alkynyl, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl, C₂-C₄-haloalkenyl, C₂-C₃-haloalkynyl, C₃-C₆-halocycloalkyl, halogen, cyano, nitro, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylamino, C₂-C₈-dialkylamino, C₃-C₆-cycloalkylamino, (C₁-C₄-alkyl)C₃-C₆-cycloalkylamino, C₂-C₄-alkylcarbonyl, C₂-C₆-alkoxycarbonyl, CO₂H, C₂-C₆-alkylaminocarbonyl, C₃-C₈-dialkylaminocarbonyl or C₃-C₆-trialkylsilyl,

n in each case independently of one another represents 0 or 1,

p in each case independently of one another represents 0, 1 or 2.

where in the case that (a) R⁵ represents hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-haloalkenyl, C₂-C₆-haloalkynyl, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio or halogen and (b) R⁸ represents hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₂-C₆-haloalkenyl, C₂-C₆-haloalkynyl, C₁-C₄-haloalkoxy, C₁-C₄-haloalkylthio, halogen, C₂-C₄-alkylcarbonyl, C₂-C₆-alkoxycarbonyl, C₂-C₆-alkylaminocarbonyl or C₃-C₈ dialkylaminocarbonyl, (c) at least one substituent selected from the group consisting of R⁶, R¹¹ and R¹² is present and (d), if R¹² is not present, at least one R⁶ or R¹¹ is different from C₂-C₆-alkylcarbonyl, C₂-C₆ alkoxycarbonyl, C₂-C₆-alkylaminocarbonyl and C₃-C₈-dialkylaminocarbonyl.

8. The composition as claimed in claim 1, 2, 3, 4, 5, 6 or 7, comprising an anthranilamide of the formula (II-I)



in which

R² represents hydrogen or C₁-C₆-alkyl,

R³ represents C₁-C₆-alkyl which is optionally substituted by a radical R⁶,

R⁴ represents C₁-C₄-alkyl, C₁-C₂-haloalkyl, C₁-C₂-haloalkoxy or halogen,

R⁵ represents hydrogen, C₁-C₄-alkyl, C₁-C₂-haloalkyl, C₁-C₂-haloalkoxy or halogen,

R⁶ represents -C(=E²)R¹⁹, -LC(=E²)R¹⁹, -C(=E²)LR¹⁹ or -LC(=E²)LR¹⁹, where each E² independently of the others represents O, S, N-R¹⁵, N-OR¹⁵, N-N(R¹⁵)₂, and each L independently of the others represents O or NR¹⁸,

R⁷ represents C₁-C₄-haloalkyl or halogen,

R⁹ represents C₁-C₂-haloalkyl, C₁-C₂-haloalkoxy, S(O)_pC₁-C₂-haloalkyl or halogen,

R¹⁵ in each case independently of one another represents hydrogen or represents in

each case optionally substituted C₁-C₆-haloalkyl or C₁-C₆-alkyl, where the substituents independently of one another may be selected from the group consisting of cyano, C₁-C₃-alkoxy, C₁-C₃-haloalkoxy, C₁-C₃-alkylthio, C₁-C₃-alkylsulfinyl, C₁-C₄-alkylsulfonyl, C₁-C₄-haloalkylthio, C₁-C₄-haloalkylsulfinyl or C₁-C₄-haloalkylsulfonyl,

R¹⁸ in each case represents hydrogen or C₁-C₄-alkyl,

R¹⁹ in each case independently of one another represents hydrogen or C₁-C₆-alkyl,

p independently of one another represents 0, 1, 2.

9. The composition as claimed in claim 1, 2, 3, 4, 5, 6, 7 or 8, comprising compounds of the formula (I) (group 1) or at least one acaricidally active compound (group 2) and at least one anthranilamide of the formula (II) in a ratio of from 500:1 to 1:50.
10. The use of a synergistically effective mixture as defined in claims 1, 2, 3, 4, 5, 6, 7 or 8 for controlling animal pests.
11. A process for preparing pesticides, characterized in that a synergistically effective mixture as defined in claims 1, 2, 3, 4, 5, 6, 7 or 8 is mixed with extenders and/or surfactants.
12. A composition substantially as herein described with reference to and as exemplified by any one of the compositions in the table on pages 25 to 28 of the accompanying specification.
13. The use of a composition as claimed in claim 12 for controlling insect pests.