




(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/03/28	(51) Cl.Int./Int.Cl. <i>C07K 16/00</i> (2006.01), <i>A61K 39/395</i> (2006.01), <i>A61P 35/00</i> (2006.01), <i>A61P 37/04</i> (2006.01), <i>C07K 16/28</i> (2006.01), <i>C07K 16/46</i> (2006.01), <i>C07K 14/735</i> (2006.01)
(87) Date publication PCT/PCT Publication Date: 2018/10/04	(71) Demandeur/Applicant: LYVGEN BIOPHARMA HOLDINGS LIMITED, KY
(85) Entrée phase nationale/National Entry: 2019/09/26	(72) Inventeur/Inventor: WANG, JIEYI, US
(86) N° demande PCT/PCT Application No.: US 2018/024872	(74) Agent: SMART & BIGGAR LLP
(87) N° publication PCT/PCT Publication No.: 2018/183520	
(30) Priorité/Priority: 2017/03/28 (US62/477,661)	

(54) Titre : AGENTS THERAPEUTIQUES ET METHODES D'AMELIORATION DE REPONSES IMMUNITAIRES DANS UN MICRO-ENVIRONNEMENT TUMORAL
(54) Title: THERAPEUTIC AGENTS AND METHODS FOR ENHANCING IMMUNE RESPONSES IN TUMOR MICROENVIRONMENT



(57) **Abrégé/Abstract:**
Human IgG1, IgG2, and IgG4 mutants having mutations in the hinge domain and exhibiting altered binding activity to Fcγ receptors such as FcγRIIB (CD32B). Also provided herein are methods for selectively activating immune responses in a subject using a therapeutic agent capable of targeting both an immune cell surface receptor and FcγRIIB.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
04 October 2018 (04.10.2018)



(10) International Publication Number
WO 2018/183520 A1

(51) International Patent Classification:

C07K 16/00 (2006.01) C07K 16/28 (2006.01)
C07K 16/24 (2006.01)

(21) International Application Number:

PCT/US2018/024872

(22) International Filing Date:

28 March 2018 (28.03.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/477,661 28 March 2017 (28.03.2017) US

(71) Applicant: LYVGEN BIOPHARMA HOLDINGS LIMITED [DE/—]; PO Box 31119, Grand Cayman, KY11-1205 (KY).

(72) Inventor; and

(71) Applicant: WANG, Jieyi [US/US]; 2317 Hastings Drive, Belmont, CA 94002 (US).

(74) Agent: CHEN, Yahua; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210-2206 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,

MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: THERAPEUTIC AGENTS AND METHODS FOR ENHANCING IMMUNE RESPONSES IN TUMOR MICROENVIRONMENT




FIG. 1A

(57) Abstract: Human IgG1, IgG2, and IgG4 mutants having mutations in the hinge domain and exhibiting altered binding activity to Fcγ receptors such as FcγRIIB (CD32B). Also provided herein are methods for selectively activating immune responses in a subject using a therapeutic agent capable of targeting both an immune cell surface receptor and FcγRIIB.

Therapeutic Agents and Methods for Enhancing Immune Responses in Tumor Microenvironment

RELATED APPLICATION

5 This application claims the benefit under 35 U.S.C. § 119(e) of U.S. provisional application number 62/477,661, filed on March 28, 2017, which is incorporated by reference herein in its entirety.

BACKGROUND OF THE INVENTION

10 Fc receptors (FcR) are a family of immune cell surface proteins capable of binding to the Fc portion of antibodies. There are several different types of Fc receptors, including Fc γ receptors, Fc α receptors, Fc ϵ receptors, and neonatal Fc receptors (FcRn), which have different binding activities to IgG, IgA, IgE, and IgG antibodies, respectively. The Fc γ receptor subfamily includes Fc γ RI (CD64), Fc γ RIIA (CD32a), Fc γ RIIB (CD32b), Fc γ RIIB (CD32c), Fc γ RIIIA (CD16a), and Fc γ RIIIB (CD16b). Fc γ RI has high binding affinity to
15 IgG1 and IgG3 antibodies, while the other Fc γ Rs have low binding affinity to IgG antibodies.

Different types of Fc receptors play different roles in the immune system. For example, Fc γ RIII receptors, expressed on NK cells and macrophages, bind to antibodies that are attached to infected cells or invading pathogens and trigger antibody-mediated
20 phagocytosis (ADCP) or antibody-dependent cell-mediated cytotoxicity (ADCC) of the immune cells, thereby leading to elimination of the infected cells or invading pathogens. On the other hand, Fc γ RII receptors, expressed on B cells and dendritic cells, can down regulate the activity of the immune cells when binding to IgG antibodies.

Therapies involving activated immune cells are promising approaches for eliminating
25 diseased cells such as cancer cells. However, such therapeutic approaches often raise safety concerns. For example, overly activated immune cells would lead to undesired cytotoxicity, causing tissue damage. It is therefore of great interest to develop new immune therapies that are effective and safe.

SUMMARY OF THE INVENTION

The present disclosure is based, at least in part, on the development of various Fc region variants of IgG1, IgG2, or IgG4 molecules that exhibit desired Fc receptor binding activity and/or selectivity. Such Fc variants can be used in constructing Fc-containing proteins such as antibodies capable of selectively modulating immune responses.

Accordingly, one aspect of the present disclosure provides a protein (*e.g.*, an antibody) comprising a variant Fc region, which comprises a mutation (*e.g.*, an amino acid residue substitution, an insertion, a deletion, or a combination thereof) at any of positions 218-329 (*e.g.*, 218-328) as compared to a wild-type parent Fc region. The Fc numbering system used herein is according to the EU index as known in the art.

In some embodiments, the mutation comprises one or more of the following: (a) mutation at one or more of positions 219-225; (b) an insertion between position 218 and position 219, or at position 236; (c) an amino acid residue substitution at one or more positions of 233-235; (d) an insertion at any of positions 220-223; (e) a deletion at one or more of positions 236-238; and (f) an amino acid substitution at one or more of positions 267, 273, 328, and 329 (*e.g.*, S267E, V273E, L328F, P329G, or a combination thereof). In some instances, the mutation comprises an insertion and optionally a deletion. For example, the mutation in the variant Fc region may comprise (i) a mutation at one or more of positions 234-238, and optionally (ii) a mutation at one or more of positions 267, 273, 328, and 329. In some examples, the mutation of (i) can be a deletion, for example, a deletion of one or more of positions 234-238 (*e.g.*, 236-238). Alternatively or in addition, the mutation of (ii) can be an amino acid residue substitution, for example, amino acid residue substitutions S267E, V273E, L328F, P329G, or a combination thereof.

In some embodiments, the wild-type parent Fc region is of an IgG1 molecule. A protein comprising an Fc variant derived from an IgG1 molecule (*e.g.*, a human IgG1 molecule) may comprise a mutation comprising one or more of the following: (a) an insertion between position 228 and position 229; (b) a mutation at one or more of positions 220-225; (c) an amino acid substitution at positions 234 and 235; (d) a deletion at one or more of positions 236-238; and (e) an amino acid substitution at one or more positions 267, 273, 328, and 329 (*e.g.*, S267E, V273E, L328F, and/or P329G). In one example, the mutation comprises (i) a deletion at one or more of positions 236-238 (*e.g.*, 236, 237, 238, or

a combination thereof), and optionally (ii) an amino acid residue substitution at one or more of positions 267, 273, and 328. For example, the mutation may comprise a deletion of position 236 or a deletion of positions 236-238. Optionally, the mutation may further comprise an amino acid residue substitution at one or more of positions 267, 273, and 268, which may be S267E, V273E, L328F, or a combination thereof. In one example, the mutation comprises a deletion at one or more of positions 236-238 and the amino acid substitution of S267E. Exemplary variant Fc regions derived from human IgG1 may be one of G1m1, G1m2, G1m-2, G1m-4, G1m5, G1m7, G1m8, G1m9, G1m15, G1m17, G1m18, G1m19, G1m25, G1m27, G1m28, G1m29, G1mAA, and G1mAG.

In other embodiments, the wild-type parent Fc region is of an IgG2 molecule (*e.g.*, a human IgG2 molecule). A protein comprising an Fc variant derived from an IgG2 molecule (*e.g.*, a human IgG2 molecule) may comprise a mutation comprising one or more of the following: (a) a mutation at one or more positions of 219-225; (b) an amino acid substitution at one or more of positions 233-235; (c) an insertion between position 218 and 219, or at position 236; (d) a deletion of one or more of positions 237 and 238; and (e) an amino acid residue substitution at one or more of positions 267, 273, and 328 (*e.g.*, S267E, V273E, L328F, or a combination thereof). In some examples, the mutation is at one or more of positions 233-238, 267, 273, and 328. For example, the mutation may comprise (i) a deletion at one or more of positions 236, 237, and 238 (*e.g.*, 237 and/or 238), and optionally (ii) an amino acid residue substitution at one or more of positions 267, 273, and 328 (*e.g.*, S267E, L328F, or a combination thereof). In some examples, the mutation comprises an addition at position 236. In other examples, the mutation may comprise an amino acid residue substitution at one or more of positions 233-238, *e.g.*, P233E, V234F or V234V, A235L, A235S, or a combination thereof. Alternatively or in addition, the mutation further comprises an amino acid residue substitution at one or more positions 267, 273, and 328, which may be S267E, V273E, and/or L328F. In some particularly examples, the variant Fc region derived from IgG2 can be one of G2m1, G2m-1, G2m2, G2m-4, G2m5, G2m7, G2m8, G2m9, G2m10, G2m15, G2m17, G2m18, G2m19, G2m20, G2m27, G2m27, and G2m28.

In yet other embodiments, the wild-type parent Fc is of an IgG4 molecule (*e.g.*, a human IgG4 molecule). The Fc variant derived from an IgG4 molecule (*e.g.*, a human IgG4) may comprise the S228P substitution. Alternatively or in addition, the Fc variant derived

from an IgG4 molecule such as a human IgG4 molecule may have a mutation comprising one or more of the following: (a) a mutation at one or more of positions 219-225; (b) an amino acid residue substitution at one or more positions 234 and 235; (c) a deletion at one or more of positions 236-238; and (d) an amino acid residue substitution at one or more of positions 267, 273, and 328 (S267E, V273F, L328F, or a combination thereof). In some examples, the Fc variants of IgG4 may comprise the S228P amino acid residue substitution and a deletion at one or more of positions 236-238. Such an Fc variant may further comprise an amino acid substitution at one or more of positions 267, 273, and 328 (*e.g.*, L328F). In some instances, the Fc variant may comprise the S228P substitution and a further mutation at one or more positions 234-238, 267, 273, and 328. For example, the mutation may comprise a deletion of one or more of position 235-237, or comprise an amino acid residue substitution at position 235, 236, or both (*e.g.*, F234V, K235S, or a combination thereof). Alternatively or in addition, the mutation further comprises an amino acid residue substitution at one or more positions 267, 273, and 328. Examples include S267E, V273E, L328F, or a combination thereof. In some particular examples, the variant Fc region is one of G4m1, G4m-1, G4m2, G4m-2, G4m3, G4m4, G4m5, G4m7, G4m8, G4m9, G4m10, G4m17, G4m18, G4m19, G4m20, G4m25, G4m27, G4m28, G4m29, G4m30, and G4mPE.

Any of the variant Fc regions described herein may exhibit an enhanced binding activity and/or an enhanced selectivity to FcγRIIB as compared with the parent wild-type Fc region. Alternatively, the variant Fc regions described herein may have low or no binding activity to any of the FcγR receptors. Alternatively or in addition, the variant Fc region binds FcRn.

Also described herein is a protein that comprises a variant Fc region of an IgG2 or IgG4 molecule (*e.g.*, human IgG2 or human IgG4 molecule). Such a variant Fc region may comprise a mutation at position 267, position 273, position 328, or a combination thereof as compared to a wild-type parent IgG2 or IgG4 Fc region. The numbering is according to the EU index.

In another aspect, provided herein is a pharmaceutical composition, comprising a protein that contains any of the variant Fc regions as described herein and a pharmaceutically

acceptable carrier. Such a pharmaceutical composition may be used to selectively activate an immune response in a subject.

In yet another aspect, the present disclosure provides a method for selectively activating an immune response in a subject, the method comprising administering to a subject
5 in need thereof an effective amount of a therapeutic agent, wherein the therapeutic agent comprises a first moiety that binds an immune cell receptor and a second moiety that binds Fc γ RIIB. In some examples, the immune cell receptor is a tumor necrosis factor receptor superfamily (TNFSF) member. Examples include, but are not limited to, FAS, TNFRSF12A, 4-1BB/CD137, TNFRSF13B, TNFRSF13C, CD27/TNFRSF7, CD30/TNFRSF8,
10 CD40/TNFRSF5, DR3/TNFRSF25, DR4/TNFRSF10A, DR5/TNFRSF10B, DR6/TNFRSF21, GITR/TNFRSF18, HVEM/TNFRSF14, LT β R, OX40/TNFRSF4, TROY/TNFRSF19, RELT/TNFRSF19L, TNFRSF12A, TNFRSF13B, TL1A/TNFSF15, TNFRSF17, TNFRSF1A, TNFRSF11B, RANK/TNFRSF11A, TNFRSF11B, NGFR, EDA2R, and TNFRSF1B.

15 In some embodiments the therapeutic agent can be an antibody (*e.g.*, an agonist antibody), which may be an IgG1, IgG2, or IgG4 molecule (*e.g.*, a human IgG1, human IgG2, or human IgG4 molecule).

In some embodiments, the therapeutic agent may selective bind Fc γ RIIB. For example, the therapeutic agent may contain a variant Fc region having a selective binding
20 affinity to Fc γ RIIB (*e.g.*, any of the variant Fc regions described herein). Alternatively or in addition, the variant Fc region may have an enhanced binding activity to Fc γ RIIB as compared with a wild-type parent Fc region.

In other embodiments, the therapeutic agent may be a bispecific antibody comprising a first moiety, which is a first antigen-binding fragment specific to the immune cell receptor,
25 and a second moiety, which is a second antigen-binding fragment specific to Fc γ RIIB.

In any of the methods described herein, the subject can be a human patient having or suspected of having a cancer. Exemplary cancers include lung cancer, stomach cancer, liver cancer, breast cancer, skin cancer, pancreatic cancer, brain cancer, prostate cancer, bladder cancer, colorectal cancer, sarcoma, bone cancer, lymphoma and a hematological cancer.

30 Also within the scope of the present disclosure are pharmaceutical compositions for

use in enhancing immune responses, comprising a protein that contains any of the variant Fc regions as described herein and a pharmaceutically acceptable carrier, and uses of such proteins for manufacturing the medicament for use in achieving the intended therapeutic purposes, for example, cancer treatment.

The details of one or more embodiments of the invention are set forth in the description below. Other features or advantages of the present invention will be apparent from the following drawings and detailed description of several embodiments, and also from the appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGs. 1A-1W are charts showing binding activity of various IgG variants as indicated to different types of Fc γ receptors expressed on CHO-K1 cells at the various concentrations as indicated. The concentrations of each IgG variant, from left to right, are 0 μ g/ml, 0.3 μ g/ml, 1 μ g/ml, 3 μ g/ml, and 10 μ g/ml in FIGs. 1A-1C, 1E-J, 1L-1M, 1P-1R, and 1T-1V. For FIGs. 1D, 1K, 1O, 1S, and 1w the bars, from left to right, correspond to the following concentrations of IgG variant: 0.3 μ g/ml, 1 μ g/ml, 3 μ g/ml, and 10 μ g/ml.

FIGs. 2A-2C are charts showing stimulation of human CD8⁺ T cells in co-culture with parental or Fc γ R expressing cells by a number of IgG variants as indicated by IFN- γ secretion. The groups, from left to right, correspond to: no OKT, 0.01 μ g/ml, 0.03 μ g/ml, 0.1 μ g/ml, and 0.3 μ g/ml, in FIG. 2A. In FIGs. 2B and 2C, the groups, from left to right, correspond to 0.01 μ g/ml, 0.03 μ g/ml, and 0.1 μ g/ml.

FIG. 3 is a chart showing murine T cell activation by a number of IgG molecules in the presence of anti-CD3 and/or anti-4-1BB antibodies.

FIGs. 4A-4B are charts showing *in vivo* anti-tumor effects and the induction of serum ALT levels by anti-murine CD137 antibodies of different IgG isoforms.

FIGs. 5A-5H are charts showing the antibody concentration of IgG variants in plasma after a single intravenous injection of 1-3 mg/kg. Male C57BL/6 mice were used, and four mice were assayed per time point. The charts of FIG. 5A show the plasma concentrations G1m27 after an IV dose of 3 mg/kg (top graph) and of G4m10 after an IV dose of 1.45 mg/kg (bottom graph). The charts of FIG. 5B show the plasma concentrations of G1m2 after an IV dose of 3 mg/kg (top graph) and of G4m2 after an IV dose of 3 mg/kg (bottom graph). The

charts of FIG. 5C show the plasma concentrations of G1m29 after an IV dose of 3 mg/kg (top graph) and of G4m7 after an IV dose of 3 mg/kg (bottom graph). The charts of FIG. 5D show the plasma concentrations of G1m28 after an IV dose of 3 mg/kg (top graph) and of G4 after an IV dose of 3 mg/kg. The charts of FIG. 5E show the plasma concentrations of
 5 G4m28 after an IV dose of 1.45 mg/kg (top graph) and of G4m29 after an IV dose of 1.5 mg/kg (bottom graph). The charts of FIG. 5F show the plasma concentrations of G4m1 after an IV dose of 3 mg/kg (top graph) and of G4m27 after an IV dose of 1.45 mg/kg (bottom graph). FIG. 5G shows the plasma concentrations of G2m19 after an IV dose of 3 mg/kg (top graph) and of G2m1 after an IV dose of 3 mg/kg (bottom graph). FIG. 5H shows the plasma
 10 concentrations of G2 (wild type) after an IV dose 3 mg/kg (top graph) and of G1Maa after an IV dose of 3 mg/kg (bottom graph).

DETAILED DESCRIPTION OF THE INVENTION

Agonistic anti-TNFR antibodies that are capable of binding to FcγRIIB (CD32B)
 15 showed enhanced activity in mouse models and mouse IgG1 isoform showed the best anti-tumor efficacy. These results indicate that agonistic antibodies having selective binding activity to the FcγRIIB (CD32B) would be expected to have better agonistic activity in stimulating immune responses. Further, it was observed that certain anti-tumor agonist antibodies showed differential liver toxicity, suggesting a novel approach for designing
 20 therapeutic agonist antibodies with enhanced therapeutic index (*e.g.*, high treatment efficacy and low toxicity).

Accordingly, described herein are approaches for designing IgG molecules (*e.g.*, IgG1, IgG2, and IgG4 molecules such as human IgG1, human IgG2, and human IgG4 molecules) containing an Fc variant having selective binding activity to FcγRIIB (CD32B) as
 25 relative to other Fc receptors such as FcγRIII (CD16), and cancer treatment methods, which involve the use of therapeutic agents capable of cross-linking an immune cell receptor such as a receptor of the tumor necrosis factor receptor superfamily (TNFSF) and FcγRIIB.

I. Fc variants

Described herein are Fc variants having enhanced selectivity to Fc γ RIIB relative to its wild-type counterpart. An Fc fragment having selectivity to Fc γ RIIB, selectively binding to Fc γ RIIB, or specifically binding to Fc γ RIIB is a term well understood in the art. A molecule is said to exhibit “selective binding” or “specific binding” if it reacts more frequently, more rapidly, with greater duration and/or with greater affinity with a particular target antigen (*e.g.*, an Fc γ RIIB receptor) than it does with alternative targets (*e.g.*, Fc γ RIII receptors). An Fc fragment “specifically binds” to an Fc receptor if it binds with greater affinity, avidity, more readily, and/or with greater duration than it binds to other Fc receptors. For example, an Fc fragment that specifically (or preferentially) binds to Fc γ RIIB is an Fc fragment that binds this Fc receptor with greater affinity, avidity, more readily, and/or with greater duration than it binds to other Fc receptors. It is also understood with this definition that, for example, an Fc fragment that selectively or specifically binds to a first Fc receptor may or may not specifically or preferentially bind to a second Fc receptor. As such, “selective binding,” “specific binding” or “preferential binding” does not necessarily require (although it can include) exclusive binding. In some examples, an Fc fragment that “selectively binds,” or “specifically binds” to a target Fc receptor (*e.g.*, Fc γ RIIB) may not bind to other Fc receptors (*i.e.*, binding not detectable by routine methods). Relative binding affinities of IgG1, IgG2, and IgG4 to different Fc receptors are given in Table 1 below.

Table 1. Relative Binding Affinities of Human and Mouse Immunoglobulins to Fc Receptors

Human				Mouse			
Fc γ R	IgG1	IgG2	IgG4	Fc γ R	IgG1	IgG2a	IgG2b
I	++++	-	++++	I	-	++++	++++
IIa (H131)	+++	++	++	III	++	++	++
IIa (R131)	+++	+	++				
IIb	++	+/-	++	IIb	+++	++	+++
IIIa (V158)	+++	+	++	IV	-	+++	++
IIIa (F158)	++	+/-	++				

The Fc variants described herein may have enhanced selectivity to Fc γ RIIB relative to their wild-type counterparts (the wild-type parent Fc region in which mutations are introduced to produce the Fc variants). The relative binding activity to Fc γ RIIB *versus* another Fc receptor (*e.g.*, Fc γ RIII) of such an Fc variant is higher than the relative binding

activity to Fc γ RIIB *versus* the other Fc receptor (*e.g.*, Fc γ RIII) of the wild-type counterpart. The Fc variant may have enhanced binding activity to Fc γ RIIB and/or decreased binding activity to another Fc receptor, for example, Fc γ RIII. In some embodiments, the Fc variants described herein may have decreased binding activity to both Fc γ RIIB and another Fc
5 receptor (for example, Fc γ RIII); however, the level of decreased binding activity to the other Fc receptor (*e.g.*, Fc γ RIII) is greater than the level of decreased binding activity to Fc γ RIIB.

In some embodiments, an Fc variant as described herein has a suitable binding affinity for Fc γ RIIB, *e.g.*, enhanced as compared with the wild-type parent Fc from which the Fc variant is derived. As used herein, “binding affinity” refers to the apparent association
10 constant or K_A . The K_A is the reciprocal of the dissociation constant (K_D). The Fc variant described herein may have a binding affinity (K_D) of at least 10^{-5} , 10^{-6} , 10^{-7} , 10^{-8} , 10^{-9} , 10^{-10} M, or lower for Fc γ RIIB. An increased binding affinity corresponds to a decreased K_D . Higher affinity binding of an Fc fragment for a first Fc receptor relative to a second Fc receptor can be indicated by a higher K_A (or a smaller numerical value K_D) for binding the
15 first Fc receptor than the K_A (or numerical value K_D) for binding the second Fc receptor. In such cases, the Fc variant has specificity for the first Fc receptor relative to the second Fc receptor. In some embodiments, the Fc variants described herein have a higher binding affinity (a higher K_A or smaller K_D) to Fc γ RIIB as compared to the binding affinity to Fc γ RIII (either Fc γ RIIIA or Fc γ RIIIB). Differences in binding affinity (*e.g.*, for specificity or other
20 comparisons) can be at least 1.5, 2, 3, 4, 5, 10, 15, 20, 37.5, 50, 70, 80, 91, 100, 500, 1000, 10,000 or 10^5 fold.

Binding affinity (or binding specificity) can be determined by a variety of methods including equilibrium dialysis, equilibrium binding, gel filtration, ELISA, surface plasmon resonance, or spectroscopy (*e.g.*, using a fluorescence assay). See also Examples below.
25 Exemplary conditions for evaluating binding affinity are in HBS-P buffer (10 mM HEPES pH7.4, 150 mM NaCl, 0.005% (v/v) Surfactant P20). These techniques can be used to measure the concentration of bound binding protein as a function of target protein concentration. The concentration of bound binding protein ([Bound]) is generally related to the concentration of free target protein ([Free]) by the following equation:

$$[\text{Bound}] = [\text{Free}] / (K_d + [\text{Free}])$$

It is not always necessary to make an exact determination of K_A , though, since sometimes it is sufficient to obtain a quantitative measurement of affinity, e.g., determined using a method such as ELISA or FACS analysis, is proportional to K_A , and thus can be used for comparisons, such as determining whether a higher affinity is, e.g., 2-fold higher, to obtain a qualitative measurement of affinity, or to obtain an inference of affinity, e.g., by activity in a functional assay (an *in vitro* or *in vivo* assay).

In some embodiments, the Fc variants described herein may be designed by mutating one or more amino acid residues in the wild-type of human IgG1, IgG2, or IgG4 Fc fragments in light of the amino acid residues in the corresponding mouse IgG, for example, mouse IgG1. A sequence comparison of human and mouse IgGs is provided below (SEQ ID Nos: 67-69, 64, and 65, from top to bottom, each representing a combination of fragments 211-245, 260-278, and 320-332 of the corresponding Fc region):

	21-	22-	23-	24-	...26-	27-	...32-	33-
15	12345678	9012345	6789	0123456789	012345...	7 89012	3 45678...	01234567 8 9012
	<u>Upper</u>		<u>Core</u>	<u>Lower</u>	CH2			
	hIgG1	VDKKVEPK-SCDKTHT CPPC PAPELLGGPSVFLFPP...TCVVVDVSHEDPEVKFNWY...KCKVSNKALPAPI						
	hIgG2	VDKTVERK-CC-V-E- CPPC PAPPVA-GPSVFLFPP...TCVVVDVSHEDPEVQFNWY...KCKVSNKGLPAPI						
	hIgG4	VDKRVESKYG---PP CPSC PAPEFLGGPSVFLFPP...TCVVVDVSQEDPEVQFNWY...KCKVSNKGLPSSI						
20	mIgG1	VDKKIVPR-DC--G CKPCIC TVPEVS---SVFIFPP...TCVVVDISKDDPEVQFSWF...KCRVNSAAFPAPI						
	mIgG2a	VDKKIEPRGPTIKP CPPCKC PAPNLLGGPSVFIFFPP...TCVVVDVSEDDPDVQISWF...KCKVNNKDLPAPI						

In some embodiments, the Fc variants described herein is a human IgG1, G2, or G4 Fc variants comprising one or more mutations (e.g., amino acid substitutions, deletions, or additions) in the hinge domain of an Fc fragment. Human IgGs contain a core motif of CPPC or CPSC in the hinge domain (positions 226-229 according to the EU index). Positions 216 to 225 are deemed as the upper portion of the hinge domain and positions 230-238 are deemed as the lower portion of the hinge domain. The numbering system used herein, unless explicitly indicated, is according to the EU index. In some examples, the one or more mutations can be located in the upper portion of the hinge domain. Alternatively or in addition, the one or more mutations can be located in the lower portion of the hinge domain.

The mutations to a human IgG Fc can be made according to the corresponding amino acid residues in the hinge domain of mouse IgG1. For example, mouse IgG1 does not

contain the GGP motif at positions 236-238. Accordingly, one or more of the residues in this GGP motif can be deleted from a human IgG1, IgG2, or IgG4 Fc fragment to produce the Fc variants described herein.

Alternatively or in addition, the human Fc variants may contain one or more amino acid substitutions in the upper portion, in the lower portion, or both of the hinge domain. For example, the Fc variant may comprise one or more amino acid substitutions at one or more of positions 233, 234, 235, and/or 236. Such an amino acid substitution may be in combination with the deletion of one or more of the GGP motif (236-238) noted herein. These mutations may be introduced into a human IgG2 or IgG4 Fc fragment to produce the Fc variants described herein. In some examples, the Fc variants described herein contains a deletion at one or more of the positions 236-238 (*e.g.*, 236, 237, 238 or any combination thereof)

Any of the mutations in the hinge domain described herein may be in combination with a mutation (*e.g.*, amino acid substitutions) at one or more positions that are involved in interaction with an Fc receptor. Such positions include, but are not limited to, positions 267, 273, 328, and/or 329. Exemplary amino acid substitutions at those positions include S261E, V271E, L328F, and P329G. Fc variants derived from IgG2 or IgG4 molecules that contain one or more mutations at positions 267, 273, 328, and/or 329 are also within the scope of the present disclosure. Such mutations may include amino acid substitutions at one or more of these positions, for example, S261E, V271E, L328F, and/or P329G.

In some embodiments, an Fc variant described herein may comprise an amino acid sequence at least 85% identical (*e.g.*, 90%, 95%, 98%, 99%, or above) to that of its wild-type counterpart (*e.g.*, the Fc fragment of wild-type human IgG1, IgG2, or IgG4 described herein). The “percent identity” of two amino acid sequences is determined using the algorithm of Karlin and Altschul Proc. Natl. Acad. Sci. USA 87:2264-68, 1990, modified as in Karlin and Altschul Proc. Natl. Acad. Sci. USA 90:5873-77, 1993. Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. J. Mol. Biol. 215:403-10, 1990. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to the protein molecules of interest. Where gaps exist between two sequences, Gapped BLAST can be utilized as described in Altschul et al., Nucleic Acids Res. 25(17):3389-3402, 1997. When

utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used.

In one example, the amino acid residue substitutions in an Fc variant described herein are conservative amino acid residue substitutions. As used herein, a “conservative amino acid substitution” refers to an amino acid substitution that does not alter the relative charge or size characteristics of the protein in which the amino acid substitution is made. Variants can be prepared according to methods for altering polypeptide sequence known to one of ordinary skill in the art such as are found in references which compile such methods, *e.g.*, Molecular Cloning: A Laboratory Manual, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or Current Protocols in Molecular Biology, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. Conservative substitutions of amino acids include substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D.

Provided below is a sequence alignment showing exemplary positions where mutations can be introduced into hIgG1, hIgG2, and IgG4 to produce various Fc variants for the present disclosure.

Sequence alignment of human IgG1 variants relative to wild-type human IgG1 (SEQ ID NOs:70-88, from top to bottom):

		21-	22-	23-	24-	...26-	27-	...32-	33-																																																													
		12345678	9012345	678901234567890123456789012345...01234567	8901234567	345678	9012345678	9012345678	9012																																																													
		<u>Upper</u>			<u>Core</u>			<u>Lower</u>		CH2																																																												
25	hIgG1	V	D	K	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	G	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I
	hIgG1m1	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I					
	hIgG1m2	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
	hIgG1m-2	K	K	V	E	P	K	-	C	C	-	V	-	E	-	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I		
	hIgG1m-4	K	K	V	E	P	K	-	G	...	PP	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I						
30	hIgG1m5	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
	hIgG1m7	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
	hIgG1m8	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
	hIgG1m9	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
	hIgG1m15	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I					
35	hIgG1m17	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I					
	hIgG1m18	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I					
	hIgG1m19	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I					
	hIgG1m25	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
	hIgG1m27	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
40	hIgG1m28	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
	hIgG1m29	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I			
	hIgG1mAA	K	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I		
	hIgG1mAG	K	K	V	E	P	K	-	S	C	D	K	T	H	T	CP	P	C	P	A	P	E	L	L	...	G	P	S	V	F	L	F	P	P	...	T	C	V	V	V	D	V	S	H	E	D	P	E	V	K	F	N	W	Y	...	K	C	K	V	S	N	K	A	L	P	A	P	I		

5

Sequence alignment of human IgG2 variants relative to wild-type human IgG2 (SEQ ID NOs: 89-105, from top to bottom):

10		21-	22-	23-	24-	...26-	27-	...32-	33-
		12345678	9012345	6789 0123456789012345...	0123456	7 89012	3 45678...	01234567	8 9012
		<u>Upper</u>		<u>Core</u>	<u>Lower</u>	CH2			
	hIgG2	VDKT	VERK-CC-V-E-	CPPC	PAPPVA-GPSVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNKGLPAPI		
	hIgG2m1	TVERK-CC-V-E-	CPPC	PAPPVA- ~~---~~	SVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNKGLPAPI		
	hIgG2m-1	KTVERK-SC	DKTHT	CPPC	PAPPVA-GPSVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNKGLPAPI		
15	hIgG2m2	TVERK-CC-V-E-	CPPC	PAPP FLG	GPSVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNKGLPAPI		
	hIgG2m-4	KTVERK YG	~~---~~	PP	CPPC	PAPPVA-GPSVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNKGLPAPI	
	hIgG2m5	TVERK-CC-V-E-	CPPC	PAPPVA-GPSVFLFPP...	TCVVVDVSHEDPE E	QFNWY...	KCKVSNKGLPAPI		
	hIgG2m7	TVERK-CC-V-E-	CPPC	PAPPVA-GPSVFLFPP...	TCVVVDV E	HEDPEVQFNWY...	KCKVSNKGLPAPI		
	hIgG2m8	TVERK-CC-V-E-	CPPC	PAPPVA-GPSVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNK G	F PAPI		
20	hIgG2m9	TVERK-CC-V-E-	CPPC	PAPPVA-GPSVFLFPP...	TCVVVDV E	HEDPEVQFNWY...	KCKVSNK G	F PAPI	
	hIgG2m10	TVERK-CC-V-E-	CPPC	PAP EVS	~~---~~	SVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNK G	F PAPI
	hIgG2m15	TVERK-CC-V-E-	CPPC	PAPPVA- ~~---~~	SVFLFPP...	TCVVVDVSHEDPE E	QFNWY...	KCKVSNKGLPAPI	
	hIgG2m17	TVERK-CC-V-E-	CPPC	PAPPVA- ~~---~~	SVFLFPP...	TCVVVDV E	HEDPEVQFNWY...	KCKVSNKGLPAPI	
	hIgG2m18	TVERK-CC-V-E-	CPPC	PAPPVA- ~~---~~	SVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNK G	F PAPI	
25	hIgG2m19	TVERK-CC-V-E-	CPPC	PAPPVA- ~~---~~	SVFLFPP...	TCVVVDV E	HEDPEVQFNWY...	KCKVSNK G	F PAPI
	hIgG2m20	TVERK-CC-V-E-	CPPC	PAPPVA- ~~---~~	PSVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNKGLPAPI		
	hIgG2m27	TVERK-CC-V-E-	CPPC	PAPPVA- ~~---~~	PSVFLFPP...	TCVVVDV E	HEDPEVQFNWY...	KCKVSNKGLPAPI	
	hIgG2m28	TVERK-CC-V-E-	CPPC	PAPPVA- ~~---~~	PSVFLFPP...	TCVVVDVSHEDPEVQFNWY...	KCKVSNK G	F PAPI	

30

Sequence alignment of human IgG4 variants relative to wild-type human IgG4 (SEQ ID NOs: 106-117, 66, and 118-127, from top to bottom)

		21-	22-	23-	24-	...26-	27-	...32-	33-
35		12345678	9012345	6789 0123456789012345...	0123456	7 89012	3 45678...	01234567	8 9012
		Upper		Core	Lower		CH2		
	hIgG4	VDKR	VESKYG----	PP CP	SCPAPEFLGGPSVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNKGLPSSI		
	hIgG4m1	KR	VESKYG----	PP CP	SCPAPEFL	SVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNKGLPSSI	
40	hIgG4m-1	KR	VESK SCDKTHT	CPPC	PAPEFLGGPSVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNKGLPSSI		
	hIgG4m2	KR	VESKYG----	PP CP	SCPAPEFL	GPSVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNKGLPSSI	
	hIgG4m-2	KR	VESK- CC-V-E	CPPC	PAPEFLGGPSVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNKGLPSSI		
	hIgG4m3	KR	VESKYG----	PP CP	SCPAPEFLGGPSVFLFPP...	TCVVVDVSQEDPE E QFNWY...	KCKVSNKGLPSSI		
	hIgG4m4	KR	VESKYG----	PP CP	SCPAPEFL	SVFLFPP...	TCVVVDVSQEDPE E QFNWY...	KCKVSNKGLPSSI	
45	hIgG4m5	KR	VESKYG----	PP CP	SCPAPEFL	GPSVFLFPP...	TCVVVDVSQEDPE E QFNWY...	KCKVSNKGLPSSI	
	hIgG4m7	KR	VESKYG----	PP CP	SCPAPEFLGGPSVFLFPP...	TCVVVDV E QEDPEVQFNWY...	KCKVSNKGLPSSI		
	hIgG4m8	KR	VESKYG----	PP CP	SCPAPEFLGGPSVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNK G FPSI		
	hIgG4m9	KR	VESKYG----	PP CP	SCPAPEFLGGPSVFLFPP...	TCVVVDV E QEDPEVQFNWY...	KCKVSNK G FPSI		
	hIgG4m10	KR	VESKYG----	PP CP	SCPAPE VS	SVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNK G FPSI	
50	hIgG4m15	KR	VESKYG----	PP CP	SCPAPEFL	SVFLFPP...	TCVVVDVSQEDPE E QFNWY...	KCKVSNKGLPSSI	
	hIgG4m17	KR	VESKYG----	PP CP	SCPAPEFL	SVFLFPP...	TCVVVDV E QEDPEVQFNWY...	KCKVSNKGLPSSI	
	hIgG4m18	KR	VESKYG----	PP CP	SCPAPEFL	SVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNK G FPSI	
	hIgG4m19	KR	VESKYG----	PP CP	SCPAPEFL	SVFLFPP...	TCVVVDV E QEDPEVQFNWY...	KCKVSNK G FPSI	
	hIgG4m20	KR	VESKYG----	PP CP	SCPAPEFLG	SVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNKGLPSSI	
55	hIgG4m25	KR	VESKYG----	PP CP	SCPAPEFL	GPSVFLFPP...	TCVVVDVSQEDPE E QFNWY...	KCKVSNKGLPSSI	
	hIgG4m27	KR	VESKYG----	PP CP	SCPAPEFL	GPSVFLFPP...	TCVVVDV E QEDPEVQFNWY...	KCKVSNKGLPSSI	
	hIgG4m28	KR	VESKYG----	PP CP	SCPAPEFL	GPSVFLFPP...	TCVVVDVSQEDPEVQFNWY...	KCKVSNK G FPSI	

hIgG4m29 KRVESKYG---PP**CP**PCPAPEFLGPSVFLFPP...TCVVVDV**E**QEDPEVQFNWY...KCKVSNKG**E**PSI
hIgG4m30 KRVESKYG---PP**CP**PCPAPEFLGPSVFLFPP...TCVVVDVSQEDPEVQFNWY...KCKVSNKGLPSSI
hIgG4mPE KRVESKYG---PP**CP**PCPAPEF**E**GGPSVFLFPP...TCVVVDVSQEDPEVQFNWY...KCKVSNKGLPSSI

5

The amino acid sequences of wild-type human IgG1, IgG2, and IgG4 Fc fragments, and a number of exemplary hIgG1, hIgG2, and hIgG4 Fc variants are provided below:

Amino acid sequence of wild-type human IgG1 Fc fragment:

10 VDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNH
YTQKSLSLSPGK (SEQ ID NO: 1)

Amino acid sequence of wild-type human IgG2 Fc fragment:

15 VDKTVERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKT
KPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTTTPMLDSGDSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHNYTQK
SLSLSPGK (SEQ ID NO: 2)

Amino acid sequence of wild-type human IgG4 Fc fragment:

20 VDKRVESKYGPPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAK
TKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSRLTVDKSRWQEGNVFSQSVMEALHNHNYTQ
KSLSLSLGK (SEQ ID NO: 3)

Amino acid sequence of human IgG4 S228P Fc variant:

25 VDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAK
TKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSRLTVDKSRWQEGNVFSQSVMEALHNHNYTQ
KSLSLSLGK (SEQ ID NO: 4)

30 **Amino acid sequences of exemplary human IgG1 Fc variants:**

G1m1:

VDKKVEPKSCDKTHTCPPCPAPELLSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK
TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHNYTQ
35 KSLSLSPGK (SEQ ID NO: 5)

G1m2:

VDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN
AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHNY
40 TQKSLSLSPGK (SEQ ID NO: 6)

G1m-2:

WO 2018/183520

PCT/US2018/024872

VDKKVEPKCCVECPPCPAPELLSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKP
REEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
SLSPGK (SEQ ID NO: 7)

5

Glm-4:

VDKKVEPKYGPPCPPCPAPELLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT
KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQK
SLSLSPGK (SEQ ID NO: 8)

10

Glm5:

VDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEEKFNWYVDGVEVH
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNH
YTQKSLSLSPGK (SEQ ID NO: 9)

15

Glm7:

VDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVKFNWYVDGVEVH
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNH
YTQKSLSLSPGK (SEQ ID NO: 10)

20

Glm8:

VDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAFFAPIEKTISKAKGQPREPQVYTLPPSREEMTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNH
YTQKSLSLSPGK (SEQ ID NO: 11)

25

Glm9:

VDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVKFNWYVDGVEVH
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAFFAPIEKTISKAKGQPREPQVYTLPPSREEMTK
NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNH
YTQKSLSLSPGK (SEQ ID NO: 12)

30

Glm15:

VDKKVEPKSCDKTHTCPPCPAPELLSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEEKFNWYVDGVEVHNAK
TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQ
KSLSLSPGK (SEQ ID NO: 13)

35

40

Glm17:

VDKKVEPKSCDKTHTCPPCPAPELLSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVKFNWYVDGVEVHNAK
TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQ
KSLSLSPGK (SEQ ID NO: 14)

45

Glm18:

5 VDKKVEPKSCDKTHTCPPCPAPELLSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK
TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAFPAIEKTIISKAKGQPREPQVYTLPPSREEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHYTQ
KSLSLSPGK (SEQ ID NO: 15)

Glm19:

10 VDKKVEPKSCDKTHTCPPCPAPELLSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVKFNWYVDGVEVHNAK
TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAFPAIEKTIISKAKGQPREPQVYTLPPSREEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHYTQ
KSLSLSPGK (SEQ ID NO: 16)

Glm25:

15 VDKKVEPKSCDKTHTCPPCPAPELLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK
AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHY
TQKSLSLSPGK (SEQ ID NO: 17)

Glm27:

20 VDKKVEPKSCDKTHTCPPCPAPELLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVKFNWYVDGVEVHNAK
AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHY
TQKSLSLSPGK (SEQ ID NO: 18)

25 *Glm28:*

VDKKVEPKSCDKTHTCPPCPAPELLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK
AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAFPAIEKTIISKAKGQPREPQVYTLPPSREEMTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHY
TQKSLSLSPGK (SEQ ID NO: 19)

30

Glm29:

VDKKVEPKSCDKTHTCPPCPAPELLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVKFNWYVDGVEVHNAK
AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAFPAIEKTIISKAKGQPREPQVYTLPPSREEMTKN
QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHY
35 TQKSLSLSPGK (SEQ ID NO: 20)

GlmAA:

VDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN
40 NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHY
YTQKSLSLSPGK (SEQ ID NO: 21)

GlmAG:

VDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK
NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVYTLPPSREEMTKN
45 NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMHEALHNHY
YTQKSLSLSPGK (SEQ ID NO: 22)

Amino acid sequences of exemplary human IgG2 Fc variants:***G2m1:***

VDKTVERKCCVECPPCPAPPVASVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKP
 5 REEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLT
 CLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSL
 SLSPGK (SEQ ID NO: 23)

G2m-1:

VDKTVERKSCDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHN
 10 AKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKN
 QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHY
 TQKSLSLSPGK (SEQ ID NO: 24)

G2m2:

VDKTVERKCCVECPPCPAPPFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAK
 TKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQV
 SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQ
 KSLSLSPGK (SEQ ID NO: 25)

20

G2m-4:

VDKTVERKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKT
 KPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS
 LTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQK
 25 SLSLSPGK (SEQ ID NO: 26)

G2m5:

VDKTVERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEEQFNWYVDGVEVHNAKT
 KPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS
 30 LTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQK
 SLSLSPGK (SEQ ID NO: 27)

G2m7:

VDKTVERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVQFNWYVDGVEVHNAKT
 35 KPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS
 LTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQK
 SLSLSPGK (SEQ ID NO: 28)

G2m8:

VDKTVERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKT
 40 KPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS
 LTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQK
 SLSLSPGK (SEQ ID NO: 29)

G2m9:

VDKTVERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVQFNWYVDGVEVHNAKT
 KPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS

WO 2018/183520

PCT/US2018/024872

LTCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQK
SLSLSPGK (SEQ ID NO: 30)

5

G2m10:

VDKTVKCCVECPPCPAPEVSSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKP
REEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGFPAIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLT
10 CLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
SLSPGK (SEQ ID NO: 31)

G2m15:

VDKTVKCCVECPPCPAPPVASVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEEQFNWYVDGVEVHNAKTKP
15 REEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
SLSPGK (SEQ ID NO: 32)

G2m17:

VDKTVKCCVECPPCPAPPVASVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKP
20 REEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
SLSPGK (SEQ ID NO: 33)

25 ***G2m18:***

VDKTVKCCVECPPCPAPPVASVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKP
REEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGFPAIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
SLSPGK (SEQ ID NO: 34)

30

G2m19:

VDKTVKCCVECPPCPAPPVASVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKP
REEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGFPAIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL
35 SLSPGK (SEQ ID NO: 35)

G2m20:

VDKTVKCCVECPPCPAPPVAPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTK
PREEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSL
40 TCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKS
LSLSPGK (SEQ ID NO: 36)

G2m27:

VDKTVKCCVECPPCPAPPVAPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTK
45 PREEQFNSTFRVVSFLTIVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKS
LSLSPGK (SEQ ID NO: 37)

G2m28:

VDKTVERKCCVECPPCPAPPVAPSVFLFPPKPKDTLMISRTPEVTCVVVDVEHEDPEVQFNWYVDGVEVHNAKTK
 PREEQFNSTFRVSVSLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSL
 5 TCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKS
 LSLSPGK (SEQ ID NO: 38)

Amino acid sequences of exemplary human IgG4 Fc variants:***G4m1:***

10 VDKRVESKYGPPCPPCPAPEFLSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKP
 REEQFNSTYRVVSVSLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT
 CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL
 SLSLGK (SEQ ID NO: 39)

G4m-1:

15 VDKRVESKSCDKTHTPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVE
 VHNAKTKPREEQFNSTYRVVSVSLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEM
 TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALH
 NHYTQKSLSLSLGK (SEQ ID NO: 40)

20

G4m2:

VDKRVESKYGPPCPPCPAPEFLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKT
 KPREEQFNSTYRVVSVSLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVS
 LTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQK
 25 SLSLSLGK (SEQ ID NO: 41)

G4m-2:

VDKRVESKCCVEPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHN
 AKTKPREEQFNSTYRVVSVSLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKN
 30 QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHY
 TQKSLSLSLGK (SEQ ID NO: 42)

G4m3:

VDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEEQFNWYVDGVEVHNAK
 35 TKPREEQFNSTYRVVSVSLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
 SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQ
 KSLSLSLGK (SEQ ID NO: 43)

G4m4:

40 VDKRVESKYGPPCPPCPAPEFLSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEEQFNWYVDGVEVHNAKTKP
 REEQFNSTYRVVSVSLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT
 CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL
 SLSLGK (SEQ ID NO: 44)

G4m5:

45 VDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEEQFNWYVDGVEVHNAK
 TKPREEQFNSTYRVVSVSLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV

SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQ
KSLSLSLGK (SEQ ID NO: 45)

G4m7:

5 VDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEQEDPEVQFNWYVDGVEVHNAK
TKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQ
KSLSLSLGK (SEQ ID NO: 46)

10 *G4m8:*

VDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAK
TKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGFPSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQ
KSLSLSLGK (SEQ ID NO: 47)

15

G4m9:

VDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEQEDPEVQFNWYVDGVEVHNAK
TKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGFPSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQ
20 KSLSLSLGK (SEQ ID NO: 48)

G4m10:

VDKRVESKYGPPCPPCPAPEVSSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKP
REEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGFPSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT
25 CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL
SLSLGK (SEQ ID NO: 49)

G4m15

VDKRVESKYGPPCPPCPAPEFLSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEEQFNWYVDGVEVHNAKTKP
30 REEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL
SLSLGK (SEQ ID NO: 128)

G4m17:

35 VDKRVESKYGPPCPPCPAPEFLSVFLFPPKPKDTLMISRTPEVTCVVVDVEQEDPEVQFNWYVDGVEVHNAKTKP
REEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL
SLSLGK (SEQ ID NO: 50)

40 *G4m18:*

VDKRVESKYGPPCPPCPAPEFLSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKP
REEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGFPSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSL
SLSLGK (SEQ ID NO: 51)

45

G4m19:

VDKRVESKYGPPCPPCPAPEFLSVFLFPPKPKDTLMISRTPEVTCVVVDVEQEDPEVQFNWYVDGVEVHNAKTKP

REEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGFPSSEIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT
CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQKSL
LSLSLGK (SEQ ID NO: 52)

5 *G4m20:*

VDKRVESKYGPPCPPCPAPEFLGSPVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTK
PREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQKS
LSLSLGK (SEQ ID NO: 53)

10

G4m25:

VDKRVESKYGPPCPPCPAPEFLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEEQFNWYVDGVEVHNAKT
KPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQK
SLSLSLGK (SEQ ID NO: 54)

15

G4m27:

VDKRVESKYGPPCPPCPAPEFLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEQEDPEVQFNWYVDGVEVHNAKT
KPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQK
LSLSLGK (SEQ ID NO: 55)

20

G4m28:

VDKRVESKYGPPCPPCPAPEFLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKT
KPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGFPSSEIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQK
LSLSLGK (SEQ ID NO: 56)

25

G4m29:

VDKRVESKYGPPCPPCPAPEFLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVEQEDPEVQFNWYVDGVEVHNAKT
KPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGFPSSEIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQK
LSLSLGK (SEQ ID NO: 57)

30

35 *G4m30:*

VDKRVESKYGPPCPPCPAPEFLPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTK
PREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQKS
LSLSLGK (SEQ ID NO: 58)

40

G4mPE:

VDKRVESKYGPPCPPCPAPEFEGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAK
TKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQ
KSLSLSLGK (SEQ ID NO: 59)

45

The Fc variants described herein may exhibit an enhanced binding activity to FcγRIIB

as compared with the wild-type counterpart. Examples include G2m2, G2m5, G2m7, G2m8, G2m9, G1m7, G1m9, and G4m7. Alternatively or in addition, the Fc variants may have an enhanced selectivity to Fc γ RIIB as compared with their wild-type counterparts, for example, G1m15, G1m17, G1m18, G1m19, G1m27, G1m28, G1m29, G4m1, G4m2, G4m7, G4m8, G4m9, G4m25, G4m27, and G4m28. These Fc variants can be used for constructing therapeutic agents described herein capable of cross-linking immune receptors and Fc γ RIIB receptor.

Alternatively, certain Fc variants as described herein may have low or no binding activity to any Fc γ R. Examples include G1m2, G1m25, G4m5, G4m18, G4m19, and G4m20. Such Fc variants may retain the binding activity to FcRn. Therapeutic agents (*e.g.*, antibodies) containing such Fc variants would have low or no activity to activate Fc γ receptors and may have reduced toxicity relative to their wild-type counterparts.

The changes of binding affinity/specificity of the exemplary Fc variants as compared with their wild-type counterparts are provided in Tables 2-4 below. “N/A” indicates no data available. When the binding activity of an Fc variant is found to be “no change” as compared with the wild-type counterpart, it means that there is no significant variation of the binding activity between the Fc variant and the wild-type counterpart as indicated by the same assay under the same experimental conditions. When the binding activity of an Fc variant is “increased” or “decreased” as relative to its wild-type counterpart means that the binding activity of the Fc variant is higher or lower than that of the wild-type counterpart as determined by the same assay under the same experimental conditions and the variation is significant (*e.g.*, biologically significant) as known to those skilled in the art. When the binding activity of an Fc variant is “slightly increased” or “slightly decreased” as relative to its wild-type counterpart means that the binding activity of the Fc variant is higher or lower than that of the wild-type counterpart as determined by the same assay under the same experimental conditions and the variation is statistically significant but to a limited level (*e.g.*, up to 10%).

Table 2. Fc γ R Binding Activity of Human IgG1 Mutants As Relative to Wild-Type Human IgG1

	Changes of Binding Activity Relative to Wild-Type Counterparts
--	--

IgG1 Mutant	FcγRI	FcγRIIA(H131)	FcγRIIA(R131)	FcγRIIB	FcγRIIC	FcγRIII
G1m1	Decreased	Decreased	Decreased	Decreased	Decreased	Decreased
G1m2	Decreased	Decreased	Decreased	Decreased	Decreased	Decreased
G1m-2	No change	N/A	N/A	No change	N/A	No change
G1m-4	No change	N/A	N/A	No change	N/A	No change
G1m5	Increased	Increased	Increased	No change	Increased	Slightly decreased
G1m7	Increased	Increased	Increased	Increased	Increased	Slightly decreased
G1m8	Increased	Slightly decreased	No change	Increased	Slightly increased	Slightly decreased
G1m9	Increased	Decreased	Increased	Increased	Increased	No change
G1m15	Decreased	Decreased	Decreased	Decreased	Decreased	Decreased
G1m17	Decreased	Decreased	Decreased	Decreased	Decreased	Decreased
G1m18	Decreased	Decreased	No change	Decreased	Decreased	Decreased
G1m19	Decreased	Increased	Increased	Increased	Increased	Decreased
G1m25	Decreased	Decreased	Decreased	Decreased	Decreased	Decreased
G1m27	Decreased	Decreased	Decreased	Increased	Decreased	Decreased
G1m28	Decreased	Decreased	Decreased	Decreased	Decreased	Decreased
G1m29	Decreased	Decreased	Increased	Increased	Increased	Decreased
G1mAA	Decreased	N/A	Decreased	Decreased	Decreased	Decreased
G1mAG	Decreased	N/A	Decreased	Decreased	Decreased	Decreased

Table 3. FcγR Binding Activity of Human IgG2 Mutants As Relative to Wild-Type Human IgG2

5

	Change of Binding Activity Relative to Wild-Type Counterpart					
IgG2 Mutant	FcγRI	FcγRIIA(H131)	FcγRIIA(R131)	FcγRIIB	FcγRIIC	FcγRIII
G2m1	No change	Decreased	Decreased	Decreased	No change	No change
G2m2	Increased	Decreased	No change	Increased	No change	No change
G2m10	No change	Decreased	No change	Slightly decreased	No change	No change
G2m5	Slightly increased	Decreased	No change	Decreased	No change	No change
G2m7	No change	Slightly increased	Increased	Increased	Slightly increased	No change
G2m8	No change	Decreased	No change	No change	No change	No change
G2m9	No change	No change	Increased	Increased	Increased	No change
G2m-1	Increased	N/A	N/A	Decreased	N/A	No change
G2m-4	Increased	N/A	N/A	Decreased	N/A	No change
G2m15	No change	N/A	N/A	Decreased	N/A	No change
G2m17	No change	N/A	Increased	No change	Increased	No change
G2m18	Slightly increased	N/A	No change	Decreased	No change	No change
G2m19	No change	N/A	Increased	Increased	No change	No change
G2m20	No change	N/A	Increased	Decreased	No change	No change
G2m27	Slightly increased	N/A	N/A	Increased	N/A	No change

G2m28	No change	N/A	N/A	Increased	N/A	No change
-------	-----------	-----	-----	-----------	-----	-----------

Table 4. FcγR Binding Activity of Human IgG4 Mutants As Relative to Wild-Type Human IgG4

	Change of Binding Activity Relative to Wild-Type Counterpart					
IgG4 Mutant	FcγRI	FcγRIIA(H131)	FcγRIIA(R131)	FcγRIIB	FcγRIIC	FcγRIII
G4m1	Decreased	Increased	No change	Slightly decreased	Slightly increased	No change
G4m2	Decreased	No change	Decreased	Slightly decreased	No change	No change
G4m10	Decreased	No change	Decreased	Decreased	No change	No change
G4m20	Decreased	No change	Decreased	Decreased	No change	No change
G4m3	Decreased	No change	No change	Decreased	No change	No change
G4m7	No change	Increased	Increased	Increased	Increased	Slightly increased
G4m8	No change	No change	Increased	Slightly increased	No change	No change
G4m9	No change	No change	Increased	Increased	Increased	No change
G4m17	Decreased	No change	Increased	Increased	Increased	Increased
G4m18	Decreased	No change	No change	Decreased	No change	Increased
G4m19	Decreased	No change	No change	Decreased	No change	Slightly increased
G4m25	Decreased	No change	Increased	Slightly decreased	No change	Increased
G4m27	Decreased	No change	No change	Decreased	No change	Slightly increased
G4m28	Decreased	No change	Slightly increased	Increased	Increased	No change
G4m29	Decreased	No change	Increased	Increased	Increased	Slightly increased
G4m4	Decreased	No change	No change	Decreased	No change	No change
G4m-1	No change	N/A	N/A	No change	N/A	No change
G4m-2	No change	N/A	N/A	Decreased	N/A	No change
G4mPE	Decreased	N/A	N/A	Slightly decreased	No change	No change
G4m30	Decreased	N/A	No change	Decreased	No change	No change

5

An Fc variant as described herein can be designed following the guidance provided herein and produced via routine recombinant technology. Its binding affinity and specificity to

various Fc receptors can be determined via routine methods. See also Examples below.

II. Therapeutic Agents Capable of Cross-linking Immune Receptors and FcγRIIB

Also provided herein are therapeutic agents that can cross-link immune receptors and FcγRIIB receptor. Such therapeutic agents are expected to possess enhanced agonistic effects to activate immune cells such as T cells and NK cells in certain microenvironment such as cancer and tumor draining lymph nodes, so as to elicit or enhance immune responses against invading pathogens or disease cells.

(i) Binding Moieties

The therapeutic agents described herein contain at least two binding moieties. One binding moiety is capable of binding to an immune cell receptor expressed on cell surface, for example, receptors of the TNF superfamily. Examples include Fas receptor/FAS, TWEAK receptor/TNFRSF12A, 4-1BB/TNFRSF9/CD137, TACI/TNFRSF13B, BAFF R/TNFRSF13C, CD27/TNFRSF7, CD30/TNFRSF8, CD40/TNFRSF5, DR3/TNFRSF25, DR4/TNFRSF10A, DR5/TNFRSF10B, DR6/TNFRSF21, GITR/TNFRSF18, HVEM/TNFRSF14, Lymphotoxin-beta receptor/LTβR, OX40/TNFRSF4, TROY/TNFRSF19, RELT/TNFRSF19L, TNFRSF12A, TACI/TNFRSF13B, TL1A/TNFRSF15, TNFRSF17, TNFR1/TNFRSF1A, TNFRSF11B, RANK/TNFRSF11A, TR1/TNFRSF11B, NGFR, EDA2R and TNFR2/TNFRSF1B. In some instances, the binding moiety to the immune cell receptor is an antibody-binding fragment (*e.g.*, a Fab fragment or a single-chain antibody fragment).

The other binding moiety in the therapeutic agent is capable of binding to FcγRIIB. In some instances, this binding moiety has a selective binding activity to FcγRIIB relative to another Fc receptor. Such a binding moiety can be any of the Fc variants described herein, which selectively or specifically binds FcγRIIB.

In some embodiments, the therapeutic agent that cross-links an immune cell receptor and FcγRIIB can be an antibody specific to the immune cell receptor and comprises an Fc fragment that has selective binding activity to FcγRIIB, *e.g.*, those described herein. For example, the therapeutic agent can be an antibody that binds any of the TNF superfamily members known in the art or listed here (*e.g.*, CD137) and have a hIgG1, hIgG2, or hIgG4 Fc

variant as described herein. In some examples, the antibody is an agonistic antibody.

In some embodiments, the therapeutic agent is a bi-specific antibody comprising one antigen-binding moiety specific to the immune cell receptor as described herein and another antigen-binding moiety specific to an Fc γ receptor such as Fc γ RIIB.

5

(ii) *Preparation of Therapeutic Agents*

Antibodies capable of binding to both an immune cell receptor such as a TNF superfamily member and Fc γ RIIB as described herein can be made as follows. Antibodies binding to the immune cell receptor can be prepared by any method known in the art. See, 10 for example, Harlow and Lane, (1998) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York.

In some embodiments, antibodies specific to a target receptor (*e.g.*, CD137) or an extracellular domain thereof can be made by the conventional hybridoma technology. The full-length target receptor or a fragment thereof, optionally coupled to a carrier protein such 15 as KLH, can be used to immunize a host animal for generating antibodies binding to that antigen. The route and schedule of immunization of the host animal are generally in keeping with established and conventional techniques for antibody stimulation and production, as further described herein. General techniques for production of mouse, humanized, and human antibodies are known in the art and are described herein. It is contemplated that any 20 mammalian subject including humans or antibody producing cells therefrom can be manipulated to serve as the basis for production of mammalian, including human hybridoma cell lines. Typically, the host animal is inoculated intraperitoneally, intramuscularly, orally, subcutaneously, intraplantar, and/or intradermally with an amount of immunogen, including as described herein.

25 Hybridomas can be prepared from the lymphocytes and immortalized myeloma cells using the general somatic cell hybridization technique of Kohler, B. and Milstein, C. (1975) *Nature* 256:495-497 or as modified by Buck, D. W., et al., *In Vitro*, 18:377-381 (1982). Available myeloma lines, including but not limited to X63-Ag8.653 and those from the Salk Institute, Cell Distribution Center, San Diego, Calif., USA, may be used in the hybridization. 30 Generally, the technique involves fusing myeloma cells and lymphoid cells using a fusogen such as polyethylene glycol, or by electrical means well known to those skilled in the art.

After the fusion, the cells are separated from the fusion medium and grown in a selective growth medium, such as hypoxanthine-aminopterin-thymidine (HAT) medium, to eliminate unhybridized parent cells. Any of the media described herein, supplemented with or without serum, can be used for culturing hybridomas that secrete monoclonal antibodies. As another
5 alternative to the cell fusion technique, EBV immortalized B cells may be used to produce the anti-immune cell receptor monoclonal antibodies described herein. The hybridomas are expanded and subcloned, if desired, and supernatants are assayed for anti-immunogen activity by conventional immunoassay procedures (*e.g.*, radioimmunoassay, enzyme immunoassay, or fluorescence immunoassay).

10 Hybridomas that may be used as source of antibodies encompass all derivatives, progeny cells of the parent hybridomas that produce monoclonal antibodies capable of modulating the activity of the target immune cell receptor. Hybridomas that produce such antibodies may be grown *in vitro* or *in vivo* using known procedures. The monoclonal antibodies may be isolated from the culture media or body fluids, by conventional
15 immunoglobulin purification procedures such as ammonium sulfate precipitation, gel electrophoresis, dialysis, chromatography, and ultrafiltration, if desired. Undesired activity if present, can be removed, for example, by running the preparation over adsorbents made of the immunogen attached to a solid phase and eluting or releasing the desired antibodies off the immunogen. Immunization of a host animal with a target antigen or a fragment
20 containing the target amino acid sequence conjugated to a protein that is immunogenic in the species to be immunized, *e.g.*, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl₂,
25 or R₁N=C=NR, where R and R₁ are different alkyl groups, can yield a population of antibodies (*e.g.*, monoclonal antibodies).

If desired, an antibody (monoclonal or polyclonal) of interest (*e.g.*, produced by a hybridoma) may be sequenced and the polynucleotide sequence may then be cloned into a vector for expression or propagation. The sequence encoding the antibody of interest may be
30 maintained in vector in a host cell and the host cell can then be expanded and frozen for future use. In an alternative, the polynucleotide sequence may be used for genetic

manipulation to “humanize” the antibody or to improve the affinity (affinity maturation), or other characteristics of the antibody. For example, the constant region may be engineered to more resemble human constant regions to avoid immune response if the antibody is used in clinical trials and treatments in humans. It may be desirable to genetically manipulate the antibody sequence to obtain greater affinity to the target antigen and greater efficacy in inhibiting or activating the activity of the immune cell receptor. It will be apparent to one of skill in the art that one or more polynucleotide changes can be made to the antibody and still maintain its binding specificity to the target receptor.

In other embodiments, fully human antibodies can be obtained by using commercially available mice that have been engineered to express specific human immunoglobulin proteins. Transgenic animals that are designed to produce a more desirable (*e.g.*, fully human antibodies) or more robust immune response may also be used for generation of humanized or human antibodies. Examples of such technology are Xenomouse^{RTM} from Amgen, Inc. (Fremont, Calif.) and HuMAb-Mouse^{RTM} and TC MouseTM from Medarex, Inc. (Princeton, N.J.). In another alternative, antibodies may be made recombinantly by phage display or yeast technology. *See*, for example, U.S. Pat. Nos. 5,565,332; 5,580,717; 5,733,743; and 6,265,150; and Winter et al., (1994) *Annu. Rev. Immunol.* 12:433-455.

Alternatively, antibody library technology, such as the phage display technology (McCafferty et al., (1990) *Nature* 348:552-553), yeast display technology, or mammalian cell display technology, can be used to isolated antibodies such as human antibodies specific to a target immune receptor.

Antigen-binding fragments of an intact antibody (full-length antibody) can be prepared via routine methods. For example, F(ab')₂ fragments can be produced by pepsin digestion of an antibody molecule, and Fab fragments that can be generated by reducing the disulfide bridges of F(ab')₂ fragments.

Methods for constructing humanized antibodies are also well known in the art. *See*, *e.g.*, Queen et al., *Proc. Natl. Acad. Sci. USA*, 86:10029-10033 (1989). In one example, variable regions of VH and VL of a parent non-human antibody are subjected to three-dimensional molecular modeling analysis following methods known in the art. Next, framework amino acid residues predicted to be important for the formation of the correct CDR structures are identified using the same molecular modeling analysis. In parallel,

human VH and VL chains having amino acid sequences that are homologous to those of the parent non-human antibody are identified from any antibody gene database using the parent VH and VL sequences as search queries. Human VH and VL acceptor genes are then selected.

5 The CDR regions within the selected human acceptor genes can be replaced with the CDR regions from the parent non-human antibody or functional variants thereof. When necessary, residues within the framework regions of the parent chain that are predicted to be important in interacting with the CDR regions (see above description) can be used to substitute for the corresponding residues in the human acceptor genes.

10 Once an antibody capable of binding to a target immune cell receptor is obtained, its antigen-binding fragment can be conjugated to a suitable Fc fragment of a suitable IgG isoform, which can selectively binds FcγRIIB, for example, any of the Fc variants described herein *via* routine recombinant technology. In some instances, the antibody is first investigated for its agonistic effect to activate the immune cell receptor to which it binds.

15 Such an agonistic antibody can be selected for making therapeutic agent described herein to enhance the agonistic effects.

 The resultant antibody molecules can be produced *via* routine recombinant technology as exemplified below. Nucleic acids encoding the heavy and light chain of an antibody as described herein can be cloned into one expression vector, each nucleotide sequence being in operable linkage to a suitable promoter. In one example, each of the nucleotide sequences encoding the heavy chain and light chain is in operable linkage to a distinct promoter. Alternatively, the nucleotide sequences encoding the heavy chain and the light chain can be in operable linkage with a single promoter, such that both heavy and light chains are expressed from the same promoter. When necessary, an internal ribosomal entry site (IRES) can be inserted between the heavy chain and light chain encoding sequences.

 In some examples, the nucleotide sequences encoding the two chains of the antibody are cloned into two vectors, which can be introduced into the same or different cells. When the two chains are expressed in different cells, each of them can be isolated from the host cells expressing such and the isolated heavy chains and light chains can be mixed and incubated under suitable conditions allowing for the formation of the antibody.

Generally, a nucleic acid sequence encoding one or all chains of an antibody can be cloned into a suitable expression vector in operable linkage with a suitable promoter using methods known in the art. For example, the nucleotide sequence and vector can be contacted, under suitable conditions, with a restriction enzyme to create complementary ends on each molecule that can pair with each other and be joined together with a ligase. Alternatively, synthetic nucleic acid linkers can be ligated to the termini of a gene. These synthetic linkers contain nucleic acid sequences that correspond to a particular restriction site in the vector. The selection of expression vectors/promoter would depend on the type of host cells for use in producing the antibodies.

A variety of promoters can be used for expression of the antibodies described herein, including, but not limited to, cytomegalovirus (CMV) intermediate early promoter, a viral LTR such as the *Rous sarcoma* virus LTR, HIV-LTR, HTLV-1 LTR, the simian virus 40 (SV40) early promoter, *E. coli* lac UV5 promoter, and the herpes simplex tk virus promoter.

Regulatable promoters can also be used. Such regulatable promoters include those using the lac repressor from *E. coli* as a transcription modulator to regulate transcription from lac operator-bearing mammalian cell promoters [Brown, M. et al., *Cell*, 49:603-612 (1987)], those using the tetracycline repressor (tetR) [Gossen, M., and Bujard, H., *Proc. Natl. Acad. Sci. USA* 89:5547-5551 (1992); Yao, F. et al., *Human Gene Therapy*, 9:1939-1950 (1998); Shockelt, P., et al., *Proc. Natl. Acad. Sci. USA*, 92:6522-6526 (1995)]. Other systems include FK506 dimer, VP16 or p65 using astradiol, RU486, diphenol murislerone, or rapamycin. Inducible systems are available from Invitrogen, Clontech and Ariad.

Regulatable promoters that include a repressor with the operon can be used. In one embodiment, the lac repressor from *E. coli* can function as a transcriptional modulator to regulate transcription from lac operator-bearing mammalian cell promoters [M. Brown et al., *Cell*, 49:603-612 (1987)]; Gossen and Bujard (1992); [M. Gossen et al., *Natl. Acad. Sci. USA*, 89:5547-5551 (1992)] combined the tetracycline repressor (tetR) with the transcription activator (VP 16) to create a tetR-mammalian cell transcription activator fusion protein, tTa (tetR-VP 16), with the tetO-bearing minimal promoter derived from the human cytomegalovirus (hCMV) major immediate-early promoter to create a tetR-tet operator system to control gene expression in mammalian cells. In one embodiment, a tetracycline inducible switch is used. The tetracycline repressor (tetR) alone, rather than the tetR-

mammalian cell transcription factor fusion derivatives can function as potent trans-modulator to regulate gene expression in mammalian cells when the tetracycline operator is properly positioned downstream for the TATA element of the CMVIE promoter (Yao et al., Human Gene Therapy). One particular advantage of this tetracycline inducible switch is that it does
5 not require the use of a tetracycline repressor-mammalian cells transactivator or repressor fusion protein, which in some instances can be toxic to cells (Gossen et al., Natl. Acad. Sci. USA, 89:5547-5551 (1992); Shockett et al., Proc. Natl. Acad. Sci. USA, 92:6522-6526 (1995)), to achieve its regulatable effects.

Additionally, the vector can contain, for example, some or all of the following: a
10 selectable marker gene, such as the neomycin gene for selection of stable or transient transfectants in mammalian cells; enhancer/promoter sequences from the immediate early gene of human CMV for high levels of transcription; transcription termination and RNA processing signals from SV40 for mRNA stability; SV40 polyoma origins of replication and ColE1 for proper episomal replication; internal ribosome binding sites (IRESes), versatile
15 multiple cloning sites; and T7 and SP6 RNA promoters for in vitro transcription of sense and antisense RNA. Suitable vectors and methods for producing vectors containing transgenes are well known and available in the art.

Examples of polyadenylation signals useful to practice the methods described herein include, but are not limited to, human collagen I polyadenylation signal, human collagen II
20 polyadenylation signal, and SV40 polyadenylation signal.

One or more vectors (*e.g.*, expression vectors) comprising nucleic acids encoding any of the antibodies may be introduced into suitable host cells for producing the antibodies. The host cells can be cultured under suitable conditions for expression of the antibody or any polypeptide chain thereof. Such antibodies or polypeptide chains thereof can be recovered by
25 the cultured cells (*e.g.*, from the cells or the culture supernatant) via a conventional method, *e.g.*, affinity purification. If necessary, polypeptide chains of the antibody can be incubated under suitable conditions for a suitable period of time allowing for production of the antibody.

In some embodiments, methods for preparing an antibody described herein involve a
30 recombinant expression vector that encodes both the heavy chain and the light chain of an antibody as described herein. The recombinant expression vector can be introduced into a

suitable host cell (*e.g.*, a dhfr- CHO cell) by a conventional method, *e.g.*, calcium phosphate-mediated transfection. Positive transformant host cells can be selected and cultured under suitable conditions allowing for the expression of the two polypeptide chains that form the antibody, which can be recovered from the cells or from the culture medium. When
5 necessary, the two chains recovered from the host cells can be incubated under suitable conditions allowing for the formation of the antibody.

In one example, two recombinant expression vectors are provided, one encoding the heavy chain of the anti-immune cell receptor antibody and the other encoding the light chain of the same antibody. Both of the two recombinant expression vectors can be introduced into
10 a suitable host cell (*e.g.*, dhfr- CHO cell) by a conventional method, *e.g.*, calcium phosphate-mediated transfection. Alternatively, each of the expression vectors can be introduced into a suitable host cells. Positive transformants can be selected and cultured under suitable conditions allowing for the expression of the polypeptide chains of the antibody. When the two expression vectors are introduced into the same host cells, the antibody produced therein
15 can be recovered from the host cells or from the culture medium. If necessary, the polypeptide chains can be recovered from the host cells or from the culture medium and then incubated under suitable conditions allowing for formation of the antibody. When the two expression vectors are introduced into different host cells, each of them can be recovered from the corresponding host cells or from the corresponding culture media. The two
20 polypeptide chains can then be incubated under suitable conditions for formation of the antibody.

Standard molecular biology techniques are used to prepare the recombinant expression vector, transfect the host cells, select for transformants, culture the host cells and recovery of the antibodies from the culture medium. For example, some antibodies can be
25 isolated by affinity chromatography with a Protein A or Protein G coupled matrix.

The bioactivity of the antibodies described herein can be verified using assays known in the art or described herein.

(iii) *Pharmaceutical Compositions*

30 The antibodies as described herein can be mixed with a pharmaceutically acceptable carrier (excipient) to form a pharmaceutical composition for use in treating a target disease.

“Acceptable” means that the carrier must be compatible with the active ingredient of the composition (and preferably, capable of stabilizing the active ingredient) and not deleterious to the subject to be treated. Pharmaceutically acceptable excipients (carriers) including buffers, which are well known in the art. See, e.g., Remington: The Science and Practice of Pharmacy 20th Ed. (2000) Lippincott Williams and Wilkins, Ed. K. E. Hoover.

The pharmaceutical compositions to be used in the present methods can comprise pharmaceutically acceptable carriers, excipients, or stabilizers in the form of lyophilized formulations or aqueous solutions. (Remington: The Science and Practice of Pharmacy 20th Ed. (2000) Lippincott Williams and Wilkins, Ed. K. E. Hoover). Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations used, and may comprise buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEENTM, PLURONICSTM or polyethylene glycol (PEG).

In some examples, the pharmaceutical composition described herein comprises liposomes containing the antibodies (or the encoding nucleic acids) which can be prepared by methods known in the art, such as described in Epstein, et al., Proc. Natl. Acad. Sci. USA 82:3688 (1985); Hwang, et al., Proc. Natl. Acad. Sci. USA 77:4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556. Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter.

The antibodies, or the encoding nucleic acid(s), may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are known in the art, see, e.g., Remington, The Science and Practice of Pharmacy 20th Ed. Mack Publishing (2000).

In other examples, the pharmaceutical composition described herein can be formulated in sustained-release format. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinyl alcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ -ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), sucrose acetate isobutyrate, and poly-D-(-)-3-hydroxybutyric acid.

The pharmaceutical compositions to be used for in vivo administration must be sterile. This is readily accomplished by, for example, filtration through sterile filtration membranes. Therapeutic antibody compositions are generally placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

The pharmaceutical compositions described herein can be in unit dosage forms such as tablets, pills, capsules, powders, granules, solutions or suspensions, or suppositories, for oral, parenteral or rectal administration, or administration by inhalation or insufflation.

For preparing solid compositions such as tablets, the principal active ingredient can be mixed with a pharmaceutical carrier, e.g., conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g., water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a non-toxic pharmaceutically acceptable salt thereof. When referring to these preformulation

compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing
5 from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer that serves to resist
10 disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

Suitable surface-active agents include, in particular, non-ionic agents, such as
15 polyoxyethylenesorbitans (*e.g.*, TweenTM 20, 40, 60, 80 or 85) and other sorbitans (*e.g.*, SpanTM 20, 40, 60, 80 or 85). Compositions with a surface-active agent will conveniently comprise between 0.05 and 5% surface-active agent, and can be between 0.1 and 2.5%. It will be appreciated that other ingredients may be added, for example mannitol or other pharmaceutically acceptable vehicles, if necessary.

Suitable emulsions may be prepared using commercially available fat emulsions, such
20 as IntralipidTM, LiposynTM, InfonutrolTM, LipofundinTM and LipiphysanTM. The active ingredient may be either dissolved in a pre-mixed emulsion composition or alternatively it may be dissolved in an oil (*e.g.*, soybean oil, safflower oil, cottonseed oil, sesame oil, corn oil or almond oil) and an emulsion formed upon mixing with a phospholipid (*e.g.* egg
25 phospholipids, soybean phospholipids or soybean lecithin) and water. It will be appreciated that other ingredients may be added, for example glycerol or glucose, to adjust the tonicity of the emulsion. Suitable emulsions will typically contain up to 20% oil, for example, between 5 and 20%. The fat emulsion can comprise fat droplets between 0.1 and 1.0 μm , particularly 0.1 and 0.5 μm , and have a pH in the range of 5.5 to 8.0.

The emulsion compositions can be those prepared by mixing an antibody with Intralipid™ or the components thereof (soybean oil, egg phospholipids, glycerol and water).

Pharmaceutical compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as set out above. In some embodiments, the compositions are administered by the oral or nasal respiratory route for local or systemic effect.

Compositions in preferably sterile pharmaceutically acceptable solvents may be nebulized by use of gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device may be attached to a face mask, tent or intermittent positive pressure breathing machine. Solution, suspension or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

III. Therapeutic Applications

Any of the therapeutic agents (e.g., antibodies) capable of cross-linking an immune cell receptor and FcγRIIB described herein are useful for enhancing immune responses against invading pathogens and/or diseased cells such as cancer cells.

To practice the method disclosed herein, an effective amount of the pharmaceutical composition described herein can be administered to a subject (e.g., a human) in need of the treatment via a suitable route, such as intravenous administration, e.g., as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, inhalation or topical routes. Commercially available nebulizers for liquid formulations, including jet nebulizers and ultrasonic nebulizers are useful for administration. Liquid formulations can be directly nebulized and lyophilized powder can be nebulized after reconstitution.

Alternatively, the antibodies as described herein can be aerosolized using a fluorocarbon formulation and a metered dose inhaler, or inhaled as a lyophilized and milled powder.

The subject to be treated by the methods described herein can be a mammal, more preferably a human. Mammals include, but are not limited to, farm animals, sport animals,

pets, primates, horses, dogs, cats, mice and rats. In some instances, the subject is a human patient having or at risk for a cell-mediated disease or disorder, such as cancer including but not limited to lung cancer, stomach cancer, liver cancer, breast cancer, skin cancer, pancreatic cancer, brain cancer, prostate cancer, bladder cancer, or colorectal cancer.

5 As used herein, “an effective amount” refers to the amount of each active agent required to confer therapeutic effect on the subject, either alone or in combination with one or more other active agents. In some embodiments, the therapeutic effect is modulating (*e.g.*, activating) the target immune receptor, thereby triggering or enhancing immune responses mediated by the receptor. Determination of whether an amount of the antibody achieved the
10 therapeutic effect would be evident to one of skill in the art. Effective amounts vary, as recognized by those skilled in the art, depending on the particular condition being treated, the severity of the condition, the individual patient parameters including age, physical condition, size, gender and weight, the duration of the treatment, the nature of concurrent therapy (if any), the specific route of administration and like factors within the knowledge and expertise
15 of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose of the individual components or combinations thereof be used, that is, the highest safe dose according to sound medical judgment.

Empirical considerations, such as the half-life, generally will contribute to the
20 determination of the dosage. For example, antibodies that are compatible with the human immune system, such as humanized antibodies or fully human antibodies, may be used to prolong half-life of the antibody and to prevent the antibody being attacked by the host's immune system. Frequency of administration may be determined and adjusted over the course of therapy, and is generally, but not necessarily, based on treatment and/or suppression
25 and/or amelioration and/or delay of a target disease/disorder. Alternatively, sustained continuous release formulations of an antibody may be appropriate. Various formulations and devices for achieving sustained release are known in the art.

In one example, dosages for an antibody as described herein may be determined empirically in individuals who have been given one or more administration(s) of the
30 antibody. Individuals are given incremental dosages of the antagonist. To assess efficacy of the antagonist, an indicator of the disease/disorder can be followed.

Generally, for administration of any of the therapeutic agents such as antibodies described herein, an initial candidate dosage can be about 2 mg/kg. For the purpose of the present disclosure, a typical daily dosage might range from about any of 0.1 µg/kg to 3 µg/kg to 30 µg/kg to 300 µg/kg to 3 mg/kg, to 30 mg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of symptoms occurs or until sufficient therapeutic levels are achieved to alleviate a target disease or disorder, or a symptom thereof. An exemplary dosing regimen comprises administering an initial dose of about 2 mg/kg, followed by a weekly maintenance dose of about 1 mg/kg of the antibody, or followed by a maintenance dose of about 1 mg/kg every other week. However, other dosage regimens may be useful, depending on the pattern of pharmacokinetic decay that the practitioner wishes to achieve. For example, dosing from one-four times a week is contemplated. In some embodiments, dosing ranging from about 3 µg/mg to about 2 mg/kg (such as about 3 µg/mg, about 10 µg/mg, about 30 µg/mg, about 100 µg/mg, about 300 µg/mg, about 1 mg/kg, and about 2 mg/kg) may be used. In some embodiments, dosing frequency is once every week, every 2 weeks, every 4 weeks, every 5 weeks, every 6 weeks, every 7 weeks, every 8 weeks, every 9 weeks, or every 10 weeks; or once every month, every 2 months, or every 3 months, or longer. The progress of this therapy is easily monitored by conventional techniques and assays. The dosing regimen (including the antibody used) can vary over time.

In some embodiments, for an adult patient of normal weight, doses ranging from about 0.3 to 5.00 mg/kg may be administered. In some examples, the dosage of the therapeutic agents such as antibodies described herein can be 10 mg/kg. The particular dosage regimen, *i.e.*, dose, timing and repetition, will depend on the particular individual and that individual's medical history, as well as the properties of the individual agents (such as the half-life of the agent, and other considerations well known in the art).

For the purpose of the present disclosure, the appropriate dosage of an antibody as described herein will depend on the specific antibody, antibodies, and/or non-antibody peptide (or compositions thereof) employed, the type and severity of the disease/disorder, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antagonist, and the discretion of the

attending physician. Typically the clinician will administer an antibody, until a dosage is reached that achieves the desired result. In some embodiments, the desired result is a decrease in thrombosis. Methods of determining whether a dosage resulted in the desired result would be evident to one of skill in the art. Administration of one or more antibodies
5 can be continuous or intermittent, depending, for example, upon the recipient's physiological condition, whether the purpose of the administration is therapeutic or prophylactic, and other factors known to skilled practitioners. The administration of an antibody may be essentially continuous over a preselected period of time or may be in a series of spaced dose, e.g., either before, during, or after developing a target disease or disorder.

10 As used herein, the term “treating” refers to the application or administration of a composition including one or more active agents to a subject, who has a target disease or disorder, a symptom of the disease/disorder, or a predisposition toward the disease/disorder, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disorder, the symptom of the disease, or the predisposition toward the disease or disorder.

15 Alleviating a target disease/disorder includes delaying the development or progression of the disease, or reducing disease severity. Alleviating the disease does not necessarily require curative results. As used therein, “delaying” the development of a target disease or disorder means to defer, hinder, slow, retard, stabilize, and/or postpone progression of the disease. This delay can be of varying lengths of time, depending on the history of the disease
20 and/or individuals being treated. A method that “delays” or alleviates the development of a disease, or delays the onset of the disease, is a method that reduces probability of developing one or more symptoms of the disease in a given time frame and/or reduces extent of the symptoms in a given time frame, when compared to not using the method. Such comparisons are typically based on clinical studies, using a number of subjects sufficient to give a
25 statistically significant result.

“Development” or “progression” of a disease means initial manifestations and/or ensuing progression of the disease. Development of the disease can be detectable and assessed using standard clinical techniques as well known in the art. However, development also refers to progression that may be undetectable. For purpose of this disclosure,
30 development or progression refers to the biological course of the symptoms. “Development”

includes occurrence, recurrence, and onset. As used herein “onset” or “occurrence” of a target disease or disorder includes initial onset and/or recurrence.

In some embodiments, the antibodies described herein are administered to a subject in need of the treatment at an amount sufficient to activate the activity of the target receptor by
5 at least 20% (*e.g.*, 30%, 40%, 50%, 60%, 70%, 80%, 90% or greater) *in vivo*.

Conventional methods, known to those of ordinary skill in the art of medicine, can be used to administer the pharmaceutical composition to the subject, depending upon the type of disease to be treated or the site of the disease. This composition can also be administered via other conventional routes, *e.g.*, administered orally, parenterally, by inhalation spray,
10 topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term “parenteral” as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional, and intracranial injection or infusion techniques. In addition, it can be administered to the subject via injectable depot routes of administration such as using 1-, 3-, or 6-month depot
15 injectable or biodegradable materials and methods. In some examples, the pharmaceutical composition is administered intraocularly or intravitreally.

Injectable compositions may contain various carriers such as vegetable oils, dimethylacetamide, dimethylformamide, ethyl lactate, ethyl carbonate, isopropyl myristate, ethanol, and polyols (glycerol, propylene glycol, liquid polyethylene glycol, and the like).

20 For intravenous injection, water soluble antibodies can be administered by the drip method, whereby a pharmaceutical formulation containing the antibody and a physiologically acceptable excipient is infused. Physiologically acceptable excipients may include, for example, 5% dextrose, 0.9% saline, Ringer’s solution or other suitable excipients.

Intramuscular preparations, *e.g.*, a sterile formulation of a suitable soluble salt form of the
25 antibody, can be dissolved and administered in a pharmaceutical excipient such as Water-for-Injection, 0.9% saline, or 5% glucose solution.

The particular dosage regimen, *i.e.*, dose, timing and repetition, used in the method described herein will depend on the particular subject and that subject's medical history.

In some embodiments, more than one antibody, or a combination of an antibody and
30 another suitable therapeutic agent, may be administered to a subject in need of the treatment.

The antibody can also be used in conjunction with other agents that serve to enhance and/or complement the effectiveness of the agents.

Treatment efficacy for a target disease/disorder can be assessed by methods well-known in the art.

5 The therapeutic agent described herein may be utilized in conjunction with other types of therapy for the target disease such as cancer. Additional anti-cancer therapy includes chemotherapy, surgery, radiation, gene therapy, and so forth. When a second therapeutic agent is used, such an agent can be administered simultaneously or sequentially (in any order) with the therapeutic agent described herein that cross-links an immune cell receptor and
10 Fc γ RIIB.

When co-administered with an additional therapeutic agent, suitable therapeutically effective dosages for each agent may be lowered due to the additive action or synergy.

The treatments of the disclosure can be combined with other immunomodulatory treatments such as, *e.g.*, therapeutic vaccines (including but not limited to GVAX, DC-based
15 vaccines, *etc.*), or checkpoint inhibitors (including but not limited to agents that block CTLA4, PD1, LAG3, TIM3, *etc.*). Alternatively, the treatment of the present disclosure can be combined with a chemotherapeutic agent, for example, pyrimidine analogs (5-fluorouracil, floxuridine, capecitabine, gemcitabine and cytarabine), purine analogs, folate antagonists and related inhibitors (mercaptopurine, thioguanine, pentostatin and 2-chlorodeoxyadenosine
20 (cladribine)); antiproliferative/antimitotic agents including natural products such as vinca alkaloids (vinblastine, vincristine, and vinorelbine), microtubule disruptors such as taxane (paclitaxel, docetaxel), vincristin, vinblastin, nocodazole, epothilones and navelbine, epidipodophyllotoxins (etoposide, teniposide), DNA damaging agents (actinomycin, amsacrine, anthracyclines, bleomycin, busulfan, camptothecin, carboplatin, chlorambucil,
25 cisplatin, cyclophosphamide, cytoxan, dactinomycin, daunorubicin, doxorubicin, epirubicin, hexamethylnelamineoxaliplatin, iphosphamide, melphalan, merchlorhtamine, mitomycin, mitoxantrone, nitrosourea, plicamycin, procarbazine, taxol, taxotere, teniposide, triethylenethiophosphoramidate and etoposide (VP16)); antibiotics such as dactinomycin (actinomycin D), daunorubicin, doxorubicin (adriamycin), idarubicin, anthracyclines,
30 mitoxantrone, bleomycins, plicamycin (mithramycin) and mitomycin; enzymes (L-

asparaginase which systemically metabolizes L-asparagine and deprives cells which do not have the capacity to synthesize their own asparagine); antiplatelet agents; antiproliferative/antimitotic alkylating agents such as nitrogen mustards (mechlorethamine, cyclophosphamide and analogs, melphalan, chlorambucil), ethylenimines and methylmelamines (hexamethylmelamine and thiotepa), alkyl sulfonates-busulfan, nitrosoureas (carmustine (BCNU) and analogs, streptozocin), trazenes-dacarbazine (DTIC); antiproliferative/antimitotic antimetabolites such as folic acid analogs (methotrexate); platinum coordination complexes (cisplatin, carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide; hormones, hormone analogs (estrogen, tamoxifen, goserelin, bicalutamide, nilutamide) and aromatase inhibitors (letrozole, anastrozole); anticoagulants (heparin, synthetic heparin salts and other inhibitors of thrombin); fibrinolytic agents (such as tissue plasminogen activator, streptokinase and urokinase), aspirin, dipyridamole, ticlopidine, clopidogrel, abciximab; antimigratory agents; antisecretory agents (breveldin); immunosuppressives (cyclosporine, tacrolimus (FK-506), sirolimus (rapamycin), azathioprine, mycophenolate mofetil); anti-angiogenic compounds (*e.g.*, TNP-470, genistein, bevacizumab) and growth factor inhibitors (*e.g.*, fibroblast growth factor (FGF) inhibitors); angiotensin receptor blocker; nitric oxide donors; anti-sense oligonucleotides; antibodies (trastuzumab); cell cycle inhibitors and differentiation inducers (tretinoin); mTOR inhibitors, topoisomerase inhibitors (doxorubicin (adriamycin), amsacrine, camptothecin, daunorubicin, dactinomycin, eniposide, epirubicin, etoposide, idarubicin and mitoxantrone, topotecan, irinotecan), corticosteroids (cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisone, and prednisolone); growth factor signal transduction kinase inhibitors; mitochondrial dysfunction inducers and caspase activators; and chromatin disruptors.

For examples of additional useful agents see also Physician's Desk Reference, 59^{sup}.th edition, (2005), Thomson P D R, Montvale N.J.; Gennaro et al., Eds. Remington's The Science and Practice of Pharmacy 20^{sup}.th edition, (2000), Lippincott Williams and Wilkins, Baltimore Md.; Braunwald et al., Eds. Harrison's Principles of Internal Medicine, 15^{sup}.th edition, (2001), McGraw Hill, NY; Berkow et al., Eds. The Merck Manual of Diagnosis and Therapy, (1992), Merck Research Laboratories, Rahway N.J.

IV. Kits

The present disclosure also provides kits for use in enhancing the desired immune responses using any of the therapeutic agents described herein, for example, anti-TNF antibodies containing an Fc variant described herein that selectively binds Fc γ RIIB.

In some embodiments, the kit can comprise instructions for use in accordance with any of the methods described herein. The included instructions can comprise a description of administration of the therapeutic agent to treat, delay the onset, or alleviate a target disease as those described herein. The kit may further comprise a description of selecting an individual suitable for treatment based on identifying whether that individual has the target disease. In still other embodiments, the instructions comprise a description of administering a therapeutic agent such as an antibody to an individual at risk of the target disease.

The instructions relating to the use of the therapeutic agent generally include information as to dosage, dosing schedule, and route of administration for the intended treatment. The containers may be unit doses, bulk packages (*e.g.*, multi-dose packages) or sub-unit doses. Instructions supplied in the kits of the invention are typically written instructions on a label or package insert (*e.g.*, a paper sheet included in the kit), but machine-readable instructions (*e.g.*, instructions carried on a magnetic or optical storage disk) are also acceptable.

The label or package insert indicates that the composition is used for treating, delaying the onset and/or alleviating a target disease or disorder such as cancer. Instructions may be provided for practicing any of the methods described herein.

The kits of this invention are in suitable packaging. Suitable packaging includes, but is not limited to, vials, bottles, jars, flexible packaging (*e.g.*, sealed Mylar or plastic bags), and the like. Also contemplated are packages for use in combination with a specific device, such as an inhaler, nasal administration device (*e.g.*, an atomizer) or an infusion device such as a minipump. A kit may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The container may also have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is the therapeutic agent as those described herein.

Kits may optionally provide additional components such as buffers and interpretive information. Normally, the kit comprises a container and a label or package insert(s) on or associated with the container. In some embodiments, the invention provides articles of manufacture comprising contents of the kits described above.

General Techniques

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, *Molecular Cloning: A Laboratory Manual*, second edition (Sambrook, et al., 1989) Cold Spring Harbor Press; *Oligonucleotide Synthesis* (M. J. Gait, ed., 1984); *Methods in Molecular Biology*, Humana Press; *Cell Biology: A Laboratory Notebook* (J. E. Cellis, ed., 1998) Academic Press; *Animal Cell Culture* (R. I. Freshney, ed., 1987); *Introduction to Cell and Tissue Culture* (J. P. Mather and P. E. Roberts, 1998) Plenum Press; *Cell and Tissue Culture: Laboratory Procedures* (A. Doyle, J. B. Griffiths, and D. G. Newell, eds., 1993-8) J. Wiley and Sons; *Methods in Enzymology* (Academic Press, Inc.); *Handbook of Experimental Immunology* (D. M. Weir and C. C. Blackwell, eds.); *Gene Transfer Vectors for Mammalian Cells* (J. M. Miller and M. P. Calos, eds., 1987); *Current Protocols in Molecular Biology* (F. M. Ausubel, et al., eds., 1987); *PCR: The Polymerase Chain Reaction*, (Mullis, et al., eds., 1994); *Current Protocols in Immunology* (J. E. Coligan et al., eds., 1991); *Short Protocols in Molecular Biology* (Wiley and Sons, 1999); *Immunobiology* (C. A. Janeway and P. Travers, 1997); *Antibodies* (P. Finch, 1997); *Antibodies: a practical approach* (D. Catty., ed., IRL Press, 1988-1989); *Monoclonal antibodies: a practical approach* (P. Shepherd and C. Dean, eds., Oxford University Press, 2000); *Using antibodies: a laboratory manual* (E. Harlow and D. Lane (Cold Spring Harbor Laboratory Press, 1999); *The Antibodies* (M. Zanetti and J. D. Capra, eds., Harwood Academic Publishers, 1995).

Without further elaboration, it is believed that one skilled in the art can, based on the above description, utilize the present invention to its fullest extent. The following specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever. All publications cited herein are

incorporated by reference for the purposes or subject matter referenced herein.

Example 1. Engineering human IgG1, IgG2 and IgG4 for selective FcγR2B/CD32B binding

To verify the impact of mutations in the hinge domain of human IgG1, IgG2, and IgG4 on binding activity to FcγR2b/CD32B, the human Fc variants G1m1, G1m2, G2m1, G2m2, G2m10, G4m1, G4m2, G4m10, and G4m20 disclosed herein were linked to the VH fragment of an anti-CD137 antibody and the corresponding IgG1, G2, and G4 CH1 fragments to produce a full-length IgG heavy chain. These mutant IgG heavy chains were cloned, co-expressed with the light chain of the anti-CD137 antibody, and purified using standard molecular biology and antibody protocols. The amino acid sequences of the VH-CH1 fragment (VH domain italicized) and the light chain are provided below:

VH-CH1 (IgG1):

EVQLVQSGAEVKKPGESLRISCKGSGYSFSTYWISWVRQMPGKGLEWMGKIYPGDSYTNYSFSFQGGQVTISADKS
ISTAYLQWSSLKASDTAMYYCARGYGIFDYWGQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP
EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTK (SEQ ID NO: 60)

VH-CH1 (IgG2):

EVQLVQSGAEVKKPGESLRISCKGSGYSFSTYWISWVRQMPGKGLEWMGKIYPGDSYTNYSFSFQGGQVTISADKS
ISTAYLQWSSLKASDTAMYYCARGYGIFDYWGQGTLVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFP
EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSNFGTQTYTCNVDPHKPSNTK (SEQ ID NO: 61)

VH-CH1 (IgG4):

EVQLVQSGAEVKKPGESLRISCKGSGYSFSTYWISWVRQMPGKGLEWMGKIYPGDSYTNYSFSFQGGQVTISADKS
ISTAYLQWSSLKASDTAMYYCARGYGIFDYWGQGTLVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFP
EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYTCNVDPHKPSNTK (SEQ ID NO: 62)

Light Chain:

SYELTQPPSVSVSPGQTASITCSGDNIGDQYAHWYQQKPGQSPVLVIYQDKNRPSGIPERFSGSNSGNTATLTIS
GTQAMDEADYYCATYTGFGLAVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAW
KADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSQRSYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO: 63)

These IgG mutants were tested for binding to a panel of human Fc receptors (FcRs), including FcRn, using the Octet Red96 System (ForteBio, Model# Red96). Human FcR proteins were purchased commercially (FcRn, MednaBio, E1032; FcRI, MednaBio, E1031; FcRIIA, MednaBio, E1033; FcRIIB/C, MednaBio, E1034; FcRIIIA-F158, MednaBio, E1036;

FcRIIIA-V158, MednaBio, E1035; FcRIIIB, MednaBio, E1037). All Fcγ receptor assays were performed at pH 7.2; all FcRn assays were performed at pH 6. The Fcγ receptors were loaded onto anti-penta-HIS1K (ForteBio, Cat#18-5122) biosensors at a concentration of 20 μg/mL. The loaded biosensors was then be dipped into an 8-point 1:3 dilution series of the test IgG molecules (including controls and the mutants described herein) in PBS with 0.1% BSA, 0.02% Tween-20 (pH 7.2) at a starting concentration of 300 nM. Kinetic analysis was performed using a 1:1 binding model and global fitting. The results are shown in Tables 5 and 6 below.

10

Table 5. FcγR Binding Activity of IgG1 and IgG2 Mutants to Various Fc Receptors

		IgG1	IgG2	IgG4	G1m1	G1m2	G2m1	G2m2	G2m10
FcγRI	KD (M)	5.6E-9	No	4.8E-09	2.0E-07	No	No	2.0E-09	No
	kon(1/Ms)	2.2E+05	No	3.6E+05	6.5E+05	No	No	4.4E+05	No
	kdis(1/s)	1.2E-03	No	1.7E-03	1.3E-01	No	No	9.1E-04	No
FcγRIIA	KD (M)	5.6E-08	8.9E-08	4.0E-07	4.4E-07	No	No	6.8E-08	1.70E-05
	kon(1/Ms)	5.7E+05	4.4E+05	1.4E+05	2.5E+05	No	No	8.4E+05	2.10E+04
	kdis(1/s)	3.2E-02	3.9E-02	5.5E-02	1.1E-01	No	No	5.7E-02	3.50E-01
FcγRIIB	KD (M)	1.3E-07	No	4.0E-07	No	No	No	1.6E-07	No
	kon(1/Ms)	1.2E+05	No	1.4E+05	No	No	No	6.2E+05	No
	kdis(1/s)	1.6E-02	No	5.0E-02	No	No	No	9.9E-02	No
FcγRIIIA(F)	KD (M)	1.2E-07	No	No	3.0E-07	No	No	2.5E-07	No
	kon(1/Ms)	1.7E+05	No	No	3.5E+05	No	No	2.8E+05	No
	kdis(1/s)	2.0E-02	No	No	1.0E-01	No	No	7.0E-02	No
FcγRIIIA(V)	KD (M)	6.0E-08	No	5.5E-07	No	No	No	1.6E-07	No
	kon(1/Ms)	4.5E+05	No	1.7E+05	No	No	No	2.1E+05	No
	kdis(1/s)	2.7E-02	No	9.2E-02	No	No	No	3.3E-02	No
FcγRIIIB	KD (M)	1.3E-07	No	No	No	No	Low	No	No
	kon(1/Ms)	2.1E+05	No	No	No	No	Low	No	No
	kdis(1/s)	2.7E-02	No	No	No	No	Low	No	No
FcRn	KD (M)	1.6E-08	1.0E-08	1.8E-08	1.1E-08	1.3E-08	5.5E-09	1.0E-08	2.10E-08
	kon(1/Ms)	4.8E+05	9.9E+05	6.4E+05	7.0E+05	7.1E+05	1.7E+05	9.0E+05	4.20E+05
	kdis(1/s)	8.0E-03	9.9E-03	1.2E-02	7.8E-03	9.3E-03	9.6E-04	9.1E-03	8.80E-03

15

Table 6. FcγR Binding Activity of IgG4 Mutants

		IgG1	IgG2	IgG4	G4m1	G4m2	G4m10	G4m20
FcγRI	KD (M)	5.6E-9	No	4.8E-09	1.3E-07	No	No	No
	kon(1/Ms)	2.2E+05	No	3.6E+05	7.4E+05	No	No	No
	kdis(1/s)	1.2E-03	No	1.7E-03	9.7E-02	No	No	No
FcγRIIA	KD (M)	5.6E-08	8.9E-08	4.0E-07	4.5E-07	No	8.30E-08	No
	kon(1/Ms)	5.7E+05	4.4E+05	1.4E+05	2.4E+05	No	4.50E+05	No
	kdis(1/s)	3.2E-02	3.9E-02	5.5E-02	1.1E-01	No	3.70E-02	No
FcγRIIB	KD (M)	1.3E-07	No	4.0E-07	2.5E-07	1.5E-07	No	No
	kon(1/Ms)	1.2E+05	No	1.4E+05	4.2E+05	3.1E+05	No	No
	kdis(1/s)	1.6E-02	No	5.0E-02	1.1E-01	4.6E-02	No	No
FcγRIIA(F)	KD (M)	1.2E-07	No	No	No	No	No	No
	kon(1/Ms)	1.7E+05	No	No	No	No	No	No
	kdis(1/s)	2.0E-02	No	No	No	No	No	No
FcγRIIA(V)	KD (M)	6.0E-08	No	5.5E-07	No	No	No	No
	kon(1/Ms)	4.5E+05	No	1.7E+05	No	No	No	No
	kdis(1/s)	2.7E-02	No	9.2E-02	No	No	No	No
FcγRIIB	KD (M)	1.3E-07	No	No	No	No	No	No
	kon(1/Ms)	2.1E+05	No	No	No	No	No	No
	kdis(1/s)	2.7E-02	No	No	No	No	No	No
FcRn	KD (M)	1.6E-08	1.0E-08	1.8E-08	1.9E-08	9.8E-09	1.10E-08	1.9E-08
	kon(1/Ms)	4.8E+05	9.9E+05	6.4E+05	6.9E+05	1.4E+06	5.10E+05	5.1E+05
	kdis(1/s)	8.0E-03	9.9E-03	1.2E-02	1.3E-02	1.3E-02	5.50E-03	9.8E-03

Certain IgG mutants tested in this study, for example, G4m2, showed selective binding activity to FcγRIIB (CD32B), and also maintained binding activity to FcRn, which is important for the half-life of the IgG molecule *in vivo*. Others, such as G1m2, G2m1 or G4m20 lost binding to all FcγRs.

Example 2. Additional IgG1, IgG2, and IgG4 Mutants and Binding Activities to Various Fc Receptors

The human Fc variants G1m5, G1m7, G1m8, G1m9, G2m5, G2m7, G2m8, G2m9, G4m5, G4m7, G4m8, and G4m9 disclosed herein were linked to the VH fragment of an anti-CD137 antibody and the corresponding IgG1, G2, and G4 CH1 fragments to produce a full-length IgG heavy chain. See Example 1 above. These mutant IgG heavy chains were cloned, co-expressed with the light chain of the anti-CD137 antibody (see Example 1 above), and

purified using standard molecular biology and antibody protocols. The results are shown in Tables 7-9 below.

Table 7. FcR Binding Activity of IgG1 Mutants

			IgG1	IgG2	IgG4	G1m5	G1m7	G1m8	G1m9	G1mAA
	KD (M)		5.6E-9	No	4.8E-09	1.6E-09	2.5E-09	2.5E-09	3.1E-09	2.35E-07
FcγRI	kon(1/Ms)		2.2E+05	No	3.6E+05	3.0E+05	6.3E+05	6.3E+05	5.1E+05	1.44E+05
	kdis(1/s)		1.2E-03	No	1.7E-03	4.7E-04	1.6E-03	1.6E-03	1.6E-03	3.37E-02
	KD (M)		5.6E-08	8.9E-08	4.0E-07	6.7E-07	2.9E-07	2.7E-07	6.8E-07	8.00E-07
FcγRIIA	kon(1/Ms)		5.7E+05	4.4E+05	1.4E+05	3.4E+05	3.8E+05	6.1E+05	1.8E+05	3.10E+05
	kdis(1/s)		3.2E-02	3.9E-02	5.5E-02	2.3E-01	1.1E-01	1.7E-01	1.3E-01	2.47E-01
	KD (M)		1.3E-07	No	4.0E-07	6.4E-07	1.1E-07	4.2E-07	2.5E-08	6.56E-07
FcγRIIB	kon(1/Ms)		1.2E+05	No	1.4E+05	2.8E+05	2.8E+05	2.7E+05	2.0E+05	2.38E+05
	kdis(1/s)		1.6E-02	No	5.0E-02	1.8E-01	3.2E-02	1.1E-01	5.1E-03	1.56E-01
	KD (M)		1.2E-07	No	No	No	1.4E-07	2.6E-07	1.8E-07	1.50E-06
FcγRIIIA(F)	kon(1/Ms)		1.7E+05	No	No	No	3.5E+05	2.2E+05	5.2E+05	1.16E+05
	kdis(1/s)		2.0E-02	No	No	No	5.0E-02	5.7E-02	9.1E-02	1.75E-01
	KD (M)		6.0E-08	No	5.5E-07	2.9E-07	8.7E-08	2.3E-07	8.6E-08	4.56E-07
FcγRIIIA(V)	kon(1/Ms)		4.5E+05	No	1.7E+05	2.5E+05	4.4E+05	3.2E+05	1.2E+06	9.23E+04
	kdis(1/s)		2.7E-02	No	9.2E-02	7.3E-02	3.9E-02	7.4E-02	1.0E-01	4.20E-02
	KD (M)		1.3E-07	No	No	No	No	No	No	5.56E-07
FcγRIIIB	kon(1/Ms)		2.1E+05	No	No	No	No	No	No	2.58E+05
	kdis(1/s)		2.7E-02	No	No	No	No	No	No	1.44E-01
	KD (M)		1.6E-08	1.0E-08	1.8E-08	2.2E-08	1.5E-08	1.65E-08	1.91E-08	1.88E-08
FcRn	kon(1/Ms)		4.8E+05	9.9E+05	6.4E+05	3.5E+05	6.7E+05	6.73E+05	6.07E+05	8.03E+05
	kdis(1/s)		8.0E-03	9.9E-03	1.2E-02	7.7E-03	1.0E-02	1.11E-02	1.16E-02	1.50E-02

Table 8. FcR Binding Activity of IgG2 Mutants

		IgG1	IgG2	IgG4	G2m5	G2m7	G2m8	G2m9
FcγRI	KD (M)	5.6E-9	No	4.8E-09	No	No	No	No
	kon(1/Ms)	2.2E+05	No	3.6E+05	No	No	No	No
	kdis(1/s)	1.2E-03	No	1.7E-03	No	No	No	No
FcγRIIA	KD (M)	5.6E-08	8.9E-08	4.0E-07	2.30E-07	1.20E-07	9.30E-08	2.80E-07
	kon(1/Ms)	5.7E+05	4.4E+05	1.4E+05	6.00E+05	2.30E+05	9.90E+05	1.10E+05
	kdis(1/s)	3.2E-02	3.9E-02	5.5E-02	1.40E-01	2.80E-02	9.20E-02	3.00E-02
FcγRIIB	KD (M)	1.3E-07	No	4.0E-07	3.10E-07	1.70E-07	5.00E-07	5.20E-08
	kon(1/Ms)	1.2E+05	No	1.4E+05	3.90E+05	3.10E+05	5.10E+05	9.70E+04
	kdis(1/s)	1.6E-02	No	5.0E-02	1.20E-01	5.20E-02	2.60E-01	5.00E-03
FcγRIIIA(F)	KD (M)	1.2E-07	No	No	No	No	No	No
	kon(1/Ms)	1.7E+05	No	No	No	No	No	No
	kdis(1/s)	2.0E-02	No	No	No	No	No	No
FcγRIIIA(V)	KD (M)	6.0E-08	No	5.5E-07	No	No	No	No
	kon(1/Ms)	4.5E+05	No	1.7E+05	No	No	No	No
	kdis(1/s)	2.7E-02	No	9.2E-02	No	No	No	No
FcγRIIIB	KD (M)	1.3E-07	No	No	No	No	No	No
	kon(1/Ms)	2.1E+05	No	No	No	No	No	No
	kdis(1/s)	2.7E-02	No	No	No	No	No	No
FcRn	KD (M)	1.6E-08	1.0E-08	1.8E-08	1.40E-08	5.70E-09	6.30E-09	2.20E-08
	kon(1/Ms)	4.8E+05	9.9E+05	6.4E+05	8.00E+05	9.10E+05	8.50E+05	2.70E+05
	kdis(1/s)	8.0E-03	9.9E-03	1.2E-02	1.10E-02	5.20E-03	5.30E-03	5.80E-03

Table 9. FcR Binding Activity of IgG4 Mutants

			IgG1	IgG2	IgG4	G4m5	G4m7	G4m8	G4m9	G4m30	G4PE
	KD (M)		5.6E-9	No	4.8E-09	N.D.	2.10E-09	3.80E-09	1.70E-08	5.28E-07	6.05E-07
FcγRI	kon(1/Ms)		2.2E+05	No	3.6E+05	N.D.	5.40E+05	7.20E+05	4.40E+05	3.65E+05	1.02E+05
	kdis(1/s)		1.2E-03	No	1.7E-03	N.D.	1.10E-03	2.80E-03	7.40E-03	1.93E-01	6.17E-02
FcγRIIA	KD (M)		5.6E-08	8.9E-08	4.0E-07	N.D.	2.10E-07	2.60E-07	No	9.77E-07	5.45E-08
	kon(1/Ms)		5.7E+05	4.4E+05	1.4E+05	N.D.	1.60E+05	3.90E+05	No	2.40E+05	4.69E+05
	kdis(1/s)		3.2E-02	3.9E-02	5.5E-02	N.D.	3.40E-02	1.00E-01	No	2.34E-01	2.56E-02
FcγRIIB	KD (M)		1.3E-07	No	4.0E-07	N.D.	4.50E-08	8.90E-08	1.40E-07	2.22E-06	3.07E-08
	kon(1/Ms)		1.2E+05	No	1.4E+05	N.D.	4.40E+05	5.30E+05	9.40E+04	5.36E+04	4.01E+05
	kdis(1/s)		1.6E-02	No	5.0E-02	N.D.	2.00E-02	4.80E-02	1.30E-02	1.18E-01	1.23E-02
FcγRIIIA (F)	KD (M)		1.2E-07	No	No	N.D.	No	No	No	2.84E-07	1.45E-07
	kon(1/Ms)		1.7E+05	No	No	N.D.	No	No	No	1.06E+06	4.68E+05
	kdis(1/s)		2.0E-02	No	No	N.D.	No	No	No	3.03E-01	6.78E-02
FcγRIIIA (V)	KD (M)		6.0E-08	No	5.5E-07	N.D.	No	No	No	2.17E-07	2.02E-07
	kon(1/Ms)		4.5E+05	No	1.7E+05	N.D.	No	No	No	1.42E+06	2.96E+05
	kdis(1/s)		2.7E-02	No	9.2E-02	N.D.	No	No	No	3.09E-01	5.96E-02
FcγRIIIB	KD (M)		1.3E-07	No	No	N.D.	No	No	No	2.16E-07	1.32E-07
	kon(1/Ms)		2.1E+05	No	No	N.D.	No	No	No	1.37E+06	4.37E+05
	kdis(1/s)		2.7E-02	No	No	N.D.	No	No	No	2.97E-01	5.79E-02
FcRn	KD (M)		1.6E-08	1.0E-08	1.8E-08	N.D.	2.30E-08	2.00E-08	7.50E-08	7.61E-09	2.19E-08
	kon(1/Ms)		4.8E+05	9.9E+05	6.4E+05	N.D.	5.20E+05	6.10E+05	2.30E+05	1.75E+06	6.98E+05
	kdis(1/s)		8.0E-03	9.9E-03	1.2E-02	N.D.	1.20E-02	1.20E-02	1.70E-02	1.33E-02	1.52E-02

Example 3. Exemplary IgG Mutants Having Combined Mutations to Enhance Selective Binding to FcγRIIB (CD32B)

The human Fc variants G1m15, G1m17, G1m18, G1m19, G2m25, G2m27, G2m28, G4m15, G4m17, G4m18, and G4m19, G4m25, G4m27, G4m28, and G4m29 disclosed herein were linked to the VH fragment of an anti-CD137 antibody and the corresponding IgG1, G2, and G4 CH1 fragments to produce a full-length IgG heavy chain. See Example 1 above. These Fc variants contain a combination of one or mutations in the hinge domain and one or more mutations in the CH2 and/or CH3 domains. These mutant IgG heavy chains were cloned, co-expressed with the light chain of the anti-CD137 antibody (see Example 1 above), and purified using standard molecular biology and antibody protocols. The results are shown in Tables 10-12 below.

Example 4. Determination of Human IgG Mutants Binding Activity to Cellular FcγR

To determine the binding activity of human IgG mutants to cellular Fc receptors, CHO cells were genetically engineered to express human FcγRs (FcγRI, FcγRIIA(H131), FcγRIIA(R131), FcγRIIB, FcγRIIC, and FcγRIII) using a lentivirus delivery system as known in the art.

IgG Fc mutants, including G1m-2, G1m-4, G1mAA, G1mAG, G2m-1 G2m-4, G2m15, G2m17, G2m17, G2m18, G2m19, G2m20, G2m27, G2m28, G4m-1, G4m-2, G430, and G4PE (amino acid sequences provided above) were designed and constructed following the disclosures herein. These IgG mutants contain mutations in either the upper hinge domain or the lower hinge domain as indicated.

For FACS analysis of the IgG mutants' binding to different FcγRs, FcγR overexpressing CHO cells were harvested using trypsin-EDTA and were suspended in cold staining buffer (3% BSA in PBS). Test IgG mutants, which were diluted in staining buffer, were added into the cells. The mixture was incubated 4°C for 2 hours, and then washed twice with cold staining buffer and re-suspended in PE-labeled anti-human IgG followed by incubation at 4°C for 2 hours. The mixture was washed twice with staining buffer and re-suspended in 2% PFA in PBS for FACS.

Table 10. FcR Binding Activity of Human IgG1 Mutants

		IgG1	G1m15	G1m17	G1m18	G1m19	G1m25	G1m27	G1m28	G1m29
FcγRI	KD (M)	5.6E-9	1.2E-07	8.8E-08	2.6E-07	1.2E-07	No	No	2.4E-07	3.7E-08
	kon(1/Ms)	2.2E+05	5.0E+05	4.6E+05	8.3E+05	7.1E+05	No	No	6.4E+05	2.6E+05
	kdis(1/s)	1.2E-03	6.0E-02	4.1E-02	2.1E-01	8.6E-02	No	No	1.6E-01	9.8E-03
FcγRIIA	KD (M)	5.6E-08	4.8E-07	2.5E-07	1.4E-07	8.1E-08	No	No	No	5.3E-07
	kon(1/Ms)	5.7E+05	2.3E+05	2.4E+05	2.0E+05	2.4E+05	No	No	No	2.6E+05
	kdis(1/s)	3.2E-02	1.1E-01	5.9E-02	2.9E-02	1.9E-02	No	No	No	1.4E-01
FcγRIIB	KD (M)	1.3E-07	7.0E-07	1.8E-07	6.7E-07	1.1E-07	No	4.1E-07	4.1E-07	4.6E-08
	kon(1/Ms)	1.2E+05	5.4E+05	2.4E+05	5.8E+05	2.6E+05	No	2.6E+05	4.8E+05	2.5E+05
	kdis(1/s)	1.6E-02	3.8E-01	4.3E-02	3.9E-01	2.8E-02	No	1.0E-01	2.0E-01	1.1E-02
FcγRIIIA (F)	KD (M)	1.2E-07	No	No	No	No	No	No	No	No
	kon(1/Ms)	1.7E+05	No	No	No	No	No	No	No	No
	kdis(1/s)	2.0E-02	No	No	No	No	No	No	No	No
FcγRIIIA (V)	KD (M)	6.0E-08	No	No	No	No	No	No	No	No
	kon(1/Ms)	4.5E+05	No	No	No	No	No	No	No	No
	kdis(1/s)	2.7E-02	No	No	No	No	No	No	No	No
FcγRIIIB	KD (M)	1.3E-07	No	No	No	No	No	No	No	No
	kon(1/Ms)	2.1E+05	No	No	No	No	No	No	No	No
	kdis(1/s)	2.7E-02	No	No	No	No	No	No	No	No
FcRn	KD (M)	1.6E-08	2.1E-08	1.07E-08	1.28E-08	1.34E-08	1.5E-08	1.6E-08	1.7E-08	1.7E-08
	kon(1/Ms)	4.8E+05	6.0E+05	7.15E+05	6.30E+05	7.70E+05	6.2E+05	6.5E+05	6.5E+05	7.6E+05
	kdis(1/s)	8.0E-03	1.3E-02	7.64E-03	8.03E-03	1.03E-02	9.3E-03	1.0E-02	1.1E-02	1.3E-02

Table 11. FcR Binding Activity of Human IgG2 Mutants

		IgG2	G2m17	G2m19	G2m20	G2m28
FcγRI	KD (M)	No	2.53E-07	No	No	6.61E-08
	kon(1/Ms)	No	1.09E+06	No	No	1.49E+06
	kdis(1/s)	No	2.77E-01	No	No	9.80E-02
FcγRIIA	KD (M)	8.9E-08	4.22E-07	3.36E-07	8.29E-07	1.91E-07
	kon(1/Ms)	4.4E+05	6.48E+05	2.49E+05	1.99E+04	4.70E+05
	kdis(1/s)	3.9E-02	2.74E-01	8.36E-02	1.65E-02	8.98E-02
FcγRIIB	KD (M)	No	2.36E-07	2.40E-07	5.17E-07	2.56E-07
	kon(1/Ms)	No	3.61E+05	1.62E+05	1.76E+05	1.81E+05
	kdis(1/s)	No	8.48E-02	3.87E-02	9.15E-02	4.64E-02
FcγRIIIA (F)	KD (M)	No	1.68E-07	No	9.17E-07	5.27E-07
	kon(1/Ms)	No	1.56E+06	No	1.43E+04	4.87E+05
	kdis(1/s)	No	2.62E-01	No	1.30E-02	2.57E-01
FcγRIIIA (V)	KD (M)	No	1.53E-07	No	1.26E-06	No
	kon(1/Ms)	No	1.34E+06	No	1.18E+04	No
	kdis(1/s)	No	2.05E-01	No	1.49E-02	No
FcγRIIIB	KD (M)	No	1.17E-07	No	No	No
	kon(1/Ms)	No	1.74E+06	No	No	No
	kdis(1/s)	No	2.04E-01	No	No	No
FcRn	KD (M)	1.0E-08	7.46E-09	3.54E-08	5.76E-08	2.64E-08
	kon(1/Ms)	19.9+E05	1.87E+06	2.65E+05	2.12E+05	7.37E+05
	kdis(1/s)	9.9E+03	1.39E-02	9.41E-03	1.22E-02	1.95E-02

Table 12. FcR Binding Activity of Human IgG4 Mutants

		IgG4	G4m15	G4m17	G4m18	G4m19	G4m25	G4m27	G4m28	G4m29
FcγRI	KD (M)	4.8E-09	N.D.	No	Low	Low	4.6E-08	6.6E-08	5.6E-08	4.4E-08
	kon(1/Ms)	3.6E+05	N.D.	No	Low	Low	3.8E+05	3.7E+05	4.9E+05	5.2E+05
	kdis(1/s)	1.7E-03	N.D.	No	Low	Low	1.7E-02	2.5E-02	2.7E-02	2.3E-02
FcγRIIA	KD (M)	4.0E-07	N.D.	6.0E-07	Low	Low	6.6E-07	Low	1.8E-06	2.6E-07
	kon(1/Ms)	1.4E+05	N.D.	4.0E+04	Low	Low	5.7E+04	Low	3.1E+04	1.0E+05
	kdis(1/s)	5.5E-02	N.D.	2.4E-02	Low	Low	3.8E-02	Low	5.5E-02	2.7E-02
FcγRIIB	KD (M)	4.0E-07	N.D.	2.8E-07	No	Low	2.4E-07	3.1E-07	1.8E-07	7.8E-08
	kon(1/Ms)	1.4E+05	N.D.	6.8E+04	No	Low	1.3E+05	1.3E+05	1.9E+05	1.7E+05
	kdis(1/s)	5.0E-02	N.D.	1.9E-02	No	Low	3.0E-02	3.9E-02	3.4E-02	1.3E-02
FcγRIIA (F)	KD (M)	No	N.D.	No	No	No	No	No	No	No
	kon(1/Ms)	No	N.D.	No	No	No	No	No	No	No
	kdis(1/s)	No	N.D.	No	No	No	No	No	No	No
FcγRIIA (V)	KD (M)	5.5E-07	N.D.	No	No	No	No	No	No	No
	kon(1/Ms)	1.7E+05	N.D.	No	No	No	No	No	No	No
	kdis(1/s)	9.2E-02	N.D.	No	No	No	No	No	No	No
FcγRIIIB	KD (M)	No	N.D.	No	No	No	No	No	No	No
	kon(1/Ms)	No	N.D.	No	No	No	No	No	No	No
	kdis(1/s)	No	N.D.	No	No	No	No	No	No	No
FcRn	KD (M)	1.8E-08	N.D.	5.2E-08	1.1E-08	1.3E-08	1.9E-08	2.3E-08	2.0E-08	2.3E-08
	kon(1/Ms)	6.4E+05	N.D.	2.1E+05	6.1E+05	6.9E+05	4.8E+05	4.5E+05	5.8E+05	5.6E+05
	kdis(1/s)	1.2E-02	N.D.	1.1E-02	6.8E-03	9.0E-03	9.0E-03	1.0E-02	1.2E-02	1.3E-02

As shown in FIGs. 1A-1W, a number of human IgG1, IgG2, and IgG4 mutants showed binding activity to FcγRs expressed on the cell surface. Qualitative summaries of the changes of the mutants' Fcγ binding activity relative to the wild-type counterpart are provided in Tables 2-4 above.

5
Example 5. Human IgG Mutants Capable of Binding to Cellular FcγRIIB Showed Enhanced Agonist Activity

To verify that binding to cellular FcγR2B would enhance agonist activity of the IgG mutants, a co-culture assay was developed, which involves CHO cells expressing FcγR2B and human CD8 positive T cells. CHO-FcγRIIB cells were plated at 2×10^4 /well in 96-well cell culture plate and were incubated overnight. To isolate human CD8 positive T cells, fresh blood from healthy donors were mixed with equal volume of DPBS gently. The blood sample was then placed on top of Ficoll underlay and centrifuged 30 min at 1000g at RT without brake. The buffy coat containing PBMC was harvested into a new tube and was washed with DPBS. CD8 positive T cells were isolated from the PBMC using EasySep™ Human CD8+ T Cell Isolation Kit (Stemcell #17953) according to the kit manual. The isolated CD8+ T cells suspended in RPMI media were added to the plates with CHO-FcγRIIB cells. The anti-human CD3 antibody OKT3 was added to a final concentration of 0.1ug/ml followed by addition of test antibodies diluted at desired concentrations. The plates were cultured for 3 days, and then the culture supernatants were harvested for measurement of IFNγ concentration by ELISA using Human IFN-gamma ELISA Ready-SET-GO kit (EBIOSCIENCE, #88-7316-88).

As shown in FIG. 2A-2C, a number of tested IgG mutants stimulated human CD8⁺ cells in the presence of CHO-FcγRIIB cells as evidenced by the secretion of IFN-γ.

Example 6. Effect of IgG Isotypes *in vitro* and *in vivo* in Mouse Models

Two different antibodies against mouse 4-1BB (CD137), LOB12.3 and 3H3, were examined in detail. These two antibodies were cloned and expressed as various mouse IgG isotypes including murine IgG1, a murine IgG1 N297A mutant known to have diminished ability to bind FcRs (as a negative control), and murine IgG2a.

Murine CD8 positive T cells were stimulated by the antibodies in a solution. In a typical *in vitro* co-stimulation assay of murine CD8 T cells, antibody LOB12.3 was not able to stimulate the T cells as measure by interferon gamma secretion, whereas 3H3 antibody exhibited agonistic activity regardless of the antibody isotype, as shown in FIG. 3.

5 *In vivo* antitumor activity and liver toxicity were examined. The activity profile of these two antibodies and their isotypes were vastly different. In a murine syngeneic CT26 colorectal cancer model, both LOB12.3 and 3H3 showed robust antitumor activity including complete rejection of tumors, although the optimal isotype for each antibody is distinct. FIG. 4A-4B. LOB12.3 showed no obvious co-stimulatory activity *in vitro* but was efficacious *in*
10 *vivo* when it was in murine IgG1 isotype, supporting that FcγR2B mediated cross-linking would result in efficient agonistic activity. 3H3 is intrinsically agonistic and its activity was not dependent on Fc-mediated cross-linking. In fact, 3H3 in Fc-null mutant was most efficacious.

15 An unexpected observation was the differential effect on liver toxicity of these antibodies. Both LOB12.3 and 3H3 were efficacious in inducing tumor rejection but only 3H3 showed liver toxicity as determined by increased liver enzyme ALT in serum samples of the treated animals (FIG. 4B).

20 These experimental data, for the first time, suggested a potential utility of selecting and engineering agonistic antibodies for increased therapeutic index of co-stimulation agonist antibodies.

Example 7. Pharmacokinetic Study of Chimeric Antibodies

25 *In vivo* studies were performed to study the pharmacokinetics of selected chimeric antibodies. Antibodies formulated in PBS were administered to C57BL/6 mice (6-7 weeks old, 19-20 g, male) via tail vein injection. Doses were 1-3 mg/kg (n=4 mice per group).

30 Blood samples were taken prior to administration of the antibodies, and 1h, 2h, 4h, 8h, 1d, 2d, 3d, 5d, 8d, 11d, 15d and 21d following administration. 10 uL blood of per time point was extracted and added to 40uL of a PBS-BSA solution. The sample was then mixed well and centrifuged at 2000 g for 5 minutes at 4°C. The supernatant was put on dry ice immediately after collection and stored at approximately -70°C until analysis. Antibody concentrations in the blood samples were determined by ELISA as described below briefly.

CD137 protein (human CD137-His tag protein (Sino Biological Inc. Cat#10377-H08H-100) or rhesus monkey CD137-His (Sino Biological Inc. Cat#90305-K08H-100)) was diluted in PBS to 1 µg/ml and used to coat an ELISA plate (Corning, Cat#9018, high binding) at a concentration of 50 µl/well. The plate was incubated overnight at 4 °C. The plate was
 5 then decanted and washed with PBS-T, and 200 µl/well assay diluent (1xPBS/1% BSA/0.05% Tween-20/0.05% proclin 300) was added. After a three hour incubation at room temperature, the plate was washed with PBS-T three times. The samples, at appropriate dilutions, along with known concentrations of antibody standards diluted in assay diluent to 0, 0.000003, 0.00003, 0.0003, 0.003, 0.03, and 0.3 µg/ml (or approximately 0, 0.00002,
 10 0.0002, 0.002, 0.02, 0.2 and 2 nM), were then added to the plate (50 µl/well). The plate was incubated for one hour at 37°C and then washed three times with PBS-T. Anti-human IgG-HRP conjugate (Bethyl Cat#A80-319P) at a 1:10,000 dilution was added to the plate (100 µl/well). The plate was incubated for 0.5 hour at 37°C, followed by washing with PBS-T three times. The TMB substrate solution was added (100 µl/well). The color was allowed to
 15 develop for 8 minutes before it was stopped with 100 µl/well 2N H₂SO₄. Absorbance at 450nm was determined with an ELISA reader, and antibody concentrations were calculated from the standard curve dose responses.

FIGs. 5A-5G show the plasma antibody concentration of the chimeric antibodies after a single intravenous injection of 1-3 mg/kg. The pharmacokinetics parameters of these
 20 antibodies examined are listed in Table 13 below, which indicates that most, if not all, of the mutations examined did not alter the normal IgG PK parameters.

Table 13. Summary of Pharmacokinetics of IgG Variants Examined in Mice

PK parameters	Unit	G4	G4m2	G4m7	G1m27	G1m28	G1m29	G4m27	G4m28
CL	mL/day/kg	10.70	8.39	9.62	8.06	10.7	10.30	9.98	8.95
V _{ss}	mL/kg	144	146	130	152	142	142	122	126
V1	mL/kg	58.9	58.7	53.6	56.8	58.4	63.7	53.3	61.1
Alpha t _{1/2}	day	0.1760	0.119	0.161	0.116	0.184	0.2850	0.306	0.295
Beta t _{1/2}	day	9.9	12.3	10.2	13.40	9.54	9.9	8.9	10.3
AUC	day*µg/mL	287	359	324	374	282	294	146	164
MRT	day	13.8	17.5	14.4	19.0	13.4	13.8	12.2	14.3

PK parameters	Unit	G4m29	G1m2	G4m10	G2m19	G4m1	G1mAA	G2	G2m1
CL	mL/day/kg	10.70	8.58	9.47	7.19	6.24	7.71	6.28	8.41
V _{ss}	mL/kg	147	173	116	127	91	125	118	122
V1	mL/kg	66.1		56.6	49.7	39.2	52.6	49.2	53.4
Alpha t _{1/2}	day	0.256		0.151	0.182	0.099	0.0919	0.108	0.123
Beta t _{1/2}	day	9.99	14.8	8.9	13.1	10.40	11.5	13.7	10.3
AUC	day*µg/mL	142	365.0	158	431	487	394	497	374
MRT	day	13.9	21	12.7	18.5	14.7	16.4	19.6	14.6

OTHER EMBODIMENTS

5 All of the features disclosed in this specification may be combined in any combination. Each feature disclosed in this specification may be replaced by an alternative feature serving the same, equivalent, or similar purpose. Thus, unless expressly stated otherwise, each feature disclosed is only an example of a generic series of equivalent or similar features.

10 From the above description, one skilled in the art can easily ascertain the essential characteristics of the present invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, other embodiments are also within the claims.

What Is Claimed Is:

1. A protein comprising a variant Fc region, wherein the variant Fc region comprises a mutation at any of positions 218-329 as compared to a wild-type parent Fc region, and wherein the numbering is according to the EU index.

2. The protein of claim 1, wherein the mutation is at any of positions 218-328.

3. The protein of claim 1 or claim 2, wherein the mutation is an amino acid residue substitution, an insertion, a deletion, or a combination thereof.

4. The protein of claim 1, wherein the mutation comprises one or more of the following:

- (a) a mutation at one or more of positions 219-225;
- (b) an insertion between position 218 and position 219, or at position 236;
- (c) an amino acid residue substitution at one or more positions of 233-235;
- (d) an insertion at any of positions 220-223;
- (e) a deletion at one or more of positions 236-238; and
- (f) an amino acid substitution at one or more of positions 267, 273, 328, and 329.

5. The protein of claim 4, wherein the mutation of (a) comprises an insertion, a deletion, an amino acid residue substitution, or a combination thereof.

6. The protein of any one of claims 1-3, wherein the mutation comprises (i) a mutation at one or more of positions 234-238, and (ii) a mutation at one or more of positions 267, 273, 328, and 329.

7. The protein of claim 4, wherein (i) is a deletion and (ii) is an amino acid residue substitution.

8. The protein of claim 7, wherein (i) comprises a deletion of one or more of positions 236-238.

9. The protein of claim 7 or claim 8, wherein (ii) comprises amino acid residue substitutions S267E, V273E, L328F, P329G, or a combination thereof.

10. The protein of any one of claims 1-9, wherein the parent Fc region is of an IgG1 molecule.

11. The protein of claim 10, wherein the mutation comprises one or more of the following:

- (a) an insertion between position 228 and position 229;
- (b) a mutation at one or more of positions 220-225;
- (c) an amino acid substitution at positions 234 and 235;
- (d) a deletion at one or more of positions 236-238; and
- (e) an amino acid substitution at one or more positions 267, 273, 328, and 329.

12. The protein of claim 11, wherein the mutation comprises (i) a deletion at one or more of positions 236-238.

13. The protein of claim 12, wherein the mutation further comprises (ii) an amino acid residue substitution at one or more of positions 267, 273, and 328.

14. The protein of claim 13, wherein the amino acid residue substitution is S267E.

15. The protein of claim 13, wherein the variant Fc region is selected from the group consisting of G1m1, G1m2, G1m-2, G1m-4, G1m5, G1m7, G1m8, G1m9, G1m15, G1m17, G1m18, G1m19, G1m25, G1m27, G1m28, G1m29, G1mAA, and G1mAG.

16. The protein of any one of claims 1-9, wherein the parent Fc region is of an IgG2 molecule.

17. The protein of claim 16, wherein the mutation comprises one of the following:

- (a) a mutation at one or more positions of 219-225;
- (b) an amino acid substitution at one or more of positions 233-235;
- (c) an insertion between position 218 and 219, or at position 236;
- (d) a deletion of one or more of positions 237 and 238; and
- (e) an amino acid residue substitution at one or more of positions 267, 273, and 328.

18. The protein of claim 14 or claim 15, wherein the mutation comprises (i) a deletion at one or more of positions 237 and 238.

19. The protein of claim 18, wherein the mutation further comprises (ii) an amino acid residue substitution at one or more of positions 267, 273, and 328.

20. The protein of claim 19, wherein the amino acid residue substitution is S267E, L328F, or a combination thereof.

21. The protein of claim 19, wherein the variant Fc region is selected from the group consisting of G2m1, G2m-1, G2m2, G2m-4, G2m5, G2m7, G2m8, G2m9, G2m10, G2m15, G2m17, G2m18, G2m19, G2m20, G2m27, G2m27, and G2m28.

22. The protein of any one of claims 1-9, wherein the parent Fc is of an IgG4 molecule.

23. The protein of claim 22, wherein the mutation comprises the S228P amino acid residue substitution, and one or more of the following:

- (a) a mutation at one or more of positions 219-225;
- (b) an amino acid residue substitution at one or more positions 234 and 235;

(c) a deletion at one or more of positions 236-238; and

(d) an amino acid residue substitution at one or more of positions 267, 273, and 328.

24. The protein of claim 23, wherein the mutation comprises (i) the S228P amino acid residue substitution, and (ii) a deletion at one or more of positions 236-238.

25. The protein of claim 24, wherein the mutation further comprises (iii) an amino acid residue substitution at one or more of positions 267, 273, and 328.

26. The protein of claim 25, wherein the amino acid residue substitution of L328F.

27. The protein of claim 24, wherein the variant Fc region is selected from the group consisting of G4m1, G4m-1, G4m2, G4m-2, G4m3, G4m4, G4m5, G4m7, G4m8, G4m9, G4m10, G4m17, G4m18, G4m19, G4m20, G4m25, G4m27, G4m28, G4m29, G4m30, and G4mPE.

28. The protein of any one of claims 1-27, wherein the variant Fc region exhibits an enhanced binding affinity to Fc γ RIIB as compared with the parent Fc region.

29. The protein of any one of claims 1-28, wherein the variant Fc region exhibits an enhanced selectivity to Fc γ RIIB as compared with the parent Fc region.

30. The protein of any one of claims 1-29, wherein the variant Fc region exhibits low or no binding activity to any Fc γ R.

31. The protein of any one of claims 1-30, wherein the variant Fc region binds FcRn.

32. The protein of any one of claims 1-31, wherein the protein is an antibody.

33. A protein comprising a variant Fc region of an IgG2 or IgG4 molecule,
5 wherein the variant Fc region comprises a mutation at position 267, position 273, position
328, or a combination thereof as compared to a wild-type parent IgG2 or IgG4 Fc region, and
wherein the numbering is according to the EU index.

34. The protein of claim 33, wherein the IgG2 or IgG4 molecule is a human IgG2
10 or human IgG4 molecule.

35. A pharmaceutical composition, comprising a protein of any one of claims 1-34
and a pharmaceutically acceptable carrier.

15 36. A method for selectively activate an immune response in a subject, the method
comprising administering to a subject in need thereof an effective amount of a therapeutic
agent, wherein the therapeutic agent comprises a first moiety that binds an immune cell
receptor and a second moiety that binds FcγRIIB.

20 37. The method of claim 36, wherein the immune cell receptor is a member of the
tumor necrosis factor receptor superfamily (TNFRSF).

38. The method of claim 37, wherein the member of TNFRSF is selected from the
group consisting of FAS, TNFRSF12A, 4-1BB/CD137, TNFRSF13B, TNFRSF13C,
25 CD27/TNFRSF7, CD30/TNFRSF8, CD40/TNFRSF5, DR3/TNFRSF25,
DR4/TNFRSF10A, DR5/TNFRSF10B, DR6/TNFRSF21, GITR/TNFRSF18,
HVEM/TNFRSF14, LTβR, OX40/TNFRSF4, TROY/TNFRSF19, RELT/TNFRSF19L,
TNFRSF12A, TNFRSF13B, TL1A/TNFRSF15, TNFRSF17, TNFRSF1A, TNFRSF11B,
RANK/TNFRSF11A, TNFRSF11B, NGFR, EDA2R, and TNFRSF1B.

39. The method of any one of claims 36-38, wherein the therapeutic agent is an antibody.

40. The method of claim 39, wherein the antibody is an agonist antibody.

5 41. The method of claim 39 or claim 40, wherein the antibody is an IgG1, IgG2, or IgG4 molecule.

42. The method of any one of claims 36-41, wherein the therapeutic agent selectively binds Fc γ RIIB.

10 43. The method of any one claims 36-42, wherein the therapeutic agent contains a variant Fc region having an enhanced binding affinity to Fc γ RIIB, and/or an enhanced binding selectivity as compared with a wild-type counterpart.

15 44. The method of claim 43, wherein the variant Fc region selectively binds human Fc γ RIIB.

20 45. The method of any one of claims 36-44, wherein the therapeutic agent is a bispecific antibody, which comprises a first antigen-binding fragment that is specific to the immune cell receptor, and a second antigen-binding fragment that is specific to Fc γ RIIB.

46. The method of any one of claims 36-45, wherein the subject is a human patient having or suspected of having a cancer.

25 47. The method of claim 46, wherein the cancer is selected from the group consisting of lung cancer, stomach cancer, liver cancer, breast cancer, skin cancer, pancreatic cancer, brain cancer, prostate cancer, bladder cancer, colorectal cancer, sarcoma, bone cancer, lymphoma and a hematological cancer.

48. A method for selectively activate an immune response in a subject, the method comprising administering to a subject in need thereof an effective amount of the pharmaceutical composition of claim 35.

1/23




FIG. 1A






FIG. 1B

2/23



3/23




FIG. 1E




FIG. 1F

4/23




FIG. 1G




FIG. 1H

5/23




FIG. 1I




FIG. 1J

6/23




FIG. 1K




FIG. 1L

7/23




FIG. 1M




FIG. 1N

8/23




FIG. 10




FIG. 1P

9/23




FIG. 1Q




FIG. 1R

10/23




FIG. 1S




FIG. 1T




FIG. 1U




FIG. 1V

12/23




FIG. 1W

13/23




FIG. 2A




FIG. 2B

14/23




FIG. 2C




FIG. 3

15/23




FIG. 4A




FIG. 4B

16/23




FIG. 5A

17/23





FIG. 5B

18/23




FIG. 5C

19/23





FIG. 5D

20/23





FIG. 5E

21/23





FIG. 5F

22/23





FIG. 5G

23/23




FIG. 5H




FIG. 1A