



US 20070134796A1

(19) **United States**

(12) **Patent Application Publication**
Holmes et al.

(10) **Pub. No.: US 2007/0134796 A1**

(43) **Pub. Date: Jun. 14, 2007**

(54) **TARGETED INTEGRATION AND
EXPRESSION OF EXOGENOUS NUCLEIC
ACID SEQUENCES**

(22) Filed: **Jul. 26, 2006**

Related U.S. Application Data

(75) Inventors: **Michael C. Holmes**, Oakland, CA
(US); **Fyodor Urnov**, Point Richmond,
CA (US); **Philip D. Gregory**, Orinda,
CA (US); **Edward J. Rebar**, El Cerrito,
CA (US); **Sean M. Brennan**, San
Ramon, CA (US)

(60) Provisional application No. 60/702,394, filed on Jul.
26, 2005. Provisional application No. 60/721,054,
filed on Sep. 26, 2005.

Publication Classification

Correspondence Address:
ROBINS & PASTERNAK
1731 EMBARCADERO ROAD
SUITE 230
PALO ALTO, CA 94303 (US)

(51) **Int. Cl.**
C12N 15/09 (2006.01)
(52) **U.S. Cl.** **435/455; 435/320.1**

(57) **ABSTRACT**

Disclosed herein are methods and compositions for targeted
integration of a exogenous sequence into a predetermined
target site in a genome for use, for example, in protein
expression and gene inactivation.

(73) Assignee: **Sangamo BioSciences, Inc.**

(21) Appl. No.: **11/493,423**

FIGURE 1

MetGlyPheLeuLysLeuIle (SEQ ID NO:2)
 CTGCCGCCGCGCGCGCGGCGTCATGGGGTTCTTGAAACTGATT
 GACGCGCGGCGCGGCGCGGCGAGTACCCCAAGACTTTGACTAA (SEQ ID NO:1)
 5 10 15 20 25 30 35 40

FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6

<u>TAGTCCTGCAGGTTTAAACGAATTCGCCCTTCTCAGCAAGCGTGAGCTCA</u>	50
<u>GGTCTCCCCCGCCTCCTTGAACCTCAAGAACTGCTCTGACTCCGCCCAGC</u>	100
<u>AACAACTCCTCCGGGGATCTGGTCCGCAGGAGCAAGTGTTTGTGTGTTGCC</u>	150
<u>ATGCAACAAGAAAAGGGGGCGGAGGCACCACGCCAGTCGTCAGCTCGCTC</u>	200
<u>CTCGTATACGCAACATCAGTCCCCGCCCTGGTCCCACTCCTGCCGGAAG</u>	250
<u>GCGAAGATCCCGTTAGGCCTGGACGTATTCTCGCGACATTTGCCGGTCGC</u>	300
<u>CCGGCTTGCACTGCGGCGTTTCCCGCGCGGGCTACCTCAGTTCTCGGGCG</u>	350
<u>TACGGCGCGGCCTGTCCTACTGCTGCCGGCGCCGCGGCCGTCAT<u>aagaaq</u></u>	400
<u><u>c</u>TTCTGAACTGATTGAAGGGCGAATTCGCGGCCGCTAAATTCAATTCTG</u>	450
<u>CCCTATAGTGAGT</u>	

(SEQ ID NO: 6)

FIGURE 7

TyrLysAsnSerAspAsnAspLysVal (SEQ ID NO:8)
 CTTCCAACCTTTCTCCTCTAGGTTACAAGAACTCGGATAATGATAAAGTCC
 GAAGGTTGGAAAGAGGAGATCCATGTTCTTGAGCCTATTACTATTTCAGG (SEQ ID NO:7)

| | | | | | | | | | |
 5 10 15 20 25 30 35 40 45 50

FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12

<u>TAGTCCTGCAGGTTTAAACGAATTCGCCCTTTCCTCTAGGTAAAGAAtT</u>	50
<u>CcGAcAAcGATAAAGTCCAGAAGTGCAGCCACTATCTATTCCCTGAAGAA</u>	100
ATCACTTCTGGCTGTCAGTTGCAAAAAAAGGAGATCCACCTCTACCAAAC	150
ATTTGTTGTTTCAGCTCCAGGACCCACGGGAACCCAGGAGACAGGCCACAC	200
AGATGCTAAAACCTGCAGAATCTGGGTAATTTGGAAAGAAAGGGTCAAGAG	250
ACCAGGGATACTGTGGGACATTGGAGTCTACAGAGTAGTGTTCTTTTATC	300
ATAAGGGTACATGGGCAGAAAAGAGGAGGTAGGGGATCATGATGGGAAGG	350
GAGGAGGTATTAGGGGCACTACCTTCAGGATCCTGACTTGTCTAGGCCAG	400
GGGAATGACCACATATGCACACATATCTCCAGTGATCCCCTGGGCTCCAG	450
AGAACCTAACACTTCACAAACTGAGTGAATCCCAGCTAGAACTGAACTGG	500
<u>AACAACAGATTCTTGAACCACTGTTTGGAGCACTTGGTGCAGTACCGGAC</u>	550
<u>TAAGGGCGAATTCGCGGCCGCTAAATTCAATTCGCCCTATAGTGAGTCGT</u>	600
<u>ATTACAATTCACTGGCCGTCGTTT</u>	

FIGURE 13

TACTGATGGTATGGGGCCAAGAGATATATCTTAGAGGGAGGGCTGAGGGT	50
TTGAAGTCCAACCTCCTAAGCCAGTGCCAGAAGAGCCAAGGACAGGTACGG	100
CTGTCATCACTTAGACCTCACCCGTGTGGAGCCACACCCTAGGGTTGGCCA	150
ATCTACTCCCAGGAGCAGGGAGGGCAGGAGCCAGGGCTGGGCATAAAAGT	200
CAGGGCAGAGCCATCTATTGCTTACATTTGCTTCTGACACAACCTGTGTTC	250
ACTAGCAACCTCAAACAGACACCATGGTGCATCTGACTCCTGAGGAGAAG	300
TCTGCCGTTACTGCCCTGTGGGGCAAGGTGAACGTGGATGAAGTTGGTGG	350
TGAGGCCCTGGGCAGGTTGGTATCAAGGTTACAAGACAGGTTTAAGGAGA	400
CCAATAGAAACTGGGCATGTGGAGACAGAGAAGACTCTTGGGTTTCTGAT	450
AGGCACTGACTCTCTCTGCCTATTGGTCTATTTTCCCACCCTTAGGCTGC	500
TGGTGGTCTACCCTTGGACCCAGAGGTTCTTTGAGTCCTTTGGGGATCTG	550
TCCACTCCTGATGCTGTTATGGGCAACCCTAAGGTGAAGGCTCATGGCAA	600
GAAAGTGCTCGGTGCCTTTAGTGATGGCCTGGCTCACCTGGACAACCTCA	650
AGGGCACCTTTGCCACACTGAGTGAGCTGCACTGTGACAAGCTGCACGTG	700

FIGURE 14




FIGURE 15




FIGURE 16




FIGURE 17




FIGURE 18

TGCTTACCAAGCTGTGATTCCAAATATTACGTAAATACACTTGCAAAGGA	50
GGATGTTTTTAGTAGCAATTTGTACTGATGGTATGGGGCCAAGAGATATA	100
TCTTAGAGGGAGGGCTGAGGGTTTGAAGTCCAACCTCCTAAGCCAGTGCCA	150
GAAGAGCCAAGGACAGGTACGGCTGTCATCACTTAGACCTCACCCCTGTGG	200
AGCCACACCCTAGGGTTGGCCAATCTACTCCCAGGAGCAGGGAGGGCAGG	250
AGCCAGGGCTGGGCATAAAAGTCAGGGCAGAGCCATCTATTGCTTACATT	300
TGCTTCTGACACAACCTGTGTTCCTAGCAACCTCAAACAGACACCATGGT	350
GCATCTGACTCCTGAGGAGAAGTCTG <u>gCGTTAgTGCCCgaattccgAtcG</u>	400
<u>TcAACcac</u>	

FIGURE 19

5-8

CACGTTTCGTGTTTCGGAGCCGCTTTAACCCACTCTGTGGAAG
GTGCAAAGCACAAGCCTCGGCGAAATTGGGTGAGACACCTTC

5-10

FIGURE 20

1 MAPKKKRKVG IHGVPAAAE RPFQCRICMR NFS**RSDNLSE** HIRTHTGEKP FACDICGRKF
61 **ARNAHRINHT** KIHTGSQKPF QCRICMRNFS **RSDTLSEHIR** THTGEKPFAC DICGRKFAAR
121 **STRTTHTKIH** LRQKDAARGS QLVKSELEEK KSELRHKLKY VPHEYIELIE IARNSTQDRI
181 LEMKVMEFFM KVGYRGKHL GGSRKPDGAI YTVGSPIDYG VIVDTKAYSG GYNLPIGQAD
241 EM**Q**RYVEENQ TRNKHINPNE WWKVYPSSVT EFKFLFVSGH FKGNYKAQLT RLNHITNCNG
301 AVLSVEELLI GGEMIKAGTL TLEEVRRKFN NGEINF

FIGURE 21

```
1  MAPKKKRKVG  IHGVPAAMAE  RPFQCRICMR  NFSSRSDLSR  HIRTHTGEKP  FACDICGRKF
61 ADSSNRKTHT  KIHTGGGGSQ  KPFQCRICMR  NFSSRSDLSV  HIRTHTGEKP  FACDICGRKF
121 ADRSNRITHT  KIHRLQKDAA  RGSQLVKSEL  EEKKSELRHK  LKYVPHEYIE  LIEIARNSTQ
181 DRILEMKVME  FFMKVYGYRG  KHLGGSRKPD  GAIYTVGSPI  DYGVIVDTKA  YSGGYNLPIG
241 QADEMQRYVE  ENQTRNKHIN  PNEWWKVYPS  SVTEFKFLFV  SGHFKGNYKA  QLTRLNHITN
301 CNGAVLSVEE  LLIGGEMIKA  GTLTLEEVRR  KFNNGEINF
```

FIGURE 22

CGAATTCTGCAGTCGACGGTACCGCGGGCCCCGGGATCCACCGGTCGCCACCATGGTGAG
CAAGGGCGAGGAGCTGTTACCGGGGTGGTGCCATCCTGGTCGAGCTGGACGGCGACG
TAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAG
CTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCCGTGCCCTGGCCCACCCTCGT
GACCACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGC
ACGACTTCTTCAAGTCCGCCATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTC
AAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGT
GAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACA
AGCTGGAGTACAAC TACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAAC
GGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGC
CGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACC
ACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATG
GTCCTGCTGGAGTTCGTGACCGCCGCGGGGATCACTCTCGGCATGGACGAGCTGTACAA
GTAAAGCGGCCGCGACTCTAGATCATAATC

FIGURE 23

CGAATTCTGCAGTCGACGGTACCGCGGGCCCCGGGATCCACCGGTGCGCCACCATGGTGAG
CAAGGGCGAGGAGCTGTTACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACG
TAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAG
CTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCCGTGCCCTGGCCCACCCTCGT
GACCACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCTAACACGAAGCAGCAC
GACTTCTTCAAGTCCGCCATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAA
GGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGA
ACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAG
CTGGAGTACAAC TACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGG
CATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCG
ACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCAC
TACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGT
CCTGCTGGAGTTCGTGACCGCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAGT
AAAGCGGCCGCGACTCTAGATCATAATC

FIGURE 24





FIGURE 25



FIGURE 26




FIGURE 27




FIGURE 28




FIGURE 29

GGCGAGGAGCTGTTACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAA
CGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGA
CCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCCGTGCCCTGGCCCCACCCTCGTGACC
ACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGA
CTTCTTCAAGTCCGCCATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGG
ACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAAC
CGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCT
GGAGTACAAC TACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGGCA
TCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGAC
CACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTA
CCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCC
TGCTGGAGTTCGTGACCGCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAGTAA
AGCGGCCGCGACTCTAGATCATAATC

FIGURE 30




FIGURE 31




FIGURE 32

GGCGAGGAGCTGTTACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAA
CGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGA
CCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCCGTGCCCTGGCCCCACCCTCGTGACC
ACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGA
CTTCTTCAAGTCCGCCATGCCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGG
ACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAAC
CGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCT
GGAGTACAACACTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGGCA
TCAAGGTGAAGTTCAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGAC
CACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTA
CCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCC
TGCTGGAGTTCGTGACCGCCCGGGGATCACTCTCGGCATGGACGAGCTGTACAAGTAA
AGCGGCCGCTCGAGTCTAGAGGGCCCGTTTAAACCCGCTGATCAGCCTCGACTGTGCCT
TCTAGTTGCCAGCCATCTGTTGTTGCCCCCTCCCCCGTGCCTTCCTTGACCCTGGAAGG
TGCCACTCCCCTGTCTTTTCTAATAAAATGAGGAAATTGCATCGCATTTGTCTGAGTA
GGTGTCAATTCTATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAA
GACAATAGCAGGCATGCTGGGGATGCGGTGGGCTCTATGGCTTCTGAGGCGGAAAGAAC
CAGCTGGGGCTCTAGGGGGTATCCCCACGCGCCCTGTAGCGGCGCATTAAGCGCGGCGG
GTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCGCTCCT
TTCGCTTTCTTCCCTTCCTTTCTCGCCACGTTGCGCCGGCTTTCCCCGTCAAGCTCTAAA
TCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAAAAC
TTGATTAGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCT
TTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAAACTGGAACAACACT
CAACCCTATCTCGGTCTATTCTTTGATTTATAAGGGATTTTGCCGATTTCGGCCTATT
GGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAATTAATTCTGTGGAATGTGT
GTCAGTTAGGGTGTGGAAGTCCCCAGGCTCCCCAGCAGGCAGAAGTATGCA

FIGURE 33




FIGURE 34




FIGURE 35




FIGURE 36




FIGURE 38




FIGURE 39



FIGURE 40

sca-29b:

MAERPFQCRICMRNFSQSGDLTRHIRTHTGEKPFACDICGRKFATSANLSRHT
KIHTGGGGSQKPFQCRICMRNFSDRSALSRHIRTHTGEKPFACDICGRKFAQS
GHLSRHTKIH (SEQ ID NO:22)

sca-36a:

MAERPFQCRICMRNFSRSQTRKTHIRTHTGEKPFACDICGRKFAQKRNRTKHT
KIHTGSQKPFQCRICMRNFSDRSALSRHIRTHTGEKPFACDICGRKFAQSGNLA
RHTKIH (SEQ ID NO:23)

sca-36b:

MAERPFQCRICMRNFSTSGSLSRHIRTHTGEKPFACDICGRKFADRSDLSRHT
KIHTGGGGSQKPFQCRICMRNFSDRSALSRHIRTHTGEKPFACDICGRKFAQS
GNLARHTKIH (SEQ ID NO:24)

sca-36c:

MAERPFQCRICMRNFSTSSSLSRHIRTHTGEKPFACDICGRKFADRSDLSRHTK
IHTGGGGSQKPFQCRICMRNFSDRSALSRHIRTHTGEKPFACDICGRKFAQSG
NLARHTKIH (SEQ ID NO:25)

FIGURE 41




FIGURE 42




FIGURE 43





FIGURE 44




59D

-tAAAGCGGCTCCGaaccacATTATCCGAGTTc-
-aTTTCGCCGAGGcTtggtgTAATAGGCTCAAg-
IL2-1

(SEQ ID NO: 211)

FIGURE 45



59D IL2-1
 -tAAAGCGGCTCCGaaccacAACTCGGATAATg-
 -aTTTCGCCGAGGCTtggtgTTGAGCCTATTAc-

(SEQ ID NO: 212)

FIGURE 46




FIGURE 47




FIGURE 48




FIGURE 49




FIGURE 50




FIGURE 51




FIGURE 52




FIGURE 53

```
CTCGGTTTCC CTAACCCAAT CCAGCCAGTA CCTGAGCATTT G::::::::: ::::::::::
CTCGGTTTCC CTAACCCAAT CCAGCCAGTA CCTGAGCATTT GTGGCACCTT CCAGGGTCAA

::::::::: ::::::::::: ::::::::::: ::::::::::: ::::::::::: :::::::::::
GGAAGGCACG GGGGAGGGGC AAACAACAGA TGGCTGGCAA CTAGAAGGCA CAGTCGAGGC

::::::::: ::::::::::: ::::::::::: ::::::::::: ::::::::::: :::::::::::
TGATCAGCGG TTTAAACTTA AGCTTGGTAC CGAGCTCGGA TCCACTTTGA CGTCAATGGG

::::::::: ::::::::::GCC AGGGAAGGTC TCCGTTCTTG CCGATGCCCA TATTCTGGGA(SEQ ID NO:249)
TGGAGTATTT ACGGTAAACC AGGGAAGGTC TCCGTTCTTG CCGATGCCCA TATTCTGGGA(SEQ ID NO:250)
```

FIGURE 54

Wild-type *FokI* cleavage half-domain

QLVKSELEEKKSELRHKLKYVPHEYIELIEIARNSTQDRILEMKVMEFFMKVYGYRGKH
LGGSRKPDGAIYTVGSPIDYGVIVDTKAYSGGYNLPIGQADEMQRYVEENQTRNKHINP
NEWWKVYPSSVTEFKFLFVSGHFKGNYKAQLTRLNHITNCNGAVLSVEELLIGGEMIKA
GTLTLEEVRKFNNGEINF (SEQ ID NO:251)

E490K

QLVKSELEEKKSELRHKLKYVPHEYIELIEIARNSTQDRILEMKVMEFFMKVYGYRGKH
LGGSRKPDGAIYTVGSPIDYGVIVDTKAYSGGYNLPIGQADEMQRYVKENQTRNKHINP
NEWWKVYPSSVTEFKFLFVSGHFKGNYKAQLTRLNHITNCNGAVLSVEELLIGGEMIKA
GTLTLEEVRKFNNGEINF (SEQ ID NO:252)

X2 (E490K:I538K)

QLVKSELEEKKSELRHKLKYVPHEYIELIEIARNSTQDRILEMKVMEFFMKVYGYRGKH
LGGSRKPDGAIYTVGSPIDYGVIVDTKAYSGGYNLPIGQADEMQRYVKENQTRNKHINP
NEWWKVYPSSVTEFKFLFVSGHFKGNYKAQLTRLNHKTNCNGAVLSVEELLIGGEMIKA
GTLTLEEVRKFNNGEINF (SEQ ID NO:253)

X3A (Q486E:E490K:I538K)

QLVKSELEEKKSELRHKLKYVPHEYIELIEIARNSTQDRILEMKVMEFFMKVYGYRGKH
LGGSRKPDGAIYTVGSPIDYGVIVDTKAYSGGYNLPIGQADEMERYVKENQTRNKHINP
NEWWKVYPSSVTEFKFLFVSGHFKGNYKAQLTRLNHKTNCNGAVLSVEELLIGGEMIKA
GTLTLEEVRKFNNGEINF (SEQ ID NO:254)

X3B (Q486I:E490K:I538K)

QLVKSELEEKKSELRHKLKYVPHEYIELIEIARNSTQDRILEMKVMEFFMKVYGYRGKH
LGGSRKPDGAIYTVGSPIDYGVIVDTKAYSGGYNLPIGQADEMIRYVKENQTRNKHINP
NEWWKVYPSSVTEFKFLFVSGHFKGNYKAQLTRLNHKTNCNGAVLSVEELLIGGEMIKA
GTLTLEEVRKFNNGEINF (SEQ ID NO:255)

TARGETED INTEGRATION AND EXPRESSION OF EXOGENOUS NUCLEIC ACID SEQUENCES

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] The present application claims the benefit of U.S. Provisional Application No. 60/702,394, filed Jul. 26, 2005 and U.S. Provisional Application No. 60/721,054, filed Sep. 26, 2005, both of which disclosures are hereby incorporated by reference in their entireties herein.

TECHNICAL FIELD

[0002] The present disclosure is in the fields of genome engineering, gene targeting, targeted chromosomal integration and protein expression.

BACKGROUND

[0003] A major area of interest in genome biology, especially in light of the determination of the complete nucleotide sequences of a number of genomes, is the targeted alteration of genome sequences. To provide but one example, sickle cell anemia is caused by mutation of a single nucleotide pair in the human β -globin gene. Thus, the ability to convert the endogenous genomic copy of this mutant nucleotide pair to the wild-type sequence in a stable fashion and produce normal β -globin would provide a cure for sickle cell anemia, as would introduction of a functional β -globin gene into a genome containing a mutant β -globin gene.

[0004] Attempts have been made to alter genomic sequences in cultured cells by taking advantage of the natural phenomenon of homologous recombination. See, for example, Capecchi (1989) *Science* 244:1288-1292; U.S. Pat. Nos. 6,528,313 and 6,528,314. If a polynucleotide has sufficient homology to the genomic region containing the sequence to be altered, it is possible for part or all of the sequence of the polynucleotide to replace the genomic sequence by homologous recombination. However, the frequency of homologous recombination under these circumstances is extremely low. Moreover, the frequency of insertion of the exogenous polynucleotide at genomic locations that lack sequence homology exceeds the frequency of homologous recombination by several orders of magnitude.

[0005] The introduction of a double-stranded break into genomic DNA, in the region of the genome bearing homology to an exogenous polynucleotide, has been shown to stimulate homologous recombination at this site by several thousand-fold in cultured cells. Rouet et al. (1994) *Mol. Cell. Biol.* 14:8096-8106; Chouluka et al. (1995) *Mol. Cell. Biol.* 15:1968-1973; Donoho et al. (1998) *Mol. Cell. Biol.* 18:4070-4078. See also Johnson et al. (2001) *Biochem. Soc. Trans.* 29:196-201; and Yanez et al. (1998) *Gene Therapy* 5:149-159. In these methods, DNA cleavage in the desired genomic region was accomplished by inserting a recognition site for a meganuclease (i.e., an endonuclease whose recognition sequence is so large that it does not occur, or occurs only rarely, in the genome of interest) into the desired genomic region.

[0006] However, meganuclease cleavage-stimulated homologous recombination relies on either the fortuitous presence of, or the directed insertion of, a suitable meganuclease recognition site in the vicinity of the genomic region

to be altered. Since meganuclease recognition sites are rare (or nonexistent) in a typical mammalian genome, and insertion of a suitable meganuclease recognition site is plagued with the same difficulties as associated with other genomic alterations, these methods are not broadly applicable.

[0007] Thus, there remain needs for compositions and methods for targeted alteration of sequences in any genome and for compositions and methods for targeted introduction of exogenous sequences into a genome.

SUMMARY

[0008] The present disclosure provides method and compositions for expressing the product of an exogenous nucleic acid sequence (i.e. a protein or a RNA molecule) in a cell. The exogenous nucleic acid sequence can comprise, for example, one or more genes or cDNA molecules, or any type of coding or noncoding sequence, and is introduced into the cell such that it is integrated into the genome of the cell in a predetermined region of interest. Integration of the exogenous nucleic acid sequence is facilitated by targeted double-strand cleavage of the genome in the region of interest. Cleavage is targeted to a particular site through the use of fusion proteins comprising a zinc finger binding domain, which can be engineered to bind any sequence of choice in the region of interest, and a cleavage domain or a cleavage half-domain. Such cleavage stimulates integration of exogenous polynucleotide sequences at or near the cleavage site. Said integration of exogenous sequences can proceed through both homology-dependent and homology-independent mechanisms.

[0009] Also provided are methods and compositions for modulating the expression of an endogenous cellular gene by targeted integration (either homology-dependent or homology-independent) of one or more exogenous sequences. Such exogenous sequences can include, for example, transcriptional control sequences such as promoters and enhancers. Modulation can include transcriptional activation (e.g., enhancement of transcription by, for example, insertion of a promoter and/or enhancer sequence) and transcriptional repression (e.g., functional "knock-out" by, for example, inserting an exogenous sequence into an endogenous transcriptional regulatory sequence, inserting a sequence facilitating transcriptional repression, or inserting a sequence that interrupts a coding region).

[0010] Also provided are methods and compositions for targeted insertion of an exogenous sequence into a genome, by either homology-dependent or homology-independent mechanisms, wherein the exogenous sequence does not express a product or modulate expression of an endogenous gene. For example, a recognition sequence for a sequence-specific DNA-cleaving enzyme can be introduced at a predetermined location in a genome so that targeted cleavage by the cleaving enzyme, at the predetermined location in the genome, can be accomplished. Exemplary DNA-cleaving enzymes include, but are not limited to, restriction enzymes, meganucleases and homing endonucleases.

[0011] In one aspect, disclosed herein is a method for expressing the product of an exogenous nucleic acid sequence in a cell, the method comprising: (a) expressing a first fusion protein in the cell, the first fusion protein comprising a first zinc finger binding domain and a first cleavage half-domain, wherein the first zinc finger binding

domain has been engineered to bind to a first target site in a region of interest in the genome of the cell; (b) expressing a second fusion protein in the cell, the second fusion protein comprising a second zinc finger binding domain and a second cleavage half domain, wherein the second zinc finger binding domain binds to a second target site in the region of interest in the genome of the cell, wherein the second target site is different from the first target site; and (c) contacting the cell with a polynucleotide comprising an exogenous nucleic acid sequence and a first nucleotide sequence that is homologous to a first sequence in the region of interest; wherein binding of the first fusion protein to the first target site, and binding of the second fusion protein to the second target site, positions the cleavage half-domains such that the genome of the cell is cleaved in the region of interest, thereby resulting in integration of the exogenous sequence into the genome of the cell in the region of interest and expression of the product of the exogenous sequence.

[0012] The exogenous nucleic acid sequence may comprise a cDNA and/or a promoter. In other embodiments, the exogenous nucleic acid sequence encodes a siRNA. The first nucleotide sequence may be identical to the first sequence in the region of interest.

[0013] In certain embodiments, the polynucleotide further comprises a second nucleotide sequence that is homologous to a second sequence in the region of interest. The second nucleotide sequence may be identical to the second sequence in the region of interest. Furthermore, in embodiments comprising first and second nucleotide sequences, the first nucleotide sequence may be identical to the first sequence in the region of interest and the second nucleotide sequence may be homologous but non-identical to a second sequence in the region of interest. In any of the methods described herein, the first and second nucleotide sequences flank the exogenous sequence.

[0014] In certain embodiments, the polynucleotide is a plasmid. In other embodiments, the polynucleotide is a linear DNA molecule.

[0015] In any of the methods described herein, the region of interest is in an accessible region of cellular chromatin, a chromosome, and/or a gene (e.g., a gene comprising a mutation such as a point mutation, a substitution, a deletion, an insertion, a duplication, an inversion and/or a translocation). In certain embodiments, the exogenous nucleic acid sequence comprises the wild-type sequence of the gene. In other embodiments, the exogenous nucleic acid sequence comprises a portion of the wild-type sequence of the gene. In still other embodiments, the exogenous nucleic acid sequence comprises a cDNA copy of a transcription product of the gene.

[0016] In any of the methods described herein, the region of interest is in a region of the genome that is not essential for viability. In other embodiments, the region of interest is in a region of the genome that is transcriptionally active. The region of interest is in a region of the genome that is transcriptionally active and not essential for viability (e.g., the human Rosa26 genome, the human homologue of the murine Rosa26 gene, or a CCR5 gene).

[0017] In another aspect, provided herein is a method for integrating an exogenous sequence into a region of interest in the genome of a cell, the method comprising: (a) express-

ing a first fusion protein in the cell, the first fusion protein comprising a first zinc finger binding domain and a first cleavage half-domain, wherein the first zinc finger binding domain has been engineered to bind to a first target site in a region of interest in the genome of the cell; (b) expressing a second fusion protein in the cell, the second fusion protein comprising a second zinc finger binding domain and a second cleavage half domain, wherein the second zinc finger binding domain binds to a second target site in the region of interest in the genome of the cell, wherein the second target site is different from the first target site; and (c) contacting the cell with a polynucleotide comprising an exogenous nucleic acid sequence; wherein binding of the first fusion protein to the first target site, and binding of the second fusion protein to the second target site, positions the cleavage half-domains such that the genome of the cell is cleaved in the region of interest, thereby resulting in integration of the exogenous sequence into the genome of the cell in the region of interest. In certain embodiments, the integration inactivates gene expression in the region of interest. The exogenous nucleic acid sequence may comprise, for example, a sequence of between 1 and 50 nucleotides in length. Furthermore, the exogenous nucleic acid sequence may encode a detectable amino acid sequence. The region of interest may be in an accessible region of cellular chromatin.

[0018] In any of the methods described herein, the first and second cleavage half-domains are from a Type IIS restriction endonuclease, for example, FokI or StlI. Furthermore, in any of the methods described herein, at least one of the fusion proteins may comprise an alteration in the amino acid sequence of the dimerization interface of the cleavage half-domain.

[0019] In any of the methods described herein, the cell can be a mammalian cell, for example, a human cell. Furthermore, the cell may be arrested in the G2 phase of the cell cycle.

[0020] The present subject matter thus includes, but is not limited to, the following embodiments:

[0021] 1. A method for expressing the product of an exogenous nucleic acid sequence in a cell, the method comprising:

[0022] (a) expressing a first fusion protein in the cell, the first fusion protein comprising a first zinc finger binding domain and a first cleavage half-domain, wherein the first zinc finger binding domain has been engineered to bind to a first target site in a region of interest in the genome of the cell;

[0023] (b) expressing a second fusion protein in the cell, the second fusion protein comprising a second zinc finger binding domain and a second cleavage half domain, wherein the second zinc finger binding domain binds to a second target site in the region of interest in the genome of the cell, wherein the second target site is different from the first target site; and

[0024] (c) contacting the cell with a polynucleotide comprising an exogenous nucleic acid sequence;

[0025] wherein binding of the first fusion protein to the first target site, and binding of the second fusion protein to the second target site, positions the cleavage half-domains such that the genome of the cell is cleaved in

the region of interest, thereby resulting in integration of the exogenous sequence into the genome of the cell in the region of interest and expression of the product of the exogenous sequence.

- [0026] 2. The method according to 1, wherein the exogenous nucleic acid sequence comprises a cDNA.
- [0027] 3. The method according to 1, wherein the exogenous sequence comprises a promoter.
- [0028] 4. The method according to 1, wherein the polynucleotide further comprises a first nucleotide sequence that is identical to a first sequence in the region of interest.
- [0029] 5. The method according to 4, wherein the polynucleotide further comprises a second nucleotide sequence that is identical to a second sequence in the region of interest.
- [0030] 6. The method according to 1, wherein the polynucleotide further comprises a first nucleotide sequence that is homologous but non-identical to a first sequence in the region of interest.
- [0031] 7. The method according to 6, wherein the polynucleotide further comprises a second nucleotide sequence that is homologous but non-identical to a second sequence in the region of interest.
- [0032] 8. The method according to 1, wherein the polynucleotide further comprises a first nucleotide sequence that is identical to a first sequence in the region of interest and a second nucleotide sequence that is homologous but non-identical to a second sequence in the region of interest.
- [0033] 9. The method according to 5, wherein the first and second nucleotide sequences flank the exogenous sequence.
- [0034] 10. The method according to 7, wherein the first and second nucleotide sequences flank the exogenous sequence.
- [0035] 11. The method according to 8, wherein the first and second nucleotide sequences flank the exogenous sequence.
- [0036] 12. The method of 1, wherein the polynucleotide is a plasmid.
- [0037] 13. The method of 1, wherein the polynucleotide is a linear DNA molecule.
- [0038] 14. The method according to 1, wherein the region of interest is in an accessible region of cellular chromatin.
- [0039] 15. The method of 1, wherein the region of interest is in a region of the genome that is not essential for viability.
- [0040] 16. The method of 1, wherein the region of interest is in a region of the genome that is transcriptionally active.
- [0041] 17. The method of 1, wherein the region of interest is in a region of the genome that is transcriptionally active and not essential for viability.

- [0042] 18. The method according to 17, wherein the region of interest is the human Rosa26 gene.
- [0043] 19. The method according to 17, wherein the region of interest is the human homologue of the murine Rosa26 gene.
- [0044] 20. The method according to 1, wherein the first and second cleavage half-domains are from a Type IIS restriction endonuclease.
- [0045] 21. The method according to 20, wherein the Type IIS restriction endonuclease is selected from the group consisting of FokI and StsI.
- [0046] 22. The method according to 1, wherein the region of interest is in a chromosome.
- [0047] 23. The method according to 1, wherein the region of interest comprises a gene.
- [0048] 24. The method according to 13, wherein the gene comprises a mutation.
- [0049] 25. The method according to 14, wherein the mutation is selected from the group consisting of a point mutation, a substitution, a deletion, an insertion, a duplication, an inversion and a translocation.
- [0050] 26. The method according to 24, wherein the exogenous nucleic acid sequence comprises the wild-type sequence of the gene.
- [0051] 27. The method according to 24, wherein the exogenous nucleic acid sequence comprises a portion of the wild-type sequence of the gene.
- [0052] 28. The method according to 24, wherein the exogenous nucleic acid sequence comprises a cDNA copy of a transcription product of the gene.
- [0053] 29. The method according to 1, wherein the exogenous nucleic acid sequence encodes a siRNA.
- [0054] 30. The method according to 1, wherein the cell is arrested in the G2 phase of the cell cycle.
- [0055] 31. The method according to 1, wherein at least one of the fusion proteins comprises an alteration in the amino acid sequence of the dimerization interface of the cleavage half-domain.
- [0056] 32. The method according to 1, wherein the cell is a mammalian cell.
- [0057] 33. The method according to 32, wherein the cell is a human cell.

BRIEF DESCRIPTION OF THE DRAWINGS

[0058] FIG. 1 shows the nucleotide sequence, in double-stranded form, of a portion of the human hSMC1L1 gene encoding the amino-terminal portion of the protein (SEQ ID NO: 1) and the encoded amino acid sequence (SEQ ID NO: 2). Target sequences for the hSMC1-specific ZFPs are underlined (one on each DNA strand).

[0059] FIG. 2 shows a schematic diagram of a plasmid encoding a ZFP-FokI fusion for targeted cleavage of the hSMC1 gene.

[0060] FIG. 3A-D show a schematic diagram of the hSMC1 gene. FIG. 3A shows a schematic of a portion of the

human X chromosome which includes the hSMC1 gene. FIG. 3B shows a schematic of a portion of the hSMC1 gene including the upstream region (left of +1), the first exon (between +1 and the right end of the arrow labeled "SMC1 coding sequence") and a portion of the first intron. Locations of sequences homologous to the initial amplification primers and to the chromosome-specific primer (see Table 3) are also provided. FIG. 3C shows the nucleotide sequence of the human X chromosome in the region of the SMC1 initiation codon (SEQ ID NO: 3), the encoded amino acid sequence (SEQ ID NO: 4), and the target sites for the SMC1-specific zinc finger proteins. FIG. 3D shows the sequence of the corresponding region of the donor molecule (SEQ ID NO: 5), with differences between donor and chromosomal sequences underlined. Sequences contained in the donor-specific amplification primer (Table 3) are indicated by double underlining.

[0061] FIG. 4 shows a schematic diagram of the hSMC1 donor construct.

[0062] FIG. 5 shows PCR analysis of DNA from transfected HEK293 cells. From left, the lanes show results from cells transfected with a plasmid encoding GFP (control plasmid), cells transfected with two plasmids, each of which encodes one of the two hSMC1-specific ZFP-FokI fusion proteins (ZFPs only), cells transfected with two concentrations of the hSMC1 donor plasmid (donor only), and cells transfected with the two ZFP-encoding plasmids and the donor plasmid (ZFPs+ donor). See Example 1 for details.

[0063] FIG. 6 shows the nucleotide sequence of an amplification product derived from a mutated hSMC1 gene (SEQ ID NO:6) generated by targeted homologous recombination. Sequences derived from the vector into which the amplification product was cloned are single-underlined, chromosomal sequences not present in the donor molecule are indicated by dashed underlining (nucleotides 32-97), sequences common to the donor and the chromosome are not underlined (nucleotides 98-394 and 402-417), and sequences unique to the donor are double-underlined (nucleotides 395-401). Lower-case letters represent sequences that differ between the chromosome and the donor.

[0064] FIG. 7 shows the nucleotide sequence of a portion of the human IL2R γ gene comprising the 3' end of the second intron and the 5' end of third exon (SEQ ID NO:7) and the amino acid sequence encoded by the displayed portion of the third exon (SEQ ID NO:8). Target sequences for the second pair of IL2R γ -specific ZFPs are underlined. See Example 2 for details.

[0065] FIG. 8 shows a schematic diagram of a plasmid encoding a ZFP-FokI fusion for targeted cleavage of IL2R γ gene.

[0066] FIG. 9A-D show a schematic diagram of the IL2R γ gene. FIG. 9A. shows a schematic of a portion of the human X chromosome which includes the IL2R γ gene. FIG. 9B shows a schematic of a portion of the IL2R γ gene including a portion of the second intron, the third exon and a portion of the third intron. Locations of sequences homologous to the initial amplification primers and to the chromosome-specific primer (see Table 5) are also provided. FIG. 9C shows the nucleotide sequence of the human X chromosome in the region of the third exon of the IL2R γ gene (SEQ ID NO: 9), the encoded amino acid sequence (SEQ ID NO: 10),

and the target sites for the first pair of IL2R γ -specific zinc finger proteins. FIG. 9D shows the sequence of the corresponding region of the donor molecule (SEQ ID NO: 11), with differences between donor and chromosomal sequences underlined. Sequences contained in the donor-specific amplification primer (Table 5) are indicated by double overlining.

[0067] FIG. 10 shows a schematic diagram of the IL2R γ donor construct.

[0068] FIG. 11 shows PCR analysis of DNA from transfected K652 cells. From left, the lanes show results from cells transfected with two plasmids, each of which encodes one of a pair of IL2R γ -specific ZFP-FokI fusion proteins (ZFPs only, lane 1), cells transfected with two concentrations of the IL2R γ donor plasmid (donor only, lanes 2 and 3), and cells transfected with the two ZFP-encoding plasmids and the donor plasmid (ZFPs+ donor, lanes 4-7). Each of the two pairs of IL2R γ -specific ZFP-FokI fusions were used (identified as "pair 1" and "pair 2") and use of both pairs resulted in production of the diagnostic amplification product (labeled "expected chimeric product" in the Figure). See Example 2 for details.

[0069] FIG. 12 shows the nucleotide sequence of an amplification product derived from a mutated IL2R γ gene (SEQ ID NO: 12) generated by targeted homologous recombination. Sequences derived from the vector into which the amplification product was cloned are single-underlined, chromosomal sequences not present in the donor molecule are indicated by dashed underlining (nucleotides 460-552), sequences common to the donor and the chromosome are not underlined (nucleotides 32-42 and 59-459), and a stretch of sequence containing nucleotides which distinguish donor sequences from chromosomal sequences is double-underlined (nucleotides 44-58). Lower-case letters represent nucleotides whose sequence differs between the chromosome and the donor.

[0070] FIG. 13 shows the nucleotide sequence of a portion of the human beta-globin gene encoding segments of the core promoter, the first two exons and the first intron (SEQ ID NO: 13). A missense mutation changing an A (in boldface and underlined) at position 5212541 on Chromosome 11 (BLAT, UCSC Genome Bioinformatics site) to a T results in sickle cell anemia. A first zinc finger/FokI fusion protein was designed such that the primary contacts were with the underlined 12-nucleotide sequence AAGGTGAACGTG (nucleotides 305-316 of SEQ ID NO: 13), and a second zinc finger/FokI fusion protein was designed such that the primary contacts were with the complement of the underlined 12-nucleotide sequence CCGTTACTGCCC (nucleotides 325-336 of SEQ ID NO:13).

[0071] FIG. 14 is a schematic diagram of a plasmid encoding ZFP-FokI fusion for targeted cleavage of the human beta globin gene.

[0072] FIG. 15 is a schematic diagram of the cloned human beta globin gene showing the upstream region, first and second exons, first intron and primer binding sites.

[0073] FIG. 16 is a schematic diagram of the beta globin donor construct, pCR4-TOPO-HBBdonor.

[0074] FIG. 17 shows PCR analysis of DNA from cells transfected with two pairs of β -globin-specific ZFP

nucleases and a beta globin donor plasmid. The panel on the left is a loading control in which the initial amp 1 and initial amp 2 primers (Table 7) were used for amplification. In the experiment shown in the right panel, the “chromosome-specific and “donor-specific” primers (Table 7) were used for amplification. The leftmost lane in each panel contains molecular weight markers and the next lane shows amplification products obtained from mock-transfected cells. Remaining lanes, from left to right, show amplification product from cells transfected with: a GFP-encoding plasmid, 100 ng of each ZFP/FokI-encoding plasmid, 200 ng of each ZFP/FokI-encoding plasmid, 200 ng donor plasmid, 600 ng donor plasmid, 200 ng donor plasmid +100 ng of each ZFP/FokI-encoding plasmid, and 600 ng donor plasmid +200 ng of each ZFP/FokI-encoding plasmid.

[0075] FIG. 18 shows the nucleotide sequence of an amplification product derived from a mutated beta-globin gene (SEQ ID NO: 14) generated by targeted homologous recombination. Chromosomal sequences not present in the donor molecule are indicated by dashed underlining (nucleotides 1-72), sequences common to the donor and the chromosome are not underlined (nucleotides 73-376), and a stretch of sequence containing nucleotides which distinguish donor sequences from chromosomal sequences is double-underlined (nucleotides 377-408). Lower-case letters represent nucleotides whose sequence differs between the chromosome and the donor.

[0076] FIG. 19 shows the nucleotide sequence of a portion of the fifth exon of the Interleukin-2 receptor gamma chain (IL-2R γ) gene (SEQ ID NO: 15). Also shown (underlined) are the target sequences for the 5-8 and 5-10 ZFP/FokI fusion proteins. See Example 5 for details.

[0077] FIG. 20 shows the amino acid sequence of the 5-8 ZFP/FokI fusion targeted to exon 5 of the human IL-2R γ gene (SEQ ID NO: 16). Amino acid residues 1-17 contain a nuclear localization sequence (NLS, underlined); residues 18-130 contain the ZFP portion, with the recognition regions of the component zinc fingers shown in boldface; the ZFP-FokI linker (ZC linker, underlined) extends from residues 131 to 140 and the FokI cleavage half-domain begins at residue 141 and extends to the end of the protein at residue 336. The residue that was altered to generate the Q486E mutation is shown underlined and in boldface.

[0078] FIG. 21 shows the amino acid sequence of the 5-10 ZFP/FokI fusion targeted to exon 5 of the human IL-2R γ gene (SEQ ID NO: 17). Amino acid residues 1-17 contain a nuclear localization sequence (NLS, underlined); residues 18-133 contain the ZFP portion, with the recognition regions of the component zinc fingers shown in boldface; the ZFP-FokI linker (ZC linker, underlined) extends from residues 134 to 143 and the FokI cleavage half-domain begins at residue 144 and extends to the end of the protein at residue 339. The residue that was altered to generate the E490K mutation is shown underlined and in boldface.

[0079] FIG. 22 shows the nucleotide sequence of the enhanced Green Fluorescent Protein gene (SEQ ID NO: 18) derived from the *Aequorea victoria* GFP gene (Tsien (1998) *Ann. Rev. Biochem.* 67:509-544). The ATG initiation codon, as well as the region which was mutagenized, are underlined.

[0080] FIG. 23 shows the nucleotide sequence of a mutant defective eGFP gene (SEQ ID NO: 19). Binding sites for

ZFP-nucleases are underlined and the region between the binding sites corresponds to the region that was modified.

[0081] FIG. 24 shows the structures of plasmids encoding Zinc Finger Nucleases targeted to the eGFP gene.

[0082] FIG. 25 shows an autoradiogram of a 10% acrylamide gel used to analyze targeted DNA cleavage of a mutant eGFP gene by zinc finger endonucleases. See Example 8 for details.

[0083] FIG. 26 shows the structure of plasmid pcDNA4/TO/GFPmut (see Example 9).

[0084] FIG. 27 shows levels of eGFPmut mRNA, normalized to GAPDH mRNA, in various cell lines obtained from transfection of human HEK293 cells. Light bars show levels in untreated cells; dark bars show levels in cell that had been treated with 2 ng/ml doxycycline. See Example 9 for details.

[0085] FIG. 28 shows the structure of plasmid pCR(R)4-TOPO-GFPdonor5. See Example 10 for details.

[0086] FIG. 29 shows the nucleotide sequence of the eGFP insert in pCR(R)4-TOPO-GFPdonor5 (SEQ ID NO:20). The insert contains sequences encoding a portion of a non-modified enhanced Green Fluorescent Protein, lacking an initiation codon. See Example 10 for details.

[0087] FIG. 30 shows a FACS trace of T18 cells transfected with plasmids encoding two ZFP nucleases and a plasmid encoding a donor sequence, that were arrested in the G2 phase of the cell cycle 24 hours post-transfection with 100 ng/ml nocodazole for 48 hours. The medium was replaced and the cells were allowed to recover for an additional 48 hours, and gene correction was measured by FACS analysis. See Example 11 for details.

[0088] FIG. 31 shows a FACS trace of T18 cells transfected with plasmids encoding two ZFP nucleases and a plasmid encoding a donor sequence, that were arrested in the G2 phase of the cell cycle 24 hours post-transfection with 0.2 μ M vinblastine for 48 hours. The medium was replaced and the cells were allowed to recover for an additional 48 hours, and gene correction was measured by FACS analysis. See Example 11 for details.

[0089] FIG. 32 shows the nucleotide sequence of a 1,527 nucleotide eGFP insert in pCR(R)4-TOPO (SEQ ID NO:21). The sequence encodes a non-modified enhanced Green Fluorescent Protein lacking an initiation codon. See, Example 13 for details.

[0090] FIG. 33 shows a schematic diagram of an assay used to measure the frequency of editing of the endogenous human IL-2R γ gene. See, Example 14 for details.

[0091] FIG. 34 shows autoradiograms of acrylamide gels used in an assay to measure the frequency of editing of an endogenous cellular gene by targeted cleavage and homologous recombination. The lane labeled “GFP” shows assay results from a control in which cells were transfected with an eGFP-encoding vector; the lane labeled “ZFPs only” shows results from another control experiment in which cells were transfected with the two ZFP/nuclease-encoding plasmids (50 ng of each) but not with a donor sequence. Lanes labeled “donor only” show results from a control experiment in which cells were transfected with 1 μ g of donor plasmid but not with the ZFP/nuclease-encoding plasmids. In the experimental lanes, 50Z refers to cells transfected with 50 ng of

each ZFP/nuclease expression plasmid, 100Z refers to cells transfected with 100 ng of each ZFP/nuclease expression plasmid, 0.5D refers to cells transfected with 0.5 μ g of the donor plasmid, and 1D refers to cells transfected with 1.0 μ g of the donor plasmid. “+” refers to cells that were exposed to 0.2 μ M vinblastine; “-” refer to cells that were not exposed to vinblastine. “wt” refers to the fragment obtained after BsrBI digestion of amplification products obtained from chromosomes containing the wild-type chromosomal IL-2R γ gene; “rflp” refers to the two fragments (of approximately equal molecular weight) obtained after BsrBI digestion of amplification products obtained from chromosomes containing sequences from the donor plasmid which had integrated by homologous recombination.

[0092] FIG. 35 shows an autoradiographic image of a four-hour exposure of a gel used in an assay to measure targeted recombination at the human IL-2R γ locus in K562 cells. “wt” identifies a band that is diagnostic for chromosomal DNA containing the native K562 IL-2R γ sequence; “rflp” identifies a doublet diagnostic for chromosomal DNA containing the altered IL-2R γ sequence present in the donor DNA molecule. The symbol “+” above a lane indicates that cells were treated with 0.2 μ M vinblastine; the symbol “-” indicates that cells were not treated with vinblastine. The numbers in the “ZFP+ donor” lanes indicate the percentage of total chromosomal DNA containing sequence originally present in the donor DNA molecule, calculated using the “peak finder, automatic baseline” function of Molecular Dynamics’ ImageQuant v. 5.1 software as described in Ch. 8 of the manufacturer’s manual (Molecular Dynamics ImageQuant User’s Guide; part 218-415). “Untr” indicates untransfected cells. See Example 15 for additional details.

[0093] FIG. 36 shows an autoradiographic image of a four-hour exposure of a gel used in an assay to measure targeted recombination at the human IL-2R γ locus in K562 cells. “wt” identifies a band that is diagnostic for chromosomal DNA containing the native K562 IL-2R γ sequence; “rflp” identifies a band that is diagnostic for chromosomal DNA containing the altered IL-2R γ sequence present in the donor DNA molecule. The symbol “+” above a lane indicates that cells were treated with 0.2 μ M vinblastine; the symbol “-” indicates that cells were not treated with vinblastine. The numbers beneath the “ZFP+ donor” lanes indicate the percentage of total chromosomal DNA containing sequence originally present in the donor DNA molecule, calculated as described in Example 35. See Example 15 for additional details.

[0094] FIG. 37 shows an autoradiogram of a four-hour exposure of a DNA blot probed with a fragment specific to the human IL-2R γ gene. The arrow to the right of the image indicates the position of a band corresponding to genomic DNA whose sequence has been altered by homologous recombination. The symbol “+” above a lane indicates that cells were treated with 0.2 μ M vinblastine; the symbol “-” indicates that cells were not treated with vinblastine. The numbers beneath the “ZFP+ donor” lanes indicate the percentage of total chromosomal DNA containing sequence originally present in the donor DNA molecule, calculated as described in Example 35. See Example 15 for additional details.

[0095] FIG. 38 shows autoradiographic images of gels used in an assay to measure targeted recombination at the

human IL-2R γ locus in CD34⁺ human bone marrow cells. The left panel shows a reference standard in which the stated percentage of normal human genomic DNA (containing a MaeII site) was added to genomic DNA from Jurkat cells (lacking a MaeII site), the mixture was amplified by PCR to generate a radiolabelled amplification product, and the amplification product was digested with MaeII. “wt” identifies a band representing undigested DNA, and “rflp” identifies a band resulting from MaeII digestion.

[0096] The right panel shows results of an experiment in which CD34⁺ cells were transfected with donor DNA containing a BsrBI site and plasmids encoding zinc finger-FokI fusion endonucleases. The relevant genomic region was then amplified and labeled, and the labeled amplification product was digested with BsrBI. “GFP” indicates control cells that were transfected with a GFP-encoding plasmid; “Donor only” indicates control cells that were transfected only with donor DNA, and “ZFP+ Donor” indicates cells that were transfected with donor DNA and with plasmids encoding the zinc finger/FokI nucleases. “wt” identifies a band that is diagnostic for chromosomal DNA containing the native IL-2R γ sequence; “rflp” identifies a band that is diagnostic for chromosomal DNA containing the altered IL-2R γ sequence present in the donor DNA molecule. The rightmost lane contains DNA size markers. See Example 16 for additional details.

[0097] FIG. 39 shows an image of an immunoblot used to test for Ku70 protein levels in cells transfected with Ku70-targeted siRNA. The T7 cell line (Example 9, FIG. 27) was transfected with two concentrations each of siRNA from two different siRNA pools (see Example 18). Lane 1: 70 ng of siRNA pool D; Lane 2: 140 ng of siRNA pool D; Lane 3: 70 ng of siRNA pool E; Lane 4: 140 ng of siRNA pool E. “Ku70” indicates the band representing the Ku70 protein; “TFIIB” indicates a band representing the TFIIB transcription factor, used as a control.

[0098] FIG. 40 shows the amino acid sequences of four zinc finger domains targeted to the human β -globin gene: sca-29b (SEQ ID NO:22); sca-36a (SEQ ID NO:23); sca-36b (SEQ ID NO:24) and sca-36c (SEQ ID NO:25). The target site for the sca-29b domain is on one DNA strand, and the target sites for the sca-36a, sca-36b and sca-36c domains are on the opposite strand. See Example 20.

[0099] FIG. 41 shows results of an in vitro assay, in which different combinations of zinc finger-FokI fusion nucleases (ZFNs) were tested for sequence-specific DNA cleavage. The lane labeled “U” shows a sample of the DNA template. The next four lanes show results of incubation of the DNA template with each of four β -globin-targeted ZFNs (see Example 20 for characterization of these ZFNs). The rightmost three lanes show results of incubation of template DNA with the sca-29b ZFN and one of the sca-36a, sca-36b or sca-36c ZFNs (all of which are targeted to the strand opposite that to which sca-29b is targeted).

[0100] FIG. 42 shows levels of eGFP mRNA in T18 cells (bars) as a function of doxycycline concentration (provided on the abscissa). The number above each bar represents the percentage correction of the eGFP mutation, in cells transfected with donor DNA and plasmids encoding eGFP-targeted zinc finger nucleases, as a function of doxycycline concentration.

[0101] FIG. 43A-C show schematic diagrams of different fusion protein configurations. FIG. 43A shows two fusion

proteins, in which the zinc finger domain is nearest the N-terminus and the FokI cleavage half-domain is nearest the C-terminus, binding to DNA target sites on opposite strands whose 5' ends are proximal to each other. FIG. 43B shows two fusion proteins, in which the FokI cleavage half-domain is nearest the N-terminus and the zinc finger domain is nearest the C-terminus, binding to DNA target sites on opposite strands whose 3' ends are proximal to each other. FIG. 43C shows a first protein in which the FokI cleavage half-domain is nearest the N-terminus and the zinc finger domain is nearest the C-terminus, binding to DNA target sites on the same strand, in which the target site for the first protein is upstream (i.e. to the 5' side) of the binding site for the second protein.

[0102] In all examples, three-finger proteins are shown binding to nine-nucleotide target sites. 5' and 3' polarity of the DNA strands is shown, and the N-termini of the fusion proteins are identified.

[0103] FIG. 44 is an autoradiogram of an acrylamide gel in which cleavage of a model substrate by zinc finger endonucleases was assayed. Lane 1 shows the migration of uncleaved substrate. Lane 2 shows substrate after incubation with the IL2-1R zinc finger/FokI fusion protein. Lane 3 shows substrate after incubation with the 5-9DR zinc finger/FokI fusion protein. Lane 4 shows substrate after incubation with both proteins. Approximate sizes (in base pairs) of the substrate and its cleavage products are shown to the right of the image. Below the image, the nucleotide sequence (SEQ ID NO:211) of the portion of the substrate containing the binding sites for the 5-9D and IL2-1 zinc finger binding domains is shown. The binding sites are identified and indicated by underlining.

[0104] FIG. 45 is an autoradiogram of an acrylamide gel in which cleavage of a model substrate by zinc finger endonucleases was assayed. Lane 1 shows the migration of uncleaved substrate. Lane 2 shows substrate after incubation with the IL2-1C zinc finger/FokI fusion protein. Lane 3 shows substrate after incubation with the IL2-1R zinc finger/FokI fusion protein. Lane 4 shows substrate after incubation with the 5-9DR zinc finger/FokI fusion protein. Lane 5 shows substrate after incubation with both the IL2-1R and 5-9DR fusion proteins. Lane 6 shows substrate after incubation with both the IL2-1C and 5-9DR proteins. Approximate sizes (in base pairs) of the substrate and its cleavage products are shown to the right of the image. Below the image, the nucleotide sequence (SEQ ID NO:212) of the portion of the substrate containing the binding sites for the 5-9D and IL2-1 zinc finger binding domains is shown. The binding sites are identified and indicated by underlining.

[0105] FIG. 46 is a schematic diagram of a plasmid containing mutant eGFP coding sequences containing an insertion of sequences from exon 5 of the IL-2R γ gene. See Example 29 for details.

[0106] FIG. 47 shows an autoradiographic image of a gel in which amplification products of DNA from transfected K562 cells were incubated with the restriction enzyme Stu I. Headings above the lane indicate DNA from cells transfected with a GFP-encoding plasmid (GFP); DNA from cells transfected with a vector encoding the 5-8G and 5-9D ZFP/FokI fusion proteins (ZFNs); DNA from cells trans-

fectected with a plasmid containing a 12-nucleotide pair exogenous sequence (including a StuI recognition site) flanked on either side by 750 nucleotide pairs of sequence homologous to exon 5 of the IL-2R γ gene, wherein the two exon 5-homologous sequences are adjacent to one another in the wild-type IL-2R γ gene (donor); and DNA from cells transfected with a vector encoding the 5-8G and 5-9D ZFP/FokI fusion proteins and a plasmid containing a 12-nucleotide pair exogenous sequence (including a StuI recognition site) flanked on either side by 750 nucleotide pairs of sequence homologous to exon 5 and adjacent regions of the IL-2R γ gene, wherein the two exon 5-homologous sequences are adjacent to one another in the wild-type IL-2R γ gene (ZFNs+ donor). Bands arising from chromosomes containing wild-type IL-2R sequences ("WT") and chromosomes into which exogenous sequences have been integrated ("patch") are indicated. The rightmost lane contains molecular weight markers. See also Example 33.

[0107] FIG. 48 shows images of gels in which amplification products of DNA from transfected K562 cells were analyzed. Headings above the lane indicate DNA from cells transfected with a vector encoding the 5-8G and 5-9D ZFP/FokI fusion proteins (ZFNs); DNA from cells transfected with a plasmid containing a 720 nucleotide pair open reading frame encoding eGFP (donor 1); DNA from cells transfected with a plasmid containing a 924 nucleotide pair sequence that included an eGFP open reading frame and a downstream polyadenylation signal (donor 2); DNA from cells transfected with a vector encoding the 5-8G and 5-9D ZFP/FokI fusion proteins and with a plasmid containing a 720 nucleotide pair open reading frame encoding eGFP (ZFNs +donor 1) and DNA from cells transfected with a vector encoding the 5-8G and 5-9D ZFP/FokI fusion proteins and with a plasmid containing a 924 nucleotide pair sequence that included an eGFP open reading frame and a downstream polyadenylation signal (ZFNs+ donor 2). The leftmost and rightmost lanes of the top panel contains molecular weight markers. The top panel is a photograph of an ethidium bromide-stained gel; the bottom panel shows an autoradiogram of a gel from a separate experiment in which labeled amplification products were analyzed. See also Example 34.

[0108] FIG. 49 is a schematic diagram describing an experiment in which a "therapeutic half-gene" was introduced into the endogenous human IL-2R γ gene. The top line represents chromosomal IL-2R γ sequences, and the middle line represents donor sequences, with exons indicated by boxes and introns by horizontal lines. Numbers inside the boxes identify the exons of the IL-2R γ gene, with "5" representing the fifth exon of the chromosomal IL-2R gene, "5u" representing the upstream portion of the fifth exon, "5d" representing the downstream portion of the fifth exon and "5d(m)" representing the downstream portion of the fifth exon containing several silent sequence changes (i.e., changes that do not alter the encoded amino acid sequence). Diagonal lines demarcate regions of homology between donor and chromosomal sequences. The bottom line shows the expected product of homologous recombination, in which exons 5d(m), 6, 7, and 8 are inserted within the fifth exon of the chromosomal gene. See also Example 35.

[0109] FIG. 50 shows an autoradiogram of a gel in which amplification products of DNA from transfected K562 cells were analyzed. Headings above the lane indicate DNA from

control cells transfected with a vector encoding green fluorescent protein ("GFP") and from experimental cells transfected with a vector encoding the 5-8G and 5-9D ZFP/FokI fusion proteins and a plasmid containing a 720-nucleotide cDNA construct containing part of exon 5 and exons 6, 7 and 8 of the IL-2R γ gene, flanked on either side by 750 nucleotide pairs of sequence homologous to exon 5 and surrounding regions of the IL-2R γ gene, wherein the two exon 5-homologous sequences are adjacent to one another in the wild-type IL-2R γ gene (ZFNs+ donor). Bands arising from chromosomes containing wild-type IL-2R sequences ("WT") and chromosomes into which exogenous sequences have been integrated ("ORF") are indicated. See also Example 35.

[0110] FIG. 51 shows the design and results of an experiment in which a 7.7 kbp antibody expression construct was inserted into the endogenous chromosomal IL-2R γ gene. The upper portion of the figure is a schematic diagram showing the result of homology-dependent targeted integration of a 7.7 kilobase pair expression construct (shaded) into exon 5 of the endogenous chromosomal IL-2R γ gene. Arrows indicate the locations and polarities of amplification primers used to detect the junctions between exogenous and endogenous sequences that result from targeted integration.

[0111] The lower portion of the figure shows photographs of ethidium bromide-stained gels in which amplification products were analyzed. The left panel shows products of cellular DNA that was amplified using primers that detect the upstream junction (Primer set A) and the right panel shows products of cellular DNA that was amplified using primers that detect the downstream junction (Primer set B). DNA samples used as templates for amplification are identified below the gel, as follows: DNA from cells transfected with a vector encoding green fluorescent protein (GFP); DNA from cells transfected only with the donor DNA molecule containing the 7.7 kbp expression construct (don.); DNA from cells transfected with a vector encoding the 5-8G and 5-9D ZFP/FokI fusion proteins and with the donor DNA molecule (ZFNs+ donor). The topology of the donor DNA (circular or linear) is also indicated. See also Example 36.

[0112] FIG. 52 is an autoradiogram of a gel in which amplification products from the CHO DHFR gene were analyzed for mismatches using a Cel-1 assay. Amplification products were obtained from wild-type CHO cell DNA (W) or DNA from CHO cells that had been treated with zinc finger nucleases (Mu), and were then exposed to Cel-1 nuclease (+) or not (-), as indicated above the gel. To the right of the gel, bands indicative of wild-type DHFR sequences (WT) and mutant DHFR sequences containing a 157-nucleotide insertion (Mutant) are indicated. See Example 37 for details.

[0113] FIG. 53 shows a portion of the nucleotide sequence of the CHO dihydrofolate reductase (DHFR) gene (upper lines) and a portion of the nucleotide sequence of a mutant DHFR gene generated by targeted homology-independent integration of exogenous sequences (lower lines). Target sequences for the zinc finger nucleases described in Table 28 are boxed; changes from wild-type sequence are underlined. See also Example 37.

[0114] FIG. 54 shows the amino acid sequences of the wild-type FokI cleavage half-domain and of several mutant cleavage half-domains containing alterations in the amino

acid sequence of the dimerization interface. Positions at which the sequence was altered (amino acids 486, 490 and 538) are underlined.

DETAILED DESCRIPTION

[0115] Disclosed herein are compositions and methods useful for targeted cleavage of cellular chromatin and for targeted alteration of a cellular nucleotide sequence, e.g., by targeted cleavage followed by non-homologous end joining (with or without an exogenous sequence inserted therebetween) or by targeted cleavage followed by homologous recombination between an exogenous polynucleotide (comprising one or more regions of homology with the cellular nucleotide sequence) and a genomic sequence. Genomic sequences include those present in chromosomes, episomes, organellar genomes (e.g., mitochondria, chloroplasts), artificial chromosomes and any other type of nucleic acid present in a cell such as, for example, amplified sequences, double minute chromosomes and the genomes of endogenous or infecting bacteria and viruses. Genomic sequences can be normal (i.e., wild-type) or mutant; mutant sequences can comprise, for example, insertions, deletions, translocations, rearrangements, and/or point mutations. A genomic sequence can also comprise one of a number of different alleles.

[0116] Compositions useful for targeted cleavage and recombination include fusion proteins comprising a cleavage domain (or a cleavage half-domain) and a zinc finger binding domain, polynucleotides encoding these proteins and combinations of polypeptides and polypeptide-encoding polynucleotides. A zinc finger binding domain can comprise one or more zinc fingers (e.g., 2, 3, 4, 5, 6, 7, 8, 9 or more zinc fingers), and can be engineered to bind to any genomic sequence. Thus, by identifying a target genomic region of interest at which cleavage or recombination is desired, one can, according to the methods disclosed herein, construct one or more fusion proteins comprising a cleavage domain (or cleavage half-domain) and a zinc finger domain engineered to recognize a target sequence in said genomic region. The presence of such a fusion protein (or proteins) in a cell will result in binding of the fusion protein(s) to its (their) binding site(s) and cleavage within or near said genomic region. Moreover, if an exogenous polynucleotide homologous to the genomic region is also present in such a cell, homologous recombination occurs at a high rate between the genomic region and the exogenous polynucleotide.

General

[0117] Practice of the methods, as well as preparation and use of the compositions disclosed herein employ, unless otherwise indicated, conventional techniques in molecular biology, biochemistry, chromatin structure and analysis, computational chemistry, cell culture, recombinant DNA and related fields as are within the skill of the art. These techniques are fully explained in the literature. See, for example, Sambrook et al. *MOLECULAR CLONING: A LABORATORY MANUAL*, Second edition, Cold Spring Harbor Laboratory Press, 1989 and Third edition, 2001; Ausubel et al., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John Wiley & Sons, New York, 1987 and periodic updates; the series *METHODS IN ENZYMOLOGY*, Academic Press, San Diego; Wolffe, *CHROMATIN*

STRUCTURE AND FUNCTION, Third edition, Academic Press, San Diego, 1998; METHODS IN ENZYMOLOGY, Vol. 304, "Chromatin" (P.M. Wassarman and A. P. Wolffe, eds.), Academic Press, San Diego, 1999; and METHODS IN MOLECULAR BIOLOGY, Vol. 119, "Chromatin Protocols" (P.B. Becker, ed.) Humana Press, Totowa, 1999.

Definitions

[0118] The terms "nucleic acid," "polynucleotide," and "oligonucleotide" are used interchangeably and refer to a deoxyribonucleotide or ribonucleotide polymer, in linear or circular conformation, and in either single- or double-stranded form. For the purposes of the present disclosure, these terms are not to be construed as limiting with respect to the length of a polymer. The terms can encompass known analogues of natural nucleotides, as well as nucleotides that are modified in the base, sugar and/or phosphate moieties (e.g., phosphorothioate backbones). In general, an analogue of a particular nucleotide has the same base-pairing specificity; i.e., an analogue of A will base-pair with T.

[0119] The terms "polypeptide," "peptide" and "protein" are used interchangeably to refer to a polymer of amino acid residues. The term also applies to amino acid polymers in which one or more amino acids are chemical analogues or modified derivatives of a corresponding naturally-occurring amino acids.

[0120] "Binding" refers to a sequence-specific, non-covalent interaction between macromolecules (e.g., between a protein and a nucleic acid). Not all components of a binding interaction need be sequence-specific (e.g., contacts with phosphate residues in a DNA backbone), as long as the interaction as a whole is sequence-specific. Such interactions are generally characterized by a dissociation constant (K_d) of 10^{-6} M⁻¹ or lower. "Affinity" refers to the strength of binding: increased binding affinity being correlated with a lower K_d .

[0121] A "binding protein" is a protein that is able to bind non-covalently to another molecule. A binding protein can bind to, for example, a DNA molecule (a DNA-binding protein), an RNA molecule (an RNA-binding protein) and/or a protein molecule (a protein-binding protein). In the case of a protein-binding protein, it can bind to itself (to form homodimers, homotrimers, etc.) and/or it can bind to one or more molecules of a different protein or proteins. A binding protein can have more than one type of binding activity. For example, zinc finger proteins have DNA-binding, RNA-binding and protein-binding activity.

[0122] A "zinc finger DNA binding protein" (or binding domain) is a protein, or a domain within a larger protein, that binds DNA in a sequence-specific manner through one or more zinc fingers, which are regions of amino acid sequence within the binding domain whose structure is stabilized through coordination of a zinc ion. The term zinc finger DNA binding protein is often abbreviated as zinc finger protein or ZFP.

[0123] Zinc finger binding domains can be "engineered" to bind to a predetermined nucleotide sequence. Non-limiting examples of methods for engineering zinc finger proteins are design and selection. A designed zinc finger protein is a protein not occurring in nature whose design/composition results principally from rational criteria. Rational criteria for design include application of substitution rules and comput-

erized algorithms for processing information in a database storing information of existing ZFP designs and binding data. See, for example, U.S. Pat. Nos. 6,140,081; 6,453,242; and 6,534,261; see also WO 98/53058; WO 98/53059; WO 98/53060; WO 02/016536 and WO 03/016496.

[0124] A "selected" zinc finger protein is a protein not found in nature whose production results primarily from an empirical process such as phage display, interaction trap or hybrid selection. See e.g., US 5,789,538; US 5,925,523; US 6,007,988; US 6,013,453; US 6,200,759; WO 95/19431; WO 96/06166; WO 98/53057; WO 98/54311; WO 00/27878; WO 01/60970 WO 01/88197 and WO 02/099084.

[0125] The term "sequence" refers to a nucleotide sequence of any length, which can be DNA or RNA; can be linear, circular or branched and can be either single-stranded or double stranded. The term "donor sequence" refers to a nucleotide sequence that is inserted into a genome. A donor sequence can be of any length, for example between 2 and 10,000 nucleotides in length (or any integer value therebetween or thereabove), preferably between about 100 and 1,000 nucleotides in length (or any integer therebetween), more preferably between about 200 and 500 nucleotides in length.

[0126] A "homologous, non-identical sequence" refers to a first sequence which shares a degree of sequence identity with a second sequence, but whose sequence is not identical to that of the second sequence. For example, a polynucleotide comprising the wild-type sequence of a mutant gene is homologous and non-identical to the sequence of the mutant gene. In certain embodiments, the degree of homology between the two sequences is sufficient to allow homologous recombination therebetween, utilizing normal cellular mechanisms. Two homologous non-identical sequences can be any length and their degree of non-homology can be as small as a single nucleotide (e.g., for correction of a genomic point mutation by targeted homologous recombination) or as large as 10 or more kilobases (e.g., for insertion of a gene at a predetermined ectopic site in a chromosome). Two polynucleotides comprising the homologous non-identical sequences need not be the same length. For example, an exogenous polynucleotide (i.e., donor polynucleotide) of between 20 and 10,000 nucleotides or nucleotide pairs can be used.

[0127] Techniques for determining nucleic acid and amino acid sequence identity are known in the art. Typically, such techniques include determining the nucleotide sequence of the mRNA for a gene and/or determining the amino acid sequence encoded thereby, and comparing these sequences to a second nucleotide or amino acid sequence. Genomic sequences can also be determined and compared in this fashion. In general, identity refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Two or more sequences (polynucleotide or amino acid) can be compared by determining their percent identity. The percent identity of two sequences, whether nucleic acid or amino acid sequences, is the number of exact matches between two aligned sequences divided by the length of the shorter sequences and multiplied by 100. An approximate alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2:482-489 (1981). This algorithm can

be applied to amino acid sequences by using the scoring matrix developed by Dayhoff, *Atlas of Protein Sequences and Structure*, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA, and normalized by Gribskov, *Nucl. Acids Res.* 14(6):6745-6763 (1986). An exemplary implementation of this algorithm to determine percent identity of a sequence is provided by the Genetics Computer Group (Madison, Wis.) in the "BestFit" utility application. The default parameters for this method are described in the Wisconsin Sequence Analysis Package Program Manual, Version 8 (1995) (available from Genetics Computer Group, Madison, Wis.). A preferred method of establishing percent identity in the context of the present disclosure is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrock, and distributed by IntelliGenetics, Inc. (Mountain View, Calif.). From this suite of packages the Smith-Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects sequence identity. Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is BLAST, used with default parameters. For example, BLASTN and BLASTP can be used using the following default parameters: genetic code=standard; filter=none; strand=both; cutoff=60; expect=10; Matrix=BLOSUM62; Descriptions=50 sequences; sort by=HIGH SCORE; Databases=non-redundant, GenBank+EMBL+DDBJ+PDB+GenBank CDS translations+Swiss protein+Spupdate+PIR. Details of these programs can be found on the internet. With respect to sequences described herein, the range of desired degrees of sequence identity is approximately 80% to 100% and any integer value therebetween. Typically the percent identities between sequences are at least 70-75%, preferably 80-82%, more preferably 85-90%, even more preferably 92%, still more preferably 95%, and most preferably 98% sequence identity.

[0128] Alternatively, the degree of sequence similarity between polynucleotides can be determined by hybridization of polynucleotides under conditions that allow formation of stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s), and size determination of the digested fragments. Two nucleic acid, or two polypeptide sequences are substantially homologous to each other when the sequences exhibit at least about 70%-75%, preferably 80%-82%, more preferably 85%-90%, even more preferably 92%, still more preferably 95%, and most preferably 98% sequence identity over a defined length of the molecules, as determined using the methods above. As used herein, substantially homologous also refers to sequences showing complete identity to a specified DNA or polypeptide sequence. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *Nucleic Acid Hybridization: A Practical Approach*, editors B. D. Hames and S. J. Higgins, (1985) Oxford; Washington, D.C.; IRL Press).

[0129] Selective hybridization of two nucleic acid fragments can be determined as follows. The degree of sequence identity between two nucleic acid molecules affects the efficiency and strength of hybridization events between such molecules. A partially identical nucleic acid sequence will at least partially inhibit the hybridization of a completely identical sequence to a target molecule. Inhibition of hybridization of the completely identical sequence can be assessed using hybridization assays that are well known in the art (e.g., Southern (DNA) blot, Northern (RNA) blot, solution hybridization, or the like, see Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, (1989) Cold Spring Harbor, N.Y.). Such assays can be conducted using varying degrees of selectivity, for example, using conditions varying from low to high stringency. If conditions of low stringency are employed, the absence of non-specific binding can be assessed using a secondary probe that lacks even a partial degree of sequence identity (for example, a probe having less than about 30% sequence identity with the target molecule), such that, in the absence of non-specific binding events, the secondary probe will not hybridize to the target.

[0130] When utilizing a hybridization-based detection system, a nucleic acid probe is chosen that is complementary to a reference nucleic acid sequence, and then by selection of appropriate conditions the probe and the reference sequence selectively hybridize, or bind, to each other to form a duplex molecule. A nucleic acid molecule that is capable of hybridizing selectively to a reference sequence under moderately stringent hybridization conditions typically hybridizes under conditions that allow detection of a target nucleic acid sequence of at least about 10-14 nucleotides in length having at least approximately 70% sequence identity with the sequence of the selected nucleic acid probe. Stringent hybridization conditions typically allow detection of target nucleic acid sequences of at least about 10-14 nucleotides in length having a sequence identity of greater than about 90-95% with the sequence of the selected nucleic acid probe. Hybridization conditions useful for probe/reference sequence hybridization, where the probe and reference sequence have a specific degree of sequence identity, can be determined as is known in the art (see, for example, *Nucleic Acid Hybridization: A Practical Approach*, editors B. D. Hames and S. J. Higgins, (1985) Oxford; Washington, D.C.; IRL Press).

[0131] Conditions for hybridization are well-known to those of skill in the art. Hybridization stringency refers to the degree to which hybridization conditions disfavor the formation of hybrids containing mismatched nucleotides, with higher stringency correlated with a lower tolerance for mismatched hybrids. Factors that affect the stringency of hybridization are well-known to those of skill in the art and include, but are not limited to, temperature, pH, ionic strength, and concentration of organic solvents such as, for example, formamide and dimethylsulfoxide. As is known to those of skill in the art, hybridization stringency is increased by higher temperatures, lower ionic strength and lower solvent concentrations.

[0132] With respect to stringency conditions for hybridization, it is well known in the art that numerous equivalent conditions can be employed to establish a particular stringency by varying, for example, the following factors: the length and nature of the sequences, base composition of the

various sequences, concentrations of salts and other hybridization solution components, the presence or absence of blocking agents in the hybridization solutions (e.g., dextran sulfate, and polyethylene glycol), hybridization reaction temperature and time parameters, as well as, varying wash conditions. The selection of a particular set of hybridization conditions is selected following standard methods in the art (see, for example, Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, (1989) Cold Spring Harbor, N.Y.).

[0133] "Recombination" refers to a process of exchange of genetic information between two polynucleotides. For the purposes of this disclosure, "homologous recombination (HR)" refers to the specialized form of such exchange that takes place, for example, during repair of double-strand breaks in cells. This process requires nucleotide sequence homology, uses a "donor" molecule to template repair of a "target" molecule (i.e., the one that experienced the double-strand break), and is variously known as "non-crossover gene conversion" or "short tract gene conversion," because it leads to the transfer of genetic information from the donor to the target. Without wishing to be bound by any particular theory, such transfer can involve mismatch correction of heteroduplex DNA that forms between the broken target and the donor, and/or "synthesis-dependent strand annealing," in which the donor is used to resynthesize genetic information that will become part of the target, and/or related processes. Such specialized HR often results in an alteration of the sequence of the target molecule such that part or all of the sequence of the donor polynucleotide is incorporated into the target polynucleotide.

[0134] "Cleavage" refers to the breakage of the covalent backbone of a DNA molecule. Cleavage can be initiated by a variety of methods including, but not limited to, enzymatic or chemical hydrolysis of a phosphodiester bond. Both single-stranded cleavage and double-stranded cleavage are possible, and double-stranded cleavage can occur as a result of two distinct single-stranded cleavage events. DNA cleavage can result in the production of either blunt ends or staggered ends. In certain embodiments, fusion polypeptides are used for targeted double-stranded DNA cleavage.

[0135] A "cleavage domain" comprises one or more polypeptide sequences which possesses catalytic activity for DNA cleavage. A cleavage domain can be contained in a single polypeptide chain or cleavage activity can result from the association of two (or more) polypeptides.

[0136] A "cleavage half-domain" is a polypeptide sequence which, in conjunction with a second polypeptide (either identical or different) forms a complex having cleavage activity (preferably double-strand cleavage activity).

[0137] "Chromatin" is the nucleoprotein structure comprising the cellular genome. Cellular chromatin comprises nucleic acid, primarily DNA, and protein, including histones and non-histone chromosomal proteins. The majority of eukaryotic cellular chromatin exists in the form of nucleosomes, wherein a nucleosome core comprises approximately 150 base pairs of DNA associated with an octamer comprising two each of histones H2A, H2B, H3 and H4; and linker DNA (of variable length depending on the organism) extends between nucleosome cores. A molecule of histone H1 is generally associated with the linker DNA. For the purposes of the present disclosure, the term "chromatin" is

meant to encompass all types of cellular nucleoprotein, both prokaryotic and eukaryotic. Cellular chromatin includes both chromosomal and episomal chromatin.

[0138] A "chromosome," is a chromatin complex comprising all or a portion of the genome of a cell. The genome of a cell is often characterized by its karyotype, which is the collection of all the chromosomes that comprise the genome of the cell. The genome of a cell can comprise one or more chromosomes.

[0139] An "episome" is a replicating nucleic acid, nucleoprotein complex or other structure comprising a nucleic acid that is not part of the chromosomal karyotype of a cell. Examples of episomes include plasmids and certain viral genomes.

[0140] An "accessible region" is a site in cellular chromatin in which a target site present in the nucleic acid can be bound by an exogenous molecule which recognizes the target site. Without wishing to be bound by any particular theory, it is believed that an accessible region is one that is not packaged into a nucleosomal structure. The distinct structure of an accessible region can often be detected by its sensitivity to chemical and enzymatic probes, for example, nucleases.

[0141] A "target site" or "target sequence" is a nucleic acid sequence that defines a portion of a nucleic acid to which a binding molecule will bind, provided sufficient conditions for binding exist. For example, the sequence 5'-GAATTC-3' is a target site for the Eco RI restriction endonuclease.

[0142] An "exogenous" molecule is a molecule that is not normally present in a cell, but can be introduced into a cell by one or more genetic, biochemical or other methods. "Normal presence in the cell" is determined with respect to the particular developmental stage and environmental conditions of the cell. Thus, for example, a molecule that is present only during embryonic development of muscle is an exogenous molecule with respect to an adult muscle cell. Similarly, a molecule induced by heat shock is an exogenous molecule with respect to a non-heat-shocked cell. An exogenous molecule can comprise, for example, a functioning version of a malfunctioning endogenous molecule or a malfunctioning version of a normally-functioning endogenous molecule.

[0143] An exogenous molecule can be, among other things, a small molecule, such as is generated by a combinatorial chemistry process, or a macromolecule such as a protein, nucleic acid, carbohydrate, lipid, glycoprotein, lipoprotein, polysaccharide, any modified derivative of the above molecules, or any complex comprising one or more of the above molecules. Nucleic acids include DNA and RNA, can be single- or double-stranded; can be linear, branched or circular; and can be of any length. Nucleic acids include those capable of forming duplexes, as well as triplex-forming nucleic acids. See, for example, U.S. Pat. Nos. 5,176,996 and 5,422,251. Proteins include, but are not limited to, DNA-binding proteins, transcription factors, chromatin remodeling factors, methylated DNA binding proteins, polymerases, methylases, demethylases, acetylases, deacetylases, kinases, phosphatases, integrases, recombinases, ligases, topoisomerases, gyrases and helicases.

[0144] An exogenous molecule can be the same type of molecule as an endogenous molecule, e.g., an exogenous

protein or nucleic acid. For example, an exogenous nucleic acid can comprise an infecting viral genome, a plasmid or episome introduced into a cell, or a chromosome that is not normally present in the cell. Methods for the introduction of exogenous molecules into cells are known to those of skill in the art and include, but are not limited to, lipid-mediated transfer (i.e., liposomes, including neutral and cationic lipids), electroporation, direct injection, cell fusion, particle bombardment, calcium phosphate co-precipitation, DEAE-dextran-mediated transfer and viral vector-mediated transfer.

[0145] By contrast, an “endogenous” molecule is one that is normally present in a particular cell at a particular developmental stage under particular environmental conditions. For example, an endogenous nucleic acid can comprise a chromosome, the genome of a mitochondrion, chloroplast or other organelle, or a naturally-occurring episomal nucleic acid. Additional endogenous molecules can include proteins, for example, transcription factors and enzymes.

[0146] A “fusion” molecule is a molecule in which two or more subunit molecules are linked, preferably covalently. The subunit molecules can be the same chemical type of molecule, or can be different chemical types of molecules. Examples of the first type of fusion molecule include, but are not limited to, fusion proteins (for example, a fusion between a ZFP DNA-binding domain and a cleavage domain) and fusion nucleic acids (for example, a nucleic acid encoding the fusion protein described supra). Examples of the second type of fusion molecule include, but are not limited to, a fusion between a triplex-forming nucleic acid and a polypeptide, and a fusion between a minor groove binder and a nucleic acid.

[0147] Expression of a fusion protein in a cell can result from delivery of the fusion protein to the cell or by delivery of a polynucleotide encoding the fusion protein to a cell, wherein the polynucleotide is transcribed, and the transcript is translated, to generate the fusion protein. Trans-splicing, polypeptide cleavage and polypeptide ligation can also be involved in expression of a protein in a cell. Methods for polynucleotide and polypeptide delivery to cells are presented elsewhere in this disclosure.

[0148] A “gene,” for the purposes of the present disclosure, includes a DNA region encoding a gene product (see *infra*), as well as all DNA regions which regulate the production of the gene product, whether or not such regulatory sequences are adjacent to coding and/or transcribed sequences. Accordingly, a gene includes, but is not necessarily limited to, promoter sequences, terminators, translational regulatory sequences such as ribosome binding sites and internal ribosome entry sites, enhancers, silencers, insulators, boundary elements, replication origins, matrix attachment sites and locus control regions.

[0149] “Gene expression” refers to the conversion of the information, contained in a gene, into a gene product. A gene product can be the direct transcriptional product of a gene (e.g., mRNA, tRNA, rRNA, antisense RNA, ribozyme, structural RNA or any other type of RNA) or a protein produced by translation of a mRNA. Gene products also include RNAs which are modified, by processes such as capping, polyadenylation, methylation, and editing, and proteins modified by, for example, methylation, acetylation, phosphorylation, ubiquitination, ADP-ribosylation, myristylation, and glycosylation.

[0150] “Modulation” of gene expression refers to a change in the activity of a gene. Modulation of expression can include, but is not limited to, gene activation and gene repression.

[0151] “Eucaryotic” cells include, but are not limited to, fungal cells (such as yeast), plant cells, animal cells, mammalian cells and human cells.

[0152] A “region of interest” is any region of cellular chromatin, such as, for example, a gene or a non-coding sequence within or adjacent to a gene, in which it is desirable to bind an exogenous molecule. Binding can be for the purposes of targeted DNA cleavage and/or targeted recombination. A region of interest can be present in a chromosome, an episome, an organellar genome (e.g., mitochondrial, chloroplast), or an infecting viral genome, for example. A region of interest can be within the coding region of a gene, within transcribed non-coding regions such as, for example, leader sequences, trailer sequences or introns, or within non-transcribed regions, either upstream or downstream of the coding region. A region of interest can be as small as a single nucleotide pair or up to 2,000 nucleotide pairs in length, or any integral value of nucleotide pairs.

[0153] The terms “operative linkage” and “operatively linked” (or “operably linked”) are used interchangeably with reference to a juxtaposition of two or more components (such as sequence elements), in which the components are arranged such that both components function normally and allow the possibility that at least one of the components can mediate a function that is exerted upon at least one of the other components. By way of illustration, a transcriptional regulatory sequence, such as a promoter, is operatively linked to a coding sequence if the transcriptional regulatory sequence controls the level of transcription of the coding sequence in response to the presence or absence of one or more transcriptional regulatory factors. A transcriptional regulatory sequence is generally operatively linked in *cis* with a coding sequence, but need not be directly adjacent to it. For example, an enhancer is a transcriptional regulatory sequence that is operatively linked to a coding sequence, even though they are not contiguous.

[0154] With respect to fusion polypeptides, the term “operatively linked” can refer to the fact that each of the components performs the same function in linkage to the other component as it would if it were not so linked. For example, with respect to a fusion polypeptide in which a ZFP DNA-binding domain is fused to a cleavage domain, the ZFP DNA-binding domain and the cleavage domain are in operative linkage if, in the fusion polypeptide, the ZFP DNA-binding domain portion is able to bind its target site and/or its binding site, while the cleavage domain is able to cleave DNA in the vicinity of the target site.

[0155] A “functional fragment” of a protein, polypeptide or nucleic acid is a protein, polypeptide or nucleic acid whose sequence is not identical to the full-length protein, polypeptide or nucleic acid, yet retains the same function as the full-length protein, polypeptide or nucleic acid. A functional fragment can possess more, fewer, or the same number of residues as the corresponding native molecule, and/or can contain one or more amino acid or nucleotide substitutions. Methods for determining the function of a nucleic acid (e.g., coding function, ability to hybridize to another nucleic acid) are well-known in the art. Similarly, methods

for determining protein function are well-known. For example, the DNA-binding function of a polypeptide can be determined, for example, by filter-binding, electrophoretic mobility-shift, or immunoprecipitation assays. DNA cleavage can be assayed by gel electrophoresis. See Ausubel et al., *supra*. The ability of a protein to interact with another protein can be determined, for example, by co-immunoprecipitation, two-hybrid assays or complementation, both genetic and biochemical. See, for example, Fields et al. (1989) *Nature* 340:245-246; U.S. Pat. No. 5,585,245 and PCT WO 98/44350.

Target Sites

[0156] The disclosed methods and compositions include fusion proteins comprising a cleavage domain (or a cleavage half-domain) and a zinc finger domain, in which the zinc finger domain, by binding to a sequence in cellular chromatin (e.g., a target site or a binding site), directs the activity of the cleavage domain (or cleavage half-domain) to the vicinity of the sequence and, hence, induces cleavage in the vicinity of the target sequence. As set forth elsewhere in this disclosure, a zinc finger domain can be engineered to bind to virtually any desired sequence. Accordingly, after identifying a region of interest containing a sequence at which cleavage or recombination is desired, one or more zinc finger binding domains can be engineered to bind to one or more sequences in the region of interest. Expression of a fusion protein comprising a zinc finger binding domain and a cleavage domain (or of two fusion proteins, each comprising a zinc finger binding domain and a cleavage half-domain), in a cell, effects cleavage in the region of interest.

[0157] Selection of a sequence in cellular chromatin for binding by a zinc finger domain (e.g., a target site) can be accomplished, for example, according to the methods disclosed in co-owned U.S. Pat. No. 6,453,242 (Sep. 17, 2002), which also discloses methods for designing ZFPs to bind to a selected sequence. It will be clear to those skilled in the art that simple visual inspection of a nucleotide sequence can also be used for selection of a target site. Accordingly, any means for target site selection can be used in the claimed methods.

[0158] Target sites are generally composed of a plurality of adjacent target subsites. A target subsite refers to the sequence (usually either a nucleotide triplet, or a nucleotide quadruplet that can overlap by one nucleotide with an adjacent quadruplet) bound by an individual zinc finger. See, for example, WO 02/077227. If the strand with which a zinc finger protein makes most contacts is designated the target strand "primary recognition strand," or "primary contact strand," some zinc finger proteins bind to a three base triplet in the target strand and a fourth base on the non-target strand. A target site generally has a length of at least 9 nucleotides and, accordingly, is bound by a zinc finger binding domain comprising at least three zinc fingers. However binding of, for example, a 4-finger binding domain to a 12-nucleotide target site, a 5-finger binding domain to a 15-nucleotide target site or a 6-finger binding domain to an 18-nucleotide target site, is also possible. As will be apparent, binding of larger binding domains (e.g., 7-, 8-, 9-finger and more) to longer target sites is also possible.

[0159] It is not necessary for a target site to be a multiple of three nucleotides. For example, in cases in which cross-strand interactions occur (see, e.g., U.S. Pat. No. 6,453,242

and WO 02/077227), one or more of the individual zinc fingers of a multi-finger binding domain can bind to overlapping quadruplet subsites. As a result, a three-finger protein can bind a 10-nucleotide sequence, wherein the tenth nucleotide is part of a quadruplet bound by a terminal finger, a four-finger protein can bind a 13-nucleotide sequence, wherein the thirteenth nucleotide is part of a quadruplet bound by a terminal finger, etc.

[0160] The length and nature of amino acid linker sequences between individual zinc fingers in a multi-finger binding domain also affects binding to a target sequence. For example, the presence of a so-called "non-canonical linker," "long linker" or "structured linker" between adjacent zinc fingers in a multi-finger binding domain can allow those fingers to bind subsites which are not immediately adjacent. Non-limiting examples of such linkers are described, for example, in U.S. Pat. No. 6,479,626 and WO 01/53480. Accordingly, one or more subsites, in a target site for a zinc finger binding domain, can be separated from each other by 1, 2, 3, 4, 5 or more nucleotides. To provide but one example, a four-finger binding domain can bind to a 13-nucleotide target site comprising, in sequence, two contiguous 3-nucleotide subsites, an intervening nucleotide, and two contiguous triplet subsites.

[0161] Distance between sequences (e.g., target sites) refers to the number of nucleotides or nucleotide pairs intervening between two sequences, as measured from the edges of the sequences nearest each other.

[0162] In certain embodiments in which cleavage depends on the binding of two zinc finger domain/cleavage half-domain fusion molecules to separate target sites, the two target sites can be on opposite DNA strands. In other embodiments, both target sites are on the same DNA strand.

Zinc Finger Binding Domains

[0163] A zinc finger binding domain comprises one or more zinc fingers. Miller et al. (1985) *EMBO J* 4:1609-1614; Rhodes (1993) *Scientific American February*:56-65; U.S. Pat. No. 6,453,242. Typically, a single zinc finger domain is about 30 amino acids in length. Structural studies have demonstrated that each zinc finger domain (motif) contains two beta sheets (held in a beta turn which contains the two invariant cysteine residues) and an alpha helix (containing the two invariant histidine residues), which are held in a particular conformation through coordination of a zinc atom by the two cysteines and the two histidines.

[0164] Zinc fingers include both canonical C_2H_2 zinc fingers (i.e., those in which the zinc ion is coordinated by two cysteine and two histidine residues) and non-canonical zinc fingers such as, for example, C_3H zinc fingers (those in which the zinc ion is coordinated by three cysteine residues and one histidine residue) and C_4 zinc fingers (those in which the zinc ion is coordinated by four cysteine residues). See also WO 02/057293.

[0165] Zinc finger binding domains can be engineered to bind to a sequence of choice. See, for example, Beerli et al. (2002) *Nature Biotechnol.* 20:135-141; Pabo et al. (2001) *Ann. Rev. Biochem.* 70:313-340; Isalan et al. (2001) *Nature Biotechnol.* 19:656-660; Segal et al. (2001) *Curr. Opin. Biotechnol.* 12:632-637; Choo et al. (2000) *Curr. Opin. Struct. Biol.* 10:411-416. An engineered zinc finger binding domain can have a novel binding specificity, compared to a

naturally-occurring zinc finger protein. Engineering methods include, but are not limited to, rational design and various types of selection. Rational design includes, for example, using databases comprising triplet (or quadruplet) nucleotide sequences and individual zinc finger amino acid sequences, in which each triplet or quadruplet nucleotide sequence is associated with one or more amino acid sequences of zinc fingers which bind the particular triplet or quadruplet sequence. See, for example, co-owned U.S. Pat. Nos. 6,453,242 and 6,534,261.

[0166] Exemplary selection methods, including phage display and two-hybrid systems, are disclosed in U.S. Pat. Nos. 5,789,538; 5,925,523; 6,007,988; 6,013,453; 6,410,248; 6,140,466; 6,200,759; and 6,242,568; as well as WO 98/37186; WO 98/53057; WO 00/27878; WO 01/88197 and GB 2,338,237.

[0167] Enhancement of binding specificity for zinc finger binding domains has been described, for example, in co-owned WO 02/077227.

[0168] Since an individual zinc finger binds to a three-nucleotide (i.e., triplet) sequence (or a four-nucleotide sequence which can overlap, by one nucleotide, with the four-nucleotide binding site of an adjacent zinc finger), the length of a sequence to which a zinc finger binding domain is engineered to bind (e.g., a target sequence) will determine the number of zinc fingers in an engineered zinc finger binding domain. For example, for ZFPs in which the finger motifs do not bind to overlapping subsites, a six-nucleotide target sequence is bound by a two-finger binding domain; a nine-nucleotide target sequence is bound by a three-finger binding domain, etc. As noted herein, binding sites for individual zinc fingers (i.e., subsites) in a target site need not be contiguous, but can be separated by one or several nucleotides, depending on the length and nature of the amino acids sequences between the zinc fingers (i.e., the inter-finger linkers) in a multi-finger binding domain.

[0169] In a multi-finger zinc finger binding domain, adjacent zinc fingers can be separated by amino acid linker sequences of approximately 5 amino acids (so-called "canonical" inter-finger linkers) or, alternatively, by one or more non-canonical linkers. See, e.g., co-owned U.S. Pat. Nos. 6,453,242 and 6,534,261. For engineered zinc finger binding domains comprising more than three fingers, insertion of longer ("non-canonical") inter-finger linkers between certain of the zinc fingers may be preferred as it may increase the affinity and/or specificity of binding by the binding domain. See, for example, U.S. Pat. No. 6,479,626 and WO 01/53480. Accordingly, multi-finger zinc finger binding domains can also be characterized with respect to the presence and location of non-canonical inter-finger linkers. For example, a six-finger zinc finger binding domain comprising three fingers (joined by two canonical inter-finger linkers), a long linker and three additional fingers (joined by two canonical inter-finger linkers) is denoted a 2x3 configuration. Similarly, a binding domain comprising two fingers (with a canonical linker therebetween), a long linker and two additional fingers (joined by a canonical linker) is denoted a 2x2 protein. A protein comprising three two-finger units (in each of which the two fingers are joined by a canonical linker), and in which each two-finger unit is joined to the adjacent two finger unit by a long linker, is referred to as a 3x2 protein.

[0170] The presence of a long or non-canonical inter-finger linker between two adjacent zinc fingers in a multi-finger binding domain often allows the two fingers to bind to subsites which are not immediately contiguous in the target sequence. Accordingly, there can be gaps of one or more nucleotides between subsites in a target site; i.e., a target site can contain one or more nucleotides that are not contacted by a zinc finger. For example, a 2x2 zinc finger binding domain can bind to two six-nucleotide sequences separated by one nucleotide, i.e., it binds to a 13-nucleotide target site. See also Moore et al. (2001a) Proc. Natl. Acad. Sci. USA 98:1432-1436; Moore et al. (2001b) Proc. Natl. Acad. Sci. USA 98:1437-1441 and WO 01/53480.

[0171] As mentioned previously, a target subsite is a three- or four-nucleotide sequence that is bound by a single zinc finger. For certain purposes, a two-finger unit is denoted a binding module. A binding module can be obtained by, for example, selecting for two adjacent fingers in the context of a multi-finger protein (generally three fingers) which bind a particular six-nucleotide target sequence. Alternatively, modules can be constructed by assembly of individual zinc fingers. See also WO 98/53057 and WO 01/53480.

Cleavage Domains

[0172] The cleavage domain portion of the fusion proteins disclosed herein can be obtained from any endonuclease or exonuclease. Exemplary endonucleases from which a cleavage domain can be derived include, but are not limited to, restriction endonucleases and homing endonucleases. See, for example, 2002-2003 Catalogue, New England Biolabs, Beverly, Mass.; and Belfort et al. (1997) *Nucleic Acids Res.* 25:3379-3388. Additional enzymes which cleave DNA are known (e.g., S1 Nuclease; mung bean nuclease; pancreatic DNase I; micrococcal nuclease; yeast HO endonuclease; see also Linn et al. (eds.) *Nucleases*, Cold Spring Harbor Laboratory Press, 1993). One or more of these enzymes (or functional fragments thereof) can be used as a source of cleavage domains and cleavage half-domains.

[0173] Similarly, a cleavage half-domain (e.g., fusion proteins comprising a zinc finger binding domain and a cleavage half-domain) can be derived from any nuclease or portion thereof, as set forth above, that requires dimerization for cleavage activity. In general, two fusion proteins are required for cleavage if the fusion proteins comprise cleavage half-domains. Alternatively, a single protein comprising two cleavage half-domains can be used. The two cleavage half-domains can be derived from the same endonuclease (or functional fragments thereof), or each cleavage half-domain can be derived from a different endonuclease (or functional fragments thereof). In addition, the target sites for the two fusion proteins are preferably disposed, with respect to each other, such that binding of the two fusion proteins to their respective target sites places the cleavage half-domains in a spatial orientation to each other that allows the cleavage half-domains to form a functional cleavage domain, e.g., by dimerizing. Thus, in certain embodiments, the near edges of the target sites are separated by 5-8 nucleotides or by 15-18 nucleotides. However any integral number of nucleotides or nucleotide pairs can intervene between two target sites (e.g., from 2 to 50 nucleotides or more). In general, the point of cleavage lies between the target sites.

[0174] Restriction endonucleases (restriction enzymes) are present in many species and are capable of sequence-

specific binding to DNA (at a recognition site), and cleaving DNA at or near the site of binding. Certain restriction enzymes (e.g., Type IIS) cleave DNA at sites removed from the recognition site and have separable binding and cleavage domains. For example, the Type IIS enzyme Fok I catalyzes double-stranded cleavage of DNA, at 9 nucleotides from its recognition site on one strand and 13 nucleotides from its recognition site on the other. See, for example, U.S. Pat. Nos. 5,356,802; 5,436,150 and 5,487,994; as well as Li et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:4275-4279; Li et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:2764-2768; Kim et al. (1994a) *Proc. Natl. Acad. Sci. USA* 91:883-887; Kim et al. (1994b) *J. Biol. Chem.* 269:31,978-31,982. Thus, in one embodiment, fusion proteins comprise the cleavage domain (or cleavage half-domain) from at least one Type IIS restriction enzyme and one or more zinc finger binding domains, which may or may not be engineered.

[0175] An exemplary Type IIS restriction enzyme, whose cleavage domain is separable from the binding domain, is Fok I. This particular enzyme is active as a dimer. Bitinaite et al. (1998) *Proc. Natl. Acad. Sci. USA* 95: 10,570-10,575. Accordingly, for the purposes of the present disclosure, the portion of the Fok I enzyme used in the disclosed fusion proteins is considered a cleavage half-domain. Thus, for targeted double-stranded cleavage and/or targeted replacement of cellular sequences using zinc finger-Fok I fusions, two fusion proteins, each comprising a FokI cleavage half-domain, can be used to reconstitute a catalytically active cleavage domain. Alternatively, a single polypeptide molecule containing a zinc finger binding domain and two Fok I cleavage half-domains can also be used. Parameters for targeted cleavage and targeted sequence alteration using zinc finger-Fok I fusions are provided elsewhere in this disclosure.

[0176] A cleavage domain or cleavage half-domain can be any portion of a protein that retains cleavage activity, or that retains the ability to multimerize (e.g., dimerize) to form a functional cleavage domain.

[0177] Exemplary Type IIS restriction enzymes are listed in Table 1. Additional restriction enzymes also contain separable binding and cleavage domains, and these are contemplated by the present disclosure. See, for example, Roberts et al. (2003) *Nucleic Acids Res.* 31:418-420.

TABLE 1

Some Type IIS Restriction Enzymes		
Aar I	BsrB I	SspD5 I
Ace III	BsrD I	Sth132 I
Aci I	BstF5 I	Sts I
Alo I	Btr I	TspDT I
Bae I	Bts I	TspGW I
Bbr7 I	Cdi I	Tth111 II
Bbv I	CjeP I	UbaP I
Bbv II	Drd II	Bsa I
BbvC I	Eci I	BsmB I
Bcc I	Eco31 I	
Bce83 I	Eco57 I	
BceA I	Eco57M I	
Beef I	Esp3 I	
Bcg I	Fau I	
BciV I	Fin I	
Bfi I	Fok I	
Bin I	Gdi II	
Bmg I	Gsu I	

TABLE 1-continued

Some Type IIS Restriction Enzymes	
Bpu10 I	Hga I
BsaX I	Hin4 II
Bsb I	Hph I
BscA I	Ksp632 I
BscG I	Mbo II
BseR I	Mly I
BseY I	Mme I
Bsi I	Mnl I
Bsm I	Pfl1108 I
BsmA I	Ple I
BsmF I	Ppi I
Bsp24 I	Psr I
BspG I	RleA I
BspM I	Sap I
BspNC I	SfaN I
Bsr I	Sim I

Zinc Finger Domain-cleavage Domain Fusions

[0178] Methods for design and construction of fusion proteins (and polynucleotides encoding same) are known to those of skill in the art. For example, methods for the design and construction of fusion protein comprising zinc finger proteins (and polynucleotides encoding same) are described in co-owned U.S. Pat. Nos. 6,453,242 and 6,534,261. In certain embodiments, polynucleotides encoding such fusion proteins are constructed. These polynucleotides can be inserted into a vector and the vector can be introduced into a cell (see below for additional disclosure regarding vectors and methods for introducing polynucleotides into cells).

[0179] In certain embodiments of the methods described herein, a fusion protein comprises a zinc finger binding domain and a cleavage half-domain from the Fok I restriction enzyme, and two such fusion proteins are expressed in a cell. Expression of two fusion proteins in a cell can result from delivery of the two proteins to the cell; delivery of one protein and one nucleic acid encoding one of the proteins to the cell; delivery of two nucleic acids, each encoding one of the proteins, to the cell; or by delivery of a single nucleic acid, encoding both proteins, to the cell. In additional embodiments, a fusion protein comprises a single polypeptide chain comprising two cleavage half domains and a zinc finger binding domain. In this case, a single fusion protein is expressed in a cell and, without wishing to be bound by theory, is believed to cleave DNA as a result of formation of an intramolecular dimer of the cleavage half-domains.

[0180] In certain embodiments, the components of the fusion proteins (e.g., ZFP-Fok I fusions) are arranged such that the zinc finger domain is nearest the amino terminus of the fusion protein, and the cleavage half-domain is nearest the carboxy-terminus. This mirrors the relative orientation of the cleavage domain in naturally-occurring dimerizing cleavage domains such as those derived from the Fok I enzyme, in which the DNA-binding domain is nearest the amino terminus and the cleavage half-domain is nearest the carboxy terminus. In these embodiments, dimerization of the cleavage half-domains to form a functional nuclease is brought about by binding of the fusion proteins to sites on opposite DNA strands, with the 5' ends of the binding sites being proximal to each other. See FIG. 43A.

[0181] In additional embodiments, the components of the fusion proteins (e.g., ZFP-Fok I fusions) are arranged such

that the cleavage half-domain is nearest the amino terminus of the fusion protein, and the zinc finger domain is nearest the carboxy-terminus. In these embodiments, dimerization of the cleavage half-domains to form a functional nuclease is brought about by binding of the fusion proteins to sites on opposite DNA strands, with the 3' ends of the binding sites being proximal to each other. See FIG. 43B.

[0182] In yet additional embodiments, a first fusion protein contains the cleavage half-domain nearest the amino terminus of the fusion protein, and the zinc finger domain nearest the carboxy-terminus, and a second fusion protein is arranged such that the zinc finger domain is nearest the amino terminus of the fusion protein, and the cleavage half-domain is nearest the carboxy-terminus. In these embodiments, both fusion proteins bind to the same DNA strand, with the binding site of the first fusion protein containing the zinc finger domain nearest the carboxy terminus located to the 5' side of the binding site of the second fusion protein containing the zinc finger domain nearest the amino terminus. See FIG. 43C.

[0183] In the disclosed fusion proteins, the amino acid sequence between the zinc finger domain and the cleavage domain (or cleavage half-domain) is denoted the "ZC linker." The ZC linker is to be distinguished from the inter-finger linkers discussed above. For the purposes of determining the length of a ZC linker, the zinc finger structure described by Pabo et al. (2001) *Ann. Rev. Biochem.* 70:313-340 is used:

X-X-C-X₂₋₄-C-X₁₂-H-X₃₋₅-H (SEQ ID NO:201)

[0184] In this structure, the first residue of a zinc finger is the amino acid located two residues amino-terminal to the first conserved cysteine residue. In the majority of naturally-occurring zinc finger proteins, this position is occupied by a hydrophobic amino acid (usually either phenylalanine or tyrosine). In the disclosed fusion proteins, the first residue of a zinc finger will thus often be a hydrophobic residue, but it can be any amino acid. The final amino acid residue of a zinc finger, as shown above, is the second conserved histidine residue.

[0185] Thus, in the disclosed fusion proteins having a polarity in which the zinc finger binding domain is amino-terminal to the cleavage domain (or cleavage half-domain), the ZC linker is the amino acid sequence between the second conserved histidine residue of the C-terminal-most zinc finger and the N-terminal-most amino acid of the cleavage domain (or cleavage half-domain). For example, in certain fusion proteins whose construction is exemplified in the Examples section, the N-terminal-most amino acid of a cleavage half-domain is a glutamine (Q) residue corresponding to amino acid number 384 in the FokI sequence of Looney et al. (1989) *Gene* 80:193-208.

[0186] For fusion proteins having a polarity in which the cleavage domain (or cleavage half-domain) is amino-terminal to the zinc finger binding domain, the ZC linker is the amino acid sequence between the C-terminal-most amino acid residue of the cleavage domain (or half-domain) and the first residue of the N-terminal-most zinc finger of the zinc finger binding domain (i.e., the residue located two residues upstream of the first conserved cysteine residue). In certain

exemplary fusion proteins, the C-terminal-most amino acid of a cleavage half-domain is a phenylalanine (F) residue corresponding to amino acid number 579 in the FokI sequence of Looney et al. (1989) *Gene* 80:193-208.

[0187] The ZC linker can be any amino acid sequence. To obtain optimal cleavage, the length of the ZC linker and the distance between the target sites (binding sites) are interrelated. See, for example, Smith et al. (2000) *Nucleic Acids Res.* 28:3361-3369; Bibikova et al. (2001) *Mol. Cell. Biol.* 21:289-297, noting that their notation for linker length differs from that given here. For example, for ZFP-Fok I fusions in which the zinc finger binding domain is amino-terminal to the cleavage half-domain, and having a ZC linker length of four amino acids as defined herein (and denoted LO by others), optimal cleavage occurs when the binding sites for the fusion proteins are located 6 or 16 nucleotides apart (as measured from the near edge of each binding site). See Example 4.

Methods for Targeted Cleavage

[0188] The disclosed methods and compositions can be used to cleave DNA at a region of interest in cellular chromatin (e.g., at a desired or predetermined site in a genome, for example, in a gene, either mutant or wild-type). For such targeted DNA cleavage, a zinc finger binding domain is engineered to bind a target site at or near the predetermined cleavage site, and a fusion protein comprising the engineered zinc finger binding domain and a cleavage domain is expressed in a cell. Upon binding of the zinc finger portion of the fusion protein to the target site, the DNA is cleaved near the target site by the cleavage domain. The exact site of cleavage can depend on the length of the ZC linker.

[0189] Alternatively, two fusion proteins, each comprising a zinc finger binding domain and a cleavage half-domain, are expressed in a cell, and bind to target sites which are juxtaposed in such a way that a functional cleavage domain is reconstituted and DNA is cleaved in the vicinity of the target sites. In one embodiment, cleavage occurs between the target sites of the two zinc finger binding domains. One or both of the zinc finger binding domains can be engineered.

[0190] For targeted cleavage using a zinc finger binding domain-cleavage domain fusion polypeptide, the binding site can encompass the cleavage site, or the near edge of the binding site can be 1, 2, 3, 4, 5, 6, 10, 25, 50 or more nucleotides (or any integral value between 1 and 50 nucleotides) from the cleavage site. The exact location of the binding site, with respect to the cleavage site, will depend upon the particular cleavage domain, and the length of the ZC linker. For methods in which two fusion polypeptides, each comprising a zinc finger binding domain and a cleavage half-domain, are used, the binding sites generally straddle the cleavage site. Thus the near edge of the first binding site can be 1, 2, 3, 4, 5, 6, 10, 25 or more nucleotides (or any integral value between 1 and 50 nucleotides) on one side of the cleavage site, and the near edge of the second binding site can be 1, 2, 3, 4, 5, 6, 10, 25 or more nucleotides (or any integral value between 1 and 50 nucleotides) on the other side of the cleavage site. Methods for mapping cleavage sites in vitro and in vivo are known to those of skill in the art.

[0191] Thus, the methods described herein can employ an engineered zinc finger binding domain fused to a cleavage

domain. In these cases, the binding domain is engineered to bind to a target sequence, at or near which cleavage is desired. The fusion protein, or a polynucleotide encoding same, is introduced into a cell. Once introduced into, or expressed in, the cell, the fusion protein binds to the target sequence and cleaves at or near the target sequence. The exact site of cleavage depends on the nature of the cleavage domain and/or the presence and/or nature of linker sequences between the binding and cleavage domains. In cases where two fusion proteins, each comprising a cleavage half-domain, are used, the distance between the near edges of the binding sites can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 25 or more nucleotides (or any integral value between 1 and 50 nucleotides). Optimal levels of cleavage can also depend on both the distance between the binding sites of the two fusion proteins (See, for example, Smith et al. (2000) *Nucleic Acids Res.* 28:3361-3369; Bibikova et al. (2001) *Mol. Cell. Biol.* 21:289-297) and the length of the ZC linker in each fusion protein.

[0192] For ZFP-FokI fusion nucleases, the length of the linker between the ZFP and the FokI cleavage half-domain (i.e., the ZC linker) can influence cleavage efficiency. In one experimental system utilizing a ZFP-FokI fusion with a ZC linker of 4 amino acid residues, optimal cleavage was obtained when the near edges of the binding sites for two ZFP-FokI nucleases were separated by 6 base pairs. This particular fusion nuclease comprised the following amino acid sequence between the zinc finger portion and the nuclease half-domain:

HQRTHQNKKQLV

(SEQ ID NO:26)

in which the two conserved histidines in the C-terminal portion of the zinc finger and the first three residues in the FokI cleavage half-domain are underlined. Accordingly, the ZC linker sequence in this construct is QNKK. Bibikova et al. (2001) *Mol. Cell. Biol.* 21:289-297. The present inventors have constructed a number of ZFP-FokI fusion nucleases having a variety of ZC linker lengths and sequences, and analyzed the cleavage efficiencies of these nucleases on a series of substrates having different distances between the ZFP binding sites. See Example 4.

[0193] In certain embodiments, the cleavage domain comprises two cleavage half-domains, both of which are part of a single polypeptide comprising a binding domain, a first cleavage half-domain and a second cleavage half-domain. The cleavage half-domains can have the same amino acid sequence or different amino acid sequences, so long as they function to cleave the DNA.

[0194] Cleavage half-domains may also be provided in separate molecules. For example, two fusion polypeptides may be introduced into a cell, wherein each polypeptide comprises a binding domain and a cleavage half-domain. The cleavage half-domains can have the same amino acid sequence or different amino acid sequences, so long as they function to cleave the DNA. Further, the binding domains bind to target sequences which are typically disposed in such a way that, upon binding of the fusion polypeptides, the two cleavage half-domains are presented in a spatial orientation to each other that allows reconstitution of a cleavage domain (e.g., by dimerization of the half-domains), thereby positioning the half-domains relative to each other to form a

functional cleavage domain, resulting in cleavage of cellular chromatin in a region of interest. Generally, cleavage by the reconstituted cleavage domain occurs at a site located between the two target sequences. One or both of the proteins can be engineered to bind to its target site.

[0195] The two fusion proteins can bind in the region of interest in the same or opposite polarity, and their binding sites (i.e., target sites) can be separated by any number of nucleotides, e.g., from 0 to 200 nucleotides or any integral value therebetween. In certain embodiments, the binding sites for two fusion proteins, each comprising a zinc finger binding domain and a cleavage half-domain, can be located between 5 and 18 nucleotides apart, for example, 5-8 nucleotides apart, or 15-18 nucleotides apart, or 6 nucleotides apart, or 16 nucleotides apart, as measured from the edge of each binding site nearest the other binding site, and cleavage occurs between the binding sites.

[0196] The site at which the DNA is cleaved generally lies between the binding sites for the two fusion proteins. Double-strand breakage of DNA often results from two single-strand breaks, or "nicks," offset by 1, 2, 3, 4, 5, 6 or more nucleotides, (for example, cleavage of double-stranded DNA by native Fok I results from single-strand breaks offset by 4 nucleotides). Thus, cleavage does not necessarily occur at exactly opposite sites on each DNA strand. In addition, the structure of the fusion proteins and the distance between the target sites can influence whether cleavage occurs adjacent a single nucleotide pair, or whether cleavage occurs at several sites. However, for many applications, including targeted recombination and targeted mutagenesis (see infra) cleavage within a range of nucleotides is generally sufficient, and cleavage between particular base pairs is not required.

[0197] As noted above, the fusion protein(s) can be introduced as polypeptides and/or polynucleotides. For example, two polynucleotides, each comprising sequences encoding one of the aforementioned polypeptides, can be introduced into a cell, and when the polypeptides are expressed and each binds to its target sequence, cleavage occurs at or near the target sequence. Alternatively, a single polynucleotide comprising sequences encoding both fusion polypeptides is introduced into a cell. Polynucleotides can be DNA, RNA or any modified forms or analogues or DNA and/or RNA.

[0198] To enhance cleavage specificity, additional compositions may also be employed in the methods described herein. For example, single cleavage half-domains can exhibit limited double-stranded cleavage activity. In methods in which two fusion proteins, each containing a three-finger zinc finger domain and a cleavage half-domain, are introduced into the cell, either protein specifies an approximately 9-nucleotide target site. Although the aggregate target sequence of 18 nucleotides is likely to be unique in a mammalian genome, any given 9-nucleotide target site occurs, on average, approximately 23,000 times in the human genome. Thus, non-specific cleavage, due to the site-specific binding of a single half-domain, may occur. Accordingly, the methods described herein contemplate the use of a dominant-negative mutant of a cleavage half-domain such as Fok I (or a nucleic acid encoding same) that is expressed in a cell along with the two fusion proteins. The dominant-negative mutant is capable of dimerizing but is unable to cleave, and also blocks the cleavage activity of a half-domain to which it is dimerized. By providing the

dominant-negative mutant in molar excess to the fusion proteins, only regions in which both fusion proteins are bound will have a high enough local concentration of functional cleavage half-domains for dimerization and cleavage to occur. At sites where only one of the two fusion proteins is bound, its cleavage half-domain forms a dimer with the dominant negative mutant half-domain, and undesirable, non-specific cleavage does not occur.

[0199] Three catalytic amino acid residues in the Fok I cleavage half-domain have been identified: Asp 450, Asp 467 and Lys 469. Bitinaite et al. (1998) *Proc. Natl. Acad. Sci. USA* 95: 10,570-10,575. Thus, one or more mutations at one of these residues can be used to generate a dominant negative mutation. Further, many of the catalytic amino acid residues of other Type IIS endonucleases are known and/or can be determined, for example, by alignment with Fok I sequences and/or by generation and testing of mutants for catalytic activity.

Dimerization Domain Mutations in the Cleavage Half-domain

[0200] Methods for targeted cleavage which involve the use of fusions between a ZFP and a cleavage half-domain (such as, e.g., a ZFP/FokI fusion) require the use of two such fusion molecules, each generally directed to a distinct target sequence. Target sequences for the two fusion proteins can be chosen so that targeted cleavage is directed to a unique site in a genome, as discussed above. A potential source of reduced cleavage specificity could result from homodimerization of one of the two ZFP/cleavage half-domain fusions. This might occur, for example, due to the presence, in a genome, of inverted repeats of the target sequences for one of the two ZFP/cleavage half-domain fusions, located so as to allow two copies of the same fusion protein to bind with an orientation and spacing that allows formation of a functional dimer.

[0201] One approach for reducing the probability of this type of aberrant cleavage at sequences other than the intended target site involves generating variants of the cleavage half-domain that minimize or prevent homodimerization. Preferably, one or more amino acids in the region of the half-domain involved in its dimerization are altered. In the crystal structure of the FokI protein dimer, the structure of the cleavage half-domains is reported to be similar to the arrangement of the cleavage half-domains during cleavage of DNA by FokI. Wah et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:10564-10569. This structure indicates that amino acid residues at positions 483 and 487 play a key role in the dimerization of the FokI cleavage half-domains. The structure also indicates that amino acid residues at positions 446, 447, 479, 483, 484, 486, 487, 490, 491, 496, 498, 499, 500, 531, 534, 537, and 538 are all close enough to the dimerization interface to influence dimerization. Accordingly, amino acid sequence alterations at one or more of the aforementioned positions will likely alter the dimerization properties of the cleavage half-domain. Such changes can be introduced, for example, by constructing a library containing (or encoding) different amino acid residues at these positions and selecting variants with the desired properties, or by rationally designing individual mutants. In addition to preventing homodimerization, it is also possible that some of these mutations may increase the cleavage efficiency above that obtained with two wild-type cleavage half-domains.

[0202] Accordingly, alteration of a FokI cleavage half-domain at any amino acid residue which affects dimerization can be used to prevent one of a pair of ZFP/FokI fusions from undergoing homodimerization which can lead to cleavage at undesired sequences. Thus, for targeted cleavage using a pair of ZFP/FokI fusions, one or both of the fusion proteins can comprise one or more amino acid alterations that inhibit self-dimerization, but allow heterodimerization of the two fusion proteins to occur such that cleavage occurs at the desired target site. In certain embodiments, alterations are present in both fusion proteins, and the alterations have additive effects; i.e., homodimerization of either fusion, leading to aberrant cleavage, is minimized or abolished, while heterodimerization of the two fusion proteins is facilitated compared to that obtained with wild-type cleavage half-domains. See Example 5.

Methods for Targeted Alteration of Genomic Sequences and Targeted Recombination

[0203] Also described herein are methods of replacing a genomic sequence (e.g., a region of interest in cellular chromatin) with a homologous non-identical sequence (i.e., targeted recombination). Previous attempts to replace particular sequences have involved contacting a cell with a polynucleotide comprising sequences bearing homology to a chromosomal region (i.e., a donor DNA), followed by selection of cells in which the donor DNA molecule had undergone homologous recombination into the genome. The success rate of these methods is low, due to poor efficiency of homologous recombination and a high frequency of non-specific insertion of the donor DNA into regions of the genome other than the target site.

[0204] The present disclosure provides methods of targeted sequence alteration characterized by a greater efficiency of targeted recombination and a lower frequency of non-specific insertion events. The methods involve making and using engineered zinc finger binding domains fused to cleavage domains (or cleavage half-domains) to make one or more targeted double-stranded breaks in cellular DNA. Because double-stranded breaks in cellular DNA stimulate cellular repair mechanisms several thousand-fold in the vicinity of the cleavage site, such targeted cleavage allows for the alteration or replacement (via homology-directed repair) of sequences at virtually any site in the genome.

[0205] In addition to the fusion molecules described herein, targeted replacement of a selected genomic sequence also requires the introduction of the replacement (or donor) sequence. The donor sequence can be introduced into the cell prior to, concurrently with, or subsequent to, expression of the fusion protein(s). The donor polynucleotide contains sufficient homology to a genomic sequence to support homologous recombination (or homology-directed repair) between it and the genomic sequence to which it bears homology. Approximately 25, 50 100, 200, 500, 750, 1,000, 1,500, 2,000 nucleotides or more of sequence homology between a donor and a genomic sequence (or any integral value between 10 and 2,000 nucleotides, or more) will support homologous recombination therebetween. Donor sequences can range in length from 10 to 5,000 nucleotides (or any integral value of nucleotides therebetween) or longer. It will be readily apparent that the donor sequence is typically not identical to the genomic sequence that it replaces. For example, the sequence of the donor polynucle-

otide can contain one or more single base changes, insertions, deletions, inversions or rearrangements with respect to the genomic sequence, so long as sufficient homology with chromosomal sequences is present. Alternatively, a donor sequence can contain a non-homologous sequence flanked by two regions of homology. Additionally, donor sequences can comprise a vector molecule containing sequences that are not homologous to the region of interest in cellular chromatin. Generally, the homologous region(s) of a donor sequence will have at least 50% sequence identity to a genomic sequence with which recombination is desired. In certain embodiments, 60%, 70%, 80%, 90%, 95%, 98%, 99%, or 99.9% sequence identity is present. Any value between 1% and 100% sequence identity can be present, depending upon the length of the donor polynucleotide.

[0206] A donor molecule can contain several, discontinuous regions of homology to cellular chromatin. For example, for targeted insertion of sequences not normally present in a region of interest, said sequences can be present in a donor nucleic acid molecule and flanked by regions of homology to sequence in the region of interest.

[0207] To simplify assays (e.g., hybridization, PCR, restriction enzyme digestion) for determining successful insertion of the donor sequence, certain sequence differences may be present in the donor sequence as compared to the genomic sequence. Preferably, if located in a coding region, such nucleotide sequence differences will not change the amino acid sequence, or will make silent amino acid changes (i.e., changes which do not affect the structure or function of the protein). The donor polynucleotide can optionally contain changes in sequences corresponding to the zinc finger domain binding sites in the region of interest, to prevent cleavage of donor sequences that have been introduced into cellular chromatin by homologous recombination.

[0208] The donor polynucleotide can be DNA or RNA, single-stranded or double-stranded and can be introduced into a cell in linear or circular form. If introduced in linear form, the ends of the donor sequence can be protected (e.g., from exonucleolytic degradation) by methods known to those of skill in the art. For example, one or more dideoxynucleotide residues are added to the 3' terminus of a linear molecule and/or self-complementary oligonucleotides are ligated to one or both ends. See, for example, Chang et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:4959-4963; Nehls et al. (1996) *Science* 272:886-889. Additional methods for protecting exogenous polynucleotides from degradation include, but are not limited to, addition of terminal amino group(s) and the use of modified internucleotide linkages such as, for example, phosphorothioates, phosphoramidates, and O-methyl ribose or deoxyribose residues. A polynucleotide can be introduced into a cell as part of a vector molecule having additional sequences such as, for example, replication origins, promoters and genes encoding antibiotic resistance. Moreover, donor polynucleotides can be introduced as naked nucleic acid, as nucleic acid complexed with an agent such as a liposome or poloxamer, or can be delivered by viruses (e.g., adenovirus, AAV, herpesvirus, retrovirus, lentivirus).

[0209] Without being bound by one theory, it appears that the presence of a double-stranded break in a cellular sequence, coupled with the presence of an exogenous DNA molecule having homology to a region adjacent to or sur-

rounding the break, activates cellular mechanisms which repair the break by transfer of sequence information from the donor molecule into the cellular (e.g., genomic or chromosomal) sequence; i.e., by a processes of homology-directed repair, also known as "gene conversion." Applicants' methods advantageously combine the powerful targeting capabilities of engineered ZFPs with a cleavage domain (or cleavage half-domain) to specifically target a double-stranded break to the region of the genome at insertion of exogenous sequences is desired.

[0210] For alteration of a chromosomal sequence, it is not necessary for the entire sequence of the donor to be copied into the chromosome, as long as enough of the donor sequence is copied to effect the desired sequence alteration.

[0211] The efficiency of insertion of donor sequences by homologous recombination is inversely related to the distance, in the cellular DNA, between the double-stranded break and the site at which recombination is desired. In other words, higher homologous recombination efficiencies are observed when the double-stranded break is closer to the site at which recombination is desired. In cases in which a precise site of recombination is not predetermined (e.g., the desired recombination event can occur over an interval of genomic sequence), the length and sequence of the donor nucleic acid, together with the site(s) of cleavage, are selected to obtain the desired recombination event. In cases in which the desired event is designed to change the sequence of a single nucleotide pair in a genomic sequence, cellular chromatin is cleaved within 10,000 nucleotides on either side of that nucleotide pair. In certain embodiments, cleavage occurs within 1,000, 500, 200, 100, 90, 80, 70, 60, 50, 40, 30, 20, 10, 5, or 2 nucleotides, or any integral value between 2 and 1,000 nucleotides, on either side of the nucleotide pair whose sequence is to be changed.

[0212] As detailed above, the binding sites for two fusion proteins, each comprising a zinc finger binding domain and a cleavage half-domain, can be located 5-8 or 15-18 nucleotides apart, as measured from the edge of each binding site nearest the other binding site, and cleavage occurs between the binding sites. Whether cleavage occurs at a single site or at multiple sites between the binding sites is immaterial, since the cleaved genomic sequences are replaced by the donor sequences. Thus, for efficient alteration of the sequence of a single nucleotide pair by targeted recombination, the midpoint of the region between the binding sites is within 10,000 nucleotides of that nucleotide pair, preferably within 1,000 nucleotides, or 500 nucleotides, or 200 nucleotides, or 100 nucleotides, or 50 nucleotides, or 20 nucleotides, or 10 nucleotides, or 5 nucleotide, or 2 nucleotides, or one nucleotide, or at the nucleotide pair of interest.

[0213] In certain embodiments, a homologous chromosome can serve as the donor polynucleotide. Thus, for example, correction of a mutation in a heterozygote can be achieved by engineering fusion proteins which bind to and cleave the mutant sequence on one chromosome, but do not cleave the wild-type sequence on the homologous chromosome. The double-stranded break on the mutation-bearing chromosome stimulates a homology-based "gene conversion" process in which the wild-type sequence from the homologous chromosome is copied into the cleaved chromosome, thus restoring two copies of the wild-type sequence.

[0214] Methods and compositions are also provided that may enhance levels of targeted recombination including, but not limited to, the use of additional ZFP-functional domain fusions to activate expression of genes involved in homologous recombination, such as, for example, members of the RAD52 epistasis group (e.g., Rad50, Rad51, Rad51B, Rad51C, Rad51D, Rad52, Rad54, Rad54B, Mre11, XRCC2, XRCC3), genes whose products interact with the aforementioned gene products (e.g., BRCA1, BRCA2) and/or genes in the NBS1 complex. Similarly ZFP-functional domain fusions can be used, in combination with the methods and compositions disclosed herein, to repress expression of genes involved in non-homologous end joining (e.g., Ku70/80, XRCC4, poly(ADP ribose) polymerase, DNA ligase 4). See, for example, Yanez et al. (1998) *Gene Therapy* 5:149-159; Hoeijmakers (2001) *Nature* 411:366-374; Johnson et al. (2001) *Biochem. Soc. Trans.* 29:196-201; Tauchi et al. (2002) *Oncogene* 21:8967-8980. Methods for activation and repression of gene expression using fusions between a zinc finger binding domain and a functional domain are disclosed, for example, in co-owned U.S. Pat. Nos. 6,534,261; 6,824,978 and 6,933,113. Additional repression methods include the use of antisense oligonucleotides and/or small interfering RNA (siRNA or RNAi) targeted to the sequence of the gene to be repressed.

[0215] As an alternative to or, in addition to, activating expression of gene products involved in homologous recombination, fusions of these protein (or functional fragments thereof) with a zinc finger binding domain targeted to the region of interest, can be used to recruit these proteins (recombination proteins) to the region of interest, thereby increasing their local concentration and further stimulating homologous recombination processes. Alternatively, a polypeptide involved in homologous recombination as described above (or a functional fragment thereof) can be part of a triple fusion protein comprising a zinc finger binding domain, a cleavage domain (or cleavage half-domain) and the recombination protein (or functional fragment thereof). Additional proteins involved in gene conversion and recombination-related chromatin remodeling, which can be used in the aforementioned methods and compositions, include histone acetyltransferases (e.g., Esa1p, Tip60), histone methyltransferases (e.g., Dot1p), histone kinases and histone phosphatases.

[0216] The p53 protein has been reported to play a central role in repressing homologous recombination (HR). See, for example, Valerie et al., (2003) *Oncogene* 22:5792-5812; Janz et al. (2002) *Oncogene* 21:5929-5933. For example, the rate of HR in p53-deficient human tumor lines is 10,000-fold greater than in primary human fibroblasts, and there is a 100-fold increase in HR in tumor cells with a non-functional p53 compared to those with functional p53. Mekeel et al. (1997) *Oncogene* 14:1847-1857. In addition, overexpression of p53 dominant negative mutants leads to a 20-fold increase in spontaneous recombination. Bertrand et al. (1997) *Oncogene* 14:1117-1122. Analysis of different p53 mutations has revealed that the roles of p53 in transcriptional transactivation and G1 cell cycle checkpoint control are separable from its involvement in HR. Saintigny et al. (1999) *Oncogene* 18:3553-3563; Boehden et al. (2003) *Oncogene* 22:4111-4117. Accordingly, downregulation of p53 activity can serve to increase the efficiency of targeted homologous recombination using the methods and compositions disclosed herein. Any method for downregulation of

p53 activity can be used, including but not limited to cotransfection and overexpression of a p53 dominant negative mutant or targeted repression of p53 gene expression according to methods disclosed, e.g., in co-owned U.S. Pat. No. 6,534,261.

[0217] Further increases in efficiency of targeted recombination, in cells comprising a zinc finger/nuclease fusion molecule and a donor DNA molecule, are achieved by blocking the cells in the G₂ phase of the cell cycle, when homology-driven repair processes are maximally active. Such arrest can be achieved in a number of ways. For example, cells can be treated with e.g., drugs, compounds and/or small molecules which influence cell-cycle progression so as to arrest cells in G₂ phase. Exemplary molecules of this type include, but are not limited to, compounds which affect microtubule polymerization (e.g., vinblastine, nocodazole, Taxol), compounds that interact with DNA (e.g., cisplatin(II) diamine dichloride, Cisplatin, doxorubicin) and/or compounds that affect DNA synthesis (e.g., thymidine, hydroxyurea, L-mimosine, etoposide, 5-fluorouracil). Additional increases in recombination efficiency are achieved by the use of histone deacetylase (HDAC) inhibitors (e.g., sodium butyrate, trichostatin A) which alter chromatin structure to make genomic DNA more accessible to the cellular recombination machinery.

[0218] Additional methods for cell-cycle arrest include overexpression of proteins which inhibit the activity of the CDK cell-cycle kinases, for example, by introducing a cDNA encoding the protein into the cell or by introducing into the cell an engineered ZFP which activates expression of the gene encoding the protein. Cell-cycle arrest is also achieved by inhibiting the activity of cyclins and CDKs, for example, using RNAi methods (e.g., U.S. Pat. No. 6,506,559) or by introducing into the cell an engineered ZFP which represses expression of one or more genes involved in cell-cycle progression such as, for example, cyclin and/or CDK genes. See, e.g., co-owned U.S. Pat. No. 6,534,261 for methods for the synthesis of engineered zinc finger proteins for regulation of gene expression.

[0219] Alternatively, in certain cases, targeted cleavage is conducted in the absence of a donor polynucleotide (preferably in S or G₂ phase), and recombination occurs between homologous chromosomes.

Methods to Screen for Cellular Factors that Facilitate Homologous Recombination

[0220] Since homologous recombination is a multi-step process requiring the modification of DNA ends and the recruitment of several cellular factors into a protein complex, the addition of one or more exogenous factors, along with donor DNA and vectors encoding zinc finger-cleavage domain fusions, can be used to facilitate targeted homologous recombination. An exemplary method for identifying such a factor or factors employs analyses of gene expression using microarrays (e.g., Affymetrix Gene Chip® arrays) to compare the mRNA expression patterns of different cells. For example, cells that exhibit a higher capacity to stimulate double strand break-driven homologous recombination in the presence of donor DNA and zinc finger-cleavage domain fusions, either unaided or under conditions known to increase the level of gene correction, can be analyzed for their gene expression patterns compared to cells that lack such capacity. Genes that are upregulated or downregulated

in a manner that directly correlates with increased levels of homologous recombination are thereby identified and can be cloned into any one of a number of expression vectors. These expression constructs can be co-transfected along with zinc finger-cleavage domain fusions and donor constructs to yield improved methods for achieving high-efficiency homologous recombination. Alternatively, expression of such genes can be appropriately regulated using engineered zinc finger proteins which modulate expression (either activation or repression) of one or more these genes. See, e.g., co-owned U.S. Pat. No. 6,534,261 for methods for the synthesis of engineered zinc finger proteins for regulation of gene expression.

[0221] As an example, it was observed that the different clones obtained in the experiments described in Example 9 and FIG. 27 exhibited a wide-range of homologous recombination frequencies, when transfected with donor DNA and plasmids encoding zinc finger-cleavage domain fusions. Gene expression in clones showing a high frequency of targeted recombination can thus be compared to that in clones exhibiting a low frequency, and expression patterns unique to the former clones can be identified.

[0222] As an additional example, studies using cell cycle inhibitors (e.g., nocodazole or vinblastine, see e.g., Examples 11, 14 and 15) showed that cells arrested in the G2 phase of the cell cycle carried out homologous recombination at higher rates, indicating that cellular factors responsible for homologous recombination may be preferentially expressed or active in G2. One way to identify these factors is to compare the mRNA expression patterns between the stably transfected HEK 293 cell clones that carry out gene correction at high and low levels (e.g., clone T18 vs. clone T7). Similar comparisons are made between these cell lines in response to compounds that arrest the cells in G2 phase. Candidate genes that are differentially expressed in cells that carry out homologous recombination at a higher rate, either unaided or in response to compounds that arrest the cells in G2, are identified, cloned, and re-introduced into cells to determine whether their expression is sufficient to re-capitulate the improved rates. Alternatively, expression of said candidate genes is activated using engineered zinc finger transcription factors as described, for example, in co-owned U.S. Pat. No. 6,534,261.

Expression Vectors

[0223] A nucleic acid encoding one or more ZFPs or ZFP fusion proteins can be cloned into a vector for transformation into prokaryotic or eukaryotic cells for replication and/or expression. Vectors can be prokaryotic vectors, e.g., plasmids, or shuttle vectors, insect vectors, or eukaryotic vectors. A nucleic acid encoding a ZFP can also be cloned into an expression vector, for administration to a plant cell, animal cell, preferably a mammalian cell or a human cell, fungal cell, bacterial cell, or protozoal cell.

[0224] To obtain expression of a cloned gene or nucleic acid, sequences encoding a ZFP or ZFP fusion protein are typically subcloned into an expression vector that contains a promoter to direct transcription. Suitable bacterial and eukaryotic promoters are well known in the art and described, e.g., in Sambrook et al., *Molecular Cloning, A Laboratory Manual* (2nd ed. 1989; 3rd ed., 2001); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and *Current Protocols in Molecular Biology*

(Ausubel et al., supra. Bacterial expression systems for expressing the ZFP are available in, e.g., *E. coli*, *Bacillus sp.*, and *Salmonella* (Palva et al., *Gene* 22:229-235 (1983)). Kits for such expression systems are commercially available. Eukaryotic expression systems for mammalian cells, yeast, and insect cells are well known by those of skill in the art and are also commercially available.

[0225] The promoter used to direct expression of a ZFP-encoding nucleic acid depends on the particular application. For example, a strong constitutive promoter is typically used for expression and purification of ZFP. In contrast, when a ZFP is administered in vivo for gene regulation, either a constitutive or an inducible promoter is used, depending on the particular use of the ZFP. In addition, a preferred promoter for administration of a ZFP can be a weak promoter, such as HSV TK or a promoter having similar activity. The promoter typically can also include elements that are responsive to transactivation, e.g., hypoxia response elements, Gal4 response elements, lac repressor response element, and small molecule control systems such as tet-regulated systems and the RU-486 system (see, e.g., Gossen & Bujard, *PNAS* 89:5547 (1992); Oligino et al., *Gene Ther.* 5:491-496 (1998); Wang et al., *Gene Ther.* 4:432-441 (1997); Neering et al., *Blood* 88:1147-1155 (1996); and Rendahl et al., *Nat. Biotechnol.* 16:757-761 (1998)). The MNDU3 promoter can also be used, and is preferentially active in CD34⁺ hematopoietic stem cells.

[0226] In addition to the promoter, the expression vector typically contains a transcription unit or expression cassette that contains all the additional elements required for the expression of the nucleic acid in host cells, either prokaryotic or eukaryotic. A typical expression cassette thus contains a promoter operably linked, e.g., to a nucleic acid sequence encoding the ZFP, and signals required, e.g., for efficient polyadenylation of the transcript, transcriptional termination, ribosome binding sites, or translation termination. Additional elements of the cassette may include, e.g., enhancers, and heterologous splicing signals.

[0227] The particular expression vector used to transport the genetic information into the cell is selected with regard to the intended use of the ZFP, e.g., expression in plants, animals, bacteria, fungus, protozoa, etc. (see expression vectors described below). Standard bacterial expression vectors include plasmids such as pBR322-based plasmids, pSKF, pET23D, and commercially available fusion expression systems such as GST and LacZ. An exemplary fusion protein is the maltose binding protein, "MBP." Such fusion proteins are used for purification of the ZFP. Epitope tags can also be added to recombinant proteins to provide convenient methods of isolation, for monitoring expression, and for monitoring cellular and subcellular localization, e.g., c-myc or FLAG.

[0228] Expression vectors containing regulatory elements from eukaryotic viruses are often used in eukaryotic expression vectors, e.g., SV40 vectors, papilloma virus vectors, and vectors derived from Epstein-Barr virus. Other exemplary eukaryotic vectors include pMSG, pAV009/A+, pMTO10/A+, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the SV40 early promoter, SV40 late promoter, metallothionein promoter, murine mammary tumor virus

promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

[0229] Some expression systems have markers for selection of stably transfected cell lines such as thymidine kinase, hygromycin B phosphotransferase, and dihydrofolate reductase. High yield expression systems are also suitable, such as using a baculovirus vector in insect cells, with a ZFP encoding sequence under the direction of the polyhedrin promoter or other strong baculovirus promoters.

[0230] The elements that are typically included in expression vectors also include a replicon that functions in *E. coli*, a gene encoding antibiotic resistance to permit selection of bacteria that harbor recombinant plasmids, and unique restriction sites in nonessential regions of the plasmid to allow insertion of recombinant sequences.

[0231] Standard transfection methods are used to produce bacterial, mammalian, yeast or insect cell lines that express large quantities of protein, which are then purified using standard techniques (see, e.g., Colley et al., *J. Biol. Chem.* 264:17619-17622 (1989); *Guide to Protein Purification*, in *Methods in Enzymology*, vol. 182 (Deutscher, ed., 1990)). Transformation of eukaryotic and prokaryotic cells are performed according to standard techniques (see, e.g., Morrison, *J. Bact.* 132:349-351 (1977); Clark-Curtiss & Curtiss, *Methods in Enzymology* 101:347-362 (Wu et al., eds., 1983)).

[0232] Any of the well known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate transfection, polybrene, protoplast fusion, electroporation, ultrasonic methods (e.g., sonoporation), liposomes, microinjection, naked DNA, plasmid vectors, viral vectors, both episomal and integrative, and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (see, e.g., Sambrook et al., supra). It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one gene into the host cell capable of expressing the protein of choice.

Nucleic Acids Encoding Fusion Proteins and Delivery to Cells

[0233] Conventional viral and non-viral based gene transfer methods can be used to introduce nucleic acids encoding engineered ZFPs in cells (e.g., mammalian cells) and target tissues. Such methods can also be used to administer nucleic acids encoding ZFPs to cells in vitro. In certain embodiments, nucleic acids encoding ZFPs are administered for in vivo or ex vivo gene therapy uses. Non-viral vector delivery systems include DNA plasmids, naked nucleic acid, and nucleic acid complexed with a delivery vehicle such as a liposome or poloxamer. Viral vector delivery systems include DNA and RNA viruses, which have either episomal or integrated genomes after delivery to the cell. For a review of gene therapy procedures, see Anderson, *Science* 256:808-813 (1992); Nabel & Felgner, *TIBTECH* 11:211-217 (1993); Mitani & Caskey, *TIBTECH* 11:162-166 (1993); Dillon, *TIBTECH* 11: 167-175 (1993); Miller, *Nature* 357:455-460 (1992); Van Brunt, *Biotechnology* 6(10):1149-1154 (1988); Vigne, *Restorative Neurology and Neuroscience* 8:35-36 (1995); Kremer & Pericaudet, *British Medical Bulletin* 51(1):31-44 (1995); Haddada et al., in *Current Topics in*

Microbiology and Immunology Doerfler and Böhm (eds.) (1995); and Yu et al., *Gene Therapy* 1:13-26 (1994).

[0234] Methods of non-viral delivery of nucleic acids encoding engineered ZFPs include electroporation, lipofection, microinjection, biolistics, virosomes, liposomes, immunoliposomes, polycation or lipid:nucleic acid conjugates, naked DNA, artificial virions, and agent-enhanced uptake of DNA. Sonoporation using, e.g., the Sonitron 2000 system (Rich-Mar) can also be used for delivery of nucleic acids.

[0235] Additional exemplary nucleic acid delivery systems include those provided by Amaxa Biosystems (Cologne, Germany), Maxcyte, Inc. (Rockville, Md.) and BTX Molecular Delivery Systems (Holliston, Mass.).

[0236] Lipofection is described in e.g., US 5,049,386, US 4,946,787; and US 4,897,355 and lipofection reagents are sold commercially (e.g., Transfectam™ and Lipofectin™). Cationic and neutral lipids that are suitable for efficient receptor-recognition lipofection of polynucleotides include those of Felgner, WO 91/17424, WO 91/16024. Delivery can be to cells (ex vivo administration) or target tissues (in vivo administration).

[0237] The preparation of lipid:nucleic acid complexes, including targeted liposomes such as immunolipid complexes, is well known to one of skill in the art (see, e.g., Crystal, *Science* 270:404-410 (1995); Blaese et al., *Cancer Gene Ther.* 2:291-297 (1995); Behr et al., *Bioconjugate Chem.* 5:382-389 (1994); Remy et al., *Bioconjugate Chem.* 5:647-654 (1994); Gao et al., *Gene Therapy* 2:710-722 (1995); Ahmad et al., *Cancer Res.* 52:4817-4820 (1992); U.S. Pat. Nos. 4,186,183, 4,217,344, 4,235,871, 4,261,975, 4,485,054, 4,501,728, 4,774,085, 4,837,028, and 4,946,787).

[0238] The use of RNA or DNA viral based systems for the delivery of nucleic acids encoding engineered ZFPs take advantage of highly evolved processes for targeting a virus to specific cells in the body and trafficking the viral payload to the nucleus. Viral vectors can be administered directly to patients (in vivo) or they can be used to treat cells in vitro and the modified cells are administered to patients (ex vivo). Conventional viral based systems for the delivery of ZFPs include, but are not limited to, retroviral, lentivirus, adenoviral, adeno-associated, vaccinia and herpes simplex virus vectors for gene transfer. Integration in the host genome is possible with the retrovirus, lentivirus, and adeno-associated virus gene transfer methods, often resulting in long term expression of the inserted transgene. Additionally, high transduction efficiencies have been observed in many different cell types and target tissues.

[0239] The tropism of a retrovirus can be altered by incorporating foreign envelope proteins, expanding the potential target population of target cells. Lentiviral vectors are retroviral vectors that are able to transduce or infect non-dividing cells and typically produce high viral titers. Selection of a retroviral gene transfer system depends on the target tissue. Retroviral vectors are comprised of cis-acting long terminal repeats with packaging capacity for up to 6-10 kb of foreign sequence. The minimum cis-acting LTRs are sufficient for replication and packaging of the vectors, which are then used to integrate the therapeutic gene into the target cell to provide permanent transgene expression. Widely used

retroviral vectors include those based upon murine leukemia virus (MuLV), gibbon ape leukemia virus (GaLV), Simian Immunodeficiency virus (SIV), human immunodeficiency virus (HIV), and combinations thereof (see, e.g., Buchscher et al., *J. Virol.* 66:2731-2739 (1992); Johann et al., *J. Virol.* 66:1635-1640 (1992); Sommerfelt et al., *Virol.* 176:58-59 (1990); Wilson et al., *J. Virol.* 63:2374-2378 (1989); Miller et al., *J. Virol.* 65:2220-2224 (1991); PCT/US94/05700).

[0240] In applications in which transient expression of a ZFP fusion protein is preferred, adenoviral based systems can be used. Adenoviral based vectors are capable of very high transduction efficiency in many cell types and do not require cell division. With such vectors, high titer and high levels of expression have been obtained. This vector can be produced in large quantities in a relatively simple system. Adeno-associated virus ("AAV") vectors are also used to transduce cells with target nucleic acids, e.g., in the in vitro production of nucleic acids and peptides, and for in vivo and ex vivo gene therapy procedures (see, e.g., West et al., *Virology* 160:38-47 (1987); U.S. Pat. No. 4,797,368; WO 93/24641; Kotin, *Human Gene Therapy* 5:793-801 (1994); Muzyczka, *J. Clin. Invest.* 94:1351 (1994). Construction of recombinant AAV vectors are described in a number of publications, including U.S. Pat. No. 5,173,414; Tratschin et al., *Mol. Cell. Biol.* 5:3251-3260 (1985); Tratschin et al., *Mol. Cell. Biol.* 4:2072-2081 (1984); Hermonat & Muzyczka, *PNAS* 81:6466-6470 (1984); and Samulski et al., *J. Virol.* 63:03822-3828 (1989).

[0241] At least six viral vector approaches are currently available for gene transfer in clinical trials, which utilize approaches that involve complementation of defective vectors by genes inserted into helper cell lines to generate the transducing agent.

[0242] pLASN and MFG-S are examples of retroviral vectors that have been used in clinical trials (Dunbar et al., *Blood* 85:3048-305 (1995); Kohn et al., *Nat. Med.* 1:1017-102 (1995); Malech et al., *PNAS* 94:22 12133-12138 (1997)). PA317/pLASN was the first therapeutic vector used in a gene therapy trial. (Blaese et al., *Science* 270:475-480 (1995)). Transduction efficiencies of 50% or greater have been observed for MFG-S packaged vectors. (Ellem et al., *Immunol Immunother.* 44(1):10-20 (1997); Dranoff et al., *Hum. Gene Ther.* 1:111-2 (1997).

[0243] Recombinant adeno-associated virus vectors (rAAV) are a promising alternative gene delivery systems based on the defective and nonpathogenic parvovirus adeno-associated type 2 virus. All vectors are derived from a plasmid that retains only the AAV 145 bp inverted terminal repeats flanking the transgene expression cassette. Efficient gene transfer and stable transgene delivery due to integration into the genomes of the transduced cell are key features for this vector system. (Wagner et al., *Lancet* 351:9117 1702-3 (1998); Kearns et al., *Gene Ther.* 9:748-55 (1996)).

[0244] Replication-deficient recombinant adenoviral vectors (Ad) can be produced at high titer and readily infect a number of different cell types. Most adenovirus vectors are engineered such that a transgene replaces the Ad E1a, E1b, and/or E3 genes; subsequently the replication defective vector is propagated in human 293 cells that supply deleted gene function in trans. Ad vectors can transduce multiple types of tissues in vivo, including nondividing, differentiated cells such as those found in liver, kidney and muscle.

Conventional Ad vectors have a large carrying capacity. An example of the use of an Ad vector in a clinical trial involved polynucleotide therapy for antitumor immunization with intramuscular injection (Sterman et al., *Hum. Gene Ther.* 7:1083-9 (1998)). Additional examples of the use of adenovirus vectors for gene transfer in clinical trials include Rosenecker et al., *Infection* 24:1 5-10 (1996); Sterman et al., *Hum. Gene Ther.* 9:7 1083-1089 (1998); Welsh et al., *Hum. Gene Ther.* 2:205-18 (1995); Alvarez et al., *Hum. Gene Ther.* 5:597-613 (1997); Topf et al., *Gene Ther.* 5:507-513 (1998); Sterman et al., *Hum. Gene Ther.* 7:1083-1089 (1998).

[0245] Packaging cells are used to form virus particles that are capable of infecting a host cell. Such cells include 293 cells, which package adenovirus, and ψ 2 cells or PA317 cells, which package retrovirus. Viral vectors used in gene therapy are usually generated by a producer cell line that packages a nucleic acid vector into a viral particle. The vectors typically contain the minimal viral sequences required for packaging and subsequent integration into a host (if applicable), other viral sequences being replaced by an expression cassette encoding the protein to be expressed. The missing viral functions are supplied in trans by the packaging cell line. For example, AAV vectors used in gene therapy typically only possess inverted terminal repeat (ITR) sequences from the AAV genome which are required for packaging and integration into the host genome. Viral DNA is packaged in a cell line, which contains a helper plasmid encoding the other AAV genes, namely rep and cap, but lacking ITR sequences. The cell line is also infected with adenovirus as a helper. The helper virus promotes replication of the AAV vector and expression of AAV genes from the helper plasmid. The helper plasmid is not packaged in significant amounts due to a lack of ITR sequences. Contamination with adenovirus can be reduced by, e.g., heat treatment to which adenovirus is more sensitive than AAV.

[0246] In many gene therapy applications, it is desirable that the gene therapy vector be delivered with a high degree of specificity to a particular tissue type. Accordingly, a viral vector can be modified to have specificity for a given cell type by expressing a ligand as a fusion protein with a viral coat protein on the outer surface of the virus. The ligand is chosen to have affinity for a receptor known to be present on the cell type of interest. For example, Han et al., *Proc. Natl. Acad. Sci. USA* 92:9747-9751 (1995), reported that Moloney murine leukemia virus can be modified to express human heregulin fused to gp70, and the recombinant virus infects certain human breast cancer cells expressing human epidermal growth factor receptor. This principle can be extended to other virus-target cell pairs, in which the target cell expresses a receptor and the virus expresses a fusion protein comprising a ligand for the cell-surface receptor. For example, filamentous phage can be engineered to display antibody fragments (e.g., Fab or Fv) having specific binding affinity for virtually any chosen cellular receptor. Although the above description applies primarily to viral vectors, the same principles can be applied to nonviral vectors. Such vectors can be engineered to contain specific uptake sequences which favor uptake by specific target cells.

[0247] Gene therapy vectors can be delivered in vivo by administration to an individual patient, typically by systemic administration (e.g., intravenous, intraperitoneal, intramuscular, subdermal, or intracranial infusion) or topical application, as described below. Alternatively, vectors can be

delivered to cells *ex vivo*, such as cells explanted from an individual patient (e.g., lymphocytes, bone marrow aspirates, tissue biopsy) or universal donor hematopoietic stem cells, followed by reimplantation of the cells into a patient, usually after selection for cells which have incorporated the vector.

[0248] *Ex vivo* cell transfection for diagnostics, research, or for gene therapy (e.g., via re-infusion of the transfected cells into the host organism) is well known to those of skill in the art. In a preferred embodiment, cells are isolated from the subject organism, transfected with a ZFP nucleic acid (gene or cDNA), and re-infused back into the subject organism (e.g., patient). Various cell types suitable for *ex vivo* transfection are well known to those of skill in the art (see, e.g., Freshney et al., *Culture of Animal Cells, A Manual of Basic Technique* (3rd ed. 1994)) and the references cited therein for a discussion of how to isolate and culture cells from patients).

[0249] In one embodiment, stem cells are used in *ex vivo* procedures for cell transfection and gene therapy. The advantage to using stem cells is that they can be differentiated into other cell types *in vitro*, or can be introduced into a mammal (such as the donor of the cells) where they will engraft in the bone marrow. Methods for differentiating CD34+ cells *in vitro* into clinically important immune cell types using cytokines such as GM-CSF, IFN- γ and TNF- α are known (see Inaba et al., *J. Exp. Med.* 176:1693-1702 (1992)).

[0250] Stem cells are isolated for transduction and differentiation using known methods. For example, stem cells are isolated from bone marrow cells by panning the bone marrow cells with antibodies which bind unwanted cells, such as CD4+ and CD8+ (T cells), CD45+ (panb cells), GR-1 (granulocytes), and lad (differentiated antigen presenting cells) (see Inaba et al., *J. Exp. Med.* 176:1693-1702 (1992)).

[0251] Vectors (e.g., retroviruses, adenoviruses, liposomes, etc.) containing therapeutic ZFP nucleic acids can also be administered directly to an organism for transduction of cells *in vivo*. Alternatively, naked DNA can be administered. Administration is by any of the routes normally used for introducing a molecule into ultimate contact with blood or tissue cells including, but not limited to, injection, infusion, topical application and electroporation. Suitable methods of administering such nucleic acids are available and well known to those of skill in the art, and, although more than one route can be used to administer a particular composition, a particular route can often provide a more immediate and more effective reaction than another route.

[0252] Methods for introduction of DNA into hematopoietic stem cells are disclosed, for example, in U.S. Pat. No. 5,928,638. Vectors useful for introduction of transgenes into hematopoietic stem cells, e.g., CD34+ cells, include adenovirus Type 35.

[0253] Vectors suitable for introduction of transgenes into immune cells (e.g., T-cells) include non-integrating lentivirus vectors. See, for example, Ory et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:11382-11388; Dull et al. (1998) *J. Virol.* 72:8463-8471; Zuffery et al. (1998) *J. Virol.* 72:9873-9880; Follenzi et al. (2000) *Nature Genetics* 25:217-222.

[0254] Pharmaceutically acceptable carriers are determined in part by the particular composition being adminis-

tered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions available, as described below (see, e.g., *Remington's Pharmaceutical Sciences*, 17th ed., 1989).

[0255] DNA constructs may be introduced into the genome of a desired plant host by a variety of conventional techniques. For reviews of such techniques see, for example, Weissbach & Weissbach *Methods for Plant Molecular Biology* (1988, Academic Press, N.Y.) Section VIII, pp. 421-463; and Grierson & Corey, *Plant Molecular Biology* (1988, 2d Ed.), Blackie, London, Ch. 7-9. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using biolistic methods, such as DNA particle bombardment (see, e.g., Klein et al (1987) *Nature* 327:70-73). Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. *Agrobacterium tumefaciens*-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, for example Horsch et al (1984) *Science* 233:496-498, and Fraley et al (1983) *Proc. Nat'l. Acad. Sci. USA* 80:4803. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria using binary T DNA vector (Bevan (1984) *Nuc. Acid Res.* 12:8711-8721) or the co-cultivation procedure (Horsch et al (1985) *Science* 227:1229-1231). Generally, the *Agrobacterium* transformation system is used to engineer dicotyledonous plants (Bevan et al (1982) *Ann. Rev. Genet.* 16:357-384; Rogers et al (1986) *Methods Enzymol.* 118:627-641). The *Agrobacterium* transformation system may also be used to transform, as well as transfer, DNA to monocotyledonous plants and plant cells. See Hernalsteen et al (1984) *EMBO J* 3:3039-3041; Hooykass-Van Slogteren et al (1984) *Nature* 311:763-764; Grimsley et al (1987) *Nature* 325:1677-179; Boulton et al (1989) *Plant Mol. Biol.* 12:31-40.; and Gould et al (1991) *Plant Physiol.* 95:426-434.

[0256] Alternative gene transfer and transformation methods include, but are not limited to, protoplast transformation through calcium-, polyethylene glycol (PEG)- or electroporation-mediated uptake of naked DNA (see Paszkowski et al. (1984) *EMBO J* 3:2717-2722, Potrykus et al. (1985) *Molec. Gen. Genet.* 199:169-177; Fromm et al. (1985) *Proc. Nat. Acad. Sci. USA* 82:5824-5828; and Shimamoto (1989) *Nature* 338:274-276) and electroporation of plant tissues (D' Halluin et al. (1992) *Plant Cell* 4:1495-1505). Additional methods for plant cell transformation include microinjection, silicon carbide mediated DNA uptake (Kaeppeler et al. (1990) *Plant Cell Reporter* 9:415-418), and microprojectile bombardment (see Klein et al. (1988) *Proc. Nat. Acad. Sci. USA* 85:4305-4309; and Gordon-Kamm et al. (1990) *Plant Cell* 2:603-618).

[0257] The disclosed methods and compositions can be used to insert exogenous sequences into a predetermined location in a plant cell genome. This is useful inasmuch as expression of an introduced transgene into a plant genome depends critically on its integration site. Accordingly, genes encoding, e.g., nutrients, antibiotics or therapeutic mol-

ecules can be inserted, by targeted recombination, into regions of a plant genome favorable to their expression.

[0258] Transformed plant cells which are produced by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the transformed genotype and thus the desired phenotype. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans, et al., "Protoplasts Isolation and Culture" in *Handbook of Plant Cell Culture*, pp. 124-176, Macmillian Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, pollens, embryos or parts thereof. Such regeneration techniques are described generally in Klee et al (1987) *Ann. Rev. of Plant Phys.* 38:467-486.

[0259] Nucleic acids introduced into a plant cell can be used to confer desired traits on essentially any plant. A wide variety of plants and plant cell systems may be engineered for the desired physiological and agronomic characteristics described herein using the nucleic acid constructs of the present disclosure and the various transformation methods mentioned above. In preferred embodiments, target plants and plant cells for engineering include, but are not limited to, those monocotyledonous and dicotyledonous plants, such as crops including grain crops (e.g., wheat, maize, rice, millet, barley), fruit crops (e.g., tomato, apple, pear, strawberry, orange), forage crops (e.g., alfalfa), root vegetable crops (e.g., carrot, potato, sugar beets, yam), leafy vegetable crops (e.g., lettuce, spinach); flowering plants (e.g., petunia, rose, chrysanthemum), conifers and pine trees (e.g., pine fir, spruce); plants used in phytoremediation (e.g., heavy metal accumulating plants); oil crops (e.g., sunflower, rape seed) and plants used for experimental purposes (e.g., *Arabidopsis*). Thus, the disclosed methods and compositions have use over a broad range of plants, including, but not limited to, species from the genera *Asparagus*, *Avena*, *Brassica*, *Citrus*, *Citrullus*, *Capsicum*, *Cucurbita*, *Daucus*, *Glycine*, *Hordeum*, *Lactuca*, *Lycopersicon*, *Malus*, *Manihot*, *Nicotiana*, *Oryza*, *Persea*, *Pisum*, *Pyrus*, *Prunus*, *Raphanus*, *Secale*, *Solanum*, *Sorghum*, *Triticum*, *Vitis*, *Vigna*, and *Zea*.

[0260] One of skill in the art will recognize that after the expression cassette is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

[0261] A transformed plant cell, callus, tissue or plant may be identified and isolated by selecting or screening the engineered plant material for traits encoded by the marker genes present on the transforming DNA. For instance, selection may be performed by growing the engineered plant material on media containing an inhibitory amount of the antibiotic or herbicide to which the transforming gene construct confers resistance. Further, transformed plants and plant cells may also be identified by screening for the activities of any visible marker genes (e.g., the β -glucuronidase, luciferase, B or C1 genes) that may be present on

the recombinant nucleic acid constructs. Such selection and screening methodologies are well known to those skilled in the art.

[0262] Physical and biochemical methods also may be used to identify plant or plant cell transformants containing inserted gene constructs. These methods include but are not limited to: 1) Southern analysis or PCR amplification for detecting and determining the structure of the recombinant DNA insert; 2) Northern blot, S1 RNase protection, primer-extension or reverse transcriptase-PCR amplification for detecting and examining RNA transcripts of the gene constructs; 3) enzymatic assays for detecting enzyme or ribozyme activity, where such gene products are encoded by the gene construct; 4) protein gel electrophoresis, Western blot techniques, immunoprecipitation, or enzyme-linked immunoassays, where the gene construct products are proteins. Additional techniques, such as in situ hybridization, enzyme staining, and immunostaining, also may be used to detect the presence or expression of the recombinant construct in specific plant organs and tissues. The methods for doing all these assays are well known to those skilled in the art.

[0263] Effects of gene manipulation using the methods disclosed herein can be observed by, for example, northern blots of the RNA (e.g., mRNA) isolated from the tissues of interest. Typically, if the amount of mRNA has increased, it can be assumed that the corresponding endogenous gene is being expressed at a greater rate than before. Other methods of measuring gene and/or CYP74B activity can be used. Different types of enzymatic assays can be used, depending on the substrate used and the method of detecting the increase or decrease of a reaction product or by-product. In addition, the levels of and/or CYP74B protein expressed can be measured immunochemically, i.e., ELISA, RIA, EIA and other antibody based assays well known to those of skill in the art, such as by electrophoretic detection assays (either with staining or western blotting). The transgene may be selectively expressed in some tissues of the plant or at some developmental stages, or the transgene may be expressed in substantially all plant tissues, substantially along its entire life cycle. However, any combinatorial expression mode is also applicable.

[0264] The present disclosure also encompasses seeds of the transgenic plants described above wherein the seed has the transgene or gene construct. The present disclosure further encompasses the progeny, clones, cell lines or cells of the transgenic plants described above wherein said progeny, clone, cell line or cell has the transgene or gene construct.

Delivery Vehicles

[0265] An important factor in the administration of polypeptide compounds, such as ZFP fusion proteins, is ensuring that the polypeptide has the ability to traverse the plasma membrane of a cell, or the membrane of an intracellular compartment such as the nucleus. Cellular membranes are composed of lipid-protein bilayers that are freely permeable to small, nonionic lipophilic compounds and are inherently impermeable to polar compounds, macromolecules, and therapeutic or diagnostic agents. However, proteins and other compounds such as liposomes have been described, which have the ability to translocate polypeptides such as ZFPs across a cell membrane.

[0266] For example, “membrane translocation polypeptides” have amphiphilic or hydrophobic amino acid subsequences that have the ability to act as membrane-translocating carriers. In one embodiment, homeodomain proteins have the ability to translocate across cell membranes. The shortest internalizable peptide of a homeodomain protein, Antennapedia, was found to be the third helix of the protein, from amino acid position 43 to 58 (see, e.g., Prochiantz, *Current Opinion in Neurobiology* 6:629-634 (1996)). Another subsequence, the h (hydrophobic) domain of signal peptides, was found to have similar cell membrane translocation characteristics (see, e.g., Lin et al., *J. Biol. Chem.* 270:14255-14258 (1995)).

[0267] Examples of peptide sequences which can be linked to a protein, for facilitating uptake of the protein into cells, include, but are not limited to: an 11 amino acid peptide of the tat protein of HIV; a 20 residue peptide sequence which corresponds to amino acids 84-103 of the p16 protein (see Fahraeus et al., *Current Biology* 6:84 (1996)); the third helix of the 60-amino acid long homeodomain of Antennapedia (Derossi et al., *J. Biol. Chem.* 269:10444 (1994)); the h region of a signal peptide such as the Kaposi fibroblast growth factor (K-FGF) h region (Lin et al., supra); or the VP22 translocation domain from HSV (Elliot & O'Hare, *Cell* 88:223-233 (1997)). Other suitable chemical moieties that provide enhanced cellular uptake may also be chemically linked to ZFPs. Membrane translocation domains (i.e., internalization domains) can also be selected from libraries of randomized peptide sequences. See, for example, Yeh et al. (2003) *Molecular Therapy* 7(5):S461, Abstract #1191.

[0268] Toxin molecules also have the ability to transport polypeptides across cell membranes. Often, such molecules (called “binary toxins”) are composed of at least two parts: a translocation/binding domain or polypeptide and a separate toxin domain or polypeptide. Typically, the translocation domain or polypeptide binds to a cellular receptor, and then the toxin is transported into the cell. Several bacterial toxins, including *Clostridium perfringens* iota toxin, diphtheria toxin (DT), *Pseudomonas* exotoxin A (PE), pertussis toxin (PT), *Bacillus anthracis* toxin, and pertussis adenylate cyclase (CYA), have been used to deliver peptides to the cell cytosol as internal or amino-terminal fusions (Arora et al., *J. Biol. Chem.*, 268:3334-3341 (1993); Perelle et al., *Infect. Immun.*, 61:5147-5156 (1993); Stenmark et al., *J. Cell Biol.* 113:1025-1032 (1991); Donnelly et al., *PNAS* 90:3530-3534 (1993); Carbonetti et al., *Abstr. Annu. Meet. Am. Soc. Microbiol.* 95:295 (1995); Sebo et al., *Infect. Immun.* 63:3851-3857 (1995); Klimpel et al., *PNAS U.S.A.* 89:10277-10281 (1992); and Novak et al., *J. Biol. Chem.* 267:17186-17193 (1992)).

[0269] Such peptide sequences can be used to translocate ZFPs across a cell membrane. ZFPs can be conveniently fused to or derivatized with such sequences. Typically, the translocation sequence is provided as part of a fusion protein. Optionally, a linker can be used to link the ZFP and the translocation sequence. Any suitable linker can be used, e.g., a peptide linker.

[0270] The ZFP can also be introduced into an animal cell, preferably a mammalian cell, via a liposomes and liposome derivatives such as immunoliposomes. The term “liposome” refers to vesicles comprised of one or more concentrically

ordered lipid bilayers, which encapsulate an aqueous phase. The aqueous phase typically contains the compound to be delivered to the cell, i.e., a ZFP.

[0271] The liposome fuses with the plasma membrane, thereby releasing the drug into the cytosol. Alternatively, the liposome is phagocytosed or taken up by the cell in a transport vesicle. Once in the endosome or phagosome, the liposome either degrades or fuses with the membrane of the transport vesicle and releases its contents.

[0272] In current methods of drug delivery via liposomes, the liposome ultimately becomes permeable and releases the encapsulated compound (in this case, a ZFP) at the target tissue or cell. For systemic or tissue specific delivery, this can be accomplished, for example, in a passive manner wherein the liposome bilayer degrades over time through the action of various agents in the body. Alternatively, active drug release involves using an agent to induce a permeability change in the liposome vesicle. Liposome membranes can be constructed so that they become destabilized when the environment becomes acidic near the liposome membrane (see, e.g., *PNAS* 84:7851 (1987); *Biochemistry* 28:908 (1989)). When liposomes are endocytosed by a target cell, for example, they become destabilized and release their contents. This destabilization is termed fusogenesis. Dioleoylphosphatidylethanolamine (DOPE) is the basis of many “fusogenic” systems.

[0273] Such liposomes typically comprise a ZFP and a lipid component, e.g., a neutral and/or cationic lipid, optionally including a receptor-recognition molecule such as an antibody that binds to a predetermined cell surface receptor or ligand (e.g., an antigen). A variety of methods are available for preparing liposomes as described in, e.g., Szoka et al., *Ann. Rev. Biophys. Bioeng.* 9:467 (1980), U.S. Pat. Nos. 4,186,183, 4,217,344, 4,235,871, 4,261,975, 4,485,054, 4,501,728, 4,774,085, 4,837,028, 4,235,871, 4,261,975, 4,485,054, 4,501,728, 4,774,085, 4,837,028, 4,946,787, PCT Publication No. WO 91/17424, Deamer & Bangham, *Biochim. Biophys. Acta* 443:629-634 (1976); Fralley, et al., *PNAS* 76:3348-3352 (1979); Hope et al., *Biochim. Biophys. Acta* 812:55-65 (1985); Mayer et al., *Biochim. Biophys. Acta* 858:161-168 (1986); Williams et al., *PNAS* 85:242-246 (1988); *Liposomes (Ostro (ed.), 1983, Chapter 1)*; Hope et al., *Chem. Phys. Lip.* 40:89 (1986); Gregoriadis, *Liposome Technology* (1984) and Lasic, *Liposomes: from Physics to Applications* (1993)). Suitable methods include, for example, sonication, extrusion, high pressure/homogenization, microfluidization, detergent dialysis, calcium-induced fusion of small liposome vesicles and ether-fusion methods, all of which are known to those of skill in the art.

[0274] In certain embodiments, it is desirable to target liposomes using targeting moieties that are specific to a particular cell type, tissue, and the like. Targeting of liposomes using a variety of targeting moieties (e.g., ligands, receptors, and monoclonal antibodies) has been described. See, e.g., U.S. Pat. Nos. 4,957,773 and 4,603,044.

[0275] Examples of targeting moieties include monoclonal antibodies specific to antigens associated with neoplasms, such as prostate cancer specific antigen and MAGE. Tumors can also be diagnosed by detecting gene products resulting from the activation or over-expression of oncogenes, such as ras or c-erbB2. In addition, many tumors express antigens normally expressed by fetal tissue, such as


the alphafetoprotein (AFP) and carcinoembryonic antigen (CEA). Sites of viral infection can be diagnosed using various viral antigens such as hepatitis B core and surface antigens (HBVc, HBVs) hepatitis C antigens, Epstein-Barr virus antigens, human immunodeficiency type-1 virus (HIV1) and papilloma virus antigens. Inflammation can be detected using molecules specifically recognized by surface molecules which are expressed at sites of inflammation such as integrins (e.g., VCAM-1), selectin receptors (e.g., ELAM-1) and the like.

[0276] Standard methods for coupling targeting agents to liposomes can be used. These methods generally involve incorporation into liposomes of lipid components, e.g., phosphatidylethanolamine, which can be activated for attachment of targeting agents, or derivatized lipophilic compounds, such as lipid derivatized bleomycin. Antibody targeted liposomes can be constructed using, for instance, liposomes which incorporate protein A (see Renneisen et al., *J. Biol. Chem.*, 265:16337-16342 (1990) and Leonetti et al., *PNAS* 87:2448-2451 (1990)).

Dosages

[0277] For therapeutic applications, the dose administered to a patient, or to a cell which will be introduced into a patient, in the context of the present disclosure, should be sufficient to effect a beneficial therapeutic response in the patient over time. In addition, particular dosage regimens can be useful for determining phenotypic changes in an experimental setting, e.g., in functional genomics studies, and in cell or animal models. The dose will be determined by the efficacy and K_d of the particular ZFP employed, the nuclear volume of the target cell, and the condition of the patient, as well as the body weight or surface area of the patient to be treated. The size of the dose also will be determined by the existence, nature, and extent of any adverse side-effects that accompany the administration of a particular compound or vector in a particular patient.

[0278] The maximum therapeutically effective dosage of ZFP for approximately 99% binding to target sites is calculated to be in the range of less than about 1.5×10^5 to 1.5×10^6 copies of the specific ZFP molecule per cell. The number of ZFPs per cell for this level of binding is calculated as follows, using the volume of a HeLa cell nucleus (approximately $1000 \mu\text{m}^3$ or 10^{-12} L; *Cell Biology*, (Altman & Katz, eds. (1976)). As the HeLa nucleus is relatively large, this dosage number is recalculated as needed using the volume of the target cell nucleus. This calculation also does not take into account competition for ZFP binding by other sites. This calculation also assumes that essentially all of the ZFP is localized to the nucleus. A value of $100 \times K_d$ is used to calculate approximately 99% binding of to the target site, and a value of $10 \times K_d$ is used to calculate approximately 90% binding of to the target site. For this example, $K_d = 25 \text{ nM}$



i.e., $\text{DNA} + \text{protein} \leftrightarrow \text{DNA:protein complex}$

$$K_d = \frac{[\text{DNA}][\text{protein}]}{[\text{DNA:protein complex}]}$$

when 50% of ZFP is bound, $K_d = [\text{protein}]$

-continued

So when $[\text{protein}] = 25 \text{ nM}$ and the nucleus volume is 10^{-12} L

$[\text{protein}] =$

$$(25 \times 10^{-9} \text{ moles/L})(10^{-12} \text{ L/nucleus})(6 \times 10^{23} \text{ molecules/mole}) =$$

15,000 molecules/nucleus for 50% binding

When 99% target is bound; $100 \times K_d = [\text{protein}]$

$$100 \times K_d = [\text{protein}] = 2.5 \mu\text{M}$$

$$(2.5 \times 10^{-6} \text{ moles/L})(10^{-12} \text{ L/nucleus})(6 \times 10^{23} \text{ molecules/mole}) =$$

about 1,500,000 molecule per

nucleus for 99% binding of target site.

[0279] The appropriate dose of an expression vector encoding a ZFP can also be calculated by taking into account the average rate of ZFP expression from the promoter and the average rate of ZFP degradation in the cell. In certain embodiments, a weak promoter such as a wild-type or mutant HSV TK promoter is used, as described above. The dose of ZFP in micrograms is calculated by taking into account the molecular weight of the particular ZFP being employed.

[0280] In determining the effective amount of the ZFP to be administered in the treatment or prophylaxis of disease, the physician evaluates circulating plasma levels of the ZFP or nucleic acid encoding the ZFP, potential ZFP toxicities, progression of the disease, and the production of anti-ZFP antibodies. Administration can be accomplished via single or divided doses.

Pharmaceutical Compositions and Administration

[0281] ZFPs and expression vectors encoding ZFPs can be administered directly to the patient for targeted cleavage and/or recombination, and for therapeutic or prophylactic applications, for example, cancer, ischemia, diabetic retinopathy, macular degeneration, rheumatoid arthritis, psoriasis, HIV infection, sickle cell anemia, Alzheimer's disease, muscular dystrophy, neurodegenerative diseases, vascular disease, cystic fibrosis, stroke, and the like. Examples of microorganisms that can be inhibited by ZFP gene therapy include pathogenic bacteria, e.g., *chlamydia*, rickettsial bacteria, mycobacteria, *staphylococci*, *streptococci*, *pneumococci*, *meningococci* and *conococci*, *klebsiella*, *proteus*, *seratia*, *pseudomonas*, *legionella*, *diphtheria*, *salmonella*, *bacilli*, *cholera*, *tetanus*, *botulism*, *anthrax*, *plague*, *leptospirosis*, and Lyme disease bacteria; infectious fungus, e.g., *Aspergillus*, *Candida* species; protozoa such as sporozoa (e.g., *Plasmodia*), rhizopods (e.g., *Entamoeba*) and flagellates (*Trypanosoma*, *Leishmania*, *Trichomonas*, *Giardia*, etc.); viral diseases, e.g., hepatitis (A, B, or C), herpes virus (e.g., VZV, HSV-1, HSV-6, HSV-II, CMV, and EBV), HIV, Ebola, adenovirus, influenza virus, flaviviruses, echovirus, rhinovirus, coxsackie virus, coronavirus, respiratory syncytial virus, mumps virus, rotavirus, measles virus, rubella virus, parvovirus, vaccinia virus, HTLV virus, dengue virus, papillomavirus, poliovirus, rabies virus, and arboviral encephalitis virus, etc.

[0282] Administration of therapeutically effective amounts is by any of the routes normally used for introduc-

ing ZFP into ultimate contact with the tissue to be treated. The ZFPs are administered in any suitable manner, preferably with pharmaceutically acceptable carriers. Suitable methods of administering such modulators are available and well known to those of skill in the art, and, although more than one route can be used to administer a particular composition, a particular route can often provide a more immediate and more effective reaction than another route.

[0283] Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions that are available (see, e.g., *Remington's Pharmaceutical Sciences*, 17th ed. 1985)).

[0284] The ZFPs, alone or in combination with other suitable components, can be made into aerosol formulations (i.e., they can be "nebulized") to be administered via inhalation. Aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like.

[0285] Formulations suitable for parenteral administration, such as, for example, by intravenous, intramuscular, intradermal, and subcutaneous routes, include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The disclosed compositions can be administered, for example, by intravenous infusion, orally, topically, intraperitoneally, intravesically or intrathecally. The formulations of compounds can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials. Injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.

Applications

[0286] The disclosed methods and compositions for targeted cleavage can be used to induce mutations in a genomic sequence, e.g., by cleaving at two sites and deleting sequences in between, by cleavage at a single site followed by non-homologous end joining, cleaving at one or two sites with insertion of an exogenous sequence between the breaks and/or by cleaving at a site so as to remove one or two or a few nucleotides. Targeted cleavage can also be used to create gene knock-outs (e.g., for functional genomics or target validation) and to facilitate targeted insertion of a sequence into a genome (i.e., gene knock-in); e.g., for purposes of cell engineering or protein overexpression. Insertion can be by means of replacements of chromosomal sequences through homologous recombination or by targeted integration, in which a new sequence (i.e., a sequence not present in the region of interest), flanked by sequences homologous to the region of interest in the chromosome, is inserted at a predetermined target site.

[0287] The same methods can also be used to replace a wild-type sequence with a mutant sequence, or to convert one allele to a different allele.

[0288] Targeted cleavage of infecting or integrated viral genomes can be used to treat viral infections in a host.

Additionally, targeted cleavage of genes encoding receptors for viruses can be used to block expression of such receptors, thereby preventing viral infection and/or viral spread in a host organism. Targeted mutagenesis of genes encoding viral receptors (e.g., the CCR5 and CXCR4 receptors for HIV) can be used to render the receptors unable to bind to virus, thereby preventing new infection and blocking the spread of existing infections. Non-limiting examples of viruses or viral receptors that may be targeted include herpes simplex virus (HSV), such as HSV-1 and HSV-2, varicella zoster virus (VZV), Epstein-Barr virus (EBV) and cytomegalovirus (CMV), HHV6 and HHV7. The hepatitis family of viruses includes hepatitis A virus (HAV), hepatitis B virus (HBV), hepatitis C virus (HCV), the delta hepatitis virus (HDV), hepatitis E virus (HEV) and hepatitis G virus (HGV). Other viruses or their receptors may be targeted, including, but not limited to, Picomaviridae (e.g., polioviruses, etc.); Caliciviridae; Togaviridae (e.g., rubella virus, dengue virus, etc.); Flaviviridae; Coronaviridae; Reoviridae; Bimaviridae; Rhabdoviridae (e.g., rabies virus, etc.); Filoviridae; Paramyxoviridae (e.g., mumps virus, measles virus, respiratory syncytial virus, etc.); Orthomyxoviridae (e.g., influenza virus types A, B and C, etc.); Bunyaviridae; Arenaviridae; Retroviridae; lentiviruses (e.g., HTLV-I; HTLV-II; HIV-1 (also known as HTLV-III, LAV, ARV, hTLR, etc.) HIV-II); simian immunodeficiency virus (SIV), human papillomavirus (HPV), influenza virus and the tick-borne encephalitis viruses. See, e.g. *Virology*, 3rd Edition (W. K. Joklik ed. 1988); *Fundamental Virology*, 2nd Edition (B. N. Fields and D. M. Knipe, eds. 1991), for a description of these and other viruses. Receptors for HIV, for example, include CCR-5 and CXCR-4.

[0289] In similar fashion, the genome of an infecting bacterium can be mutagenized by targeted DNA cleavage followed by non-homologous end joining, to block or ameliorate bacterial infections.

[0290] The disclosed methods for targeted recombination can be used to replace any genomic sequence with a homologous, non-identical sequence. For example, a mutant genomic sequence can be replaced by its wild-type counterpart, thereby providing methods for treatment of e.g., genetic disease, inherited disorders, cancer, and autoimmune disease. In like fashion, one allele of a gene can be replaced by a different allele using the methods of targeted recombination disclosed herein.

[0291] Exemplary genetic diseases include, but are not limited to, achondroplasia, achromatopsia, acid maltase deficiency, adenosine deaminase deficiency (OMIM No.102700), adrenoleukodystrophy, aicardi syndrome, alpha-1 antitrypsin deficiency, alpha-thalassemia, androgen insensitivity syndrome, apert syndrome, arrhythmogenic right ventricular, dysplasia, ataxia telangiectasia, Barth syndrome, beta-thalassemia, blue rubber bleb nevus syndrome, canavan disease, chronic granulomatous diseases (CGD), cri du chat syndrome, cystic fibrosis, dercun's disease, ectodermal dysplasia, Fanconi anemia, fibrodysplasia ossificans progressive, fragile X syndrome, galactosemia, Gaucher's disease, generalized gangliosidosis (e.g., GM1), hemochromatosis, the hemoglobin C mutation in the 6th codon of beta-globin (HbC), hemophilia, Huntington's disease, Hurler Syndrome, hypophosphatase, Klinefelter syndrome, Krabbes Disease, Langer-Giedion Syndrome, leukocyte adhesion deficiency (LAD, OMIM No. 116920), leukodys-

trophy, long QT syndrome, Marfan syndrome, Moebius syndrome, mucopolysaccharidosis (MPS), nail patella syndrome, nephrogenic diabetes insipidus, neurofibromatosis, Neimann-Pick disease, osteogenesis imperfecta, porphyria, Prader-Willi syndrome, progeria, Proteus syndrome, retinoblastoma, Rett syndrome, Rubinstein-Taybi syndrome, Sanfilippo syndrome, severe combined immunodeficiency (SCID), Shwachman syndrome, sickle cell disease (sickle cell anemia), Smith-Magenis syndrome, Stickler syndrome, Tay-Sachs disease, Thrombocytopenia Absent Radius (TAR) syndrome, Treacher Collins syndrome, trisomy, tuberous sclerosis, Turner's syndrome, urea cycle disorder, von Hippel-Landau disease, Waardenburg syndrome, Williams syndrome, Wilson's disease, Wiskott-Aldrich syndrome, X-linked lymphoproliferative syndrome (XLP, OMIM No. 308240).

[0292] Additional exemplary diseases that can be treated by targeted DNA cleavage and/or homologous recombination include acquired immunodeficiencies, lysosomal storage diseases (e.g., Gaucher's disease, GM1, Fabry disease and Tay-Sachs disease), mucopolysaccharidosis (e.g. Hunter's disease, Hurler's disease), hemoglobinopathies (e.g., sickle cell diseases, HbC, α -thalassemia, β -thalassemia) and hemophilias.

[0293] In certain cases, alteration of a genomic sequence in a pluripotent cell (e.g., a hematopoietic stem cell) is desired. Methods for mobilization, enrichment and culture of hematopoietic stem cells are known in the art. See for example, U.S. Pat. Nos. 5,061,620; 5,681,559; 6,335,195; 6,645,489 and 6,667,064. Treated stem cells can be returned to a patient for treatment of various diseases including, but not limited to, SCID and sickle-cell anemia.

[0294] In many of these cases, a region of interest comprises a mutation, and the donor polynucleotide comprises the corresponding wild-type sequence. Similarly, a wild-type genomic sequence can be replaced by a mutant sequence, if such is desirable. For example, overexpression of an oncogene can be reversed either by mutating the gene or by replacing its control sequences with sequences that support a lower, non-pathologic level of expression. As another example, the wild-type allele of the ApoA1 gene can be replaced by the ApoA1 Milano allele, to treat atherosclerosis. Indeed, any pathology dependent upon a particular genomic sequence, in any fashion, can be corrected or alleviated using the methods and compositions disclosed herein.

[0295] Targeted cleavage and targeted recombination can also be used to alter non-coding sequences (e.g., regulatory sequences such as promoters, enhancers, initiators, terminators, splice sites) to alter the levels of expression of a gene product. Such methods can be used, for example, for therapeutic purposes, functional genomics and/or target validation studies.

[0296] The compositions and methods described herein also allow for novel approaches and systems to address immune reactions of a host to allogeneic grafts. In particular, a major problem faced when allogeneic stem cells (or any type of allogeneic cell) are grafted into a host recipient is the high risk of rejection by the host's immune system, primarily mediated through recognition of the Major Histocompatibility Complex (MHC) on the surface of the engrafted cells. The MHC comprises the HLA class I protein(s) that

function as heterodimers that are comprised of a common β subunit and variable α subunits. It has been demonstrated that tissue grafts derived from stem cells that are devoid of HLA escape the host's immune response. See, e.g., Coffman et al. *J Immunol* 151, 425-35. (1993); Markmann et al. *Transplantation* 54, 1085-9. (1992); Koller et al. *Science* 248, 1227-30. (1990). Using the compositions and methods described herein, genes encoding HLA proteins involved in graft rejection can be cleaved, mutagenized or altered by recombination, in either their coding or regulatory sequences, so that their expression is blocked or they express a non-functional product. For example, by inactivating the gene encoding the common β subunit gene ($\beta 2$ microglobulin) using ZFP fusion proteins as described herein, HLA class I can be removed from the cells to rapidly and reliably generate HLA class I null stem cells from any donor, thereby reducing the need for closely matched donor/recipient MHC haplotypes during stem cell grafting.

[0297] Inactivation of any gene (e.g., the $\beta 2$ microglobulin gene) can be achieved, for example, by a single cleavage event, by cleavage followed by non-homologous end joining, by cleavage at two sites followed by joining so as to delete the sequence between the two cleavage sites, by targeted recombination of a missense or nonsense codon into the coding region, or by targeted recombination of an irrelevant sequence (i.e., a "stuffer" sequence) into the gene or its regulatory region, so as to disrupt the gene or regulatory region.

[0298] Targeted modification of chromatin structure, as disclosed in co-owned WO 01/83793, can be used to facilitate the binding of fusion proteins to cellular chromatin.

[0299] In additional embodiments, one or more fusions between a zinc finger binding domain and a recombinase (or functional fragment thereof) can be used, in addition to or instead of the zinc finger-cleavage domain fusions disclosed herein, to facilitate targeted recombination. See, for example, co-owned U.S. Pat. No. 6,534,261 and Akopian et al. (2003) *Proc. Natl. Acad. Sci. USA* 100:8688-8691.

[0300] In additional embodiments, the disclosed methods and compositions are used to provide fusions of ZFP binding domains with transcriptional activation or repression domains that require dimerization (either homodimerization or heterodimerization) for their activity. In these cases, a fusion polypeptide comprises a zinc finger binding domain and a functional domain monomer (e.g., a monomer from a dimeric transcriptional activation or repression domain). Binding of two such fusion polypeptides to properly situated target sites allows dimerization so as to reconstitute a functional transcription activation or repression domain.

Targeted Integration

[0301] As disclosed above, the methods and compositions set forth herein can be used for targeted integration of exogenous sequences into a region of interest in the genome of a cell. Targeted integration of an exogenous sequence at a double-strand break in a genome can occur by both homology-dependent and homology-independent mechanisms.

[0302] As noted above, in certain embodiments, targeted integration by both homology-dependent and homology-independent mechanisms involves insertion of an exogenous sequence between the ends generated by cleavage. The

exogenous sequence inserted can be any length, for example, a relatively short "patch" sequence of between 1 and 50 nucleotides in length (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45 or 50 nucleotide sequence).

[0303] In cases in which targeted integration is homology-dependent, a donor nucleic acid or donor sequence comprises an exogenous sequence together with one or more sequences that are either identical, or homologous but non-identical, with a predetermined genomic sequence (i.e., a target site). In certain embodiments two of the identical sequences or two of the homologous but non-identical sequences (or one of each) are present, flanking the exogenous sequence. An exogenous sequence (or exogenous nucleic acid or exogenous polynucleotide) is one that contains a nucleotide sequence that is not normally present in the region of interest.

[0304] Exemplary exogenous sequences include, but are not limited to, cDNAs, promoter sequences, enhancer sequences, epitope tags, marker genes, cleavage enzyme recognition sites and various types of expression constructs. Marker genes include, but are not limited to, sequences encoding proteins that mediate antibiotic resistance (e.g., ampicillin resistance, neomycin resistance, G418 resistance, puromycin resistance), sequences encoding colored or fluorescent or luminescent proteins (e.g., green fluorescent protein, enhanced green fluorescent protein, red fluorescent protein, luciferase), and proteins which mediate enhanced cell growth and/or gene amplification (e.g., dihydrofolate reductase). Epitope tags include, for example, one or more copies of FLAG, His, myc, Tap, HA or any detectable amino acid sequence.

[0305] Protein expression constructs include, but are not limited to, cDNAs and transcriptional control sequences in operative linkage with cDNA sequences. Transcriptional control sequences include promoters, enhancers and insulators. Additional transcriptional and translational regulatory sequences which can be included in expression constructs include, e.g., internal ribosome entry sites, sequences encoding 2A peptides and polyadenylation signals. An exemplary protein expression construct is an antibody expression construct comprising a sequence encoding an antibody heavy chain and a sequence encoding an antibody light chain, each sequence operatively linked to a promoter (the promoters being the same or different) and either or both sequences optionally operatively linked to an enhancer (and, in the case of both coding sequences being linked to enhancers, the enhancers being the same or different).

[0306] Cleavage enzyme recognition sites include, for example, sequences recognized by restriction endonucleases, homing endonucleases and/or meganucleases. Targeted integration of a cleavage enzyme recognition site (by either homology-dependent or homology-independent mechanisms) is useful for generating cells whose genome contains only a single site that can be cleaved by a particular enzyme. Contacting such cells with an enzyme that recognizes and cleaves at the single site facilitates subsequent targeted integration of exogenous sequences (by either homology-dependent or homology-independent mechanisms) and/or targeted mutagenesis at the site that is cleaved.

[0307] One example of a cleavage enzyme recognition site is that recognized by the homing endonuclease I-SceI, which has the following sequence:

TAGGGATAACAGGGTAAT (SEQ ID NO:213)

See, for example, U.S. Pat. No. 6,833,252. Additional exemplary homing endonucleases include I-CeuI, PI-PspI, PI-Sce, I-SceIV, I-Csml, I-PanI, I-SceII, I-PpoI, I-SceIII, I-CreI, I-TevI, I-TevII and I-TevIII. Their recognition sequences are known. See also U.S. Pat. No. 5,420,032; Belfort et al. (1997) *Nucleic Acids Res.* 25:3379-3388; Dujon et al. (1989) *Gene* 82:115-118; Perler et al. (1994) *Nucleic Acids Res.* 22, 1125-1127; Jasin (1996) *Trends Genet.* 12:224-228; Gimble et al. (1996) *J. Mol. Biol.* 263:163-180; Argast et al. (1998) *J. Mol. Biol.* 280:345-353 and the New England Biolabs catalogue.

[0308] Although the cleavage specificity of most homing endonucleases is not absolute with respect to their recognition sites, the sites are of sufficient length that a single cleavage event per mammalian-sized genome can be obtained by expressing a homing endonuclease in a cell containing a single copy of its recognition site. It has also been reported that cleavage enzymes can be engineered to bind non-natural target sites. See, for example, Chevalier et al. (2002) *Molec. Cell* 10:895-905; Epinat et al. (2003) *Nucleic Acids Res.* 31:2952-2962; Ashworth et al. (2006) *Nature* 441:656-659.

[0309] Previous methods for obtaining targeted recombination and integration using homing endonucleases suffered from the problem that targeted insertion of the recognition site is extremely inefficient, requiring laborious screening to identify cells that contained the recognition site inserted at the desired location. The present methods surmount these problems by allowing highly-efficient targeted integration (either homology-dependent or homology-independent) of a recognition site for a DNA-cleaving enzyme.

[0310] In certain embodiments, targeted integration is used to insert a RNA expression construct, e.g., sequences responsible for regulated expression of micro RNA or siRNA. Promoters, enhancers and additional transcription regulatory sequences, as described above, can also be incorporated in a RNA expression construct.

[0311] In embodiments in which targeted integration occurs by a homology-dependent mechanism, the donor sequence contains sufficient homology, in the regions flanking the exogenous sequence, to support homology-directed repair of a double-strand break in a genomic sequence, thereby inserting the exogenous sequence at the genomic target site. Therefore, the donor nucleic acid can be of any size sufficient to support integration of the exogenous sequence by homology-dependent repair mechanisms (e.g., homologous recombination). Without wishing to be bound by any particular theory, the regions of homology flanking the exogenous sequence are thought to provide the broken chromosome ends with a template for re-synthesis of the genetic information at the site of the double-stranded break.

[0312] Targeted integration of exogenous sequences, as disclosed herein, can be used to generate cells and cell lines for protein expression. See, for example, co-owned U.S. Patent Application Publication No. 2006/0063231 (the dis-

closure of which is hereby incorporated by reference herein, in its entirety, for all purposes). For optimal expression of one or more proteins encoded by exogenous sequences integrated into a genome, the chromosomal integration site should be compatible with high-level transcription of the integrated sequences, preferably in a wide range of cell types and developmental states. However, it has been observed that transcription of integrated sequences varies depending on the integration site due to, among other things, the chromatin structure of the genome at the integration site. Accordingly, genomic target sites that support high-level transcription of integrated sequences are desirable. In certain embodiments, it will also be desirable that integration of exogenous sequences not result in ectopic activation of one or more cellular genes (e.g., oncogenes). On the other hand, in the case of integration of promoter and/or enhancer sequences, ectopic expression may be desired.

[0313] For certain embodiments, it is desirable that an integration site is not present in an essential gene (e.g., a gene essential for cell viability), so that inactivation of said essential gene does not result from integration of the exogenous sequences. On the other hand, if the intent is to disable gene function (i.e., create a gene “knock-out”) targeted integration of an exogenous sequence to disrupt an endogenous gene is an effective method. In these cases, the exogenous sequence can be any sequence capable of blocking transcription of the endogenous gene or of generating a non-functional translation product, for example a short patch of amino acid sequence, which is optionally detectable (see above). In certain embodiments, the exogenous sequences can comprise a marker gene (described above), allowing selection of cells that have undergone targeted integration.

[0314] Non-limiting examples of chromosomal regions that do not encode an essential gene and support high-level transcription of sequences integrated therein (“safe harbor” integration sites) include the Rosa26 and CCR5 loci.

[0315] The Rosa26 locus has been identified in the murine genome. Zambrowicz et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:3789-3794. The sequence of mouse Rosa26 mRNA was compared to data from a human cDNA screen (Strausberg et al. (2002) *Proc. Natl. Acad. Sci. USA* 99:16899-16903), and a homologous human transcript was detected by the present inventors. Accordingly, the human homologue of Rosa26 can be used as a target site for integration of exogenous sequences into the genome of human cells and cell lines, using the methods and compositions disclosed herein.

[0316] CCR5 genomic sequences (including allelic variants such as CCR5-Δ32) are well known in the art. See, e.g., Liu et al. (1996) *Cell* 367-377.

[0317] Additional genomic target sites supporting high-level transcription of integrated sequences can be identified as regions of open chromatin or “accessible regions” as described, for example in co-owned U.S. Patent Application Publications 2002/0064802 (May 30, 2002) and 2002/0081603 (Jun. 27, 2002).

[0318] The presence of a double-stranded break in a genomic sequence facilitates not only homology-dependent integration of exogenous sequences (i.e., homologous recombination) but also homology-independent integration of exogenous sequences into the genome at the site of the

double-strand break. Accordingly, the compositions and methods disclosed herein can be used for targeted cleavage of a genomic sequence, followed by non-homology-dependent integration of an exogenous sequence at or near the targeted cleavage site. For example, a cell can be contacted with one or more ZFP-cleavage domain (or cleavage half-domain) fusion proteins engineered to cleave in a region of interest in a genome as described herein (or one or more polynucleotides encoding such fusion proteins), and a polynucleotide comprising an exogenous sequence lacking homology to the region of interest, to obtain a cell in which all or a portion of the exogenous sequence is integrated in the region of interest.

[0319] The methods of targeted integration (i.e., insertion of an exogenous sequence into a genome), both homology-dependent and -independent, disclosed herein can be used for a number of purposes. These include, but are not limited to, insertion of a gene or cDNA sequence into the genome of a cell to enable expression of the transcription and/or translation products of the gene or cDNA by the cell. For situations in which a disease or pathology can result from one of a plurality of mutations (e.g., multiple point mutations spread across the sequence of the gene), targeted integration (either homology-dependent or homology-independent) of a cDNA copy of the wild-type gene is particularly effective. For example, such a wild-type cDNA is inserted into an untranslated leader sequence or into the first exon of a gene upstream of all known mutations. In certain integrants, in which translational reading frame is preserved, the result is that the wild-type cDNA is expressed and its expression is regulated by the appropriate endogenous transcriptional regulatory sequences. In additional embodiments, such integrated cDNA sequences can include transcriptional (and/or translational) termination signals disposed downstream of the wild-type cDNA and upstream of the mutant endogenous gene. In this way, a wild-type copy of the disease-causing gene is expressed, and the mutant endogenous gene is not expressed. In other embodiments, a portion of a wild-type cDNA is inserted into the appropriate region of a gene (for example, a gene in which disease-causing mutations are clustered).

EXAMPLES

Example 1

Editing of a Chromosomal hSMC1L1 Gene by Targeted Recombination

[0320] The hSMC1L1 gene is the human orthologue of the budding yeast gene structural maintenance of chromosomes 1. A region of this gene encoding an amino-terminal portion of the protein which includes the Walker ATPase domain was mutagenized by targeted cleavage and recombination. Cleavage was targeted to the region of the methionine initiation codon (nucleotides 24-26, FIG. 1), by designing chimeric nucleases, comprising a zinc finger DNA-binding domain and a FokI cleavage half-domain, which bind in the vicinity of the codon. Thus, two zinc finger binding domains were designed, one of which recognizes nucleotides 23-34 (primary contacts along the top strand as shown in FIG. 1), and the other of which recognizes nucleotides 5-16 (primary contacts along the bottom strand). Zinc finger proteins were designed as described in co-owned U.S. Pat. Nos. 6,453,242

and 6,534,261. See Table 2 for the amino acid sequences of the recognition regions of the zinc finger proteins.

[0321] Sequences encoding each of these two ZFP binding domains were fused to sequences encoding a FokI cleavage half-domain (amino acids 384-579 of the native FokI sequence; Kita et al. (1989) *J. Biol. Chem.* 264:5751-5756), such that the encoded protein contained FokI sequences at the carboxy terminus and ZFP sequences at the amino terminus. Each of these fusion sequences was then cloned in a modified mammalian expression vector pcDNA3 (FIG. 2).

TABLE 2

Zinc Finger Designs for the hSMC1L1 Gene				
Target sequence	F1	F2	F3	F4
CATGGGGTTCCT (SEQ ID NO:27)	RSHDLIE (SEQ ID NO: 28)	TSSSLSR (SEQ ID NO:29)	RSDHLST (SEQ ID NO:30)	TNSNRIT (SEQ ID NO:31)
GCGGCGCCGGCG (SEQ ID NO:32)	RSDDLRS (SEQ ID NO:33)	RSDDRKT (SEQ ID NO:34)	RSEDLIR (SEQ ID NO:35)	RSDTLRS (SEQ ID NO:36)

Note:

The zinc finger amino acid sequences shown above (in one-letter code) represent residues -1 through +6, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein, and Finger F4 is closest to the carboxy terminus.

[0322] A donor DNA molecule was obtained as follows. First, a 700 base pair fragment of human genomic DNA representing nucleotides 52415936-52416635 of the “-” strand of the X chromosome (UCSC human genome release July, 2003), which includes the first exon of the human hSMC1L1 gene, was amplified, using genomic DNA from HEK293 cells as template. Sequences of primers used for amplification are shown in Table 3 (“Initial amp 1” and “Initial amp 2”). The PCR product was then altered, using standard overlap extension PCR methodology (see, e.g., Ho, et al. (1989) *Gene* 77:51-59), resulting in replacement of the sequence ATGGGG (nucleotides 24-29 in FIG. 1) to ATAA-GAAGC. This change resulted in conversion of the ATG codon (methionine) to an ATA codon (isoleucine) and replacement of GGG (nucleotides 27-29 in FIG. 1) by the sequence AGAAGC, allowing discrimination between donor-derived sequences and endogenous chromosomal sequences following recombination. A schematic diagram of the hSMC1 gene, including sequences of the chromosomal DNA in the region of the initiation codon, and sequences in the donor DNA that differ from the chromosomal sequence, is given in FIG. 3. The resulting 700 base pair donor fragment was cloned into pCR4BluntTopo, which does not contain any sequences homologous to the human genome. See FIG. 4.

[0323] For targeted mutation of the chromosomal hSMC1L1 gene, the two plasmids encoding ZFP-FokI fusions and the donor plasmid were introduced into 1×10⁶ HEK293 cells by transfection using Lipofectamine 2000® (Invitrogen). Controls included cells transfected only with the two plasmids encoding the ZFP-FokI fusions, cells transfected only with the donor plasmid and cells transfected with a control plasmid (pEGFP-N1, Clontech). Cells were cultured in 5% CO₂ at 37° C. At 48 hours after transfection, genomic DNA was isolated from the cells, and 200 ng was used as template for PCR amplification, using one primer complementary to a region of the gene outside of its region

of homology with the donor sequences (nucleotides 52416677-52416701 on the “-” STRAND of the X chromosome; UCSC July 2003), and a second primer complementary to a region of the donor molecule into which distinguishing mutations were introduced. Using these two primers, an amplification product of 400 base pairs will be obtained from genomic DNA if a targeted recombination event has occurred. The sequences of these primers are given in Table 3 (labeled “chromosome-specific” and “donor-specific,” respectively). Conditions for amplification

were: 94° C., 2 min, followed by 40 cycles of 94° C., 30 sec, 60° C., 1 min, 72° C., 1 min; and a final step of 72° C., 7 min.

[0324] The results of this analysis (FIG. 5) indicate that a 400 base pair amplification product (labeled “Chimeric DNA” in the Figure) was obtained only with DNA extracted from cells which had been transfected with the donor plasmid and both ZFP-FokI plasmids.

TABLE 3

Amplification Primers for the hSMC1L1 Gene	
Initial amp 1	AGCAACAACCTCCCGGGGATC (SEQ ID NO:37)
Initial amp 2	TTCCAGACGCGACTCTTTGGC (SEQ ID NO: 38)
Chromosome-specific	CTCAGCAAGCGTGAGCTCAGGTCTC (SEQ ID NO: 39)
Donor-specific	CAATCAGTTTCAGGAAGCTTCTT (SEQ ID NO: 40)
Outside 1	CTCAGCAAGCGTGAGCTCAGGTCTC (SEQ ID NO: 41)
Outside 2	GGGGTCAAGTAAGGCTGGGAAGC (SEQ ID NO: 42)

[0325] To confirm this result, two additional experiments were conducted. First, the amplification product was cloned into pCR4Blunt-Topo (Invitrogen) and its nucleotide sequence was determined. As shown in FIG. 6 (SEQ ID NO: 6), the amplified sequence obtained from chromosomal DNA of cells transfected with the two ZFP-FokI-encoding plasmids and the donor plasmid contains the AAGAAGC sequence that is unique to the donor (nucleotides 395-401 of the sequence presented in FIG. 6) covalently linked to chromosomal sequences not present in the donor molecule

(nucleotides 32-97 of FIG. 6), indicating that donor sequences have been recombined into the chromosome. In particular, the G→A mutation converting the initiation codon to an isoleucine codon is observed at position 395 in the sequence.

[0326] In a second experiment, chromosomal DNA from cells transfected only with donor plasmid, cells transfected with both ZFP-FokI fusion plasmids, cells transfected with the donor plasmid and both ZFP-FokI fusion plasmids or cells transfected with the EGFP control plasmid was used as template for amplification, using primers complementary to sequences outside of the 700-nucleotide region of homology between donor and chromosomal sequences (identified as “Outside 1” and “Outside 2” in Table 3). The resulting amplification product was purified and used as template for a second amplification reaction using the donor-specific and chromosome-specific primers described above (Table 3). This amplification yielded a 400 nucleotide product only from cells transfected with the donor construct and both ZFP-FokI fusion constructs, a result consistent with the replacement of genomic sequences by targeted recombination in these cells.

protein designed to bind to nucleotides 29-40 (primary contacts along the top strand as shown in FIG. 7) and a zinc finger protein designed to bind to nucleotides 8-20 (primary contacts along the bottom strand). The second pair (third and fourth rows of Table 4) comprises two zinc finger proteins, the first of which recognizes nucleotides 23-34 (primary contacts along the top strand as shown in FIG. 7) and the second of which recognizes nucleotides 8-16 (primary contacts along the bottom strand). Zinc finger proteins were designed as described in co-owned U.S. Pat. Nos. 6,453,242 and 6,534,261. See Table 4 for the amino acid sequences of the recognition regions of the zinc finger proteins.

[0328] Sequences encoding the ZFP binding domains were fused to sequences encoding a FokI cleavage half-domain (amino acids 384-579 of the native FokI sequence, Kita et al., supra), such that the encoded protein contained FokI sequences at the carboxy terminus and ZFP sequences at the amino terminus. Each of these fusion sequences was then cloned in a modified mammalian expression vector pcDNA3. See FIG. 8 for a schematic diagram of the constructs.

TABLE 4

Zinc Finger Designs for the IL2R γ Gene				
Target sequence	F1	F2	F3	F4
AAC TCGGATAAT (SEQ ID NO:43)	DRSTLIE (SEQ ID NO:44)	SSSNLSR (SEQ ID NO:45)	RSDDL SK (SEQ ID NO:46)	DNSNRIK (SEQ ID NO:47)
TAGAGG aGAAAGG (SEQ ID NO:48)	RSDNLSN (SEQ ID NO:49)	TSSSRIN (SEQ ID NO:50)	RSDHLSQ (SEQ ID NO:51)	RNADRKT (SEQ ID NO:52)
TACAAGAACTCG (SEQ ID NO:53)	RSDDL SK (SEQ ID NO:54)	DNSNRIK (SEQ ID NO:55)	RSDALSV (SEQ ID NO:56)	DNANRTK (SEQ ID NO:57)
GGAGAAAGG (SEQ ID NO:58)	RSDHLTQ (SEQ ID NO:59)	QSGNLAR (SEQ ID NO:60)	RSDHLSR (SEQ ID NO:61)	

Note:

The zinc finger amino acid sequences shown above (in one-letter code) represent residues -1 through +6, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein.

Example 2

Editing of a Chromosomal IL2R γ Gene by Targeted Recombination

[0327] The IL-2R γ gene encodes a protein, known as the “common cytokine receptor gamma chain,” that functions as a subunit of several interleukin receptors (including IL-2R, IL-4R, IL-7R, IL-9R, IL-15R and IL-21R). Mutations in this gene, including those surrounding the 5' end of the third exon (e.g. the tyrosine 91 codon), can cause X-linked severe combined immunodeficiency (SCID). See, for example, Puck et al. (1997) Blood 89:1968-1977. A mutation in the tyrosine 91 codon (nucleotides 23-25 of SEQ ID NO: 7; FIG. 7), was introduced into the IL2R γ gene by targeted cleavage and recombination. Cleavage was targeted to this region by designing two pairs of zinc finger proteins. The first pair (first two rows of Table 4) comprises a zinc finger

[0329] A donor DNA molecule was obtained as follows. First, a 700 base pair fragment of human DNA corresponding to positions 69196910-69197609 on the “-” strand of the X chromosome (UCSC, July 2003), which includes exon 3 of the IL2R γ gene, was amplified, using genomic DNA from K562 cells as template. See FIG. 9. Sequences of primers used for amplification are shown in Table 5 (labeled initial amp 1 and initial amp 2). The PCR product was then altered via standard overlap extension PCR methodology (Ho, et al., supra) to replace the sequence TACAAGAACTCGGATAAT (SEQ ID NO:62) with the sequence TAAAAGAATTCCGACAAC (SEQ ID NO:63). This replacement results in the introduction of a point mutation at nucleotide 25 (FIG. 7), converting the tyrosine 91 codon TAC to a TAA termination codon and enables discrimination between donor-derived and endogenous chromosomal sequences following recombination, because of differences in the sequences downstream of codon 91. The resulting 700

base pair fragment was cloned into pCR4BluntTopo which does not contain any sequences homologous to the human genome. See FIG. 10.

[0330] For targeted mutation of the chromosomal IL2R γ gene, the donor plasmid, along with two plasmids each encoding one of a pair of ZFP-FokI fusions, were introduced into 2 $\times$ 10⁶ K652 cells using mixed lipofection/electroporation (Amaxa). Each of the ZFP/FokI pairs (see Table 4) was tested in separate experiments. Controls included cells transfected only with two plasmids encoding ZFP-FokI fusions, and cells transfected only with the donor plasmid. Cells were cultured in 5% CO₂ at 37° C. At 48 hours after transfection, genomic DNA was isolated from the cells, and 200 ng was used as template for PCR amplification, using one primer complementary to a region of the gene outside of its region of homology with the donor sequences (nucleotides 69196839-69196863 on the “+” strand of the X chromosome; UCSC, July 2003), and a second primer complementary to a region of the donor molecule into which distinguishing mutations were introduced (see above) and whose sequence therefore diverges from that of chromosomal DNA. See Table 5 for primer sequences, labeled “chromosome-specific” and “donor-specific,” respectively. Using these two primers, an amplification product of 500 bp is obtained from genomic DNA in which a targeted recombination event has occurred. Conditions for amplification were: 94° C., 2 min, followed by 35 cycles of 94° C., 30 sec, 62° C., 1 min, 72° C., 45 sec; and a final step of 72° C., 7 min.

[0331] The results of this analysis (FIG. 11) indicate that an amplification product of the expected size (500 base pairs) is obtained with DNA extracted from cells which had been transfected with the donor plasmid and either of the pairs of ZFP-FokI-encoding plasmids. DNA from cells transfected with plasmids encoding a pair of ZFPs only (no donor plasmid) did not result in generation of the 500 bp product, nor did DNA from cells transfected only with the donor plasmid.

TABLE 5

Amplification Primers for the IL2R γ Gene	
Initial amp 1	TGTCGAGTACATGAATTGCACTTGG (SEQ ID NO:64)
Initial amp 2	TTAGGTTCTCTGGAGCCCAGGG (SEQ ID NO:65)

TABLE 5-continued

Amplification Primers for the IL2R γ Gene	
Chromosome-specific	CTCCAAACAGTGGTTCAAGAATCTG (SEQ ID NO:66)
Donor-specific	TCCTCTAGGTAAAGAATTCCGACAAC (SEQ ID NO:67)

[0332] To confirm this result, the amplification product obtained from the experiment using the second pair of ZFP/FokI fusions was cloned into pCR4Blunt-Topo (Invitrogen) and its nucleotide sequence was determined. As shown in FIG. 12 (SEQ ID NO: 12), the sequence consists of a fusion between chromosomal sequences and sequences from the donor plasmid. In particular, the G to A mutation converting tyrosine 91 to a stop codon is observed at position 43 in the sequence. Positions 43-58 contain nucleotides unique to the donor; nucleotides 32-42 and 59-459 are sequences common to the donor and the chromosome, and nucleotides 460-552 are unique to the chromosome. The presence of donor-unique sequences covalently linked to sequences present in the chromosome but not in the donor indicates that DNA from the donor plasmid was introduced into the chromosome by homologous recombination.

Example 3

Editing of a Chromosomal P-globin Gene by Targeted Recombination

[0333] The human beta globin gene is one of two gene products responsible for the structure and function of hemoglobin in adult human erythrocytes. Mutations in the beta-globin gene can result in sickle cell anemia. Two zinc finger proteins were designed to bind within this sequence, near the location of a nucleotide which, when mutated, causes sickle cell anemia. FIG. 13 shows the nucleotide sequence of a portion of the human beta-globin gene, and the target sites for the two zinc finger proteins are underlined in the sequence presented in FIG. 13. Amino acid sequences of the recognition regions of the two zinc finger proteins are shown in Table 6. Sequences encoding each of these two ZFP binding domains were fused to sequences encoding a FokI cleavage half-domain, as described above, to create engineered ZFP-nucleases that targeted the endogenous beta globin gene. Each of these fusion sequences was then cloned in the mammalian expression vector pcDNA3.1 (FIG. 14).

TABLE 6

Zinc Finger Designs for the beta-globin Gene				
Target sequence	F1	F2	F3	F4
GGGCAGTAACGG (SEQ ID NO: 68)	RSDHLSE (SEQ ID NO:69)	QSANRTK (SEQ ID NO:70)	RSDNLSA (SEQ ID NO:71)	RSQNRTR (SEQ ID NO:72)

TABLE 6-continued

<u>Zinc Finger Designs for the beta-globin Gene</u>				
Target sequence	F1	F2	F3	F4
AAGGTGAACGTG (SEQ ID NO:73)	RSDSLSR (SEQ ID NO:74)	DSSNRKT (SEQ ID NO:75)	RSDSLSA (SEQ ID NO:76)	RNDNRKT (SEQ ID NO:77)

Note:

The zinc finger amino acid sequences shown above (in one-letter code) represent residues -1 through +6, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein, and Finger F4 is closest to the carboxy terminus.

[0334] A donor DNA molecule was obtained as follows. First, a 700 base pair fragment of human genomic DNA corresponding to nucleotides 5212134- 5212833 on the “-” strand of Chromosome 11 (BLAT, UCSC Human Genome site) was amplified by PCR, using genomic DNA from K562 cells as template. Sequences of primers used for amplification are shown in Table 7 (labeled initial amp 1 and initial amp 2). The resulting amplified fragment contains sequences corresponding to the promoter, the first two exons and the first intron of the human beta globin gene. See FIG. 15 for a schematic illustrating the locations of exons 1 and 2, the first intron, and the primer binding sites in the beta globin sequence. The cloned product was then further modified by PCR to introduce a set of sequence changes between nucleotides 305-336 (as shown in FIG. 13), which replaced the sequence CCGTTACTGCCCTGTGGGGCAAGGTGAACGTG (SEQ ID NO: 78) with gCGTTAgTGC-CCGAATCCGAtcGTcAACcAc (SEQ ID NO: 79) (changes in bold). Certain of these changes (shown in lowercase) were specifically engineered to prevent the ZFP/FokI fusion proteins from binding to and cleaving the donor sequence, once integrated into the chromosome. In addition, all of the sequence changes enable discrimination between donor and endogenous chromosomal sequences following recombination. The resulting 700 base pair fragment was cloned into pCR4-TOPO, which does not contain any sequences homologous to the human genome (FIG. 16).

[0335] For targeted mutation of the chromosomal beta globin gene, the two plasmids encoding ZFP-FokI fusions and the donor plasmid (pCR4-TOPO-HBBdonor) were introduced into 1×10⁶ K562 cells by transfection using Nucleofector™ Solution (Amaxa Biosystems). Controls included cells transfected only with 100 ng (low) or 200 ng (high) of the two plasmids encoding the ZFP-FokI fusions, cells transfected only with 200 ng (low) or 600 ng (high) of the donor plasmid, cells transfected with a GFP-encoding plasmid, and mock transfected cells. Cells were cultured in RPMI Medium 1640 (Invitrogen), supplemented with 10% fetal bovine serum (FBS) (Hyclone) and 2 mM L-glutamine. Cells were maintained at 37° C. in an atmosphere of 5% CO₂. At 72 hours after transfection, genomic DNA was isolated from the cells, and 200 ng was used as template for PCR amplification, using one primer complementary to a region of the gene outside of its region of homology with the donor sequences (nucleotides 5212883-5212905 on the “-” strand of chromosome 11), and a second primer complementary to a region of the donor molecule into which distinguishing mutations were introduced into the donor sequence (see supra). The sequences of these primers are

given in Table 7 (labeled “chromosome-specific” and “donor-specific,” respectively). Using these two primers, an amplification product of 415 base pairs will be obtained from genomic DNA if a targeted recombination event has occurred. As a control for DNA loading, PCR reactions were also carried out using the Initial amp 1 and Initial amp 2 primers to ensure that similar levels of genomic DNA were added to each PCR reaction. Conditions for amplification were: 95° C., 2 min, followed by 40 cycles of 95° C., 30 sec, 60° C., 45 sec, 68° C., 2 min; and a final step of 68° C., 10 min.

[0336] The results of this analysis (FIG. 17) indicate that a 415 base pair amplification product was obtained only with DNA extracted from cells which had been transfected with the “high” concentration of donor plasmid and both ZFP-FokI plasmids, consistent with targeted recombination of donor sequences into the chromosomal beta-globin locus.

TABLE 7

Amplification Primers for the human beta globin gene	
Initial amp 1	TACTGATGGTATGGGGCCAAGAG (SEQ ID NO:80)
Initial amp 2	CACGTGCAGCTTGTCACAGTGC (SEQ ID NO:81)
Chromosome-specific	TGCTTACCAAGCTGTGATTCCA (SEQ ID NO:82)
Donor-specific	GGTTGACGATCGGAATTC (SEQ ID NO:83)

[0337] To confirm this result, the amplification product was cloned into pCR4-TOPO (Invitrogen) and its nucleotide sequence was determined. As shown in FIG. 18 (SEQ ID NO: 14), the sequence consists of a fusion between chromosomal sequences not present on the donor plasmid and sequences unique to the donor plasmid. For example, two C→G mutations which disrupt ZFP-binding are observed at positions 377 and 383 in the sequence. Nucleotides 377-408 represent sequence obtained from the donor plasmid containing the sequence changes described above; nucleotides 73-376 are sequences common to the donor and the chromosome, and nucleotides 1-72 are unique to the chromosome. The covalent linkage of donor-specific and chromosome-specific sequences in the genome confirms the successful recombination of the donor sequence at the correct locus within the genome of K562 cells.

Example 4

ZFP-FokI Linker (ZC Linker) Optimization

[0338] In order to test the effect of ZC linker length on cleavage efficiency, a four-finger ZFP binding domain was fused to a FokI cleavage half-domain, using ZC linkers of various lengths. The target site for the ZFP is 5'-AACTCG-GATAAT-3' (SEQ ID NO:84) and the amino acid sequences of the recognition regions (positions -1 through +6 with respect to the start of the alpha-helix) of each of the zinc fingers were as follows (wherein F1 is the N-most, and F4 is the C-most zinc finger):

F1: DRSTLIE	(SEQ ID NO:85)
F2: SSSNLSR	(SEQ ID NO:86)
F3: RSDDLK	(SEQ ID NO:87)
F4: DNSNRRIK	(SEQ ID NO:88)

[0339] ZFP-FokI fusions, in which the aforementioned ZFP binding domain and a FokI cleavage half-domain were separated by 2, 3, 4, 5, 6, or 10 amino acid residues, were constructed. Each of these proteins was tested for cleavage of substrates having an inverted repeat of the ZFP target site, with repeats separated by 4, 5, 6, 7, 8, 9, 12, 15, 16, 17, 22, or 26 basepairs.

[0340] The amino acid sequences of the fusion constructs, in the region of the ZFP-FokI junction (with the ZC linker sequence underlined), are as follows:

10-residue linker	HTKIHLROKDA <u>ARG</u> SQLV (SEQ ID NO:89)
6-residue linker	HTKIHLROK <u>GS</u> SQLV (SEQ ID NO:90)
5-residue linker	HTKIHLRO <u>GS</u> SQLV (SEQ ID NO:91)
4-residue linker	HTKIHLR <u>GS</u> SQLV (SEQ ID NO:92)
3-residue linker	HTKIHL <u>GS</u> SQLV (SEQ ID NO:93)
2-residue linker	HTKIHL <u>GS</u> SQLV (SEQ ID NO:94)

[0341] The sequences of the various cleavage substrates, with the ZFP target sites underlined, are as follows:

4bp separation	CTAGCATTATCCGAGTTACACA <u>AACTCGGATAATG</u> CTAGGATCGTAATAGGCTCAATGTGTTGAGCCTA TTACGATC (SEQ ID NO:95)
5bp separation	CTAGCATTATCCGAGTTCACACA <u>AACTCGGATAATG</u> CTAGGATCGTAATAGGCTCAAGTGTGTTGAGCCTA TTACGATC (SEQ ID NO:96)
6bp separation	CTAGGCATTATCCGAGTTCACACA <u>AACTCGGATAA</u> TGACTAGGATCCGTAATAGGCTCAAGTGGTGTG AGCCTATTACTGATC (SEQ ID NO:97)
7bp separation	CTAGCATTATCCGAGTTCACACA <u>AACTCGGATA</u> ATGCTAGGATCGTAATAGGCTCAAGTGTGTTG AGCCTATTACGATC (SEQ ID NO:98)

-continued

8bp separation	CTAGCATTATCCGAGTTCACACACA <u>AACTCGGAT</u> AATGCTAGGATCGTAATAGGCTCAAGTGGTGTGT TGAGCCTATTACGATC (SEQ ID NO:99)
9bp separation	CTAGCATTATCCGAGTTCACACACA <u>AACTCGGA</u> TAATGCTAGGATCGTAATAGGCTCAAGTGTGTGT GTTGAGCCTATTACGATC (SEQ ID NO:100)
12bp separation	CTAGCATTATCCGAGTTCACACCAACACA <u>AACTC</u> GGATAATGCTAGGATCGTAATAGGCTCAAGTGGT GGTTGTGTGAGCCTATTACGATC (SEQ ID NO:101)
15bp separation	CTAGCATTATCCGAGTTCACACCAACCACACA <u>AA</u> CTCGGATAATGCTAGGATCGTAATAGGCTCAAGT GGTGGTGGTGTGTTGAGCCTATTACGATC (SEQ ID NO:102)
16bp separation	CTAGCATTATCCGAGTTCACACCAACCACACCA <u>AA</u> ACTCGGATAATGCTAGGATCGTAATAGGCTCAAG TGGTGGTGGTGTGTTGAGCCTATTACGATC (SEQ ID NO:103)
17bp separation	CTAGCATTATCCGAGTTCACACCAACCACACCA <u>AA</u> AACTCGGATAATGCTAGGATCGTAATAGGCTCAAG GTTGGTGGTGGTGTGTTGAGCCTATTACGATC (SEQ ID NO:104)
22bp separation	CTAGCATTATCCGAGTTCACACCAACCACACCA <u>AA</u> AACACA <u>AACTCGGATAATGCTAGGATCGTAATAGG</u> CTCAAAGTTGGTGGTGGTGGTGGTGGTGGTGGTGG TATTACGATC (SEQ ID NO:105)
26bp separation	CTAGCATTATCCGAGTTCACACCAACCACACCA <u>AA</u> AACACCACA <u>AACTCGGATAATGCTAGGATCGTAA</u> TAGGCTCAAGTGGTGGTGGTGGTGGTGGTGGTGGTGG GTTGAGCCTATTACGATC (SEQ ID NO:106)

[0342] Plasmids encoding the different ZFP-FokI fusion proteins (see above) were constructed by standard molecular biological techniques, and an in vitro coupled transcription/translation system was used to express the encoded proteins. For each construct, 200 ng linearized plasmid DNA was incubated in 20 μ L TnT mix and incubated at 30° C. for 1 hour and 45 minutes. TnT mix contains 100 μ L TnT lysate (Promega, Madison, Wis.) with 4 μ L T7 RNA polymerase (Promega)+2 μ L Methionine (1 mM)+2.5 μ L ZnCl₂ (20 mM).

[0343] For analysis of DNA cleavage by the different ZFP-FokI fusions, 1 μ L of the coupled transcription/translation reaction mixture was combined with approximately 1 ng DNA substrate (end-labeled with ³²P using T4 polynucleotide kinase), and the mixture was diluted to a final volume of 19 μ L with FokI Cleavage Buffer. FokI Cleavage buffer contains 20 mM Tris-HCl pH 8.5, 75 mM NaCl, 10 μ M ZnCl₂, 1 mM DTT, 5% glycerol, 500 μ g/ml BSA. The mixture was incubated for 1 hour at 37° C. 6.5 μ L of FokI buffer, also containing 8 mM MgCl₂, was then added and incubation was continued for one hour at 37° C. Protein was extracted by adding 10 μ L phenol-chloroform solution to each reaction, mixing, and centrifuging to separate the phases. Ten microliters of the aqueous phase from each reaction was analyzed by electrophoresis on a 10% polyacrylamide gel.

[0344] The gel was subjected to autoradiography, and the cleavage efficiency for each ZFP-FokI fusion/substrate pair was calculated by quantifying the radioactivity in bands corresponding to uncleaved and cleaved substrate, summing to obtain total radioactivity, and determining the percentage of the total radioactivity present in the bands representing cleavage products.

[0345] The results of this experiment are shown in Table 8. This data allows the selection of a ZC linker that provides optimum cleavage efficiency for a given target site separation. This data also allows the selection of linker lengths that allow cleavage at a selected pair of target sites, but discriminate against cleavage at the same or similar ZFP target sites that have a separation that is different from that at the intended cleavage site.

TABLE 8

DNA cleavage efficiency for various ZC linker lengths and various binding site separations*						
	2-residue	3-residue	4-residue	5-residue	6-residue	10-residue
4 bp	74%	81%	74%	12%	6%	4%
5 bp	61%	89%	92%	80%	53%	40%
6 bp	78%	89%	95%	91%	93%	76%
7 bp	15%	55%	80%	80%	70%	80%
8 bp	0%	0%	8%	11%	22%	63%
9 bp	2%	6%	23%	9%	13%	51%
12 bp	8%	12%	22%	40%	69%	84%
15 bp	73%	78%	97%	92%	95%	88%
16 bp	59%	89%	100%	97%	90%	86%
17 bp	5%	22%	77%	71%	85%	82%
22 bp	1%	3%	5%	8%	18%	58%
26 bp	1%	2%	35%	36%	84%	78%

*The columns represent different ZFP-FokI fusion constructs with the indicated number of residues separating the ZFP and the FokI cleavage half-domain. The rows represent different DNA substrates with the indicated number of basepairs separating the inverted repeats of the ZFP target site.

[0346] For ZFP-FokI fusions with four residue linkers, the amino acid sequence of the linker was also varied. In separate constructs, the original LRGS linker sequence (SEQ ID NO:107) was changed to LGGS (SEQ ID NO:108), TGGS (SEQ ID NO:109), GGGS (SEQ ID NO:110), LPGS (SEQ ID NO: 111), LRKS (SEQ ID NO:112), and LRWS (SEQ ID NO:113); and the resulting fusions were tested on substrates having a six-basepair separation between binding sites. Fusions containing the LGGS (SEQ ID NO:108) linker sequence were observed to cleave more efficiently than those containing the original LRGS sequence(SEQ ID NO:107). Fusions containing the LRKS(SEQ ID NO:112) and LRWS(SEQ ID NO:113) sequences cleaved with less efficiency than the LRGS sequence(SEQ ID NO:107), while the cleavage efficiencies of the remaining fusions were similar to that of the fusion comprising the original LRGS sequence(SEQ ID NO:107).

Example 5

Increased Cleavage Specificity Resulting from Alteration of the FokI Cleavage Half-domain in the Dimerization Interface

[0347] A pair of ZFP/FokI fusion proteins (denoted 5-8 and 5-10) were designed to bind to target sites in the fifth exon of the IL-2R γ gene, to promote cleavage in the region between the target sites. The relevant region of the gene, including the target sequences of the two fusion proteins, is shown in FIG. 19. The amino acid sequence of the 5-8 protein is shown in FIG. 20, and the amino acid sequence of the 5-10 protein is shown in FIG. 21. Both proteins contain a 10 amino acid ZC linker. With respect to the zinc finger portion of these proteins, the DNA target sequences, as well as amino acid sequences of the recognition regions in the zinc fingers, are given in Table 9.

TABLE 9

Zinc Finger Designs for the IL2R γ Gene					
Fusion	Target sequence	F1	F2	F3	F4
5-8	ACTCTGTGGAAG (SEQ ID NO:114)	RSDNLSE (SEQ ID NO:115)	RNAHRIN (SEQ ID NO:116)	RSDTLSE (SEQ ID NO:117)	ARSTRIT (SEQ ID NO:118)
5-10	AACACGaAACGTG (SEQ ID NO:119)	RSDSLSR (SEQ ID NO:120)	DSSNRKT (SEQ ID NO:121)	RSDSLSV (SEQ ID NO:122)	DRSNRIT (SEQ ID NO:123)

Note:

The zinc finger amino acid sequences shown above (in one-letter code) represent residues -1 through +6, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein.

[0348] The ability of this pair of fusion proteins to catalyze specific cleavage of DNA between their target sequences (see FIG. 19) was tested in vitro using a labeled DNA template containing the target sequence and assaying for the presence of diagnostic digestion products. Specific cleavage was obtained when both proteins were used (Table 10, first row). However, the 5-10 fusion protein (comprising a wild-type FokI cleavage half-domain) was also capable of aberrant cleavage at a non-target site in the absence of the 5-8 protein (Table 10, second row), possibly due to self-dimerization.

[0349] Accordingly, 5-10 was modified in its FokI cleavage half-domain by converting amino acid residue 490 from glutamic acid (E) to lysine (K). (Numbering of amino acid residues in the FokI protein is according to Wah et al., supra.) This modification was designed to prevent homodimerization by altering an amino acid residue in the dimerization interface. The 5-10 (E490K) mutant, unlike the parental 5-10 protein, was unable to cleave at aberrant sites in the absence of the 5-8 fusion protein (Table 10, Row 3). However, the 5-10 (E490K) mutant, together with the 5-8 protein, catalyzed specific cleavage of the substrate (Table 10, Row 4). Thus, alteration of a residue in the cleavage half-domain of 5-10, that is involved in dimerization, prevented aberrant cleavage by this fusion protein due to self-dimerization. An E490R mutant also exhibits lower levels of homodimerization than the parent protein.

[0350] In addition, the 5-8 protein was modified in its dimerization interface by replacing the glutamine (Q) residue at position 486 with glutamic acid (E). This 5-8 (Q486E) mutant was tested for its ability to catalyze targeted cleavage in the presence of either the wild-type 5-10 protein or the 5-10 (E490K) mutant. DNA cleavage was not observed when the labeled substrate was incubated in the presence of both 5-8 (Q486E) and wild-type 5-10 (Table 10, Row 5). However, cleavage was obtained when the 5-8 (Q486E) and 5-10 (E490K) mutants were used in combination (Table 10, Row 6).

[0351] These results indicate that DNA cleavage by a ZFP/FokI fusion protein pair, at regions other than that defined by the target sequences of the two fusion proteins, can be minimized or abolished by altering the amino acid sequence of the cleavage half-domain in one or both of the fusion proteins.

TABLE 10

DNA cleavage by ZFP/FokI fusion protein pairs containing wild-type and mutant cleavage half-domains			
	ZFP 5-8 binding domain	ZFP 5-10 binding domain	DNA cleavage
1	Wild-type FokI	Wild-type FokI	Specific
2	Not present	Wild-type FokI	Non-specific
3	Not present	FokI E490K	None
4	Wild-type FokI	FokI E490K	Specific

TABLE 10-continued

DNA cleavage by ZFP/FokI fusion protein pairs containing wild-type and mutant cleavage half-domains			
	ZFP 5-8 binding domain	ZFP 5-10 binding domain	DNA cleavage
5	FokI Q486E	Wild-type FokI	None
6	FokI Q486E	FokI E490K	Specific

Note:

Each row of the table presents results of a separate experiment in which ZFP/FokI fusion proteins were tested for cleavage of a labeled DNA substrate. One of the fusion proteins contained the 5-8 DNA binding domain, and the other fusion protein contained the 5-10 DNA binding domain (See Table 9 and FIG. 19). The cleavage half-domain portion of the fusion proteins was as indicated in the Table. Thus, the entries in the ZFP 5-8 column indicate the type of FokI cleavage domain fused to ZFP 5-8; and the entries in the ZFP 5-10 column indicates the type of FokI cleavage domain fused to ZFP 5-10. For the FokI cleavage half-domain mutants, the number refers to the amino acid residue in the FokI protein; the letter preceding the number refers to the amino acid present in the wild-type protein and the letter following the number denotes the amino acid to which the wild-type residue was changed in generating the modified protein. 'Not present' indicates that the entire ZFP/FokI fusion protein was omitted from that particular experiment. The DNA substrate used in this experiment was an approximately 400 bp PCR product containing the target sites for both ZFP 5-8 and ZFP 5-10. See FIG. 19 for the sequences and relative orientation of the two target sites.

Example 6

Generation of a Defective Enhanced Green Fluorescent Protein (eGFP) Gene

[0352] The enhanced Green Fluorescent Protein (eGFP) is a modified form of the Green Fluorescent Protein (GFP; see, e.g., Tsien (1998) *Ann. Rev. Biochem.* 67:509-544) containing changes at amino acid 64 (phe to leu) and 65 (ser to thr). Heim et al. (1995) *Nature* 373:663-664; Cormack et al. (1996) *Gene* 173:33-38. An eGFP-based reporter system was constructed by generating a defective form of the eGFP gene, which contained a stop codon and a 2-bp frameshift mutation. The sequence of the eGFP gene is shown in FIG. 22. The mutations were inserted by overlapping PCR mutagenesis, using the Platinum® Taq DNA Polymerase High Fidelity kit (Invitrogen) and the oligonucleotides GFP-Bam, GFP-Xba, stop sense2, and stop anti2 as primers (oligonucleotide sequences are listed below in Table 11). GFP-Bam and GFP-Xba served as the external primers, while the primers stop sense2 and stop anti2 served as the internal primers encoding the nucleotide changes. The peGFP-N1 vector (BD Biosciences), encoding a full-length eGFP gene, was used as the DNA template in two separate amplification reactions, the first utilizing the GFP-Bam and stop anti2 oligonucleotides as primers and the second using the GFP-Xba and stop sense2 oligonucleotides as primers. This generated two amplification products whose sequences overlapped. These products were combined and used as template in a third amplification reaction, using the external GFP-Bam and GFP-Xba oligonucleotides as primers, to regenerate a modified eGFP gene in which the sequence GACCACAT (SEQ ID NO: 124) at nucleotides 280-287 was replaced with the sequence TAACAC (SEQ ID NO: 125). The PCR conditions for all amplification reactions were as follows: the template was initially denatured for 2 minutes at 94 degrees and followed by 25 cycles of amplification by incubating the reaction for 30 sec. at 94 degrees C., 45 sec.

at 46 degrees C., and 60 sec. at 68 degrees C. A final round of extension was carried out at 68 degrees C. for 10 minutes. The sequence of the final amplification product is shown in FIG. 23. This 795 bp fragment was cloned into the pCR(R)4-TOPO vector using the TOPO-TA cloning kit (Invitrogen) to generate the pCR(R)4-TOPO-GFPmut construct.

TABLE 11

<u>Oligonucleotide sequences for GFP</u>	
Oligo	sequence 5'–3'
GFP-Bam	CGAATTCTGCGAGTCGAC (SEQ ID NO:126)
GFP-Xha	GATTATGATCTAGAGTCG (SEQ ID NO:127)
stop sense 2	AGCCGCTACCCCTAACACGAAGCAG (SEQ ID NO:128)
stop anti 2	CTGCTTCGTGTTAGGGGTAGCGGCT (SEQ ID NO:129)

Example 7

Design and Assembly of Zinc Finger Nucleases
Targeting eGFP

[0353] Two three-finger ZFPs were designed to bind a region of the mutated GFP gene (Example 6) corresponding to nucleotides 271-294 (numbering according to FIG. 23). The binding sites for these proteins occur in opposite orientation with 6 base pairs separating the two binding sites. See FIG. 23. ZFP 287A binds nucleotides 271-279 on the non-coding strand, while ZFP 296 binds nucleotides 286-294 on the coding strand. The DNA target and amino acid sequence for the recognition regions of the ZFPs are listed below, and in Table 12:

TABLE 12

<u>Zinc finger designs for the GFP gene</u>				
Protein	Target sequence	F1	F2	F3
287A	GGGGTAGCGg (SEQ ID NO:136)	RSDDLTR (SEQ ID NO:137)	QSGALAR (SEQ ID NO:138)	RSDHLNR (SEQ ID NO:139)
296S	GAAGCAGCA (SEQ ID NO:140)	QSGSLTR (SEQ ID NO:141)	QSGDLTR (SEQ ID NO:142)	QSGNLAR (SEQ ID NO:143)

Note:

The zinc finger amino acid sequences shown above (in one-letter code) represent residues -1 through +6, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein, and Finger F3 is closest to the carboxy terminus.

287A:

F1 (GCGg) RSDDLTR (SEQ ID NO:130)

F2 (GTA) QSGALAR (SEQ ID NO:131)

F3 (GGG) RSDHLNR (SEQ ID NO:132)

296S:

F1 (GCA) QSGSLTR (SEQ ID NO:133)

F2 (GCA) QSGDLTR (SEQ ID NO:134)

F3 (GAA) QSGNLAR (SEQ ID NO:135)

[0354] Sequences encoding these proteins were generated by PCR assembly (e.g., U.S. Pat. No. 6,534,261), cloned between the KpnI and BamHI sites of the pcDNA3.1 vector

(Invitrogen), and fused in frame with the catalytic domain of the FokI endonuclease (amino acids 384-579 of the sequence of Looney et al. (1989) *Gene* 80:193-208). The resulting constructs were named pcDNA3.1-GFP287-FokI and pcDNA3.1-GFP296-FokI (FIG. 24).

Example 8

Targeted in Vitro DNA Cleavage by Designed Zinc
Finger Nucleases

[0355] The pCR(R)4-TOPO-GFPmut construct (Example 6) was used to provide a template for testing the ability of the 287 and 296 zinc finger proteins to specifically recognize their target sites and cleave this modified form of eGFP in vitro.

[0356] A DNA fragment containing the defective eGFP-encoding insert was obtained by PCR amplification, using the T7 and T3 universal primers and pCR(R)4-TOPO-GFPmut as template. This fragment was end-labeled using γ -³²P-ATP and T4 polynucleotide kinase. Unincorporated nucleotide was removed using a microspin G-50 column (Amersham).

[0357] An in vitro coupled transcription/translation system was used to express the 287 and 296 zinc finger nucleases described in Example 7. For each construct, 200 ng linearized plasmid DNA was incubated in 20 μ L TnT mix and incubated at 30° C. for 1 hour and 45 minutes. TnT mix contains 100 μ L TnT lysate (which includes T7 RNA polymerase, Promega, Madison, Wis.) supplemented with 2 μ L Methionine (1 mM) and 2.5 μ L ZnCl₂ (20 mM).

[0358] For analysis of DNA cleavage, aliquots from each of the 287 and 296 coupled transcription/translation reaction mixtures were combined, then serially diluted with cleavage buffer. Cleavage buffer contains 20 mM Tris-HCl pH 8.5, 75 mM NaCl, 10 mM MgCl₂, 10 μ M ZnCl₂, 1 mM DTT, 5%

glycerol, 500 μ g/ml BSA. 5 μ L of each dilution was combined with approximately 1 ng DNA substrate (end-labeled with ³²P using T4 polynucleotide kinase as described

above), and each mixture was further diluted to generate a 20 μ l cleavage reaction having the following composition: 20 mM Tris-HCl pH 8.5, 75 mM NaCl, 10 mM MgCl₂, 10 μ M ZnCl₂, 1 mM DTT, 5% glycerol, 500 μ g/ml BSA. Cleavage reactions were incubated for 1 hour at 37° C. Protein was extracted by adding 10 μ l phenol-chloroform solution to each reaction, mixing, and centrifuging to separate the phases. Ten microliters of the aqueous phase from each reaction was analyzed by electrophoresis on a 10% polyacrylamide gel.

[0359] The gel was subjected to autoradiography, and the results of this experiment are shown in FIG. 25. The four left-most lanes show the results of reactions in which the final dilution of each coupled transcription/translation reaction mixture (in the cleavage reaction) was 1/156.25, 1/31.25, 1/12.5 and 1/5, respectively, resulting in effective volumes of 0.032, 0.16, 0.4, and 1 μ l, respectively of each coupled transcription/translation reaction. The appearance of two DNA fragments having lower molecular weights than the starting fragment (lane labeled "uncut control" in FIG. 25) is correlated with increasing amounts of the 287 and 296 zinc finger endonucleases in the reaction mixture, showing that DNA cleavage at the expected target site was obtained.

Example 9

Generation of Stable Cell Lines Containing an Integrated Defective eGFP Gene

[0360] A DNA fragment encoding the mutated eGFP, eGFPmut, was cleaved out of the pCR(R)4-TOPO-GFPmut vector (Example 6) and cloned into the HindIII and NotI sites of pcDNA4/TO, thereby placing this gene under control of a tetracycline-inducible CMV promoter. The resulting plasmid was named pcDNA4/TO/GFPmut (FIG. 26). T-Rex 293 cells (Invitrogen) were grown in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen) supplemented with 10% Tet-free fetal bovine serum (FBS) (HyClone). Cells were plated into a 6-well dish at 50% confluence, and two wells were each transfected with pcDNA4/TO/GFPmut. The cells were allowed to recover for 48 hours, then cells from both wells were combined and split into 10 $\times$ 15-cm² dishes in selective medium, i.e., medium supplemented with 400 μ g/ml Zeocin (Invitrogen). The medium was changed every 3 days, and after 10 days single colonies were isolated and expanded further. Each clonal line was tested individually for doxycycline(dox)-inducible expression of the eGFPmut gene by quantitative RT-PCR (TaqMan®).

[0361] For quantitative RT-PCR analysis, total RNA was isolated from dox-treated and untreated cells using the High Pure Isolation Kit (Roche Molecular Biochemicals), and 25 ng of total RNA from each sample was subjected to real time quantitative RT-PCR to analyze endogenous gene expression, using TaqMan® assays. Probe and primer sequences are shown in Table 13. Reactions were carried out on an ABI 7700 SDS machine (PerkinElmer Life Sciences) under the following conditions. The reverse transcription reaction was performed at 48° C. for 30 minutes with MultiScribe reverse transcriptase (PerkinElmer Life Sciences), followed by a 10-minute denaturation step at 95° C. Polymerase chain reaction (PCR) was carried out with AmpliGold DNA polymerase (PerkinElmer Life Sciences) for 40 cycles at 95° C. for 15 seconds and 60° C. for 1 minute. Results were analyzed using the SDS version 1.7 software and are shown

in FIG. 27, with expression of the eGFPmut gene normalized to the expression of the human GAPDH gene. A number of cell lines exhibited doxycycline-dependent expression of eGFP; line 18 (T18) was chosen as a model cell line for further studies.

TABLE 13

Oligonucleotides for mRNA analysis	
Oligonucleotide	Sequence
eGFP primer 1 (5T)	CTGCTGCCCCACAACCA (SEQ ID NO:144)
eGFP primer 2 (3T)	CCATGTGATCGCGCTTCTC (SEQ ID NO:145)
eGFP probe	CCCAGTCCGCCCTGAGCAAAGA (SEQ ID NO:146)
GAPDH primer 1	CCATGTTTCGTCATGGGTGTA (SEQ ID NO:147)
GAPDH primer 2	CATGGACTGTGGTCATGAGT (SEQ ID NO:148)
GAPDH probe	TCCTGCACCACCAACTGCTTAGCA (SEQ ID NO:149)

Example 10

Generation of a Donor Sequence for Correction of a Defective Chromosomal eGFP Gene

[0362] A donor construct containing the genetic information for correcting the defective eGFPmut gene was constructed by PCR. The PCR reaction was carried out as described above, using the peGFP-NI vector as the template. To prevent background expression of the donor construct in targeted recombination experiments, the first 12 bp and start codon were removed from the donor by PCR using the primers GFPnostart and GFP-Xba (sequences provided in Table 14). The resulting PCR fragment (734 bp) was cloned into the pCR(R)4-TOPO vector, which does not contain a mammalian cell promoter, by TOPO-TA cloning to create pCR(R)4-TOPO-GFPdonor5 (FIG. 28). The sequence of the eGFP insert of this construct (corresponding to nucleotides 64-797 of the sequence shown in FIG. 22) is shown in FIG. 29 (SEQ ID NO:20).

TABLE 14

Oligonucleotides for construction of donor molecule	
Oligonucleotide	Sequence 5'-3'
GFPnostart	GGCGAGGAGCTGTTTAC (SEQ ID NO:150)
GFP-Xba	GATTATGATCTAGAGTCG (SEQ ID NO:151)

Example 11

Correction of a Mutation in an Integrated Chromosomal eGFP Gene by Targeted Cleavage and Recombination

[0363] The T18 stable cell line (Example 9) was transfected with one or both of the ZFP-FokI expression plasmid

(pcDNA3.1-GFP287-FokI and pcDNA3.1-GFP296-FokI, Example 7) and 300 ng of the donor plasmid pCR(R)4-TOPO-GFPdonor5 (Example 10) using LipofectAMINE 2000 Reagent (Invitrogen) in Opti-MEM I reduced serum medium, according to the manufacturer's protocol. Expression of the defective chromosomal eGFP gene was induced 5-6 hours after transfection by the addition of 2 ng/ml doxycycline to the culture medium. The cells were arrested in the G2 phase of the cell cycle by the addition, at 24 hours post-transfection, of 100 ng/ml Nocodazole (FIG. 30) or 0.2 μ M Vinblastine (FIG. 31). G2 arrest was allowed to continue for 24-48 hours, and was then released by the removal of the medium. The cells were washed with PBS and the medium was replaced with DMEM containing tetracycline-free FBS and 2 ng/ml doxycycline. The cells were allowed to recover for 24-48 hours, and gene correction efficiency was measured by monitoring the number of cells exhibiting eGFP fluorescence, by fluorescence-activated cell sorting (FACS) analysis. FACS analysis was carried out using a Beckman-Coulter EPICS XL-MCL instrument and System II Data Acquisition and Display software, version 2.0. eGFP fluorescence was detected by excitation at 488 nm with an argon laser and monitoring emissions at 525 nm (x-axis). Background or autofluorescence was measured by monitoring emissions at 570 nm (y-axis). Cells exhibiting high fluorescent emission at 525 nm and low emission at 570 nm (region E) were scored positive for gene correction.

[0364] The results are summarized in Table 15 and FIGS. 30 and 31. FIGS. 30 and 31 show results in which T18 cells were transfected with the pcDNA3.1-GFP287-FokI and pcDNA3.1-GFP296-FokI plasmids encoding ZFP nucleases and the pCR(R)4-TOPO-GFPdonor5 plasmid, eGFP expression was induced with doxycycline, and cells were arrested in G2 with either nocodazole (FIG. 30) or vinblastine (FIG. 31). Both figures show FACS traces, in which cells exhibiting eGFP fluorescence are represented in the lower right-hand portion of the trace (identified as Region E, which is the portion of Quadrant 4 underneath the curve). For transfected cells that had been treated with nocodazole, 5.35% of the cells exhibited GFP fluorescence, indicative of correction of the mutant chromosomal eGFP gene (FIG. 30), while 6.7% of cells treated with vinblastine underwent eGFP gene correction (FIG. 31). These results are summarized, along with additional control experiments, in Rows 1-8 of Table 15.

[0365] In summary, these experiments show that, in the presence of two ZFP nucleases and a donor sequence, approximately 1% of treated cells underwent gene correction, and that this level of correction was increased 4-5 fold by arresting treated cells in the G2 phase of the cell cycle.

TABLE 15

Correction of a defective chromosomal eGFP gene		
Expt.	Treatment ¹	Percent cells with corrected eGFP gene ²
1	300 ng donor only	0.01
2	100 ng ZFP 287 + 300 ng donor	0.16
3	100 ng ZFP 296 + 300 ng donor	0.6
4	50 ng ZFP 287 + 50 ng ZFP 296 + 300 ng donor	1.2
5	as 4 + 100 ng/ml nocodazole	5.35

TABLE 15-continued

Correction of a defective chromosomal eGFP gene		
Expt.	Treatment ¹	Percent cells with corrected eGFP gene ²
6	as 4 + 0.2 μ M vinblastine	6.7
7	no donor, no ZFP, 100 ng/ml nocodazole	0.01
8	no donor, no ZFP, 0.2 μ M vinblastine	0.0
9	100 ng ZFP287/Q486E + 300 ng donor	0.0
10	100 ng ZFP296/E490K + 300 ng donor	0.01
11	50 ng 287/Q486E + 50 ng 296/E490K + 300 ng donor	0.62
12	as 11 + 100 ng/ml nocodazole	2.37
13	as 11 + 0.2 μ M vinblastine	2.56

Notes:

¹T18 cells, containing a defective chromosomal eGFP gene, were transfected with plasmids encoding one or two ZFP nucleases and/or a donor plasmid encoding a nondefective eGFP sequence, and expression of the chromosomal eGFP gene was induced with doxycycline. Cells were optionally arrested in G2 phase of the cell cycle after eGFP induction. FACS analysis was conducted 5 days after transfection.

²The number is the percent of total fluorescence exhibiting high emission at 525 nm and low emission at 570 nm (region E of the FACS trace).

Example 12

Correction of a Defective Chromosomal Gene Using Zinc Finger Nucleases with Sequence Alterations in the Dimerization Interface

[0366] Zinc finger nucleases whose sequences had been altered in the dimerization interface were tested for their ability to catalyze correction of a defective chromosomal eGFP gene. The protocol described in Example 11 was used, except that the nuclease portion of the ZFP nucleases (i.e., the FokI cleavage half-domains) were altered as described in Example 5. Thus, an E490K cleavage half-domain was fused to the GFP96 ZFP domain (Table 12), and a Q486E cleavage half-domain was fused to the GFP287 ZFP (Table 12).

[0367] The results are shown in Rows 9-11 of Table 15 and indicate that a significant increase in the frequency of gene correction was obtained in the presence of two ZFP nucleases having alterations in their dimerization interfaces, compared to that obtained in the presence of either of the nucleases alone. Additional experiments, in which T18 cells were transfected with donor plasmid and plasmids encoding the 287/Q486E and 296/E490K zinc finger nucleases, then arrested in G2 with nocodazole or vinblastine, showed a further increase in frequency of gene correction, with over 2% of cells exhibiting eGFP fluorescence, indicative of a corrected chromosomal eGFP gene (Table 15, Rows 12 and 13).

Example 13

Effect of Donor Length on Frequency of Gene Correction

[0368] In an experiment similar to those described in Example 11, the effect of the length of donor sequence on frequency of targeted recombination was tested. T18 cells were transfected with the two ZFP nucleases, and eGFP expression was induced with doxycycline, as in Example 11. Cells were also transfected with either the pCR(R)4-TOPO-

GFPdonor5 plasmid (FIG. 28) containing a 734 bp eGFP insert (FIG. 29) as in Example 11, or a similar plasmid containing a 1527 bp sequence insert (FIG. 32) homologous to the mutated chromosomal eGFP gene. Additionally, the effect of G2 arrest with nocodazole on recombination frequency was assessed.

[0369] In a second experiment, donor lengths of 0.7, 1.08 and 1.5 kbp were compared. T18 cells were transfected with 50 ng of the 287-FokI and 296-FokI expression plasmids (Example 7, Table 12) and 500 ng of a 0.7 kbp, 1.08 kbp, or 1.5 kbp donors, as described in Example 11. Four days after transfection, cells were assayed for correction of the defective eGFP gene by FACS, monitoring GFP fluorescence.

[0370] The results of these two experiments, shown in Table 16, show that longer donor sequence increases the frequency of targeted recombination (and, hence, of gene correction) and confirm that arrest of cells in the G2 phase of the cell cycle also increases the frequency of targeted recombination.

TABLE 16

Effect of donor length and cell-cycle arrest on targeted recombination frequency			
Experiment 1		Experiment 2	
Nocodazole concentration:			
Donor length (kb)	0 ng/ml	100 ng/ml	—
0.7	1.41	5.84	1.2
1.08	not done	not done	2.2
1.5	2.16	8.38	2.3

Note:
Numbers represent percentage of total fluorescence in Region E of the FACS trace (see Example 11) which is an indication of the fraction of cells that have undergone targeted recombination to correct the defective chromosomal eGFP gene.

Example 14

Editing of the Endogenous Human IL-2R γ Gene by Targeted Cleavage and Recombination Using Zinc Finger Nucleases

[0371] Two expression vectors, each encoding a ZFP-nuclease targeted to the human IL-2R γ gene, were constructed. Each ZFP-nuclease contained a zinc finger protein-based DNA binding domain (see Table 17) fused to the nuclease domain of the type IIS restriction enzyme FokI (amino acids 384-579 of the sequence of Looney et al. (1989) *Gene* 80:193-208) via a four amino acid ZC linker (see Example 4). The nucleases were designed to bind to positions in exon 5 of the chromosomal IL-2R γ gene surrounding codons 228 and 229 (a mutational hotspot in the gene) and to introduce a double-strand break in the DNA between their binding sites.

TABLE 17

Zinc Finger Designs for exon 5 of the IL2R γ Gene				
Target sequence	F1	F2	F3	F4
ACTCTGTGGAAG (SEQ ID NO:152) 5-8G	RSDNLSV (SEQ ID NO:153)	RNAHIRN (SEQ ID NO:154)	RSDTLSE (SEQ ID NO:155)	ARSTRTN (SEQ ID NO:156)
AAAGCGGCTCCG (SEQ ID NO:157) 5-9D	RSDTLSE (SEQ ID NO:158)	ARSTRTT (SEQ ID NO:159)	RSDSLSK (SEQ ID NO:160)	QRSNLKV (SEQ ID NO:161)

Note:
The zinc finger amino acid sequences shown above (in one-letter code) represent residues -1 through +6, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein.

[0372] The complete DNA-binding portion of each of the chimeric endonucleases was as follows:

(SEQ ID NO:152)
Nuclease targeted to ACTCTGTGGAAG
(SEQ ID NO:162)
MAERPFQCRICMRINFSRSDNLSVHIRTHTGKPFACDICGRKFAARNAHR
INHTKIHTGSQKPFQCRICMRNFSRSDTLSEHIRTHTGKPFACDICGRK
FAARSTRTNHTKIHLRGS
(SEQ ID NO:157)
Nuclease targeted to AAAGCGGCTCCG
(SEQ ID NO:163)
MAERPFQCRICMRNFSRSDTLSEHIRTHTGKPFACDICGRKFAARSTRT
THTKIHTGSQKPFQCRICMRNFSRSDSLSKHIRTHTGKPFACDICGRKF
AQRSLNKVHTKIHLRGS

[0373] Human embryonic kidney 293 cells were transfected (Lipofectamine 2000; Invitrogen) with two expression constructs, each encoding one of the ZFP-nucleases described in the preceding paragraph. The cells were also transfected with a donor construct carrying as an insert a 1,543 bp fragment of the IL2R γ locus corresponding to positions 69195166-69196708 of the “minus” strand of the

X chromosome (UCSC human genome release July 2003), in the pCR4Blunt Topo (Invitrogen) vector. The IL-2R γ insert sequence contained the following two point mutations in the sequence of exon 5 (underlined):

F R V R S R F N P L C G S (SEQ ID NO:164)
 TTTCGTGTTTCGGAGCCGTTTAACCCGCTCTGTGGAAGT (SEQ ID NO:165)

[0374] The first mutation (CGC→CGG) does not change the amino acid sequence (upper line) and serves to adversely affect the ability of the ZFP-nuclease to bind to the donor DNA, and to chromosomal DNA following recombination. The second mutation (CCA→CCG) does not change the amino acid sequence and creates a recognition site for the restriction enzyme BsrBI.

[0375] Either 50 or 100 nanograms of each ZFP-nuclease expression construct and 0.5 or 1 microgram of the donor construct were used in duplicate transfections. The following control experiments were also performed: transfection with an expression plasmid encoding the eGFP protein; transfection with donor construct only; and transfection with plasmids expressing the ZFP nucleases only. Twenty four hours after transfection, vinblastine (Sigma) was added to 0.2 μ M final concentration to one sample in each set of duplicates, while the other remained untreated. Vinblastine affects the cell's ability to assemble the mitotic spindle and therefore acts as a potent G₂ arresting agent. This treatment was performed to enhance the frequency of targeting because the homology-directed double-stranded break repair pathway is more active than non-homologous end-joining in the G₂ phase of the cell cycle. Following a 48 hr period of treatment with 0.2 μ M vinblastine, growth medium was replaced, and the cells were allowed to recover from vinblastine treatment for an additional 24 hours. Genomic DNA was then isolated from all cell samples using the DNEasy Tissue Kit (Qiagen). Five hundred nanograms of genomic DNA from each sample was then assayed for frequency of gene targeting, by testing for the presence of a new BsrBI site in the chromosomal IL-2R γ locus, using the assay described schematically in FIG. 33.

[0376] In brief, 20 cycles of PCR were performed using the primers shown in Table 18, each of which hybridizes to the chromosomal IL-2R γ locus immediately outside of the region homologous to the 1.5 kb donor sequence. Twenty microcuries each of α -³²P-dCTP and α -³²P-dATP were included in each PCR reaction to allow detection of PCR products. The PCR reactions were desalted on a G-50 column (Amersham), and digested for 1 hour with 10 units of BsrBI (New England Biolabs). The digestion products were resolved on a 10% non-denaturing polyacrylamide gel (BioRad), and the gel was dried and autoradiographed (FIG. 34). In addition to the major PCR product, corresponding to the 1.55 kb amplified fragment of the IL2R γ locus ("wt" in FIG. 34), an additional band ("rflp" in FIG. 34) was observed in lanes corresponding to samples from cells that were transfected with the donor DNA construct and both ZFP-nuclease constructs. This additional band did not appear in any of the control lanes, indicating that ZFP nuclease-facilitated recombination of the BsrBI RFLP-containing donor sequence into the chromosome occurred in this experiment.

[0377] Additional experiments, in which trace amounts of a RFLP-containing IL-2R γ DNA sequence was added to human genomic DNA (containing the wild-type IL-2R γ gene), and the resultant mixture was amplified and subjected

to digestion with a restriction enzyme which cleaves at the RFLP, have indicated that as little as 0.5% RFLP-containing sequence can be detected quantitatively using this assay.

TABLE 18

Oligonucleotides for analysis of the human IL-2R γ gene	
Oligonucleotide	Sequence
Ex5_1.5detF1	GATTC AACCAGACAGATAGAAGG (SEQ ID NO:166)
Ex5_1.5detR1	TTACTGTCTCATCCTTTACTCC (SEQ ID NO:167)

Example 15

Targeted Recombination at the IL-2R γ Locus in K562 Cells

[0378] K562 is a cell line derived from a human chronic myelogenous leukemia. The proteins used for targeted cleavage were FokI fusions to the 5-8G and 5-9D zinc finger DNA-binding domains (Example 14, Table 17). The donor sequence was the 1.5 kbp fragment of the human IL-2R γ gene containing a BsrBI site introduced by mutation, described in Example 14.

[0379] K562 cells were cultured in RPMI Medium 1640 (Invitrogen), supplemented with 10% fetal bovine serum (FBS) (Hyclone) and 2 mM L-glutamine. All cells were maintained at 37° C. in an atmosphere of 5% CO₂. These cells were transfected by Nucleofection™ (Solution V, Program T16) (Amaxa Biosystems), according to the manufacturers' protocol, transfecting 2 million cells per sample. DNAs for transfection, used in various combinations as described below, were a plasmid encoding the 5-8G ZFP-FokI fusion endonuclease, a plasmid encoding the 5-9D ZFP-FokI fusion endonuclease, a plasmid containing the donor sequence (described above and in Example 14) and the pGFP-N1 vector (BD Biosciences) used as a control.

[0380] In the first experiment, cells were transfected with various plasmids or combinations of plasmids as shown in Table 19.

TABLE 19

Sample #	p-eGFP-N1	p5-8G	p5-9D	donor	vinblastine
1	5 μ g	—	—	—	—
2	—	—	—	50 μ g	—
3	—	—	—	50 μ g	yes
4	—	10 μ g	10 μ g	—	—
5	—	5 μ g	5 μ g	25 μ g	—

TABLE 19-continued

Sample #	p-eGFP-N1	p5-8G	p5-9D	donor	vinblastine
6	—	5 µg	5 µg	25 µg	yes
7	—	7.5 µg	7.5 µg	25 µg	—
8	—	7.5 µg	7.5 µg	25 µg	yes
9	—	7.5 µg	7.5 µg	50 µg	—
10	—	7.5 µg	7.5 µg	50 µg	yes

[0381] Vinblastine-treated cells were exposed to 0.2 µM vinblastine at 24 hours after transfection for 30 hours. The cells were collected, washed twice with PBS, and re-growth plated in growth medium. Cells were harvested 4 days after transfection for analysis of genomic DNA.

[0382] Genomic DNA was extracted from the cells using the DNEasy kit (Qiagen). One hundred nanograms of genomic DNA from each sample were used in a PCR reaction with the following primers:

(SEQ ID NO:168)
Exon 5 forward: GCTAAGGCCAAGAAAGTAGGGCTAAAG

(SEQ ID NO:169)
Exon 5 reverse: TTCCTTCCATCACCAACCTCTTG

[0383] These primers amplify a 1,669 bp fragment of the X chromosome corresponding to positions 69195100-69196768 on the “—” strand (UCSC human genome release July 2003) that contain exon 5 of the IL2Rγ gene. Amplification of genomic DNA which has undergone homologous recombination with the donor DNA yields a product containing a BsrBI site; whereas the amplification product of genomic DNA which has not undergone homologous recombination with donor DNA will not contain this restriction site.

[0384] Ten microcuries each of a-³²PdCTP and a-³²PdATP were included in each amplification reaction to allow visualization of reaction products. Following 20 cycles of PCR, the reaction was desalted on a Sephadex G-50 column (Pharmacia), and digested with 10 Units of BsrBI (New England Biolabs) for 1 hour at 37° C. The reaction was then resolved on a 10% non-denaturing PAGE, dried, and exposed to a PhosphorImager screen.

[0385] The results of this experiment are shown in FIG. 35. When cells were transfected with the control GFP plasmid, donor plasmid alone or the two ZFP-encoding plasmids in the absence of donor, no BsrBI site was present in the amplification product, as indicated by the absence of the band marked “rflp” in the lanes corresponding to these samples in FIG. 35. However, genomic DNA of cells that were transfected with the donor plasmid and both ZFP-encoding plasmids contained the BsrBI site introduced by homologous recombination with the donor DNA (band labeled “rflp”). Quantitation of the percentage of signal represented by the RFLP-containing DNA, shown in FIG. 35, indicated that, under optimal conditions, up to 18% of all IL-2Rγ genes in the transfected cell population were altered by homologous recombination.

[0386] A second experiment was conducted according to the protocol just described, except that the cells were expanded for 10 days after transfection. DNAs used for transfection are shown in Table 20.

TABLE 20

Sample #	p-eGFP-N1	p5-8G	p5-8D	donor	vinblastine
1	50 µg	—	—	—	—
2	—	—	—	50 µg	—
3	—	—	—	50 µg	yes
4	—	7.5 µg	7.5 µg	—	—
5	—	5 µg	5 µg	25 µg	—
6	—	5 µg	5 µg	25 µg	yes
7	—	7.5 µg	7.5 µg	50 µg	—
8	—	7.5 µg	7.5 µg	50 µg	yes

[0387] Analysis of BsrBI digestion of amplified DNA, shown in FIG. 36, again demonstrated that up to 18% of IL-2Rγ genes had undergone sequence alteration through homologous recombination, after multiple rounds of cell division. Thus, the targeted recombination events are stable.

[0388] In addition, DNA from transfected cells in this second experiment was analyzed by Southern blotting. For this analysis, twelve micrograms of genomic DNA from each sample were digested with 100 units EcoRI, 50 units BsrBI, and 40 units of DpnI (all from New England Biolabs) for 12 hours at 37° C. This digestion generates a 7.7 kbp EcoRI fragment from the native IL-2Rγ gene (lacking a BsrBI site) and fragments of 6.7 and 1.0 kbp from a chromosomal IL-2Rγ gene whose sequence has been altered, by homologous recombination, to include the BsrBI site. DpnI, a methylation-dependent restriction enzyme, was included to destroy the dam-methylated donor DNA. Unmethylated K562 cell genomic DNA is resistant to DpnI digestion.

[0389] Following digestion, genomic DNA was purified by phenol-chloroform extraction and ethanol precipitation, resuspended in TE buffer, and resolved on a 0.8% agarose gel along with a sample of genomic DNA digested with EcoRI and SphI to generate a size marker. The gel was processed for alkaline transfer following standard procedure and DNA was transferred to a nylon membrane (Schleicher and Schuell). Hybridization to the blot was then performed by using a radiolabelled fragment of the IL-2Rγ locus corresponding to positions 69198428-69198769 of the “—” strand of the X chromosome (UCSC human genome July 2003 release). This region of the gene is outside of the region homologous to donor DNA. After hybridization, the membrane was exposed to a PhosphorImager plate and the data quantitated using Molecular Dynamics software. Alteration of the chromosomal IL-2Rγ sequence was measured by analyzing the intensity of the band corresponding to the EcoRI-BsrBI fragment (arrow next to autoradiograph; BsrBI site indicated by filled triangle in the map above the autoradiograph).

[0390] The results, shown in FIG. 37, indicate up to 15% of chromosomal IL-2Rγ sequences were altered by homologous recombination, thereby confirming the results obtained by PCR analysis that the targeted recombination event was stable through multiple rounds of cell division. The Southern blot results also indicate that the results shown in FIG. 36 do not result from an amplification artifact.

Example 16

Targeted Recombination at the IL-2R γ Locus in
CD34-positive Hematopoietic Stem Cells

[0391] Genetic diseases (e.g., severe combined immune deficiency (SCID) and sickle cell anemia) can be treated by homologous recombination-mediated correction of the specific DNA sequence alteration responsible for the disease. In certain cases, maximal efficiency and stability of treatment would result from correction of the genetic defect in a pluripotent cell. To this end, this example demonstrates alteration of the sequence of the IL-2R γ gene in human CD34-positive bone marrow cells. CD34⁺ cells are pluripotent hematopoietic stem cells which give rise to the erythroid, myeloid and lymphoid lineages.

[0392] Bone marrow-derived human CD34 cells were purchased from AllCells, LLC and shipped as frozen stocks. These cells were thawed and allowed to stand for 2 hours at 37° C. in an atmosphere of 5% CO₂ in RPMI Medium 1640 (Invitrogen), supplemented with 10% fetal bovine serum (FBS) (Hyclone) and 2 mM L-glutamine. Cell samples (1×10⁶ or 2×10⁶ cells) were transfected by Nucleofection™ (amaxa biosystems) using the Human CD34 Cell Nucleofector™ Kit, according to the manufacturers' protocol. After transfection, cells were cultured in RPMI Medium 1640 (Invitrogen), supplemented with 10% FBS, 2 mM L-glutamine, 100 ng/ml granulocyte-colony stimulating factor (G-CSF), 100 ng/ml stem cell factor (SCF), 100 ng/ml thrombopoietin (TPO), 50 ng/ml Flt3 Ligand, and 20 ng/ml Interleukin-6 (IL-6). The caspase inhibitor zVAD-FMK (Sigma-Aldrich) was added to a final concentration of 40 μ M in the growth medium immediately after transfection to block apoptosis. Additional caspase inhibitor was added 48 hours later to a final concentration of 20 μ M to further prevent apoptosis. These cells were maintained at 37° C. in an atmosphere of 5% CO₂ and were harvested 3 days post-transfection.

[0393] Cell numbers and DNAs used for transfection are shown in Table 21.

TABLE 21

Sample	# cells	p-eGFP-N1 ¹	Donor ²	p5-8G ³	p5-9D ³
1	1 × 10 ⁶	5 μ g	—	—	—
2	2 × 10 ⁶	—	50 μ g	—	—
3	2 × 10 ⁶	—	50 μ g	7.5 μ g	7.5 μ g

¹This is a control plasmid encoding an enhanced green fluorescent protein.

²The donor DNA is a 1.5 kbp fragment containing sequences from exon 5 of the IL-2R γ gene with an introduced BsrBI site (see Example 14).

³These are plasmids encoding FokI fusions with the 5-8G and 5-9D zinc finger DNA binding domains (see Table 17).

[0394] Genomic DNA was extracted from the cells using the MasterPure DNA Purification Kit (Epicentre). Due to the presence of glycogen in the precipitate, accurate quantitation of this DNA used as input in the PCR reaction is impossible; estimates using analysis of ethidium bromide-stained agarose gels indicate that ca. 50 ng genomic DNA was used in each sample. Thirty cycles of PCR were then performed

using the following primers, each of which hybridizes to the chromosomal IL-2R γ locus immediately outside of the region homologous to the 1.5 kb donor:

(SEQ ID NO:170)
ex5_1.5detF3 GCTAAGGCCAAGAAAGTAGGGCTAAAG

(SEQ ID NO:171)
ex5_1.5detR3 TTCCTTCCATCACCAAACCTCTTGT

[0395] Twenty microcuries each of α -³²PdCTP and α -³²PdATP were included in each PCR reaction to allow detection of PCR products. To provide an in-gel quantitation reference, the existence of a spontaneously occurring SNP in exon 5 of the IL-2R γ gene in Jurkat cells was exploited: this SNP creates a RFLP by destroying a MaeII site that is present in normal human DNA. A reference standard was therefore created by adding 1 or 10 nanograms of normal human genomic DNA (obtained from Clontech, Palo Alto, Calif.) to 100 or 90 ng of Jurkat genomic DNA, respectively, and performing the PCR as described above. The PCR reactions were desalted on a G-50 column (Amersham), and digested for 1 hour with restriction enzyme: experimental samples were digested with 10 units of BsrBI (New England Biolabs); the "reference standard" reactions were digested with MaeII. The digestion products were resolved on a 10% non-denaturing PAGE (BioRad), the gel dried and analyzed by exposure to a PhosphorImager plate (Molecular Dynamics).

[0396] The results are shown in FIG. 38. In addition to the major PCR product, corresponding to the 1.6 kb fragment of the IL2R γ locus ("wt" in the right-hand panel of FIG. 38), an additional band (labeled "rflp") was observed in lanes corresponding to samples from cells that were transfected with plasmids encoding both ZFP-nucleases and the donor DNA construct. This additional band did not appear in the control lanes, consistent with the idea that ZFP-nuclease assisted gene targeting of exon 5 of the common gamma chain gene occurred in this experiment.

[0397] Although accurate quantitation of the targeting rate is complicated by the proximity of the RFLP band to the wild-type band; the targeting frequency was estimated, by comparison to the reference standard (left panel), to be between 1-5%.

Example 17

Donor-target Homology Effects

[0398] The effect, on frequency of homologous recombination, of the degree of homology between donor DNA and the chromosomal sequence with which it recombines was examined in T18 cell line, described in Example 9. This line contains a chromosomally integrated defective eGFP gene, and the donor DNA contains sequence changes, with respect to the chromosomal gene, that correct the defect.

[0399] Accordingly, the donor sequence described in Example 10 was modified, by PCR mutagenesis, to generate a series of ~700 bp donor constructs with different degrees of non-homology to the target. All of the modified donors contained sequence changes that corrected the defect in the chromosomal eGFP gene and contained additional silent mutations (DNA mutations that do not change the sequence

of the encoded protein) inserted into the coding region surrounding the cleavage site. These silent mutations were intended to prevent the binding to, and cleavage of, the donor sequence by the zinc finger-cleavage domain fusions, thereby reducing competition between the intended chromosomal target and the donor plasmid for binding by the chimeric nucleases. In addition, following homologous recombination, the ability of the chimeric nucleases to bind and re-cleave the newly-inserted chromosomal sequences (and possibly stimulating another round of recombination, or causing non-homologous end joining or other double-strand break-driven alterations of the genome) would be minimized.

[0400] Four different donor sequences were tested. Donor 1 contains 8 mismatches with respect to the chromosomal defective eGFP target sequence, Donor 2 has 10 mismatches, Donor 3 has 6 mismatches, and Donor 5 has 4 mismatches. Note that the sequence of donor 5 is identical to wild-type eGFP sequence, but contains 4 mismatches with respect to the defective chromosomal eGFP sequence in the T18 cell line. Table 22 provides the sequence of each donor between nucleotides 201-242. Nucleotides that are divergent from the sequence of the defective eGFP gene integrated into the genome of the T18 cell line are shown in bold and underlined. The corresponding sequences of the defective chromosomal eGFP gene (GFP mut) and the normal eGFP gene (GFP wt) are also shown.

TABLE 22

Donor	Sequence	SEQ ID NO.
Donor 1	CTTCAGCCGCTATCC AGACCACATGAACA CACGA CTTCTT	172
Donor 2	CTTCAGCCGGTATCC AGACCACATGAACA CATGA CTTCTT	173
Donor 3	CTTCAGCCGCTACCC AGACCACATGAACA CAGCACGA CTTCTT	174
Donor 5	CTTCAGCCGCTACCC AGACCACATGAAG CAGCACGA CTTCTT	175
GFP mut	CTTCAGCCGCTACCCCTAACAC--GAAGCAGCACGA CTTCTT	176
GFP wt	CTTCAGCCGCTACCC AGACCACATGAAG CAGCACGA CTTCTT	177

[0401] The T18 cell line was transfected, as described in Example 11, with 50 ng of the 287-FokI and 296-FokI expression constructs (Example 7 and Table 12) and 500 ng of each donor construct. FACS analysis was conducted as described in Example 11.

[0402] The results, shown in Table 23, indicate that a decreasing degree of mismatch between donor and chromosomal target sequence (i.e., increased homology) results in an increased frequency of homologous recombination as assessed by restoration of GFP function.

TABLE 23¹

Donor	# mismatches	Percent cells with corrected eGFP gene ²
Donor 2	10	0.45%
Donor 1	8	0.53%
Donor 3	6	0.89%
Donor 5	4	1.56%

¹T18 cells, containing a defective chromosomal eGFP gene, were transfected with plasmids encoding two ZFP nucleases and with donor plasmids encoding a nondefective eGFP sequence having different numbers of sequence mismatches with the chromosomal target sequence. Expression of the chromosomal eGFP gene was induced with doxycycline and FACS analysis was conducted 5 days after transfection.

²The number is the percent of total fluorescence exhibiting high emission at 525 nm and low emission at 570 nm (region E of the FACS trace).

[0403] The foregoing results show that levels of homologous recombination are increased by decreasing the degree of target-donor sequence divergence. Without wishing to be bound by any particular theory or to propose a particular mechanism, it is noted that greater homology between donor and target could facilitate homologous recombination by increasing the efficiency by which the cellular homologous recombination machinery recognizes the donor molecule as a suitable template. Alternatively, an increase in donor homology to the target could also lead to cleavage of the donor by the chimeric ZFP nucleases. A cleaved donor could help facilitate homologous recombination by increasing the rate of strand invasion or could aid in the recognition of the cleaved donor end as a homologous stretch of DNA during homology search by the homologous recombination machinery. Moreover, these possibilities are not mutually exclusive.

Example 18

Preparation of siRNA

[0404] To test whether decreasing the cellular levels of proteins involved in non-homologous end joining (NHEJ) facilitates targeted homologous recombination, an experiment in which levels of the Ku70 protein were decreased through siRNA inhibition was conducted. siRNA molecules targeted to the Ku70 gene were generated by transcription of Ku70 cDNA followed by cleavage of double-stranded transcript with Dicer enzyme.

[0405] Briefly, a cDNA pool generated from 293 and U2OS cells was used in five separate amplification reactions, each using a different set of amplification primers specific to the Ku70 gene, to generate five pools of cDNA fragments (pools A-E), ranging in size from 500-750 bp. Fragments in each of these five pools were then re-amplified using primers containing the bacteriophage T7 RNA polymerase promoter element, again using a different set of primers for each cDNA pool. cDNA generation and PCR reactions were performed using the Superscript Choice cDNA system and Platinum Taq High Fidelity Polymerase (both from Invitrogen, Carlsbad, Calif.), according to manufacturers protocols and recommendations.

[0406] Each of the amplified DNA pools was then transcribed in vitro with bacteriophage T7 RNA polymerase to generate five pools (A-E) of double stranded RNA (dsRNA), using the RNAMAXX in vitro transcription kit (Stratagene,

San Diego, Calif.) according to the manufacturer's instructions. After precipitation with ethanol, the RNA in each of the pools was resuspended and cleaved in vitro using recombinant Dicer enzyme (Stratagene, San Diego, Calif.) according to the manufacturer's instructions. 21-23 bp siRNA products in each of the five pools were purified by a two-step method, first using a Microspin G-25 column (Amersham), followed by a Microcon YM-100 column (Amicon). Each pool of siRNA products was transiently transfected into the T7 cell line using Lipofectamone 2000®.

[0407] Western blots to assay the relative effectiveness of the siRNA pools in suppressing Ku70 O expression were performed approximately 3 days post-transfection. Briefly, cells were lysed and disrupted using RIPA buffer (Santa Cruz Biotechnology), and homogenized by passing the lysates through a QIAshredder (Qiagen, Valencia, Calif.). The clarified lysates were then treated with SDS PAGE sample buffer (with β mercaptoethanol used as the reducing agent) and boiled for 5 minutes. Samples were then resolved on a 4-12% gradient NUPAGE gel and transferred onto a PVDF membrane. The upper portion of the blot was exposed to an anti-Ku70 antibody (Santa Cruz sc-5309) and the lower portion exposed to an anti-TF IIB antibody (Santa Cruz sc-225, used as an input control). The blot was then exposed to horseradish peroxidase-conjugated goat anti-mouse secondary antibody and processed for electrochemiluminescent (ECL) detection using a kit from Pierce Chemical Co. according to the manufacturer's instructions.

[0408] FIG. 39 shows representative results following transfection of two of the siRNA pools (pools D and E) into T7 cells. Transfection with 70 ng of siRNA E results in a significant decrease in Ku70 protein levels (FIG. 39, lane 3).

Example 19

Increasing the Frequency of Homologous Recombination by Inhibition of Expression of a Protein Involved in Non-Homologous End Joining

[0409] Repair of a double-stranded break in genomic DNA can proceed along two different cellular pathways; homologous recombination (HR) or non-homologous end joining (NHEJ). Ku70 is a protein involved in NHEJ, which binds to the free DNA ends resulting from a double-stranded break in genomic DNA. To test whether lowering the intracellular concentration of a protein involved in NHEJ increases the frequency of HR, small interfering RNAs (siRNAs), prepared as described in Example 18, were used to inhibit expression of Ku70 mRNA, thereby lowering levels of Ku70 protein, in cells co-transfected with donor DNA and with plasmids encoding chimeric nucleases.

[0410] For these experiments, the T7 cell line (see Example 9 and FIG. 27) was used. These cells contain a chromosomally-integrated defective eGFP gene, but have been observed to exhibit lower levels of targeted homologous recombination than the T18 cell line used in Examples 11-13.

[0411] T7 cells were transfected, as described in Example 11, with either 70 or 140 ng of one of two pools of dicer product targeting Ku70 (see Example 18). Protein blot analysis was performed on extracts derived from the transfected cells to determine whether the treatment of cells with

siRNA resulted in a decrease in the levels of the Ku70 protein (see previous Example). FIG. 39 shows that levels of the Ku70 protein were reduced in cells that had been treated with 70 ng of siRNA from pool E.

[0412] Separate cell samples in the same experiment were co-transfected with 70 or 140 ng of siRNA (pool D or pool E) along with 50 ng each of the 287-FokI and 296-FokI expression constructs (Example 7 and Table 12) and 500 ng of the 1.5 kbp GFP donor (Example 13), to determine whether lowering Ku70 levels increased the frequency of homologous recombination. The experimental protocol is described in Table 24. Restoration of eGFP activity, due to homologous recombination, was assayed by FACS analysis described in Example 11.

TABLE 24

Expt. #	Donor ¹	ZFNs ²	SiRNA ³	% correction ⁴
1	500 ng	—	—	0.05
2	—	50 ng each	—	0.01
3	500 ng	50 ng each	—	0.79
4	500 ng	50 ng each	70 ng pool D	0.68
5	500 ng	50 ng each	140 ng pool D	0.59
6	500 ng	50 ng each	70 ng pool E	1.25
7	500 ng	50 ng each	140 ng pool E	0.92

¹A plasmid containing a 1.5 kbp sequence encoding a functional eGFP protein which is homologous to the chromosomally integrated defective eGFP gene

²Plasmids encoding the eGFP-targeted 287 and 296 zinc finger protein/FokI fusion endonucleases

³See Example 18

⁴Percent of total fluorescence exhibiting high emission at 525 nm and low emission at 570 nm (region E of the FACS trace, see Example 11).

[0413] The percent correction of the defective eGFP gene in the transfected T7 cells (indicative of the frequency of targeted homologous recombination) is shown in the right-most column of Table 24. The highest frequency of targeted recombination is observed in Experiment 6, in which cells were transfected with donor DNA, plasmids encoding the two eGFP-targeted fusion nucleases and 70 ng of siRNA Pool E. Reference to Example 18 and FIG. 39 indicates that 70 ng of Pool E siRNA significantly depressed Ku70 protein levels. Thus, methods that reduce cellular levels of proteins involved in NHEJ can be used as a means of facilitating homologous recombination.

Example 20

Zinc Finger-FokI Fusion Nucleases Targeted to the Human β -globin Gene

[0414] A number of four-finger zinc finger DNA binding domains, targeted to the human β -globin gene, were designed and plasmids encoding each zinc finger domain, fused to a FokI cleavage half-domain, were constructed. Each zinc finger domain contained four zinc fingers and recognized a 12 bp target site in the region of the human β -globin gene encoding the mutation responsible for Sickle Cell Anemia. The binding affinity of each of these proteins to its target sequence was assessed, and four proteins exhibiting strong binding (sca-r29b, sca-36a, sca-36b, and sca-36c) were used for construction of FokI fusion endonucleases.

[0415] The target sites of the ZFP DNA binding domains, aligned with the sequence of the human β -globin gene, are

shown below. The translational start codon (ATG) is in bold and underlined, as is the A-T substitution causing Sickle Cell Anemia.

sca-36a GAAGTCTGCCGT (SEQ ID NO:178)

sca-36b GAAGTCtGCCGTT (SEQ ID NO:179)

sca-36c GAAGTCtGCCGTT (SEQ ID NO:180)

CAAACAGACACC**AT**GGTGCATCTG (SEQ ID NO:181)

ACTCCTG**TGG**GAGAAGTCTGCCGTT

ACTGGTTTGTCTGTGGTACACGT

AGACTGAGGAC**A**CCTCTTCAGACG

GCAATGAC

sca-r29b ACGTAGaCTGAGG (SEQ ID NO:182)

[0416] Amino acid sequences of the recognition regions of the zinc fingers in these four proteins are shown in Table 25. The complete amino acid sequences of these zinc finger domains are shown in FIG. 40. The sca-36a domain recognizes a target site having 12 contiguous nucleotides (shown in upper case above), while the other three domain recognize a thirteen nucleotide sequence consisting of two six-nucleotide target sites (shown in upper case) separated by a single nucleotide (shown in lower case). Accordingly, the sca-r29b, sca-36b and sca-36c domains contain a non-canonical inter-finger linker having the amino acid sequence TGGGGSQKP (SEQ ID NO: 183) between the second and the third of their four fingers.

sca-r29b, sca-36a, sca-36b, and sca-36c protein were incubated in an in vitro transcription/translation system. See Example 4. A total of 2 ul of the TNT reaction (2 ul of a single reaction when a single protein was being assayed or 1 ul of each reaction when a pair of proteins was being assayed) was added to 13 ul of the cleavage buffer mix and 3 ul of labeled probe (~1 ng/ul). The probe was end-labeled with ³²P using polynucleotide kinase. This reaction was incubated for 1 hour at room temperature to allow binding of the ZFNs. Cleavage was stimulated by the addition of 8 ul of 8 mM MgCl₂, diluted in cleavage buffer, to a final concentration of approximately 2.5 mM. The cleavage reaction was incubated for 1 hour at 37° C. and stopped by the addition of 11 ul of phenol/chloroform. The DNA was isolated by phenol/chloroform extraction and analyzed by gel electrophoresis, as described in Example 4. As a control, 3 ul of probe was analyzed on the gel to mark the migration of uncut DNA (labeled “U” in FIG. 41).

[0419] The results are shown in FIG. 41. Incubation of the target DNA with any single zinc finger/FokI fusion resulted in no change in size of the template DNA. However, the combination of the sca-r29b nuclease with either of the sca-36b or sca-36c nucleases resulted in cleavage of the target DNA, as evidenced by the presence of two shorter DNA fragments (rightmost two lanes of FIG. 41).

TABLE 25

ZFP	F1	F2	F3	F4
sca-r29b	QSGDLTR (SEQ ID NO:184)	TSANLSR (SEQ ID NO:185)	DRSALSR (SEQ ID NO:186)	QSGHLSR (SEQ ID NO:187)
sca-36a	RSQTRKT (SEQ ID NO:188)	QKRNRTK (SEQ ID NO:189)	DRSALSR (SEQ ID NO:190)	QSGNLAR (SEQ ID NO:191)
sca-36b	TSGSLSR (SEQ ID NO:192)	DRSDLR (SEQ ID NO:193)	DRSALSR (SEQ ID NO:194)	QSGNLAR (SEQ ID NO:195)
sca-36c	TSSSLSR (SEQ ID NO: 196)	DRSDLR (SEQ ID NO:197)	DRSALSR (SEQ ID NO:198)	QSGNLAR (SEQ ID NO:199)

Example 21

In Vitro Cleavage of a DNA Target Sequence by
β-globin-targeted ZFP/FokI Fusion Endonucleases

[0417] Fusion proteins containing a FokI cleavage half-domain and one the four ZFP DNA binding domains described in the previous example were tested for their ability to cleave DNA in vitro with the predicted sequence specificity. These ZFP domains were cloned into the pcDNA3.1 expression vector via KpnI and BamHI sites and fused in-frame to the FokI cleavage domain via a 4 amino acid ZC linker, as described above. A DNA fragment containing 700 bp of the human β-globin gene was cloned from genomic DNA obtained from K562 cells. The isolation and sequence of this fragment was described in Example 3, supra.

[0418] To produce fusion endonucleases (ZFNs) for the in vitro assay, circular plasmids encoding FokI fusions to

Example 22

ZFP/FokI Fusion Endonucleases, Targeted to the
β-globin Gene, Tested in a Chromosomal GFP
Reporter System

[0420] A DNA fragment containing the human P-globin gene sequence targeted by the ZFNs described in Example 20 was synthesized and cloned into a SpeI site in an eGFP reporter gene thereby, disrupting eGFP expression. The fragment contained the following sequence, in which the nucleotide responsible for the sickle cell mutation is in bold and underlined):

(SEQ ID NO:200)

CTAGACACCATGGTGCATCTGACTCCTG**TGG**GAGAAGTCTGCCGTACTGC

CCTAG

[0421] This disrupted eGFP gene containing inserted β -globin sequences was cloned into pcDNA4/TO (Invitrogen, Carlsbad, Calif.) using the HindIII and NotI sites, and the resulting vector was transfected into HEK293 TRex cells (Invitrogen). Individual stable clones were isolated and grown up, and the clones were tested for targeted homologous recombination by transfecting each of the sca-36 proteins (sca-36a, sca-36b, sca-36c) paired with sca-29b (See Example 20 and Table 25 for sequences and binding sites of these chimeric nucleases). Cells were transfected with 50 ng of plasmid encoding each of the ZFNs and with 500 ng of the 1.5-kb GFP Donor (Example 13). Five days after transfection, cells were tested for homologous recombination at the inserted defective eGFP locus. Initially, cells were examined by fluorescence microscopy for eGFP function. Cells exhibiting fluorescence were then analyzed quantitatively using a FACS assay for eGFP fluorescence, as described in Example 11.

[0422] The results showed that all cell lines transfected with sca-29b and sca-36a were negative for eGFP function, when assayed by fluorescence microscopy. Some of the lines transfected with sca-29b paired with either sca-36b or sca-36c were positive for eGFP expression, when assayed by fluorescence microscopy, and were therefore further analyzed by FACS analysis. The results of FACS analysis of two of these lines are shown in Table 26, and indicate that zinc finger nucleases targeted to β -globin sequences are capable of catalyzing sequence-specific double-stranded DNA cleavage to facilitate homologous recombination in living cells.

TABLE 26

Cell line	DNA transfected:				% corr. ¹
	sca-29b	sca-36a	sca-36b	sca-36c	
#20	+	+			0
	+		+		0.08
	+			+	0.07
#40	+	+			0
	+		+		0.18
	+			+	0.12

¹Percent of total fluorescence exhibiting high emission at 525 nm and low emission at 570 nm (region E of the FACS trace, see Example 11).

Example 23

Effect of Transcription Level on Targeted Homologous Recombination

[0423] Since transcription of a chromosomal DNA sequence involves alterations in its chromatin structure (generally to make the transcribed sequences more accessible), it is possible that an actively transcribed gene might be a more favorable substrate for targeted homologous recombination. This idea was tested using the T18 cell line (Example 9) which contains chromosomal sequences encoding a defective eGFP gene whose transcription is under the control of a doxycycline-inducible promoter.

[0424] Separate samples of T18 cells were transfected with plasmids encoding the eGFP-targeted 287 and 296 zinc finger/FokI fusion proteins (Example 7) and a 1.5 kbp donor DNA molecule containing sequences that correct the defect in the chromosomal eGFP gene (Example 9). Five hours

after transfection, transfected cells were treated with different concentrations of doxycycline, then eGFP mRNA levels were measured 48 hours after addition of doxycycline. eGFP fluorescence at 520 nm (indicative of targeted recombination of the donor sequence into the chromosome to replace the inserted β -globin sequences) was measured by FACS at 4 days after transfection.

[0425] The results are shown in FIG. 42. Increasing steady-state levels of eGFP mRNA normalized to GAPDH mRNA (equivalent, to a first approximation, to the rate of transcription of the defective chromosomal eGFP gene) are indicated by the bars. The number above each bar indicate the percent of cells exhibiting eGFP fluorescence. The results show that increasing transcription rate of the target gene is accompanied by higher frequencies of targeted recombination. This suggests that targeted activation of transcription (as disclosed, e.g. in co-owned U.S. Pat. Nos. 6,534,261 and 6,607,882) can be used, in conjunction with targeted DNA cleavage, to stimulate targeted homologous recombination in cells.

Example 24

Generation of a Cell Line Containing a Mutation in the IL-2R γ Gene

[0426] K562 cells were transfected with plasmids encoding the 5-8GL0 and the 5-9DL0 zinc finger nucleases (ZFNs) (see Example 14; Table 17) and with a 1.5 kbp DraI donor construct. The DraI donor is comprised of a sequence with homology to the region encoding the 5th exon of the IL2R γ gene, but inserts an extra base between the ZFN-binding sites to create a frameshift and generate a DraI site.

[0427] 24 hours post-transfection, cells were treated with 0.2 μ M vinblastine (final concentration) for 30 hours. Cells were washed three times with PBS and re-plated in medium. Cells were allowed to recover for 3 days and an aliquot of cells were removed to perform a PCR-based RFLP assay, similar to that described in Example 14, testing for the presence of a DraI site. It was determined the gene correction frequency within the population was approximately 4%.

[0428] Cells were allowed to recover for an additional 2 days and 1600 individual cells were plated into 40 $\times$ 96-well plates in 100 μ l of medium.

[0429] The cells are grown for about 3 weeks, and cells homozygous for the DraI mutant phenotype are isolated. The cells are tested for genome modification (by testing for the presence of a DraI site in exon 5 of the IL-2R γ gene) and for levels of IL-2R γ mRNA (by real-time PCR) and protein (by Western blotting) to determine the effect of the mutation on gene expression. Cells are tested for function by FACS analysis.

[0430] Cells containing the DraI frameshift mutation in the IL-2R γ gene are transfected with plasmids encoding the 5-8GL0 and 5-9DL0 fusion proteins and a 1.5 kb BsrBI donor construct (Example 14) to replace the DraI frameshift mutation with a sequence encoding a functional protein. Levels of homologous recombination greater than 1% are obtained in these cells, as measured by assaying for the presence of a BsrBI site as described in Example 14. Recovery of gene function is demonstrated by measuring mRNA and protein levels and by FACS analysis.

Example 25

ZFP/FokI Fusion Endonucleases with Different Polarities

[0431] A vector encoding a ZFP/FokI fusion, in which the ZFP domain was N-terminal to the FokI domain, was constructed. The ZFP domain, denoted IL2-1, contained four zinc fingers, and was targeted to the sequence AACTCG-GATAAT (SEQ ID NO:202), located in the third exon of the IL-2R γ gene. The amino acid sequences of the recognition regions of the zinc fingers are given in Table 27.

sequences, was also constructed. The IL2-1 four-finger zinc finger domain was inserted, as a KpnI/BamHI fragment, into a vector encoding a fusion protein containing a NLS, the KOX-1 repression domain, EGFP and a FLAG epitope tag, that had been digested with KpnI and BamHI to release the EGFP-encoding sequences. This generated a vector containing sequences encoding, from N-terminus to C-terminus, a NLS (from the SV40 large T-Antigen), a KOX repression domain, the IL2-1 zinc finger domain and a FLAG epitope tag. This construct was then digested with EcoRI and KpnI to release the NLS- and KOX-encoding sequences, and an

TABLE 27

Zinc Finger Design of IL2-1 binding domain				
Target sequence	F1 (AAT)	F2 (GAT)	F3 (TCG)	F4 (AAC)
AACTCGGATAAT (SEQ ID NO:203)	DRSTLIE (SEQ ID NO:204)	SSSNSLR (SEQ ID NO:205)	RSDDLK (SEQ ID NO:206)	DNSNRIK (SEQ ID NO:207)

Note:
The DNA target sequence is shown in the left-most column. The remaining columns show the amino acid sequences (in one-letter code) of residues -1 through +6 of each of the four zinc fingers, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein. The three-nucleotide subsite bound by each finger is shown in the top row adjacent to the finger designation.

[0432] Sequences encoding this zinc finger domain were joined to sequences encoding the cleavage half-domain of the FokI restriction endonuclease (amino acids 384-579 according to Looney et al. (1989) *Gene* 80:193-208) such that a four amino acid linker was present between the ZFP domain and the cleavage half-domain (i.e., a four amino acid ZC linker). The FokI cleavage half-domain was obtained by PCR amplification of genomic DNA isolated from the bacterial strain *Planomicrobium okeanokoites* (ATCC 33414) using the following primers:

(SEQ ID NO:208)
5'-GGATCCCACTAGTCAAAAGTGAAC
(SEQ ID NO:209)
5'-CTCGAGTTAAAGTTTATCTCGCCG.

[0433] The PCR product was digested with BamHI and XhoI (sites underlined in sequences shown above) and then ligated with a vector fragment prepared from the plasmid pcDNA-nls-ZFP1656-VP16-flag after BamHI and XhoI digestion. The resulting construct, pcDNA-nls-ZFP1656-FokI, encodes a fusion protein containing, from N-terminus to C-terminus, a SV40 large T antigen-derived nuclear localization signal (NLS, Kalderon et al. (1984) *Cell* 39:499-509), ZFP1656, and a FokI cleavage half-domain, in a pcDNA3.1 (Invitrogen, Carlsbad, Calif.) vector backbone. This construct was digested with KpnI and BamHI to release the ZFP1656-encoding sequences, and a KpnI/BamHI fragment encoding the IL2-1 zinc finger binding domain was inserted by ligation. The resulting construct (pIL2-1C) encodes a fusion protein comprising, from N- to C-terminus, a nuclear localization signal, the four-finger IL2-1 zinc finger binding domain and a FokI cleavage half-domain, with a four amino acid ZC linker.

[0434] A vector encoding a ZFP/FokI fusion protein, in which the FokI sequences were N-terminal to the ZFP

EcoRI/KpnI fragment (generated by PCR using, as template, a vector encoding FokI) encoding amino acids 384-579 of the FokI restriction enzyme and a NLS was inserted. The resulting construct, pIL2-1R encodes a fusion protein containing, from N-terminus to C-terminus, a FokI cleavage half-domain, a NLS, and the four-finger IL2-1 ZFP binding domain. The ZC linker in this construct is 21 amino acids long and includes the seven amino acid nuclear localization sequence (PKKKRKV; SEQ ID NO: 210).

[0435] The 5-9D zinc finger domain binds the 12-nucleotide target sequence AAAGCGGCTCCG (SEQ ID NO: 157) located in the fifth exon of the IL-2R γ gene. See Example 14 (Table 17). Sequences encoding the 5-9D zinc finger domain were inserted into a vector to generate a FokI/ZFP fusion, in which the FokI sequences were N-terminal to the ZFP sequences. To make this construct, the pIL2-1R plasmid described in the previous paragraph was digested with KpnI and BamHI to release a fragment containing sequences encoding the IL2-1 zinc finger binding domain, and a KpnI/BamHI fragment encoding the 5-9D zinc finger binding domain was inserted in its place. The resulting construct, p5-9DR, encodes a fusion protein containing, from N-terminus to C-terminus, a FokI cleavage half-domain, a NLS, and the four-finger 5-9D zinc finger binding domain. The ZC linker in this construct is 22 amino acids long and includes the seven amino acid nuclear localization sequence (PKKKRKV; SEQ ID NO: 210).

[0436] See co-owned U.S. Pat. Nos. 6,453,242 and 6,534,261 for additional details of vector construction.

Example 26

Construction of Synthetic Substrates for DNA Cleavage

[0437] The target sequences bound by the IL2-1 and 5-9D fusion proteins described above were introduced into

double-stranded DNA fragments in a variety of orientations, to test the cleavage ability of zinc finger/FokI fusion proteins having an altered polarity in which the FokI domain is N-terminal to the ZFP domain. In template 1, the 5-9D target site is present in one strand and the IL2-1 target site is present on the complementary strand, with the 3' ends of the binding sites being proximal to each other and separated by six intervening nucleotide pairs. In template 2, the 5-9D and IL2-1 target sites are present on the same DNA strand, with the 3' end of the 5-9D binding site separated by six nucleotide pairs from the 5' end of the IL2-1 binding site.

[0438] DNA fragments of approximately 442 base pairs, containing the sequences described above, were obtained as amplification products of plasmids into which the templates had been cloned. The IL2-1 and 5-9D target sites were located within these fragments such that double-stranded DNA cleavage between the two target sites would generate DNA fragments of approximately 278 and 164 base pairs. Amplification products were radioactively labeled by transfer of orthophosphate from γ -32P-ATP using T4 polynucleotide kinase.

Example 27

Targeted DNA Cleavage with Zinc Finger/FokI Fusions Having Altered Polarity

[0439] The IL2-1C, IL2-1R and 5-9DR fusion proteins were obtained by incubating plasmids encoding these proteins in a TNT coupled reticulocyte lysate (Promega, Madison, Wis.). Cleavage reactions were conducted in 23 μ l of a mixture containing 1 μ l of TNT reaction for each fusion protein, 1 μ l labeled digestion substrate and 20 μ l cleavage buffer. Cleavage buffer was prepared by adding 1 μ l of 1M dithiothreitol and 50 μ l of bovine serum albumin (10 mg/ml) to 1 ml of 20 mM Tris-Cl, pH 8.5, 75 mM NaCl, 10 μ M ZnCl₂, 5% (v/v) glycerol. Cleavage reactions were incubated at 37° C. for 2 hours, then shaken with 13 μ l phenol/chloroform/isoamyl alcohol (25:24:1). After centrifugation, 10 μ l of the aqueous phase was analyzed on a 10% polyacrylamide gel. Radioactivity in the gel was detected using a Phosphorimager (Molecular Dynamics) and quantitated using ImageQuant software (Molecular Dynamics).

[0440] FIG. 44 shows the results obtained using two chimeric nucleases having a NH₂-FokI domain-zinc finger domain-COOH polarity to cleave a substrate in which the binding sites for the two chimeric nucleases are located on opposite strands and the 3' ends of the binding sites are proximal to each other and separated by six nucleotide pairs. Incubation of the substrate with either of the IL2-1R or 5-9DR nucleases alone does not result in cleavage of the substrate (compare lanes 2 and 3 with lane 1), while incubation of both nucleases results in almost complete cleavage of the DNA substrate at the intended target site (lane 4).

[0441] FIG. 45 shows the ability of a first chimeric nuclease having a NH₂-zinc finger domain-FokI domain-COOH polarity, and a second chimeric nuclease having a NH₂-FokI domain-zinc finger domain-COOH polarity, to cleave a substrate in which the binding sites for the two chimeric nucleases are located on the same strand, and the 3' end of the first binding site is proximal to the 5' end of the second binding site and separated from it by six nucleotide pairs. Only the combination of the 5-9DR and the IL2-1C

nucleases (i.e. each nuclease having a different polarity) was successful in cleaving the substrate having both target sites on the same strand (compare lane 6 with lanes 1-5).

Example 28

Chimeric Nucleases with Different ZC Linker Lengths

[0442] Two sets of fusion proteins with different ZC linker lengths, in which the FokI domain is amino terminal to the ZFP domain, were designed. The FokI domain is amino acids 384-579 according to Looney et al. (1989) *Gene* 80:193-208. The ZFP domain was selected from the IL1-2 (Table 27), 5-8G (Table 17) and 5-9D (Table 17) domains. The first set had the structure NH₂-NLS-FokI-ZFP-Flag-COOH. In this set, proteins having ZC linker lengths of 13, 14, 18, 19, 28 and 29 amino acids were designed. The second set had the structure NH₂-FokI-NLS-ZFP-Flag-COOH and proteins with ZC linkers of 21, 22, 23, 24, 28, 29, 38 and 39 amino acids were designed. Note that, in the second set, the NLS is part of the ZC linker. Plasmids encoding these fusion proteins are also constructed.

[0443] Model DNA sequences were designed to test the cleavage activity of these fusion proteins and to determine optimal ZC linker lengths as a function of distance between the target sites for the two fusion proteins. The following sequences were designed:

- [0444] 1. 5-9D target site and IL2-1 target site on opposite strands
- [0445] 2. 5-9D target site and IL2-1 target site on same strand
- [0446] 3. 5-9D target site and 5-8G target site on opposite strands
- [0447] 4. 5-9D target site and 5-8G target site on same strand

[0448] For each of these four pairs of target sites, sequences are constructed in which the separation between the two target sites is 4, 5, 6 or 7 base pairs.

[0449] These sequences are introduced into labeled substrates as described in Example 26 and are used to test the various fusion proteins described in this example for their ability to cleave DNA, according to the methods described in Example 27.

Example 29

Construction of a Stable Cell Line Containing an Integrated Defective eGFP Reporter Gene

[0450] An eGFP (enhanced green fluorescent protein) coding sequence containing a frameshift mutation and a fragment of exon 5 of the IL-2R γ gene, operatively linked to a tetracycline-regulated CMV promoter, was constructed as follows. A silent mutation was inserted into the eGFP coding sequences in the pEGFP-N1 vector (BD Biosciences) to create a novel SpeI site. Subsequently a one-nucleotide deletion (creating a frameshift mutation) was introduced downstream of the new SpeI site. The following sequence from exon 5 of the IL-2R γ gene, containing target sites for

the 5-8G and 5-9D zinc finger/FokI fusion proteins (described in Example 14, Table 17, supra), was inserted into the newly-introduced SpeI site:

(SEQ ID NO:214)
CTAGCTACACGTTTCGTGTTTCGGAGCGCTTTAACCCACTCTGTGGAAGT
GCTCCTAG

[0451] The resulting plasmid contained sequences encoding mutant eGFP containing a fragment of DNA sequence from exon 5 of the IL2R γ gene. This plasmid was digested with HindIII and NotI, releasing a fragment containing the mutated eGFP sequence (including the inserted IL-2R γ exon 5 sequences). This fragment was inserted into the HindIII and NotI sites of the pcDNA4/TO vector (Invitrogen), resulting in a construct in which expression of eGFP sequence is controlled by a 2 $\times$ tet-operator-regulated CMV promoter. A schematic diagram of this plasmid is shown in FIG. 46.

[0452] This construct was used to transform HEK293 TRex cells (Invitrogen), and a stable cell line containing an integrated copy of this construct was isolated. In this cell line, the eGFP coding sequences are transcribed upon addition of doxycycline, but because of the frameshift mutation and the IL-2R γ insertion, no functional protein is expressed.

Example 30

Targeted Homology-dependent Integration of a Puromycin Resistance Marker into a Chromosomal eGFP Gene

[0453] An experiment was conducted to test for integration of a puromycin resistance marker into the mutant chromosomal eGFP gene described in the preceding Example.

[0454] A promoterless donor was constructed that contained sequences encoding puromycin resistance (denoted "puro sequences"), flanked by sequences homologous to the eGFP cDNA construct, as follows. Sequences were PCR amplified from the pTRE2pur-HA vector (BD Biosciences) to generate a puro sequence with flanking SpeI sites and a consensus Kozak sequence upstream of the ATG initiation codon. Amplification primers were:

(SEQ ID NO:215)
puro-5': ACTAGTGCCGCCACCATGACCGAGTACAAGCCCA

(SEQ ID NO:216)
puro-3': ACTAGTCAGGCACCGGGCTT

[0455] This PCR fragment was cloned into the pEGFP-N1 vector containing a modified eGFP gene that encoded a novel SpeI restriction site and a frameshift mutation that prevented functional expression of the gene (see Example 29). This eGFP/Puromycin gene was cloned into the pcDNA4/TO vector, via HindIII and NotI sites, to create the vector pcDNA4/TO/GFPpuro, which also served as the positive control in experiments to obtain Puromycin resistant cells by targeted integration. In order to create a promoterless donor, the pcDNA4/TO/GFPpuro vector was PCR amplified with the following primers:

(SEQ ID NO:217)
GFP-Bam CGAATTCTGCAGTCGAC

(SEQ ID NO:218)
pcDNA42571 TGCATACTTCTGCCTGC

[0456] The resulting amplification product was Topo cloned into the pCR4-TOPO vector and its sequence was confirmed. This created a donor with 413 bp of sequence homologous to the chromosomal eGFP construct upstream of the puro sequences and 1285 bp of sequence homologous to the chromosomal eGFP construct downstream of the puro sequences.

[0457] To test for targeted integration of puro sequences, the cell line described in Example 29 was subjected to targeted DNA cleavage by zinc finger/FokI fusion proteins in the presence of the donor construct described in this example, transfected cells were selected for puromycin resistance and their chromosomal DNA was analyzed. The two zinc finger/FokI fusion proteins (ZFNs), designed to cleave within the exon 5 sequences inserted into the eGFP gene (5-8G and 5-9D) have been described in Example 14, Table 17, supra. Puromycin resistance can arise from either homology-dependent or homology-independent integration of donor sequences at the cleavage site located within the IL-2R γ sequences inserted into the eGFP coding sequences. Homology-dependent integration of the donor construct will result in replacement of IL-2R γ sequences by the puro sequences.

[0458] HEK 293 cells were grown in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen), supplemented with 10% fetal bovine serum (FBS) (Hyclone) and 2 mM L-glutamine and maintained at 37° C. in an atmosphere of 5% CO₂. To test for targeted integration of the puro sequences, cells were transfected with 50 ng of each ZFN-encoding plasmid and 500 ng of donor plasmid. In negative control experiments, cells were transfected either with 50 ng of each ZFN-encoding plasmid or 500 ng of donor plasmid. As a positive control, 500 ng of the pcDNA4/TO/GFPpuro vector was transfected into HEK293 cells. Cells were transfected using LipofectAMINE 2000 Reagent (Invitrogen) in Opti-MEM I reduced serum medium. Puromycin resistance was assayed by the addition of doxycycline to 2 ng/ml (to activate transcription of the integrated sequences) and puromycin to 2 μ g/ml (final concentration) in the growth medium.

[0459] Puromycin resistant colonies were obtained only from cells that had been transfected with both ZFN-encoding plasmids and the donor plasmid. Twenty-four clonal populations were isolated, subjected to >6-weeks of selection, then analyzed by PCR for a targeted integration event. The following PCR primers were used to detect whether a targeted integration event occurred:

(SEQ ID NO:219)
CMVPuro-5' TTTGACCTCCATAGAAGACA

(SEQ ID NO:220)
CMVPuro-3' GCGCACCGTGGGCTTGCTACT

[0460] One of the primers is complementary to the exogenous puro sequences and the other is complementary to sequences in the CMV promoter present in the integrated reporter construct. Twenty-one out of 24 colonies yielded amplification products whose sizes were consistent with targeted integration of puro sequences. These fragments were cloned and their nucleotide sequences were determined. Sequence analysis indicated that eight out of the 24 clones had undergone homology-directed integration of puro sequences into the chromosomal eGFP construct, while 13 had undergone homology-independent integration of donor DNA into chromosomal sequences, accompanied by partial duplication of the puro sequences.

Example 31

Codon Optimization of Zinc Finger/FokI Fusion Proteins Targeted to Exon 5 of the IL-2R γ Gene

[0461] Fusion proteins containing the 5-8G and 5-9D zinc finger binding domains (Table 17) joined to a FokI cleavage half-domain by a 4 amino acid ZC linker (L0) have been described supra. See, e.g., Example 14 and Example 24. Polynucleotides encoding these two fusion proteins were designed so that the codons were optimized for expression in mammalian cells. The codon-optimized nucleotide sequences encoding these two fusion proteins are as follows:

5-8G L0 FokI

(SEQ ID NO:221)

```
aattcgctagcgccaccatggcccccagaagaagaggaaagtgggaatc
cacgggtaccggcgctatggccgagagggcccttcagtgctggatctg
catgcggaacttcagcggagcgacacccctgagcgtgcacatccgcacc
acacaggcgagaagccttttgcctgtgacatcttggaggagaaatttgc
cgcaagccaccggcatcggaacacacccaagatccacacggatctcagaa
gccctttcagtgagaatctgcatgagaaacttctcccggtccgacacc
tgagcgaacacatcaggacacacacggcgagaaaccccttcgctgcgac
atctgtggccgcaagtttgcgcgacgaagcaccgcacaaatcacacaaa
gattcacctgcggggatcccagctggtgaagagcgagctggaggagaaga
agtcgagctgcggcacaagctgaagtacgtgcccacgagtagcatcgag
ctgatcgagatcgccagggaacacgacccagcagcgcacccctggagatgaa
ggtgatggagttcttcatgaaggtgtacggctacaggggaaagcacctgg
gcgggaagcagaaagcctgacggcgccatctatacagtgggcagccccatc
gattacggcggtgatcgtggacacaaaaggcctacagcggcggtacaaatct
gcctatcgccggcggcagcagagatgcagagatacgtggaggagaaccaga
cccggaataagcacatcaacccccaacgagtggtggaaggtgtacccctagc
agcgtgaccgagttcaagttcctgttcgtgagcggccacttcaagggcaa
ctacaaggcccgctgaccaggctgaaccacatcaccaactgcaatggcg
ccgtgctgagcgtggaggagctgctgacggcgcgagatgatcaagcc
ggcaccctgacactggaggaggtgcggcgcaagttcaacaacggcgagat
caacttctgataac
```

5-9D L0 FokI

(SEQ ID NO:222)

```
aattcgctagcgccaccatggcccccagaagaagaggaaagtgggaatc
cacgggtaccggcgctatggccgagagggcccttcagtgctggatctg
catgcggaacttcagcaggagcgacacccctgagcgaacacatccgcacc
acacaggcgagaagccttttgcctgtgacatcttggaggagaaatttgc
cgcaagaccaccggcatcggaacacacccaagatccacacggatctcagaa
gccctttcagtgagaatctgcatgagaaacttctcccggtccgacagcc
tgagcaagcacattagacccacacggggagaaaccccttcgctgcgac
atctgtggccgcaaatgttgcgcgacgagcaacccctgaaagtgcacacaa
gattcacctgcggggatcccagctggtgaagagcgagctggaggagaaga
agtcgagctgcggcacaagctgaagtacgtgcccacgagtagcatcgag
ctgatcgagatcgccagggaacacgacccagcagcgcacccctggagatgaa
ggtgatggagttcttcatgaaggtgtacggctacaggggaaagcacctgg
gcgggaagcagaaagcctgacggcgccatctatacagtgggcagccccatc
gattacggcggtgatcgtggacacaaaaggcctacagcggcggtacaaatct
gcctatcgccggcggcagcagagatgcagagatacgtggaggagaaccaga
cccggaataagcacatcaacccccaacgagtggtggaaggtgtacccctagc
agcgtgaccgagttcaagttcctgttcgtgagcggccacttcaagggcaa
ctacaaggcccgctgaccaggctgaaccacatcaccaactgcaatggcg
```

-continued

```
ccgtgctgagcgtggaggagctgctgacggcgcgagatgatcaagcc
ggcaccctgacactggaggaggtgcggcgcaagttcaacaacggcgagat
caacttctgataac
```

Example 32

Growth and Transfection of K-562 Cells for Targeted Homologous Integration

[0462] Human K-562 erythroleukemia cells (ATCC) were cultured at 37° C. in DMEM supplemented with 10% fetal bovine serum, penicillin and streptomycin, and transfected (Nucleofector; Amaxa) with 2.5 μ g of an expression vector encoding two zinc finger nucleases (ZFN) designed to introduce a double-strand break at a position surrounding the codon for arginine 226 in the endogenous IL2R γ gene. The two nucleases (5-8G and 5-9D) have been described in Example 14, Table 17, supra. Sequences encoding the nucleases were separated by sequences encoding a 2A peptide. See, e.g., Szymczak et al. (2004) *Nature Biotechnol.* 22:589-594. At the same time, the cells were transfected with either 25 or 50 μ g of a donor DNA plasmid carrying a 1.5 kb DNA stretch of IL2R γ chromosomal DNA sequence centered on exon 5 (Urnov et al. (2005) *Nature* 435:646-651), interrupted by the DNA sequence to be inserted (see examples 33-36 below). Seventy two hours after transfection, genomic DNA was isolated (DNEasy; Qiagen) and cell genotype at the IL2R γ locus was determined by PCR of the exon 5-containing stretch of the X chromosome, using primers that anneal outside of the 1.5 kb region of donor homology and generate a 1.6 kilobase pair amplification product from the wild-type IL2R γ sequence (Urnov et al., supra). PCR products were analyzed by gel electrophoresis and, where indicated, by restriction digestion. Control samples included: (1) cells transfected with a GFP-encoding expression vector, (2) cells transfected solely with the expression vector encoding the ZFNs, and (3) cells transfected solely with the donor DNA molecule.

Example 33

Targeted Homology-dependent Integration of a 12-nucleotide Exogenous Sequence into the Endogenous IL-2R γ Gene

[0463] Cell growth and transfection were conducted as described in Example 32. The donor DNA molecule was engineered to contain a 12 nucleotide pair sequence tag containing a novel diagnostic recognition site for the restriction enzyme StuI. Cellular DNA was isolated and used as a template for amplification as described in Example 32, then digested with StuI. As shown in FIG. 47, all control samples carried chromosomes yielding amplification products that were resistant to cleavage by the restriction enzyme. In contrast, 15% of all amplification products in the cell sample transfected both with the donor DNA molecule and the ZFN expression construct were sensitive to the restriction enzyme, indicating integration of the donor DNA. Direct nucleotide sequence determination of the chromosome-derived PCR product confirmed that integration was homology-dependent.

Example 34

Targeted Homology-dependent Integration of
Exogenous Open Reading Frames into the
Endogenous IL-2R γ Gene

[0464] Cell growth and transfection were conducted as described in Example 32. In this experiment, two different donor DNA molecules were used. Donor DNA molecule #1 was engineered to contain the entire 720 bp ORF of enhanced green fluorescent protein (eGFP) flanked by sequences homologous to the chromosomal IL2R γ locus (see Example 32). Donor DNA molecule #2 contained a 924 bp sequence consisting of the entire eGFP ORF followed by a polyadenylation signal; this sequence was flanked by IL2R γ -homologous sequences (see Example 32). Following transfection, cellular DNA was isolated and used as a template for amplification as described in Example 32. As shown in FIG. 48, all control samples carried chromosomes yielding PCR products of wild-type size (~1.6 kb). In contrast, 3-6% of all chromosomes in the cell samples transfected both with the ORF-carrying donors and the ZFN expression construct yielded amplification products that were larger than the wild-type-chromosome-derived PCR product, and the difference in size was consistent with the notion that ZFN-driven targeted integration of the eGFP ORFs had occurred. Direct nucleotide sequence determination of the chromosome-derived PCR products confirmed this observation and also indicated that integration was homology-dependent.

Example 35

Targeted Homology-dependent Integration of an
Exogenous "Therapeutic Half-gene" into the
Endogenous IL-2R γ Gene

[0465] Cell growth and transfection were conducted as described in Example 32. The donor DNA molecule consisted of a 720 nucleotide-pair partial IL2R γ cDNA containing the downstream portion of exon 5, and complete copies of exons 6, 7 and 8, (including a translation termination codon and a polyadenylation signal within exon 8). These cDNA sequences were flanked, on one side, by sequences homologous to the upstream portion of exon 5 and the adjoining portion of intron 4 and, on the other side, by sequences homologous to the downstream portion of exon 5 and the adjoining portion of intron 5 (see FIG. 49). Because two copies of the downstream portion of exon 5 are present in the donor construct, and to ensure that recombination occurred in the copy adjacent to exon 8, several silent sequence changes were introduced into the copy adjacent to exon 6. These changes did not alter the coding potential of the exogenous exon 5 sequences, but introduced sufficient non-homology with chromosomal sequences to prevent use of these sequences for the initial homing event in the repair of the break. Thus, integration of the donor construct at the site targeted by the nucleases can be used to correct any IL2R γ mutation in exons 6, 7 or 8 and in the downstream portion of exon 5 contained in the donor construct.

[0466] Following transfection, cellular DNA was isolated and used as a template for amplification as described in Example 32. As shown in FIG. 50, a control sample in which cells were transfected with a GFP-encoding plasmid contained chromosomes yielding PCR products of wild-type

size only (~1.6 kb). In contrast, 6% of all chromosomes in the cell samples transfected both with the therapeutic half-gene-carrying donor and the ZFN expression construct were larger than the wild-type-chromosome-derived PCR product, and the difference in size was consistent with the notion that ZFN-driven targeted integration of the "therapeutic half-gene" had occurred. Direct nucleotide sequence determination of the larger PCR product confirmed that homology-dependent integration of the donor construct had occurred.

Example 36

Targeted Homology-dependent Integration of an
Exogenous 7.7 Kilobase Pair Expression Construct
into the Endogenous IL-2R γ Gene

[0467] Cell growth and transfection were conducted as described in Example 32. A donor DNA molecule was constructed that contained a 7.7 kbp antibody expression construct flanked by sequences homologous to IL2R γ exon 5 and adjacent sequences (See Example 32). In this experiment, two topological forms of the donor were used: a plasmid donor, in which the vector backbone abuts an insert with two homology arms interrupted by the expression construct ("circular"); and a linear donor, which contains two homology arms interrupted by the expression construct ("linear").

[0468] DNA was isolated from transfected cells as described in Example 32. The DNA was analyzed by PCR, using two primer pairs designed to detect the junction between integrated exogenous sequences and endogenous IL-2R γ exon 5 sequences. Thus, for each primer pair, one of the primers was complementary to endogenous exon 5 sequence, and the other primer was complementary to the expression construct (see upper portion of FIG. 51). As shown in the lower portion of FIG. 51, no PCR product was observed in control samples transfected only with donor DNA. In contrast, PCR products of the expected size were observed in cell samples transfected with donor DNA (either linear or circular) and the ZFN-encoding plasmid. Critically, primer sets specific for both ends of the expression construct yielded identical results, consistent with the notion that ZFN-driven targeted integration of the exogenous 7.7 kb sequence has occurred. Nucleotide sequence determination of the amplification products confirmed that homology-dependent integration had occurred.

Example 37

Targeted, Homology-independent Integration of
Exogenous Sequences at an Endogenous
Chromosomal Locus

[0469] A pair of zinc finger/FokI fusion proteins were constructed to bind to two target sites, separated by six nucleotide pairs, in the Chinese hamster dihydrofolate reductase (DHFR) gene and cleave the gene between the two target sites. The nucleotide sequences of the target sites, and the amino acid sequences of the recognition regions of the fusion proteins, are shown in Table 28.

TABLE 28

<u>Zinc Finger Designs for the CHO DHFR gene</u>				
Target sequence	F1	F2	F3	F4
GGAAGGTCTCCG (SEQ ID NO:223)	RSDTLSE (SEQ ID NO:224)	NNRDRTK (SEQ ID NO:225)	RSDHLSA (SEQ ID NO:226)	QSGHLRS (SEQ ID NO:227)
AATGCTCAGGTA (SEQ ID NO:228)	QSGALAR (SEQ ID NO:229)	RSDNLRE (SEQ ID NO:230)	QSSDLRS (SEQ ID NO:231)	TSSNRKT (SEQ ID NO:232)

Note:

The DNA target sequence is shown in the left-most column. The remaining columns show the amino acid sequences (in one-letter code) of residues -1 through +6 of each of the four zinc fingers, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein.

[0470] Chinese hamster ovary (CHO) cells were cultured in adherent medium (DMEM+10% FBS supplemented with 2 mM L-glutamine plus non-essential amino acids) at 37° C. 3×10^5 cells were grown to 70% confluence in 12-well plates and transiently transfected (using Lipofectamine 2000®) with 100 ng each of the two fusion protein-encoding plasmids. 24 hours after transfection, 20 μ M vinblastine was added to the growth medium. Medium was replaced 24 hours after addition of vinblastine. 24 hours after replacement of medium, cellular DNA was purified (Qiagen) and DHFR gene sequences surrounding the target sites were amplified by PCR. Primers were designed such that DNA from cells containing a wild-type DHFR gene were expected to yield a 383 nucleotide pair amplification product. Unexpectedly, two amplification products were obtained, one of the expected size and another approximately 150 nucleotide pairs larger.

[0471] To determine if mutations had been induced at the cleavage site, the amplification product was analyzed using a Cel-1 assay, in which the amplification product is denatured and renatured, followed by treatment with the mismatch-specific Cel-1 nuclease. See, for example, Olejowski et al. (1998) *Nucleic Acids res.* 26:4597-4602; Qui et al. (2004) *BioTechniques* 36:702-707; Yeung et al. (2005) *BioTechniques* 38:749-758. The results of the Cel-1 assay (FIG. 52) showed that, in addition to small mismatches in the reannealed products resulting from non-homologous end-joining at the cleavage site (indicated by the presence of two low molecular weight bands in rightmost lane of FIG. 52), a larger insertion had also occurred (indicated by the presence of a high molecular weight band, identified as "Mutant," in lanes 3 and 5 of FIG. 52). This corroborated the observation of a larger amplification product described above.

[0472] The nucleotide sequences of the two amplification products described above were determined, to characterize the nature of the insertion, and are shown in FIG. 53. The sequence shown on the top line (SEQ ID NO:233) is the wild-type DHFR sequence, while the sequence shown on the bottom line (SEQ ID NO:234) consists of the DHFR sequence containing, at the cleavage site, an insertion of 157 base pairs and a deletion of a single nucleotide pair. Further analysis revealed that the inserted 157 base pairs correspond to a portion of the vector plasmid encoding the zinc finger/FokI fusion proteins. Moreover, when uptake of fluorescent methotrexate was assayed, cells containing this mutation

showed 53% mean methotrexate uptake, compared to wild-type CHO cells, consistent with loss of function of one copy of the DHFR gene.

[0473] Thus, targeted, homology-independent integration of exogenous vector sequences occurred at the site of targeted cleavage in the DHFR gene, resulting in the generation of a heterozygous DHFR⁻ mutant cell line.

Example 38

Multiple Mutations in the FokI Dimerization
Domain for Targeted Cleavage and
Homology-directed Repair of an Endogenous Gene

[0474] Additional sequence alterations were introduced into the mutagenized FokI cleavage half-domain described in Example 5, in which residue 490 was converted from glutamic acid to lysine (E490K), to provide further improvement in its cleavage specificity. In one embodiment (mutant X2), amino acid 538 was converted from isoleucine to lysine (I538K). In further embodiments, amino acid 486 of the X2 mutant was converted from glutamine to glutamic acid (Q486E) to generate the X3A mutant, or from glutamine to isoleucine (Q486I) to generate the X3B mutant. The amino acid sequences of the E490K, X2, X3A and X3B mutants, compared to the amino acid sequence of the wild-type FokI cleavage half-domain, are presented in FIG. 54.

[0475] Plasmids were constructed in which sequences encoding these mutant cleavage half-domains were fused to sequences encoding the 5-8G and 5-9D zinc finger domains (see Example 14). Various combinations of these mutants were then assayed for their ability to stimulate homology-directed repair of a double-stranded break in exon 5 of the IL-2R γ gene, in the presence of a donor DNA sequence containing a BsrBI site. The assay system and procedures described in Example 15 were used, except that cells were not treated with vinblastine and 20 Units of BsrBI was used for digestion.

[0476] After a 48 hour exposure of the gel, the Phosphorimager screen was read and the intensity of the RFLP-derived and wild-type bands were quantified using ImageQuant software (MolecularDynamics). The intensity of the RFLP-derived band, as percentage of the total radioactivity for the wild-type and RFLP bands, is given in Table 29. The results indicates that the X3 FokI mutant functions significantly better when paired with the Q486E mutant than when paired with a second copy of itself.

TABLE 29

Homology-directed alteration of an endogenous gene using zinc finger/FokI fusion proteins containing mutations in the FokI dimerization interface*			
Sample	5-8	5-9	% GC
1	WT (1 ug)	WT (1 ug)	2.6
2	WT (2.5 ug)	WT (2.5 ug)	<1
3	WT (5 ug)	WT (5 ug)	1.5
4	WT (7.5 ug)	WT (7.5 ug)	<1
5	X3 (1 ug)	Q486E (1 ug)	4.1
6	X3 (2.5 ug)	Q486E (2.5 ug)	4.3
7	X3 (5 ug)	Q486E (5 ug)	8.6
8	X3 (7.5 ug)	Q486E (7.5 ug)	3.6
9	X3 (5 ug)	X3 (5 ug)	0
10	Q486E (5 ug)	Q486E (5 ug)	2.3

*K562 cells were transfected with plasmids encoding two zinc finger/FokI fusion proteins and with a plasmid containing a donor DNA sequence homologous to exon 5 of the IL-2R γ gene but containing a sequence change resulting in the presence of a BsrBI site. The second and third columns identify the nature of the FokI cleavage half-domain in the 5-8 and 5-9 zinc finger fusion proteins, as follows: WT (wild-type FokI cleavage half-domain); Q486E mutant cleavage half-domain (containing a single amino acid change compared to wild-type, described in Example 5); X3 mutant cleavage half-domain (containing three amino acid changes compared to wild-type, shown in FIG. 54). “% GC” refers to fraction of total amplification product that is cleaved by BsrBI, as measured by radioactivity in BsrBI digestion products.

Example 39

Zinc Finger/FokI Fusion Proteins with Multiple Mutations in the FokI Dimerization Domain Tested with a Chromosomal GFP Reporter Gene Assay

[0477] The cell line described in Example 29, containing a chromosomally integrated mutant eGFP coding sequence operatively linked to a tetracycline-regulated CMU promoter, was used in experiments to test different combinations of zinc finger/FokI fusion proteins (ZFNs) containing amino acid sequence alterations in the dimerization interface of the FokI cleavage half-domain. See Example 38 and FIG. 54. The exogenous donor DNA construct, previously described in Example 13 and FIG. 32, contained a 1527 nucleotide pair insert homologous to wild-type eGFP coding sequences. It was constructed by amplification of eGFP sequences using the following primers:

(SEQ ID NO:235)
GFPnostart GGCGAGGAGCTGTTCAC
(SEQ ID NO:236)
pcDNA42571 TGCATACTTCTGCCTGC

The amplification product was topo cloned into the pCR4-TOPO vector to generate the donor construct, denoted pCR4-TOPO_GFPDonor_1.5KB. Targeted, homology-directed integration of this donor sequence will result in replacement of the mutant chromosomal eGFP sequences with wild-type eGFP sequences and doxycycline-inducible expression of functional eGFP.

[0478] Cells containing the chromosomally integrated mutant eGFP sequences (described above) were grown in Dulbecco’s modified Eagle’s medium (DMEM) (Invitrogen), supplemented with 10% fetal bovine serum (FBS) (Hyclone) and 2 mM L-glutamine and maintained at 37° C. in an atmosphere of 5% CO₂. Cells were transfected using

LipofectAMINE 2000 Reagent (Invitrogen) in Opti-MEM I reduced serum medium. Cells were transfected with plasmids encoding the ZFNs only (5 ng each), donor plasmid only (500 ng), or plasmids encoding the ZFNs (5 ng each) +donor plasmid (500 ng). Expression of the chromosomal eGFP coding sequences was activated by the addition of 2 ng/ml doxycycline (final concentration) to the growth medium 5 hours post-transfection. The cells were harvested 3 days post-transfection and assayed by flow cytometry for eGFP expression. The results, shown in Table 30, indicate the fusion proteins containing mutations in the dimerization interface function more effectively to promote homology-directed repair in the presence of a different cleavage half-domain, suggesting they are less prone to homodimerization.

TABLE 30

Homology-directed alteration of a mutant chromosomal eGFP gene using zinc finger/FokI fusion proteins containing mutations in the FokI dimerization interface*						
5-8:						
5-9	WT	Q486E	E490K	X2	X3A	X3B
WT	0.66	0.38	0.61	0.40	0.70	0.18
Q486E	0.26	0.14	0.54	0.53	0.50	0.23
E490K	0.58	0.42	0.30	0.01	0.02	0.03
X2	0.14	0.55	0.07	0.01	0.01	0.01
X3A	0.43	0.43	0.02	0.01	0.03	0.01
X3B	0.19	0.33	0.06	0.02	0.03	0.02

*Cells were transfected with a donor construct and two ZFN expression constructs: one expressing a 5-8 zinc finger binding domain fused to a FokI cleavage half-domain and the other expressing a 5-9 zinc finger binding domain fused to a FokI cleavage half-domain. The nature of the cleavage half-domain fused to the 5-8 zinc finger binding domain is given across the top row; the nature of the cleavage half-domain fused to the 5-9 zinc finger binding domain is given down the leftmost column. Numbers indicate percentage of cells exhibiting eGFP fluorescence for each pair of ZFNs tested.

Example 40

Targeted Homology-dependent Integration of a 41-nucleotide Exogenous Sequence into the Endogenous CCR-5 Gene

[0479] Growth and transfection of K562 cells were conducted as described in Example 32. Cells were transfected with 2.5 μ g of a construct encoding two zinc finger/FokI fusion proteins (separated by a 2A peptide sequence) in which the zinc finger domains (7568 and 7296) were designed to bind target sites in the human CCR-5 gene, and 50 μ g of donor construct (see below). The target sites for the zinc finger domains (boxed) are separated by 5 nucleotide pairs, as shown below.

(SEQ ID NO:237)
7568
5' CTGGTCATCCTCATCCTGATAAACTGCAAAAGCT
GACAGTAGGAGTAGGACTATTGACGTTTCCGA 5'
7296

[0480] The nucleotide sequences of the target sites, and the amino acid sequences of the recognition regions of the zinc finger domains, are shown in Table 31.

TABLE 31

<u>Zinc Finger Designs for the human CCR-5 gene</u>				
Target sequence	F1	F2	F3	F4
GATGAGGATGAC (SEQ ID NO:238)	DRSNLSR (SEQ ID NO:239)	TSANLSR (SEQ ID NO:240)	RSDNLSR (SEQ ID NO:241)	TSANLSR (SEQ ID NO:242)
AAACTGCAAAAG (SEQ ID NO:243)	RSDHLSR (SEQ ID NO:244)	QNANRIT (SEQ ID NO:245)	RSDVLSR (SEQ ID NO:246)	QRNHRIT (SEQ ID NO:247)

Note:

The DNA target sequence is shown in the left-most column. The remaining columns show the amino acid sequences (in one-letter code) of residues -1 through +6 of each of the four zinc fingers, with respect to the start of the alpha-helical portion of each zinc finger. Finger F1 is closest to the amino terminus of the protein.

[0481] The donor DNA molecule comprised a ~2 kilobase-pair portion of the human CCR-5 gene engineered to contain a 41 nucleotide pair sequence tag containing a novel diagnostic recognition site for the restriction enzyme BglI. The donor molecule was constructed by mutagenizing the CCR-5 gene fragment to create a XbaI site, and introducing the 41-nucleotide tag into that XbaI site. As a result, the 41-nucleotide tag was flanked by approximately 0.5 kilobase pairs of CCR-5 sequence on one side and approximately 1.5 kilobase pairs of CCR-5 sequence on the other. This sequence is shown below, with the 41-nucleotide pair tag shown in upper case and the BglI site underlined.

(SEQ ID NO:248)

gttgtcaaagcttcattcactccatggtgctatagagcacaagattttat
 ttggtgagatggtgctttcatgaattccccacagagccaagctctcca
 tctagtggacagggagctagcagcaaaccttccctcactacaaaactt
 cattgcttggccaaaagagagtttaattcaatgtagacatctatgtaggc
 aattaaaaacctattgatgtataaaacagtttgcatcattgagggcaac
 taaatacattctaggactttataaaagatcactttttatgtacagag
 gtggaacaagatggattatcaagtgtcaagtccaatctatgacatcaatt
 attatacatcgagccctgccccaaaataatgtgaagcaaatcgagcc
 cgctcctgctcctcgctcactcactggtgttcatctttgtttgtggg
 caacatgctggtcatcctcatctagaTCAGTGAGTATGCCCTGATGGCGT
 CTGGACTGGATGCTCGTctagataaaactgcaaaaggctgaagagcatga
 ctgacatctacctgctcaacctggccatctctgacctgtttttccttctt
 actgtcccttctgtggctcactatgctgcccagtgaggacttttgaaa
 tacaatgtgtcaactcttgacagggctctattttataggcttcttctctg
 gaatcttcttcatcctcctgacaatcgataggtacctggtgctgctc
 catgctgtgtttgcttttaaaagccaggacggtcacctttggggtggtgac
 aagtgtgatcacttgggtggtggtgtgttgcgtctctcccgaatca
 tctttaccagatctcaaaaagaaggtcttctattacacctgcagctctcat
 ttccatacagtcagtatcaattctggaagaatttcagacattaaagat
 agtcatcttggggtggtcctgcccgtgctgtgcatggtcatctgctact

-continued

cggggaatcctaaaaactctgcttcggtgtcgaaatgagaagaagaggcac
 agggctgtgaggcttatcttcaccatcatgattgtttatcttctctctg
 ggctccctacaacattgtccttctcctgaacaccttccaggaattctttg
 gcctgaataaattgcagtagctctaacaggttgaccaagctatgcaggtg
 acagagactcttgggatgacgcactgctgcatcaaccccatcatctatgc
 ctttctcggggagaagttcagaaactacctcttagtcttcttccaaaagc
 acattgcaaaacgcttctgcaaatgctgttctatcttccagcaagaggct
 cccgagcgagcaagctcagtttacacccgatccactggggagcaggaaat
 atctgtgggctgtgtgacacggactcaagtgggctggtgacctcagag
 ttgtgcacatggcttagttttcatacacagcctgggctgggggtggggtg
 ggagaggctcttttttaaaaggaagttactgttatagagggtctaagattc
 atccatttatttggcatctgtttaagtagattagatcttttaagcccat
 caattatagaaaagccaaatcaaatatgttgatgaaaaatagcaaccttt
 ttatctcccttcacatgcatcaagttattgacaaactctcccttctactc
 cgaaagttccttatgtatattttaaagaaagcctcagagaattgctgatt
 cttgagtttagtgatctgaacagaaataccaaaattatttcagaaatgta
 caactttttacctagtagacaaggcaacatataggtgtgaaatgtgtttaa
 acaggtctttgtctgtgctatggggagaaaagacatgaatatgattagtaa
 agaaatgacacttttcatgtgtgatttc

[0482] Six days after transfection, cellular DNA was isolated and used as a template for amplification as described in Example 32, then digested with BglI. In DNA from cells transfected with both the donor construct and the construct encoding two zinc finger/FokI fusion proteins, approximately 1% of the amplification products were cleaved by BglI, indicative of targeted insertion of the sequence tag into the CCR-5 gene. DNA from untransfected cells, cells that were transfected only with the donor construct, and cells that were transfected only with the construct encoding two zinc finger/FokI fusion proteins did not yield amplification products that were cleaved by BglI. It is significant that the targeted insertion of this 41-nucleotide sequence tag generates a frameshift mutation in the CCR-5 gene, thereby inactivating gene function, including its function as a receptor for HIV.

[0483] All patents, patent applications and publications mentioned herein are hereby incorporated by reference, in their entireties, for all purposes.

[0484] Although disclosure has been provided in some detail by way of illustration and example for the purposes of

clarity of understanding, it will be apparent to those skilled in the art that various changes and modifications can be practiced without departing from the spirit or scope of the disclosure. Accordingly, the foregoing descriptions and examples should not be construed as limiting.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 253

<210> SEQ ID NO 1

<211> LENGTH: 44

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(44)

<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 1

ctgccgcgcg cgccgcggcc gtcattgggt tcctgaaact gatt

44

<210> SEQ ID NO 2

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Gly Phe Leu Lys Leu Ile

1

5

<210> SEQ ID NO 3

<211> LENGTH: 47

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(47)

<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 3

ctgccgcgcg cgccgcggcc gtcattgggt tcctgaaact gattgag

47

<210> SEQ ID NO 4

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Met Gly Phe Leu Lys Leu Ile Glu

1

5

<210> SEQ ID NO 5

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Sequence of donor molecule; modified fragment of human SMCI gene

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(50)

<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 5

-continued

ctgccgccgg cgccgcggcc gtcataagaa gcttcctgaa actgattgag 50

<210> SEQ ID NO 6
<211> LENGTH: 463
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Mutated human SMC1 gene

<400> SEQUENCE: 6

tagtcctgca ggtttaaacg aattcgccct tctcagcaag cgtgagctca ggtctcccc 60
gcctccttga acctcaagaa ctgctctgac tccgcccgagc aacaactcct ccgggggatct 120
ggtcgcgagg agcaagtgtt tgttgttgcc atgcaacaag aaaagggggc ggaggcacca 180
cgccagtcgt cagctcgctc ctcgatatac caacatcagt ccccgcccct ggtcccactc 240
ctgccggaag gcgaagatcc cgtaggcctt ggacgtattc tcgcgacatt tgccgggtcg 300
ccggcttgca ctgcggcggt tcccgcgcgg gctacctcag ttctcgggcg tacggcgcg 360
cctgtcctac tgctgcggc gccgcggcg tcataagaag cttcctgaaa ctgattgaag 420
ggcgaattcg cgccgcgtaa attcaattcg ccctatagt agt 463

<210> SEQ ID NO 7
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(50)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 7

cttccaacct ttctcctcta ggtacaagaa ctcgataat gataaagtcc 50

<210> SEQ ID NO 8
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Tyr Lys Asn Ser Asp Asn Asp Lys Val
1 5

<210> SEQ ID NO 9
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

gttcctcttc cttccaacct ttctcctcta ggtacaagaa ctcgataat gataaagtc 59

<210> SEQ ID NO 10
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Tyr Lys Asn Ser Asp Asn Asp Lys Val
1 5

-continued

<210> SEQ ID NO 11
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sequence of donor molecule; modified fragment
of human SMC1 gene

<400> SEQUENCE: 11

gttctctcttc cttccaacct ttctcctcta ggtaaaagaa ttccgacaac gataaagtc 59

<210> SEQ ID NO 12
<211> LENGTH: 624
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Mutated human IL-2Rgamma gene

<400> SEQUENCE: 12

tagtcttgca ggtttaaacg aattcgccct ttcctctagg taaaagaatt ccgacaacga 60
taaaagtcag aagtgcagcc actatctatt ccctgaagaa atcacttctg gctgtcagtt 120
gcaaaaaaag gagatccacc tctaccaaac atttggtggt cagctccagg acccacggga 180
acccaggaga caggccacac agatgctaaa actgcagaat ctgggtaatt tggaaagaaa 240
gggtcaagag accagggata ctgtgggaca ttggagtcta cagagtagtg ttcttttatac 300
ataagggtag atgggcagaa aagaggaggt aggggatcat gatgggaagg gaggaggtat 360
taggggcact accttcagga tcctgacttg tctaggccag gggaatgacc acatatgcac 420
acatatctcc agtgatcccc tgggctccag agaacctaac acttcacaaa ctgagtgaat 480
cccagctaga actgaactgg aacaacagat tcttgaacca ctgtttggag cacttggtgc 540
agtaccggac taaggcgcaa ttcgcggccg ctaaattcaa ttcgccctat agtgagtcgt 600
attacaattc actggccgtc gttt 624

<210> SEQ ID NO 13
<211> LENGTH: 700
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

tactgatggt atggggccaa gagatatatc ttagaggag ggctgagggt ttgaagtcca 60
actcctaagc cagtgccaga agagccaagg acaggtacgg ctgtcatcac ttagacctca 120
ccctgtggag ccacacccta gggttggcca atctactccc aggagcaggg agggcaggag 180
ccagggtcgg gcataaaagt cagggcagag ccatctattg cttacatttg cttctgacac 240
aactgtgttc actagcaacc tcaaacagac accatggtgc atctgactcc tgaggagaag 300
tctgccgtta ctgccctgtg gggcaagggt aacgtggatg aagttggtgg tgaggccctg 360
ggcaggttg tatcaagggt acaagacagg ttaaggaga ccaatagaaa ctgggcatgt 420
ggagacagag aagactcttg ggtttctgat aggcactgac tctctctgcc tattgtcta 480
ttttccacc cttaggctgc tgggtggtcta cccttgacc cagaggttct ttgagtcctt 540
tggggatctg tccactcctg atgctgttat gggcaaccct aaggtgaagg ctcattggcaa 600
gaaagtgtc ggtgccttta gtgatggcct ggctcacctg gacaacctca agggcacctt 660
tgccacactg agtgagctgc actgtgacaa gctgcacgtg 700

-continued

```
<210> SEQ ID NO 14
<211> LENGTH: 408
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Mutated human beta-globin gene

<400> SEQUENCE: 14

tgcttaccaa gctgtgattc caaatattac gtaaatacac ttgcaaagga ggatgttttt      60
agtagcaatt tgtactgatg gtatggggcc aagagatata tcttagaggg agggctgagg      120
gtttgaagtc caactcctaa gccagtgcca gaagagccaa ggacaggtac ggctgtcatt      180
acttagacct caccctgtgg agccacaccc tagggttggc caatctactc ccaggagcag      240
ggagggcagg agccaggggt gggcataaaa gtcagggcag agccatctat tgcttacatt      300
tgcttctgac acaactgtgt tcactagcaa cctcaaacag acaccatggt gcactctgact      360
cctgaggaga agtctggcgt tagtgcccga attccgatcg tcaaccac      408
```

```
<210> SEQ ID NO 15
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Portion of 5th exon of human IL-2Rgamma gene
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(42)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 15
```

cacgtttcgt gttcggagcc gctttaaccc actctgtgga ag 42

```
<210> SEQ ID NO 16
<211> LENGTH: 336
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sequence of the 5-8 ZFP/FokI fusion targeted
to exon 5 of the human IL-2Rgamma gene
```

<400> SEQUENCE: 16

Met Ala Pro Lys Lys Lys Arg Lys Val Gly Ile His Gly Val Pro Ala
1 5 10 15

Ala Met Ala Glu Arg Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe
20 25 30

Ser Arg Ser Asp Asn Leu Ser Glu His Ile Arg Thr His Thr Gly Glu
35 40 45

Lys Pro Phe Ala Cys Asp Ile Cys Gly Arg Lys Phe Ala Arg Asn Ala
50 55 60

His Arg Ile Asn His Thr Lys Ile His Thr Gly Ser Gln Lys Pro Phe
65 70 75 80

Gln Cys Arg Ile Cys Met Arg Asn Phe Ser Arg Ser Asp Thr Leu Ser
85 90 95

Glu His Ile Arg Thr His Thr Gly Glu Lys Pro Phe Ala Cys Asp Ile
100 105 110

Cys Gly Arg Lys Phe Ala Ala Arg Ser Thr Arg Thr Thr His Thr Lys
115 120 125

Ile His Leu Arg Gln Lys Asp Ala Ala Arg Gly Ser Gln Leu Val Lys

-continued

130	135	140
Ser Glu Leu Glu Glu Lys Lys Ser Glu Leu Arg His Lys Leu Lys Tyr	150	155
145		160
Val Pro His Glu Tyr Ile Glu Leu Ile Glu Ile Ala Arg Asn Ser Thr	165	170
		175
Gln Asp Arg Ile Leu Glu Met Lys Val Met Glu Phe Phe Met Lys Val	180	185
		190
Tyr Gly Tyr Arg Gly Lys His Leu Gly Gly Ser Arg Lys Pro Asp Gly	195	200
		205
Ala Ile Tyr Thr Val Gly Ser Pro Ile Asp Tyr Gly Val Ile Val Asp	210	215
		220
Thr Lys Ala Tyr Ser Gly Gly Tyr Asn Leu Pro Ile Gly Gln Ala Asp	225	230
		235
Glu Met Gln Arg Tyr Val Glu Glu Asn Gln Thr Arg Asn Lys His Ile	245	250
		255
Asn Pro Asn Glu Trp Trp Lys Val Tyr Pro Ser Ser Val Thr Glu Phe	260	265
		270
Lys Phe Leu Phe Val Ser Gly His Phe Lys Gly Asn Tyr Lys Ala Gln	275	280
		285
Leu Thr Arg Leu Asn His Ile Thr Asn Cys Asn Gly Ala Val Leu Ser	290	295
		300
Val Glu Glu Leu Leu Ile Gly Gly Glu Met Ile Lys Ala Gly Thr Leu	305	310
		315
Thr Leu Glu Glu Val Arg Arg Lys Phe Asn Asn Gly Glu Ile Asn Phe	325	330
		335

<210> SEQ ID NO 17

<211> LENGTH: 339

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Sequence of the 5-10 ZFP/FokI fusion targeted to exon 5 of the human IL-2Rgamma gene

<400> SEQUENCE: 17

Met Ala Pro Lys Lys Lys Arg Lys Val Gly Ile His Gly Val Pro Ala	1	5	10	15
Ala Met Ala Glu Arg Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe	20	25	30	
Ser Arg Ser Asp Ser Leu Ser Arg His Ile Arg Thr His Thr Gly Glu	35	40	45	
Lys Pro Phe Ala Cys Asp Ile Cys Gly Arg Lys Phe Ala Asp Ser Ser	50	55	60	
Asn Arg Lys Thr His Thr Lys Ile His Thr Gly Gly Gly Ser Gln	65	70	75	80
Lys Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser Arg Ser Asp	85	90	95	
Ser Leu Ser Val His Ile Arg Thr His Thr Gly Glu Lys Pro Phe Ala	100	105	110	
Cys Asp Ile Cys Gly Arg Lys Phe Ala Asp Arg Ser Asn Arg Ile Thr	115	120	125	
His Thr Lys Ile His Leu Arg Gln Lys Asp Ala Ala Arg Gly Ser Gln	130	135	140	

-continued

Leu	Val	Lys	Ser	Glu	Leu	Glu	Glu	Lys	Lys	Ser	Glu	Leu	Arg	His	Lys
145				150						155					160
Leu	Lys	Tyr	Val	Pro	His	Glu	Tyr	Ile	Glu	Leu	Ile	Glu	Ile	Ala	Arg
			165						170					175	
Asn	Ser	Thr	Gln	Asp	Arg	Ile	Leu	Glu	Met	Lys	Val	Met	Glu	Phe	Phe
			180					185					190		
Met	Lys	Val	Tyr	Gly	Tyr	Arg	Gly	Lys	His	Leu	Gly	Gly	Ser	Arg	Lys
	195						200					205			
Pro	Asp	Gly	Ala	Ile	Tyr	Thr	Val	Gly	Ser	Pro	Ile	Asp	Tyr	Gly	Val
	210					215					220				
Ile	Val	Asp	Thr	Lys	Ala	Tyr	Ser	Gly	Gly	Tyr	Asn	Leu	Pro	Ile	Gly
225				230						235					240
Gln	Ala	Asp	Glu	Met	Gln	Arg	Tyr	Val	Glu	Glu	Asn	Gln	Thr	Arg	Asn
			245						250					255	
Lys	His	Ile	Asn	Pro	Asn	Glu	Trp	Trp	Lys	Val	Tyr	Pro	Ser	Ser	Val
			260				265						270		
Thr	Glu	Phe	Lys	Phe	Leu	Phe	Val	Ser	Gly	His	Phe	Lys	Gly	Asn	Tyr
	275						280					285			
Lys	Ala	Gln	Leu	Thr	Arg	Leu	Asn	His	Ile	Thr	Asn	Cys	Asn	Gly	Ala
	290					295					300				
Val	Leu	Ser	Val	Glu	Glu	Leu	Leu	Ile	Gly	Gly	Glu	Met	Ile	Lys	Ala
305				310					315						320
Gly	Thr	Leu	Thr	Leu	Glu	Glu	Val	Arg	Arg	Lys	Phe	Asn	Asn	Gly	Glu
			325					330						335	

Ile Asn Phe

<210> SEQ ID NO 18

<211> LENGTH: 797

<212> TYPE: DNA

<213> ORGANISM: Aequorea victoria

<400> SEQUENCE: 18

cgaattctgc agtcgacggt accgcggggc cgggatccac cggtcgccac catggtgagc	60
aaggcgagg agctgttcac cggggtggtg cccatcctgg tcgagctgga cggcgacgta	120
aacggccaca agttcagcgt gtccggcgag ggcgaggcg atgccaccta cggcaagctg	180
accctgaagt tcatctgcac caccggcaag ctgcccgtgc cctggccac cctcgtgacc	240
accctgacct acggcgtgca gtgcttcagc cgctaccccg accacatgaa gcagcacgac	300
ttcttcaagt ccgcatgcc cgaaggctac gtccaggagc gcaccatctt cttcaaggac	360
gacggcaact acaagaccg cgccgaggtg aagttcgagg gcgacacct ggtgaaccgc	420
atcgagctga agggcatcga cttcaaggag gacggcaaca tcctggggca caagctggag	480
tacaactaca acagccacaa cgtctatatc atggccgaca agcagaagaa cggcatcaag	540
gtgaacttca agatccgcc caacatcgag gacggcagcg tgcagctcgc cgaccactac	600
cagcagaaca ccccatcgg cgacggcccc gtgctgctgc ccgacaacca ctacctgagc	660
accagtcgg cctgagcaa agaccccaac gagaagcgcg atcacatggt cctgctggag	720
ttcgtgaccg ccgccgggat cactctcggc atggacgagc tgtacaagta aagcggccgc	780
gactctagat cataatc	797

<210> SEQ ID NO 19

-continued

```

<211> LENGTH: 795
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Mutant defective synthetic eGFP gene

<400> SEQUENCE: 19
cgaattctgc agtcgacggt accgcgggcc cgggatccac cggtcgccac catggtgagc    60
aagggcgagg agctgttcac cggggtggtg cccatcctgg tcgagctgga cggcgacgta    120
aacggccaca agttcagcgt gtccggcgag ggcgaggcg atgccaccta cggcaagctg    180
accctgaagt tcatctgcac caccggcaag ctgcccgtgc cctggcccac cctcgtgacc    240
accctgacct acggcgctga gtgcttcagc cgctaccctt aacacgaagc agcacgactt    300
cttcaagtcc gccatgcccc aaggctacgt ccaggagcgc accatcttct tcaaggacga    360
cggcaactac aagaccgcgc ccgaggtgaa gtctgagggc gacaccctgg tgaaccgcat    420
cgagctgaag ggcatcgact tcaaggagga cggcaacatc ctggggcaca agctggagta    480
caactacaac agccacaacg tctatatcat ggccgacaag cagaagaacg gcatcaaggt    540
gaacttcaag atccgccaca acatcgagga cggcagcgtg cagctcgccg accactacca    600
gcagaacacc cccatcggcg acggccccgt gctgctgccc gacaaccact acctgagcac    660
ccagtccgcc ctgagcaaa accccaacga gaagcgcgat cacatggtcc tgctggagtt    720
cgtgaccgcc gccgggatca ctctcgcat ggacgagctg tacaagtaa gcggccgcga    780
ctctagatca taatc                                         795

```

```

<210> SEQ ID NO 20
<211> LENGTH: 734
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic eGFP insert in pCR(R)4-TOPO-GFPdonor5

```

```

<400> SEQUENCE: 20
ggcgaggagc tgttcaccgg ggtggtgccc atcctggtcg agctggacgg cgacgtaaac    60
ggccacaagt tcagcgtgtc cggcgagggc gagggcgatg ccacctacgg caagctgacc    120
ctgaagtta tctgcaccac cggcaagctg cccgtgccct ggcccacct cgtgaccacc    180
ctgacctacg gcgtgcatgt cttcagccgc taccctgacc acatgaagca gcacgacttc    240
ttcaagtccg ccatgcccca aggtacgtc caggagcgca ccattcttct caaggacgac    300
ggcaactaca agaccgcgc cgaggtgaag ttcgaggcg acaccctggt gaaccgcac    360
gagctgaagg gcatcgactt caaggaggac ggcaacatcc tggggcaca gctggagtac    420
aactacaaca gccacaacgt ctatatcatg gccgacaagc agaagaacgg catcaaggtg    480
aacttcaaga tccgccaca catcgaggac ggacgcgtgc agctcgccga ccactaccag    540
cagaacaccc ccacgcgca cggccccgtg ctgctgccc acaaccacta cctgagcacc    600
cagtcgcccc tgagcaaaga ccccaacgag aagcgcgatc acatggtcct gctggagttc    660
gtgaccgccg ccgggatcac tctcgcatg gacgagctgt acaagtaaag cggccgcgac    720
tctagatcat aatc                                         734

```

```

<210> SEQ ID NO 21
<211> LENGTH: 1527
<212> TYPE: DNA

```

-continued

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic eGFP insert in pCR(R)4-TOPO

<400> SEQUENCE: 21

```

ggcgaggagc tgttcaccgg ggtggtgccc atcctggtcg agctggacgg cgacgtaaac    60
ggccacaagt tcagcgtgtc cggcgagggc gagggcgatg ccacctacgg caagctgacc    120
ctgaagttca tctgcaccac cggcaagctg cccgtgccct ggcccaccct cgtgaccacc    180
ctgacctacg gcgtgcagtg cttcagccgc taccgccgacc acatgaagca gcacgacttc    240
ttcaagtccg ccattgcccga aggctacgtc caggagcgca ccattcttctt caaggacgac    300
ggcaactaca agaccgcgcg cgaggtgaag ttcgagggcg acaccctggt gaaccgcac    360
gagctgaagg gcatcgactt caaggaggac ggcaacatcc tggggcacia gctggagtac    420
aactacaaca gccacaacgt ctatatcatg gccgacaagc agaagaacgg catcaagggtg    480
aacttcaaga tccgccacaa catcgaggac ggagcgctgc agctcgccga ccactaccag    540
cagaacaccc ccattcgccga cggccccgtg ctgctgcccg acaaccacta cctgagcacc    600
cagtccgccc tgagcaaaga ccccaacgag aagcgcgac acatggctct gctggagttc    660
gtgaccgccc ccgggatcac tctcgcatg gacgagctgt acaagtaaag cggccgctcg    720
agtctagagg gcccgtttaa acccgctgat cagcctcgac tgtgccttct agttgccagc    780
catctgttgt ttgccctcc cccgtgcctt ccttgacctt ggaagggtgc actccactg    840
tcctttccta ataaaatgag gaaattgcat cgcattgtct gactaggtgt cattctattc    900
tgggggggtg ggtggggcag gacagcaagg gggaggattg ggaagacaat agcaggcatg    960
ctgggggatgc ggtgggctct atggcttctg aggcggaaag aaccagctgg ggctctaggg   1020
ggtatcccca cgcgcctgt agcggcgcat taagcgcggc ggggtgtggtg gttacgcgca   1080
gcgtgaccgc tacacttgcc agcgcctag cgcgcgctcc ttctgcttcc tcccttccct   1140
ttctcgccac gttcgccggc ttccccgc cagctctaaa tcgggggctc cctttagggt   1200
tccgatttag tgctttacgg cacctcgacc ccaaaaaact tgattagggt gatggttcac   1260
gtagtgggcc atcgccctga tagacggtt ttccgccctt gacgttgagg tccacgttct   1320
ttaatagtgg actcttgttc caaactggaa caacactcaa ccctatctcg gtctattctt   1380
ttgatttata agggattttg ccgatttcgg cctattggtt aaaaaatgag ctgatttaac   1440
aaaaatttaa cgcgaattaa ttctgtggaa tgtgtgtcag ttaggtgtg gaaagtcccc   1500
aggctcccca gcaggcagaa gtatgca                                     1527

```

<210> SEQ ID NO 22

<211> LENGTH: 116

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger domain targeted to the human B-globin gene

<400> SEQUENCE: 22

```

Met Ala Glu Arg Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser
1           5           10          15

```

```

Gln Ser Gly Asp Leu Thr Arg His Ile Arg Thr His Thr Gly Glu Lys
20          25          30

```

```

Pro Phe Ala Cys Asp Ile Cys Gly Arg Lys Phe Ala Thr Ser Ala Asn

```

-continued

```

      35              40              45
Leu Ser Arg His Thr Lys Ile His Thr Gly Gly Gly Gly Ser Gln Lys
  50              55              60

Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser Asp Arg Ser Ala
  65              70              75              80

Leu Ser Arg His Ile Arg Thr His Thr Gly Glu Lys Pro Phe Ala Cys
      85              90              95

Asp Ile Cys Gly Arg Lys Phe Ala Gln Ser Gly His Leu Ser Arg His
  100              105              110

Thr Lys Ile His
  115

<210> SEQ ID NO 23
<211> LENGTH: 113
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger domain targeted to the
      human B-globin gene

<400> SEQUENCE: 23

Met Ala Glu Arg Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser
  1              5              10              15

Arg Ser Gln Thr Arg Lys Thr His Ile Arg Thr His Thr Gly Glu Lys
      20              25              30

Pro Phe Ala Cys Asp Ile Cys Gly Arg Lys Phe Ala Gln Lys Arg Asn
  35              40              45

Arg Thr Lys His Thr Lys Ile His Thr Gly Ser Gln Lys Pro Phe Gln
  50              55              60

Cys Arg Ile Cys Met Arg Asn Phe Ser Asp Arg Ser Ala Leu Ser Arg
  65              70              75              80

His Ile Arg Thr His Thr Gly Glu Lys Pro Phe Ala Cys Asp Ile Cys
      85              90              95

Gly Arg Lys Phe Ala Gln Ser Gly Asn Leu Ala Arg His Thr Lys Ile
  100              105              110

His

<210> SEQ ID NO 24
<211> LENGTH: 116
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger domain targeted to the
      human B-globin gene

<400> SEQUENCE: 24

Met Ala Glu Arg Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser
  1              5              10              15

Thr Ser Gly Ser Leu Ser Arg His Ile Arg Thr His Thr Gly Glu Lys
      20              25              30

Pro Phe Ala Cys Asp Ile Cys Gly Arg Lys Phe Ala Asp Arg Ser Asp
  35              40              45

Leu Ser Arg His Thr Lys Ile His Thr Gly Gly Gly Gly Ser Gln Lys
  50              55              60

Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser Asp Arg Ser Ala
  65              70              75              80

```

-continued

Leu Ser Arg His Ile Arg Thr His Thr Gly Glu Lys Pro Phe Ala Cys
85 90 95
Asp Ile Cys Gly Arg Lys Phe Ala Gln Ser Gly Asn Leu Ala Arg His
100 105 110
Thr Lys Ile His
115

<210> SEQ ID NO 25
<211> LENGTH: 116
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger domain targeted to the
human B-globin gene

<400> SEQUENCE: 25

Met Ala Glu Arg Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser
1 5 10 15
Thr Ser Ser Ser Leu Ser Arg His Ile Arg Thr His Thr Gly Glu Lys
20 25 30
Pro Phe Ala Cys Asp Ile Cys Gly Arg Lys Phe Ala Asp Arg Ser Asp
35 40 45
Leu Ser Arg His Thr Lys Ile His Thr Gly Gly Gly Gly Ser Gln Lys
50 55 60
Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser Asp Arg Ser Ala
65 70 75 80
Leu Ser Arg His Ile Arg Thr His Thr Gly Glu Lys Pro Phe Ala Cys
85 90 95
Asp Ile Cys Gly Arg Lys Phe Ala Gln Ser Gly Asn Leu Ala Arg His
100 105 110
Thr Lys Ile His
115

<210> SEQ ID NO 26
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Type of synthetic fusion ZFP-FokI nuclease

<400> SEQUENCE: 26

His Gln Arg Thr His Gln Asn Lys Lys Gln Leu Val
1 5 10

<210> SEQ ID NO 27
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human SMC1L1 gene for zinc
finger design

<400> SEQUENCE: 27

catgggggttc ct

12

<210> SEQ ID NO 28
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
SMC1L1 gene

<400> SEQUENCE: 28

Arg Ser His Asp Leu Ile Glu
1 5

<210> SEQ ID NO 29
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
SMC1L1 gene

<400> SEQUENCE: 29

Thr Ser Ser Ser Leu Ser Arg
1 5

<210> SEQ ID NO 30
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
SMC1L1 gene

<400> SEQUENCE: 30

Arg Ser Asp His Leu Ser Thr
1 5

<210> SEQ ID NO 31
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
SMC1L1 gene

<400> SEQUENCE: 31

Thr Asn Ser Asn Arg Ile Thr
1 5

<210> SEQ ID NO 32
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human SMC1L1 gene for zinc
finger design

<400> SEQUENCE: 32

gcggcgccgg cg

12

<210> SEQ ID NO 33
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
SMC1L1 gene

<400> SEQUENCE: 33

Arg Ser Asp Asp Leu Ser Arg

-continued

1 5

<210> SEQ ID NO 34
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
SMC1L1 gene

<400> SEQUENCE: 34

Arg Ser Asp Asp Arg Lys Thr
1 5

<210> SEQ ID NO 35
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
SMC1L1 gene

<400> SEQUENCE: 35

Arg Ser Glu Asp Leu Ile Arg
1 5

<210> SEQ ID NO 36
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
SMC1L1 gene

<400> SEQUENCE: 36

Arg Ser Asp Thr Leu Ser Arg
1 5

<210> SEQ ID NO 37
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the human SMC1L1 Gene

<400> SEQUENCE: 37

agcaacaact cctccgggga tc 22

<210> SEQ ID NO 38
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the human SMC1L1 Gene

<400> SEQUENCE: 38

ttccagacgc gactctttgg c 21

<210> SEQ ID NO 39
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the human SMC1L1 Gene

-continued

<400> SEQUENCE: 39

ctcagcaagc gtgagctcag gtctc

25

<210> SEQ ID NO 40

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Amplification primer for the human SMC1L1 Gene

<400> SEQUENCE: 40

caatcagttt caggaagctt ctt

23

<210> SEQ ID NO 41

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Amplification primer for the human SMC1L1 Gene

<400> SEQUENCE: 41

ctcagcaagc gtgagctcag gtctc

25

<210> SEQ ID NO 42

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Amplification primer for the human SMC1L1 Gene

<400> SEQUENCE: 42

ggggtcaagt aaggctggga agc

23

<210> SEQ ID NO 43

<211> LENGTH: 12

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma gene for zinc finger design

<400> SEQUENCE: 43

aactcggata at

12

<210> SEQ ID NO 44

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human IL-2Rgamma gene

<400> SEQUENCE: 44

Asp Arg Ser Thr Leu Ile Glu

1

5

<210> SEQ ID NO 45

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human IL-2Rgamma gene

-continued

<400> SEQUENCE: 45

Ser Ser Ser Asn Leu Ser Arg
1 5

<210> SEQ ID NO 46

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 46

Arg Ser Asp Asp Leu Ser Lys
1 5

<210> SEQ ID NO 47

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 47

Asp Asn Ser Asn Arg Ile Lys
1 5

<210> SEQ ID NO 48

<211> LENGTH: 13

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma gene for
zinc finger design

<400> SEQUENCE: 48

tagaggagaa agg

13

<210> SEQ ID NO 49

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 49

Arg Ser Asp Asn Leu Ser Asn
1 5

<210> SEQ ID NO 50

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 50

Thr Ser Ser Ser Arg Ile Asn
1 5

<210> SEQ ID NO 51

-continued

<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 51

Arg Ser Asp His Leu Ser Gln
1 5

<210> SEQ ID NO 52
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 52

Arg Asn Ala Asp Arg Lys Thr
1 5

<210> SEQ ID NO 53
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma gene for
zinc finger design

<400> SEQUENCE: 53

tacaagaact cg

12

<210> SEQ ID NO 54
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 54

Arg Ser Asp Asp Leu Ser Lys
1 5

<210> SEQ ID NO 55
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 55

Asp Asn Ser Asn Arg Ile Lys
1 5

<210> SEQ ID NO 56
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

-continued

<400> SEQUENCE: 56

Arg Ser Asp Ala Leu Ser Val
1 5

<210> SEQ ID NO 57

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 57

Asp Asn Ala Asn Arg Thr Lys
1 5

<210> SEQ ID NO 58

<211> LENGTH: 9

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma gene for
zinc finger design

<400> SEQUENCE: 58

ggagaaaagg

9

<210> SEQ ID NO 59

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 59

Arg Ser Asp His Leu Thr Gln
1 5

<210> SEQ ID NO 60

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 60

Gln Ser Gly Asn Leu Ala Arg
1 5

<210> SEQ ID NO 61

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma gene

<400> SEQUENCE: 61

Arg Ser Asp His Leu Ser Arg
1 5

<210> SEQ ID NO 62

-continued

<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence used to obtain donor DNA molecule

<400> SEQUENCE: 62
tacaagaact cggataat 18

<210> SEQ ID NO 63
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence used to obtain donor DNA molecule

<400> SEQUENCE: 63
taaaagaatt ccgacaac 18

<210> SEQ ID NO 64
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primers for the human IL-2Rgamma Gene

<400> SEQUENCE: 64
tgtcgagtac atgaattgca cttgg 25

<210> SEQ ID NO 65
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primers for the human IL-2Rgamma Gene

<400> SEQUENCE: 65
ttaggttctc tggagcccag gg 22

<210> SEQ ID NO 66
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primers for the human IL-2Rgamma Gene

<400> SEQUENCE: 66
ctccaaacag tggttcaaga atctg 25

<210> SEQ ID NO 67
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primers for the human IL-2Rgamma Gene

<400> SEQUENCE: 67
tcctctaggt aaagaattcc gacaac 26

-continued

<210> SEQ ID NO 68
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human beta-globin gene for
zinc finger design

<400> SEQUENCE: 68

gggcagtaac gg

12

<210> SEQ ID NO 69
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger designs for the human
beta-globin gene

<400> SEQUENCE: 69

Arg Ser Asp His Leu Ser Glu
1 5

<210> SEQ ID NO 70
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger designs for the human
beta-globin gene

<400> SEQUENCE: 70

Gln Ser Ala Asn Arg Thr Lys
1 5

<210> SEQ ID NO 71
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger designs for the human
beta-globin gene

<400> SEQUENCE: 71

Arg Ser Asp Asn Leu Ser Ala
1 5

<210> SEQ ID NO 72
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger designs for the human
beta-globin gene

<400> SEQUENCE: 72

Arg Ser Gln Asn Arg Thr Arg
1 5

<210> SEQ ID NO 73
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human beta-globin gene for

-continued

zinc finger design

<400> SEQUENCE: 73

aaggtgaacg tg

12

<210> SEQ ID NO 74

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger designs for the human
beta-globin gene

<400> SEQUENCE: 74

Arg Ser Asp Ser Leu Ser Arg
1 5

<210> SEQ ID NO 75

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger designs for the human
beta-globin gene

<400> SEQUENCE: 75

Asp Ser Ser Asn Arg Lys Thr
1 5

<210> SEQ ID NO 76

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger designs for the human
beta-globin gene

<400> SEQUENCE: 76

Arg Ser Asp Ser Leu Ser Ala
1 5

<210> SEQ ID NO 77

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger designs for the human
beta-globin gene

<400> SEQUENCE: 77

Arg Asn Asp Asn Arg Lys Thr
1 5

<210> SEQ ID NO 78

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence used in the process to
obtain a donor DNA molecule

<400> SEQUENCE: 78

ccgttactgc cctgtggggc aaggtgaacg tg

32

-continued

<210> SEQ ID NO 79
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence used in the process to
obtain a donor DNA molecule

<400> SEQUENCE: 79

gcgttagtgc ccgaattccg atcgtcaacc ac 32

<210> SEQ ID NO 80
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the human beta
globin gene

<400> SEQUENCE: 80

tactgatggt atggggccaa gag 23

<210> SEQ ID NO 81
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the human beta
globin gene

<400> SEQUENCE: 81

cacgtgcagc ttgtcacagt gc 22

<210> SEQ ID NO 82
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the human beta
globin gene

<400> SEQUENCE: 82

tgcttaccaa gctgtgattc ca 22

<210> SEQ ID NO 83
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the human beta
globin gene

<400> SEQUENCE: 83

ggttgacgat cggaattc 18

<210> SEQ ID NO 84
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic target site for ZFP

<400> SEQUENCE: 84

aactcggata at 12

-continued

<210> SEQ ID NO 85
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger F1

<400> SEQUENCE: 85

Asp Arg Ser Thr Leu Ile Glu
1 5

<210> SEQ ID NO 86
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger F2

<400> SEQUENCE: 86

Ser Ser Ser Asn Leu Ser Arg
1 5

<210> SEQ ID NO 87
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger F3

<400> SEQUENCE: 87

Arg Ser Asp Asp Leu Ser Lys
1 5

<210> SEQ ID NO 88
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger F4

<400> SEQUENCE: 88

Asp Asn Ser Asn Arg Ile Lys
1 5

<210> SEQ ID NO 89
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic fusion construct in the region of
the ZFP-FokI junction

<400> SEQUENCE: 89

His Thr Lys Ile His Leu Arg Gln Lys Asp Ala Ala Arg Gly Ser Gln
1 5 10 15

Leu Val

<210> SEQ ID NO 90
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic fusion construct in the region of
the ZFP-FokI junction

-continued

<400> SEQUENCE: 90

His Thr Lys Ile His Leu Arg Gln Lys Gly Ser Gln Leu Val
1 5 10

<210> SEQ ID NO 91

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic fusion construct in the region of the ZFP-FokI junction

<400> SEQUENCE: 91

His Thr Lys Ile His Leu Arg Gln Gly Ser Gln Leu Val
1 5 10

<210> SEQ ID NO 92

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic fusion construct in the region of the ZFP-FokI junction

<400> SEQUENCE: 92

His Thr Lys Ile His Leu Arg Gly Ser Gln Leu Val
1 5 10

<210> SEQ ID NO 93

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic fusion construct in the region of the ZFP-FokI junction

<400> SEQUENCE: 93

His Thr Lys Ile His Leu Gly Ser Gln Leu Val
1 5 10

<210> SEQ ID NO 94

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic fusion construct in the region of the ZFP-FokI junction

<400> SEQUENCE: 94

His Thr Lys Ile His Gly Ser Gln Leu Val
1 5 10

<210> SEQ ID NO 95

<211> LENGTH: 38

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP target site

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(38)

<223> OTHER INFORMATION: STRANDEDNESS: Double

<400> SEQUENCE: 95

-continued

ctagcattat ccgagttaca caactcggat aatgctag 38

<210> SEQ ID NO 96
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(39)
<223> OTHER INFORMATION: STRANDEDNESS: Double

<400> SEQUENCE: 96

ctagcattat ccgagttcac acaactcgga taatgctag 39

<210> SEQ ID NO 97
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(42)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 97

ctaggcatta tccgagttca ccacaactcg gataatgact ag 42

<210> SEQ ID NO 98
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(41)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 98

ctagcattat ccgagttcac acacaactcg gataatgcta g 41

<210> SEQ ID NO 99
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(42)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 99

ctagcattat ccgagttcac cacacaactc ggataatgct ag 42

<210> SEQ ID NO 100
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(43)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 100

ctagcattat ccgagttcac acacacaact cggataatgc tag 43

<210> SEQ ID NO 101
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(46)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 101

ctagcattat ccgagttcac caccaacaca actcggataa tgctag 46

<210> SEQ ID NO 102
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(49)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 102

ctagcattat ccgagttcac caccaaccac acaactcggg taatgctag 49

<210> SEQ ID NO 103
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(50)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 103

ctagcattat ccgagttcac caccaaccac accaactcggg ataatgctag 50

<210> SEQ ID NO 104
<211> LENGTH: 51
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(51)
<223> OTHER INFORMATION: STRANDEDNESS: double

-continued

<400> SEQUENCE: 104

ctagcattat ccgagttcaa ccaccaacca caccaactcg gataatgcta g 51

<210> SEQ ID NO 105

<211> LENGTH: 56

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(56)

<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 105

ctagcattat ccgagttcaa ccaccaacca caccaacaca actcggataa tgctag 56

<210> SEQ ID NO 106

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic cleavage substrate with the ZFP
target site

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(60)

<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 106

ctagcattat ccgagttcaa ccaccaacca caccaacacc accaactcgg ataatgctag 60

<210> SEQ ID NO 107

<211> LENGTH: 4

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic linker for ZFP-FokI fusion

<400> SEQUENCE: 107

Leu Arg Gly Ser

1

<210> SEQ ID NO 108

<211> LENGTH: 4

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic linker for ZFP-FokI fusion

<400> SEQUENCE: 108

Leu Gly Gly Ser

1

<210> SEQ ID NO 109

<211> LENGTH: 4

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic linker for ZFP-FokI fusion

<400> SEQUENCE: 109

Thr Gly Gly Ser

1

-continued

<210> SEQ ID NO 110
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic linker for ZFP-FokI fusion

<400> SEQUENCE: 110

Gly Gly Gly Ser
1

<210> SEQ ID NO 111
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic linker for ZFP-FokI fusion

<400> SEQUENCE: 111

Leu Pro Gly Ser
1

<210> SEQ ID NO 112
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic linker for ZFP-FokI fusion

<400> SEQUENCE: 112

Leu Arg Lys Ser
1

<210> SEQ ID NO 113
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic linker for ZFP-FokI fusion

<400> SEQUENCE: 113

Leu Arg Trp Ser
1

<210> SEQ ID NO 114
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma gene for zinc finger design

<400> SEQUENCE: 114

actctgtgga ag

12

<210> SEQ ID NO 115
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human IL-2Rgamma Gene

<400> SEQUENCE: 115

-continued

Arg Ser Asp Asn Leu Ser Glu
1 5

<210> SEQ ID NO 116
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma Gene

<400> SEQUENCE: 116

Arg Asn Ala His Arg Ile Asn
1 5

<210> SEQ ID NO 117
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma Gene

<400> SEQUENCE: 117

Arg Ser Asp Thr Leu Ser Glu
1 5

<210> SEQ ID NO 118
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma Gene

<400> SEQUENCE: 118

Ala Arg Ser Thr Arg Thr Thr
1 5

<210> SEQ ID NO 119
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma gene for
zinc finger design

<400> SEQUENCE: 119

aacacgaaac gtg

13

<210> SEQ ID NO 120
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma Gene

<400> SEQUENCE: 120

Arg Ser Asp Ser Leu Ser Arg
1 5

<210> SEQ ID NO 121
<211> LENGTH: 7

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma Gene

<400> SEQUENCE: 121

Asp Ser Ser Asn Arg Lys Thr
1 5

<210> SEQ ID NO 122
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma Gene

<400> SEQUENCE: 122

Arg Ser Asp Ser Leu Ser Val
1 5

<210> SEQ ID NO 123
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
IL-2Rgamma Gene

<400> SEQUENCE: 123

Asp Arg Ser Asn Arg Ile Thr
1 5

<210> SEQ ID NO 124
<211> LENGTH: 8
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sequence replaced in the synthetic eGFP gene

<400> SEQUENCE: 124

gaccacat 8

<210> SEQ ID NO 125
<211> LENGTH: 6
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Replacement sequence in the modified synthetic
eGFP gene

<400> SEQUENCE: 125

taacac 6

<210> SEQ ID NO 126
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide sequences for GFP

<400> SEQUENCE: 126

cgaattctgc agtcgac 17

-continued

<210> SEQ ID NO 127
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide sequences for GFP

<400> SEQUENCE: 127

gattatgatac tagagtcg 18

<210> SEQ ID NO 128
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide sequences for GFP

<400> SEQUENCE: 128

agccgctacc cctaacacga agcag 25

<210> SEQ ID NO 129
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide sequences for GFP

<400> SEQUENCE: 129

ctgcttcgtg ttaggggtag cggct 25

<210> SEQ ID NO 130
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger protein design for eGFP
gene

<400> SEQUENCE: 130

Arg Ser Asp Asp Leu Thr Arg
1 5

<210> SEQ ID NO 131
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger protein design for eGFP
gene

<400> SEQUENCE: 131

Gln Ser Gly Ala Leu Ala Arg
1 5

<210> SEQ ID NO 132
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger protein design for eGFP
gene

<400> SEQUENCE: 132

Arg Ser Asp His Leu Ser Arg

-continued

1 5

<210> SEQ ID NO 133
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger protein design for eGFP
gene

<400> SEQUENCE: 133

Gln Ser Gly Ser Leu Thr Arg
1 5

<210> SEQ ID NO 134
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger protein design for eGFP
gene

<400> SEQUENCE: 134

Gln Ser Gly Asp Leu Thr Arg
1 5

<210> SEQ ID NO 135
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger protein design for eGFP
gene

<400> SEQUENCE: 135

Gln Ser Gly Asn Leu Ala Arg
1 5

<210> SEQ ID NO 136
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of synthetic GFP gene for
zinc finger design

<400> SEQUENCE: 136

ggggtagcgg

10

<210> SEQ ID NO 137
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for GFP gene

<400> SEQUENCE: 137

Arg Ser Asp Asp Leu Thr Arg
1 5

<210> SEQ ID NO 138
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic zinc finger design for GFP gene

<400> SEQUENCE: 138

Gln Ser Gly Ala Leu Ala Arg
1 5

<210> SEQ ID NO 139

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for GFP gene

<400> SEQUENCE: 139

Arg Ser Asp His Leu Ser Arg
1 5

<210> SEQ ID NO 140

<211> LENGTH: 9

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Target sequence of synthetic GFP gene for
zinc finger design

<400> SEQUENCE: 140

gaagcagca

9

<210> SEQ ID NO 141

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for GFP gene

<400> SEQUENCE: 141

Gln Ser Gly Ser Leu Thr Arg
1 5

<210> SEQ ID NO 142

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for GFP gene

<400> SEQUENCE: 142

Gln Ser Gly Asp Leu Thr Arg
1 5

<210> SEQ ID NO 143

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for GFP gene

<400> SEQUENCE: 143

Gln Ser Gly Asn Leu Ala Arg
1 5

<210> SEQ ID NO 144

<211> LENGTH: 17

<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide for mRNA analysis

<400> SEQUENCE: 144
ctgctgcccg acaacca 17

<210> SEQ ID NO 145
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide for mRNA analysis

<400> SEQUENCE: 145
ccatgtgata gcgcttctc 19

<210> SEQ ID NO 146
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide for mRNA analysis

<400> SEQUENCE: 146
cccagtcgc cctgagcaaa ga 22

<210> SEQ ID NO 147
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide for mRNA analysis

<400> SEQUENCE: 147
ccatgttcgt catgggtgtg a 21

<210> SEQ ID NO 148
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide for mRNA analysis

<400> SEQUENCE: 148
catggactgt ggtcatgagt 20

<210> SEQ ID NO 149
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide for mRNA analysis

<400> SEQUENCE: 149
tcctgcacca ccaactgctt agca 24

<210> SEQ ID NO 150
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide for construction of donor molecule

-continued

<400> SEQUENCE: 150

ggcgaggagc tgttcac

17

<210> SEQ ID NO 151

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide for construction of donor molecule

<400> SEQUENCE: 151

gattatgata tagagtcg

18

<210> SEQ ID NO 152

<211> LENGTH: 12

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma gene for zinc finger design

<400> SEQUENCE: 152

actctgtgga ag

12

<210> SEQ ID NO 153

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for exon 5 of the human IL-2Rgamma Gene

<400> SEQUENCE: 153

Arg Ser Asp Asn Leu Ser Val

1

5

<210> SEQ ID NO 154

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for exon 5 of the human IL-2Rgamma Gene

<400> SEQUENCE: 154

Arg Asn Ala His Arg Ile Asn

1

5

<210> SEQ ID NO 155

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for exon 5 of the human IL-2Rgamma Gene

<400> SEQUENCE: 155

Arg Ser Asp Thr Leu Ser Glu

1

5

<210> SEQ ID NO 156

<211> LENGTH: 7

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for exon 5 of
the human IL-2Rgamma Gene

<400> SEQUENCE: 156

Ala Arg Ser Thr Arg Thr Asn
1 5

<210> SEQ ID NO 157
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma gene for
zinc finger design

<400> SEQUENCE: 157

aaagcggctc cg

12

<210> SEQ ID NO 158
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for exon 5 of
the human IL-2Rgamma Gene

<400> SEQUENCE: 158

Arg Ser Asp Thr Leu Ser Glu
1 5

<210> SEQ ID NO 159
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for exon 5 of
the human IL-2Rgamma Gene

<400> SEQUENCE: 159

Ala Arg Ser Thr Arg Thr Thr
1 5

<210> SEQ ID NO 160
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for exon 5 of
the human IL-2Rgamma Gene

<400> SEQUENCE: 160

Arg Ser Asp Ser Leu Ser Lys
1 5

<210> SEQ ID NO 161
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for exon 5 of
the human IL-2Rgamma Gene

<400> SEQUENCE: 161

-continued

Gln Arg Ser Asn Leu Lys Val
1 5

<210> SEQ ID NO 162
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric ZFP endonuclease targeted to human
IL-2Rgamma Gene

<400> SEQUENCE: 162

Met Ala Glu Arg Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser
1 5 10 15
Arg Ser Asp Asn Leu Ser Val His Ile Arg Thr His Thr Gly Glu Lys
20 25 30
Pro Phe Ala Cys Asp Ile Cys Gly Arg Lys Phe Ala Arg Asn Ala His
35 40 45
Arg Ile Asn His Thr Lys Ile His Thr Gly Ser Gln Lys Pro Phe Gln
50 55 60
Cys Arg Ile Cys Met Arg Asn Phe Ser Arg Ser Asp Thr Leu Ser Glu
65 70 75 80
His Ile Arg Thr His Thr Gly Glu Lys Pro Phe Ala Cys Asp Ile Cys
85 90 95
Gly Arg Lys Phe Ala Ala Arg Ser Thr Arg Thr Asn His Thr Lys Ile
100 105 110
His Leu Arg Gly Ser
115

<210> SEQ ID NO 163
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chimeric ZFP endonuclease targeted to human
IL-2Rgamma Gene

<400> SEQUENCE: 163

Met Ala Glu Arg Pro Phe Gln Cys Arg Ile Cys Met Arg Asn Phe Ser
1 5 10 15
Arg Ser Asp Thr Leu Ser Glu His Ile Arg Thr His Thr Gly Glu Lys
20 25 30
Pro Phe Ala Cys Asp Ile Cys Gly Arg Lys Phe Ala Ala Arg Ser Thr
35 40 45
Arg Thr Thr His Thr Lys Ile His Thr Gly Ser Gln Lys Pro Phe Gln
50 55 60
Cys Arg Ile Cys Met Arg Asn Phe Ser Arg Ser Asp Ser Leu Ser Lys
65 70 75 80
His Ile Arg Thr His Thr Gly Glu Lys Pro Phe Ala Cys Asp Ile Cys
85 90 95
Gly Arg Lys Phe Ala Gln Arg Ser Asn Leu Lys Val His Thr Lys Ile
100 105 110
His Leu Arg Gly Ser
115

<210> SEQ ID NO 164

-continued

<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Human IL-2Rgamma insert sequence

<400> SEQUENCE: 164

Phe Arg Val Arg Ser Arg Phe Asn Pro Leu Cys Gly Ser
1 5 10

<210> SEQ ID NO 165
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Exon 5 of human IL-2Rgamma insert sequence

<400> SEQUENCE: 165

tttcgtgttc ggagccggtt taaccgcctc tgtggaagt 39

<210> SEQ ID NO 166
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide for analysis of the human
IL-2Rgamma gene

<400> SEQUENCE: 166

gattcaacca gacagataga agg 23

<210> SEQ ID NO 167
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide for analysis of the human
IL-2Rgamma gene

<400> SEQUENCE: 167

ttactgtctc atcctttact cc 22

<210> SEQ ID NO 168
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Exon 5 forward primer

<400> SEQUENCE: 168

gctaaggcca agaaagtagg gctaaag 27

<210> SEQ ID NO 169
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Exon 5 reverse primer

<400> SEQUENCE: 169

ttccttccat caccaaacc tcttg 25

<210> SEQ ID NO 170
<211> LENGTH: 27

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer targeted to human
IL-2Rgamma locus

<400> SEQUENCE: 170

gctaaggcca agaaagtagg gctaaag 27

<210> SEQ ID NO 171
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer targeted to human
IL-2Rgamma locus

<400> SEQUENCE: 171

ttccttccat caccaaacc tcttg 25

<210> SEQ ID NO 172
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Donor sequence; modified synthetic GFP sequence

<400> SEQUENCE: 172

cttcagccgc tatccagacc acatgaaaca acacgacttc tt 42

<210> SEQ ID NO 173
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Donor sequence; modified synthetic GFP sequence

<400> SEQUENCE: 173

cttcagccgc tatccagacc acatgaaaca acatgacttc tt 42

<210> SEQ ID NO 174
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Donor sequence; modified synthetic GFP sequence

<400> SEQUENCE: 174

cttcagccgc taccagacc acatgaaaca gcacgacttc tt 42

<210> SEQ ID NO 175
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Donor sequence; modified synthetic GFP sequence

<400> SEQUENCE: 175

cttcagccgc taccagacc acatgaagca gcacgacttc tt 42

<210> SEQ ID NO 176
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic GFP mut sequence

<400> SEQUENCE: 176

cttcagccgc taccctaac acgaagcagc acgacttctt 40

<210> SEQ ID NO 177
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic GFP wt sequence

<400> SEQUENCE: 177

cttcagccgc taccctgacc acatgaagca gcaagacttc tt 42

<210> SEQ ID NO 178
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target site of the ZFP DNA binding domain for human beta-globin gene

<400> SEQUENCE: 178

gaagtctgcc gt 12

<210> SEQ ID NO 179
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target site of the ZFP DNA binding domain for human beta-globin gene

<400> SEQUENCE: 179

gaagtctgcc gtt 13

<210> SEQ ID NO 180
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target site of the ZFP DNA binding domain for human beta-globin gene

<400> SEQUENCE: 180

gaagtctgcc gtt 13

<210> SEQ ID NO 181
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sequence of human beta-globin gene
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(52)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 181

caaacagaca ccatggtgca tctgactcct gtggagaagt ctgccgttac tg 52

<210> SEQ ID NO 182

-continued

<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target site of the ZFP DNA binding domain for
human beta-globin gene

<400> SEQUENCE: 182

acgtagactg agg

13

<210> SEQ ID NO 183
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Non-canonical inter-finger linker

<400> SEQUENCE: 183

Thr Gly Gly Gly Gly Ser Gln Lys Pro
1 5

<210> SEQ ID NO 184
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 184

Gln Ser Gly Asp Leu Thr Arg
1 5

<210> SEQ ID NO 185
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 185

Thr Ser Ala Asn Leu Ser Arg
1 5

<210> SEQ ID NO 186
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 186

Asp Arg Ser Ala Leu Ser Arg
1 5

<210> SEQ ID NO 187
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 187

-continued

Gln Ser Gly His Leu Ser Arg
1 5

<210> SEQ ID NO 188
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 188

Arg Ser Gln Thr Arg Lys Thr
1 5

<210> SEQ ID NO 189
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 189

Gln Lys Arg Asn Arg Thr Lys
1 5

<210> SEQ ID NO 190
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 190

Asp Arg Ser Ala Leu Ser Arg
1 5

<210> SEQ ID NO 191
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 191

Gln Ser Gly Asn Leu Ala Arg
1 5

<210> SEQ ID NO 192
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 192

Thr Ser Gly Ser Leu Ser Arg
1 5

<210> SEQ ID NO 193

-continued

<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 193

Asp Arg Ser Asp Leu Ser Arg
1 5

<210> SEQ ID NO 194
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 194

Asp Arg Ser Ala Leu Ser Arg
1 5

<210> SEQ ID NO 195
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 195

Gln Ser Gly Asn Leu Ala Arg
1 5

<210> SEQ ID NO 196
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 196

Thr Ser Ser Ser Leu Ser Arg
1 5

<210> SEQ ID NO 197
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 197

Asp Arg Ser Asp Leu Ser Arg
1 5

<210> SEQ ID NO 198
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

-continued

<400> SEQUENCE: 198

Asp Arg Ser Ala Leu Ser Arg
1 5

<210> SEQ ID NO 199

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Recognition region of ZFP for human
beta-globin gene

<400> SEQUENCE: 199

Gln Ser Gly Asn Leu Ala Arg
1 5

<210> SEQ ID NO 200

<211> LENGTH: 55

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: DNA fragment containing human beta-globin gene

<400> SEQUENCE: 200

ctagacacca tgggtgcatct gactcctgtg gagaagtctg ccgttactgc cctag 55

<210> SEQ ID NO 201

<211> LENGTH: 27

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger structure

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(2)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (4)..(7)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

and up to 2 of them can be present or absent

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (9)..(20)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (22)..(26)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

and up to 2 of them can be present or absent

<400> SEQUENCE: 201

Xaa Xaa Cys Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10 15Xaa Xaa Xaa Xaa His Xaa Xaa Xaa Xaa His
20 25

<210> SEQ ID NO 202

<211> LENGTH: 12

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma ZFP domain

<400> SEQUENCE: 202

-continued

aactcggata at 12

<210> SEQ ID NO 203
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human IL-2Rgamma binding domain for zinc finger design

<400> SEQUENCE: 203

aactcggata at 12

<210> SEQ ID NO 204
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design of human IL-2Rgamma binding domain

<400> SEQUENCE: 204

Asp Arg Ser Thr Leu Ile Glu
1 5

<210> SEQ ID NO 205
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design of human IL-2Rgamma binding domain

<400> SEQUENCE: 205

Ser Ser Ser Asn Ser Leu Arg
1 5

<210> SEQ ID NO 206
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design of human IL-2Rgamma binding domain

<400> SEQUENCE: 206

Arg Ser Asp Asp Leu Ser Lys
1 5

<210> SEQ ID NO 207
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design of human IL-2Rgamma binding domain

<400> SEQUENCE: 207

Asp Asn Ser Asn Arg Ile Lys
1 5

<210> SEQ ID NO 208
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer used in PCR to obtain FokI
cleavage half-domain

<400> SEQUENCE: 208

ggatcccaac tagtcaaaag tgaac 25

<210> SEQ ID NO 209
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer used in PCR to obtain FokI
cleavage half-domain

<400> SEQUENCE: 209

ctcgagttaa aagtttatct cgccg 25

<210> SEQ ID NO 210
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic nuclear localization sequence

<400> SEQUENCE: 210

Pro Lys Lys Lys Arg Lys Val
1 5

<210> SEQ ID NO 211
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sequence of portion of substrate containing
binding sites for 5-9D and human IL-2Rgamma zinc finger binding
domains
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(32)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 211

taaagcggct ccgaaccaca ttatccgagt tc 32

<210> SEQ ID NO 212
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sequence of portion of substrate containing
binding sites for 5-9D and human IL-2Rgamma zinc finger binding
domains
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(32)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 212

taaagcggct ccgaaccaca actcggataa tg 32

<210> SEQ ID NO 213
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic endonuclease I-SceI

<400> SEQUENCE: 213

tagggataac agggtaat 18

<210> SEQ ID NO 214

<211> LENGTH: 58

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Sequence from exon 5 of the human IL-2Rgamma gene

<400> SEQUENCE: 214

ctagctacac gtttcgtggt cggagccgct ttaaccact ctgtggaagt gtcctag 58

<210> SEQ ID NO 215

<211> LENGTH: 34

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic amplification primer

<400> SEQUENCE: 215

actagtgccg ccaccatgac cgagtacaag ccca 34

<210> SEQ ID NO 216

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic amplification primer

<400> SEQUENCE: 216

actagtcagg caccgggctt 20

<210> SEQ ID NO 217

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic PCR primer used to amplify the pcDNA4/TO/GFPpuro vector

<400> SEQUENCE: 217

cgaattctgc agtcgac 17

<210> SEQ ID NO 218

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic PCR primer used to amplify the pcDNA4/TO/GFPpuro vector

<400> SEQUENCE: 218

tgcatattc tgctctgc 17

<210> SEQ ID NO 219

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic PCR primer

-continued

<400> SEQUENCE: 219

tttgacctcc atagaagaca 20

<210> SEQ ID NO 220

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic PCR primer

<400> SEQUENCE: 220

gcgcaccgtg ggcttgact 20

<210> SEQ ID NO 221

<211> LENGTH: 1014

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Codon-optimized nucleotide sequence encoding
ZFN 5-8D targeted to exon 5 of human IL-2Rgamma gene

<400> SEQUENCE: 221

aattcgctag cgccaccatg gcccccaaga agaagaggaa agtgggaatc cacggggtag 60
ccgccgctat ggccgagagg cccttcacgt gtcggatctg catgcggaac ttcagccgga 120
gcgacaacct gagcgtgcac atccgcaccc acacaggcga gaagcctttt gcctgtgaca 180
tttgtgggag gaaatttgcc cgcaacgccc accgcatcaa ccacaccaag atccacaccg 240
gatctcagaa gcccttttcag tgcagaatct gcattgagaaa cttctcccggt tccgacaccc 300
tgagcgaaca catcaggaca cacaccggcg agaaaccctt cgcttgcgac atctgtggcc 360
gcaagtttgc cgccagaagc acccgacaaa atcacacaaa gattcacctg cggggatccc 420
agctggtgaa gagcgagctg gaggagaaga agtccgagct gcggcacaag ctgaagtacg 480
tgccccacga gtacatcgag ctgatcgaga tcgccaggaa cagcacccag gaccgcatcc 540
tgagatgaa ggtgatggag ttcttcatga aggtgtacgg ctacagggga aagcacctgg 600
gcggaagcag aaagcctgac ggcgcatct atacagtggg cagccccatc gattacggcg 660
tgatcgtgga cacaaggcc tacagcggcg gctacaatct gcctatcggc caggccgacg 720
agatgcagag atacgtggag gagaaccaga cccggaataa gcacatcaac cccaacgagt 780
ggtggaaggt gtaccctagc agcgtgaccg agttcaagtt cctgttcgtg agcggccact 840
tcaagggcaa ctacaaggcc cagctgacca ggctgaacca catcaccaac tgcaatggcg 900
ccgtgctgag cgtggaggag ctgctgatcg gcggcgagat gatcaaagcc ggcaccctga 960
cactggagga ggtgcggcgc aagttcaaca acggcgagat caacttctga taac 1014

<210> SEQ ID NO 222

<211> LENGTH: 1014

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Codon-optimized nucleotide sequence encoding
ZFN 5-9D targeted to exon 5 of human IL-2Rgamma gene

<400> SEQUENCE: 222

aattcgctag cgccaccatg gcccccaaga agaagaggaa agtgggaatc cacggggtag 60
ccgccgctat ggccgagagg cccttcacgt gtcggatctg catgcggaac ttcagcagga 120

-continued

```

gcgacaccct gagcgaacac atccgcaccc acacaggcga gaagcctttt gcctgtgaca    180
tttgtgggag gaaatttgcc gccagaagca ccgcgacaac ccacaccaag atccacaccg    240
gatctcagaa gcccttttcag tgcagaatct gcatgagaaa cttctcccggt tccgacagcc    300
tgagcaagca cattaggacc cacaccgggg agaaaccctt cgctgtcgac atctgtggcc    360
gcaaatttgc ccagcgagc aacctgaaag tgcacacaaa gattcacctg cggggatccc    420
agctggtgaa gagcgagctg gaggagaaga agtccgagct gcggcacaag ctgaagtacg    480
tgccccacga gtacatcgag ctgatcgaga tcgccaggaa cagcaccag gaccgcatcc    540
tggagatgaa ggtgatggag ttcttcatga aggtgtacgg ctacagggga aagcacctgg    600
gcggaagcag aaagcctgac ggcgccatct atacagtggg cagccccatc gattacggcg    660
tgatcgtgga ccaaaggcc tacagcggcg gctacaatct gcctatcggc caggccgacg    720
agatgcagag atacgtggag gagaaccaga ccggaataa gcacatcaac cccaacgagt    780
ggtggaaggt gtaccctagc agcgtgaccg agttcaagtt cctgttcgtg agcggccact    840
tcaagggcaa ctacaaggcc cagctgacca ggctgaacca catcaccaac tgcaatggcg    900
ccgtgctgag cgtggaggag ctgctgatcg gcggcgagat gatcaaagcc ggcaccctga    960
cactggagga ggtgcggcgc aagttcaaca acggcgagat caacttctga taac        1014

```

```

<210> SEQ ID NO 223
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of CHO DHFR gene for zinc
                        finger design

```

<400> SEQUENCE: 223

```

ggaaggtctc cg                                12

```

```

<210> SEQ ID NO 224
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the CHO DHFR
                        gene

```

<400> SEQUENCE: 224

```

Arg Ser Asp Thr Leu Ser Glu
1              5

```

```

<210> SEQ ID NO 225
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the CHO DHFR
                        gene

```

<400> SEQUENCE: 225

```

Asn Asn Arg Asp Arg Thr Lys
1              5

```

```

<210> SEQ ID NO 226
<211> LENGTH: 7
<212> TYPE: PRT

```

-continued

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the CHO DHFR gene

<400> SEQUENCE: 226

Arg Ser Asp His Leu Ser Ala
1 5

<210> SEQ ID NO 227

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the CHO DHFR gene

<400> SEQUENCE: 227

Gln Ser Gly His Leu Ser Arg
1 5

<210> SEQ ID NO 228

<211> LENGTH: 12

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Target sequence of CHO DHFR gene for zinc finger design

<400> SEQUENCE: 228

aatgctcagg ta

12

<210> SEQ ID NO 229

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the CHO DHFR gene

<400> SEQUENCE: 229

Gln Ser Gly Ala Leu Ala Arg
1 5

<210> SEQ ID NO 230

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the CHO DHFR gene

<400> SEQUENCE: 230

Arg Ser Asp Asn Leu Arg Glu
1 5

<210> SEQ ID NO 231

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic zinc finger design for the CHO DHFR gene

<400> SEQUENCE: 231

-continued

Gln Ser Ser Asp Leu Ser Arg
1 5

<210> SEQ ID NO 232
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the CHO DHFR
gene

<400> SEQUENCE: 232

Thr Ser Ser Asn Arg Lys Thr
1 5

<210> SEQ ID NO 233
<211> LENGTH: 84
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Wild-type CHO DHFR sequence

<400> SEQUENCE: 233

ctcggtttcc ctaacccaat ccagccagta cctgagcatt ggccagggaa ggtctccgtt 60

cttgccgatg cccatattct ggga 84

<210> SEQ ID NO 234
<211> LENGTH: 240
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic mutated DHFR sequence

<400> SEQUENCE: 234

ctcggtttcc ctaacccaat ccagccagta cctgagcatt gtggcacctt ccagggtcaa 60

ggaaggcacg ggggaggggc aaacaacaga tggctggcaa ctagaaggca cagtcgaggc 120

tgatcagcgg tttaaactta agcttggtag cgagctcgga tccactttga cgtcaatggg 180

tggagtattt acggtaaacc agggaaggtc tccgttcttg ccgatgccca tattctggga 240

<210> SEQ ID NO 235
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer used in amplification of
eGFP sequences

<400> SEQUENCE: 235

ggcggaggagc tgttcac 17

<210> SEQ ID NO 236
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer used in amplification of
eGFP sequences

<400> SEQUENCE: 236

tgcatacttc tgctctgc 17

-continued

<210> SEQ ID NO 237
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sites for zinc finger domains in human CCR-5 gene
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(35)
<223> OTHER INFORMATION: STRANDEDNESS: double

<400> SEQUENCE: 237

ctgggtcatcc tcatcctgat aaactgcaaa aggct 35

<210> SEQ ID NO 238
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence in human CCR-5 gene for zinc finger design

<400> SEQUENCE: 238

gatgaggatg ac 12

<210> SEQ ID NO 239
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human CCR-5 gene

<400> SEQUENCE: 239

Asp Arg Ser Asn Leu Ser Arg
1 5

<210> SEQ ID NO 240
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human CCR-5 gene

<400> SEQUENCE: 240

Thr Ser Ala Asn Leu Ser Arg
1 5

<210> SEQ ID NO 241
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human CCR-5 gene

<400> SEQUENCE: 241

Arg Ser Asp Asn Leu Ala Arg
1 5

<210> SEQ ID NO 242
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
CCR-5 gene

<400> SEQUENCE: 242

Thr Ser Ala Asn Leu Ser Arg
1 5

<210> SEQ ID NO 243
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Target sequence of human CCR-5 gene for zinc
finger design

<400> SEQUENCE: 243

aaactgcaaa ag

12

<210> SEQ ID NO 244
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
CCR-5 gene

<400> SEQUENCE: 244

Arg Ser Asp His Leu Ser Glu
1 5

<210> SEQ ID NO 245
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
CCR-5 gene

<400> SEQUENCE: 245

Gln Asn Ala Asn Arg Ile Thr
1 5

<210> SEQ ID NO 246
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
CCR-5 gene

<400> SEQUENCE: 246

Arg Ser Asp Val Leu Ser Glu
1 5

<210> SEQ ID NO 247
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic zinc finger design for the human
CCR-5 gene

<400> SEQUENCE: 247

Gln Arg Asn His Arg Thr Thr

-continued

1	5	
 <210> SEQ ID NO 248 <211> LENGTH: 1928 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Donor DNA molecule; modified fragment of human CCR-5 gene <400> SEQUENCE: 248		
gttgtcaaa	cttcattcac	tccatggtgc
ggtgctttca	tgaattcccc	caacagagcc
gcagcaaa	ttcccttcac	tacaaaactt
atgtagacat	ctatgtaggc	aattaaaaac
ggagggcaac	taaatacatt	ctaggacttt
gtggaacaag	atggattatc	aagtggtcaag
ggagccctgc	caaaaaatca	atgtgaagca
ctcactgggtg	ttcatctttg	gttttgtggg
tgagtatgcc	ctgatggcgt	ctggactgga
aagagcatga	ctgacatcta	cctgctcaac
actgtccct	tctgggctca	ctatgctgcc
caactcttga	cagggtctta	ttttataggc
ctgacaatcg	ataggtagct	ggctgtcgtc
gtcacctttg	gggtgggtgac	aagtggtgac
ccaggaatca	tctttaccag	atctcaaaaa
tttccatata	gtcagtatca	attctggaag
gggtgggtcc	tgccgtgctg	tgtcatggtc
cttcgggtgc	gaaatgagaa	gaagaggcac
attgtttatt	ttctcttctg	ggctccctac
gaattctttg	gcctgaataa	ttgcagtagc
acagagactc	ttgggatgac	gcaactgctg
gagaagtcca	gaaactacct	cttagtcttc
aaatgctgtt	ctattttcca	gcaagaggct
tccactgggg	agcaggaaat	atctgtgggc
ccagttagag	ttgtgcacat	ggcttagttt
ggagaggctc	tttttaaaag	gaagttagct
ttggcatctg	tttaaagtag	attagatctt
aaaaatgttt	gatgaaaaat	agcaaccttt
gacaaactct	cccttcactc	cgaaagtctc
attgctgatt	cttgagtcca	gtgatctgaa
caacttttta	cctagtacaa	ggcaacatat
gtcttgctat	ggggagaaaa	gacatgaata

-continued

gtgatttc

1928

<210> SEQ ID NO 249
 <211> LENGTH: 196
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Wild-type FokI cleavage half-domain

<400> SEQUENCE: 249

Gln Leu Val Lys Ser Glu Leu Glu Glu Lys Lys Ser Glu Leu Arg His
 1 5 10 15
 Lys Leu Lys Tyr Val Pro His Glu Tyr Ile Glu Leu Ile Glu Ile Ala
 20 25 30
 Arg Asn Ser Thr Gln Asp Arg Ile Leu Glu Met Lys Val Met Glu Phe
 35 40 45
 Phe Met Lys Val Tyr Gly Tyr Arg Gly Lys His Leu Gly Gly Ser Arg
 50 55 60
 Lys Pro Asp Gly Ala Ile Tyr Thr Val Gly Ser Pro Ile Asp Tyr Gly
 65 70 75 80
 Val Ile Val Asp Thr Lys Ala Tyr Ser Gly Gly Tyr Asn Leu Pro Ile
 85 90 95
 Gly Gln Ala Asp Glu Met Gln Arg Tyr Val Glu Glu Asn Gln Thr Arg
 100 105 110
 Asn Lys His Ile Asn Pro Asn Glu Trp Trp Lys Val Tyr Pro Ser Ser
 115 120 125
 Val Thr Glu Phe Lys Phe Leu Phe Val Ser Gly His Phe Lys Gly Asn
 130 135 140
 Tyr Lys Ala Gln Leu Thr Arg Leu Asn His Ile Thr Asn Cys Asn Gly
 145 150 155 160
 Ala Val Leu Ser Val Glu Glu Leu Leu Ile Gly Gly Glu Met Ile Lys
 165 170 175
 Ala Gly Thr Leu Thr Leu Glu Glu Val Arg Arg Lys Phe Asn Asn Gly
 180 185 190
 Glu Ile Asn Phe
 195

<210> SEQ ID NO 250
 <211> LENGTH: 196
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic cleavage half-domains

<400> SEQUENCE: 250

Gln Leu Val Lys Ser Glu Leu Glu Glu Lys Lys Ser Glu Leu Arg His
 1 5 10 15
 Lys Leu Lys Tyr Val Pro His Glu Tyr Ile Glu Leu Ile Glu Ile Ala
 20 25 30
 Arg Asn Ser Thr Gln Asp Arg Ile Leu Glu Met Lys Val Met Glu Phe
 35 40 45
 Phe Met Lys Val Tyr Gly Tyr Arg Gly Lys His Leu Gly Gly Ser Arg
 50 55 60
 Lys Pro Asp Gly Ala Ile Tyr Thr Val Gly Ser Pro Ile Asp Tyr Gly
 65 70 75 80

-continued

```

Val Ile Val Asp Thr Lys Ala Tyr Ser Gly Gly Tyr Asn Leu Pro Ile
      85                      90                      95
Gly Gln Ala Asp Glu Met Gln Arg Tyr Val Lys Glu Asn Gln Thr Arg
      100                    105                    110
Asn Lys His Ile Asn Pro Asn Glu Trp Trp Lys Val Tyr Pro Ser Ser
      115                    120                    125
Val Thr Glu Phe Lys Phe Leu Phe Val Ser Gly His Phe Lys Gly Asn
      130                    135                    140
Tyr Lys Ala Gln Leu Thr Arg Leu Asn His Ile Thr Asn Cys Asn Gly
      145                    150                    155                    160
Ala Val Leu Ser Val Glu Glu Leu Leu Ile Gly Gly Glu Met Ile Lys
      165                    170                    175
Ala Gly Thr Leu Thr Leu Glu Glu Val Arg Arg Lys Phe Asn Asn Gly
      180                    185                    190
Glu Ile Asn Phe
      195

```

```

<210> SEQ ID NO 251
<211> LENGTH: 196
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage half-domains

```

```

<400> SEQUENCE: 251

```

```

Gln Leu Val Lys Ser Glu Leu Glu Glu Lys Lys Ser Glu Leu Arg His
1          5          10          15
Lys Leu Lys Tyr Val Pro His Glu Tyr Ile Glu Leu Ile Glu Ile Ala
20         25         30
Arg Asn Ser Thr Gln Asp Arg Ile Leu Glu Met Lys Val Met Glu Phe
35         40         45
Phe Met Lys Val Tyr Gly Tyr Arg Gly Lys His Leu Gly Gly Ser Arg
50         55         60
Lys Pro Asp Gly Ala Ile Tyr Thr Val Gly Ser Pro Ile Asp Tyr Gly
65         70         75         80
Val Ile Val Asp Thr Lys Ala Tyr Ser Gly Gly Tyr Asn Leu Pro Ile
85         90         95
Gly Gln Ala Asp Glu Met Gln Arg Tyr Val Lys Glu Asn Gln Thr Arg
100        105        110
Asn Lys His Ile Asn Pro Asn Glu Trp Trp Lys Val Tyr Pro Ser Ser
115        120        125
Val Thr Glu Phe Lys Phe Leu Phe Val Ser Gly His Phe Lys Gly Asn
130        135        140
Tyr Lys Ala Gln Leu Thr Arg Leu Asn His Lys Thr Asn Cys Asn Gly
145        150        155        160
Ala Val Leu Ser Val Glu Glu Leu Leu Ile Gly Gly Glu Met Ile Lys
165        170        175
Ala Gly Thr Leu Thr Leu Glu Glu Val Arg Arg Lys Phe Asn Asn Gly
180        185        190
Glu Ile Asn Phe
195

```

```

<210> SEQ ID NO 252

```

-continued

<211> LENGTH: 196
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage half-domains

<400> SEQUENCE: 252

Gln Leu Val Lys Ser Glu Leu Glu Glu Lys Lys Ser Glu Leu Arg His
1 5 10 15
Lys Leu Lys Tyr Val Pro His Glu Tyr Ile Glu Leu Ile Glu Ile Ala
20 25 30
Arg Asn Ser Thr Gln Asp Arg Ile Leu Glu Met Lys Val Met Glu Phe
35 40 45
Phe Met Lys Val Tyr Gly Tyr Arg Gly Lys His Leu Gly Gly Ser Arg
50 55 60
Lys Pro Asp Gly Ala Ile Tyr Thr Val Gly Ser Pro Ile Asp Tyr Gly
65 70 75 80
Val Ile Val Asp Thr Lys Ala Tyr Ser Gly Gly Tyr Asn Leu Pro Ile
85 90 95
Gly Gln Ala Asp Glu Met Glu Arg Tyr Val Lys Glu Asn Gln Thr Arg
100 105 110
Asn Lys His Ile Asn Pro Asn Glu Trp Trp Lys Val Tyr Pro Ser Ser
115 120 125
Val Thr Glu Phe Lys Phe Leu Phe Val Ser Gly His Phe Lys Gly Asn
130 135 140
Tyr Lys Ala Gln Leu Thr Arg Leu Asn His Lys Thr Asn Cys Asn Gly
145 150 155 160
Ala Val Leu Ser Val Glu Glu Leu Leu Ile Gly Gly Glu Met Ile Lys
165 170 175
Ala Gly Thr Leu Thr Leu Glu Glu Val Arg Arg Lys Phe Asn Asn Gly
180 185 190
Glu Ile Asn Phe
195

<210> SEQ ID NO 253
<211> LENGTH: 196
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic cleavage half-domains

<400> SEQUENCE: 253

Gln Leu Val Lys Ser Glu Leu Glu Glu Lys Lys Ser Glu Leu Arg His
1 5 10 15
Lys Leu Lys Tyr Val Pro His Glu Tyr Ile Glu Leu Ile Glu Ile Ala
20 25 30
Arg Asn Ser Thr Gln Asp Arg Ile Leu Glu Met Lys Val Met Glu Phe
35 40 45
Phe Met Lys Val Tyr Gly Tyr Arg Gly Lys His Leu Gly Gly Ser Arg
50 55 60
Lys Pro Asp Gly Ala Ile Tyr Thr Val Gly Ser Pro Ile Asp Tyr Gly
65 70 75 80
Val Ile Val Asp Thr Lys Ala Tyr Ser Gly Gly Tyr Asn Leu Pro Ile
85 90 95
Gly Gln Ala Asp Glu Met Ile Arg Tyr Val Lys Glu Asn Gln Thr Arg

-continued

100					105					110						
Asn	Lys	His	Ile	Asn	Pro	Asn	Glu	Trp	Trp	Lys	Val	Tyr	Pro	Ser	Ser	
115					120					125						
Val	Thr	Glu	Phe	Lys	Phe	Leu	Phe	Val	Ser	Gly	His	Phe	Lys	Gly	Asn	
130					135					140						
Tyr	Lys	Ala	Gln	Leu	Thr	Arg	Leu	Asn	His	Lys	Thr	Asn	Cys	Asn	Gly	
145					150					155					160	
Ala	Val	Leu	Ser	Val	Glu	Glu	Leu	Leu	Ile	Gly	Gly	Glu	Met	Ile	Lys	
165					170					175						
Ala	Gly	Thr	Leu	Thr	Leu	Glu	Glu	Val	Arg	Arg	Lys	Phe	Asn	Asn	Gly	
180					185					190						
Glu	Ile	Asn	Phe													
195																

What is claimed is:

1. A method for expressing the product of an exogenous nucleic acid sequence in a cell, the method comprising:

- (a) expressing a first fusion protein in the cell, the first fusion protein comprising a first zinc finger binding domain and a first cleavage half-domain, wherein the first zinc finger binding domain has been engineered to bind to a first target site in a region of interest in the genome of the cell;
- (b) expressing a second fusion protein in the cell, the second fusion protein comprising a second zinc finger binding domain and a second cleavage half domain, wherein the second zinc finger binding domain binds to a second target site in the region of interest in the genome of the cell, wherein the second target site is different from the first target site; and
- (c) contacting the cell with a polynucleotide comprising an exogenous nucleic acid sequence and a first nucleotide sequence that is homologous to a first sequence in the region of interest;

wherein binding of the first fusion protein to the first target site, and binding of the second fusion protein to the second target site, positions the cleavage half-domains such that the genome of the cell is cleaved in the region of interest, thereby resulting in integration of the exogenous sequence into the genome of the cell in the region of interest and expression of the product of the exogenous sequence.

2. The method according to claim 1, wherein the polynucleotide further comprises a second nucleotide sequence that is homologous to a second sequence in the region of interest.

3. The method according to claim 2, wherein the first and second nucleotide sequences flank the exogenous sequence.

4. The method according to claim 1, wherein the region of interest is in a region of the genome that is transcriptionally active and not essential for viability.

5. The method according to claim 4, wherein the region of interest is the human homologue of the murine Rosa26 gene.

6. The method according to claim 4, wherein the region of interest is a CCR5 gene.

7. The method according to claim 1, wherein the first and second cleavage half-domains are from a Type IIS restriction endonuclease.

8. The method according to claim 1, wherein the region of interest comprises a gene.

9. The method according to claim 8, wherein the gene comprises a mutation.

10. The method according to claim 9, wherein the exogenous nucleic acid sequence comprises the wild-type sequence of the gene.

11. The method according to claim 9, wherein the exogenous nucleic acid sequence comprises a portion of the wild-type sequence of the gene.

12. The method according to claim 9, wherein the exogenous nucleic acid sequence comprises a cDNA copy of a transcription product of the gene.

13. The method according to claim 1, wherein the cell is arrested in the G2 phase of the cell cycle.

14. The method according to claim 1, wherein at least one of the fusion proteins comprises an alteration in the amino acid sequence of the dimerization interface of the cleavage half-domain.

15. A method for integrating an exogenous sequence into a region of interest in the genome of a cell, the method comprising:

- (a) expressing a first fusion protein in the cell, the first fusion protein comprising a first zinc finger binding domain and a first cleavage half-domain, wherein the first zinc finger binding domain has been engineered to bind to a first target site in a region of interest in the genome of the cell;
- (b) expressing a second fusion protein in the cell, the second fusion protein comprising a second zinc finger binding domain and a second cleavage half domain, wherein the second zinc finger binding domain binds to a second target site in the region of interest in the genome of the cell, wherein the second target site is different from the first target site; and
- (c) contacting the cell with a polynucleotide comprising an exogenous nucleic acid sequence;

wherein binding of the first fusion protein to the first target site, and binding of the second fusion protein to

the second target site, positions the cleavage half-domains such that the genome of the cell is cleaved in the region of interest, thereby resulting in integration of the exogenous sequence into the genome of the cell in the region of interest.

16. The method according to claim 15, wherein the integration inactivates gene expression in the region of interest.

17. The method according to claim 15, wherein the exogenous nucleic acid sequence comprises a sequence of between 1 and 50 nucleotides in length.

18. The method according to claim 15, wherein the exogenous sequence comprises a cleavage enzyme recognition site.

19. The method according to claim 18, wherein the cleavage enzyme is a meganuclease.

20. The method according to claim 19, wherein the meganuclease is I-SceI.

21. The method according to claim 19, wherein the meganuclease has been engineered to bind a non-natural target site.

22. The method according to claim 15, wherein the first and second cleavage half-domains are from a Type IIS restriction endonuclease.

23. The method according to claim 15, wherein the cell is arrested in the G2 phase of the cell cycle.

24. The method according to claim 15, wherein at least one of the fusion proteins comprises an alteration in the amino acid sequence of the dimerization interface of the cleavage half-domain.

* * * * *