



(51) International Patent Classification:

A23L 33/135 (2016.01) A23C 9/12 (2006.01)

(21) International Application Number:

PCT/EP2017/061270

(22) International Filing Date:

11 May 2017 (11.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

16169215.7 11 May 2016 (11.05.2016) EP
16197223.7 04 November 2016 (04.11.2016) EP

(71) Applicant: **CHR. HANSEN A/S** [DK/DK]; Boege Alle
10-12, 2970 Hoersholm (DK).

(72) Inventors: **GILLELADEN, Christian**; Enghavevej 55,
1 sal, 1674 Copenhagen V (DK). **TRIHAAS, Jeorgos**;
Egholmvej 2, 3310 Oelsted (DK). **JANZEN, Thomas**;
Byvangen 4, 2700 Broenshoej (DK). **OEHRSTROEM**
RUNGE, Mette; Piletoften 18, 3070 Snekkersten (DK).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR,
KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))
— with (an) indication(s) in relation to deposited biological
material furnished under Rule 13bis separately from the
description (Rules 13bis.4(d)(i) and 48.2(a)(viii))

(54) Title: LACTIC ACID BACTERIA FOR A HEAT-TREATED FOOD PRODUCT FOR STORAGE AT AMBIENT TEMPERATURE

(57) Abstract: Process for producing an ambient storage food product comprising providing a food product with a pH of between 3.4 and 4.4, subjecting the food product to a heat treatment to obtain a heat treated food product, adding aseptically to the heat treated food product one or more of ambient storage lactic acid bacteria strains to obtain an ambient storage food product, and storing the ambient storage food product at ambient temperature for a period of time, wherein the ambient storage lactic acid bacteria strain is selected from the group consisting of strains, (i) wherein the strain is capable of retaining viability at the end of 150 days at a temperature of 25°C, and (ii) wherein the pH at most decreases 0.8 units during the storage period, and (iii) wherein the strain is selected from the group consisting of *Lactobacillus paracasei*, *Lactobacillus rhamnosus*, *Lactobacillus fermentum* and *Lactobacillus delbrueckii* subsp. *bulgaricus*, and mutants and variants thereof.



LACTIC ACID BACTERIA FOR A HEAT-TREATED FOOD PRODUCT FOR STORAGE AT AMBIENT TEMPERATURE

FIELD OF THE INVENTION

5

The present invention relates to lactic acid bacteria, which are suitable for adding to a heat-treated food product with a pH of between 3.4 and 4.4 to be stored at ambient temperature.

10 BACKGROUND OF THE INVENTION

In recent years fermented dairy products, such as yogurts, which can be stored, transported, handled and consumed in non-refrigerated conditions, i.e. at ambient temperature, for several months have become widely used. Such
15 yogurts allow the consumer to carry the yogurt with him/her for a period of time without the need for refrigeration in the same manner as is possible for a number of beverages, and hence such yogurts provide a significant convenience advantage for the consumer. In order to obtain such a long-term shelf life at ambient temperature, the yogurt has been heat-treated after completion of the
20 fermentation process to kill or at least inhibit further growth of the bulk of the lactic acid bacteria used in the fermentation process. The heat-treatment may e.g. be a pasteurization process or an Ultra High Temperature (UHT) process. Such yogurts are sometimes referred to as Post Pasteurization Yogurt or as Ambient Yogurt.

25

Post Pasteurization Yogurt products contain no or only few viable lactic acid bacteria. However, it is desired that Post Pasteurization Yogurt products contain lactic acid bacteria and or probiotic bacteria in order to provide the consumers with the various benefits of such bacteria, e.g. health and food supplement
30 benefits. Of course the addition of live bacteria to Post Pasteurization Yogurt products to be stored at ambient temperatures introduces the technical problem that the bacteria will propagate to an extent where the yogurt will spoil, e.g. by a decrease of pH. In the prior art this technical challenge has been addressed in a number of different ways. For example, the bacteria cultures to the Post
35 Pasteurization Yogurt products have been added in the to the Post Pasteurization

Yogurt products in the form of spores. Also, the bacteria cultures for addition to the Post Pasteurization Yogurt products have been added in the form of powdered, dried, freeze-dried, coated or encapsulated cultures. Furthermore, the bacteria cultures for addition to the Post Pasteurization Yogurt products have
5 been inactivated e.g. by irradiation, microwave treatment, antibiotics, mild pasteurization, chemical agents (inhibitor) or adjustment of pH, water activity or temperature.

WO2009/116864 discloses a dairy product containing spores of probiotic
10 bacteria, wherein the dairy product can be stored at non-refrigerated temperatures for an extended period of time.

WO2004/069156 discloses food products containing probiotic bacteria, which has been inactivated by irradiation, microwave treatment, antibiotics, mild
15 pasteurization and chemical agents (inhibitor).

EP-B1-1 289 380 discloses a food product, such as dairy products, containing non-viable Lactobacillus bacteria. The Lactobacillus may be rendered non-viable e.g. by mild heat-treatment, pH adjustment or water activity adjustment.
20

EP-B1-1 514 553 discloses a double-coated lactic acid bacteria powder with high survival rate in the human body, wherein the lactic acid bacteria have been coated doubly by a protein and a polysaccharide.

25 CN101323850 discloses a process of producing microcapsules of Lactobacillus helveticus in microencapsulated form having a strong heat resistance.

EP-B1-0 555 618 discloses a dietary product containing lyophilized lactic acid bacteria.
30

CN102492643 discloses a Lactobacillus rhamnosus strain GRX19 and its application in a starter culture for producing a fermented milk product containing live Lactobacillus bacteria. The fermented milk product is subjected to heat treatment, e.g. at 70-75 °C for 15-20 seconds, and the Lactobacillus strain is
35 resistant to said heat treatment in that a fraction of the bacteria, e.g. 10×10^7

CFU/ml survives the heat treatment. After heat treatment the heat-treated product is filled aseptically into a container and stored at room temperature for e.g. 30 days.

- 5 WO2015/169928 discloses a liquid dairy composition suitable for making a foamed dairy product, wherein the composition is shelf-stable under ambient storage conditions, has a pH of between 3.8 and 4.4 and comprises fermented milk, up to 0.12 % hydrolyzed whey protein, up to 5 % fat, and up to 1 % high methylester pectin.

10

US20100009034 discloses a process of preparing a fermented milk beverage keeping high viable cell count at ambient temperature comprising performing a milk fermentation using a conventional starter culture of lactic acid bacteria, diluting, mixing and sterilizing, and adding *Lactobacillus rhamnosus* ATCC 53103

- 15 to the mixed milk beverage at aseptic conditions.

US20100015285 discloses a process of preparing a direct-acidified milk beverage keeping high viable cell count at ambient temperature comprising performing direct acidification by adjusting the pH to 4.0-4.5 to obtain acidified

20 milk beverage, sterilizing, and adding *Lactobacillus rhamnosus* ATCC 53103 together with 0.01-0.3 % growth promoting factors, e.g. a carbohydrate, to the mixed milk beverage at aseptic conditions.

- There is a need for developing improved Post Pasteurization Yogurt products
- 25 containing viable lactic acid bacteria.

SUMMARY OF THE INVENTION

- In one aspect the present invention relates to a process for producing an
- 30 ambient storage food product comprising providing a food product with a pH of between 3.4 and 4.4, subjecting the food product to a heat treatment so as to reduce the level of bacteria to no more than 1×10^2 CFU per g to obtain a heat treated food product, adding aseptically to the heat treated food product one or more of ambient storage lactic acid bacteria strains in a total amount of
- 35 at least 1.0×10^3 CFU per g to obtain an ambient storage food product, and

storing the ambient storage food product at ambient temperature for a period of time,

wherein the ambient storage lactic acid bacteria strain is selected from the group consisting of strains,

- 5 (i) wherein the strain when added in an amount of 2.5×10^7 CFU per g to a fermented milk test product in the form of yogurt obtained by fermentation with a starter culture containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii bulgaricus* at a temperature of 43 °C to a pH of 4.3, which has been heat treated at 75 °C for 30 seconds, is capable of retaining viability in an
- 10 amount of at least 1.0×10^3 CFU/g at the end of a storage period of the test product of 150 days at a temperature of 25 °C, and
- (ii) wherein the pH of the test product at most decreases 0.8 units during the storage period, and
- (iii) wherein the strain is selected from the group consisting of *Lactobacillus*
- 15 *paracasei*, *Lactobacillus rhamnosus*, *Lactobacillus fermentum* and *Lactobacillus delbrueckii* subsp. *bulgaricus*, and mutants and variants thereof.

The present invention is based on the unexpected experimental finding that bacterial strains of *Lactobacillus paracasei* and *Lactobacillus rhamnosus* when

- 20 added to a yogurt product subjected to storage at ambient temperature is capable of retaining viability at a certain level without decreasing the pH level in any significant degree for a period of at least 150 days. This is a surprising finding, since in a milk substrate lactic acid bacteria will grow on the carbohydrate source available while reducing the pH until the pH reaches a level, where the bacteria
- 25 is unable to live, so in general lactic acid bacteria in a milk substrate will either be in a growing pH-reducing state or be dead.

In another aspect the present invention relates to a lactic acid bacteria strain for use in an ambient storage food product, wherein the product has a pH of

- 30 between 3.4 and 4.4, wherein the product contains at least 1.0×10^3 CFU per g of the strain, wherein the ambient storage food product is stored at ambient temperature for a period of time, and
- wherein the lactic acid bacteria strain is selected from the group consisting of strains,

- 35 (i) wherein the strain when added in an amount of 2.5×10^7 CFU per g to a

fermented milk test product in the form of yogurt obtained by fermentation with a starter culture containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii bulgaricus* at a temperature of 43 °C to a pH of 4.3, which has been heat treated at 75 °C for 30 seconds, is capable of retaining viability in an amount of at least 1.0×10^3 CFU/g at the end of a storage period of the test product of 150 days at a temperature of 25 °C, and

(ii) wherein the pH of the test product at most decreases 0.8 units during the storage period, and

(iii) wherein the strain is selected from the group consisting of *Lactobacillus paracasei*, *Lactobacillus rhamnosus*, *Lactobacillus fermentum* and *Lactobacillus delbrueckii* subsp. *bulgaricus*, and mutants and variants thereof.

DETAILED DISCLOSURE OF THE INVENTION

15 Process for producing an ambient storage food product

The present invention relates to a process for producing an ambient storage food product comprising providing a food product with a pH of between 3.4 and 4.4, subjecting the food product to a heat treatment so as to reduce the level of bacteria to no more than 1×10^2 CFU per g to obtain a heat treated food product, adding aseptically to the heat treated food product one or more of ambient storage lactic acid bacteria strains in a total amount of at least 1.0×10^3 CFU per g to obtain an ambient storage food product, and storing the ambient storage food product at ambient temperature for a period of time, wherein the ambient storage lactic acid bacteria strain is selected from the group consisting of strains,

(i) wherein the strain when added in an amount of 2.5×10^7 CFU per g to a fermented milk test product in the form of yogurt obtained by fermentation with a starter culture containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii bulgaricus* at a temperature of 43 °C to a pH of 4.3, which has been heat treated at 75 °C for 30 seconds, is capable of retaining viability in an amount of at least 1.0×10^3 CFU/g at the end of a storage period of the test product of 150 days at a temperature of 25 °C, and

(ii) wherein the pH of the test product at most decreases 0.8 units during the storage period, and

(iii) wherein the strain is selected from the group consisting of *Lactobacillus paracasei*, *Lactobacillus rhamnosus*, *Lactobacillus fermentum* and *Lactobacillus delbrueckii* subsp. *bulgaricus*, and mutants and variants thereof.

- 5 In a preferred embodiment, the ambient storage strain is capable of retaining viability in an amount of at least 1.0×10^3 CFU/g, preferably at least 5.0×10^3 CFU/g, more preferably at least 1.0×10^4 CFU/g, more preferably at least 5.0×10^4 CFU/g and most preferably at least 1.0×10^5 CFU/g at the end of the storage period.

10

Preferably, the pH at most decreases 0.7, preferably 0.6, preferably 0.5, preferably 0.4, preferably 0.3, and most preferably 0.2 during the storage period.

- 15 In a preferred embodiment of the invention the strain when added in an amount of 2.5×10^7 CFU per g to the test product increases to an amount of at least 5.0×10^7 CFU per g, preferably 7.5×10^7 CFU per g, and most preferably 1.0×10^8 CFU per g.

- 20 Preferably, the increase of the amount of cells occurs within 45 days of the addition of the strain to the test product, preferably within 40 days, preferably within 35 days, preferably within 30 days, preferably within 25 days, preferably within 20 days, and most preferably within 15 days. Preferably, the amount of cells reaches a maximum within 45 days of the addition of the strain to the test
25 product, preferably within 40 days, preferably within 35 days, preferably within 30 days, preferably within 25 days, preferably within 20 days, and most preferably within 15 days.

- In a particular embodiment of the invention, the ambient storage food product is
30 stored at ambient temperature for a period of at least one day, preferably at least 2 days, more preferably at least 3 days, more preferably at least 4 days, more preferably at least 5 days, more preferably at least 6 days, more preferably at least 7 days, more preferably at least 8 days, more preferably at least 9 days, and most preferably at least 10 days.

35

In a preferred embodiment of the invention, the *Lactobacillus rhamnosus* strain is a lactose-fermenting strain. The terms "lactose-fermenting" is used in the context of the present invention to characterize a strain which has the ability to use lactose either partially or completely as a source for cell growth or
5 maintaining cell viability.

In a preferred embodiment of the invention, the *Lactobacillus paracasei* strain is selected from the group consisting of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus paracasei* strain CRL 431,
10 CHCC23026 deposited as ATCC 55544, and *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389.

In a preferred embodiment of the invention, the *Lactobacillus rhamnosus* strain is selected from the group consisting of *Lactobacillus rhamnosus* strain
15 CHCC5366 deposited as DSM23035 and *Lactobacillus rhamnosus* LGG[®] deposited as ATCC53103. Preferably, the *Lactobacillus rhamnosus* strain is *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035.

In a preferred embodiment of the invention, the *Lactobacillus delbrueckii* subsp.
20 *bulgaricus* strain is *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910. In a preferred embodiment of the invention, the *Lactobacillus delbrueckii* subsp. *bulgaricus* is a lactose-deficient strain.

In a preferred embodiment of the invention, the *Lactobacillus fermentum* strain
25 is *Lactobacillus fermentum* strain CHCC14591 deposited as DSM32086.

In a particular embodiment of the invention the strain is selected from one of the following groups: The group consisting of *Lactobacillus paracasei*, *Lactobacillus rhamnosus*, and *Lactobacillus fermentum*; the group consisting of
30 *Lactobacillus paracasei*, *Lactobacillus rhamnosus*, and *Lactobacillus delbrueckii* subsp. *bulgaricus*; the group consisting of *Lactobacillus paracasei*, *Lactobacillus fermentum*, and *Lactobacillus delbrueckii* subsp. *bulgaricus*; the group consisting of *Lactobacillus rhamnosus*, *Lactobacillus fermentum*, and *Lactobacillus delbrueckii* subsp. *bulgaricus*.

In a particular embodiment of the invention the strain is selected from one of the following groups: The group consisting of *Lactobacillus paracasei* and *Lactobacillus rhamnosus*; the group consisting of *Lactobacillus paracasei* and *Lactobacillus fermentum*; the group consisting of *Lactobacillus paracasei* and
5 *Lactobacillus delbrueckii* subsp. *bulgaricus*; the group consisting of *Lactobacillus rhamnosus* and *Lactobacillus fermentum*, the group consisting of *Lactobacillus rhamnosus*, and *Lactobacillus delbrueckii* subsp. *bulgaricus*; the group consisting of *Lactobacillus fermentum*, and *Lactobacillus delbrueckii* subsp. *bulgaricus*.

- 10 A particular aspect of the invention relates to a lactic acid bacteria strain, wherein the strain is selected from the group consisting of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544,
15 *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, *Lactobacillus rhamnosus* LGG[®] deposited as ATCC53103, *Lactobacillus fermentum* CHCC14591 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and variants thereof. In particular, the strain is selected from the group consisting of any
20 subset of the seven members of the group indicated in the preceding sentence.

- A particular aspect of the invention relates to a lactic acid bacteria strain, wherein the strain is selected from the group consisting of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572,
25 *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544, *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, *Lactobacillus fermentum* CHCC14591 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and
30 variants thereof. This group does not include *Lactobacillus rhamnosus* LGG[®] deposited as ATCC53103, which is lactose-deficient (unable to ferment lactose).

- A particular aspect of the invention relates to a lactic acid bacteria strain, wherein the strain is selected from the group consisting of *Lactobacillus*
35 *paracasei* strain LC-02, CHCC6272, deposited as DSM16572,

Lactobacillus rhamnosus strain CHCC5366 deposited as DSM23035,
Lactobacillus paracasei strain CRL 431, CHCC23026 deposited as ATCC 55544,
Lactobacillus paracasei strain CHCC5584 deposited as DSM32389, and
Lactobacillus rhamnosus LGG[®] deposited as ATCC53103, and mutants and
5 variants thereof.

A particular aspect of the invention relates to a lactic acid bacteria strain,
wherein the strain is selected from the group consisting of *Lactobacillus*
paracasei strain LC-02, CHCC6272, deposited as DSM16572,
10 *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035,
Lactobacillus paracasei strain CRL 431, CHCC23026 deposited as ATCC 55544,
Lactobacillus paracasei strain CHCC5584 deposited as DSM32389, and
Lactobacillus delbrueckii subsp. *bulgaricus* strain CHCC18944 deposited as
DSM28910, and mutants and variants thereof.

15

A particular aspect of the invention relates to a lactic acid bacteria strain,
wherein the strain is selected from the group consisting of *Lactobacillus*
paracasei strain LC-02, CHCC6272, deposited as DSM16572,
Lactobacillus rhamnosus strain CHCC5366 deposited as DSM23035,
20 *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544,
and *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, and
mutants and variants thereof.

A particular aspect of the invention relates to a lactic acid bacteria strain,
25 wherein the strain is selected from the group consisting of *Lactobacillus*
paracasei strain LC-02, CHCC6272, deposited as DSM16572,
Lactobacillus rhamnosus strain CHCC5366 deposited as DSM23035,
Lactobacillus paracasei strain CRL 431, CHCC23026 deposited as ATCC 55544,
and *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, and
30 mutants and variants thereof. The five strains of this aspect of the invention are
all lactose-positive.

A particular aspect of the invention relates to a lactic acid bacteria strain,
wherein the strain is selected from the group consisting of *Lactobacillus*
35 *paracasei* strain LC-01, CHCC2115, deposited as DSM19465,

Lactobacillus rhamnosus strain CHCC12697 deposited as DSM24616, and
Lactobacillus paracasei strain CHCC7155 deposited as DSM18875.

In a particular embodiment of the process of the invention, the ambient storage
5 strain is lactose-fermenting, i.e. capable of fermenting lactose to form lactic acid
and hence lower the pH of the food product. It is surprising that the lactose-
fermenting strains of the present invention is suitable as strains for adding to
ambient storage food products, since it will be expected that such strains during
long term storage of the food product at ambient temperature will cause an
10 undesired reduction of the pH of the food product.

In a particular embodiment of the process of the invention, the food product
with a pH of between 3.4 and 4.4 is a starter culture fermented milk product
provided by fermentation of a milk substrate using a starter culture of lactic acid
15 bacteria to obtain the starter culture fermented milk product.

In a particular embodiment of the process of the invention, the starter culture
fermented milk product has a protein content of more than 5.1 % (w/w).

20 In a particular embodiment of the process of the invention, the starter culture
fermented milk product is not subjected to diluting.

It is surprising that the lactose-fermenting strains of the present invention is
suitable as strains for adding to a ambient storage food product in the form of
25 an undiluted starter culture fermented milk product, because such undiluted
product has a higher concentration of lactose than a diluted product.

In the following the process of the invention is described in more detail in
relation to the process for producing an ambient storage fermented milk
30 product.

The starter culture may be any conventional starter culture of lactic acid
bacteria, including single strain culture and culture blends, used for producing a
specific type of fermented milk product. In a preferred embodiment of the above
35 process of the invention, the fermentation is carried out so as to obtain a pH of

between 3.0 and 5.0, preferably between 3.9 and 4.8, more preferably between 4.0 and 4.6 and most preferably between 4.1 and 4.4.

The heat treatment so as to reduce the level of bacteria of the starter culture to
5 no more than 1.0×10^2 CFU per g fermented milk is preferably carried out
by subjecting the starter culture fermented milk product to a temperature of
between 50 °C and 90 °C, preferably between 60 °C and 85 °C, more preferably
between 65 °C and 82 °C, and most preferably between 70 °C and 80 °C. The
heat treatment is preferably carried out for a period of between 10 seconds and
10 180 seconds, preferably between 12 seconds and 120 seconds, more preferably
between 14 seconds and 90 seconds, more preferably between 16 seconds and
60 seconds, more preferably between 18 seconds and 50 seconds and most
preferably between 20 and 40 seconds. Preferably, the level of bacteria of the
starter culture is reduced to no more than 1.0×10^1 CFU per g fermented
15 milk, more preferably 0 CFU per g.

Lactic acid bacteria strain for use in an ambient storage food product

In one aspect the present invention relates to a lactic acid bacteria strain for use
20 in an ambient storage food product, wherein the product has a pH of between
3.4 and 4.4, wherein the product contains at least 1.0×10^3 CFU per g of the
strain, wherein the ambient storage food product is stored at ambient
temperature for a period of time, and
wherein the lactic acid bacteria strain is selected from the group consisting of
25 strains,

(i) wherein the strain when added in an amount of 2.5×10^7 CFU per g to a
fermented milk test product in the form of yogurt obtained by fermentation with
a starter culture containing *Streptococcus thermophilus* and *Lactobacillus*
delbrueckii bulgaricus at a temperature of 43 °C to a pH of 4.3, which has been
30 heat treated at 75 °C for 30 seconds, is capable of retaining viability in an
amount of at least 1.0×10^3 CFU/g at the end of a storage period of the test
product of 150 days at a temperature of 25 °C, and

(ii) wherein the pH of the test product at most decreases 0.8 units during the
storage period, and
35 (iii) wherein the strain is selected from the group consisting of *Lactobacillus*

paracasei, *Lactobacillus rhamnosus*, *Lactobacillus fermentum* and *Lactobacillus delbrueckii* subsp. *bulgaricus*, and mutants and variants thereof.

In a particular embodiment of the strain of the present invention, the product is
5 a chemically acidified product.

In a particular embodiment of the strain of the present invention, the product is a fermented milk product obtained by fermentation of a milk substrate using a starter culture of lactic acid bacteria, wherein the product contains no more than
10 1×10^2 CFU of the starter culture per g and at least 1×10^3 CFU per g of the ambient storage lactic acid bacteria strain.

In a particular embodiment of the strain of the present invention, the product is a fermented milk product obtained by fermentation of a milk substrate using a
15 starter culture of lactic acid bacteria, wherein the fermented milk product after fermentation has been subjected to a heat treatment so as to reduce the level of bacteria of the starter culture to no more than 1×10^2 CFU per g, and wherein after the heat treatment the ambient storage strain of claim 1 has been added aseptically to the heat treated product in an amount of at least
20 1.0×10^3 CFU per g. Preferably, the ambient storage strain of the invention has been added aseptically to the heat treated product in an amount of at least 1.0×10^4 CFU per g, more preferably at least 1.0×10^5 CFU per g, more preferably at least 1.0×10^6 CFU per g, more preferably at least 1.0×10^7 CFU per g, and most preferably at least 1.0×10^8 CFU per g.
25

In a particular embodiment of the strain of the present invention, the strain is selected from the group consisting of of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026
30 deposited as ATCC 55544, *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, *Lactobacillus rhamnosus* LGG[®] deposited as ATCC53103, *Lactobacillus fermentum* CHCC14591 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and variants thereof.

In a particular embodiment of the strain of the present invention, the strain is selected from the group consisting of of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026
5 deposited as ATCC 55544, *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, *Lactobacillus fermentum* CHCC14591 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and variants thereof.

- 10 The strain of the invention may be formulated in a composition comprising one or more of the strain according to the invention. Thus, in one embodiment the strain of the invention is formulated in a composition containing a single strain according to the invention. In another embodiment the strain is formulated in a composition containing two or more strains of the invention. The composition
15 may be in the form of freeze-dried or frozen granules.

Ambient storage food product

- In a preferred embodiment of the invention, the ambient storage food product is
20 selected from the group consisting of fermented milk products, chemically acidified milk products, fruit beverage, fermented cereal products, chemically acidified cereal products, soy milk products and any mixture thereof.

- The fermented milk product typically contains protein in a level of between 2.0
25 % by weight to 3.5 % by weight. The fermented milk product may also be a low protein product with a protein level of between 1.0 % by weight and 2.0 % by weight. Alternatively, the fermented milk product may be a high protein product with a protein level of above 3.5 % by weight, preferably above 5.1 % by weight. In a particular embodiment of the fermented milk product of the
30 invention the product is a mixture of a fermented milk product and a cereal product, e.g. an oat product, wherein the cereal product may be a fermented cereal product, e.g. a fermented oat product.

- In a particular embodiment of the invention, the ambient storage food product is
35 a fermented cereal product. The fermented cereal product may be prepared by

milling the grains of a cereal biological source material to produce a cereal flour, which is then subjected to fermentation. The fermentation of the cereal flour may be carried out using the same lactic acid bacteria (starter culture) as used for fermentation of a milk substrate as described elsewhere in this application.

5

In a particular embodiment of the invention, the ambient storage food product is a fruit beverage. The fruit beverages may further contain e.g. oat, soy, almond, whey and/or non-fermented milk, e.g. in the form of milk powder. In a particular embodiment, the fruit beverages of the invention do not contain dairy

10 components, such as milk. In another particular embodiment of the fruit beverage of the invention, the fruit beverage further contains a fermented milk product.

In another embodiment of the invention the ambient storage food product of the
15 invention is a chemically acidified product. The acidification may be carried out using any acidifying agent suitable for adding to food products, such as lactic acid, citric acid, fruit juice, fruit pulp and fruit compound. In a particular embodiment, the ambient storage food product is milk acidified with fruit juice.

20 In a particular embodiment of the invention, the ambient storage food product is a chemically acidified cereal product. The chemically acidified cereal product may be prepared by milling the grains of a cereal biological source material to produce a cereal flour, which is then used to produce an aqueous suspension, and the pH of the said suspension is then adjusted to a desired level. In a
25 particular embodiment, the ambient storage food product is a cereal food product acidified with a fruit beverage.

Any combination of the above-described elements, aspects and embodiments in all possible variations thereof is encompassed by the invention unless otherwise
30 indicated herein or otherwise clearly contradicted by context.

DEFINITIONS

In connection with the present invention the terms and expressions listed below
35 have the following meaning:

The expression "heat treatment" means any treatment using any temperature, for any period of time and by any means or equipment, which inactivates at least a portion of the bacteria of the starter culture. In this connection the term
5 "inactivate" means any stop, reduction or inhibition of growth of the bacteria, e.g. cell lysing.

The expression "ambient storage" means storage at ambient temperature. The expression "ambient temperature" means the temperature of the surroundings,
10 e.g. room temperature. For example, the ambient temperature may be between 5 °C and 40 °C, more particularly between 10 °C and 35 °C, more particularly between 15 °C and 30 °C, and most particularly between 18 °C and 27 °C. The ambient temperature may be controlled, i.e. the temperature is the same over the course of one full day (24 hours), or it may be uncontrolled, i.e. it varies
15 over the course of one full day (24 hours).

The expression "viability" means the bacteria is capable of exhibiting growth (forming a colony) on an MRS agar plate incubated at anaerobic conditions at 37 °C for 3 days. The MRS agar has the following composition (g/l):
20 Peptone: 10.0
Beef extract: 10.0
Yeast extract: 5.0
Dextrose: 20.0
Polysorbate 80: 1.0
25 Ammonium Citrate: 2.0
Sodium Acetate: 5.0
Magnesium Sulfate: 0.1
Manganese Sulfate: 0.05
Dipotassium Phosphate: 2.0
30 Agar: 15.0

The expression "ambient storage lactic acid bacteria strain" means a lactic acid bacteria strain, which when added to a fermented milk product is suitable for ambient storage for a period of time.

The expression "starter culture fermented milk product" means a fermented milk product, which contains the starter culture used to ferment the milk.

The expression "heat treated fermented milk product" means a fermented milk
5 product, which has been subjected to heat treatment.

The expression "ambient storage fermented milk product" means a fermented milk product, which is suitable for ambient storage for a period of time.

- 10 The expression "lactic acid bacteria" designates a gram-positive, microaerophilic or anaerobic bacterium, which ferments sugars with the production of acids including lactic acid as the predominantly produced acid, acetic acid and propionic acid. The industrially most useful lactic acid bacteria are found within the order "Lactobacillales" which includes *Lactococcus spp.*, *Streptococcus spp.*,
15 *Lactobacillus spp.*, *Leuconostoc spp.*, *Pseudoleuconostoc spp.*, *Pediococcus spp.*, *Brevibacterium spp.*, *Enterococcus spp.* and *Propionibacterium spp.* These are frequently used as food cultures alone or in combination with other lactic acid bacteria.
- 20 Lactic acid bacteria, including bacteria of the species *Lactobacillus sp.* and *Lactococcus sp.*, are normally supplied to the dairy industry either as frozen or freeze-dried cultures for bulk starter propagation or as so-called "Direct Vat Set" (DVS) cultures, intended for direct inoculation into a fermentation vessel or vat for the production of a dairy product, such as a fermented milk product or a
25 cheese. Such lactic acid bacterial cultures are in general referred to as "starter cultures" or "starters".

The term "milk" is to be understood as the lacteal secretion obtained by milking of any mammal, such as cows, sheep, goats, buffaloes or camels. In a preferred
30 embodiment, the milk is cow's milk. The term milk also includes protein/fat solutions made of plant materials, e.g. soy milk.

The term "milk substrate" may be any raw and/or processed milk material that can be subjected to fermentation according to the method of the invention.
35 Thus, useful milk substrates include, but are not limited to, solutions/-

suspensions of any milk or milk like products comprising protein, such as whole or low fat milk, skim milk, buttermilk, reconstituted milk powder, condensed milk, dried milk, whey, whey permeate, lactose, mother liquid from crystallization of lactose, whey protein concentrate, or cream. Obviously, the
5 milk substrate may originate from any mammal, e.g. being substantially pure mammalian milk, or reconstituted milk powder.

Prior to fermentation, the milk substrate may be homogenized and pasteurized according to methods known in the art.

10

"Homogenizing" as used herein means intensive mixing to obtain a soluble suspension or emulsion. If homogenization is performed prior to fermentation, it may be performed so as to break up the milk fat into smaller sizes so that it no longer separates from the milk. This may be accomplished by forcing the milk at
15 high pressure through small orifices.

"Pasteurizing" as used herein means treatment of the milk substrate to reduce or eliminate the presence of live organisms, such as microorganisms. Preferably, pasteurization is attained by maintaining a specified temperature for a specified
20 period of time. The specified temperature is usually attained by heating. The temperature and duration may be selected in order to kill or inactivate certain bacteria, such as harmful bacteria. A rapid cooling step may follow.

"Fermentation" in the methods of the present invention means the conversion of
25 carbohydrates into alcohols or acids through the action of a microorganism. Preferably, fermentation in the methods of the invention comprises conversion of lactose to lactic acid.

Fermentation processes to be used in production of dairy products are well
30 known and the person of skill in the art will know how to select suitable process conditions, such as temperature, oxygen, amount and characteristics of microorganism(s) and process time. Obviously, fermentation conditions are selected so as to support the achievement of the present invention, i.e. to obtain a dairy product in solid (such as a cheese) or liquid form (such as a fermented
35 milk product).

In the present context, the term "mutant" should be understood as a strain derived from a strain of the invention by means of e.g. genetic engineering, radiation and/or chemical treatment, and/or selection, adaptation, screening, etc. It is preferred that the mutant is a functionally equivalent mutant, e.g. a mutant that has substantially the same, or improved, properties with respect to suitability for ambient storage as the mother strain. Such a mutant is a part of the present invention. Especially, the term "mutant" refers to a strain obtained by subjecting a strain of the invention to any conventionally used mutagenization treatment including treatment with a chemical mutagen such as ethane methane sulphonate (EMS) or N-methyl-N'-nitro-N-nitroguanidine (NTG), UV light or to a spontaneously occurring mutant. A mutant may have been subjected to several mutagenization treatments (a single treatment should be understood one mutagenization step followed by a screening/selection step), but it is presently preferred that no more than 20, no more than 10, or no more than 5, treatments are carried out. In a presently preferred mutant, less than 1%, or less than 0.1%, less than 0.01%, less than 0.001% or even less than 0.0001% of the nucleotides in the bacterial genome have been changed (such as by replacement, insertion, deletion or a combination thereof) compared to the mother strain.

In the present context, the term "variant" should be understood as a strain which is functionally equivalent to a strain of the invention, e.g. having substantially the same, or improved, properties with respect to suitability for ambient storage as the mother strain. Such variants, which may be identified using appropriate screening techniques, are a part of the present invention.

Preferably, the "mutant" and "variant" of the strain of the present invention has the property that when added in an amount of 2.5×10^7 CFU per g to a fermented milk test product in the form of yogurt obtained by fermentation with a starter culture containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii bulgaricus* at a temperature of 43 °C to a pH of 4.3, which has been heat treated at 75 °C for 30 seconds, the strain is capable of retaining viability in an amount of at least 1.0×10^3 CFU/g at the end of a storage period of the test product of 150 days at a temperature of 25 °C, and wherein the pH at most

decreases 0.8 units during the storage period.

Preferably, the "mutant" and "variant" of the strain of the present invention has less than 25, more preferably less than 10, more preferably less than 9, more
5 preferably less than 8, more preferably less than 7, more preferably less than 6, more preferably less than 5, more preferably less than 4, more preferably less than 3, more preferably less than 2 mutations in the amino acid sequence of one or more of the proteins of the strain. In this connection the term "mutation" means a mutation selected from the group consisting of a substitution, a
10 deletion and an insertion.

The use of the terms "a" and "an" and "the" and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise
15 indicated herein or clearly contradicted by context. The terms "comprising", "having", "including" and "containing" are to be construed as open-ended terms (i.e., meaning "including, but not limited to,") unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless
20 otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., "such as") provided herein, is intended merely to
25 better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

30 The expression "fermented milk product" means a food or feed product wherein the preparation of the food or feed product involves fermentation of a milk substrate with a lactic acid bacteria. "Fermented milk product" as used herein includes but is not limited to products such as thermophilic fermented milk products, e.g. yogurt, mesophilic fermented milk products, e.g. sour cream and
35 buttermilk, cheese as well as fermented whey.

The term "thermophile" herein refers to microorganisms that thrive best at temperatures above 43°C. The industrially most useful thermophilic bacteria include *Streptococcus* spp. and *Lactobacillus* spp. The term "thermophilic fermentation" herein refers to fermentation at a temperature above about 35°C, such as between about 35°C to about 45°C. The term "thermophilic fermented milk product" refers to fermented milk products prepared by thermophilic fermentation of a thermophilic starter culture and include such fermented milk products as set-yogurt, stirred-yogurt and drinking yogurt, e.g. Yakult.

10

The term "mesophile" herein refers to microorganisms that thrive best at moderate temperatures (15°C-40°C). The industrially most useful mesophilic bacteria include *Lactococcus* spp. and *Leuconostoc* spp. The term "mesophilic fermentation" herein refers to fermentation at a temperature between about 22°C and about 35°C. The term "mesophilic fermented milk product" refers to fermented milk products prepared by mesophilic fermentation of a mesophilic starter culture and include such fermented milk products as buttermilk, sour milk, cultured milk, smetana, sour cream, Kefir and fresh cheese, such as quark, tvarog and cream cheese.

20

The term "cheese" is understood to encompass any cheese, including hard, semi-hard and soft cheeses, such as cheeses of the following types: Cottage, Feta, Cheddar, Parmesan, Mozzarella, Emmentaler, Danbo, Gouda, Edam, Feta-type, blue cheeses, brine cheeses, Camembert and Brie. The person skilled in the art knows how to convert the coagulum into cheese, methods can be found in the literature, see e.g. Kosikowski, F. V., and V. V. Mistry, "Cheese and Fermented Milk Foods", 1997, 3rd Ed. F. V. Kosikowski, L. L. C. Westport, CT. As used herein, a cheese which has a NaCl concentration below 1.7% (w/w) is referred to as a "low-salt cheese".

30

In the present context the term "fruit juice" refers to the liquid naturally contained in fruit prepared by mechanically squeezing or macerating fresh fruits without the presence of heat and solvents. The "fruit juice" may consist of juice from one type of fruit or a mixture of more than one type of fruit.

35

The term "fruit drink" in the present context refers to a beverage having a fruit juice content of between 0 to 29%.

The term "nectar" in the present context refers to a beverage having a fruit juice
5 content of between 30 to 99% fruit juice.

In the present context the term "puree" refers to fruits prepared by grounding, pressing and/or straining into the consistency of a thick liquid or a soft paste without the presence of heat and solvents. "Puree" is made of 100% fruit as
10 opposed to being made from just the juice of the fruit.

In the present context the term "fruit beverage" refers to a beverage comprising fruit juice, fruit concentrate and/or fruit puree. The term "fruit beverage" covers "fruit juice", "fruit drink" and "nectar" as defined herein. The "fruit beverage"
15 may be either one containing pulp, or one from which the pulp has been removed by such an operation as centrifugation.

The term "adding aseptically" means without introducing or introducing a minimum of any microorganism other than the ambient storage lactic acid
20 bacteria.

The term "cereal product" means any product obtained from a cereal or grain biological source material, including oat, corn, barley, rye, buckwheat, wheat and rice.

25

The term "lactose-deficient" is used in the context of the present invention to characterize LAB which either partially or completely lost the ability to use lactose as a source for cell growth or maintaining cell viability. Such LAB are capable of metabolizing one or several carbohydrates selected from sucrose,
30 galactose and/or glucose or another fermentable carbohydrate. Since these carbohydrates are not naturally present in milk in sufficient amounts to support fermentation by lactose deficient mutants, it will be necessary to add these carbohydrates to the milk. Lactose deficient and partially deficient LAB can be characterized as white colonies on a medium containing lactose and X-Gal.

35

The expression "X.Xx10expYY" and "X.XEYY", both mean $X.X \times 10^{YY}$, and the two said expressions are used interchangeably.

The expression "CFU" means Colony Forming Units.

5

SPECIFIC ITEMS OF THE INVENTION

1. An ambient storage lactic acid bacteria strain, wherein the strain when added in an amount of 2.5×10^7 CFU per g to a fermented milk test product in the form of yogurt obtained by fermentation with a starter culture containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii bulgaricus* at a temperature of 43 °C to a pH of 4.3, which has been heat treated at 75 °C for 30 seconds, is capable of retaining viability in an amount of at least 1.0×10^3 CFU/g at the end of a storage period of the test product of 150 days at a temperature of 25 °C, and wherein the pH at most decreases 0.8 units during the storage period, and wherein the strain is selected from the group consisting of *Lactobacillus paracasei*, *Lactobacillus rhamnosus*, *Lactobacillus fermentum* and *Lactobacillus delbrueckii* subsp. *bulgaricus*, and mutants and variants thereof.

20

2. Strain according to claim 1, wherein the strain is capable of retaining viability in an amount of at least 1.0×10^3 CFU/g, preferably at least 5.0×10^3 CFU/g, more preferably at least 1.0×10^4 CFU/g, more preferably at least 5.0×10^4 CFU/g and most preferably at least 1.0×10^5 CFU/g at the end of the storage period.

25

3. Strain according to claim 1 or 2, wherein the pH at most decreases 0.7, preferably 0.6, preferably 0.5, preferably 0.4, preferably 0.3, and most preferably 0.2 during the storage period.

30

4. Strain according to any of the preceding claims, wherein the strain when added in an amount of 2.5×10^7 CFU per g to the test product increase to an amount of at least 5.0×10^7 CFU per g, preferably 7.5×10^7 CFU per g, and most preferably 1.0×10^8 CFU per g.

35

5. Strain according to claim 4, wherein the increase of the amount of cells occurs within 45 days of the addition of the strain to the test product, preferably within 40 days, preferably within 35 days, preferably within 30 days, preferably within 25 days, preferably within 20 days, and most preferably within 15 days.

5

6. Strain according to claim 4 or 5, wherein the amount of cells reaches a maximum within 45 days of the addition of the strain to the test product, preferably within 40 days, preferably within 35 days, preferably within 30 days, preferably within 25 days, preferably within 20 days, and most preferably within 15 days.

10

7. Composition comprising one or more of the ambient storage lactic acid bacteria strains according to any of claim 1-6.

15 8. Ambient storage food product with a pH of between 3.4 and 4.4, wherein the product contains at least 1.0×10^3 CFU per g of the ambient storage strain of claim 1.

9. Food product according to claim 8, wherein product is a chemically acidified product.

20

10. Food product according to claim 8, wherein the product is a fermented milk product obtained by fermentation of a milk substrate using a starter culture of lactic acid bacteria, wherein the product contains no more than 1×10^2 CFU of the starter culture per g and at least 1×10^3 CFU per g of the ambient storage strains of claim 1.

25

11. Food product according to claim 8, wherein the product is a fermented milk product obtained by fermentation of a milk substrate using a starter culture of lactic acid bacteria, wherein the fermented milk product after fermentation has been subjected to a heat treatment so as to reduce the level of bacteria of the starter culture to no more than 1×10^2 CFU per g, and wherein after the heat treatment the ambient storage strain of claim 1 has been added aseptically to the heat treated product in an amount of at least 1.0×10^3 CFU per g.

35

12. Process for producing an ambient storage food product comprising providing a food product with a pH of between 3.4 and 4.4, subjecting the food product to a heat treatment so as to reduce the level of bacteria to no more than 1×10^2 CFU per g to obtain a heat treated food product, and adding
5 aseptically to the heat treated food product one or more of the ambient storage lactic acid bacteria strains of claim 1 in a total amount of at least 1.0×10^3 CFU per g to obtain an ambient storage food product.

13. Process for producing an ambient storage fermented milk product
10 comprising fermentation of a milk substrate using a starter culture of lactic acid bacteria to obtain a starter culture fermented milk product, subjecting the starter culture fermented milk product to a heat treatment so as to reduce the level of bacteria of the starter culture to no more than 1×10^2 CFU per g to obtain a heat treated fermented milk product, and adding aseptically to the heat
15 treated fermented milk product one or more of the ambient storage lactic acid strains of claim 1 in a total amount of at least 1.0×10^3 CFU per g to obtain an ambient storage fermented milk product.

14. Use of an ambient storage lactic acid bacteria strain according to claim 1 for
20 adding aseptically to a heat treated food product in a total amount of at least 1.0×10^3 CFU per g, wherein the heat treated food product has a pH of between 3.4 and 4.4 and has been subjected to a heat treatment so as to reduce the level of starter culture bacteria to no more than 1×10^2 CFU per g.

25

15. A lactic acid bacteria strain, wherein the strain is selected from the group consisting of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544,
30 *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, *Lactobacillus rhamnosus* LGG® deposited as ATCC53103*, *Lactobacillus fermentum* CHCC14591 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and variants thereof.

35

EXAMPLES

Example 1: Testing of ambient storage suitability of *Lactobacillus***5 *paracasei* and *Lactobacillus rhamnosus***Milk substrate

Fat level 2.8%*

Protein level 2.8%*

10 Lactose 3.0 %

Sucrose 5.0 % (added)

Modified Starch E1442 Cargill type 75720 1.50 %

Pectin type LMA CPKelco type LM 106 AS-YA 0.25 %

Gellan Gum type Kelcogel YSS 0.05 %

15 * Level in final product, i.e. after heat treatment, addition of the ambient storage strain and storage for 150 days.

Starter cultureYoFlex® starter culture type YF-L904 containing the two strains *Streptococcus*

20 *thermophilus* and *Lactobacillus delbrueckii* spp. *bulgaricus*.

Ambient storage strains tested*Lactobacillus paracasei* strain LC-02, CHCC6272 deposited as DSM16572.*Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035.

25 *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544.

Lactobacillus acidophilus La-5, CHCC2169, deposited as DSM13241.*Bifidobacterium animalis* BB-12, CHCC5445, deposited as DSM15954*Lactobacillus helveticus*, (mixture of 4 strains).*Lactococcus lactis* DN-224, CHCC3955, deposited as DSM11037.

30

Procedure for producing test product

1. Dispersing the dry ingredients into the milk

2. Resting for 3 hours with gentle stirring

3. Heating the milk until a temperature of 65°C is reached

35 4. Homogenization at 150 Bar

5. Heat treatment to 95°C for 5 min.
6. Cooling to fermentation temperature 43°C
7. Pump the milk into fermentation vat
8. Inoculation of YoFlex Culture type YF-L904.
- 5 9. Fermentation until pH reaches 4.30.
10. Break the curd and stir until smooth structure is obtained
11. Heat treatment at 75°C for 30 sec.
12. Cooling to 25°C

10 Procedure for testing ambient storage strains

13. Aseptic filling into 100 ml. sterile containers
14. Inoculation of strains & cultures
15. Storage for 6 months at room temperature 25°C

15 Method of cell counting

1. *L. paracasei* (LC-02, CHCC6272), *L. rhamnosus* (CHCC5366), *L. paracasei* (CRL 431, CHCC23026), *L. acidophilus* (La-5, CHCC2169), and *L. helveticus*

Samples from the test product were grown on MRS agar plates incubated at

20 anaerobic conditions at 37 °C for 3 days.

2. *B. animalis*, BB-12 (CHCC5445)

Samples from the test product were grown on MRS agar plates containing 10 % cysteine hydrochloride incubated at anaerobic conditions at 37 °C for 3 days.

25

3. *Lactococcus lactis*, DN-224 (CHCC3955)

Samples from the test product were grown on M-17 agar plates incubated at aerobic conditions at 30 °C for 3 days.

30

Results

Table 1: Cell Counts (cfu/g) for storage at 5 °C

Test strain	Day 4/5	Day 14	Day 35	Day 56
<i>L. paracasei</i> , LC-02, CHCC6272	5.4E07	2.9E07	1.7E07	2.5E07
<i>L. rhamnosus</i> CHCC5366	1.0E07	9.3E06	1.3E07	1.8E07
<i>L. paracasei</i> , CRL 431, CHCC23026	1.6E07	1.6E07	1.3E07	1.5E07
<i>L. acidophilus</i> , La-5, CHCC2169	7.2E06	5.0E04	1.4E04	1.4E03
<i>B. animalis</i> , BB- 12, CHCC5445	1.4E07	8.3E06	4.6E06	1.5E06
<i>L. helveticus</i> Mixture of 4 strains	ND	7.9E06	6.4E05	<1.0E03
<i>Lactococcus</i> <i>lactis</i> , DN-224, CHCC3955	ND	1.0E06	1.1E05	1.0E03

5 ND: No Data

Table 2: Cell Counts (cfu/g) for storage at 25 °C

Test strain	Day 0	Day 1	Day 14	Day 35	Day 56	Day 90	Day 150
<i>L. paracasei</i> LC-02, CHCC6272	7.2E06	1.4E07	3.7E08	3.4E08	2.1E08	5.0E06	5.9E04
<i>L. rhamnosus</i> CHCC5366	2.5E07	4.9E07	9.0E07	1.6E08	4.2E07	2.2E06	5.0E03
<i>L. paracasei</i> , CRL 431, CHCC23026	1.1E07	1.9E07	1.3E08	1.5E08	7.9E07	2.1E07	1.2E05
<i>L. acidophilus</i> La-5, CHCC2169	ND	ND	<1.0E02	<1.0E01	ND	ND	ND
<i>B. animalis</i> BB-12, CHCC5445	ND	ND	4.0E05	<1.0E03	ND	ND	ND
<i>L. helveticus</i> Mixture of 4 strains	ND	ND	3.6E07	9.4E07	1.5E06	<1.0E01	ND
<i>L. lactis</i> , DN-224 CHCC3955	ND	ND	<1.0E03	ND	ND	ND	ND

ND: No Data

Table 3: Cell Counts (cfu/g) for storage at 25 °C

Test yogurt starter culture*	Day 14	Day 35	Day 56	Day 90
TC-X11	Lb: 4.0E02 St: 5.7E04	ND ND	ND ND	ND ND
CH-1	Lb: 2.3E04 St: <1.0E03	Lb: 2.5E02 St: 4.0E01	Lb: 2.5E02 St: 4.0E01	Lb: <10E01 St: <10E01
Acidifix	Lb: 5.0E01 St: <1.0E03	ND ND	ND ND	ND ND
Yoflex Advance 1.2	Lb: 1.6E04 St: 5.0E04	Lb: <1.0E02 St: <1.0E02	ND ND	ND ND
Yoflex Premium 1.0	Lb: 8.0E02 St: 1.0E05	ND ND	ND ND	ND ND
Yoflex Mild 2.0	Lb: 2.0E02 St: <1.0E03	ND ND	ND ND	ND ND

ND: No Data

- 5 All yogurt starter cultures are commercial products from Chr. Hansen A/S in the form of blends of *Streptococcus thermophilus* (St) and *Lactobacillus delbrueckii* subsp. *bulgaricus* (Lb).

For the three strains stored at 25 °C and followed until 150 days, i.e. *L.*

- 10 *paracasei*, LC-02, CHCC6272, *L. rhamnosus* CHCC5366 and *L. paracasei*, CRL 431, CHCC23026, the pH was 3.55 for all three strains.

- As will appear from the results shown in Table 1, after 56 days at a standard refrigerated reference temperature of 5 °C the three strains *L. paracasei*, LC-02, CHCC6272, *L. rhamnosus* CHCC5366 and *L. paracasei*, CRL 431, CHCC23026 maintained a cell count of above 1x10exp07, and *B. animalis*, BB-12, CHCC5445 also maintained a high cell count of above 1x10exp06 (CFU/g), whereas the three strains *L. acidophilus*, La-5, CHCC2169, *L. helveticus* (mixture of 4 strains), and *Lactococcus lactis*, DN-224, CHCC3955 only maintained a cell count
- 20 1.4x10exp03 (CFU/g) or lower.

As will appear from the results shown in Table 2, after 150 days at a temperature of 25 °C (ambient temperature) the three strains *L. paracasei*, LC-02, CHCC6272, *L. rhamnosus* CHCC5366 and *L. paracasei*, CRL 431, CHCC23026 maintained a cell count of 5×10^3 CFU/g or higher. In comparison, for the 4 other strains tested the cell count at ambient temperature dropped quickly to a level of below 1.0×10^3 CFU/g. Thus, the three strains *L. paracasei*, LC-02, CHCC6272, *L. rhamnosus* CHCC5366 and *L. paracasei*, CRL 431, CHCC23026 have superior properties with respect to long term storage at ambient temperature while at the same time maintaining a pH level of 3.55. A pH level of 3.55 is acceptable for a number of fermented milk products. Also, as will appear from Table 2, the cell count has a characteristic temporal profile, wherein it initially increases to reach a maximum at 14 days after which the cell count starts to decrease slowly until 56 days after which the rate of decrease of the cell count starts to increase towards 150 days. With respect to the acidification profile of the samples, it is surprising that it was possible to maintain a pH level of 3.55 in view of the fact that the said 3 strains are all lactose fermenting and that the samples contain lactose.

As will appear from Table 3, the cell counts for all the *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus* strains of the different yogurt starter cultures dropped quickly to low levels.

Example 2: Testing of ambient storage suitability of *Lactobacillus paracasei* and *Lactobacillus delbrueckii* subsp. *bulgaricus*

25

Milk substrate

Fat level 2.7%*

Protein level 2.9%*

Lactose 3.0 %

30 Sucrose 7.0 % (added)

Modified Starch E1442 Cargill type 75720 1.50 %

Pectin type LMA CPKelco type LM 106 AS-YA 0.30 %

Gellan Gum type Kelcogel YSS 0.05 %

* Level in final product, i.e. after heat treatment, addition of the ambient

35 storage strain and storage for 150 days.

Starter culture

YoFlex® starter culture type YF-L904 containing the two strains *Streptococcus thermophilus* and *Lactobacillus delbrueckii* spp. *bulgaricus*.

5

Ambient storage strains tested

Lactobacillus paracasei strain CHCC5584 deposited as DSM32389.

Lactobacillus delbrueckii subsp. *bulgaricus* CHCC18944 deposited as DSM28910.

20 other strains selected from the following species:

10 *Lactobacillus helveticus*

Lactobacillus plantarum

Streptococcus thermophilus

Lactococcus lactis

Pediococcus acidilactici

15

Procedure for producing test product

16. Dispersing the dry ingredients into the milk

17. Resting for 3 hours with gentle stirring

18. Heating the milk until a temperature of 65°C is reached

20 19. Homogenization at 150 Bar

20. Heat treatment to 95°C for 5 min.

21. Cooling to fermentation temperature 43°C

22. Pump the milk into fermentation vat

23. Inoculation of YoFlex Culture type YF-L904.

25 24. Fermentation until pH reaches 4.30.

25. Break the curd and stir until smooth structure is obtained

26. Heat treatment at 75°C for 20 sec.

27. Cooling to 25°C

30 Procedure for testing ambient storage strains

28. Aseptic filling into 100 ml. sterile containers

29. Inoculation of strains & cultures

30. Storage for 6 months at room temperature 25°C

35

Method of cell counting

Cell counting was carried out using an automated liquid handling process on a Hamilton apparatus. The process involves plating the sample to be assayed on a suitable agar substrate layer at a number of sequential dilutions and counting the number of colonies for the first dilution of the sequence for which at least one colony grows.

Results

10

Among the 22 tested strains only 5 strains had a cell count of more than 1.0×10^3 CFU/g after 6 months. Among the said 5 strains with a cell count of more than 1.0×10^3 CFU/g after 6 months, only two samples had a pH above 3.5:

15

Lactobacillus paracasei strain CHCC5584 deposited as DSM32389.

Lactobacillus delbrueckii subsp. *bulgaricus* CHCC18944 deposited as DSM28910.

Table 4: Cell Counts (cfu/g) for storage at 25 °C

Test strain	0 months	2 months	4 months	6 months
<i>L. paracasei</i> CHCC5584	1.00E06	1.00E06	8.00E02	1.00E04
pH	4.23	3.54	3.60	3.51
<i>L. delbrueckii</i> subsp. <i>bulgaricus</i> CHCC18944	3.70E05	7.00E05	1.00E06	5.50E05
pH	4.19	3,97	3.70	3.51

20

Example 3: Testing of ambient storage suitability of three *Lactobacillus* species

Experimental plan

5

The object of the present experiment is to test the effect on the survivability of four *Lactobacillus* strains of the following parameters:

- pH of heat-treated yogurt base: 4.0 and pH 4.3 were tested.
- 10 • CFU dosage level: 10000 CFU and 100000 CFU were tested.
- Level of sucrose in milk base: 0, 3 and 7 % sucrose were tested.

Starter culture

- 15 YoFlex® starter culture type YF-L904 containing the two strains *Streptococcus thermophilus* and *Lactobacillus delbrueckii* spp. *bulgaricus*.

Milk substrate

20 Table 5: Composition of milk substrate

	%	g
3.5 % fat milk	74.3	89160.0
Water	23.45	28140.0
Modified starch - Roquette	1.5	1800
Pectin	0.12	144.0
Gelan gum	0.03	36.0
Whey Protein Concentrate (WPC) 80 – Nutrilac YO-7830	0.60	720.0
	100.0	120000

Procedure for producing test product

1. De-aeration of milk base at 350-400 mbar.
- 25 2. Homogenization at 60 °C at 150 bar.
3. Heat treatment at 95 °C for 5 min. in plate pasteurizer. Cool to 5 °C.

4. Inoculation with culture
5. Fermentation temperature: 43 °C.
6. Final heat treatment: 74 °C for 20 seconds in holding tube.
7. Packing in flow bench into sterile cups.

5

Ambient storage strains tested

Lactobacillus paracasei strain CHCC6272 deposited as DSM16572.

10 *Lactobacillus fermentum* strain CHCC14591 deposited as DSM32084.

Lactobacillus rhamnosus strain LGG[®] deposited as ATCC53103.

Analysis

15 CFU plating by RAPID-CFU: Development (every 2 months).

Carbohydrate and small acid content by HPLC: Every 2 months.

pH measurement by change of color indicator (hue): Development (every 1 month).

20 *CFU determination*

CFU is determined by a procedure involving automated plating using a Hamilton robot.

25 *pH measurement*

The pH was measured by means of a measurement of change in color. As color indicator Bromocresol Purple / Green was used in a 50/50 mixture. The color indicator solution was prepared as follows: A 1000 ml volumetric flask was filled
30 partly with milli-Q water, and 1.00 g Bromocresol Green and 1.0 g Bromocresol Purple was transferred to the flask. 2 ml 4N NaOH was added to the flask, which was then stirred and filled to 1000 ml with milli-Q water.

The color intensity of the samples was read by a flatbed scanner. A change in
35 hue of 1 unit corresponds approximately to a change in pH of 0.05 pH units.

Results

Table 6: CFU counts and pH development during 4 months of storage of heat treated ambient yogurt product (test product) with a start pH of 4.0 and a CFU dosage of 10000

Test strain	0 months			2 months			4 months		
Sucrose (%)	0	3	7	0	3	7	0	3	7
No added test strain	0	0	0	0	0	0	0	0	0
pH	138.9	138.9	138.9	140.9	140.7	140.1	139.0	141.6	140.5
<i>L. paracasei</i> CHCC6272	7.0 E05	4.5 E05	6.5 E05	1.0 E06	1.0 E06	1.0 E06	1.4 E07	3.5 E06	4.0 E06
pH	139.4	139.0	138.8	134.9	134.4	134.9	140.6	140.0	138.9
<i>L. fermentum</i> CHCC14591	1.75 E07	1.65 E07	1.2 E07	5.5 E05	1.0 E06	5.5 E05	7.0 E05	7.0 E05	1.5 E05
pH	139.5	138.8	138.8	139.7	141.3	140.2	140.7	138.2	140.2
<i>L. rhamnosus</i> LGG®	2.55 E06	2.5 E06	8.0 E05	1.0 E06	ND	1.0 E06	1.6 E07	1.6 E07	6.5 E06
pH	139.3	138.8	138.7	131.6	ND	134.9	130.9	121.6	131.6

Table 7: CFU counts and pH development during 4 months of storage of heat treated ambient yogurt product (test product) with a start pH of 4.3 and a CFU dosage of 10000

Test strain	0 months			2 months			4 months		
Sucrose (%)	0	3	7	0	3	7	0	3	7
No added test strain	0	0	0	0	0	0	0	0	0
pH	145.7	145.3	144.7	146.4	147.1	146.6	142.6	145.0	145.3
<i>L. paracasei</i> CHCC6272	1.0 E05	7.0 E04	7.5 E04	1.0 E06	1.0 E06	1.0 E06	1.05 E07	1.45 E07	1.0 E07
pH	146.3	145.6	145.3	137.7	136.6	136.9	147.3	148.1	146.4
<i>L. fermentum</i> CHCC14591	1.55 E06	3.15 E06	4.5 E05	5.05 E05	5.5 E05	1.0 E06	9.5 E04	1.95 E05	3.7 E05
pH	146.3	145.7	145.3	147.3	147.5	147.1	147.2	150.1	149.3
<i>L. rhamnosus</i> LGG®	1.5 E04	3.0 E04	4.5 E04	1.0 E06	1.0 E06	1.0 E06	1.2 E07	1.45 E07	1.0 E07
pH	146.3	145.7	145.1	137.5	136.9	137.5	134.7	133.4	136.2

Table 8: CFU counts and pH development during 4 months of storage of heat treated ambient yogurt product (test product) with a start pH of 4.0 and a CFU dosage of 100000

Test strain	0 months			2 months			4 months		
Sucrose (%)	0	3	7	0	3	7	0	3	7
No added test strain	0	0	0	0	0	0	0	0	0
pH	140.3	139.3	139.3	139.9	140.2	141.0	136.6	138.6	138.5
<i>L. paracasei</i> CHCC6272	5.0 E06	2.0 E06	3.5 E06	1.0 E06	1.0 E06	1.0 E06	1.4 E07	9.0 E06	7.0 E06
pH	140.0	139.8	139.7	134.4	134.1	135.0	136.6	138.6	138.5
<i>L. fermentum</i> CHCC14591	1.5 E07	1.25 E07	8.5 E06	5.5 E05	5.5 E06	5.5 E05	7.0 E05	6.0 E06	6.1 E06
pH	140.7	139.9	139.8	136.8	140.6	140.5	139.7	140.5	140.4
<i>L. rhamnosus</i> LGG®	4.5 E06	2.0 E06	1.1 E06	1.0 E06	1.0 E06	1.0 E06	1.3 E07	1.35 E07	4.0 E06
pH	139.9	139.7	139.5	134.8	135.8	136.8	134.9	135.7	142.6

Table 9: CFU counts and pH development during 4 months of storage of heat treated ambient yogurt product (test product) with a start pH of 4.3 and a CFU dosage of 100000

Test strain	0 months			2 months			4 months		
Sucrose (%)	0	3	7	0	3	7	0	3	7
No added test strain	0	0	0	0	0	0	0	0	0
pH	144.4	144.3	143.9	143.9	143.0	143.0	141.7	141.5	140.0
<i>L. paracasei</i> CHCC6272	8.0 E05	7.5 E05	5.5 E05	1.0 E06	1.0 E06	1.0 E06	1.1 E07	1.1 E07	8.0 E06
pH	144.5	144.1	143.5	135.7	135.2	136.0	141.7	141.5	140.0
<i>L. fermentum</i> CHCC14591	1.0 E06	6.5 E06	1.0 E06	1.0 E06	1.0 E05	1.0 E06	2.5 E04	1.1 E06	3.5 E04
pH	144.4	144.4	143.9	143.3	144.4	143.9	139.5	143.7	145.3
<i>L. rhamnosus</i> LGG®	4.5 E05	6.0 E05	6.5 E05	1.0 E06	1.0 E06	1.0 E06	1.0 E06	2.0 E06	8.0 E05
pH	144.8	144.1	144.1	135.9	136.4	137.5	135.1	135.1	138.0

5

As will appear from Tables 6-9, for all sample combinations of 1) the *Lactobacillus* strain to be tested, 2) pH of heat-treated yogurt base, 3) CFU dosage level and 4) level of sucrose in the heat-treated yogurt, the level of CFU/g was between 2.5E04 and 1.6E07 after 4 months of storage.

10

Comparing yogurt base starting pH: For both levels of CFU dosage and no added sucrose, a yogurt base starting pH of 4.0 results in higher CFU counts after 4 months than a yogurt base starting pH of 4.3 for all tested strains. For both levels of CFU dosage and 3 % added sucrose, a yogurt base starting pH of 4.0

15 results in higher CFU counts after 4 months than a yogurt base starting pH of 4.3 for the *L. rhamnosus* and *L. fermentum* strains. For a CFU dosage of 10000 CFU/g and a level of 7 % added sucrose, a yogurt base starting pH of 4.3 results

in higher CFU counts after 4 months than a yogurt base starting pH of 4.0 for all tested strains.

Comparing CFU dosage levels: For the *L. rhamnosus* strain, the CFU dosage
5 level of 10000 CFU/g for both yogurt base starting pH results in higher CFU
counts after 4 months than a CFU dosage level of 100000 CFU/g for all levels of
added sucrose. For an added sucrose level of 7 % and a yogurt base starting pH
of 4.3, a CFU dosage level of 10000 CFU/g results in higher CFU counts after 4
months than a CFU dosage level of 100000 CFU/g for all tested strains.

10

Comparing the levels of added sucrose: For a CFU dosage of 10000 CFU/g and a
yogurt base starting pH of 4.0, a sucrose level of 0 % results in higher CFU
counts after 4 months than sucrose levels of 3 and 7 % (however, for the *L.*
rhamnosus and *L. fermentum* strains and a sucrose level of 3 % the CFU counts
15 are the same as those for 0%). For a CFU dosage of 100000 CFU/g and a yogurt
base starting pH of 4.3, a sucrose level of 3 % results in higher CFU counts after
4 months than sucrose levels of 0 and 7 %.

In order to achieve a high survivability of *Lactobacillus* bacteria during long term
20 storage of ambient temperature heat-treated yogurt, it is believed that it is
essential to achieve a moderate, balanced growth of bacteria. Thus, if the
bacteria inoculated into the yogurt base have limiting growth conditions, the
bacteria culture will not be able to sustain itself and the cell count will decrease
in an unacceptable extent. In this case, the decrease of the yogurt pH will only
25 be small due to the limited bacteria growth. On the other hand, if the bacteria
inoculated into the yogurt base have growth promoting conditions, the bacteria
culture will enter into a phase of high growth resulting in a strong pH drop,
which will eventually stop of bacteria growth and potentially inactivate or lyse
the bacteria cells. Also, such a strong pH drop is undesired as it might render
30 the taste of the yogurt too acidic.

In conclusion, it is believed that that it is essential to achieve a moderate,
balanced growth of bacteria. A number of factors are important to achieve such
a moderate, balanced growth of bacteria. Among these factors are the
35 parameters tested in this Example as listed above. In particular, the starting pH

of the yogurt base is important, because it influences both the level of bacteria growth and the final pH reached during storage. The CFU dosage level is important because it determines in the peak level of bacteria reached during storage and hence the level of pH drop. Finally, it is common practice to add
5 sucrose to heat-treated, ambient yogurt as a sweetener. The sucrose level is important for the survivability of bacteria during long term storage because it is believed that sucrose will degrade during storage to form glucose and fructose hence forming a carbohydrate source for growth of bacteria and causing increased growth of the bacteria.

10

As will appear from the results of the present Example, both the yogurt base starting pH levels of 4.0 and 4.3 result in high CFU counts of between $2.5E04$ and $1.6E07$ after 4 months of storage. Whether a yogurt base starting pH level of 4.0 or 4.3 results in a higher cell count after 4 months depend on the other
15 factors tested as described in detail above.

As will appear from the results of the present Example, both the CFU dosage levels of 10000 and 100000 CFU/g result in high CFU counts of between $2.5E04$ and $1.6E07$ after 4 months of storage. Whether a CFU dosage level of 10000 or
20 100000 CFU/g results in a higher cell count after 4 months depend on the other factors tested as described in detail above.

As will appear from the results of the present Example, both the levels of added sucrose of 0, 3 and 7 % result in high CFU counts of between $2.5E04$ and
25 $1.6E07$ after 4 months of storage. Whether a level of added sucrose of 0, 3 or 7 % results in a higher cell count after 4 months depend on the other factors tested as described in detail above.

Example 4: Testing of ambient storage suitability of two *Lactobacillus*
30 **species in low pH fermented milk product (Yakult)**

Fermented milk product base

In this Example a commercial Yakult product with a low pH (pH 3.8) was used as a base for adding ambient storage strains for the testing of their survivability,
35 post-acidification and growth during long term storage at ambient temperature.

Ingredients: Water, sugar, syrup, Skimmed Milk Powder, glucose, *Lactobacillus paracasei* (live lactic acid bacteria > 3.0E08 CFU/g), and flavor.

Nutrition value: The commercial product per 100 ml contained 1.2 g protein, 0 g fat and 15.4 g carbohydrate.

5

Ambient storage strains tested

Lactobacillus fermentum strain CHCC14591 deposited as DSM32084.

Lactobacillus rhamnosus strain LGG[®] deposited as ATCC53103.

10 Procedure for producing test product

The commercial Yakult samples were in a first step pasteurized at 74 °C for 20 seconds in a Mini Thermisation Unit (MTU). The pasteurized samples were then filled into sterile 100 ml cups in a LAF bench. The cups were then inoculated with the ambient storage test strains at a level of 1.0E06/ml and stored at 25 °C for

15 3 months.

Results

Table 10

	Month	CFU/ml	pH
<i>L. rhamnosus</i> strain LGG [®]	0	3.70E07	3.55
	1.5	1.08E07	
	3	1.30E06	3.15
<i>L. fermentum</i> strain CHCC14591	0	1.05E07	3.5
	1.5	3.70E07	
	3	9.80E07	3.4

20 As will appear from Table 10, the cell count was very high after 3 months of storage at 25 °C for both strains tested, i.e. it was higher than the inoculated level. For the *L. fermentum* strain CHCC14591 the pH only dropped 0.1 units during 3 months of storage at 25 °C, which must be considered to be a very low level of post-acidification at 25 °C, in particular in view of the fact that the cell

25 count has increased in this period. For the *L. rhamnosus* strain LGG[®], the pH drop during 3 months of storage at 25 °C was somewhat higher, i.e. 0.4 units, which is acceptable for some fermented milk products.

DEPOSITS AND EXPERT SOLUTION

The Applicant requests that a sample of the deposited microorganism should be made available only to an expert approved by the Applicant.

5

The strains of the invention have been deposited previously and have the following accession numbers:

Lactobacillus paracasei strain LC-02, CHCC6272, deposited as DSM16572 on
10 2016-05-10 at Deutsche Sammlung von Mikroorganismen und Zellkulturen
GmbH (DSMZ), Inhoffenstr. 7B, D-38124 Braunschweig.

Lactobacillus rhamnosus strain CHCC5366, deposited as DSM23035 on
2012-02-02 at Deutsche Sammlung von Mikroorganismen und Zellkulturen
15 GmbH (DSMZ), Inhoffenstr. 7B, D-38124 Braunschweig.

Lactobacillus acidophilus La-5, CHCC2169, deposited as DSM13241 on
2003-09-30 at Deutsche Sammlung von Mikroorganismen und Zellkulturen
GmbH Maschroder Weg 1b, D-38124 Braunschweig.

20

Bifidobacterium animalis BB-12, CHCC5445, deposited as DSM15954 on
2003-09-30 at Deutsche Sammlung von Mikroorganismen und Zellkulturen
GmbH Maschroder Weg 1b, D-38124 Braunschweig.

25 *Lactococcus lactis* DN-224, CHCC3955, deposited as DSM11037 on 1996-06-26
at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
Maschroder Weg 1b, D-38124 Braunschweig.

Lactobacillus paracasei strain CHCC5584 deposited as DSM32389 on 2016-11-02
30 at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH,
Inhoffenstr. 7B, D-38124 Braunschweig.

Lactobacillus fermentum strain CHCC14591 deposited as DSM32086 on 2015-
07-16 at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH,
35 Inhoffenstr. 7B, D-38124 Braunschweig.

Lactobacillus delbrueckii subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910 on 2014-06-12 at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Inhoffenstr. 7B, D-38124 Braunschweig.

5

Lactobacillus paracasei strain LC-01, CHCC2115, deposited as DSM19465 on 2007-06-27 at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Inhoffenstr. 7B, D-38124 Braunschweig.

- 10 *Lactobacillus rhamnosus* strain CHCC12697 deposited as DSM24616 on 2011-03-01 at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Inhoffenstr. 7B, D-38124 Braunschweig.

Lactobacillus paracasei strain CHCC7155 deposited as DSM18875 on 2006-12-19

- 15 at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Inhoffenstr. 7B, D-38124 Braunschweig.

The deposits were made according to the Budapest Treaty on the international recognition of the deposit of microorganisms for the purposes of patent

- 20 procedure.

The Applicant requests that a sample of the deposited microorganism should be made available only to an expert approved by the Applicant.

25

CLAIMS

1. Process for producing an ambient storage food product comprising providing a food product with a pH of between 3.4 and 4.4, subjecting the food product to a
5 heat treatment so as to reduce the level of bacteria to no more than 1×10^2 CFU per g to obtain a heat treated food product, adding aseptically to the heat treated food product one or more of ambient storage lactic acid bacteria strains in a total amount of at least 1.0×10^3 CFU per g to obtain an ambient storage food product, and storing the ambient storage food product at ambient
10 temperature for a period of time,
wherein the ambient storage lactic acid bacteria strain is selected from the group consisting of strains,
(i) wherein the strain when added in an amount of 2.5×10^7 CFU per g to a fermented milk test product in the form of yogurt obtained by fermentation with
15 a starter culture containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii bulgaricus* at a temperature of 43 °C to a pH of 4.3, which has been heat treated at 75 °C for 30 seconds, is capable of retaining viability in an amount of at least 1.0×10^3 CFU/g at the end of a storage period of the test product of 150 days at a temperature of 25 °C, and
20 (ii) wherein the pH of the test product at most decreases 0.8 units during the storage period, and
(iii) wherein the strain is selected from the group consisting of *Lactobacillus paracasei*, *Lactobacillus rhamnosus*, *Lactobacillus fermentum* and *Lactobacillus delbrueckii* subsp. *bulgaricus*, and mutants and variants thereof.
25
2. Process according to claim 1, wherein the ambient storage lactic acid bacteria strain is capable of retaining viability in an amount of at least 1.0×10^3 CFU/g, preferably at least 5.0×10^3 CFU/g, more preferably at least 1.0×10^4 CFU/g, more preferably at least 5.0×10^4 CFU/g and most
30 preferably at least 1.0×10^5 CFU/g at the end of the storage period of the test product of 150 days.
3. Process according to claim 1 or 2, wherein the pH at most decreases 0.7, preferably 0.6, preferably 0.5, preferably 0.4, preferably 0.3, and most
35 preferably 0.2 during the storage period of the test product of 150 days.

4. Process according to any of the preceding claims, wherein the strain when added in an amount of 2.5×10^7 CFU per g to the test product increases to an amount of at least 5.0×10^7 CFU per g, preferably 7.5×10^7 CFU per g, and most preferably 1.0×10^8 CFU per g.

5. Process according to claim 4, wherein the increase of the amount of cells occurs within 45 days of the addition of the strain to the test product, preferably within 40 days, preferably within 35 days, preferably within 30 days, preferably within 25 days, preferably within 20 days, and most preferably within 15 days.

6. Process according to claim 4 or 5, wherein the amount of cells reaches a maximum within 45 days of the addition of the strain to the test product, preferably within 40 days, preferably within 35 days, preferably within 30 days, preferably within 25 days, preferably within 20 days, and most preferably within 15 days.

7. Process according to any of claims 1-6, wherein the ambient storage food product is stored at ambient temperature for a period of at least one day, preferably at least 2 days, more preferably at least 3 days, more preferably at least 4 days, more preferably at least 5 days, more preferably at least 6 days, more preferably at least 7 days, more preferably at least 8 days, more preferably at least 9 days, and most preferably at least 10 days.

8. Process according to any of claims 1-7, wherein the ambient storage lactic acid bacteria strain is selected from the group consisting of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544, *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, *Lactobacillus rhamnosus* LGG® deposited as ATCC53103, *Lactobacillus fermentum* CHCC14591 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and variants thereof.

9. Process according to any of claims 1-7, wherein *Lactobacillus rhamnosus* strain is a lactose-fermenting strain.

10. Process according to claim 9, wherein the ambient storage lactic acid
5 bacteria strain is selected from the group consisting of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544, *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, *Lactobacillus fermentum* CHCC14591
10 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and variants thereof.

11. Process according to any of claims 1-10, wherein the food product with a pH of between 3.4 and 4.4 is a starter culture fermented milk product provided by
15 fermentation of a milk substrate using a starter culture of lactic acid bacteria to obtain the starter culture fermented milk product.

12. Process according to any of claims 1-11, wherein the starter culture fermented milk product has a protein content of more than 5.1 % (w/w).
20

13. Process according to any of claims 1-12, wherein the starter culture fermented milk product is not subjected to diluting.

14. Lactic acid bacteria strain for use in an ambient storage food product,
25 wherein the product has a pH of between 3.4 and 4.4, wherein the product contains at least 1.0×10^3 CFU per g of the strain, wherein the ambient storage food product is stored at ambient temperature for a period of time, and wherein the lactic acid bacteria strain is selected from the group consisting of strains,

30 (i) wherein the strain when added in an amount of 2.5×10^7 CFU per g to a fermented milk test product in the form of yogurt obtained by fermentation with a starter culture containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii bulgaricus* at a temperature of 43 °C to a pH of 4.3, which has been heat treated at 75 °C for 30 seconds, is capable of retaining viability in an
35 amount of at least 1.0×10^3 CFU/g at the end of a storage period of the test

product of 150 days at a temperature of 25 °C, and

(ii) wherein the pH of the test product at most decreases 0.8 units during the storage period, and

(iii) wherein the strain is selected from the group consisting of *Lactobacillus*
5 *paracasei*, *Lactobacillus rhamnosus*, *Lactobacillus fermentum* and *Lactobacillus*
delbrueckii subsp. *bulgaricus*, and mutants and variants thereof.

15. Strain according to claim 14, wherein the product is a chemically acidified product.

10

16. Strain according to claim 14, wherein the product is a fermented milk product obtained by fermentation of a milk substrate using a starter culture of lactic acid bacteria, wherein the product contains no more than 1×10^2 CFU of the starter culture per g and at least 1×10^3 CFU per g of the ambient

15 storage lactic acid bacteria strain.

17. Strain according to claim 14, wherein the product is a fermented milk product obtained by fermentation of a milk substrate using a starter culture of lactic acid bacteria, wherein the fermented milk product after fermentation has
20 been subjected to a heat treatment so as to reduce the level of bacteria of the starter culture to no more than 1×10^2 CFU per g, and wherein after the heat treatment the ambient storage strain of claim 1 has been added aseptically to the heat treated product in an amount of at least 1.0×10^3 CFU per g.

25 18. Strain according to any of claims 14-17, wherein the strain is selected from the group consisting of of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544, *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389,
30 *Lactobacillus rhamnosus* LGG® deposited as ATCC53103, *Lactobacillus fermentum* CHCC14591 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and variants thereof.

35 19. Strain according to any of claims 14-17, wherein the strain is selected from

the group consisting of of *Lactobacillus paracasei* strain LC-02, CHCC6272, deposited as DSM16572, *Lactobacillus rhamnosus* strain CHCC5366 deposited as DSM23035, *Lactobacillus paracasei* strain CRL 431, CHCC23026 deposited as ATCC 55544, *Lactobacillus paracasei* strain CHCC5584 deposited as DSM32389, 5 *Lactobacillus fermentum* CHCC14591 deposited as DSM32086, and *Lactobacillus delbrueckii* subsp. *bulgaricus* strain CHCC18944 deposited as DSM28910, and mutants and variants thereof.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/061270

A. CLASSIFICATION OF SUBJECT MATTER
INV. A23L33/135 A23C9/12
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A23L A23C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2010/009034 A1 (LING HAIBO [CN]) 14 January 2010 (2010-01-14)	1-8, 11-18
Y	the whole document	9,10
X	US 2015/079057 A1 (HORNBAEK TINA [DK] ET AL) 19 March 2015 (2015-03-19)	14-19
Y	abstract the whole document	1-13
X	US 2013/195917 A1 (ESKESEN DORTE [DK]) 1 August 2013 (2013-08-01)	14-19
Y	the whole document	1-13
X	US 2011/150852 A1 (CHAMBAUD ISABELLE [FR] ET AL) 23 June 2011 (2011-06-23)	14-19
Y	Entire document, in particular para. [0035] and Table II	1-13
	----- -/--	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

19 July 2017

Date of mailing of the international search report

27/07/2017

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Florey, Marianne

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2017/061270

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2015/193459 A1 (CHR HANSEN AS [DK]) 23 December 2015 (2015-12-23)	14-19
Y	the whole document -----	1-13
X,P	WO 2017/037052 A1 (CHR HANSEN AS [DK]) 9 March 2017 (2017-03-09)	14-19
Y,P	the whole document -----	1-13
X	José Manuel Barroso: "DURCHFÜHRUNGSVERORDNUNG (EU) Nr. 451/2012 DER KOMMISSION", Amtsblatt der Europäischen Union, 30 May 2012 (2012-05-30), pages 55-63, XP055310236, Retrieved from the Internet: URL: http://www.feedcheck.de/Verordnungen/Verordnungen2012/Texte/Menuepunkte/451-2012.pdf [retrieved on 2016-10-13]	14-19
Y	page L 140/60, line 1 -----	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2017/061270

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
US 2010009034	A1	14-01-2010	CN	101273736	A	01-10-2008
			CN	101273737	A	01-10-2008
			CN	101273738	A	01-10-2008
			US	2010009034	A1	14-01-2010
			US	2010015285	A1	21-01-2010
			WO	2008116355	A1	02-10-2008
			WO	2008116356	A1	02-10-2008

US 2015079057	A1	19-03-2015	AR	090631	A1	26-11-2014
			CN	104284975	A	14-01-2015
			DK	2836587	T3	08-05-2017
			EA	201491847	A1	29-05-2015
			EP	2836587	A1	18-02-2015
			HK	1206391	A1	08-01-2016
			JP	2015515273	A	28-05-2015
			KR	20140148479	A	31-12-2014
			US	2015079057	A1	19-03-2015
			WO	2013153074	A1	17-10-2013

US 2013195917	A1	01-08-2013	CN	103220921	A	24-07-2013
			CN	105288616	A	03-02-2016
			DK	2627198	T3	28-09-2015
			EP	2627198	A1	21-08-2013
			HK	1182902	A1	15-01-2016
			KR	20130124948	A	15-11-2013
			US	2013195917	A1	01-08-2013
			WO	2012049301	A1	19-04-2012

US 2011150852	A1	23-06-2011	AR	071330	A1	09-06-2010
			BR	PI0911420	A2	04-08-2015
			CA	2721704	A1	29-10-2009
			CN	102066550	A	18-05-2011
			EP	2271744	A2	12-01-2011
			ES	2410263	T3	01-07-2013
			FR	2930264	A1	23-10-2009
			JP	5649564	B2	07-01-2015
			JP	2011517570	A	16-06-2011
			KR	20110009152	A	27-01-2011
			RU	2010146956	A	27-05-2012
			US	2011150852	A1	23-06-2011
			WO	2009130423	A2	29-10-2009
			ZA	201007510	B	31-08-2011

WO 2015193459	A1	23-12-2015	AU	2015276091	A1	15-12-2016
			CN	106714565	A	24-05-2017
			EA	201692283	A1	30-06-2017
			US	2017135360	A1	18-05-2017
			WO	2015193459	A1	23-12-2015

WO 2017037052	A1	09-03-2017	NONE			
