



US008486000B2

(12) **United States Patent**
Coroneo

(10) **Patent No.:** **US 8,486,000 B2**
(45) **Date of Patent:** **Jul. 16, 2013**

(54) **OCULAR PRESSURE REGULATION**

(75) Inventor: **Minas Theodore Coroneo**, Sydney (AU)

(73) Assignee: **Transcend Medical, Inc.**, Menlo Park,
CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/579,330**

(22) PCT Filed: **Nov. 12, 2004**

(86) PCT No.: **PCT/AU2004/001566**

§ 371 (c)(1),
(2), (4) Date: **Aug. 8, 2006**

(87) PCT Pub. No.: **WO2005/046782**

PCT Pub. Date: **May 26, 2005**

(65) **Prior Publication Data**

US 2007/0088242 A1 Apr. 19, 2007

(51) **Int. Cl.**
A61M 5/00 (2006.01)

(52) **U.S. Cl.**
USPC **604/8; 604/9**

(58) **Field of Classification Search**
USPC **604/8–10, 521**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,767,759 A 10/1973 Wichterle
3,788,327 A * 1/1974 Donowitz et al. **604/247**
3,915,172 A 10/1975 Wichterle

4,037,604 A 7/1977 Newkirk
4,402,681 A 9/1983 Haas et al.
4,457,757 A 7/1984 Moltano **604/294**
4,521,210 A 6/1985 Wong
4,554,918 A 11/1985 White **604/10**
4,604,087 A 8/1986 Joseph
4,634,418 A 1/1987 Binder **604/8**

(Continued)

FOREIGN PATENT DOCUMENTS

EP 0228185 7/1987
EP 1184010 3/2002

(Continued)

OTHER PUBLICATIONS

Classen, L., et al., "A histopathologic and immunohistochemical
analysis of the filtration bleb after unsuccessful glaucoma seton
implantation," American Journal of Ophthalmology, 122:205-212,
(1996).

(Continued)


Primary Examiner — Philip R Wiest

(74) *Attorney, Agent, or Firm* — Fred C. Hernandez; Mintz,
Levin, Cohn, Ferris, Glovsky and Popeo, P.C.

(57) **ABSTRACT**

This invention comprises a flexible ocular device for implan-
tation into the eye formed of a biocompatible elastomeric
material, foldable to a diameter of 1.5 mm or less, comprising
a fluid drainage tube having at one end a foldable plate
adapted to located the device on the inner surface of the sclera
in a suprachoroidal space formed by cyclodialysis, said drain-
age tube opening on to the disc at one end and opening to the
anterior chamber when implanted into the eye at its other end,
so as to provide aqueous pressure regulation. Also provided
are methods for the treatment of glaucoma utilizing the flex-
ible ocular device, and an ocular pressure spike shunt.

4 Claims, 3 Drawing Sheets



US 8,486,000 B2

Page 2

U.S. PATENT DOCUMENTS

4,722,724 A	2/1988	Schocket		6,626,858 B2	9/2003	Lynch et al.	
4,750,901 A	6/1988	Molteno		6,638,239 B1	10/2003	Bergheim et al.	
4,787,885 A	11/1988	Binder	604/8	6,648,283 B2	11/2003	Chase et al.	
4,826,478 A	5/1989	Schocket	604/8	6,666,841 B2	12/2003	Gharib et al.	
4,846,172 A	7/1989	Berlin		6,676,607 B2	1/2004	de Juan, Jr. et al.	
4,863,457 A	9/1989	Lee		6,699,210 B2	3/2004	Williams et al.	604/8
4,886,488 A	12/1989	White	604/9	6,699,211 B2	3/2004	Savage	604/9
4,900,300 A	2/1990	Lee		6,719,750 B2	4/2004	Varner et al.	604/289
4,946,436 A	8/1990	Smith		6,726,664 B2	4/2004	Yaron et al.	604/265
4,968,296 A	11/1990	Ritch et al.	604/8	6,730,056 B1	5/2004	Ghaem et al.	
5,041,081 A	8/1991	Odrich	604/9	6,736,791 B1	5/2004	Tu et al.	
5,071,408 A	12/1991	Ahmed	606/108	6,741,666 B1	5/2004	Henry et al.	
5,073,163 A	12/1991	Lippman		6,780,164 B2	8/2004	Bergheim et al.	
5,092,837 A	3/1992	Ritch et al.	604/8	6,783,544 B2	8/2004	Lynch et al.	
5,127,901 A	7/1992	Odrich	604/9	6,827,699 B2	12/2004	Lynch et al.	
5,171,213 A	12/1992	Price, Jr.	604/9	6,827,700 B2	12/2004	Lynch et al.	
5,178,604 A	1/1993	Baerveldt et al.	604/8	6,881,197 B1	4/2005	Nigam	604/8
5,180,362 A	1/1993	Worst	604/8	6,881,198 B2	4/2005	Brown	604/8
5,300,020 A *	4/1994	L'Esperance, Jr.	604/9	6,939,298 B2	9/2005	Brown et al.	
5,338,291 A	8/1994	Speckman et al.	604/9	6,955,656 B2	10/2005	Bergheim et al.	
5,342,370 A	8/1994	Simon et al.	606/107	6,962,573 B1	11/2005	Wilcox	604/9
5,346,464 A	9/1994	Camras	604/9	6,966,888 B2	11/2005	Cullen et al.	
5,370,607 A	12/1994	Memmen	604/8	6,969,384 B2	11/2005	de Juan, Jr. et al.	
5,372,577 A	12/1994	Ungerleider		6,981,958 B1	1/2006	Gharib et al.	
5,397,300 A	3/1995	Baerveldt et al.	604/8	6,989,007 B2	1/2006	Shadduck	606/4
5,433,701 A	7/1995	Rubinstein	604/8	7,041,077 B2	5/2006	Shields	
5,443,505 A	8/1995	Wong et al.	623/4.1	7,090,681 B2	8/2006	Weber et al.	
5,454,746 A	10/1995	Guegan et al.	446/72	7,094,225 B2	8/2006	Tu et al.	
5,476,445 A	12/1995	Baerveldt et al.	604/8	7,135,009 B2	11/2006	Tu et al.	
5,558,629 A	9/1996	Baerveldt et al.	604/8	7,160,264 B2	1/2007	Lisk, Jr. et al.	
5,558,630 A	9/1996	Fisher		7,163,543 B2	1/2007	Smedley et al.	
5,601,094 A	2/1997	Reiss	128/899	7,186,232 B1	3/2007	Smedley et al.	
5,626,558 A	5/1997	Suson	604/8	7,192,412 B1	3/2007	Zhou et al.	
5,626,559 A *	5/1997	Solomon	604/9	7,195,774 B2	3/2007	Carvalho et al.	
5,651,782 A	7/1997	Simon et al.	606/1	7,207,965 B2	4/2007	Simon	
5,676,944 A	10/1997	Alvarado et al.	424/93.71	7,220,238 B2	5/2007	Lynch et al.	
5,702,414 A	12/1997	Richter et al.	606/166	7,273,475 B2	9/2007	Tu et al.	
5,704,907 A	1/1998	Nordquist et al.	604/8	7,297,130 B2	11/2007	Bergheim et al.	
5,713,844 A	2/1998	Peyman		7,331,984 B2	2/2008	Tu et al.	
5,741,292 A	4/1998	Mendius	606/191	7,431,710 B2	10/2008	Tu et al.	
5,743,868 A *	4/1998	Brown et al.	604/8	7,488,303 B1	2/2009	Haffner et al.	
5,752,928 A	5/1998	De Roulhac et al.		7,857,782 B2	12/2010	Tu et al.	
5,807,302 A *	9/1998	Wandel	604/8	2001/0025150 A1	9/2001	De Juan, Jr. et al.	
5,868,697 A	2/1999	Richter et al.	604/8	2002/0013546 A1	1/2002	Grieshaber et al.	
5,882,327 A	3/1999	Jacob	604/8	2002/0013572 A1	1/2002	Berlin	
5,893,837 A	4/1999	Eagles et al.	604/9	2002/0072673 A1	6/2002	Yamamoto et al.	600/452
5,968,058 A	10/1999	Richter et al.	606/166	2002/0087111 A1	7/2002	Ethier et al.	
6,007,510 A	12/1999	Nigam	604/8	2002/0111608 A1	8/2002	Baerveldt et al.	
6,007,511 A	12/1999	Prywes	604/9	2002/0128613 A1	9/2002	Nakayama	604/294
6,019,786 A	2/2000	Thompson		2002/0133168 A1	9/2002	Smedley et al.	
6,050,970 A	4/2000	Baerveldt	604/28	2002/0143284 A1	10/2002	Tu et al.	
6,077,299 A	6/2000	Adelberg et al.	623/24	2002/0177856 A1	11/2002	Richter et al.	606/108
6,102,045 A	8/2000	Nordquist et al.	128/898	2002/0193725 A1	12/2002	Odrich	604/8
6,142,969 A	11/2000	Nigam	604/8	2003/0055372 A1	3/2003	Lynch et al.	604/8
6,186,974 B1	2/2001	Allan et al.		2003/0060752 A1	3/2003	Bergheim et al.	
6,203,513 B1	3/2001	Yaron et al.	604/9	2003/0097151 A1	5/2003	Smedley et al.	
6,221,078 B1	4/2001	Bylsma	606/107	2003/0097171 A1	5/2003	Elliott	
6,251,090 B1	6/2001	Avery et al.	604/9	2003/0135149 A1	7/2003	Cullen et al.	
6,261,256 B1	7/2001	Ahmed	604/9	2003/0181848 A1	9/2003	Bergheim et al.	
6,264,668 B1	7/2001	Prywes	606/167	2003/0187384 A1	10/2003	Bergheim et al.	
6,331,313 B1	12/2001	Wong et al.		2003/0187385 A1	10/2003	Bergheim et al.	604/27
6,375,642 B1	4/2002	Grieshaber et al.		2003/0208163 A1	11/2003	Yaron et al.	
6,383,219 B1	5/2002	Telandro et al.		2003/0229303 A1	12/2003	Haffner et al.	
6,450,984 B1	9/2002	Lynch et al.		2003/0236483 A1	12/2003	Ren	604/8
6,464,724 B1	10/2002	Lynch et al.		2003/0236484 A1	12/2003	Lynch et al.	
6,468,283 B1 *	10/2002	Richter et al.	606/108	2004/0024345 A1	2/2004	Gharib et al.	
6,471,666 B1	10/2002	Odrich		2004/0050392 A1	3/2004	Tu et al.	128/898
6,471,777 B1	10/2002	Kobayashi et al.	118/500	2004/0073156 A1	4/2004	Brown	
6,508,779 B1	1/2003	Suson	604/8	2004/0088048 A1	5/2004	Richter et al.	623/4.1
6,510,600 B2	1/2003	Yaron et al.	29/428	2004/0092856 A1	5/2004	Dahan	
6,524,275 B1	2/2003	Lynch et al.		2004/0102729 A1	5/2004	Haffner et al.	604/8
6,533,768 B1 *	3/2003	Hill	604/521	2004/0111050 A1	6/2004	Smedley et al.	
6,537,568 B2	3/2003	Olejnik et al.		2004/0127843 A1	7/2004	Tu et al.	
6,544,208 B2	4/2003	Ethier et al.	604/8	2004/0147870 A1	7/2004	Burns et al.	
6,544,249 B1	4/2003	Yu et al.	604/521	2004/0148022 A1	7/2004	Eggleston	
6,558,342 B1	5/2003	Yaron et al.	604/9	2004/0193095 A1	9/2004	Shadduck	604/8
6,589,203 B1	7/2003	Mitrev	604/27	2004/0193262 A1	9/2004	Shadduck	623/4.1
6,595,945 B2	7/2003	Brown		2004/0210181 A1	10/2004	Vass et al.	604/8
				2004/0210185 A1	10/2004	Tu et al.	

2004/0216749	A1	11/2004	Tu et al.	
2004/0225250	A1	11/2004	Yablonski	604/8
2004/0236343	A1	11/2004	Taylor et al.	606/108
2004/0249333	A1	12/2004	Bergheim et al.	
2004/0254517	A1	12/2004	Quiroz-Mercado et al.	604/8
2004/0254519	A1	12/2004	Tu et al.	
2004/0254520	A1	12/2004	Porteous et al.	
2004/0254521	A1	12/2004	Simon	604/8
2004/0260228	A1	12/2004	Lynch et al.	
2005/0008673	A1	1/2005	Snyder et al.	
2005/0049578	A1	3/2005	Tu et al.	
2005/0090806	A1	4/2005	Lynch et al.	604/523
2005/0090807	A1	4/2005	Lynch et al.	
2005/0107734	A1	5/2005	Coroneo	
2005/0119601	A9	6/2005	Lynch et al.	604/8
2005/0119636	A1	6/2005	Haffner et al.	604/500
2005/0119737	A1	6/2005	Bene et al.	
2005/0125003	A1	6/2005	Pinchuk et al.	
2005/0143817	A1	6/2005	Hunter et al.	623/11.11
2005/0149080	A1	7/2005	Hunter et al.	606/155
2005/0175663	A1	8/2005	Hunter et al.	424/423
2005/0181011	A1	8/2005	Hunter et al.	424/423
2005/0181977	A1	8/2005	Hunter et al.	514/2
2005/0182350	A1	8/2005	Nigam	604/8
2005/0191331	A1	9/2005	Hunter et al.	424/423
2005/0192527	A1	9/2005	Gharib et al.	604/8
2005/0197613	A1	9/2005	Sniegowski et al.	
2005/0209549	A1	9/2005	Bergheim et al.	
2005/0209550	A1	9/2005	Bergheim et al.	
2005/0232972	A1	10/2005	Odrich	
2005/0244462	A1	11/2005	Farooq	
2005/0250788	A1	11/2005	Tu et al.	
2005/0266047	A1	12/2005	Tu et al.	
2005/0267397	A1	12/2005	Bhalla	
2005/0267398	A1	12/2005	Protopsaltis et al.	
2005/0271704	A1	12/2005	Tu et al.	
2005/0273033	A1	12/2005	Grahn et al.	604/9
2005/0277864	A1	12/2005	Haffner et al.	604/8
2005/0283108	A1	12/2005	Savage	
2005/0288617	A1	12/2005	Yaron et al.	
2005/0288619	A1	12/2005	Gharib et al.	
2006/0020248	A1	1/2006	Prescott	
2006/0032507	A1	2/2006	Tu et al.	
2006/0036207	A1	2/2006	Koonmen et al.	
2006/0069340	A1	3/2006	Simon	
2006/0074375	A1	4/2006	Bergheim et al.	
2006/0084907	A1	4/2006	Bergheim et al.	
2006/0116626	A1	6/2006	Smedley et al.	
2006/0155238	A1	7/2006	Shields	
2006/0173397	A1	8/2006	Tu et al.	
2006/0195055	A1	8/2006	Bergheim et al.	
2006/0195056	A1	8/2006	Bergheim et al.	
2006/0200113	A1	9/2006	Haffner et al.	
2006/0235367	A1	10/2006	Takahima et al.	604/541
2006/0241580	A1	10/2006	Mittelstein et al.	
2006/0241749	A1	10/2006	Tu et al.	
2006/0276739	A1	12/2006	Brown	
2007/0010827	A1	1/2007	Tu et al.	
2007/0088432	A1	4/2007	Solovay et al.	
2007/0106235	A1	5/2007	Coroneo	
2007/0106236	A1	5/2007	Coroneo	
2007/0112292	A1	5/2007	Tu et al.	
2007/0118147	A1	5/2007	Smedley et al.	
2007/0149915	A1	6/2007	Yablonski	
2007/0191863	A1	8/2007	de Juan, Jr. et al.	
2007/0276315	A1	11/2007	Haffner et al.	
2007/0276316	A1	11/2007	Haffner et al.	
2007/0282244	A1	12/2007	Tu et al.	
2007/0282245	A1	12/2007	Tu et al.	
2007/0293807	A1	12/2007	Lynch et al.	
2008/0015488	A1	1/2008	Tu et al.	
2008/0045878	A1	2/2008	Bergheim et al.	
2008/0195027	A1	8/2008	Coroneo	
2008/0200860	A1	8/2008	Tu et al.	
2008/0228127	A1	9/2008	Burns et al.	
2008/0234624	A2	9/2008	Bergheim et al.	
2009/0036819	A1	2/2009	Tu et al.	
2010/0274259	A1	10/2010	Yaron et al.	
2011/0028884	A1	2/2011	Coroneo	

2011/0087149	A1	4/2011	Coroneo
2011/0087150	A1	4/2011	Coroneo
2011/0087151	A1	4/2011	Coroneo

FOREIGN PATENT DOCUMENTS

EP	1310222	5/2003
EP	1473004	11/2004
EP	1477146	11/2004
EP	1418868	3/2008
EP	1977724	10/2008
EP	2027837	2/2009
RU	2018289 C1	8/1994
RU	2056818 C1	3/1996
RU	2074686 C1	3/1997
RU	2074687 C1	3/1997
RU	2157678 C1	10/2000
WO	89/00869	2/1989
WO	91/12046	8/1991
WO	92/19294	11/1992
WO	94/09721	5/1994
WO	94/09837	5/1994
WO	94/13234	6/1994
WO	95/08310	3/1995
WO	96/20742	7/1996
WO	96/36377	11/1996
WO	98/23237	6/1998
WO	98/30181	7/1998
WO	99/26567	6/1999
WO	00/64511	11/2000
WO	01/78656	10/2001
WO	00/06223	11/2001
WO	02/36052	5/2002
WO	02/070045	9/2002
WO	02/074052	9/2002
WO	02/080811	10/2002
WO	02/080829	10/2002
WO	02/087418	11/2002
WO	02/087479	11/2002
WO	02/089699	11/2002
WO	02/102274	12/2002
WO	03/015659	2/2003
WO	03/015667	2/2003
WO	03/041622	5/2003
WO	03/073968	9/2003
WO	03/099175	12/2003
WO	2004/026347	4/2004
WO	2004/043231	5/2004
WO	2004/056294	7/2004
WO	2004/060219	7/2004
WO	2004/062469	7/2004
WO	2004/110391	12/2004
WO	2005/016418	2/2005
WO	2005/046516	5/2005
WO	2005/046782	5/2005
WO	2005/055873	6/2005
WO	2005/107664	11/2005
WO	2005/107845	11/2005
WO	2006/012421	2/2006
WO	2006/036715	4/2006
WO	2007/087061	8/2007
WO	2007/115259	10/2007
WO	2008/061043	5/2008

OTHER PUBLICATIONS

Cohen, V.M., et al., "First day post-operative review following uncomplicated phacoemulsification," Eye, 12(4):634-636, (1998).
Derwent English abstract for EP 1184010, published Mar. 6, 2002 entitled: "Drainage unit for an eye, consists of a hollow line, a distribution member, and a pressure relief valve which only allows water to leave the eye chamber above a certain pressure," Accession Nbr. 12409716 [351].
Dinakaran, S., et al., "Is the first post-operative day review necessary following uncomplicated phacoemulsification surgery?" Eye, 14(3A):364-366, (2000).
Einmahl, S., et al., "Evaluation of a novel biomaterial in the suprachoroidal space of the rabbit eye," Investigative Ophthalmology and Visual Sciences, 43:1533-1539, (2002).

- Gross, R.L., et al., "Surgical therapy of chronic glaucoma in aphakia and pseudophakia," *Ophthalmology*, 95:1195-1201, (1988).
- Hildebrand, G.D., et al., "Efficacy of anterior chamber decompression in controlling early intraocular pressure spikes after uneventful phacoemulsification," *Journal of Cataract and Refractive Surgery*, 29:1087-1092, (2003).
- Howorth, D.J., "Feasibility study for a micromachined glaucoma drainage device," Cranfield University School of industrial and manufacturing science MSc Thesis Academic Year 2001-2002 Sep. 13, 2002.
- Hylton, C. and A. Robin et al., "Update on prostaglandin analogs," *Current Opinion in Ophthalmology*, 14:65-69, (2003).
- Jordan, J. "A novel approach to suprachoroidal drainage for the surgical treatment of intractable glaucoma," *Journal of Glaucoma*, 15:200-205, (2006).
- Lee et al. "Magnetic resonance imaging of the aqueous flow in eyes implanted with the trabeculo-suprachoroidal suprachoroidal glaucoma seton," *Investigative Ophthalmology and Visual Sciences* 33:948 (1992).
- Mehta, K.R., "The suprachoroidal hema wedge in glaucoma surgery," *American Academy of Ophthalmology meeting 1977* pp. 144.
- Rosenberg, L. and T. Krupin, *The Glaucomas*, Second Edition, Chapter 88, Eds., Ritch, R., et al., Mosby, St. Louis, MO., pp. 1783-1807, (1986).
- "The Solx Gold Micro-shunt (GMS) Treatment," Solx clinical literature hand-out from Industry Show, Feb. 2006.
- Srinivasan, R. et al., "Microbial contamination of the anterior chamber during phacoemulsification," *Journal of Cataract and Refractive Surgery*, 28:2173-2176, (2002).
- Toris, C., et al., "Aqueous humor dynamics in the aging human eye," *American Journal of Ophthalmology*, 127:407-412, (1999).
- Yablonski, M.E., "Some thoughts on the pressure dependence of uveoscleral flow," *Journal of Glaucoma*, 12(1):90-92, (2003).
- Yablonski, M.E., "Trabeculectomy with Internal Tube Shunt: a novel glaucoma surgery," *Journal of Glaucoma*, 14:91-97, (2005).
- Fuchs E. "Detachment of the choroid inadvertently during cataract surgery," *von Graefes Arch Ophthalmol*, 51:199-224 (1900) [German with English Language abstract].
- Heine I. "Cyclodialysis, a new glaucoma operation," *Dtsch Med Wochenschr*, 31:824-826 (1905) [German with English Language abstract].
- Karlen et al., "Deep sclerectomy with collagen implant: medium term results" *Br. J. Ophthalmol*, 83(1):6-11 (1999).
- Kozlov et al. "Nonpenetrating deep sclerectomy with collagen," *Eye Microsurgery* 3:44-46 (1990). [Russian with English translation].
- Bick, *Arch Ophthalmol*. Oct. 1949; 42(4): 373-88.
- Bietti "The present state of the use of plastics in eye surgery" *Acta Ophthalmol (Copenh)* 1955; 33(4):337-70.
- Emi et al. "Hydrostatic pressure of the suprachoroidal space" *Invest. Ophthalmol. Visual Sci.* 30(2):233-238 (1989).
- Gills "Cyclodialysis implants" *South Med J.* 1967 60(7):692-5.
- Gills et al. "Action of cyclodialysis utilizing an implant studied by manometry in a human eye" *Exp Eye Res* 1967; 6:75-78.
- Jordan et al. *J. Glaucoma* 2006; 15:200-205.
- Klemm et al. "Experimental use of space-retaining substances with extended duration: functional and morphological results" *Graefes Arch Clin Exp Ophthalmol Sep.* 1995; 233(9):592-7.
- Krejci "Cyclodialysis with hydroxymethyl methacrylate capillary strip (HCS). Animal experiments with a new approach in glaucoma drainage surgery" *Ophthalmologica* 1972; 164(2):113-21.
- Losche "Proposals for improvement of cyclodialysis" *Klin Monatsblätter Augenheilkd Augenarztl Fortbild* 1952, 121(6):715-6 [German].
- Nesterov AP et al. "Surgical stimulation of the uveoscleral outflow. Experimental studies on enucleated human eyes" *Acta Ophthalmol (Copenh) Jun.*; 57(3):409-17 (1979).
- Ozdamar et al. "Suprachoroidal seton implantation in refractory glaucoma: a novel surgical technique" *J. Glaucoma Aug.* 2003; 12(4):354-9.
- Pinnas et al. "Cyclodialysis with teflon tube implants" *Am J. Ophthalmol Nov.* 1969; 68(5):879-883.
- Row H. (1934) *Arch. Ophthalmol* 12, 325.
- Troncosco UM Cyclodialysis with insertion of metal implant in treatment of glaucoma Preliminary report *Arch. Ophth.* 23:270, 1940.
- Barsky et al. "Evaluation of absorbable gelatin film (Gelfilm) in cyclodialysis clefts" *Arch. Ophth.* 60(6): 1044-1052, 1958.
- Brown et al., "Internal Sclerectomy for Glaucoma Filtering Surgery with an Automated Trephine," *Archives of Ophthalmology*, vol. 105, Jan. 1987, 133-136.
- Burchfield Jc, Kass Ma, Wax MB. Primary valve malfunction of the Krupin eye valve with disk. *J Glaucoma. Jun.* 1997;6(3):152-6.
- Chiou et al. "Ultrasound biomicroscopy of eyes undergoing deep sclerectomy with collagen implant" *Br J Ophthalmol* 80 (1996), pp. 541-544.
- Chylack LT, Bellows AR. Molecular sieving in suprachoroidal fluid formation in man. *Invest Ophthalmol Vis Sci* 17: 420, 1978.
- Collaborative Normal-Tension Study Group. Comparison of glaucomatous progression between untreated patients with normal-tension glaucoma and patients with therapeutically reduced intraocular pressures. *Am J Ophthalmol* 1998;126:487-97.
- Demailly et al. "Non-penetrating deep sclerectomy (NPDS) with or without collagen device (CD) in primary open-angle glaucoma: middle-term retrospective study" *International Ophthalmology* 20: 131-140, 1997.
- Draeger "Chirurgische Maßnahmen bei kongenitalem Glaukom" (Surgical Interventions in Congenital Glaucoma) *Klin Monatsbl Augenheilkd* 1993; 202(5): 425-427 [Article in German with English summary included].
- Ellis, RA "A Reduction of Intraocular Pressure Using Plastics in Surgery" *Am J Ophth.* 50; 1960, 733-742.
- Fanous MM, Cohn RA. Propionibacterium endophthalmitis following Molteno tube repositioning. *J Glaucoma. Aug.* 1997;6(4):201-2.
- Gills, "Cyclodialysis Implants in Human Eyes" *Am J Ophth* 61:1966,841-846.
- Goldberg "Management of Uncontrolled Glaucoma With the Molteno System" *Australian and New Zealand Journal of Ophthalmology* 1987; 15: 97-107.
- Gordon MO, Kass. MA, for the Ocular Hypertension Treatment Study Group. The Ocular Hypertension Treatment Study. Design and baseline description of the participants. *Arch Ophthalmol* 1999:573-83.
- Harper SL, Foster CS. Intraocular lens explantation in uveitis. *Int Ophthalmol Clin.* 2000 Winter; 40(1):107-16.
- Harrington "Cataract and glaucoma. Management of the coexistent conditions and a description of a new operation combining lens extraction with reverse cyclodialysis." *Am J Ophthalmol.* May 1966;61(5 Pt 2):1134-40.
- Heijl A, Leske MC, Bengtsson B, et al for the Early Manifest Glaucoma Trial Group. Reduction of intraocular pressure and glaucoma progression. Results from the Early Manifest Glaucoma Trial. *Arch Ophthalmol* 2002;120:1268-79.
- Javitt JC, Chiang YP. Preparing for managed competition. Utilization of ambulatory eye care visits to ophthalmologists. *Arch Ophthalmol* 1993;111:1034-5.
- Jay JL, Allan D. The benefit of early trabeculectomy versus conventional management in primary open-angle glaucoma relative to severity of disease. *Eye* 1989; 3:528-35.
- Kass MA, Heuer DK, Higginbotham EJ, et al for the Ocular Hypertension Treatment Study Group. The Ocular Hypertension Treatment Study. A randomized trial determines that topical ocular hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Arch Ophthalmol* 2002;120:701-13.
- Klemm et al. "Die Ultraschallbiomikroskopie als Kriterium der Funktionsprüfung des suprachoroidalen Spaltes nach kammerwinkelchirurgischen Eingriffen (Ultrasound Biomicroscopic Imaging for Assessment of the Suprachoroidal Cleft after Angle Surgery)" *Klinische Monatsblätter für Augenheilkunde* 1997; 210: 74-77 [Article in German with English summary included].
- Krejci L. "Microdrainage of anterior chamber of eye glaucoma operation using hydon capillary drain." *Acta Univ Carol Med Monogr.* 1974;(61):1-90.
- Kupfer "Studies on intraocular pressure. I. A technique for polyethylene tube implantation into the anterior chamber of the rabbit." *Arch Ophthalmol.* Apr. 1961;65:565-70.

- La Rocca "Gonioplasty in Glaucoma" A Preliminary Report" Br J Ophth 46:1962, 404-415.
- Law et al., "Retinal Complications After Aqueous Shunt Surgical Procedures for Glaucoma" Arch Ophthal.; Dec. 1996; vol. 114:1473-1480.
- Lee Ky. Trabeculo-suprachoroidal shunt for treating recalcitrant and secondary glaucoma. Presented at the American Academy of Ophthalmology Annual Meeting, Anaheim, CA, 1991.
- Leske MC, Heijl A, Hussein M, et al for the Early Manifest Glaucoma Trial Group. Factors for glaucoma progression and the effect of treatment. The Early Manifest Glaucoma Trial. Arch Ophthalmol 2003;121:48-56.
- Lichter PR, Musch DC, Gillespie BW, et al and the CIGTS Study Group. Interim clinical outcomes in the Collaborative Initial Glaucoma Treatment Study comparing initial treatment randomized to medications or surgery. Ophthalmology 2001;108:1943-53.
- Marx et al., "Use of the Ganciclovir Implant in the Treatment of Recurrent Cytomegalovirus Retinitis" Arch Ophthal.; Jul. 1996; vol. 114:815-820.
- McPherson "Combined Trabeculotomy and Cataract Extraction as a Single Operation" Tr. Am. Ophth. Soc., vol. LXXIV, 1976; 251-260.
- Migdal C, Gregory W, Hitchings R. Long term functional outcome after early surgery compared with laser and medicine in open-angle glaucoma. Ophthalmology 1994;101:1651-7.
- Miglior S, Zeyen T, Pfeiffer N, et al for the European Glaucoma Prevention Study Group. The European Glaucoma Prevention Study design and baseline description of the participants. Ophthalmology 2002;109:1612-21.
- Miki, MD et al., "Intraocular Cannula for Continuous, Chronic Drug Delivery-Histopathic Observations and Function" Arch Ophthal.; May 1985; vol. 103:712-717.
- Molteno et al. "Long tube implants in the management of glaucoma" South African Medical Journal, Jun. 26, 1976;50(27):1062-6.
- Molteno et al. "The Vicryl tie technique for inserting a draining implant in the treatment of secondary glaucoma." Australian and New Zealand Journal of Ophthalmology 1986; 14: 343-354.
- Moses RA "Detachment of ciliary body-anatomical and physical considerations" Investigative Ophthalmology & Visual Science, Assoc. For Research in Vision and Ophthalmology, US, vol. 4, No. 5, Oct. 1, 1965, 935-941.
- Nguyen et al., "Complications of Baerveldt Glaucoma Drainage Implants" Arch Ophthal.; May 1998; vol. 116:571-575.
- O'Brien et al. "Cyclodialysis" Arch Ophthal. 1949;42(5):606-619.
- Portney GL, "Silicone elastomer implantation cyclodialysis." Arch Ophthalmol 1973; 89: 10-12.
- Pruett et al., "The Fishmouth Phenomenon-II. Wedge Scleral Buckling" Arch Ophthal.; Oct. 1977; vol. 95:1782-1787.
- Qadeer "Acrylic Gonio-Subconjunctival Plates in Glaucoma Surgery" Br J Ophthalmol. Jun. 1954; 38(6): 353-356.
- Quigley HA, Vitale S. Models of open-angle glaucoma prevalence and incidence in the United States. Invest Ophthalmol Vis Sci 1997; 38:83-91.
- Richards et al. "Artificial Drainage Tubes for Glaucoma" Am J Ophth 60:1965,405-408.
- Ritch R, Shields MB, Krupin T. The Glaucomas. St. Louis: Mosby, 1996; 337-343).
- Sampimon "A New Approach to Filtering Glaucoma Surgery" Ophthalmologica (Basel) 151: 1966, 637-644.
- Schappert S. Office visits for glaucoma: United States, 1991-92. Advance data from vital and health statistics. vol. 262. Hyattsville, Md: National Center for Health Statistics, 1995.
- Shaffer RN, Weiss DI. Concerning cyclodialysis and hypotony. Arch Ophthalmol 68: 25, 1962.
- Sommer A, Tielsch JM, Katz J, et al. Racial differences in the cause-specific prevalence of blindness in east Baltimore. N Engl J Med 1991;325:1412-7.
- Sourdille et al. "Reticulated hyaluronic acid implant in non-perforating trabecular surgery." J Cataract Refract Surg 25: 332-339. (1999):.
- Suguro K, Toris CB, Pederson JE. Uveoscleral outflow following cyclodialysis in the monkey eye using a fluorescent tracer. Invest Ophthalmol Vis Sci 1985: 26, 810.
- The Advanced Glaucoma Intervention Study (AGIS): 7. The relationship between control of intraocular pressure and visual field deterioration. The AGIS Investigators. Am J Ophthalmol 2000;130:429-40.
- The Glaucoma Laser Trial (GLT). 2. Results of argon laser trabeculoplasty versus topical medicines. The Glaucoma Laser Trial Research Group. Ophthalmology 1990;97:1403-13.
- The Glaucoma Laser Trial (GLT) and Glaucoma Laser Trial Follow-up Study: 7. Results. Am J Ophthalmol 1995;120:718-31.
- Tielsch JM, Sommer A, Katz J, et al. Racial variations in the prevalence of primary open-angle glaucoma. The Baltimore Eye Survey. JAMA 1991;266:369-74.
- Toris et al. "Effect of intraocular pressure on uveoscleral outflow following cyclodialysis in the monkey eye." Investigative Ophthalmology & Visual Science. 26 (1985) 1745-1749.
- Torts CB. Extravascular albumin concentration of the uvea. Invest Ophthalmol Vis Sci 1990; 31:43.
- Zhou et al. "A trabecular bypass flow hypothesis" J Glaucoma. 14(1):74-83 (2005).
- Grant, W.M. , MD, Further Studies on Facility of Flow Through the Trabecular Meshwork, A.M.A. Archives of Ophthalmology, Oct. 1958, vol. 60, pp. 523-533.
- Hoskins, et al., "Aqueous Humor Outflow", Becker-Shaffer's Diagnosis and Therapy of the Glaucomas, 6th Edition, Chapter 4, pp. 41-66, 1989.
- Olsen, Timothy W., et al., Cannulation of the Suprachoroidal Space: A Novel Drug Delivery Methodology to the Posterior Segment, American Journal of Ophthalmology, vol. 142, No. 5, Nov. 2006, pp. 777-787.e2.
- Rohen, Johannes W., Anatomy of the Aqueous Outflow Channels, Glaucoma, vol. 1, Chapter 14, pp. 277-296, Edited by J.E. Cairns, Grune & Stratton, Harcourt Brace Jovanovich Publishers, 1986.
- Rowan, Patrick J., MD, Combined Cyclodialysis and Cataract Surgery, Ophthalmic Surgery and Lasers, Dec. 1998, vol. 29, No. 12, pp. 962-968 (9 pages).
- Troncoso, Manuel U., Tantalum implants for inducing hypotny, Am Journal of Ophthalmology, vol. 32(4):499-508 (1949).
- Wagner, Justin A., et al., Characterization of Uveoscleral Outflow in Enucleated Porcine Eyes Perfused under Constant Pressure, Invest Ophthalmol Vis Sci., Published in edited form in Sep. 2004, vol. 45, Issue 9, pp. 3203-3206.

* cited by examiner

FIGURE 1




FIGURE 2





FIGURE 3



1

OCULAR PRESSURE REGULATION

This Application is a 371 of PCT/AU2004/001566 filed Nov. 12, 2004; the disclosure of which is incorporated herein by reference.

FIELD OF THE INVENTION

This invention is directed to therapeutic methods and devices for the treatment of glaucoma. In particular, this invention is concerned with the use of a shunt or drain for the treatment of glaucoma. In another aspect this invention is concerned with ocular pressure spikes shunts and use of the same in ocular surgery.

BACKGROUND OF THE INVENTION

The glaucomas are a common group of blinding conditions usually associated with elevated intraocular pressure. This elevated pressure in the eye may be regarded as a disorder of the drainage system of the eye which gives rise to the glaucomas.

Aqueous humor of the eye ("aqueous") is a flowing liquid fluid (composed of sodium, chloride, bicarb, amino acids, glucose, ascorbic acid, and water) that is actively secreted by the ciliary body and flows out past the iris into the anterior chamber (are between the lens/iris and the cornea). The aqueous drains out through angle formed by the iris and the sclera into a meshwork call the trabeculum, and from there into the canal of Schlem and then into the episcleral veins. Uveosclera drainage also occurs. Normal intraocular pressure (IOP) of aqueous in anterior chamber is between 10 and 20 mm Hg. Prolonged IOPs of greater than 21 mm Hg are associated with damage to optic nerve fibres.

In some cases of glaucoma the cause can be found: the trabecular meshwork becomes blocked by pigment or membrane. In other cases, blockage is due to a closure of the angle between the iris and the cornea. This angle type of glaucoma is referred to as "angle-closure glaucoma". In the majority of glaucoma cases, however, called "open angle glaucoma", the cause is unknown.

Elevated intraocular pressure results in the death of retinal ganglion cells (which convey retinal information to the brain) resulting in a characteristic pattern of loss of the field of vision, progressing to tunnel vision and blindness if left untreated.

Treatment of glaucoma consists predominantly of methods to lower the intraocular pressure (pharmacological, trabecular meshwork laser and surgery to drain fluid from the eye). More recently protection of the retinal ganglion cells by neuroprotective agents has been attempted.

Although pharmacological treatments of glaucoma have improved, they have important implications for the patient's quality of life, have compliance issues which are important in the elderly (in whom glaucoma is prevalent), expose the patient of glaucoma to side effects, and over a lifetime are costly.

Surgery for glaucoma treatment is usually a trabeculectomy in which a fistula is created to drain fluid from the anterior chamber to the subconjunctival space near the limbus, creating a bulge in the conjunctiva known as a bleb. Frequently scarring occurs and attempts to counter this with antimetabolites such as Mitomycin C have met with some success. In recalcitrant cases, glaucoma implants, drainage, shunt or valve devices have been developed eg Molteno (U.S. Pat. No. 4,457,757), Krupin (U.S. Pat. No. 5,454,746) and Baerveldt (U.S. Pat. No. 5,178,604). These suffer from simi-

2

lar problems of scarring (Classen L, Kivela T, Tarkkanen "A Histopathologic and immunohistochemical analysis of the filtration bleb after unsuccessful glaucoma seton implantation" *Am J Ophthalmol*, 1996;122:205-12) around the external opening of the tube devices in the subconjunctival space—the development of a large number of these devices is testament to the fact that many fail in the longer term. In these devices a drainage tube is located in the anterior chamber and is in fluid communication with the sclera or a surgically created subconjunctival space.

Whereas cataract surgery has been revolutionized in the last two decades, improvements in glaucoma surgery have been slower. Antifibrotic agents have improved the success rate of conventional filtration surgery (trabeculectomy), but with increased bleb leaks, blebitis, endophthalmitis and hypotensive maculopathy. Glaucoma shunts have had limited success in eyes that have "failed" multiple standard procedures. However complications with malpositioned tubes, erosion & strabismus persist. A considerable issue is the lack of reproducibility and predictability in achieving the desired target intraocular pressure (IOP). Final IOP is largely determined by healing which can be unpredictable—in view of vast biological variations, it is impossible to predict which eyes will rapidly scar causing failure and which will fail to heal resulting in prolonged post-operative hypotony. Scarring remains a significant problem in all these external drainage proposals, where aqueous drains into the conjunctiva, or surgical chambers in the sclera.

The introduction of a new class of antiglaucoma drugs, the prostaglandin analogues has resulted in acknowledgment of the importance of the uveoscleral pathway in drainage of fluid from the eye (Hylton C, Robin A L "Update on prostaglandin analogs" *Curr Opin Ophthalmol*, 2003;14:65-9). Uveoscleral flow where aqueous humor flows through the interstitium of the ciliary muscle into the suprachoroidal space (a potential space between the choroids and sclera) and out through the sclera into the connective tissue of the orbit may account for 54% of outflow young healthy humans (Toris C B, Yablonsid M E, Wang Y L, Camras C B "Aqueous humor dynamics in the aging human eye" *Am J Ophthalmol*, 1999;127:407-12).

Cyclodialysis, the separation of the ciliary body from the scleral spur and underlying sclera, creates free communication between the anterior chamber and the suprachoroidal space and enhances uveoscleral flow. It has long been known that cyclodialysis can cause a profound reduction of intraocular pressure—initially (Fuchs E. "Detachment of the choroid inadvertently during cataract surgery" [German] *von Graefes Arch Ophthalmol*, 1900;51:199-224) cyclodialysis was recognized as a complication of cataract surgery. Deliberate creation of a cyclodialysis cleft for treating elevated intraocular pressure in uncontrolled glaucoma was first described as a surgical procedure in 1905 (Heine I. "Cyclodialysis, a new glaucoma operation" [German] *Dtsch Med Wochenschr*, 1905;31:824-826. Since such clefts can heal and close spontaneously a number of devices have been used to keep them open, including platinum wire, horse hair, magnesium strips, tantalum foil, supramid, gelatin film, Teflon, silicone and polymethylmethacrylate (Rosenberg L F, Krupin T. "Implants in glaucoma surgery" Chapter 88, *The Glaucomas*, Ritch R, Shields B M, Krupin T Eds. 2nd Edition Mosby St Louis 1986) and Hema (Mehta K R. "The suprachoroidal Hema wedge in glaucoma surgery" American Academy of Ophthalmology meeting 1977, pp 144). However the success rate of such approaches has been low (as low as 15%, Rosenberg & Krupin *ibid* and Gross R L, Feldman R M, Spaeth G L, et al "Surgical therapy of chronic glaucoma in aphalia and pseudophakia" *Ophthalmology*, 1988;95:1195-201). Failure

was due to uncontrolled low pressure (hypotony) with consequential macular edema, bleeding (hyphema) and inadequate pressure control.

The device and method of a first aspect of this invention takes advantage of the methods used in cataract surgery to develop a minimally invasive glaucoma procedure—thus small, self sealing incisions and materials that are biocompatible and foldable so that they fit through small openings will reduce surgical trauma and time. The controlled draining of aqueous into the suprachoroidal space according to this invention provides some predictability of outcome and overcomes scarring problems that have plagued glaucoma implants in the past.

The most frequent complication following modern cataract surgery with phacoemulsification, requiring specific treatment is elevated intraocular pressure (Cohen V M, Demetria H, Jordan K, Lamb R J, Vivian A J.: First day post-operative review following uncomplicated phacoemulsification” *Eye*, 1998; 12 (Pt 4):634-6, and Dinakaran S, Desai S P, Raj P S. “Is the first post-operative day review necessary following uncomplicated phacoemulsification surgery?” *Eye*, 2000 June; 14 (Pt 3A):364-6. The increase may be marked and typically peaks at 5 to 7 hours before returning to near normal levels in 1 to 3 days (Hildebrand G D, Wickremasinghe S S, Tranos P G, Harris M L, Little B C. “Efficacy of anterior chamber decompression in controlling early intraocular pressure spikes after uneventful phacoemulsification” *J Cataract Refract Surg.*, 2003; 29:1087-92). Such pressure spikes can cause pain and may increase the risk of sight-threatening complications such as retinal vascular occlusion, increases loss of visual field in advanced glaucoma and ischemic optic neuropathy—effects in otherwise healthy eyes are unknown (Hildebrand G D et al, *ibid*).

A number of prophylactic treatments are used with limited success—these include intracameral carbachol or acetylcholine, topical timolol, dorzolamide, aproclonidine, latanoprost and systemic acetazolamide (see Hildebrand G D et al, *ibid*). This also exposes the patient to the risk of drug side effects, increased cost and it has been postulated that reducing the flow of aqueous humor post surgery prolongs the residence time of bacteria that frequently (46.3% of cases) contaminate the anterior chamber during surgery (Srinivasan R, Tiroumal S, Kanungo R, Natarajan M K. “Microbial contamination of the anterior chamber during phacoemulsification” *J Cataract Refract Surg*, 2002; 28:2173-6.). This may increase the risk of endophthalmitis one of the most devastating sequelae of intraocular surgery, since the bacteria are not being “flushed out” of the eye by the normal production of aqueous humour, the secretion of which has been suppressed by the drugs. Another technique is to decompress the anterior chamber by applying pressure to the posterior lip of the paracentesis wound at the appropriate time. This requires surveillance and could increase the risk of infection. Another aspect of this invention hereinafter described overcomes these problems.

SUMMARY OF THE INVENTION

According to the present invention there is provided a flexible ocular device for implantation into the eye formed of a biocompatible elastomeric material, foldable to a diameter of 1.5 mm or less, comprising a fluid drainage tube having at one end a foldable plate adapted to locate the device on the inner surface of the sclera in a suprachoroidal space formed by cyclodialysis, said drainage tube opening onto the disc at one end and opening to the anterior chamber when implanted into the eye at its other end, so as to provide aqueous pressure regulation.

Preferably the fluid drainage tube has a diameter selected to provide predetermined resistance to aqueous humor flow, for example a pressure of 10 mm Hg or less. Alternatively said tube contains a valve so as to regulate pressure of the aqueous chamber at a predetermined level, for example at no less than 10 mm Hg.

In accordance with another embodiment of this invention there is provided a method for treating glaucoma which comprises:

providing a flexible ocular device formed of a biocompatible elastomeric material foldable to a diameter of 1.5 mm or less, comprising a fluid drainage tube having at one end a foldable plate adapted to locate the device on the inner surface of the sclera and at its other end being open so as to allow fluid communication through said tube;

forming a small self-sealing incision at the juncture of the cornea and sclera of the eye opening into the anterior chamber;

filling the anterior chamber with a viscoelastic substance; introducing the foldable ocular device into a suprachoroidal space formed by cyclodialysis via a hollow cannula, wherein said plate locates the device on the inner surface of the sclera in the suprachoroidal space, and said drainage tube is located in the anterior chamber of the eye so as to provide aqueous humor pressure regulation; and thereafter removing said cannula and viscoelastic material from the eye.

In another aspect there is provided an ocular pressure spike shunt for insertion into an ocular paracentesis incision port following ocular surgery, comprising a flexible fluid transfer tube formed of biocompatible material, preferably biocompatible elastomeric material, so as to allow paracentesis incision closure around said tube, having an inner end and an outer end, a tubular lumen disposed between said inner end and said outer end to allow fluid communication through said tube, said lumen containing a valve for controlling pressure in the eye following ocular surgery, which valve opens permitting fluid flow through said tube when a predetermined pressure is exceeded, said shunt being configured such that on insertion into a paracentesis port said outer end is substantially flush with the surface of the cornea, and said inner end opens into the anterior chamber of the eye.

In another aspect there is provided a method for preventing ocular pressure spikes following ocular surgery wherein a paracentesis incision port is formed in the eye during said surgery, comprising introducing an ocular pressure spike shunt into said paracentesis port at the conclusion of ocular surgery, said shunt comprising a flexible fluid transfer tube formed of biocompatible material, preferably biocompatible elastomeric material, so as to allow paracentesis incision closure around said tube, having an inner end and an outer end, a tubular lumen disposed between said inner end and said outer end to allow fluid communication through said tube, said lumen containing a valve for controlling pressure in the eye following ocular surgery, which valve opens permitting fluid flow through said tube when a predetermined pressure is exceeded, said shunt being configured such that on insertion into a paracentesis port said outer end is substantially flush with the surface of the cornea, and said inner end protrudes into the anterior chamber of the eye.

DESCRIPTION OF THE FIGURES

FIG. 1 shows a diagrammatic representation of a side sectional view of suprachoroidal shunt insertion using an injector.

5

FIG. 2 shows a diagrammatic representation of a side sectional view of an eye showing the unfolded plate portion of the device and a cannula introducing said device across the anterior chamber at 180° to the site of insertion.

FIG. 3 shows a diagrammatic representation of an eye containing a pressure spike shunt inserted into a paracentesis port.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS OF THE INVENTION

The ocular device according to the present invention is implanted in a patient's eye using minimally invasive surgery techniques, adopted from modern cataract surgery.

The ocular device is formed from a biocompatible elastomeric material. Preferably, the device is made of soft surgical grade polymeric material, such as silicon or acrylic material such that the device is foldable and may be rolled up for insertion via a cannula. FIG. 1 shows a proximal end of a cannula forming a cyclodialysis. The folded device may be introduced via such a cannula. The elastomeric material is selected to be sufficiently soft that it does not erode delicate underlying choroid material when inserted into the eye. Such material and ocular lenses formed therefrom are well known and used in cataract surgery.

Sutures are not required to hold the device in place once surgically introduced into the eye, as the foldable plate is adapted to locate the device on the inner surface of the sclera in a suprachoroidal space formed by cyclodialysis (FIG. 2). Preferably, the plate is of a disc-like shape which matches the curvature of the eye once unfolded. FIG. 2 depicts an unfolded disc (connected tube not shown) after cannula introduction across the anterior chamber (transcarnal). Alternatively, any plate-like configuration which locates the device on the inner surface of the sclera in the suprachoroidal space may be used, such as for example a rectangular foldable plate. Preferably the plate diameter is from 0.05 to 6 mm, and preferably the plate thickness is from 12.5μ to 250μ. The fluid drainage tube of the ocular device is preferably integral with the plate, and is attached at one end to the plate, preferably at the periphery of the plate. Alternatively, the tube may be microwelded or otherwise fixed to the plate. Fabrication techniques well known in production of intraocular foldable lenses are preferably used in this invention. The tube has a hollow lumen, and is preferably of a length from about 1 mm to 4 mm. Preferred diameters of the tubing comprise an outer diameter of 400-1000μ, and preferably the inner diameter is from 50 to 500μ.

The diameter of the tube may be selected so as to provide a resistance to aqueous humor flow of predetermined pressure, preferably being a pressure less than 10 mm Hg. This enables the pressure of the aqueous to be regulated in a controlled manner, providing relief from excess ocular pressure associated with glaucoma, with avoidance of hypotony (uncontrolled low pressure). Alternatively, the tube may contain a valve, for example disposed at the end of the tube opening onto the disc so as to regulate ocular pressure at a predetermined level. Preferably, the valve prevents aqueous flow through the tube at a pressure of less than 10 mm Hg. Examples of valves which may be used include a slit valve. The drainage stops altogether if the pressure drops to a predetermined threshold level controlled by the valve.

The flexible foldable nature of the device according to the present invention enables well established techniques used in cataract surgery to be employed in the treatment of glaucoma. The device according to the present invention may be folded into a cannula and introduced for location into the eye.

6

Intraocular surgery techniques allow a paracentesis (opening onto the anterior chamber from without at the juncture of the cornea and sclera—the limbus) to be performed and the anterior chamber filled with viscoelastic substance. A cyclodialysis instrument is introduced via the paracentesis, with the paracentesis preferably being carried out 180° from the insertion site. A cyclodialysis is carried out, for example by advancing an instrument tip into the angle between the ciliary body and sclera so as to create a cyclodialysis. This is preferably carried out with direct visualisation via gonioscopy lens viewed through an operating microscope. A surgical gonioscopy lens is preferably placed on the cornea while the cyclodialysis is carried out.

The rolled up ocular device is introduced through a cannula, for example using an introducer such as used in cataract surgery or other ocular surgery, from which the device can be detached by pressing a plunger into the introducer when the device has been inserted into the suprachoroidal space created by the cyclodialysis. The tubing of the device is positioned into the interior chamber, and the plate unfolds in the suprachoroidal space to locate the device in the eye. Because of its size, the device cannot fall through the opening through which it was introduced into the suprachoroidal space by the cyclodialysis. The plate therefore keeps the tube in the appropriate position in the anterior chamber allowing controlled aqueous drainage and providing an effective treatment for elevated ocular pressure.

The pressure spike shunt is designed to fit snugly in a paracentesis port that is routinely made during cataract or other ocular surgery. The tubing will not distort the port and there will be no leakage around the port. The outer end of the tube will sit flush on the surface of the cornea—the inner aspect of the tube will preferably just protrude into the anterior chamber—tube length will generally be 1-2 mm and tube diameter is preferably from 0.4-1.2 mm. The tube will contain the same valvular device as contained in the ocular device described above and will open when the intraocular pressure exceeds a predetermined level, preferably 10 mm Hg. At normal ocular pressure the valve will be closed, closing said tube to any fluid communication. FIG. 3 shows a shunt located in a paracentesis port. In most cases the shunt will be removed and discarded at the first post-operative dressing.

The shunt may be inserted into a paracentesis port, or one or more ports, using, for example, a punctum plug inserting instrument such as described in U.S. Pat. No. 5,741,292.

This invention will now be described with reference to the following examples.

EXAMPLE 1

Fresh whole porcine eyes were taken and mounted in a temperature controlled (37°) perfusion chamber. The eyes were perfused with Balanced Salt Solution via a 30 gauge needle inserted via a paracentesis into the anterior chamber. A peristaltic pump was used at a flow rate of 2 μl/min. Intraocular pressure was continuously monitored via a second paracentesis.

Typically intraocular pressures stabilized at 10-15 mm Hg and fell with time (the "washout effect", as glycosan aminoglycans are washed out of the trabecular meshwork with time). Creation of a cyclodialysis (initially with a small spatula, then viscoelastic injection to enlarge the area of detachment of the ciliary body from the sclera) with or without insertion of the device in the cyclodialysis cleft (silicone tubing, length 3 mm, external diameter—1 mm, plate diam-

eter 3 mm) resulted in lower intraocular pressures (below 10 mm Hg) on reperfusion at the same perfusion rate as control eyes.

EXAMPLE 2

Adequate anesthesia is provided to the eye of a glaucoma patient prepared for intraocular surgery. A paracentesis (opening into anterior chamber from without at the junction of the cornea and sclera—the limbus) is performed and the anterior chamber is filled with a viscoelastic substance. A surgical gonioscopy lens is placed on the cornea (or anterior segment endoscope is used) and a cyclodialysis instrument is introduced via the paracentesis—the paracentesis is carried out 180° away from the planned implant insertion site. The cyclodialysis instrument tip is advanced into the angle and pushed into the space between the ciliary body and sclera creating a cycodialysis—this is carried out with direct visualization via the gonioscopy lens viewed through an operating microscope. In order to minimize bleeding, the area in the angle (anterior ciliary body face and overlying trabecular meshwork) can be lasered either preoperatively or at the time of surgery to ablate surface blood vessels).

Through an opening at the tip of the cyclodialysis instrument viscoelastic is inserted to further create a space in the suprachoroidal space. The implant is then introduced—the device is rolled up in the same manner as an ultrathin intraocular lens. The ocular device is attached to an introducer from which it is detached by pushing a plunger in the introducer when the implant is inserted into the suprachoroidal space created by the cyclodialysis instrument and viscoelastic. The tubing is then positioned into the anterior chamber and may be cut to size. The plate unfolds in the suprachoroidal space and because of its size cannot fall through the opening through which it was introduced into the suprachoroidal space. The plate therefore keeps the tube in an appropriate position. The valve is then flushed (with a cannula inserted via the paracentesis) via the tube opening in the anterior chamber. Viscoelastic is then removed from the anterior chamber and antibiotics, steroids and a dressing applied to the eye.

EXAMPLE 3

Fresh whole porcine eyes were taken and mounted in a temperature controlled (37°) perfusion chamber as in Example 1. The eyes were perfused with Balanced Salt Solution via a 30 gauge needle inserted via a paracentesis into the anterior chamber. A peristaltic pump was used at a flow rate of 2 μ l/min. Intraocular pressure was continuously monitored via a second paracentesis.

Typically intraocular pressures stabilized at 10-15 mmHg and fell with time (the “washout effect, as glycoaminoglycans are washed out of the trabecular meshwork with time). Silicone tubing, length 3 mm, external diameter 1 mm was introduced into one paracentesis port. One end of the port (outer end) was flush with the cornea and the inner end of the port extended slightly into the anterior chamber. Intraocular pressure did not exceed 10 mm Hg.

The invention claimed is:

1. An ocular pressure spike shunt for insertion into an ocular paracentesis incision port following ocular surgery, comprising a flexible fluid transfer tube formed of biocompatible material so as to allow paracentesis incision closure

around said tube, having a distal region on an inner surface of a cornea and a proximal end on an outer surface of the cornea, a tubular portion having a lumen, the tubular portion disposed between said distal region and said proximal end to allow fluid communication through said tube, wherein said distal region and proximal end both have an enlarged diameter relative to a diameter of a central section of said shunt, and wherein said distal region is cone-shaped such that the diameter of the distal region gradually reduces moving in a distal direction and a distal-most end of the distal region is curved, said lumen containing a slit valve for controlling pressure in the eye following ocular surgery, which valve opens permitting fluid flow through said tube when a predetermined pressure is exceeded, said shunt being configured such that on insertion into a paracentesis port said proximal end is flush with the outer surface of the cornea and a portion of the enlarged diameter of the distal region is positioned flat against the inner surface of the cornea, and said distal region opens into the anterior chamber of the eye, wherein the tube is removable from the eye, the proximal end having a foldable plate that lies flush with the surface of the cornea when the pressure spike shunt is implanted, the foldable plate being perpendicular to the tubular portion, wherein the ocular pressure spike shunt, including the foldable plate, is foldable to a diameter of 1.5 mm or less.

2. A shunt according to claim 1 wherein said predetermined pressure is 10 mm Hg.

3. An ocular pressure spike shunt as in claim 1, wherein the tubular portion has a length equal to a thickness of the cornea.

4. A method for preventing ocular pressure spikes following ocular surgery wherein a paracentesis incision port is formed in the eye during said surgery, comprising introducing an ocular pressure spike shunt into said paracentesis port at the conclusion of ocular surgery, said shunt comprising a flexible fluid transfer tube formed of biocompatible material so as to allow paracentesis incision closure around said tube, having a distal region on an inner surface of a cornea and a proximal end on an outer surface of the cornea, a tubular portion having a lumen, the tubular portion disposed between said distal region and said proximal end to allow fluid communication through said tube, wherein said distal region and proximal end both have an enlarged diameter relative to a diameter of a central section of said shunt, and wherein said distal region is cone-shaped such that the diameter of the distal region gradually reduces moving in a distal direction and a distal-most end of the distal region is curved, said lumen containing a slit valve for controlling pressure in the eye following ocular surgery, which valve opens permitting fluid flow through said tube when a predetermined pressure is exceeded, said shunt being configured such that on insertion into a paracentesis port said proximal end is substantially flush with the outer surface of the cornea and a portion of the enlarged diameter of the distal region is positioned flat against the inner surface of the cornea, and said distal region extends into the anterior chamber of the eye, wherein the tube is removable from the eye, the proximal end having a foldable plate that lies flush with the surface of the cornea when the pressure spike shunt is implanted, the foldable plate being perpendicular to the tubular portion, wherein the ocular pressure spike shunt, including the foldable plate, is foldable to a diameter of 1.5 mm or less.

* * * * *