



(86) Date de dépôt PCT/PCT Filing Date: 2010/05/07
(87) Date publication PCT/PCT Publication Date: 2010/11/11
(45) Date de délivrance/Issue Date: 2023/03/14
(85) Entrée phase nationale/National Entry: 2011/11/07
(86) N° demande PCT/PCT Application No.: US 2010/034026
(87) N° publication PCT/PCT Publication No.: 2010/129861
(30) Priorités/Priorities: 2009/05/08 (US61/176,594);
2010/03/25 (US61/317,350)

(51) Cl.Int./Int.Cl. *C12N 15/113* (2010.01),
A61K 31/713 (2006.01), *A61K 48/00* (2006.01),
C12N 15/12 (2006.01), *C12N 15/63* (2006.01)
(72) Inventeurs/Inventors:
COLLARD, JOSEPH, US;
KHORKOVA SHERMAN, OLGA, US
(73) Propriétaire/Owner:
CURNA, INC., US
(74) Agent: NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : TRAITEMENT DE MALADIES LIEES A LA FAMILLE DE LA DYSTROPHINE PAR INHIBITION DU PRODUIT
ANTISENS NATUREL DE TRANSCRIPTION VERS LA FAMILLE DE LA DMD
(54) Title: TREATMENT OF DYSTROPHIN FAMILY RELATED DISEASES BY INHIBITION OF NATURAL ANTISENSE
TRANSCRIPT TO DMD FAMILY

(57) **Abrégé/Abstract:**

The present invention relates to antisense oligonucleotides that modulate the expression of and/or function of Dystrophin family, in particular, by targeting natural antisense polynucleotides of Dystrophin family. The invention also relates to the identification of these antisense oligonucleotides and their use in treating diseases and disorders associated with the expression of DMD family.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
11 November 2010 (11.11.2010)

PCT

(10) International Publication Number
WO 2010/129861 A3

(51) International Patent Classification:

C12N 15/113 (2010.01) A61K 31/713 (2006.01)
C12N 15/63 (2006.01) A61K 48/00 (2006.01)

(21) International Application Number:

PCT/US2010/034026

(22) International Filing Date:

7 May 2010 (07.05.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/176,594 8 May 2009 (08.05.2009) US
61/317,350 25 March 2010 (25.03.2010) US

(71) Applicant (for all designated States except US): **CUR-NA, INC.** [US/US]; 555 Heritage Drive, Suite 110, Jupiter, Florida 33458 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **COLLARD, Joseph** [US/US]; 1004, Brooks Lane, Delray Beach, Florida 33483 (US). **KHORKOVA SHERMAN, Olga** [US/US]; 18288 SE, Heritage Drive, Tequesta, Florida 33469 (US).

(74) Agent: **GUISE, Jeffrey**; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, CA 94304 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,

CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

12 May 2011

(54) Title: TREATMENT OF DYSTROPHIN FAMILY RELATED DISEASES BY INHIBITION OF NATURAL ANTISENSE TRANSCRIPT TO DMD FAMILY

(57) Abstract: The present invention relates to antisense oligonucleotides that modulate the expression of and/or function of Dystrophin family, in particular, by targeting natural antisense polynucleotides of Dystrophin family. The invention also relates to the identification of these antisense oligonucleotides and their use in treating diseases and disorders associated with the expression of DMD family.



WO 2010/129861 A3

**TREATMENT OF DYSTROPHIN FAMILY RELATED DISEASES BY INHIBITION OF NATURAL
ANTISENSE TRANSCRIPT TO DMD FAMILY
FIELD OF THE INVENTION**

5

[0002] Embodiments of the invention comprise oligonucleotides modulating expression and/or function of DMD family and associated molecules.

BACKGROUND

10 [0003] DNA-RNA and RNA-RNA hybridization are important to many aspects of nucleic acid function including DNA replication, transcription, and translation. Hybridization is also central to a variety of technologies that either detect a particular nucleic acid or alter its expression. Antisense nucleotides, for example, disrupt gene expression by hybridizing to target RNA, thereby interfering with RNA splicing, transcription, translation, and replication. Antisense DNA has the added feature that DNA-RNA hybrids serve as a substrate for digestion by ribonuclease H, an activity
15 that is present in most cell types. Antisense molecules can be delivered into cells, as is the case for oligodeoxynucleotides (ODNs), or they can be expressed from endogenous genes as RNA molecules. The FDA recently approved an antisense drug, VITRAVENE™ (for treatment of cytomegalovirus retinitis), reflecting that antisense has therapeutic utility.

SUMMARY

20 [0004] This Summary is provided to present a summary of the invention to briefly indicate the nature and substance of the invention. It is submitted with the understanding that it will not be used to interpret or limit the scope or meaning of the claims.

[0005] In one embodiment, the invention provides methods for inhibiting the action of a natural antisense transcript by using antisense oligonucleotide(s) targeted to any region of the natural antisense transcript resulting in up-regulation of
25 the corresponding sense gene. It is also contemplated herein that inhibition of the natural antisense transcript can be achieved by siRNA, ribozymes and small molecules, which are considered to be within the scope of the present invention.

[0006] One embodiment provides a method of modulating function and/or expression of an DMD family polynucleotide in patient cells or tissues in vivo or in vitro comprising contacting said cells or tissues with an antisense
30 oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to a reverse complement of a polynucleotide comprising 5 to 30 consecutive nucleotides within nucleotides 1 to 378 of SEQ ID NO: 3, 1 to 294 of SEQ ID NO: 4, 1 to 686 of SEQ ID NO: 5, 1 to 480 of SEQ ID NO: 6 and 1 to 501 of SEQ

ID NO: 7 (Figure 3) thereby modulating function and/or expression of the DMD family polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0007] In another preferred embodiment, an oligonucleotide targets a natural antisense sequence of DMD family polynucleotides, for example, nucleotides set forth in SEQ ID NO: 3 to 7, and any variants, alleles, homologs, mutants, derivatives, fragments and complementary sequences thereto. Examples of antisense oligonucleotides are set forth as SEQ ID NOS: 8 to 22 (Figure 4 and 5).

[0008] Another embodiment provides a method of modulating function and/or expression of an DMD family polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising contacting said cells or tissues with an antisense oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to a reverse complement of the an antisense of the DMD family polynucleotide; thereby modulating function and/or expression of the DMD family polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0009] Another embodiment provides a method of modulating function and/or expression of an DMD family polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising contacting said cells or tissues with an antisense oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to an antisense oligonucleotide to an DMD family antisense polynucleotide; thereby modulating function and/or expression of the DMD family polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0010] In a preferred embodiment, a composition comprises one or more antisense oligonucleotides which bind to sense and/or antisense DMD family polynucleotides.

[0011] In another preferred embodiment, the oligonucleotides comprise one or more modified or substituted nucleotides.

[0012] In another preferred embodiment, the oligonucleotides comprise one or more modified bonds.

[0013] In yet another embodiment, the modified nucleotides comprise modified bases comprising phosphorothioate, methylphosphonate, peptide nucleic acids, 2'-O-methyl, fluoro- or carbon, methylene or other locked nucleic acid (LNA) molecules. Preferably, the modified nucleotides are locked nucleic acid molecules, including α -L-LNA.

[0014] In another preferred embodiment, the oligonucleotides are administered to a patient subcutaneously, intramuscularly, intravenously or intraperitoneally.

[0015] In another preferred embodiment, the oligonucleotides are administered in a pharmaceutical composition. A treatment regimen comprises administering the antisense compounds at least once to patient; however, this treatment can be modified to include multiple doses over a period of time. The treatment can be combined with one or more other types of therapies.

[0016] In another preferred embodiment, the oligonucleotides are encapsulated in a liposome or attached to a carrier molecule (e.g. cholesterol, TAT peptide).

[0017] Other aspects are described *infra*.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018] Figure 1:

Figure 1A is a graph of real time PCR results showing the fold change + standard deviation in DMD family mRNA after treatment of 518A2 cells with phosphorothioate oligonucleotides introduced using Lipofectamine 2000, as compared to control. Real time PCR results show that the levels of DMD family mRNA in 518A2 cells are significantly increased 48 h after treatment with two of the siRNAs designed to DMD family antisense BG208074. Bars denoted as CUR-0636 to CUR-0654, correspond to samples treated with SEQ ID NOS: 8 to 17 respectively.

Figure 1B is a graph of real time PCR results showing the fold change + standard deviation in DMD family mRNA after treatment of 518A2 cells with phosphorothioate oligonucleotides introduced using Lipofectamine 2000, as compared to control. Treatment with siRNAs to other antisense molecules, BF838561, BF768753 and BF950643, did not elevate DMD family mRNA levels. Bars denoted as CUR-0638, CUR-0648, CUR-0646 and CUR-0652 correspond to samples treated with SEQ ID NOS: 9, 14, 13 and 16 respectively.

Figure 1C is a graph of real time PCR results showing the fold change + standard deviation in UTRN mRNA after treatment of MCF-7 cells with phosphorothioate oligonucleotides introduced using Lipofectamine 2000, as compared to control. Bars denoted as CUR-1443 to CUR-1447 correspond to samples treated with SEQ ID NOS: 18 to 22 respectively.

[0019] Figure 2 shows

SEQ ID NO: 1: Homo sapiens Dystrophin family, transcript variant Dp427m, mRNA (NCBI Accession No.: NM_004006).

SEQ ID NO: 2: Homo sapiens Utrophin (UTRN), mRNA. (NCBI Accession No.: NM_007124)

[0020] Figure 3 shows

SEQ ID NO: 3: Natural DMD family antisense sequence (BF838561)

SEQ ID NO: 4: Natural DMD family antisense sequence (BG208074)

SEQ ID NO: 5: Natural DMD family antisense sequence (BF950643)

SEQ ID NO: 6: Natural DMD family antisense sequence (BF768753)

SEQ ID NO: 7: Natural UTRN antisense sequence (ENST00000431309).

[0021] Figure 4 shows DMD antisense oligonucleotides, SEQ ID NOS: 8 to 17. 'r' indicates RNA.

[0022] Figure 5 shows the UTRN antisense oligonucleotides, SEQ ID NOS: 18 to 22. * indicates phosphothioate bond.

[0023] Figure 6 shows the DMD sense oligonucleotides, SEQ ID NOS: 23 to 32. The sense oligonucleotides SEQ ID NO: 23 to 32 are the reverse complements of the antisense oligonucleotides SEQ ID NO: 8 to 17 respectively. 'r' indicates RNA.

DETAILED DESCRIPTION

[0024] Several aspects of the invention are described below with reference to example applications for illustration. It should be understood that numerous specific details, relationships, and methods are set forth to provide a full understanding of the invention. One having ordinary skill in the relevant art, however, will readily recognize that the invention can be practiced without one or more of the specific details or with other methods. The present invention is not limited by the ordering of acts or events, as some acts may occur in different orders and/or concurrently with other acts or events. Furthermore, not all illustrated acts or events are required to implement a methodology in accordance with the present invention.

[0025] All genes, gene names, and gene products disclosed herein are intended to correspond to homologs from any species for which the compositions and methods disclosed herein are applicable. Thus, the terms include, but are not limited to genes and gene products from humans and mice. It is understood that when a gene or gene product from a particular species is disclosed, this disclosure is intended to be exemplary only, and is not to be interpreted as a limitation unless the context in which it appears clearly indicates. Thus, for example, for the genes disclosed herein, which in some embodiments relate to mammalian nucleic acid and amino acid sequences are intended to encompass homologous and/or orthologous genes and gene products from other animals including, but not limited to other mammals, fish, amphibians, reptiles, and birds. In preferred embodiments, the genes or nucleic acid sequences are human.

Definitions

[0026] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention. As used herein, the singular forms "a", "an" and "the" are intended to include the plural forms as well, unless the context clearly indicates otherwise. Furthermore, to the extent that the terms "including", "includes", "having", "has", "with", or variants thereof are used in either the detailed description and/or the claims, such terms are intended to be inclusive in a manner similar to the term "comprising."

[0027] The term "about" or "approximately" means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. For example, "about" can mean within 1 or more than 1 standard deviation, per the practice in the art. Alternatively, "about" can mean a range of up to 20%, preferably up to 10%, more preferably up to 5%, and more preferably still up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value. Where particular values are described in the application and claims, unless otherwise stated the term "about" meaning within an acceptable error range for the particular value should be assumed.

[0028] As used herein, the term "mRNA" means the presently known mRNA transcript(s) of a targeted gene, and any further transcripts which may be elucidated.

[0029] By "antisense oligonucleotides" or "antisense compound" is meant an RNA or DNA molecule that binds to another RNA or DNA (target RNA, DNA). For example, if it is an RNA oligonucleotide it binds to another RNA target by means of RNA-RNA interactions and alters the activity of the target RNA (Eguchi *et al.*, (1991) *Ann. Rev. Biochem.* 60, 631-652). An antisense oligonucleotide can upregulate or downregulate expression and/or function of a particular polynucleotide. The definition is meant to include any foreign RNA or DNA molecule which is useful from a therapeutic, diagnostic, or other viewpoint. Such molecules include, for example, antisense RNA or DNA molecules, interference RNA (RNAi), micro RNA, decoy RNA molecules, siRNA, enzymatic RNA, therapeutic editing RNA and agonist and antagonist RNA, antisense oligomeric compounds, antisense oligonucleotides, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other oligomeric compounds that hybridize to at least a portion of the target nucleic acid. As such, these compounds may be introduced in the form of single-stranded, double-stranded, partially single-stranded, or circular oligomeric compounds.

[0030] In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. The term "oligonucleotide", also includes linear or circular oligomers of natural and/or modified monomers or linkages, including deoxyribonucleosides, ribonucleosides, substituted and alpha-anomeric forms thereof, peptide nucleic acids (PNA), locked nucleic acids (LNA), phosphorothioate, methylphosphonate, and the like. Oligonucleotides are capable of specifically binding to a target polynucleotide by way of a regular pattern of monomer-to-monomer interactions, such as Watson-Crick type of base pairing, Hoogsteen or reverse Hoogsteen types of base pairing, or the like.

[0031] The oligonucleotide may be "chimeric", that is, composed of different regions. In the context of this invention "chimeric" compounds are oligonucleotides, which contain two or more chemical regions, for example, DNA region(s), RNA region(s), PNA region(s) etc. Each chemical region is made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotides compound. These oligonucleotides typically comprise at least one region wherein the oligonucleotide is modified in order to exhibit one or more desired properties. The desired properties of the oligonucleotide include, but are not limited, for example, to increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. Different regions of the oligonucleotide may therefore have different properties. The chimeric oligonucleotides of the present invention can be formed as mixed structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide analogs as described above.

[0032] The oligonucleotide can be composed of regions that can be linked in "register", that is, when the monomers are linked consecutively, as in native DNA, or linked via spacers. The spacers are intended to constitute a covalent "bridge" between the regions and have in preferred cases a length not exceeding about 100 carbon atoms. The spacers may carry different functionalities, for example, having positive or negative charge, carry special nucleic acid binding

properties (intercalators, groove binders, toxins, fluorophors etc.), being lipophilic, inducing special secondary structures like, for example, alanine containing peptides that induce alpha-helices.

[0033] As used herein "DMD family", "Dystrophin family" and "dystrophin-related protein family", "dystrophin gene family" are inclusive of all family members, mutants, alleles, fragments, species, coding and noncoding sequences, sense and antisense polynucleotide strands, etc.

[0034] As used herein, the words Dystrophin, DMD, BMD, CMD3B, DXS142, DXS164, DXS206, DXS230, DXS239, DXS268, DXS269, DXS270 and DXS272 are used interchangeably in the present application.

[0035] As used herein, the words Utrophin, UTRN, DMDL, DRP, DRP1, Dystrophin-related protein 1, FLJ23678, are used interchangeably in the present application.

[0036] As used herein, the words "dystrophin related protein 2", "dystrophin-related protein 2" and DRP2 are used interchangeably in the present application.

[0037] As used herein, the words Dystrobrevin α -dystrobrevin, β -dystrobrevin, DTNA and DTNB are used interchangeably in the present application.

[0038] As used herein, the term "oligonucleotide specific for" or "oligonucleotide which targets" refers to an oligonucleotide having a sequence (i) capable of forming a stable complex with a portion of the targeted gene, or (ii) capable of forming a stable duplex with a portion of a mRNA transcript of the targeted gene. Stability of the complexes and duplexes can be determined by theoretical calculations and/or in vitro assays. Exemplary assays for determining stability of hybridization complexes and duplexes are described in the Examples below.

[0039] As used herein, the term "target nucleic acid" encompasses DNA, RNA (comprising premRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA, coding, noncoding sequences, sense or antisense polynucleotides. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds, which specifically hybridize to it, is generally referred to as "antisense". The functions of DNA to be interfered include, for example, replication and transcription. The functions of RNA to be interfered, include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of an encoded product or oligonucleotides.

[0040] RNA interference "RNAi" is mediated by double stranded RNA (dsRNA) molecules that have sequence-specific homology to their "target" nucleic acid sequences (Caplen, N. J., *et al.* (2001) *Proc. Natl. Acad. Sci. USA* 98:9742-9747). In certain embodiments of the present invention, the mediators are 5-25 nucleotide "small interfering" RNA duplexes (siRNAs). The siRNAs are derived from the processing of dsRNA by an RNase enzyme known as Dicer (Bernstein, E., *et al.* (2001) *Nature* 409:363-366). siRNA duplex products are recruited into a multi-protein

siRNA complex termed RISC (RNA Induced Silencing Complex). Without wishing to be bound by any particular theory, a RISC is then believed to be guided to a target nucleic acid (suitably mRNA), where the siRNA duplex interacts in a sequence-specific way to mediate cleavage in a catalytic fashion (Bernstein, E., *et al.* (2001) *Nature* 409:363-366; Boutfa, A., *et al.* (2001) *Curr. Biol.* 11:1776-1780). Small interfering RNAs that can be used in accordance with the present invention can be synthesized and used according to procedures that are well known in the art and that will be familiar to the ordinarily skilled artisan. Small interfering RNAs for use in the methods of the present invention suitably comprise between about 1 to about 50 nucleotides (nt). In examples of non limiting embodiments, siRNAs can comprise about 5 to about 40 nt, about 5 to about 30 nt, about 10 to about 30 nt, about 15 to about 25 nt, or about 20-25 nucleotides.

[0041] Selection of appropriate oligonucleotides is facilitated by using computer programs that automatically align nucleic acid sequences and indicate regions of identity or homology. Such programs are used to compare nucleic acid sequences obtained, for example, by searching databases such as GenBank or by sequencing PCR products. Comparison of nucleic acid sequences from a range of species allows the selection of nucleic acid sequences that display an appropriate degree of identity between species. In the case of genes that have not been sequenced, Southern blots are performed to allow a determination of the degree of identity between genes in target species and other species. By performing Southern blots at varying degrees of stringency, as is well known in the art, it is possible to obtain an approximate measure of identity. These procedures allow the selection of oligonucleotides that exhibit a high degree of complementarity to target nucleic acid sequences in a subject to be controlled and a lower degree of complementarity to corresponding nucleic acid sequences in other species. One skilled in the art will realize that there is considerable latitude in selecting appropriate regions of genes for use in the present invention.

[0042] By "enzymatic RNA" is meant an RNA molecule with enzymatic activity (Cech, (1988) *J. American. Med. Assoc.* 260, 3030-3035). Enzymatic nucleic acids (ribozymes) act by first binding to a target RNA. Such binding occurs through the target binding portion of an enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through base pairing, and once bound to the correct site, acts enzymatically to cut the target RNA.

[0043] By "decoy RNA" is meant an RNA molecule that mimics the natural binding domain for a ligand. The decoy RNA therefore competes with natural binding target for the binding of a specific ligand. For example, it has been shown that over-expression of HIV trans-activation response (TAR) RNA can act as a "decoy" and efficiently binds HIV tat protein, thereby preventing it from binding to TAR sequences encoded in the HIV RNA (Sullenger *et al.* (1990) *Cell*, 63, 601- 608). This is meant to be a specific example. Those in the art will recognize that this is but one example, and other embodiments can be readily generated using techniques generally known in the art.

[0044] As used herein, the term "monomers" typically indicates monomers linked by phosphodiester bonds or analogs thereof to form oligonucleotides ranging in size from a few monomeric units, e.g., from about 3-4, to about several

hundreds of monomeric units. Analogs of phosphodiester linkages include: phosphorothioate, phosphorodithioate, methylphosphonates, phosphoroselenoate, phosphoramidate, and the like, as more fully described below.

[0045] The term "nucleotide" covers naturally occurring nucleotides as well as nonnaturally occurring nucleotides. It should be clear to the person skilled in the art that various nucleotides which previously have been considered "non-naturally occurring" have subsequently been found in nature. Thus, "nucleotides" includes not only the known purine and pyrimidine heterocycles-containing molecules, but also heterocyclic analogues and tautomers thereof. Illustrative examples of other types of nucleotides are molecules containing adenine, guanine, thymine, cytosine, uracil, purine, xanthine, diaminopurine, 8-oxo- N6-methyladenine, 7-deazaxanthine, 7-deazaguanine, N4,N4-ethanocytosin, N6,N6-ethano-2,6- diaminopurine, 5-methylcytosine, 5-(C3-C6)-alkynylcytosine, 5-fluorouracil, 5-bromouracil, pseudoisocytosine, 2-hydroxy-5-methyl-4-triazolopyridin, isocytosine, isoguanin, inosine and the "non-naturally occurring" nucleotides described in Benner *et al.*, U.S. Pat No. 5,432,272. The term "nucleotide" is intended to cover every and all of these examples as well as analogues and tautomers thereof. Especially interesting nucleotides are those containing adenine, guanine, thymine, cytosine, and uracil, which are considered as the naturally occurring nucleotides in relation to therapeutic and diagnostic application in humans. Nucleotides include the natural 2'-deoxy and 2'-hydroxyl sugars, e.g., as described in Kornberg and Baker, *DNA Replication*, 2nd Ed. (Freeman, San Francisco, 1992) as well as their analogs.

[0046] "Analog" in reference to nucleotides includes synthetic nucleotides having modified base moieties and/or modified sugar moieties (see e.g., described generally by Scheit, *Nucleotide Analogs*, John Wiley, New York, 1980; Freier & Altmann, (1997) *Nucl. Acid. Res.*, 25(22), 4429- 4443; Toulmé, J.J., (2001) *Nature Biotechnology* 19:17-18; Manoharan M., (1999) *Biochemica et Biophysica Acta* 1489:117-139; Freier S. M., (1997) *Nucleic Acid Research*, 25:4429-4443, Uhlman, E., (2000) *Drug Discovery & Development*, 3: 203-213, Herdewin P., (2000) *Antisense & Nucleic Acid Drug Dev.*, 10:297-310; 2'-O, 3'-C-linked [3.2.0] bicycloarabinonucleosides (see e.g. N.K Christensen., *et al.*, (1998) *J. Am. Chem. Soc.*, 120: 5458-5463; Prakash TP, Bhat B. (2007) *Curr Top Med Chem.* 7(7):641-9; Cho EJ, *et al.* (2009) *Annual Review of Analytical Chemistry*, 2, 241-264). Such analogs include synthetic nucleotides designed to enhance binding properties, e.g., duplex or triplex stability, specificity, or the like.

[0047] As used herein, "hybridization" means the pairing of substantially complementary strands of oligomeric compounds. One mechanism of pairing involves hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases (nucleotides) of the strands of oligomeric compounds. For example, adenine and thymine are complementary nucleotides which pair through the formation of hydrogen bonds. Hybridization can occur under varying circumstances.

[0048] An antisense compound is "specifically hybridizable" when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a modulation of function and/or activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic

acid sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and under conditions in which assays are performed in the case of in vitro assays.

[0049] As used herein, the phrase "stringent hybridization conditions" or "stringent conditions" refers to conditions under which a compound of the invention will hybridize to its target sequence, but to a minimal number of other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances and in the context of this invention, "stringent conditions" under which oligomeric compounds hybridize to a target sequence are determined by the nature and composition of the oligomeric compounds and the assays in which they are being investigated. In general, stringent hybridization conditions comprise low concentrations ($<0.15M$) of salts with inorganic cations such as Na^{++} or K^{++} (i.e., low ionic strength), temperature higher than $20^{\circ}C - 25^{\circ}C$, below the T_m of the oligomeric compound:target sequence complex, and the presence of denaturants such as formamide, dimethylformamide, dimethyl sulfoxide, or the detergent sodium dodecyl sulfate (SDS). For example, the hybridization rate decreases 1.1% for each 1% formamide. An example of a high stringency hybridization condition is 0.1X sodium chloride-sodium citrate buffer (SSC)/0.1% (w/v) SDS at $60^{\circ}C$ for 30 minutes.

[0050] "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides on one or two oligomeric strands. For example, if a nucleobase at a certain position of an antisense compound is capable of hydrogen bonding with a nucleobase at a certain position of a target nucleic acid, said target nucleic acid being a DNA, RNA, or oligonucleotide molecule, then the position of hydrogen bonding between the oligonucleotide and the target nucleic acid is considered to be a complementary position. The oligomeric compound and the further DNA, RNA, or oligonucleotide molecule are complementary to each other when a sufficient number of complementary positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of precise pairing or complementarity over a sufficient number of nucleotides such that stable and specific binding occurs between the oligomeric compound and a target nucleic acid.

[0051] It is understood in the art that the sequence of an oligomeric compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. Moreover, an oligonucleotide may hybridize over one or more segments such that intervening or adjacent segments are not involved in the hybridization event (e.g., a loop structure, mismatch or hairpin structure). The oligomeric compounds of the present invention comprise at least about 70%, or at least about 75%, or at least about 80%, or at least about 85%, or at least about 90%, or at least about 95%, or at least about 99% sequence complementarity to a target region within the target nucleic acid sequence to which they are targeted. For example, an antisense compound in which 18 of 20 nucleotides of the antisense compound are complementary to a target region, and would therefore specifically hybridize, would represent 90 percent complementarity. In this example, the remaining noncomplementary nucleotides may be clustered or interspersed with

complementary nucleotides and need not be contiguous to each other or to complementary nucleotides. As such, an antisense compound which is 18 nucleotides in length having 4 (four) noncomplementary nucleotides which are flanked by two regions of complete complementarity with the target nucleic acid would have 77.8% overall complementarity with the target nucleic acid and would thus fall within the scope of the present invention. Percent complementarity of an antisense compound with a region of a target nucleic acid can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul *et al.*, (1990) *J. Mol. Biol.*, 215, 403-410; Zhang and Madden, (1997) *Genome Res.*, 7, 649-656). Percent homology, sequence identity or complementarity, can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), using default settings, which uses the algorithm of Smith and Waterman (*Adv. Appl. Math.*, (1981) 2, 482-489).

[0052] As used herein, the term "Thermal Melting Point (T_m)" refers to the temperature, under defined ionic strength, pH, and nucleic acid concentration, at which 50% of the oligonucleotides complementary to the target sequence hybridize to the target sequence at equilibrium. Typically, stringent conditions will be those in which the salt concentration is at least about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short oligonucleotides (e.g., 10 to 50 nucleotide). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

[0053] As used herein, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene.

[0054] The term "variant," when used in the context of a polynucleotide sequence, may encompass a polynucleotide sequence related to a wild type gene. This definition may also include, for example, "allelic," "splice," "species," or "polymorphic" variants. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or an absence of domains. Species variants are polynucleotide sequences that vary from one species to another. Of particular utility in the invention are variants of wild type gene products. Variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes that give rise to variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

[0055] The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs,) or single base

mutations in which the polynucleotide sequence varies by one base. The presence of SNPs may be indicative of, for example, a certain population with a propensity for a disease state, that is susceptibility versus resistance.

[0056] Derivative polynucleotides include nucleic acids subjected to chemical modification, for example, replacement of hydrogen by an alkyl, acyl, or amino group. Derivatives, e.g., derivative oligonucleotides, may comprise non-naturally-occurring portions, such as altered sugar moieties or inter-sugar linkages. Exemplary among these are phosphorothioate and other sulfur containing species which are known in the art. Derivative nucleic acids may also contain labels, including radionucleotides, enzymes, fluorescent agents, chemiluminescent agents, chromogenic agents, substrates, cofactors, inhibitors, magnetic particles, and the like.

[0057] A "derivative" polypeptide or peptide is one that is modified, for example, by glycosylation, pegylation, phosphorylation, sulfation, reduction/alkylation, acylation, chemical coupling, or mild formalin treatment. A derivative may also be modified to contain a detectable label, either directly or indirectly, including, but not limited to, a radioisotope, fluorescent, and enzyme label.

[0058] As used herein, the term "animal" or "patient" is meant to include, for example, humans, sheep, elks, deer, mule deer, minks, mammals, monkeys, horses, cattle, pigs, goats, dogs, cats, rats, mice, birds, chicken, reptiles, fish, insects and arachnids.

[0059] "Mammal" covers warm blooded mammals that are typically under medical care (e.g., humans and domesticated animals). Examples include feline, canine, equine, bovine, and human, as well as just human.

[0060] "Treating" or "treatment" covers the treatment of a disease-state in a mammal, and includes: (a) preventing the disease-state from occurring in a mammal, in particular, when such mammal is predisposed to the disease-state but has not yet been diagnosed as having it; (b) inhibiting the disease-state, e.g., arresting its development; and/or (c) relieving the disease-state, e.g., causing regression of the disease state until a desired endpoint is reached. Treating also includes the amelioration of a symptom of a disease (e.g., lessen the pain or discomfort), wherein such amelioration may or may not be directly affecting the disease (e.g., cause, transmission, expression, etc.).

[0061] As used herein, "cancer" refers to all types of cancer or neoplasm or malignant tumors found in mammals, including, but not limited to: leukemias, lymphomas, melanomas, carcinomas and sarcomas. The cancer manifests itself as a "tumor" or tissue comprising malignant cells of the cancer. Examples of tumors include sarcomas and carcinomas such as, but not limited to: fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung

carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma. Additional cancers which can be treated by the disclosed composition according to the invention include but not limited to, for example, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, neuroblastoma, breast cancer, ovarian cancer, lung cancer, rhabdomyosarcoma, primary thrombocytosis, primary macroglobulinemia, small-cell lung tumors, primary brain tumors, stomach cancer, colon cancer, malignant pancreatic insulanoma, malignant carcinoid, urinary bladder cancer, premalignant skin lesions, testicular cancer, lymphomas, thyroid cancer, neuroblastoma, esophageal cancer, genitourinary tract cancer, malignant hypercalcemia, cervical cancer, endometrial cancer, adrenal cortical cancer, and prostate cancer.

[0062] As used herein, a "muscle disease or disorder" includes but is not limited to muscular dystrophy (MD), a muscle-wasting disease, inflammatory myopathy or myositis (including, for example, dermatomyositis, polymyositis, inclusion body myotosis), myotubular myopathy, nemaline myopathy, a desmin related myopathy, Marfan myopathy, a mitochondrial myopathy etc.. As used herein, muscular dystrophy refers to a group of genetic, hereditary muscle diseases that cause progressive muscle weakness and which may be characterized by progressive skeletal muscle weakness, defects in muscle proteins, and the death of muscle cells and tissue. Muscular dystrophy includes, but is not limited to Duchenne (DMD family), Becker (BMD), Spinal Muscular Atrophy, Spinal bulbar muscular atrophy, dystrophinopathy, sarcoglycanopathy, limb girdle muscular dystrophy (LGMD), congenital muscular dystrophy (CMD), facioscapulohumeral (FSHD), myotonic, oculopharyngeal, distal, and Emery- Dreifuss.

[0063] "Neurological disease or disorder" refers to any disease or disorder of the nervous system and/or visual system.

"Neurological disease or disorder" include disease or disorders that involve the central nervous system (brain, brainstem and cerebellum), the peripheral nervous system (including cranial nerves), and the autonomic nervous system (parts of which are located in both central and peripheral nervous system). Examples of neurological disorders include but are not limited to, headache, stupor and coma, dementia, seizure, sleep disorders, trauma, infections, neoplasms, neuroophthalmology, movement disorders, demyelinating diseases, spinal cord disorders, and disorders of peripheral nerves, muscle and neuromuscular junctions. Addiction and mental illness, include, but are not limited to, bipolar disorder and schizophrenia, are also included in the definition of neurological disorder. The following is a list of several neurological disorders, symptoms, signs and syndromes that can be treated using compositions and methods according to the present invention: acquired epileptiform aphasia; acute disseminated encephalomyelitis; adrenoleukodystrophy; age-related macular degeneration; agenesis of the corpus callosum; agnosia; Aicardi syndrome; Alexander disease; Alpers' disease; alternating hemiplegia; Vascular dementia; amyotrophic lateral sclerosis; anencephaly; Angelman syndrome; angiomatosis; anoxia; aphasia; apraxia; arachnoid cysts; arachnoiditis; Arnold-Chiari malformation; arteriovenous malformation; Asperger syndrome; ataxia telegiectasia; attention deficit hyperactivity disorder; autism; autonomic dysfunction; back pain; Batten disease; Behcet's disease; Bell's palsy; benign

essential blepharospasm; benign focal; amyotrophy; benign intracranial hypertension; Binswanger's disease; blepharospasm; Bloch Sulzberger syndrome; brachial plexus injury; brain abscess; brain injury; brain tumors (including glioblastoma multiforme); spinal tumor; Brown-Sequard syndrome; Canavan disease; carpal tunnel syndrome; causalgia; central pain syndrome; central pontine myelinolysis; cephalic disorder; cerebral aneurysm; cerebral arteriosclerosis; cerebral atrophy; cerebral gigantism; cerebral palsy; Charcot-Marie-Tooth disease; chemotherapy-induced neuropathy and neuropathic pain; Chiari malformation; chorea; chronic inflammatory demyelinating polyneuropathy; chronic pain; chronic regional pain syndrome; Coffin Lowry syndrome; coma, including persistent vegetative state; congenital facial diplegia; corticobasal degeneration; cranial arteritis; craniosynostosis; Creutzfeldt-Jakob disease; cumulative trauma disorders; Cushing's syndrome; cytomegalic inclusion body disease; cytomegalovirus infection; dancing eyes-dancing feet syndrome; DandyWalker syndrome; Dawson disease; De Morsier's syndrome; Dejerine-Klumpke palsy; dementia; dermatomyositis; diabetic neuropathy; diffuse sclerosis; dysautonomia; dysgraphia; dyslexia; dystonias; early infantile epileptic encephalopathy; empty sella syndrome; encephalitis; encephaloceles; encephalotrigeminal angiomatosis; epilepsy; Erb's palsy; essential tremor; Fabry's disease; Fahr's syndrome; fainting; familial spastic paralysis; febrile seizures; Fisher syndrome; Friedreich's ataxia; fronto-temporal dementia and other "tauopathies"; Gaucher's disease; Gerstmann's syndrome; giant cell arteritis; giant cell inclusion disease; globoid cell leukodystrophy; Guillain-Barre syndrome; HTLV-1-associated myelopathy; Hallervorden-Spatz disease; head injury; headache; hemifacial spasm; hereditary spastic paraplegia; hereditary ataxia; hereditary polyneuritis; herpes zoster ophthalmicus; herpes zoster; Hirayama syndrome; HIV-associated dementia and neuropathy (also neurological manifestations of AIDS); holoprosencephaly; Huntington's disease and other polyglutamine repeat diseases; hydranencephaly; hydrocephalus; hypercortisolism; hypoxia; immune-mediated encephalomyelitis; inclusion body myositis; incontinentia pigmenti; infantile phytanic acid storage disease; infantile Refsum disease; infantile spasms; inflammatory myopathy; intracranial cyst; intracranial hypertension; Joubert syndrome; Keams-Sayre syndrome; Kennedy disease; Kinsbourne syndrome; Klippel Feil syndrome; Krabbe disease; Kugelberg-Welander disease; kuru; Lafora disease; Lambert-Eaton myasthenic syndrome; Landau-Kleffner syndrome; lateral medullary (Wallenberg) syndrome; learning disabilities; Leigh's disease; Lennox-Gustaut syndrome; Lesch-Nyhan syndrome; leukodystrophy; Lewy body dementia; Lissencephaly; locked-in syndrome; Lou Gehrig's disease (i.e., motor neuron disease or amyotrophic lateral sclerosis); lumbar disc disease; Lyme disease--neurological sequelae; Machado-Joseph disease; macrencephaly; megalencephaly; Melkersson-Rosenthal syndrome; Meniere's disease; meningitis; Menkes disease; metachromatic leukodystrophy; microcephaly; migraine; Miller Fisher syndrome; mini-strokes; mitochondrial myopathies; Mobius syndrome; monomelic amyotrophy; motor neuron disease; Moyamoya disease; mucopolysaccharidoses; multi-infarct dementia; multifocal motor neuropathy; multiple sclerosis and other demyelinating disorders; multiple system atrophy with postural hypotension; p muscular dystrophy; myasthenia gravis; myelinoclastic diffuse sclerosis; myoclonic encephalopathy of infants; myoclonus; myopathy; myotonia congenital;

narcolepsy; neurofibromatosis; neuroleptic malignant syndrome; neurological manifestations of AIDS; neurological sequelae of lupus; neuromyotonia; neuronal ceroid lipofuscinosis; neuronal migration disorders; Niemann-Pick disease; O'Sullivan-McLeod syndrome; occipital neuralgia; occult spinal dysraphism sequence; Ohtahara syndrome; olivopontocerebellar atrophy; opsoclonus myoclonus; optic neuritis; orthostatic hypotension; overuse syndrome;

5 paresthesia; Neurodegenerative disease or disorder (Parkinson's disease, Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis (ALS), dementia, multiple sclerosis and other diseases and disorders associated with neuronal cell death); paramyotonia congenital; paraneoplastic diseases; paroxysmal attacks; Parry Romberg syndrome; Pelizaeus-Merzbacher disease; periodic paralyses; peripheral neuropathy; painful neuropathy and neuropathic pain; persistent vegetative state; pervasive developmental disorders; photic sneeze reflex; phytanic acid storage disease;

10 Pick's disease; pinched nerve; pituitary tumors; polymyositis; porencephaly; post-polio syndrome; postherpetic neuralgia; postinfectious encephalomyelitis; postural hypotension; Prader- Willi syndrome; primary lateral sclerosis; prion diseases; progressive hemifacial atrophy; progressive multifocal leukoencephalopathy; progressive sclerosing poliodystrophy; progressive supranuclear palsy; pseudotumor cerebri; Ramsay-Hunt syndrome (types I and II); Rasmussen's encephalitis; reflex sympathetic dystrophy syndrome; Refsum disease; repetitive motion disorders;

15 repetitive stress injuries; restless legs syndrome; retrovirus-associated myelopathy; Rett syndrome; Reye's syndrome; Saint Vitus dance; Sandhoff disease; Schilder's disease; schizencephaly; septo-optic dysplasia; shaken baby syndrome; shingles; Shy-Drager syndrome; Sjogren's syndrome; sleep apnea; Soto's syndrome; spasticity; spina bifida; spinal cord injury; spinal cord tumors; spinal muscular atrophy; Stiff-Person syndrome; stroke; Sturge-Weber syndrome; subacute sclerosing panencephalitis; subcortical arteriosclerotic encephalopathy; Sydenham chorea; syncope; syringomyelia;

20 tardive dyskinesia; Tay-Sachs disease; temporal arteritis; tethered spinal cord syndrome; Thomsen disease; thoracic outlet syndrome; Tic Douloureux; Todd's paralysis; Tourette syndrome; transient ischemic attack; transmissible spongiform encephalopathies; transverse myelitis; traumatic brain injury; tremor; trigeminal neuralgia; tropical spastic paraparesis; tuberous sclerosis; vascular dementia (multi-infarct dementia); vasculitis including temporal arteritis; Von Hippel-Lindau disease; Wallenberg's syndrome; Werdnig-Hoffman disease; West syndrome; whiplash; Williams

25 syndrome; Wildon's disease; and Zellweger syndrome. As used herein, a "neuromuscular disease or disorder" refers to any disease adversely affecting both nervous elements (brain, spinal cord, peripheral nerve) and muscle (striated or smooth muscle), including but not limited to involuntary movement disorders, dystonias, spinal cord injury or disease, multiple sclerosis, myasthenia gravis, Parkinson's disease, Amyotrophic Lateral Sclerosis (ALS), Huntington's disease, Lambert-Eaton Myasthenic Syndrome (LES), Congenital Myasthenic Syndromes (CMS), Charcot-Marie-Tooth

30 Disease (CMT), Dejerine-Sottas Disease (DS), Creutzfeldt-Jakob disease, Friedreich's Ataxia, muscular dystrophy, spasticity from cerebral palsy and stroke.

[0064] A cardiovascular disease or disorder includes those disorders that can either cause ischemia or are caused by reperfusion of the heart. Examples include, but are not limited to, atherosclerosis, coronary artery disease,

granulomatous myocarditis, chronic myocarditis (non-granulomatous), primary hypertrophic cardiomyopathy, peripheral artery disease (PAD), stroke, angina pectoris, myocardial infarction, cardiovascular tissue damage caused by cardiac arrest, cardiovascular tissue damage caused by cardiac bypass, cardiogenic shock, and related conditions that would be known by those of ordinary skill in the art or which involve dysfunction of or tissue damage to the heart or vasculature, especially, but not limited to, tissue damage related to DMD family activation. CVS diseases include, but are not limited to, atherosclerosis, granulomatous myocarditis, myocardial infarction, myocardial fibrosis secondary to valvular heart disease, myocardial fibrosis without infarction, primary hypertrophic cardiomyopathy, and chronic myocarditis (non-granulomatous).

[0065] As used herein, "cardiomyopathy" refers to any disease or dysfunction of the myocardium (heart muscle) in which the heart is abnormally enlarged, thickened and/or stiffened. As a result, the heart muscle's ability to pump blood is usually weakened. The disease or disorder can be, for example, inflammatory, metabolic, toxic, infiltrative, fibroplastic, hematological, genetic, or unknown in origin. Such cardiomyopathies may result from a lack of oxygen. Other diseases include those that result from myocardial injury which involves damage to the muscle or the myocardium in the wall of the heart as a result of disease or trauma. Myocardial injury can be attributed to many things such as, but not limited to, cardiomyopathy, myocardial infarction, or congenital heart disease. Specific cardiac disorders to be treated also include congestive heart failure, ventricular or atrial septal defect, congenital heart defect or ventricular aneurysm. The cardiac disorder may be pediatric in origin. Cardiomyopathy includes but is not limited to, cardiomyopathy (dilated, hypertrophic, restrictive, arrhythmogenic and unclassified cardiomyopathy), sporadic dilated cardiomyopathy, X-linked Dilated Cardiomyopathy (XLDC), acute and chronic heart failure, right heart failure, left heart failure, biventricular heart failure, congenital heart defects, mitral valve stenosis, mitral valve insufficiency, aortic valve stenosis, aortic valve insufficiency, tricuspidal valve stenosis, tricuspidal valve insufficiency, pulmonic valve stenosis, pulmonic valve insufficiency, combined valve defects, myocarditis, acute myocarditis, chronic myocarditis, viral myocarditis, diastolic heart failure, systolic heart failure, diabetic heart failure and accumulation diseases.

Polynucleotide and Oligonucleotide Compositions and Molecules

[0066] *Targets:* In one embodiment, the targets comprise nucleic acid sequences of Dystrophin family, including without limitation sense and/or antisense noncoding and/or coding sequences associated with DMD family.

[0067] Dystrophin has been known since 1987, (Hoffman *et al.* (1987), *Cell*. 51:509-517) to be the protein that is deficient in Duchenne muscular dystrophy (DMD). This is an elongated protein present at the cytoplasmic surface of the vertebrate muscle cell membrane (Hoffman *et al.* (1987), *Cell*. 51:509-517). Three other dystrophin-related proteins, i.e., DRP₁ (dystrophin-related protein Type 1, or utrophin), DRP₂ (dystrophin-related protein Type 2), and dystrobrevins, have also been identified as products of different genes (Wang *et al.* (1998) *Hum Mol Genet.* 7:581-588). Dystrophins and utrophin have been detected in muscle of other mammalian species (Pons *et al.* (1994) *Circulation*. 90:369-374; Wang *et al.* (1998) *Hum Mol Genet.* 7:581-588; Rafael *et al.* (2000) *Hum Mol Genet.* 9:1357-1367) and

also in other tissues, such as the electric organ, that are derived from skeletal muscle (Chang *et al.* (1989) *J Biol Chem.* 264:20831-20834), and in nerves (Rivier *et al.* (1999a) *Histochem J.* 31:425-432, Rivier *et al.* (1999b) *J Muscle Res Cell Motil.* 20:305-314) of *T. marmorata*. Dystrophin in mammal skeletal muscle interacts with an associated protein complex (DAPC) to form a link between the cytoskeleton and the extracellular matrix (Ibraghimov-Beskrovnaya *et al.* (1992) *Nature.* 355:696-702). This complex consists of two dystroglycans (α - and β -) (Deys *et al.* (1995) *J Biol Chem.* 270:25956-25959), sarcoglycans (α -, β -, γ -, δ -, and ϵ) that are complexed with sarcospan (Nigro *et al.* (1996) *Hum Mol Genet.* 5:1179-1186; Crosbie *et al.* (1997) *J Biol Chem.* 272:31221-31224; McNally *et al.* (1998) *FEBS Lett.* 422:27-32), and three syntrophins (α -, β 1-, β 2-) in muscle tissues (Ahn *et al.* (1996) *J Biol Chem* 271:2724-2730). However, new isoforms of syntrophins (γ 1- and γ 2-) have also been reported and were found expressed as brain-specific protein (Piluso *et al.* (2000) *J Biol Chem.* 275:15851-15860). Deficiency or variations in some associated proteins generate a different muscle pathology, but the pathogenesis of all of these related muscular dystrophies is still unclear. This may be due to the heterogeneity of the data recorded, e.g., due to the existence of four dystrophin-related proteins that all share homology with dystrophin's cysteine-rich and C-terminal domains and also because of the muscle type analyzed.

[0068] Dystrophin is a rod-shaped cytoplasmic protein, and a vital part of a protein complex that connects the cytoskeleton of a muscle fiber to the surrounding extracellular matrix through the cell membrane. This complex is variously known as the costamere or the dystrophin-associated protein complex. Many muscle proteins, such as α -dystrobrevin, syncoilin, synemin, sarcoglycan, dystroglycan, and sarcospan, colocalize with dystrophin at the costamere.

[0069] Its deficiency is one of the root causes of muscular dystrophy. Normal tissue contains small amounts of dystrophin (about 0.002% of total muscle protein), but its absence leads to both DMD family and fibrosis, a condition of muscle hardening. A different mutation of the same gene causes defective dystrophin, leading to Becker's muscular dystrophy (BMD). Thus, it would be of great therapeutic value to modulate the expression and/or function of dystrophin in cells, tissues or organs of patients in need of such treatment.

[0070] Dystrophin family, the largest known human gene, encodes a protein called dystrophin. There are many different versions of dystrophin, some of which are specific to certain cell types. Dystrophin is located chiefly in muscles used for movement (skeletal muscles) and the muscles of the heart (cardiac muscles). Small amounts of the protein are present in nerve cells in the brain.

[0071] In skeletal and cardiac muscles, dystrophin is part of a protein complex that strengthens muscle fibers and protects them from injury as muscles contract and relax. The dystrophin complex acts as an anchor, connecting each muscle cell's structural framework (cytoskeleton) with the lattice of proteins and other molecules outside the cell. The dystrophin complex may also play a role in cell signaling by interacting with proteins that send and receive chemical signals.

[0072] Little is known about the function of dystrophin in nerve cells and without wishing to be bound by theory, dystrophin could be important for the normal structure and function of synapses.

[0073] Duchenne and Becker muscular dystrophy are caused by mutations in the DMD family gene.

[0074] Muscular dystrophy (MD) refers to a group of genetic disorders whose major symptom is muscle wasting.

5 There are two major forms of MD, differing in severity and age of onset. In Duchenne muscular dystrophy, symptoms are noticeable in early childhood and quickly become debilitating. Becker muscular dystrophy, on the other hand, is of later onset and less severe. Both forms of MD are caused by mutations in the dystrophin gene, a large (2.6Mb) gene comprised of 97 exons. The dystrophin protein plays an important structural role as part of a large complex in muscle fiber membranes. When dystrophin is missing or non-functional, the entire complex is compromised, leading to
10 degeneration of muscle tissue. When the ability to regenerate the muscle is exhausted, muscle wasting occurs.

[0075] Mutations in the DMD family gene also cause a form of heart disease called X-linked dilated cardiomyopathy. This condition enlarges and weakens the cardiac muscle, preventing it from pumping blood efficiently. Although dilated cardiomyopathy is a sign of Duchenne and Becker muscular dystrophy, the isolated X-linked form of this heart condition is not associated with weakness and wasting of skeletal muscles.

15 [0076] Utrophin is a 395 kDa protein encoded by multiexonic 1 Mb UTRN gene located on chromosome 6q24 (Pearce, *et al.* (1993) *Hum Mol Gene.* 2: 1765 1772). The structure of the gene bears large similarities to that of the dystrophin gene. In contrast to dystrophin, the utrophin gene has a long 5' untranslated region, split over two exons, and it is preceeded by an HTF-island. In mouse, the gene maps to chromosome 10. Utrophin is distributed throughout the sarcolemma in fetal and regenerating muscle, but is down-regulated in normal adult muscle and is restricted to the
20 myotendinous and neuromuscular junctions (Blake *et al.*, 1996). In the dystrophin deficient mdx mouse, utrophin levels in muscle remain elevated soon after birth compared with normal mice; once the utrophin levels have decreased to the adult levels (about 1 week after birth) the first signs of muscle fibre necrosis are detected. However, there is evidence to suggest that in the small calibre muscles, continual increased levels of utrophin can interact with the DGC complex (or an antigenically related complex) at the sarcolemma thus preventing loss of the complex with the result that these
25 muscles appear normal. There is also a substantial body of evidence demonstrating that utrophin is capable of localising to the sarcolemma in normal muscle. During fetal muscle development there is increased utrophin expression, localised to the sarcolemma, up until 18 weeks in the human and 20 days gestation in the mouse. After this time the utrophin sarcolemmal staining steadily decreases to the significantly lower adult levels shortly before birth where utrophin is localised almost exclusively to the NMJ. The decrease in utrophin expression coincides with increased expression of
30 dystrophin. See reviews (Ibraghimov Beskrovnaya, *et al.* (1992) *Nature* 355, 696 702, Blake, *et al.* (1994) *Trends in Cell Biol.* 4: 19 23, Tinsley, *et al.* (1993) *Curr Opin Genet Dev.* 3: 484 90).

[0077] DRP2 is predicted to resemble certain short C-terminal isoforms of dystrophin and dystrophin-related protein 1 (DRP1 or utrophin). DRP2 is a relatively small protein, encoded in man by a 45 kb gene localized to Xq22. It is expressed principally in the brain and spinal cord, and is similar in overall structure to the Dp116 dystrophin isoform.

[0078] Dystrobrevin, a member of the dystrophin family of proteins, was originally identified from the *Torpedo californica* electric organ as an 87-kDa phosphoprotein associated with the cytoplasmic face of the postsynaptic membrane (Wagner K. R. *et al.* (1993) *Neuron*, 10:511--522; Carr C. *et al.* (1989) *J Cell Biol.* 109:1753--1764). It has been postulated that the 87-kDa protein plays a role in synapse formation or stability because it copurifies with acetylcholine receptors from the electric organ membranes. In mammalian skeletal muscle, dystrophin is found in association with several integral and peripheral membrane proteins, forming a complex known as the dystrophin glycoprotein complex (DGC) (Ervasti J M. *et al.* (1991) *Cell*, 66:1121--1131; Ibraghimov-Beskrovnaya O. *et al.* (1992) *Nature* (London), ; Yoshida M. *et al.* (1990) *J Biochem (Tokyo)*, 108:748--752).

[0079] In preferred embodiments, antisense oligonucleotides are used to prevent or treat diseases or disorders associated with DMD family members. Exemplary Dystrophin family mediated diseases and disorders which can be treated with cell/tissues regenerated from stem cells obtained using the antisense compounds comprise: a muscle disease or disorder (e.g., Muscular dystrophy including Duchenne's muscular dystrophy, Becker's muscular dystrophy, Spinal bulbar muscular atrophy, dystrophinopathy, sarcoglycanopathy, limb girdle muscular dystrophy, congenital muscular dystrophy, congenital myopathy, distal myopathy, Symptomatic form of muscular dystrophy of Duchenne and Becker in female carriers myotonic syndrome etc.; a muscle-wasting disease), a neurological disease or disorder (including a neuromuscular disease or disorder e.g., dystonia, myoclonus-dystonia syndrome, etc.) a disease or disorder associated with altered level of dystrophin or dystrophin DAPC-complex, Left ventricular noncompaction, cancer, a cardiovascular disease or disorder, cardiomyopathy (e.g., sporadic dilated cardiomyopathy, X-linked Dilated Cardiomyopathy (XLDC) etc.), atherosclerosis a cytoskeletal disorder, congenital stationary night blindness and loss of hearing.

[0080] In a preferred embodiment, the oligonucleotides are specific for polynucleotides of DMD family, which includes, without limitation noncoding regions. The DMD family targets comprise variants of DMD family; mutants of DMD family, including SNPs; noncoding sequences of DMD family; alleles, fragments and the like. Preferably the oligonucleotide is an antisense RNA molecule.

[0081] In accordance with embodiments of the invention, the target nucleic acid molecule is not limited to DMD family polynucleotides alone but extends to any of the isoforms, receptors, homologs, non-coding regions and the like of DMD family.

[0082] In another preferred embodiment, an oligonucleotide targets a natural antisense sequence (natural antisense to the coding and non-coding regions) of DMD family targets, including, without limitation, variants, alleles, homologs,

mutants, derivatives, fragments and complementary sequences thereto. Preferably the oligonucleotide is an antisense RNA or DNA molecule.

[0083] In another preferred embodiment, the oligomeric compounds of the present invention also include variants in which a different base is present at one or more of the nucleotide positions in the compound. For example, if the first nucleotide is an adenine, variants may be produced which contain thymidine, guanosine, cytidine or other natural or unnatural nucleotides at this position. This may be done at any of the positions of the antisense compound. These compounds are then tested using the methods described herein to determine their ability to inhibit expression of a target nucleic acid.

[0084] In some embodiments, homology, sequence identity or complementarity, between the antisense compound and target is from about 50% to about 60%. In some embodiments, homology, sequence identity or complementarity, is from about 60% to about 70%. In some embodiments, homology, sequence identity or complementarity, is from about 70% to about 80%. In some embodiments, homology, sequence identity or complementarity, is from about 80% to about 90%. In some embodiments, homology, sequence identity or complementarity, is about 90%, about 92%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100%.

[0085] An antisense compound is specifically hybridizable when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a loss of activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences under conditions in which specific binding is desired. Such conditions include, i.e., physiological conditions in the case of in vivo assays or therapeutic treatment, and conditions in which assays are performed in the case of in vitro assays.

[0086] An antisense compound, whether DNA, RNA, chimeric, substituted etc, is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

[0087] In another preferred embodiment, targeting of DMD family including without limitation, antisense sequences which are identified and expanded, using for example, PCR, hybridization etc., one or more of the sequences set forth as SEQ ID NO: 3 to 7, and the like, modulate the expression or function of DMD family. In one embodiment, expression or function is up-regulated as compared to a control. In another preferred embodiment, expression or function is down-regulated as compared to a control.

[0088] In another preferred embodiment, oligonucleotides comprise nucleic acid sequences set forth as SEQ ID NOS: 8 to 22 including antisense sequences which are identified and expanded, using for example, PCR, hybridization etc. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds

and the like. Examples of modified bonds or internucleotide linkages comprise phosphorothioate, phosphorodithioate or the like. In another preferred embodiment, the nucleotides comprise a phosphorus derivative. The phosphorus derivative (or modified phosphate group) which may be attached to the sugar or sugar analog moiety in the modified oligonucleotides of the present invention may be a monophosphate, diphosphate, triphosphate, alkylphosphate, 5 alkanephosphate, phosphorothioate and the like. The preparation of the above-noted phosphate analogs, and their incorporation into nucleotides, modified nucleotides and oligonucleotides, *per se*, is also known and need not be described here.

[0089] The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals 10 and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

[0090] In embodiments of the present invention oligomeric antisense compounds, particularly oligonucleotides, bind to target nucleic acid molecules and modulate the expression and/or function of molecules encoded by a target gene. 15 The functions of DNA to be interfered comprise, for example, replication and transcription. The functions of RNA to be interfered comprise all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The functions may be up-regulated or inhibited depending on the functions desired.

[0091] The antisense compounds, include, antisense oligomeric compounds, antisense oligonucleotides, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other oligomeric compounds that 20 hybridize to at least a portion of the target nucleic acid. As such, these compounds may be introduced in the form of single-stranded, double-stranded, partially single-stranded, or circular oligomeric compounds.

[0092] Targeting an antisense compound to a particular nucleic acid molecule, in the context of this invention, can be a multistep process. The process usually begins with the identification of a target nucleic acid whose function is to be 25 modulated. This target nucleic acid may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target nucleic acid encodes Dystrophin family.

[0093] The targeting process usually also includes determination of at least one target region, segment, or site within 30 the target nucleic acid for the antisense interaction to occur such that the desired effect, e.g., modulation of expression, will result. Within the context of the present invention, the term "region" is defined as a portion of the target nucleic acid having at least one identifiable structure, function, or characteristic. Within regions of target nucleic acids are

segments. "Segments" are defined as smaller or sub-portions of regions within a target nucleic acid. "Sites," as used in the present invention, are defined as positions within a target nucleic acid.

[0094] In a preferred embodiment, the antisense oligonucleotides bind to the natural antisense sequences of Dystrophin family and modulate the expression and/or function of Dystrophin family (SEQ ID NO: 1 and 2). Examples of antisense sequences include SEQ ID NOS: 3 to 22.

[0095] In another preferred embodiment, the antisense oligonucleotides bind to one or more segments of Dystrophin family polynucleotides and modulate the expression and/or function of Dystrophin family. The segments comprise at least five consecutive nucleotides of the Dystrophin family sense or antisense polynucleotides.

[0096] In another preferred embodiment, the antisense oligonucleotides are specific for natural antisense sequences of Dystrophin family wherein binding of the oligonucleotides to the natural antisense sequences of Dystrophin family modulate expression and/or function of Dystrophin family.

[0097] In another preferred embodiment, oligonucleotide compounds comprise sequences set forth as SEQ ID NOS: 8 to 22, antisense sequences which are identified and expanded, using for example, PCR, hybridization etc. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds and the like. Examples of modified bonds or internucleotide linkages comprise phosphorothioate, phosphorodithioate or the like. In another preferred embodiment, the nucleotides comprise a phosphorus derivative. The phosphorus derivative (or modified phosphate group) which may be attached to the sugar or sugar analog moiety in the modified oligonucleotides of the present invention may be a monophosphate, diphosphate, triphosphate, alkylphosphate, alkanephosphate, phosphorothioate and the like. The preparation of the above-noted phosphate analogs, and their incorporation into nucleotides, modified nucleotides and oligonucleotides, per se, is also known and need not be described here.

[0098] Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes has a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG; and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). Eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA transcribed from a gene encoding Dystrophin family, regardless of the sequence(s) of such codons. A translation termination codon (or "stop codon") of a gene may

have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively).

[0099] The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon. Consequently, the "start codon region" (or "translation initiation codon region") and the "stop codon region" (or "translation termination codon region") are all regions that may be targeted effectively with the antisense compounds of the present invention.

[00100] The open reading frame (ORF) or "coding region", which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Within the context of the present invention, a targeted region is the intragenic region encompassing the translation initiation or termination codon of the open reading frame (ORF) of a gene.

[00101] Another target region includes the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA (or corresponding nucleotides on the gene). Still another target region includes the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA (or corresponding nucleotides on the gene). The 5' cap site of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap site. Another target region for this invention is the 5' cap region.

[00102] Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. In one embodiment, targeting splice sites, i.e., intron-exon junctions or exon-intron junctions, is particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular splice product is implicated in disease. An aberrant fusion junction due to rearrangement or deletion is another embodiment of a target site. mRNA transcripts produced via the process of splicing of two (or more) mRNAs from different gene sources are known as "fusion transcripts". Introns can be effectively targeted using antisense compounds targeted to, for example, DNA or pre-mRNA.

[00103] In another preferred embodiment, the antisense oligonucleotides bind to coding and/or non-coding regions of a target polynucleotide and modulate the expression and/or function of the target molecule.

[00104] In another preferred embodiment, the antisense oligonucleotides bind to natural antisense polynucleotides and modulate the expression and/or function of the target molecule.

[00105] In another preferred embodiment, the antisense oligonucleotides bind to sense polynucleotides and modulate the expression and/or function of the target molecule.

5 [00106] Alternative RNA transcripts can be produced from the same genomic region of DNA. These alternative transcripts are generally known as "variants". More specifically, "pre-mRNA variants" are transcripts produced from the same genomic DNA that differ from other transcripts produced from the same genomic DNA in either their start or stop position and contain both intronic and exonic sequence.

10 [00107] Upon excision of one or more exon or intron regions, or portions thereof during splicing, pre-mRNA variants produce smaller "mRNA variants". Consequently, mRNA variants are processed pre-mRNA variants and each unique pre-mRNA variant must always produce a unique mRNA variant as a result of splicing. These mRNA variants are also known as "alternative splice variants". If no splicing of the pre-mRNA variant occurs then the pre-mRNA variant is identical to the mRNA variant.

15 [00108] Variants can be produced through the use of alternative signals to start or stop transcription. Pre-mRNAs and mRNAs can possess more than one start codon or stop codon. Variants that originate from a pre-mRNA or mRNA that use alternative start codons are known as "alternative start variants" of that pre-mRNA or mRNA. Those transcripts that use an alternative stop codon are known as "alternative stop variants" of that pre-mRNA or mRNA. One specific type of alternative stop variant is the "polyA variant" in which the multiple transcripts produced result from the alternative selection of one of the "polyA stop signals" by the transcription machinery, thereby producing transcripts that terminate
20 at unique polyA sites. Within the context of the invention, the types of variants described herein are also embodiments of target nucleic acids.

[00109] The locations on the target nucleic acid to which the antisense compounds hybridize are defined as at least a 5-nucleotide long portion of a target region to which an active antisense compound is targeted.

25 [00110] While the specific sequences of certain exemplary target segments are set forth herein, one of skill in the art will recognize that these serve to illustrate and describe particular embodiments within the scope of the present invention. Additional target segments are readily identifiable by one having ordinary skill in the art in view of this disclosure.

[00111] Target segments 5-100 nucleotides in length comprising a stretch of at least five (5) consecutive nucleotides selected from within the illustrative preferred target segments are considered to be suitable for targeting as well.

30 [00112] Target segments can include DNA or RNA sequences that comprise at least the 5 consecutive nucleotides from the 5'-terminus of one of the illustrative preferred target segments (the remaining nucleotides being a consecutive stretch of the same DNA or RNA beginning immediately upstream of the 5'-terminus of the target segment and continuing until the DNA or RNA contains about 5 to about 100 nucleotides). Similarly preferred target segments are

represented by DNA or RNA sequences that comprise at least the 5 consecutive nucleotides from the 3'-terminus of one of the illustrative preferred target segments (the remaining nucleotides being a consecutive stretch of the same DNA or RNA beginning immediately downstream of the 3'-terminus of the target segment and continuing until the DNA or RNA contains about 5 to about 100 nucleotides). One having skill in the art armed with the target segments
5 illustrated herein will be able, without undue experimentation, to identify further preferred target segments.

[00113] Once one or more target regions, segments or sites have been identified, antisense compounds are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

[00114] In embodiments of the invention the oligonucleotides bind to an antisense strand of a particular target. The
10 oligonucleotides are at least 5 nucleotides in length and can be synthesized so each oligonucleotide targets overlapping sequences such that oligonucleotides are synthesized to cover the entire length of the target polynucleotide. The targets also include coding as well as non coding regions.

[00115] In one embodiment, it is preferred to target specific nucleic acids by antisense oligonucleotides. Targeting an antisense compound to a particular nucleic acid, is a multistep process. The process usually begins with the
15 identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a non coding polynucleotide such as for example, non coding RNA (ncRNA).

[00116] RNAs can be classified into (1) messenger RNAs (mRNAs), which are translated into proteins, and (2) non-protein-coding RNAs (ncRNAs). ncRNAs comprise microRNAs, antisense transcripts and other Transcriptional Units
20 (TU) containing a high density of stop codons and lacking any extensive "Open Reading Frame". Many ncRNAs appear to start from initiation sites in 3' untranslated regions (3'UTRs) of protein-coding loci. ncRNAs are often rare and at least half of the ncRNAs that have been sequenced by the FANTOM consortium seem not to be polyadenylated. Most researchers have for obvious reasons focused on polyadenylated mRNAs that are processed and exported to the cytoplasm. Recently, it was shown that the set of non-polyadenylated nuclear RNAs may be very large, and that many
25 such transcripts arise from so-called intergenic regions (Cheng, J. *et al.* (2005) *Science* 308 (5725), 1149-1154; Kapranov, P. *et al.* (2005). *Genome Res* 15 (7), 987-997). The mechanism by which ncRNAs may regulate gene expression is by base pairing with target transcripts. The RNAs that function by base pairing can be grouped into (1) cis encoded RNAs that are encoded at the same genetic location, but on the opposite strand to the RNAs they act upon and therefore display perfect complementarity to their target, and (2) trans-encoded RNAs that are encoded at a
30 chromosomal location distinct from the RNAs they act upon and generally do not exhibit perfect base-pairing potential with their targets.

[00117] Without wishing to be bound by theory, perturbation of an antisense polynucleotide by the antisense oligonucleotides described herein can alter the expression of the corresponding sense messenger RNAs. However, this

regulation can either be discordant (antisense knockdown results in messenger RNA elevation) or concordant (antisense knockdown results in concomitant messenger RNA reduction). In these cases, antisense oligonucleotides can be targeted to overlapping or non-overlapping parts of the antisense transcript resulting in its knockdown or sequestration. Coding as well as non-coding antisense can be targeted in an identical manner and that either category is capable of regulating the corresponding sense transcripts – either in a concordant or discordant manner. The strategies that are employed in identifying new oligonucleotides for use against a target can be based on the knockdown of antisense RNA transcripts by antisense oligonucleotides or any other means of modulating the desired target.

[00118] *Strategy 1:* In the case of discordant regulation, knocking down the antisense transcript elevates the expression of the conventional (sense) gene. Should that latter gene encode for a known or putative drug target, then knockdown of its antisense counterpart could conceivably mimic the action of a receptor agonist or an enzyme stimulant.

[00119] *Strategy 2:* In the case of concordant regulation, one could concomitantly knock down both antisense and sense transcripts and thereby achieve synergistic reduction of the conventional (sense) gene expression. If, for example, an antisense oligonucleotide is used to achieve knockdown, then this strategy can be used to apply one antisense oligonucleotide targeted to the sense transcript and another antisense oligonucleotide to the corresponding antisense transcript, or a single energetically symmetric antisense oligonucleotide that simultaneously targets overlapping sense and antisense transcripts.

[00120] According to the present invention, antisense compounds include antisense oligonucleotides, ribozymes, external guide sequence (EGS) oligonucleotides, siRNA compounds, single- or double-stranded RNA interference (RNAi) compounds such as siRNA compounds, and other oligomeric compounds which hybridize to at least a portion of the target nucleic acid and modulate its function. As such, they may be DNA, RNA, DNA-like, RNA-like, or mixtures thereof, or may be mimetics of one or more of these. These compounds may be single-stranded, doublestranded, circular or hairpin oligomeric compounds and may contain structural elements such as internal or terminal bulges, mismatches or loops. Antisense compounds are routinely prepared linearly but can be joined or otherwise prepared to be circular and/or branched. Antisense compounds can include constructs such as, for example, two strands hybridized to form a wholly or partially double-stranded compound or a single strand with sufficient self-complementarity to allow for hybridization and formation of a fully or partially double-stranded compound. The two strands can be linked internally leaving free 3' or 5' termini or can be linked to form a continuous hairpin structure or loop. The hairpin structure may contain an overhang on either the 5' or 3' terminus producing an extension of single stranded character. The double stranded compounds optionally can include overhangs on the ends. Further modifications can include conjugate groups attached to one of the termini, selected nucleotide positions, sugar positions or to one of the internucleoside linkages. Alternatively, the two strands can be linked via a non-nucleic acid moiety or linker group. When formed from only one strand, dsRNA can take the form of a self-complementary hairpin-type

molecule that doubles back on itself to form a duplex. Thus, the dsRNAs can be fully or partially double stranded. Specific modulation of gene expression can be achieved by stable expression of dsRNA hairpins in transgenic cell lines, however, in some embodiments, the gene expression or function is up regulated. When formed from two strands, or a single strand that takes the form of a self-complementary hairpin-type molecule doubled back on itself to form a duplex, the two strands (or duplex-forming regions of a single strand) are complementary RNA strands that base pair in Watson-Crick fashion.

[00121] Once introduced to a system, the compounds of the invention may elicit the action of one or more enzymes or structural proteins to effect cleavage or other modification of the target nucleic acid or may work via occupancy-based mechanisms. In general, nucleic acids (including oligonucleotides) may be described as "DNA-like" (i.e., generally having one or more 2'-deoxy sugars and, generally, T rather than U bases) or "RNA-like" (i.e., generally having one or more 2'-hydroxyl or 2'-modified sugars and, generally U rather than T bases). Nucleic acid helices can adopt more than one type of structure, most commonly the A- and B-forms. It is believed that, in general, oligonucleotides which have B-form-like structure are "DNA-like" and those which have A-formlike structure are "RNA-like." In some (chimeric) embodiments, an antisense compound may contain both A- and B-form regions.

[00122] In another preferred embodiment, the desired oligonucleotides or antisense compounds, comprise at least one of: antisense RNA, antisense DNA, chimeric antisense oligonucleotides, antisense oligonucleotides comprising modified linkages, interference RNA (RNAi), short interfering RNA (siRNA); a micro, interfering RNA (miRNA); a small, temporal RNA (stRNA); or a short, hairpin RNA (shRNA); small RNA-induced gene activation (RNAa); small activating RNAs (saRNAs), or combinations thereof.

[00123] dsRNA can also activate gene expression, a mechanism that has been termed "small RNA-induced gene activation" or RNAa. dsRNAs targeting gene promoters induce potent transcriptional activation of associated genes. RNAa was demonstrated in human cells using synthetic dsRNAs, termed "small activating RNAs" (saRNAs). It is currently not known whether RNAa is conserved in other organisms.

[00124] Small double-stranded RNA (dsRNA), such as small interfering RNA (siRNA) and microRNA (miRNA), have been found to be the trigger of an evolutionary conserved mechanism known as RNA interference (RNAi). RNAi invariably leads to gene silencing via remodeling chromatin to thereby suppress transcription, degrading complementary mRNA, or blocking protein translation. However, in instances described in detail in the examples section which follows, oligonucleotides are shown to increase the expression and/or function of the Dystrophin family polynucleotides and encoded products thereof. dsRNAs may also act as small activating RNAs (saRNA). Without wishing to be bound by theory, by targeting sequences in gene promoters, saRNAs would induce target gene expression in a phenomenon referred to as dsRNA-induced transcriptional activation (RNAa).

[00125] In a further embodiment, the "preferred target segments" identified herein may be employed in a screen for additional compounds that modulate the expression of Dystrophin family polynucleotides. "Modulators" are those

compounds that decrease or increase the expression of a nucleic acid molecule encoding Dystrophin family and which comprise at least a 5-nucleotide portion that is complementary to a preferred target segment. The screening method comprises the steps of contacting a preferred target segment of a nucleic acid molecule encoding sense or natural antisense polynucleotides of Dystrophin family with one or more candidate modulators, and selecting for one or more candidate modulators which decrease or increase the expression of a nucleic acid molecule encoding Dystrophin family polynucleotides, e.g. SEQ ID NOS: 8 to 22. Once it is shown that the candidate modulator or modulators are capable of modulating (e.g. either decreasing or increasing) the expression of a nucleic acid molecule encoding Dystrophin family polynucleotides, the modulator may then be employed in further investigative studies of the function of Dystrophin family polynucleotides, or for use as a research, diagnostic, or therapeutic agent in accordance with the present invention.

[00126] Targeting the natural antisense sequence preferably modulates the function of the target gene. For example, the DMD family gene (e.g. accession number NM_004006 and NM_007124, Fig. 2). In a preferred embodiment, the target is an antisense polynucleotide of the DMD family gene. In a preferred embodiment, an antisense oligonucleotide targets sense and/or natural antisense sequences of Dystrophin family polynucleotides (e.g. accession number NM_004006 and NM_007124, Fig. 2), variants, alleles, isoforms, homologs, mutants, derivatives, fragments and complementary sequences thereto. Preferably the oligonucleotide is an antisense molecule and the targets include coding and noncoding regions of antisense and/or sense DMD family polynucleotides.

[00127] The preferred target segments of the present invention may be also be combined with their respective complementary antisense compounds of the present invention to form stabilized double-stranded (duplexed) oligonucleotides.

[00128] Such double stranded oligonucleotide moieties have been shown in the art to modulate target expression and regulate translation as well as RNA processing via an antisense mechanism. Moreover, the double-stranded moieties may be subject to chemical modifications (Fire *et al.*, (1998) *Nature*, 391, 806-811; Timmons and Fire, (1998) *Nature*, 395, 854; Timmons *et al.*, (2001) *Gene*, 263, 103-112; Tabara *et al.*, (1998) *Science*, 282, 430-431; Montgomery *et al.*, (1998) *Proc. Natl. Acad. Sci. USA*, 95, 15502-15507; Tuschl *et al.*, (1999) *Genes Dev.*, 13, 3191-3197; Elbashir *et al.*, (2001) *Nature*, 411, 494-498; Elbashir *et al.*, (2001) *Genes Dev.*, 15, 188-200). For example, such double-stranded moieties have been shown to inhibit the target by the classical hybridization of antisense strand of the duplex to the target, thereby triggering enzymatic degradation of the target (Tijsterman *et al.*, (2002) *Science*, 295, 694-697).

[00129] In a preferred embodiment, an antisense oligonucleotide targets Dystrophin family polynucleotides (e.g. accession number NM_004006 and NM_007124), variants, alleles, isoforms, homologs, mutants, derivatives, fragments and complementary sequences thereto. Preferably the oligonucleotide is an antisense molecule.

[00130] In accordance with embodiments of the invention, the target nucleic acid molecule is not limited to Dystrophin family alone but extends to any of the isoforms, receptors, homologs and the like of Dystrophin family molecules.

[00131] In another preferred embodiment, an oligonucleotide targets a natural antisense sequence of DMD family polynucleotides, for example, polynucleotides set forth as SEQ ID NO: 3 to 7, and any variants, alleles, homologs, mutants, derivatives, fragments and complementary sequences thereto. Examples of antisense oligonucleotides are set forth as SEQ ID NOS: 8 to 22.

[00132] In one embodiment, the oligonucleotides are complementary to or bind to nucleic acid sequences of Dystrophin family antisense, including without limitation noncoding sense and/or antisense sequences associated with Dystrophin family polynucleotides and modulate expression and/or function of Dystrophin family molecules.

[00133] In another preferred embodiment, the oligonucleotides are complementary to or bind to nucleic acid sequences of DMD family natural antisense, set forth as SEQ ID NO: 3 to 7 and modulate expression and/or function of DMD family molecules.

[00134] In a preferred embodiment, oligonucleotides comprise sequences of at least 5 consecutive nucleotides of SEQ ID NOS: 8 to 22 and modulate expression and/or function of Dystrophin family molecules.

[00135] The polynucleotide targets comprise DMD family, including family members thereof, variants of DMD family; mutants of DMD family, including SNPs; noncoding sequences of DMD family; alleles of DMD family; species variants, fragments and the like. Preferably the oligonucleotide is an antisense molecule.

[00136] In another preferred embodiment, the oligonucleotide targeting Dystrophin family polynucleotides, comprise: antisense RNA, interference RNA (RNAi), short interfering RNA (siRNA); micro interfering RNA (miRNA); a small, temporal RNA (stRNA); or a short, hairpin RNA (shRNA); small RNA-induced gene activation (RNAa); or, small activating RNA (saRNA).

[00137] In another preferred embodiment, targeting of Dystrophin family polynucleotides, e.g. SEQ ID NO: 3 to 7, modulates the expression or function of these targets. In one embodiment, expression or function is up-regulated as compared to a control. In another preferred embodiment, expression or function is down-regulated as compared to a control.

[00138] In another preferred embodiment, antisense compounds comprise sequences set forth as SEQ ID NOS: 8 to 22. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds and the like.

[00139] In another preferred embodiment, SEQ ID NOS: 8 to 22 comprise one or more LNA nucleotides.

[00140] The modulation of a desired target nucleic acid can be carried out in several ways known in the art. For example, antisense oligonucleotides, siRNA etc. Enzymatic nucleic acid molecules (e.g., ribozymes) are nucleic acid molecules capable of catalyzing one or more of a variety of reactions, including the ability to repeatedly cleave other

separate nucleic acid molecules in a nucleotide base sequence-specific manner. Such enzymatic nucleic acid molecules can be used, for example, to target virtually any RNA transcript (Zaug *et al.*, 324, *Nature* 429 1986; Cech, 260 *JAMA* 3030, 1988; and Jefferies *et al.*, 17 *Nucleic Acids Research* 1371, 1989).

[00141] Because of their sequence-specificity, trans-cleaving enzymatic nucleic acid molecules show promise as therapeutic agents for human disease (Usman & McSwiggen, (1995) *Ann. Rep. Med. Chem.* 30, 285-294; Christoffersen and Marr, (1995) *J. Med. Chem.* 38, 2023-2037). Enzymatic nucleic acid molecules can be designed to cleave specific RNA targets within the background of cellular RNA. Such a cleavage event renders the mRNA non-functional and abrogates protein expression from that RNA. In this manner, synthesis of a protein associated with a disease state can be selectively inhibited.

[00142] In general, enzymatic nucleic acids with RNA cleaving activity act by first binding to a target RNA. Such binding occurs through the target binding portion of a enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through complementary base pairing, and once bound to the correct site, acts enzymatically to cut the target RNA. Strategic cleavage of such a target RNA will destroy its ability to direct synthesis of an encoded protein. After an enzymatic nucleic acid has bound and cleaved its RNA target, it is released from that RNA to search for another target and can repeatedly bind and cleave new targets.

[00143] Several approaches such as in vitro selection (evolution) strategies (Orgel, (1979) *Proc. R. Soc. London*, B 205, 435) have been used to evolve new nucleic acid catalysts capable of catalyzing a variety of reactions, such as cleavage and ligation of phosphodiester linkages and amide linkages, (Joyce, (1989) *Gene*, 82, 83-87; Beaudry *et al.*, (1992) *Science* 257, 635-641; Joyce, (1992) *Scientific American* 267, 90-97; Breaker *et al.*, (1994) *TIBTECH* 12, 268; Bartel *et al.*, (1993) *Science* 261:1411- 1418; Szostak, (1993) *TIBS* 17, 89-93; Kumar *et al.*, (1995) *FASEB J.*, 9, 1183; Breaker, (1996) *Curr. Op. Biotech.*, 7, 442).

[00144] The development of ribozymes that are optimal for catalytic activity would contribute significantly to any strategy that employs RNA-cleaving ribozymes for the purpose of regulating gene expression. The hammerhead ribozyme, for example, functions with a catalytic rate (k_{cat}) of about 1 min⁻¹ in the presence of saturating (10 mM) concentrations of Mg²⁺ cofactor. An artificial "RNA ligase" ribozyme has been shown to catalyze the corresponding self-modification reaction with a rate of about 100 min⁻¹. In addition, it is known that certain modified hammerhead ribozymes that have substrate binding arms made of DNA catalyze RNA cleavage with multiple turn-over rates that approach 100 min⁻¹. Finally, replacement of a specific residue within the catalytic core of the hammerhead with certain nucleotide analogues gives modified ribozymes that show as much as a 10-fold improvement in catalytic rate. These findings demonstrate that ribozymes can promote chemical transformations with catalytic rates that are significantly greater than those displayed in vitro by most natural self-cleaving ribozymes. It is then possible that the structures of

certain selfcleaving ribozymes may be optimized to give maximal catalytic activity, or that entirely new RNA motifs can be made that display significantly faster rates for RNA phosphodiester cleavage.

[00145] Intermolecular cleavage of an RNA substrate by an RNA catalyst that fits the "hammerhead" model was first shown in 1987 (Uhlenbeck, O. C. (1987) *Nature*, 328: 596-600). The RNA catalyst was recovered and reacted with multiple RNA molecules, demonstrating that it was truly catalytic.

[00146] Catalytic RNAs designed based on the "hammerhead" motif have been used to cleave specific target sequences by making appropriate base changes in the catalytic RNA to maintain necessary base pairing with the target sequences (Haseloff and Gerlach, (1988) *Nature*, 334, 585; Walbot and Bruening, (1988) *Nature*, 334, 196; Uhlenbeck, O. C. (1987) *Nature*, 328: 596-600; Koizumi, M., *et al.* (1988) *FEBS Lett.*, 228: 228-230). This has allowed use of the catalytic RNA to cleave specific target sequences and indicates that catalytic RNAs designed according to the "hammerhead" model may possibly cleave specific substrate RNAs in vivo. (see Haseloff and Gerlach, (1988) *Nature*, 334, 585; Walbot and Bruening, (1988) *Nature*, 334, 196; Uhlenbeck, O. C. (1987) *Nature*, 328: 596-600).

[00147] RNA interference (RNAi) has become a powerful tool for modulating gene expression in mammals and mammalian cells. This approach requires the delivery of small interfering RNA (siRNA) either as RNA itself or as DNA, using an expression plasmid or virus and the coding sequence for small hairpin RNAs that are processed to siRNAs. This system enables efficient transport of the pre-siRNAs to the cytoplasm where they are active and permit the use of regulated and tissue specific promoters for gene expression.

[00148] In a preferred embodiment, an oligonucleotide or antisense compound comprises an oligomer or polymer of ribonucleic acid (RNA) and/or deoxyribonucleic acid (DNA), or a mimetic, chimera, analog or homolog thereof. This term includes oligonucleotides composed of naturally occurring nucleotides, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally occurring portions which function similarly. Such modified or substituted oligonucleotides are often desired over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for a target nucleic acid and increased stability in the presence of nucleases.

[00149] According to the present invention, the oligonucleotides or "antisense compounds" include antisense oligonucleotides (e.g. RNA, DNA, mimetic, chimera, analog or homolog thereof), ribozymes, external guide sequence (EGS) oligonucleotides, siRNA compounds, single- or double-stranded RNA interference (RNAi) compounds such as siRNA compounds, saRNA, aRNA, and other oligomeric compounds which hybridize to at least a portion of the target nucleic acid and modulate its function. As such, they may be DNA, RNA, DNA-like, RNA-like, or mixtures thereof, or may be mimetics of one or more of these. These compounds may be single-stranded, double-stranded, circular or hairpin oligomeric compounds and may contain structural elements such as internal or terminal bulges, mismatches or loops. Antisense compounds are routinely prepared linearly but can be joined or otherwise prepared to be circular and/or branched. Antisense compounds can include constructs such as, for example, two strands hybridized to form a

wholly or partially double-stranded compound or a single strand with sufficient self-complementarity to allow for hybridization and formation of a fully or partially double-stranded compound. The two strands can be linked internally leaving free 3' or 5' termini or can be linked to form a continuous hairpin structure or loop. The hairpin structure may contain an overhang on either the 5' or 3' terminus producing an extension of single stranded character. The double stranded compounds optionally can include overhangs on the ends. Further modifications can include conjugate groups attached to one of the termini, selected nucleotide positions, sugar positions or to one of the internucleoside linkages. Alternatively, the two strands can be linked via a non-nucleic acid moiety or linker group. When formed from only one strand, dsRNA can take the form of a self-complementary hairpin-type molecule that doubles back on itself to form a duplex. Thus, the dsRNAs can be fully or partially double stranded. Specific modulation of gene expression can be achieved by stable expression of dsRNA hairpins in transgenic cell lines (Hammond *et al.*, (1991) *Nat. Rev. Genet.*, 2, 110-119; Matzke *et al.*, (2001) *Curr. Opin. Genet. Dev.*, 11, 221-227; Sharp, (2001) *Genes Dev.*, 15, 485-490). When formed from two strands, or a single strand that takes the form of a self-complementary hairpin-type molecule doubled back on itself to form a duplex, the two strands (or duplex-forming regions of a single strand) are complementary RNA strands that base pair in Watson-Crick fashion.

[00150] Once introduced to a system, the compounds of the invention may elicit the action of one or more enzymes or structural proteins to effect cleavage or other modification of the target nucleic acid or may work via occupancy-based mechanisms. In general, nucleic acids (including oligonucleotides) may be described as "DNA-like" (i.e., generally having one or more 2'-deoxy sugars and, generally, T rather than U bases) or "RNA-like" (i.e., generally having one or more 2'-hydroxyl or 2'-modified sugars and, generally U rather than T bases). Nucleic acid helices can adopt more than one type of structure, most commonly the A- and B-forms. It is believed that, in general, oligonucleotides which have B-form-like structure are "DNA-like" and those which have A-formlike structure are "RNA-like." In some (chimeric) embodiments, an antisense compound may contain both A- and B-form regions.

[00151] The antisense compounds in accordance with this invention can comprise an antisense portion from about 5 to about 80 nucleotides (i.e. from about 5 to about 80 linked nucleosides) in length. This refers to the length of the antisense strand or portion of the antisense compound. In other words, a single-stranded antisense compound of the invention comprises from 5 to about 80 nucleotides, and a double-stranded antisense compound of the invention (such as a dsRNA, for example) comprises a sense and an antisense strand or portion of 5 to about 80 nucleotides in length. One of ordinary skill in the art will appreciate that this comprehends antisense portions of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80 nucleotides in length, or any range therewithin.

[00152] In one embodiment, the antisense compounds of the invention have antisense portions of 10 to 50 nucleotides in length. One having ordinary skill in the art will appreciate that this embodies oligonucleotides having antisense

portions of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides in length, or any range therewithin. In some embodiments, the oligonucleotides are 15 nucleotides in length.

[00153] In one embodiment, the antisense or oligonucleotide compounds of the invention have antisense portions of 12 or 13 to 30 nucleotides in length. One having ordinary skill in the art will appreciate that this embodies antisense compounds having antisense portions of 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 nucleotides in length, or any range therewithin.

[00154] In another preferred embodiment, the oligomeric compounds of the present invention also include variants in which a different base is present at one or more of the nucleotide positions in the compound. For example, if the first nucleotide is an adenosine, variants may be produced which contain thymidine, guanosine or cytidine at this position. This may be done at any of the positions of the antisense or dsRNA compounds. These compounds are then tested using the methods described herein to determine their ability to inhibit expression of a target nucleic acid.

[00155] In some embodiments, homology, sequence identity or complementarity, between the antisense compound and target is from about 40% to about 60%. In some embodiments, homology, sequence identity or complementarity, is from about 60% to about 70%. In some embodiments, homology, sequence identity or complementarity, is from about 70% to about 80%. In some embodiments, homology, sequence identity or complementarity, is from about 80% to about 90%. In some embodiments, homology, sequence identity or complementarity, is about 90%, about 92%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100%.

[00156] In another preferred embodiment, the antisense oligonucleotides, such as for example, nucleic acid molecules set forth in SEQ ID NOS: 3 to 22 comprise one or more substitutions or modifications. In one embodiment, the nucleotides are substituted with locked nucleic acids (LNA).

[00157] In another preferred embodiment, the oligonucleotides target one or more regions of the nucleic acid molecules sense and/or antisense of coding and/or non-coding sequences associated with DMD family and the sequences set forth as SEQ ID NOS: 1, 2 and 3 to 7. The oligonucleotides are also targeted to overlapping regions of SEQ ID NOS: 1, 2 and 3 to 7.

[00158] Certain preferred oligonucleotides of this invention are chimeric oligonucleotides. "Chimeric oligonucleotides" or "chimeras," in the context of this invention, are oligonucleotides which contain two or more chemically distinct regions, each made up of at least one nucleotide. These oligonucleotides typically contain at least one region of modified nucleotides that confers one or more beneficial properties (such as, for example, increased nuclease resistance, increased uptake into cells, increased binding affinity for the target) and a region that is a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of antisense modulation of gene expression.

Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art. In one preferred embodiment, a chimeric oligonucleotide comprises at least one region modified to increase target binding affinity, and, usually, a region that acts as a substrate for RNase H. Affinity of an oligonucleotide for its target (in this case, a nucleic acid encoding ras) is routinely determined by measuring the T_m of an oligonucleotide/target pair, which is the temperature at which the oligonucleotide and target dissociate; dissociation is detected spectrophotometrically. The higher the T_m , the greater is the affinity of the oligonucleotide for the target.

[00159] Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotides mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures comprise, but are not limited to, US patent nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922.

[00160] In another preferred embodiment, the region of the oligonucleotide which is modified comprises at least one nucleotide modified at the 2' position of the sugar, most preferably a 2'-O-alkyl, 2'-O-alkyl-O-alkyl or 2'-fluoro-modified nucleotide. In other preferred embodiments, RNA modifications include 2'-fluoro, 2'-amino and 2' O-methyl modifications on the ribose of pyrimidines, abasic residues or an inverted base at the 3' end of the RNA. Such modifications are routinely incorporated into oligonucleotides and these oligonucleotides have been shown to have a higher T_m (i.e., higher target binding affinity) than 2'-deoxyoligonucleotides against a given target. The effect of such increased affinity is to greatly enhance RNAi oligonucleotide inhibition of gene expression. RNase H is a cellular endonuclease that cleaves the RNA strand of RNA:DNA duplexes; activation of this enzyme therefore results in cleavage of the RNA target, and thus can greatly enhance the efficiency of RNAi inhibition. Cleavage of the RNA target can be routinely demonstrated by gel electrophoresis. In another preferred embodiment, the chimeric oligonucleotide is also modified to enhance nuclease resistance. Cells contain a variety of exo- and endo-nucleases which can degrade nucleic acids. A number of nucleotide and nucleoside modifications have been shown to make the oligonucleotide into which they are incorporated more resistant to nuclease digestion than the native oligodeoxynucleotide. Nuclease resistance is routinely measured by incubating oligonucleotides with cellular extracts or isolated nuclease solutions and measuring the extent of intact oligonucleotide remaining over time, usually by gel electrophoresis. Oligonucleotides which have been modified to enhance their nuclease resistance survive intact for a longer time than unmodified oligonucleotides. A variety of oligonucleotide modifications have been demonstrated to enhance or confer nuclease resistance. Oligonucleotides which contain at least one phosphorothioate modification are presently more preferred. In some cases, oligonucleotide modifications which enhance target binding affinity are also,

independently, able to enhance nuclease resistance. Some desirable modifications can be found in De Mesmaeker *et al.* (1995) *Acc. Chem. Res.*, 28:366-374.

[00161] Specific examples of some preferred oligonucleotides envisioned for this invention include those comprising modified backbones, for example, phosphorothioates, phosphotriesters, methyl phosphonates, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages. Most preferred are oligonucleotides with phosphorothioate backbones and those with heteroatom backbones, particularly CH₂--NH--O--CH₂, CH₂--N(CH₃)--O--CH₂ [known as a methylene(methylimino) or MMI backbone], CH₂--O--N(CH₃)--CH₂, CH₂--N(CH₃)--N(CH₃)--CH₂ and O--N(CH₃)--CH₂--CH₂ backbones, wherein the native phosphodiester backbone is represented as O--P--O--CH₂. The amide backbones disclosed by De Mesmaeker *et al.* (1995) *Acc. Chem. Res.* 28:366-374 are also preferred. Also preferred are oligonucleotides having morpholino backbone structures (Summerton and Weller, U.S. Pat. No. 5,034,506). In other preferred embodiments, such as the peptide nucleic acid (PNA) backbone, the phosphodiester backbone of the oligonucleotide is replaced with a polyamide backbone, the nucleotides being bound directly or indirectly to the aza nitrogen atoms of the polyamide backbone (Nielsen *et al.* (1991) *Science* 254, 1497). Oligonucleotides may also comprise one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH, SH, SCH₃, F, OCN, OCH₃, OCH₃, O(CH₂)_nCH₃, O(CH₂)_nNH₂ or O(CH₂)_nCH₃ where n is from 1 to about 10; C₁ to C₁₀ lower alkyl, alkoxyalkoxy, substituted lower alkyl, alkaryl or aralkyl; Cl; Br; CN; CF₃; OCF₃; O--, S--, or N-alkyl; O--, S--, or N-alkenyl; SOCH₃; SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy [2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl)] (Martin *et al.*, (1995) *Helv. Chim. Acta*, 78, 486). Other preferred modifications include 2'-methoxy (2'-O--CH₃), 2'-propoxy (2'-OCH₂CH₂CH₃) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyls in place of the pentofuranosyl group.

[00162] Oligonucleotides may also include, additionally or alternatively, nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleotides include adenine (A), guanine (G), thymine (T), cytosine (C) and uracil (U). Modified nucleotides include nucleotides found only infrequently or transiently in natural nucleic acids, e.g., hypoxanthine, 6-methyladenine, 5-Me pyrimidines, particularly 5-methylcytosine (also referred to as 5-methyl-2' deoxycytosine and often referred to in the art as 5-Me-C), 5-hydroxymethylcytosine (HMC), glycosyl HMC and gentobiosyl HMC, as well as synthetic nucleotides, e.g., 2-aminoadenine, 2-(methylamino)adenine, 2-(imidazolylalkyl)adenine, 2-(aminoalkylamino)adenine or other

heterosubstituted alkyladenines, 2-thiouracil, 2-thiothymine, 5- bromouracil, 5-hydroxymethyluracil, 8-azaguanine, 7-deazaguanine, N6 (6-aminohexyl)adenine and 2,6-diaminopurine. (Kornberg, A., DNA Replication, W. H. Freeman & Co., San Francisco, 1980, pp75-77; Gebeyehu, G., (1987) *et al. Nucl. Acids Res.* 15:4513). A "universal" base known in the art, e.g., inosine, may be included. 5-Me-C substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C. (Sanghvi, Y. S., in Crooke, S. T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions.

[00163] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety, a cholesteryl moiety (Letsinger *et al.*, (1989) *Proc. Natl. Acad. Sci. USA* 86, 6553), cholic acid (Manoharan *et al.* (1994) *Bioorg. Med. Chem. Let.* 4, 1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660, 306; Manoharan *et al.* (1993) *Bioorg. Med. Chem. Let.* 3, 2765), a thiocholesterol (Oberhauser *et al.*, (1992) *Nucl. Acids Res.* 20, 533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras *et al.* *EMBO J.* 1991, 10, 111; Kabanov *et al.* (1990) *FEBS Lett.* 259, 327; Svinarchuk *et al.* (1993) *Biochimie* 75, 49), a phospholipid, e.g., dihexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero- 3-H-phosphonate (Manoharan *et al.* (1995) *Tetrahedron Lett.* 36, 3651; Shea *et al.* (1990) *Nucl. Acids Res.* 18, 3777), a polyamine or a polyethylene glycol chain (Manoharan *et al.* (1995) *Nucleosides & Nucleotides*, 14, 969), or adamantane acetic acid (Manoharan *et al.* (1995) *Tetrahedron Lett.* 36, 3651). Oligonucleotides comprising lipophilic moieties, and methods for preparing such oligonucleotides are known in the art, for example, U.S. Pat. Nos. 5,138,045, 5,218,105 and 5,459,255.

[00164] It is not necessary for all positions in a given oligonucleotide to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single oligonucleotide or even at within a single nucleoside within an oligonucleotide. The present invention also includes oligonucleotides which are chimeric oligonucleotides as hereinbefore defined.

[00165] In another embodiment, the nucleic acid molecule of the present invention is conjugated with another moiety including but not limited to abasic nucleotides, polyether, polyamine, polyamides, peptides, carbohydrates, lipid, or polyhydrocarbon compounds. Those skilled in the art will recognize that these molecules can be linked to one or more of any nucleotides comprising the nucleic acid molecule at several positions on the sugar, base or phosphate group.

[00166] The oligonucleotides used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including Applied Biosystems. Any other means for such synthesis may also be employed; the actual synthesis of the oligonucleotides is well within the talents of one of ordinary skill in the art. It is also well known to use similar techniques to prepare other oligonucleotides such as the phosphorothioates and alkylated derivatives. It is also well known to use similar techniques and commercially available modified amidites and controlled-pore glass (CPG)

products such as biotin, fluorescein, acridine or psoralen-modified amidites and/or CPG (available from Glen Research, Sterling VA) to synthesize fluorescently labeled, biotinylated or other modified oligonucleotides such as cholesterol-modified oligonucleotides.

[00167] In accordance with the invention, use of modifications such as the use of LNA monomers to enhance the potency, specificity and duration of action and broaden the routes of administration of oligonucleotides comprised of current chemistries such as MOE, ANA, FANA, PS etc (Uhlman, *et al.* (2000) *Current Opinions in Drug Discovery & Development* Vol. 3 No 2). This can be achieved by substituting some of the monomers in the current oligonucleotides by LNA monomers. The LNA modified oligonucleotide may have a size similar to the parent compound or may be larger or preferably smaller. It is preferred that such LNA-modified oligonucleotides contain less than about 70%, more preferably less than about 60%, most preferably less than about 50% LNA monomers and that their sizes are between about 5 and 25 nucleotides, more preferably between about 12 and 20 nucleotides.

[00168] Preferred modified oligonucleotide backbones comprise, but not limited to, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates comprising 3'alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

[00169] Representative United States patents that teach the preparation of the above phosphorus containing linkages comprise, but are not limited to, US patent nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5, 177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455, 233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563, 253; 5,571,799; 5,587,361; and 5,625,050.

[00170] Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These comprise those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

[00171] Representative United States patents that teach the preparation of the above oligonucleosides comprise, but are not limited to, US patent nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264, 562; 5,

264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596, 086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623, 070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439.

[00172] In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds comprise, but are not limited to, US patent nos. 5,539,082; 5,714,331; and 5,719,262.

Further teaching of PNA compounds can be found in Nielsen, *et al.* (1991) *Science* 254, 1497-1500.

[00173] In another preferred embodiment of the invention the oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular- CH₂-NH-O-CH₂-, -CH₂-N (CH₃)-O-CH₂-known as a methylene (methylimino) or MMI backbone, - CH₂-O-N (CH₃)-CH₂-, -CH₂N(CH₃)-N(CH₃) CH₂-and-O-N(CH₃)-CH₂-CH₂- wherein the native phosphodiester backbone is represented as-O-P-O-CH₂- of the above referenced US patent no. 5,489,677, and the amide backbones of the above referenced US patent no. 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced US patent no. 5,034,506.

[00174] Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C to CO alkyl or C2 to CO alkenyl and alkynyl. Particularly preferred are O (CH₂)_n OmCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON(CH₂)_nCH₃)₂ where n and m can be from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C to CO, (lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification comprises 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2- methoxyethyl) or 2'-MOE) (Martin *et al.*, (1995) *Helv. Chim. Acta*, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification comprises 2'-dimethylaminoethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples herein below, and 2'-

dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH₂-O-CH₂-N(CH₂)₂.

[00175] Other preferred modifications comprise 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-O-CH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures comprise, but are not limited to, US patent nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920.

[00176] Oligonucleotides may also comprise nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleotides comprise the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleotides comprise other synthetic and natural nucleotides such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudo-uracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine.

[00177] Further, nucleotides comprise those disclosed in United States Patent No. 3,687,808, those disclosed in 'The Concise Encyclopedia of Polymer Science And Engineering', pages 858-859, Kroschwitz, J.L., ed. John Wiley & Sons, 1990, those disclosed by Englisch *et al.*, 'Angewandte Chemie, International Edition', 1991, 30, page 613, and those disclosed by Sanghvi, Y.S., Chapter 15, 'Antisense Research and Applications', pages 289-302, Crooke, S.T. and Lebleu, B. ca., CRC Press, 1993. Certain of these nucleotides are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These comprise 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and 0-6 substituted purines, comprising 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds, 'Antisense Research and Applications', CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[00178] Representative United States patents that teach the preparation of the above noted modified nucleotides as well as other modified nucleotides comprise, but are not limited to, US patent nos. 3,687,808, as well as 4,845,205;

5,130,302; 5,134,066; 5,175, 273; 5, 367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,596,091; 5,614,617; 5,750,692, and 5,681,941.

[00179] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates, which enhance the activity, cellular distribution, or cellular uptake of the oligonucleotide.

[00180] Such moieties comprise but are not limited to, lipid moieties such as a cholesterol moiety (Letsinger *et al.*, (1989) *Proc. Natl. Acad. Sci. USA*, 86, 6553-6556), cholic acid (Manoharan *et al.*, (1994) *Bioorg. Med. Chem. Lett.*, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan *et al.*, (1992) *Ann. N. Y. Acad. Sci.*, 660, 306-309; Manoharan *et al.*, (1993) *Bioorg. Med. Chem. Lett.*, 3, 2765-2770), a thiocholesterol (Oberhauser *et al.*, (1992) *Nucl. Acids Res.*, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Kabanov *et al.*, (1990) *FEBS Lett.*, 259, 327-330; Svinarchuk *et al.*, (1993) *Biochimie* 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan *et al.*, (1995) *Tetrahedron Lett.*, 36, 3651-3654; Shea *et al.*, (1990) *Nucl. Acids Res.*, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan *et al.*, (1995) *Nucleosides & Nucleotides*, 14, 969-973), or adamantane acetic acid (Manoharan *et al.*, (1995) *Tetrahedron Lett.*, 36, 3651-3654), a palmitoyl moiety (Mishra *et al.*, (1995) *Biochim. Biophys. Acta*, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl- α -oxycholesterol moiety (Crooke *et al.*, (1996) *J. Pharmacol. Exp. Ther.*, 277, 923-937).

[00181] Representative United States patents that teach the preparation of such oligonucleotides conjugates comprise, but are not limited to, US patent nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552, 538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486, 603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762, 779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082, 830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5, 245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391, 723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5, 565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599, 928 and 5,688,941.

[00182] *Drug discovery:* The compounds of the present invention can also be applied in the areas of drug discovery and target validation. The present invention comprehends the use of the compounds and preferred target segments identified herein in drug discovery efforts to elucidate relationships that exist between Dystrophin family polynucleotides and a disease state, phenotype, or condition. These methods include detecting or modulating Dystrophin family polynucleotides comprising contacting a sample, tissue, cell, or organism with the compounds of the present invention, measuring the nucleic acid or protein level of Dystrophin family polynucleotides and/or a related phenotypic or chemical endpoint at some time after treatment, and optionally comparing the measured value to a non-

treated sample or sample treated with a further compound of the invention. These methods can also be performed in parallel or in combination with other experiments to determine the function of unknown genes for the process of target validation or to determine the validity of a particular gene product as a target for treatment or prevention of a particular disease, condition, or phenotype.

5 *Assessing Up-regulation or Inhibition of Gene Expression:*

[00183] Transfer of an exogenous nucleic acid into a host cell or organism can be assessed by directly detecting the presence of the nucleic acid in the cell or organism. Such detection can be achieved by several methods well known in the art. For example, the presence of the exogenous nucleic acid can be detected by Southern blot or by a polymerase chain reaction (PCR) technique using primers that specifically amplify nucleotide sequences associated with the nucleic acid. Expression of the exogenous nucleic acids can also be measured using conventional methods including gene expression analysis. For instance, mRNA produced from an exogenous nucleic acid can be detected and quantified using a Northern blot and reverse transcription PCR (RT-PCR).

[00184] Expression of RNA from the exogenous nucleic acid can also be detected by measuring an enzymatic activity or a reporter protein activity. For example, antisense modulatory activity can be measured indirectly as a decrease or increase in target nucleic acid expression as an indication that the exogenous nucleic acid is producing the effector RNA. Based on sequence conservation, primers can be designed and used to amplify coding regions of the target genes. Initially, the most highly expressed coding region from each gene can be used to build a model control gene, although any coding or non coding region can be used. Each control gene is assembled by inserting each coding region between a reporter coding region and its poly(A) signal. These plasmids would produce an mRNA with a reporter gene in the upstream portion of the gene and a potential RNAi target in the 3' non-coding region. The effectiveness of individual antisense oligonucleotides would be assayed by modulation of the reporter gene. Reporter genes useful in the methods of the present invention include acetohydroxyacid synthase (AHAS), alkaline phosphatase (AP), beta galactosidase (LacZ), beta glucuronidase (GUS), chloramphenicol acetyltransferase (CAT), green fluorescent protein (GFP), red fluorescent protein (RFP), yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), horseradish peroxidase (HRP), luciferase (Luc), nopaline synthase (NOS), octopine synthase (OCS), and derivatives thereof. Multiple selectable markers are available that confer resistance to ampicillin, bleomycin, chloramphenicol, gentamycin, hygromycin, kanamycin, lincomycin, methotrexate, phosphinothricin, puromycin, and tetracycline. Methods to determine modulation of a reporter gene are well known in the art, and include, but are not limited to, fluorometric methods (e.g. fluorescence spectroscopy, Fluorescence Activated Cell Sorting (FACS), fluorescence microscopy), antibiotic resistance determination.

[00185] DMD family protein and mRNA expression can be assayed using methods known to those of skill in the art and described elsewhere herein. For example, immunoassays such as the ELISA can be used to measure protein levels.

DMD family antibodies for ELISAs are available commercially, e.g., from Abnova, (Walnut, CA), Abcam, Cambridge, MA.

[00186] In embodiments, DMD family expression (e.g., mRNA or protein) in a sample (e.g., cells or tissues in vivo or in vitro) treated using an antisense oligonucleotide of the invention is evaluated by comparison with DMD family expression in a control sample. For example, expression of the protein or nucleic acid can be compared using methods known to those of skill in the art with that in a mock-treated or untreated sample. Alternatively, comparison with a sample treated with a control antisense oligonucleotide (e.g., one having an altered or different sequence) can be made depending on the information desired. In another embodiment, a difference in the expression of the DMD family protein or nucleic acid in a treated vs. an untreated sample can be compared with the difference in expression of a different nucleic acid (including any standard deemed appropriate by the researcher, e.g., a housekeeping gene) in a treated sample vs. an untreated sample.

[00187] Observed differences can be expressed as desired, e.g., in the form of a ratio or fraction, for use in a comparison with control. In embodiments, the level of DMD family mRNA or protein, in a sample treated with an antisense oligonucleotide of the present invention, is increased or decreased by about 1.25-fold to about 10-fold or more relative to an untreated sample or a sample treated with a control nucleic acid. In embodiments, the level of DMD family mRNA or protein is increased or decreased by at least about 1.25-fold, at least about 1.3-fold, at least about 1.4-fold, at least about 1.5-fold, at least about 1.6-fold, at least about 1.7-fold, at least about 1.8-fold, at least about 2-fold, at least about 2.5-fold, at least about 3-fold, at least about 3.5-fold, at least about 4-fold, at least about 4.5-fold, at least about 5-fold, at least about 5.5-fold, at least about 6-fold, at least about 6.5-fold, at least about 7-fold, at least about 7.5-fold, at least about 8-fold, at least about 8.5-fold, at least about 9-fold, at least about 9.5-fold, or at least about 10-fold or more.

Kits, Research Reagents, Diagnostics, and Therapeutics

[00188] The compounds of the present invention can be utilized for diagnostics, therapeutics, and prophylaxis, and as research reagents and components of kits. Furthermore, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes or to distinguish between functions of various members of a biological pathway.

[00189] For use in kits and diagnostics and in various biological systems, the compounds of the present invention, either alone or in combination with other compounds or therapeutics, are useful as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

[00190] As used herein the term "biological system" or "system" is defined as any organism, cell, cell culture or tissue that expresses, or is made competent to express products of the Dystrophin family genes. These include, but are not limited to, humans, transgenic animals, cells, cell cultures, tissues, xenografts, transplants and combinations thereof.

[00191] As one non limiting example, expression patterns within cells or tissues treated with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds that affect expression patterns.

[00192] Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazina and Vilo, (2000) *FEBS Lett.*, 480, 17-24; Celis, *et al.*, (2000) *FEBS Lett.*, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, *et al.*, (2000) *Drug Discov. Today*, 5, 415- 425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, (1999) *Methods Enzymol.*, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, *et al.*, (2000) *Proc. Natl. Acad. Sci. U.S.A.*, 97, 1976-81), protein arrays and proteomics (Celis, *et al.*, (2000) *FEBS Lett.*, 480, 2-16; Jungblut, *et al.*, *Electrophoresis*, 1999, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, *et al.*, *FEBS Lett.*, 2000, 480, 2-16; Larsson, *et al.*, *J. Biotechnol.*, 2000, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, *et al.*, (2000) *Anal. Biochem.* 286, 91-98; Larson, *et al.*, (2000) *Cytometry* 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, (2000) *Curr. Opin. Microbiol.* 3, 316-21), comparative genomic hybridization (Carulli, *et al.*, (1998) *J. Cell Biochem. Suppl.*, 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, (1999) *Eur. J. Cancer*, 35, 1895-904) and mass spectrometry methods (To, Comb. (2000) *Chem. High Throughput Screen*, 3, 235-41).

[00193] The compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding Dystrophin family. For example, oligonucleotides that hybridize with such efficiency and under such conditions as disclosed herein as to be effective Dystrophin family modulators are effective primers or probes under conditions favoring gene amplification or detection, respectively. These primers and probes are useful in methods requiring the specific detection of nucleic acid molecules encoding Dystrophin family and in the amplification of said nucleic acid molecules for detection or for use in further studies of Dystrophin family. Hybridization of the antisense oligonucleotides, particularly the primers and probes, of the invention with a nucleic acid encoding Dystrophin family can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabeling of the oligonucleotide, or any other suitable detection means. Kits using such detection means for detecting the level of Dystrophin family in a sample may also be prepared.

[00194] The specificity and sensitivity of antisense are also harnessed by those of skill in the art for therapeutic uses. Antisense compounds have been employed as therapeutic moieties in the treatment of disease states in animals, including humans. Antisense oligonucleotide drugs have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that antisense compounds can be useful

therapeutic modalities that can be configured to be useful in treatment regimes for the treatment of cells, tissues and animals, especially humans.

[00195] For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of Dystrophin family polynucleotides is treated by administering antisense compounds in accordance with this invention. For example, in one non-limiting embodiment, the methods comprise the step of administering to the animal in need of treatment, a therapeutically effective amount of Dystrophin family modulator. The Dystrophin family modulators of the present invention effectively modulate the activity of the Dystrophin family or modulate the expression of the Dystrophin family protein. In one embodiment, the activity or expression of Dystrophin family in an animal is inhibited by about 10% as compared to a control. Preferably, the activity or expression of Dystrophin family in an animal is inhibited by about 30%. More preferably, the activity or expression of Dystrophin family in an animal is inhibited by 50% or more. Thus, the oligomeric compounds modulate expression of Dystrophin family mRNA by at least 10%, by at least 50%, by at least 25%, by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 75%, by at least 80%, by at least 85%, by at least 90%, by at least 95%, by at least 98%, by at least 99%, or by 100% as compared to a control.

[00196] In one embodiment, the activity or expression of Dystrophin family and/or in an animal is increased by about 10% as compared to a control. Preferably, the activity or expression of Dystrophin family in an animal is increased by about 30%. More preferably, the activity or expression of Dystrophin family in an animal is increased by 50% or more. Thus, the oligomeric compounds modulate expression of Dystrophin family mRNA by at least 10%, by at least 50%, by at least 25%, by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 75%, by at least 80%, by at least 85%, by at least 90%, by at least 95%, by at least 98%, by at least 99%, or by 100% as compared to a control.

[00197] For example, the reduction of the expression of Dystrophin family may be measured in serum, blood, adipose tissue, liver or any other body fluid, tissue or organ of the animal. Preferably, the cells contained within said fluids, tissues or organs being analyzed contain a nucleic acid molecule encoding Dystrophin family peptides and/or the Dystrophin family protein itself.

[00198] The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of a compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the compounds and methods of the invention may also be useful prophylactically.

Conjugates

[00199] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates that enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. These moieties or conjugates can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter

molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugate groups include cholesterols, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve uptake, enhance resistance to degradation, and/or strengthen sequence-specific hybridization with the target nucleic acid. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve uptake, distribution, metabolism or excretion of the compounds of the present invention. Representative conjugate groups are disclosed in International Patent Application No. PCT/US92/09196, filed Oct. 23, 1992, and U.S. Pat. No. 6,287,860.

Conjugate moieties include, but are not limited to, lipid moieties such as a cholesterol moiety, cholic acid, a thioether, e.g., hexyl-5- tritylthiol, a thiocholesterol, an aliphatic chain, e.g., dodecandiol or undecyl residues, a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-Hphosphonate, a polyamine or a polyethylene glycol chain, or adamantane acetic acid, a palmityl moiety, or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety. Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic.

[00200] Representative United States patents that teach the preparation of such oligonucleotides conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941.

Formulations

[00201] The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as forexample, liposomes, receptor-targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption.

Representative United States patents that teach the preparation of such uptake, distribution and/or absorption-assisting formulations include, but are not limited to, U.S. Pat. Nos. 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,165; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170;

5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756.

[00202] Although, the antisense oligonucleotides do not need to be administered in the context of a vector in order to modulate a target expression and/or function, embodiments of the invention relates to expression vector constructs for the expression of antisense oligonucleotides, comprising promoters, hybrid promoter gene sequences and possess a strong constitutive promoter activity, or a promoter activity which can be induced in the desired case.

[00203] In an embodiment, invention practice involves administering at least one of the foregoing antisense oligonucleotides with a suitable nucleic acid delivery system. In one embodiment, that system includes a non-viral vector operably linked to the polynucleotide. Examples of such nonviral vectors include the oligonucleotide alone (e.g. any one or more of SEQ ID NOS: 8 to 22) or in combination with a suitable protein, polysaccharide or lipid formulation.

[00204] Additionally suitable nucleic acid delivery systems include viral vector, typically sequence from at least one of an adenovirus, adenovirus-associated virus (AAV), helper-dependent adenovirus, retrovirus, or hemagglutinin virus of Japan-liposome (HVJ) complex. Preferably, the viral vector comprises a strong eukaryotic promoter operably linked to the polynucleotide e.g., a cytomegalovirus (CMV) promoter.

[00205] Additionally preferred vectors include viral vectors, fusion proteins and chemical conjugates. Retroviral vectors include Moloney murine leukemia viruses and HIV-based viruses. One preferred HIV-based viral vector comprises at least two vectors wherein the gag and pol genes are from an HIV genome and the env gene is from another virus. DNA viral vectors are preferred. These vectors include pox vectors such as orthopox or avipox vectors, herpesvirus vectors such as a herpes simplex I virus (HSV) vector [Geller, A.I. *et al.*, (1995) *J. Neurochem*, 64: 487; Lim, F., *et al.*, in *DNA Cloning: Mammalian Systems*, D. Glover, Ed. (Oxford Univ. Press, Oxford England) (1995); Geller, A.I. *et al.*, (1993) *Proc Natl. Acad. Sci. U.S.A.*:90 7603; Geller, A.I., *et al.*, (1990) *Proc Natl. Acad. Sci USA*: 87:1149], Adenovirus Vectors (LeGal LaSalle *et al.*, *Science*, 259:988 (1993); Davidson, *et al.*, (1993) *Nat. Genet.* 3: 219; Yang, *et al.*, (1995) *J. Virol.* 69: 2004) and Adeno-associated Virus Vectors (Kaplit, M.G., *et al.*, (1994) *Nat. Genet.* 8:148).

[00206] The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof.

[00207] The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto. For oligonucleotides, preferred examples of pharmaceutically acceptable salts and their uses are further described in U.S. Pat. No. 6,287,860.

[00208] The present invention also includes pharmaceutical compositions and formulations that include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration.

[00209] For treating tissues in the central nervous system, administration can be made by, e.g., injection or infusion into the cerebrospinal fluid. Administration of antisense RNA into cerebrospinal fluid is described, e.g., in U.S. Pat. App. Pub. No. 2007/0117772, "Methods for slowing familial ALS disease progression."

[00210] When it is intended that the antisense oligonucleotide of the present invention be administered to cells in the central nervous system, administration can be with one or more agents capable of promoting penetration of the subject antisense oligonucleotide across the blood-brain barrier. Injection can be made, e.g., in the entorhinal cortex or hippocampus. Delivery of neurotrophic factors by administration of an adenovirus vector to motor neurons in muscle tissue is described in, e.g., U.S. Pat. No. 6,632,427, "Adenoviral-vector-mediated gene transfer into medullary motor neurons."

Delivery of vectors directly to the brain, e.g., the striatum, the thalamus, the hippocampus, or the substantia nigra, is known in the art and described, e.g., in U.S. Pat. No. 6,756,523, "Adenovirus vectors for the transfer of foreign genes into cells of the central nervous system particularly in brain."

Administration can be rapid as by injection or made over a period of time as by slow infusion or administration of slow release formulations.

[00211] The subject antisense oligonucleotides can also be linked or conjugated with agents that provide desirable pharmaceutical or pharmacodynamic properties. For example, the antisense oligonucleotide can be coupled to any substance, known in the art to promote penetration or transport across the blood-brain barrier, such as an antibody to the transferrin receptor, and administered by intravenous injection. The antisense compound can be linked with a viral vector, for example, that makes the antisense compound more effective and/or increases the transport of the antisense compound across the blood-brain barrier. Osmotic blood brain barrier disruption can also be accomplished by, e.g., infusion of sugars including, but not limited to, meso erythritol, xylitol, D(+) galactose, D(+) lactose, D(+) xylose, dulcitol, myo-inositol, L(-) fructose, D(-) mannitol, D(+) glucose, D(+) arabinose, D(-) arabinose, cellobiose, D(+) maltose, D(+) raffinose, L(+) rhamnose, D(+) melibiose, D(-) ribose, adonitol, D(+) arabitol, L(-) arabitol, D(+) fucose, L(-) fucose, D(-) lyxose, L(+) lyxose, and L(-) lyxose, or amino acids including, but not limited to, glutamine, lysine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glycine, histidine, leucine, methionine, phenylalanine,

proline, serine, threonine, tyrosine, valine, and taurine. Methods and materials for enhancing blood brain barrier penetration are described, e.g., in U. S. Patent No. 4,866,042, "Method for the delivery of genetic material across the blood brain barrier," 6,294,520, "Material for passage through the blood-brain barrier," and 6,936,589, "Parenteral delivery systems."

- 5 [00212] The subject antisense compounds may be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, for example, liposomes, receptor-targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. For example, cationic lipids may be included in the formulation to facilitate oligonucleotide uptake. One such composition shown to facilitate uptake is LIPOFECTIN (available from GIBCO-BRL, Bethesda, MD).
- 10 [00213] Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.
- 15 [00214] The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.
- 20 [00215] The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances that increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.
- 25 [00216] Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, foams and liposome-containing formulations. The pharmaceutical compositions and formulations of the present invention may comprise one or more penetration enhancers, carriers, excipients or other active or inactive ingredients.
- [00217] Emulsions are typically heterogeneous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter. Emulsions may contain additional components in addition to the dispersed
- 30 phases, and the active drug that may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Microemulsions are included as an embodiment of the present invention. Emulsions and their uses are well known in the art and are further described in U.S. Pat. No. 6,287,860.

[00218] Formulations of the present invention include liposomal formulations. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers. Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior that contains the composition to be delivered. Cationic liposomes are positively charged liposomes that are believed to interact with negatively charged DNA molecules to form a stable complex. Liposomes that are pH-sensitive or negatively-charged are believed to entrap DNA rather than complex with it. Both cationic and noncationic liposomes have been used to deliver DNA to cells.

[00219] Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids. When incorporated into liposomes, these specialized lipids result in liposomes with enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome comprises one or more glycolipids or is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. Liposomes and their uses are further described in U.S. Pat. No. 6,287,860.

[00220] The pharmaceutical formulations and compositions of the present invention may also include surfactants. The use of surfactants in drug products, formulations and in emulsions is well known in the art. Surfactants and their uses are further described in U.S. Pat. No. 6,287,860.

[00221] In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs. Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating nonsurfactants. Penetration enhancers and their uses are further described in U.S. Pat. No. 6,287,860.

[00222] One of skill in the art will recognize that formulations are routinely designed according to their intended use, i.e. route of administration.

[00223] Preferred formulations for topical administration include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoyl-phosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoyl-phosphatidyl ethanolamine DOTMA).

[00224] For topical or other administration, oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed

to lipids, in particular to cationic lipids. Preferred fatty acids and esters, pharmaceutically acceptable salts thereof, and their uses are further described in U.S. Pat. No. 6,287,860.

[00225] Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts and fatty acids and their uses are further described in U.S. Pat. No. 6,287,860.

Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-ecetyl ether. Oligonucleotides of the invention may be delivered orally, in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents and their uses are further described in U.S. Pat. No. 6,287,860.

[00226] Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions that may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

[00227] Certain embodiments of the invention provide pharmaceutical compositions containing one or more oligomeric compounds and one or more other chemotherapeutic agents that function by a non-antisense mechanism.

Examples of such chemotherapeutic agents include but are not limited to cancer chemotherapeutic drugs such as daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bischloroethyl- nitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, ansacrine, chlorambucil, methyleyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5- azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclo-phosphoramidate, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in

compositions of the invention. Combinations of antisense compounds and other non-antisense drugs are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

[00228] In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. For example, the first target may be a particular antisense sequence of Dystrophin family, and the second target may be a region from another nucleotide sequence. Alternatively, compositions of the invention may contain two or more antisense compounds targeted to different regions of the same Dystrophin family nucleic acid target. Numerous examples of antisense compounds are illustrated herein and others may be selected from among suitable compounds known in the art. Two or more combined compounds may be used together or sequentially.

Dosing:

[00229] The formulation of therapeutic compositions and their subsequent administration (dosing) is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC₅₀s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 µg to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 µg to 100 g per kg of body weight, once or more daily, to once every 20 years.

[00230] In embodiments, a patient is treated with a dosage of drug that is at least about 1, at least about 2, at least about 3, at least about 4, at least about 5, at least about 6, at least about 7, at least about 8, at least about 9, at least about 10, at least about 15, at least about 20, at least about 25, at least about 30, at least about 35, at least about 40, at least about 45, at least about 50, at least about 60, at least about 70, at least about 80, at least about 90, or at least about 100 mg/kg body weight. Certain injected dosages of antisense oligonucleotides are described, e.g., in U.S. Pat. No. 7,563,884, "Antisense modulation of PTP1B expression."

[00231] While various embodiments of the present invention have been described above, it should be understood that they have been presented by way of example only, and not limitation. Numerous changes to the disclosed embodiments can be made in accordance with the disclosure herein without departing from the spirit or scope of the invention. Thus, the breadth and scope of the present invention should not be limited by any of the above described embodiments.

By citation of various references in this document, Applicants do not admit any particular reference is "prior art" to their invention. Embodiments of inventive compositions and methods are illustrated in the following examples.

EXAMPLES

[00233] The following non-limiting Examples serve to illustrate selected embodiments of the invention. It will be appreciated that variations in proportions and alternatives in elements of the components shown will be apparent to those skilled in the art and are within the scope of embodiments of the present invention.

10 *Example 1: Design of antisense oligonucleotides specific for a nucleic acid molecule antisense to a Dystrophin family and/or a sense strand of Dystrophin family polynucleotide*

[00234] As indicated above the term "oligonucleotide specific for" or "oligonucleotide targets" refers to an oligonucleotide having a sequence (i) capable of forming a stable complex with a portion of the targeted gene, or (ii) capable of forming a stable duplex with a portion of an mRNA transcript of the targeted gene.

15 [00235] Selection of appropriate oligonucleotides is facilitated by using computer programs that automatically align nucleic acid sequences and indicate regions of identity or homology. Such programs are used to compare nucleic acid sequences obtained, for example, by searching databases such as GenBank or by sequencing PCR products. Comparison of nucleic acid sequences from a range of species allows the selection of nucleic acid sequences that display an appropriate degree of identity between species. In the case of genes that have not been sequenced, Southern
20 blots are performed to allow a determination of the degree of identity between genes in target species and other species. By performing Southern blots at varying degrees of stringency, as is well known in the art, it is possible to obtain an approximate measure of identity. These procedures allow the selection of oligonucleotides that exhibit a high degree of complementarity to target nucleic acid sequences in a subject to be controlled and a lower degree of complementarity to corresponding nucleic acid sequences in other species. One skilled in the art will realize that there is considerable
25 latitude in selecting appropriate regions of genes for use in the present invention.

[00236] An antisense compound is "specifically hybridizable" when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a modulation of function and/or activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the
30 case of in vivo assays or therapeutic treatment, and under conditions in which assays are performed in the case of in vitro assays

[00237] The hybridization properties of the oligonucleotides described herein can be determined by one or more in vitro assays as known in the art. For example, the properties of the oligonucleotides described herein can be obtained

by determination of binding strength between the target natural antisense and a potential drug molecules using melting curve assay.

[00238] The binding strength between the target natural antisense and a potential drug molecule (Molecule) can be estimated using any of the established methods of measuring the strength of intermolecular interactions, for example, a melting curve assay.

[00239] Melting curve assay determines the temperature at which a rapid transition from double-stranded to single-stranded conformation occurs for the natural antisense/Molecule complex. This temperature is widely accepted as a reliable measure of the interaction strength between the two molecules.

[00240] A melting curve assay can be performed using a cDNA copy of the actual natural antisense RNA molecule or a synthetic DNA or RNA nucleotide corresponding to the binding site of the Molecule. Multiple kits containing all necessary reagents to perform this assay are available (e.g. Applied Biosystems Inc. MeltDoctor kit). These kits include a suitable buffer solution containing one of the double strand DNA (dsDNA) binding dyes (such as ABI HRM dyes, SYBR Green, SYTO, etc.). The properties of the dsDNA dyes are such that they emit almost no fluorescence in free form, but are highly fluorescent when bound to dsDNA.

[00241] To perform the assay the cDNA or a corresponding oligonucleotide are mixed with Molecule in concentrations defined by the particular manufacturer's protocols. The mixture is heated to 95 °C to dissociate all pre-formed dsDNA complexes, then slowly cooled to room temperature or other lower temperature defined by the kit manufacturer to allow the DNA molecules to anneal. The newly formed complexes are then slowly heated to 95 °C with simultaneous continuous collection of data on the amount of fluorescence that is produced by the reaction. The fluorescence intensity is inversely proportional to the amounts of dsDNA present in the reaction. The data can be collected using a real time PCR instrument compatible with the kit (e.g. ABI's StepOne Plus Real Time PCR System or LightTyper instrument, Roche Diagnostics, Lewes, UK).

[00242] Melting peaks are constructed by plotting the negative derivative of fluorescence with respect to temperature ($-d(\text{Fluorescence})/dT$) on the y-axis against temperature (x-axis) using appropriate software (for example LightTyper (Roche) or SDS Dissociation Curve, ABI). The data is analyzed to identify the temperature of the rapid transition from dsDNA complex to single strand molecules. This temperature is called T_m and is directly proportional to the strength of interaction between the two molecules. Typically, T_m will exceed 40 °C.

Example 2: Modulation of DMD family polynucleotides

Treatment of 518A2 cells with antisense oligonucleotides

[00243] 518A2 cells obtained from Albert Einstein-Montefiore Cancer Center, NY were grown in growth media (MEM/EBSS (Hyclone cat #SH30024, or Mediatech cat # MT-10-010-CV) +10% FBS (Mediatech cat# MT35- 011-CV)+ penicillin/streptomycin (Mediatech cat# MT30-002-CI)) at 37°C and 5% CO₂. One day before the experiment the cells were replated at the density of 1.5×10^5 /ml into 6 well plates and incubated at 37°C and 5% CO₂. On the day

of the experiment the media in the 6 well plates was changed to fresh growth media. All antisense oligonucleotides were diluted to the concentration of 20 μ M. Two μ l of this solution was incubated with 400 μ l of Opti-MEM media (Gibco cat#31985-070) and 4 μ l of Lipofectamine 2000 (Invitrogen cat# 11668019) at room temperature for 20 min and applied to each well of the 6 well plates with 518A2 cells. A Similar mixture including 2 μ l of water instead of the oligonucleotide solution was used for the mock-transfected controls. After 3-18 h of incubation at 37°C and 5% CO₂ the media was changed to fresh growth media. 48 h after addition of antisense oligonucleotides the media was removed and RNA was extracted from the cells using SV Total RNA Isolation System from Promega (cat # Z3105) or RNeasy Total RNA Isolation kit from Qiagen (cat# 74181) following the manufacturers' instructions. 600 ng of RNA was added to the reverse transcription reaction performed using Verso cDNA kit from Thermo Scientific (cat#AB1453B) or High Capacity cDNA Reverse Transcription Kit (cat# 4368813 as described in the manufacturer's protocol. The cDNA from this reverse transcription reaction was used to monitor gene expression by real time PCR using ABI Taqman Gene Expression Mix (cat#4369510) and primers/probes designed by ABI (Applied Biosystems Taqman Gene Expression Assay: Hs00187805_m1 by Applied Biosystems Inc., Foster City CA). The following PCR cycle was used: 50°C for 2 min, 95°C for 10 min, 40 cycles of (95°C for 15 seconds, 60°C for 1 min) using StepOne Plus Real Time PCR Machine (Applied Biosystems).

Fold change in gene expression after treatment with antisense oligonucleotides was calculated based on the difference in 18S-normalized dCt values between treated and mock-transfected samples.

[00244] *Results:* Real time PCR results show that the levels of DMD family mRNA in 518A2 cells are significantly increased 48 h after treatment with two of the siRNAs designed to DMD family antisense BG208074. Treatment with siRNAs to other antisense molecules, BF838561, BF768753 and BF950643, did not elevate DMD family mRNA levels (Fig 1A and B).

Treatment of MCF-7 cells with antisense oligonucleotides

[00245] MCF-7 cells from ATCC (cat# HTB-22) were grown in growth media (MEM/EBSS (Hyclone cat #SH30024, or Mediatech cat # MT-10-010-CV) +10% FBS (Mediatech cat# MT35-011-CV)+ penicillin/streptomycin (Mediatech cat# MT30-002-CI)) at 37°C and 5% CO₂. One day before the experiment the cells were replated at the density of 1.5×10^5 /ml into 6 well plates and incubated at 37°C and 5% CO₂. On the day of the experiment the media in the 6 well plates was changed to fresh growth media. All antisense oligonucleotides were diluted to the concentration of 20 μ M. Two μ l of this solution was incubated with 400 μ l of Opti-MEM media (Gibco cat#31985-070) and 4 μ l of Lipofectamine 2000 (Invitrogen cat# 11668019) at room temperature for 20 min and applied to each well of the 6 well plates with MCF-7 cells. A Similar mixture including 2 μ l of water instead of the oligonucleotide solution was used for the mock-transfected controls. After 3-18 h of incubation at 37°C and 5% CO₂ the media was changed to fresh growth media. 48 h after addition of antisense oligonucleotides the media was removed and RNA was extracted from the cells using SV Total RNA Isolation System from Promega (cat # Z3105) or RNeasy Total RNA Isolation kit from Qiagen

(cat# 74181) following the manufacturers' instructions. 600 ng of RNA was added to the reverse transcription reaction performed using Verso cDNA kit from Thermo Scientific (cat#ABI453B) or High Capacity cDNA Reverse Transcription Kit (cat# 4368813) as described in the manufacturer's protocol. The cDNA from this reverse transcription reaction was used to monitor gene expression by real time PCR using ABI Taqman Gene Expression Mix (cat#4369510) and primers/probes designed by ABI (Applied Biosystems Taqman Gene Expression Assay: Hs01126016_m1 by Applied Biosystems Inc., Foster City CA). The following PCR cycle was used: 50°C for 2 min, 95°C for 10 min, 40 cycles of (95°C for 15 seconds, 60°C for 1 min) using Mx4000 thermal cycler (Stratagene) or StepOne Plus Real Time PCR Machine (Applied Biosystems).

[00246] Fold change in gene expression after treatment with antisense oligonucleotides was calculated based on the difference in 18S-normalized dCt values between treated and mock-transfected samples.

[00247] *Results:* Real time PCR results show that the levels of UTRN mRNA in MCF-7 cells are significantly increased 48 h after treatment with siRNAs designed to UTRN family antisense ENST00000431309.

[00248] Although the invention has been illustrated and described with respect to one or more implementations, equivalent alterations and modifications will occur to others skilled in the art upon the reading and understanding of this specification and the annexed drawings. In addition, while a particular feature of the invention may have been disclosed with respect to only one of several implementations, such feature may be combined with one or more other features of the other implementations as may be desired and advantageous for any given or particular application.

[00249] The Abstract of the disclosure will allow the reader to quickly ascertain the nature of the technical disclosure. It is submitted with the understanding that it will not be used to interpret or limit the scope or meaning of the following claims.

CLAIMS:

1. An antisense oligonucleotide of 10-30 nucleotides in length targeting a region of a natural antisense polynucleotide of a Dystrophin family polynucleotide, said antisense oligonucleotide upregulating the expression of the Dystrophin family polynucleotide in patient cells or tissues *in vivo* or *in vitro*, wherein the natural antisense polynucleotide comprises the nucleotide sequence as set forth in any one of SEQ ID NOs: 3-7.
2. The antisense oligonucleotide of claim 1, wherein said antisense oligonucleotide upregulates the expression of the Dystrophin family polynucleotide in patient cells or tissues *in vivo* or *in vitro* with respect to a control.
3. The antisense oligonucleotide of claim 1 or 2, wherein the antisense oligonucleotide targets a region of the natural antisense polynucleotide wherein the region is 10-30 nucleotides in length.
4. The antisense oligonucleotide of any one of claims 1-3, wherein the Dystrophin family polynucleotide comprises the nucleotide sequence depicted in SEQ ID NO: 1 or SEQ ID NO: 2.
5. The antisense oligonucleotide of any one of claims 1-4, wherein the antisense oligonucleotide comprises one or more modifications selected from: at least one modified sugar moiety, at least one modified internucleoside linkage, at least one modified nucleotide, and combinations thereof.
6. The antisense oligonucleotide of claim 5, wherein the one or more modifications comprise at least one modified sugar moiety selected from: a 2'-O-methoxyethyl modified sugar moiety, a 2'-methoxy modified sugar moiety, a 2'-O-alkyl modified sugar moiety, a bicyclic sugar moiety, and combinations thereof.
7. The antisense oligonucleotide of claim 5, wherein the one or more modifications comprise at least one modified internucleoside linkage selected from: a phosphorothioate, 2'-O-methoxyethyl (MOE), 2-fluoro, alkylphosphonate, phosphorodithioate, alkylphosphonothioate, phosphoramidate, carbamate, carbonate, phosphate triester, acetamidate, carboxymethyl ester, and combinations thereof.

8. The antisense oligonucleotide of claim 5, wherein the one or more modifications comprise at least one modification selected from: a peptide nucleic acid (PNA), a locked nucleic acid (LNA), an arabino-nucleic acid (ANA), and combinations thereof.
9. The antisense oligonucleotide of any one of claims 1-8, said antisense oligonucleotide comprising the nucleotide sequence set forth in SEQ ID NO: 18 or 19.
10. A composition for upregulating the expression of the Dystrophin family polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising at least one antisense oligonucleotide as defined in any one of claims 1-9 and a carrier.
11. A method of modulating a function of and/or the expression of a Dystrophin family polynucleotide in an isolated cell comprising:
contacting said cell with at least one antisense oligonucleotide 10-30 nucleotides in length targeting a region of a natural antisense polynucleotide of the Dystrophin family polynucleotide, said antisense oligonucleotide upregulating the expression of the Dystrophin family polynucleotide in said cell, wherein the natural antisense polynucleotide comprises the nucleotide sequence as set forth in any one of SEQ ID NOs: 3-7.
12. The method of claim 11, wherein said at least one antisense oligonucleotide upregulates the expression of the Dystrophin family polynucleotide in said cell with respect to a control.
13. The method of claim 11 or 12, wherein the at least one antisense oligonucleotide targets the natural antisense polynucleotide of the Dystrophin family polynucleotide of 10-30 nucleotides in length.
14. The method of any one of claims 11-13, wherein the at least one antisense oligonucleotide is complementary to the natural antisense sequence of the Dystrophin family polynucleotide of 10-30 nucleotides in length.
15. The method of any one of claims 11-14, wherein the Dystrophin family polynucleotide comprises the nucleotide sequence depicted in SEQ ID NO: 1 or SEQ ID NO: 2.
16. The method of any one of claims 11-15, wherein the at least one antisense oligonucleotide comprises one or more modifications selected from: at least one modified sugar moiety, at least one modified internucleoside linkage, at least one modified nucleotide, and combinations thereof.

17. The method of claim 16, wherein the one or more modifications comprise at least one modified sugar moiety selected from: a 2'-O-methoxyethyl modified sugar moiety, a 2'-methoxy modified sugar moiety, a 2'-O-alkyl modified sugar moiety, a bicyclic sugar moiety, and combinations thereof.

18. The method of claim 16, wherein the one or more modifications comprise at least one modified internucleoside linkage selected from: a phosphorothioate, 2'-O-methoxyethyl (MOE), 2-fluoro, alkylphosphonate, phosphorodithioate, alkylphosphonothioate, phosphoramidate, carbamate, carbonate, phosphate triester, acetamidate, carboxymethyl ester, and combinations thereof.

19. The method of claim 16, wherein the one or more modifications comprise at least one modification selected from: a peptide nucleic acid (PNA), a locked nucleic acid (LNA), an arabino-nucleic acid (ANA), and combinations thereof.

20. The method of any one of claims 11-19, said antisense oligonucleotide comprising the nucleotide sequence set forth in SEQ ID NO: 18 or 19.

21. A method of modulating a function of and/or the expression of a Dystrophin family polynucleotide in an isolated cell comprising:
contacting said cell with at least one antisense oligonucleotide of any one of claims 1-9, or the composition of claim 10, said at least one antisense oligonucleotide or composition upregulating the expression of the Dystrophin family polynucleotide in said cell.

22. Use of at least one antisense oligonucleotide 10-30 nucleotides in length targeting a region of a natural antisense polynucleotide of a Dystrophin family polynucleotide, for upregulating the expression of the Dystrophin family polynucleotide in a patient cell or tissue, said at least one antisense oligonucleotide upregulating the expression of the Dystrophin family polynucleotide in said cell or tissue, wherein the natural antisense polynucleotide comprises the nucleotide sequence as set forth in any one of SEQ ID NOs: 3-7.

23. Use of at least one antisense oligonucleotide 10-30 nucleotides in length targeting a region of a natural antisense polynucleotide of a Dystrophin family polynucleotide, in the manufacture of a medicament for upregulating the expression of the Dystrophin family polynucleotide in a patient cell or tissue, said at least one antisense oligonucleotide upregulating the expression of the Dystrophin family polynucleotide in said cell or tissue,

wherein the natural antisense polynucleotide comprises the nucleotide sequence as set forth in any one of SEQ ID NOs: 3-7.

24. The use of claim 22 or 23, wherein said at least one antisense oligonucleotide upregulates the expression of the Dystrophin family polynucleotide in said cell or tissue with respect to a control.

25. The use of any one of claims 22-24, wherein the at least one antisense oligonucleotide targets the natural antisense polynucleotide of the Dystrophin family polynucleotide of 10-30 nucleotides in length.

26. The use of any one of claims 22-25, wherein the at least one antisense oligonucleotide is complementary to the natural antisense polynucleotide of the Dystrophin family polynucleotide of 10-30 nucleotides in length.

27. The use of any one of claims 22-26, wherein the Dystrophin family polynucleotide comprises the nucleotide sequence depicted in SEQ ID NO: 2.

28. The use of any one of claims 22-27, wherein the at least one antisense oligonucleotide comprises one or more modifications selected from: at least one modified sugar moiety, at least one modified internucleoside linkage, at least one modified nucleotide, and combinations thereof.

29. The use of claim 28, wherein the one or more modifications comprise at least one modified sugar moiety selected from: a 2'-O-methoxyethyl modified sugar moiety, a 2'-methoxy modified sugar moiety, a 2'-O-alkyl modified sugar moiety, a bicyclic sugar moiety, and combinations thereof.

30. The use of claim 28, wherein the one or more modifications comprise at least one modified internucleoside linkage selected from: a phosphorothioate, 2'-O-methoxyethyl (MOE), 2-fluoro, alkylphosphonate, phosphorodithioate, alkylphosphonothioate, phosphoramidate, carbamate, carbonate, phosphate triester, acetamidate, carboxymethyl ester, and combinations thereof.

31. The use of claim 28, wherein the one or more modifications comprise at least one modification selected from: a peptide nucleic acid (PNA), a locked nucleic acid (LNA), an arabino-nucleic acid (ANA), and combinations thereof.

32. The use of any one of claims 22-26, wherein the Dystrophin family polynucleotide comprises the nucleotide sequence depicted in SEQ ID NO: 1.

33. The use of any one of claims 22-31, said antisense oligonucleotide comprising the nucleotide sequence set forth in SEQ ID NO: 18 or 19.

34. Use of at least one antisense oligonucleotide of any one of claims 1-9 or the composition of claim 10, for upregulating the expression of the Dystrophin family polynucleotide in a patient cell or tissue, said at least one antisense oligonucleotide or composition upregulating the expression of the Dystrophin family polynucleotide in said cell or tissue.

35. Use of at least one antisense oligonucleotide of any one of claims 1-9 or the composition of claim 10, in the manufacture of a medicament for upregulating the expression of the Dystrophin family polynucleotide in a patient cell or tissue, said at least one antisense oligonucleotide or composition upregulating the expression of the Dystrophin family polynucleotide in said cell or tissue.

FIGURE 1A





FIGURE 1B



2/8
FIGURE 1C




FIGURE 2

(SEQ ID NO: 1)

>gi|238018044|ref|NM_004006| Homo sapiens dystrophin (DMD), transcript variant Dp427m, mRNA

TCCTGGCATCAGTTACTGTGTTGACTCACTCAGTGTGGGATCACTCACTTTCCCCCTACAGGACTCAGATCTGGGAGGCAATTACCTTCGGA
GAAAAACGAATAGGAAAACTGAAGTGTACTTTTTTAAAGCTGCTGAAGTTTGTGGTTTCTCATTGTTTTTAAGCCTACTGGAGCAATAA
AGTTTGAAGAACTTTTACCAGGTTTTTTTTATCGCTGCTTGTATATACACTTTTCAAATGCTTTGGTGGGAAGAAGTAGAGGACTGTTATGA
AAGAGAAGATGTTCAAAGAAAAACATTCACAAAATGGGTAAATGCACAATTTTCTAAGTTTGGGAAGCAGCATATTGAGAACCCTCTTCAGTGA
CCTACAGGATGGGAGGCGCTCCTAGACCTCCTCGAAGGCCTGACAGGGCAAAAAGTCCAAAAAGAAAAAGGATCCACAAGAGTTTCATGCCCT
GAACAATGTCAACAAGGCACTGCGGTTTTGTCAGAACATAATGTTGATTAGTGAATATTGGAAGTACTGACATCGTAGATGGAAATCATAA
ACTGACTCTTGGTTTGATTGGAAATATAATCCTCCACTGGCAGGTCAAAAATGTAATGAAAAATATCATGGCTGGATTGCAACAAACCAACAG
TGAAAAGATTCTCTGAGCTGGTCCGACAATCAACTCGTAATTATCCACAGGTTAATGTAATCAACTTCACCACCAGCTGGTCTGATGGCCT
GGCTTTGAATGCTCTCATCCATAGTCATAGGCCAGACCTATTTGACTGGAATAGTGTGGTTTGCCAGCAGTCAGCCACACAACGACTGGAACA
TGCATTCAACATCGCCAGATATCAATTAGGCATAGAGAACTACTCGATCCTGAAGATGTTGATACCACTATCCAGATAAGAAGTCCATCTT
AATGTACATCACATCACTCTTCCAAGTTTTGCTCAACAAGTGAAGCCATCCAGGAAGTGGAAATGTTGCCAAGGCCACCTAAAGT
GACTAAAGAAGAACATTTTCAGTTACATCATCAATGCACTATTCTCAACAGATCACGGTCAGTCTAGCACAGGGATATGAGAGAACTCTTC
CCCTAAGCCTCGATTCAAGAGCTATGCCTACACACAGGCTGCTTATGTACCACCTCTGACCCTACACGGAGCCCATTTCTTCACAGCATTT
GGAAGCTCCTGAAGACAAGTCATTTGGCAGTTTCATTGATGGAGAGTGAAGTAAACCTGGACCGTTATCAAAACAGCTTTAGAAGAAGTATTATC
GTGGCTTCTTCTGCTGAGGACACATTGCAAGCACAAAGGAGAGATTTCTAATGATGTGGAAGTGGTGAAAGACCAGTTTCATACTCATGAGGG
GTACATGATGGATTGACAGCCCATCAGGGCCGGTTGGTAATATTCTACAATTGGGAAGTAAGCTGATTGGAACAGGAAAAATTATCAGAAGA
TGAAGAAATGAAGTACAAGAGCAGATGAATCTCTAAATTCAAGATGGGAATGCCTCAGGGTAGCTAGCATGGAACCAAGCAATTTTACA
TAGAGTTTAAATGGATCTCCAGAATCAGAACTGAAGAGTTGAATGACTGGCTAAACAAAAACAGAAGAAAGAACCAAGGAAATGGAGGAAGA
GCCTCTTGGACCTGATCTTGAAGACCTAAACGCCAAGTACAACAACATAAGGTGCTTCAAGAAGATCTAGAACAAGAACAACTCAGGGTCAA
TTCTCTCACTCACATGGTGGTGGTAGTTGATGAATCTAGTGGAGATCAGCAACTGCTGCTTTGGAAGAACAACTTAAGGTATTGGGAGATCG
ATGGGCAAAATCTGTAGATGGACAGAAGACCGCTGGGTTCTTTTACAAGACATCCTTCTCAAATGGCAACGCTTACTGAAGAACAGTGCCT
TTTTAGTGCATGGCTTTCAGAAAAAGAGATGTCAGTGAACAAGATTACACAACCTGGCTTTAAAGATCAAAATGAATGTTATCAAGTCTTCA
AAAAGTGGCCGTTTTTAAAGCGGATCTAGAAAAAGAAAAAGCAATCCATGGGCAAACTGTATTCACTCAAAACAAGATCTTCTTTCAACTGAA
GAATAAGTCAGTGACCCAGAAGACGGAAGCATGGCTGGATAACTTTGCCCGGTGTTGGGATAAATTAGTCCAAAACTTGAAAGAGTACAGC
ACAGATTTTACAGGCTGTCAACCACTCAGCCATCACTAACACAGACAACTGTAATGGAAACAGTAACTACGGTGACCAACAGGGAACAGAT
CCTGGTAAAGCATGCTCAAGAGGAACCTTCCACCACCCTCCCAAAAGAAAGAGGCAGATTACTGTGGATTCTGAAATTAGGAAAAGGTTGGA
TGTTGATATAACTGAACCTTACAGCTGGATTACTCGCTCAGAAGCTGTGTTGCAGAGTCTGAAATTGCAATCTTTTCGGAAGGAAGGCAACTT
CTCAGACTTAAAGAAAAAGTCAATGCCATAGAGCGAGAAAAAGCTGAGAAGTTCAGAAAACTGCAAGATGCCAGCAGATCAGCTCAGGCCCT
GGTGGACAGATGGTGAATGAGGGTGTAAATGCAGATAGCATCAAAACAGCCTCAGAACAACTGAACAGCCGGTGGATCGAATTCTGCCAGTT
GCTAAGTGAGAGACTTAATCGCTGGAGTATCAGAACACATCATCGCTTTCTATAATCAGCTACAACAATTGGAGCAGATGACAACTACTGCT
TGAAAAGTGGTTGAAAAATCCAAACCCACCCATCAGAGCCAAACAGCAATTAAGTCAAGTAAAGTAAAGTAAAGTAAAGTAAAGTAAAGTAAAGT
ATCAGGTCTTCAACCTCAAATTGAACGATTAAAAATTAAGCATAGCCCTGAAAGAGAAAGGACAAGGACCCATGTTCTCTGGATGCAGACTT
TGTGGCCTTTTCAAATCATTTTAAAGCAAGTCTTTCTGATGTGCAGGCCAGAGAGAAAGAGCTACAGACAATTTTGGACACTTGTCCACCAAT
GCGCTATCAGGAGACCATGAGTGCATCAGGACATGGGTCCAGCAGTCAGAACCAAACTCTCCATACCTCAACTTAGTGTACCCGACTATGA
AATCATGGAGCAGAGACTCGGGGAATTGCAGGCTTACAAAGTTCTCTGCAAGAGCAACAAAGTGGCCTATACTATCTCAGCACCCTGTGAA
AGAGATGTGCAAGAAAGCGCCCTCTGAAATTAGCCGGAAATATCAATCAGAATTTGAAGAAATTGAGGGACGCTGGAAGAAAGCTCTCTCCCA

GCTGGTTGAGCAATTGTCAAAGCTAGAGGAGCAAAATGAATAAACTCCGAAAAATTCAGAAATCACATACAAACCTTGAAAGAAATGGATGGCTGA
 AGTTGATGTTTTTCTGAAGGAGGAATGGCTGCCCCCTGGGGATTCAGAAATTCATAAAAGCAGCTGAAACAGTGGCAGATTTTTAGTCACTGA
 TATTCAGACAATTCAGCCAGTCTAAACAGTGTCAATGAAGGTGGGCGAGAAGATAAAGAAATGAAGCAGAGCTAGAGTTTGCTTCGAGACTTGA
 GACAGAACTCAAAGAACTTAACACTCAGTGGGATCAGATGTGCCAACAGGTCTATGCCAGAAAGGAGGCCCTTGAAAGGAGGTTTGGAGAAAAC
 TGTAAAGCTCCAGAAAGATCTATCAGAGATGACAGATGGATGACACAGGCTGAAGAAAGATATCTTGAGAGAGATTTTGAATATAAACTCC
 AGATGAATTACAGAAAGCAGTTGAAGAGATGAAGAGAGCTAAAGAAAGAGGCCCAACAAAAGAAAGCGAAAGTGAAGTCTCTACTGAGTCTGT
 AAAATGTTGTCATAGCTCAAGCTCCACCTGTAGCACAGAGGCCCTTAAAGAAAGAACTTGAAACTCTAACCCAACTACCAGTGGCTCTGCAC
 TAGGCTGAATGGGAAATGCAAGACTTTGGAAAGAGTTTGGGCATGTTGGCATGAGTTATTGTCTACTTGGAGAAAGCAAAACAGTGGCTAAA
 TGAAGTAGAATTTAACTTAAACCTGAAACATTCCTGGCGGAGCTGAGGAAATCTCTGAGGTGCTAGATTCAGTTGAAATTTGATGGC
 ACATTCAGAGGATAACCCAAATCAGATTGCGATATTGGCACAGACCTTAACAGATGGCGGAGTCTATGGATGAGCTAATCAATGAGGAATCTGA
 GACATTTAATTTCTGTTGGAGGGAATACATGAAGAGGCTGTAAAGAGGCAAAAGTTGCTTGAACAGAGCATCCAGTCTGCCAGGAGACTGA
 AAAATCTTACACTTAATCCAGAGTCCCTCACATTCATTGACAAAGCAGTTGGCAGCTTATATTGACAGACAAGGTGGACGCACTGCAATGCC
 TCAGGAAGCCAGAAAATCCAATCTGATTTGACAGTCAATGAGATCAGTTTGAAGAAATGAAGAAACATAATCAGGGGAAGAGGGCTGCCCA
 AAGAGTCTGTCTCAGATTGATGTTGCACAGAAAAATTACAAGATGTCCTCATGAAGTTTCGATTTATCCAGAAACCAAGCCATTTTGAGCA
 GCGTCTACAGAAAGTAAAGATGATTTAGATGAAGTGAAGATGCACTTGGCTGCAATTGCAACAAAGAGTGTGGAACAGGAAGTAACTACAGTC
 ACAGCTAATCATTGTGTGAATCTGTATAAAGTCTGAGTGAAGTGAAGTCTGAAGTGAAGTGGTGAAGAACTGGAGCTCAGATTGTACA
 GAAAAGCAGAGCGGAAATCCCAAGAACTTGATGAAGAGTAAACAGCTTTGAAATTTGCATTATAATGAGCTGGGAGCAAGGTAAACAGAAAG
 AAAGCAACAGTTGGGAAATGCTTGAATTTGCTCCGTAAGATGCAAGAAAGAAATGAATGCTTGCAGAAATGGCTGGCAGCTACAGATATGGA
 ATTGACAAAGAGATCAGCAGTTGAAGGAATGCCCTAGTAATTTGGATTCTGAAGTTGGCTGGGGAAGGGCTACTCAAAAAGAGATTGAGAAACA
 GAAGGTGCACTGAGAGATACACAGAGGTAGGAGAGGCCCTTGAAGAACTTTTGGGCAAGAGGAGAGCTTTGGTGGAGATAACTCAGTCT
 TCTGAATGAATCACTGATAGCTGTCACTCCCGAGCAGAAAGATGGTTAAATCTTTTGGTGGAAATACAGAAACAGATGGAATCTTTGACCA
 GATGTGGACCAACATCAAGGTGGATCTTCAAGCTGACACACTTTTGGATGAATGACAGAAAGAAACCCACAGTAAAGAAAGTGGCT
 TAAGCGTTTAAAGGCAGAACTGAATGACATACGCCCAAGGTGGACTCTACAGCTGACCAAGCAGCAAACTTGATGGCAACCGCGGTGACCA
 CTGCGAGGAATAGTAGAGGCCCAATCTCAGAGCTCAACCATCGATTTCAGAGCCATTTCACACAGAAATTAAGACTGGAAAGGCCCTCCATTCC
 TTTGAAGGAATTTGGAGCAGTTTAACTCAGATATACAAAATTTGCTTGAACCACTGGAGGCTGAAATTCAGCAGGGGGTGAATCTGAAGAGGA
 AGACTTCAATGAAGATATGAATGAAGACAAATGAGGTACTGTAAAGCAATTTGCAAGAGAGGAGACAACTTACAAACAAAGATCAGAGATGA
 GAGAAAGCAGAGGAAATAAAGATAAAGACAGCAGCTTTACAGACAAAACATAATGCTCTCAAGGATTTGAGGCTCTCAAGAAAGAAAGAGGC
 TCTAGAAATTTCTCACTAGTGGTATCAGTACAGAGGCGAGGCTGATGATCTCTGAAATGCTTGGATGACATTGAAAAAAATTAGCCAGCCT
 ACCTGAGGCCAGAGATGAAGAGGAAATAAAGGAAATGATCGGGAATTCAGAGAAAGAAAGAGAGGAGCTGAATGCACTGCGTAGGCCAAGCTGA
 GGGCTTGTCTGAGGATGGGGGCCCAATGGCAGTGGAGCCAACTCAGATCCAGCTCAGCAAGCGCTGGCGGGAAATTTGAGAGCAATTTGCTCA
 GTTTGCAAGACTCAACTTTGCAAAATTCACACTGTCCGTGAAGAAACGATGATGGTGAATGACTGAAGACATGCCCTTGGAAATTTCTTATGT
 GCTTCTCTATTGACTGAAATCACTCATGTCTCAAGGCCCTATTAGAGCTGGAAACAACTTCTCAATGCTCTGAGCTCTGTGCTGAAGGA
 CTTTGAAGATCTCTTGAAGCAAGAGGAGTCTCTGAAGAAATATAAAGATAGTCTACAAACAAAGCTCAGGTGGGATTGACATTATTCAATAGCAA
 GAAGACAGCAGCATTGCAAAAGTGCAACGCCCTGTGGAAGGGGTGAAGCTACAGGAAGCTCTCTCCAGCTTGATTTCCAATGGGAAAGATTA
 CAAAATGTACAAGGACCGACAAAGGGCGATTGACAGATCTGTTGAGAAATGGCGGCTTTTCATTATGATATAAAGATATTTAATCAGTGGCT
 AACAGAACTGAACAGTTTCTCAGAAAGACCAAAATCTCAGAAATTTGGAACTATGCTAAATCAAAATGGTATCTTAAGGAATCTCAGGATGG
 CATTGGGAGCTGGGAACTGTTGTCAGAACATTGAATGCACTGGGGAAGAAATAATTCAGCAATCTCAGAAACAGCTTCCAGATTTCTACA
 GGAAAAATTTGGCAAGCTGAATCTGCGGTGGCAGGAGTCTGCAACAGCTGTCAGACAGAAAAAGAGGCTAGAGAAACAAAGAAATATCTT
 GTCAGAAATTTCAAGAGATTTAATGAATTTGTTTTTGTGTTGGAGGAAGCAGATACATTTGCTAGTATCCCTCTGAGCTGGAAGAGAGCA
 CCAACTAAAGAAAGCTTGAGCAAGTCAAGTTACTGCTGGAAGATTTGCCCTGGCGCAGGCAATTTCTCAAACAATTAATGAACTGGAGG
 ACCCGTCTTGTAGTGTCTCCATAAGCCAGAGAGCAAGATTAACCTGAAATTAAGCTCAAGCAGACAAATCTCCAGTGGATAAAGGTTTC
 CAGAGCTTACTGCAAGAACAGAGCAAAATTAAGCTCAAAATAAAGACCTTTGGGACGCTTGAAGAAAGAGCTTCAAGACCTTGAAGAGCAGTT
 AAATCATCTGCTGCTGTGTTTATCTCTATTAGGAATCAGTTGGAAATTTATAACCAACCAAAACCAAGAGGACCAATTTGACGTTTCAAGAAAC
 TGAATAGCAGTTCAAGCTAAACAAACCGGATGTGGAAGAGATTTTGTCTAAGAGGCGAGCTTTGTACAAGCAAAACCAAGCCACTCAGCCACT
 GAAGAGGAAGTTAGAAGATCTGAGCTCTGAGTGAAGGCGGTAACCGTTTACTTCAGAGCTGAGGGGCAAGCAGGCTGACCTAGCTCTGAG
 ACTGACCACTATTGGAGGCTCTCTACTCAGACTGTACTCTGCTGACACAACTGTGTTTACTAAGGAACCTGCCATCTCCAACTAGAAAT
 GCCATCTCTCTTGAATTTGGAGGTACTGCTGTGGCAGATTTCAACCGGCTTGGACAGAACTTACCGACTGGCTTTCTCTGCTTGAATCAAGT
 TATAAATCAGAGAGGTTGATGTTGGGTGACCTTGAGGATATCAACAGATGATCATCAAGCAGAGGCAACAATGCAGGATTTGGAACAGAG
 GCTTCCCGAGTTGGAGAACTCAATACCGCTGCCCAAAATTTGAAGAACAGAGCAGCAATCAAGAGGCTAGAGCAATCAATACCGATCGAAT
 TGAAAGAAATCAGAACTCAGTGGATGAAGTACAAGAACACCTTCAGAACCGGAGGCAACAGTTGAATGAATGTTAAAGGATTAACACAATG
 GCTGGAAGCTAAGGAAGAGCTGAGCAGGCTTTAGGACAGGCCAGAGCCAGCTTGAGTCAATGGAAGGAGGCTCCCTATACAGTAGATGCAAT
 CCAAGAAATCAGAGAAACCAAGCAGTTGGCCAAAGACCTCCCGCAGCTGGCAGCAAAATGTAGATCTGGCAAAATGCTTGGCCCTGAAACT
 TCTCCGGGATTTATCTGCAAGATGATACAGAAAGTCCACATGATAACAGAGAAATCAATGCCCTTTGGAGAGGCAATTCATAAAGGGTGAG
 TGAGCGAGAGGCTGCTTTGGAAGAAACTCATAGATTACTGCAACAGCTTCCCGCTGGAGCTGGAAAGGTTTCTTGGCTGGCTTACAGAGCTGA
 AACAACTGCCAATGTCTACAGGATGCTACCCGTAAGGAAGGCTCCTAGAGACTCCAGGGAGTAAAGAGCTGATGAACAAATGGCAAGA
 CCTCCAGCTGAAATTTGAAGCTCACACAGATGTTTATCACAACTGGATGAAACAGCCCAAAATFCTGAGATCCCTGGAAGGTTCCGATGA
 TGCAGTCTGTTAGATCAAGACCTTGAAGATGCAATTTCAAGTGGAGTGAATTCGGAAAGAGTCTCTCAACATTGAGTCCCTTTGGAGGC
 CAGTTCTGACAGTGAAGCGTCTGCACCTTTCTCTGCAGGAATCTCTGTTGTGGCTACAGCTGAAAGATGATGAATTAAGCGCGGAGGCACT
 TATTGGAGCGGACTTTCCAGCAGTTTCAAGAGCAGAAAGATGTACATAGGGCTTCAAGAGGGAAATGGAAGAACTTAAAGAACTGTAAATCATGAG
 TACTCTTGAGACTGTACGAATATTTCTGACASAGCAGCTTTTGAAGGATAGAGAAATCTACCAGGAGCCAGAGAGCTGCTCTGAGGA
 GAGAGCCAGAGATGTCACTCGGCTTCTACGAAAGCAGGCTGAGGAGGCTCAATGACTGAGTGGGAAATTTGAACCTGCACTCCGCTGACTGGCA
 GAGAAATAGATGAGACCTTGAAGAGACTCCAGGAATTCAGAGAGCCAGGATGAGCTGCACTTCAAGCTGGCCAGAGCTGAGTGAATCAA
 GGGATCTTGGCAGCCGTTGGGCGATCTCTCATTTGACTCTCTCAAGATCACTTCAGAGAAAGTCAAGGCACCTTCAGAGGAGAAATTCGCGCTCT
 GAAAGAGAACGTGAGCCAGCTCAATGACCTGCTCGCCAGCTTACCACTTTGGGCAATTCAGCTCTCACCGTATAACCTCAGCACTCTGGAAGA
 CTTGAACACCAGATGGAAGCTTCTGCAAGTGGCGCTCGAGGACCGAGTCAAGCAGTGCATGAAGCCACAGGCACTTTGGTCCAGCACTCTCA

GCACCTTTCCTTTCCACGTCCTGTCGCCGGTCCCTGGGAGAGAGCCATCTCGCCAAACAAAGTGCCTTACTATATCAACCACGAGACTCAAACAAC
TTGCTTGGGACCATCCAAAATGCACAGAGCTCTACAGCTCTTAGCTGACCTTGATATATGTCAGATTCTCAGCTTATAGGACTGCCATGAAACT
CCGAGAGACTCGCAGAAAGGCCCTTTTCTTGGATCTCTTTGAGCTCTGTCAGCTGCTATGTGATGCTTTGGACAGCAGCACRACCTCAAGCAAAATGACCA
GCCCATGGATATCTTCGAGATTAHTAATTTGTTTGACCACTTTTATGACCGCTGGAGCAAGAGGACCAAAATTTGGTCACGCTCCCTCTCTG
CGTGGATATGTGCTGAACTGGCTGCTGAATGTTTATGATACGGGACGAACAGGGAGGATCCGCTGTCTCTCTTTTAAAACTGGCATCATTTT
CCTGTGTAAAGCAGATTTGGAGACAGATACAGATACCTTTTCAAGCAAGTGGCAAGTTCAACAGGATTTTGTGACAGCGCGAGGCTGGGCGT
CCTTCTGCTATTAATCTATCCAAATTCGAAGCAGTTGGGTGAAGTTGCATCCTTTGGGGGCGAGTAACTTTGAGCCAAAGTCTCCGGAGCTGCTT
CCAATTTGGCTAATAAATGAAGCAGAGATCTGAAGCGGCCCTCTCTCTAGACTGGATGAGACTGGAAACCCAGCTCATGGTGTGGCTGCCCTCTCT
GCACAGAGTGGCTGCTGCAGAAACTGCCAAGCATCAGGCCAAATGTAACTCTGCAAAAGATGTGCCAATCATTTGGATTACGATACAGGAGTCT
AAAGCACTTTAATTATGACATCTGCCAAAGCTGCTTTTTTCTGGTCGAGTTGCAAAAGGCCATAAAATGCATATCCATGCTGGGAATATT
CACTCCGACTACATCAGGAGAGATGTTCTGAGACTTTGGCAAGGTACTAAAAACAAATTTGGAACAAAAGGTATTTTGGCAAGCATCCCGG
AATGGGCTACCTGCCAGTCGAGACTGCTTTAGAGGGGGACAACTGGAAACTCCCGTTACTCTGATCAAACTCTGGCCAGTAGATTTCTGGATC
TGCTCTGGTCCCTCAGCTTTTACACAGATGATACACTACACCGATTGAACATTTAGCTAGCAGGCTAGCAGAAATGGAAACAGCAGATGGATC
TTATCTAAATGATAGCATCTCTCTTAATGAGAGCATAGATGATGAACATTTGTTAACTCCAGCATTACTGCCAAAGTTTGAACCAGGACTCCCG
CCTGAGCCAGCCTCGTAGCTCTGCCAGATCTTGATTTCTTTAGAGAGTGGAGAAAGAGGGGAGCTAGAGAGAACTCTAGCAGATCTTGAGGA
AGAAAACAGGAAATCGCAAGCAGAAATGACCGCTCTAAAGCAGCAGCAGCAACATAAAGGCTGTGCCCACTGCCGCTCCCTCTGAAATGAT
GCCCCCTCTCCCCAGAGTCCCCGGGATGCTGAGCTCATGTCTGAGGCCCAAGCTACTGCTGCAACCAAAAGCCCGCTGGAAAGCCAGGATGCA
AATCCTCGAAGACCAATAAAACAGCTGGAGTCAAGTTTACACAGGCTAAGGCAGCTGTGGAGCAACCCAGGACAGAGGCCAAAGTGAATGG
CACAACGGTGTCTCTCTCTTACCTCTCTACAGAGGTCCGACAGCAGCTCAGCCTATGCTGCTCCGAGTGGTTGGCAGTCAAACCTTCGAGCTC
CAHGGGTGAGGAAGATCTTCTCAGTCTCTCCCAAGGACACAGGACAGGCTTAGAGCAGGTTGATGGAGCAACTCAACAACCTCTTCCCTAGTTC
AAGAGGAAGAAATACCCCTGGAAAGCCAAATGAGAGAGGACCAAACTGAGGAAGTCTTTTCCACATCCGAGATGATTTGGGCAGAGCGATGGAG
TCTTATGATACAGTCATGACAGATGAGAGAGGACAGAAATTAATGTTTACAACCTCTGATCTCCCGCATGGTTTATATATGATTCATACAA
AAGAGGATAGACACTAAGAGTTTACAGAAATAAATCTATATTTTGTGAAGGGTAGTGGTATTATATCTGTATGATTTTCACTAGTTTCTAAGT
CTGTTATTTGTTTTGTAAACATGGCAGGTTTTACAGCTCTATGCAATTTGTACAAAAGTTTATAAGAAAACCTACATGTAAATCTTGATAGCT
AATAACTTGGCATTTCTTTATATGGAGCGCATTTTGGGTTGTTTAAAAATTTATAACAGTTATAAAGAAAGATTGTAACTAAAGTGTGCTT
ATAAANAAGATCTTTTATAAAGACCCCTAAAAACAAAACAAACACACACACACACACACACACACACACACACACACACACACACACAC
ATGTTTTGACATCTTTTGGGCTGATATCCATATGAAATCTAGCTAGGCTTTTTCTTTTTTGTGATATAAAGATAGAGACTCTCTTACCAACACA
CCAAATGACTACTACACACTGCTCAATTTGAGAACTGTCAGCTGAGTGGGGCAGGCTTGAGTTTTCAATTTTATATATCTAATGTCTATAGTA
TATAAATACTATAGTTATATAGATAAAGAGATAGCAATTTCTATAGACTGACTTTTTCCATTTTTTAAATGTTTATGTTCATCTCTAATAGAA
AGAAATACTTCTAGTCACTCATCGGCTTACCTGCTTGGCTCTAGAAATGGAATTTTTTCCCGGAGCCGGAAGCCAGGAGGAAACCTACACCAAC
TAAACATTTGTCTACAGCTCCAGATTTTCTCTTTTTTAAACAACTTTTCCACTGACACAGAAAGTAAAGTAAAGTTTGGATTTTTTAAAGG
AACATGTGAATGAATACACAGGACTTTATATATCAGAGTAGTAATCGGTGGTGGTGGTGAATTCATTGATTGATTGATGATTCAGCTTCTG
CTGCTAGCAATGCCACGATTTAGATTTAATGATGCTTCAGTGGAAATCAATCAGAAAGGTATTTCTGACCTTGTGAACATCAGAAGGTATTTTT
AATCCCAAGCAGTASCAGAGCAGTATAGGGCTGGAGGGCTATGGAATCCAGGCCATCCCTGTGAAGGAGTAGGCCACTCTTTAAGTGAAG
GATTTGGATGATTTGTTCATAATACATAAAGTTTCTCTGTAATTTACAACATAAAATTTATTTATGCTTCTCTCACAGTCAAAAGGAACTGGGTGGTT
GGTTTTTGTGCTTTTTAGATTTATGTGCCATGTGGGATCAGTTTTTAAATGCCACAGCATAATTTAAATAAATAAATACTTTGGGAAAA
GGTGTAAAGCAGTAGCCCCATCACATTTGTGATACTGACAGGTATCAACCCAGAAGCCCATGAAGTGTGTTTTCCATCCTTTGCATTTCTCTGC
GAGTAGTTCCACACAGGTTTTGTAAGTACCTAAGAAAGAGGCCAAATTTGATTCAAATGTTTCAAAAAAACCCCTTCTTGGTGGATTAGCAGGTT
AAATATATAAACAAACAAACAAAAATTTGCTCAAAAAAGAGGAGGAAAGCTCAAGAGGAAAGGCTAAGCACTGTGATGAGGAAAGCTTTTACTCTT
TCATGCCATTTTATTTCTTTTGTATTTTAAATCATTCATTCATAGATACACCCGCTGTGACCTATAAATTTTGCAAATCTGTGTACTCTGACA
TCAAGTGTAAATGACTTTTTGGAGAGTGGGCTGACATCAGTAGTGAATTAGCTTTTGGAGAGTGGGTTTTGTCCATTTATTAATTAATTAATTAAT
AACATCAACACAGGCTTCTCATGCTATTTCTTACTTCACTTTTGGTTTTTGGGCTGCTCTGATAAATTTGTCACACCTGAGTTCACAGCTTCAACA
CTGTGCAATTCGCTTATTTTCTTTTTCTTTTATATCTTTCTTTTTCTCTATAATTTTCAAAAGGAAACCCAAAGCTCTAAGGTAAACAAAT
TACCAAAATACATGAAGATTTGGTTTTTCTCTTCTGATTTTTTCTTTATGTGACGCTGACCTTTTTCTTTACCCAGGATTTTAAACACTCA
GATTTAAACAAAGGGGTTACTTTTACATCTTACATGAAGATTTTAAAGTAAGTAAGTTTCAATCTAAAACTCAGAGGTAAATAGAGTGCATTAATTA
TTTTGTTTTAAATCTTTTTGTTTTTCTTTTACACACATTAGCTCTGGAGTGAGTCTGTCTATAATTTTGAACAAAAAATTCAGAGCTTTATTTGCT
GCATTTTAAACATAATTAATTTGGACATTTATTTCTGTGTTGCTTCTTTATAACCAACCGAGTATTAAGCTGTAAATCATATGTAACTGAAGCA
TAAACATCACATGGCATTTTTTTGTCATTTGTTTTTTCAGGATCTGAGCTTCTACTTGAAGTATCATAAATATATTTGTGTTTTTAAACCAACACTGTAA
CATTTACGAATTTATTTTTTAAACTTCAGTTTTTACTGCAATTTTCCACACATATCAGACTTCACCAAAATATATGCTTACTATTTGTATTAATG
ACTGCTTTTACTGTGATCTCTCAATAAAGCAGCGCATTTATGTTAC

(SEO ID NO: 2)

>gi|110611227|ref|NM_007124| Homo sapiens utrophin (UTRN), mRNA

GTATTGATGTC AAGCTGA ACCCATCGTAGGAAGTTGAAAGCCTTAGAAGAGGAGCACTTGGTAAAGTTTTGGATTATCTTGAAGACTCTGGCAAGA
TGGCCAAGTATGGAGAACATGAAGCCAGTCTCTGACAATGGGCAGAACGAATTCAGTGATATCATTAAGTCCAGATCTGATGAACACAATGAAG
TACAGAAGAAAACCTTTACCAAAATGGATAAATGCTCCGATTTCAAAGAGTGGGAAACCACCCATCAATGATATCTTTCACAGACCTCAAGAGATG
CAAGGAAGCTATTGGATCTCTTAGAAGCCCTCAGAGGAACATCACTGCCAAAGGAACGTCGTTCCACAAGGGTACATGCCCTTAAATAAAGCTCA
GAGAGTGCTCGAGGTTTTCATACAGAAACAATGTGGAAATTAGTGAATATAGGGGGAACTGACATTTGTGGATGGAAATCAACAACTGBCCTTTGG
GGTTACTTTGGAGCATCACTTTTCCACTGGCAGGTGAAAGATGTCTATGAAGGATGTCTATGTGCGAAGCTGCAGCAGACGGAACAGTGAAGAGATCC
TGCTCAGCTGGGTCGCTCAGACCACCAAGGCCCTACAGCCCAAGTCAAGGCTCTCAACTTCACCAACAGCTGGACAGATGGAGCTCGCCTTTAATG
CTGTCTCTCCACCGACATAAACCCTGATCTCTTCAGCTGGGATAAAGATGTGTCAAAATGTGTCAACAAATTAGAGAGACTTGAACATTCCTCTCAGCAAGG
CTCAAACTTATTGGGAATTGAAAGCTGTTAGATCTGAAGATGTTGCCGTTTCAGCTCTGACACAGAAATCCATAATTTATGTTTAACT
CTTTGTTTGAGGCTCTACCTCAGCAAGTCAACCAATAGACGCCCTCCGGTGAGGTAGAGACACCHCCCAAGGAAATATAAAAAGGAATGTGAAGAG

[illegible]

ATAAAGCTTCCAGTTTCCAGATATGAGAACAGCAATTACAGAAAAAATTGGAAAGGGTCAAGAACCCAGTGGGATGGCACCAGCATGGCGTTGAGC
 TAAGACAGCAGCAGCTTGGAGGATGATTATTGACAGTCTTCAGTGGGATGACCATAGGGAGGAGACTGAAGAACTGATGAGAAAAATATGAGG
 CTCGACTCTATATTCTTCAGCAASCCCGACGGGATCCACTCACCACAACTAATCTGTATACCAAACTACTGTTTCAAGAACTEGGTCCTGGAG
 ATGGTATCTTCATGGCGTTCCGATACCTCTGCAGAACTCTCGAGGAAATATGGGAGTGATGACACAAGGAATGTGAAGAAAAACACAGAGT
 ACTTAAAAACATCATGGATCAATCTCACAACAAAGTATTCCTGACAGACAGAACGCCCTGGAGGGCTGAGTGGAGGACGGTGCAGGGCTCTCGCA
 GAGATCTGGAAAACTTCTGAACTGGATCCAGAAGCAGAGACCAAGTGAATGTGCTTGTGGATGGCTCTCACTGGGAGAAATGTCTCTCAGG
 ATAGTATCTTGGCCAGGGAACTCAAAACAGCAGATGCAGGACATCCAGGCACAAATTTGATGCCCCACAATGACATATTTAAAGCACTTGCACGGAA
 ACAGGCAGAAATGTTAAAAAGCTTTGGGAAATTTCTGAAGAGGCTACTATGCTTCAACATCGACTGGATGATATGAACCAAGATGGAATGACT
 TAAAAGCAAAATCTGCTAGCATCAGGGGCCATTTGGAGGCCAGGCTGAGAACTGGAACAGGTTGCTGATGTCTTTAGAGAACTGATCAAAAT
 GGCTGAATATGAAGATGAAGAGCTTAAGAAACAAATGCCATTATGGAGGAGATGTTCCAGCCTTACAGCTCCAGTATGACCAITGTAAAGGCC
 TGAGACGGGAGTTAAAGGAGAAAGAAATATCTGCTCGAATGTGTCSACAGGGCCCGAGTTTTCTTGGCTGATCAGGCCAATTGAGGGCCCTG
 AAGAGCCAAAGAAAGACCTACATTCAAAAACAGAAATTAACCTCTGAGGAGAGAGGCCAAAGATTCGCAAGGCCATGGCCAAACAGTCTTCTG
 AAGTCAAAAGAAAAATGGGAAAGTCTAAATGCTGTAACTAGCAATTTGCCAAAAGCAAGTGGACAAGGCATTGGAGAAACTCAGAGACCTGCAGG
 GAGCTATGGATGACCTGGAGCGTGCATGCAAGGAGGCAGAGTCCGTGGCGAATGGCTGGAAGCCCTGGGAGAGCTTACTCAITGACTTCGCTGC
 AGGATCAGATTGAAGAAATCATGGCAATTTACAGAGAAATTTGCCCAATCAACTTTAAAGTTAAAGCGGTGAATGATTATCCAGTCAGCTGT
 CTCACCTTGACCTGCATCCCTCTCTAAAGATGTCTCGCCAGCTAGATGACCTTAATATGCGATGGAACTTTTACAGGTTTCTGTGGATGATC
 GGCTTAAACAGCTTCAGGAAGGCCACAGAGATTTTGGACCATCTCTCAGCAATTTCTCTCTACGTCAGTCCAGCTGGCGTGGCAAGATCCA
 TTTCCATATAATAAGTGGCTTATTACATCAACATCAACACAGACCCCTGTTGGGACCATCTTAAATGACCGAAGCTTTTCAATCCCTTG
 CTGACCTGAATAATGTACGTTTTCTGCTACCGTACAGCAATCAAACTCCGAAGACTACAAAAGCACTATGTTTGGATCTCTTAGAGTTGA
 GTACAACAATGAATTTTCAACAGCAGCAAGTTGAACCAAAATGACCAGCTCTCTCAGTGTTCCAGATGTCTCACTCAACTGTCTGACAACAATT
 ATGATGGACTTGAGCAAAATGCTAAGSACCTGGTCAAGCTTCCACTCTGTGTGATATGTGTCTCAATTTGGTGGCTCAATGTCTATGACACGG
 GTCGAAGCTGGAAAAATTAGAGCTGCAGAGCTCTGAAGATTTGGATGATGCTCTCTCCCAAGGCTCTCTTGGAAAGAAATCTCTTTT
 AGGAAGTTGCAGGGCCACAGAAATGTGTGACAGAGGCAGCTGGGCTGTTACTTTCATGATGCCATCCAGATCCCGGGCAGCTAGGTGAAG
 TAGCAGCTTTTGGAGCAGTAAATATTGAGCTAGTGTTCGACAGTGTCTTCCACACGAATAACAATAAACCGAATAAGTGTGAAGAGTTTA
 TAGATTGGATGCATTTGAACCCACAGTCCATGGTTTTGGCTCCAGTTTTACATCGAGTGGCAGCAGCGGAGACTCGAAAACATCAGGCCAAT
 GCAACATCTGTAAAGAAATTTCCAAATTTGTGGCTTCAGGTATAGAAGCTTTAAGCAATTTAACTATGATGTCTGCCAGATTTGTTCTTTCCG
 TCTGAACAGCAAAAGTTCACAAATTAATTTACCCAAATGGTGGAAATTTGATACCTCAACACATCTGGGGAAGATGTACAGAGACTTCACAAAGG
 TACTTAAGAACAGTTTCAGGTGGAAGAAGTACTTTGCCAAACACCCCTCGACTTGGTTACCTGCTTCCAGACAGTTCTTGAAGGTGACAACT
 TAGAGACTCTTATCAGCTCATCAGTATGTGGCCAGAGCACTATGACCCCTCACAATCTCTCACTGTTTCTATGATGACACCCATTCAAGAA
 TAGAACAAATATGCCACAGCACTGGGCCAGATGGAAAGGACTAATGGGTCTTTTCTCACTGATAGCAGCTCCACCACAGGAAGTGTGGAAGAGC
 AGCAGCCCTCATCTAGCAGTATTTGCCAAACACTCGAGAGGAGTCCCGAGTGAGCCAGCCGCGAGAGCCAGCTCAGATCTGAAGTCAGTAG
 AGAGGGAAGAACCTGAGAACTGGAGAGGATCATCTGACCTCGAGGAGCAAAAGAAATCTACAGGTGGAGTATGAGCAGCTGGAAGCACC
 AGCAGCTTCCGAAGGGGGCTCCCTGTGGTTTACCAGCAGAGTGCATTAATCTCTCCCATCACACGCTCTGAGGATTCAGAACTTTATAGCAGAAG
 CAAAACCTCTCAGGCAGCACAAGAGTTCGGCTGGAGGCTAGGATGCAGATTTTGAAGATCACAATAAACAGCTGGAGTCTCAGCTCCACCGCC
 TCCGACAGCTGCTGGAGCAGCTGAATCTGATTTCCGGAATCAATGGTGTTTTCCCATGCGCTTCTCTCAGCAATCTGCACTGAGCTACTCGC
 TTGATCCAGATGGCTCCGCCCCACAGTTTCCACAGGCGAGGGGAGGAGCTCTGCCCCCACCAGCAGCAGGATCTCAAGGAGG
 TCATGAGCAGATTTACAGCAGCTTTCCATCTTGTCTGCCCAAAATGTTTCCGAGGCGACAGGCAATGTGAAGTATTCATCTCCGCCCCAACAA
 GTTTCTGACGTACAGTGTGTCCTTTTTCAGCAATGCCAATTTCAAGTTCCATTAAATCAGAAGCTCCATGGCTCTCTGGCCCCAGGATGTTG
 AGTGTGACTGTGTGTCTACGGAAGAGTAAACACTGACTATCCAAAGAGAAATGGATATTTGTTTTTATAATAACCATATATTATTGTT
 TTCTTCTCTCTCTCTATGCAAGTGTAAATTAATGAACAGAGAGCTATTTGGAAATGGTAATACATTTGTCACGGATTTCTATAATGTATACA
 GCATTGGGAAGTGGGTGGGGGCTTTCTAATATGATACCGTCTTTTAAATAACTATGACAAAGCTTACATGAAGATTTAGAGACCACTTTACA
 TTTTACATTTCTCTGCTGTTTCATATTAACCTTGACCAATTAATTTCAATTTTCTCTTGACTCTTTTACCACAATGTTTTGGTTATTTATAAT
 TTATCAGCCATATGTTTATCAGCCATATAACCACTAGATCCCAATAGATCCATGTATTTGTTTCCGTGATTTGGCCACATTAATAAATTCA
 TAAATTTCAATCAAAATATCTTATATATACACATATGGTTTAAAGCTACAGCCCTGTGTATGCGCTTTAACTTTATTTGACGTTGCCCACTTA
 CTCTTTGCTGACCACTTGGATAACCGTAATAAAAAATCCTATAAGCCTAAATGGCATTTCTTTTGGGATATTTTCTCTGCATTTTATTCCTT
 TTTATATAAGTAGGAATTAATTAATTTATTTATGCTTAATCTATTTGATAAAGAGACTACATTAATAAATCTCAAGATCATATTAACAA
 AGGTTGCCCACTTGACCATATTTCAATTTGACACAGAAACAAATTTAGTACAACCTTCTAGTTCCCATGTCTTGATTTTCACTATTACA
 TGCACAGCAGACCTTTACCTATTGTGATACCAAGACACATCAETGTCTTTGGTTCCCTTCAAGAGAAATTTTATTTGTTGTTTGTATTTTCAA
 GTCTTAAATAGTTCTTGAACCTCTAGTTCTTTCTTGTGTAAGCAGACACATTTAGTGCAAGGCTTAATTTTACCTTTCCGGTGAAGAT
 CAGATGTTTTTATACCTTCACTTGGATCAATATATTTGGAAGAAATGTTTATCAAAAGTCTATGTCACTGCTTCTACAGAAGATGAATTA
 TGCTTAGTGATGGTACCTCCACCTACATCTTTTGGAGTCAATTAATATGATTTTGGTTTACGTTCTGATTTAACTTAATTTGATTCAG
 TTTAAACATGTTACTTAATTAGCAATGTAGAGGAACCAAAAAAGGTGAAAATATATGTTTTGATTCAACCTTAAGACATATAAAGCATAA
 AGACATTTTAACTTTGGGTTCTCTTAGCTGGGATCTGGCCAGAGGAGGCTTAAAGTTAGAAATGCTATTTATTTTAAATAGGTTGGGTGG
 GTTGGGGGCAAGGGTGTCTATTTGACAGCAGATATTTTGAAGAAAGAAATTTGTTTTATATAAAGGAAAGCCATGACCACTTTCTAC
 CTCAGATCCATCTTCATCCATTGCAATGGAAACTGCTTTATGCTGCTGCAAGTCTGAAGTCTAGAGCTTTTATCAGGCCATGTCTACCCAA
 GAAAGCACTATTTTAAAGAAAAACAATTCCTGAGCTCTCAACTCCAAGTTGTAGATTTGGTGTCTCTCTTCTTACTTTTAAAGTCAAT
 GTGTTAATTTTCTTCTGCTGATTTGTTATGCAAAATGTCCCTCTATCTGCTATTAAGAAAGCTACGTAAGAAACACTACATTTGTACCTCT
 AAGTAATAATAAATAAAGAAATATATTCAGTAACAATGGGAAGTAAATGTAGTTCTTTTGAATATGTGGTAAAGAACTAATCACAGA
 CIATCATCTAATCTGGTTACATATTTGATTTTTCATCTGAATAAAGTAATTTTAAACAAAAA

FIGURE 3

Natural antisense sequence (BF838561): SEQ ID NO: 3

AGGTTAGAGAATCGCTTGAACAGGGAGGAGGAGGTcGCAGCAAGCTGAGATCGGGCCACTGCACCTCCACCCTGGGTGAGAGAGTGGAGACTCC
ATCTCAAAAAACAAACAAACAAAAATTAAAAAAGAAAAACAGTGGCTCAAAATGCATAATGGCTTGCCTAGAGCATAACAGGGAAAAAGG
AAAATCAGATGCTTTTASAGACTGATACACTGAAACTATTGGAGCGTTCTGTAAATCAATAAATCCTAAACCTACCAATACAAATGGAAT
TATCATGCAGCCAAATCCAGTAAATATGGTAGCTGGAAGTAATGATAGTGATTGTAAAAACTATGGCTGGGCATGGTGGCTCATGCCGTGAAT
CCGAGC

Natural antisense sequence (BG208074): SEQ ID NO: 4

ctcaggTTCTCTGCCCTGCACATCtctTTTCATCTGGCTCCTTTACATGTCCAGATGAAAGAATTATCTCACCATTCCGGCTCAACAAGAAATTC
GTCCCAGACAGACAAAAAGTTTAAAAATATTTTATTAACATTAACCTGGACTTTTCGTAGAAGAAATAAAGCTGAGAACGTCATATTTTCAGC
AGCTTCAATACTTGTCTTCTGATTGAATCATAAATTTtATTACTTTTACTTGTCTTCTACAATGACAAAGTACCCAAAGCTAAATGTGAATCT
CTAGTAGGTCgGAAA

Natural antisense sequence (BF950643): SEQ ID NO: 5

cgctcgaagctcatcAAAAATATTgATTGAATGACAGTTAAAGAGCCTTTACTCTATTTTTAATTTTCCAAATAACATATATGTGTACATATTTTT
TAAAAATATGGAGAGATTCAGCGCTTTTGTCTTGTCTTACTTTGGTAGGTTAATGCCGTGATGTTTCATCTTTTCCAGTTCAACAGGCTGGGTGA
GTGGTATGGTACCATAAAAAAGGATGATCAGCTGCAAAATCAAGACCTTCAATCTCTACTTCCATTCCCTCTAACAAACAGCCATGGCATGTTGT
CATTTAATCTCGAGAGAGCTTCATTTCCACATCTTCAGAAAGACAGAACAAATGCTAGATGTACTTGCATGTATTCCAAATAGTGATTGGCTTTC
AGACCTTCTCTCTCTCCCATTTGAATAGACATTAGCCCTCATGAGTGCACCTCCgCACCTGTACTCTCCCATACACGATTCTGCATATAGCCAC
CAAATGTAACTTTCTAATATGCAAAACATAAAAAAABCTTCTCTGGTTTTTTTGAACACATTCTCTCAATTGAGCAAGGAATCATCCATCCA
CATTTGATCCCCAGTCCACCTTTCCCGCATCCTTTTCAATTTTTTGCCAAATGCCCCATATCCCAAACCTCCAAatgagcttgagggaagtgagggt
acgactcgaattcgtatcaggcatagtggtctgga

Natural antisense sequence (BF768753): SEQ ID NO: 6

ctcaacgctTTATACTCACATCCAATTATACTTCGCATCCAATTATACTTCATATACAAAGTATAGCTCTCCAGTTTGTGTAGTCCTCATC
ACTCCCTATCCTTTTTTATACGCAACCTCATCTCAGTTCCTTATGTTATGTTTCTGGCTTCTACAAATAGGCATTTGGATTTGGTGGCCCCCGGG
GTCTTGATCACAGCTTCTCAAATTAATATGGATACAGCCCACTGGTGATTCTGATTTAAATTCACGGGGTTGGGTGGAAGAAATTCAGTTT
TAACCAGCTTCCAGGTGAAACCAATGTTGCTGGGTCTATGGATCACACTTTGAATGACAAAGCTTCCAGGTAAGAAATATTATTTATTGAAAAACC
ACTGCTAGTCTACTTAAAGGTATGTGTTCGATCTATTAGTTACTACTGCTgAAAACTCAGAAGATATAGATTTATGTGACACTGGCAGGGAAGTA
TAAAGcgttgagggg

Natural antisense sequence (ENST00000431309): SEQ ID NO: 7

CCCCATCAATGGGAGAGGCTCTGTTATGGGAAGACTGTATTGAATTAAGAGTTAATATCAGGCTGAGCTCAGTGGCTCATACCTGTAAATCCC
AGCACTTTGGGAGGTTGATGCTATTACTATCTCTACTTTACTCAGGCACAGAGCAGCTAAGCCATTGTCCAGGTTCCACAGAGCTAGCAGGTG
GCCATGCTTACACCCACAGACCTGATACTGTCTGAACCATTCCTAAAGTATTTCTACAGAGAGTACTAATAAGATTTTAAATCCTGACAG
GATTTTTCTTATTATCTTTTCAAATATGTGTAAATTAATACCATGCCATCCTTGATATCTAACTCATTTTAGAGATAAAATGCTTACCCT
AAATACCATCATATAATCTATATTTCTATAAOCCTACAGCAATATTGCAGGTTAGAATCCAAAGTGTGCAGAAATATTTGTGAAGATCAGTGTA
GATAATACTGAAATGACAGTGTGAGGAAGAAGGTA

FIGURE 4

Sequence ID	Antisense Sequence Name	Sequence
SEQ ID NO:8	CUR-0636	rUrCrUrCrArCrUrCrUrCrUrCrArCrCrCrArGrGrGrUrGrGrArGrUrG
SEQ ID NO:9	CUR-0638	zArGrCrUrGrCrUrGrArArArUrArUrGrArCrGrUrUrCrUrCrArGrCrU
SEQ ID NO:10	CUR-0640	zArGrGrCrArUrUrArArCrCrUrArCrCrArArArGrUrArArArGrCrArA
SEQ ID NO:11	CUR-0642	rGrArArArGrUrUrArCrArUrUrUrGrGrUrGrGrCrUrArUrArUrGrCrA
SEQ ID NO:12	CUR-0644	rUrArCrArGrCrArArArCrUrGrGrArGrArGrCrUrUrArUrArCrUrUrU
SEQ ID NO:13	CUR-0646	rUrUrCrCrArUrUrUrGrUrArUrUrGrGrUrArGrGrUrUrUrArGrGrArU
SEQ ID NO:14	CUR-0648	zCrGrGrArArUrGrGrUrGrArGrArUrArArUrUrCrUrUrCrArUrCrU
SEQ ID NO:15	CUR-0650	rCrGrUrGrArArUrUrUrArArArUrCrArGrArArUrCrArCrArGrGrU
SEQ ID NO:16	CUR-0652	rGrUrArArCrUrArArUrArGrArUrCrGrArArCrArCrArUrArCrUrU
SEQ ID NO:17	CUR-0654	zArGrGrGrArArUrGrGrArArGrUrArGrArGrArUrGrArArGrGrUrC

FIGURE 5

Sequence ID	Antisense Sequence Name	Sequence
SEQ ID NO:18	CUR-1443	G*C*C*A*C*C*T*G*C*T*A*G*C*T*C*T*G*T*G*A
SEQ ID NO:19	CUR-1444	T*A*C*C*T*T*C*T*T*C*C*T*C*A*T*C*A*C*T
SEQ ID NO:20	CUR-1445	G*G*A*A*T*T*A*C*A*G*G*T*A*T*S*A*G*C*A*C
SEQ ID NO:21	CUR-1446	G*G*T*T*C*A*G*A*C*A*G*T*A*T*C*A*G*G*T*G*C
SEQ ID NO:22	CUR-1447	T*C*C*C*A*T*A*A*C*A*G*A*C*C*T*T*C*T*C*C*C

FIGURE 6

Sequence ID	Sense Sequence Name	Sequence
SEQ ID NO:23	CUR-0636	rCtUrCrCrArCrCpCrUfGrGrGrUrGrArGrArGrArGrUrGrAGA
SEQ ID NO:24	CUR-0638	rCrUrGrArGrArArCrGrUrCrArUrArUrUrCrArGrCrArGCT
SEQ ID NO:25	CUR-0640	rGrCrUrUrUrArCrUrUrUrGrGrUrArGrGrUrArArUrGrCCT
SEQ ID NO:26	CUR-0642	rCrArUrArUrArGrCrCrArCrCrArArArUrGrUrArArCrUrUTC
SEQ ID NO:27	CUR-0644	rArGrUrArUrArArGrCrUrCrUrCrCrArGrUrUrUrGrCrUrGTA
SEQ ID NO:29	CUR-0646	rCrCrUrArArArCrCrUrArCrCrArArUrArArArUrGrGAA
SEQ ID NO:29	CUR-0648	rArUrGrArArArGrArArUrUrArUrCrUrCrArCrCrArUrUrCCG
SEQ ID NO:30	CUR-0650	rCrUrGrGrUrGrArUrUrCrUrGrArUrUrUrArArUrUrCrACG
SEQ ID NO:31	CUR-0652	rGrGrUrArUrGrUrGrUrCrGrArUrCrUrArUrUrArGrUrUAC
SEQ ID NO:32	CUR-0654	rCrCrUrUrCrArArUrCrUrCrUrArCrUrUrCrCrArUrUrCrCCT